

Biotechnology for Biofuels

Enhancing oil production and harvest by combining the marine alga *Nannochloropsis oceanica* and the oleaginous fungus *Mortierella elongata* --Manuscript Draft--

Manuscript Number:	BBIO-D-18-00127R1											
Full Title:	Enhancing oil production and harvest by combining the marine alga <i>Nannochloropsis oceanica</i> and the oleaginous fungus <i>Mortierella elongata</i>											
Article Type:	Research											
Section/Category:	Development of algal feedstocks											
Funding Information:	<table> <tr> <td>Office of Science (DE-FG02-91ER20021)</td><td>Dr. Christoph Benning</td></tr> <tr> <td>Office of Science (DE-FC02-07ER64494)</td><td>Dr. Christoph Benning</td></tr> <tr> <td>Office of Science (DE-SC0018409)</td><td>Dr. Gregory Bonito</td></tr> <tr> <td>National Science Foundation (DEB 1737898)</td><td>Dr. Gregory Bonito</td></tr> <tr> <td>National Science Foundation (NSF DBI-1358474)</td><td>Dr. Christoph Benning</td></tr> </table>		Office of Science (DE-FG02-91ER20021)	Dr. Christoph Benning	Office of Science (DE-FC02-07ER64494)	Dr. Christoph Benning	Office of Science (DE-SC0018409)	Dr. Gregory Bonito	National Science Foundation (DEB 1737898)	Dr. Gregory Bonito	National Science Foundation (NSF DBI-1358474)	Dr. Christoph Benning
Office of Science (DE-FG02-91ER20021)	Dr. Christoph Benning											
Office of Science (DE-FC02-07ER64494)	Dr. Christoph Benning											
Office of Science (DE-SC0018409)	Dr. Gregory Bonito											
National Science Foundation (DEB 1737898)	Dr. Gregory Bonito											
National Science Foundation (NSF DBI-1358474)	Dr. Christoph Benning											
Abstract:	<p>Background: Although microalgal biofuels have potential advantages over conventional fossil fuels, high production costs limit their application in the market. We developed bio-flocculation and incubation methods for the marine alga <i>Nannochloropsis oceanica</i> CCMP1779 and the oleaginous fungus <i>Mortierella elongata</i> AG77 resulting in increased oil productivity.</p> <p>Results: Grown separately and then combining the cells the <i>M. elongata</i> mycelium could efficiently capture <i>N. oceanica</i> due to an intricate cellular interaction between the two species leading to bio-flocculation. Use of a high-salt culture medium induced accumulation of triacylglycerol (TAG) and enhanced the content of polyunsaturated fatty acids (PUFAs) including arachidonic acid (ARA) and docosahexaenoic acid (DHA) in <i>M. elongata</i>. To increase TAG productivity in the alga, we developed an effective reduced nitrogen supply regime based on ammonium in environmental photobioreactors (ePBRs). Under optimized conditions, <i>N. oceanica</i> produced high levels of TAG that could be indirectly monitored by following chlorophyll content. Combining <i>N. oceanica</i> and <i>M. elongata</i> to initiate bio-flocculation yielded high levels of TAG and total fatty acids, ~15% and 22% of total dry weight (DW), respectively, as well as high levels of PUFAs. Genetic engineering <i>N. oceanica</i> for higher TAG content in nutrient-replete medium was accomplished by overexpressing DGTT5, a gene encoding the type II acyl-CoA:diacylglycerol acyltransferase 5. Combined with bio-flocculation this approach led to increased production of TAG under nutrient replete conditions (~10% of DW) compared to the wild type (~6% of DW).</p> <p>Conclusions: The combined use of <i>M. elongata</i> and <i>N. oceanica</i> with available genomes and genetic engineering tools for both species opens up new avenues to improve bio-fuel productivity and allows the engineering of polyunsaturated fatty acids.</p> <p>Keywords: Microalgae, Filamentous fungi, Flocculation, Cell wall interaction, Biofuel, Nitrogen starvation, Polyunsaturated fatty acid, Triacylglycerol</p>											
Corresponding Author:	Christoph Benning Michigan State University UNITED STATES											
Corresponding Author Secondary Information:												
Corresponding Author's Institution:	Michigan State University											
Corresponding Author's Secondary Institution:												
First Author:	Zhi-Yan Du											
First Author Secondary Information:												

Order of Authors:	Zhi-Yan Du
	Jonathan Alvaro
	Brennan Hyden
	Krzysztof Zienkiewicz
	Nils Benning
	Agnieszka Zienkiewicz
	Gregory Bonito
	Christoph Benning
Order of Authors Secondary Information:	
Response to Reviewers:	The authors' response letter has been included as a supplementary file.

[Click here to view linked References](#)

Enhancing oil production and harvest by combining the marine alga *Nannochloropsis oceanica* and the oleaginous fungus *Mortierella elongata*

Zhi-Yan Du^{1,2}, Jonathan Alvaro², Brennan Hyden¹, Krzysztof Zienkiewicz^{2,3} Nils Benning², Agnieszka Zienkiewicz^{2,5} Gregory Bonito^{4,5} and Christoph Benning^{1,2,5,6*}

¹Department of Energy-Plant Research Laboratory, Michigan State University, East Lansing, MI 48824, USA; ²Department of Biochemistry and Molecular Biology, Michigan State University, East Lansing, Michigan 48824, USA; ³Department of Plant Biochemistry, Albrecht-von-Haller-Institute for Plant Sciences, Georg-August-University, 37073 Goettingen, Germany; ⁴Department of Plant, Soil and Microbial Sciences, Michigan State University, East Lansing, Michigan 48824, USA; ⁵Great Lakes Bioenergy Research Center, Michigan State University, East Lansing, Michigan 48824, USA; ⁶Department of Plant Biology, Michigan State University, East Lansing, Michigan 48824, USA

*Address correspondence to Christoph Benning

Tel: 1-517-355-1609; E-mail: benning@msu.edu

Key words: *Nannochloropsis*, *Mortierella*, biofuel, bio-flocculation, polyunsaturated fatty acid, triacylglycerol, photobioreactor

Abstract (350)

Background: Although microalgal biofuels have potential advantages over conventional fossil fuels, high production costs limit their application in the market. We developed bio-flocculation and incubation methods for the marine alga *Nannochloropsis oceanica* CCMP1779 and the oleaginous fungus *Mortierella elongata* AG77 resulting in increased oil productivity.

Results: Grown separately and then combining the cells the *M. elongata* mycelium could efficiently capture *N. oceanica* due to an intricate cellular interaction between the two species leading to bio-flocculation. Use of a high-salt culture medium induced accumulation of triacylglycerol (TAG) and enhanced the content of polyunsaturated fatty

acids (PUFAs) including arachidonic acid (ARA) and docosahexaenoic acid (DHA) in *M. elongata*. To increase TAG productivity in the alga, we developed an effective reduced nitrogen supply regime based on ammonium in environmental photobioreactors (ePBRs). Under optimized conditions, *N. oceanica* produced high levels of TAG that could be indirectly monitored by following chlorophyll content. Combining *N. oceanica* and *M. elongata* to initiate bio-flocculation yielded high levels of TAG and total fatty acids, ~15% and 22% of total dry weight (DW), respectively, as well as high levels of PUFAs. Genetic engineering *N. oceanica* for higher TAG content in nutrient-replete medium was accomplished by overexpressing *DGTT5*, a gene encoding the type II acyl-CoA:diacylglycerol acyltransferase 5. Combined with bio-flocculation this approach led to increased production of TAG under nutrient replete conditions (~10% of DW) compared to the wild type (~6% of DW).

Conclusions: The combined use of *M. elongata* and *N. oceanica* with available genomes and genetic engineering tools for both species opens up new avenues to improve bio-fuel productivity and allows the engineering of polyunsaturated fatty acids.

Keywords: Microalgae, Filamentous fungi, Flocculation, Cell wall interaction, Biofuel, Nitrogen starvation, Polyunsaturated fatty acid, Triacylglycerol

Introduction

Plant and algal oils are some of the most energy-dense naturally occurring compounds that can be used as feedstocks for biofuel products. Microalgae have been considered as promising sustainable feedstock for supplanting fossil fuels since the 1970s. Advantages of microalgae over other biofuel feedstocks include high oil yield, short generation times, low agricultural land requirements, reduced fresh water needs, and reduced greenhouse gas emissions during algal cultivation [1-3]. To improve lipid productivity of microalgae, many studies have been conducted in model microalgae such as *Chlamydomonas reinhardtii* [4, 5], and biotechnologically relevant species such as *Nannochloropsis* [6-9]. *Nannochloropsis* species have become a focus for lipid and biofuel research, because these marine microalgae grow fast in open ponds or photobioreactors, and can be grown in seawater with high yields of lipid - up to 60% of dry weight (DW) [10-12]. In addition, *Nannochloropsis* is enriched in high-value polyunsaturated fatty acids (PUFAs) such as

1
2
3
4 omega-3 eicosapentenoic acid (EPA), and it has a small and compact haploid genome (~30
5 Mbp). In recent years, the genomes of many species and strains of *Nannochloropsis*
6 including *N. gaditana* and *N. oceanica* have been sequenced [13-15] and genetic
7 engineering methods have been developed for gene disruption, i.e. CRISPR-Cas9 [8, 16,
8 17], and to generate overproduction systems for triacylglycerols (TAGs) and other target
9 molecules [18, 19].
10

11
12 In spite of these apparent advantages, the high cost of microalgal-based fuel
13 production prevents its application in the market [20-22]. The major barriers for the cost-
14 effective production of microalgal biofuels include: (1) high cost for harvesting microalgae;
15 (2) low oil content and suboptimal composition; (3) high cost of lipid extraction; and (4)
16 impasses in sustainable nutrient supply. Among these barriers harvesting microalgae is
17 particularly challenging because of the small cell size (typically 2-20 μm) and low density
18 (0.3-5 g/L) of microalgae, which can account for up to 50% of the total cost of biofuel
19 products [23-25]. Traditional harvesting methods include chemical flocculation using
20 multivalent cations such as metal salts and cationic polymers to neutralize the negative
21 charge on the surface of microalgal cell walls, filtration for relatively large algae (>70 μm),
22 sedimentation/floatation for species that either fall out of suspension or float without
23 sufficient mixing, thermal drying, and centrifugation, which has a high cost and energy
24 consumption [24-26].
25

26
27 Recently, bio-flocculation of microalgae with living materials such as bacteria and
28 fungi has gained interest because of their high efficiency and relative low energy
29 requirement [23, 27-29]. In addition, many bio-flocculants can be cultured with nutrients
30 from industrial wastes [29, 30]. The gram-positive bacterium *Solibacillus silvestris* can
31 efficiently flocculate *N. oceanica* DUT01 without the addition of chemical flocculants or
32 high-energy inputs. The bacterial flocculant exhibits no growth effect on the algal cells and
33 it can be reused for lower cost in harvesting [31]. In addition, some bio-flocculants such as
34 filamentous fungi *Aspergillus fumigatus* and *Mucor circinelloides* can accumulate 10-15%
35 lipid per DW which can increase the total lipid yield [30]. However, these filamentous
36 fungi are also human pathogens, making them unsuitable for use as bio-flocculants.
37

38
39 Here we screened oleaginous fungi for their ability to flocculate *N. oceanica*
40 CCMP1779, a marine alga with a sequenced genome, rapidly growing molecular
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

engineering tool kit, and the ability to produce high levels of TAG [9, 13, 18]. We discovered that *N. oceanica* could be efficiently flocculated by *M. elongata* AG77, a soil fungus that accumulates high levels of TAG and arachidonic acid (ARA), and for which a sequenced genome is available [32]. We provide detailed insights into the physical interaction between *N. oceanica* and *M. elongata* AG77 and test approaches for increasing TAG content in *N. oceanica* by optimizing growth conditions and genetic engineering approaches in combination with bio-flocculation to harvest algal cells.

Results

***N. oceanica* cells are captured by the *M. elongata* mycelium.**

Fungi were incubated in potato dextrose broth (PDB). Fungal mycelium (~3 times of algal biomass) was added to the *N. oceanica* culture containing log-phase cells in f/2 medium in shaker flasks. After 6-d co-cultivation with *M. elongata*, *N. oceanica* cells aggregated in dense green clumps along the mycelium of the fungus (Fig. 1A). The interaction of *N. oceanica* with filamentous fungi appeared specific to *M. elongata*, as it was not observed in co-culture with *Morchella americana* 3668S (Fig. 1). Differential interference contrast (DIC) light microscopy showed dense numbers of *N. oceanica* cells attached to the *M. elongata* mycelium (Fig. 1C); in comparison, mycelium of *M. americana* hardly captured any algal cells (Fig. 1D). Three *Mortierella* strains, *M. elongata* AG77, *M. elongata* NVP64, and *M. gamsii* GBAus22 were used to test flocculation efficiency for harvesting of *N. oceanica* with *M. americana* as a negative control. All three *Mortierella* isolates aggregated ~10% of algal cells after 2-h co-culture and up to ~15% after 12 h (Fig. 1E). After 6-d co-cultivation, *M. elongata* AG77 and NVP64 captured ~60% of algal cells *M. gamsii* GBAus 22 captured ~25%. The short period of co-cultivation with fungi did not appear to affect the morphology of the algal cells and did not significantly change their diameter (Fig. 1F).

Physical interaction between the cell walls of *N. oceanica* and *Mortierella* fungi.

Scanning electron microscopy (SEM) was performed to investigate the physical interaction between *N. oceanica* and *M. elongata* strains AG77 (Fig. 2A) and NVP64 (Fig. 2B). Low magnification images (Fig. 2, top panels) showed an aggregation of algal cells around the

1
2
3
4 fungal mycelium as seen in the light micrographs (Fig. 1C). Higher magnification images
5 displayed details of the physical interaction between the alga and fungi (Fig. 2, middle and
6 bottom panels). Similar to the cell wall structure of *N. gaditana* [33], *N. oceanica* has
7 extensions on the outer layer of the cell wall, which are attached to the rugged surface of
8 the fungal hyphae; irregular tube-like structures are formed between the algal and fungal
9 cell walls, which very likely contribute to anchoring the algal cells to the mycelium. The
10 *M. americana* strain 3668S, which has much thicker hyphae (10-20 μm in diameter) than
11 the *M. elongata* strains AG77 and NVP64 ($< 2 \mu\text{m}$), showed no obvious capture of *N.*
12 *oceanica* cells (Fig. 2C) or flocculation.

13 14 15 16 17 18 19 20 21 22 23 **Flocculation of *N. oceanica* with *Mortierella* fungi increases the yield of TAG and** 24 **PUFAs.**

25
26 *Mortierella* fungi are known to produce high amounts of TAG and PUFAs including ARA
27 [34, 35]. Indeed, numerous lipid droplets were observed in both *Mortierella* and *Morchella*
28 fungi tested for alga flocculation (Fig. 3A-D). In contrast, *N. oceanica* had fewer and
29 smaller lipid droplets when grown in nutrient-sufficient f/2 medium with or without fungi
30 (Fig. 3E-I) in shaker flasks. Lipids were extracted and separated by thin-layer
31 chromatography (TLC) and fatty acid methyl esters were quantified by gas
32 chromatography and flame ionization detection (GC-FID) to determine the lipid and fatty
33 acid composition. As shown in Table 1, *M. elongata* AG77 and *M. gamsii* GBAus22 had
34 much higher content of TAG, ARA, total PUFAs and total fatty acids but less EPA
35 compared to *N. oceanica*, which affects the final yield of these compounds in the alga-
36 fungus aggregate. *N. oceanica* TAG is mainly composed of saturated and monounsaturated
37 fatty acids such as C16:0 and C16:1 (Fig. 4A), whereas *Mortierella* fungi have more
38 PUFAs, especially ARA (Fig. 4B). *N. oceanica* has more EPA in total lipid than in TAG
39 (Fig. 4A), and the alga-fungus aggregate contains ~10% ARA and ~7% EPA of total lipid
40 (Fig. 4C).

41
42
43
44
45
46
47
48
49
50
51
52
53
54 Compared to regular PDB fungal growth medium, f/2 medium has a high salt
55 concentration and an elevated pH (=7.6) and lacks sugar [36]. Thus, *M. elongata* AG77
56 and *M. gamsii* GBAus22 were incubated in different media in shaker flasks to test the
57 impact on lipid metabolism of high pH (PDB medium, pH7.6), high pH and high salinity
58
59
60
61
62
63
64
65

(f/2+1% sugar), and high pH and high salinity with sugar starvation (f/2 medium). These adverse conditions generally increased the TAG and total lipid content of *M. elongata* AG77 and *M. gamsii* GBAus22, especially under high salinity condition (PDB pH7.6 compared to f/2+1% sugar) (Additional file 1: Table S1). Compared to *M. gamsii* GBAus22, *M. elongata* AG77 showed a significant increase in TAG and total lipid under high pH (PDB, from pH5.3 to 7.6), and a lower increase in total lipid, and slight decrease in TAG, upon sugar starvation (f/2+1% sugar compared to f/2) (Additional file 1: Table S1). These adverse conditions reduced the content of ARA and total PUFAs in *M. gamsii* GBAus22, while EPA increased upon high pH but decreased under high salinity and sugar starvation (Additional file 1: Table S1). In contrast, *M. elongata* AG77 had increased content of ARA and PUFAs in response to sugar starvation but these fatty acids decreased under high pH and high salinity conditions; EPA of *M. elongata* AG77 was decreased under all stress conditions compared to regular growth condition (Additional file 1: Table S1).

Increasing TAG content in *N. oceanica* cells using ammonium as the N source.

It has been reported that TAG is the major compound for transitory carbon storage in *N. oceanica* cells grown under light/dark cycles [37]. However, the TAG content was relatively low when cells were grown under regular conditions [13, 38]. Indeed, *N. oceanica* cells produced fewer and smaller lipid droplets than the fungi during incubation in f/2 medium visible in confocal micrographs (Fig. 3). To increase TAG yield in *N. oceanica*, two approaches were employed: nutrient deprivation and genetic engineering. N deprivation is one of the most efficient ways to promote TAG synthesis in microalgae [2, 7, 39]. Following 120-h N deprivation in shaker flasks, TAG accumulated in *N. oceanica* accounted for up to ~70% of the total lipid fraction (Additional file 2: Figure S1A), which is over 20% of DW (Additional file 2: Figure S1B). The content of TAG quickly increased following N deprivation and decreased following N resupply, indicating that *N. oceanica* cells are very sensitive to N supply (Additional file 2: Figure S1). Under laboratory conditions, N deprivation of algal cultures can be performed by centrifugation to pellet the algal cells, followed by washes and resuspension in N-depleted medium. However, this approach is not practical during scale up for industrial purposes.

Thus, we developed a limited N supply culturing method for large-volume cultures to

1
2
3
4 induce TAG accumulation largely without compromising growth and biomass yields. To
5
6 mimic natural cultivation conditions for *N. oceanica*, such as an open-pond system, we
7
8 used for these experiments environmental photobioreactors (ePBRs). We grew the algal
9
10 cells under varying light (0 to 2,000 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$) under long-day (14/10 h
11
12 light/dark) cycles (Additional file 3: Figure S2), and we sparged with 5% CO_2 at 0.37 L
13
14 min^{-1} for 2 min per h at 23°C primarily to adjust the pH in the reactor vessel as previously
15
16 described in [40]. Illumination in the ePBR is provided by a high power white LED light
17
18 on top of a conical culture vessel (total height of 27 cm) containing 330 mL of algal culture
19
20 (20 cm in depth), which was designed to simulate pond depths from 5 to 25 cm [40]. We
21
22 substituted several N sources in the f/2 medium for the incubation of *N. oceanica*
23
24 containing set amounts of either ammonium, nitrate, or urea. Compared to nitrate and urea,
25
26 *N. oceanica* grew faster in the f/2- NH_4Cl medium (Additional file 4: Figure S3A),
27
28 consistent with previous observations for *Nannochloropsis* sp. [41] and *Nannochloropsis*
29
30 *gaditana* [42]. The DW of *N. oceanica* cells per L was also higher in the f/2- NH_4Cl culture
31
32 after 7-d incubation in the ePBR (Additional file 4: Figure S3B). Intriguingly, the cells
33
34 grown in f/2- NH_4Cl medium turned from vivid green to yellow following 7-d incubation
35
36 once they reached stationary phase (Additional file 3: Figure S2A), indicative of
37
38 chlorophyll degradation in the algal cells. Lipid analysis by TLC (Fig. 5A) and GC-FID
39
40 (Fig. 5B) demonstrated TAG accumulation during the period of days 2 to 8 after the culture
41
42 reached stationary phase (incubation time S2 to S8), which was inversely correlated with
43
44 chlorophyll content, while cell density and dry weight remained at similar levels during
45
46 this period (Additional file 4: Figure S3C, D). Previously, to prevent carbon limitation,
47
48 NaHCO_3 was added to *N. oceanica* cultures in shaker flasks [43]. Here using the ePBRs,
49
50 we noticed that the addition of NaHCO_3 prevented acidification of the cultures, which were
51
52 sparged with 5% CO_2 (Additional file 5: Figure S4A). *N. oceanica* cells accumulated more
53
54 TAG upon acidification in the culture medium without NaHCO_3 supply, especially from
55
56 S6 to S8, compared to the NaHCO_3 culture (Additional file 5: Figure S4A, B).

57 58 59 **Increasing the yield of TAG and total fatty acid in alga-fungus aggregates.**

60
61 *N. oceanica* cells were inoculated to $\sim 1 \times 10^6 \text{ mL}^{-1}$ in f/2- NH_4Cl medium in the ePBR and
62
63 then grown to stationary phase. The cultures were further incubated for 8 days to allow the
64
65

1
2
3
4 algal cells accumulate high levels of TAG as described above. Following this prolonged
5 incubation in the ePBR, *N. oceanica* cells had a high TAG content, as shown by direct
6 quantification (Fig. 5B) and confocal microscopy (Fig. 6A). These high-oil algal cells were
7 then transferred from the ePBR to flasks and ~ 3 times biomass of fungal mycelium was
8 added. This led to the formation of algal fungal aggregates during 6 days of co-cultivation
9 in flask cultures. We did not attempt to co-cultivate algae and fungi in the ePBRs to avoid
10 fungal contamination of the ePBRs and aggregation disruption by the constant stir bar
11 motion. Following prolonged incubation in the ePBR, *N. oceanica* cells showed a high
12 TAG content, as determined by direct quantification (Fig. 5B) and confocal microscopy
13 (Fig. 6A). Following a subsequent co-cultivation of algal and fungal cells for 6 days, the
14 algal-fungal aggregates were then collected by mesh filtration for further microscopy and
15 lipid analysis. Both alga and fungus contained a high number of lipid droplets (Fig. 6B),
16 and were enriched in TAG (~15% of DW) and total fatty acid (~22% of DW) (Fig. 6C),
17 which is much higher than using f/2 medium grown algal cells that produced 6.2% TAG
18 and 16.9% total fatty acid of DW (Table I) with the co-cultivation of fungi. The overall oil
19 productivity was increased from ~0.25 g/L/d TAG and ~0.67 g/L/d total fatty acid, when
20 the algae were grown in f/2 medium in shaker flasks followed by co-cultivation with fungal
21 cells, to 0.59 g/L/d TAG and 0.86 g/L/d total fatty acid, when algae were grown in f/2-
22 NH₄Cl medium in the ePBR followed by co-cultivation with fungal cells. In addition, the
23 oil from these algal-fungal aggregates contained ~10% ARA in TAG and total fatty acid
24 (Fig. 6C).

25
26 Genetic engineering was also performed to increase TAG content in *N. oceanica* by
27 overexpressing *DGTT5* under the control of a strong elongation factor gene (*EF*) promoter
28 (Additional file 6: Figure S5). Two strains (*DGTT5ox3* and *ox6*) obtained from nuclear
29 transformation displayed a strong increase in the *DGTT5* expression compared to the wild
30 type and empty-vector control by quantitative RT-PCR (Additional file 7: Figure S6A).
31 The construct generated *DGTT5* fused to the cerulean fluorescent protein, and the presence
32 of the cerulean protein in *DGTT5ox3* and *ox6* was confirmed by confocal microscopy
33 (Additional file 7: Figure S6B). Subsequent BODIPY staining showed that the *DGTT5oxs*
34 lines had more lipid droplets compared to the wild type under regular growth condition
35 (Additional file 7: Figure S6C), which was further confirmed by quantitative lipid analysis
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

(Additional file 7: Figure S6C). Because the elongation factor promotor was inhibited by N starvation, the *DGTT*oxs cells were not incubated in f/2-NH₄Cl medium but grown in regular f/2 medium, and then log phase cells of *DGTT*5oxs were co-cultivated with ~ 3 times biomass of *M. elongata* AG77 and TAG yield of the algal-fungal aggregates were determined. The *DGTT*5oxs had increased TAG content (~10% of DW) compared to the wild type (~6% of DW) (Additional file 7: Figure S6E), which was less productive than the cells cultivated in a low N-medium (~15% of DW).

Fatty acid and TAG synthesis pathways in *M. elongata* AG77.

The genome of *N. oceanica* CCMP1779 has been sequenced and analyzed for the presence of metabolic pathway genes for PUFA and TAG biosynthesis [13], information used in the genetic engineering for increased EPA content [18]. For *Mortierella* fungi, nuclear transformation methods were established [44, 45], and the *M. elongata* AG77 genome has been sequenced and annotated [32], but lipid metabolic pathways have not yet been reconstructed. Thus, we applied the genome browser and BLAST tools from the JGI fungal genome portal MycoCosm to predict fatty acid, PUFA, and TAG synthesis pathways for *M. elongata* AG77. The fatty acid synthesis pathway (Fig. 7A) was predicted according to gene candidates (Additional file 8: Table S2) and previous reports on eukaryotic fatty acid pathways [46, 47]. *M. elongata* AG77 has a type-I fatty acid synthase with a similar domain organization as found in yeast (Fig. 7B) [46]. Nine elongases and twelve desaturases were identified within the *M. elongata* AG77 genome for PUFA synthesis, including a $\Delta 15$ fatty acid desaturase (FAD) for EPA synthesis (Fig. 7C, Additional file 8: Table S2). Three DGATs and one PDAT (phospholipid:diacylglycerol acyltransferase) were present in the *M. elongata* AG77 genome, which is similar to what was reported for *M. alpina* [47].

Discussion

Microalgal biofuel is a renewable and sustainable energy source that can be a supplement or alternative to fossil fuels [1, 2, 26]. Although microalgal biofuel has many competitive advantages, current high costs associated with algal biofuel production limit its commercialization [20-22]. To overcome the major challenges in algal biofuel production, including high costs of harvesting, lipid extraction, and nutrient supply, as well as low oil

content in algae, we developed a method to harvest the oleaginous marine alga *N. oceanica* through bio-flocculation with an oleaginous fungus *M. elongata* AG77. By optimizing the incubation conditions, we have increased TAG biofuel yields.

Bio-flocculation of *N. oceanica* by *Mortierella* fungi and physical interactions between the partners.

Physical interaction and symbioses between fungi and algae naturally occur, such as in lichens, where autotrophic algae co-exist with fungi and develop a mutualistic relationship [48]. In fact, about one-fifth of all known extant fungal species are symbiotically associated with photobionts such as algae and cyanobacteria in lichenized forms [49]. Taking advantage of known algal-fungal interactions, filamentous fungi have been tested for harvesting microalgae in recent years [23, 29]. However, although some of the tested fungal strains including *Aspergillus nomius* and *A. fumigatus* can very efficiently flocculate algae, they are not enriched in oil such that they dilute the algal oil in the resulting biomass [50, 51]. In addition, some fungal strains reported to flocculate algae such as *A. fumigatus* and *M. circinelloides* are animal pathogens and can threaten human health [28, 30].

In this study, we sought to use non-pathogenic oleaginous fungi to test their potential for bio-flocculation of microalgae. Amongst the screened fungi, *M. elongata* AG77 can be used to efficiently harvest *N. oceanica* cells (Fig. 1A). *M. elongata* AG77, which grows in soil and associates with plant roots [52], produces TAG and PUFAs including ARA and EPA (Table 1). *N. oceanica* cells were captured by the *M. elongata* AG77 mycelium. With the addition of ~3 fold algal biomass, *M. elongata* AG77 can flocculate ~65% of algal cells after 6-d co-cultivation in flask cultures and was the most efficient isolate tested (Fig. 1C, E). Other studies have shown high efficiency of algal flocculation by fungi [30, 50, 51]. However, the exact nature of the interaction between algae and fungi remains to be determined. Previous studies have suggested that the fungal cell wall has a positive charge that likely attracts and neutralizes the negatively charged algal cells [30, 50, 51], similar to chemical flocculation methods using metal salts and cationic polymers. Fungal cell walls contain chitin, which has a strong positive charge, and the algal flocculation efficiency was correlated with the chitin content (12-16% of DW) in the five fungal strains tested for harvesting *Nannochloropsis* sp. [50]. However, zeta potential measurements that represent

the surface charges do not necessarily correlate with flocculation efficiency [30, 50], suggesting that the flocculation of algae by fungi is not only dependent on the ionic attraction. In this study, we investigated details of the physical interaction between *N. oceanica* and fungi with SEM. Micrographs show *N. oceanica* cells trapped within the mesh-like mycelium of *M. elongata* AG77 and NVP64 (Fig. 2A, B). High magnification images (x 20,000 to x 250,000) show that *N. oceanica* has extensions along the outer layer of the cell wall, which appear to attach algal cells to the rugged surface of the fungal cell wall. Similar extension structures have also been observed in *N. gaditana* cells, which are connected to a thin algaenan layer of the *N. gaditana* cell wall, based on quick-freeze deep-etch EM and analyses of cell wall compositions [33]. Intriguingly, some fiber-like extensions from the *N. oceanica* cell walls seem to fuse with the surface of the fungal cell wall and form irregular-tube-like structures that appear to anchor the algal cells to the mycelium (Fig. 2A, B). However, the *Morchella* strain 3668S that cannot flocculate *N. oceanica*, is also not able to capture *N. oceanica* cells in its mycelium (Fig. 2C). Considering that both *Mortierella* and *Morchella* fungi have high chitin contents in their cell wall [53, 54], *M. elongata* AG77 and NVP64 and *M. americana* 3668S likely have positive surface charges. A major physical difference between the strains is that *M. americana* 3668S has much thicker hyphae than the *M. elongata* strains, suggesting that the size and structure of the mycelium may be important for the physical attraction of algal cells, but fungal exudates and chemotaxis could also underlie efficient bio-flocculation.

TAG and PUFAs are increased in *Mortierella* grown in f/2 medium.

The oleaginous fungi tested in this study produce a large number of lipid droplets, when incubated in regular PDB medium in flasks cultures as shown by confocal microscopy (Fig. 3A-D). To flocculate *N. oceanica* cells, fungal mycelium was added into the algal culture that contains high concentration of sea salt at pH 7.6 but lacks a sugar supply. The combined stress conditions presented by f/2 medium elevated the TAG and PUFA content in *M. elongata* AG77 (Additional file 1: Table S1) and decreased its growth, such that it would not outgrow the algal cells. In fact, *N. oceanica* cells had normal diameter (Fig. 1F) and low amount of oil when co-cultivated with the fungi (Fig. 3F-I), similar to the cells grown alone grown in f/2 medium (Fig. 3E).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

M. elongata AG77 is a representative strain of *Mortierella*, a widespread genus of soil fungi, which are industrially important for their production of ARA-rich oil [34, 35]. *Mortierella* fungi grow fast in nutrient-rich media and studies have shown that *M. elongata* CBS 121.71 and *M. alpina* SD003 can also grow in wastewater [55, 56], while a related species *Umbelopsis isabellina* can grow on sewage sludge [57], suggesting low-cost incubation methods of growing *M. elongata* AG77 for the bio-flocculation of algae.

Increasing TAG in *N. oceanica* cells through genetic engineering and by optimizing incubation conditions.

In recent years, intensive studies have been performed on *Nannochloropsis* because of its potential for biofuel production [7, 10-12]. Besides its high TAG content, *Nannochloropsis* is enriched in PUFAs such as EPA, a high-value omega-3 fatty acid [18]. However, *Nannochloropsis* species, such as *N. oceanica*, do not have high oil content under favorable growth conditions (Fig. 3E; Additional file 2: Figure S1) [13, 37]. Thus, although the *N. oceanica*- *M. elongata* AG77 co-culture generated a favorable yield of ARA and EPA, TAG production was only ~6% of DW- lower than the ~10% TAG *M. elongata* AG77 alone (Table 1, Fig. 4). *Nannochloropsis* accumulates high levels of TAG and total fatty acids in response to stress conditions such as N deprivation (Additional file 2: Figure S1) and following the addition of the ribosome inhibitor cycloheximide [8]. Similarly, TAG and total lipid stimulation by environmental stresses have been commonly observed in the other microalgae such as green algae and diatoms [2, 7]. TAG is usually used as the feedstock for biodiesel and can be converted into high-energy fatty acid methyl esters (FAMES) by transesterification with glycerol as a valuable side-product [25, 58]. In addition, TAG is a non-polar lipid stored in lipid droplets and is more readily extractable with organic solvent than the polar membrane lipids [59]. Thus, increasing TAG content can simplify the lipid extraction and transesterification, and therefore reduce the cost of processing. To increase TAG content in *N. oceanica*, we developed an N starvation method for *N. oceanica* using ammonium as the N source. We tested the growth of *N. oceanica* with three N sources - ammonium, nitrate, and urea - in ePBRs that can mimic natural growth conditions such as open-pond system (Additional file 3: Figure S2) [40]. *N. oceanica* cells grow significantly faster and have higher biomass in the ammonium medium

1
2
3
4 than the nitrate and urea media, following the same period of incubation (Additional file 4:
5 Figure S3). In addition, *N. oceanica* cells consumed ammonium faster than nitrate and
6 approached N starvation during the week after the culture reached stationary phase (Fig. 5;
7 Additional file 3: Figure S2A), consistent with the previous observation that
8 *Nannochloropsis* sp. grow faster in ammonium medium with a significantly higher uptake
9 rate of ammonium than nitrate [41]. N deprivation is one of the most efficient ways to
10 stimulate TAG accumulation in microalgae. However, it is not feasible to centrifuge the
11 cells and change the medium at industrial scales, as one can do in the laboratory.
12 Considering that *N. oceanica* cells are very sensitive to the N supply (Additional file 2:
13 Figure S1), feeding ammonium to the culture with limited amount of N, or in multiple-low
14 doses, can allow rapid cell growth and provides a means to control the abundance of N in
15 the culture. The N supply can be stopped as the culture reaches the stationary phase such
16 that the cells exhaust their N supply and enter N starvation, while the total cell count and
17 biomass are not compromised (Additional file 4: Figure S3C, D).

18
19 We have developed a quick and simple method to indirectly monitor TAG content
20 for *N. oceanica* by using chlorophyll as an indicator. Chlorophyll content is inversely
21 correlated with TAG content during nutrient deprivation (Fig. 5B). In fact, thylakoid
22 membrane degradation, in parallel with TAG accumulation, has been commonly observed
23 in microalgae and plant vegetative tissues. These combined processes can reduce
24 photosynthesis and photooxidative stress, sequester toxic compounds such as free fatty
25 acids, and save the acyl chains for latter resynthesis of membranes when conditions
26 improve [7, 39, 60].

27
28 We also noticed that culture acidification caused by CO₂ supply led to faster and
29 higher TAG accumulation in *N. oceanica* cells (Additional file 5: Figure S4). The described
30 N starvation method should be applicable for most microalgal species, especially for the
31 wild-type and transgenic strains optimized for growth rate, biomass, and stress resistance,
32 but which have low oil content under regular growth condition. Under the right conditions,
33 TAG accumulation occurs once the culture reaches stationary phase/high biomass. The N-
34 starvation method we describe will not reduce cell growth or biomass, which has been
35 commonly observed for genetically engineered high TAG strains [8, 9, 19]. On the contrary,
36 the higher the biomass and cell concentration, the faster the algal cells deplete their N

source and generated more TAG. Taking advantage of the high TAG *N. oceanica* cells produced by the N-starvation method (Fig. 6B), we obtained high yields of TAG (~15% of DW) and total fatty acids (~22% of DW) in the *N. oceanica*-AG77 aggregate (Fig. 6B, C).

Potential of genetic engineering of PUFAs in *N. oceanica* and *M. elongata* AG77.

Both *N. oceanica* CCMP1779 and *M. elongata* AG77 generate ARA and EPA: ARA is low in *N. oceanica* but high in AG77, whereas EPA is high in *N. oceanica* but low in AG77 (Fig. 4A, B). Besides ARA and EPA, *M. elongata* AG77 also produces ~5% DHA (Fig. 4B). Thus, the complimentary production of ARA, EPA, and DHA, high-value products with important health benefits [61], could increase the value of the oil produced by aggregates of *N. oceanica* and *M. elongata* AG77. In addition, the genome of *N. oceanica* CCMP1779 and its BLAST and gene browser tools are available [13], and genetic engineering methods such as CRISPR-Cas9 and overexpression systems have been established for *N. oceanica* [9, 16, 18]. Recent studies have successfully increased the EPA production by the overexpression of $\Delta 5$ and $\Delta 12$ FADs in *N. oceanica* [18] and induced TAG production under normal growth condition by overexpressing *N. oceanica* *DGTT5* [9].

Transformation methods for *Mortierella* fungi such as *M. alpina* have been developed [45, 62], and the genome of *M. elongata* AG77 is available [32]. In this study, we analyzed the genome data and predicted the PUFA and TAG synthesis pathways (Fig. 7), based on the gene candidates involved in the processes (Additional file 8: Table S2), which will facilitate genetic engineering on PUFA and TAG products in *M. elongata* AG77 in the future.

Conclusions

N. oceanica CCMP1779 and *M. elongata* AG77 are model species of algae and fungi, respectively. Both genera contain many other oleaginous strains (e.g. *N. gaditana* and *M. alpina*). The bio-flocculation method demonstrated in this study could be performed with other strains leading to synergistic increases in TAG content of the biomass. The N starvation and seawater medium can also be used on transgenic strains with high TAG or high biomass traits, which has the potential to increase oil yield and reduce the cost of oil

1
2
3
4 extraction and transesterification. *N. oceanica* grows in seawater and *M. elongata* AG77
5 can be incubated with wastewater and sewage that most likely can reduce the cost of
6 nutrient supply. Overall, the combination of *N. oceanica* and *M. elongata* AG77 has the
7 potential to overcome some of the barriers for the commercialization of microalgal/fungal
8 biofuels and biomaterials.
9
10
11
12
13
14

15 **Methods**

16 **Materials and growth condition**

17
18 The marine alga *Nannochloropsis oceanica* CCMP1779 was obtained from the Provasoli-
19 Guillard National Center for Culture of Marine Phytoplankton. *N. oceanica* *DGTT5*-
20 overexpressing strains *DGTT5ox3* and *DGTT5ox6* were generated and examined by
21 quantitative RT-PCR as previously described [9]. The alga was grown in f/2 medium
22 containing 2.5 mM NaNO₃, 0.036 mM NaH₂PO₄, 0.106 mM Na₂SiO₃, 0.012 mM FeCl₃,
23 0.012 mM Na₂EDTA, 0.039 μM CuSO₄, 0.026 μM Na₂MoO₄, 0.077 μM ZnSO₄, 0.042
24 μM CoCl₂, 0.91 μM MnCl₂, 0.3 μM thiamine HCl/vitamin B1, 2.05 nM biotin, 0.37 nM
25 cyanocobalamin/vitamin B12 [63] and 20 mM sodium bicarbonate and 15 mM Tris buffer
26 (pH 7.6) to prevent carbon limitation [43]. The cells were grown in batch cultures in two
27 systems: shaker flask with f/2 medium (under ~80 μmole photons m⁻² s⁻¹ at 23°C) or in
28 ePBRs (Lucker et al., 2014) with f/2-NH₄Cl (2.5 mM NH₄Cl replacing 2.5 mM NaNO₃) or
29 f/2-urea (2.5 mM urea replacing 2.5 mM NaNO₃) medium with varying light as indicated
30 (Additional file 3: Figure S2, 0 to 2,000 μmol photons m⁻² s⁻¹ under diurnal 14/10 h
31 light/dark cycle) at 23°C and sparged with air enriched to 5% CO₂ at 0.37 L min⁻¹ for 2
32 min per h. For prolonged-incubation in the ePBR, *N. oceanica* cells were inoculated to ~1
33 x 10⁶ mL⁻¹ in f/2-NH₄Cl medium and grown to stationary phase. The cultures were further
34 incubated for 8 days to increase TAG content.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50 *Mortierella* fungi *M. elongata* AG77, *M. elongata* NVP64, and *M. gamsii* GBAus22
51 isolates were isolated from soil samples collected in North Carolina (AG77), Michigan
52 (NVP64), USA, and Australia (GBAus22). *Morchella americana* 3668S was obtained
53 from the USDA NRRL Agriculture Research Station. Fungal samples were incubated in
54 PDB medium (12 g/L potato dextrose broth and 1 g/L yeast extract, pH5.3) at 23°C. For
55 the algal-fungal co-cultivation, fungal mycelia were briefly blended into small pieces (~1
56
57
58
59
60
61
62
63
64
65

cm) with a sterilized blender and were collected by centrifugation (3,000 g for 3 min) after 24-h recovery in PDB medium. The samples were washed twice with f/2 or f/2- NH₄Cl medium and resuspended in 5-10 mL of the respective medium. One third of the samples were used for determining dry biomass: 1 mL culture was transferred and filtered with pre-dried and -weighed Whatman GF/C filters and dried overnight at 80°C. The remaining fungal mycelia were added to the *N. oceanica* culture (~3 times to algal biomass) for 6-d co-cultivation on a shaker (~60 rpm) under continuous light (~80 μmol photons m⁻² s⁻¹) at 23°C. Cell size and concentration of *N. oceanica* cultures were calculated with a Z2 Coulter Counter (Beckman). The bio-flocculation efficiency of *N. oceanica* cells using fungal mycelium was determined by the cell density of uncaptured algal cells compared to that of an algal culture control, to which no fungus was added.

Light Microscopy

Interactions between the algal and fungal cells were examined by light microscopy using an inverted microscope with DIC function (DMi8, Leica). DIC images were taken of the algae-fungi aggregates after 6 d co-cultivation.

Scanning electron microscopy

SEM was performed to investigate the physical interaction between *N. oceanica* and fungi at the Center for Advanced Microscopy of Michigan State University (CAM, MSU). Algae-fungi aggregates were collected after 6-d co-culture of the alga *N. oceanica* with *M. elongata* (AG77 and NVP64) or *M. americana* 3668S and were fixed in 4% (v/v) glutaraldehyde solution, followed by drying in a critical point dryer (Model 010, Balzers Union). The samples were then mounted on aluminum stubs with high vacuum carbon tabs (SPI Supplies), and were coated with osmium using a NEOC-AT osmium coater (Meiwafosis). The samples were observed with a JSM-7500F scanning electron microscope (Japan Electron Optics Laboratories).

Confocal microscopy

Confocal microscopy was carried out to visualize and briefly quantify lipid droplets in the alga and fungi. The samples were stained with 10 μg mL⁻¹ BODIPY 493/503

(ThermoFisher Scientific) in PBS buffer for ~30 min at 23°C. After two washes with PBS buffer, the samples were observed using an Olympus Spectral FV1000 microscope at CAM, MSU. An argon (488 nm) laser and a solid-state laser (556 nm) were used for BODIPY (emission, 510 to 530 nm) and chloroplast (emission, 655 to 755 nm) fluorescence. *N. oceanica* DGTT5 fused to the cerulean fluorescent protein was overproduced using the EF promotor as previously published [9]. The presence of the fluorescent protein in the *DGTT5ox* strains was detected by confocal microscopy (emission 420-440 nm) using a LSM 510 Meta Confocal Laser Scanning Microscope (Zeiss).

Lipid extraction and analysis

For lipid extraction, log phase *N. oceanica* cells grown in f/2 medium were collected by centrifugation (4,000 g for 5 min). To test lipid content in different media, *Mortierella* fungi grown in PDB medium were washed twice with different media: PDB medium, pH7.6; f/2 medium with 1% glucose; f/2 medium. The cells were incubated in the respective medium for 48 h and were subsequently collected for lipid extraction by centrifugation (3,000 g for 3 min). For total lipid extraction, algae-fungi aggregates were collected by mesh filtration and frozen in liquid nitrogen prior to grinding with mortar and pestle. The fine powders were transferred to a pre-weighed and -frozen glass tube and total lipids were extracted with methanol-chloroform-88% formic acid (1:2:0.1 by volume) on a multi-tube vortexer (1,500 g for ~20 min; Benchmark Scientific), followed by addition of 0.5 volume of 1 M KCl and 0.2 M H₃PO₄. After phase separation by centrifugation (2,000 g for 3 min), total lipids were collected for TAG separation and fatty acid analysis. The solids were dried at 80°C over night to provide the non-lipid biomass.

TAG was separated by TLC using G60 silica gel TLC plates (Machery-Nagel) developed with petroleum ether-diethyl ether-acetic acid (80:20:1 by volume). An internal standard of 5 µg of tridecanoic acid (C13:0) or pentadecanoic acid (C15:0) was added to each tube containing TAG or total lipid. FAMES were then prepared with 1 M methanolic HCl at 80°C for 25 min, and were phase separated with hexane and 0.9% NaCl and nitrogen-dried and resuspended in ~50 µL of hexane. Gas chromatography and flame ionization detection (Agilent) were used to quantify the FAMES in TAG and total lipid as

previously described [64]. Dry weight of algae-fungi biomass was obtained by summing up non-lipid and total lipid mass.

Chlorophyll measurement

N. oceanica cells were collected by centrifugation from 1 mL culture aliquots during prolonged-incubation in the ePBRs. Chlorophyll of the pelleted cells was extracted with 900 µL of acetone:DMSO (3:2, v/v) for 20 min with agitation at 23°C, and measured with an Uvikon 930 spectrophotometer (Kontron) [65].

Generation of *N. oceanica* DGTT5 overexpression strains

The NoDGTT5 sequence was amplified by PCR using the primers: forward, 5'-ATGACGCCGCAAGCCGACATCACCAGCAAGACGA-3'; reverse, 5'-CTCAATGGACAACGGGCGCGTCTCCCACTCC-3', followed by gel purification with E.Z.N.A. Gel Extraction Kit (OMEGA Biotek) and ligated into the the pnoC ox cerulean hyg vector (Additional file 6: Figure S5A) to obtain the final vector pnoC ox DGTT5 cerulean hyg (Additional file 6: Figure S5B) for nuclear transformation. *N. oceanica* CCMP1779 cells were transformed by electroporation as previously described [13].

Prediction of fatty acid and TAG pathways

The sequenced genome of *M. elongata* AG77 [32] was annotated for genes and proteins likely involved in the synthesis of fatty acids, PUFAs, and TAGs using by BLAST searches against KOG and KEGG databases at the JGI fungal genome portal MycoCosm *M. elongata* AG77 v2.0 and by comparison to previously published annotations of lipid pathways of *Mortierella alpina* [47].

Accession numbers

Sequence data presented in this article can be found in the genome of *N. oceanica* CCMP1779 at the JGI database (<https://genome.jgi.doe.gov/Nanoce1779/Nanoce1779.home.html>). Gene ID of *N. oceanica* DGTT5, CCMP1779_3915; accession number, KY273672. Sequence data of *M. elongata* AG77 can be found at the JGI fungal genome portal MycoCosm

(<https://genome.jgi.doe.gov/Morel2/Morel2.home.html>). Transcript IDs and protein IDs of the genes are listed in the Additional file 8: Table S2.

Additional files

Additional file 1: Table S1. Lipid and fatty acid contents of *Mortierella* fungi incubated in different media (mg g⁻¹ total dry weight).

Additional file 2: Figure S1. Triacylglycerol content in *N. oceanica* cells.

Additional file 3: Figure S2. Incubation of *N. oceanica* cells in the environmental photobioreactor (ePBR).

Additional file 4: Figure S3. Cell growth and biomass in the environmental photobioreactor (ePBR).

Additional file 5: Figure S4. Triacylglycerol accumulation during prolonged-incubation in f/2 medium supplemented with or without sodium bicarbonate.

Additional file 6: Figure S5. Maps of the plasmids used for the generation of *N. oceanica* *DGTT5*-overexpressing strains.

Additional file 7: Figure S6. Increasing triacylglycerol content in *N. oceanica* by the overexpression of *N. oceanica* *DGTT5* encoding acyl-CoA:diacylglycerol acyltransferase *DGTT5*.

Additional file 8: Table S2. Predicted genes and proteins involved in fatty acid and glycerolipid synthesis in *M. elongata* AG77.

Abbreviations

ARA: arachidonic acid; *DGTT5*: a gene encoding the type II acyl-CoA:diacylglycerol acyltransferase 5; DHA: docosahexaenoic acid; DW: dry weight; *EF*: elongation factor gene; EPA: eicosapentenoic acid; ePBR: environmental photobioreactor; FAMES: fatty acid methyl esters; GC-FID: gas chromatography and flame ionization detection; PDAT: phospholipid:diacylglycerol acyltransferase; PDB: potato dextrose broth; PUFAs: polyunsaturated fatty acids; S2 to S8: days 2 to 8 after the culture reached stationary phase; SEM: scanning electron microscopy; TAG: triacylglycerol; TLC: thin layer chromatography.

Authors' contributions

ZD, GB, and CB designed the experiments. ZD performed the experiments, analyzed the data, and wrote the first draft of the article; JA contributed to the initial tests of bio-flocculation; BH contributed to growth and lipid experiments using ePBR; KZ generated the DGTT5 strains; NB contributed to the lipid analysis and biomass determination; AZ contributed to the data analysis of lipid-related genes and proteins and prediction of lipid pathways. ZD, GB, and CB coordinated the writing and edited the drafts of the article. All authors read and approved the final manuscript.

Author details

¹Department of Energy-Plant Research Laboratory, Michigan State University, East Lansing, MI 48824, USA; ²Department of Biochemistry and Molecular Biology, Michigan State University, East Lansing, Michigan 48824, USA; ³Department of Plant Biochemistry, Albrecht-von-Haller-Institute for Plant Sciences, Georg-August-University, 37073 Goettingen, Germany; ⁴Department of Plant, Soil and Microbial Sciences, Michigan State University, East Lansing, Michigan 48824, USA; ⁵Great Lakes Bioenergy Research Center, Michigan State University, East Lansing, Michigan 48824, USA; ⁶Department of Plant Biology, Michigan State University, East Lansing, Michigan 48824, USA

Acknowledgements

The authors thank Ben Luckner and David Kramer (MSU) for providing the ePBRs, Melinda Frame and Carol Flegler (CAM, MSU) for the assistance with confocal and scanning electron microscopy, Eric Poliner (MSU) for providing the pnox ox Cerulean hyg vector, Natalie Vande Pol (MSU) and Kerry O'Donnell (NRRL) for *M. americana* 3668S.

Competing interests

The authors declare that they have no conflict of interests.

Availability of data and materials

All data generated or analyzed during this study are included in this published article and its supplementary information files. All materials are available from the corresponding author.

Ethical approval and consent to participate

All authors consented to the publication of this work.

Funding

This work was supported in part by a grant from 507 the Chemical Sciences, Geosciences, and Biosciences Division, OZce of Basic Energy Sciences, OZce of Science, U.S. Department of Energy (DE-FG02-91ER20021) to C.B. This work was also funded in part by the Department of Energy Great Lakes Bioenergy Research Center (DOE Office of Science BER DE-FC02-07ER64494) grant to C.B. and BER DE-SC0018409 to G.B. G.B. also acknowledges the National Science Foundation DEB 1737898 for financial support. G.B. and C.B. acknowledge support from AgBioResearch. K.Z. received funding from the People Programme (Marie Curie Actions) of the European Union's Seventh Framework Programme FP7/2007-2013/ under REA grant agreement no [627266] supporting K.Z. The EU is not liable for any use that may be made of the information contained therein. J.A. and B.H were participants in the Plant Genomics @ MSU Research Experience for Undergraduates Program funded by NSF DBI-1358474.

Table 1. Lipid contents of different strains grown in f/2 medium in shaker flasks (mg g⁻¹ total dry weight).

TAG, triacylglycerol; ARA, arachidonic acid (20:4); EPA, eicosapentaenoic acid (20:5); PUFAs, polyunsaturated fatty acids (e.g. 18:2, 18:3, 20:4 and 20:5). *M. elongata* AG77/*M. gamsii* GBAus22 & *N. oceanica*, co-cultivation of the alga and fungi in f/2 medium for 6 d in shaker flasks. Results are the average of three biological replicates with standard deviations.

Strains	Total fatty acid	TAG	ARA	EPA	Total PUFAs
<i>N. oceanica</i>	118.7 ± 18.4	15.1 ± 2.3	3.1 ± 0.5	17.0 ± 2.6	21.5 ± 3.3
<i>M. elongata</i> AG77	238.8 ± 14.5	94.6 ± 4.5	42.4 ± 2.3	4.3 ± 0.5	89.1 ± 4.8
<i>M. gamsii</i> GBAus 22	178.0 ± 23.9	54.9 ± 3.9	29.3 ± 2.1	1.7 ± 0.5	66.1 ± 2.2
<i>M. elongata</i> AG77 & <i>N. oceanica</i>	168.5 ± 8.9	62.1 ± 3.0	16.3 ± 1.1	12.0 ± 0.9	46.5 ± 3.7
<i>M. gamsii</i> GBAus22 & <i>N. oceanica</i>	163.3 ± 10.5	42.0 ± 9.5	17.5 ± 1.7	9.0 ± 1.4	36.1 ± 6.1

FIG LEGENDS

Figure 1. Harvesting *Nannochloropsis oceanica* by bio-flocculation with *Mortierella* fungi.

A and B, Co-culture of *N. oceanica* (*Noc*) with *M. elongata* AG77 A, or *Morchella americana* 3668S B. Fungal mycelium collected from PDB medium was added to a log-phase *Noc* culture grown in flasks and both were incubated for 6 d in flasks containing f/2 medium. The red arrow indicates green aggregates formed by AG77 mycelium and attached *Noc* cells. C, *Noc* cells attached to AG77 mycelium as shown by differential interference contrast (DIC) microscopy. D, No obvious attachment of *Noc* cells on the 3668S mycelium. E, Bio-flocculation efficiency for harvesting *Noc* cells by cocultivation with *Mortierella elongata* AG77, *M. elongata* NVP64, and *M. gamsii* GBAus22 in f/2 medium. The bio-flocculation efficiency was determined by the cell density of uncaptured cells compared to that of a no-fungus *Noc* culture control. *Morchella* 3668S culture was used as a negative control. The results are the average of five biological replicates and error bars indicate standard deviation. Asterisks indicate significant differences relative to the 2h co-cultures by paired-sample Student's t-test (* $P \leq 0.05$; ** $P \leq 0.01$). F, Measurement of *Noc* cell size (diameter) in the *Noc* culture and alga-fungus co-culture.

Figure 2. Interaction between *N. oceanica* and *Mortierella* mycelium co-cultivated in f/2 medium. Fungal mycelium was added to log-phase *Noc* culture grown in flasks and then the mixture was incubated for 6 d in f/2 medium. Scanning electron microscopy was performed to investigate the interaction between *N. oceanica* (*Noc*) cells and *M. elongata* AG77 (A) and NVP64 (B). *Noc* cells are attached to the fungal mycelium as shown in the top panel. Higher magnification micrographs show that *Noc* cells have a highly structured cell wall with protrusions, with which they attach to the rough surface of the fungal cell wall. The red arrowheads indicate tube-like structures that connect the algal and fungal cell walls. (c) *Morchella americana* 3668S mycelium collected from *Noc*-3668S culture after 6-day co-cultivation.

Figure 3. *Mortierella* fungi have more oil droplets than *N. oceanica* in f/2 medium. A to D, Confocal micrographs showing lipid droplets in the fungal mycelium grown in flasks containing PDB medium. Green fluorescence indicates lipid droplets stained with

BODIPY. E, Lipid droplets in log-phase *N. oceanica* (*Noc*) cells grown in flasks containing f/2 medium. Red, autofluorescence of *Noc* chloroplast. F to I, Lipid droplets in the algal and fungal cells after 6-d co-cultivation in f/2 medium. Fungal mycelium was added to the log-phase *Noc* culture for 6-d co-culture in flasks. AG77, *M. elongata* AG77; NVP64, *M. elongata* NVP64; GBAus22, *M. gamsii* GBAus22; 3668S, *Morchella americana* 3668S.

Figure 4. Fatty acid profiling of triacylglycerol (TAG) and total lipid in *Mortierella* fungi, *N. oceanica*, and algae-fungi aggregates after 6-d co-cultivation. A, Fatty acid assays of triacylglycerol and total lipid in log-phase *N. oceanica* grown in shaker flask containing f/2 medium. Fatty acids are indicated with number of carbons : number of double bonds. Results are the average of five biological replicates with error bars indicating standard deviations (n = 5). B, Fatty acid assays of *M. elongata* AG77 incubated in f/2 medium. Fungal mycelium was collected from PDB medium and then washed and added to f/2 medium and incubated for 6 d. n = 5. C, Fatty acids of the algae-fungi aggregates after 6-d co-cultivation in flasks containing f/2 medium. n = 5.

Figure 5. Chlorophyll as proxy of triacylglycerol accumulation. A, Analysis of triacylglycerol (TAG) by thin layer chromatography (TLC). Algal cells were incubated in environmental photobioreactor (ePBR) containing f/2-NH₄Cl medium after inoculation to stationary phase and then prolonged incubation. Red arrowheads indicate the TAG bands. S1 to S8, day 1 to 8 after the cells reached stationary phase; control, TAG standard. B, Inverse correlation of chlorophyll content and TAG-to-total-lipid ratio following prolonged incubation in the ePBR containing f/2-NH₄Cl medium. TAG and total lipid were subjected to transesterification reaction and the resulting fatty acid methyl esters were quantified by GC-FID. r^2 , correlation coefficient; n = 4.

Figure 6. Increasing triacylglycerol (TAG) content in *N. oceanica* using limited ammonium as nitrogen source. A, *N. oceanica* (*Noc*) cells produce large lipid droplets during prolonged incubation in the environmental photobioreactor (ePBR) containing f/2-NH₄Cl medium. *Noc* cells grow fast in f/2-NH₄Cl medium and suffer from nutrient limitation after being for 8 d in the stationary phase, when the confocal micrographs were

taken. Green fluorescence indicates lipid droplets stained with BODIPY, while red fluorescence represents autofluorescence of *Noc* chloroplasts. B, Lipid droplet staining of *M. elongata* AG77 and *Noc* cells after 6-d co-cultivation in shaker flasks following the incubation of the algae in ePBRs. C, Fatty acid (FA) analyses of triacylglycerol and total lipid in the alga-fungus aggregate as shown in (B). Biomass ratio of TAG or total FA relative to the total cell dry weight (DW). n = 5.

Figure 7. Predicted fatty acid/lipid pathways in *M. elongata* AG77. Proteins likely involved in the synthesis of fatty acids (FA), polyunsaturated fatty acids (PUFA), and triacylglycerol (TAG) are identified in the sequenced genome of *M. elongata* AG77 at the JGI fungal genome portal MycoCosm (Additional file 8: Table S2). A, FA synthesis. ACP, acyl carrier protein; AT, acetyltransferase; MPT, malonyl/palmitoyl transferase; ACSL, acyl-CoA synthetase; KS, β -ketoacyl synthase; ER, β -enoyl reductase; DH, dehydratase; KR, β -ketoacyl reductase. B, Linear domain organization of fatty acid synthase (FASN) of *M. elongata* AG77. PPT, phosphopantetheine transferase. C, PUFA synthesis. ELOVL, fatty acid elongase; FAD, fatty acid desaturase. Fatty acids are designated by the number of total carbon:the number of double bonds. The position of specific double bonds is indicated either from the carboxyl end (Δ) or from the methyl end (ω). D, TAG synthesis. ALDH, aldehyde dehydrogenase; ADH, alcohol dehydrogenase; GK, glycerol kinase; GPDH, glycerol-3-phosphate dehydrogenase; GPAT, glycerol-3-phosphate acyltransferase; PlsC, 1-acyl-*sn*-glycerol-3-phosphate acyltransferase; LPIN, phosphatidate phosphatase LPIN; PAP, phosphatidate phosphatase 2; Dgk, diacylglycerol kinase; DGAT, diacylglycerol acyltransferase; PDAT, phospholipid diacylglycerol acyltransferase.

References

1. Georgianna DR, Mayfield SP: **Exploiting diversity and synthetic biology for the production of algal biofuels.** *Nature* 2012, **488**(7411):329-335.
2. Hu Q, Sommerfeld M, Jarvis E, Ghirardi M, Posewitz M, Seibert M, Darzins A: **Microalgal triacylglycerols as feedstocks for biofuel production: perspectives and advances.** *Plant J* 2008, **54**(4):621-639.
3. Miranda AF, Taha M, Wrede D, Morrison P, Ball AS, Stevenson T, Mouradov A: **Lipid production in association of filamentous fungi with genetically modified cyanobacterial cells.** *Biotechnol Biofuels* 2015, **8**:179.
4. Li-Beisson Y, Beisson F, Riekhof W: **Metabolism of acyl-lipids in *Chlamydomonas reinhardtii*.** *Plant J* 2015, **82**(3):504-522.
5. Merchant SS, Kropat J, Liu B, Shaw J, Warakanont J: **TAG, you're it! *Chlamydomonas* as a reference organism for understanding algal triacylglycerol accumulation.** *Curr Opin Biotechnol* 2012, **23**(3):352-363.
6. Ma XN, Chen TP, Yang B, Liu J, Chen F: **Lipid Production from *Nannochloropsis*.** *Mar Drugs* 2016, **14**(4).
7. Du ZY, Benning C: **Triacylglycerol Accumulation in Photosynthetic Cells in Plants and Algae.** *Subcell Biochem* 2016, **86**:179-205.
8. Ajjawi I, Verruto J, Aqui M, Soriaga LB, Coppersmith J, Kwok K, Peach L, Orchard E, Kalb R, Xu W *et al*: **Lipid production in *Nannochloropsis gaditana* is doubled by decreasing expression of a single transcriptional regulator.** *Nat Biotechnol* 2017, **35**(7):647-652.
9. Zienkiewicz K, Zienkiewicz A, Poliner E, Du ZY, Vollheyde K, Herrfurth C, Marmon S, Farre EM, Feussner I, Benning C: ***Nannochloropsis*, a rich source of diacylglycerol acyltransferases for engineering of triacylglycerol content in different hosts.** *Biotechnol Biofuels* 2017, **10**:8.
10. Ma Y, Wang Z, Yu C, Yin Y, Zhou G: **Evaluation of the potential of 9 *Nannochloropsis* strains for biodiesel production.** *Bioresour Technol* 2014, **167**:503-509.
11. Rodolfi L, Chini Zittelli G, Bassi N, Padovani G, Biondi N, Bonini G, Tredici MR: **Microalgae for oil: strain selection, induction of lipid synthesis and outdoor mass cultivation in a low-cost photobioreactor.** *Biotechnol Bioeng* 2009, **102**(1):100-112.
12. Liu JY, Song YM, Qiu W: **Oleaginous microalgae *Nannochloropsis* as a new model for biofuel production: Review & analysis.** *Renewable and Sustainable Energy Reviews* 2017, **72**:154-162.
13. Vieler A, Wu G, Tsai CH, Bullard B, Cornish AJ, Harvey C, Reca IB, Thornburg C, Achawanantakun R, Buehl CJ *et al*: **Genome, functional gene annotation, and nuclear transformation of the heterokont oleaginous alga *Nannochloropsis oceanica* CCMP1779.** *PLoS Genet* 2012, **8**(11):e1003064.
14. Radakovits R, Jinkerson RE, Fuerstenberg SI, Tae H, Settlage RE, Boore JL, Posewitz MC: **Draft genome sequence and genetic transformation of the oleaginous alga *Nannochloropsis gaditana*.** *Nat Commun* 2012, **3**:686.
15. Wang D, Ning K, Li J, Hu J, Han D, Wang H, Zeng X, Jing X, Zhou Q, Su X *et al*: ***Nannochloropsis* genomes reveal evolution of microalgal oleaginous traits.** *PLoS Genet* 2014, **10**(1):e1004094.

16. Wang Q, Lu Y, Xin Y, Wei L, Huang S, Xu J: **Genome editing of model oleaginous microalgae *Nannochloropsis* spp. by CRISPR/Cas9.** *Plant J* 2016, **88**(6):1071-1081.
17. Poliner E, Takeuchi T, Du ZY, Benning C, Farre E: **Non-transgenic marker-free gene disruption by an episomal CRISPR system in the oleaginous microalga, *Nannochloropsis oceanica* CCMP1779.** *ACS Synth Biol* 2018, in press.
18. Poliner E, Pulman JA, Zienkiewicz K, Childs K, Benning C, Farre EM: **A toolkit for *Nannochloropsis oceanica* CCMP1779 enables gene stacking and genetic engineering of the eicosapentaenoic acid pathway for enhanced long-chain polyunsaturated fatty acid production.** *Plant Biotechnol J* 2018, **16**(1):298-309.
19. Xin Y, Lu Y, Lee YY, Wei L, Jia J, Wang Q, Wang D, Bai F, Hu H, Hu Q *et al*: **Producing Designer Oils in Industrial Microalgae by Rational Modulation of Co-evolving Type-2 Diacylglycerol Acyltransferases.** *Mol Plant* 2017, **10**(12):1523-1539.
20. Gao Y, Gregor C, Liang Y, Tang D, Tweed C: **Algae biodiesel - a feasibility report.** *Chem Cent J* 2012, **6 Suppl 1**:S1.
21. Gerber LN, Tester JW, Beal CM, Huntley ME, Sills DL: **Target Cultivation and Financing Parameters for Sustainable Production of Fuel and Feed from Microalgae.** *Environ Sci Technol* 2016, **50**(7):3333-3341.
22. Huang GH, Chen F, Wei D, Zhang XW, Chen G: **Biodiesel production by microalgal biotechnology.** *Applied Energy* 2010, **87**:38-46.
23. Gultom SO, Hu B: **Review of Microalgae Harvesting via Co-Pelletization with Filamentous Fungus.** *Energies* 2013, **6**:5921-5939.
24. Leite GB, Abdelaziz AE, Hallenbeck PC: **Algal biofuels: challenges and opportunities.** *Bioresour Technol* 2013, **145**:134-141.
25. Aguirre AM, Bassi A, Saxena P: **Engineering challenges in biodiesel production from microalgae.** *Crit Rev Biotechnol* 2013, **33**(3):293-308.
26. Rocca S AA, Giuntoli J, Marelli L. Sci. Tech. Res. Rep., Off. Eur. Union, Luxembourg: **Biofuels from algae: technology options, energy balance and GHG emissions; Insights from a literature review.** 2015, **EUR 27582**.
27. Miranda AF, Ramkumar N, Andriotis C, Holtkemeier T, Yasmin A, Rochfort S, Wlodkowic D, Morrison P, Roddick F, Spangenberg G *et al*: **Applications of microalgal biofilms for wastewater treatment and bioenergy production.** *Biotechnol Biofuels* 2017, **10**:120.
28. Rajendran A, Hu B: **Mycoalgae biofilm: development of a novel platform technology using algae and fungal cultures.** *Biotechnol Biofuels* 2016, **9**:112.
29. Padmaperuma G, Kapoore RV, Gilmour DJ, Vaidyanathan S: **Microbial consortia: a critical look at microalgae co-cultures for enhanced biomanufacturing.** *Crit Rev Biotechnol* 2017:1-14.
30. Muradov N, Taha M, Miranda AF, Wrede D, Kadali K, Gujar A, Stevenson T, Ball AS, Mouradov A: **Fungal-assisted algal flocculation: application in wastewater treatment and biofuel production.** *Biotechnol Biofuels* 2015, **8**:24.
31. Wan C, Zhao XQ, Guo SL, Asraful Alam M, Bai FW: **Bioflocculant production from *Solibacillus silvestris* W01 and its application in cost-effective harvest of marine microalga *Nannochloropsis oceanica* by flocculation.** *Bioresour Technol* 2013, **135**:207-212.

32. Uehling J, Gryganskyi A, Hameed K, Tschaplinski T, Misztal PK, Wu S, Desiro A, Vande Pol N, Du Z, Zienkiewicz A *et al*: **Comparative genomics of *Mortierella elongata* and its bacterial endosymbiont *Mycoavidus cysteinexigens***. *Environ Microbiol* 2017, **19**(8):2964-2983.
33. Scholz MJ, Weiss TL, Jinkerson RE, Jing J, Roth R, Goodenough U, Posewitz MC, Gerken HG: **Ultrastructure and composition of the *Nannochloropsis gaditana* cell wall**. *Eukaryot Cell* 2014, **13**(11):1450-1464.
34. Sakuradani E, Ando A, Ogawa J, Shimizu S: **Improved production of various polyunsaturated fatty acids through filamentous fungus *Mortierella alpina* breeding**. *Appl Microbiol Biotechnol* 2009, **84**(1):1-10.
35. Ji XJ, Ren LJ, Nie ZK, Huang H, Ouyang PK: **Fungal arachidonic acid-rich oil: research, development and industrialization**. *Crit Rev Biotechnol* 2014, **34**(3):197-214.
36. Guillard RRL (ed.): **Culture of phytoplankton for feeding marine invertebrates**. . New York, USA.: Plenum Press 1975.
37. Poliner E, Panchy N, Newton L, Wu G, Lapinsky A, Bullard B, Zienkiewicz A, Benning C, Shiu SH, Farre EM: **Transcriptional coordination of physiological responses in *Nannochloropsis oceanica* CCMP1779 under light/dark cycles**. *Plant J* 2015, **83**(6):1097-1113.
38. Jia j, Han DX, Gerken HG, Li YT, Sommerfeld M, Hu Q, Xu J: **Molecular mechanisms for photosynthetic carbon partitioning into storage neutral lipids in *Nannochloropsis oceanica* under nitrogen-depletion conditions** *Algal research* 2015, **7**:66-77.
39. Zienkiewicz K, Du ZY, Ma W, Vollheyde K, Benning C: **Stress-induced neutral lipid biosynthesis in microalgae - Molecular, cellular and physiological insights**. *Biochim Biophys Acta* 2016, **1861**(9 Pt B):1269-1281.
40. Lucker BF, Hall CC, Zegarac R, Kramer DM: **The environmental photobioreactor (ePBR): An algal culturing platform for simulating dynamic natural environments**. . *Algal research* 2014, **6**:242-249.
41. Hii YS, Soo CL, Chuah TS, Mohd-Azmi A, Abol-Munafi A: **Interactive effect of ammonia and nitrate on the nitrogen uptake by *Nannochloropsis* sp.** . *Journal of Sustainability Science and Management* 2011, **6**:60-68.
42. Ren M, Ogden K: **Cultivation of *Nannochloropsis gaditana* on Mixtures of Nitrogen Sources Environ. Prog. Sustainable Energy**. *Environmental Progress & Sustainable Energy* 2013, **33**:DOI 10.1002/ep.
43. Vieler A, Brubaker SB, Vick B, Benning C: **A lipid droplet protein of *Nannochloropsis* with functions partially analogous to plant oleosins**. *Plant Physiol* 2012, **158**(4):1562-1569.
44. Takeno S, Sakuradani E, Tomi A, Inohara-Ochiai M, Kawashima H, Shimizu S: **Transformation of oil-producing fungus, *Mortierella alpina* 1S-4, using Zeocin, and application to arachidonic acid production**. *J Biosci Bioeng* 2005, **100**(6):617-622.
45. Ando A, Sakuradani E, Horinaka K, Ogawa J, Shimizu S: **Transformation of an oleaginous zygomycete *Mortierella alpina* 1S-4 with the carboxin resistance gene conferred by mutation of the iron-sulfur subunit of succinate dehydrogenase**. *Curr Genet* 2009, **55**(3):349-356.

46. Leibundgut M, Maier T, Jenni S, Ban N: **The multienzyme architecture of eukaryotic fatty acid synthases.** *Curr Opin Struct Biol* 2008, **18**(6):714-725.
47. Wang L, Chen W, Feng Y, Ren Y, Gu Z, Chen H, Wang H, Thomas MJ, Zhang B, Berquin IM *et al*: **Genome characterization of the oleaginous fungus *Mortierella alpina*.** *PLoS One* 2011, **6**(12):e28319.
48. !!! INVALID CITATION !!!
49. Lutzoni F, Pagel M, Reeb V: **Major fungal lineages are derived from lichen symbiotic ancestors.** *Nature* 2001, **411**(6840):937-940.
50. Talukder MMR, Das, P., Wu, J.C.: **Immobilization of microalgae on exogenous fungal mycelium: A promising separation method to harvest both marine and freshwater microalgae.** *Biochem Eng J* 2014(91):53-57.
51. Wrede D, Taha M, Miranda AF, Kadali K, Stevenson T, Ball AS, Mouradov A: **Co-cultivation of fungal and microalgal cells as an efficient system for harvesting microalgal cells, lipid production and wastewater treatment.** *PLoS One* 2014, **9**(11):e113497.
52. Bonito G, Hameed K, Ventura K, Krishnan R, Schadt JC, Vilgalys R: **Isolating a functionally relevant guild of fungi from the root microbiome of *Populus*.** *Fungal Ecol* 2016, **22**:35-42.
53. Manocha MS: **Cell surface characteristics of *Mortierella* species and their interaction with a rnycoparasite** *Canadian Journal of Microbiology* 1984, **30**:290-298.
54. Ruiz-Herrera J, Osorio E: **Isolation and chemical analysis of the cell wall of *Morchella* sp.** *Antonie Van Leeuwenhoek* 1974, **40**(1):57-64.
55. Fernandes BS, Fernandes Messias MC, Carvalho PO, Zaiat M, da Cruz Pradella JG: **Data of added-value lipid production, Arachidonic acid, among other lipids by *Mortierella elongata*, using low cost simulated wastewater.** *Data Brief* 2017, **14**:255-259.
56. Song X, Ma Z, Tan Y, Zhang H, Cui Q: **Wastewater recycling technology for fermentation in polyunsaturated fatty acid production.** *Bioresour Technol* 2017, **235**:79-86.
57. Liu Z, Ruan Z, Xiao Y, Yi Y, Tang YJ, Liao W, Liu Y: **Integration of sewage sludge digestion with advanced biofuel synthesis.** *Bioresour Technol* 2013, **132**:166-170.
58. Durrett TP, Benning C, Ohlrogge J: **Plant triacylglycerols as feedstocks for the production of biofuels.** *Plant J* 2008, **54**(4):593-607.
59. Ghasemi Naghdi F, Gonzalez Gonzalez LM, Chan W, Schenk PM: **Progress on lipid extraction from wet algal biomass for biodiesel production.** *Microb Biotechnol* 2016, **9**(6):718-726.
60. Du ZY, Lucker BF, Zienkiewicz K, Miller TE, Zienkiewicz A, Sears BB, Kramer DM, Benning C: **Galactoglycerolipid Lipase PGD1 Is Involved in Thylakoid Membrane Remodeling in Response to Adverse Environmental Conditions in *Chlamydomonas*.** *Plant Cell* 2018.
61. Cuellar-Bermudez SP, Aguilar-Hernandez I, Cardenas-Chavez DL, Ornelas-Soto N, Romero-Ogawa MA, Parra-Saldivar R: **Extraction and purification of high-value metabolites from microalgae: essential lipids, astaxanthin and phycobiliproteins.** *Microb Biotechnol* 2015, **8**(2):190-209.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
62. Mackenzie DA, Wongwathanarat P, Carter AT, Archer DB: **Isolation and use of a homologous histone H4 promoter and a ribosomal DNA region in a transformation vector for the oil-producing fungus *Mortierella alpina***. *Appl Environ Microbiol* 2000, **66**(11):4655-4661.
63. Andersen RA, Berges JA, Harrison PJ, Watanabe MM: **Appendix A. Algal Culturing Techniques**. . San Diego: Elsevier Academic Press; 2005.
64. Liu B, Vieler A, Li C, Jones AD, Benning C: **Triacylglycerol profiling of microalgae *Chlamydomonas reinhardtii* and *Nannochloropsis oceanica***. *Bioresour Technol* 2013, **146**:310-316.
65. Du ZY, Lucker BF, Zienkiewicz K, Miller TE, Zienkiewicz A, Sears BB, Kramer DM, Benning C: **Galactoglycerolipid Lipase PGD1 Is Involved in Thylakoid Membrane Remodeling in Response to Adverse Environmental Conditions in *Chlamydomonas***. *Plant Cell* 2018, **30**(2):447-465.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6







Figure 7




Click here to access/download
Supplementary Material
Additional File 1_ST1.pdf







Click here to access/download
Supplementary Material
Additional File 2_FS1.pdf





Click here to access/download
Supplementary Material
Additional File 3_FS2.pdf








Click here to access/download
Supplementary Material
Additional File 4_FS3.pdf








Click here to access/download
Supplementary Material
Additional File 5_FS4.pdf








Click here to access/download
Supplementary Material
Additional File 6_FS5.pdf








Click here to access/download
Supplementary Material
Additional File 7_FS6.pdf





Click here to access/download
Supplementary Material
Additional File 8_TS2.pdf







[Click here to access/download](#)

Supplementary Material

Response to Reviewers_F.pdf





[Click here to access/download](#)

Supplementary Material

Manuscript Nannofungi_Revision_tracked_R7.docx





Click here to access/download
Supplementary Material
Supplemental data_tracked_R4.pdf



Dear Editor,

Thank you for your helpful comments. Please find enclosed the revised version of our manuscript prepared by the first author and me. We made edits as suggested by the reviewers and explained in the uploaded response letter. I also included a tracked version of the revised manuscript and a tracked version of the combined supplemental material both for review purposes only.

Thanks for handling this manuscript.

With best regards, Christoph Benning