



Vortex Axisymmetrization, Inviscid Damping, and Vorticity Depletion in the Linearized 2D Euler Equations

Jacob Bedrossian¹ · Michele Coti Zelati² · Vlad Vicol^{3,4}

Received: 24 July 2018 / Accepted: 29 January 2019

© Springer Nature Switzerland AG 2019

Abstract

Coherent vortices are often observed to persist for long times in turbulent 2D flows even at very high Reynolds numbers and are observed in experiments and computer simulations to potentially be asymptotically stable in a weak sense for the 2D Euler equations. We consider the incompressible 2D Euler equations linearized around a radially symmetric, strictly monotone decreasing vorticity distribution. For sufficiently regular data, we prove the inviscid damping of the θ -dependent radial and angular velocity fields with the optimal rates $\|u^r(t)\| \lesssim \langle t \rangle^{-1}$ and $\|u^\theta(t)\| \lesssim \langle t \rangle^{-2}$ in the appropriate radially weighted L^2 spaces. We moreover prove that the vorticity weakly converges back to radial symmetry as $t \rightarrow \infty$, a phenomenon known as *vortex axisymmetrization* in the physics literature, and characterize the dynamics in higher Sobolev spaces. Furthermore, we prove that the θ -dependent angular Fourier modes in the vorticity are ejected from the origin as $t \rightarrow \infty$, resulting in faster inviscid damping rates than those possible with passive scalar evolution. This non-local effect is called *vorticity depletion*. Our work appears to be the first to find vorticity depletion relevant for the dynamics of vortices.

Keywords Euler equations · Inviscid damping · Vortex axisymmetrization · Vorticity depletion · Phase mixing · Scattering

Jacob Bedrossian was partially supported by NSF CAREER Grant Nos. DMS-1552826, NSF DMS-1413177, and a Alfred P. Sloan research fellowship. Additionally, the research was supported in part by NSF RNMS #1107444 (Ki-Net). Michele Coti Zelati was partially supported by NSF Grant DMS-1713886. Vlad Vicol was partially supported by the NSF CAREER Grant DMS-1652134, and an Alfred P. Sloan research fellowship.

✉ Vlad Vicol
vvicol@math.princeton.edu; vicol@cims.nyu.edu

Extended author information available on the last page of the article

Contents

1	Introduction and Statements of Results	...
1.1	Background	...
1.2	Statement of Main Results	...
2	Preliminaries and Outline of the Proof	...
2.1	Skew-Symmetric Structure, Neutral Modes, and Contour Integral Representation	...
2.2	Outline of the Proof for $k = 1$...
2.3	Outline of the Proof for $k \geq 2$...
2.3.1	Depletion Trick and Contour Integral Decomposition	...
2.3.2	Vanishing for $z \notin I_\alpha$ as $\varepsilon \rightarrow 0$...
2.3.3	Green's Function for the Rayleigh Problem for $z \in I_\alpha$ as $\varepsilon \rightarrow 0$...
2.3.4	Representation Formulas and Boundedness for $(r\partial_r)^j f_1$ and $(r\partial_r)^j f_2$...
3	Dynamics of the $k = 1$ Mode	...
3.1	An Explicit Representation of the Vorticity Profile	...
3.2	Vorticity Depletion and Inviscid Damping	...
4	The Homogeneous Rayleigh Problem for $k \geq 2$...
4.1	Existence and Uniqueness of Solutions	...
4.1.1	An Auxiliary Weight	...
4.1.2	Existence and Uniqueness of \tilde{P}	...
4.2	Convergence of $(r\partial_r)^j \tilde{P}$...
4.3	Analysis of $\partial_G \tilde{P}$...
4.4	Analysis of $\partial_{r_c} \tilde{P}$...
4.5	Convergence in Optimal Weights	...
4.5.1	The Case $r_c \leq 1$...
4.5.2	The Case $r_c > 1$...
4.6	Further Properties for the Real Solution	...
4.6.1	The Function Q_0 and Its Properties	...
4.6.2	The Function Q_∞ and Its Properties	...
5	The Inhomogeneous Rayleigh Problem for $k \geq 2$...
5.1	Estimates on the Wronskian	...
5.2	Estimates on H_0 and H_∞	...
6	Representation Formulas and Estimates on $(r\partial_r)^j f_1$, and $(r\partial_r)^j f_2$...
6.1	Recursion Relations for Derivatives of f_1 and f_2	...
6.2	Integral Operators Appearing in the Derivatives of X and Y	...
6.3	Iterated Integral Operators	...
6.3.1	Recursion Scheme for Integral Operators	...
6.3.2	Estimates on Iterated Integral Kernels	...
6.4	Vorticity Decomposition	...
A	Preliminary Technical Lemmas	...
B	Boundedness and Convergence of Integral Operators	...
B.1	Two Singular Integrals and One Delta Distribution	...
B.1.1	Convergence Away from the Diagonal $s_0 = r_c$...
B.1.2	Identifying the Leading Order Operator Near the $s_0 = r_c$ Diagonal	...
B.1.3	Convergence of the Operator with $s_0 = r_c$...
B.1.4	A Useful Product Formula for Weights	...
B.2	The Remaining Combinations of Iterated Operators	...
C	Properties of H_0 and H_∞	...
D	Vanishing for $k \geq 2$ Outside I_α	...
D.1	Estimates Near $c = u(0)$...
D.1.1	Proof of (D.5)	...
D.1.2	Proof of (D.6)	...
D.1.3	Proof of (D.7)	...
D.2	Estimates Near $c = 0$...
D.2.1	Proof of (D.8)	...
D.2.2	Proof of (D.9)	...
D.3	Vanishing of f_E	...

D.4 Vanishing of f_S
D.5 Vanishing of Higher Derivatives
D.5.1 Proof for $j \in \{1, \dots, k-1\}$
D.5.2 Proof for $j = k$
References

1 Introduction and Statements of Results

In polar coordinates $(r, \theta) \in [0, \infty) \times \mathbb{T}$, the two-dimensional Euler equations in vorticity formulation read

$$\partial_t \tilde{\omega} + \tilde{u}^r \partial_r \tilde{\omega} + \frac{\tilde{u}^\theta}{r} \partial_\theta \tilde{\omega} = 0, \quad (1.1)$$

where the velocity vector $\tilde{\mathbf{u}} = (\tilde{u}^r, \tilde{u}^\theta)$ is recovered from the vorticity $\tilde{\omega}$ by means of the streamfunction $\tilde{\psi}$, via the relations

$$\begin{aligned} \tilde{\omega} &= -\frac{1}{r} \partial_\theta \tilde{u}^r + \frac{1}{r} \partial_r (r \tilde{u}^\theta), \quad \tilde{\mathbf{u}} = \left(\frac{1}{r} \partial_\theta \tilde{\psi}, -\partial_r \tilde{\psi} \right), \\ & - \left(\partial_{rr} + \frac{1}{r} \partial_r + \frac{1}{r^2} \partial_{\theta\theta} \right) \tilde{\psi} = \tilde{\omega}. \end{aligned}$$

Any radially symmetric configuration $\Omega = \Omega(r)$ is a stationary solution of (1.1) and the above relations simplify to

$$\Omega = \frac{1}{r} \partial_r (rU), \quad U = -\partial_r \Psi, \quad - \left(\partial_{rr} + \frac{1}{r} \partial_r \right) \Psi = \Omega. \quad (1.2)$$

In what follows, we denote

$$u(r) = \frac{U(r)}{r} = -\frac{\partial_r \Psi(r)}{r}, \quad \beta(r) = -\frac{\partial_r \Omega(r)}{r}. \quad (1.3)$$

Writing $\tilde{\omega}(t, r, \theta) = \omega(t, r, \theta) + \Omega(r)$ and dropping terms quadratic in ω gives the linearized 2D Euler equations which are the main object of study in this paper:

$$\partial_t \omega + u(r) \partial_\theta \omega - \beta(r) \partial_\theta \psi = 0, \quad (1.4a)$$

$$- \left(\partial_{rr} + \frac{1}{r} \partial_r + \frac{1}{r^2} \partial_{\theta\theta} \right) \psi = \omega, \quad (1.4b)$$

$$\omega(0, r, \theta) = \omega^{in}(r, \theta). \quad (1.4c)$$

By expanding the solution ω to (1.4) as a Fourier series in the angular θ variable, namely

$$\omega(t, r, \theta) = \sum_{k \in \mathbb{Z}} \omega_k(t, r) e^{ik\theta}, \quad \psi(t, r, \theta) = \sum_{k \in \mathbb{Z}} \psi_k(t, r) e^{ik\theta}, \quad (1.5)$$

we perform a k -by- k analysis of the linearized equations, since for any integer k we have that

$$\partial_t \omega_k + iku(r)\omega_k - ik\beta(r)\psi_k = 0, \quad (1.6a)$$

$$-\Delta_k \psi_k := -\left(\partial_{rr} + \frac{1}{r}\partial_r - \frac{k^2}{r^2}\right)\psi_k = \omega_k. \quad (1.6b)$$

Note that $\omega_0(t, r) = \omega_0(0, r)$ (i.e., the θ -average of the solution), and therefore we restrict to $k \neq 0$ without loss of generality. Similarly, by reality we only consider $k \geq 1$ without loss of generality.

1.1 Background

The stability of vortices is one of the most fundamental problems in the theory of hydrodynamic stability and has been considered by many authors, starting with Kelvin [62] and Orr [54] and continuing to present day in both mathematics and physics (see e.g. [2,18,33,34,36,40,43,49,57] and the references therein). Nonlinear stability results in weighted L^2 spaces (of the vorticity) are available using energy-Casimir methods [29], however, these results do not provide a clear description of the long-time dynamics. Experimental observations, computer simulations, and formal asymptotics (see e.g. [2,3,36,57] and the references therein) suggest that a vortex subjected to a sufficiently small disturbance might return to radial symmetry as $t \rightarrow \infty$ in a weak sense. Specifically, it is observed that the vorticity in the angle-dependent modes is stirred up around the vortex into a spiral pattern (sometimes called ‘filamentation’) and eventually the angle-dependent modes weakly converge to zero as $t \rightarrow \infty$. This weak convergence is called *vortex axisymmetrization* and is thought to be relevant to understanding coherent vortices in 2D turbulence [17], atmospheric dynamics [51,58], and various settings in plasma physics [22,57,71]. Our paper appears to be the first mathematically rigorous confirmation of this behavior for vortices in the linearized 2D Euler equations and the first paper to obtain an accurate prediction of the decay rates for general initial data.

When studying the stability of the planar Couette flow, Orr [54] identified another effect associated with vorticity mixing: the strong convergence (in L^2) of the velocity field to equilibrium as $t \rightarrow \infty$. This effect is now often called *inviscid damping*. Further studies of the 2D Euler equations linearized around planar shear flows were made by Case [20], Dikii [27], and Stepin [60]. More recently, the linearized problem was revisited in [45] and optimal rates were deduced by Zillinger [73,74] for shear flows close to Couette flow and later by Wei et al. [66] for more general strictly monotone shear flows in a channel. See also [24,75] for inviscid damping of Taylor–Couette in an annulus and [69] for inviscid damping in stratified shear flows. The recent works of Wei et al. [67,68] deduce optimal inviscid damping rates for the some shear flows with non-degenerate critical points. This latter works also confirmed the predictions of Bouchet and Morita [16] that the linearized 2D Euler equations can have a faster inviscid damping rate than if the vorticity were evolving under passive scalar dynamics. We prove a similar effect here as well; see the discussion

following Theorem 1.1 for more details. Finally, see [44] for an approach to the problem which is well-suited to treating general problems but obtains less precise decay estimates.

Inviscid damping is closely related to Landau damping in the Vlasov equations, which arises in the kinetic theory of plasmas and galactic dynamics. Landau damping involves the rapid decay of the self-generated electric field in a plasma in the absence of any dissipative mechanisms and was first predicted by Landau in the Vlasov equations linearized around a homogeneous Maxwellian [41]. The predictions matched with experiments [47] and many works on the linearized Vlasov equations followed (see [25,35,55,63] and the references therein). In Vlasov, the decay is caused by the mixing of particles traveling at different velocities whereas in 2D Euler it is caused by the mixing of vorticity. Due to the special structure of the Vlasov equations, inviscid damping for the linearized 2D Euler equations (with the exception of the Couette flow [39,54]) is significantly harder than the linearized theory of Landau damping near *homogeneous* equilibria (and in general the rates are much faster—on $\mathbb{T}^n \times \mathbb{R}^n$ the decay can be exponential).

All of the above mentioned works on inviscid and Landau damping only apply to the *linearized* Euler or Vlasov equations (respectively). The work of [19] first demonstrated the existence of (analytic) Landau damping solutions to the nonlinear Vlasov equations (see also [37]). Later, Mouhot and Villani [52] demonstrated that on $\mathbb{T}^n \times \mathbb{R}^n$, all perturbations small enough in a sufficiently regular Gevrey class give rise to nonlinear dynamics that matches the linearized dynamics (and in particular, rapid Landau damping). This work was followed subsequently by a variety of others on Landau damping in nonlinear Vlasov and related models (see e.g. [9,15,31,32,70] and the references therein). In [14], it was shown that Orr’s inviscid damping predictions for Couette flow hold also for the nonlinear 2D Euler equations in $\mathbb{T} \times \mathbb{R}$, provided the perturbation is small in a sufficiently regular Gevrey class. At sufficiently low regularities, it was proved in [45,46] that linearized and nonlinear dynamics do not necessarily agree (for both 2D Euler and Vlasov). High regularity does not play a special role in the linear theory, however, it was shown in [5] that on $\mathbb{T} \times \mathbb{R}$ one cannot (in general) extend the linearized theory of Landau damping to the nonlinear Vlasov equations in *any* Sobolev space (however, see [9]). This is due to the *plasma echoes*, a nonlinear oscillation observed in experiments in [48]. The work of [5] showed the existence of solutions to Vlasov with arbitrarily many, arbitrarily small, plasma echoes. Similar nonlinear echoes are observed in experiments on vortices in the 2D Euler equations [71,72] (see also the analyses of [64,65]). Hence, we expect the linear and nonlinear regularity requirements to be drastically different. For this reason, it is important to study the linearized Euler equations in as high regularity as possible (preferably Gevrey class), and determine if such high regularity can be propagated in a suitable sense (see Remark 1.4 below). If the answer is ‘no’, then it might be possible to introduce nonlinear instabilities even for Gevrey or analytic data in the nonlinear equations.

Mixing involves a transfer of vorticity from large to small scales. When there is diffusion present, it has been shown that this can enhance the dissipative time-scale. For example, Kelvin showed in [39] that x -dependent modes of the linearized Couette flow in $\mathbb{T} \times \mathbb{R}$ decay on a time-scale like $O(v^{-1/3})$ as opposed to the natural heat equation

time-scale of $O(\nu^{-1})$ (denoting ν to be the inverse Reynolds number). This effect has been called the ‘shear-diffuse mechanism’, ‘relaxation enhancement’, or ‘enhanced dissipation’ and has been studied many times in linear and some nonlinear settings in both mathematics [4,6–8,10–13,23,26,33,38,43,67] and physics [1,30,42,56]. Thus far, it has also proved challenging to obtain enhanced dissipation estimates on the linearized Navier–Stokes. Ideally, the goal is to obtain both simultaneously on Navier–Stokes; it seems the best result in this direction for non-trivial shear flows is [67]. See [26,33,43] and the references therein for the most recent results on the 2D Navier–Stokes equations linearized around the Oseen vortex.

1.2 Statement of Main Results

Throughout the article, we assume the following conditions on the background vortex:

- (V1) $0 \leq \Omega(r) \lesssim \langle r \rangle^{-6}$;
 (V2) $|(r\partial_r)^j \Omega(r)| \lesssim_j \langle r \rangle^{-6}$ for all $j \geq 0$;
 (V3) $\partial_r \Omega(r) < 0, \forall r > 0$,

Among the above assumptions we single out that Ω is assumed to be smooth, and the fact that Ω is assumed to be *strictly* monotone. Indeed, strict monotonicity is a well-known sufficient condition for the linearized stability (see also Remark 1.11 below). We additionally take the following orthogonality condition on the initial condition of (1.4):

$$\int_0^\infty \omega_{\pm 1}^{in}(r) r^2 dr = 0, \quad (1.7)$$

which removes the neutral modes that arise due to the translation invariance (see Lemma 1.7). Recall that for smooth functions ω , the asymptotic expansion of $\omega_k(r)$ at the origin is necessarily in the form (see e.g. [3])

$$\omega_k(r) \sim r^k \left(a_0 + a_1 r^2 + a_2 r^4 + \dots \right) \text{ as } r \rightarrow 0. \quad (1.8)$$

Let χ be a smooth cut-off which is 1 for $r < 1/2$ and 0 for $r > 3/4$ and denote

$$\omega_{k,0}^{in} = \lim_{r \rightarrow 0} \frac{\omega_k^{in}(r)}{r^k}, \quad F_k(r) = \omega_k^{in}(r) \sqrt{r} - \frac{\beta(r)}{\beta(0)} \chi(r) r^{k+1/2} \omega_{k,0}^{in}. \quad (1.9)$$

By (1.8), for smooth ω_k^{in} , we have $F_k(r) \sim r^{k+2+1/2}$ as $r \rightarrow 0$ and thus we may use a stronger weight for F_k at the origin than for ω_k^{in} . To state our main result, for $\delta > 0$ we define smooth weights $w_{\psi,\delta}$, $w_{f,\delta}$, $w_{F,\delta}$ and corresponding L^2 -norms which satisfy the following asymptotics:

$$w_{\psi,\delta}(r) \approx \min \left\{ r^{k+1/2-\delta}, r^{-k+1/2+\delta} \right\}, \quad \|g\|_{L^2_{\psi,\delta}} := \left(\int_0^\infty \frac{|g(r)|^2}{w_{\psi,\delta}^2(r)} dr \right)^{1/2}, \quad (1.10a)$$

$$w_{f,\delta}(r) \approx \min \left\{ r^{k+1/2-\delta}, r^{-k+1/2-6+\delta} \right\}, \quad \|g\|_{L^2_{f,\delta}} := \left(\int_0^\infty \frac{|g(r)|^2}{w_{f,\delta}^2(r)} dr \right)^{1/2}, \quad (1.10b)$$

$$w_{F,\delta}(r) \approx \min \left\{ r^{k+1+2-\delta}, r^{-k+1-6+\delta} \right\}, \quad \|g\|_{L^2_{F,\delta}} := \left(\int_0^\infty \frac{|g(r)|^2}{w_{F,\delta}^2(r)} dr \right)^{1/2}. \quad (1.10c)$$

We assume that $(r\partial_r)^j w_{*,\delta}$ satisfies the same upper bounds as $w_{*,\delta}$ (up to a j -dependent constant). Noting that (V3) ensures $\beta > 0$, we also make use of the L^2 -space normed by

$$\|g\|_{L^2_\beta} := \left(\int_0^\infty \frac{|g(r)|^2}{\beta(r)} r dr \right)^{1/2},$$

which is the natural energy space for the equations (see Lemma 2.2). The main result of the paper is as follows.

Theorem 1.1 *Let $k \neq 0$, and assume (V1)–(V3). For all $1 \gg \delta \gg \eta_0 > 0$ and any smooth $\omega_k^{in} \in L^2_\beta \cap L^2_{f,\delta}$ satisfying the orthogonality condition (1.7), the solution ω, ψ to (1.4) obeys the following inviscid damping estimates*

$$\begin{aligned} \|\psi_k(t)\|_{L^2_{\psi,\delta}} &\lesssim_{\delta,\eta_0} \frac{1}{\langle kt \rangle^2} \left(k^{5+\eta_0} |\omega_{k,0}^{in}| + \sum_{j=0}^2 k^{6-2j+\eta_0} \|(r\partial_r)^j F_k\|_{L^2_{F,\delta/4}} \right) \\ \langle kt \rangle \|ru_k^r(t)\|_{L^2_{\psi,\delta}} &+ \|ru_k^\theta(t)\|_{L^2_{\psi,\delta}} \\ &\lesssim_{\delta,\eta_0} \frac{1}{\langle kt \rangle} \left(k^{5+\eta_0} |\omega_{k,0}^{in}| + \sum_{j=0}^2 k^{6-2j+\eta_0} \|(r\partial_r)^j F_k\|_{L^2_{F,\delta/4}} \right). \end{aligned} \quad (1.11)$$

Furthermore, there exist $f_{k;1}(t, r)$ and $f_{k;2}(t, r)$ such that

$$\omega_k(t, r) = e^{-iktu(r)} f_{k;1}(t, r) + e^{-iktu(r)} f_{k;2}(t, r), \quad (1.12)$$

and the following vorticity depletion estimates hold

$$\|\sqrt{r}(r\partial_r)^n f_{k;1}(t)\|_{L^2_{F,\delta}} \lesssim_{\delta,n,\eta_0} \left(k^{2n+1+\eta_0} |\omega_{k,0}^{in}| + \sum_{j=0}^n k^{2(n-j)+\eta_0} \|(r\partial_r)^j F_k\|_{L^2_{F,\delta/4}} \right), \quad (1.13a)$$

$$\begin{aligned} \|(r\partial_r)^n f_{k;2}(t)\|_{L^2_{f,\delta}} \\ \lesssim_{\delta,n,\eta_0} \frac{1}{\langle kt \rangle} \left(k^{2n+4+\eta_0} |\omega_{k,0}^{in}| + \sum_{j=0}^n k^{2(n-j)+3+\eta_0} \|(r\partial_r)^j F_k\|_{L^2_{F,\delta/4}} \right), \end{aligned} \quad (1.13b)$$

for all $0 \leq n \leq \max(2, k)$ in (1.13a) and for all $0 \leq n \leq \max(1, k-1)$ in (1.13b).

Remark 1.2 By density we can extend the results to cover any $\omega_k^{in} \in L^2$ (satisfying (1.7)) for which the norms appearing on the right-hand sides above are finite.

Remark 1.3 The L^2 norms we are using in Theorem 1.1, namely (1.10), are natural in light of (1.8) and are well-suited for studying vorticity depletion. However, these norms are quite strong at the origin (and infinity). Note that $\|\Delta_k^{-1}\|_{L_{f,\delta}^2 \rightarrow L_{\psi,\delta}^2} \approx k^{-1}$ (as opposed to k^{-2} as one might expect), which explains why some of the powers of k in Theorem 1.1 are slightly higher than might be at first expected. Similarly, note that F_k contains information about the second term in the expansion (1.8).

Remark 1.4 The correct analogue of propagation of regularity for mixing problems is the regularity of $e^{iktu(r)}\omega_k(t, r)$, the object which measures the difference between the passive scalar and full linearized (or nonlinear) dynamics. Regularity of this object is often studied in dispersive equations and it is sometimes called the ‘profile’; see e.g. [14] for more discussions (note that regularity of this type was called ‘gliding regularity’ in [52]). Understanding regularity of the profile plays a major role in all of the works involving nonlinear inviscid/Landau damping [5–8,10,14,15,19,52] including those which obtain results in Sobolev spaces [9,31]. Theorem 1.1 deduces higher regularity of the vorticity profile than is necessary to prove the (1.11), at least for $k \geq 3$. However, as regularity plays a crucial role in the nonlinear theory, it seems appropriate to study it as carefully as possible in the linear problem. This goal has motivated many of the primary aspects of our approach.

Remark 1.5 Because in this work we were only concerned with obtaining finite Sobolev regularity of the vorticity profile, we have not carefully quantified how the constants in (1.13) depend on n . This is sufficient for any fixed Sobolev space of interest, but e.g. for the end-point cases such as $n = k - 1$ and $n = k$, we have not quantified the rate at which the constants grow in n as $n = k \rightarrow \infty$, an issue which becomes important at infinite regularity.

A direct consequence of (1.13) is the following weak convergence result which shows that the solution weakly converges back to a radially symmetric vortex.

Corollary 1.6 (Vortex axisymmetrization). *For all $k \neq 0$, $\omega_k(t, r) \rightarrow 0$ in $L_{f,\delta/2}^2$ as $t \rightarrow \pm\infty$.*

The above corollary is due to the bound (1.11), (1.13) and interpolation. Another direct corollary of Theorem 1.1 shows that the vorticity behaves as a passive scalar evolution in the limit $t \rightarrow \pm\infty$ (the analogue of ‘scattering’ in dispersive equations):

Corollary 1.7 (Scattering to passive scalar evolution). *There exists $\omega_{k,\pm\infty} \in L_{f,\delta}^2$ such that*

$$\lim_{t \rightarrow \pm\infty} \|e^{iktu}\omega_k(t) - \omega_{k,\pm\infty}\|_{L_{f,\delta}^2} = 0.$$

Using (1.12) and (1.13) we only get a weak convergence result. To upgrade to the stated strong convergence, we read a bound on $\partial_t(f_{k;1}(t, r) + f_{k;2}(t, r))$ directly from

equation (1.6a) in the correct weighted space in view of the decay of β (see Lemma 2.1). This, combined with the strong decay of $f_{k;2}(t, r)$ in (1.13b), implies the desired strong convergence.

Remark 1.8 If we additionally proved that $(r\partial_r)^j f_{k,1}(t)$ converged as $t \rightarrow \pm\infty$ in $L^2_{f,\delta}$, then of course scattering in stronger spaces would follow immediately. This seems likely with some mild technical refinements of our method, but we did not pursue this direction.

Remark 1.9 Vortices with a Gaussian distribution of vorticity constitute an important class that satisfies our assumptions (V1)–(V3). Specifically, we can consider vortices with

$$\Omega_{\lambda,L}(r) = \frac{\lambda}{4\pi L^2} \exp\left(-\frac{r^2}{4L^2}\right), \quad \Psi_{\lambda,L}(r) = -\frac{\lambda}{2\pi} \int_0^r \frac{1}{r} \left[1 - \exp\left(-\frac{r^2}{4L^2}\right)\right] dr$$

having length scale $L > 0$ and total circulation $\lambda > 0$. In view of the notation introduced in (1.3), we can compute

$$u_{\lambda,L}(r) = \frac{\lambda}{2\pi} \frac{1}{r^2} \left[1 - \exp\left(-\frac{r^2}{4L^2}\right)\right], \quad \beta_{\lambda,L}(r) = \frac{\lambda}{8\pi L^4} \exp\left(-\frac{r^2}{4L^2}\right). \quad (1.14)$$

Remark 1.10 The restriction $j \leq \max(k, 2)$ and the loss of k^2 per $r\partial_r$ derivative (as opposed to k) are due to difficulties specific to the vortex case. We expect that our methods can easily be adapted to get boundedness of $e^{iktu(y)}\omega_k(y)$ in H^s for all $s \geq 0$ for strictly monotone shear flows on $\mathbb{T} \times \mathbb{R}$. Our methods may also be able to shed further light on higher derivatives of $e^{iktu(y)}\omega_k(y)$ in the case of a channel $\mathbb{T} \times [-1, 1]$ (see [66, 73]).

Remark 1.11 The strict monotonicity (V3) plays a crucial role. See e.g. the studies [2, 36, 53, 59] showing various kinds of pathologies in non-strictly monotone vortices, including embedded neutral modes (as occur e.g. in Rankine vortices) and non-normal algebraic instabilities. See also the recent nonlinear counter examples to inviscid damping around a vortex constructed in [21] without monotonicity.

The angular velocity of the background vortex $u(r)$ satisfies $u'(0) = 0$, which indicates that the mixing is weak at the vortex core. For well-localized data supported near the origin, one can show that the passive scalar evolution (e.g. $\omega_k(t, r) = e^{-iktu(r)}\omega_k^{in}(r)$) generally cannot give rise to inviscid damping faster than $\|u^r(t)\| \lesssim \langle t \rangle^{-1/2}$ and $\|u^\theta(t)\| \lesssim \langle t \rangle^{-1}$ (see the proof of Lemma 2.15 below for more details). Hence, the non-local term in (1.6a) improves the inviscid damping rate in (1.11). A related effect was conjectured for shear flows with non-degenerate critical points (e.g. points such that $u'(y_c) = 0$ but $u''(y_c) \neq 0$) by Bouchet and Morita [16]. Bouchet and Morita predict that vorticity will be ejected from the critical point, allowing the break-down of the mixing there to have less effect than naively predicted. Specifically,

Bouchet and Morita predict that the vorticity should behave as

$$\omega_k(t, y) = e^{-iktu(y)} \omega_{k,\infty}(y) + O(t^{-\gamma}) \text{ for some } \gamma > 0 \text{ such that } \omega_{k,\infty}(y_c) = 0. \quad (1.15)$$

In [67,68], the authors prove that indeed the inviscid damping for such shear flows can be faster than passive scalar. The authors directly study the evolution of the stream-function via methods somewhat different from our approach (though a number of common themes exist); our methods and theirs each have their own advantages and disadvantages. Specifically, our methods obtain more precise information about the vorticity directly, and thus the inviscid damping (1.11) is a straightforward consequence of our vorticity decomposition (1.13) (see Lemma 2.15). A vortex analogue of the depletion effect (1.15) (more carefully quantified) is described by (1.13):

$$\begin{aligned} \omega_k(t, r) &= e^{-iktu(r)} f_{k,1}(t, r) + O\left(\frac{r^k}{t}\right) \quad \text{as } r \rightarrow 0, \\ f_{k,1}(t, r) &= O(r^{k+2}) \quad \text{as } r \rightarrow 0. \end{aligned}$$

Although a hint of the vorticity depletion effect can be seen in the numerics and formal asymptotics of [3], our work appears to be the first to precisely connect this depletion effect to vortex dynamics.

Physically, vorticity depletion in the vortex seems related potentially to the centrifugal force that will tend to move vorticity away from the vortex core. This effect is over-powered by inviscid damping away from the vortex core, however near the core, there is apparently a balance that allows to alter the decay rate. See Remark 2.5 for a brief discussion of how it is deduced mathematically.

In what follows, we denote $\langle r \rangle = \sqrt{1 + r^2}$. We use the notation $f \lesssim g$ if there exists a constant $C > 0$ such that $f \leq Cg$ (and analogously $f \gtrsim g$) and $f \approx g$ if $f \lesssim g$ and $f \gtrsim g$. We use the notation $f \lesssim_{a,b,\dots} g$ to emphasize that the constant C depends on a, b, \dots . The implicit constants will *never* depend on the quantities c, k, r, ε , or ω_k^{in} or similar variables except where otherwise noted (see below for the appearance of these quantities). Finally, we let χ be a smooth, non-negative function which satisfies $\chi(r) = 1$ for $|r| < 1/2$ and 0 for $|r| > 3/4$.

2 Preliminaries and Outline of the Proof

2.1 Skew-Symmetric Structure, Neutral Modes, and Contour Integral Representation

The following lemma is a basic consequence of the Biot–Savart law for radially symmetric functions (recall $u(r) = (2\pi)^{-1} r^{-2} \int_0^r \Omega(s) s ds$).

Lemma 2.1 (Basic properties of the vortex). *Every $\Omega(r)$ satisfying conditions (VI)–(V3) satisfies the following:*

- $\beta(r) > 0$ for all $r \geq 0$, $u'(r) < 0$ for all $r \in (0, \infty)$, $u(r) > 0$ for all $r \geq 0$, and $u'(0) = 0$;
- $|(r\partial_r)^j \beta(r)| \lesssim_j \langle r \rangle^{-8}$ for all $j \geq 0$;
- $|(r\partial_r)^j u(r)| \lesssim_j \langle r \rangle^{-2}$ for all $j \geq 0$;
- for $r \leq 1$ there holds $u'(r) \approx -r$ and for $r \geq 1$ there holds $u'(r) \approx -r^{-3}$;
- there holds the identity

$$\beta(r) + u''(r) + \frac{3u'(r)}{r} = 0, \quad \forall r \in (0, \infty). \quad (2.1)$$

We rewrite (1.6a) in terms of the vorticity alone as

$$\partial_t \omega_k + ikL_k \omega_k = 0, \quad L_k := u(r) + \beta(r)\Delta_k^{-1}. \quad (2.2)$$

A key observation is the following, which is a straightforward calculation via Schur's lemma.

Lemma 2.2 *The operator $L_k : L_\beta^2 \rightarrow L_\beta^2$ is bounded and self-adjoint with respect to the inner product*

$$\langle g_1, g_2 \rangle_\beta := \int_0^\infty g_1(r) \overline{g_2(r)} \frac{r}{\beta(r)} dr.$$

It follows that the L_β^2 norm is a conserved quantity:

$$\int_0^\infty \frac{|\omega_k(t, r)|^2}{\beta(r)} r dr = \int_0^\infty \frac{|\omega_k^{in}(r)|^2}{\beta(r)} r dr, \quad \forall t \in \mathbb{R}. \quad (2.3)$$

Remark 2.3 This conserved quantity is the quadratic variation of the Casimir used in the energy-Casimir method of nonlinear stability [29].

The next lemma regards the neutral modes that arises due to translation invariance. Note that the conservation law in Lemma 2.4 below is equivalent to the conservation of $\int_{\mathbb{R}^2} x \omega dx$ under the Euler flow.

Lemma 2.4 (Neutral modes and orthogonality condition (1.7)). *When $k = \pm 1$, we have the conservation law:*

$$\frac{d}{dt} \int_0^\infty \omega_{\pm 1}(t, r) r^2 dr = 0. \quad (2.4)$$

Moreover, in view of (2.1), it is easy to check that

$$\psi_S(r) = ru(r), \quad \omega_S(r) = r\beta(r) \quad (2.5)$$

is a steady state for (1.6a) with $k = \pm 1$ (this is also pointed out in e.g. [3]). In particular, $\omega_S(r) = r\beta(r)$ is a neutral eigenmode for $L_{\pm 1}$.

Proof of Lemma 2.4 Indeed, dropping the time dependence and the indices $k = \pm 1$, a straightforward calculation (note that the boundary terms vanish due to L_β^2 conservation) shows that

$$\begin{aligned} \int_0^\infty [u(r)\omega(r) - \beta(r)\psi(r)] r^2 dr &= - \int_0^\infty \left[r^2 u(r) \partial_{rr} \psi(r) + r u(r) \partial_r \psi(r) \right. \\ &\quad \left. - u(r) \psi(r) + r^2 \beta(r) \psi(r) \right] dr \\ &= - \int_0^\infty \left[r^2 u''(r) \psi(r) + 3r u'(r) \psi(r) \right. \\ &\quad \left. + r^2 \beta(r) \psi(r) \right] dr = 0, \end{aligned}$$

where we made use of (2.1) in the last equality. Hence, from (1.6a) we infer (2.4). That (2.5) is a steady state is also simple consequence of (2.1). \square

As a result of the self-adjointness of L_k , (1.6a) falls under essentially the same framework as Schrödinger operators. Hence, for all $\omega_k^{in} \in L_\beta^2$, the solution to (2.2) can be represented via the formula (see [61, Proposition 1.9])

$$\omega_k(t, r) = e^{-ikL_k t} \omega_k^{in}(r) = \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2\pi i} \int_{\mathbb{R}} e^{-ikct} \left[(c - i\varepsilon - L_k)^{-1} - (c + i\varepsilon - L_k)^{-1} \right] \omega_k^{in}(r) dc. \quad (2.6)$$

Using (2.2), the function

$$\Delta_k \tilde{\Phi}_k = (z - L_k)^{-1} \omega_k^{in}(r) \quad r \in [0, \infty), z \in \mathbb{C}$$

satisfies the so-called *inhomogeneous Rayleigh equation* (explicitly writing out Δ_k):

$$\left(\partial_{rr} + \frac{1}{r} \partial_r - \frac{k^2}{r^2} \right) \tilde{\Phi}_k + \frac{\beta(r)}{u(r) - z} \tilde{\Phi}_k = \frac{\omega_k^{in}(r)}{u(r) - z}. \quad (2.7)$$

Note that for $z = c \pm i\varepsilon$ with $\varepsilon > 0$ (2.7) is a smooth perturbation of Δ_k and hence we will not have qualitative smoothness or local-integrability problems for $\varepsilon > 0$. Such difficulties arise only in the limit $\varepsilon \rightarrow 0$. By replacing $\tilde{\Phi}_k$ with

$$\Phi_k = \sqrt{r} \tilde{\Phi}_k$$

to eliminate the first-order derivative term, we obtain

$$\text{RAY}_z \Phi_k(r, z) = \frac{\omega_k^{in}(r) \sqrt{r}}{u(r) - z}, \quad \text{RAY}_z := \partial_{rr} + \frac{1/4 - k^2}{r^2} + \frac{\beta(r)}{u(r) - z}, \quad (2.8)$$

supplemented with the boundary conditions that Φ_k vanishes as $r \rightarrow 0$ and ∞ (asymptotic analysis shows that $\Phi_k(r, z) \sim r^{k+1/2}$ as $r \rightarrow 0$ and $\Phi_k(r, z) \sim r^{1/2-k}$ as $r \rightarrow \infty$ provided that $z \neq u(0)$).

In what follows, normally we will set $z = c \pm i\varepsilon$ and suppress the c, ε dependence to write

$$\text{RAY}_{\pm} := \partial_{rr} + \frac{1/4 - k^2}{r^2} + \frac{\beta(r)}{u(r) - c \mp i\varepsilon}.$$

Finally, from (2.6) we deduce that

$$\begin{aligned} \omega_k(t, r) &= \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2\pi i} \int_{\mathbb{R}} e^{-ikct} \left[\frac{\omega_k^{in}(r) - \beta(r) \tilde{\Phi}_k(r, c - i\varepsilon)}{u(r) - c + i\varepsilon} \right. \\ &\quad \left. - \frac{\omega_k^{in}(r) - \beta(r) \tilde{\Phi}_k(r, c + i\varepsilon)}{u(r) - c - i\varepsilon} \right] dc \\ &= e^{-iku(r)t} \omega_k^{in}(r) + \frac{\beta(r)}{2\pi i \sqrt{r}} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}} e^{-ikct} \left[\frac{\Phi_k(r, c + i\varepsilon)}{u(r) - c - i\varepsilon} \right. \\ &\quad \left. - \frac{\Phi_k(r, c - i\varepsilon)}{u(r) - c + i\varepsilon} \right] dc. \end{aligned} \quad (2.9)$$

Remark 2.5 The vorticity depletion mechanism for both shear flows and radial vortices is related to the second order singularity in the Rayleigh operator (2.8) at the critical point $r_c = u^{-1}(c)$ and at $r = 0$. However, unlike the shear flow case, here an additional challenge arises due to the collision of r_c with the singularity in Δ_k at $r = 0$.

In this work, vorticity depletion is isolated through a detailed analysis of the fundamental solution of the Rayleigh equation (2.8), especially evident in the estimate on the Wronskian, which has a clear singularity near the origin (see Lemma 5.2). This, in turn, is deduced with a detailed analysis of the behavior near the ‘critical layer’ of the fundamental solution of (2.8) as $r_c \rightarrow 0$ that depends heavily on the detailed monotonicity properties deduced in Sect. 4.6 below. As it is such a subtle effect to capture mathematically, we are currently unaware of any simple heuristic or easily checkable structural condition that could help to find where a similar behavior could be expected in other settings in fluid mechanics or kinetic theory.

2.2 Outline of the Proof for $k = 1$

In this case, the proof of Theorem 1.1 is based on the explicit formulas

$$\begin{aligned} f_{1;1}(r) &= \omega_1^{in}(r) + \frac{\beta(r)}{r^2 u'(r)} \int_0^r \omega_1^{in}(\rho) \rho^2 d\rho, \\ f_{1;2}(r, t) &= r \beta(r) \int_r^\infty e^{i(u(r) - u(\rho))t} f_{1;1}(\rho) \frac{d\rho}{\rho u'(\rho)}, \end{aligned}$$

which can be obtained thanks to the explicit solution of the homogeneous Rayleigh problem for $k = 1$:

$$\phi(r, z) = r^{3/2}(u - z); \quad (2.10)$$

(c.f. (2.5) and see Sect. 3.1). The vorticity depletion effect is encoded in the property that

$$f_{1;1}(r) = O(r^3) \quad \text{as } r \rightarrow 0$$

instead of just an $O(r)$ behavior, while

$$f_{1;2}(t, r) = O\left(\frac{r}{\langle t \rangle}\right) \quad \text{as } r \rightarrow 0,$$

which vanishes slower at the origin, but decays in time. The rigorous arguments needed to complete the proof of Theorem 1.1 for $k = 1$ are carried out in Sect. 3.2.

2.3 Outline of the Proof for $k \geq 2$

The case when $k \geq 2$ presents fundamental differences compared to the case when $k = 1$. To simplify notation, we omit the dependence on k of the functions involved except when it is relevant.

2.3.1 Depletion Trick and Contour Integral Decomposition

The first step of the proof is to isolate the asymptotic expansion at the origin from the rest of the profile. For this, we will apply the following trick, in which we remove a harmonic function (with a smooth cutoff) from Φ . Besides the function F in (1.9), we define

$$Y(r, z) := \Phi(r, z) - \frac{\chi(r)r^{k+1/2}}{\beta(0)}\omega_{k,0}^{in}, \quad (2.11)$$

$$F_*(r) := -\frac{1}{\beta(0)}\left((2k+1)\chi'(r)r^{k-1/2} + \chi''(r)r^{k+1/2}\right)\omega_{k,0}^{in}, \quad (2.12)$$

where we recall that χ is a smooth, non-negative function which satisfies $\chi(r) = 1$ for $|r| < 1/2$ and 0 for $|r| > 3/4$. From (2.8), we deduce that

$$\text{RAY}_z Y = \frac{F(r)}{u(r) - z} + F_*(r). \quad (2.13)$$

Going back to (2.11), we have from (2.9) that the *profile*

$$f(t, r) = e^{iku(r)t}\omega(t, r),$$

satisfies

$$f(t, r) = \frac{F(r)}{\sqrt{r}} + \frac{\beta(r)}{2\pi i \sqrt{r}} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}} e^{ik(u(r)-c)t} \left[\frac{Y(r, c+i\varepsilon)}{u(r)-c-i\varepsilon} - \frac{Y(r, c-i\varepsilon)}{u(r)-c+i\varepsilon} \right] dc. \quad (2.14)$$

Next, we sub-divide the integrals in (2.14) in several natural pieces. First, we isolate the contributions near and far from the spectrum (2.14) via the smooth cut-off function

$$\chi_\sigma(c) = \begin{cases} 1 & -R_\delta/2 \leq c \leq u(0) + 1/2 \\ 0 & c < -R_\delta \\ 0 & c > u(0) + 3/4, \end{cases} \quad (2.15)$$

where $R_\delta > 0$ will be fixed later (depending on δ), so that

$$\begin{aligned} f(t, r) = & \frac{F(r)}{\sqrt{r}} + \frac{\beta(r)}{2\pi i \sqrt{r}} \lim_{\varepsilon \rightarrow 0^+} \int_{-R_\delta}^{u(0)+1} e^{ik(u(r)-c)t} \left[\frac{Y(r, c+i\varepsilon)}{u(r)-c-i\varepsilon} \right. \\ & \left. - \frac{Y(r, c-i\varepsilon)}{u(r)-c+i\varepsilon} \right] \chi_\sigma(c) dc \\ & + \frac{\beta(r)}{2\pi i \sqrt{r}} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}} e^{ik(u(r)-c)t} \left[\frac{Y(r, c+i\varepsilon)}{u(r)-c-i\varepsilon} \right. \\ & \left. - \frac{Y(r, c-i\varepsilon)}{u(r)-c+i\varepsilon} \right] (1 - \chi_\sigma(c)) dc. \end{aligned} \quad (2.16)$$

Further, define

$$r_c := u^{-1}(c), \quad c \in (0, u(0)). \quad (2.17)$$

The region $r \sim r_c$ is known in the classical fluid mechanics literature as the *critical layer* [28]. Near the points $c = 0$ and $c = u(0)$ there are a variety of subtleties in the resolvent. This can be expected due to the change in the nature of the singularities in the Rayleigh equation (2.8) at these points. Since the first-order singular point at the critical layer merges with the second order singularities at $r = 0, \infty$, the influence of ε in (2.8) will be felt much more, hence, we need to pass to the limit $\varepsilon \rightarrow 0$ in a *non-uniform* way over the spectrum $c \in [0, u(0)]$. From Lemma 2.1 and considering the points where $(u(0) - c)^2 \approx \varepsilon^2$ and $c^2 \approx \varepsilon^2$, we see that the natural place to divide the complex plane is along the curves $\varepsilon \approx r_c^2$ for $r_c \ll 1$ and $\varepsilon \approx r_c^{-2}$ for $r_c \gg 1$. A small, but crucial, point is that we can afford some flexibility in this boundary. Let $\alpha \in (0, \delta)$ be a parameter chosen sufficiently small in the sequel depending only on δ (from Theorem 1.1). Define the set

$$I_\alpha = \left\{ z = c \pm i\varepsilon \in \mathbb{C} : c \in (0, u(0)), \ k^5 \varepsilon \leq \min(r_c^{2+\alpha}, r_c^{-2-\alpha}) \right\} \quad (2.18)$$

and the associated smooth cut-off function

$$\chi_I(r_c) = \chi \left(\frac{r_c}{(k^5 \varepsilon)^{-\frac{1}{2+\alpha}}} \right) \left[1 - \chi \left(\frac{r_c}{(k^5 \varepsilon)^{\frac{1}{2+\alpha}}} \right) \right].$$

Then, we further divide the contour integral by

$$\begin{aligned}
 f(t, r) = & \frac{F(r)}{\sqrt{r}} + \frac{\beta(r)}{2\pi i \sqrt{r}} \lim_{\varepsilon \rightarrow 0^+} \int_0^{u(0)} e^{ik(u(r)-c)t} \left[\frac{Y(r, c + i\varepsilon)}{u(r) - c - i\varepsilon} \right. \\
 & \left. - \frac{Y(r, c - i\varepsilon)}{u(r) - c + i\varepsilon} \right] \chi_I(r_c) dc \\
 & + \frac{\beta(r)}{2\pi i \sqrt{r}} \lim_{\varepsilon \rightarrow 0^+} \int_{-R_\delta}^{u(0)+1} e^{ik(u(r)-c)t} \left[\frac{Y(r, c + i\varepsilon)}{u(r) - c - i\varepsilon} \right. \\
 & \left. - \frac{Y(r, c - i\varepsilon)}{u(r) - c + i\varepsilon} \right] \chi_\sigma(c) (1 - \chi_I(r_c)) dc \\
 & + \frac{\beta(r)}{2\pi i \sqrt{r}} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}} e^{ik(u(r)-c)t} \left[\frac{Y(r, c + i\varepsilon)}{u(r) - c - i\varepsilon} \right. \\
 & \left. - \frac{Y(r, c - i\varepsilon)}{u(r) - c + i\varepsilon} \right] (1 - \chi_\sigma(c)) dc. \tag{2.19}
 \end{aligned}$$

The first term will be thought of as “close to the spectrum”, whereas the latter two terms will be considered “far from the spectrum”. Given the singular integrals in the representation formula, the two relevant quantities appearing are

$$X(r, c, \varepsilon) = Y(r, c + i\varepsilon) - Y(r, c - i\varepsilon), \tag{2.20a}$$

$$A(r, c, \varepsilon) = Y(r, c + i\varepsilon) + Y(r, c - i\varepsilon) = X + 2Y(r, c - i\varepsilon) \tag{2.20b}$$

so that from (2.19) we write:

$$\begin{aligned}
 f(t, r) = & \frac{F(r)}{\sqrt{r}} + \frac{\beta(r)}{2\pi i \sqrt{r}} \lim_{\varepsilon \rightarrow 0^+} \int_0^{u(0)} \frac{i\varepsilon e^{ik(u(r)-c)t}}{(u(r) - c)^2 + \varepsilon^2} \chi_I(r_c) A(r, c, \varepsilon) dc \\
 & + \frac{\beta(r)}{2\pi i \sqrt{r}} \lim_{\varepsilon \rightarrow 0^+} \int_0^{u(0)} \frac{(u(r) - c) e^{ik(u(r)-c)t}}{(u(r) - c)^2 + \varepsilon^2} \chi_I(r_c) X(r, c, \varepsilon) dc \\
 & + \frac{\beta(r)}{2\pi i \sqrt{r}} \lim_{\varepsilon \rightarrow 0^+} \int_{-R_\delta}^{u(0)+1} e^{ik(u(r)-c)t} \left[\frac{Y(r, c + i\varepsilon)}{u(r) - c - i\varepsilon} \right. \\
 & \left. - \frac{Y(r, c - i\varepsilon)}{u(r) - c + i\varepsilon} \right] \chi_\sigma(c) (1 - \chi_I(r_c)) dc \\
 & + \frac{\beta(r)}{2\pi i \sqrt{r}} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}} e^{ik(u(r)-c)t} \left[\frac{Y(r, c + i\varepsilon)}{u(r) - c - i\varepsilon} \right. \\
 & \left. - \frac{Y(r, c - i\varepsilon)}{u(r) - c + i\varepsilon} \right] (1 - \chi_\sigma(c)) dc. \tag{2.21}
 \end{aligned}$$

See Fig. 1 for a summary of how the limiting procedure in (2.21) is carried out below.

There is one additional decomposition necessary in order to see the vorticity depletion effect—the decomposition in f_1 and f_2 . While $F(r)r^{-1/2}$ has better decay at the origin than $\omega_k^{in}(r)$, it is clear that for, e.g. $r_c = 1$, both $A(r, c, \varepsilon)r^{-1/2}$ and

$X(r, c, \varepsilon)r^{-1/2}$ can be expected to have the same asymptotic expansion at the origin as the solution of the Laplace equation (specifically, $\sim r^k$). Hence, at any fixed t , clearly f cannot have better decay at the origin (qualitatively speaking) than $\omega_k^{in}(r)$. We instead deduce that the leading asymptotic expansion is decaying in time. We divide the contribution from X in two pieces, by means of

$$\chi_1(r, r_c) = [1 - \chi(r/2)\chi(r/r_c)]\chi_I(r_c), \quad \chi_2(r, r_c) = \chi(r/2)\chi(r/r_c)\chi_I(r_c).$$

Note that $\chi(r/2)\chi(r/r_c) = 1$ when $r \leq 1$ and $r_c \geq 2r$. In this way, we can rewrite (2.21) as

$$f(t, r) = \lim_{\varepsilon \rightarrow 0} [f_1^\varepsilon(t, r) + f_2^\varepsilon(t, r) + f_S^\varepsilon(t, r) + f_E^\varepsilon(t, r)], \quad (2.22)$$

where

$$f_1^\varepsilon(t, r) = \frac{F(r)}{\sqrt{r}} + \frac{\beta(r)}{2\pi i \sqrt{r}} \int_0^{u(0)} \frac{i\varepsilon e^{ik(u(r)-c)t}}{(u(r)-c)^2 + \varepsilon^2} \chi_I(r_c) A(r, c, \varepsilon) dc \\ + \frac{\beta(r)}{2\pi i \sqrt{r}} \int_0^{u(0)} \frac{(u(r)-c)e^{ik(u(r)-c)t}}{(u(r)-c)^2 + \varepsilon^2} \chi_1(r, r_c) X(r, c, \varepsilon) dc, \quad (2.23)$$

$$f_2^\varepsilon(t, r) = \frac{\beta(r)}{2\pi i \sqrt{r}} \int_0^{u(0)} \frac{(u(r)-c)e^{ik(u(r)-c)t}}{(u(r)-c)^2 + \varepsilon^2} \chi_2(r, r_c) X(r, c, \varepsilon) dc, \quad (2.24)$$




Fig. 1 This figure summarizes the limiting procedure used to treat (2.21). Region III represents I_α . The contribution from this region is further decomposed into f_1^ε and f_2^ε (see (2.23) and (2.24) below) which converge to the decomposition in (1.13). The limiting procedure is done by constructing the Green's function for RAY_z and making analyses of the resulting integral operators (carried out in Sects. 4–6 together with “Appendix B”). In regions I and IV we apply energy estimates on RAY_z to prove these contributions vanish (carried out in “Appendix D”). Here $\delta > 0$ is traded to gain the flexibility to take $\alpha > 0$. In region II, we apply a compactness-contradiction argument with a second order comparison principle that shows these contributions also must vanish (carried out in “Appendix D.1.2”)

$$f_S^\varepsilon(t, r) = \frac{\beta(r)}{2\pi i \sqrt{r}} \int_{-R_\delta}^{u(0)+1} e^{ik(u(r)-c)t} \left[\frac{Y(r, c+i\varepsilon)}{u(r)-c-i\varepsilon} - \frac{Y(r, c-i\varepsilon)}{u(r)-c+i\varepsilon} \right] \\ \times \chi_\sigma(c)(1-\chi_I(r_c))dc, \quad (2.25)$$

$$f_E^\varepsilon(t, r) = \frac{\beta(r)}{2\pi i \sqrt{r}} \int_{\mathbb{R}} e^{ik(u(r)-c)t} \left[\frac{Y(r, c+i\varepsilon)}{u(r)-c-i\varepsilon} - \frac{Y(r, c-i\varepsilon)}{u(r)-c+i\varepsilon} \right] (1-\chi_\sigma(c))dc. \quad (2.26)$$

In order to prove Theorem 1.1 we moreover need to express $(r\partial_r)^j f$ for $j \leq k$. In what follows, denote ∂_G as:

$$\partial_G := \frac{1}{u'(r)} \partial_r + \partial_c. \quad (2.27)$$

It is important here that u' does not vanish. The significance of this derivative, which has the form $\partial_z + \partial_c$ with $z = u(r)$, is the following: formally integrating by parts in c assuming that no boundary terms appear gives:

$$r\partial_r \left(\int_0^\infty h(u(r)-c)B(r, c)dc \right) = \int_0^\infty h(u(r)-c)ru'(r)\partial_G B(r, c)dc.$$

The G refers to the terminology of “good derivative” in [66], where an analogous derivative arises for a similar reason. Iterating gives,

$$(r\partial_r)^j f_1^\varepsilon(t, r) = (r\partial_r)^j F(r) + \frac{1}{2\pi i} \int_0^\infty \frac{2i\varepsilon u'(r_c) e^{itk(u(r)-u(r_c))}}{(u(r)-c)^2 + \varepsilon^2} \\ \times (ru'(r)\partial_G)^j \left(\chi_I(r_c) \frac{\beta(r)}{\sqrt{r}} A(r, c, \varepsilon) \right) dr_c \quad (2.28a)$$

$$+ \frac{1}{2\pi i} \int_0^\infty e^{itk(u(r)-u(r_c))} \frac{(u(r)-c)u'(r_c)}{(u(r)-c)^2 + \varepsilon^2} \\ \times (ru'(r)\partial_G)^j \left(\chi_1(r, r_c) \frac{\beta(r)}{\sqrt{r}} X(r, c, \varepsilon) \right) dr_c, \quad (2.28b)$$

$$(r\partial_r)^j f_2^\varepsilon(t, r) = \frac{1}{2\pi i} \int_0^\infty e^{itk(u(r)-u(r_c))} \frac{(u(r)-c)u'(r_c)}{(u(r)-c)^2 + \varepsilon^2} \\ \times (ru'(r)\partial_G)^j \left(\chi_2(r, r_c) \frac{\beta(r)}{\sqrt{r}} X(r, c, \varepsilon) \right) dr_c. \quad (2.28c)$$

This formula will be used below to obtain higher derivative estimates on f and f_j^ε . Finally, in order to complete our characterization of vorticity depletion, we obtain a decay estimate on $(r\partial_r)^j f_2^\varepsilon$ like $O(t^{-1})$. For this, we will integrate by parts in r_c :

$$(r\partial_r)^j f_2^\varepsilon(t, r) = -\frac{1}{2\pi kt} \int_0^\infty e^{itk(u(r)-u(r_c))} \partial_{r_c} \left(\frac{(u(r)-c)}{(u(r)-c)^2 + \varepsilon^2} \right. \\ \left. \times (ru'(r)\partial_G)^j \left(\chi_2(r, r_c) \frac{\beta(r)}{\sqrt{r}} X(r, c, \varepsilon) \right) \right) dr_c. \quad (2.29)$$

Notice that in the formulas above, near $r = 0$, the derivatives landing on X or A will be roughly $O(r^{2\ell})\partial_G^\ell$; we will see that each ∂_G ‘costs’ r^{-2} near the origin. Indeed, we have the following observation regarding F and F_* :

Lemma 2.6 *There holds*

$$\left\| \partial_G^\ell F \right\|_{L_{F,\delta+2\ell}^2} \lesssim_\ell \sum_{m=1}^{\ell} \left\| (r\partial_r)^m F \right\|_{L_{F,\delta}^2}, \quad \left\| \partial_G^\ell F_* \right\|_{L_{F,\delta+2\ell}^2} \lesssim_\ell k^{1+\ell} \left| \omega_{k,0}^{in} \right|.$$

2.3.2 Vanishing for $z \notin I_\alpha$ as $\varepsilon \rightarrow 0$

The contributions of f_S^ε and f_E^ε (2.25) and (2.26) vanish as $\varepsilon \rightarrow 0$. Hence, the only relevant values of the spectral parameter c are those contained in the interval $[0, u(0)]$, the range of u , which is the continuous spectrum of the operator L_k in (2.2). The precise statement is contained in the following theorem.

Theorem 2.7 *Assume $k \geq 2$, and let $j \in \{0, \dots, k\}$ be a fixed integer, $\alpha < \delta/8$, and δ sufficiently small. Then for all $t \in \mathbb{R}$,*

$$\lim_{\varepsilon \rightarrow 0} \left[\left\| (r\partial_r)^j f_E^\varepsilon(ts, \cdot) \right\|_{L_{f,\delta}^2} + \left\| (r\partial_r)^j f_S^\varepsilon(t, \cdot) \right\|_{L_{f,\delta}^2} \right] = 0.$$

The proof of this theorem is contained in “Appendix D”. The main ingredient is a set of careful energy estimates on a slightly more generalized version of the Rayleigh problem (2.13), as stated in Theorem D.1. These estimates allow to trade some $\delta > 0$ for freedom to choose $\alpha > 0$ in (2.18). The estimates are then used in an iterative way, to bootstrap from the initial L^2 bound up to the k -th derivative. Indeed, the functions X, Y, A and their ∂_G derivatives satisfy an equation of the type

$$\text{RAY}_\pm \partial_G^{j+1} Y(\cdot, c \pm i\varepsilon) = \frac{F_{j+1}^\pm}{u(r) - c \mp i\varepsilon} + R_{j+1}, \quad (2.30)$$

where R_j and F_j depend on F, F_* and lower order derivatives of Y . The difficulties we face are summarized below.

◇ The energy estimates depend on the region where c ranges, and, in turn, on the asymptotic expansion of u near $r = 0$ and $r = \infty$. Hence, the bounds are necessarily different and have to be treated on a case-by-case basis. The non-uniformity in which $\varepsilon \rightarrow 0$ discussed in the previous section plays a key role. Moreover, bounds have to encode the correct integrability in c , as the formula (2.25) deals with the endpoints $c = 0, u(0)$, while (2.26) requires integrability as $c \rightarrow \pm\infty$.

◇ While in most cases a (weighted) energy estimate for (2.30) is obtainable by multiplication by Y_ε and integration by parts, the case near $c = 0$ requires a contradiction argument. Due to compactness, a failure of the energy estimate would imply the existence of a localized solution to $\text{RAY}_\pm \phi = 0$, which is ruled out by a second order comparison principle against the homogeneous solution at $k = 1$ associated with (2.5).

- ◇ In the iterative process, R_j contains coefficients (that depend on u and β) that are very singular and that require a gain of r^2 at $r = 0$. This is related to the “cost” of taking ∂_G derivatives, discussed above. This loss can be handled up to $k - 1$ derivatives, by keeping track of the weight correction allowed in the energy estimates (the parameter γ in Theorem D.1). The case $j = 0$ is carried out in detail in Sects. D.3–D.4 in “Appendix”, while the generalization to derivatives up to $k - 1$ is handled in Sect. D.5.1 in “Appendix”.
- ◇ The k -th derivative is handled directly by expanding the Rayleigh operator in (2.30) and exploiting the elliptic regularity due to the second order derivative present in RAY_\pm (see Sect. D.5.2 in “Appendix”).

2.3.3 Green’s Function for the Rayleigh Problem for $z \in I_\alpha$ as $\varepsilon \rightarrow 0$

2.3.3.1 Homogeneous Rayleigh Problem While for $k = 1$ the exact solution (2.5) allows the construction of the Green’s function in a fairly direct way (for all $z \in \mathbb{C}$), the picture in the case $k \geq 2$ is complicated by the lack of explicit formulae for the solution to the homogeneous Rayleigh problem $\text{RAY}_\pm \phi = 0$. In Sect. 4, we derive the existence of a unique solution ϕ of the form

$$\phi(r, z) = P(r, z)(u(r) - z), \quad z \in I_\alpha,$$

which satisfies $P(r_c, z) = 1, \partial_r P(r_c, z) = 0$. The function P also encodes the behavior of ϕ away from the critical layer (essentially, the precise asymptotics as $r \rightarrow 0$ and $r \rightarrow \infty$). Theorem 4.1 treats the general case when $z \in I_\alpha$, while Theorem 4.2 focuses on further properties when $z \in (0, u(0))$ is real-valued. The convergence estimates are stated in Theorem 4.3. The proofs are articulated in different steps.

◇ Existence and uniqueness of P is proved through an auxiliary function \tilde{P} , related to P via

$$P(r, z) = \left(\frac{r}{r_c}\right)^{3/2} \tilde{P}(r, z),$$

which satisfies a proper integral equation, treatable under the contraction mapping principle in weighted L^∞ -based spaces (note this step shares some similarity with [66]). In this way, existence, uniqueness, and the nearly correct behavior at 0 and ∞ (up to a small correction) in both r and r_c is obtained at once, along with suitable bounds (Proposition 4.6). Note that the definition of \tilde{P} is informed by the exact solution (2.10) in the case $k = 1$.

◇ When $z \in (0, u(0))$ is real, further monotonicity properties of P , are available (see Theorem 4.2). Of extreme importance is the fact that ϕ only vanishes at the critical layer, when $r = r_c$, along with the correct k -dependence of the estimates involved.

◇ In order to prove that the complex solution $\phi(r, z)$ only vanishes at the critical layer for every $z \in I_\alpha$, we prove convergence estimates for $P(r, c \pm i\varepsilon) - P(r, c)$ and its various derivatives, in the correct L^∞ -weighted spaces. This is carried out in various steps. In Sect. 4.2, we use again the function \tilde{P} , and we derive sub-optimal (in both r and r_c) convergence estimates. Similarly, we treat $\partial_G \tilde{P}$ near the critical layer in Sect. 4.3 and $\partial_{r_c} \tilde{P}$ in Sect. 4.4. The optimality in r is then obtained in Sect. 4.5. In both

cases, factors related to r_c^2 and r_c^{-2} appear in the convergence estimates, due to the nature of the singularities at $r, r_c = 0, \infty$. This is the primary practical motivation behind the definition of the region I_α , and the subsequent non-uniform passage to the limit as $\varepsilon \rightarrow 0$. Indeed, only for $z \in I_\alpha$ are we able to deduce that $\phi(r, z) \neq 0$ for all $r \neq r_c$, a key property (as we see from (2.31) below).

2.3.3.2 Inhomogeneous Rayleigh Problem In order to construct a Green's function for RAY_z from (2.8), we again use reduction of order. For $z \in I_\alpha \setminus \text{Ran } u$, we define the two homogeneous solutions which each satisfy one of the boundary conditions via:

$$H_0(r, z) = -\phi(r, z) \int_0^r \frac{1}{\phi(s, z)^2} ds, \quad H_\infty(r, z) = \phi(r, z) \int_r^\infty \frac{1}{\phi(s, z)^2} ds. \quad (2.31)$$

Note that H_0 and H_∞ are well defined by absolutely convergent integrals for $z \in I_\alpha \setminus \text{Ran } u$ and are solutions to the homogeneous Rayleigh equation (4.1), whose Wronskian is

$$M(z) := H_0(r, z) \partial_r H_\infty(r, z) - H_\infty(r, z) \partial_r H_0(r, z) = \int_0^\infty \frac{1}{\phi^2(s, z)} ds. \quad (2.32)$$

One of the crucial lemmas is Lemma 5.2, which provides the following lower bound uniformly for $z \in I_\alpha$,

$$|M(c \pm i\varepsilon)| \gtrsim k \max(r_c^{-3}, r_c^5).$$

Note that the singularities at $r_c \rightarrow 0$ and $r_c \rightarrow \infty$ are in fact a gain. These gains are crucial for obtaining the vorticity depletion, and arise from the detailed asymptotic analysis of ϕ around the critical layer. Hence, the Green's function for the Rayleigh operator for $z \in I_\alpha \setminus \text{Ran } u$ is

$$\mathcal{G}(r, r', z) = \frac{1}{M(z)} \begin{cases} H_0(r, z) H_\infty(r', z), & r < r', \\ H_0(r', z) H_\infty(r, z), & r > r'. \end{cases} \quad (2.33)$$

In addition to the lower bound on M , precise estimates on H_0 and H_∞ follow from our study of $\phi(r, z)$ (see Sect. 5) as well as convergence as $\varepsilon \rightarrow 0$. Due to the apparently singular integrals that appear as $\varepsilon \rightarrow 0$ in (2.31), it is not obvious that $H_0(r, c \pm i\varepsilon)$ and $H_\infty(r, c \pm i\varepsilon)$ converge, but cancellations for $\varepsilon > 0$ ensure we have well-defined, log-Lipschitz limiting functions $H_0(r, c \pm i0)$, $H_\infty(r, c \pm i0)$ (as expected from shear flows [28]). See Sect. 5 for more details.

The Green's function gives us representation formulas for $Y^\pm := Y(r, c \pm i\varepsilon)$ and $X := X(r, c, \varepsilon)$ for $\varepsilon > 0$ (see Lemma 6.3),

$$\begin{aligned} Y^\pm &= \int_0^\infty \mathcal{G}(r, s, c \pm i\varepsilon) \frac{u(s) - c}{(u(s) - c)^2 + \varepsilon^2} F(s) ds \\ &\quad \pm \int_0^\infty \mathcal{G}(r, s, c \pm i\varepsilon) \frac{i\varepsilon}{(u(s) - c)^2 + \varepsilon^2} F(s) ds + \int_0^\infty \mathcal{G}(r, s, c \pm i\varepsilon) F_*(s) ds \end{aligned} \quad (2.34a)$$

$$\begin{aligned}
X &= \int_0^\infty B_{X\delta;\varepsilon}^{(1)}(r, s, c) \frac{2i\varepsilon}{(u(s) - c)^2 + \varepsilon^2} F(s) ds \\
&+ \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u(s_0) - c)^2 + \varepsilon^2} B_{XS;\varepsilon}^{(1)}(r, s_0, c) B_{XS;\varepsilon}^{(2)}(s_0, s, c) ds_0 \right) \\
&\times \frac{u(s) - c}{(u(s) - c)^2 + \varepsilon^2} F(s) ds \\
&+ \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u(s_0) - c)^2 + \varepsilon^2} B_{XG;\varepsilon}^{(1)}(r, s_0, c) B_{XG;\varepsilon}^{(2)}(s_0, s, c) ds_0 \right) F_*(s) ds,
\end{aligned} \tag{2.34b}$$

where

$$\begin{aligned}
B_{X\delta;\varepsilon}^{(1)}(r, s, c) &= \mathcal{G}(r, s, c + i\varepsilon) \\
&+ \int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u(s_0) - c)^2 + \varepsilon^2} \mathcal{G}(r, s_0, c + i\varepsilon) \mathcal{G}(s_0, s, c - i\varepsilon) ds_0, \\
B_{XG;\varepsilon}^{(1)} &= B_{XS;\varepsilon}^{(1)}(r, s_0, c) = \mathcal{G}(r, s_0, c + i\varepsilon).
\end{aligned}$$

2.3.4 Representation Formulas and Boundedness for $(r\partial_r)^j f_1$ and $(r\partial_r)^j f_2$

The details of the below discussion are given in Sect. 6 below.

2.3.4.1 Iteration scheme and representation formulas for ∂_G^j derivatives for $j \leq k - 1$ From (2.28), we see that a key step in the proof of Theorem 1.1 is estimating $\partial_G^j X$ and $\partial_G^j A$ for $z \in I_\alpha$. The crucial property of ∂_G derivatives is that they vanish on functions of $u - c$, and hence, the commutator $[\partial_G, \text{RAY}_z]$ is not more singular than RAY_z itself at the critical layer (see (6.2) below). As a result, we are able to use an iteration scheme of the following general form to control higher derivatives; this iteration scheme is one of the insights for obtaining higher regularity of the profile.

Lemma 2.8 (Iteration lemma for $\partial_G^j X$ and $\partial_G^j Y$ for $\varepsilon > 0$). *For $F_j, R_j, R_j^x, \mathcal{E}_j$ defined below in Lemma 6.1 we have the iteration*

$$\begin{aligned}
\text{RAY}_+ \partial_G^{j+1} X &= \frac{2i\varepsilon}{(u - c)^2 + \varepsilon^2} \left(F_{j+1}^- - \beta \partial_G^{j+1} Y^- \right) + \frac{\mathcal{E}_{j+1}}{u - c - i\varepsilon} + R_{j+1}^x \\
\text{RAY}_\pm \partial_G^{j+1} Y^\pm &= \frac{F_{j+1}^\pm}{u - c \mp i\varepsilon} + R_{j+1}^\pm.
\end{aligned}$$

The quantities $R_j, \mathcal{E}_j, R_j^x$, and F_j depend only on $\partial_G^\ell X$ and $\partial_G^\ell Y$ for $0 \leq \ell \leq j - 1$ (as well as u, β, F and F_*).

Using the recursion scheme described in (2.35) and Fubini's theorem, it is not hard to formally verify the following Proposition by induction, which allows to directly express $\partial_G^j X$ and $\partial_G^j Y$ in terms of F and F_* in a form essentially the same as (2.34) except with much more complicated kernels. See Lemmas 6.4–6.6 below.

Proposition 2.9 *For all $j \leq k - 1$, there hold representation formulas of the general form for various kernels B and weights w for $\varepsilon > 0$:*

$$\begin{aligned} \partial_G^j Y &= \sum_{\ell=0}^j \int_0^\infty B_{YS;j,\ell}(r, s, c) \frac{u(s) - c}{(u(s) - c)^2 + \varepsilon^2} w_{S;j,\ell}(s) \partial_G^\ell F(s) ds \\ &+ \sum_{\ell=0}^j \int_0^\infty B_{Y\delta;j,\ell}(r, s, c) \frac{2i\varepsilon}{(u(s) - c)^2 + \varepsilon^2} w_{\delta;j,\ell}(s) \partial_G^\ell F(s) ds \\ &+ \sum_{\ell=0}^j \int_0^\infty B_{YG;j,\ell}(r, s, c) w_{G;j,\ell}(s) \partial_G^\ell F_*(s) ds \\ &+ \text{Similar terms with different } B, w, \end{aligned} \quad (2.36)$$

where by “Similar terms with different B, w ” we mean terms with exactly the same formal structure, except with different B kernels and weights w (however, all of the omitted terms will share the same estimates). Similarly, for various kernels B and weights w we have a similar representation formula for $\varepsilon > 0$:

$$\begin{aligned} \partial_G^j X &= \sum_{\ell=0}^j \int_0^\infty B_{X\delta 1;j,\ell}^{(1)}(r, s, c) \frac{2i\varepsilon}{(u(s) - c)^2 + \varepsilon^2} w_{X\delta 1;j,\ell}(s) \partial_G^\ell F(s) ds \\ &+ \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u(s_0) - c)^2 + \varepsilon^2} B_{X\delta;j,\ell}^{(1)}(r, s_0, c) B_{X\delta;j,\ell}^{(2)}(s_0, s, c) ds_0 \right) \\ &\times \frac{2i\varepsilon}{(u(s) - c)^2 + \varepsilon^2} w_{X\delta 2;j,\ell}(s) \partial_G^\ell F(s) ds \\ &+ \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u(s_0) - c)^2 + \varepsilon^2} B_{XS;j,\ell}^{(1)}(r, s_0, c) B_{XS;j,\ell}^{(2)}(s_0, s, c) ds_0 \right) \\ &\times \frac{(u(s) - c)}{(u(s) - c)^2 + \varepsilon^2} w_{XS;j,\ell}(s) \partial_G^\ell F(s) ds \\ &+ \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u(s_0) - c)^2 + \varepsilon^2} B_{XG;j,\ell}^{(1)}(r, s_0, c) B_{XG;j,\ell}^{(2)}(s_0, s, c) ds_0 \right) \\ &\times w_{XG;j,\ell}(s) \partial_G^\ell F_*(s) ds \\ &+ \text{Similar terms with different } B, w. \end{aligned} \quad (2.37)$$

Furthermore, in each term above, the weights satisfy an estimate of the following form for some $0 \leq \ell' \leq j - \ell$ and all $m \geq 0$ (different ℓ' for each term),

$$|(s \partial_s)^m w_{*;j,\ell}(s)| \lesssim_m \max(s^{-2\ell'}, s^{2\ell'}). \quad (2.38)$$

Along with (2.36) and (2.37), the proof of Proposition 2.9 derives also an associated recursion for the various B kernels appearing above (see Lemmas 6.4–6.6). That is, the kernels $B_{*,j,\ell}^{(*)}$ are determined from $B_{*,j-1,m}^{(*)}$ via a few canonical integral operators involving the Green's function \mathcal{G} . A crucial idea of our method is to use these recursion formulae on the B 's, together with precise estimates on \mathcal{G} , to obtain precise estimates on all possible B kernels by induction. This method allows us to treat all of $\partial_G^j X$ and $\partial_G^j Y$ for $j \leq k-1$ *simultaneously* (as discussed below, the method only stops due to the $\max(r^{-2}, r^2)$ losses coming from ∂_G). Define the bounding functions

$$\begin{aligned} \mathbb{K}(r, s, c) := & \mathbf{1}_{r_c > 1} + \mathbf{1}_{r_c \leq 1} \left(\mathbf{1}_{s < r < r_c} + \mathbf{1}_{s < r_c < r < 1} \frac{r_c^2}{r^2} + \mathbf{1}_{s < r_c < 1 < r} r_c^2 \right. \\ & + \mathbf{1}_{r_c < s < r < 1} \frac{s^2}{r^2} + \mathbf{1}_{r_c < s < 1 < r} s^2 + \mathbf{1}_{1 < s < r} \\ & + \mathbf{1}_{r < s < r_c} + \mathbf{1}_{r < r_c < s < 1} \frac{r_c^2}{s^2} + \mathbf{1}_{r < r_c < 1 < s} r_c^2 + \mathbf{1}_{r_c < r < s < 1} \frac{r^2}{s^2} \\ & \left. + \mathbf{1}_{r_c < r < 1 < s} r^2 + \mathbf{1}_{1 < r < s} \right), \end{aligned} \quad (2.39a)$$

$$\mathcal{B}(r, s) := \left(\mathbf{1}_{s < r} \frac{s^{k-1/2}}{r^{k-1/2}} + \mathbf{1}_{r < s} \frac{r^{k+1/2}}{s^{k+1/2}} \right) \langle s \rangle^4 \quad (2.39b)$$

$$\mathcal{L}_{J,\ell}(r, s) := k^J \max \left(\frac{1}{r^2}, \frac{1}{s^2}, r^2, s^2 \right)^\ell. \quad (2.39c)$$

The full properties and the estimates obtained on the kernels are laid out in Definitions 6.7 and 6.8 below. The main result in Sect. 6.3.1 is the following.

Proposition 2.10 *For $j \leq k-1$, each of the kernels $B_{X*,j,\ell}^{(1)}$ appearing in (2.36) and (2.37) is Suitable $(2\ell'', \ell'' + \eta, \gamma)$ of type I for some $\gamma \in (0, 1)$, $1 \gg \eta > 0$, and some integer $\ell'' \geq 0$ (difference for each kernel). For $j \leq k-1$, each of the kernels $B_{X*,j,\ell}^{(2)}$ and $B_{Y*,j,\ell}$ appearing in (2.36) and (2.37) is Suitable $(2\ell'', \ell'' + \eta, \gamma)$ of type II for some $\gamma \in (0, 1)$, $1 \gg \eta > 0$, and some integer $\ell'' \geq 0$ (difference for each kernel). In particular, each satisfies the uniform-in- ε boundedness:*

$$\begin{aligned} \left| B_{*,j,\ell}^{(1)}(r, s, c) \right| & \lesssim_\eta |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{2\ell'', \ell'' + \eta}(r, s) \\ \left| B_{*,j,\ell}^{(2)}(r, s, c) \right| & \lesssim_\eta |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{2\ell'', \ell'' + \eta}(r, s), \end{aligned}$$

and each of the kernels satisfies analogous log-Lipschitz regularity estimates and convergence estimates as $\varepsilon \rightarrow 0$ (see Lemmas 6.10–6.14). Finally, all terms in Proposition 2.9 satisfy the additional constraint

$$\ell + \ell' + \ell'' \leq j, \quad \ell + \ell' + \ell''_1 + \ell''_2 \leq j. \quad (2.40)$$

Remark 2.11 The gains encoded by \mathbb{K} , the fact that in certain regions of (r, s, c) -space \mathbb{K} is much smaller than 1, are what ultimately allows us to deduce the vorticity depletion effect and are inherited from precise estimates on \mathcal{G} . The losses encoded by \mathcal{L} are inherited from the $\max(r^{-2}, r^2)$ losses inherent in ∂_G derivatives. It is crucial that the losses in \mathcal{L} do not depend directly on r_c .

Remark 2.12 The constraint (2.40) arises due to the fact that each application of ∂_G to (2.35) can land either on X, Y, F, F_* , or on the fixed coefficients that depend only on the background vortex. Each application will lose $\max(r^{-2}, r^2)$, and hence there are $\max(r^{-2}, r^2)^j$ powers to distribute between different factors that the kernels and the weights account for.

2.3.4.2 Extension to ∂_G^k or $r_c \partial_{r_c} \partial_G^j$ with $j \leq k - 1$ It is not clear how to obtain estimates for even a single r_c derivative, e.g. $\partial_{r_c} \partial_G^j$: the commutator $[\partial_{r_c}, \text{RAY}_z]$ is too singular near r_c to use an approach similar to the one we used on ∂_G . Moreover, the arguments of Proposition 2.10 break down at $j = k$ due to the singularities in the right-hand side of (2.35) at zero and infinity (encoded by the constraint (2.40)). In order to overcome this difficulty, first notice that while the commutator $[\partial_r, \text{RAY}_z]$ is too singular to use an approach like what we used on ∂_G , we should nevertheless expect to be able to estimate $\partial_r \partial_G^j Y$ by elliptic regularity. Indeed, away from the critical layer, it is a straightforward extension of our methods to show directly that ∂_r derivatives of (2.36) and (2.37) should not be significantly worse than the $\partial_G^j X$ and $\partial_G^j Y$ themselves. The next observation is that, just as ∂_G arises when taking ∂_r derivatives in (2.28), similarly, ∂_G arises when taking ∂_{r_c} (or ∂_G) derivatives of (2.37) and (2.36) (see Sect. 6.4 for details). Therefore, while it seems intractable to build a reasonable iteration scheme for taking multiple ∂_r and ∂_{r_c} , it turns out we can take a *single* $\partial_r, \partial_{r_c}$ away from the critical layer, or a single additional ∂_G derivative near the critical layer, of (2.36) and (2.37). For f_2 , we only need ∂_{r_c} away from $r \sim r_c$ and hence this will be sufficient. For f_1 , away from $r \sim r_c$ we write

$$ru'(r)\partial_G = r\partial_r + \frac{ru'(r)}{r_c u'(r_c)} r_c \partial_{r_c},$$

and estimate these two derivatives separately (whereas for $r \sim r_c$ we naturally leave the derivative as is). See Sect. 6.4 for details.

2.3.4.3 Convergence and boundedness of $(r\partial_r)^j f_1^\varepsilon$ and $(r\partial_r)^j f_2^\varepsilon$ Next, our goal is to pass to the limit $\varepsilon \rightarrow 0$ and obtain L^2 bounds on $(r\partial_r)^j f_1$ and $(r\partial_r)^j f_2$. The first proposition gives convergence in the weaker weighted space $L_{f,\delta}^2$ and boundedness of the limit in the weighted space which is r^{-2} stronger at the origin.

Proposition 2.13 *For all $j \leq k$, we have the convergence of $(r\partial_r)^j f_1^\varepsilon(t, r)$ to a limit $(r\partial_r)^j f_1(t, r)$ in the norm:*

$$\lim_{\varepsilon \rightarrow 0} \left\| (r\partial_r)^j (f_1^\varepsilon - f_1) \right\|_{L_{f,\delta}^2} = 0.$$

Moreover, there holds the uniform-in- t bounds for $n \leq k$ in the stronger weighted space: for all $\eta > 0$,

$$\left\| \sqrt{r} (r \partial_r)^n f_1 \right\|_{L^2_{F,\delta}} \lesssim_{n,\delta,\eta} k^{2n+1} \left| \omega_{k,0}^{in} \right| + \sum_{j=0}^n k^{2(n-j)+\eta} \left\| (r \partial_r)^j F \right\|_{L^2_{F,\delta/4}}.$$

Similarly, we prove the requisite decay $O(t^{-1})$ of f_2 in the natural $L^2_{f,\delta}$ space.

Proposition 2.14 *For all $j \leq k-1$, we have the convergence of $(r \partial_r)^j f_2^\varepsilon(t, r)$ to a limit $(r \partial_r)^j f_2(t, r)$ in $L^2_{f,\delta}$:*

$$\lim_{\varepsilon \rightarrow 0} \left\| (r \partial_r)^j (f_2^\varepsilon - f_2) \right\|_{L^2_{f,\delta}} = 0.$$

Furthermore, there holds the following decay estimate for $n \leq k-1$: for all $\eta > 0$,

$$\left\| (r \partial_r)^n f_2 \right\|_{L^2_{f,\delta}} \lesssim_\delta \frac{1}{\langle kt \rangle} |k|^{2n+4+\eta} \left| \omega_{k,0}^{in} \right| + \frac{1}{\langle kt \rangle} \sum_{j=0}^{n+1} k^{2(n-j)+3+\eta} \left\| (r \partial_r)^j F \right\|_{L^2_{F,\delta/4}}.$$

Propositions 2.13 and 2.14 give the vorticity depletion characterization in (1.13). Combining (2.28) and Proposition 2.9, the proof of Propositions 2.13 and 2.14 reduces to passing to the limit as $\varepsilon \rightarrow 0$ in operators of the type arising in Proposition 2.9 (and obviously bounding the limiting operators) with the B 's satisfying a list of properties such as those alluded to in Proposition 2.10 (the actual list is much longer; see Sect. 6.3.1). This involves a number of very technical decompositions over (in general) four variables r, s_0, r_c, s adapted to the various asymptotic behaviors near the origin, infinity, and the critical layer. The various Hölder regularity properties of the kernels becomes important for passing to the limit in the iterated singular integral operators arising. The details are carried out in “Appendix B”.

Finally, via the Biot–Savart law, integration by parts in r , and the Hilbert–Schmidt lemma, Propositions 2.13 and 2.14, directly imply Theorem 1.1, as the next lemma shows.

Lemma 2.15 (Vorticity depletion implies optimal inviscid damping). *The vorticity depletion estimates (1.13) imply the inviscid damping estimates (1.11).*

Proof of Lemma 2.15 We may without loss of generality consider the case $|kt| \geq 1$. Denote $G_k(r, \rho)$ to be the Green's function for the Laplacian restricted to the k -th angular Fourier mode, i.e.

$$G_k(r, \rho) = \frac{\rho}{k} \begin{cases} \frac{\rho^k}{r^k} & \rho \leq r \\ \frac{r^k}{\rho^k} & \rho \geq r. \end{cases}$$

Using the decomposition of ω_k as in (1.12), we have

$$\begin{aligned}\psi_k(t, r) &= \int_0^\infty G_k(r, \rho) e^{-iktu(\rho)} f_{k;1}(t, \rho) d\rho + \int_0^\infty G_k(r, \rho) e^{-iktu(\rho)} f_{k;2}(t, \rho) d\rho \\ &:= \psi_{k;1}(t, r) + \psi_{k;2}(t, r).\end{aligned}$$

It is convenient to denote

$$\sup_{t \geq 0} \sum_{n=0}^2 k^{-n} \left\| \sqrt{r} (r \partial_r)^n f_{k;1}(t) \right\|_{L^2_{F,\delta}} + \sup_{t \geq 0} \langle kt \rangle \sum_{n=0}^1 k^{-n} \left\| (r \partial_r)^n f_{k;2}(t) \right\|_{L^2_{f,\delta}} = M_0$$

where by (1.13) we have that M_0 is bounded in terms of the datum ω_k^{in} . For the contribution from $\psi_{k;2}$, we integrate by parts and obtain

$$\begin{aligned}\langle kt \rangle \langle kt \rangle \frac{|\psi_{k;2}(t, r)|}{w_{\psi,2\delta}(r)} &= \langle kt \rangle \left| \int_0^\infty \frac{-i e^{-iktu(\rho)}}{w_{\psi,2\delta}(r)} \partial_\rho \left(\frac{1}{u'(\rho)} G_k(r, \rho) f_{k;2}(t, \rho) \right) d\rho \right| \\ &\leq \int_0^\infty T_2(r, \rho) \frac{\langle kt \rangle (|f_{k;2}(t, \rho)| + k^{-1} |(\rho \partial_\rho) f_{k;2}(t, \rho)|)}{w_{f,\delta}(\rho)} d\rho,\end{aligned}$$

where

$$T_2(r, \rho) := \frac{w_{f,\delta}(\rho) G_k(r, \rho)}{\rho |u'(\rho)| w_{\psi,2\delta}(r)} \left(\frac{|(\rho \partial_\rho) G_k(r, \rho)|}{G_k(r, \rho)} + \frac{k \rho |u''(\rho)|}{|u'(\rho)|} + k \right).$$

Using that (recall the strong decay imposed on f (1.10b) at infinity),

$$\begin{aligned}\|T_2(r, \rho)\|_{L^2(dr d\rho)} &\lesssim k \left\| \frac{w_{f,\delta}(\rho) G_k(r, \rho)}{\rho |u'(\rho)| w_{\psi,2\delta}(r)} \right\|_{L^2(dr d\rho)} \\ &\lesssim \left\| \frac{w_{f,\delta}(\rho) \min \left\{ \frac{\rho^k}{r^k}, \frac{r^k}{\rho^k} \right\}}{|u'(\rho)| w_{\psi,2\delta}(r)} \right\|_{L^2(dr d\rho)} \lesssim 1,\end{aligned}$$

we obtain,

$$\|\psi_{k;2}(t)\|_{L^2_{\psi,2\delta}} \lesssim \frac{M_0}{\langle kt \rangle \langle kt \rangle},$$

as desired. Similarly, for the $\psi_{k;1}$ contribution, we integrate by parts in ρ twice, keeping track of the boundary terms arising from the second derivative of the Green's function. We arrive at

$$\begin{aligned}(kt)^2 \psi_{k;1}(t, r) &= 2e^{-iktu(r)} \frac{f_{k;1}(t, r)}{(u'(r))^2} \\ &\quad - \int_0^\infty e^{-iktu(\rho)} \partial_\rho \left(\frac{1}{u'(\rho)} \partial_\rho \left(\frac{G_k(r, \rho)}{u'(\rho)} f_{k;1}(t, \rho) \right) \right) d\rho\end{aligned}$$

from which we deduce

$$\begin{aligned} & (kt)^2 \frac{|\psi_{k;1}(t, r)|}{w_{\psi, 2\delta}(r)} \\ & \leq \frac{2w_{F, \delta}(r)}{w_{\psi, 2\delta}(r)\sqrt{r}(u'(r))^2} \frac{\sqrt{r}|f_{k;1}(t, r)|}{w_{F, \delta}(r)} \\ & \quad + \int_0^\infty T_1(r, \rho) \frac{\sqrt{\rho} \left(|f_{k;1}(t, \rho)| + k^{-1} |(\rho \partial_\rho) f_{k;1}(t, \rho)| + k^{-2} |(\rho \partial_\rho)^2 f_{k;1}(t, \rho)| \right)}{w_{F, \delta}(\rho)} d\rho \end{aligned}$$

where

$$\begin{aligned} T_1(r, \rho) &= \frac{w_{F, \delta}(\rho) G_k(r, \rho)}{\rho^2 (u'(\rho))^2 \sqrt{\rho} w_{\psi, 2\delta}(r)} \left(\frac{|(\rho \partial_\rho)^2 G_k(r, \rho)|}{G_k(r, \rho)} \right. \\ &\quad \left. + k \left(3 + \frac{3\rho |u''(\rho)|}{|u'(\rho)|} \right) \frac{|\rho \partial_\rho G_k(r, \rho)|}{G_k(r, \rho)} + k^2 \right. \\ &\quad \left. + \frac{k^2 \rho^2 |u'''(\rho)|}{|u'(\rho)|} + \frac{3k^2 \rho^2 (u''(\rho))^2}{(u'(\rho))^2} + \frac{3k^2 \rho |u''(\rho)|}{|u'(\rho)|} \right) \\ &\lesssim k \frac{w_{F, \delta}(\rho) \min \left\{ \frac{\rho^k}{r^k}, \frac{r^k}{\rho^k} \right\}}{\rho^2 (u'(\rho))^2 \sqrt{\rho} w_{\psi, 2\delta}(r)}. \end{aligned}$$

Therefore, using that

$$\left\| \frac{w_{F, \delta}(r)}{w_{\psi, 2\delta}(r)\sqrt{r}(u'(r))^2} \right\|_{L^\infty(dr)} + \left\| \frac{w_{F, \delta}(\rho) \min \left\{ \frac{\rho^k}{r^k}, \frac{r^k}{\rho^k} \right\}}{\rho^{2+1/2} (u'(\rho))^2 w_{\psi, 2\delta}(r)} \right\|_{L^2(dr d\rho)} \lesssim 1$$

which may be checked directly, we obtain that

$$\|\psi_{k;1}(t)\|_{L^2_{\psi, 2\delta}} \lesssim \frac{M_0}{k^2 t^2}$$

which is the desired estimate. The inviscid damping of the velocity field follows in a similar manner from the Biot–Savart law, or by noting that $r(u_k^r, u_k^\theta) = (ik, -r\partial_r)\psi_k$, and we omit these details to avoid redundancy. \square

This completes the proof of Theorem 1.1.

3 Dynamics of the $k = 1$ Mode

In this section we give the proof of Theorem 1.1 for the mode $k = 1$. For this, we derive an equivalent formulation of (2.9) as follows. From (1.6a) we obtain that

$$\partial_t \left(e^{iku(r)t} \omega_k \right) = e^{iku(r)t} ik\beta(r) \psi_k.$$

An integration over time then yields

$$e^{iku(r)t} \omega_k(t, r) = \omega_k^{in}(r) + ik\beta(r) \int_0^t e^{iku(r)\tau} \psi_k(\tau, r) d\tau.$$

Now, by writing the evolution equation satisfied by ψ_k , we infer that

$$\psi_k(t, r) = \frac{\beta(r)}{2\pi i \sqrt{r}} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}} e^{-ikct} [\Phi_k(r, c - i\varepsilon) - \Phi_k(r, c + i\varepsilon)] dc,$$

and therefore

$$e^{iku(r)t} \omega_k(t, r) = \omega_k^{in}(r) + \frac{\beta(r)}{2\pi i \sqrt{r}} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}} \frac{1 - e^{ik(u(r)-c)t}}{u(r) - c} [\Phi_k(r, c + i\varepsilon) - \Phi_k(r, c - i\varepsilon)] dc. \quad (3.1)$$

We remark that this is roughly the form used in [66] (though the contour integral is set up slightly differently).

3.1 An Explicit Representation of the Vorticity Profile

The case $k = 1$ is special, because the homogeneous Rayleigh problem has an explicit solution

$$\phi(r, z) = (u(r) - z)r^{3/2} \text{ solves } \text{RAY}_z \phi = 0.$$

This fact may be verified by a direct computation, in view of (2.1). Additionally, this special solution has the property that

$$\lim_{r \rightarrow 0} \phi(r, z) = 0$$

and thus it may be used directly in the construction of the Green's function for RAY_z . This fact is yet another special property of $k = 1$. Using the reduction of order technique (see e.g. [50]), one obtains another independent solution to $\text{RAY}_z = 0$

$$H_\infty(r, z) = \phi(r, z) \int_r^\infty \frac{ds}{\phi(s, z)^2},$$

which vanishes as $r \rightarrow \infty$. One verifies that the Wronskian of these two solutions is

$$\phi \partial_r H_\infty - \partial_r \phi H_\infty = -1,$$

and thus we may directly combine ϕ and H_∞ to obtain the Green's function for the $k = 1$ Rayleigh operator:

$$G_1(r, \rho, z) = \begin{cases} -\phi(r, z)H_\infty(\rho, z), & r < \rho, \\ -\phi(\rho, z)H_\infty(r, z), & r > \rho. \end{cases} \quad (3.2)$$

The upshot of (3.2) is that the solution of the inhomogeneous Rayleigh problem for $k = 1$,

$$\text{RAY}_z \Phi_1 = \frac{\omega_1^{in}(r)\sqrt{r}}{u(r) - z}$$

for $z \in \mathbb{C}$, is given by

$$\Phi_1(r, z) = \int_0^\infty \mathcal{G}_1(r, \rho, z) \frac{\omega_1^{in}(\rho)\sqrt{\rho}}{u(\rho) - z} d\rho. \quad (3.3)$$

In order to derive a formula for the vorticity profile

$$f_1(t, r) = e^{iu(r)t} \omega_1(t, r)$$

we appeal to the representation formula (3.1) which yields

$$f_1(t, r) = \omega_1^{in}(r) + \frac{\beta(r)}{2\pi i \sqrt{r}} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}} \frac{1 - e^{i(u(r)-c)t}}{u(r) - c} [\Phi_1(r, c + i\varepsilon) - \Phi_1(r, c - i\varepsilon)] dc. \quad (3.4)$$

Further, setting $z = c \pm i\varepsilon$ in (3.3) and using essentially that $\int_0^\infty \omega_1^{in}(r)r^2 dr = 0$, we may pass to the $\varepsilon \rightarrow 0^+$ limit in the contour integral of (3.4), to obtain

$$f_1(t, r) = f_{1;1}(r) + f_{1;2}(r, t), \quad (3.5a)$$

$$f_{1;1}(r) = \omega_1^{in}(r) + \frac{\beta(r)}{r^2 u'(r)} \int_0^r \omega_1^{in}(\rho) \rho^2 d\rho, \quad (3.5b)$$

$$f_{1;2}(r, t) = r\beta(r) \int_r^\infty e^{i(u(r)-u(\rho))t} f_{1;1}(\rho) \frac{d\rho}{\rho u'(\rho)}. \quad (3.5c)$$

where as usual we used the notation $r_c = u^{-1}(c)$ for $c \in \text{Ran}(u) = (0, u(0)]$. The proof of the convergence as $\varepsilon \rightarrow 0^+$ of the expression in (3.4) to the expression in (3.5) is rather tedious, but direct. We thus omit these details. Alternatively, one may directly verify (by plugging in) that the vorticity

$$\omega_1(t, r) = e^{-iu(r)t} f_1(t, r) \quad (3.6)$$

obeys

$$\partial_t \omega_1 + iu(r)\omega_1 - i\beta(r)\psi_1 = 0, \quad -\Delta_1 \psi_1 = \omega_1, \quad (3.7)$$

which is what we are after in the first place.

3.2 Vorticity Depletion and Inviscid Damping

A few comments are in order concerning the decomposition (3.5). Although $f_{1;1}(r)$ is time independent, and thus it does not decay with t , it is unusually small near the center of the vortex, that is

$$f_{1;1}(r) = O(r^3) \quad \text{as } r \rightarrow 0,$$

instead of just an $O(r)$ behavior. To see this, using the notation in (1.9) for $k = 1$, one expands

$$\omega_1^{in}(r) = \omega_{1,0}^{in} r + O(r^3), \quad \text{as } r \rightarrow 0, \quad (3.8)$$

and uses the precise Taylor series for u and β near $r = 0$ (note that $\beta(0) = -4u''(0)$ from (2.1)). Inserting this in (3.5b) shows that the coefficient $\omega_{1,0}^{in}$ of r cancels out, leading to the $O(r^3)$ behavior (recall (1.8)). This is the vorticity depletion due to the non-locality of the linear equation. Moreover, if ω_1^{in} is compactly supported away from $r = 0$, the same holds for $f_{1;1}$. On the other hand, for the time dependent contribution to f_1 we have the asymptotics

$$f_{1;2}(t, r) = O\left(\frac{r}{\langle t \rangle}\right) \quad \text{as } r \rightarrow 0,$$

which vanishes only $O(r)$ as $r \rightarrow 0$, but instead decays in time.

We now make this intuition rigorous. Using the notation of (1.9), we rewrite

$$\omega_1^{in}(r) = r^{-1/2} F_1(r) + r \chi(r) \frac{\beta(r)}{\beta(0)} \omega_{1,0}^{in} \quad (3.9)$$

where $\omega_{1,0}^{in} = \lim_{r \rightarrow 0} r^{-1} \omega_1^{in}(r)$ (recall (1.8)), and by definition we have that

$$F_1(r) \sim \begin{cases} r^{3/2+2}, & \text{for } r \leq 1, \\ r^{1/2} \omega_1^{in}(r), & \text{for } r > 1. \end{cases} \quad (3.10)$$

The important observation is that (3.10) is precisely consistent with the definition of the weight $w_{F,\delta}$ in (1.10c), when $k = 1$. Inserting (3.9) in (3.5b) and using (2.1), it follows after a short computation that

$$\begin{aligned} f_{1;1}(r) &= r^{-1/2} F_1(r) + \frac{\beta(r)}{r^2 u'(r)} \int_0^r \rho^{3/2} F_1(\rho) d\rho \\ &\quad + \omega_{1,0}^{in} \frac{\beta(r)}{\beta(0) r^2 u'(r)} \int_0^r \rho^3 u'(\rho) \chi'(\rho) d\rho. \end{aligned} \quad (3.11)$$

We note that the last term on the right side of (3.11) vanishes identically for $r \leq 1/2$, by the definition of the smooth cut-off function χ , and behaves as $\beta(r)r$ for $r \geq 1$. It

follows from (3.10)–(3.11) and the definition (1.10c) that

$$\|\sqrt{r}f_{1;1}\|_{L^2_{F,\delta}} \lesssim |\omega_{1;0}^{in}| + \|F_1\|_{L^2_{F,\delta/2}}. \quad (3.12)$$

In a similar way, using that the operator $r\partial_r$ is scale invariant, we may apply derivatives to (3.11) and obtain that

$$\|\sqrt{r}(r\partial_r)f_{1;1}\|_{L^2_{F,\delta}} \lesssim |\omega_{1;0}^{in}| + \|F_1\|_{L^2_{F,\delta/2}} + \|(r\partial_r)F_1\|_{L^2_{F,\delta/2}}$$

and

$$\|\sqrt{r}(r\partial_r)^2f_{1;1}\|_{L^2_{F,\delta}} \lesssim |\omega_{1;0}^{in}| + \|F_1\|_{L^2_{F,\delta/2}} + \|(r\partial_r)F_1\|_{L^2_{F,\delta/2}} + \|(r\partial_r)^2F_1\|_{L^2_{F,\delta/2}}$$

which completes the proof of the first half of (1.13).

We obtain similar estimates for $f_{1;2}$ by using (3.5c), the already established bounds for $f_{1;1}$, and integration by parts. For instance, for the weighted L^2 estimate on $f_{1;2}$ we have

$$\begin{aligned} itf_{1;2}(r) &= r\beta(r) \int_r^\infty \frac{-1}{u'(\rho)} \partial_\rho \left(e^{i(u(r)-u(\rho))t} \right) f_{1;1}(\rho) \frac{d\rho}{\rho u'(\rho)} \\ &= \int_0^\infty \mathbf{1}_{\rho>r} r\beta(r) e^{i(u(r)-u(\rho))t} \frac{w_{F,\delta/2}(\rho)}{\rho^{5/2}(u'(\rho))^2} \frac{\sqrt{\rho}(\rho\partial_\rho)f_{1;1}(\rho)}{w_{F,\delta/2}(\rho)} d\rho \\ &\quad - \int_0^\infty \mathbf{1}_{\rho>r} r\beta(r) e^{i(u(r)-u(\rho))t} \frac{w_{F,\delta/2}(\rho)(u'(\rho) + 2\rho u''(\rho))}{\rho^{5/2}(u'(\rho))^3} \frac{\sqrt{\rho}f_{1;1}(\rho)}{w_{F,\delta/2}(\rho)} d\rho \\ &\quad + \sqrt{r}\beta(r) \frac{w_{F,\delta/2}(r)}{r(u'(r))^2} \frac{\sqrt{r}f_{1;1}(r)}{w_{F,\delta/2}(r)}. \end{aligned} \quad (3.13)$$

Upon dividing by $w_{f,\delta}(r)$ and using that

$$\begin{aligned} \left\| \frac{\sqrt{r}\beta(r)}{w_{f,\delta}(r)} \frac{w_{F,\delta/2}(r)}{r(u'(r))^2} \right\|_{L_r^\infty} &\lesssim 1, \\ \left\| \mathbf{1}_{\rho>r} \frac{r\beta(r)}{w_{f,\delta}(r)} \frac{w_{F,\delta/2}(\rho)}{\rho^{5/2}(u'(\rho))^2} \right\|_{L_r^2 L_\rho^2} &\lesssim 1, \\ \left\| \mathbf{1}_{\rho>r} \frac{r\beta(r)}{w_{f,\delta}(r)} \frac{w_{F,\delta/2}(\rho)(u'(\rho) + 2\rho u''(\rho))}{\rho^{5/2}(u'(\rho))^3} \right\|_{L_r^2 L_\rho^2} &\lesssim 1, \end{aligned}$$

and, in view of the already established bounds on $f_{1;1}$, we conclude that

$$\begin{aligned} t\|f_{1;2}\|_{L^2_{f,\delta}} &\lesssim \|\sqrt{r}f_{1;1}\|_{L^2_{F,\delta/2}} + \|\sqrt{r}(r\partial_r)f_{1;1}\|_{L^2_{F,\delta/2}} \lesssim |\omega_{1;0}^{in}| + \|F_1\|_{L^2_{F,\delta/2}} \\ &\quad + \|(r\partial_r)F_1\|_{L^2_{F,\delta/2}}. \end{aligned}$$

In order to bound the weighted $r\partial_r$ norm of $f_{1;2}$, we apply an $r\partial_r$ derivative (which preserves scale) to (3.13) and then divide by $w_{f,\delta}$. Similar $L_r^2 L_\rho^2$ bounds hold for the kernel which arises, and the norm of $\sqrt{r}(r\partial_r)^2 f_{1;1}$ enters the calculation (but we have it already bounded in the suitable norm). We omit these computational details. This concludes the proof of the second half of (1.13).

The proof of inviscid damping for the stream function now directly follows from Lemma 2.15 and the bounds established on $f_{1;1}$ and $f_{1;2}$. We note that for $k = 1$ the stream function has a particularly nice formula

$$\begin{aligned}\psi_1(t, r) &= -\frac{\partial_t + iu(r)}{i\beta(r)} \left(e^{-iu(r)t} f_1(t, r) \right) \\ &= -\frac{e^{-iu(r)t}}{i\beta(r)} \partial_t f_{1;2}(t, r) = -r \int_r^\infty (u(r) - u(r_c)) e^{-itu(r_c)} \frac{f_{1;1}(r_c)}{r_c u'(r_c)} dr_c,\end{aligned}\quad (3.14)$$

which could also have been used to obtain inviscid damping, by integrating twice by parts in r_c , as was done in [68].

4 The Homogeneous Rayleigh Problem for $k \geq 2$

We consider here the homogeneous version of Eq. (2.8), namely

$$\partial_{rr}\phi + \left(\frac{1/4 - k^2}{r^2} + \frac{\beta(r)}{u(r) - z} \right) \phi = 0, \quad (4.1)$$

for $k \geq 2$. In this section we construct a specific solution for $z \in \mathbb{C}$ in a neighborhood of the spectrum and study various properties. Most of the estimates exploit the following weight, defined as

$$w_\phi(r, r_c) = \begin{cases} (r/r_c)^{-k+1/2}, & r < 1, \\ (r/r_c)^{k+1/2}, & r > 2, \end{cases} \quad w_{\phi,\gamma}(r, r_c) = \begin{cases} (r/r_c)^{-k+1/2-\gamma}, & r < 1, \\ (r/r_c)^{k+1/2+\gamma}, & r > 2, \end{cases} \quad (4.2)$$

using also the notation $L_{\phi,\gamma}^2$ as in (1.10a). In what follows, we will often use the the following smooth cutoffs:

$$\chi_c(r, c) = \chi\left(\frac{k(r-r_c)}{r_c}\right) \chi_{\neq}(r, c) = 1 - \chi_c. \quad (4.3a)$$

Note that $|\partial_G \chi_c| \lesssim \max(r_c^{-2}, r_c^2) \left| \chi'_c\left(\frac{k(r-r_c)}{r_c}\right) \right|$ and χ'_c is supported away from the critical layer; analogous observations hold also for $r\partial_r \chi_c$ and $r_c \partial_{r_c} \chi_c$.

Theorem 4.1 (Homogeneous solutions for $k \geq 2$). *Assume $k \geq 2$, let $\varepsilon_0 \in (0, 1/2)$, and define*

$$D_{\varepsilon_0} = (0, \infty) \times ((0, u(0)) + i(-\varepsilon_0, \varepsilon_0)) \subset (0, \infty) \times \mathbb{C}. \quad (4.4)$$

There exists a unique solution $\phi : D_{\varepsilon_0} \rightarrow \mathbb{C}$ to (4.1) of the form

$$\phi(r, z) = P(r, z)(u(r) - z) \quad (4.5)$$

with $P : D_{\varepsilon_0} \rightarrow \mathbb{C}$ such that $P(r_c, z) = 1$ and $\partial_r P(r_c, z) = 0$ for every $z \in ((0, u(0)) + i(-\varepsilon_0, \varepsilon_0))$. Moreover, ϕ is continuously differentiable with respect to r and r_c .

The proof of this theorem is carried out throughout Sect. 4. Let Q_0, Q_∞ be defined via the identities

$$\phi(r, z) = \left(\frac{r}{r_c}\right)^{1/2-k} Q_0(r, z)(u(r) - z), \quad \forall r \in (0, r_c], \quad (4.6)$$

and

$$\phi(r, z) = \left(\frac{r}{r_c}\right)^{1/2+k} Q_\infty(r, z)(u(r) - z), \quad \forall r \in (r_c, \infty). \quad (4.7)$$

Moreover, the functions

$$B_0(r, z) = (k+1)Q_0(r, z) - r\partial_r Q_0(r, z), \quad r \leq r_c, \quad (4.8)$$

and

$$B_\infty(r, z) = (k-1)Q_\infty(r, z) + r\partial_r Q_\infty(r, z), \quad r \geq r_c, \quad (4.9)$$

will play an important role. When z is real and belongs to the interval $(0, u(0))$, more can be deduced.

Theorem 4.2 (Further properties for the real solution). *Assume that $z = c \in (0, u(0))$, and let $\phi(r, c)$ be the unique solution to the problem posed in (4.5). Then*

$$Q_0(r, c) > 0, \quad \partial_r Q_0(r, c) > 0, \quad \forall r \in (0, r_c], \quad (4.10)$$

and

$$Q_\infty(r, c) > 0, \quad \partial_r Q_\infty(r, c) < 0, \quad \forall r \in (r_c, \infty). \quad (4.11)$$

Moreover,

$$Q_0(r, c), Q_\infty(r, c) \approx 1, \quad \|r\partial_r Q_0(\cdot, c)\|_{L^\infty(0, r_c)}, \|r\partial_r Q_\infty(\cdot, c)\|_{L^\infty(r_c, \infty)} \lesssim k. \quad (4.12)$$

Finally, there is a constant δ_0 depending only on u such that for all $r_c \in (0, \infty)$, there holds for all $|r - r_c| < \delta_0 r_c$ (uniformly in k and r_c),

$$k^2 \frac{r^{2k-1}}{r_c^{2k}} |r_c - r| \lesssim B_0(r, c) \leq \frac{k^2 - 1}{r} |r_c - r|, \quad \forall r \in (0, r_c], \quad (4.13)$$

and

$$k^2 \frac{r^{2k-1}}{r_c^{2k}} |r_c - r| \lesssim B_\infty(r, c) \leq \frac{k^2 - 1}{r_c} |r - r_c|, \quad \forall r \in [r_c, \infty). \quad (4.14)$$

As a consequence, the theorem proves that, when the spectral parameter $c \in (0, u(0))$, the only point at which P vanishes is the critical layer, while otherwise P is bounded away from 0 uniformly, thanks to (4.12). Combining this with suitable convergence estimates deduced below, we deduce non-vanishing properties of the complex solution as well. However, due to extra factors of $r_c^{2+\varkappa}$ and $r_c^{-2-\varkappa}$ (see e.g. Lemmas 4.11–4.14), this information will only be available in the domain I_α given in (2.18). Eventually, \varkappa will be chosen to be a multiple of α .

Theorem 4.3 *On the domain $(r, c) \in (0, \infty) \times I_\alpha$, we have*

$$\|(r \partial_r)^j P\|_{L_\phi^\infty} \lesssim_\alpha k^j, \quad j = 0, 1, \quad (4.15)$$

$$\|(u(r) - z)r^2 \partial_{rr} P\|_{L_{\phi, \alpha/2}^\infty} \lesssim_\alpha k^3, \quad (4.16)$$

$$\|r_c \partial_{r_c} P\|_{L_\phi^\infty} \lesssim k^3, \quad (4.17)$$

$$\|(r \partial_r)^j (P(r, c \pm i\varepsilon) - P(r, c))\|_{L_\phi^\infty} \lesssim_\alpha \varepsilon^{\eta_\alpha}, \quad j = 0, 1, \quad (4.18)$$

$$\|r_c \partial_{r_c} (P(r, c \pm i\varepsilon) - P(r, c))\|_{L_\phi^\infty} \lesssim_\alpha \varepsilon^{\eta_\alpha}, \quad (4.19)$$

$$\|r \partial_r r_c \partial_{r_c} (P(r, c \pm i\varepsilon) - P(r, c)) \chi_\neq\|_{L_\phi^\infty} \lesssim_\alpha \varepsilon^{\eta_\alpha}, \quad (4.20)$$

where χ_\neq is defined in (4.3), and we have the (uniform in ε) Lipschitz bound

$$\frac{|r \partial_r P(r, c \pm i\varepsilon)|}{w_{\phi, \alpha/2}(r, r_c)} \lesssim_\alpha k^3 \frac{|r - r_c|}{r_c}, \quad (4.21)$$

and convergence estimate

$$\|(u(r) - c)r^2 \partial_{rr} (P(r, c \pm i\varepsilon) - P(r, c))\|_{L_{\phi, \alpha/2}^\infty} \lesssim_\alpha \varepsilon^{\frac{2}{2+\alpha}}. \quad (4.22)$$

Finally, for every $r \in (0, \infty)$ and $r_c \in I_\alpha$ such that

$$|r - r_c| > \frac{r_c}{k},$$

there holds (uniformly in ε)

$$\frac{|r \partial_r r_c \partial_{r_c} P(r, c \pm i\varepsilon)|}{w_{\phi, \alpha/2}(r, r_c)} \lesssim_\alpha k^4, \quad (4.23)$$

while if

$$|r - r_c| \leq \frac{r_c}{k},$$

we have the pointwise bounds (uniformly in ε)

$$\begin{aligned} \min\{r_c^2, r_c^{-2}\} \frac{|\partial_G P(r, c \pm i\varepsilon)|}{w_{\phi, \alpha/2}(r, r_c)} &\lesssim_\alpha k^2, \\ \min\{r_c^2, r_c^{-2}\} \frac{|r \partial_r \partial_G P(r, c \pm i\varepsilon)|}{w_{\phi, \alpha/2}(r, r_c)} &\lesssim_\alpha k^4 \frac{|r - r_c|}{r_c}, \end{aligned} \quad (4.24)$$

and the convergence estimates

$$\min\{r_c^2, r_c^{-2}\} \frac{|(r \partial_r)^j \partial_G (P(r, c \pm i\varepsilon) - P(r, c))|}{w_{\phi, \alpha/2}(r, r_c)} \lesssim_\alpha \varepsilon^{\eta_\alpha}, \quad j = 0, 1. \quad (4.25)$$

Remark 4.4 The estimates on P of the above Theorem 4.3 can also be written in terms on Q_0, Q_∞ . Of particular importance are

$$\left\| (r \partial_r)^j (Q_\bullet(r, c \pm i\varepsilon) - Q_\bullet(r, c)) \right\|_{L^\infty} \lesssim_\alpha \varepsilon^{\eta_\alpha}, \quad (4.26)$$

$$\left\| r_c \partial_{r_c} (Q_\bullet(r, c \pm i\varepsilon) - Q_\bullet(r, c)) \right\|_{L^\infty} \lesssim_\alpha \varepsilon^{\eta_\alpha}, \quad (4.27)$$

$$\left\| r \partial_r r_c \partial_{r_c} (Q_\bullet(r, c \pm i\varepsilon) - Q_\bullet(r, c)) \chi_{\neq} \right\|_{L^\infty} \lesssim_\alpha \varepsilon^{\eta_\alpha}, \quad (4.28)$$

where $\eta_\alpha = \frac{\alpha}{2(2+\alpha)}$ and $\bullet = 0, \infty$. From (4.26) combined with (4.12), we infer that

$$|Q_\bullet(r, z)| \approx_\alpha 1, \quad \|r \partial_r Q_\bullet(\cdot, z)\|_{L^\infty} \lesssim_\alpha k. \quad (4.29)$$

The first estimate is crucially stating that ϕ vanishes only at the critical layer, when $r = r_c$.

The proof of Theorem 4.3 combines Lemmas 4.11–4.14 and Remark 4.15 below for $\varkappa = \alpha/2$ with the definition of I_α . Once (4.26) is established, the bounds (4.15) and (4.29) follow from (4.12) and (4.6)–(4.7), while (4.22) is precisely (4.72). The rest are stated in an equivalent way in Propositions 4.6 and 4.9.

4.1 Existence and Uniqueness of Solutions

Fix $z \in \mathbb{C}$ be such that $c = \operatorname{Re} z \in (0, u(0))$. We will denote by $\phi_1 = \phi_1(r, z)$ the homogeneous solution for the $k = 1$ Rayleigh problem devised in (2.10) (see Sect.

3.1), namely

$$\phi_1(r, z) = \left(\frac{r}{r_c}\right)^{3/2} (u(r) - z), \quad (4.30)$$

appropriately normalized at the critical layer. We will look for a solution ϕ to (4.1) of the form

$$\phi(r, z) = \phi_1(r, z) \tilde{P}(r, z), \quad (4.31)$$

and set up a contraction mapping argument for \tilde{P} , which satisfies

$$\partial_r \left(\phi_1^2 \partial_r \tilde{P} \right) + \frac{1 - k^2}{r^2} \phi_1^2 \tilde{P} = 0, \quad (4.32)$$

subject to the boundary conditions

$$\tilde{P}(r_c, z) = 1, \quad \partial_r \tilde{P}(r_c, z) = 0. \quad (4.33)$$

Integrating (4.32) twice and using (4.33), we infer that

$$\partial_r \tilde{P}(r, z) = -\frac{k^2 - 1}{r^3 (u(r) - z)^2} \int_r^{r_c} s (u(s) - z)^2 \tilde{P}(s, z) ds \quad (4.34)$$

and

$$\tilde{P}(r, z) = 1 + (k^2 - 1) \int_r^{r_c} \frac{1}{\rho^3 (u(\rho) - z)^2} \int_\rho^{r_c} s (u(s) - z)^2 \tilde{P}(s, z) ds d\rho =: 1 + \mathcal{I}_z[\tilde{P}]. \quad (4.35)$$

The above expression will be useful to set up a proper fixed point scheme to deduce existence and uniqueness of \tilde{P} . For further reference, we can take a ∂_{r_c} of the above expression, taking into account that $z = u(r_c) \pm i\varepsilon$, obtaining

$$\begin{aligned} \partial_{r_c} \tilde{P}(r, z) &= \mathcal{I}_z[\partial_{r_c} \tilde{P}] + 2(k^2 - 1) \int_r^{r_c} \frac{u'(r_c)}{\rho^3 (u(\rho) - z)^3} \int_\rho^{r_c} s (u(s) - z)^2 \tilde{P}(s, z) ds d\rho \\ &\quad - 2(k^2 - 1) \int_r^{r_c} \frac{u'(r_c)}{\rho^3 (u(\rho) - z)^2} \int_\rho^{r_c} s (u(s) - z) \tilde{P}(s, z) ds d\rho \\ &\quad - \varepsilon^2 (k^2 - 1) r_c \int_r^{r_c} \frac{1}{\rho^3 (u(\rho) - z)^2} d\rho \\ &= \mathcal{I}_z[\partial_{r_c} \tilde{P}] + \mathcal{I}_1 + \mathcal{I}_2 + \mathcal{I}_3. \end{aligned} \quad (4.36)$$

Notice that the above expression is valid for $\varepsilon > 0$. Using integration by parts, we derive the equivalent formula

$$\begin{aligned} \partial_{r_c} \tilde{P}(r, z) = & \mathcal{T}_z[\partial_{r_c} \tilde{P}] + (k^2 - 1)u'(r_c) \left[\frac{1}{r^3 u'(r)(u(r) - z)^2} \int_r^{r_c} \rho(u(\rho) - z)^2 \tilde{P}(\rho, z) d\rho \right. \\ & + \int_r^{r_c} \partial_\rho \left(\frac{1}{\rho^3 u'(\rho)} \right) \frac{1}{(u(\rho) - z)^2} \int_\rho^{r_c} s(u(s) - z)^2 \tilde{P}(s, z) ds d\rho \\ & + \int_r^{r_c} \frac{1}{\rho^3 (u(\rho) - z)^2} \int_\rho^{r_c} \partial_s \left(\frac{s}{u'(s)} \right) (u(s) - z)^2 \tilde{P}(s, z) ds d\rho \\ & \left. + \int_r^{r_c} \frac{1}{\rho^3 (u(\rho) - z)^2} \int_\rho^{r_c} \frac{s}{u'(s)} (u(s) - z)^2 \partial_s \tilde{P}(s, z) ds d\rho \right]. \end{aligned} \quad (4.37)$$

This is the formulation that we use also at $\varepsilon = 0$ as no singular integral appears here (see Sect. 4.4 below).

4.1.1 An Auxiliary Weight

With the convention adopted in (2.17), we define an auxiliary weight $\tilde{w} = \tilde{w}(r, r_c)$ to solve the ODE

$$-\partial_{rr} \tilde{w} - \frac{3}{r} \partial_r \tilde{w} + \frac{1}{A^2 - 1} \frac{1}{r^2} \tilde{w} = 0, \quad \tilde{w}(r_c, r_c) = 1, \quad \partial_r \tilde{w}(r_c, r_c) = 0, \quad (4.38)$$

where $A > k$ is a parameter that we leave unspecified at the moment (one should think of A as close to k). In fact, we can solve the above ODE explicitly, to find

$$\tilde{w}(r, r_c) = \frac{A+1}{2A} \left(\frac{r}{r_c} \right)^{A-1} + \frac{A-1}{2A} \left(\frac{r}{r_c} \right)^{-A-1}. \quad (4.39)$$

Notice that, given a fixed $r_c > 0$, \tilde{w} attains its minimum at $r = r_c$ and

$$\tilde{w}(r, r_c) \geq 1, \quad \forall r > 0. \quad (4.40)$$

The following properties of \tilde{w} will prove useful later.

Lemma 4.5 *Let $c \in (0, u(0))$, $A > k$ and \tilde{w} be given by (4.39). Then*

$$\int_r^{r'} \frac{1}{\rho^3} \int_\rho^{r_c} s \tilde{w}(s, r_c) ds d\rho = \frac{1}{A^2 - 1} (\tilde{w}(r, r_c) - \tilde{w}(r', r_c)), \quad \forall r, r' > 0. \quad (4.41)$$

Furthermore, for any $b \neq \pm A$ and for $A = k + \varkappa$ with $\varkappa \in (0, 1)$, we have

$$\left| \int_r^{r_c} s^b \tilde{w}(s, r_c) ds \right| \lesssim_b \frac{r^{1+b}}{k} w(r, r_c), \quad \forall r > 0. \quad (4.42)$$

Moreover

$$\left| \int_r^{r_c} s \tilde{w}(s, r_c) ds \right| \lesssim \frac{r^2}{r_c} |r - r_c| \tilde{w}(r, r_c), \quad \forall r > 0, \quad (4.43)$$

and

$$\left| \int_r^{r_c} s^b \tilde{w}(s, r_c) ds \right| \lesssim_b \max\{r, r_c\}^b |r - r_c| \tilde{w}(r, r_c), \quad \forall r > 0. \quad (4.44)$$

Proof of Lemma 4.5 Equation (4.41) is obtained by explicitly computing the integrals from formula (4.39). Turning to (4.42)–(4.43), we first observe that

$$\begin{aligned} \int_r^{r_c} s^b \tilde{w}(s, r_c) ds &= \frac{A+1}{2A(A+b)} r^{b+1} \left[\left(\frac{r}{r_c} \right)^{-b-1} - \left(\frac{r}{r_c} \right)^{A-1} \right] \\ &\quad - \frac{A-1}{2A(A-b)} r^{b+1} \left[\left(\frac{r}{r_c} \right)^{-b-1} - \left(\frac{r}{r_c} \right)^{-A-1} \right]. \end{aligned} \quad (4.45)$$

By considering the different cases $r \leq r_c$ and $r > r_c$ and the different ranges of $b \neq \pm A$ with respect to A , (4.42) follows immediately from the definition (4.39). Also, (4.44) is obvious from (4.40). Regarding (4.43), we use the fact that if $r \leq r_c$, then applying the mean value theorem to the function $(0, 1] \ni x \mapsto x^{2A}$ and using (4.39), we have

$$\begin{aligned} \left| \int_r^{r_c} s \tilde{w}(s, r_c) ds \right| &= -\frac{r^2}{2A} \left(\frac{r}{r_c} \right)^{-A-1} \left[\left(\frac{r}{r_c} \right)^{2A} - 1 \right] \leq \frac{r^2}{2A} \left(\frac{r}{r_c} \right)^{-A-1} 2A \left| \frac{r}{r_c} - 1 \right| \\ &\leq \frac{2A}{A-1} \frac{r^2}{r_c} |r - r_c| \tilde{w}(r, r_c). \end{aligned} \quad (4.46)$$

A similar computation, applied to the function $[1, \infty) \ni x \mapsto x^{-2A}$, also show that if $r > r_c$, then

$$\left| \int_r^{r_c} s \tilde{w}(s, r_c) ds \right| \leq \frac{2A}{A+1} \frac{r^2}{r_c} |r - r_c| \tilde{w}(r, r_c). \quad (4.47)$$

The two estimates can be grouped together as in (4.42), concluding the proof. \square

4.1.2 Existence and Uniqueness of \tilde{P}

We begin with proving existence, uniqueness and some regularity for (4.35). It is clear that existence and uniqueness of \tilde{P} is equivalent to existence and uniqueness of P in Theorem 4.1. Moreover, all the properties on \tilde{P} translate into properties of P , since

$$P(r, z) = \left(\frac{r}{r_c} \right)^{3/2} \tilde{P}(r, z). \quad (4.48)$$

Proposition 4.6 *Let $\varepsilon_0 \in (0, 1/2)$ and D_{ε_0} as in Theorem 4.1, and suppose \tilde{w} is given by (4.39) with $A = k + \varkappa$. Then (4.35) has a unique solution $\tilde{P} \in L^\infty_{\tilde{w}}(D_{\varepsilon_0})$ such that*

$$\|(r\partial_r)^j \tilde{P}\|_{L^\infty_{\tilde{w}}} := \|\tilde{w}^{-1}(r\partial_r)^j \tilde{P}\|_{L^\infty} \lesssim_{\varkappa} k^{1+j}, \quad j = 0, 1, \quad (4.49)$$

and

$$\|(u(r) - z)r^2\partial_{rr}\tilde{P}\|_{L^\infty_{\tilde{w}}} \lesssim_{\varkappa} k^3. \quad (4.50)$$

Moreover

$$\frac{|r\partial_r\tilde{P}(r, z)|}{\tilde{w}(r, r_c)} \lesssim_{\varkappa} k^3 \frac{|r - r_c|}{r_c}, \quad (4.51)$$

for any $r, r_c > 0$.

Thus, thanks to (4.48), Proposition 4.6 immediately proves the bound

$$\|(r\partial_r)^j P\|_{L^\infty_{\phi, \varkappa}} \lesssim_{\varkappa} k^{1+j}, \quad j = 0, 1. \quad (4.52)$$

Proposition 4.6 is based on the contraction properties of the operator \mathcal{T}_z in (4.35). This approach can be viewed as a refinement of an analogous argument in [66] to the more complicated vortex case.

Lemma 4.7 *If $Z \in L^\infty(D_{\varepsilon_0})$, there holds*

$$\|\mathcal{T}_z[\tilde{w}Z]\|_{L^\infty_{\tilde{w}}} \leq \frac{k^2 - 1}{A^2 - 1} \|Z\|_{L^\infty}, \quad (4.53)$$

where \tilde{w} is given by (4.39).

Proof of Lemma 4.7 By linearity, we can assume that $\|Z\|_{L^\infty} = 1$. In view of (4.35), (4.30), and the monotonicity of u (hence $|u(s) - z| \leq |u(\rho) - z|$ in the integral), we have the immediate bound

$$\left| \tilde{w}^{-1} \mathcal{T}_z[\tilde{w}Z] \right| \leq \frac{k^2 - 1}{\tilde{w}(r, r_c)} \left| \int_r^{r_c} \frac{1}{\rho^3} \int_\rho^{r_c} s \tilde{w}(s, r_c) ds d\rho \right|.$$

From (4.41), we readily obtain

$$\left| \tilde{w}^{-1} \mathcal{T}_z[\tilde{w}Z] \right| \leq \frac{k^2 - 1}{A^2 - 1} \frac{\tilde{w}(r, r_c) - 1}{\tilde{w}(r, r_c)}, \quad (4.54)$$

which implies (4.53). \square

We can now proceed with the proof of the main result of this section.

Proof of Proposition 4.6 The existence and uniqueness of \tilde{P} follows from the contraction mapping principle. Indeed, from (4.53) and the fact that \tilde{w} is bounded below (see (4.40)), the operator $1 + \mathcal{T}_z[\cdot] : L_{\tilde{w}}^\infty \rightarrow L_{\tilde{w}}^\infty$ is a contraction whenever $A^2 > k^2$. Moreover, from (4.35) and (4.53), the unique solution \tilde{P} satisfies

$$|\tilde{w}^{-1} \tilde{P}(r, z)| \leq \frac{1}{\tilde{w}(r, r_c)} + \left| \tilde{w}^{-1} \mathcal{T}_z[\tilde{w} \tilde{w}^{-1} \tilde{P}] \right| \leq 1 + \frac{k^2 - 1}{A^2 - 1} \|\tilde{P}\|_{L_{\tilde{w}}^\infty}, \quad (4.55)$$

and (4.49) with $j = 0$ follows. We now turn to the second part of (4.49). In light of (4.34), we have that

$$|\tilde{w}^{-1} r \partial_r \tilde{P}(r, z)| \leq \frac{k^2 - 1}{r^2 \tilde{w}(r, r_c)} \|\tilde{P}\|_{L_{\tilde{w}}^\infty} \left| \int_r^{r_c} s \tilde{w}(s, r_c) ds \right|. \quad (4.56)$$

Thus, taking advantage of (4.42), we deduce that

$$|\tilde{w}^{-1} r \partial_r \tilde{P}(r, z)| \lesssim \frac{k^2 - 1}{k} \|\tilde{P}\|_{L_{\tilde{w}}^\infty}, \quad (4.57)$$

and the claim follows by combining the above estimate with (4.49) with $j = 0$. We now prove (4.50). Firstly, notice that we can read $\partial_{rr} \tilde{P}$ directly from (4.32), obtaining

$$\partial_{rr} \tilde{P}(r, z) = -\frac{3}{r} \partial_r \tilde{P} - \frac{2u'(r)}{u(r) - z} \partial_r \tilde{P} - (k^2 - 1) \frac{\tilde{P}(r, z)}{r^2}. \quad (4.58)$$

Hence,

$$\|(u(r) - z) r^2 \partial_{rr} \tilde{P}(r, z)\|_{L_{\tilde{w}}^\infty} \lesssim \|(u(r) - z) r \partial_r \tilde{P}\|_{L_{\tilde{w}}^\infty} + (k^2 - 1) \|\tilde{P}\|_{L_{\tilde{w}}^\infty},$$

and (4.50) follows from (4.49). Finally, (4.51) is a consequence of (4.43) and (4.56). The proof is concluded. \square

4.2 Convergence of $(r \partial_r)^j \tilde{P}$

As a first step towards the proof of the convergence estimates in Theorem 4.3, we deduce a convergence estimate on \tilde{P} . These estimates are relevant near the critical layer, whereas near $r = 0, \infty$ improve the estimates further below.

Proposition 4.8 *Under the assumptions in Theorem 4.1, and with \tilde{w} given by (4.39) with $A > k \geq 2$, there holds*

$$\|\min\{r_c^2, r_c^{-2}\} (r \partial_r)^j (\tilde{P}(r, c \pm i\varepsilon) - \tilde{P}(r, c))\|_{L_{\tilde{w}}^\infty} \lesssim \varepsilon k^{2+j}, \quad j = 0, 1, \quad (4.59)$$

and

$$\|\min\{r_c^2, r_c^{-2}\} (u(r) - c) r^2 \partial_{rr} (\tilde{P}(r, c \pm i\varepsilon) - \tilde{P}(r, c))\|_{L_{\tilde{w}}^\infty} \lesssim \varepsilon k^4, \quad (4.60)$$

for every $\varepsilon \in (0, \varepsilon_0)$.

Proof of Proposition 4.8 We begin by showing the case $j = 0$. By setting $z = c \pm i\varepsilon$, we use (4.35) to deduce that

$$\begin{aligned} \tilde{P}(r, c) - \tilde{P}(r, z) &= \mp i\varepsilon(k^2 - 1) \int_r^{r_c} \frac{1}{\rho^3} \frac{2(u(\rho) - c) \mp i\varepsilon}{(u(\rho) - z)^2(u(\rho) - c)^2} \\ &\quad \int_\rho^{r_c} s(u(s) - c)^2 \tilde{P}(s, c) ds d\rho \end{aligned} \quad (4.61)$$

$$\begin{aligned} &\pm i\varepsilon(k^2 - 1) \int_r^{r_c} \frac{1}{\rho^3(u(\rho) - z)^2} \\ &\quad \int_\rho^{r_c} s[2(u(s) - c) \mp i\varepsilon] \tilde{P}(s, c) ds d\rho \end{aligned} \quad (4.62)$$

$$+ \mathcal{T}_z[\tilde{P}(r, c) - \tilde{P}(r, z)]. \quad (4.63)$$

We now use Lemma 4.5 and (4.49) several times. Let us consider the case when $r < r_c$ only, since the other case is analogous. Define $b_k = 1 - 1/k > 0$. Then,

$$\begin{aligned} |(4.61) + (4.62)| &\lesssim \varepsilon k^2 \|\tilde{P}\|_{L_w^\infty} \int_{b_k r_c}^{r_c} \frac{1}{\rho^3} \frac{1}{|u(\rho) - z|} \int_\rho^{r_c} s \tilde{w}(s, r_c) ds d\rho \\ &\quad + \varepsilon k^2 \|\tilde{P}\|_{L_w^\infty} \int_r^{b_k r_c} \frac{1}{\rho^3} \frac{1}{|u(\rho) - z|} \int_\rho^{r_c} s \tilde{w}(s, r_c) ds d\rho \\ &\lesssim \varepsilon k^2 \|\tilde{P}\|_{L_w^\infty} \frac{1}{|u'(r_c)| r_c} \int_{b_k r_c}^{r_c} \frac{1}{\rho} \tilde{w}(\rho, r_c) d\rho \\ &\quad + \varepsilon k \|\tilde{P}\|_{L_w^\infty} \frac{1}{|u(b_k r_c) - u(r_c)|} \int_r^{b_k r_c} \frac{1}{\rho} \tilde{w}(\rho, r_c) d\rho \\ &\lesssim \varepsilon k^2 \frac{\tilde{w}(r, r_c)}{\min\{r_c^2, r_c^{-2}\}}. \end{aligned} \quad (4.64)$$

Hence, using (4.53) to control (4.63), we arrive at (4.59) with $j = 0$. We now deal with the similar convergence estimate for the derivative of \tilde{P} , namely the case $j = 1$. Use (4.34) to get

$$\frac{r \partial_r (\tilde{P}(r, c) - \tilde{P}(r, z))}{k^2 - 1} = \pm i\varepsilon \frac{2(u(r) - c) \mp i\varepsilon}{r^2(u(r) - z)^2(u(r) - c)^2} \int_r^{r_c} s(u(s) - c)^2 \tilde{P}(s, c) ds \quad (4.65)$$

$$\mp i\varepsilon \frac{1}{r^2(u(r) - z)^2} \int_r^{r_c} s[2(u(s) - c) \mp i\varepsilon] \tilde{P}(s, c) ds \quad (4.66)$$

$$- \frac{1}{r^2(u(r) - z)^2} \int_r^{r_c} s(u(s) - z)^2 [\tilde{P}(s, c) - \tilde{P}(s, z)] ds. \quad (4.67)$$

In the same way as above, we find

$$|(4.65) + (4.66)| \lesssim \varepsilon \|\tilde{P}\|_{L_w^\infty} \frac{1}{r^2 |u(r) - z|} \left| \int_r^{r_c} s \tilde{w}(s, r_c) ds \right| \lesssim \varepsilon k \frac{w(r, r_c)}{\min\{r_c^2, r_c^{-2}\}},$$

while appealing to (4.59) with $j = 0$ we also infer that

$$\begin{aligned} |(4.67)| &\lesssim \|\min\{r_c^2, r_c^{-2}\}(\tilde{P}(r, c \pm i\varepsilon) - \tilde{P}(r, c))\|_{L_w^\infty} \\ &\quad \frac{1}{r^2 \min\{r_c^2, r_c^{-2}\}} \left| \int_r^{r_c} s \tilde{w}(s, r_c) ds \right| \lesssim \varepsilon k \frac{w(r, r_c)}{\min\{r_c^2, r_c^{-2}\}}, \end{aligned} \quad (4.68)$$

and (4.59) follows. Concerning (4.60), note that we can read $\partial_{rr} \tilde{P}$ directly from (4.32), obtaining

$$\begin{aligned} \partial_{rr}(\tilde{P}(r, z) - \tilde{P}(r, c)) &= -\frac{3}{r} \partial_r(\tilde{P}(r, z) - \tilde{P}(r, c)) \\ &\quad \pm i\varepsilon \frac{2u'(r)}{(u(r) - z)(u(r) - c)} \partial_r \tilde{P}(r, z) \\ &\quad - \frac{2u'(r)}{u(r) - z} \partial_r(\tilde{P}(r, z) - \tilde{P}(r, c)) \\ &\quad + (k^2 - 1) \frac{\tilde{P}(r, z) - \tilde{P}(r, c)}{r^2}. \end{aligned}$$

and hence using the boundedness of u we obtain

$$\begin{aligned} &\|\min\{r_c^2, r_c^{-2}\}(u(r) - c)r^2 \partial_{rr}(\tilde{P}(r, c \pm i\varepsilon) - \tilde{P}(r, c))\|_{L_w^\infty} \\ &\lesssim \|\min\{r_c^2, r_c^{-2}\}r \partial_r(\tilde{P}(r, z) - \tilde{P}(r, c))\|_{L_w^\infty} \\ &\quad + \varepsilon \left\| \min\{r_c^2, r_c^{-2}\}r^2 \frac{u'(r)}{u(r) - z} \partial_r \tilde{P}(r, z) \right\|_{L_w^\infty} \\ &\quad + k^2 \|\min\{r_c^2, r_c^{-2}\}(\tilde{P}(r, z) - \tilde{P}(r, c))\|_{L_w^\infty}. \end{aligned} \quad (4.69)$$

Given (4.59), we only need to treat the second term above. Using (4.51), if $|r - r_c| < r_c/2$, we obtain

$$\frac{|u'(r)r}{|u(r) - z|} |\tilde{w}^{-1} r \partial_r \tilde{P}(r, z)| \lesssim k^3 \frac{|u'(r)r}{|u(r) - z|} \frac{|r - r_c|}{r_c} \lesssim k^3, \quad (4.70)$$

while if $|r - r_c| \geq r_c/2$, since $|u(r) - u(r_c)| \gtrsim \min\{r_c^2, r_c^{-2}\}$, we have

$$\frac{|u'(r)r}{|u(r) - z|} |\tilde{w}^{-1} r \partial_r \tilde{P}(r, z)| \lesssim \frac{|u'(r)r}{\min\{r_c^2, r_c^{-2}\}} \|r \partial_r \tilde{P}\|_{L_w^\infty} \lesssim \frac{k^2}{\min\{r_c^2, r_c^{-2}\}}.$$

Thus, we can plug in the above estimates in (4.69) and use (4.59) to conclude the proof. \square

As a consequence of (4.48), we deduce

$$\|\min\{r_c^2, r_c^{-2}\}(r\partial_r)^j(P(r, c \pm i\varepsilon) - P(r, c))\|_{L_{\phi, \varkappa}^\infty} \lesssim_{\varkappa} \varepsilon k^{2+j}, \quad j = 0, 1, \quad (4.71)$$

and

$$\|\min\{r_c^2, r_c^{-2}\}(u(r) - c)r^2\partial_{rr}(P(r, c \pm i\varepsilon) - P(r, c))\|_{L_{\phi, \varkappa}^\infty} \lesssim \varepsilon k^4. \quad (4.72)$$

Note that the last estimate is exactly (4.22). We now deal with ∂_G and ∂_{r_c} derivatives.

4.3 Analysis of $\partial_G \tilde{P}$

In this section, we analyze more carefully the behavior of \tilde{P} near the critical layer, by making use of the differential operator ∂_G defined in (2.27).

Proposition 4.9 *Let \tilde{w} be given by (4.39) with $A = k + \varkappa$, and let us fix $r_c \in (0, \infty)$. For every $r \in (0, \infty)$ such that*

$$|r - r_c| \leq \frac{r_c}{k},$$

we have the pointwise bounds

$$\min\{r_c^2, r_c^{-2}\} \frac{|\partial_G \tilde{P}(r, z)|}{\tilde{w}(r, r_c)} \lesssim_{\varkappa} k^2, \quad (4.73a)$$

$$\min\{r_c^2, r_c^{-2}\} \frac{|r\partial_r \partial_G \tilde{P}(r, z)|}{\tilde{w}(r, r_c)} \lesssim_{\varkappa} k^4 \frac{|r - r_c|}{r_c}, \quad (4.73b)$$

and the convergence estimates

$$(\min\{r_c^2, r_c^{-2}\})^2 \frac{|(r\partial_r)^j \partial_G(\tilde{P}(r, c \pm i\varepsilon) - \tilde{P}(r, c))|}{\tilde{w}(r, r_c)} \lesssim_{\varkappa} k^{3+j}\varepsilon, \quad j = 0, 1. \quad (4.74)$$

Proof of Proposition 4.9 A direct computation from (4.34) yields

$$\begin{aligned} \partial_r \partial_G \tilde{P}(r, z) &= \frac{u''(r)}{(u'(r))^2} \partial_r \tilde{P}(r, z) + \frac{3(k^2 - 1)}{r^4 u'(r)(u(r) - z)^2} \int_r^{r_c} s(u(s) - z)^2 \tilde{P}(s, z) ds \\ &\quad - \frac{k^2 - 1}{r^3 (u(r) - z)^2} \int_r^{r_c} \partial_s \left(\frac{s}{u'(s)} \right) (u(s) - z)^2 \tilde{P}(s, z) ds \\ &\quad - \frac{k^2 - 1}{r^3 (u(r) - z)^2} \int_r^{r_c} s(u(s) - z)^2 \partial_G \tilde{P}(s, z) ds. \end{aligned} \quad (4.75)$$

Thus, integrating on (r, r_c) and noting from (4.33) and (4.37) that $\partial_G \tilde{P}(r_c, z) = 0$, we find

$$\begin{aligned} \partial_G \tilde{P}(r, z) = & - \int_r^{r_c} \frac{u''(\rho)}{(u'(\rho))^2} \partial_r \tilde{P}(\rho, z) d\rho \\ & - \int_r^{r_c} \frac{3(k^2 - 1)}{\rho^4 u'(\rho)(u(\rho) - z)^2} \int_\rho^{r_c} s(u(s) - z)^2 \tilde{P}(s, z) ds d\rho \\ & + \int_r^{r_c} \frac{k^2 - 1}{\rho^3 (u(\rho) - z)^2} \int_\rho^{r_c} \partial_s \left(\frac{s}{u'(s)} \right) (u(s) - z)^2 \tilde{P}(s, z) ds d\rho \\ & + \mathcal{T}_z[\partial_G \tilde{P}]. \end{aligned} \quad (4.76)$$

Assuming $|r - r_c| < r_c/k$, we now bound each term on the right-hand side above, tacitly using Lemma 4.5 and the fact that $|u'(r)| \approx \min\{r, r^{-3}\}$. For the first term, we exploit (4.49) to get

$$\begin{aligned} \left| \int_r^{r_c} \frac{u''(\rho)}{(u'(\rho))^2} \partial_r \tilde{P}(\rho, z) d\rho \right| & \lesssim \|r \partial_r \tilde{P}\|_{L^\infty_{\tilde{w}}} \left| \int_r^{r_c} \frac{u''(\rho)}{(u'(\rho))^2 \rho} \tilde{w}(\rho, r_c) d\rho \right| \\ & \lesssim k \frac{\tilde{w}(r, r_c)}{\min\{r_c^2, r_c^{-2}\}}, \end{aligned} \quad (4.77)$$

where the factor of k is due to integration of \tilde{w} . Similarly

$$\left| \int_r^{r_c} \frac{3(k^2 - 1)}{\rho^4 u'(\rho)(u(\rho) - z)^2} \int_\rho^{r_c} s(u(s) - z)^2 \tilde{P}(s, z) ds d\rho \right| \lesssim k \frac{\tilde{w}(r, r_c)}{\min\{r_c^2, r_c^{-2}\}},$$

and

$$\left| \int_r^{r_c} \frac{k^2 - 1}{\rho^3 (u(\rho) - z)^2} \int_\rho^{r_c} \partial_s \left(\frac{s}{u'(s)} \right) (u(s) - z)^2 \tilde{P}(s, z) ds d\rho \right| \lesssim k \frac{\tilde{w}(r, r_c)}{\min\{r_c^2, r_c^{-2}\}}.$$

For the last term, we use (4.41) to deduce that

$$\begin{aligned} |\mathcal{T}_z[\partial_G \tilde{P}]| & = \left| \int_r^{r_c} \frac{k^2 - 1}{\rho^3 (u(\rho) - z)^2} \int_\rho^{r_c} s(u(s) - z)^2 \partial_G \tilde{P}(s, z) ds d\rho \right| \\ & \leq \frac{k^2 - 1}{A^2 - 1} \tilde{w}(r, r_c) \sup_r \frac{|\partial_G \tilde{P}(r, z)|}{\tilde{w}(r, r_c)}. \end{aligned} \quad (4.78)$$

The fact that the right-hand side above is finite for all $\varepsilon > 0$ follows from general ODE theory, since $\partial_G \tilde{P}$ satisfies essentially a perturbation of Laplace's equation. Collecting the above estimate, we obtain

$$\min\{r_c^2, r_c^{-2}\} \left[\frac{\partial_G \tilde{P}(r, z)}{\tilde{w}(r, r_c)} - \frac{k^2 - 1}{A^2 - 1} \sup_r \frac{|\partial_G \tilde{P}(r, z)|}{\tilde{w}(r, r_c)} \right] \lesssim k,$$

and we easily arrive at (4.73a). We now turn to (4.74) with $j = 0$. From (4.76), we infer that

$$\partial_G \tilde{P}(r, c) - \partial_G \tilde{P}(r, z) = \mathcal{T}_c[\partial_G \tilde{P}(r, c)] - \mathcal{T}_z[\partial_G \tilde{P}(r, z)] + \mathcal{V}_1 + (k^2 - 1)[\mathcal{V}_2 + \mathcal{V}_3],$$

where

$$\begin{aligned} \mathcal{V}_1 &= \int_r^{r_c} \frac{u''(\rho)}{(u'(\rho))^2} \partial_r [\tilde{P}(\rho, z) - \tilde{P}(\rho, c)] d\rho, \\ \mathcal{V}_2 &= \pm 3i\varepsilon \int_r^{r_c} \frac{2(u(\rho) - c) \mp i\varepsilon}{\rho^4 u'(\rho)(u(\rho) - z)^2 (u(\rho) - c)^2} \int_\rho^{r_c} s(u(s) - c)^2 \tilde{P}(s, c) ds d\rho \\ &\quad \mp 3i\varepsilon \int_r^{r_c} \frac{1}{\rho^4 u'(\rho)(u(\rho) - z)^2} \int_\rho^{r_c} s[2(u(s) - c) \mp i\varepsilon] \tilde{P}(s, c) ds d\rho \\ &\quad - 3 \int_r^{r_c} \frac{1}{\rho^4 u'(\rho)(u(\rho) - z)^2} \int_\rho^{r_c} s(u(s) - z)^2 [\tilde{P}(s, c) - \tilde{P}(s, z)] ds d\rho, \\ \mathcal{V}_3 &= \pm i\varepsilon \int_r^{r_c} \frac{2(u(\rho) - c) \mp i\varepsilon}{\rho^3 (u(\rho) - c)^2 u(\rho) - z)^2} \int_\rho^{r_c} \partial_s \left(\frac{s}{u'(s)} \right) (u(s) - c)^2 \tilde{P}(s, c) ds d\rho \\ &\quad \mp i\varepsilon \int_r^{r_c} \frac{1}{\rho^3 (u(\rho) - c)^2} \int_\rho^{r_c} \partial_s \left(\frac{s}{u'(s)} \right) [2(u(s) - c) \mp i\varepsilon] \tilde{P}(s, c) ds d\rho \\ &\quad + \int_r^{r_c} \frac{1}{\rho^3 (u(\rho) - c)^2} \int_\rho^{r_c} \partial_s \left(\frac{s}{u'(s)} \right) (u(s) - c)^2 [\tilde{P}(s, c) - \tilde{P}(s, z)] ds d\rho. \end{aligned} \quad (4.79)$$

Arguing as in (4.77) and appealing to (4.59), we find that

$$\begin{aligned} \min\{r_c^2, r_c^{-2}\} |\mathcal{V}_1| &\lesssim \frac{1}{k} \|\min\{r_c^2, r_c^{-2}\} r \partial_r (\tilde{P}(r, z) - \tilde{P}(r, c))\|_{L_w^\infty} \frac{\tilde{w}(r, r_c)}{\min\{r_c^2, r_c^{-2}\}} \\ &\lesssim \varepsilon k^2 \frac{\tilde{w}(r, r_c)}{\min\{r_c^2, r_c^{-2}\}}. \end{aligned}$$

Turning to \mathcal{V}_2 , we find

$$\begin{aligned} \min\{r_c^2, r_c^{-2}\} |\mathcal{V}_2| &\lesssim \varepsilon \min\{r_c^2, r_c^{-2}\} \|\tilde{P}\|_{L_w^\infty} \int_r^{r_c} \frac{1}{\rho^4 |u'(\rho)| |u(\rho) - z|} \\ &\quad \times \int_\rho^{r_c} s \tilde{w}(s, r_c) ds d\rho \\ &\quad + \|\min\{r_c^2, r_c^{-2}\} (\tilde{P}(\rho, c) - \tilde{P}(\rho, z))\|_{L_w^\infty} \int_r^{r_c} \frac{1}{\rho^4 |u'(\rho)|} \\ &\quad \times \int_\rho^{r_c} s \tilde{w}(s, r_c) ds d\rho \\ &\lesssim \frac{\varepsilon}{k} \frac{\|\tilde{P}\|_{L_w^\infty}}{\min\{r_c^2, r_c^{-2}\}} \tilde{w}(r, r_c) \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{k^2} \frac{\|\min\{r_c^2, r_c^{-2}\}(\tilde{P}(\rho, c) - \tilde{P}(\rho, z))\|_{L^\infty_{\tilde{w}}}}{\min\{r_c^2, r_c^{-2}\}} \tilde{w}(r, r_c) \\
& \lesssim \varepsilon \frac{\tilde{w}(r, r_c)}{\min\{r_c^2, r_c^{-2}\}}.
\end{aligned} \tag{4.80}$$

The \mathcal{V}_3 contribution is estimated as in \mathcal{V}_2 with the same bound. Lastly,

$$\begin{aligned}
\frac{\mathcal{I}_c[\partial_G \tilde{P}(r, c)] - \mathcal{I}_z[\partial_G \tilde{P}(r, z)]}{k^2 - 1} &= \mp i\varepsilon \int_r^{r_c} \frac{2(u(\rho) - c) \mp i\varepsilon}{\rho^3(u(\rho) - c)^2(u(\rho) - z)^2} \\
&\quad \times \int_\rho^{r_c} s(u(s) - c)^2 \partial_G \tilde{P}(s, c) ds d\rho \\
&\quad \pm i\varepsilon \int_r^{r_c} \frac{1}{\rho^3(u(\rho) - z)^2} \\
&\quad \times \int_\rho^{r_c} s[2(u(s) - c) \mp i\varepsilon] \partial_G \tilde{P}(s, c) ds d\rho \\
&\quad + \frac{\mathcal{I}_z[\partial_G \tilde{P}(r, z) - \partial_G \tilde{P}(r, c)]}{k^2 - 1}
\end{aligned} \tag{4.81}$$

The first two terms are analogous to \mathcal{V}_2 , estimated in (4.80), while we deal with the last term as in (4.78). Hence, appealing to (4.73a) we find

$$\begin{aligned}
& \frac{\mathcal{I}_c[\partial_G \tilde{P}](r, c) - \mathcal{I}_z[\partial_G \tilde{P}](r, z)}{k^2 - 1} - \frac{1}{A^2 - 1} \tilde{w}(r, r_c) \sup_s \frac{|\partial_G \tilde{P}(s, c) - \partial_G \tilde{P}(s, z)|}{\tilde{w}(s, r_c)} \\
& \lesssim \varepsilon k^2 \frac{\tilde{w}(r, r_c)}{\min\{r_c^2, r_c^{-2}\}}
\end{aligned}$$

Collecting all of the above

$$\begin{aligned}
& (\min\{r_c^2, r_c^{-2}\})^2 \left[\frac{|\partial_G \tilde{P}(r, c) - \partial_G \tilde{P}(r, z)|}{\tilde{w}(r, r_c)} - \frac{k^2 - 1}{A^2 - 1} \sup_s \frac{|\partial_G \tilde{P}(s, c) - \partial_G \tilde{P}(s, z)|}{\tilde{w}(s, r_c)} \right] \\
& \lesssim_{\varepsilon} \varepsilon k^2,
\end{aligned}$$

from which (4.74) with $j = 0$ easily follows. We now go back (4.75), and use (4.44), (4.49), (4.51) and (4.73a) to deduce (4.73b) by the methods above. From (4.75),

$$r \partial_r \partial_G \tilde{P}(r, c) - r \partial_r \partial_G \tilde{P}(r, z) = \mathcal{W}_1 + (k^2 - 1) [\mathcal{W}_2 + \mathcal{W}_3 + \mathcal{W}_4] \tag{4.82}$$

where

$$\begin{aligned}
\mathcal{W}_1 &= \frac{u''(r)}{(u'(r))^2} r \partial_r [\tilde{P}(r, c) - \tilde{P}(r, z)] \\
\mathcal{W}_2 &= \mp 3i\varepsilon \frac{2(u(r) - c) \mp i\varepsilon}{r^3 u'(r)(u(r) - c)^2(u(r) - z)^2} \int_r^{r_c} s(u(s) - c)^2 \tilde{P}(s, c) ds
\end{aligned}$$

$$\begin{aligned}
& \pm 3i\varepsilon \frac{1}{r^3 u'(r)(u(r) - z)^2} \int_r^{r_c} s [2(u(s) - c) \mp i\varepsilon] \tilde{P}(s, c) ds \\
& + \frac{3}{r^3 u'(r)(u(r) - z)^2} \int_r^{r_c} s (u(s) - z)^2 [\tilde{P}(s, c) - \tilde{P}(s, z)] ds \\
\mathcal{W}_3 = & \pm i\varepsilon \frac{2(u(r) - c) \mp i\varepsilon}{r^2(u(r) - c)^2(u(r) - z)^2} \int_r^{r_c} \partial_s \left(\frac{s}{u'(s)} \right) (u(s) - c)^2 \tilde{P}(s, c) ds \\
& \mp i\varepsilon \frac{1}{r^2(u(r) - z)^2} \int_r^{r_c} \partial_s \left(\frac{s}{u'(s)} \right) [2(u(s) - c) \mp i\varepsilon] \tilde{P}(s, c) ds \\
& - \frac{1}{r^2(u(r) - z)^2} \int_r^{r_c} \partial_s \left(\frac{s}{u'(s)} \right) (u(s) - z)^2 [\tilde{P}(s, c) - \tilde{P}(s, z)] ds \\
\mathcal{W}_4 = & \pm i\varepsilon \frac{2(u(r) - c) \mp i\varepsilon}{r^2(u(r) - c)^2(u(r) - z)^2} \int_r^{r_c} s (u(s) - c)^2 \partial_G \tilde{P}(s, c) ds \\
& \mp i\varepsilon \frac{1}{r^2(u(r) - z)^2} \int_r^{r_c} s [2(u(s) - c) \mp i\varepsilon] \partial_G \tilde{P}(s, c) ds \\
& - \frac{1}{r^2(u(r) - z)^2} \int_r^{r_c} s (u(s) - z)^2 \partial_G [\tilde{P}(s, c) - \tilde{P}(s, z)] ds
\end{aligned}$$

Arguing as above, we see that \mathcal{W}_1 is analogous to \mathcal{V}_1 , without a gain in k due to the absence of the integral, so that

$$\min\{r_c^2, r_c^{-2}\} |\mathcal{W}_1| \lesssim \varepsilon k^3 \frac{\tilde{w}(r, r_c)}{\min\{r_c^2, r_c^{-2}\}}.$$

Similarly, $\mathcal{W}_2, \mathcal{W}_3$ and \mathcal{W}_4 resemble \mathcal{V}_2 , provided we take into account the bounds (4.49), (4.59), (4.73), and (4.74) with $j = 0$, so that

$$\min\{r_c^2, r_c^{-2}\} |\mathcal{W}_2 + \mathcal{W}_3 + \mathcal{W}_4| \lesssim_{\varkappa} \varepsilon k^2 \frac{\tilde{w}(r, r_c)}{\min\{r_c^2, r_c^{-2}\}}.$$

The proof of Proposition 4.9 is now complete. \square

4.4 Analysis of $\partial_{r_c} \tilde{P}$

We proceed with the analysis of $\partial_{r_c} \tilde{P}$, relying on (4.36) and (4.37). The main result of this section reads as follows.

Proposition 4.10 (The ∂_{r_c} derivative). *Let \tilde{w} be given by (4.39) with $A = k + \varkappa$. Then,*

$$\|r_c \partial_{r_c} \tilde{P}\|_{L_{\tilde{w}}^\infty} \lesssim_{\varkappa} k^3, \quad (4.83)$$

and

$$\|\min\{r_c^2, r_c^{-2}\} r_c \partial_{r_c} (\tilde{P}(r, c \pm i\varepsilon) - \tilde{P}(r, c))\|_{L_{\tilde{w}}^\infty} \lesssim_{\varkappa} k^4 \varepsilon. \quad (4.84)$$

Moreover, for every $r_c > 0$ and every $r > 0$ such that

$$|r - r_c| > \frac{r_c}{k},$$

there holds

$$\frac{|r \partial_r r_c \partial_{r_c} \tilde{P}(r, z)|}{\tilde{w}(r, r_c)} \lesssim_{\mathcal{X}} k^4, \quad (4.85)$$

and

$$\min\{r_c^2, r_c^{-2}\} \frac{|r \partial_r r_c \partial_{r_c} \tilde{P}(r, c \pm i\varepsilon) - \tilde{P}(r, c)|}{\tilde{w}(r, r_c)} \lesssim_{\mathcal{X}} k^5 \varepsilon. \quad (4.86)$$

Proof of Proposition 4.10 The proof heavily relies on Lemma 4.5 and Propositions 4.6, 4.8 and 4.9, but the arguments are very similar to those used earlier. Moreover, we shall only deal with the case

$$|r - r_c| > \frac{r_c}{k}, \quad (4.87)$$

even for (4.83) and (4.84), since in the other case

$$r_c \partial_{r_c} = u'(r_c) r_c \partial_G - \frac{u'(r_c) r_c}{u'(r)} \partial_r \approx \min\{r_c^2, r_c^{-2}\} \partial_G + r \partial_r,$$

and hence the result follows from the respective bounds and convergence estimates on ∂_G and $r \partial_r$. For the sake of brevity, we consider only the case $r < r_c$, which from (4.87) implies that $r < b_k r_c$ with $b_k = 1 - 1/k > 0$. To prove (4.83), we need to show that

$$\frac{|r_c \partial_{r_c} \tilde{P}(r, z)|}{\tilde{w}(r, r_c)} \lesssim k^3, \quad \forall r, r_c > 0, \quad |r - r_c| > \frac{r_c}{k}.$$

We use (4.36), multiply by r_c and bound each term. As in (4.78),

$$|\mathcal{I}_z[r_c \partial_{r_c} \tilde{P}]| \leq \frac{k^2 - 1}{A^2 - 1} \tilde{w}(r, r_c) \sup_s \frac{|\partial_{r_c} \tilde{P}(s, z)|}{\tilde{w}(s, r_c)}.$$

The fact that the right-hand side above is finite for all $\varepsilon > 0$ follows from general ODE theory, since $\partial_{r_c} \tilde{P}$ satisfies an equation that is essentially a perturbation of Laplace's equation. Moreover,

$$\begin{aligned} r_c |\mathcal{I}_1 + \mathcal{I}_2| &\lesssim k^2 \|\tilde{P}\|_{L_w^\infty} |u'(r_c)| r_c \int_{b_k r_c}^{r_c} \frac{1}{\rho^3 |u(\rho) - z|} \int_\rho^{r_c} s \tilde{w}(s, r_c) ds d\rho \\ &\quad + k^2 \|\tilde{P}\|_{L_w^\infty} |u'(r_c)| r_c \int_r^{b_k r_c} \frac{1}{\rho^3 |u(\rho) - z|} \int_\rho^{r_c} s \tilde{w}(s, r_c) ds d\rho \\ &\lesssim k^3 \int_r^{r_c} \frac{\tilde{w}(\rho, r_c)}{\rho} d\rho \lesssim k^2 \tilde{w}(r, r_c), \end{aligned} \quad (4.88)$$

and

$$r_c |\mathcal{I}_3| \lesssim k^2 r_c^2 \int_r^{r_c} \frac{1}{\rho^3} d\rho \lesssim k^2 \frac{r_c^2}{r^2} \lesssim k^2 \tilde{w}(r, r_c).$$

Hence, (4.83) is a consequence of the above estimates. Combining (4.59) and (4.74), we obtain (4.84) when we restrict to the domain $|r - r_c| \leq r_c/k$. On the same region, to prove (4.84) we aim to show:

$$\min\{r_c^2, r_c^{-2}\} \frac{|r_c \partial_{r_c} (\tilde{P}(r, c \pm i\varepsilon) - \tilde{P}(r, c))|}{\tilde{w}(r, r_c)} \lesssim_{\mathcal{K}} k^4 \varepsilon. \quad (4.89)$$

Again, just consider the case when $0 < r < b_k r_c$. A combination of (4.36) and (4.37) allows us to write

$$\begin{aligned} \partial_{r_c} \tilde{P}(r, z) = & \mathcal{T}[\partial_{r_c} \tilde{P}] + (k^2 - 1)u'(r_c) \left[2 \int_r^{b_k r_c} \frac{1}{\rho^3(u(\rho) - z)^3} \right. \\ & \times \int_\rho^{r_c} s(u(s) - z)^2 \tilde{P}(s, z) ds d\rho \\ & - 2 \int_r^{b_k r_c} \frac{1}{\rho^3(u(\rho) - z)^2} \int_\rho^{r_c} s(u(s) - z) \tilde{P}(s, z) ds d\rho \\ & - \varepsilon^2 \frac{r_c}{u'(r_c)} \int_r^{b_k r_c} \frac{1}{\rho^3(u(\rho) - z)^2} d\rho \\ & + \frac{1}{(b_k r_c)^3 u'(b_k r_c)(u(b_k r_c) - z)^2} \int_{b_k r_c}^{r_c} \rho(u(\rho) - z)^2 \tilde{P}(\rho, z) d\rho \\ & + \int_{b_k r_c}^{r_c} \partial_\rho \left(\frac{1}{\rho^3 u'(\rho)} \right) \frac{1}{(u(\rho) - z)^2} \int_\rho^{r_c} s(u(s) - z)^2 \tilde{P}(s, z) ds d\rho \\ & + \int_{b_k r_c}^{r_c} \frac{1}{\rho^3(u(\rho) - z)^2} \int_\rho^{r_c} \partial_s \left(\frac{s}{u'(s)} \right) (u(s) - z)^2 \tilde{P}(s, z) ds d\rho \\ & \left. + \int_{b_k r_c}^{r_c} \frac{1}{\rho^3(u(\rho) - z)^2} \int_\rho^{r_c} \frac{s}{u'(s)} (u(s) - z)^2 \partial_s \tilde{P}(s, z) ds d\rho \right]. \end{aligned}$$

From this, we write

$$\partial_{r_c} \tilde{P}(r, c) - \partial_{r_c} \tilde{P}(r, z) = \mathcal{T}[\partial_{r_c} \tilde{P}(r, c)] - \mathcal{T}[\partial_{r_c} \tilde{P}(r, z)] + (k^2 - 1)u'(r_c) \sum_{i=1}^7 \mathcal{U}_i,$$

where

$$\begin{aligned} \mathcal{U}_1 = & \mp 2i\varepsilon \int_r^{b_k r_c} \frac{3(u(\rho) - c)^2 \mp 3i\varepsilon(u(\rho) - c) - \varepsilon^2}{\rho^3(u(\rho) - c)^3(u(\rho) - z)^3} \int_\rho^{r_c} s(u(s) - c)^2 \tilde{P}(s, c) ds d\rho \\ & \pm 2i\varepsilon \int_r^{b_k r_c} \frac{1}{\rho^3(u(\rho) - z)^3} \int_\rho^{r_c} s[2(u(s) - c) \mp i\varepsilon] \tilde{P}(s, c) ds d\rho \end{aligned}$$

$$\begin{aligned}
& + 2 \int_r^{b_{kr_c}} \frac{1}{\rho^3(u(\rho) - z)^3} \int_\rho^{r_c} s(u(s) - z)^2 [\tilde{P}(s, c) - \tilde{P}(s, z)] ds d\rho, \\
\mathcal{U}_2 = & \pm 2i\varepsilon \int_r^{b_{kr_c}} \frac{2(u(\rho) - c) \mp i\varepsilon}{\rho^3(u(\rho) - c)^2(u(\rho) - z)^2} \int_\rho^{r_c} s(u(s) - c) \tilde{P}(s, c) ds d\rho \\
& \mp 2i\varepsilon \int_r^{b_{kr_c}} \frac{1}{\rho^3(u(\rho) - z)^2} \int_\rho^{r_c} s \tilde{P}(s, c) ds d\rho \\
& - 2 \int_r^{b_{kr_c}} \frac{1}{\rho^3(u(\rho) - z)^2} \int_\rho^{r_c} s(u(s) - c) [\tilde{P}(s, c) - \tilde{P}(z, c)] ds d\rho, \\
\mathcal{U}_3 = & \varepsilon^2 \frac{r_c}{u'(r_c)} \int_r^{b_{kr_c}} \frac{1}{\rho^3(u(\rho) - z)^2} d\rho, \\
\mathcal{U}_4 = & \mp i\varepsilon \frac{2(u(b_{kr_c}) - c) \mp i\varepsilon}{(b_{kr_c})^3 u'(b_{kr_c})(u(b_{kr_c}) - z)^2(u(b_{kr_c}) - c)^2} \int_{b_{kr_c}}^{r_c} \rho(u(\rho) - c)^2 \tilde{P}(\rho, c) d\rho \\
& \pm i\varepsilon \frac{1}{(b_{kr_c})^3 u'(b_{kr_c})(u(b_{kr_c}) - z)^2} \int_{b_{kr_c}}^{r_c} \rho [2(u(\rho) - c) \mp i\varepsilon] \tilde{P}(\rho, c) d\rho \\
& + \frac{1}{(b_{kr_c})^3 u'(b_{kr_c})(u(b_{kr_c}) - z)^2} \int_{b_{kr_c}}^{r_c} \rho(u(\rho) - z)^2 [\tilde{P}(\rho, c) - \tilde{P}(\rho, z)] d\rho, \\
\mathcal{U}_5 = & \mp i\varepsilon \int_{b_{kr_c}}^{r_c} \partial_\rho \left(\frac{1}{\rho^3 u'(\rho)} \right) \frac{2(u(\rho) - c) \mp i\varepsilon}{(u(\rho) - z)^2(u(\rho) - c)^2} \int_\rho^{r_c} s(u(s) - c)^2 \tilde{P}(s, c) ds d\rho \\
& \pm i\varepsilon \int_{b_{kr_c}}^{r_c} \partial_\rho \left(\frac{1}{\rho^3 u'(\rho)} \right) \frac{1}{(u(\rho) - z)^2} \int_\rho^{r_c} s [2(u(s) - c) \mp i\varepsilon] \tilde{P}(s, c) ds d\rho \\
& + \int_{b_{kr_c}}^{r_c} \partial_\rho \left(\frac{1}{\rho^3 u'(\rho)} \right) \frac{1}{(u(\rho) - z)^2} \int_\rho^{r_c} s(u(s) - z)^2 [\tilde{P}(s, c) - \tilde{P}(s, z)] ds d\rho, \\
\mathcal{U}_6 = & \mp i\varepsilon \int_{b_{kr_c}}^{r_c} \frac{2(u(\rho) - c) \mp i\varepsilon}{\rho^3(u(\rho) - z)^2(u(\rho) - c)^2} \int_\rho^{r_c} \partial_s \left(\frac{s}{u'(s)} \right) (u(s) - c)^2 \tilde{P}(s, c) ds d\rho \\
& \pm i\varepsilon \int_{b_{kr_c}}^{r_c} \frac{1}{\rho^3(u(\rho) - z)^2} \int_\rho^{r_c} \partial_s \left(\frac{s}{u'(s)} \right) [2(u(s) - c) \mp i\varepsilon] \tilde{P}(s, c) ds d\rho \\
& + \int_{b_{kr_c}}^{r_c} \frac{1}{\rho^3(u(\rho) - z)^2} \int_\rho^{r_c} \partial_s \left(\frac{s}{u'(s)} \right) (u(s) - z)^2 [\tilde{P}(s, c) - \tilde{P}(s, z)] ds d\rho, \\
\mathcal{U}_7 = & \mp i\varepsilon \int_{b_{kr_c}}^{r_c} \frac{2(u(\rho) - c) \mp i\varepsilon}{\rho^3(u(\rho) - z)^2(u(\rho) - c)^2} \int_\rho^{r_c} \frac{s}{u'(s)} (u(s) - c)^2 \partial_s \tilde{P}(s, c) ds d\rho \\
& \pm i\varepsilon \int_{b_{kr_c}}^{r_c} \frac{1}{\rho^3(u(\rho) - z)^2} \int_\rho^{r_c} \frac{s}{u'(s)} [2(u(s) - c) \mp i\varepsilon] \partial_s \tilde{P}(s, c) ds d\rho \\
& + \int_{b_{kr_c}}^{r_c} \frac{1}{\rho^3(u(\rho) - z)^2} \int_\rho^{r_c} \frac{s}{u'(s)} (u(s) - z)^2 [\partial_s \tilde{P}(s, c) - \partial_s \tilde{P}(s, z)] ds d\rho.
\end{aligned}$$

Bounding these terms essentially relies repeatedly on Lemma 4.5, Propositions 4.6 and 4.8. Note that

$$\begin{aligned}
& \frac{\mathcal{T}_c[\partial_{r_c} \tilde{P}(r, c)] - \mathcal{T}_c[\partial_{r_c} \tilde{P}(r, z)]}{k^2 - 1} \\
& = \mp i\varepsilon \int_r^{r_c} \frac{2(u(\rho) - c) \mp i\varepsilon}{\rho^3(u(\rho) - c)^2(u(\rho) - z)^2} \int_\rho^{r_c} s(u(s) - c)^2 \partial_{r_c} \tilde{P}(s, c) ds d\rho
\end{aligned}$$

$$\begin{aligned} & \pm i\varepsilon \int_r^{r_c} \frac{1}{\rho^3(u(\rho) - z)^2} \int_\rho^{r_c} s [2(u(s) - c) \mp i\varepsilon] \partial_{r_c} \tilde{P}(s, c) ds d\rho \\ & + \frac{\mathcal{T}_z[\partial_{r_c} \tilde{P}(r, c) - \partial_{r_c} \tilde{P}(r, z)]}{k^2 - 1}. \end{aligned} \quad (4.90)$$

For the first two terms in (4.90) we obtain the bound

$$\begin{aligned} & \varepsilon \frac{\|r_c \partial_{r_c} \tilde{P}\|_{L_w^\infty}}{r_c} \int_{b_k r_c}^{r_c} \frac{1}{\rho^3 |u(\rho) - z|} \int_\rho^{r_c} s \tilde{w}(s, r_c) ds d\rho \\ & + \varepsilon \frac{\|r_c \partial_{r_c} \tilde{P}\|_{L_w^\infty}}{r_c} \int_r^{b_k r_c} \frac{1}{\rho^3 |u(\rho) - z|} \int_\rho^{r_c} s \tilde{w}(s, r_c) ds d\rho \\ & \lesssim \varepsilon \frac{k^3}{r_c^2} \int_{b_k r_c}^{r_c} \frac{|\rho - r_c|}{\rho |u(\rho) - z|} \tilde{w}(\rho, r_c) d\rho + \varepsilon \frac{k^2}{r_c} \frac{1}{|u(b_k r_c) - u(r_c)|} \int_r^{b_k r_c} \frac{\tilde{w}(\rho, r_c)}{\rho} d\rho \\ & \lesssim \varepsilon \frac{k^3}{r_c^2 |u'(r_c)|} \int_{b_k r_c}^{r_c} \frac{\tilde{w}(\rho, r_c) d\rho}{\rho} + \varepsilon \frac{k^2}{r_c} \frac{1}{|u(b_k r_c) - u(r_c)|} \int_r^{b_k r_c} \frac{\tilde{w}(\rho, r_c)}{\rho} d\rho \\ & \lesssim \varepsilon \frac{k^2}{r_c} \frac{\tilde{w}(r, r_c)}{|u'(r_c)| r_c} \lesssim k^2 \varepsilon \frac{\tilde{w}(r, r_c)}{r_c \min\{r_c^2, r_c^{-2}\}}, \end{aligned}$$

while for the last term we use (4.54) to obtain

$$\begin{aligned} \frac{\mathcal{T}_z[\partial_{r_c} \tilde{P}(r, c) - \partial_{r_c} \tilde{P}(r, z)]}{k^2 - 1} & \leq \sup_r |\partial_{r_c} \tilde{P}(r, z) - \partial_{r_c} \tilde{P}(r, c)| \\ & \quad \times \int_r^{r_c} \frac{1}{\rho^3} \int_\rho^{r_c} s \tilde{w}(s, r_c) ds d\rho \\ & \leq \frac{1}{A^2 - 1} \frac{\tilde{w}(r, r_c)}{r_c} \sup_r \frac{|r_c \partial_{r_c} \tilde{P}(r, z) - r_c \partial_{r_c} \tilde{P}(r, c)|}{\tilde{w}(r, r_c)}. \end{aligned}$$

Concerning the \mathcal{U}_i 's, for the first three we have

$$\begin{aligned} |\mathcal{U}_1 + \mathcal{U}_2| & \lesssim \varepsilon \|\tilde{P}\|_{L_w^\infty} \int_r^{b_k r_c} \frac{1}{\rho^3 |u(\rho) - z|^2} \int_\rho^{r_c} s \tilde{w}(s, r_c) ds d\rho \\ & \quad + \|\min\{r_c^2, r_c^{-2}\}(\tilde{P}(s, c) - \tilde{P}(s, z))\|_{L_w^\infty} \frac{1}{\min\{r_c^2, r_c^{-2}\}} \\ & \quad \times \int_r^{b_k r_c} \frac{1}{\rho^3 |u(\rho) - z|} \int_\rho^{r_c} s \tilde{w}(s, r_c) ds d\rho \\ & \lesssim \varepsilon \frac{1}{|u(b_k r_c) - u(r_c)|^2} \int_r^{b_k r_c} \frac{\tilde{w}(\rho, r_c)}{\rho} d\rho \\ & \quad + \varepsilon k \frac{1}{\min\{r_c^2, r_c^{-2}\}} \frac{1}{|u(b_k r_c) - u(r_c)|} \int_r^{b_k r_c} \frac{\tilde{w}(\rho, r_c)}{\rho} d\rho \end{aligned}$$

$$\lesssim \varepsilon k \frac{\tilde{w}(r, r_c)}{(\min\{r_c^2, r_c^{-2}\})^2},$$

and

$$\begin{aligned} |\mathcal{U}_3| &\lesssim \varepsilon \frac{r_c}{|u'(r_c)|} \int_r^{b_k r_c} \frac{1}{\rho^3 |u(\rho) - z|} d\rho \lesssim \varepsilon \frac{r_c}{|u'(r_c)| |u(b_k r_c) - u(r_c)|} \int_r^{b_k r_c} \frac{1}{\rho^3} d\rho \\ &\lesssim \varepsilon k \frac{\tilde{w}(r, r_c)}{(\min\{r_c^2, r_c^{-2}\})^2}. \end{aligned}$$

Concerning the others, we only show how to deal with \mathcal{U}_4 and \mathcal{U}_5 , as \mathcal{U}_6 and \mathcal{U}_7 are treated similarly. We have

$$\begin{aligned} |\mathcal{U}_4| &\lesssim \varepsilon \|\tilde{P}\|_{L_w^\infty} \frac{1}{|u'(r_c)| r_c^3 |u(b_k r_c) - u(r_c)|} \int_{b_k r_c}^{r_c} \rho \tilde{w}(\rho, r_c) d\rho \\ &\quad + \|\min\{r_c^2, r_c^{-2}\}(\tilde{P}(s, c) - \tilde{P}(s, z))\|_{L_w^\infty} \frac{1}{|u'(r_c)| r_c^3 \min\{r_c^2, r_c^{-2}\}} \int_{b_k r_c}^{r_c} \rho \tilde{w}(\rho, r_c) d\rho \\ &\lesssim \varepsilon k \frac{\tilde{w}(r, r_c)}{(\min\{r_c^2, r_c^{-2}\})^2}, \end{aligned}$$

and, considering that $|\partial_\rho (\rho^3 u'(\rho))^{-1}| \approx 1/\min\{\rho^5, \rho\}$, we also deduce that

$$\begin{aligned} |\mathcal{U}_5| &\lesssim \varepsilon \|\tilde{P}\|_{L_w^\infty} \int_{b_k r_c}^{r_c} \frac{1}{\min\{\rho^5, \rho\}} \frac{1}{|u(\rho) - z|} \int_\rho^{r_c} s \tilde{w}(s, r_c) ds d\rho \\ &\quad + \|\min\{r_c^2, r_c^{-2}\}(\tilde{P}(r, c) - \tilde{P}(r, z))\|_{L_w^\infty} \frac{1}{\min\{r_c^2, r_c^{-2}\}} \int_{b_k r_c}^{r_c} \frac{1}{\min\{\rho^5, \rho\}} \\ &\quad \times \int_\rho^{r_c} s \tilde{w}(s, r_c) ds d\rho \\ &\lesssim \varepsilon k \frac{1}{\min\{r_c^2, r_c^{-2}\}} \int_{b_k r_c}^{r_c} \frac{1}{\min\{\rho^3, \rho^{-1}\}} \tilde{w}(\rho, r_c) d\rho \lesssim \varepsilon k \frac{\tilde{w}(r, r_c)}{(\min\{r_c^2, r_c^{-2}\})^2}. \end{aligned}$$

Collecting all of the above, (4.89) follows. Going back to (4.34), we also infer that

$$\begin{aligned} \frac{r \partial_r r_c \partial_{r_c} \tilde{P}(r, z)}{k^2 - 1} &= -2 \frac{u'(r_c) r_c}{r^2 (u(r) - z)^3} \int_r^{r_c} s (u(s) - z)^2 \tilde{P}(s, z) ds + \varepsilon^2 \frac{r_c^2}{r^2 (u(r) - z)^2} \\ &\quad + 2 \frac{u'(r_c) r_c}{r^2 (u(r) - z)^2} \int_r^{r_c} s (u(s) - z) \tilde{P}(s, z) ds \\ &\quad - \frac{1}{r^2 (u(r) - z)^2} \int_r^{r_c} s (u(s) - z)^2 r_c \partial_{r_c} \tilde{P}(s, z) ds \end{aligned}$$

Hence, we argue as above to obtain (for $|r - r_c| \geq r_c/k$),

$$\begin{aligned} \left| \frac{rr_c \partial_r \partial_{r_c} \tilde{P}(r, z)}{k^2 - 1} \right| &\lesssim \|\tilde{P}\|_{L_w^\infty} \frac{\min\{r_c^2, r_c^{-2}\}}{r^2 |u(b_k r_c) - u(r_c)|} \int_r^{r_c} s \tilde{w}(s, r_c) ds + \frac{r_c^2}{r^2} \\ &\quad + \|r_c \partial_{r_c} \tilde{P}\|_{L_w^\infty} \frac{1}{r^2} \int_r^{r_c} s \tilde{w}(s, r_c) ds \\ &\lesssim k \|\tilde{P}\|_{L_w^\infty} \tilde{w}(r, r_c) + \tilde{w}(r, r_c) \\ &\quad + \frac{1}{k} \|r_c \partial_{r_c} \tilde{P}\|_{L_w^\infty} \tilde{w}(r, r_c) \lesssim k^2 \tilde{w}(r, r_c), \end{aligned}$$

and (4.85) follows. Lastly, the proof of (4.86) is simpler than the proof of (4.89), and is hence omitted. \square

4.5 Convergence in Optimal Weights

With the convergence estimates of the previous sections at hand, we now aim to show the validity of the convergence estimate in Theorem 4.3. The proof utilizes the Green's function of the operator

$$\partial_{rr} + \frac{1/4 - k^2}{r^2},$$

which is explicitly given by

$$\mathcal{L}(r, \rho) = -\frac{1}{2k} \min\left(\frac{\rho}{r}, \frac{r}{\rho}\right)^k \sqrt{r} \sqrt{\rho}, \quad r, \rho > 0, \quad (4.91)$$

as we treat the Rayleigh problem as a perturbation of the Laplacian. We split the proof in different cases.

4.5.1 The Case $r_c \leq 1$

All the estimates are pointwise in r_c , and hence the norms and spaces here are involving only the variable r . We begin by optimizing the weight near the origin.

Lemma 4.11 *Let $j = 0, 1$. There exists a universal constant $\zeta \in (0, 1/4)$ such that, for all $r_c \leq 1$ and $\varkappa \in (0, 1)$, there hold*

$$\left\| r^{k-1/2} (r \partial_r)^j P(r, c \pm i\varepsilon) \right\|_{L^\infty(0, \zeta r_c)} \lesssim_{\varkappa, \zeta} \frac{\varepsilon}{r_c^2} k^{1+j} r_c^{k-1/2}, \quad (4.92)$$

and

$$\left\| r^{k-1/2} (r \partial_r)^j (P(r, c \pm i\varepsilon) - P(r, c)) \right\|_{L^\infty(0, \zeta r_c)} \lesssim_{\varkappa, \zeta} \frac{\varepsilon}{r_c^2} k^{2+j} r_c^{k-1/2}. \quad (4.93)$$

Proof of Lemma 4.11 The proofs of (4.92) and (4.93) are very similar, so we focus on the more challenging (4.93). Note that the norm appearing on the left-hand side of (4.93) (and (4.92)) is a priori finite since RAY_z is a regular perturbation of Laplace's equation near $r \sim 0$.

Let us first consider the case $j = 0$. If $r \in (0, \zeta r_c)$, then (4.52) and (4.71) imply that

$$r^{k-1/2} |(r\partial_r)^j P(r, c \pm i\varepsilon)| \lesssim_{\mathcal{X}} k^{1+j} r_c^{k-1/2} \frac{r_c^{\mathcal{X}}}{r^{\mathcal{X}}}, \quad j = 0, 1, \quad (4.94)$$

$$r^{k-1/2} |(r\partial_r)^j (P(r, c \pm i\varepsilon) - P(r, c))| \lesssim_{\mathcal{X}} \varepsilon k^{2+j} r_c^{k-1/2} \frac{1}{r_c^{2-\mathcal{X}} r^{\mathcal{X}}}, \quad j = 0, 1. \quad (4.95)$$

Let $\chi = \chi(r/a)$ be a smooth cut-off function at some scale $a > 0$ to be determined. Define

$$g_\varepsilon(r, r_c) = \chi(r/a) [\phi(r, c \pm i\varepsilon) - \phi(r, c)]$$

and compute

$$\begin{aligned} \left(\partial_{rr} + \frac{1/4 - k^2}{r^2} \right) g_\varepsilon &= - \frac{\beta(r)}{u(r) - c \mp i\varepsilon} g_\varepsilon \\ &\quad - \frac{\pm i\varepsilon \beta(r)}{(u(r) - c \mp i\varepsilon)(u(r) - c)} \chi(r/a) \phi(r, c) \\ &\quad - \frac{2}{a} \partial_r \chi \partial_r (\phi(r, c \pm i\varepsilon) - \phi(r, c)) \\ &\quad - \frac{1}{a^2} \partial_{rr} \chi (\phi(r, c \pm i\varepsilon) - \phi(r, c)). \end{aligned}$$

Using (4.91), we then have

$$\begin{aligned} g_\varepsilon(r, r_c) &= - \int_0^\infty \mathcal{L}(r, \rho) \frac{\beta(\rho)}{u(\rho) - c \mp i\varepsilon} g_\varepsilon(\rho, r_c) d\rho \\ &\quad - \int_0^\infty \mathcal{L}(r, \rho) \frac{\pm i\varepsilon \beta(\rho)}{(u(\rho) - c \mp i\varepsilon)(u(\rho) - c)} \chi(\rho/a) \phi(\rho, c) d\rho \\ &\quad - \int_0^\infty \mathcal{L}(r, \rho) \left(\frac{2}{a} (\partial_\rho \chi) (\partial_\rho (\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c))) \right. \\ &\quad \left. + \frac{1}{a^2} (\partial_{\rho\rho} \chi) (\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c)) \right) d\rho \\ &= \sum_{\ell=1}^4 J_\ell. \end{aligned} \quad (4.96)$$

Since $\beta(\rho) \lesssim 1$ and $|u'(r_c)| \approx r_c$ as $r_c \rightarrow 0$, we have for $\zeta \in (0, 1/4)$ that

$$|u(\rho) - c \mp i\varepsilon| \gtrsim r_c^2, \quad \forall \rho \in (0, \zeta r_c).$$

Hence, by choosing $a = \frac{\zeta}{2} r_c$, with $\zeta \ll 1$, we obtain

$$\begin{aligned} r^{k-1/2} |J_1(r)| &\leq \frac{1}{2k} \int_0^r \rho \left| \frac{\beta(\rho)}{u(\rho) - c \mp i\varepsilon} \rho^{k-1/2} g_\varepsilon(\rho, r_c) \right| d\rho \\ &\quad + \frac{r^{2k}}{2k} \int_r^{2a} \rho^{-2k+1} \left| \frac{\beta(\rho)}{u(\rho) - c \mp i\varepsilon} \rho^{k-1/2} g_\varepsilon(\rho, r_c) \right| d\rho \\ &\leq \frac{1}{2} \left\| r^{k-1/2} g_\varepsilon(r, r_c) \right\|_{L^\infty(0, \zeta r_c)}. \end{aligned} \quad (4.97)$$

Turning to J_2 , we use (4.94) to obtain

$$\begin{aligned} r^{k-1/2} |J_2(r)| &\leq \frac{\varepsilon}{2k} \int_0^r \rho \left| \frac{\beta(\rho)}{u(\rho) - c \mp i\varepsilon} \rho^{k-1/2} P(\rho, c) \right| d\rho \\ &\quad + \varepsilon \frac{r^{2k}}{2k} \int_r^{2a} \rho^{-2k+1} \left| \frac{\beta(\rho)}{u(\rho) - c \mp i\varepsilon} \rho^{k-1/2} P(\rho, c) \right| d\rho \\ &\lesssim \frac{\varepsilon}{k r_c^2} \int_0^{2a} \rho \left| \rho^{k-1/2} P(\rho, c) \right| d\rho \\ &\lesssim \frac{\varepsilon}{r_c^{2-\varkappa}} r_c^{k-1/2} \int_0^{2a} \rho^{1-\varkappa} d\rho \lesssim_{\varkappa, \zeta} \varepsilon r_c^{k-1/2}. \end{aligned} \quad (4.98)$$

For J_3 , from (4.94)–(4.95) we obtain the pointwise bound for $\rho \in (0, \zeta r_c)$

$$\begin{aligned} |\rho \partial_\rho (\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c))| &\lesssim_\zeta r_c^2 |P(\rho, c \pm i\varepsilon) - P(\rho, c)| \\ &\quad + r_c^2 |\rho \partial_\rho (P(\rho, c \pm i\varepsilon) - P(\rho, c))| \\ &\quad + \varepsilon |\rho \partial_\rho P(\rho, c \pm i\varepsilon)| \\ &\lesssim_{\varkappa, \zeta} \varepsilon k^3 \frac{r_c^{k-1/2}}{r_c^{k-1/2}} \frac{r_c^\varkappa}{r_c^\varkappa}. \end{aligned}$$

Hence, arguing as above,

$$\begin{aligned} r^{k-1/2} |J_3(r)| &\leq \frac{1}{ak} \int_0^{2a} |\rho \partial_\rho (\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c))| \rho^{k-1/2} d\rho \\ &\lesssim_{\varkappa, \zeta} \frac{\varepsilon k^2}{a} r_c^{k-1/2} r_c^\varkappa \int_0^{2a} \rho^{-\varkappa} d\rho \lesssim_{\varkappa, \zeta} \varepsilon k^2 r_c^{k-1/2}. \end{aligned} \quad (4.99)$$

Finally, by (4.94) and (4.95) we obtain the pointwise bounds

$$|\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c)| \rho^{k-1/2} \lesssim_{\varkappa, \zeta} \varepsilon k^2 r_c^{k-1/2} \frac{r_c^\varkappa}{r_c^\varkappa}$$

and therefore

$$\begin{aligned} r^{k-1/2} |J_4(r)| &\leq \frac{1}{a^2 k} \int_0^{2a} |\rho(\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c))| \rho^{k-1/2} d\rho \\ &\lesssim_{\varkappa, \zeta} \frac{\varepsilon k}{a^2} r_c^{k-1/2} r_c^\varkappa \int_0^{2a} \rho^{1-\varkappa} d\rho \lesssim_{\varkappa, \zeta} \varepsilon k r_c^{k-1/2}, \end{aligned} \quad (4.100)$$

Hence, collecting (4.96)–(4.100), we arrive at

$$\left\| r^{k-1/2} g_\varepsilon(r, r_c) \right\|_{L^\infty(0, \zeta r_c)} \lesssim_{\varkappa, \zeta} \varepsilon k^2 r_c^{k-1/2}.$$

Note that g_ε and $\phi(r, c \pm i\varepsilon) - \phi(r, c)$ coincide in this region, and (4.93) is recovered from the definition of P , together with the inequality $|u(r) - c| \gtrsim r_c^2$ and a further application of (4.94). Finally, the case $j = 1$ follows immediately. Indeed, taking and $r \partial_r$ derivative of (4.96), we simply notice that

$$r \partial_r \mathcal{L}(r, \rho) = c_k \mathcal{L}(r, \rho), \quad c_k = \begin{cases} -(k-1/2), & \rho \leq r, \\ k+1/2, & \rho > r. \end{cases} \quad (4.101)$$

Therefore, the result follows in the exact same way, by using the estimates on g_ε derived above. \square

The interval $(\zeta r_c, R)$, for $R \gg 1$, independent of $r_c \leq 1$, is treated already by (4.71), which implies that for $r \in (\zeta r_c, r_c)$ there holds,

$$r^{k-1/2} |(r \partial_r)^j (P(r, c \pm i\varepsilon) - P(r, c))| \lesssim_{\varkappa, \zeta} \varepsilon k^{2+j} r_c^{k-1/2} \frac{1}{r_c^2}, \quad (4.102)$$

while if $r \in (r_c, R)$ there holds

$$r^{-k-1/2} |(r \partial_r)^j (P(r, c \pm i\varepsilon) - P(r, c))| \lesssim_{\varkappa, R} \varepsilon k^{2+j} r_c^{-k-1/2} \frac{1}{r_c^{2+\varkappa}}. \quad (4.103)$$

Finally, we need to correct the weight at infinity.

Lemma 4.12 *Let $j = 0, 1$. There exists a universal constant $R > 2$ such that, for all $r_c \leq 1$ and any $\varkappa \in (0, 1)$, there hold*

$$\left\| r^{-k-1/2} (r \partial_r)^j P(r, c \pm i\varepsilon) \right\|_{L^\infty(R, \infty)} \lesssim_{\varkappa, R} \frac{\varepsilon}{r_c^{2+\varkappa}} k^{1+j} r_c^{-k-1/2}$$

and

$$\left\| r^{-k-1/2} (r \partial_r)^j (P(r, c \pm i\varepsilon) - P(r, c)) \right\|_{L^\infty(R, \infty)} \lesssim_{\varepsilon, R} \frac{\varepsilon}{r_c^{2+\varkappa}} k^{2+j} r_c^{-k-1/2}.$$

Proof of Lemma 4.12 Again, we only treat the case $j = 0$. From (4.52) and (4.71), if $r \in (R, \infty)$ we have that

$$r^{-k-1/2} |(r \partial_r)^j P(r, c \pm i\varepsilon)| \lesssim_{\varepsilon} k^{1+j} r_c^{-k-1/2} \frac{r^\varkappa}{r_c^\varkappa}, \quad j = 0, 1, \quad (4.104)$$

$$r^{-k-1/2} |(r \partial_r)^j (P(r, c \pm i\varepsilon) - P(r, c))| \lesssim_{\varepsilon} \varepsilon k^{2+j} r_c^{-k-1/2} \frac{r^\varkappa}{r_c^{2+\varkappa}}, \quad j = 0, 1. \quad (4.105)$$

Let $\chi = \chi(r/R)$ be a smooth cut-off function, with $R > 0$ to be determined, and define

$$g_\varepsilon(r, r_c) = \chi(r/R) [\phi(r, c \pm i\varepsilon) - \phi(r, c)].$$

As in the proof of Lemma 4.11,

$$\begin{aligned} g_\varepsilon(r, r_c) &= - \int_0^\infty \mathcal{L}(r, \rho) \frac{\beta(\rho)}{u(\rho) - c \mp i\varepsilon} g_\varepsilon(\rho, r_c) d\rho \\ &\quad - \int_0^\infty \mathcal{L}(r, \rho) \frac{\pm i\varepsilon \beta(\rho)}{(u(\rho) - c \mp i\varepsilon)(u(\rho) - c)} \chi(\rho/R) \phi(\rho, c) d\rho \\ &\quad - \int_0^\infty \mathcal{L}(r, \rho) \left(\frac{2}{R} \partial_\rho \chi(\rho/R) \partial_\rho (\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c)) \right. \\ &\quad \left. - \frac{1}{R^2} \partial_{\rho\rho} \chi(\rho/R) (\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c)) \right) d\rho \\ &= \sum_{\ell=1}^4 J_\ell. \end{aligned} \quad (4.106)$$

For the first term, we use Lemma 2.1 that $|\beta(\rho)| \lesssim \langle \rho \rangle^{-6}$ and that $|u(\rho) - c \mp i\varepsilon| \gtrsim 1$ (thanks to our choice of $R > 2$ and $r_c \leq 1$) to obtain

$$\begin{aligned} r^{-k-1/2} |J_1(r)| &\leq r^{-k-1/2} \left| \int_0^\infty \mathcal{L}(r, \rho) \frac{\beta(\rho)}{u(\rho) - c \mp i\varepsilon} g_\varepsilon(\rho, r_c) d\rho \right| \\ &\lesssim \frac{1}{k} \int_R^\infty \frac{\rho \beta(\rho)}{|u(\rho) - u(r_c) \mp i\varepsilon|} \rho^{-k-1/2} |g_\varepsilon(\rho, r_c)| d\rho \\ &\lesssim R^{-6} \left\| r^{-k-1/2} g_\varepsilon(r, r_c) \right\|_{L^\infty(R, \infty)} \leq \frac{1}{2} \left\| r^{-k-1/2} g_\varepsilon(r, r_c) \right\|_{L^\infty(R, \infty)}, \end{aligned} \quad (4.107)$$

provided $R \gg 1$ is big enough. Regarding J_2 , from (4.104) we infer that

$$\begin{aligned} r^{-k-1/2} |J_2(r)| &\leq r^{-k-1/2} \left| \int_0^\infty \mathcal{L}(r, \rho) \frac{\pm i \varepsilon \beta(\rho)}{(u(\rho) - c \mp i \varepsilon)(u(\rho) - c)} \chi(\rho/R) \phi(\rho, c) d\rho \right| \\ &\lesssim \frac{\varepsilon}{k} \int_R^\infty \rho \beta(\rho) \rho^{-k-1/2} |P(\rho, c)| d\rho \\ &\lesssim_{\varkappa} \frac{\varepsilon r_c^{-k-1/2}}{r_c^{\varkappa}} \int_R^\infty \rho^{1+\varkappa} \rho^{-6} d\rho \lesssim_{R, \varkappa} \frac{\varepsilon}{r_c^{\varkappa}} r_c^{-k-1/2}. \end{aligned} \quad (4.108)$$

The terms involving $\partial_\rho \chi$ and $\partial_{\rho\rho} \chi$ are estimated similarly as in Lemma 4.11, except for the weight $r^{-k-1/2}$, and the fact that $\partial_\rho \chi$ and $\partial_{\rho\rho} \chi$ are supported in interval $[R, 2R]$. Using that u is bounded, $|u'(\rho)| \approx \rho^{-3}$ and (4.104)–(4.105), we obtain for $r_c < 1$ that

$$\begin{aligned} |\rho \partial_\rho (\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c))| &\lesssim \frac{1}{\rho^2} |P(\rho, c \pm i\varepsilon) - P(\rho, c)| \\ &\quad + |\rho \partial_\rho (P(\rho, c \pm i\varepsilon) - P(\rho, c))| \\ &\quad + \varepsilon |\rho \partial_\rho P(\rho, c \pm i\varepsilon)| \\ &\lesssim_{\varkappa} \varepsilon k^3 \frac{r_c^{-k-1/2}}{\rho^{-k-1/2}} \frac{\rho^{\varkappa}}{r_c^{2+\varkappa}}, \end{aligned}$$

and

$$\begin{aligned} |\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c)| &\lesssim |P(\rho, c \pm i\varepsilon) - P(\rho, c)| + \varepsilon |P(\rho, c \pm i\varepsilon)| \\ &\lesssim_{\varkappa} \varepsilon k^2 \frac{r_c^{-k-1/2}}{\rho^{-k-1/2}} \frac{\rho^{\varkappa}}{r_c^{2+\varkappa}}. \end{aligned}$$

Hence, we arrive at

$$\begin{aligned} r^{-k-1/2} |J_3(r)| &\leq \frac{1}{Rk} \int_R^{2R} |\rho \partial_\rho (\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c))| \rho^{-k-1/2} d\rho \\ &\lesssim_{\varkappa, R} \frac{\varepsilon}{r_c^{2+\varkappa}} k^2 r_c^{-k-1/2}, \end{aligned} \quad (4.109)$$

and

$$\begin{aligned} r^{-k-1/2} |J_4(r)| &\leq \frac{1}{R^2 k} \int_R^{2R} \rho |\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c)| \rho^{-k-1/2} d\rho \\ &\lesssim_{\varkappa, R} \frac{\varepsilon}{r_c^{2+\varkappa}} k r_c^{-k-1/2}. \end{aligned} \quad (4.110)$$

We then collect (4.106)–(4.110) to deduce that

$$\left\| r^{-k-1/2} g_\varepsilon(r, r_c) \right\|_{L^\infty(R, \infty)} \lesssim_{\varkappa, R} \frac{\varepsilon}{r_c^{2+\varkappa}} k^2 r_c^{-k-1/2},$$

since $r_c \leq 1$. Since, in this region, g_ε and $P(r, c \pm i\varepsilon) - P(r, c)$ satisfy the same estimates, the proof for $j = 0$ is over, while the case $j = 1$ follows again as in the previous lemma. \square

4.5.2 The Case $r_c > 1$

We now turn our attention to the case $r_c > 1$. Again, we will split in different cases. The proofs are similar as in the previous section, so we will only highlight the main differences. Since $r_c > 1$, (4.52) and (4.71) entail the following estimates in the case $r_c > 1$ (notice that the splitting of the interval $(0, \infty)$ slightly differs from the case $r_c \leq 1$):

$\diamond r \in (0, \zeta)$:

$$r^{k-1/2} |(r\partial_r)^j P(r, c \pm i\varepsilon)| \lesssim_{\varkappa} k^{1+j} r_c^{k-1/2} \frac{r_c^{\varkappa}}{r^{\varkappa}}, \quad j = 0, 1, \quad (4.111)$$

$$r^{k-1/2} |(r\partial_r)^j (P(r, c \pm i\varepsilon) - P(r, c))| \lesssim_{\varkappa} \varepsilon k^{2+j} r_c^{k-1/2} \frac{r_c^{2+\varkappa}}{r^{\varkappa}}, \quad j = 0, 1. \quad (4.112)$$

$\diamond r \in (\zeta, r_c)$:

$$r^{k-1/2} |(r\partial_r)^j (P(r, c \pm i\varepsilon) - P(r, c))| \lesssim_{\varkappa, \zeta} \varepsilon k^{2+j} r_c^{k-1/2} r_c^{2+\varkappa}, \quad j = 0, 1. \quad (4.113)$$

$\diamond r \in (r_c, Rr_c)$:

$$r^{-k-1/2} |(r\partial_r)^j (P(r, c \pm i\varepsilon) - P(r, c))| \lesssim_{\varkappa, R} \varepsilon k^{2+j} r_c^{-k-1/2} r_c^2, \quad j = 0, 1. \quad (4.114)$$

$\diamond r \in (Rr_c, \infty)$:

$$r^{-k-1/2} |(r\partial_r)^j P(r, c \pm i\varepsilon)| \lesssim_{\varkappa} k^{1+j} r_c^{-k-1/2} \frac{r_c^{\varkappa}}{r^{\varkappa}}, \quad j = 0, 1 \quad (4.115)$$

$$r^{-k-1/2} |(r\partial_r)^j (P(r, c \pm i\varepsilon) - P(r, c))| \lesssim_{\varkappa} \varepsilon k^{2+j} r_c^{-k-1/2} r_c^{\varkappa} r_c^{2-\varkappa}, \quad j = 0, 1. \quad (4.116)$$

We begin with the case $r \in (0, \zeta)$.

Lemma 4.13 *Let $j = 0, 1$. There exists a universal constant $\zeta \in (0, 1/4)$ such that, for all $r_c > 1$ and $\varkappa \in (0, 1)$, there hold*

$$\left\| r^{k-1/2} (r\partial_r)^j P(r, c \pm i\varepsilon) \right\|_{L^\infty(0, \zeta)} \lesssim_{\varkappa, \zeta} \varepsilon r_c^{2+\varkappa} k^{1+j} r_c^{k-1/2}.$$

and

$$\left\| r^{k-1/2} (r\partial_r)^j (P(r, c \pm i\varepsilon) - P(r, c)) \right\|_{L^\infty(0, \zeta)} \lesssim_{\varkappa, \zeta} \varepsilon r_c^{2+\varkappa} k^{2+j} r_c^{k-1/2}. \quad (4.117)$$

Proof of Lemma 4.13 The proof is similar to that of Lemma 4.11, so we only treat the case $j = 0$. As in Lemma 4.11, the norms appearing in this lemma are a priori finite. For a cut-off function $\chi = \chi(r/\zeta)$, the function

$$g_\varepsilon(r, r_c) = \chi(r/\zeta) [\phi(r, c \pm i\varepsilon) - \phi(r, c)]$$

can be written as

$$\begin{aligned} g_\varepsilon(r, r_c) &= - \int_0^\infty \mathcal{L}(r, \rho) \frac{\beta(\rho)}{u(\rho) - c \mp i\varepsilon} g_\varepsilon(\rho, r_c) d\rho \\ &\quad - \int_0^\infty \mathcal{L}(r, \rho) \frac{\pm i\varepsilon \beta(\rho)}{(u(\rho) - c \mp i\varepsilon)(u(\rho) - c)} \chi(\rho/\zeta) \phi(\rho, c) d\rho \\ &\quad - \int_0^\infty \mathcal{L}(r, \rho) \left(\frac{2}{\zeta} (\partial_\rho \chi) (\partial_\rho (\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c))) \right. \\ &\quad \left. + \frac{1}{\zeta^2} (\partial_{\rho\rho} \chi) (\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c)) \right) d\rho \\ &= \sum_{\ell=1}^4 J_\ell. \end{aligned} \quad (4.118)$$

Since $\beta(\rho) \lesssim 1$ and $|u(\rho) - c \mp i\varepsilon| \gtrsim 1$ in this regime, we choose $\zeta \ll 1$ to deduce that

$$r^{k-1/2} |J_1(r)| \leq \frac{1}{2} \left\| r^{k-1/2} g_\varepsilon(r, r_c) \right\|_{L^\infty(0, \zeta)}. \quad (4.119)$$

For J_2 , we use (4.111) to obtain

$$r^{k-1/2} |J_2(r)| \lesssim \frac{\varepsilon}{k} \int_0^\zeta \rho \left| \rho^{k-1/2} P(\rho, c) d\rho \right| d\rho \lesssim_{\varkappa, \zeta} \varepsilon r_c^\varkappa r_c^{k-1/2}, \quad (4.120)$$

For J_3 , we preliminary note that

$$\begin{aligned} \partial_\rho (\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c)) &= u'(\rho) (P(\rho, c \pm i\varepsilon) - P(\rho, c)) + (u(\rho) \\ &\quad - u(r_c)) \partial_\rho (P(\rho, c \pm i\varepsilon) - P(\rho, c)) \mp i\varepsilon \partial_\rho P(\rho, c \pm i\varepsilon), \end{aligned}$$

so that from (4.111)–(4.112) we obtain the pointwise bound for $\rho \in (0, \zeta)$

$$|\rho \partial_\rho (\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c))| \lesssim_{\varkappa, \zeta} \varepsilon k^3 \frac{r_c^{k-1/2}}{\rho^{k-1/2}} \frac{r_c^{2+\varkappa}}{\rho^\varkappa}.$$

Hence, arguing as above,

$$r^{k-1/2} |J_3(r)| \leq \frac{1}{\zeta k} \int_0^\zeta |\rho \partial_\rho (\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c))| \rho^{k-1/2} d\rho \lesssim_{\varkappa, \zeta} \varepsilon r_c^{2+\varkappa} k^2 r_c^{k-1/2}. \quad (4.121)$$

On the other hand, due to (4.111) and (4.112) we obtain the pointwise bounds

$$|\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c)|\rho^{k-1/2} \lesssim_{\varkappa, \zeta} \varepsilon k^2 r_c^{k-1/2} \frac{r_c^{2+\varkappa}}{\rho^\varkappa},$$

and therefore

$$r^{k-1/2} |J_4(r)| \leq \frac{1}{\zeta^2 k} \int_0^\zeta |\rho(\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c))| \rho^{k-1/2} d\rho \lesssim_{\varkappa, \zeta} \varepsilon r_c^{2+\varkappa} k r_c^{k-1/2}, \quad (4.122)$$

Hence, collecting (4.118)–(4.122), we arrive at

$$\left\| r^{k-1/2} g_\varepsilon(r, r_c) \right\|_{L^\infty(0, \zeta r_c)} \lesssim_{\varkappa, \zeta} \varepsilon r_c^{2+\varkappa} k^2 r_c^{k-1/2}.$$

Since, in this region, g_ε and $P(r, c \pm i\varepsilon) - P(r, c)$ satisfy the same estimates, the proof is over. \square

Also in this case, the regime $r \in (\zeta, Rr_c)$ for any $R \geq 1$ is already contained in (4.113) and (4.114). For $r \in (Rr_c, \infty)$, we follow the ideas in Lemma 4.12.

Lemma 4.14 *Let $j = 0, 1$. There exists a universal constant $R > 2$ such that, for all $r_c > 1$ and any $\varkappa \in (0, 1)$, there hold*

$$\left\| r^{-k-1/2} (r \partial_r)^j P(r, c \pm i\varepsilon) \right\|_{L^\infty(Rr_c, \infty)} \lesssim_{\varkappa, R} \varepsilon r_c^2 k^{1+j} r_c^{-k-1/2}.$$

and

$$\left\| r^{-k-1/2} (r \partial_r)^j (P(r, c \pm i\varepsilon) - P(r, c)) \right\|_{L^\infty(Rr_c, \infty)} \lesssim_{\varkappa, R} \varepsilon r_c^2 k^{2+j} r_c^{-k-1/2}.$$

Proof of Lemma 4.14 We use a cut-off of the form $\chi = \chi(r/Rr_c)$, with $R > 0$ to be determined, and define

$$g_\varepsilon(r, r_c) = \chi(r/Rr_c) [\phi(r, c \pm i\varepsilon) - \phi(r, c)].$$

Hence,

$$\begin{aligned} g_\varepsilon(r, r_c) &= - \int_0^\infty \mathcal{L}(r, \rho) \frac{\beta(\rho)}{u(\rho) - c \mp i\varepsilon} g_\varepsilon(\rho, r_c) d\rho \\ &\quad - \int_0^\infty \mathcal{L}(r, \rho) \frac{\pm i\varepsilon \beta(\rho)}{(u(\rho) - c \mp i\varepsilon)(u(\rho) - c)} \chi(\rho/Rr_c) \phi(\rho, c) d\rho \\ &\quad - \int_0^\infty \mathcal{L}(r, \rho) \left(\frac{2}{Rr_c} \partial_\rho \chi \partial_\rho (\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c)) \right. \\ &\quad \left. - \frac{1}{R^2 r_c^2} \partial_{\rho\rho} \chi (\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c)) \right) d\rho \\ &= \sum_{\ell=1}^4 J_\ell. \end{aligned} \quad (4.123)$$

Since $|u'(\rho)| \approx \rho^{-3}$ as $\rho \rightarrow \infty$, in this region we have that $|u(Rr_c) - u(r_c)| \gtrsim Rr_c^{-2}$. Hence, since $r_c > 1$, we can choose $R \gg 1$, independent of r_c , to have

$$\begin{aligned} r^{-k-1/2} |J_1(r)| &\lesssim \frac{1}{k} \int_{Rr_c}^{\infty} \frac{\rho \beta(\rho)}{|u(\rho) - u(r_c) \mp i\varepsilon|} \rho^{-k-1/2} |g_\varepsilon(\rho, r_c)| d\rho \\ &\lesssim \frac{r_c^2 \langle Rr_c \rangle^{-6}}{R} \left\| r^{-k-1/2} g_\varepsilon(r, r_c) \right\|_{L^\infty(Rr_c, \infty)} \\ &\leq \frac{1}{2} \left\| r^{-k-1/2} g_\varepsilon(r, r_c) \right\|_{L^\infty(Rr_c, \infty)}. \end{aligned} \quad (4.124)$$

For J_2 , from (4.115) we infer that,

$$r^{-k-1/2} |J_2(r)| \lesssim \frac{\varepsilon}{k} \int_{Rr_c}^{\infty} \rho \beta(\rho) \rho^{-k-1/2} |P(\rho, c)| d\rho \lesssim_{R, \varkappa} \varepsilon r_c^{-k-1/2}. \quad (4.125)$$

The terms involving $\partial_\rho \chi$ and $\partial_{\rho\rho} \chi$ are estimated similarly as in Lemma 4.11, except for the weight $r^{-k-1/2}$, and the fact that $\partial_\rho \chi$ and $\partial_{\rho\rho} \chi$ are supported in interval $[Rr_c, 2Rr_c]$.

Regarding J_3 and J_4 , using that $|u'(\rho)| \approx \rho^{-3}$ and (4.115)–(4.116) for $\rho \in (Rr_c, 2Rr_c)$ we obtain

$$|\rho \partial_\rho (\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c))| \lesssim_{\varkappa, R} \varepsilon k^3 \frac{r_c^{-k-1/2}}{\rho^{-k-1/2}},$$

and

$$|\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c)| \lesssim_{\varkappa, R} \varepsilon k^2 \frac{r_c^{-k-1/2}}{\rho^{-k-1/2}},$$

so that

$$\begin{aligned} r^{-k-1/2} |J_3(r)| &\leq \frac{1}{Rr_c k} \int_{Rr_c}^{2Rr_c} |\rho \partial_\rho (\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c))| \rho^{-k-1/2} d\rho \\ &\lesssim_{\varkappa, R} \varepsilon k^2 r_c^{-k-1/2}, \end{aligned} \quad (4.126)$$

and

$$\begin{aligned} r^{-k-1/2} |J_4(r)| &\leq \frac{1}{R^2 r_c^2 k} \int_{Rr_c}^{2Rr_c} \rho |\phi(\rho, c \pm i\varepsilon) - \phi(\rho, c)| \rho^{-k-1/2} d\rho \\ &\lesssim_{\varkappa, R} \varepsilon k r_c^{-k-1/2}. \end{aligned} \quad (4.127)$$

We then collect (4.123)–(4.127) to deduce that

$$\left\| r^{-k-1/2} g_\varepsilon(r, r_c) \right\|_{L^\infty(Rr_c, \infty)} \lesssim_{\varkappa, R} \varepsilon k^2 r_c^{-k-1/2},$$

since $r_c > 1$. Since, in this region, g_ε and $P(r, c \pm i\varepsilon) - P(r, c)$ differ by a factor proportional to r_c^2 , the proof is over. \square

Remark 4.15 The proof carries over to optimize the weights for $r_c \partial_{r_c} \tilde{P}$, for $j = 0, 1$. Indeed, from (4.1) we see that

$$\text{RAY}_z r_c \partial_{r_c} \phi = \frac{\beta(r) u'(r_c) r_c}{u(r) - z} \phi.$$

The extra singularity on the right-hand side is in fact innocuous. Indeed, Lemmas 4.11, 4.12, 4.13 and 4.14 are relevant in regions that are far from the critical layer, in which $|u(r) - z| \gtrsim \min\{r_c^2, r_c^{-2}\}$. Since $|u'(r_c)| r_c \approx \min\{r_c^2, r_c^{-2}\}$ as well, the contribution of the singular denominator cancels. Note that for $r \partial_r r_c \partial_{r_c} \tilde{P}$ we can simply differentiate the Green's function \mathcal{L} . However, we can only deduce information away from the critical layer, due to (4.86). We therefore arrive at the conclusion of the proof of Theorem 4.1.

4.6 Further Properties for the Real Solution

In this section, we prove Theorem 4.2.

4.6.1 The Function Q_0 and Its Properties

With Q_0 defined as in (4.6). From (4.1) we have that Q_0 obeys the second order equation (note that Q_0 is C^2 for $r < r_c$),

$$\begin{aligned} r(u(r) - c) \partial_{rr} Q_0 + \left(2ru'(r) + (1 - 2k)(u(r) - c) \right) \partial_r Q_0 \\ = 2(k + 1)u'(r)Q_0, \quad r < r_c, \end{aligned} \quad (4.128)$$

or, similarly,

$$\begin{aligned} \partial_r(r^2(u(r) - c) \partial_r Q_0) + \left(r^2 u'(r) - (1 + 2k)r(u(r) - c) \right) \partial_r Q_0 \\ = 2(k + 1)ru'(r)Q_0, \quad r > r_c. \end{aligned} \quad (4.129)$$

Note that we may rewrite (4.128) as

$$(u(r) - c) \left(r \partial_{rr} Q_0 + (1 - 2k) \partial_r Q_0 \right) = -2u'(r) \left(r \partial_r Q_0 - (k + 1) Q_0 \right). \quad (4.130)$$

Also, according to (4.30), (4.31) and (4.33), we have that

$$Q_0(r_c, c) = 1 \quad \text{and} \quad \partial_r Q_0(r_c, c) = (k + 1) r_c^{-1}. \quad (4.131)$$

Observe also that that (4.128) and (4.131) imply

$$\partial_{rr} Q_0(r_c, c) = \frac{(4k-1)\partial_r Q(r_c, c)}{3r_c} = \frac{(4k-1)(k+1)}{3r_c^2}. \quad (4.132)$$

We begin by proving (4.10). The fact that $Q_0(r, c) > 0$ for all $r \in (0, r_c]$ is clear from (4.35) and continuity in r , so we only need to prove monotonicity. From (4.131) it follows that $\partial_r Q_0(r_c, c) > 0$. Assume for the sake of contradiction that there exists a first point $r_* \in (0, r_c)$ (meaning closest to r_c) such that $\partial_r Q_0(r_*, c) = 0$.

Note that on $(r_*, r_c]$ we have $\partial_r Q_0 > 0$, and thus also $r^2(u(r) - c)\partial_r Q_0(r, c) > 0$ on (r_*, r_c) . By the minimality of r_* , we have that $r^2(u(r) - c)\partial_r Q_0(r, c)$ attains the value 0 for the first time (from r_c towards 0), and thus we must have

$$\partial_r(r^2(u(r) - c)\partial_r Q_0(r, c)) \Big|_{r=r_*} \geq 0. \quad (4.133)$$

On the other hand, evaluating (4.129) at $r = r_*$, and using that $\partial_r Q_0(r_*, c) = 0$ we obtain that

$$\partial_r(r^2(u(r) - c)\partial_r Q_0(r, c)) \Big|_{r=r_*} = (2(k+1)r_*u'(r_*))Q_0(r_*, c).$$

Since u is monotone decreasing, we immediately arrive at a contradiction with (4.133), and (4.10) follows.

To establish (4.12), we rely on the following lemma.

Lemma 4.16 *For $k \geq 2$, define the function let B_0 be defined as in (4.8). Then we have $B_0(r, c) \geq 0$ for all $r \in (0, r_c]$. As a consequence,*

$$r\partial_{rr} Q_0 + (1-2k)\partial_r Q_0 < 0 \quad (4.134)$$

on $(0, r_c]$. Moreover $B_0(r_c, c) = 0$ and the upper bound stated in (4.13) holds.

Proof of Lemma 4.16 Note that according to (4.131) we have that $B_0(r_c, c) = 0$. Then, upon differentiating, appealing to (4.130), we obtain that

$$(k-1)\partial_r Q_0 + \partial_r B_0 = (2k-1)\partial_r Q_0 - r\partial_{rr} Q_0 = \frac{-2u'(r)}{u(r)-c} B_0.$$

holds on $(0, r_c)$. Switching the order of the terms, we arrive at

$$-\partial_r((u(r)-c)^2 B_0) = (k-1)(u(r)-c)^2 \partial_r Q_0$$

which we may integrate from r to r_c and obtain

$$(u(r)-c)^2 B_0(r, c) = (k-1) \int_r^{r_c} (u(s)-c)^2 \partial_r Q_0(s, c) ds \geq 0, \quad (4.135)$$

appealing to that fact that $\partial_r Q_0(r, c) > 0$. Also, (4.134) follows immediately for $r \in (0, r_c)$, while for $r = r_c$ it requires (4.131) and (4.132). Finally, the fact that $B_0(r_c, c) = 0$ is a simple computation using (4.131). Now, from (4.8), the monotonicity of Q_0 and the positivity of B_0 , we have the upper bound in (4.12) for $r \partial_r Q_0$ as

$$0 \leq \partial_r Q_0(r, c) \leq (k+1) \frac{Q_0(r, c)}{r} \leq (k+1) \frac{Q_0(r_c, c)}{r} = \frac{k+1}{r}$$

while from (4.135) and the fact that u is decreasing we deduce that

$$\begin{aligned} (u(r) - c)^2 B_0(r, c) &\leq (k-1)(u(r) - c)^2 \int_r^{r_c} \partial_r Q_0(s, c) ds \\ &\leq (k^2 - 1)(u(r) - c)^2 \log(r_c/r), \end{aligned} \quad (4.136)$$

and the proof is complete. \square

Now, from (4.134), we obtain that

$$\partial_r \log(\partial_r Q_0) \leq \partial_r (\log r^{2k-1})$$

holds on $(0, r_c]$. Here we used that on $(0, r_c]$ we have $\partial_r Q_0(r, c) > 0$. Integrating the above from r to r_c , we infer

$$\partial_r Q_0(r, c) \geq \partial_r Q_0(r_c, c) \frac{r^{2k-1}}{r_c^{2k-1}}. \quad (4.137)$$

Inserting the above information in (4.135), which holds for all $r > 0$, we infer that

$$\begin{aligned} (k+1)Q_0(r, c) - r \partial_r Q_0(r, c) &= B_0(r, c) \\ &\geq \frac{(k-1)\partial_r Q_0(r_c, c)}{r_c^{2k-1}(u(r) - c)^2} \int_r^{r_c} (u(s) - c)^2 s^{2k-1} ds. \end{aligned} \quad (4.138)$$

In the above inequality, we drop the term $-r \partial_r Q_0(r, c) < 0$, and moreover (again by monotonicity) we have that $Q_0(r, c)$ converges, as $r \rightarrow 0$, to some limit $Q_0(0, c)$. Therefore, we have a lower bound on $Q_0(0, c)$ which is

$$Q_0(0, c) \geq \frac{(k-1)\partial_r Q_0(r_c, c)}{r_c^{2k-1}(k+1)(u(0) - c)^2} \int_0^{r_c} (u(s) - c)^2 s^{2k-1} ds. \quad (4.139)$$

We then arrive at the lower bound in (4.12) through a further use of the boundary condition (4.131). Regarding the upper bound, going back to (4.137) and re-arranging, we have an explicit lower bound on $\partial_r Q_0$, namely

$$\partial_r Q_0(r_c, c) r^{2k-1} < r_c^{2k-1} \partial_r Q_0(r, c).$$

By the fundamental theorem of calculus and using this lower bound, for any $s \in (0, r_c)$ we have

$$\begin{aligned} Q_0(r_c, c) - Q_0(s, c) &= \int_s^{r_c} \partial_r Q(r, c) dr > \partial_r Q_0(r_c, c) r_c^{1-2k} \int_s^{r_c} r^{2k-1} dr \\ &= \frac{\partial_r Q_0(r_c, c)}{2k r_c^{2k-1}} \left[r_c^{2k} - s^{2k} \right]. \end{aligned}$$

Hence, passing to the limit as $s \rightarrow 0$, we get the upper bound

$$Q_0(r_c, c) - \frac{\partial_r Q_0(r_c, c) r_c}{2k} > Q_0(0, c).$$

The left-hand side however, using (4.131) is

$$Q_0(r_c, c) - \frac{\partial_r Q_0(r_c, c) r_c}{2k} = 1 - \frac{k+1}{2k} = \frac{k-1}{2k},$$

from which the upper bound in (4.12) follows. The lower bound in (4.13) is proven in the next lemma.

Lemma 4.17 *There is a constant δ_0 depending only on u such that for all $r_c \in (0, \infty)$, there holds for all $|r - r_c| < \delta_0 r_c$ (uniformly in k and r_c),*

$$B_0(r, c) \gtrsim k^2 \frac{r^{2k-1}}{r_c^{2k}} (r_c - r), \quad \text{for } r \leq r_c.$$

Proof of Lemma 4.17 Since we are assuming that $|r - r_c| < \delta_0 r_c$, for δ_0 sufficiently small (depending only on u''), we have from Taylor's theorem that $u(r) - u(r_c) \approx u'(r_c)(r - r_c)$, and hence from (4.138) there holds

$$B_0(r, c) \gtrsim \frac{k^2 - 1}{r_c^{2k} |r - r_c|^2} \int_r^{r_c} (s - r_c)^2 s^{2k-1} ds.$$

This integral is explicitly computed via integration by parts:

$$\begin{aligned} \int_r^{r_c} (s - r_c)^2 s^{2k-1} ds &= -\frac{1}{2k} |r - r_c|^2 r^{2k} - \frac{2}{2k} \int_r^{r_c} (s - r_c) s^{2k} ds \\ &= -\frac{1}{2k} |r - r_c|^2 r^{2k} + \frac{2}{2k(2k+1)} (r - r_c) r^{2k+1} \\ &\quad + \frac{2}{2k(2k+1)(2k+2)} \left(r_c^{2k+2} - r^{2k+2} \right). \end{aligned}$$

By Taylor's theorem, there exists some $\zeta \in (r, r_c)$ such that

$$\begin{aligned} r_c^{2k+2} - r^{2k+2} &= (2k+2)r^{2k+1}(r_c - r) + \frac{1}{2}(2k+2)(2k+1)r^{2k}(r_c - r)^2 \\ &\quad + \frac{1}{3}(2k+2)(2k+1)2k\zeta^{2k-1}(r_c - r)^3, \end{aligned}$$

from which the conclusion follows since $\zeta \geq r$. \square

4.6.2 The Function Q_∞ and Its Properties

Analogous to (4.128), Q_∞ obeys the second order equation

$$r(u(r) - c)\partial_{rr}Q_\infty + \left(2ru'(r) + (1+2k)(u(r) - c)\right)\partial_rQ_\infty = -2(k-1)u'(r)Q_\infty, \quad (4.140)$$

or

$$\begin{aligned} \partial_r(r^2(u(r) - c)\partial_rQ_\infty) + \left(r^2u'(r) + (2k-1)r(u(r) - c)\right)\partial_rQ_\infty \\ = -2(k-1)ru'(r)Q_\infty. \end{aligned} \quad (4.141)$$

Note that we may rewrite (4.140) as

$$(u(r) - c)\left(r\partial_{rr}Q_\infty + (1+2k)\partial_rQ_\infty\right) = -2u'(r)\left(r\partial_rQ_\infty + (k-1)Q_\infty\right). \quad (4.142)$$

As conditions at the critical layer we have

$$Q_\infty(r_c, c) = 1, \quad \partial_rQ_\infty(r_c, c) = -(k-1)r_c^{-1}, \quad \partial_{rr}Q_\infty(r_c, c) = \frac{(4k+1)(k-1)}{3r_c^2}. \quad (4.143)$$

Proceeding as in the previous section, it is not hard to verify (4.11). Turning to (4.12), we have a lemma similar to Lemma 4.16 which we state without proof.

Lemma 4.18 *For $k \geq 2$, define the function let B_∞ be defined as in (4.9). Then we have $B_\infty(r, c) \geq 0$ for all $r \in [r_c, \infty)$. As a consequence,*

$$r\partial_{rr}Q_\infty + (1+2k)\partial_rQ_\infty < 0 \quad (4.144)$$

on $[r_c, \infty)$. Moreover $B_\infty(r_c, c) = 0$ and there holds the bound (4.14).

From (4.144) and arguing as in the previous section, we infer that

$$\begin{aligned} (k-1)Q_\infty(r, c) + r\partial_r Q_\infty(r, c) &= B_\infty(r, c) \\ &\geq -\frac{(k+1)\partial_r Q_\infty(r_c, c)r_c^{2k+1}}{(u(r) - c)^2} \int_{r_c}^r \frac{(u(s) - c)^2}{s^{2k+1}} ds. \end{aligned}$$

which implies the lower bound in (4.12). The upper bound follows similarly, while the lower bound for B_∞ in (4.14) is similar to that of B_0 .

5 The Inhomogeneous Rayleigh Problem for $k \geq 2$

Recall from (2.31), (2.32), and (2.33) how the reduction of order technique is used to derive two linearly independent homogeneous solutions (each satisfying one of the boundary conditions) H_0 , H_∞ , and their Wronskian M , and hence the Green's function for (2.8). In this section we will lay out a few technical results regarding M , H_0 , H_∞ , and of course by extension, \mathcal{G} . The properties of P (equivalently Q_0 and Q_∞) deduced in Sect. 4 are crucial.

Lemma 5.1 (Complex integral expansion). *For $\varepsilon > 0$ and $c \pm i\varepsilon = z \in I_\alpha$, define the following quantities,*

$$E(r, z) = \frac{1}{u'(r)} \partial_r \left(\frac{1}{u'(r)P(r, z)^2} \right) \quad (5.1a)$$

$$R_{a,b}^\varepsilon(z) = \int_a^b \frac{u'(r)(u(r) - c)}{(u(r) - c)^2 + \varepsilon^2} E(r, z) dr \quad (5.1b)$$

$$E_{a,b}^\varepsilon(z) = \int_a^b \frac{\varepsilon u'(r)}{(u(r) - c)^2 + \varepsilon^2} E(r, z) dr. \quad (5.1c)$$

Then, we have, for $z = c \pm i\varepsilon$,

$$H_0(r, z) = -\phi(s, z) \int_0^r \frac{1}{\phi^2(s, z)} ds = \frac{1}{u'(r)P(r, z)} - \phi(r, z) (R_{0,r}^\varepsilon(z) \pm iE_{0,r}^\varepsilon(z)) \quad (5.2a)$$

$$H_\infty(r, z) = \phi(s, z) \int_r^\infty \frac{1}{\phi^2(s, z)} ds = \frac{1}{u'(r)P(r, z)} + \phi(r, z) (R_{r,\infty}^\varepsilon(z) \pm iE_{r,\infty}^\varepsilon(z)). \quad (5.2b)$$

Similarly, there holds

$$M(z) = \int_0^\infty \frac{1}{\phi^2(s, z)} ds = R_{0,\infty}^\varepsilon(z) \pm iE_{0,\infty}^\varepsilon(z).$$

Proof of Lemma 5.1 The lemma follows by integration by parts in the complex integral:

$$\begin{aligned} \int_a^b \frac{1}{\phi(r, z)^2} dr &= \int_a^b \frac{1}{(u-z)^2 P(r, z)^2} dz = - \frac{1}{(u(r)-z)u'(r)P^2(r, z)} \Big|_a^b \\ &\quad + \int_a^b \frac{u'(r)(u-z)}{(u(r)-c)^2 + \varepsilon^2} E(r, z) dr. \end{aligned}$$

Note that boundary terms vanish when $a = 0$ or $b = \infty$ by the asymptotic behavior of P (Theorem 4.3). \square

Denote the formal limits of the above quantities:

$$\begin{aligned} E_{0,r}(c) &:= \pi E(r_c, c) \mathbf{1}_{r_c < r}, & E_{r,\infty}(c) &:= \pi E(r_c, c) \mathbf{1}_{r_c > r}, \\ R_{0,r}(c) &:= p.v. \int_0^r \frac{u'}{u-c} E(s, c) ds, & R_{r,\infty}(c) &:= p.v. \int_r^\infty \frac{u'}{u-c} E(s, c) ds. \end{aligned}$$

Here we are defining (and in the remainder of the paper as well)

$$p.v. \int_0^\infty \frac{u'(s)}{u(s)-c} g(s, c) ds := \lim_{\varepsilon' \rightarrow 0} \int_{|u(s)-c| \geq \varepsilon'} \frac{u'(s)}{u(s)-c} g(s, c) ds. \quad (5.3)$$

For $r < r_c$, by (2.1) (recall the definition of B_0 from Lemma 4.16)

$$\begin{aligned} E(r, z) &= \frac{\beta(r) Q_0(r, z) + 2 \frac{u'(r)}{r} ((1+k) Q_0(r, z) - r \partial_r Q_0(r, z))}{(u'(r))^3 Q_0^3(r, z) (r_c/r)^{2k-1}} \\ &= \frac{\beta(r) Q_0(r, z) + 2 \frac{u'(r)}{r} B_0(r, z)}{(u'(r))^3 Q_0^3(r, z) (r_c/r)^{2k-1}}, \end{aligned} \quad (5.4)$$

whereas for $r > r_c$ there holds (recall Lemma 4.18)

$$\begin{aligned} E(r, z) &= \frac{\beta Q_\infty(r, z) - 2 \frac{u'}{r} ((k-1) Q_\infty(r, z) + r \partial_r Q_\infty(r, z))}{(u'(r))^3 Q_\infty^3(r, z) (r/r_c)^{1+2k}} \\ &= \frac{\beta Q_\infty(r, z) - 2 \frac{u'}{r} B_\infty(r, z)}{(u'(r))^3 Q_\infty^3(r, z) (r/r_c)^{1+2k}}. \end{aligned} \quad (5.5)$$

Recall from Lemma 4.18 $B_\infty \geq 0$ and that $B_\infty(r_c, c) = 0$, and hence $E(r, c) < 0$ for $r \geq r_c$. However, from Lemma 4.16 $B_0 \geq 0$ and $B_0(r_c, c)$, and hence E is *not* sign definite for $r < r_c$. Finally, we note that there holds

$$E(r_c, c) = \frac{\beta(r_c)}{(u'(r_c))^3}.$$

5.1 Estimates on the Wronskian

First, we must deduce lower bounds on the Wronskian $M(c \pm i\varepsilon)$. Moreover, the gain as $r_c \rightarrow 0$ obtained in this lemma is crucial for deducing vorticity depletion, because it improves the bounds on \mathbb{K} in certain regions. This proof requires a delicate use of the monotonicity properties deduced on Q_0 and Q_∞ in Sect. 4. In particular, the lack of sign-definiteness of $E(r, c)$ on each side of the critical layer presents a complication for ruling out cancellations in the singular integrals which define $M(z)$.

Lemma 5.2 (Wronskian lower bounds). *For all $z \in I_\alpha$ with $z = c \pm i\varepsilon$ with ε sufficiently small (depending only on α and k), there holds the following lower bound,*

$$|M(c \pm i\varepsilon)| \gtrsim_\alpha k \max(r_c^{-3}, r_c^5), \quad (5.6)$$

and if we write $M(c \pm i0) = R_{0,\infty}(c) \pm iE_{0,\infty}(c)$, then the following uniform convergence holds for some sufficiently small $\eta > 0$

$$|M(c \pm i\varepsilon) - M(c \pm i0)| \lesssim \varepsilon^\eta \max(r_c^{-3}, r_c^5).$$

Proof of Lemma 5.2 For $\varepsilon > 0$ we have,

$$|M(c \pm i\varepsilon)|^2 = |R_{0,\infty}^\varepsilon(c \pm i\varepsilon)|^2 + |E_{0,\infty}^\varepsilon(c \pm i\varepsilon)|^2 \mp 2\operatorname{Im} E_{0,\infty}^\varepsilon \overline{R_{0,\infty}}(c \pm i\varepsilon). \quad (5.7)$$

We first study $E_{0,\infty}^\varepsilon(r, z)$ for $z = c \pm i\varepsilon$. First, for all ε sufficiently small and all sufficiently small $\eta > 0$, we prove

$$E_{0,\infty}^\varepsilon(z) = \frac{\pi\beta(r_c)}{(u'(r_c))^3} + \mathcal{O}\left(\varepsilon^\eta \max(r_c^5, r_c^{-3})\right). \quad (5.8)$$

Indeed, write

$$\begin{aligned} E_{0,\infty}^\varepsilon(z) &= \int_0^\infty \frac{\varepsilon u'(s)}{(u(s) - c)^2 + \varepsilon^2} E(s, c) ds \\ &\quad + \int_0^\infty \frac{\varepsilon u'(s)}{(u(s) - c)^2 + \varepsilon^2} (E(s, z) - E(s, c)) ds. \end{aligned} \quad (5.9)$$

Further decompose:

$$\begin{aligned} \int_0^\infty \frac{\varepsilon u'(s)}{(u(s) - c)^2 + \varepsilon^2} E(s, c) ds &= \int_0^\infty \frac{\varepsilon u'(s)}{(u(s) - c)^2 + \varepsilon^2} (\chi_c + \chi_{\neq}) E(s, c) ds \\ &= E_{0,\infty}^{\varepsilon;0c} + E_{0,\infty}^{\varepsilon;0\neq}. \end{aligned}$$

Further expand:

$$E_{0,\infty}^{\varepsilon;0c} = \frac{\beta(r_c)}{(u'(r_c))^3} \int_0^\infty \frac{\varepsilon u'(s)}{(u(s)-c)^2 + \varepsilon^2} \chi_c ds \\ + \int_0^\infty \frac{\varepsilon u'(s)}{(u(s)-c)^2 + \varepsilon^2} \chi_c \left(\frac{\beta(r_c)}{(u'(r_c))^3} - E(s, c) \right) ds.$$

By Theorem 4.3 (and (5.4), (5.5), and Lemma A.1),

$$\chi_c \left| \frac{\beta(r_c)}{(u'(r_c))^3} - E(s, c) \right| \lesssim \chi_c k^2 \frac{|r - r_c|}{r^2 |u'(r)|^2} \approx \chi_c k^2 |r - r_c| \max\left(\frac{1}{r_c^4}, r_c^4\right) \quad (5.10)$$

and hence by (A.5), for all $\eta > 0$ sufficiently small,

$$E_{0,\infty}^{\varepsilon;0c} = \frac{\beta(r_c)}{(u'(r_c))^3} + \mathcal{O}(\varepsilon^\eta \max(r_c^{-3}, r_c^5)).$$

Whereas, away from the critical layer there holds by (A.1),

$$\left| E_{0,\infty}^{\varepsilon;0\neq} \right| \lesssim \varepsilon k \max\left(\frac{1}{r_c^2}, r_c^2\right) \int_0^\infty \frac{|u'(s)|}{|u-c|} \chi_{\neq} \max\left(\frac{1}{s^3}, s^5\right) \min\left(\frac{s^{2k-1}}{r_c^{2k-1}}, \frac{r_c^{2k+1}}{s^{2k+1}}\right) ds \quad (5.11)$$

$$\lesssim \varepsilon k \max\left(\frac{1}{r_c^2}, r_c^2\right) \max(r_c^{-3}, r_c^5), \quad (5.12)$$

which goes into the error by the definition of I_α (2.18). This completes the estimate of the first term in (5.9). To control the latter term, we first decompose

$$\int_0^\infty \frac{\varepsilon u'(s)}{(u(s)-c)^2 + \varepsilon^2} (E(s, z) - E(s, c)) ds \quad (5.13)$$

$$= \int_0^\infty \frac{\varepsilon u'(s)}{(u(s)-c)^2 + \varepsilon^2} \frac{\beta(s)}{(u'(s))^3} \left(\frac{1}{P^2(s, z)} - \frac{1}{P^2(s, c)} \right) ds \quad (5.14)$$

$$+ \int_0^\infty \frac{\varepsilon u'(s)}{(u(s)-c)^2 + \varepsilon^2} \left(\left(E(s, z) - \frac{\beta(s)}{(u'(s))^3 P^2(s, z)} \right) \right. \\ \left. - \left(E(s, c) - \frac{\beta(s)}{(u'(s))^3 P^2(s, c)} \right) \right) ds \quad (5.15)$$

$$= E_{0,\infty}^{\varepsilon;10} + E_{0,\infty}^{\varepsilon;11}. \quad (5.16)$$

Then, by Theorem 4.3, for all $z \in I_\alpha$ and all sufficiently small $\eta > 0$ (using the same argument as (5.12))

$$\left| E_{0,\infty}^{\varepsilon;10} \right| \lesssim \varepsilon^\eta \int_0^\infty \frac{\varepsilon |u'(s)|}{(u(s)-c)^2 + \varepsilon^2} \max\left(\frac{1}{s^3}, s^3\right) \min\left(\frac{s^{2k-1}}{r_c^{2k-1}}, \frac{r_c^{2k+1}}{s^{2k+1}}\right) ds \\ \lesssim \varepsilon^\eta \max(r_c^3, r_c^{-3}).$$

The vanishing of $E_{0,\infty}^{\varepsilon;11}$ follows in a similar manner (due to the lack of convergence of (5.10) as $\varepsilon \rightarrow 0$, we use a trick to combine convergence and uniform Hölder regularity from (5.10); see e.g. the proofs of Lemmas A.4 or A.4 for details). This completes the proof of (5.8), which gives us a lower bound on $|E_{0,\infty}^{\varepsilon}(z)|$ as well as an upper bound on its contribution to the error in (5.7).

It is clear from (5.8) that we will need to make a detailed analysis on $R_{0,\infty}(z)$ to obtain (5.6). Indeed, $\beta(r_c)(u'(r_c))^{-3}$ potentially goes to zero very rapidly as $r_c \rightarrow \infty$ unless we impose stringent lower bounds on β as $r_c \rightarrow 0$ (note we wish to include the case $\beta(r) \sim e^{-r^2}$). Moreover, we would like to gain the power of k as well.

As above, we divide $R_{0,\infty}^{\varepsilon}$ based on the decompositions of $E_{0,\infty}^{\varepsilon}$ in (5.4) and (5.5):

$$\begin{aligned} R_{0,\infty}(c \pm i\varepsilon) &= \int_0^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \frac{\beta(s)}{(u'(s))^3 P^2(s, c \pm i\varepsilon)} ds \\ &\quad + \int_0^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \left(E(s, c \pm i\varepsilon) - \frac{\beta(s)}{(u'(s))^3 P^2(s, c \pm i\varepsilon)} \right) ds \\ &= R_{0,\infty}^{(1)}(c \pm i\varepsilon) + R_{0,\infty}^{(2)}(c \pm i\varepsilon). \end{aligned} \quad (5.17)$$

Denote the formal limit of the first term as

$$R_{0,\infty}^{(1)}(c) = p.v. \int_0^\infty \frac{u'}{u-c} \frac{\beta(s)}{(u'(s))^3 P^2(s, c)} ds.$$

Sub-divide via the critical layer:

$$\begin{aligned} R_{0,\infty}^{(1)}(c \pm i\varepsilon) &= \int_0^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} (\chi_c + \chi_{\neq}) \frac{\beta(s)}{(u'(s))^3 P^2(s, c \pm i\varepsilon)} ds \\ &= R_{0,\infty}^{(1,c)} + R_{0,\infty}^{(1,\neq)}; \end{aligned}$$

with analogous definitions also for the $\varepsilon = 0$ limit. For $R_{0,\infty}^{(1,\neq)}$ it is straightforward to verify the following from Theorem 4.3, (A.3), Lemma A.4, and $\beta(r) \lesssim \langle r \rangle^{-6}$ from Lemma 2.1, for all $z \in I_\alpha$ an all $\eta > 0$ sufficiently small (note that the integration gains k^{-1})

$$|R_{0,\infty}^{(1,\neq)}(c \pm i\varepsilon)| \lesssim \max(r_c^{-3}, r_c^3) \quad (5.18a)$$

$$|R_{0,\infty}^{(1,\neq)}(c \pm i\varepsilon) - R_{0,\infty}^{(1,\neq)}(c)| \lesssim \varepsilon^\eta \max(r_c^{-3}, r_c^3). \quad (5.18b)$$

Similarly, Theorem 4.3 and Lemma A.4 imply that for η sufficiently small there holds:

$$|R_{0,\infty}^{(1,c)}(c \pm i\varepsilon)| \lesssim \max(r_c^{-3}, r_c^3) \quad (5.19a)$$

$$|R_{0,\infty}^{(1,c)}(c \pm i\varepsilon) - R_{0,\infty}^{(1,c)}(c)| \lesssim \varepsilon^\eta \max(r_c^{-3}, r_c^3). \quad (5.19b)$$

Next, we consider the convergence and uniform estimates of the second term in (5.17), which we write as (in particular, note that since B_0 and B_∞ are Lipschitz continuous and vanish at the critical layer by Theorem 4.3),

$$R_{0,\infty}^{(2)}(c \pm i\varepsilon) = \int_0^{r_c} \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \left(2 \frac{B_0(s, z)}{(u'(s))^2 s P^3(s, z)} \right) ds \\ - \int_{r_c}^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \left(2 \frac{B_\infty(s, z)}{(u'(s))^2 s P^3(s, z)} \right) ds.$$

First, by following the same argument as that applied to $R_{0,\infty}^{(1)}(c \pm i\varepsilon)$ we deduce the analogous properties

$$\left| R_{0,\infty}^{(2)}(c \pm i\varepsilon) \right| \lesssim k \max(r_c^{-3}, r_c^5) \quad (5.20a)$$

$$\left| R_{0,\infty}^{(2)}(c \pm i\varepsilon) - R_{0,\infty}^{(2)}(c) \right| \lesssim \varepsilon^\eta \max(r_c^{-3}, r_c^5). \quad (5.20b)$$

Next, we establish the lower bound on $R_{0,\infty}^{(2)}$. From Theorem 4.2 we make the crucial observation that $R_{0,\infty}^{(2)}(c)$ is *strictly negative*. Further, the quantitative lower bounds on B_0 and B_∞ near the critical layer will allow us to deduce lower bounds. Using the negative definite signs and Theorem 4.2 (and Lemma 2.1):

$$R_{0,\infty}^{(2)}(c) \lesssim k^2 \int_{(1-\delta_0)r_c}^{r_c} \frac{u'}{u-c} \frac{1}{(u'(s))^2 s} \frac{s^{4k-2}}{r_c^{4k-1}} (r_c - s) ds \\ \lesssim -k^2 \int_{(1-\delta_0)r_c}^{r_c} \frac{1}{(u'(s))^2} \frac{s^{4k-3}}{r_c^{4k-1}} ds \lesssim -k \max(r_c^{-3}, r_c^5),$$

and hence by (5.20), we have for all $\varepsilon \geq 0$ sufficiently small,

$$R_{0,\infty}^{(2)}(c) \approx -k \max(r_c^{-3}, r_c^5) \quad (5.21a)$$

$$\left| R_{0,\infty}^{(2)}(c \pm i\varepsilon) \right| \approx k \max(r_c^{-3}, r_c^5). \quad (5.21b)$$

Putting together (5.21), (5.18), (5.19), and (5.8), for all small $\eta > 0$ we have,

$$\left| 2\operatorname{Im} E_{0,\infty}^\varepsilon \overline{R_{0,\infty}}(c \pm i\varepsilon) \right| \lesssim \varepsilon^\eta \max(r_c^{-3}, r_c^5)^2.$$

Next, we use (5.7) to deduce (5.6). For $r_c \gtrsim 1$, (5.8) is not useful, however, (5.21) together with (5.18) and (5.19) imply (5.6) for $r_c > R$ for R large enough depending only on u and universal constants. For $r_c < R$, (5.21) together with (5.18) and (5.19) imply (5.6) for $k > k_0$ for some k_0 depending only on u and universal constants. Whereas, for $r_c < R$ and $k < k_0$, the lower bound (5.6) follows from (5.8). \square

Lemma 5.3 *For $\varepsilon \ll 1$ and all $z \in I_\alpha$ there holds the following*

$$\left| r_c \partial_{r_c} M(z) \right| \lesssim k^3 \max(r_c^{-3}, r_c^5),$$

and we have that for all $z \in I_\alpha$ and $\eta > 0$ sufficiently small,

$$|r_c \partial_{r_c} M(c \pm i\varepsilon) - r_c \partial_{r_c} M(c \pm i0)| \lesssim \varepsilon^\eta k^3 \max(r_c^{-3}, r_c^5),$$

Proof of Lemma 5.3 By integration by parts

$$\begin{aligned} \frac{1}{u'(r_c)} \partial_{r_c} M(z) &= \int_0^\infty \frac{u'(s)}{(u-z)^2} \partial_G \frac{\chi_c}{u' P^2} ds + \frac{1}{u'(r_c)} \int_0^\infty \frac{1}{(u-z)^2} \partial_{r_c} \frac{\chi_{\neq}}{P^2} ds \\ &\quad + \int_0^\infty \frac{1}{(u-z)^3} \frac{\chi_{\neq}}{P^2} ds. \end{aligned}$$

From Theorem 4.3, we see that $r_c \partial_{r_c} P$ essentially satisfies the same upper bounds as P (up to powers of k), and hence the terms containing χ_{\neq} can be estimated directly using (A.1) and Lemma 2.1. Consider for example, the latter term (estimating as in (A.1)):

$$\begin{aligned} \left| \int_0^\infty \frac{1}{(u-z)^3} \frac{\chi_{\neq}}{P^2} ds \right| &\lesssim \int_0^\infty \chi_{\neq} \min \left(\frac{s^{2k-1}}{r_c^{2k-1}}, \frac{r_c^{2k+1}}{s^{2k+1}} \right) \\ &\quad \times \left(\mathbf{1}_{r_c \leq 1} \left(\mathbf{1}_{r \leq 2} \min \left(\frac{k^3}{r_c^6}, \frac{k^3}{r^6} \right) + \mathbf{1}_{r \geq 2} \right) \right. \\ &\quad \left. + \mathbf{1}_{r_c \geq 1} \left(\mathbf{1}_{r \leq 1/2} + \mathbf{1}_{r \geq 1/2} \min \left(k^3 r_c^6, k^3 r^6 \right) \right) \right) ds \\ &\lesssim k^2 \max(r_c^{-5}, r_c^7), \end{aligned} \quad (5.22)$$

which is consistent with the desired estimate (the additional power of k is lost in the term involving $\partial_{r_c} P$; see Theorem 4.3). Turn next to the term involving χ_c . Here, we integrate by parts again as in Lemma 5.1:

$$\int_0^\infty \frac{u'}{(u-z)^2} \partial_G \frac{\chi_c}{u' P^2} dr = \int_0^\infty \frac{u'}{u-z} \frac{1}{u'} \partial_r \left(\partial_G \frac{\chi_c}{u' P^2} \right) dr. \quad (5.23)$$

Note,

$$\begin{aligned} \frac{1}{u'} \partial_r \left(\partial_G \frac{\chi_c}{u' P^2} \right) &= \frac{1}{u'} \partial_r \left(\frac{\partial_G \chi_c}{u' P^2} - \frac{u'' \chi_c}{(u')^3 P^2} \right) - 2 \frac{1}{u'} \left(\partial_r \frac{\chi_c}{u'} \right) \frac{\partial_G P}{P^3} - 2 \frac{\chi_c \partial_r \partial_G P}{u' P^3} \\ &\quad + 6 \frac{\chi_c \partial_r P \partial_G P}{u' P^4}. \end{aligned} \quad (5.24)$$

Obtaining estimates on the contributions of the first two terms in (5.23) is essentially the same as the estimates made in Lemma 5.2. Hence, this is omitted for the sake of brevity. Next, consider the contribution of the term containing $\partial_r \partial_G P$. For this, from (4.24) we have,

$$|r_c u'(r_c)| \left| \int_0^\infty \frac{1}{u-z} \left(\frac{\chi_c \partial_r \partial_G P}{P^3} \right) dr \right| \lesssim k^3 \int_0^\infty \frac{\chi_c}{r_c^2 |u'(r_c)|} dr \lesssim \frac{k^2}{|r_c u'(r_c)|}.$$

For the term in (5.24) involving $r\partial_r$ we similarly use (4.21).

Next, consider the problem of convergence as $\varepsilon \rightarrow 0$. We define the expected limit as:

$$\begin{aligned} \frac{1}{u'(r_c)} \partial_{r_c} M(c \pm i0) &= p.v. \int_0^\infty \frac{u'}{u-c} \frac{1}{u'} \partial_r \left(\partial_G \frac{\chi_c}{u' P^2} \right) dr \\ &\quad \mp i\pi \left(\frac{1}{u'} \partial_r \left(\partial_G \frac{1}{u' P^2} \right) \right) (r_c, c) \\ &\quad + \frac{1}{u'(r_c)} \int_0^\infty \frac{1}{(u-c)^2} \partial_{r_c} \frac{\chi_{\neq}}{P^2} ds + \frac{1}{u'(r_c)} \int_0^\infty \frac{u'(r_c)}{(u-c)^3} \frac{\chi_{\neq}}{P^2} ds. \end{aligned}$$

Convergence of the terms involving ∂_G follows from Lemma A.4 and Theorem 4.3. Convergence away from the diagonal follows from Theorem 4.3 along with

$$\begin{aligned} \chi_{\neq} \left| \frac{1}{(u-c \mp i\varepsilon)^2} - \frac{1}{(u-c)^2} \right| &= \chi_{\neq} \left| \frac{\mp 2i\varepsilon(u-c) \pm \varepsilon^2}{(u-c \mp i\varepsilon)^2(u-c)^2} \right| \\ &\lesssim \varepsilon \max(r_c^{-2}, r_c^2) \frac{\chi_{\neq}}{|u-c|^2}, \end{aligned}$$

and the the analogous

$$\chi_{\neq} \left| \frac{1}{(u-c \mp i\varepsilon)^3} - \frac{1}{(u-c)^3} \right| \lesssim \varepsilon \max(r_c^{-2}, r_c^2) \frac{\chi_{\neq}}{|u-c|^3}.$$

From there, the estimate follows as in (5.22) and the assumption that $z \in I_\alpha$. \square

5.2 Estimates on H_0 and H_∞

Next, we outline the basic estimates available on H_0 and H_∞ ; see “Appendix C” for proof sketches, as most are minor refinements of ideas appearing in the proofs of Lemmas 5.2 and 5.3.

Lemma 5.4 (Estimates on H_0 and H_∞). *Let $z = c \pm i\varepsilon \in I_\alpha$.*

- (a) *(explicit expansions near the critical layer) In the region $|r - r_c| < r_c/k$, there holds:*

$$H_0(r, c \pm i\varepsilon) = \frac{1}{u'(r)P(r, c \pm i\varepsilon)} - \phi(r, c \pm i\varepsilon) \left(R_{0,r}(c \pm i\varepsilon) \mp iE_{0,r}^\varepsilon(c \pm i\varepsilon) \right) \quad (5.25a)$$

$$H_\infty(r, c \pm i\varepsilon) = \frac{1}{u'(r)P(r, c \pm i\varepsilon)} + \phi(r, c \pm i\varepsilon) \left(R_{r,\infty}(c \pm i\varepsilon) \mp iE_{r,\infty}^\varepsilon(c \pm i\varepsilon) \right), \quad (5.25b)$$

and we record the following estimates: in the region $|r - r_c| < r_c/k$, we write

$$\frac{1}{u'(r)P(r, c \pm i\varepsilon)} \approx \frac{1}{u'(r_c)} \approx -\max(r_c^{-1}, r_c^3),$$

and

$$\begin{aligned} & |\phi(r, c \pm i\varepsilon) (R_{0,r}^\varepsilon(c \pm i\varepsilon) \mp iE_{0,r}^\varepsilon(c \pm i\varepsilon))| \\ & \lesssim \mathbf{1}_{r_c \leq 1} \frac{|r - r_c|}{r_c^2} \left(k + \left| \log k \left| \frac{r - r_c}{r_c} \right| \right| \right) \end{aligned} \quad (5.26a)$$

$$+ \mathbf{1}_{r_c > 1} r_c^3 \frac{|r - r_c|}{r_c} \left(k + \left| \log k \left| \frac{r - r_c}{r_c} \right| \right| \right), \quad (5.26b)$$

$$\begin{aligned} & |\phi(r, c \pm i\varepsilon) (R_{r,\infty}^\varepsilon(c \pm i\varepsilon) \mp iE_{r,\infty}^\varepsilon(c \pm i\varepsilon))| \\ & \lesssim \mathbf{1}_{r_c \leq 1} \frac{|r - r_c|}{r_c^2} \left(k + \left| \log k \left| \frac{r - r_c}{r_c} \right| \right| \right) \end{aligned} \quad (5.26c)$$

$$+ \mathbf{1}_{r_c > 1} r_c^3 \frac{|r - r_c|}{r_c} \left(k + \left| \log k \left| \frac{r - r_c}{r_c} \right| \right| \right). \quad (5.26d)$$

In particular, for $|r - r_c| \leq r_c/k$,

$$|H_0(r, c \pm i\varepsilon)| + |H_\infty(r, c \pm i\varepsilon)| \lesssim \max(r_c^{-1}, r_c^3). \quad (5.27)$$

(b) (bounds away from the critical layer) In the region $|r - r_c| \geq r_c/k$, there holds

$$\begin{aligned} |H_0(r, c \pm i\varepsilon)| & \lesssim \mathbf{1}_{r_c \leq 1} \left(\mathbf{1}_{r < r_c} \frac{r_c^{k+1/2}}{r_c^{k+3/2}} + \mathbf{1}_{r_c < r < 1} k \frac{r_c^{k+2+1/2}}{r_c^{k+2+3/2}} + \mathbf{1}_{r > 1} k \frac{r_c^{k+1/2}}{r_c^{k+3+1/2}} \right) \\ & + \mathbf{1}_{r_c > 1} \left(\mathbf{1}_{r < 1} \frac{r_c^{k+1/2}}{r_c^{k-1/2}} + \mathbf{1}_{1 < r < r_c} \frac{r_c^{k+5/2}}{r_c^{k-1/2}} + \mathbf{1}_{r > r_c} k \frac{r_c^{k+1/2}}{r_c^{k-5/2}} \right), \end{aligned} \quad (5.28a)$$

$$\begin{aligned} |H_\infty(r, c \pm i\varepsilon)| & \lesssim \mathbf{1}_{r_c \leq 1} \left(\mathbf{1}_{r < r_c} k \frac{r_c^{k-3/2}}{r_c^{k-1/2}} + \mathbf{1}_{r_c < r < 1} \frac{r_c^{k+1/2}}{r_c^{k+3/2}} + \mathbf{1}_{r > 1} \frac{r_c^{k+1/2}}{r_c^{k-1/2}} \right) \\ & + \mathbf{1}_{r_c > 1} \left(\mathbf{1}_{r < 1} k \frac{r_c^{k+5-1/2}}{r_c^{k-1/2}} + \mathbf{1}_{1 \leq r < r_c} k \frac{r_c^{k+5-1/2}}{r_c^{k+3/2}} + \mathbf{1}_{r < r_c} \frac{r_c^{k+5/2}}{r_c^{k-1/2}} \right). \end{aligned} \quad (5.28b)$$

(c) (convergence) Furthermore, if we define

$$\begin{aligned} H_0(r, c \pm i0) & = \frac{1}{u'(r)P(r, c)} - \phi(r, c) (R_{0,r}(c) \pm i\pi E(r_c) \mathbf{1}_{r_c < r}) \\ H_\infty(r, c \pm i0) & = \frac{1}{u'(r)P(r, c)} + \phi(r, c) (R_{r,\infty}(c) \pm i\pi E(r_c) \mathbf{1}_{r_c > r}), \end{aligned}$$

then we have that $H_0(r, c \pm i0)$ and $H_\infty(r, c \pm i0)$ satisfy the same pointwise estimates as the $\varepsilon > 0$ counterparts and there exists an $\eta > 0$ such that for $z \in I_\alpha$, the difference $\varepsilon^{-\eta} (H_0(r, c \pm i\varepsilon) - H_0(r, c \pm i0))$ satisfies (5.27) and (5.28a) (and analogously for H_∞).

The next lemma details analogous properties on the derivatives of H_0 and H_∞ .

Lemma 5.5 *Let $z = c \pm i\varepsilon \in I_\alpha$.*

(a) *(explicit expansions near critical layer) for ∂_r derivatives there holds for $|r - r_c| \leq r_c/k$:*

$$r \partial_r H_0(r, c \pm i\varepsilon) = r \partial_r \left(\frac{1}{u' P(r, c \pm i\varepsilon)} \right) - r \partial_r \phi \left(R_{0,r}^\varepsilon(z) \mp i E_{0,r}^\varepsilon \right) - r u'(r) P E(r, c \pm i\varepsilon), \quad (5.29a)$$

$$r \partial_r H_\infty(r, c \pm i\varepsilon) = r \partial_r \left(\frac{1}{u' P(r, c \pm i\varepsilon)} \right) + r \partial_r \phi \left(R_{r,\infty}^\varepsilon(z) \mp i E_{r,\infty}^\varepsilon \right) - r u'(r) P E(r, c \pm i\varepsilon); \quad (5.29b)$$

for ∂_G derivatives there holds (for $|r - r_c| \leq r_c/2k$):

$$\begin{aligned} H_0(r, z) &= \partial_G \left(\frac{1}{u' P(r, z)} \right) - (u - z) (\partial_G P) \left(R_{0,r}^\varepsilon(z) \mp i E_{0,r}^\varepsilon \right) \\ &\quad - \phi \int_0^r \frac{u'(s)}{(u - z)} \partial_G (\chi_c E) \, ds \\ &\quad + \frac{1}{u'(r_c)} \phi \int_0^r \chi_{\neq} \frac{u'(s) u'(r_c)}{(u - z)^2} E(s, z) - \frac{u'(s)}{(u - z)} \partial_{r_c} (\chi_{\neq} E(s, z)) \, ds. \end{aligned} \quad (5.30a)$$

$$\begin{aligned} H_\infty(r, z) &= \partial_G \left(\frac{1}{u' P(r, c \pm i\varepsilon)} \right) + (u - z) (\partial_G P) \left(R_{r,\infty}^\varepsilon(z) \mp i E_{r,\infty}^\varepsilon \right) \\ &\quad + \phi \int_r^\infty \frac{u'(s)}{(u - z)} \partial_G (\chi_c E) \, ds \\ &\quad - \frac{1}{u'(r_c)} \phi \int_r^\infty \chi_{\neq} \frac{u'(s) u'(r_c)}{(u - z)^2} E(s, z) - \frac{u'(s)}{(u - z)} \partial_{r_c} (\chi_{\neq} E(s, z)) \, ds. \end{aligned} \quad (5.30b)$$

Furthermore, for $|r - r_c| < r_c/k$, there holds the estimates:

$$\begin{aligned} &(|r \partial_r H_\infty(r, z)| + |r \partial_r H_0(r, z)|) \mathbf{1}_{|r - r_c| < r_c/k} \\ &\lesssim \max(r_c^{-1}, r_c^3) \left(k + \left| \log \frac{k |r - r_c|}{r_c} \right| \right) \end{aligned} \quad (5.31a)$$

$$\begin{aligned} &(|\partial_G H_\infty(r, z)| + |\partial_G H_0(r, z)|) \mathbf{1}_{|r - r_c| < r_c/k} \\ &\lesssim k \frac{\max(r_c^{-1}, r_c^3)}{|r_c u'(r_c)|} \end{aligned} \quad (5.31b)$$

$$(|r\partial_r\partial_G H_\infty(r, z)| + |r\partial_r\partial_G H_0(r, z)|) \mathbf{1}_{|r-r_c| < r_c/k} \\ \lesssim k \frac{\max(r_c^{-1}, r_c^3)}{|r_c u'(r_c)|} \left(k + \left| \log \frac{k|r-r_c|}{r_c} \right| \right). \quad (5.31c)$$

- (b) (*derivative bounds away from critical layer*) For $|r - r_c| \geq r_c/k$, we have that $|r - r_c| \geq r_c/k$, $k^{-1}r\partial_r H_0$, $k^{-3}r_c\partial_{r_c} H_0$, $k^{-4}r\partial_r r_c\partial_{r_c} H_0$ satisfy the estimate (5.28a) whereas $k^{-1}r\partial_r H_\infty$, $k^{-3}r_c\partial_{r_c} H_\infty$, and $k^{-4}r\partial_r r_c\partial_{r_c} H_\infty$ satisfy the estimate (5.28b).
- (c) (*convergence*) For some $\eta > 0$, for $z \in I_\alpha$ we have the following convergence: $\varepsilon^{-\eta} (H_\infty(r, c \pm i\varepsilon) - H_\infty(r, c \pm i0))$ and $\varepsilon^{-\eta} (H_0(r, c \pm i\varepsilon) - H_0(r, c \pm i0))$ satisfy (5.31) and the assertions in part (b) for ε sufficiently small.

6 Representation Formulas and Estimates on $(r\partial_r)^j f_1$, and $(r\partial_r)^j f_2$

6.1 Recursion Relations for Derivatives of f_1 and f_2

Next, our goal is to derive formulas for the derivatives of f_1 and f_2 . As discussed in Sect. 2.3.4, the first step is the iteration scheme outlined in Lemma 2.8.

Lemma 6.1 (Iteration lemma for $\partial_G^j X$ and $\partial_G^j Y$). *Set*

$$\mathcal{E}_0 = 0, \quad F_0 = F(r), \quad R_0 = F_*(r), \quad R_0^x = 0. \quad (6.1)$$

For $j \geq 0$ if we define

$$\begin{aligned} F_{j+1}^\pm &= \partial_G F_j^\pm - \frac{2u''}{(u')^2} (F_j - \beta \partial_G^j Y^\pm) - \frac{\beta'}{u'} \partial_G^j Y^\pm \\ R_{j+1}^\pm &= \partial_G R_j - \frac{2u''}{(u')^2} R_j + 2 \left(\frac{1}{4} - k^2 \right) \left(\frac{u' + ru''}{r(u')^2} \right) \frac{\partial_G^j Y^\pm}{r^2} + \partial_{rr} \left(\frac{1}{u'} \right) \partial_r \partial_G^j Y^\pm \\ \mathcal{E}_{j+1} &= \partial_G \mathcal{E}_j - \frac{2u''}{(u')^2} (\mathcal{E}_j - \beta \partial_G^j X) - \frac{\beta'}{u'} \partial_G^j X \\ R_{j+1}^x &= \partial_G R_j^x - \frac{2u''}{(u')^2} R_j^x + 2 \left(\frac{1}{4} - k^2 \right) \left(\frac{u' + ru''}{r(u')^2} \right) \frac{\partial_G^j X}{r^2} + \partial_{rr} \left(\frac{1}{u'} \right) \partial_r \partial_G^j X_j, \end{aligned}$$

then (2.35) holds for all j .

Proof of Lemma 6.1 Recall that ∂_G commutes with functions of $u - c$. Hence, the lemma follows from the following relation:

$$\begin{aligned} [\partial_G, \text{RAY}_\pm] Z &= -\partial_{rr} \left(\frac{1}{u'} \right) \partial_r Z + \frac{2u''}{(u')^2} \text{RAY}_\pm Z - 2 \left(\frac{1}{4} - k^2 \right) \left(\frac{u' + ru''}{r(u')^2} \right) \frac{Z}{r^2} \\ &\quad + \left(\frac{\beta'}{u'} - \frac{2u''\beta}{(u')^2} \right) \frac{Z}{u - z}. \end{aligned} \quad (6.2)$$

□

Lemma 6.2 For coefficients $e_{j,\ell}$, $h_{j,\ell}$, $q_{j,\ell}$, $p_{j,\ell}$, and $r_{j,\ell}$ we have

$$R_{j+1} = \sum_{\ell=0}^{j+1} e_{j+1,\ell} \partial_G^\ell F_* + \sum_{\ell=0}^j q_{j+1,\ell} \partial_r \partial_G^\ell Y^\pm + p_{j+1,\ell} \partial_G^\ell Y^\pm, \quad (6.3a)$$

$$R_{j+1}^x = \sum_{\ell=0}^j q_{j+1,\ell} \partial_r \partial_G^\ell X^\pm + p_{j+1,\ell} \partial_G^\ell X^\pm, \quad (6.3b)$$

$$F_{j+1}^\pm = \sum_{\ell=0}^{j+1} h_{j+1,\ell} \partial_G^\ell F + \sum_{\ell=0}^j r_{j+1,\ell} \partial_G^\ell Y^\pm, \quad (6.3c)$$

$$\mathcal{E}_{j+1} = \sum_{\ell=0}^j r_{j+1,\ell} \partial_G^\ell X, \quad (6.3d)$$

where the coefficients are given by the following recursion formulas: for $\ell \leq j$ (with the convention that $e_{j,-1} = h_{j,-1} = 0$),

$$\begin{aligned} e_{j+1,j+1} &= 1 \\ e_{j+1,\ell} &= \frac{1}{u'} \partial_r e_{j,\ell} + e_{j,\ell-1} - \frac{2u''}{(u')^2} e_{j,\ell} \\ h_{j+1,j+1}(r) &= 1 \\ h_{j+1,\ell}(r) &= \frac{1}{u'} \partial_r h_{j,\ell} + h_{j,\ell-1} - \frac{2u''}{(u')^2} h_{j,\ell}, \end{aligned}$$

and for $\ell \leq j-1$, (with the convention that $p_{j,-1} = q_{j,-1}, r_{j,-1} = 0$),

$$\begin{aligned} p_{j+1,j} &= p_{j,j-1} + 2 \left(\frac{1}{4} - k^2 \right) \left(\frac{u' + ru''}{r(u')^2} \right) \frac{1}{r^2} \\ p_{j+1,\ell} &= \frac{1}{u'} \partial_r p_{j,\ell} + p_{j,\ell-1} - \frac{2u''}{(u')^2} p_{j,\ell} \\ q_{j+1,j} &= q_{j,j-1} + \partial_{rr} \frac{1}{u'} \\ q_{j+1,\ell} &= \frac{1}{u'} \partial_r q_{j,\ell} + q_{j,\ell-1} - \frac{u''}{(u')^2} q_{j,\ell} \\ r_{j+1,j}(r) &= r_{j,j-1} + \frac{2u''}{(u')^2} \beta - \frac{\beta'}{u'} \\ r_{j+1,\ell}(r) &= \frac{1}{u'} \partial_r r_{j,\ell} + r_{j,\ell-1} - \frac{2u''}{(u')^2} r_{j,\ell}. \end{aligned}$$

Finally, the above coefficients satisfy the following estimates: for $\ell \leq j+1$ and $n \geq 0$,

$$\begin{aligned} |(r \partial_r)^n e_{j+1,\ell}(r)| &\lesssim_n \max(r^{-2}, r^2)^{j+1-\ell} \\ |(r \partial_r)^n h_{j+1,\ell}(r)| &\lesssim_n \max(r^{-2}, r^2)^{j+1-\ell}, \end{aligned}$$

and for $\ell \leq j$ and $n \geq 0$,

$$\begin{aligned} |(r\partial_r)^n q_{j+1,\ell}(r)| &\lesssim_n \max(r^{-3}, r) \max(r^{-2}, r^2)^{j-\ell} \\ |(r\partial_r)^n p_{j+1,\ell}(r)| &\lesssim_n k^2 \max(r^{-4}, 1) \max(r^{-2}, r^2)^{j-\ell} \\ |(r\partial_r)^n r_{j+1,\ell}(r)| &\lesssim_n \frac{1}{r^2 \langle r \rangle^4} \max(r^{-2}, r^2)^{j-\ell}. \end{aligned}$$

Proof of Lemma 6.2 Note

$$\begin{aligned} \partial_G(q(r)\partial_r Z) &= \left(\frac{1}{u'}\partial_r q\right)\partial_r Z + q(r)\partial_G\partial_r Z \\ &= \left(\frac{1}{u'}\partial_r q - q(r)\partial_r\left(\frac{1}{u'}\right)\right)\partial_r Z + q(r)\partial_r\partial_G Z. \end{aligned}$$

The rest follows from Lemma 2.1. \square

6.2 Integral Operators Appearing in the Derivatives of X and Y

Define the following operators which appear in the expression for $\partial_G^j Y^\pm$,

$$Z_{Y\delta}^{\pm\epsilon}[g] := \pm \int_0^\infty \mathcal{G}(r, s, c \pm i\epsilon) \frac{2i\epsilon}{(u-c)^2 + \epsilon^2} g(s, c) ds \quad (6.4a)$$

$$Z_{YS}^{\pm\epsilon}[g] := \int_0^\infty \mathcal{G}(r, s, c \pm i\epsilon) \frac{(u-c)}{(u-c)^2 + \epsilon^2} g(s, c) ds \quad (6.4b)$$

$$Z_{YG}^{\pm\epsilon}[g] := \int_0^\infty \mathcal{G}(r, s, c \pm i\epsilon) g(s, c) ds. \quad (6.4c)$$

In an abuse of notation below, we use the definition

$$Z_{YrG}^{\pm\epsilon} \circ w Z_{Ya}^{\pm\epsilon}[g] := \int_0^\infty \mathcal{G}(r, s, c \pm i\epsilon) w(s) \partial_s Z_{Ya}^{\pm\epsilon}[g](s) ds. \quad (6.5)$$

A more complicated set of operators arises in the formula for X and its derivatives (using a similar abuse of notation):

$$\begin{aligned} Z_{X\delta}^\epsilon[g] &:= \int_0^\infty B_{X\delta;\epsilon}^{(1)}(r, s, c) \frac{2i\epsilon}{(u-c)^2 + \epsilon^2} g(s) ds \\ Z_{XS}^\epsilon[g] &:= \int_0^\infty \left(\int_0^\infty \frac{2i\epsilon\beta(s_0)}{(u-c)^2 + \epsilon^2} B_{XS;\epsilon}^{(1)}(r, s_0, c) B_{XS;\epsilon}^{(2)}(s_0, s, c) ds_0 \right) \\ &\quad \times \frac{(u-c)}{(u-c)^2 + \epsilon^2} g(s) ds \\ Z_{XG}^\epsilon[g] &:= \int_0^\infty \left(\int_0^\infty \frac{2i\epsilon\beta(s_0)}{(u-c)^2 + \epsilon^2} B_{XG;\epsilon}^{(1)}(r, s_0, c) B_{XG;\epsilon}^{(2)}(s_0, s, c) ds_0 \right) g(s) ds \end{aligned}$$

$$Z_{XrG}^\varepsilon \circ w Z_{Ya}^{\pm\varepsilon}[g] := \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} B_{XrG;\varepsilon}^{(1)}(r, s_0, c) B_{XrG;\varepsilon}^{(2)}(s_0, s, c) ds_0 \right) \\ \times w(s) \partial_s Z_{Ya}^{\pm\varepsilon}[g](s) ds. \quad (6.6a)$$

where the kernels are defined via:

$$B_{X\delta;\varepsilon}^{(1)}(r, s, c) = \mathcal{G}(r, s, c + i\varepsilon) \\ + \int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \mathcal{G}(r, s_0, c + i\varepsilon) \mathcal{G}(s_0, s, c - i\varepsilon) ds_0, \quad (6.7a)$$

$$B_{XrG;\varepsilon}^{(1)} = B_{XG;\varepsilon}^{(1)} = B_{XS;\varepsilon}^{(1)}(r, s_0, c) = \mathcal{G}(r, s_0, c + i\varepsilon) \quad (6.7b)$$

$$B_{XrG;\varepsilon}^{(2)} = B_{XG;\varepsilon}^{(2)} = B_{XS;\varepsilon}^{(2)}(s_0, s, c) = \mathcal{G}(s_0, s, c - i\varepsilon). \quad (6.7c)$$

The next lemma verifies the relevance of the above operators.

Lemma 6.3 (Representation formulas). *Let $\varepsilon > 0$ and $c \pm i\varepsilon \in I_\alpha$. There holds,*

$$Y^{\pm\varepsilon}(r, c \pm i\varepsilon) = (Z_{Y\delta}^{\pm\varepsilon} + Z_{YS}^{\pm\varepsilon})[F] + Z_{YG}^{\pm\varepsilon}[F_*], \quad (6.8a)$$

$$X(r, c, \varepsilon) = (Z_{X\delta}^\varepsilon + Z_{XS}^\varepsilon)[F] + Z_{XG}^\varepsilon[F_*], \quad (6.8b)$$

and for $j > 0$ (assuming the integrands are integrable),

$$\partial_G^j Y^\pm = (Z_{Y\delta}^{\pm\varepsilon} + Z_{YS}^{\pm\varepsilon})[F_j] + Z_{YG}^{\pm\varepsilon} \left[\sum_{\ell=0}^j e_{j,\ell} \partial_G^\ell F_* + \sum_{\ell=0}^{j-1} p_{j,\ell} \partial_G^\ell Y^\pm \right] \\ - Z_{YG}^{\pm\varepsilon} \left[\sum_{\ell=0}^{j-1} q_{j,\ell} \partial_r \partial_G^\ell Y^\pm \right] \quad (6.9)$$

$$\partial_G^j X = (Z_{X\delta}^\varepsilon + Z_{XS}^\varepsilon)[F_j^-] + Z_{XG}^\varepsilon \left[\sum_{\ell=0}^j e_{j,\ell} \partial_G^\ell F_* + \sum_{\ell=0}^{j-1} p_{j,\ell} \partial_G^\ell Y^- \right] \\ - Z_{XG}^\varepsilon \left[\sum_{\ell=0}^{j-1} q_{j,\ell} \partial_r \partial_G^\ell Y^- \right] \\ + (Z_{YS}^{+\varepsilon} + Z_{Y\delta}^{+\varepsilon})[\mathcal{E}_j] + Z_{YG}^{+\varepsilon} \left[\sum_{\ell=0}^{j-1} p_{j,\ell} \partial_G^\ell X \right] - Z_{YG}^{+\varepsilon} \left[\sum_{\ell=0}^j q_{j+1,\ell} \partial_r \partial_G^\ell X \right]. \quad (6.10)$$

Proof of Lemma 6.3 First, observe

$$\begin{aligned} \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} (F - \beta Y) &= \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} \left(F - \beta R_{AY}^{-1} \left[\frac{F}{u-c+i\varepsilon} \right] \right) \\ &\quad - \frac{2i\varepsilon\beta}{(u-c)^2 + \varepsilon^2} Z_{YG}^{-\varepsilon}[F_*], \end{aligned}$$

and by definition

$$Z_{YG}^{+\varepsilon} \left[\frac{2i\varepsilon\beta}{(u-c)^2 + \varepsilon^2} Z_{YG}^{-\varepsilon}[F_*] \right] = Z_{XG}^{\varepsilon}[F_*].$$

This completes the proof of (6.8); the cases $j > 0$ follow similarly. \square

6.3 Iterated Integral Operators

6.3.1 Recursion Scheme for Integral Operators

In this section, we will analyze the recursion algorithm derived above. By Lemma 6.2, the F_j and \mathcal{E}_j terms appearing on the RHS of Lemma 6.3 can also be expanded only in terms of X_ℓ and Y_ℓ with $\ell < j$.

We will write all possible operators appearing in the iteration scheme as variations of the original operators with kernels derived from a recursive algorithm. The operators appearing in this algorithm are of the following form:

$$\mathcal{O}_{S;\varepsilon}^{(1)}[wK](r, s_0, c) = \int_0^\infty K(s, s_0, c) \frac{(u-c)}{(u-c)^2 + \varepsilon^2} w(s) \mathcal{G}(r, s, c + i\varepsilon) ds \quad (6.11a)$$

$$\mathcal{O}_{S;\varepsilon}^{(2)}[wK](s_0, r, c) = \int_0^\infty K(s_0, s, c) \frac{(u-c)}{(u-c)^2 + \varepsilon^2} w(s) \mathcal{G}(r, s, c - i\varepsilon) ds \quad (6.11b)$$

$$\mathcal{O}_{\delta;\varepsilon}^{(1)}[wK](r, s_0, c) = \int_0^\infty K(s, s_0, c) \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w(s) \mathcal{G}(r, s, c + i\varepsilon) ds \quad (6.11c)$$

$$\mathcal{O}_{\delta;\varepsilon}^{(2)}[wK](s_0, r, c) = \int_0^\infty K(s_0, s, c) \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w(s) \mathcal{G}(r, s, c - i\varepsilon) ds \quad (6.11d)$$

$$\mathcal{O}_{G;\varepsilon}^{(1)}[wK](r, s_0, c) = \int_0^\infty K(s, s_0, c) w(s) \mathcal{G}(r, s, c + i\varepsilon) ds \quad (6.11e)$$

$$\mathcal{O}_{G;\varepsilon}^{(2)}[wK](s_0, r, c) = \int_0^\infty K(s_0, s, c) w(s) \mathcal{G}(r, s, c - i\varepsilon) ds \quad (6.11f)$$

$$\mathcal{O}_{r;\varepsilon}^{(1)}[wK](r, s_0, c) = \int_0^\infty K(s, s_0, c) w(s) \partial_s \mathcal{G}(r, s, c + i\varepsilon) ds \quad (6.11g)$$

$$\mathcal{O}_{r;\varepsilon}^{(2)}[wK](s_0, r, c) = \int_0^\infty K(s_0, s, c) w(s) \partial_s \mathcal{G}(r, s, c - i\varepsilon) ds. \quad (6.11h)$$

The proofs of the following Lemmas are given in the subsequence sections.

Lemma 6.4 *Let $a \in \{S, G, rG\}$. For each set of weights $\{w_j\}$ and ordering of operators $b_j \in \{S, \delta, G, rG\}$ there exists kernels $B_{\dots, Xa; \varepsilon}^{(1)}, B_{XS, \dots, Yb_J; \varepsilon}^{(2)}$ such that (recall abuse of notations (6.5), (6.6a))*

$$\begin{aligned} & Z_{Xa} \circ w_1 Z_{Yb_1} \circ \dots \circ w_J Z_{YS}[g](r, r_c, \varepsilon) \\ &= \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon u'(s_0)}{(u-c)^2 + \varepsilon^2} B_{Xa; \varepsilon}^{(1)}(r, s_0, c) B_{XS, \dots, YS; \varepsilon}^{(2)}(s_0, s, c) ds_0 \right) \\ & \quad \times \frac{(u-c)}{(u-c)^2 + \varepsilon^2} g(s) ds \\ & Z_{Yb_1} \circ \dots \circ w_{J-1} Z_{Yb_{J-1}} \circ w_J Z_{Xa} \circ w_{J+1} Z_{Yb_{J+1}} \circ \dots \circ w_{J'} Z_{YS}[g](r, r_c, \varepsilon) \\ &= \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon u'(s_0)}{(u-c)^2 + \varepsilon^2} B_{Yb_1, \dots, Yb_J, Xa; \varepsilon}^{(1)}(r, s_0, c) B_{Xa, Yb_{J+1}, \dots, YS; \varepsilon}^{(2)}(s_0, s, c) ds_0 \right) \\ & \quad \times \frac{(u-c)}{(u-c)^2 + \varepsilon^2} g(s) ds. \end{aligned}$$

moreover if the last YS operator is replaced by the allowable alternatives, the expression changes via:

$$\begin{aligned} YS &\mapsto YG \Rightarrow \frac{(u-c)}{(u-c)^2 + \varepsilon^2} \mapsto 1 \\ YS &\mapsto Y\delta \Rightarrow \frac{(u-c)}{(u-c)^2 + \varepsilon^2} \mapsto \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2}, \end{aligned}$$

(notice that the final operator cannot be YrG ; see Lemma 6.1). The operators are obtained by the following recursion formulas:

1. To define $B^{(2)}$ we use the recursion scheme:

$$B_{Xa, Yb_1, \dots, Yb_{J-1}, Yb_J; \varepsilon}^{(2)}(s_0, r, c) = \mathcal{O}_{b_{J-1}; \varepsilon}^{(2)}[w_J B_{Xa, Yb_1, \dots, Yb_{J-1}; \varepsilon}^{(2)}](s_0, r, c),$$

and if $J = 1$ then Xa plays the role of b_{J-1} .

2. To define $B^{(1)}$ we use the recursion scheme

$$B_{Yb_1, Yb_2, \dots, Xa; \varepsilon}^{(1)}(r, s_0, c) = \mathcal{O}_{b_1; \varepsilon}^{(1)}[w_1 B_{Yb_2, \dots, Xa; \varepsilon}^{(1)}](r, s_0, c).$$

Lemma 6.5 *For $a = \delta$, for each set of weights $\{w_j\}$, and choice of operators $b_j \in \{S, \delta, G, rG\}$ there exists kernels (depending on the weights) such that*

$$\begin{aligned} & Z_{Yb_1} \circ \dots \circ w_J Z_{Yb_{J-1}} \circ w_J Z_{X\delta}[g] \\ &= \int_0^\infty \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} B_{Yb_1, \dots, Yb_J, Xa; \varepsilon}^{(1)}(r, s, c) g(s) ds \\ & Z_{Yb_1} \circ \dots \circ w_J Z_{Yb_{J-1}} \circ w_J Z_{Xa} \circ w_{J+1} Z_{Yb_{J+1}} \circ \dots \circ w_{J+1} Z_{YS}[g] \end{aligned}$$

$$= \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} B_{Yb_1, \dots, Yb_J, Xa; \varepsilon}^{(1)}(r, s_0, c) B_{Xa, Yb_{J+1}, \dots, Yb_{J'}; \varepsilon}^{(2)}(s_0, s, c) ds_0 \right) \frac{(u-c)}{(u-c)^2 + \varepsilon^2} g(s) ds,$$

where the last operator is changed for $YS \mapsto YG, Y\delta$ then as above

$$YS \mapsto YG \Rightarrow \frac{(u-c)}{(u-c)^2 + \varepsilon^2} \mapsto 1$$

$$YS \mapsto Y\delta \Rightarrow \frac{(u-c)}{(u-c)^2 + \varepsilon^2} \mapsto \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2}.$$

The kernels are constructed via the following recursion.

1. To construct the $B^{(1)}$ kernels we use the iteration scheme

$$B_{Yb_1, Yb_2, \dots, X\delta; \varepsilon}^{(1)}(r, s_0, c) = \mathcal{O}_{b_1; \varepsilon}^{(1)}[w_1 B_{Yb_2, \dots, X\delta; \varepsilon}^{(1)}](r, s_0, c).$$

2. The first steps of $B^{(2)}$ are given by:

$$B_{X\delta, Ya; \varepsilon}^{(2)} = \mathcal{G}(s_0, r, c - i\varepsilon),$$

and to construct further kernels $B^{(2)}$ we use the recursion scheme

$$B_{X\delta, Yb_1, \dots, Yb_{J-1}, Yb_J; \varepsilon}^{(2)}(s_0, s, c) = \mathcal{O}_{b_{J-1}; \varepsilon}[w_J B_{X\delta, Yb_1, \dots, Yb_{J-1}}^{(2)}](s_0, r, c).$$

Lemma 6.6 For each set of weights $\{w_j\}$ and choice of operators $b_j \in \{S, \delta, G, rG\}$ there exist kernels $B_{XS, Yb_1, \dots, Yb_J; \varepsilon}$ (depending on the weights) such that

$$w_1 Z_{Yb_1} \circ \dots \circ w_J Z_{Yb_J}[g] = \int_0^\infty B_{Yb_1, Yb_2, \dots, Yb_J; \varepsilon}(r, s, c) \frac{(u-c)}{(u-c)^2 + \varepsilon^2} g(s) ds,$$

with the requisite change for $YS \mapsto YG, Y\delta$ as above (as usual, the final operator cannot be YrG). To construct the kernels we use the following scheme as above:

1. All of the basic kernels are given by

$$B_{Ya; \varepsilon} = \mathcal{G}(r, s, c - i\varepsilon).$$

2. Further kernels are constructed via

$$B_{Yb_1, \dots, Yb_{J-1}, Yb_J; \varepsilon}(s_0, r, c) = \mathcal{O}_{b_{J-1}; \varepsilon}^{(2)}[w_J B_{Yb_1, \dots, Yb_{J-1}; \varepsilon}].$$

6.3.2 Estimates on Iterated Integral Kernels

The next step is to use induction to deduce the requisite estimates on the B kernels appearing in Lemmas 6.4–6.6. In order to effectively pass to the limit $\varepsilon \rightarrow 0$ in (2.28), we need to prove that the kernels satisfy a variety of regularity properties. Moreover, in order to close the induction argument, a variety of additional regularity properties are required as well. These properties are outlined in Definitions 6.7 and 6.8 below.

Recall the bounding functions defined in (2.39). Further, define the following variants of the ∂_G derivatives suitable for functions of three variables:

$$\partial_G = \partial_G^{(r)} = \frac{1}{u'(r)} \partial_r + \frac{1}{u'(r_c)} \partial_{r_c} \quad (6.12a)$$

$$\partial_G^{(r,s)} = \frac{1}{u'(r)} \partial_r + \frac{1}{u'(s)} \partial_s + \frac{1}{u'(r_c)} \partial_{r_c}. \quad (6.12b)$$

Definition 6.7 (Suitable (J, ℓ, γ) kernel of type I). We say $K^\varepsilon(r, s, c)$ is a *Suitable* (J, ℓ, γ) kernel of type I if the following properties hold for $c \pm i\varepsilon \in I_\alpha$ with all constants independent of ε :

(a) uniform boundedness and regularity away from the critical layer:

$$|K^\varepsilon(r, s, c)| \lesssim |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s) \quad (6.13a)$$

$$|r \partial_r K^\varepsilon(r, s, c)| \lesssim |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s) \times \left(k + \mathbf{1}_{|r-r_c| < r_c/k} \left| \log \frac{k|r-r_c|}{r_c} \right| \right), \quad (6.13b)$$

$$|r_c \partial_{r_c} K^\varepsilon(r, s, c)| \mathbf{1}_{|s-r_c| > r_c/k} \lesssim k^3 |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s) \times \left(k + \mathbf{1}_{|r-r_c| < r_c/k} \left| \log \frac{k|r-r_c|}{r_c} \right| \right); \quad (6.13c)$$

(b) regularity near the critical layer:

$$\left| \partial_G^{(s)} K^\varepsilon(r, s, c) \right| \mathbf{1}_{|s-r_c| < r_c/k} \mathbf{1}_{|r-r_c| \geq r_c/k} \lesssim \frac{1}{|r_c u'(r_c)|} k |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \quad (6.14a)$$

$$\left| \partial_G^{(r)} K^\varepsilon(r, s, c) \right| \mathbf{1}_{|r-r_c| < r_c/k} \mathbf{1}_{|s-r_c| \geq r_c/k} \lesssim \frac{1}{|r_c u'(r_c)|} k |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \quad (6.14b)$$

$$\left| r \partial_r \partial_G^{(r)} K^\varepsilon(r, s, c) \right| \mathbf{1}_{|r-r_c| < r_c/k} \mathbf{1}_{|s-r_c| \geq r_c/k} \lesssim \frac{1}{|r_c u'(r_c)|} |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \left(k^2 + k \left| \log \frac{k|r-r_c|}{r_c} \right| \right), \quad (6.14c)$$

$$\begin{aligned} & \left| \partial_G^{(r,s)} K^\varepsilon(r, s, c) \right| \mathbf{1}_{|r-r_c| < r_c/k} \mathbf{1}_{|s-r_c| < r_c/k} \\ & \lesssim \frac{1}{|r_c u'(r_c)|} k |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \end{aligned} \quad (6.14d)$$

$$\begin{aligned} & \left| r \partial_r \partial_G^{(r,s)} K^\varepsilon(r, s, c) \right| \mathbf{1}_{|r-r_c| < r_c/k} \mathbf{1}_{|s-r_c| < r_c/k} \\ & \lesssim \frac{1}{|r_c u'(r_c)|} |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \left(k^2 + k \left| \log \frac{k|r-r_c|}{r_c} \right| \right); \end{aligned} \quad (6.14e)$$

(c) Hölder regularity in s near the critical layer: for $|s - r_c| < r_c/k$ there holds:

$$\begin{aligned} & |K^\varepsilon(r, s, c) - K^\varepsilon(r, r_c, c)| \\ & \lesssim |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \left(\frac{k|s-r_c|}{r_c} \right)^\gamma \end{aligned} \quad (6.15a)$$

$$\begin{aligned} & \left| \partial_G^{(s)} (K^\varepsilon(r, s, c) - K^\varepsilon(r, r_c, c)) \right| \mathbf{1}_{|r-r_c| \geq r_c/k} \\ & \lesssim \frac{1}{|r_c u'(r_c)|} k |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \left(\frac{k|s-r_c|}{r_c} \right)^\gamma \end{aligned} \quad (6.15b)$$

$$\left| \partial_G^{(r,s)} (K^\varepsilon(r, s, c) - K^\varepsilon(r, r_c, c)) \right| \mathbf{1}_{|r-r_c| < r_c/k} \quad (6.15c)$$

$$\lesssim \frac{1}{|r_c u'(r_c)|} k |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \left(\frac{k|s-r_c|}{r_c} \right)^\gamma \quad (6.15d)$$

$$\begin{aligned} & \left| r \partial_r (K^\varepsilon(r, s, c) - K^\varepsilon(r, r_c, c)) \right| \mathbf{1}_{|r-r_c| \geq r_c/k} \\ & \lesssim k^2 |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \left(\frac{k|s-r_c|}{r_c} \right)^\gamma \end{aligned} \quad (6.15e)$$

$$\begin{aligned} & \left| r_c \partial_{r_c} (K^\varepsilon(r, s, c) - K^\varepsilon(r, r_c, c)) \right| \mathbf{1}_{|r-r_c| \geq r_c/k} \\ & \lesssim k^3 |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \left(\frac{k|s-r_c|}{r_c} \right)^\gamma. \end{aligned} \quad (6.15f)$$

(c) convergence: there exists an $\eta > 0$ such that $\varepsilon^{-\eta} (K^\varepsilon - K^0)$ satisfies (6.13), (6.14), and (6.15) for $z \in I_\alpha$ and ε sufficiently small.

Definition 6.8 (*Suitable (J, ℓ) kernel of type II*). We say $K^\varepsilon(r, s, c)$ is a *Suitable (J, ℓ, γ) kernel of type II* if the following properties hold for $c \pm i\varepsilon \in I_\alpha$ with all constants independent of ε :

(a) uniform boundedness and regularity away from the critical layer:

$$|K^\varepsilon(r, s, c)| \lesssim |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \quad (6.16a)$$

$$|s \partial_s K^\varepsilon(r, s, c)| \quad (6.16b)$$

$$\lesssim k |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \left(k + \mathbf{1}_{|s-r_c| < r_c/k} \left| \log \frac{k|s-r_c|}{r_c} \right| \right), \quad (6.16c)$$

$$\begin{aligned}
& \left| r_c \partial_{r_c} K^\varepsilon(r, s, c) \right| \mathbf{1}_{|r-r_c| > r_c/k} \\
& \lesssim k^3 |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \left(k + \mathbf{1}_{|s-r_c| < r_c/k} \left| \log \frac{k|s-r_c|}{r_c} \right| \right); \tag{6.16d}
\end{aligned}$$

(b) regularity near the critical layer:

$$\begin{aligned}
& \left| \partial_G^{(s)} K^\varepsilon(r, s, c) \right| \mathbf{1}_{|s-r_c| < r_c/k} \mathbf{1}_{|r-r_c| \geq r_c/k} \\
& \lesssim \frac{1}{|r_c u'(r_c)|} k |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \tag{6.17a}
\end{aligned}$$

$$\left| \partial_G^{(r)} K^\varepsilon(r, s, c) \right| \mathbf{1}_{|r-r_c| < r_c/k} \mathbf{1}_{|s-r_c| \geq r_c/k} \tag{6.17b}$$

$$\lesssim \frac{1}{|r_c u'(r_c)|} k |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \tag{6.17c}$$

$$\left| s \partial_s \partial_G^{(s)} K^\varepsilon(r, s, c) \right| \mathbf{1}_{|r-r_c| < r_c/k} \mathbf{1}_{|s-r_c| \geq r_c/k} \tag{6.17d}$$

$$\lesssim \frac{1}{|r_c u'(r_c)|} |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \left(k^2 + k \left| \log \frac{k|s-r_c|}{r_c} \right| \right), \tag{6.17e}$$

$$\left| \partial_G^{(r,s)} K^\varepsilon(r, s, c) \right| \mathbf{1}_{|r-r_c| < r_c/k} \mathbf{1}_{|s-r_c| < r_c/k} \tag{6.17f}$$

$$\lesssim \frac{1}{|r_c u'(r_c)|} k |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \tag{6.17g}$$

$$\left| s \partial_s \partial_G^{(r,s)} K^\varepsilon(r, s, c) \right| \mathbf{1}_{|r-r_c| < r_c/k} \mathbf{1}_{|s-r_c| < r_c/k} \tag{6.17h}$$

$$\lesssim \frac{1}{|r_c u'(r_c)|} |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \left(k^2 + k \left| \log \frac{k|s-r_c|}{r_c} \right| \right); \tag{6.17i}$$

(c) Hölder regularity in r near critical layer: for $|r - r_c| < r_c/k$ there holds:

$$\begin{aligned}
& \left| K^\varepsilon(r, s, c) - K^\varepsilon(r_c, s, c) \right| \\
& \lesssim |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \left(\frac{k|r-r_c|}{r_c} \right)^\gamma \tag{6.18a}
\end{aligned}$$

$$\begin{aligned}
& \left| \partial_G^{(r)} (K^\varepsilon(r, s, c) - K^\varepsilon(r_c, s, c)) \right| \mathbf{1}_{|s-r_c| \geq r_c/k} \\
& \lesssim \frac{1}{|r_c u'(r_c)|} k |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \left(\frac{k|r-r_c|}{r_c} \right)^\gamma \tag{6.18b}
\end{aligned}$$

$$\begin{aligned}
& \left| \partial_G^{(r,s)} (K^\varepsilon(r, s, c) - K^\varepsilon(r_c, s, c)) \right| \mathbf{1}_{|s-r_c| < r_c/k} \\
& \lesssim \frac{1}{|r_c u'(r_c)|} k |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \left(\frac{k|r-r_c|}{r_c} \right)^\gamma \tag{6.18c}
\end{aligned}$$

$$\begin{aligned}
& \left| s \partial_s (K^\varepsilon(r, s, c) - K^\varepsilon(r_c, s, c)) \right| \mathbf{1}_{|s-r_c| \geq r_c/k} \\
& \lesssim k^2 |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J,\ell}(r, s, c) \left(\frac{k|r-r_c|}{r_c} \right)^\gamma \tag{6.18d}
\end{aligned}$$

$$\begin{aligned}
& |r_c \partial_{r_c} (K^\varepsilon(r, s, c) - K^\varepsilon(r_c, s, c))| \mathbf{1}_{|s-r_c| \geq r_c/k} \\
& \lesssim k^3 |u'(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathcal{L}_{J, \ell}(r, s, c) \left(\frac{k|r-r_c|}{r_c} \right)^\gamma. \quad (6.18e)
\end{aligned}$$

(c) convergence: there exists an $\eta > 0$ such that $\varepsilon^{-\eta} (K^\varepsilon - K^0)$ satisfies (6.16), (6.17), and (6.18) for $z \in I_\alpha$ and ε sufficiently small (depending only on k).

Remark 6.9 Note that the main differences between Definitions 6.7 and 6.8 is in the regularity requirements.

First, we prove that the Green's function is the prototypical suitable kernel of both types.

Lemma 6.10 *The Green's function $\mathcal{G}(r, s, c \pm i\varepsilon)$ is both a suitable $(0, 0, \gamma)$ kernel of type I and a suitable $(0, 0, \gamma)$ kernel of type II for all $\gamma \in (0, 1)$.*

Proof of Lemma 6.10 Consider just the case statement of Type I; Type II is exactly analogous.

Step 1: Proof of (6.13) First, consider the boundedness estimate (6.13a). From Lemmas 5.2 and 5.4 (and 2.1),

$$\begin{aligned}
\left| \frac{\mathcal{G}(r, s, c - i\varepsilon)}{u'(s)} \right| & \lesssim \max(s^{-1}, s^3) \min\left(\frac{r_c^3}{k}, \frac{1}{kr_c^5}\right) \\
& \times \left(\mathbf{1}_{r_c \leq 1} \mathbf{1}_{s < r} \left(\mathbf{1}_{s < r_c} \frac{s^{k+1/2}}{r_c^{k+3/2}} + \mathbf{1}_{r_c < s < 1} k \frac{s^{k+2+1/2}}{r_c^{k+2+3/2}} + \mathbf{1}_{s > 1} k \frac{s^{k+1/2}}{r_c^{k+3+1/2}} \right) \right. \\
& \times \left(\mathbf{1}_{r < r_c} k \frac{r_c^{k-3/2}}{r^{k-1/2}} + \mathbf{1}_{r_c < r < 1} \frac{r_c^{k+1/2}}{r^{k+3/2}} + \mathbf{1}_{r > 1} \frac{r_c^{k+1/2}}{r^{k-1/2}} \right) \\
& + \mathbf{1}_{r_c \leq 1} \mathbf{1}_{r < s} \left(\mathbf{1}_{r < r_c} \frac{r^{k+1/2}}{r_c^{k+3/2}} + \mathbf{1}_{r_c < r < 1} k \frac{r^{k+2+1/2}}{r_c^{k+2+3/2}} + \mathbf{1}_{r > 1} k \frac{r^{k+1/2}}{r_c^{k+3+1/2}} \right) \\
& \times \left(\mathbf{1}_{s < r_c} k \frac{r_c^{k-3/2}}{s^{k-1/2}} + \mathbf{1}_{r_c < s < 1} \frac{r_c^{k+1/2}}{s^{k+3/2}} + \mathbf{1}_{s > 1} \frac{r_c^{k+1/2}}{s^{k-1/2}} \right) \\
& + \mathbf{1}_{r_c > 1} \mathbf{1}_{s < r} \left(\mathbf{1}_{s < 1} \frac{s^{k+1/2}}{r_c^{k-1/2}} + \mathbf{1}_{1 < s < r_c} \frac{s^{k+5/2}}{r_c^{k-1/2}} + \mathbf{1}_{s > r_c} k \frac{s^{k+1/2}}{r_c^{k-5/2}} \right) \\
& \times \left(\mathbf{1}_{r < 1} k \frac{r_c^{k+5-1/2}}{r^{k-1/2}} + \mathbf{1}_{1 \leq r < r_c} k \frac{r_c^{k+5-1/2}}{r^{k+3/2}} + \mathbf{1}_{r_c < r} \frac{r_c^{k+5/2}}{r^{k-1/2}} \right) \\
& + \mathbf{1}_{r_c > 1} \mathbf{1}_{s > r} \left(\mathbf{1}_{r < 1} \frac{r^{k+1/2}}{r_c^{k-1/2}} + \mathbf{1}_{1 < r < r_c} \frac{r^{k+5/2}}{r_c^{k-1/2}} + \mathbf{1}_{r > r_c} k \frac{r^{k+1/2}}{r_c^{k-5/2}} \right) \\
& \times \left(\mathbf{1}_{s < 1} k \frac{r_c^{k+5-1/2}}{s^{k-1/2}} + \mathbf{1}_{1 \leq s < r_c} k \frac{r_c^{k+5-1/2}}{s^{k+3/2}} + \mathbf{1}_{r_c < s} \frac{r_c^{k+5/2}}{s^{k-1/2}} \right) \Big) \\
& \lesssim \mathbf{1}_{r_c \leq 1} \left(\mathbf{1}_{s < r < r_c} \frac{s^{k-1/2}}{r^{k-1/2}} + \mathbf{1}_{s < r_c < r < 1} \frac{s^{k-1/2}}{r^{k-1/2}} \frac{r_c^2}{r^2} + \mathbf{1}_{s < r_c < 1 < r} \frac{s^{k-1/2}}{r^{k-1/2}} r_c^2 \right. \\
& \left. + \mathbf{1}_{r_c < s < r < 1} \frac{s^{k+3/2}}{r^{k+3/2}} + \mathbf{1}_{r_c < s < 1 < r} \frac{s^{k+3/2}}{r^{k-1/2}} + \mathbf{1}_{1 < s < r} \frac{s^{k+3+1/2}}{r^{k-1/2}} \right)
\end{aligned}$$

$$\begin{aligned}
& + \mathbf{1}_{r_c \leq 1} \left(\mathbf{1}_{r < s < r_c} \frac{r^{k+1/2}}{s^{k+1/2}} + \mathbf{1}_{r < r_c < s < 1} \frac{r^{k+1/2}}{s^{k+1/2}} \frac{r_c^2}{s^2} + \mathbf{1}_{r < r_c < 1 < s} \frac{r^{k+1/2}}{s^{k+1/2}} s^4 r_c^2 \right. \\
& + \mathbf{1}_{r_c < r < s < 1} \frac{r^{k+2+1/2}}{s^{k+2+1/2}} + \mathbf{1}_{r_c < r < 1 < s} \frac{r^{k+2+1/2}}{s^{k-1/2}} s^3 + \mathbf{1}_{1 < r < s} \frac{r^{k+1/2}}{s^{k-1/2}} s^3 \Big) \\
& + \mathbf{1}_{r_c > 1} \left(\mathbf{1}_{s < r < 1} \frac{s^{k-1/2}}{r^{k-1/2}} + \mathbf{1}_{s < 1 < r < r_c} \frac{s^{k-1/2}}{r^{k+3/2}} + \mathbf{1}_{s < 1 < r_c < r} \frac{s^{k-1/2}}{r^{k-1/2}} \frac{1}{r_c^2} \right. \\
& + \mathbf{1}_{1 < s < r < r_c} \frac{s^{k-1/2}}{r^{k+3/2}} + \mathbf{1}_{1 < s < r_c < r} \frac{s^{k+3+5/2}}{r^{k-1/2}} \frac{1}{r_c^2} + \mathbf{1}_{1 < r_c < s < r} \frac{s^{k+1/2+3}}{r^{k-1/2}} \Big) \\
& + \mathbf{1}_{r_c > 1} \left(\mathbf{1}_{r < s < 1} \frac{r^{k+1/2}}{s^{k+1/2}} + \mathbf{1}_{r < 1 < s < r_c} \frac{r^{k+1/2}}{s^{k+1/2}} s^2 + \mathbf{1}_{r < 1 < r_c < s} \frac{r^{k+1/2}}{s^{k+1/2}} s^4 r_c^{-2} \right. \\
& + \mathbf{1}_{1 < r < s < r_c} \frac{r^{k+1/2}}{s^{k+1/2}} r^2 s^2 + \mathbf{1}_{1 < r < r_c < s} \frac{r^{k+1/2}}{s^{k+1/2}} s^4 r_c^{-2} \\
& \left. + \mathbf{1}_{1 < r_c < r < s} \frac{r^{k+1/2}}{s^{k+1/2}} s^4 \right) \lesssim \mathbb{K}(r, s, c) \mathcal{B}(r, s),
\end{aligned}$$

which is (6.13a). The estimate (6.13b) follows from a similar argument together with Lemma 5.5. To see the $r_c \partial_{r_c}$ control, first note that

$$\begin{aligned}
r_c \partial_{r_c} \left(\frac{\mathcal{G}(r, s, c)}{u'(s)} \right) &= \frac{1}{u'(s)M(z)} \begin{cases} \frac{-r_c \partial_{r_c} M(z)}{M(z)} H_0(r, c) H_\infty(s, c) & r < s \\ \frac{-r_c \partial_{r_c} M(z)}{M(z)} H_0(r, c) H_\infty(s, c) & s > r \end{cases} \\
&+ \frac{1}{u'(s)M(z)} \begin{cases} r_c \partial_{r_c} H_0(r, c) H_\infty(s, c) + H_0(r, c) r_c \partial_{r_c} H_\infty(s, c) & r < s \\ r_c \partial_{r_c} H_0(s, c) H_\infty(r, c) + H_0(s, c) r_c \partial_{r_c} H_\infty(r, c) & r > s. \end{cases}
\end{aligned}$$

Hence, all the lemmas in Sect. 5 together imply (6.13c) as in (6.13a).

Step 2: Proof of (6.14) Consider the case $|r - r_c| \geq r_c/k$ and $|s - r_c| < r_c/k$ as in (6.14a). Here we have,

$$\begin{aligned}
\partial_G^{(s)} \left(\frac{\mathcal{G}(r, s, c)}{u'(s)} \right) &= \frac{1}{u'(s)M(z)} \begin{cases} r_c \partial_{r_c} H_0(r, c) H_\infty(s, c) + H_0(r, c) \partial_G H_\infty(s, c) & r < s \\ \partial_G H_0(s, c) H_\infty(r, c) + H_0(s, c) r_c \partial_{r_c} H_\infty(r, c) & r > s \end{cases} \\
&- \left(\frac{u''(s)}{(u'(s))^3 M(z)} + \frac{r_c \partial_{r_c} M(z)}{M^2(z)} \right) \begin{cases} H_0(r, c) H_\infty(s, c) & r < s \\ H_0(r, c) H_\infty(s, c) & r > s. \end{cases}
\end{aligned}$$

By the lemmas in Sect. 5, we deduce (6.14a). Note further that from the lemmas in Sect. 5, we may deduce a logarithmically singular upper bound on $s \partial_s \partial_G^{(s)} \left((u'(s))^{-1} \mathcal{G}(r, s, c) \right)$. This is important both to verify that the kernel is suitable of type II but also to prove the Hölder regularity (6.15) below. The estimates (6.14b) and (6.14c) are analogous and omitted for the sake of brevity. Consider the estimates (6.14d) and (6.14e) next. We have

$$\partial_G^{(r,s)} \left(\frac{\mathcal{G}(r,s,c)}{u'(s)} \right) = \frac{1}{u'(s)M(z)} \begin{cases} \partial_G H_0(r,c)H_\infty(s,c) + H_0(r,c)\partial_G H_\infty(s,c) & r < s \\ \partial_G H_0(s,c)H_\infty(r,c) + H_0(s,c)\partial_G H_\infty(r,c) & r > s \end{cases} \\ - \left(\frac{u''(s)}{(u'(s))^3 M(z)} + \frac{r_c \partial_{r_c} M(z)}{M(z)} \right) \begin{cases} H_0(r,c)H_\infty(s,c) & r < s \\ H_0(r,c)H_\infty(s,c) & r > s. \end{cases} \quad (6.19)$$

Hence, this satisfies the desired estimates by the lemmas in Sect. 5. Similarly, we can obtain $s\partial_s$ estimates as well.

Step 3: Proof of (6.15) The inequalities (6.15a) are a consequence of the log-Lipschitz regularity of \mathcal{G} in both variables (from Lemma 5.5); we omit the details as they are straightforward. Finally, we note that the convergence stated in Definition 6.7 follows from the lemmas in Sect. 5. \square

Next, we prove that all possible integral operators arising in the iteration scheme are suitable.

Lemma 6.11 (Iterated integral operators $\mathcal{O}_{\delta;\varepsilon}^{(j)}$ and $\mathcal{O}_{S;\varepsilon}^{(j)}$). *Let $K_\varepsilon^{(1)}$ be a Suitable(J, ℓ', γ) kernel of type I and $K_\varepsilon^{(2)}$ a Suitable(J, ℓ', γ) kernel of type II. Further, suppose that*

$$|w(r)| + |r\partial_r w(r)| \lesssim \max \left(\frac{1}{r^2}, r^2 \right)^{\ell+1} \frac{1}{\langle r \rangle^6}.$$

Further, suppose $\ell + \ell' < k - 1/2$. Then, for all $\eta > 0$ and $\gamma' \in (0, \gamma)$,

- $\mathcal{O}_{S;\varepsilon}^{(1)}[wK_\varepsilon^{(1)}]$ is suitable ($J+1, \ell' + \ell + 1 + \eta, \gamma'$) of type I, $\mathcal{O}_{\delta;\varepsilon}^{(1)}[wK_\varepsilon^{(1)}]$ is suitable ($J, \ell' + \ell + 1 + \eta, \gamma'$) of type I;
- $\mathcal{O}_{S;\varepsilon}^{(2)}[wK_\varepsilon^{(2)}]$ is suitable ($J+1, \ell' + \ell + 1 + \eta, \gamma'$) of type II and $\mathcal{O}_{\delta;\varepsilon}^{(2)}[wK_\varepsilon^{(2)}]$ is suitable ($J, \ell' + \ell + 1 + \eta, \gamma'$) kernel of type II.

Lemma 6.12 (Iterated integral operators $\mathcal{O}_G^{(j)}$). *Suppose that $K_\varepsilon^{(1)}$ is a suitable (J, ℓ', γ) kernel of type I and $K_\varepsilon^{(2)}$ is suitable (J, ℓ', γ) kernel of type II. Further, suppose that*

$$|w(r)| + |r\partial_r w(r)| \lesssim k^2 \max \left(\frac{1}{r^2}, r^2 \right)^\ell \max \left(\frac{1}{r^4}, 1 \right),$$

and that $\ell + \ell' < k - 3/2$. Then, for all $\eta > 0$, and $\gamma' \in (0, \gamma)$, $\mathcal{O}_{G;\varepsilon}^{(1)}[wK_\varepsilon^{(1)}]$ is suitable ($J+2, \ell' + \ell + 1 + \eta, \gamma'$) of type I and $\mathcal{O}_{G;\varepsilon}^{(2)}[wK_\varepsilon^{(2)}]$ is suitable ($J+2, \ell' + \ell + 1 + \eta, \gamma'$) of type II.

Lemma 6.13 (Iterated integral operator $\mathcal{O}_{r;\varepsilon}^{(j)}$). *Suppose that*

$$|w(r)| + |r\partial_r w(r)| \lesssim \max \left(\frac{1}{r^2}, r^2 \right)^\ell \max(r^{-4}, 1)$$

and that $\ell + \ell' < k - 3/2$. Suppose that the kernel $K_\varepsilon^{(1)}$ is a suitable (J, ℓ', γ) kernel of type I and that $K_\varepsilon^{(2)}$ is a suitable (J, ℓ', γ) kernel of type II. Then, for all $\eta > 0$ and $\gamma' \in (0, \gamma)$, $\mathcal{O}_{r;\varepsilon}^{(1)}[wK_\varepsilon^{(1)}]$ is a suitable $(J+1, \ell' + \ell + 1 + \eta, \gamma')$ kernel of type I, and $\mathcal{O}_{r;\varepsilon}^{(2)}[wK_\varepsilon^{(2)}]$ is a suitable $(J+1, \ell' + \ell + 1 + \eta, \gamma')$ kernel of type II.

We now prove Lemmas 6.11–6.13.

Proof of Lemma 6.11 The cases of $j = 1$ and $j = 2$ are essentially the same, we will focus on $j = 1$ here. The treatment of \mathcal{O}_S is similar to, but slightly harder than, the case \mathcal{O}_δ , so we focus on the former. Define:

$$\tilde{K}(r, s_0, c) := \int_0^\infty \frac{(u-c)}{(u-c)^2 + \varepsilon^2} w(s) \mathcal{G}(r, s, c + i\varepsilon) K(s, s_0, c) ds.$$

Boundedness Estimate (6.13a) Most of the non-trivial methods involved in the proof of Lemma 6.11 appear in some form in the proof of (6.13a). Recall (4.3) and split the integral based on proximity to the critical layer:

$$\begin{aligned} \tilde{K}(r, s_0, c) &= \int_0^\infty \frac{(u-c)u'(s)}{(u-c)^2 + \varepsilon^2} (\chi_c + \chi_{\neq}) w(s) \frac{\mathcal{G}(r, s, c + i\varepsilon)}{u'(s)} K(s, s_0, c) ds \\ &=: \tilde{K}_c + \tilde{K}_{\neq}. \end{aligned} \quad (6.20)$$

First consider the problem of estimating \tilde{K}_{\neq} for $r_c \leq 1$. In the case of \tilde{K}_{\neq} , we apply Lemma 6.10 and (A.1),

$$\begin{aligned} \mathbf{1}_{r_c \leq 1} \left| \frac{1}{u'(s_0)} \tilde{K}_{\neq}(r, s_0, c) \right| &\lesssim \mathbf{1}_{r_c \leq 1} \int_0^\infty \frac{|u'(s)|}{|u-c|} \chi_{\neq} |w(s)| \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathbb{K}(s, s_0, c) \mathcal{B}(s, s_0) \mathcal{L}_{J, \ell'}(s, s_0) ds \\ &\lesssim \mathbf{1}_{r_c \leq 1} \int_0^\infty \left(\frac{k}{s^{1+2\ell} \max(s^2, r_c^2)} \mathbf{1}_{s \leq 1} + s^{2\ell-7} \mathbf{1}_{s \geq 1} \right) \\ &\quad \times \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathbb{K}(s, s_0, c) \mathcal{B}(s, s_0) \mathcal{L}_{J, \ell'}(s, s_0) ds. \end{aligned}$$

This integral is estimated by a tedious, but straightforward, calculation. Note that the requirement $\ell + \ell' < k - 1/2$ is necessary to ensure the resulting integrands are integrable at zero and infinity. The calculation is summarized via:

$$\begin{aligned} \mathbf{1}_{r_c \leq 1} \left| \frac{1}{u'(s_0)} \tilde{K}_{0, \neq}(r, s_0, c) \right| &\lesssim \mathbf{1}_{r \leq s_0 \leq r_c \leq 1} \frac{r^{k+1/2}}{s_0^{k+1/2}} \left(\frac{k}{r^{2+2(\ell+\ell')}} \right) + \mathbf{1}_{s_0 \leq r \leq r_c \leq 1} \frac{s_0^{k-1/2}}{r^{k-1/2}} \left(\frac{k}{s_0^{2+2(\ell+\ell')}} \right) \\ &\quad + \mathbf{1}_{s_0 \leq r_c \leq r \leq 1} \frac{r_c^2 s_0^{k-1/2}}{r^2 r^{k-1/2}} \left(\frac{k}{s_0^{2(\ell+\ell')+2}} \right) \end{aligned}$$

$$\begin{aligned}
& + \mathbf{1}_{s_0 \leq r_c \leq 1 \leq r} \frac{r_c^2 s_0^{k-1/2}}{r^{k-1/2}} \left(\frac{k}{s_0^{2(\ell+\ell')+2}} + \max(s_0^{-2\ell'}, r^{2\ell-2} s_0^{-2\ell'}, r^{2(\ell+\ell')-2}) \langle \log r \rangle \right) \\
& + \mathbf{1}_{0 \leq r \leq r_c \leq s_0 \leq 1} \frac{r_c^2 r^{k+1/2}}{s_0^2 s_0^{k+1/2}} \left(\frac{k}{r^{2(\ell+\ell')+2}} \right) \\
& + \mathbf{1}_{r_c \leq r \leq s_0 \leq 1} \frac{r^2 r^{k+1/2}}{s_0^2 s_0^{k+1/2}} \left(\frac{r_c^{2+2k-2\ell-2\ell'}}{r^{2k+4}} + \frac{k}{r^{2+2\ell+2\ell'}} \right) \\
& + \mathbf{1}_{r_c \leq s_0 \leq r \leq 1} \frac{s_0^{3/2+k}}{r^{3/2+k}} \left(\frac{r_c^{2+2k-2(\ell+\ell')}}{s_0^{4+2k}} + \frac{k}{s_0^{2(\ell+\ell')+2}} \right) \\
& + \mathbf{1}_{r_c \leq s_0 \leq 1 \leq r} \frac{s_0^{k+3/2}}{r^{k-1/2}} \left(\frac{r_c^{2+2k-2(\ell+\ell')}}{s_0^{2k+4}} + \frac{k}{s_0^{2+2(\ell+\ell')}} \right. \\
& \quad \left. + \max(s_0^{-2\ell'}, r^{2(\ell+\ell')-2}, r^{2\ell-2} s_0^{-2\ell'}) \langle \log r \rangle \right) \\
& + \mathbf{1}_{r \leq r_c \leq 1 \leq s_0} \frac{r^{k+1/2} r_c^2}{s_0^{k+1/2}} \langle s_0 \rangle^4 \left(\frac{k}{r^{2+2\ell}} \max\left(\frac{1}{r^{2\ell'}}, s_0^{2\ell'}\right) \right. \\
& \quad \left. + \max(1, s_0^{2(\ell+\ell')-2}) \langle \log s_0 \rangle \right) \\
& + \mathbf{1}_{r_c \leq r \leq 1 \leq s_0} \frac{r^2 r^{k+1/2}}{s_0^{k+1/2}} \langle s_0 \rangle^4 \left(\frac{r_c^{2+2k-2\ell}}{r^{4+2k}} \max(r_c^{-2\ell'}, s_0^{2\ell'}) \right. \\
& \quad \left. + \frac{k}{r^{2+2\ell}} \max(r^{-2\ell'}, s_0^{2\ell'}) + \max(1, s_0^{2(\ell+\ell')-2}) \langle \log s_0 \rangle \right) \\
& + \mathbf{1}_{r_c \leq 1 \leq r \leq s_0} \frac{r^{k+1/2}}{s_0^{k+1/2}} \langle s_0 \rangle^4 \left(\frac{r_c^{2+2k-2\ell}}{r^{2k}} \max(r_c^{-2\ell'}, s_0^{2\ell'}) \right. \\
& \quad \left. + \max(1, s_0^{2(\ell+\ell')-2}) \langle \log s_0 \rangle \right) \\
& + \mathbf{1}_{r_c \leq 1 \leq s_0 \leq r} \frac{s_0^{k-1/2}}{r^{k-1/2}} \langle s_0 \rangle^4 \left(\frac{r_c^{2+2k-2\ell}}{s_0^{2k}} \max(r_c^{-2\ell'}, s_0^{2\ell'}) \right. \\
& \quad \left. + \langle \log r \rangle \max(1, r^{2(\ell+\ell')-2}) \right).
\end{aligned}$$

A key constraint is to not lose powers of r_c^{-1} while still gaining the improvement encoded in \mathbb{K} , that is, we specifically want \mathcal{L} to be independent of r_c , so when e.g. $r_c \ll r, s_0$, we still need to get good estimates (which one observes is indeed the case using that $\ell + \ell' \leq k$). After simplification, we therefore have the following estimate for all $\eta > 0$,

$$\mathbf{1}_{r_c \leq 1} \left| \frac{1}{u'(s_0)} \tilde{K}_{0, \neq}(r, s_0, c) \right| \lesssim \mathbb{K}(r, s_0, c) \mathcal{B}(r, s_0) \mathcal{L}_{J+1, \ell+\ell'+1+\eta}(r, s_0), \quad (6.21)$$

which is the desired estimate. In the case $r_c \geq 1$, we similarly have (omitting the tedious intermediate steps):

$$\begin{aligned} \mathbf{1}_{r_c > 1} \left| \frac{1}{u'(s_0)} \tilde{K}_{0,\neq}(r, s_0, c) \right| &\lesssim \mathbf{1}_{r_c > 1} \int_0^\infty \left(\frac{1}{s^{1+2\ell}} \mathbf{1}_{s \leq 1} + \max(r_c^2, s^2) s^{2\ell-7} \mathbf{1}_{s \geq 1} \right) \\ &\quad \times \mathbb{K}(r, s, c) \mathcal{B}(r, s) \mathbb{K}(s, s_0, c) \mathcal{B}(s, s_0) \mathcal{L}_{J,\ell'}(s, s_0) ds \\ &\lesssim \mathcal{B}(r, s_0) \mathcal{L}_{J+1,\ell+\ell'+1+\eta}(r, s_0). \end{aligned} \quad (6.22)$$

This completes the treatment of the \tilde{K}_{\neq} .

Consider next the contributions of \tilde{K}_c . First consider the case that $|r - r_c| < r_c/k$. Near the critical layer we write

$$\begin{aligned} \frac{1}{u'(s_0)} \tilde{K}_c(r, s_0, c) &= w(r_c) \frac{\mathcal{G}(r, r_c, c)}{u'(r_c)} \frac{K(r_c, s_0, c)}{u'(s_0)} \int_0^\infty \chi_c \frac{(u-c)u'(s)}{(u-c)^2 + \varepsilon^2} ds \\ &\quad + \int_0^\infty \chi_c \frac{(u-c)u'(s)}{(u-c)^2 + \varepsilon^2} \left(w(s) \frac{\mathcal{G}(r, s, c)}{u'(s)} \frac{K(s, s_0, c)}{u'(s_0)} \right. \\ &\quad \left. - w(r_c) \frac{\mathcal{G}(r, r_c, c)}{u'(r_c)} \frac{K(r_c, s_0, c)}{u'(s_0)} \right) ds \\ &= \tilde{K}_{c1} + \tilde{K}_{c2}. \end{aligned}$$

For \tilde{K}_{c1} , by Definition 6.7, Lemmas 6.10, A.1 and A.3, followed by arguments similar to those to deduce (6.21) and (6.22) without losing powers of $r_c^{\pm 1}$ we have the following estimates:

$$\begin{aligned} \left| \tilde{K}_{c1}(r, s_0, c) \right| &\lesssim \max(r_c^2, r_c^{-2})^{\ell+1} \langle r_c \rangle^{-6} \mathbb{K}(r, r_c, c) \mathcal{B}(r, r_c) \\ &\quad \times \mathbb{K}(r_c, s_0, c) \mathcal{B}(r_c, s_0) \mathcal{L}_{J,\ell'}(r_c, s_0) \\ &\lesssim \mathbb{K}(r, s_0, c) \mathcal{B}(r, s_0) \mathcal{L}_{J,\ell'+\ell+1}(r, s_0). \end{aligned}$$

Next, turn to \tilde{K}_{c2} . By the logarithmic regularity in Definition 6.7, we have (using $|r_c - s| < r_c/k$ and Lemma A.1),

$$\begin{aligned} |K(s, s_0, c) - K(r_c, s_0, c)| &\lesssim \mathbb{K}(r_c, s_0, c) \mathcal{B}(r_c, s_0) \mathcal{L}_{J,\ell'}(r_c, s_0) \frac{|s - r_c|}{r_c} \\ &\quad \times \int_0^1 r_c |\partial_s K(r_c + \theta(s - r_c), s_0, c)| d\theta \\ &\lesssim \mathbb{K}(r_c, s_0, c) \mathcal{B}(r_c, s_0) \frac{|s - r_c|}{r_c} \\ &\quad \times \int_0^1 \left| k + \left| \log \frac{k\theta |s - r_c|}{r_c} \right| \right| d\theta \\ &\lesssim \mathbb{K}(r_c, s_0, c) \mathcal{B}(r_c, s_0) \mathcal{L}_{J,\ell'}(r_c, s_0) \frac{|s - r_c|}{r_c} \\ &\quad \times \left(k + \left| \log \frac{k |s - r_c|}{r_c} \right| \right). \end{aligned}$$

Analogous estimates also hold for \mathcal{G} due to Lemma 6.10 (and clearly also w). Therefore, we have (again arguing as in (6.21) and (6.22) to avoid losing powers of r_c),

$$\begin{aligned} |\tilde{K}_{c2}| &\lesssim \max(r_c^2, r_c^{-2})^{\ell+1} \langle r_c \rangle^{-6} \mathbb{K}(r, r_c, c) \mathcal{B}(r, r_c) \mathbb{K}(r_c, r_c, c) \mathcal{B}(r_c, s_0) \mathcal{L}_{J, \ell'}(r_c, s_0) \\ &\quad \times \int_0^\infty \chi_c \frac{(u-c)u'(s)}{(u-c)^2 + \varepsilon^2} \frac{|s-r_c|}{r_c} \left(k + \left| \log \frac{k|s-r_c|}{r_c} \right| \right) ds \\ &\lesssim \max(r_c^2, r_c^{-2})^{\ell+1} \langle r_c \rangle^{-6} \mathbb{K}(r, r_c, c) \mathcal{B}(r, r_c) \mathbb{K}(r_c, s_0, c) \mathcal{B}(r_c, s_0) \mathcal{L}_{J+1, \ell'}(r_c, s_0) \\ &\lesssim \mathbb{K}(r, s_0, c) \mathcal{B}(r, s_0) \mathcal{L}_{J+1, \ell'+\ell+1}(r, s_0). \end{aligned}$$

This completes the proof of (6.13a).

$r\partial_r$ estimates (6.13b) Next, we estimate $r\partial_r \tilde{K}$. By continuity of \mathcal{G} and K ,

$$\begin{aligned} r\partial_r \frac{\tilde{K}(r, s_0, c)}{u'(s_0)} &= r\partial_r H_\infty(r, c - i\varepsilon) \int_0^r \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \frac{K(s, s_0, c)}{u'(s_0)} \frac{wH_0(s, c - i\varepsilon)}{M(c - i\varepsilon)} ds \\ &\quad + r\partial_r H_0(r, c - i\varepsilon) \int_r^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \frac{K(s, s_0, c)}{u'(s_0)} \frac{wH_\infty(s, c - i\varepsilon)}{M(c - i\varepsilon)} ds. \end{aligned}$$

From the arguments used to deduce (6.13a), for $|r - r_c| \geq r_c/k$ and the lemmas of Sect. 5, we can derive (6.13b) in the same manner as (6.13a). The details are omitted for brevity. Turn to the case $|r - r_c| < r_c/k$. In this region, our goal is to deduce the logarithmic upper bound; in order to avoid losing an additional logarithm we will need to extract a cancellation. Write,

$$\begin{aligned} r\partial_r \frac{\tilde{K}(r, s_0, c)}{u'(s_0)} &= ru' R_{r, \infty}^\varepsilon(z) \int_0^r \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} K(s, s_0, c) \frac{wH_0(s, c - i\varepsilon)}{M(c - i\varepsilon)} ds \\ &\quad - ru' R_{0, r}^\varepsilon(z) \int_r^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} K(s, s_0, c) \frac{wH_\infty(s, c - i\varepsilon)}{M(c - i\varepsilon)} ds \\ &\quad + (r\partial_r H_\infty(r, z) - ru' R_{r, \infty}^\varepsilon(z)) \int_0^r \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} K(s, s_0, c) \\ &\quad \times \frac{wH_0(s, c - i\varepsilon)}{M(c - i\varepsilon)} ds \\ &\quad + (r\partial_r H_0(r, c - i\varepsilon) + ru' R_{0, r}^\varepsilon(z)) \int_r^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} K(s, s_0, c) \\ &\quad \times \frac{wH_\infty(s, c - i\varepsilon)}{M(c - i\varepsilon)} ds \\ &= \sum_{j=1}^4 T_j. \end{aligned}$$

From Lemmas 5.4 and 5.5, we deduce $r\partial_r H_\infty - ru' R_{r, \infty}^\varepsilon$ is bounded near $r \approx r_c$, and hence T_3 and T_4 are treated using techniques used above in the proof of (6.13a). Due to the inability to extract an additional cancellation, these terms are logarithmically

unbounded (since $\int_0^r \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} g(s) ds$ is singular near $r \sim r_c$ for any smooth g ; see Lemma A.3):

$$(|T_3| + |T_4|) \mathbf{1}_{|r-r_c| < r_c/k} \lesssim |u'(s_0)| \mathbb{K}(r, s_0, c) \mathcal{B}(r, s_0) \mathcal{L}_{J+1, \ell'+\ell+1+\eta}(r, s_0) \left(k + \left| \log \frac{k|r-r_c|}{r_c} \right| \right),$$

which is consistent with (6.13b). For T_1 and T_2 , first divide via

$$\begin{aligned} T_1 + T_2 &= ru' R_{r,\infty}^\varepsilon(z) \int_0^r \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_c K(s, s_0, c) \frac{wH_0(s, c-i\varepsilon)}{M(c-i\varepsilon)} ds \\ &\quad - ru' R_{0,r}^\varepsilon(z) \int_r^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_c K(s, s_0, c) \frac{wH_\infty(s, c-i\varepsilon)}{M(c-i\varepsilon)} ds \\ &\quad + ru' R_{r,\infty}^\varepsilon(z) \int_0^r \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_{\neq} K(s, s_0, c) \frac{wH_0(s, c-i\varepsilon)}{M(c-i\varepsilon)} ds \\ &\quad - ru' R_{0,r}^\varepsilon(z) \int_r^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_{\neq} K(s, s_0, c) \frac{wH_\infty(s, c-i\varepsilon)}{M(c-i\varepsilon)} ds \\ &= T_{1;c} + T_{2;c} + T_{1;\neq} + T_{2;\neq}. \end{aligned}$$

The terms $T_{1;\neq}$ and $T_{2;\neq}$ are also treated as in (6.13a) and are hence omitted for the sake of brevity (in particular, $R_{0,r}^\varepsilon$ and $R_{r,\infty}^\varepsilon$ contain a logarithmic singularity by Lemma C.1 but the integrals involving K do not due to χ_{\neq}). For the remaining terms we use the following cancellation:

$$\begin{aligned} T_{1;c} + T_{2;c} &= ru' \left(R_{r,\infty}^\varepsilon(z) - E(r_c, z) \int_r^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_c ds \right) \\ &\quad \times \int_0^r \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_c K(s, s_0, c) \frac{wH_0(s, c-i\varepsilon)}{M(c-i\varepsilon)} ds \\ &\quad - ru' \left(R_{0,r}^\varepsilon(z) - E(r_c, z) \int_0^r \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_c ds \right) \\ &\quad \times \int_r^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_c K(s, s_0, c) \frac{wH_\infty(s, c-i\varepsilon)}{M(c-i\varepsilon)} ds \\ &\quad + ru' \left(E(r_c, z) \int_r^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_c ds \right) \int_0^r \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_c \\ &\quad \times \left(K(s, s_0, c) \frac{wH_0(s, c-i\varepsilon)}{M(c-i\varepsilon)} - K(r_c, s_0, c) \frac{w(r_c)}{u'(r_c)M(c-i\varepsilon)} \right) ds \\ &\quad - ru' \left(E(r_c, z) \int_0^r \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_c ds \right) \int_r^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_c \\ &\quad \times \left(K(s, s_0, c) \frac{wH_\infty(s, c-i\varepsilon)}{M(c-i\varepsilon)} - K(r_c, s_0, c) \frac{w(r_c)}{u'(r_c)M(c-i\varepsilon)} \right) ds. \end{aligned}$$

Note that e.g.

$$\begin{aligned} R_{r,\infty}^\varepsilon(z) - E(r_c, z) \int_r^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_c ds \\ = \int_r^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} (E(s, z) - E(r_c, z)) \chi_c ds, \end{aligned}$$

does not have any logarithmic singularities by arguments used in the proof Lemma 5.2. From here, the above terms are estimated in manners analogous to the arguments in the proof of (6.13a) and are hence omitted for the sake of brevity.

Derivatives involving ∂_{r_c} First we prove (6.13c). Taking a ∂_{r_c} derivative directly yields (integrating by parts in s),

$$\begin{aligned} r_c \partial_{r_c} \tilde{K} &= \int_0^\infty r_c \partial_{r_c} \left(\frac{(u-c)u'(s)}{(u-c)^2 + \varepsilon^2} \right) w(s) \frac{\mathcal{G}(r, s, c + i\varepsilon)}{u'(s)} K(s, s_0, c) ds \\ &\quad + \int_0^\infty \chi_{\neq} \left(\frac{(u-c)u'(s)}{(u-c)^2 + \varepsilon^2} \right) w(s) r_c \partial_{r_c} \left(\frac{\mathcal{G}(r, s, c + i\varepsilon)}{u'(s)} K(s, s_0, c) \right) ds \\ &\quad + \int_0^\infty u'(r_c) r_c \left(\frac{(u-c)u'(s)}{(u-c)^2 + \varepsilon^2} \right) w(s) \partial_G^{(s)} \left(\chi_c \frac{\mathcal{G}(r, s, c + i\varepsilon)}{u'(s)} K(s, s_0, c) \right) ds \\ &= \sum_{j=1}^3 \tilde{K}_j. \end{aligned}$$

Due to Definition 6.7 and Lemma 6.10, we can apply the methods used above to prove (6.13a) to prove that (6.13c) holds for \tilde{K}_1 and \tilde{K}_2 . For \tilde{K}_3 , we need to argue that the presence of ∂_{r_c} derivatives on \mathcal{G} does not stop us from finding a similar cancellation as we used in the proof of (6.13b) above. To that end, note that for $|r - r_c| < r_c/k$:

$$r_c \partial_{r_c} H_0(r, z) = r_c u'(r_c) \partial_G^{(r)} H_0(r, z) - \frac{r_c u'(r_c)}{r u'(r)} r \partial_r H_0(r, z),$$

The former term is bounded from Lemma 5.5 and hence in the neighborhood of $r \approx r_c$, we can extract the *same* cancellation in \tilde{K}_3 as we did in (6.13b) so that we deduce only one power of logarithm is lost. We omit the repetitive details for brevity, which concludes the proof of (6.13c).

Next, consider the proof of (6.14a) (which is relevant for the region where $r \not\approx r_c$ and $s_0 \approx r_c$). In this case,

$$\begin{aligned} \partial_G^{(s_0)} \tilde{K}(r, s_0, c) &= \int_0^\infty \frac{1}{u'(r_c)} \partial_{r_c} \left(\frac{(u-c)u'(s)}{(u-c)^2 + \varepsilon^2} \right) w(s) \chi_{\neq} \frac{\mathcal{G}(r, s, c + i\varepsilon)}{u'(s)} K(s, s_0, c) ds \\ &\quad + \int_0^\infty \left(\frac{(u-c)u'(s)}{(u-c)^2 + \varepsilon^2} \right) \partial_G^{(s_0)} \left(w(s) \chi_{\neq} \frac{\mathcal{G}(r, s, c + i\varepsilon)}{u'(s)} K(s, s_0, c) \right) ds \\ &\quad + \int_0^\infty \left(\frac{(u-c)u'(s)}{(u-c)^2 + \varepsilon^2} \right) \partial_G^{(s, s_0)} \left(w(s) \left(\frac{\mathcal{G}(r, s, c + i\varepsilon)}{u'(s)} \right) \chi_c K(s, s_0, c) \right) ds, \end{aligned}$$

Notice that since $r \not\approx r_c$, we do not need to obtain additional regularity in r in order to satisfy Definition 6.7. From here, we may again apply the methods of (6.13a) to deduce the desired estimates.

Next consider (6.14b) (which holds in $r \approx r_c$ but $s_0 \not\approx r_c$) Here,

$$\begin{aligned} \partial_G^{(r)} \tilde{K} &= \int_0^\infty \frac{\chi_{\neq}}{u'(r_c)} \partial_{r_c} \left(\frac{(u-c)u'(s)}{(u-c)^2 + \varepsilon^2} \right) w(s) \frac{\mathcal{G}(r, s, c + i\varepsilon)}{u'(s)} K(s, s_0, c) ds \\ &\quad + \int_0^\infty \left(\frac{(u-c)u'(s)}{(u-c)^2 + \varepsilon^2} \right) w(s) \partial_G^{(r)} \left(\chi_{\neq} \frac{\mathcal{G}(r, s, c + i\varepsilon)}{u'(s)} K(s, s_0, c) \right) ds \\ &\quad + \int_0^\infty \left(\frac{(u-c)u'(s)}{(u-c)^2 + \varepsilon^2} \right) w(s) \partial_G^{(r,s)} \left(\chi_c \frac{\mathcal{G}(r, s, c + i\varepsilon)}{u'(s)} K(s, s_0, c) \right) ds. \end{aligned} \quad (6.23)$$

Obtaining boundedness estimates is again a straightforward adaptation of the proof of (6.13a). Next, consider obtaining (6.14c). For this we apply an $r \partial_r$ to (6.23), however, some care must be taken due to the jumps in the derivatives of \mathcal{G} :

$$\begin{aligned} r \partial_r \partial_G^{(r)} \tilde{K} &= \int_0^\infty \frac{\chi_{\neq}}{u'(r_c)} \partial_{r_c} \left(\frac{(u-c)u'(s)}{(u-c)^2 + \varepsilon^2} \right) w(s) r \partial_r \frac{\mathcal{G}(r, s, c + i\varepsilon)}{u'(s)} K(s, s_0, c) ds \\ &\quad + \int_0^\infty \left(\frac{(u-c)u'(s)}{(u-c)^2 + \varepsilon^2} \right) w(s) r \partial_r \partial_G^{(r)} \left(\chi_{\neq} \frac{\mathcal{G}(r, s, c + i\varepsilon)}{u'(s)} K(s, s_0, c) \right) ds \\ &\quad + \int_0^\infty \left(\frac{(u-c)u'(s)}{(u-c)^2 + \varepsilon^2} \right) w(s) r \partial_r \partial_G^{(r,s)} \left(\chi_c \frac{\mathcal{G}(r, s, c + i\varepsilon)}{u'(s)} K(s, s_0, c) \right) ds \\ &= \sum_{j=1}^3 \tilde{K}_j; \end{aligned} \quad (6.24)$$

note that two of the potential boundary terms from $r = s$ vanished due to the presence of χ_{\neq} and the assumption that $|r - r_c| < r_c/k$ whereas the other two boundary terms coming from $r = s$ vanished due to the symmetric structure of (6.19). The \tilde{K}_1 and \tilde{K}_2 terms are estimated in essentially the same manner as done previously for (6.13b) and are omitted for the sake of brevity. To treat \tilde{K}_3 , again we use a cancellation analogous that used in (6.13b) to avoid losses of higher powers of logarithms. This is only an issue if both derivatives land on \mathcal{G} (otherwise the situation is essentially the same as (6.13b)). Recall the identities:

$$\begin{aligned} \partial_r \partial_G H_0 &= \partial_r \partial_G \left(\frac{1}{u'P(r, c \pm i\varepsilon)} \right) - u'(\partial_G P) (R_{0,r}^\varepsilon(z) \mp iE_{0,r}^\varepsilon) \\ &\quad - (u-z)(\partial_r \partial_G P) (R_{0,r}^\varepsilon(z) \mp iE_{0,r}^\varepsilon) - (\partial_G P) u'(r) E(r, z) \\ &\quad + (u'P + (u-z)\partial_r P) \int_0^r \frac{u'(s)}{(u-z)} \partial_G (\chi_c E) ds + Pu'(r) \partial_G (\chi_c E) \\ &\quad - \frac{1}{u'(r_c)} (\partial_r \phi) \int_0^r \chi_{\neq} \frac{u'(s)u'(r_c)}{(u-z)^2} E(s, z) - \frac{u'(s)}{(u-z)} \chi_{\neq} \partial_{r_c} E(s, z) ds, \end{aligned}$$

and

$$\begin{aligned} \partial_r \partial_G H_\infty &= \partial_r \partial_G \left(\frac{1}{u' P(r, c \pm i\varepsilon)} \right) + u' (\partial_G P) (R_{r,\infty}^\varepsilon(z) \mp i E_{r,\infty}^\varepsilon) \\ &\quad + (u - z) (\partial_r \partial_G P) (R_{r,\infty}^\varepsilon(z) \mp i E_{r,\infty}^\varepsilon) + (\partial_G P) u'(r) E(r, z) \\ &\quad - (u' P + (u - z) \partial_r P) \int_r^\infty \frac{u'(s)}{(u - z)} \partial_G (\chi_c E) ds - P u'(r) \partial_G (\chi_c E) \\ &\quad + \frac{1}{u'(r_c)} (\partial_r \phi) \int_r^\infty \chi_{\neq} \frac{u'(s) u'(r_c)}{(u - z)^2} E(s, z) - \frac{u'(s)}{(u - z)} \chi_{\neq} \partial_{r_c} E(s, z) ds. \end{aligned}$$

We see that the logarithmically singular terms between $\partial_r \partial_G H_\infty$ and $\partial_r \partial_G H_0$ in \tilde{K}_3 in (6.24) have a structure similar to that of $\partial_r H_\infty$ and $\partial_r H_0$ that was exploited in the proof of (6.13b). Note that singular terms in e.g. $\partial_r \partial_G H_0$ are $u' \partial_G R_{0,r}^\varepsilon$ and $u' P \int_0^r \frac{u'}{u-z} \partial_G (\chi_c E) ds$. After these observations, the proof follows analogously and is hence omitted for the sake of brevity. This completes the proof of (6.14c). The proofs of (6.14d) and (6.14e) are straightforward variants of the techniques used on the other inequalities in (6.14) and are hence omitted for the sake of brevity.

Hölder Regularity in s_0 near r_c estimates Notice that:

$$\begin{aligned} \tilde{K}(r, s_0, c) - \tilde{K}(r, r_c, c) &= \int_0^\infty (K(s, s_0, c) \\ &\quad - K(s, r_c, c)) \frac{(u - c)}{(u - c)^2 + \varepsilon^2} w(s) \mathcal{G}(r, s, c - i\varepsilon) ds. \end{aligned}$$

Consider first obtaining (6.15a). The only difference between proving (6.13a) and proving (6.15a) is the lack of an analogous Hölder regularity estimate on $\partial_r \tilde{K}(r, s_0, c) - \partial_r \tilde{K}(r, r_c, c)$ (which was used to control $s \approx r_c$). This is dealt with via the following (see the proof of Lemma A.4 for a similar approach): for $\theta \in (0, 1)$,

$$\begin{aligned} &\left| \tilde{K}(r, s_0, c) - \tilde{K}(r, r_c, c) - \tilde{K}(r_c, s_0, c) + \tilde{K}(r_c, r_c, c) \right| \\ &\leq \left(\left| \tilde{K}(r, s_0, c) - \tilde{K}(r_c, s_0, c) \right| + \left| \tilde{K}(r, r_c, c) - \tilde{K}(r_c, r_c, c) \right| \right)^\theta \\ &\quad \times \left(\left| \tilde{K}(r, s_0, c) - \tilde{K}(r, r_c, c) \right| + \left| \tilde{K}(r_c, s_0, c) - \tilde{K}(r_c, r_c, c) \right| \right)^{1-\theta} \\ &\lesssim \left(\frac{|r - r_c|}{r_c} \right)^\theta \left(k + \left| \log \left| k \frac{r - r_c}{r_c} \right| \right| \right)^\theta \left(\frac{k |s_0 - r_c|}{r_c} \right)^{\gamma(1-\theta)}, \end{aligned}$$

and hence we choose $\gamma' = \gamma(1 - \theta)$. Other than this slight difference, the proof of (6.15a) follows as in (6.13a). The rest of the estimates in (6.15) are adapted in essentially this same way; the details are omitted for the sake of brevity.

Convergence as $\varepsilon \rightarrow 0$ As seen above, verifying the boundedness estimate contains most of the non-trivial work. Hence, for the convergence estimate, we will focus on

this. That is, if we define

$$\tilde{K}_0(r, s_0, c) = p.v. \int_0^\infty \frac{u'(s)}{u(s) - c} \frac{\mathcal{G}(r, s, c + i0)}{u'(s)} K_0(s, s_0, c) ds,$$

we are interested in obtaining an estimate of the following form for some $\eta_1, \eta_2 > 0$:

$$\left| \tilde{K}_\varepsilon(r, s_0, c) - \tilde{K}_0(r, s_0, c) \right| \lesssim_{\eta_1, \eta_2} \varepsilon^{\eta_1} \mathbb{K}(r, s_0, c) \mathcal{B}(r, s_0) \mathcal{L}_{J+1, \ell+\ell'+1+\eta_2}(r, s_0).$$

Write the difference as the following:

$$\begin{aligned} & \tilde{K}_\varepsilon(r, s_0, c) - \tilde{K}_0(r, s_0, c) \\ &= \int \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \frac{\mathcal{G}(r, s, c + i\varepsilon) - \mathcal{G}(r, s, c + i0)}{u'(s)} K_\varepsilon(s, s_0, c) ds \\ &+ \int \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \frac{\mathcal{G}(r, s, c - i0)}{u'(s)} (K_\varepsilon(s, s_0, c) - K_0(s, r_c, c)) ds \\ &+ p.v. \int \left(\frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} - \frac{u'}{u-c} \right) \frac{\mathcal{G}(r, s, c + i0)}{u'(s)} K_0(s, s_0, c) ds \\ &= \sum_{j=1}^3 T_j. \end{aligned}$$

The terms T_1 and T_2 are treated in essentially the same way as above (see Lemma 6.10 for control on $\mathcal{G}(r, s, c + i\varepsilon) - \mathcal{G}(r, s, c + i0)$). Hence, it remains to treat T_3 . Sub-divide via:

$$\begin{aligned} T_3 &= p.v. \int \left(\frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} - \frac{u'}{u-c} \right) (\chi_c + \chi_{\neq}) \frac{\mathcal{G}(r, s, c + i0)}{u'(s)} K^0(s, s_0, c) ds \\ &=: T_{3c} + T_{3\neq}. \end{aligned}$$

For $T_{3\neq}$ we apply (A.3) to obtain some decay and then we argue as in (6.13a). For T_{3c} , we apply (A.4) (which applies due to the various regularity and convergence estimates satisfied by K and \mathcal{G}).

As remarked above, the rest of the claimed inequalities are a straightforward adaptation of the above arguments and are hence omitted for the sake of brevity.

Adaptation to $\mathcal{O}_{S;\varepsilon}^{(2)}$ case The case $j = 2$ is essentially the same. As above, we begin by dividing based on the critical layer:

$$\begin{aligned} \frac{\tilde{K}^{(2)}(s_0, r, c)}{u'(r)} &:= \int_0^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} (\chi_c + \chi_{\neq}) w(s) \frac{\mathcal{G}(r, s, c + i\varepsilon)}{u'(r)} \frac{K(s_0, s, c)}{u'(s)} ds \\ &= \tilde{K}_c + \tilde{K}_{\neq}. \end{aligned} \quad (6.25)$$

The contribution from near the critical layer, \tilde{K}_c , is treated in essentially the same manner as in the $j = 1$ case and is hence omitted for brevity. Hence, we need only to

check the tedious but simple contributions of \tilde{K}_{\neq} ; we omit the intermediate steps for the sake of brevity: for all $\eta > 0$,

$$\begin{aligned} \left| \frac{1}{u'(r)} \tilde{K}_{\neq}(s_0, r, c) \right| \mathbf{1}_{r_c \leq 1} &\lesssim \mathbf{1}_{r_c \leq 1} \int_0^\infty \left(\frac{k}{\max(s^2, r_c^2) s^{1+2\ell}} \mathbf{1}_{s \leq 1} + s^{2\ell-7} \mathbf{1}_{s > 1} \right) \\ &\quad \times \chi_{\neq} \mathbb{K}(s, r, c) \mathcal{B}(s, r) \mathbb{K}(s_0, s, c) \mathcal{B}(s_0, s) ds \\ &\lesssim_{\eta} \mathbf{1}_{r_c \leq 1} \mathbb{K}(s_0, r, c) \mathcal{B}(s_0, r) \mathcal{L}_{1, \ell+\ell'+1+\eta}(s_0, r). \end{aligned}$$

Similarly, we have the contributions from $r_c > 1$: for all $\eta > 0$,

$$\begin{aligned} \left| \frac{1}{u'(r)} \tilde{K}_{\neq}(s_0, r, c) \right| \mathbf{1}_{r_c > 1} &\lesssim \mathbf{1}_{r_c > 1} \int_0^\infty \left(\frac{1}{s^{1+2\ell}} \mathbf{1}_{s \leq 1} + \max(r_c^2, s^2) s^{2\ell-7} \mathbf{1}_{s > 1} \right) \\ &\quad \times \chi_{\neq} \mathbb{K}(s, r, c) \mathcal{B}(s, r) \mathbb{K}(s_0, s, c) \mathcal{B}(s_0, s) ds \\ \left| \frac{1}{u'(r)} \tilde{K}_{\neq}(s_0, r, c) \right| \mathbf{1}_{r_c > 1} &\lesssim \mathcal{B}(s_0, r) \mathcal{L}_{1, \ell+\ell'+1+\eta}(s_0, r). \end{aligned}$$

This completes the proof of (6.16a) in Definition 6.8. As discussed above, this estimate involves most of the non-trivial work necessary to deduce the rest of Definition 6.8, and hence the remaining estimates are omitted for the sake of brevity. \square

Proof of Lemma 6.12 Consider the case $j = 1$ first. As there are no singular integrals, we do not need to separate the critical layer from the rest, and hence the calculations are a small variant of those done to estimate \tilde{K}_{\neq} in the proof of Lemma 6.11 above. As above, this is a tedious, but simple and direct, calculation, and hence we omit the intermediate steps. For the case of $r_c \leq 1$, using Definition 6.7, Lemma 6.10, and the requirement that $\ell + \ell' < k - 3/2$ (to retain integrability at zero and infinity), there holds

$$\begin{aligned} \left| \frac{1}{u'(s_0)} \tilde{K}(r, s_0, c) \right| \mathbf{1}_{r_c \leq 1} &\leq \int_0^\infty \left| u'(s) w(s) \frac{\mathcal{G}(r, s, c + i\varepsilon)}{u'(s)} \right| \left| \frac{K(s, s_0, c)}{u'(s_0)} \right| ds \\ &\lesssim \int_0^\infty \frac{k^2}{s^3} \max \left(\frac{1}{s^{2\ell}}, s^{2\ell} \right) \mathbb{K}(r, s, c) \mathcal{B}(r, s) \\ &\quad \times \mathbb{K}(s, s_0, c) \mathcal{B}(s, s_0) \mathcal{L}(s, s_0, c) ds \\ &\lesssim \mathbb{K}(r, s_0, c) \mathcal{B}(r, s_0) \mathcal{L}_{J+2, \ell+\ell'+1+\eta}(r, s_0), \end{aligned}$$

and similarly for $r_c \geq 1$:

$$\begin{aligned} \left| \frac{1}{u'(s_0)} \tilde{K}(r, s_0, c) \right| \mathbf{1}_{r_c \geq 1} &\lesssim \int_0^\infty \frac{k^2}{s^3} \max \left(\frac{1}{s^{2\ell}}, s^{2\ell} \right) \mathcal{B}(r, s) \mathcal{B}(s, s_0) \mathcal{L}(s, s_0, c) ds \\ &\lesssim \mathcal{B}(r, s_0) \mathcal{L}_{J+2, \ell+\ell'+1+\eta}(r, s_0), \end{aligned}$$

which completes the proof of (6.13a). As in the proof of Lemma 6.11, the remaining estimates in Definition 6.7 follow from straightforward variants of the proof of (6.13a), and hence we omit these arguments for brevity.

As above in the proof of Lemma 6.11, the case $j = 2$ follows in a similar manner with slightly different integrals. The repetitive details are omitted for brevity. \square

Proof of Lemma 6.13 We will consider only the $j = 2$ case; $j = 1$ is the same. For $\mathcal{O}_r^{(2)}$ (by symmetry of \mathcal{G}),

$$\frac{\tilde{K}(s_0, r, c)}{u'(r)} = \int_0^\infty \frac{1}{u'(r)} s \partial_s \mathcal{G}(s, r, c - i\varepsilon) \frac{w(s)}{s} K(s_0, s, c) ds,$$

Boundedness estimate (6.16a) From Lemma 6.10 and definition 6.8, the proof of (6.16a) is essentially the same as the corresponding estimate made on $\mathcal{O}_G^{(2)}$.

Regularity estimates First consider (6.16c). Taking an $r \partial_r$ derivative gives (using the definition of M (2.32)),

$$\begin{aligned} r \partial_r \tilde{K}(s_0, r, c) &= w(r) K(s_0, r, c) + r \partial_r H_\infty(r, c - i\varepsilon) \int_0^r K(s_0, s, c) \frac{w \partial_s H_0(s, c - i\varepsilon)}{M(c - i\varepsilon)} ds \\ &\quad + r \partial_r H_0(r, c - i\varepsilon) \int_r^\infty K(s_0, s, c) \frac{w \partial_s H_\infty(s, c - i\varepsilon)}{M(c - i\varepsilon)} ds. \end{aligned}$$

This does not present any new challenges and hence (6.16c) is deduced as in Lemma 6.11 (though significantly easier, as no delicate cancellation is necessary) and is hence omitted for the sake of brevity.

The more subtle problem is ∂_{r_c} derivatives:

$$\begin{aligned} \partial_{r_c} \frac{\tilde{K}(s_0, r, c)}{u'(r)} &= \int_0^\infty \frac{1}{u'(r)} \partial_s \partial_{r_c} \mathcal{G}(r, s, c - i\varepsilon) w(s) K(s_0, s, c) ds \\ &\quad + \int_0^\infty \frac{1}{u'(r)} \partial_s \mathcal{G}(r, s, c - i\varepsilon) w(s) \partial_{r_c} K(s_0, s, c) ds \\ &= \tilde{K}_1 + \tilde{K}_2. \end{aligned}$$

The treatment of \tilde{K}_2 is similar to the proof of (6.16a) and is hence omitted. Consider next \tilde{K}_1 . The trick is to integrate by parts so that two derivatives never land on the same kernel (note that the boundary terms vanish):

$$\begin{aligned} \tilde{K}_1 &= - \int_0^r \frac{1}{u'(r)} \partial_s (\partial_{r_c} H_\infty(r, c - i\varepsilon) H_0(s, c - i\varepsilon) \\ &\quad + H_\infty(r, c - i\varepsilon) \partial_{r_c} H_0(s, c - i\varepsilon)) (w(s) K(s_0, s, c)) ds \\ &\quad - \int_r^\infty \frac{1}{u'(r)} \partial_s (\partial_{r_c} H_0(r, c - i\varepsilon) H_\infty(s, c - i\varepsilon) \\ &\quad + H_\infty(s, c - i\varepsilon) \partial_{r_c} H_0(r, c - i\varepsilon)) (w(s) K(s_0, s, c)) ds \\ &= \int_0^r \frac{1}{u'(r)} (\partial_{r_c} H_\infty(r, c - i\varepsilon) H_0(s, c - i\varepsilon) \\ &\quad + H_\infty(r, c - i\varepsilon) \partial_{r_c} H_0(s, c - i\varepsilon)) \partial_s (w(s) K(s_0, s, c)) ds \end{aligned}$$

$$\begin{aligned}
& + \int_r^\infty \frac{1}{u'(r)} (\partial_{r_c} H_0(r, c - i\varepsilon) H_\infty(s, c - i\varepsilon) \\
& + H_\infty(s, c - i\varepsilon) \partial_{r_c} H_0(r, c - i\varepsilon)) \partial_s (w(s) K(s_0, s, c)) ds.
\end{aligned}$$

The log-boundedness (6.16d) follows from similar arguments as in Lemmas 6.12 and 6.11. Consider (6.17c) (which is relevant for $r \not\approx r_c$ and $s_0 \approx r_c$) for which we apply the same approach:

$$\begin{aligned}
\partial_G^{(s_0)} \tilde{K}(s_0, r, c) &= - \int_0^\infty \frac{1}{u'(r_c)} \partial_{r_c} \mathcal{G}(s, r, c - i\varepsilon) \partial_s (w(s) K(s_0, s, c)) ds \\
&+ \int_0^\infty \partial_s \mathcal{G}(s, r, c - i\varepsilon) w(s) \partial_G^{(s_0)} K(s_0, s, c) ds.
\end{aligned}$$

Due to the restriction $|r - r_c| \geq r_c/k$ we can apply analogous estimates as in (6.16c) to deduce (6.17c). Similar arguments deduce (6.17a) and (6.17g) which we omit for brevity.

Consider next estimate (6.17e). Some care is required due to the jumps in the derivatives of \mathcal{G} (recall this is only relevant in the case $|r - r_c| < r_c/k$ and $|s_0 - r_c| \geq r_c/k$):

$$\begin{aligned}
& r \partial_r \partial_G^{(r)} \tilde{K}(s_0, r, c) \\
&= r \partial_r \int_0^r \frac{1}{M} \left(\partial_G H_\infty(r, z) \partial_s H_0(s, z) + H_\infty(r, z) \frac{1}{u'(r_c)} \partial_{r_c} \partial_s H_0(s, z) \right. \\
&\quad \left. - \frac{\partial_{r_c} M}{M} H_\infty(r, z) \partial_s H_0(s, z) \right) w(s) K(s_0, s, c) ds \\
&\quad + r \partial_r \int_r^\infty \frac{1}{M} \left(\partial_G H_0(r, z) \partial_s H_\infty(s, z) + H_0(r, z) \frac{1}{u'(r_c)} \partial_{r_c} \partial_s H_\infty(s, z) \right. \\
&\quad \left. - \frac{\partial_{r_c} M}{M} H_\infty(r, z), \partial_s H_0(s, z) \right) w(s) K(s_0, s, c) ds \\
&\quad + r \partial_r \int_0^r \frac{1}{M} H_\infty(r, z) \partial_s H_0(s, z) w(s) \frac{1}{u'(r_c)} \partial_{r_c} K(s_0, s, c) ds \\
&\quad + r \partial_r \int_r^\infty \frac{1}{M} H_0(r, z) \partial_s H_\infty(s, z) w(s) \frac{1}{u'(r_c)} \partial_{r_c} K(s_0, s, c) ds \\
&= \frac{r}{M} \left(\partial_G H_\infty(r, z) \partial_r H_0(r, z) + H_\infty(r, z) \frac{1}{u'(r_c)} \partial_{r_c} \partial_r H_0(r, z) \right. \\
&\quad \left. - \frac{\partial_{r_c} M}{M} H_\infty(r, z) \partial_r H_0(r, z) \right) w(r) K(s_0, r, c) \\
&\quad - \frac{r}{M} \left(\partial_G H_0(r, z) \partial_r H_\infty(r, z) + H_0(r, z) \frac{1}{u'(r_c)} \partial_{r_c} \partial_r H_\infty(r, z) \right. \\
&\quad \left. - \frac{\partial_{r_c} M}{M} H_\infty(r, z), \partial_r H_0(r, z) \right) w(r) K(s_0, r, c)
\end{aligned}$$

$$\begin{aligned}
& + \frac{r}{M} H_\infty(r, z) \partial_r H_0(r, z) w(r) \frac{1}{u'(r_c)} \partial_{r_c} K(s_0, r, c) \\
& - \frac{r}{M} H_0(r, z) \partial_r H_\infty(r, z) w(r) \frac{1}{u'(r_c)} \partial_{r_c} K(s_0, r, c) \\
& + \int_0^r \frac{1}{M} \left(r \partial_r \partial_G H_\infty(r, z) \partial_s H_0(s, z) + r \partial_r H_\infty(r, z) \frac{1}{u'(r_c)} \partial_{r_c} \partial_s H_0(s, z) \right. \\
& \quad \left. - \frac{\partial_{r_c} M}{M} r \partial_r H_\infty(r, z) \partial_s H_0(s, z) \right) w(s) K(s_0, s, c) ds \\
& + \int_r^\infty \frac{1}{M} \left(r \partial_r \partial_G H_0(r, z) \partial_s H_\infty(s, z) + r \partial_r H_0(r, z) \frac{1}{u'(r_c)} \partial_{r_c} \partial_s H_\infty(s, z) \right. \\
& \quad \left. - \frac{\partial_{r_c} M}{M} r \partial_r H_\infty(r, z) \partial_s H_0(s, z) \right) w(s) K(s_0, s, c) ds \\
& + \int_0^r \frac{1}{M} r \partial_r H_\infty(r, z) \partial_s H_0(s, z) w(s) \frac{1}{u'(r_c)} \partial_{r_c} K(s_0, s, c) ds \\
& + \int_r^\infty \frac{1}{M} r \partial_r H_0(r, z) \partial_s H_\infty(s, z) w(s) \frac{1}{u'(r_c)} \partial_{r_c} K(s_0, s, c) ds \\
& = \sum_{j=1}^8 \tilde{K}_j.
\end{aligned}$$

There are several cancellations to observe. First, we observe that (recalling $|r - r_c| < r_c/k$) from Lemma 5.4,

$$\begin{aligned}
\partial_r H(r, z) - \partial_r H_\infty(r, z) &= r \partial_r \phi M(z) \\
|H_0(r, z) - H_\infty(r, z)| &\lesssim \mathbf{1}_{r_c \leq 1} \frac{|r - r_c|}{r_c^2} \left(k + \left| \log k \left| \frac{r - r_c}{r_c} \right| \right| \right) \\
&\quad + \mathbf{1}_{r_c > 1} r_c^3 \frac{|r - r_c|}{r_c} \left(k + \left| \log k \left| \frac{r - r_c}{r_c} \right| \right| \right),
\end{aligned}$$

and hence the terms $\tilde{K}_3 + \tilde{K}_4$ are only logarithmically singular at $r \approx r_c$ due to $\partial_{r_c} K$. For \tilde{K}_1 and \tilde{K}_2 we can uncover the cancellations via writing the following for $r \approx r_c$:

$$\begin{aligned}
& H_\infty(r, z) \frac{1}{u'(r_c)} \partial_{r_c} \partial_r H_0(r, z) - H_0(r, z) \frac{1}{u'(r_c)} \partial_{r_c} \partial_r H_\infty(r, z) \\
&= H_\infty(r, z) \partial_G \partial_r H_0(r, z) - H_0(r, z) \partial_G \partial_r H_\infty(r, z) \\
&\quad - \frac{1}{u'(r)} H_\infty(r, z) \partial_{rr} H_0(r, z) + \frac{1}{u'(r)} H_0(r, z) \partial_{rr} H_\infty(r, z) \\
&= H_\infty(r, z) \partial_G \partial_r H_0(r, z) - H_0(r, z) \partial_G \partial_r H_\infty(r, z) \\
&\quad - \frac{1}{u'(r)} H_\infty(r, z) \left(\frac{k^2 - 1/4}{r^2} H_0 - \frac{\beta H_0}{u - z} \right)
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{u'(r)} H_0(r, z) \left(\frac{k^2 - 1/4}{r^2} H_\infty - \frac{\beta H_\infty}{u - z} \right) \\
& = H_\infty(r, z) \partial_G \partial_r H_0(r, z) - H_0(r, z) \partial_G \partial_r H_\infty(r, z).
\end{aligned}$$

Note the commutation relation: $\partial_G \partial_r h = \partial_r \partial_G h + \frac{u''}{(u')^2} \partial_r h$. Hence, $\tilde{K}_1 + \tilde{K}_2$ is again only logarithmically singular via Lemmas 5.4 and 5.5. Finally the remaining terms \tilde{K}_5 through \tilde{K}_8 are treated using techniques used on previously made estimates in (6.17) and are hence omitted for the sake of brevity. The treatment of (6.17i) is similar and is hence omitted for brevity. This completes the estimates in (6.17).

Hölder regularity Consider next the estimates in (6.18). As in Lemma 6.11, write

$$\tilde{K}(s_0, r, c) - \tilde{K}(r_c, r, c) = \int_0^\infty (K(s_0, s, c) - K(r_c, s, c)) w(s) \partial_s \mathcal{G}(r, s, c - i\varepsilon) ds.$$

The estimate (6.18a) hence follows as in the proof of (6.16a). Consider next (6.18b) (recall $|r - r_c| \geq r_c/k$ in this case):

$$\begin{aligned}
& \partial_G^{(s_0)} \tilde{K}(s_0, r, c) - \partial_G^{(s_0)} \tilde{K}(r_c, r, c) \\
& = - \int_0^\infty \partial_G^{(s, s_0)} (w \chi_c (K(s_0, s, c) - K(r_c, s, c))) \mathcal{G}(r, s, c - i\varepsilon) ds \\
& \quad - \int_0^\infty \partial_G^{(s_0)} (w \chi_\neq (K(s_0, s, c) - K(r_c, s, c))) \partial_s \mathcal{G}(r, s, c - i\varepsilon) ds \\
& \quad + \int_0^\infty \chi_\neq (K(s_0, s, c) - K(r_c, s, c)) w(s) \frac{1}{u'(r_c)} \partial_{r_c} \partial_s \mathcal{G}(r, s, c - i\varepsilon) ds.
\end{aligned}$$

Note that no boundary terms appear (as in the proof of (6.16d) above). From here, the proof follows as in (6.16d) using the hypotheses on K . the treatment of the other inequalities in (6.18) follow via similar reductions and are hence omitted for the sake of brevity.

Convergence By the Lemmas in Sect. 5, the $r \partial_r$ derivative of H_0 and H_∞ satisfy analogous quantitative convergence estimates as H_0 and H_∞ themselves and hence the convergence as $\varepsilon \rightarrow 0$ is a straightforward consequence of arguments used previously; the details are omitted for the sake of brevity. \square

Finally, we verify that the original kernels satisfy the estimates necessary to run the iteration scheme.

Lemma 6.14 For $a \in S, \delta, G, rG$, $B_{Xa}^{(1)}$ is suitable $(0, 0)$ of Type I. For $a \in S, \delta, G, rG$, $B_{Xa}^{(2)}$ and B_{Ya} are suitable $(0, 0)$ of Type II.

Proof of Lemma 6.14 The treatment of $B_{X\delta; \varepsilon}^{(1)}$ is the only case not covered by Lemma 6.10. This follows Lemma 6.11—indeed:

$$B_{X\delta; \varepsilon}^{(1)}(r, s, c) = B_{XS; \varepsilon}^{(1)}(r, s, c) + \mathcal{O}_\delta^1[\beta B_{XS}^{(1)}],$$

and hence we may apply the lemma if we set $w(r) = \beta(r)$, and $\ell = \ell' = J = 0$. \square

Proof of Propositions 2.9 and 2.10 From Lemmas 6.1, 6.2, and (6.3), we can express all $\partial_G^j Y^\pm$ and $\partial_G^j X$ in terms of $\partial_G^\ell F$, $\partial_G^\ell F_*$, the coefficients derived in Lemma 6.2, and compositions of the integral operators in (6.4) and (6.6). Moreover, the coefficients are such that if one has ℓ'' compositions and $\partial_G^\ell F$ (or $\partial_G^\ell F_*$), then the *total* of all of the losses from all of the coefficients is ℓ' with $\ell + \ell' + \ell'' \leq j$. This condition ensures that the compositions all involve integrable functions (for $\varepsilon > 0$) and hence we iteratively apply Fubini's theorem and prove Lemmas 6.4–6.6. This, in turn, implies Proposition 2.9 and finally Lemmas 6.10–6.14 imply Proposition 2.10. \square

6.4 Vorticity Decomposition

In this section, we prove Propositions 2.13 and 2.14.

Proof of Proposition 2.13 We will first prove the lemma in the case $n = 0$, then explain how to extend to $n \leq k - 1$, and finally, to extend to $n \leq k$.

Case $n = 0$ Write

$$f_1 = \frac{F}{\sqrt{r}} + \frac{1}{2\pi i} \int_0^\infty e^{itk(u(r)-u(r_c))} \frac{2i\varepsilon u'(r_c)}{(u-c)^2 + \varepsilon^2} \chi_I \frac{\beta(r)}{\sqrt{r}} A(r, c, \varepsilon) dr_c \quad (6.26)$$

$$+ \frac{1}{2\pi i} \int_0^\infty e^{itk(u(r)-u(r_c))} \frac{(u-c)u'(r_c)}{(u-c)^2 + \varepsilon^2} \chi_I \frac{\beta(r)}{\sqrt{r}} X(r, c, \varepsilon) dr_c \quad (6.27)$$

$$= \frac{F}{\sqrt{r}} + f_{1;A} + f_{1;X}. \quad (6.28)$$

From the expansion for X , there holds

$$\begin{aligned} \sqrt{r} w_{F,\delta}^{-1} f_{1;X} &= \frac{e^{itku(r)}}{2\pi i} \int_0^\infty \frac{(u-c)u'(r_c)}{(u-c)^2 + \varepsilon^2} \chi_1(r, r_c) \frac{\beta(r) e^{-ikt c}}{w_{F,\delta}(r)} \\ &\quad \times \int_0^\infty \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} B_{X\delta;\varepsilon}(r, s, c) F(s) ds dr_c \\ &\quad + \frac{e^{itku(r)}}{2\pi i} \int_0^\infty \frac{(u-c)u'(r_c)}{(u-c)^2 + \varepsilon^2} \chi_1(r, r_c) \frac{\beta(r) e^{-ikt c}}{w_{F,\delta}(r)} \\ &\quad \times \int_0^\infty \int_0^\infty \frac{2i\varepsilon \beta(s_0)}{(u(s_0)-c)^2 + \varepsilon^2} w_{F,\delta/2}(s_0) \\ &\quad \times B_{XS;\varepsilon}^{(1)}(r, s_0, c) B_{XS;\varepsilon}^{(2)}(s_0, s', c) \frac{w_{F,\delta/4}(s')}{w_{F,\delta/2}(s_0)} \frac{(u-c)}{(u-c)^2 + \varepsilon^2} \\ &\quad \times \frac{F(s')}{w_{F,\delta/2}(s')} ds' ds_0 dr_c \\ &\quad + \frac{e^{itku(r)}}{2\pi i} \int_0^\infty \frac{(u-c)u'(r_c)}{(u-c)^2 + \varepsilon^2} \chi_1(r, r_c) \frac{\beta(r) e^{-ikt c}}{w_{F,\delta}(r)} \end{aligned}$$

$$\begin{aligned}
& \times \int_0^\infty \int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u(s_0) - c)^2 + \varepsilon^2} w_{F,\delta/2}(s_0) \\
& \times B_{XG;\varepsilon}^{(1)}(r, s_0, c) B_{XG;\varepsilon}^{(2)}(s_0, s', c) \frac{w_{F,\delta/4}(s')}{w_{F,\delta/2}(s_0)} \frac{F_*(s')}{w_{F,\delta/4}(s')} ds' ds_0 dr_c.
\end{aligned} \tag{6.29}$$

To pass to the limit, we apply Theorems B.1 and B.18 (together with Theorem B.2). Note that the requisite properties on the kernel are obtained by Lemmas 6.10 and 6.14 above. Hence, we have the strong L^2 limit:

$$\begin{aligned}
\lim_{\varepsilon \rightarrow 0} \sqrt{r} w_{F,\delta}^{-1} f_{1;X} &= e^{iktu(r)} p.v. \int_0^\infty \frac{u'(r_c)}{u - c} \chi_1(r, r_c) \frac{w_{F,\delta/2}(r_c)}{w_{F,\delta}(r)} \beta(r) \\
&\times \frac{B_{X\delta}(r, r_c, c)}{u'(r_c)} \frac{e^{-ikt c} F(r_c)}{w_{F,\delta/2}(r_c)} dr_c \\
&+ \frac{e^{itku(r)}}{2\pi i} p.v. \int_0^\infty \frac{u'(r_c)}{(u - c)} \chi_1(r, r_c) \frac{w_{F,\delta/2}(r_c)}{w_{F,\delta}(r)} \beta(r) \frac{B_{XS}^{(1)}(r, r_c, c)}{u'(r_c)} \\
&\times e^{-ikt c} \left(p.v. \int_0^\infty \beta(r_c) B_{XS}^{(2)}(r_c, s', c) \frac{w_{F,\delta/4}(s')}{w_{F,\delta/2}(r_c)} \right. \\
&\times \left. \frac{1}{(u - c)} \frac{F(s')}{w_{F,\delta/4}(s')} ds' \right) dr_c \\
&+ \frac{e^{itku(r)}}{2\pi i} p.v. \int_0^\infty \frac{u'(r_c)}{(u - c)} \chi_1(r, r_c) \frac{w_{F,\delta/2}(r_c)}{w_{F,\delta}(r)} \beta(r) \frac{B_{XG}^{(1)}(r, r_c, c)}{u'(r_c)} \\
&\times e^{-ikt c} \left(\int_0^\infty \beta(r_c) B_{XG}^{(2)}(r_c, s', c) \frac{w_{F,\delta/4}(s')}{w_{F,\delta/2}(r_c)} \frac{F_*(s')}{w_{F,\delta/4}(s')} ds' \right) dr_c.
\end{aligned}$$

A crucial point to notice is that if $r \leq 1$ then χ_1 implies that $r_c < 2r$. This is what allows to transfer the gain in r_c from \mathbb{K} to a gain in r in Theorem B.1 so that we may use the stronger $w_{F,\delta}$ as opposed to $w_{f,\delta}$. Theorems B.1 and B.18 also provide the following estimate: for all $\eta > 0$ (recall (2.12)),

$$\left\| \lim_{\varepsilon \rightarrow 0} \sqrt{r} w_{F,\delta}^{-1} f_{1;X} \right\|_{L^2} \lesssim_{\eta,\delta} k^\eta \|F\|_{L_{F,\delta/4}^2} + \|F_*\|_{L_{F,\delta/4}^2} \lesssim |k|^\eta \|F\|_{L_{F,\delta/4}^2} + |k| \left| \omega_{k,0}^{in} \right|.$$

Turn next to $f_{1;A}$ in (6.28) and expand via

$$\begin{aligned}
f_{1;A} &= \frac{1}{2\pi i} \int_0^\infty \frac{2i\varepsilon u'(r_c) e^{itk(u(r) - u(r_c))}}{(u - c)^2 + \varepsilon^2} \chi_I \frac{\beta(r)}{\sqrt{r}} (X(r, c, \varepsilon) + 2Y(r, c - i\varepsilon)) dr_c \\
&= f_{1;AX} + f_{1;AY}.
\end{aligned}$$

Further expand $f_{1;AX}$ via χ_1 and χ_2 :

$$\begin{aligned} f_{1;AX} &= \frac{1}{2\pi i} \int_0^\infty e^{ik(u(r)-u(r_c))} \frac{2i\varepsilon u'(r_c)}{(u-c)^2 + \varepsilon^2} \chi_1 \frac{\beta(r)}{\sqrt{r}} X(r, c, \varepsilon) dr_c \\ &\quad + \frac{1}{2\pi i} \int_0^\infty e^{ik(u(r)-u(r_c))} \frac{2i\varepsilon u'(r_c)}{(u-c)^2 + \varepsilon^2} \chi_2 \frac{\beta(r)}{\sqrt{r}} X(r, c, \varepsilon) dr_c \\ &= f_{1;AX1} + f_{1;AX2}. \end{aligned}$$

The contribution of $f_{1;AX1}$ is treated in the same way as $f_{1;X}$ (but with different integral operators in Theorem B.18) and is hence omitted. Next, we show that $f_{1;AX2}$ vanishes as $\varepsilon \rightarrow 0$. Indeed, expanding X as in $f_{1;AX1}$ gives

$$\begin{aligned} \sqrt{r} w_{f,\delta}^{-1} f_{1;AX2}(t, r) &= \frac{e^{iktu(r)}}{2\pi i} \int_0^\infty \frac{2i\varepsilon u'(r_c)}{(u-c)^2 + \varepsilon^2} \chi_2 \beta(r) e^{-iktu(r_c)} \\ &\quad \times \int_0^\infty \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} \frac{w_{F,\delta/4}(s)}{w_{f,\delta}(r)} B_{X\delta;\varepsilon}(r, s, c) \frac{F(s)}{w_{F,\delta/4}(s)} ds dr_c \\ &\quad + \frac{e^{iktu(r)}}{2\pi i} \int_0^\infty \frac{2i\varepsilon u'(r_c)}{(u-c)^2 + \varepsilon^2} \chi_2 e^{-iktc} \\ &\quad \times \int_0^\infty \int_0^\infty \frac{2i\varepsilon \beta(s_0)}{(u(s_0)-c)^2 + \varepsilon^2} \frac{\beta(r) w_{F,\delta/2}(s_0)}{w_{f,\delta}(r)} \\ &\quad \times B_{XS;\varepsilon}^{(1)}(r, s_0, c) B_{XS;\varepsilon}^{(2)}(s_0, s', c) \frac{w_{F,\delta/4}(s')}{w_{F,\delta/2}(s_0)} \frac{(u-c)}{(u-c)^2 + \varepsilon^2} \\ &\quad \times \frac{F(s')}{w_{F,\delta/4}(s')} ds' ds_0 dr_c \\ &\quad + \frac{e^{iktu(r)}}{2\pi i} \int_0^\infty \frac{2i\varepsilon u'(r_c)}{(u-c)^2 + \varepsilon^2} \chi_2 e^{-iktu(r_c)} \\ &\quad \times \int_0^\infty \int_0^\infty \frac{2i\varepsilon \beta(s_0)}{(u(s_0)-c)^2 + \varepsilon^2} \frac{\beta(r) w_{F,\delta/2}(s_0)}{w_{f,\delta}(r)} \\ &\quad \times B_{XG;\varepsilon}^{(1)}(r, s_0, c) B_{XG;\varepsilon}^{(2)}(s_0, s', c) \frac{w_{F,\delta/4}(s')}{w_{F,\delta/2}(s_0)} \frac{F_*(s')}{w_{F,\delta/4}(s')} ds' ds_0 dr_c. \end{aligned}$$

Therefore, Theorems B.18 and Theorem B.3 imply this term vanishes in the limit due to the support of χ_2 (note the weaker space $w_{f,\delta}$). Turn next to $f_{1;AY}$, which we similarly decompose via:

$$\begin{aligned} f_{1;AY} &= -\frac{2}{\pi i} \int_0^\infty \frac{i\varepsilon u'(r_c) e^{ik(u(r)-u(r_c))}}{(u-c)^2 + \varepsilon^2} \chi_1 \frac{\beta(r)}{\sqrt{r}} Y(r, c - i\varepsilon) dr_c \\ &\quad - \frac{2}{\pi i} \int_0^\infty \frac{i\varepsilon u'(r_c) e^{ik(u(r)-u(r_c))}}{(u-c)^2 + \varepsilon^2} \chi_2 \frac{\beta(r)}{\sqrt{r}} Y(r, c - i\varepsilon) dr_c \\ &= f_{1;AY1} + f_{1;AY2}. \end{aligned}$$

By Lemmas 6.10 and 6.14, the kernels satisfy the hypotheses necessary to apply Theorems B.18, B.3, and B.3. Therefore, we have again that $\lim_{\varepsilon \rightarrow 0} w_{f,\delta}^{-1} f_{1;AY2} = 0$ in L^2 , and that we may pass to the limit $\varepsilon \rightarrow 0$ in $f_{1;AY1}$ and deduce:

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \sqrt{r} w_{F,\delta}^{-1} f_{1;AY1} &= -p.v. \int_0^\infty \frac{w_{F,\delta/2}(s)}{w_{F,\delta}(r)} \beta(r) B_{YS}(r, s, u(r)) \frac{1}{u(s) - u(r)} \frac{F(s)}{w_{F,\delta/2}(s)} ds \\ &\quad - 2i\pi \frac{1}{w_{F,\delta}(r)} \beta(r) \frac{B_{Y\delta}(r, r, u(r))}{u'(r)} F(r) \\ &\quad - \int_0^\infty \frac{w_{F,\delta/2}(s)}{w_{F,\delta}(r)} \beta(r) B_{YG}(r, s, u(r)) \frac{F_*(s)}{w_{F,\delta/2}(s)} ds. \end{aligned}$$

Similarly, Theorem B.18 implies that we have the boundedness for all $\eta > 0$,

$$\left\| \sqrt{r} w_{F,\delta}^{-1} f_{1;AY1} \right\|_{L^2} \lesssim k^\eta \|F\|_{L^2_{F,\delta/2}} + \|F_*\|_{L^2_{F,\delta/2}} \lesssim k^\eta \|F\|_{L^2_{F,\delta/2}} + |k| \left\| \omega_{k,0}^n \right\|_{L^2_{F,\delta/2}}.$$

This completes the case $j = 0$.

Case $n \leq k - 1$ Next, turn to $(r\partial_r)^n$ for $n \leq k - 1$. From (2.28), denote the three contributions of f_1 as:

$$(r\partial_r)^n f_1^\varepsilon = (r\partial_r)^n F + f_{1;X}^\varepsilon + f_{1;A}^\varepsilon.$$

Consider first $f_{1;X}^\varepsilon$. After distributing the ∂_G derivatives there are many terms all of the general form

$$\int_0^\infty e^{itk(u(r)-u(r_c))} \frac{(u-c)u'(r_c)}{(u-c)^2 + \varepsilon^2} \frac{1}{w_{F,\delta}(r)} H(r, c, \varepsilon) \partial_G^j X(r, c, \varepsilon) dr_c,$$

for some weight H^ε satisfying the following for all $m \geq 0$ (depending on ε through χ_I):

$$\left| (r\partial_r)^m H^\varepsilon(r, c) \right| \lesssim_{n,j,m} \min(r^{2j}, r^{-7-2j}) (\chi_1 + \chi_{r \leq 1} \chi_{2r \approx r_c} + \chi_{r \approx 1} \chi_{2r < r_c}). \quad (6.30)$$

From Proposition 2.9 we have representations of the following form for a variety of complicated integral kernels:

$$\begin{aligned} &\int_0^\infty e^{itk(u(r)-u(r_c))} \frac{(u-c)u'(r_c)}{(u-c)^2 + \varepsilon^2} \frac{H^\varepsilon(r, c)}{w_{F,\delta}(r)} \partial_G^j X(r, c, \varepsilon) dr_c \\ &= \sum_{\ell=0}^j \frac{e^{itku(r)}}{2\pi i} \int_0^\infty \frac{(u-c)u'(r_c)}{(u-c)^2 + \varepsilon^2} H^\varepsilon(r, c) \frac{e^{-ikt c} w_{F,\delta/4+2\ell}(s)}{w_{F,\delta}(r)} \\ &\quad \times \int_0^\infty \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} B_{X\delta;j,\ell}(r, s, c) w_{X\delta 1;j,\ell}(s) \frac{\partial_G^\ell F(s)}{w_{F,\delta/4+2\ell}(s)} ds dr_c \\ &\quad + \sum_{\ell=0}^j \frac{e^{itku(r)}}{2\pi i} \int_0^\infty \frac{(u-c)u'(r_c)}{(u-c)^2 + \varepsilon^2} H^\varepsilon(r, c) \frac{e^{-ikt c}}{w_{F,\delta}(r)} \\ &\quad \times \int_0^\infty \int_0^\infty \frac{2i\varepsilon \beta(s_0)}{(u(s_0)-c)^2 + \varepsilon^2} B_{X\delta;j,\ell}^{(1)}(r, s_0, c) \end{aligned}$$

$$\begin{aligned}
& \times B_{X\delta;j,\ell}^{(2)}(s_0, s, c) w_{F,\delta/4+2\ell}(s) w_{X\delta 2;j,\ell}(s) \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} \frac{\partial_G^\ell F(s)}{w_{F,\delta/4+2\ell}(s)} ds ds_0 dr_c \\
& + \sum_{\ell=0}^j \frac{e^{itku(r)}}{2\pi i} \int_0^\infty \frac{(u-c)u'(r_c)}{(u-c)^2 + \varepsilon^2} H^\varepsilon(r, c) \frac{e^{-ikt c}}{w_{F,\delta}(r)} \\
& \times \int_0^\infty \int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u(s_0)-c)^2 + \varepsilon^2} B_{XS;j,\ell}^{(1)}(r, s_0, c) \\
& \times B_{XS;j,\ell}^{(2)}(s_0, s, c) w_{F,\delta/4+2\ell}(s) w_{XS;j,\ell}(s) \frac{(u-c)}{(u-c)^2 + \varepsilon^2} \frac{\partial_G^\ell F(s)}{w_{F,\delta/4+2\ell}(s)} ds ds_0 dr_c \\
& + \sum_{\ell=0}^j \frac{e^{itku(r)}}{2\pi i} \int_0^\infty \frac{(u-c)u'(r_c)}{(u-c)^2 + \varepsilon^2} H^\varepsilon(r, c) \frac{e^{-ikt c}}{w_{F,\delta}(r)} \\
& \times \int_0^\infty \int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u(s_0)-c)^2 + \varepsilon^2} B_{XG;j,\ell}^{(1)}(r, s_0, c) \\
& \times B_{XG;j,\ell}^{(2)}(s_0, s, c) w_{F,\delta/4+2\ell}(s) w_{XG;j,\ell}(s) \frac{\partial_G^\ell F_*(s)}{w_{F,\delta/4+2\ell}(s')} ds ds_0 dr_c \\
& + \text{Similar terms with different } B \text{ and } w.
\end{aligned}$$

Recall that in each term there holds for some ℓ' (different ℓ' in each term) $|w_{*,j,\ell}(s)| \lesssim \max(s^{-2\ell'}, s^{2\ell'})$. By Lemmas 6.10–6.13 (using also the recursion scheme laid out in Lemmas 6.4 and 6.5), for sufficiently small $\eta > 0$ and $\gamma > 0$, $B_{X\delta;j,\ell}$ is suitable $(2\ell'_1, \ell'_1 + \eta, \gamma)$ of type I, $B_{Xa;j,\ell}^{(1)}$ is suitable $(2\ell'_1, \ell'_1 + \eta, \gamma)$ of type I for some ℓ'_1 and $\gamma > 0$, whereas $B_{Xa;j,\ell}^{(2)}$ is suitable $(2\ell'_2, \ell'_2 + \eta, \gamma)$ of Type II for some ℓ'_2 (as above, each term may have a different ℓ'_j). In all terms there holds the inequality (where for the $B_{X\delta;j,\ell}$ terms we take $\ell'_2 = 0$):

$$\ell + \ell' + \ell'_1 + \ell'_2 \leq j.$$

Note that the total losses matches with (6.30). Therefore, for η chosen sufficiently small relative to δ , Theorems B.1 and B.18 (together with Theorems B.2 and B.3), we can pass to the limit $\varepsilon \rightarrow 0$ in the same way as we did for the case $j = 0$, giving also the L^2 bounds:

$$\begin{aligned}
& \left\| \lim_{\varepsilon \rightarrow 0} \int_0^\infty e^{itk(u(r)-u(r_c))} \frac{(u-c)u'(r_c)}{(u-c)^2 + \varepsilon^2} \frac{H(r, c, \varepsilon)}{w_{F,\delta}(r)} \partial_G^j X(r, c, \varepsilon) dr_c \right\|_{L^2} \\
& \lesssim \sum_{\ell=0}^j |k|^{2(j-\ell)+\eta} \left\| \partial_G^\ell F \right\|_{L^2_{F,\delta/4+2\ell}} + |k|^{2(j-\ell)+\eta} \left\| \partial_G^\ell F_* \right\|_{L^2_{F,\delta/4+2\ell}} \\
& \lesssim |k|^{2j+\eta+1} \left| \omega_{k,0}^{in} \right| + \sum_{\ell=0}^j |k|^{2(j-\ell)+\eta} \sum_{m=0}^\ell \left\| (r \partial_r)^m F \right\|_{L^2_{F,\delta/4}}.
\end{aligned}$$

Analogous to the case $j = 0$, the other contributions to f_1^ε are similarly; the details are omitted for brevity.

Case $n = k$ Recall that, as discussed in Remark 1.5, we do not really get useful information about the k -dependence in the case $n = k$. In the case $n = k$, the problematic terms in (2.28b) are those that contain $\partial_G^k X$ and $\partial_G^k A$; all other terms are treated as in the case $j \leq n - 1$. Hence, the terms we must consider are:

$$\tilde{f}_{1;A} := \frac{1}{2\pi i} \int_0^\infty \frac{2i\varepsilon u'(r_c) e^{ik(u(r)-u(r_c))}}{(u(r)-c)^2 + \varepsilon^2} (ru'(r))^k \chi_I(r_c) \frac{\beta(r)}{\sqrt{r}} \partial_G^k A(r, c, \varepsilon) dr_c \quad (6.31)$$

$$\tilde{f}_{1;X} := \frac{1}{2\pi i} \int_0^\infty e^{ik(u(r)-u(r_c))} \frac{(u(r)-c)u'(r_c)}{(u(r)-c)^2 + \varepsilon^2} (ru'(r))^k \chi_1(r, r_c) \frac{\beta(r)}{\sqrt{r}} \partial_G^k X(r, c, \varepsilon) dr_c. \quad (6.32)$$

The key difficulty is that we cannot use the iteration scheme to compute this derivative in the same manner as above. Consider $\tilde{f}_{1;X}$ and sub-divide based on the critical layer:

$$\begin{aligned} \tilde{f}_{1;X} &= \int_0^\infty \frac{e^{ik(u(r)-u(r_c))}}{2\pi i} \frac{(u(r)-c)u'(r_c)}{(u(r)-c)^2 + \varepsilon^2} (ru'(r))^k \chi_1 \frac{\beta(r)}{\sqrt{r}} (\chi_c(r, r_c) \\ &\quad + \chi_{\neq}(r, r_c)) \partial_G^k X(r, c, \varepsilon) dr_c \end{aligned} \quad (6.33)$$

$$= \tilde{f}_{1;X_c} + \tilde{f}_{1;X_{\neq}}. \quad (6.34)$$

On the support of $\tilde{f}_{1;X_{\neq}}$, we write

$$(ru'(r))^k \partial_G^k X = \frac{ru'(r)}{r_c u'(r_c)} (ru'(r))^{k-1} r_c \partial_{r_c} \partial_G^{k-1} X + (ru'(r))^{k-1} r \partial_r \partial_G^{k-1} X, \quad (6.35)$$

Next, we take ∂_r and ∂_{r_c} derivatives of the representation formula (2.37). Let us start with the easier ∂_r (note that $|r - r_c| \gtrsim r_c/k$ on the support of $f_{1;X_{\neq}}$). Due to Lemmas 6.10–6.13, these derivatives only land on Type I kernels:

$$\begin{aligned} \partial_r \partial_G^j X &= \sum_{\ell=0}^j \int_0^\infty \partial_r B_{X\delta;j,\ell}(r, s, c) \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w_{X\delta 1;j,\ell}(s) \partial_G^\ell F(s) ds \\ &\quad + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon \beta(s_0)}{(u-c)^2 + \varepsilon^2} \partial_r B_{X\delta;j,\ell}^{(1)}(r, s_0, c) B_{X\delta;j,\ell}^{(2)}(s_0, s, c) ds_0 \right) \\ &\quad \times \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w_{X\delta 2;j,\ell}(s) \partial_G^\ell F ds \\ &\quad + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon \beta(s_0)}{(u-c)^2 + \varepsilon^2} \partial_r B_{XS;j,\ell}^{(1)}(r, s_0, c) B_{XS;j,\ell}^{(2)}(s_0, s, c) ds_0 \right) \\ &\quad \times \frac{(u-c)}{(u-c)^2 + \varepsilon^2} w_{XS;j,\ell}(s) \partial_G^\ell F ds \\ &\quad + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon \beta(s_0)}{(u-c)^2 + \varepsilon^2} \partial_r B_{XG;j,\ell}^{(1)}(r, s_0, c) B_{XG;j,\ell}^{(2)}(s_0, s, c) ds_0 \right) \end{aligned}$$

$$\begin{aligned}
& \times w_{XG;j,\ell}(s) \partial_G^\ell F_* ds \\
& + \text{Similar terms with different } B, w.
\end{aligned} \tag{6.36}$$

By Definition 6.7, $k^{-1} \chi_{\neq}(r, r_c) r \partial_r B_{X\delta;j,\ell}(r, s, c)$ and $k^{-1} \chi_{\neq}(r, r_c) \partial_r B_{Xa;j,\ell}^{(1)}(r, s, c)$ satisfy the conditions necessary to apply Theorems B.1, B.18, B.2, and B.3 (note Remark B.4) with the same parameters as $\chi_{\neq}(r, r_c) B_{X\delta;j,\ell}$ and $\chi_{\neq}(r, r_c) \partial_r B_{Xa;j,\ell}^{(1)}$ respectively. Specifically, the $r \partial_r$ derivative does *not* incur a loss on the weights. Hence,

$$\int_0^\infty \frac{e^{itk(u(r)-u(r_c))}}{2\pi i} \frac{(u(r)-c)u'(r_c)}{(u(r)-c)^2 + \varepsilon^2} \chi_1 \frac{\beta(r)}{\sqrt{r}} \chi_{\neq}(r, r_c) (ru'(r))^{k-1} r \partial_r \partial_G^{k-1} X(r, c, \varepsilon) dr_c,$$

is treated via the same methods used to treat the case $j = k - 1$ above. Repetitive details are omitted for brevity.

Turn next to ∂_{r_c} derivatives, which are more technical. From (2.37), (still for $|r - r_c| \geq r_c/k$), we have

$$\begin{aligned}
\partial_{r_c} \partial_G^j X &= \sum_{\ell=0}^j \int_0^\infty \partial_{r_c} B_{X\delta;j,\ell}(r, s, c) \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w_{X\delta 1;j,\ell}(s) \partial_G^\ell F(s) ds \\
&+ \sum_{\ell=0}^j \int_0^\infty B_{X\delta;j,\ell}(r, s, c) \partial_{r_c} \left(\frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} \right) w_{X\delta 1;j,\ell}(s) \partial_G^\ell F(s) ds \\
&+ \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \partial_{r_c} \left(\frac{2i\varepsilon \beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) B_{X\delta;j,\ell}^{(1)}(r, s_0, c) B_{X\delta;j,\ell}^{(2)}(s_0, s, c) ds_0 \right) \\
&\times \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w_{X\delta 2;j,\ell}(s) \partial_G^\ell F ds \\
&+ \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon \beta(s_0)}{(u-c)^2 + \varepsilon^2} \partial_{r_c} B_{X\delta;j,\ell}^{(1)}(r, s_0, c) B_{X\delta;j,\ell}^{(2)}(s_0, s, c) ds_0 \right) \\
&\times \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w_{X\delta 2;j,\ell}(s) \partial_G^\ell F ds \\
&+ \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon \beta(s_0)}{(u-c)^2 + \varepsilon^2} B_{X\delta;j,\ell}^{(1)}(r, s_0, c) \partial_{r_c} B_{X\delta;j,\ell}^{(2)}(s_0, s, c) ds_0 \right) \\
&\times \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w_{X\delta 2;j,\ell}(s) \partial_G^\ell F ds \\
&+ \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon \beta(s_0)}{(u-c)^2 + \varepsilon^2} B_{X\delta;j,\ell}^{(1)}(r, s_0, c) B_{X\delta;j,\ell}^{(2)}(s_0, s, c) ds_0 \right) \\
&\times \partial_{r_c} \left(\frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} \right) w_{X\delta 2;j,\ell}(s) \partial_G^\ell F ds \\
&+ \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \partial_{r_c} \left(\frac{2i\varepsilon \beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) B_{X\delta;j,\ell}^{(1)}(r, s_0, c) B_{X\delta;j,\ell}^{(2)}(s_0, s, c) ds_0 \right)
\end{aligned}$$

$$\begin{aligned}
& \times \frac{(u-c)}{(u-c)^2 + \varepsilon^2} w_{XS;j,\ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \partial_{r_c} B_{XS;j,\ell}^{(1)}(r, s_0, c) B_{XS;j,\ell}^{(2)}(s_0, s, c) ds_0 \right) \\
& \times \frac{(u-c)}{(u-c)^2 + \varepsilon^2} w_{XS;j,\ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} B_{XS;j,\ell}^{(1)}(r, s_0, c) \partial_{r_c} B_{XS;j,\ell}^{(2)}(s_0, s, c) ds_0 \right) \\
& \times \frac{(u-c)}{(u-c)^2 + \varepsilon^2} w_{XS;j,\ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} B_{XS;j,\ell}^{(1)}(r, s_0, c) B_{XS;j,\ell}^{(2)}(s_0, s, c) ds_0 \right) \\
& \times \partial_{r_c} \left(\frac{(u-c)}{(u-c)^2 + \varepsilon^2} \right) w_{XS;j,\ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \partial_{r_c} \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) B_{XG;j,\ell}^{(1)}(r, s_0, c) B_{XG;j,\ell}^{(2)}(s_0, s, c) ds_0 \right) \\
& \times w_{XG;j,\ell}(s) \partial_G^\ell F_* ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \partial_{r_c} B_{XG;j,\ell}^{(1)}(r, s_0, c) B_{XG;j,\ell}^{(2)}(s_0, s, c) ds_0 \right) \\
& \times w_{XG;j,\ell}(s) \partial_G^\ell F_* ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} B_{XG;j,\ell}^{(1)}(r, s_0, c) \partial_{r_c} B_{XG;j,\ell}^{(2)}(s_0, s, c) ds_0 \right) \\
& \times w_{XG;j,\ell}(s) \partial_G^\ell F_* ds \\
& + \text{Similar terms with different } B, w \\
& = \sum_{n=0}^{12} T_n + \text{Similar terms with different } B, w. \tag{6.37}
\end{aligned}$$

Many of the terms permit a similar treatment, hence, let us only consider a few. As in (2.28) and Lemma 5.3 (and Sect. 6), we use $\frac{1}{u'(r_c)} \partial_{r_c} h(u(s) - u(r_c)) = -\frac{1}{u'(s)} \partial_s h(u(s) - u(r_c))$. Hence, we integrate by parts:

$$\begin{aligned}
r_c (T_0 + T_1) &= \sum_{\ell=0}^j \int_0^\infty r_c \partial_{r_c} (\chi_{\neq}(s, c) B_{X\delta;j,\ell}(r, s, c)) \\
&\quad \times \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w_{X\delta 1;j,\ell}(s) \partial_G^\ell F(s) ds \\
&\quad + \sum_{\ell=0}^j \int_0^\infty B_{X\delta;j,\ell}(r, s, c) \chi_{\neq}(s, c) r_c \partial_{r_c} \left(\frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} \right)
\end{aligned}$$

$$\begin{aligned}
& \times w_{X\delta 1; j, \ell}(s) \partial_G^\ell F(s) ds \\
& + \sum_{\ell=0}^j \int_0^\infty r_c u'(r_c) \partial_G^{(s)} \left(\chi_c B_{X\delta; j, \ell}(r, s, c) w_{X\delta 1; j, \ell}(s) \partial_G^\ell F(s) \right) \\
& \times \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} ds, \tag{6.38}
\end{aligned}$$

and note we are interested in passing to the limit in the singular integral

$$\int_0^\infty \frac{e^{itk(u(r)-u(r_c))}}{2\pi i} \frac{(u(r)-c)u'(r_c)}{(u(r)-c)^2 + \varepsilon^2} \chi_1 \frac{\beta \chi_{\neq}}{w_{F, \delta}(r) \sqrt{r}} \frac{ru'(r)}{r_c u'(r_c)} (ru'(r))^{k-1} r_c (T_0 + T_1) dr_c.$$

Due to the cutoffs in $\chi_c(s, c)$ and $\chi_{\neq}(s, c)$ in (6.38) and Definitions 6.7 and 6.8, we are still in a position to apply Theorems B.1 and B.18 and that the $r_c \partial_{r_c}$ derivatives have not changed the weights (as was the case for $r \partial_r$). Note that there is the leading ratio $(ru'(r))(r_c u'(r_c))^{-1}$. The numerator of this represents a gain in the weight in r and hence is what allows us to use the stronger weight w_F whereas the loss of $(r_c u'(r_c))^{-1}$ is balanced by the gains in \mathbb{K} . With these observations, we may hence apply Theorems B.1 and B.18 and pass to the limit $\varepsilon \rightarrow 0$, also obtaining the bounds (using Lemma 2.6):

$$\begin{aligned}
& \left\| \lim_{\varepsilon \rightarrow 0} \int_0^\infty \frac{e^{itk(u(r)-u(r_c))}}{2\pi i} \frac{(u(r)-c)u'(r_c)}{(u(r)-c)^2 + \varepsilon^2} \chi_1 \frac{\beta(r)}{w_{F, \delta}(r) \sqrt{r}} \right. \\
& \quad \left. \chi_{\neq}(r, r_c) \frac{ru'(r)}{r_c u'(r_c)} (ru'(r))^{k-1} r_c (T_0 + T_1) dr_c \right\|_{L^2} \\
& \lesssim \sum_{\ell=0}^{k-1} |k|^{2(j-\ell)+\eta+3} \left\| \partial_G^\ell F \right\|_{L^2_{F, \delta/4+2\ell}} + |k|^{2(j-\ell)+\eta+3} \left\| \partial_G^\ell F_* \right\|_{L^2_{F, \delta/4+2\ell}} \\
& \lesssim_k \left| \omega_{k,0}^n \right| + \sum_{\ell=0}^{k-1} \left\| (r \partial_r)^m F \right\|_{L^2_{F, \delta/4}};
\end{aligned}$$

(recall Remark 1.5). For the compound terms in (6.37) the picture is a little more complicated as these involve the triple derivatives appearing in Definitions 6.7 and 6.8:

$$\begin{aligned}
\frac{1}{u'(r_c)} \sum_{n=2}^5 T_n &= \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{\chi_{\neq}(s_0, c)}{u'(r_c)} \partial_{r_c} \left(\frac{2i\varepsilon \beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \quad \times B_{X\delta; j, \ell}^{(1)}(r, s_0, c) B_{X\delta; j, \ell}^{(2)}(s_0, s, c) ds_0 \Big) \\
& \quad \times \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w_{X\delta 2; j, \ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon \beta(s_0)}{(u-c)^2 + \varepsilon^2} B_{X\delta; j, \ell}^{(1)}(r, s_0, c) B_{X\delta; j, \ell}^{(2)}(s_0, s, c) ds_0 \right) \\
& \quad \times \chi_{\neq}(s, c) \frac{1}{u'(r_c)} \partial_{r_c} \left(\frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} \right) w_{X\delta 2; j, \ell}(s) \partial_G^\ell F ds
\end{aligned}$$

$$\begin{aligned}
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \frac{1}{u'(r_c)} \right. \\
& \times \partial_{r_c} \left(\chi_{\neq}(s_0, c) B_{X\delta; j, \ell}^{(1)}(r, s_0, c) \chi_{\neq}(s, c) B_{X\delta; j, \ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \times \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w_{X\delta 2; j, \ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right. \\
& \times \partial_G^{(s)} \left(\chi_{\neq}(s_0, c) B_{X\delta; j, \ell}^{(1)}(r, s_0, c) \chi_c(s, c) B_{X\delta; j, \ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \times \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w_{X\delta 2; j, \ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right. \\
& \times \left(B_{X\delta; j, \ell}^{(1)}(r, s_0, c) \chi_c(s, c) B_{X\delta; j, \ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \times \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} \frac{1}{u'(s)} \partial_s \left(w_{X\delta 2; j, \ell}(s) \partial_G^\ell F \right) ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \partial_G^{(s, s_0)} \left(\chi_c(s_0, c) B_{X\delta; j, \ell}^{(1)}(r, s_0, c) \chi_c(s, c) \right. \right. \\
& \times \left. \left. B_{X\delta; j, \ell}^{(2)}(s_0, s, c) \right) ds_0 \right) \\
& \times \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w_{X\delta 2; j, \ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right. \\
& \times \partial_G^{(s_0)} \left(\chi_c(s_0, c) B_{X\delta; j, \ell}^{(1)}(r, s_0, c) \chi_{\neq}(s, c) B_{X\delta; j, \ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \times \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w_{X\delta 2; j, \ell}(s) \partial_G^\ell F ds \tag{6.39}
\end{aligned}$$

It is crucial to note the very specific structure in (6.39): whenever $s \approx r_c$ and/or $s_0 \approx r_c$ the derivatives landing on the kernels are either $\partial_G^{(s)}$ or $\partial_G^{(s_0)}$ or ∂_G^{s, s_0} so that one never evaluates ∂_{r_c} (or ∂_r) of a kernel near the critical layer without the matching s or s_0 derivatives. A similar structure is seen also in the $\sum_{j=6}^9 T_j$ terms which are omitted for brevity. The last three terms instead have:

$$\begin{aligned}
\frac{1}{u'(r_c)} \sum_{j=10}^{12} T_j &= \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \frac{1}{u'(r_c)} \partial_{r_c} \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
&\times \chi_{\neq}(s_0, c) B_{XG; j, \ell}^{(1)}(r, s_0, c) B_{XG; j, \ell}^{(2)}(s_0, s, c) ds_0 \Big)
\end{aligned}$$

$$\begin{aligned}
& \times w_{XG;j,\ell}(s) \partial_G^\ell F_* ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \times \frac{1}{r_c} \partial_{r_c} \left(\chi_{\neq}(s_0, c) B_{XG;j,\ell}^{(1)}(r, s_0, c) B_{XG;j,\ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \times w_{XG;j,\ell}(s) \partial_G^\ell F_* ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \times \partial_G^{(s_0)} \left(\chi_c(s_0, c) B_{XG;j,\ell}^{(1)}(r, s_0, c) B_{XG;j,\ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \times w_{XG;j,\ell}(s) \partial_G^\ell F_* ds. \tag{6.40}
\end{aligned}$$

Notice that near $s_0 \approx r_c$ we are still using $\partial_G^{(s_0)}$ derivatives, despite that s can be close to the critical layer as well. Hence, the derivatives are not quite the correct form for directly using that $B_{XG;j,\ell}^{(2)}$ is suitable $(2\ell_2'', \ell_2'' + \eta, \gamma)$ of type II (for some ℓ_2'' and γ and all $\eta > 0$). However,

$$\partial_G^{(s_0)} B_{XG;j,\ell}^{(2)}(s_0, s, c) = \partial_G^{(s,s_0)} B_{XG;j,\ell}^{(2)}(s_0, s, c) - \frac{1}{u'(s)} \partial_s B_{XG;j,\ell}^{(2)}(s_0, s, c).$$

Note that the former is bounded near the critical layer whereas the latter is logarithmically singular there (see Definition 6.8). However, since there are no singular integral operators or approximately δ -functions in s in these terms, it is straightforward to verify that we may still apply Theorem B.18.

Finally, putting together (6.34) and (6.35) with the associated decompositions of (6.36), (6.37), (6.38), (6.39) (and the analogous omitted terms), and (6.40) with Definitions 6.7, 6.8 and Theorems B.1, B.18, B.2, B.3 (and Remark B.4), we may pass to the limit in $\varepsilon \rightarrow 0$ as we did in the $n = k - 1$ case. This gives us a (very complicated) representation formula for $f_{1;X\neq}$, and, in particular, the bound

$$\left\| \lim_{\varepsilon \rightarrow 0} \sqrt{r} w_{F,\delta} f_{1;X\neq} \right\|_{L^2} \lesssim_{k,\delta,\alpha} \left| \omega_{k,0}^{in} \right| + \sum_{\ell=0}^k \left\| (r \partial_r)^\ell F \right\|_{L_{F,\delta/4}^2}.$$

Next, we consider $\tilde{f}_{1;Xc}$. We directly take a $\partial_G = \partial_G^{(r)}$ derivative of (2.37) and as above, apply the usual integration by parts when $s_0 \approx r_c$ and/or $s \approx r_c$. This yields (on the support of the integrand)

$$\begin{aligned}
\partial_G \partial_G^j X &= \sum_{\ell=0}^j \int_0^\infty B_{X\delta;j,\ell}(r, s, c) \chi_{\neq}(s, c) \frac{1}{u'(r_c)} \\
&\times \partial_{r_c} \left(\frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} \right) w_{X\delta 1;j,\ell}(s) \partial_G^\ell F(s) ds
\end{aligned}$$

$$\begin{aligned}
& + \sum_{\ell=0}^j \int_0^\infty \partial_G^{(r)} (B_{X\delta;j,\ell}(r, s, c) \chi_{\neq}(s, c)) \\
& \times \left(\frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} \right) w_{X\delta1;j,\ell}(s) \partial_G^\ell F(s) ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} \right) \\
& \times \partial_G^{(r,s)} \left(B_{X\delta;j,\ell}(r, s, c) \chi_c(s, c) w_{X\delta1;j,\ell}(s) \partial_G^\ell F(s) \right) ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \chi_{\neq}(s_0, c) \frac{1}{u'(r_c)} \partial_{r_c} \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \times B_{X\delta;j,\ell}^{(1)}(r, s_0, c) B_{X\delta;j,\ell}^{(2)}(s_0, s, c) ds_0 \Big) \\
& \times \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w_{X\delta2;j,\ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \times \partial_G^{(r)} \left(\chi_{\neq}(s_0, c) B_{X\delta;j,\ell}^{(1)}(r, s_0, c) \chi_{\neq}(s, c) B_{X\delta;j,\ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \times \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w_{X\delta2;j,\ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \times \partial_G^{(r,s)} \left(\chi_{\neq}(s_0, c) B_{X\delta;j,\ell}^{(1)}(r, s_0, c) \chi_c(s, c) B_{X\delta;j,\ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \times \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w_{X\delta2;j,\ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \times \partial_G^{(r,s_0)} \left(\chi_c(s_0, c) B_{X\delta;j,\ell}^{(1)}(r, s_0, c) \chi_{\neq}(s, c) B_{X\delta;j,\ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \times \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w_{X\delta2;j,\ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \times \partial_G^{(r,s_0,s)} \left(\chi_c(s_0, c) B_{X\delta;j,\ell}^{(1)}(r, s_0, c) \chi_c(s, c) B_{X\delta;j,\ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \times \frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} w_{X\delta2;j,\ell}(s) \partial_G^\ell F ds
\end{aligned}$$

$$\begin{aligned}
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \quad \times \left(B_{X\delta;j,\ell}^{(1)}(r, s_0, c) \chi_{\neq}(s, c) B_{X\delta;j,\ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \quad \times \frac{1}{u'(r_c)} \partial_{r_c} \left(\frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} \right) w_{X\delta 2;j,\ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \quad \times \left(B_{X\delta;j,\ell}^{(1)}(r, s_0, c) \chi_c(s, c) B_{X\delta;j,\ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \quad \times \left(\frac{2i\varepsilon}{(u-c)^2 + \varepsilon^2} \right) \frac{1}{u'(s)} \partial_s \left(w_{X\delta 2;j,\ell}(s) \partial_G^\ell F \right) ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \chi_{\neq}(s_0, c) \frac{1}{u'(r_c)} \right. \\
& \quad \times \partial_{r_c} \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) B_{XS;j,\ell}^{(1)}(r, s_0, c) B_{XS;j,\ell}^{(2)}(s_0, s, c) ds_0 \Big) \\
& \quad \times \frac{(u-c)}{(u-c)^2 + \varepsilon^2} w_{XS;j,\ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \quad \times \partial_G^{(r)} \left(\chi_{\neq}(s_0, c) B_{XS;j,\ell}^{(1)}(r, s_0, c) \chi_{\neq}(s, c) B_{XS;j,\ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \quad \times \frac{(u-c)}{(u-c)^2 + \varepsilon^2} w_{XS;j,\ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \quad \times \partial_G^{(r,s)} \left(\chi_{\neq}(s_0, c) B_{XS;j,\ell}^{(1)}(r, s_0, c) \chi_c(s, c) B_{XS;j,\ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \quad \times \frac{(u-c)}{(u-c)^2 + \varepsilon^2} w_{XS;j,\ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \quad \times \partial_G^{(r,s_0)} \left(\chi_c(s_0, c) B_{XS;j,\ell}^{(1)}(r, s_0, c) \chi_{\neq}(s, c) B_{XS;j,\ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \quad \times \frac{(u-c)}{(u-c)^2 + \varepsilon^2} w_{XS;j,\ell}(s) \partial_G^\ell F ds
\end{aligned}$$

$$\begin{aligned}
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \times \partial_G^{(r, s_0, s)} \left(\chi_c(s_0, c) B_{XS; j, \ell}^{(1)}(r, s_0, c) \chi_c(s, c) B_{XS; j, \ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \times \frac{(u-c)}{(u-c)^2 + \varepsilon^2} w_{XS; j, \ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \times \left(B_{XS; j, \ell}^{(1)}(r, s_0, c) \chi_{\neq}(s, c) B_{XS; j, \ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \times \frac{1}{u'(r_c)} \partial_{r_c} \left(\frac{u-c}{(u-c)^2 + \varepsilon^2} \right) w_{XS; j, \ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \times \left(B_{XS; j, \ell}^{(1)}(r, s_0, c) \chi_c(s, c) B_{XS; j, \ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \times \left(\frac{(u-c)}{(u-c)^2 + \varepsilon^2} \right) \frac{1}{u'(s)} \partial_s \left(w_{XS; j, \ell}(s) \partial_G^\ell F \right) ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \chi_{\neq}(s_0, c) \frac{1}{u'(r_c)} \right. \\
& \times \partial_{r_c} \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) B_{XG; j, \ell}^{(1)}(r, s_0, c) B_{XG; j, \ell}^{(2)}(s_0, s, c) ds_0 \Big) \\
& \times w_{XG; j, \ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \times \partial_G^{(r)} \left(\chi_{\neq}(s_0, c) B_{XG; j, \ell}^{(1)}(r, s_0, c) B_{XG; j, \ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \times w_{XG; j, \ell}(s) \partial_G^\ell F ds \\
& + \sum_{\ell=0}^j \int_0^\infty \left(\int_0^\infty \left(\frac{2i\varepsilon\beta(s_0)}{(u-c)^2 + \varepsilon^2} \right) \right. \\
& \times \partial_G^{(r, s_0)} \left(\chi_c(s_0, c) B_{XG; j, \ell}^{(1)}(r, s_0, c) B_{XG; j, \ell}^{(2)}(s_0, s, c) \right) ds_0 \Big) \\
& \times w_{XG; j, \ell}(s) \partial_G^\ell F ds \\
& + \text{Similar terms with different } B, w.
\end{aligned}$$

We see that, although slightly more technical, the overall structure of which derivatives appear in what contributions of the integrals, is very similar to the case of $\partial_{r_c}^j X$. Hence, the arguments used above apply with no major variations and we may pass to

the limit and deduce the estimate:

$$\left\| \lim_{\varepsilon \rightarrow 0} \sqrt{r} w_{F,\delta} f_{1;Xc} \right\|_{L^2} \lesssim_{k,\delta,\alpha} |\omega_{k,0}^{in}| + \sum_{\ell=0}^k \left\| (r \partial_r)^\ell F \right\|_{L_{F,\delta/4}^2};$$

we omit the repetitive details for brevity. This completes the treatment of $\tilde{f}_{1;X}$.

Turn next to $\tilde{f}_{1;A}$. As in the cases $n \leq k-1$, we write

$$\begin{aligned} \tilde{f}_{1;A} &= \frac{1}{2\pi i} \int_0^\infty \frac{2i\varepsilon u'(r_c) e^{itk(u(r)-u(r_c))}}{(u(r)-c)^2 + \varepsilon^2} (ru'(r))^k \chi_I(r_c) \frac{\beta(r)}{\sqrt{r}} \partial_G^k (X(r, c, \varepsilon) \\ &\quad + 2Y(r, c - i\varepsilon)) dr_c \\ &= \tilde{f}_{1;AX} + \tilde{f}_{1;AY}. \end{aligned}$$

The term $\tilde{f}_{1;AX}$ is treated in essentially the same manner as $\tilde{f}_{1;X}$ and is hence omitted for the sake of brevity. Similarly, we see that the treatment of $\tilde{f}_{1;AY}$ is made via a small variant of the method used to treat the first term in (2.37). Hence, this is also omitted for the brevity. This completes the proof of Proposition 2.13. \square

Proof of Proposition 2.14 Recall from (2.29),

$$\begin{aligned} w_{f,\delta}^{-1} (r \partial_r)^n f_2^\varepsilon &= -\frac{1}{2\pi k t w_{f,\delta}(r)} \int_0^\infty e^{itk(u(r)-u(r_c))} \partial_{r_c} \left(\frac{(u(r)-c)}{(u(r)-c)^2 + \varepsilon^2} \right) \\ &\quad \times (ru'(r) \partial_G)^n \left(\chi_2(r, r_c) \frac{\beta(r)}{\sqrt{r}} X(r, c, \varepsilon) \right) dr_c \\ &\quad - \frac{1}{2\pi k t w_{f,\delta}(r)} \int_0^\infty e^{itk(u(r)-u(r_c))} \left(\frac{(u(r)-c)}{(u(r)-c)^2 + \varepsilon^2} \right) \\ &\quad \times \partial_{r_c} (ru'(r) \partial_G)^n \left(\chi_2(r, r_c) \frac{\beta(r)}{\sqrt{r}} X(r, c, \varepsilon) \right) dr_c \\ &= f_{2,a}^\varepsilon + f_{2,b}^\varepsilon. \end{aligned} \tag{6.41}$$

Due to the presence of χ_2 , the support of these integrands satisfies $r < \min(r_c/2, 1)$. In particular, the integral in r_c is not converging to a singular integral as $\varepsilon \rightarrow 0$. Analogous to the treatment of f_1^ε in the proof of Proposition 2.13 above, we may write $f_{2,a}^\varepsilon$ as the sum of terms of the general form

$$-\frac{1}{2\pi k t} \int_0^\infty e^{itk(u(r)-u(r_c))} \frac{1}{w_{f,\delta}(r)} H^\varepsilon(r, c) \partial_G^j X(r, c; \varepsilon) dr_c,$$

for weights H satisfying,

$$|H^\varepsilon(r, c)| \lesssim \mathbf{1}_{r < r_c/2} \mathbf{1}_{r \leq 1} r^{-1/2} \min(r^{2j}, r^{-2j-7}) \max(r_c^{-3}, 1).$$

From Proposition 2.37 we have an expansion as in the proof Proposition 2.13 above which by lemmas 6.10–6.14 (using also the recursion scheme laid out in Lemmas 6.4 and 6.5) satisfies similar properties. The main difference here is that we are using a weaker weight ($w_{f,\delta}$ instead of $w_{F,\delta}$) and we have lost an additional r_c^{-2} from the integration by parts in r_c . The loss in r_c is balanced by the gains in \mathbb{K} ; these were used to recover the strong weight on f_1 whereas here the gains are used to gain the r_c^{-2} necessary to allow us to integrate by parts in r_c to deduce. After this adjustment, the proof of convergence follows from Theorems B.1 and B.18 (together with Theorems B.2 and B.3) as in the proof of Proposition 2.13 and is hence omitted for brevity.

Consider next $f_{2,b}^\varepsilon$ in (6.41). The terms where ∂_{r_c} lands on χ_2 are treated as in $f_{2,a}^\varepsilon$. For terms containing $\partial_{r_c} \partial_G^j X$, we apply the same methods as in Proposition 2.13 when ∂_{r_c} derivatives were computed away from the critical layer as in (6.37). Indeed, due to χ_2 , the entire integrand in $f_{2,b}^\varepsilon$ is supported away from the critical layer and hence this is the only case we need to consider here. Hence, combining ideas in Proposition 2.13 with those used to treat $f_{2,a}^\varepsilon$ completes the desired bounds; we omit the details for brevity as they are repetitive. This completes the proof of Proposition 2.14. \square

A Preliminary Technical Lemmas

We record a few minor technical observations used several times in the proof.

Lemma A.1 *Let $r, r_c \in (0, \infty)$ and $k \geq 2$ such that $|r - r_c| \leq r_c/k$. Then*

- $\frac{k-1}{k}r_c \leq r \leq \frac{k+1}{k}r_c$;
- for all $a \in \mathbb{R}$, there exists constants c_a, C_a (depending only on a) such that $c_a r_c^{ak} \leq r^{ak} \leq C_a r_c^{ak}$.

The next lemma contains a few useful inequalities regarding u . The proof follows immediately from Lemma 2.1.

Lemma A.2 *There holds*

$$\begin{aligned} \left| \frac{\chi_{\neq} u'(r)}{u(r) - c} \right| &\lesssim \mathbf{1}_{r_c \leq 1} \left(\mathbf{1}_{r \leq 2} \min \left(\frac{kr}{r_c^2}, \frac{kr}{r^2} \right) + \frac{1}{r^3} \mathbf{1}_{r \geq 2} \right) \\ &\quad + \mathbf{1}_{r_c \geq 1} \left(\mathbf{1}_{r \leq 2r} + \mathbf{1}_{r \geq 2} \min \left(\frac{kr_c^2}{r^3}, \frac{k}{r} \right) \right), \end{aligned} \quad (\text{A.1})$$

and for $z \in I_\alpha$ there holds

$$\left| \frac{\varepsilon u'}{(u - c)^2 + \varepsilon^2} \right| \chi_{\neq} \lesssim \varepsilon^{\frac{2\alpha}{2+\alpha}} \frac{|u'|}{|u - c|} \chi_{\neq}, \quad (\text{A.2})$$

$$\left| \frac{(u - c)u'}{(u - c)^2 + \varepsilon^2} - \frac{u'}{u - c} \right| \chi_{\neq} \lesssim \varepsilon^{\frac{2\alpha}{2+\alpha}} \frac{|u'|}{|u - c|} \chi_{\neq}. \quad (\text{A.3})$$

Lemma A.3 Let χ_c be defined as in (4.3). Then, the following holds independent of c :

$$\int_0^\infty \frac{(u(s) - c)u'(s)}{(u(s) - c)^2 + \varepsilon^2} \chi_c(s, c) ds \lesssim 1, \quad (\text{A.4})$$

and for $|r - r_c| \lesssim r_c/k$ there holds

$$\begin{aligned} \int_0^r \frac{(u(s) - c)u'(s)}{(u(s) - c)^2 + \varepsilon^2} \chi_c(s, c) ds &\lesssim 1 + \left| \log \frac{k|r - r_c|}{r_c} \right| \\ \int_r^\infty \frac{(u(s) - c)u'(s)}{(u(s) - c)^2 + \varepsilon^2} \chi_c(s, c) ds &\lesssim 1 + \left| \log \frac{k|r - r_c|}{r_c} \right|. \end{aligned}$$

Proof of Lemma A.3 Consider just (A.4); the other estimates follow similarly (and are slightly easier). Integration by parts yields the following for any r :

$$\begin{aligned} \int_0^\infty \frac{(u - c)u'(s)}{(u(s) - c)^2 + \varepsilon^2} \chi_c(s, c) ds &= - \int_0^\infty \log \left((u(s) - c)^2 + \varepsilon^2 \right) \partial_s \chi_c(s, c) ds \\ &= - \int_0^\infty \left(\log \left((u(s) - c)^2 + \varepsilon^2 \right) \right. \\ &\quad \left. - \log \left((u(r) - c)^2 + \varepsilon^2 \right) \right) \partial_s \chi_c(s, c) ds \\ &= - \int_0^\infty \left(\log \frac{(u(s) - c)^2 + \varepsilon^2}{(u(r) - c)^2 + \varepsilon^2} \right) \partial_s \chi_c(s, c) ds. \end{aligned}$$

Choose $r = (1 + \frac{1}{k})r_c$ and hence, on the support of the integrand,

$$\left| \log \frac{(u(s) - c)^2 + \varepsilon^2}{(u(r) - c)^2 + \varepsilon^2} \right| \lesssim \left| \frac{(u(s) - c)^2 - (u(r) - c)^2}{(u(r) - c)^2 + \varepsilon^2} \right| \lesssim 1.$$

and hence (A.4) follows. \square

Lemma A.4 For all $0 < \gamma \leq 1$, for all η sufficiently small (depending on γ and α) and all $z \in I_\alpha$, there holds

$$\int_0^\infty \frac{\varepsilon |u'(r)|}{(u - c)^2 + \varepsilon^2} \left(\frac{|r - r_c|}{r_c} \right)^\gamma \chi_c dr \lesssim \varepsilon^\eta. \quad (\text{A.5})$$

Let $G^\varepsilon(r, c)$ be defined for $z = c \pm i\varepsilon \in I_\alpha$ and $|r - r_c| \lesssim r_c$ and (over the same range of r, r_c, ε) satisfy the following estimates (uniformly in ε) for some exponents $\gamma_i \in (0, 1]$:

$$|G^\varepsilon(r, c) - G^\varepsilon(r_c, c)| \lesssim \left(\frac{|r - r_c|}{r_c} \right)^{\gamma_0} \quad (\text{A.6})$$

$$|G^\varepsilon(r, c) - G^0(r, c)| \lesssim \varepsilon^{\gamma_1}. \quad (\text{A.7})$$

Then for all η sufficiently small (depending on α, γ_i),

$$\left| \int_0^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_c G^\varepsilon(r, c) ds - p.v. \int_0^\infty \frac{u'}{u-c} \chi_c G^0(r, c) ds \right| \lesssim \varepsilon^\eta. \quad (\text{A.8})$$

Proof of Lemma A.4 Consider first (A.5). For all $0 < p < \gamma$, by Lemmas A.1 and 2.1,

$$\begin{aligned} \int_0^\infty \frac{\varepsilon |u'(r)|}{(u-c)^2 + \varepsilon^2} \left(\frac{|r-r_c|}{r_c} \right)^\gamma \chi_c dr &\lesssim \varepsilon^p \frac{k^p}{|u'(r_c)|^p} \int_0^\infty \frac{1}{|r-r_c|^{1+p-\gamma}} dr \\ &\lesssim \varepsilon^p k^p \max\left(\frac{1}{r_c^{2p}}, r_c^{2p}\right), \end{aligned}$$

and hence (A.5) follows from the definition of I_α . Next, consider (A.8). We have

$$\begin{aligned} &\left| \int_0^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_c G^\varepsilon(r, c) ds - p.v. \int_0^\infty \frac{u'}{u-c} \chi_c G^0(r, c) ds \right| \\ &\leq \left| \int_0^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_c \left(G^\varepsilon(r, c) - G^0(r, c) \right) ds \right| \\ &\quad + \left| p.v. \int_0^\infty \left(\frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} - \frac{1}{u-c} \right) \chi_c G^0(r, c) ds \right| \\ &= T_1 + T_2. \end{aligned}$$

By the assumptions on G^ε , we have

$$\begin{aligned} &\chi_c \left| G^\varepsilon(r, c) - G^\varepsilon(r_c, c) - G^0(r, c) + G^0(r_c, c) \right| \\ &\leq \chi_c \left(\left| G^\varepsilon(r, c) - G^\varepsilon(r_c, c) \right| + \left| G^0(r, c) - G^0(r_c, c) \right| \right)^\gamma \\ &\quad \times \left(\left| G^\varepsilon(r, c) - G^0(r, c) \right| + \left| G^\varepsilon(r_c, c) - G^0(r_c, c) \right| \right)^{1-\gamma} \\ &\lesssim \varepsilon^{\gamma_1(1-\gamma)} \left(\frac{|r-r_c|}{r_c} \right)^{\gamma\gamma_0}. \end{aligned}$$

Therefore, setting $\eta = \gamma_1(1-\gamma)$ and using Lemma A.3 implies

$$\begin{aligned} T_1 &\leq \left| G^\varepsilon(r_c, c) - G^0(r_c, c) \right| \left| \int_0^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_c dr \right| \\ &\quad + \left| \int_0^\infty \frac{(u-c)u'}{(u-c)^2 + \varepsilon^2} \chi_c \left(G^\varepsilon(r, c) - G^\varepsilon(r_c, c) - G^0(r, c) + G^0(r_c, c) \right) dr \right| \\ &\lesssim \varepsilon^\eta. \end{aligned}$$

The proof of T_2 follows from noting:

$$T_2 \leq \left| G^0(r_c, c) p.v. \int_0^\infty \left(\frac{\varepsilon^2 u'}{(u-c)((u-c)^2 + \varepsilon^2)} \right) \chi_c ds \right|$$

$$+ \left| p.v. \int_0^\infty \left(\frac{\varepsilon^2 u'}{(u-c)((u-c)^2 + \varepsilon^2)} \right) \chi_c \left(G^0(r, c) - G^0(r_c, c) \right) ds \right|.$$

The latter integral is treated by an easy variant of the treatment of T_1 and is hence omitted. The former integral is estimated via

$$\begin{aligned} & \left| p.v. \int_0^\infty \left(\frac{\varepsilon^2 u'}{(u-c)((u-c)^2 + \varepsilon^2)} \right) \chi_c ds \right| \\ &= \left| \int_0^\infty \chi_c \frac{1}{2} \partial_s \left(\log((u-c)^2 + \varepsilon^2) - \log(u-c)^2 \right) ds \right| \\ &\lesssim \int_0^\infty \left| \frac{(u-c)^2 + \varepsilon^2}{(u-c)^2} - 1 \right| |\chi'_c| ds \\ &\lesssim \varepsilon^2 k^2 \max(r_c^{-4}, r_c^4), \end{aligned}$$

which completes the proof by the definition of I_α . \square

B Boundedness and Convergence of Integral Operators

B.1 Two Singular Integrals and One Delta Distribution

Our goal is to prove the convergence as $\varepsilon \rightarrow 0$ of the following “model operator”:

$$\begin{aligned} L_\varepsilon[f](r) &= \int_0^\infty \int_0^\infty \int_0^\infty \frac{(u(r) - u(r_c))u'(r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} \frac{(u(s) - u(r_c))u'(s)}{(u(s) - u(r_c))^2 + \varepsilon^2} \frac{\varepsilon u'(s_0)}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \\ &\quad \times \mathfrak{B}_{\varepsilon,1}(r, s_0, r_c) \mathfrak{B}_{\varepsilon,2}(s_0, s, r_c) f(s) ds ds_0 dr_c \end{aligned} \quad (\text{B.1})$$

in $L^2(dr)$ as $\varepsilon \rightarrow 0$, under certain assumptions on the weights $B_{\varepsilon,1}$ and $B_{\varepsilon,2}$.

Theorem B.1 *Let $\delta \in (0, \frac{1}{2})$ and assume that for some $\gamma \in (0, \frac{\delta}{4})$ we have that the functions $\mathfrak{B}_{\varepsilon,1}$ and $\mathfrak{B}_{\varepsilon,2}$ obey the conditions (B.8), (B.10), and either (B.37)–(B.41) or (B.42)–(B.43). Additionally, assume that there exists $\zeta \in (0, \gamma)$ such that conditions (B.57)–(B.58) hold, for some limiting weights $\mathfrak{B}_{0,1}$ and $\mathfrak{B}_{0,2}$. Then, if $f \in L^2$, we have that the operator $L_\varepsilon[f]$, defined in (B.1), converges as $\varepsilon \rightarrow 0$, in L^2 to the operator $L_0[f]$, defined by duality via*

$$\begin{aligned} \langle L_0[f], \varphi \rangle &= -\pi \int_0^\infty \left(p.v. \int_0^\infty \frac{u'(r)}{u(r) - u(r_c)} \frac{u'(r_c) \mathfrak{B}_{0,1}(r, r_c, r_c)}{u'(r)} \varphi(r) dr \right) \\ &\quad \times \left(p.v. \int_0^\infty \frac{u'(s)}{u(s) - u(r_c)} \frac{\mathfrak{B}_{0,2}(r_c, s, r_c)}{\langle s \rangle^\delta} (\langle s \rangle^\delta f(s)) ds \right) dr_c \end{aligned} \quad (\text{B.2})$$

and the operator L_0 is bounded from L^2 to L^2 , with norm less than k^ζ .

The first standard example of pairs of weights $\mathfrak{B}_{\varepsilon,1}$ and $\mathfrak{B}_{\varepsilon,2}$ which obey the conditions of Theorem B.1 are:

Theorem B.2 Let $0 \leq j \leq k-1$, and $0 \leq \ell, \ell_1, \ell_2$ be such that $\ell + \ell_1 + \ell_2 \leq j$. Let $0 < \zeta \leq \frac{\delta}{6}$, and $0 < \eta \leq \frac{\zeta}{4}$. Consider the weights

$$\mathfrak{B}_{\varepsilon,1}(r, s_0, r_c) = \chi_1(r, r_c) \frac{\beta(r) \min(r^2, r^{-2})^j}{w_{F,\delta}(r) u'(s_0)} B_{\ell,\varepsilon}^{(1)}(r, s_0, r_c) \quad (\text{B.3})$$

$$\mathfrak{B}_{\varepsilon,2}(s_0, s, r_c) = \frac{\beta(s_0) w_{F,\frac{\delta}{4}+2\ell}(s)}{u'(s)} B_{\ell,\varepsilon}^{(2)}(s_0, s, r_c) \quad (\text{B.4})$$

where $B_{\ell,\varepsilon}^{(1)}$ is a suitable $(2\ell_1, \ell_1 + \eta/2)$ kernel of type I or II, and $B_{\ell,\varepsilon}^{(2)}$ is a suitable $(2\ell_2, \ell_2 + \eta/2)$ kernel of type I or II. Then the conditions of Theorem B.1 are satisfied for the weights (B.3)–(B.4). The corresponding operator L_ε defined in (B.1) converges to the corresponding operator L_0 defined in (B.2), which in this case becomes

$$\begin{aligned} \langle L_0[f], \varphi \rangle = & -\pi \int_0^\infty \left(p.v. \int_0^\infty \frac{\chi_1(r, r_c)}{u(r) - u(r_c)} \frac{\beta(r) \min(r^2, r^{-2})^j}{w_{F,\delta}(r)} B_{\ell,0}^{(1)}(r, r_c, r_c) \varphi(r) dr \right) \\ & \times \left(p.v. \int_0^\infty \frac{1}{u(s) - u(r_c)} \frac{\beta(r_c) w_{F,\frac{\delta}{4}+2\ell}(s)}{\langle s \rangle^\delta} B_{\ell,0}^{(2)}(r_c, s, r_c) (\langle s \rangle^\delta f(s)) ds \right) dr_c, \end{aligned}$$

as operators from L^2 to L^2 . The operator L_0 is bounded on L^2 with norm $\lesssim k^{\zeta+2\ell_1+2\ell_2}$.

Proof of Theorem B.2 The theorem follows from Theorem B.1, upon verifying that the weights in (B.3)–(B.4) obey the needed conditions. This is done in Corollaries B.6, B.11 and B.14 below. \square

The second standard example of pairs of weights $\mathfrak{B}_{\varepsilon,1}$ and $\mathfrak{B}_{\varepsilon,2}$ which obey the conditions of Theorem B.1 are:

Theorem B.3 Let $0 \leq j \leq k-1$, and $0 \leq \ell, \ell_1, \ell_2$ be such that $\ell + \ell_1 + \ell_2 \leq j$. Let $0 < \zeta \leq \frac{\delta}{6}$, and $0 < \eta \leq \frac{\zeta}{4}$. Consider the weights

$$\mathfrak{B}_{\varepsilon,1}(r, s_0, r_c) = \chi_2(r, r_c) \frac{\beta(r) \min(r^2, r^{-2})^j}{r^{\frac{1}{2}} w_{f,\delta}(r) u'(s_0) r_c u'(r_c)} B_{\ell,\varepsilon}^{(1)}(r, s_0, r_c) \quad (\text{B.5})$$

$$\mathfrak{B}_{\varepsilon,2}(s_0, s, r_c) = \frac{\beta(s_0) w_{F,\frac{\delta}{4}+2\ell}(s)}{u'(s)} B_{\ell,\varepsilon}^{(2)}(s_0, s, r_c) \quad (\text{B.6})$$

where $B_{\ell,\varepsilon}^{(1)}$ is a suitable $(2\ell_1, \ell_1 + \eta/2)$ kernel of type I or II, and $B_{\ell,\varepsilon}^{(2)}$ is a suitable $(2\ell_2, \ell_2 + \eta/2)$ kernel of type I or II. Then the conditions of Theorem B.1 are satisfied for the weights (B.5)–(B.6). The corresponding operator L_ε defined in (B.1) converges to the corresponding operator L_0 defined in (B.2), as operators from L^2 to L^2 , and the limiting operator is bounded on this space, with norm bounded by $k^{\zeta+2\ell_1+2\ell_2}$.

Proof of Theorem B.3 We remark that the main difference between (B.3) and (B.5) is a factor proportional to

$$\frac{r^2 \langle r_c \rangle^4}{r_c^2}$$

besides the obvious difference of replacing χ_1 with $\chi_2 \approx \mathbf{1}_{r \leq 1} \mathbf{1}_{2r \leq r_c}$. The theorem follows from Theorem B.1, upon verifying that the weights in (B.5)–(B.6) obey the needed conditions. This is done in Corollary B.7, Corollary B.11, and Corollary B.15 below. \square

Remark B.4 It is clear from the proof of Theorems B.1, B.2, and B.3 that not all properties of a type I or type II kernel are used. For instance, for a type I kernel, Theorems B.2 and B.3 only use (6.13a), (6.13b), (6.15a), and the convergence as $\varepsilon \rightarrow 0$ in these inequalities, with some positive rate ε^ζ for some $\zeta > 0$. Similarly, for a type II kernel, Theorems B.2 and B.3 only use the global uniform boundedness, weighted Hölder regularity in each of the two variables not called r_c near the critical layer, and the convergence as $\varepsilon \rightarrow 0$ in these inequalities, at a positive rate.

The remainder of this section is dedicated to the proof of Theorem B.1, which is decomposed into several steps, detailed in the following subsections. In each subsection, we show that the weights (B.3)–(B.4) obey the necessary properties, so that the proof of Theorem B.2 is done concomitantly. Checking that the weights (B.5)–(B.6) obey the necessary properties is done at the end of this section, yielding the proof of Theorem B.3.

B.1.1 Convergence Away from the Diagonal $s_0 = r_c$

In this section we consider the contribution to the operator L_ε in (B.1) due to the set

$$\left\{ |s_0 - r_c| \geq \frac{r_c}{k} \right\}.$$

We first prove an abstract lemma, and then show that the available conditions on the coefficients $\mathfrak{B}_{\varepsilon,1}$ and $\mathfrak{B}_{\varepsilon,2}$ are sufficient in order to apply this lemma. Let us denote by $L_{\varepsilon,1}$ the contribution to the operator L_ε in (B.1) from $|s_0 - r_c| \geq \frac{r_c}{k}$, i.e. the operator

$$\begin{aligned} L_{\varepsilon,1}[f](r) &= \int_0^\infty \int_0^\infty \int_0^\infty \frac{(u(r) - u(r_c))u'(r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} \frac{(u(s) - u(r_c))u'(s)}{(u(s) - u(r_c))^2 + \varepsilon^2} \\ &\quad \times \frac{\varepsilon u'(s_0)}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \\ &\quad \times \mathbf{1}_{|s_0 - r_c| \geq \frac{r_c}{k}} \mathfrak{B}_{\varepsilon,1}(r, s_0, r_c) \mathfrak{B}_{\varepsilon,2}(s_0, s, r_c) f(s) \, ds ds_0 dr_c. \end{aligned} \quad (\text{B.7})$$

Lemma B.5 Assume that

$$|\mathfrak{B}_{\varepsilon,1}(r, s_0, r_c) \mathfrak{B}_{\varepsilon,2}(s_0, s, r_c)| \lesssim \mathfrak{B}_0(r, s_0, s, r_c) \quad (\text{B.8})$$

holds for some ε -independent function \mathfrak{B}_0 . In addition, define the cut-off

$$\mathbf{1}_{\text{strange}} = \mathbf{1}_{\frac{r_c}{2} \leq r \leq 1} \mathbf{1}_{r_c \leq 1} \mathbf{1}_{2r \leq s} \mathbf{1}_{2r_c \leq s} + \mathbf{1}_{r \leq 1} \mathbf{1}_{2r \leq r_c} \quad (\text{B.9})$$

and assume that \mathfrak{B}_0 obeys the bound

$$\begin{aligned} \mathfrak{B}_0(r, s_0, s, r_c) &\lesssim \left(\mathbf{1}_{\text{strange}} \frac{s^{\frac{1}{2}}}{r^{\frac{1}{2}} \langle s \rangle^{\frac{1}{2}}} + (1 - \mathbf{1}_{\text{strange}}) \right) \frac{r^{\frac{\delta}{2}} \langle s \rangle^{\frac{1}{2}}}{\langle r \rangle^{\frac{1}{2} + \frac{3\delta}{2}} \langle s_0 \rangle^{3\gamma}} \\ &\quad \times \left(\max \left\{ r^2, r^{-2}, s^2, s^{-2}, s_0^2, s_0^{-2} \right\} \right)^\eta \end{aligned} \quad (\text{B.10})$$

uniformly in r, s_0, s , and r_c , for some $\eta \in (0, \frac{\gamma}{16})$, $\gamma \in (0, \frac{\delta}{4})$ and $\delta \in (0, \frac{1}{2})$. Then, if $f \in L^2$, we have that $L_{\varepsilon,1}[f] \rightarrow 0$ as $\varepsilon \rightarrow 0$, in L^2 .

Proof of Lemma B.5 Let $\varphi \in L^2(\mathbb{R}_+)$ be arbitrary. We then have by (B.8) that

$$\begin{aligned} \left| \langle L_{\varepsilon,1}[f](r), \varphi(r) \rangle \right| &\leq \int_{\mathbb{R}_+^4} \frac{|(u(r) - u(r_c))u'(r_c)|}{(u(r) - u(r_c))^2 + \varepsilon^2} \frac{|(u(s) - u(r_c))u'(s)|}{(u(s) - u(r_c))^2 + \varepsilon^2} \\ &\quad \times \frac{\varepsilon |u'(s_0)|}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \\ &\quad \times \mathbf{1}_{|s_0 - r_c| \geq \frac{r_c}{k}} \mathfrak{B}_0(r, s_0, s, r_c) |f(s)\varphi(r)| \, ds ds_0 dr_c \\ &=: \int_{\mathbb{R}_+^4} \mathbb{J}(r, s_0, s, r_c) |f(s)\varphi(r)| \, ds ds_0 dr_c dr. \end{aligned} \quad (\text{B.11})$$

Our goal is to show that the integrand on the right side of (B.11) lies in $L^1(dr \, ds \, ds_0 \, dr_c)$, and moreover vanishes as $\varepsilon \rightarrow 0$ in this norm.

Case $r_c \geq 1$ The proof is based on the following estimate (c.f. (A.1)),

$$\frac{|u'(\rho)|}{|u(\rho) - u(t)|} \lesssim \mathbf{1}_{|\rho - t| \leq \frac{1}{10}} \frac{1}{|\rho - t|} \left(\mathbf{1}_{\rho \leq 1} \frac{\rho}{\rho + t} + \mathbf{1}_{\rho \geq 1} \right) + \mathbf{1}_{|\rho - t| \geq \frac{1}{10}} \frac{1}{\langle \rho \rangle} \quad (\text{B.12})$$

and the asymptotic description

$$|u'(\rho)| \approx \frac{\rho}{\langle \rho \rangle^4}. \quad (\text{B.13})$$

Here we use (B.12)–(B.13) to estimate

$$\begin{aligned} \frac{\mathbf{1}_{r_c \geq 1} \varepsilon^{\frac{\gamma}{3}} |u(r) - u(r_c)| |u'(r_c)|}{(u(r) - u(r_c))^2 + \varepsilon^2} &\lesssim \mathbf{1}_{r_c \geq 1} |u'(r_c)|^{\frac{\gamma}{3}} \left(\frac{|u'(r_c)|}{|u(r_c) - u(r)|} \right)^{1 - \frac{\gamma}{3}} \\ &\lesssim \frac{\mathbf{1}_{r_c \geq 1} \mathbf{1}_{|r_c - r| \leq \frac{1}{10}}}{\langle r_c \rangle^\gamma |r_c - r|^{1 - \frac{\gamma}{3}}} + \frac{\mathbf{1}_{r_c \geq 1} \mathbf{1}_{|r_c - r| \geq \frac{1}{10}}}{\langle r_c \rangle^{1 + \frac{2\gamma}{3}}} \end{aligned} \quad (\text{B.14})$$

and similarly, using that $|s - r_c| \leq \frac{1}{10} \Rightarrow s \geq r_c - \frac{1}{10} \geq \frac{9}{10}$ for $r_c \geq 1$, we obtain

$$\frac{\mathbf{1}_{r_c \geq 1} \varepsilon^{\frac{\gamma}{3}} |u(s) - u(r_c)| |u'(s)|}{(u(s) - u(r_c))^2 + \varepsilon^2} \lesssim \frac{\mathbf{1}_{r_c \geq 1} \mathbf{1}_{|s - r_c| \leq \frac{1}{10}} \mathbf{1}_{s \geq \frac{9}{10}}}{\langle s \rangle^\gamma |s - r_c|^{1 - \frac{\gamma}{3}}} + \frac{s^{\frac{\gamma}{3}} \mathbf{1}_{r_c \geq 1} \mathbf{1}_{|s - r_c| \geq \frac{1}{10}}}{\langle s \rangle^{1 + \gamma}}. \quad (\text{B.15})$$

Lastly, we have

$$\begin{aligned} & \frac{\mathbf{1}_{r_c \geq 1} \mathbf{1}_{|s_0 - r_c| \geq \frac{r_c}{k}} \varepsilon^{1 - \gamma} |u'(s_0)|}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \\ & \lesssim \mathbf{1}_{r_c \geq 1} \mathbf{1}_{|s_0 - r_c| \geq \frac{r_c}{k}} \frac{1}{|u'(s_0)|^\gamma} \left(\frac{|u'(s_0)|}{|u(s_0) - u(r_c)|} \right)^{1 + \gamma} \\ & \lesssim \frac{\mathbf{1}_{r_c \geq 1} \mathbf{1}_{|s_0 - r_c| \geq \frac{1}{10}}}{s_0^\gamma \langle s_0 \rangle^{1 - 3\gamma}} + \frac{\mathbf{1}_{r_c \geq 1} \mathbf{1}_{\frac{r_c}{k} \leq |s_0 - r_c| \leq \frac{1}{10}}}{|s_0 - r_c|^{1 + \gamma}} \left(\mathbf{1}_{s_0 \leq 1} \frac{s_0}{(s_0 + r_c)^{1 + \gamma}} + \mathbf{1}_{s_0 \geq 1} s_0^{3\gamma} \right) \\ & \lesssim \langle s_0 \rangle^{3\gamma} \left(\frac{\mathbf{1}_{r_c \geq 1} \mathbf{1}_{|s_0 - r_c| \geq \frac{1}{10}}}{s_0^\gamma \langle s_0 \rangle} + k^{2\gamma} \frac{\mathbf{1}_{r_c \geq 1} \mathbf{1}_{|s_0 - r_c| \leq \frac{1}{10}}}{|s_0 - r_c|^{1 - \gamma}} \right) \end{aligned} \quad (\text{B.16})$$

since $r_c \geq 1$. In the above inequalities, the implicit constants are independent of ε and k , but may depend on γ . From the above three estimates we arrive at (recall (B.11))

$$\begin{aligned} \mathbf{1}_{r_c \geq 1} \mathbb{J}(r, s_0, s, r_c) & \lesssim \varepsilon^{\frac{\gamma}{3}} \mathbf{1}_{r_c \geq 1} \mathbf{1}_{|s_0 - r_c| \geq \frac{r_c}{k}} \langle s_0 \rangle^{3\gamma} \mathfrak{B}_0(r, s_0, s, r_c) \\ & \times \left[\frac{\mathbf{1}_{|r_c - r| \leq \frac{1}{10}}}{\langle r_c \rangle^\gamma |r_c - r|^{1 - \frac{\gamma}{3}}} + \frac{\mathbf{1}_{|r_c - r| \geq \frac{1}{10}}}{\langle r_c \rangle^{1 + \frac{2\gamma}{3}}} \right] \\ & \times \left[\frac{\mathbf{1}_{|s - r_c| \leq \frac{1}{10}}}{\langle s \rangle^\gamma |s - r_c|^{1 - \frac{\gamma}{3}}} + \frac{s^{\frac{\gamma}{3}} \mathbf{1}_{|s - r_c| \geq \frac{1}{10}}}{\langle s \rangle^{1 + \gamma + \delta}} \right] \\ & \times \left[\frac{\mathbf{1}_{|s_0 - r_c| \geq \frac{1}{10}}}{s_0^\gamma \langle s_0 \rangle} + k^{2\gamma} \frac{\mathbf{1}_{|s_0 - r_c| \leq \frac{1}{10}}}{|s_0 - r_c|^{1 - \gamma}} \right]. \end{aligned}$$

At this stage we use assumption (B.10), noting that $\mathbf{1}_{\text{strange}} \mathbf{1}_{r_c \geq 1} = \mathbf{1}_{2r \leq r_c} \mathbf{1}_{r_c \geq 1} \mathbf{1}_{r \leq 1}$, (recall (B.9)) to obtain

$$\begin{aligned} \frac{\mathbf{1}_{r_c \geq 1} \mathbb{J}(r, s_0, s, r_c)}{\varepsilon^{\frac{\gamma}{3}}} & \lesssim \mathbf{1}_{r_c \geq 1} \mathbf{1}_{|s_0 - r_c| \geq \frac{r_c}{k}} \left[\mathbf{1}_{2r \leq r_c} \mathbf{1}_{r \leq 1} \frac{s^{\frac{1}{2}}}{r^{\frac{1}{2}} \langle s \rangle^{\frac{1}{2}}} + (1 - \mathbf{1}_{2r \leq r_c} \mathbf{1}_{r \leq 1}) \right] \\ & \times \frac{r^{\frac{\delta}{2}} \langle s \rangle^{\frac{1}{2} + \delta}}{\langle r \rangle^{\frac{1}{2} + \frac{3\delta}{2}}} \left[\frac{\mathbf{1}_{|r_c - r| \leq \frac{1}{10}}}{\langle r_c \rangle^\gamma |r_c - r|^{1 - \frac{\gamma}{3}}} + \frac{\mathbf{1}_{|r_c - r| \geq \frac{1}{10}}}{\langle r_c \rangle^{1 + \frac{2\gamma}{3}}} \right] \\ & \times \left(\max \left\{ r^2, r^{-2}, s^2, s^{-2}, s_0^2, s_0^{-2} \right\} \right)^\eta \end{aligned}$$

$$\begin{aligned} & \times \left[\frac{\mathbf{1}_{|s-r_c| \leq \frac{1}{10}}}{\langle s \rangle^{\gamma+\delta} |s-r_c|^{1-\frac{\gamma}{3}}} + \frac{s^{\frac{\gamma}{3}} \mathbf{1}_{|s-r_c| \geq \frac{1}{10}}}{\langle s \rangle^{1+\gamma+\delta}} \right] \left[\frac{\mathbf{1}_{|s_0-r_c| \geq \frac{1}{10}}}{s_0^{\gamma} \langle s_0 \rangle} \right. \\ & \left. + k^{2\gamma} \frac{\mathbf{1}_{|s_0-r_c| \leq \frac{1}{10}}}{|s_0-r_c|^{1-\gamma}} \right]. \end{aligned} \quad (\text{B.17})$$

We first note that since η is sufficiently small, we have

$$\int_{\mathbb{R}_+} \mathbf{1}_{r_c \geq 1} \mathbf{1}_{|s_0-r_c| \geq \frac{r_c}{k}} (\max\{s_0^2, s_0^{-2}\})^\eta \left[\frac{k^{2\gamma} \mathbf{1}_{|s_0-r_c| \leq \frac{1}{10}}}{|s_0-r_c|^{1-\gamma}} + \frac{\mathbf{1}_{|s_0-r_c| \geq \frac{1}{10}}}{s_0^{\gamma} \langle s_0 \rangle} \right] ds_0 \lesssim k^{2\gamma},$$

so that, after some manipulations, we arrive at

$$\begin{aligned} & \frac{\mathbf{1}_{r_c \geq 1}}{\varepsilon^{\frac{\gamma}{3}}} \int_{\mathbb{R}_+} \mathbb{J}(r, s_0, s, r_c) ds_0 \\ & \lesssim \mathbf{1}_{r_c \geq 1} \left[\mathbf{1}_{2r \leq r_c} \mathbf{1}_{r \leq 1} \frac{s^{\frac{1}{2}}}{r^{\frac{1}{2}} \langle s \rangle^{\frac{1}{2}}} + (1 - \mathbf{1}_{2r \leq r_c} \mathbf{1}_{r \leq 1}) \right] \left(\max\{r^2, r^{-2}, s^2, s^{-2}\} \right)^\eta \\ & \times \frac{r^{\frac{\delta}{2}} \langle s \rangle^{\frac{1}{2}+\delta}}{\langle r \rangle^{\frac{1}{2}+\frac{3\delta}{2}}} \left[\frac{\mathbf{1}_{|r_c-r| \leq \frac{1}{10}}}{\langle r_c \rangle^{\gamma} |r_c-r|^{1-\frac{\gamma}{3}}} + \frac{\mathbf{1}_{|r_c-r| \geq \frac{1}{10}}}{\langle r_c \rangle^{1+\frac{2\gamma}{3}}} \right] \left[\frac{\mathbf{1}_{|s-r_c| \leq \frac{1}{10}}}{\langle s \rangle^{\gamma+\delta} |s-r_c|^{1-\frac{\gamma}{3}}} \right. \\ & \left. + \frac{s^{\frac{\gamma}{3}} \mathbf{1}_{|s-r_c| \geq \frac{1}{10}}}{\langle s \rangle^{1+\gamma+\delta}} \right] \\ & \lesssim \frac{\mathbf{1}_{|r_c-r| \leq \frac{1}{10}}}{\langle r-r_c \rangle^{\gamma-2\eta} |r_c-r|^{1-\frac{\gamma}{3}}} \frac{\mathbf{1}_{|s-r_c| \leq \frac{1}{10}}}{\langle s-r_c \rangle^{\gamma-2\eta} |s-r_c|^{1-\frac{\gamma}{3}}} \\ & + \frac{1}{\langle r_c \rangle^{\frac{1}{2}+\delta-2\eta}} \frac{\mathbf{1}_{|r_c-r| \leq \frac{1}{10}}}{\langle r-r_c \rangle^{\gamma} |r_c-r|^{1-\frac{\gamma}{3}}} \frac{1}{\langle s \rangle^{\frac{1}{2}+\frac{2\gamma}{3}-2\eta}} \\ & + \frac{1}{\langle r_c \rangle^{\frac{1}{2}+\frac{2\gamma}{3}-2\eta}} \frac{(\max\{r^2, r^{-2}\})^\eta}{r^{\frac{1-\delta}{2}} \langle r \rangle^{\frac{3\delta}{2}}} \frac{\mathbf{1}_{|s-r_c| \leq \frac{1}{10}}}{\langle s-r_c \rangle^{\gamma} |s-r_c|^{1-\frac{\gamma}{3}}} \\ & + \frac{1}{\langle r_c \rangle^{1+\frac{2\gamma}{3}}} \frac{(\max\{r^2, r^{-2}\})^\eta}{r^{\frac{1-\delta}{2}} \langle r \rangle^{\frac{3\delta}{2}}} \frac{(\max\{s^2, s^{-2}\})^\eta}{\langle s \rangle^{\frac{1}{2}+\frac{2\gamma}{3}}} \\ & =: \mathbb{J}_{11}(r_c-r) \mathbb{J}_{12}(r_c-s) + \mathbb{J}_{21}(r_c) \mathbb{J}_{22}(r_c-r) \mathbb{J}_{23}(s) \\ & + \mathbb{J}_{31}(r_c) \mathbb{J}_{32}(r) \mathbb{J}_{33}(r_c-s) + \mathbb{J}_{41}(r_c) \mathbb{J}_{42}(r) \mathbb{J}_{43}(s), \end{aligned} \quad (\text{B.18})$$

where the identification of the \mathbb{J}_{ij} functions, for $1 \leq i \leq 4$, and $1 \leq j \leq 3$ is the obvious one. We then use Young's inequality and Hölder's inequality to deduce

$$\begin{aligned} & \int_{\mathbb{R}_+^3} \mathbb{J}_{11}(r_c-r) \mathbb{J}_{12}(r_c-s) |f(s)| |\varphi(r)| ds dr dr_c \\ & \leq \|\mathbb{J}_{11} * |\varphi|\|_{L^2(dr_c)} \|\mathbb{J}_{12} * |f|\|_{L^2(dr_c)} \end{aligned}$$

$$\begin{aligned} &\leq \|\mathbb{J}_{11}\|_{L^1} \|\mathbb{J}_{12}\|_{L^1} \|\varphi\|_{L^2} \|f\|_{L^2} \\ &\lesssim \|\varphi\|_{L^2} \|f\|_{L^2}, \end{aligned}$$

and similarly

$$\begin{aligned} &\int_{\mathbb{R}_+^3} \mathbb{J}_{21}(r_c) \mathbb{J}_{22}(r_c - r) \mathbb{J}_{23}(s) |f(s)| |\varphi(r)| \, ds dr dr_c \\ &\leq \|\mathbb{J}_{21}\|_{L^2} \|\mathbb{J}_{22}\|_{L^1} \|\mathbb{J}_{23}\|_{L^2} \|\varphi\|_{L^2} \|f\|_{L^2} \\ &\lesssim \|\varphi\|_{L^2} \|f\|_{L^2} \\ &\int_{\mathbb{R}_+^3} \mathbb{J}_{31}(r_c) \mathbb{J}_{32}(r) \mathbb{J}_{33}(r_c - s) |f(s)| |\varphi(r)| \, ds dr dr_c \\ &\leq \|\mathbb{J}_{31}\|_{L^2} \|\mathbb{J}_{32}\|_{L^2} \|\mathbb{J}_{33}\|_{L^1} \|\varphi\|_{L^2} \|f\|_{L^2} \\ &\lesssim \|\varphi\|_{L^2} \|f\|_{L^2} \\ &\int_{\mathbb{R}_+^3} \mathbb{J}_{41}(r_c) \mathbb{J}_{42}(r) \mathbb{J}_{43}(s) |f(s)| |\varphi(r)| \, ds dr dr_c \\ &\leq \|\mathbb{J}_{41}\|_{L^1} \|\mathbb{J}_{42}\|_{L^2} \|\mathbb{J}_{43}\|_{L^2} \|\varphi\|_{L^2} \|f\|_{L^2} \\ &\lesssim \|\varphi\|_{L^2} \|f\|_{L^2}. \end{aligned}$$

Summarizing the above estimates, we arrive at

$$\int_{\mathbb{R}_+^4} \mathbf{1}_{r_c \geq 1} \mathbb{J}(r, s_0, s, r_c) |f(s) \varphi(r)| \, ds ds_0 dr_c dr \lesssim \varepsilon^{\frac{\gamma}{3}} \|\varphi\|_{L^2} \|f\|_{L^2} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.$$

Case $r_c \leq 1$ In this case, instead of (B.12)–(B.13), we also have the improved estimate

$$\frac{|u'(r_c)|}{|u(r) - u(r_c)|} \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|r_c - r| \geq \frac{1}{10}} \lesssim r_c, \quad (\text{B.19})$$

which is useful when $r_c \ll 1$. Similar to the $r_c \leq 1$ case we obtain the bounds

$$\frac{\mathbf{1}_{r_c \leq 1} \varepsilon^{\frac{\gamma}{3}} |u(r) - u(r_c)| |u'(r_c)|}{(u(r) - u(r_c))^2 + \varepsilon^2} \lesssim \frac{r_c \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|r_c - r| \leq \frac{1}{10}}}{(r_c + r)^{1 - \frac{\gamma}{3}} |r_c - r|^{1 - \frac{\gamma}{3}}} + r_c \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|r_c - r| \geq \frac{1}{10}} \quad (\text{B.20})$$

$$\frac{\mathbf{1}_{r_c \leq 1} \varepsilon^{\frac{\gamma}{3}} |u(s) - u(r_c)| |u'(s)|}{(u(s) - u(r_c))^2 + \varepsilon^2} \lesssim \frac{s \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s - r_c| \leq \frac{1}{10}}}{(s + r_c)^{1 - \frac{\gamma}{3}} |s - r_c|^{1 - \frac{\gamma}{3}}} + \frac{s^{\frac{\gamma}{3}} \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s - r_c| \geq \frac{1}{10}}}{\langle s \rangle^{1 + \gamma}} \quad (\text{B.21})$$

$$\frac{\mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s_0 - r_c| \geq \frac{r_c}{k}} \varepsilon^{1 - \gamma} |u'(s_0)|}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \lesssim \frac{k^{2\gamma} \langle s_0 \rangle^{3\gamma} \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s_0 - r_c| \leq \frac{1}{10}}}{r_c^{2\gamma} |s_0 - r_c|^{1 - \gamma}} + \frac{\mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s_0 - r_c| \geq \frac{1}{10}}}{s_0^\gamma \langle s_0 \rangle^{1 - 3\gamma}}. \quad (\text{B.22})$$

From the above three estimates, and by using (B.10) we arrive at

$$\begin{aligned}
 & \frac{\mathbf{1}_{r_c \leq 1} \mathbb{J}(r, s_0, s, r_c)}{\varepsilon^{\frac{\gamma}{3}}} \\
 & \lesssim \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s_0 - r_c| \geq \frac{r_c}{k}} \mathfrak{B}_0(r, s_0, s, r_c) \\
 & \quad \times \left[\frac{r_c \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|r_c - r| \leq \frac{1}{10}}}{(r_c + r)^{1 - \frac{\gamma}{3}} |r_c - r|^{1 - \frac{\gamma}{3}}} + r_c \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|r_c - r| \geq \frac{1}{10}} \right] \\
 & \quad \times \left[\frac{s \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s - r_c| \leq \frac{1}{10}}}{(s + r_c)^{1 - \frac{\gamma}{3}} |s - r_c|^{1 - \frac{\gamma}{3}}} + \frac{s^{\frac{\gamma}{3}} \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s - r_c| \geq \frac{1}{10}}}{\langle s \rangle^{1 + \gamma}} \right] \\
 & \quad \times \left[\frac{k^{2\gamma} \langle s_0 \rangle^{3\gamma} \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s_0 - r_c| \leq \frac{1}{10}}}{r_c^{2\gamma} |r_c - s_0|^{1 - \gamma}} + \frac{\mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s_0 - r_c| \geq \frac{1}{10}}}{s_0^\gamma \langle s_0 \rangle^{1 - 3\gamma}} \right] \\
 & \lesssim \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s_0 - r_c| \geq \frac{r_c}{k}} \left(\mathbf{1}_{\text{strange}} \frac{s^{\frac{1}{2}}}{\langle s \rangle^{\frac{1}{2}} r^{\frac{1}{2}}} + (1 - \mathbf{1}_{\text{strange}}) \right) \frac{r^{\frac{\delta}{2}} \langle s \rangle^{\frac{1}{2}}}{\langle r \rangle^{\frac{1}{2} + \frac{3\delta}{2}}} \\
 & \quad \times \left[\frac{r_c \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|r_c - r| \leq \frac{1}{10}}}{(r_c + r)^{1 - \frac{\gamma}{3}} |r_c - r|^{1 - \frac{\gamma}{3}}} + r_c \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|r_c - r| \geq \frac{1}{10}} \right] \\
 & \quad \times \left(\max \{r^2, r^{-2}, s^2, s^{-2}, s_0^2, s_0^{-2}\} \right)^\eta \\
 & \quad \times \left[\frac{s \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s - r_c| \leq \frac{1}{10}}}{(s + r_c)^{1 - \frac{\gamma}{3}} |s - r_c|^{1 - \frac{\gamma}{3}}} + \frac{s^{\frac{\gamma}{3}} \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s - r_c| \geq \frac{1}{10}}}{\langle s \rangle^{1 + \gamma}} \right] \\
 & \quad \times \left[\frac{k^{2\gamma} \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s_0 - r_c| \leq \frac{1}{10}}}{r_c^{2\gamma} |r_c - s_0|^{1 - \gamma}} + \frac{\mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s_0 - r_c| \geq \frac{1}{10}}}{s_0^\gamma \langle s_0 \rangle} \right]. \tag{B.23}
 \end{aligned}$$

Similar to the estimates (B.17)–(B.18) for the case $r_c \leq 1$, we first integrate the s_0 dependent-part of (B.23) in s_0 to obtain that

$$\begin{aligned}
 & \int_{\mathbb{R}_+} \mathbf{1}_{r_c \leq 1} (\max \{s_0^2, s_0^{-2}\})^\eta \left| \frac{k^{2\gamma} \mathbf{1}_{\frac{r_c}{k} \leq |s_0 - r_c| \leq \frac{1}{10}}}{r_c^{2\gamma} |s_0 - r_c|^{1 - \gamma}} + \frac{\mathbf{1}_{|s_0 - r_c| \geq \frac{1}{10}}}{s_0^\gamma \langle s_0 \rangle} \right| ds_0 \\
 & \lesssim \frac{\mathbf{1}_{r_c \leq 1} k^{2\gamma}}{r_c^{2\gamma}} \int_{\mathbb{R}_+} (\max \{s_0^2, s_0^{-2}\})^\eta \left(\frac{\mathbf{1}_{|s_0 - r_c| \leq \frac{1}{10}}}{|s_0 - r_c|^{1 - \gamma}} + \frac{\mathbf{1}_{|s_0 - r_c| \geq \frac{1}{10}}}{s_0^\gamma \langle s_0 \rangle} \right) ds_0 \lesssim \frac{k^{2\gamma}}{r_c^{2\gamma}}
 \end{aligned}$$

since η is sufficiently small. Then, using that $\gamma \leq \frac{\delta}{4}$, we have the inequality

$$\begin{aligned}
 & \mathbf{1}_{r_c \leq 1} \frac{r^{\frac{\delta}{2}}}{r_c^{2\gamma}} \left(\mathbf{1}_{|r_c - r| \leq \frac{1}{10}} \frac{r_c}{(r + r_c)^{1 - \frac{\gamma}{3}} |r_c - r|^{1 - \frac{\gamma}{3}}} + r_c \mathbf{1}_{|r_c - r| \geq \frac{1}{10}} \right) \\
 & \lesssim r^{\frac{\delta}{2}} \mathbf{1}_{r_c \leq 1} \left(\frac{\mathbf{1}_{|r_c - r| \leq \frac{1}{10}}}{|r_c - r|^{1 - \frac{\gamma}{3}}} + \mathbf{1}_{|r_c - r| \geq \frac{1}{10}} \right),
 \end{aligned}$$

and since on the support of $\mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s-r_c| \leq \frac{1}{10}}$ we have $s \leq r_c + \frac{1}{10} \leq \frac{11}{10}$, it remains to consider the integral of $\varphi(r)f(s)$ multiplied by

$$\begin{aligned} & \int_{\mathbb{R}_+} \frac{\mathbf{1}_{r_c \leq 1} \mathbb{J}(r, s_0, s, r_c)}{\varepsilon^{\frac{\gamma}{3}}} ds_0 \\ & \lesssim \mathbf{1}_{r_c \leq 1} \left(\mathbf{1}_{\text{strange}} \frac{s^{\frac{1}{2}}}{\langle s \rangle^{\frac{1}{2}} r^{\frac{1}{2}}} + (1 - \mathbf{1}_{\text{strange}}) \right) \frac{r^{\frac{\delta}{2}}}{\langle r \rangle^{\frac{\delta}{2}}} \left(\max \{r^2, r^{-2}\} \right)^\eta \\ & \quad \times \left[\frac{\mathbf{1}_{|r_c-r| \leq \frac{1}{10}}}{|r_c-r|^{1-\frac{\gamma}{3}}} + \frac{\mathbf{1}_{|r_c-r| \geq \frac{1}{10}}}{\langle r \rangle^{\frac{1}{2}+\delta}} \right] \left[\frac{\mathbf{1}_{|s-r_c| \leq \frac{1}{10}}}{|s-r_c|^{1-\frac{\gamma}{3}}} + \frac{\mathbf{1}_{|s-r_c| \geq \frac{1}{10}}}{\langle s \rangle^{\frac{1}{2}+\frac{2\gamma}{3}-2\eta}} \right] \\ & \lesssim \mathbf{1}_{r_c \leq 1} \frac{1}{r^{\frac{1}{2}-\frac{\gamma}{6}} r_c^{\frac{\gamma}{6}}} \left[\frac{\mathbf{1}_{|r_c-r| \leq \frac{1}{10}}}{|r_c-r|^{1-\frac{\gamma}{3}}} + \frac{1}{\langle r \rangle^{\frac{1}{2}+\delta}} \right] \left[\frac{\mathbf{1}_{|s-r_c| \leq \frac{1}{10}}}{|s-r_c|^{\frac{1}{2}-\frac{\gamma}{3}}} + \frac{1}{\langle s \rangle^{\frac{1}{2}+\frac{2\gamma}{3}}} \right] \\ & \quad + \mathbf{1}_{r_c \leq 1} \left[\frac{\mathbf{1}_{|r_c-r| \leq \frac{1}{10}}}{|r_c-r|^{1-\frac{\gamma}{3}}} + \frac{1}{\langle r \rangle^{\frac{1}{2}+\delta}} \right] \left[\frac{\mathbf{1}_{|s-r_c| \leq \frac{1}{10}}}{|s-r_c|^{1-\frac{\gamma}{3}}} + \frac{1}{\langle s \rangle^{\frac{1}{2}+\frac{2\gamma}{3}}} \right]. \end{aligned}$$

Here we have used properties of the support of $\mathbf{1}_{\text{strange}}$ (recall (B.9)). Similar to the case $r_c \geq 1$, for the second term above (the one coming from $1 - \mathbf{1}_{\text{strange}}$) one may use Young's and Hölder's inequality to check that

$$\begin{aligned} & \int_{\mathbb{R}_+^3} \mathbf{1}_{r_c \leq 1} \left[\frac{\mathbf{1}_{|r_c-r| \leq \frac{1}{10}}}{|r_c-r|^{1-\frac{\gamma}{3}}} + \frac{1}{\langle r \rangle^{\frac{1}{2}+\delta}} \right] \left[\frac{\mathbf{1}_{|s-r_c| \leq \frac{1}{10}}}{|s-r_c|^{1-\frac{\gamma}{3}}} + \frac{1}{\langle s \rangle^{\frac{1}{2}+\frac{2\gamma}{3}}} \right] |\varphi(r)f(s)| dr ds dr_c \\ & \lesssim \|\varphi\|_{L^2} \|f\|_{L^2}. \end{aligned}$$

For the term first term (due to $\mathbf{1}_{\text{strange}}$), we first note that

$$\begin{aligned} & \sup_{r_c \leq 1} \int_{\mathbb{R}_+} |f(s)| \left[\frac{\mathbf{1}_{|s-r_c| \leq \frac{1}{10}}}{|s-r_c|^{\frac{1}{2}-\frac{\gamma}{3}}} + \frac{1}{\langle s \rangle^{\frac{1}{2}+\frac{2\gamma}{3}}} \right] ds \\ & \lesssim \|f\|_{L^2} \sup_{r_c \leq 1} \left\| \frac{\mathbf{1}_{|s-r_c| \leq \frac{1}{10}}}{|s-r_c|^{\frac{1}{2}-\frac{\gamma}{3}}} + \frac{1}{\langle s \rangle^{\frac{1}{2}+\frac{2\gamma}{3}}} \right\|_{L^2(ds)} \\ & \lesssim \|f\|_{L^2}, \end{aligned}$$

so that we only are left to bound

$$\begin{aligned} & \int_{\mathbb{R}_+^2} |\varphi(r)| \frac{\mathbf{1}_{r \leq \frac{11}{10}} \mathbf{1}_{r_c \leq 1}}{r^{\frac{1}{2}-\frac{\gamma}{6}} r_c^{\frac{\gamma}{6}}} \left[\frac{\mathbf{1}_{|r_c-r| \leq \frac{1}{10}}}{|r_c-r|^{1-\frac{\gamma}{3}}} + \frac{1}{\langle r \rangle^{\frac{1}{2}+\delta}} \right] dr dr_c \\ & \lesssim \|\varphi\|_{L^2} \left\| \frac{\mathbf{1}_{r \leq \frac{11}{10}}}{r^{\frac{1}{2}-\frac{\gamma}{6}}} \right\|_{L^2(dr)} \sup_{r \leq \frac{11}{10}} \left\| \frac{\mathbf{1}_{r_c \leq 1} \mathbf{1}_{|r-r_c| \leq \frac{1}{10}}}{r_c^{\frac{\gamma}{6}} |r-r_c|^{1-\frac{\gamma}{3}}} \right\|_{L^1(dr_c)} \end{aligned}$$

$$\begin{aligned}
& + \|\varphi\|_{L^2} \left\| \frac{1}{r^{\frac{1}{2}-\frac{\gamma}{6}} \langle r \rangle^{\frac{1}{2}+\delta}} \right\|_{L^2(dr)} \left\| \frac{\mathbf{1}_{r_c \leq 1}}{r_c^{\frac{\gamma}{6}}} \right\|_{L^1(dr_c)} \\
& \lesssim \|\varphi\|_{L^2}.
\end{aligned}$$

Combining the above, we arrive at

$$\int_{\mathbb{R}_+^4} \mathbf{1}_{r_c \leq 1} |\mathbb{J}(r, s_0, s, r_c)(f(s))\varphi(r)| \, ds ds_0 dr_c dr \lesssim \varepsilon^{\frac{\gamma}{3}} \|\varphi\|_{L^2} \|f\|_{L^2} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0$$

which is the desired estimate, and concludes the proof of the lemma. \square

Corollary B.6 *Let $0 \leq j \leq k-1$, and $0 \leq \ell, \ell_1, \ell_2$ be such that $\ell + \ell_1 + \ell_2 \leq j$. Assume that the functions $\mathfrak{B}_{\varepsilon,1}$ and $\mathfrak{B}_{\varepsilon,2}$ in (B.7) are given by (B.3)–(B.4), where $B_{\ell,\varepsilon}^{(1)}$ is a suitable $(2\ell_1, \ell_1 + \eta/2)$ kernel of type I or II, and $B_{\ell,\varepsilon}^{(2)}$ is a suitable $(2\ell_2, \ell_2 + \eta/2)$ kernel of type I or II. Then the operator $L_{\varepsilon,1}$ defined in (B.7) vanishes in $L^2(dr)$ as $\varepsilon \rightarrow 0$.*

Proof of Corollary B.6 We recall that the following estimates are available

$$\begin{aligned}
|B_{\ell,\varepsilon}^{(1)}(r, s_0, r_c)| & \lesssim |u'(s_0)| \mathcal{B}(r, s_0) \mathbb{K}(r, s_0, r_c) \mathcal{L}_{2\ell_1, \ell_1 + \eta/2}(r, s_0) \\
|B_{\ell,\varepsilon}^{(2)}(s_0, s, r_c)| & \lesssim |u'(s)| \mathcal{B}(s_0, s) \mathbb{K}(s_0, s, r_c) \mathcal{L}_{2\ell_2, \ell_2 + \eta/2}(s_0, s)
\end{aligned}$$

on \mathbb{R}_+^3 , where as before we recall the definitions

$$\begin{aligned}
\mathcal{B}(r, s) & = \left(\mathbf{1}_{s < r} \frac{s^{k-\frac{1}{2}}}{r^{k-\frac{1}{2}}} + \mathbf{1}_{s > r} \frac{r^{k+\frac{1}{2}}}{s^{k+\frac{1}{2}}} \right) \langle s \rangle^4 \\
\mathbb{K}(r, s, r_c) & = \mathbf{1}_{r_c > 1} + \mathbf{1}_{r_c \leq 1} \left(\mathbf{1}_{s < r < r_c} + \mathbf{1}_{s < r_c < r < 1} \frac{r_c^2}{r^2} + \mathbf{1}_{s < r_c < 1 < r} r_c^2 \right. \\
& \quad \left. + \mathbf{1}_{r_c < s < r < 1} \frac{s^2}{r^2} + \mathbf{1}_{r_c < s < 1 < r} s^2 + \mathbf{1}_{1 < s < r} \right. \\
& \quad \left. + \mathbf{1}_{r < s < r_c} + \mathbf{1}_{r < r_c < s < 1} \frac{r_c^2}{s^2} + \mathbf{1}_{r < r_c < 1 < s} r_c^2 + \mathbf{1}_{r_c < r < s < 1} \frac{r^2}{s^2} \right. \\
& \quad \left. + \mathbf{1}_{r_c < r < 1 < s} r^2 + \mathbf{1}_{1 < r < s} \right) \\
\mathcal{L}_{J,\ell}(r, s) & = k^J \left(\max \left\{ \frac{1}{r^2}, r^2, \frac{1}{s^2}, s^2 \right\} \right)^\ell.
\end{aligned} \tag{B.24}$$

Recalling the definition of the weights $w_{F,\delta}$ we obtain that

$$\begin{aligned}
& \mathbf{1}_{|s_0 - r_c| \geq \frac{r_c}{k}} |B_{\ell,\varepsilon}^{(1)}(r, s_0, r_c) B_{\ell,\varepsilon}^{(2)}(s_0, s, r_c)| \\
& \lesssim \mathbf{1}_{|s_0 - r_c| \geq \frac{r_c}{k}} (\mathbf{1}_{r \leq 1} \mathbf{1}_{r_c \leq 2r} + \mathbf{1}_{r \geq 1}) \left[\frac{\mathbf{1}_{r \leq 1}}{r^{k+3-2j-\delta}} + \mathbf{1}_{r \geq 1} r^{k-1-2j-\delta} \right]
\end{aligned}$$

$$\begin{aligned}
& \mathcal{L}_{2\ell_1, \ell_1+\eta/2}(r, s_0) \mathcal{L}_{2\ell_2, \ell_2+\eta/2}(s_0, s) \\
& \times \frac{1}{\langle s_0 \rangle^6} \mathcal{B}(r, s_0) \mathbb{K}(r, s_0, r_c) \left[\mathbf{1}_{s \leq 1} s^{k+3-2\ell-\frac{\delta}{4}} + \frac{\mathbf{1}_{s \geq 1}}{s^{k+5-2\ell-\frac{\delta}{4}}} \right] \mathcal{B}(s_0, s) \mathbb{K}(s_0, s, r_c) \\
& =: \mathfrak{B}_0(r, s_0, s, r_c).
\end{aligned} \tag{B.25}$$

The above defined function \mathfrak{B}_0 is explicit, and we need to verify that it obeys condition (B.10). Note that the terms due to the η corrections in \mathcal{L} are already incorporated in the $(\max\{r^2, r^{-2}, s^2, s^{-2}, s_0^2, s_0^{-2}\})^\eta$ term on the right side of (B.10), so that we ignore these factors from here on, working as if $\eta = 0$. This is done by considering the possible orderings of r, s_0, s , and r_c . It is useful to denote by

$$\begin{aligned}
\mathbb{W}(r, s, s_0) &:= \left[\frac{\mathbf{1}_{r \leq 1}}{r^{k+\frac{1}{2}}} + \mathbf{1}_{r \geq 1} r^{k-\frac{1}{2}} \right] \left[\mathbf{1}_{s \leq 1} s^{k+\frac{1}{2}} + \frac{\mathbf{1}_{s \geq 1}}{s^{k-\frac{1}{2}}} \right] \frac{\mathcal{B}(r, s_0)}{\langle s_0 \rangle^4} \frac{\mathcal{B}(s_0, s)}{\langle s \rangle^4} \\
\mathbb{L}(r, s, s_0) &:= \left[\mathbf{1}_{r \leq 1} r^{2j} + \frac{\mathbf{1}_{r \geq 1}}{r^{2j}} \right] \left[\frac{\mathbf{1}_{s \leq 1}}{s^{2\ell}} + \mathbf{1}_{s \geq 1} s^{2\ell} \right] \mathcal{L}_{2\ell_1, \ell_1}(r, s_0) \mathcal{L}_{2\ell_2, \ell_2}(s_0, s).
\end{aligned}$$

In view of Lemma B.17, we have that

$$\mathbb{W}(r, s, s_0) \mathbb{L}(r, s, s_0) \lesssim 1. \tag{B.26}$$

Estimate (B.26) requires some care in proving and we defer the proof to the Subsection B.1.4. With this notation, and using estimate (B.26), we have that

$$\begin{aligned}
\mathfrak{B}_0(r, s_0, s, r_c) &\lesssim \mathbf{1}_{|s_0-r_c| \geq \frac{r_c}{k}} (\mathbf{1}_{r \leq 1} \mathbf{1}_{r_c \leq 2r} + \mathbf{1}_{r \geq 1}) \left[\frac{\mathbf{1}_{r \leq 1}}{r^{3-\frac{1}{2}-\delta}} + \frac{\mathbf{1}_{r \geq 1}}{r^{\frac{1}{2}+\delta}} \right] \\
&\times \left[\mathbf{1}_{s \leq 1} s^{3-\frac{1}{2}-\frac{\delta}{4}} + \frac{\mathbf{1}_{s \geq 1}}{s^{\frac{3}{2}-\frac{\delta}{4}}} \right] \frac{1}{\langle s_0 \rangle^2} \mathbb{K}(r, s_0, r_c) \mathbb{K}(s_0, s, r_c).
\end{aligned} \tag{B.27}$$

Checking condition (B.10) for the above defined \mathfrak{B}_0 thus reduces to verifying the uniform boundedness of

$$\begin{aligned}
\mathbb{J}(r, s_0, s, r_c) &:= \mathbf{1}_{|s_0-r_c| \geq \frac{r_c}{k}} (\mathbf{1}_{r \leq 1} \mathbf{1}_{r_c \leq 2r} + \mathbf{1}_{r \geq 1}) \left(\mathbf{1}_{\text{strange}} \frac{r^{\frac{1}{2}} \langle s \rangle^{\frac{1}{2}}}{s^{\frac{1}{2}}} + (1 - \mathbf{1}_{\text{strange}}) \right) \\
&\times \left[\frac{\mathbf{1}_{r \leq 1}}{r^{\frac{5}{2}-\frac{\delta}{2}}} + \mathbf{1}_{r \geq 1} \right] \left[\mathbf{1}_{s \leq 1} s^{\frac{5}{2}-\frac{\delta}{4}} + \frac{\mathbf{1}_{s \geq 1}}{s^{2-\frac{3\delta}{4}}} \right] \mathbb{K}(r, s_0, r_c) \mathbb{K}(s_0, s, r_c),
\end{aligned} \tag{B.28}$$

where we have used that $3\gamma \leq 2$, and thus $\langle s_0 \rangle^{-2+3\gamma} \lesssim 1$.

Case $r_c > 1$ Note that here we are working on the support of $\mathbf{1}_{\text{strange}} = 0$, and by definition $\mathbb{K}(r, s_0, r_c) \lesssim 1$, $\mathbb{K}(s_0, s, r_c) \lesssim 1$. Thus, condition (B.28) reduces to

proving the uniform boundedness of

$$\mathbb{J}_{r_c > 1} = \left[\frac{\mathbf{1}_{\frac{1}{2} < r \leq 1}}{r^{\frac{5}{2} - \frac{\delta}{2}}} + \mathbf{1}_{r \geq 1} \right] \left[\mathbf{1}_{s \leq 1} s^{\frac{5}{2} - \frac{\delta}{4}} + \frac{\mathbf{1}_{s \geq 1}}{s^{2 - \frac{3\delta}{4}}} \right].$$

Since both of the above terms are $\lesssim 1$, so is their product, and thus

$$\mathbb{J}_{r_c > 1} \lesssim 1$$

as desired.

Case $r_c < 1$ From condition (B.10) we need to show that

$$\begin{aligned} \mathbb{J}_{r_c \leq 1} &:= \mathbf{1}_{r_c \leq 1} \left(\mathbf{1}_{\text{strange}} \frac{r^{\frac{1}{2}} \langle s \rangle^{\frac{1}{2}}}{s^{\frac{1}{2}}} + (1 - \mathbf{1}_{\text{strange}}) \right) (\mathbf{1}_{r \leq 1} \mathbf{1}_{r_c \leq 2r} + \mathbf{1}_{r \geq 1}) \\ &\quad \times \left[\frac{\mathbf{1}_{\frac{r_c}{2} \leq r \leq 1}}{r^{\frac{5}{2} - \frac{\delta}{2}}} + \mathbf{1}_{r \geq 1} \right] \left[\mathbf{1}_{s \leq 1} s^{\frac{5}{2} - \frac{\delta}{4}} + \frac{\mathbf{1}_{s \geq 1}}{s^{2 - \frac{3\delta}{4}}} \right] \mathbb{K}(r, s_0, r_c) \mathbb{K}(s_0, s, r_c) \end{aligned} \quad (\text{B.29})$$

is uniformly bounded in r, r_c, s, s_0 . As in the case $r_c > 1$, since $\mathbb{K}(s_0, s, r_c) \leq 1$, $\mathbb{K}(r, s_0, r_c) \leq 1$, it is clear that proving the uniform boundedness of the term $\mathbb{J}_{r_c \leq 1}$ defined in (B.29), resumes to checking the uniform boundedness of $\mathbb{J}_{r_c \leq 1} \mathbf{1}_{r \leq 1} \mathbf{1}_{r \leq s}$. Indeed, when $s \leq r$ the quotient $(s/r)^{5/2 - \delta/4} \lesssim 1$, and no singularity at $r \ll 1$ arises. This issue is avoided altogether if $r \geq 1$. Thus, we see that our desired estimate $\mathbb{J}_{r_c \leq 1}$ reduces to proving the uniform boundedness of

$$\begin{aligned} \mathbb{J}_{r_c \leq 1} \mathbf{1}_{\frac{r_c}{2} \leq r \leq 1} \mathbf{1}_{r \leq s} &= \mathbb{J}_{r_c \leq 1}^{(1)} + \mathbb{J}_{r_c \leq 1}^{(2)} \\ &:= \mathbf{1}_{r_c \leq 1} \left(\mathbf{1}_{\text{strange}} \frac{r^{\frac{1}{2}} \langle s \rangle^{\frac{1}{2}}}{s^{\frac{1}{2}}} + (1 - \mathbf{1}_{\text{strange}}) \right) \left[\frac{\mathbf{1}_{\frac{r_c}{2} \leq r \leq 1}}{r^{\frac{5}{2} - \frac{\delta}{2}}} + \mathbf{1}_{r \geq 1} \right] \\ &\quad \times \mathbf{1}_{s \leq 1} s^{\frac{5}{2} - \frac{\delta}{4}} \mathbb{K}(r, s_0, r_c) \mathbb{K}(s_0, s, r_c) \\ &\quad + \mathbf{1}_{r_c \leq 1} \left(\mathbf{1}_{\text{strange}} \frac{r^{\frac{1}{2}} \langle s \rangle^{\frac{1}{2}}}{s^{\frac{1}{2}}} + (1 - \mathbf{1}_{\text{strange}}) \right) \left[\frac{\mathbf{1}_{\frac{r_c}{2} \leq r \leq 1}}{r^{\frac{5}{2} - \frac{\delta}{2}}} + \frac{\mathbf{1}_{r \geq 1}}{r^{1 + \delta}} \right] \\ &\quad \times \frac{\mathbf{1}_{s \geq 1}}{s^{2 - \frac{3\delta}{4}}} \mathbb{K}(r, s_0, r_c) \mathbb{K}(s_0, s, r_c) \end{aligned} \quad (\text{B.30})$$

where the decomposition is based on $s \leq 1$ or $s \geq 1$.

When $s \leq 1$, and either $s \leq 2r_c$ or $s \leq 2r$, the quotient $(s/r)^{5/2 - \delta/4}$ is bounded by a universal constant, so we are left to consider the case $s \geq 2r$ and $s \geq 2r_c$, which is precisely the support of $\mathbf{1}_{\text{strange}}$. Therefore, the boundedness of $\mathbb{J}_{r_c \leq 1}^{(1)}$ reduces to the boundedness of

$$\mathbb{J}_{r_c \leq 1}^{(1)} \mathbf{1}_{\text{strange}} = \mathbf{1}_{\text{strange}} \mathbf{1}_{s \leq 1} \frac{s^{2-\frac{\delta}{4}}}{r^{2-\frac{\delta}{2}}} \mathbb{K}(s_0, s, r_c) \mathbb{K}(r, s_0, r_c).$$

At this stage the specific form of \mathbb{K} is useful to us. By analyzing the product $\mathbb{K}(s_0, s, r_c) \mathbb{K}(r, s_0, r_c)$ we note that

$$\begin{aligned} & \mathbf{1}_{\frac{r_c}{2} \leq r \leq s \leq 1} \mathbf{1}_{r_c \leq 1} \mathbb{K}(s_0, s, r_c) \mathbb{K}(r, s_0, r_c) \\ &= \left(\mathbf{1}_{s_0 \leq r_c \leq r \leq s \leq 1} \frac{r_c^4}{s^2 r^2} + \mathbf{1}_{r_c \leq s_0 \leq r \leq s \leq 1} \frac{s_0^4}{s^2 r^2} + \mathbf{1}_{r_c \leq r \leq s_0 \leq s \leq 1} \frac{r^2}{s^2} \right. \\ & \quad \left. + \mathbf{1}_{r_c \leq r \leq s \leq s_0 \leq 1} \frac{r^2 s^2}{s_0^4} + \mathbf{1}_{r_c \leq r \leq s \leq 1 \leq s_0} s^2 r^2 \right) \\ & \quad + \mathbf{1}_{\frac{r_c}{2} \leq r} \left(\mathbf{1}_{s_0 \leq r \leq r_c \leq s \leq 1} \frac{r_c^2}{s^2} + \mathbf{1}_{r \leq s_0 \leq r_c \leq s \leq 1} \frac{r_c^2}{s^2} + \mathbf{1}_{r \leq r_c \leq s_0 \leq s \leq 1} \frac{r^2}{s^2} \right. \\ & \quad \left. + \mathbf{1}_{r \leq r_c \leq s \leq s_0 \leq 1} \frac{s^2 r_c^2}{s_0^4} + \mathbf{1}_{r \leq r_c \leq s \leq 1 \leq s_0} s^2 r_c^2 \right) \\ & \quad + \mathbf{1}_{\frac{r_c}{2} \leq r} \left(\mathbf{1}_{s_0 \leq r \leq s \leq r_c \leq 1} + \mathbf{1}_{r \leq s_0 \leq s \leq r_c \leq 1} + \mathbf{1}_{r \leq s \leq s_0 \leq r_c \leq 1} \right. \\ & \quad \left. + \mathbf{1}_{r \leq s \leq r_c \leq s_0 \leq 1} \frac{r_c^4}{s_0^4} + \mathbf{1}_{r \leq s \leq r_c \leq 1 \leq s_0} r_c^4 \right) \\ & \lesssim \left(\mathbf{1}_{s_0 \leq r_c \leq r \leq s \leq 1} \frac{r^2}{s^2} + \mathbf{1}_{r_c \leq s_0 \leq r \leq s \leq 1} \frac{r^2}{s^2} + \mathbf{1}_{r_c \leq r \leq s_0 \leq s \leq 1} \frac{r^2}{s^2} \right. \\ & \quad \left. + \mathbf{1}_{r_c \leq r \leq s \leq s_0 \leq 1} \frac{r^2}{s^2} + \mathbf{1}_{r_c \leq r \leq s \leq 1 \leq s_0} s^2 r^2 \right) \\ & \quad + \mathbf{1}_{\frac{r_c}{2} \leq r} \left(\mathbf{1}_{s_0 \leq r \leq r_c \leq s \leq 1} \frac{r^2}{s^2} + \mathbf{1}_{r \leq s_0 \leq r_c \leq s \leq 1} \frac{r^2}{s^2} + \mathbf{1}_{r \leq r_c \leq s_0 \leq s \leq 1} \frac{r^2}{s^2} \right. \\ & \quad \left. + \mathbf{1}_{r \leq r_c \leq s \leq s_0 \leq 1} \frac{r^2}{s^2} + \mathbf{1}_{r \leq r_c \leq s \leq 1 \leq s_0} s^2 r^2 \right) \\ & \quad + \mathbf{1}_{\frac{r_c}{2} \leq r} \left(\mathbf{1}_{s_0 \leq r \leq s \leq r_c \leq 1} \frac{r^2}{s^2} + \mathbf{1}_{r \leq s_0 \leq s \leq r_c \leq 1} \frac{r^2}{s^2} + \mathbf{1}_{r \leq s \leq s_0 \leq r_c \leq 1} \frac{r^2}{s^2} \right. \\ & \quad \left. + \mathbf{1}_{r \leq s \leq r_c \leq s_0 \leq 1} \frac{r^2}{s^2} + \mathbf{1}_{r \leq s \leq r_c \leq 1 \leq s_0} r^4 \right) \\ & \lesssim \mathbf{1}_{\frac{r_c}{2} \leq r \leq s \leq 1} \mathbf{1}_{r_c \leq 1} \frac{r^2}{s^2}. \end{aligned}$$

Therefore, since $r \leq s \leq 1$, we are left with

$$\mathbb{J}_{r_c \leq 1}^{(1)} \mathbf{1}_{\text{strange}} \lesssim 1$$

which is the needed estimate.

The case $s \geq 1$ is treated similarly. When $s \leq 2r$ or $s \leq 2r_c$, then since $\frac{r_c}{2} \leq r \leq s$, r is bounded from below, and thus there is no loss of $r^{-\frac{5}{2}+\frac{\delta}{2}}$. We are left to consider the support of $\mathbf{1}_{\text{strange}}$ and bound the term

$$\mathbb{J}_{r_c \leq 1}^{(2)} \mathbf{1}_{\text{strange}} = \mathbf{1}_{\text{strange}} \mathbf{1}_{s \geq 1} \frac{1}{r^{2-\frac{\delta}{2}} s^{2-\frac{3\delta}{4}}} \mathbb{K}(s_0, s, r_c) \mathbb{K}(r, s_0, r_c)$$

Here we use that

$$\begin{aligned} & \mathbf{1}_{\frac{r_c}{2} \leq r \leq 1 < s} \mathbf{1}_{r_c \leq 1} \mathbb{K}(s_0, s, r_c) \mathbb{K}(r, s_0, r_c) \\ &= \left(\mathbf{1}_{s_0 \leq r_c \leq r \leq 1 < s} \frac{r_c^4}{r^2} + \mathbf{1}_{r_c \leq s_0 \leq r \leq 1 < s} \frac{s_0^4}{r^2} + \mathbf{1}_{r_c \leq r \leq s_0 \leq 1 < s} r^2 \right. \\ & \quad \left. + \mathbf{1}_{r_c \leq r \leq 1 < s_0 \leq s} r^2 + \mathbf{1}_{r_c \leq r \leq 1 < s \leq s_0} r^2 \right) \\ & \quad + \mathbf{1}_{\frac{r_c}{2} \leq r \leq r_c \leq 1 < s} r_c^2 \\ & \lesssim \mathbf{1}_{\frac{r_c}{2} \leq r \leq 1 < s} \mathbf{1}_{r_c \leq 1} r^2. \end{aligned} \quad (\text{B.31})$$

Therefore, it follows that

$$\mathbb{J}_{r_c \leq 1}^{(2)} \mathbf{1}_{\text{strange}} = \mathbf{1}_{\text{strange}} \mathbf{1}_{s \geq 1} \frac{r^{\frac{\delta}{2}}}{s^{2-\frac{3\delta}{4}}} \lesssim 1$$

which concludes the proof. \square

A similar result may be obtained for the weight which has $\chi_2(r, r_c)$ instead of $\chi_1(r, r_c)$, but an additional argument has to be given to control the region in which r_c is much larger than all the other parameters. We have:

Corollary B.7 *Let $0 \leq j \leq k-1$, and $0 \leq \ell, \ell_1, \ell_2$ be such that $\ell + \ell_1 + \ell_2 \leq j$. Assume that the functions $\mathfrak{B}_{\varepsilon,1}$ and $\mathfrak{B}_{\varepsilon,2}$ in (B.7) are given by (B.5)–(B.6), where $B_{\ell,\varepsilon}^{(1)}$ is a suitable $(2\ell_1, \ell_1 + \eta/2)$ kernel of type I or II, and $B_{\ell,\varepsilon}^{(2)}$ is a suitable $(2\ell_2, \ell_2 + \eta/2)$ kernel of type I or II. Then the operator $L_{\varepsilon,1}$ defined in (B.7) vanishes in $L^2(dr)$ as $\varepsilon \rightarrow 0$.*

Proof of Corollary B.7 The proof is nearly identical to the proof of Corollary B.6, so we only emphasize here the points which are different. As noted below Theorem B.3, the main difference is that a factor of $\frac{r^2 \langle r_c \rangle^4}{r_c^2}$ enters the estimates. Using the definition of $\mathbf{1}_{\text{strange}}$, we see that instead of checking the uniform boundedness of the expression in (B.28), we are left to check the uniform boundedness of the new expression

$$\begin{aligned} \mathbb{J}(r, s_0, s, r_c) &:= \mathbf{1}_{|s_0 - r_c| \geq \frac{r_c}{k}} \frac{r^2 \langle r_c \rangle^4}{r_c^2} \beta(s_0) \langle s_0 \rangle^{4+3\gamma} \mathbf{1}_{2r \leq r_c} \frac{r^{\frac{1}{2}} \langle s \rangle^{\frac{1}{2}}}{s^{\frac{1}{2}}} \frac{\mathbf{1}_{r \leq 1}}{r^{\frac{5}{2}-\frac{\delta}{2}}} \\ & \quad \times \left[\mathbf{1}_{s \leq 1} s^{\frac{5}{2}-\frac{\delta}{4}} + \frac{\mathbf{1}_{s \geq 1}}{s^{2-\frac{3\delta}{4}}} \right] \mathbb{K}(r, s_0, r_c) \mathbb{K}(s_0, s, r_c) \end{aligned}$$

$$\leq \mathbf{1}_{|s_0-r_c| \geq \frac{r_c}{k}} \frac{\langle r_c \rangle^4}{r_c^2} \beta(s_0) \langle s_0 \rangle^{4+3\gamma} \mathbf{1}_{2r \leq r_c} \mathbf{1}_{r \leq 1} r^{\frac{\delta}{2}} \\ \times \left[\mathbf{1}_{s \leq 1} s^{2-\frac{\delta}{4}} + \frac{\mathbf{1}_{s \geq 1}}{s^{2-\frac{3\delta}{4}}} \right] \mathbb{K}(r, s_0, r_c) \mathbb{K}(s_0, s, r_c).$$

Since $\mathbb{K}(r, s_0, r_c) \mathbb{K}(s_0, s, r_c) \leq 1$, the boundedness in the regions $s \leq r_c \leq 1$, and $1 \leq r_c \leq 2s$ follows immediately. Moreover, for the region $\mathbf{1}_{r_c \leq 1} \mathbf{1}_{s \geq r_c}$ we explicitly check that

$$\mathbf{1}_{r_c \leq 1} \mathbf{1}_{s \geq r_c} \mathbf{1}_{2r \leq r_c} \mathbf{1}_{r \leq 1} \frac{s^2}{r_c^2 \langle s \rangle^4} \mathbb{K}(r, s_0, r_c) \mathbb{K}(s_0, s, r_c) \\ \lesssim \mathbf{1}_{r_c \leq 1} \mathbf{1}_{s \geq r_c} \mathbf{1}_{2r \leq r_c} \mathbf{1}_{r \leq 1} \left[\mathbf{1}_{s \leq 1} \left(1 + s^4 + \mathbf{1}_{s \leq s_0} \frac{s^4}{s_0^4} \right) + \mathbf{1}_{s \geq 1} \frac{1}{s^2} \right] \lesssim 1.$$

We are left to consider the region where $\mathbf{1}_{r_c \geq 1} \mathbf{1}_{2s \leq r_c}$. If $r_c \leq 2s_0$, we can absorb the bad power of r_c^2 into and s_0^2 , and use that $\beta(s_0) \langle s_0 \rangle^{6+3\gamma} \leq 1$ to obtain the desired boundedness. However, in the case $r_c \gg 2r, 2s, 2s_0, 2$, there is nothing to make the above term, and a different argument is needed.

We recall that in the definition of $\chi_2(r, r_c)$ we have the cut-off function $\chi_I(r_c)$, which restricts our attention to $r_c \leq \varepsilon^{-\frac{1}{2+\alpha}}$ for some $\alpha > 0$. Here this information is used essentially. We start from the beginning of the proof of Lemma B.5, namely from (B.11), and focus only on the remaining region $\mathbf{1}_{r \leq 1} \mathbf{1}_{2r \leq r_c} \mathbf{1}_{r_c \geq 1} \mathbf{1}_{2s \leq r_c} \mathbf{1}_{2s_0 \leq r_c}$. We are instead left to consider the convergence as $\varepsilon \rightarrow 0$ of

$$\int_{\mathbb{R}_+^4} \frac{|(u(r) - u(r_c))u'(r_c)| |(u(s) - u(r_c))u'(s)|}{(u(r) - u(r_c))^2 + \varepsilon^2} \frac{\varepsilon |u'(s_0)|}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \mathbf{1}_{r \leq 1} \mathbf{1}_{2r \leq r_c} \mathbf{1}_{2s \leq r_c} \mathbf{1}_{2s_0 \leq r_c} \\ \times \frac{\chi_I(r_c) r^{2+2j} r_c^2 \beta(s_0) w_{F, \frac{\delta}{4}+2\ell}(s) \mathcal{B}(r, s_0) \mathbb{K}(r, s_0, r_c) \mathcal{L}_{2\ell_1, \ell_1+\eta/2}(r, s_0) \mathcal{B}(s_0, s) \mathbb{K}(s_0, s, r_c) \mathcal{L}_{2\ell_1, \ell_1+\eta/2}(s_0, s)}{w_{F, \delta}(r)} \\ \times |f(s)\varphi(r)| ds ds_0 dr_c \\ =: \int_{\mathbb{R}_+^4} \mathbb{J}(r, s_0, s, r_c) |f(s)\varphi(r)| ds ds_0 dr_c dr. \quad (\text{B.32})$$

By appealing to (B.26) and the boundedness of \mathbb{K} , similarly to (B.28) we obtain that

$$\mathbb{J}(r, s_0, s, r_c) \\ \lesssim \frac{|u'(r_c)|}{\sqrt{(u(r) - u(r_c))^2 + \varepsilon^2}} \frac{|u'(s)|}{\sqrt{(u(s) - u(r_c))^2 + \varepsilon^2}} \frac{\varepsilon |u'(s_0)|}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \\ \times \mathbf{1}_{r \leq 1} \mathbf{1}_{r_c \leq 2r} \mathbf{1}_{2r \leq r_c} \mathbf{1}_{2s \leq r_c} \mathbf{1}_{2s_0 \leq r_c} \frac{\chi_I(r_c) r_c^2 \langle s_0 \rangle^{4+\eta} s_0^{-2\eta} \beta(s_0) \left(\mathbf{1}_{s \leq 1} s^{\frac{5}{2}-\frac{\delta}{4}-2\eta} + \mathbf{1}_{s \geq 1} s^{-\frac{3}{2}+\frac{\delta}{4}+2\eta} \right)}{r^{\frac{1}{2}-\delta+2\eta}},$$

and using the definition of χ_I we obtain

$$\begin{aligned}
 & \mathbb{J}(r, s_0, s, r_c) \\
 & \lesssim \frac{|u'(r_c)|}{\sqrt{(u(r) - u(r_c))^2 + \varepsilon^2}} \frac{|u'(s)|}{\sqrt{(u(s) - u(r_c))^2 + \varepsilon^2}} \frac{\varepsilon^{\frac{\alpha}{2+\alpha}} |u'(s_0)|}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \\
 & \quad \times \mathbf{1}_{r \leq 1 \leq r_c} \mathbf{1}_{2r, 2s, 2s_0 \leq r_c} \frac{\chi_I(r_c) \langle s_0 \rangle^{4+\eta} s_0^{-2\eta} \beta(s_0) \left(\mathbf{1}_{s \leq 1} s^{\frac{5}{2} - \frac{\delta}{4} - 2\eta} + \mathbf{1}_{s \geq 1} s^{-\frac{3}{2} + \frac{\delta}{4} + 2\eta} \right)}{r^{\frac{1}{2} - \delta + 2\eta}} \\
 & \lesssim \varepsilon^{\frac{\alpha}{4(2+\alpha)}} \frac{|u'(r_c)|}{|u(r) - u(r_c)|^{1 - \frac{\alpha}{4(2+\alpha)}}} \frac{|u'(s)|}{|u(s) - u(r_c)|^{1 - \frac{\alpha}{4(2+\alpha)}}} \frac{|u'(s_0)|}{|u(s_0) - u(r_c)|^{2 - \frac{\alpha}{4(2+\alpha)}}} \\
 & \quad \times \mathbf{1}_{r \leq 1 \leq r_c} \mathbf{1}_{2r, 2s, 2s_0 \leq r_c} \frac{\langle s_0 \rangle^{4+\eta} s_0^{-2\eta} \beta(s_0) \left(\mathbf{1}_{s \leq 1} s^{\frac{5}{2} - \frac{\delta}{4} - 2\eta} + \mathbf{1}_{s \geq 1} s^{-\frac{3}{2} + \frac{\delta}{4} + 2\eta} \right)}{r^{\frac{1}{2} - \delta + 2\eta}} \\
 & \lesssim \varepsilon^{\frac{\alpha}{4(2+\alpha)}} \frac{1}{\langle r_c \rangle^{1 + \frac{\alpha}{2(2+\alpha)}}} \frac{s}{\langle s \rangle^{2 + \frac{\alpha}{4(2+\alpha)}}} \frac{s_0}{\langle s_0 \rangle^{\frac{\alpha}{2(2+\alpha)}}} \\
 & \quad \times \mathbf{1}_{r \leq 1 \leq r_c} \mathbf{1}_{2r, 2s, 2s_0 \leq r_c} \frac{\langle s_0 \rangle^{4+\eta} s_0^{-2\eta} \beta(s_0) \left(\mathbf{1}_{s \leq 1} s^{\frac{5}{2} - \frac{\delta}{4} - 2\eta} + \mathbf{1}_{s \geq 1} s^{-\frac{3}{2} + \frac{\delta}{4} + 2\eta} \right)}{r^{\frac{1}{2} - \delta + 2\eta}},
 \end{aligned}$$

by using properties of the cut-off $\mathbf{1}_{r \leq 1 \leq r_c} \mathbf{1}_{2r, 2s, 2s_0 \leq r_c}$, and estimates (B.12) and (B.19) in the region $|\rho - t| \geq \frac{1}{10}$ relevant here. As above, we first take care of the integral with respect to s_0

$$\int_0^\infty \frac{s_0^{1-2\eta} \langle s_0 \rangle^{4+2\eta} \beta(s_0)}{\langle s_0 \rangle^{\frac{\alpha}{2(2+\alpha)}}} ds_0 \lesssim 1$$

in view of the decay rate of β . To conclude the proof and obtain the desired vanishing as $\varepsilon \rightarrow 0$, we are left to show the boundedness of

$$\begin{aligned}
 & \int_{\mathbb{R}_+^3} \frac{1}{\langle r_c \rangle^{1 + \frac{\alpha}{2(2+\alpha)}}} \mathbf{1}_{r \leq 1 \leq r_c} \mathbf{1}_{2r, 2s \leq r_c} \frac{\left(\mathbf{1}_{s \leq 1} s^{\frac{7}{2} - \frac{\delta}{4} - 2\eta} + \mathbf{1}_{s \geq 1} s^{-\frac{5}{2} + \frac{\delta}{4} - \frac{\alpha}{4(2+\alpha)} + 2\eta} \right)}{r^{\frac{1}{2} - \delta + 2\eta}} \\
 & \quad \times |f(s)\varphi(r)| ds dr_c dr \\
 & \lesssim \|f\|_{L^2(ds)} \|\varphi\|_{L^2(dr)} \left\| \mathbf{1}_{s \leq 1} s^{\frac{7}{2} - \frac{\delta}{4} - 2\eta} + \mathbf{1}_{s \geq 1} s^{-\frac{5}{2} + \frac{\delta}{4} - \frac{\alpha}{4(2+\alpha)} + 2\eta} \right\|_{L^2(ds)} \\
 & \quad \times \left\| \frac{\mathbf{1}_{r \leq 1}}{r^{\frac{1}{2} - \delta + 2\eta}} \right\|_{L^2(dr)} \\
 & \lesssim \|f\|_{L^2(ds)} \|\varphi\|_{L^2(dr)}.
 \end{aligned}$$

This concludes the proof of the corollary, upon passing $\varepsilon \rightarrow 0$. \square

B.1.2 Identifying the Leading Order Operator Near the $s_0 = r_c$ Diagonal

In this section we consider the set

$$\left\{ |s_0 - r_c| \leq \frac{r_c}{k} \right\} \quad (\text{B.33})$$

and show that the contribution to the operator L_ε in (B.1) coming from the operators

$$\begin{aligned} L_{\varepsilon,2}[f](r) &= \int_0^\infty \int_0^\infty \int_0^\infty \frac{(u(r) - u(r_c))u'(r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} \frac{(u(s) - u(r_c))u'(s)}{(u(s) - u(r_c))^2 + \varepsilon^2} \frac{\varepsilon u'(s_0)}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \\ &\quad \times \mathbf{1}_{|s_0 - r_c| \leq \frac{r_c}{k}} (\mathfrak{B}_{\varepsilon,1}(r, s_0, r_c) - \mathfrak{B}_{\varepsilon,1}(r, r_c, r_c)) \mathfrak{B}_{\varepsilon,2}(s_0, s, r_c) f(s) \, ds ds_0 dr_c \end{aligned} \quad (\text{B.34})$$

$$\begin{aligned} L_{\varepsilon,3}[f](r) &= \int_0^\infty \int_0^\infty \int_0^\infty \frac{(u(r) - u(r_c))u'(r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} \frac{(u(s) - u(r_c))u'(s)}{(u(s) - u(r_c))^2 + \varepsilon^2} \frac{\varepsilon u'(s_0)}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \\ &\quad \times \mathbf{1}_{|s_0 - r_c| \leq \frac{r_c}{k}} \mathfrak{B}_{\varepsilon,1}(r, r_c, r_c) (\mathfrak{B}_{\varepsilon,2}(s_0, s, r_c) - \mathfrak{B}_{\varepsilon,2}(r_c, s, r_c)) f(s) \, ds ds_0 dr_c \end{aligned} \quad (\text{B.35})$$

vanish as $\varepsilon \rightarrow 0$ in $L^2(dr)$. The goal is to establish a result which is similar to Lemma B.5. Once achieved, such a result shows that

$$\begin{aligned} L_{\varepsilon,4}[f] &= \int_0^\infty \int_0^\infty \int_0^\infty \frac{(u(r) - u(r_c))u'(r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} \frac{(u(s) - u(r_c))u'(s)}{(u(s) - u(r_c))^2 + \varepsilon^2} \frac{\varepsilon u'(s_0)}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \\ &\quad \times \mathbf{1}_{|s_0 - r_c| \leq \frac{r_c}{k}} \mathfrak{B}_{\varepsilon,1}(r, r_c, r_c) \mathfrak{B}_{\varepsilon,2}(r_c, s, r_c) f(s) \, ds ds_0 dr_c \end{aligned} \quad (\text{B.36})$$

is the leading order operator with respect to ε in $L_\varepsilon[f]$. Indeed, we note that

$$L_\varepsilon[f] - L_{\varepsilon,4}[f] = L_{\varepsilon,1}[f] + L_{\varepsilon,2}[f] + L_{\varepsilon,r}[f]$$

where the right side vanishes in $L^2(dr)$ as $\varepsilon \rightarrow 0$.

Lemma B.8 *Let $\delta \in (0, \frac{1}{2})$ and assume that for some $\gamma \in (0, \frac{\delta}{4})$ we have that*

$$\mathbf{1}_{|s_0 - r_c| \leq \frac{r_c}{k}} \frac{r_c^\gamma |\mathfrak{B}_{\varepsilon,1}(r, s_0, r_c) - \mathfrak{B}_{\varepsilon,1}(r, r_c, r_c)|}{|s_0 - r_c|^\gamma} |\mathfrak{B}_{\varepsilon,2}(s_0, s, r_c)| \lesssim \mathfrak{B}_0(r, s_0, s, r_c) \quad (\text{B.37})$$

holds for some ε -independent function \mathfrak{B}_0 which obeys the bound (B.10). Then, if $f \in L^2(ds)$, we have that the operator $L_{\varepsilon,2}[f]$, defined in (B.34), vanishes as $\varepsilon \rightarrow 0$, in $L^2(dr)$.

Proof of Lemma B.8 The proof closely follows the proof of Lemma B.5. Similarly to (B.11), the lemma reduces to showing that the function

$$\begin{aligned} \mathbb{J}(r, s_0, s, r_c) := & \frac{|(u(r) - u(r_c))u'(r_c)|}{(u(r) - u(r_c))^2 + \varepsilon^2} \frac{|(u(s) - u(r_c))u'(s)|}{(u(s) - u(r_c))^2 + \varepsilon^2} \\ & \times \frac{\varepsilon |u'(s_0)| \mathbf{1}_{|s_0 - r_c| \leq \frac{r_c}{k}} \frac{|s_0 - r_c|^\gamma}{r_c^\gamma}}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \mathfrak{B}_0(r, s_0, s, r_c) \end{aligned}$$

obeys

$$\int_{\mathbb{R}_+^4} \mathbb{J}(r, s, s_0, r_c) |\varphi(r) f(s)| ds_0 ds dr dr_c \lesssim \varepsilon^{\frac{\gamma}{6}} \|\varphi\|_{L^2} \|f\|_{L^2}. \quad (\text{B.38})$$

The power law $\varepsilon^{\frac{\gamma}{6}}$ is rather arbitrary.

Case $r_c \geq 1$ Note that estimates (B.14) and (B.15) hold as is, since they are independent of the bound $|s_0 - r_c| \leq \frac{r_c}{k}$. Since $r_c \geq 1$ and $k \geq 2$, this restriction implies that $s_0 \geq \frac{r_c}{2} \geq \frac{1}{2}$, and we may thus replace (B.16) with

$$\begin{aligned} & \frac{\mathbf{1}_{r_c \geq 1} \mathbf{1}_{|s_0 - r_c| \leq \frac{r_c}{k}} \varepsilon^{1 - \frac{5\gamma}{6}} |s_0 - r_c|^\gamma |u'(s_0)|}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \\ & \lesssim \mathbf{1}_{r_c \geq 1} \mathbf{1}_{|s_0 - r_c| \leq \frac{r_c}{k}} \frac{|s_0 - r_c|^\gamma}{|u'(s_0)|^{\frac{5\gamma}{6}}} \left(\frac{|u'(s_0)|}{|u(s_0) - u(r_c)|} \right)^{1 + \frac{5\gamma}{6}} \\ & \lesssim \langle s_0 \rangle^{3\gamma} \left(\frac{\mathbf{1}_{r_c \geq 1} \mathbf{1}_{|s_0 - r_c| \geq \frac{1}{10}}}{k^\gamma \langle s_0 \rangle^{1 + \frac{\gamma}{3}}} + \frac{\mathbf{1}_{r_c \geq 1} \mathbf{1}_{|s_0 - r_c| < \frac{1}{10}}}{|s_0 - r_c|^{1 - \frac{\gamma}{6}}} \right). \quad (\text{B.39}) \end{aligned}$$

Estimate (B.39) is nearly identical to bound (B.16), and in particular once the $\langle s_0 \rangle^{3\gamma}$ is absorbed by the bound on \mathfrak{B}_0 , the resulting object is integrable in s_0 , with a bound that is $O(1)$ with respect to s, r_c , and r . All the following arguments in the proof of Lemma B.5, for the case $r_c \geq 1$ follow line by line in this case too. To avoid redundancy we omit these details.

Case $r_c < 1$ As before, the bounds (B.20) and (B.21) hold as is, since they are independent of the restriction $|s_0 - r_c| \leq \frac{r_c}{k}$. For $r_c \leq 1$, the later restriction implies $|s_0 - r_c| \leq \frac{1}{10}$ and also $s_0 \leq \frac{11}{10}$, and thus (B.22) needs to be replaced by

$$\frac{\mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s_0 - r_c| \leq \frac{r_c}{k}} \varepsilon^{1 - \frac{5\gamma}{6}} \frac{|s_0 - r_c|^\gamma}{r_c^\gamma} |u'(s_0)|}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \lesssim \frac{s_0^{\frac{5\gamma}{6}} \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s_0 - r_c| \leq \frac{1}{10}}}{r_c^\gamma |s_0 - r_c|^{1 - \frac{\gamma}{6}}} \lesssim \frac{\langle s_0 \rangle^{3\gamma} \mathbf{1}_{r_c \leq 1} \mathbf{1}_{|s_0 - r_c| \leq \frac{1}{10}}}{r_c^{2\gamma} |s_0 - r_c|^{1 - \frac{\gamma}{6}}}. \quad (\text{B.40})$$

In the last inequality we have used that $r_c \leq 1$, with the purpose of showing that the right side of (B.40) is nearly identical to the first term on the right side of (B.22). In

particular, integrating (B.40) with respect to s_0 we obtain the same estimate as in the proof of Lemma B.5. All the following arguments in the proof of Lemma B.5, for the case $r_c \leq 1$ follow line by line in this case too. To avoid redundancy we omit these details. \square

The proof of Lemma B.8 clearly implies, mutatis mutandi, also the following result:

Lemma B.9 *Let $\delta \in (0, \frac{1}{2})$ and assume that for some $\gamma \in (0, \frac{\delta}{4})$ we have that*

$$\mathbf{1}_{|s_0-r_c| \leq \frac{r_c}{k}} \left| \mathfrak{B}_{\varepsilon,1}(r, r_c, r_c) \right| \frac{r_c^\gamma \left| \mathfrak{B}_{\varepsilon,2}(s_0, s, r_c) - \mathfrak{B}_{\varepsilon,2}(r_c, s, r_c) \right|}{|s_0 - r_c|^\gamma} \lesssim \mathfrak{B}_0(r, s_0, s, r_c) \quad (\text{B.41})$$

holds for some ε -independent function \mathfrak{B}_0 which obeys the bound (B.10). Then, if $f \in L^2(ds)$, we have that the operator $L_{\varepsilon,3}[f]$, defined in (B.35), vanishes as $\varepsilon \rightarrow 0$ in $L^2(dr)$.

Remark B.10 We note here that assumptions (B.37) and (B.41) may be replaced with the dual pair of conditions

$$\mathbf{1}_{|s_0-r_c| \leq \frac{r_c}{k}} \left| \mathfrak{B}_{\varepsilon,1}(r, s_0, r_c) \right| \frac{r_c^\gamma \left| \mathfrak{B}_{\varepsilon,2}(s_0, s, r_c) - \mathfrak{B}_{\varepsilon,2}(r_c, s, r_c) \right|}{|s_0 - r_c|^\gamma} \lesssim \mathfrak{B}_0(r, s_0, s, r_c) \quad (\text{B.42})$$

$$\mathbf{1}_{|s_0-r_c| \leq \frac{r_c}{k}} \frac{r_c^\gamma \left| \mathfrak{B}_{\varepsilon,1}(r, s_0, r_c) - \mathfrak{B}_{\varepsilon,1}(r, r_c, r_c) \right|}{|s_0 - r_c|^\gamma} \left| \mathfrak{B}_{\varepsilon,2}(r_c, s, r_c) \right| \lesssim \mathfrak{B}_0(r, s_0, s, r_c) \quad (\text{B.43})$$

where $\mathfrak{B}_0(r, s_0, s, r_c)$ obeys (B.10).

Lastly, similarly to Corollary B.6, we have that:

Corollary B.11 *Let $0 \leq j \leq k-1$, and $0 \leq \ell, \ell_1, \ell_2$ be such that $\ell + \ell_1 + \ell_2 \leq j$. Assume that the functions $\mathfrak{B}_{\varepsilon,1}$ and $\mathfrak{B}_{\varepsilon,2}$ in (B.34) and (B.35) are given by either (B.3)–(B.4), or (B.5)–(B.6), where either $B_{\ell,\varepsilon}^{(1)}$ is a suitable $(2\ell_1, \ell_1)$ kernel of type II and $B_{\ell,\varepsilon}^{(2)}$ is a suitable $(2\ell_2, \ell_2)$ kernel of type I, or $B_{\ell,\varepsilon}^{(1)}$ is a suitable $(2\ell_1, \ell_1 + \eta/2)$ kernel of type I and $B_{\ell,\varepsilon}^{(2)}$ is a suitable $(2\ell_2, \ell_2 + \eta/2)$ kernel of type II. Then the operators $L_{\varepsilon,2}$ and $L_{\varepsilon,3}$ defined in (B.34) and (B.35), vanish in $L^2(dr)$ as $\varepsilon \rightarrow 0$.*

The proof of the corollary follows from the definition of being a kernel of type I, respectively II, and the proof of Corollaries B.6 and B.7. Indeed, the definitions of the kernel types precisely show that the conditions of Lemmas B.8 and B.9 are satisfied. Also here, we note that for the weight (B.5)–(B.6) a separate argument must be carried out in the region $\{r \leq 1 \leq r_c\} \cap \{2r, 2s, 2s_0 \leq r_c\}$, as in the proof of Corollary B.7.

B.1.3 Convergence of the Operator with $s_0 = r_c$

In this section we give the proof of Theorem B.1. In view of Lemmas B.5, B.8, and B.9, we have shown that under the conditions (B.8), (B.37), and (B.41), (see also Remark B.10 for a dual pair of conditions), we have that

$$\lim_{\varepsilon \rightarrow 0} \|L_\varepsilon - L_{\varepsilon,4}\|_{L^2 \rightarrow L^2} = 0 \quad (\text{B.44})$$

where the operator $L_{\varepsilon,4}$ is defined in (B.36), and may be rewritten as

$$\begin{aligned} L_{\varepsilon,4}[f] &= \int_0^\infty \int_0^\infty \frac{(u(r) - u(r_c))u'(r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} \frac{(u(s) - u(r_c))u'(s)}{(u(s) - u(r_c))^2 + \varepsilon^2} \\ &\quad \times \mathfrak{B}_{\varepsilon,1}(r, r_c, r_c) \mathfrak{B}_{\varepsilon,2}(r_c, s, r_c) \\ &\quad \times \left(\arctan \left(\frac{u(\frac{k+1}{k}r_c) - u(r_c)}{\varepsilon} \right) \right. \\ &\quad \left. + \arctan \left(\frac{u(r_c) - u(\frac{k-1}{k}r_c)}{\varepsilon} \right) \right) f(s) ds dr_c \end{aligned} \quad (\text{B.45})$$

upon performing the explicit integration in s_0 . Therefore, computing the limiting operator for L_ε reduces to computing the limiting operator for $L_{\varepsilon,4}$. For this purpose, we consider an arbitrary $\varphi \in L^2(dr)$ and compute as before

$$\begin{aligned} \langle L_{\varepsilon,4}[f](r), \varphi(r) \rangle &= \int_0^\infty \left(\arctan \left(\frac{u(\frac{k+1}{k}r_c) - u(r_c)}{\varepsilon} \right) + \arctan \left(\frac{u(r_c) - u(\frac{k-1}{k}r_c)}{\varepsilon} \right) \right) \\ &\quad \times \left(\int_0^\infty \frac{(u(r) - u(r_c))u'(r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} \mathfrak{B}_{\varepsilon,1}(r, r_c, r_c) \varphi(r) dr \right) \\ &\quad \times \left(\int_0^\infty \frac{(u(s) - u(r_c))u'(s)}{(u(s) - u(r_c))^2 + \varepsilon^2} \mathfrak{B}_{\varepsilon,2}(r_c, s, r_c) f(s) ds \right) dr_c \\ &=: \int_0^\infty \left(\arctan \left(\frac{u(\frac{k+1}{k}r_c) - u(r_c)}{\varepsilon} \right) \right. \\ &\quad \left. + \arctan \left(\frac{u(r_c) - u(\frac{k-1}{k}r_c)}{\varepsilon} \right) \right) \mathcal{M}_{\varepsilon,1}[\varphi](r_c) \mathcal{M}_{\varepsilon,2}[f](r_c) dr_c \end{aligned} \quad (\text{B.46})$$

where we have denoted the operators

$$\mathcal{M}_{\varepsilon,1}[g](r_c) = \int_0^\infty \frac{(u(r) - u(r_c))u'(r)}{(u(r) - u(r_c))^2 + \varepsilon^2} \frac{u'(r_c) \mathfrak{B}_{\varepsilon,1}(r, r_c, r_c)}{m(r_c)u'(r)} g(r) dr \quad (\text{B.47})$$

$$\mathcal{M}_{\varepsilon,2}[g](r_c) = \int_0^\infty \frac{(u(r) - u(r_c))u'(s)}{(u(s) - u(r_c))^2 + \varepsilon^2} m(r_c) \mathfrak{B}_{\varepsilon,2}(r_c, s, r_c) g(s) ds \quad (\text{B.48})$$

and the weight $m(r_c)$ is at our discretion. Assume for the moment that we can show for a fixed function $g \in L^2(\mathbb{R}_+)$ that we have

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \mathcal{M}_{\varepsilon,1}[g](r_c) &= \mathcal{M}_{0,1}[g](r_c) \\ &:= p.v. \int_0^\infty \frac{u'(r)}{u(r) - u(r_c)} \frac{u'(r_c) \mathfrak{B}_{0,1}(r, r_c, r_c)}{m(r_c) u'(r)} g(r) dr \quad \text{in } L^2(dr_c) \end{aligned} \quad (\text{B.49})$$

and

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \mathcal{M}_{\varepsilon,2}[g](r_c) &= \mathcal{M}_{0,2}[g](r_c) \\ &:= p.v. \int_0^\infty \frac{u'(s)}{u(s) - u(r_c)} m(r_c) \mathfrak{B}_{0,2}(r_c, s, r_c) g(s) ds \quad \text{in } L^2(dr_c) \end{aligned} \quad (\text{B.50})$$

where

$$\mathfrak{B}_{0,1}(r, r_c, r_c) = \lim_{\varepsilon \rightarrow 0} \mathfrak{B}_{\varepsilon,1}(r, r_c, r_c) \quad \text{and} \quad \mathfrak{B}_{0,2}(r_c, s, r_c) = \lim_{\varepsilon \rightarrow 0} \mathfrak{B}_{\varepsilon,2}(r_c, s, r_c) \quad (\text{B.51})$$

holds in a sense that is determined by Corollary B.13 below. If (B.49) and (B.50) hold, then by (B.46), the fact that $|\arctan(\cdot)| \leq \frac{\pi}{2}$, and the Dominated Convergence Theorem, we would have that

$$\begin{aligned} &\lim_{\varepsilon \rightarrow 0} \left| \langle L_{\varepsilon,4}[f](r), \varphi(r) \rangle + \pi \int_0^\infty \mathcal{M}_{0,1}[\varphi](r_c) \mathcal{M}_{0,2}[f](r_c) dr_c \right| \\ &\lesssim \lim_{\varepsilon \rightarrow 0} \pi \|\mathcal{M}_{\varepsilon,2}[f] - \mathcal{M}_{0,2}[f]\|_{L^2_{dr_c}} \|\mathcal{M}_{\varepsilon,1}[\varphi]\|_{L^2_{dr_c}} \\ &\quad + \lim_{\varepsilon \rightarrow 0} \pi \|\mathcal{M}_{0,2}[f]\|_{L^2_{dr_c}} \|\mathcal{M}_{\varepsilon,1}[\varphi] - \mathcal{M}_{0,1}[\varphi]\|_{L^2_{dr_c}} \\ &\quad + \lim_{\varepsilon \rightarrow 0} \int_0^\infty \left(\pi + \arctan \left(\frac{u(\frac{k+1}{k} r_c) - u(r_c)}{\varepsilon} \right) + \arctan \left(\frac{u(r_c) - u(\frac{k-1}{k} r_c)}{\varepsilon} \right) \right) \\ &\quad \times |\mathcal{M}_{0,1}[\varphi](r_c) \mathcal{M}_{0,2}[f](r_c)| dr_c \\ &= 0 \end{aligned} \quad (\text{B.52})$$

and therefore we have identified the limit in $L^2(dr)$ sense of $L_{\varepsilon,4}[f]$, finishing the proof of the theorem.

It remains to show that (B.49) and (B.50) hold under a suitable convergence condition of the weight. For this purpose it remains to consider the convergence as $\varepsilon \rightarrow 0$, in $L^2(dr_c)$, of the model operator

$$\mathcal{M}_\varepsilon[g](r_c) = \int_0^\infty \frac{(u(r) - u(r_c))u'(r)}{(u(r) - u(r_c))^2 + \varepsilon^2} \mathfrak{M}_\varepsilon(r, r_c) g(r) dr \quad (\text{B.53})$$

where in view of (B.47)–(B.48) the weight $\mathfrak{M}_\varepsilon(r, r_c)$ plays the role of either

$$\frac{u'(r_c)\mathfrak{B}_{\varepsilon,1}(r, r_c, r_c)}{\mathfrak{m}(r_c)u'(r)} \quad \text{or} \quad \mathfrak{m}(r_c)\mathfrak{B}_{\varepsilon,2}(r_c, r, r_c).$$

We prove the $L^2(dr_c)$ convergence of \mathcal{M}_ε under suitable conditions on the difference $\mathfrak{M}_\varepsilon(r, r_c) - \mathfrak{M}_0(r, r_c)$, and then check that these conditions hold for the specific weights that arise in (B.47)–(B.48). A simple-to-work-with set of assumptions are:

Lemma B.12 *Assume that there exists $0 < \zeta \leq \gamma < 1$ such that the following properties hold for all r, r_c :*

$$|\mathfrak{M}_\varepsilon(r, r_c) - \mathfrak{M}_0(r, r_c)| \lesssim \varepsilon^\zeta \mathfrak{M}(r, r_c) \quad (\text{B.54a})$$

$$|\mathfrak{M}_0(r, r_c)| \lesssim \mathfrak{M}(r, r_c) \quad (\text{B.54b})$$

$$\mathbf{1}_{|r-r_c| \leq \frac{r_c}{k}} |\mathfrak{M}_0(r, r_c) - \mathfrak{M}_0(r_c, r_c)| \lesssim \frac{k^\gamma |r - r_c|^\gamma}{r_c^\gamma} \mathfrak{M}(r, r_c) \quad (\text{B.54c})$$

where the ε -independent function $\mathfrak{M} \geq 0$ is defined by

$$\mathfrak{M}(r, r_c) = \frac{r_c^{\frac{3\zeta}{2}}}{\langle r_c \rangle^{3\zeta}} \left(\mathbf{1}_{r \leq r} \frac{r^{\frac{1-\zeta}{2}}}{r_c^{\frac{1-\zeta}{2}}} + \mathbf{1}_{r < r_c} \frac{\langle r \rangle^{\frac{1+\zeta}{2}}}{\langle r_c \rangle^{\frac{1+\zeta}{2}}} \right). \quad (\text{B.55})$$

Then, for any $g \in L^2(dr)$, the operator $\mathcal{M}_\varepsilon[g]$ defined in (B.53) converges as $\varepsilon \rightarrow 0$, in $L^2(dr_c)$, to $\mathcal{M}_0[g]$, defined by

$$\begin{aligned} \mathcal{M}_0[g](r_c) &= p.v. \int_0^\infty \frac{u'(r)}{u(r) - u(r_c)} \mathfrak{M}_0(r, r_c) g(r) dr \\ &=: \lim_{\varepsilon' \rightarrow 0} \int_{|u(r) - u(r_c)| \geq \varepsilon'} \frac{u'(r)}{u(r) - u(r_c)} \mathfrak{M}_0(r, r_c) g(r) dr \end{aligned} \quad (\text{B.56})$$

and this limiting operator is bounded on $L^2(dr_c)$, with norm bounded from above by k^ζ .

Under the assumptions (B.54a)–(B.55), the proof of the above lemma is direct. We give it here for the sake of completeness. However, before giving the proof, we state an immediate corollary, which yields the proof of Theorem B.1.

Lemma B.13 *Assume that there exists $0 < \zeta < \frac{\delta}{4}$, limiting weights $\mathfrak{B}_{0,1}(r, r_c, r_c)$, $\mathfrak{B}_{0,2}(r_c, r, r_c)$, and that we may choose a weight $\mathfrak{m}(r_c) > 0$, which obeys $r_c \partial_{r_c} \mathfrak{m}(r_c) / \mathfrak{m}(r_c) \lesssim k$, such that the following conditions hold:*

$$|\mathfrak{B}_{\varepsilon,1}(r, r_c, r_c) - \mathfrak{B}_{0,1}(r, r_c, r_c)| \lesssim \varepsilon^\zeta \frac{r(r_c)^4 \mathfrak{m}(r_c)}{\langle r \rangle^4 r_c} \mathfrak{M}(r, r_c) \quad (\text{B.57a})$$

$$|\mathfrak{B}_{0,1}(r, r_c, r_c)| \lesssim \frac{r(r_c)^4 \mathfrak{m}(r_c)}{\langle r \rangle^4 r_c} \mathfrak{M}(r, r_c) \quad (\text{B.57b})$$

$$\mathbf{1}_{|r-r_c| \leq \frac{r_c}{k}} |\mathfrak{B}_{0,1}(r, r_c, r_c) - \mathfrak{B}_{0,1}(r_c, r_c, r_c)| \lesssim \frac{k^\gamma |r - r_c|^\gamma}{r_c^\gamma} \frac{r(r_c)^4 \mathfrak{m}(r_c)}{\langle r \rangle^4 r_c} \mathfrak{M}(r, r_c) \quad (\text{B.57c})$$

and

$$|\mathfrak{B}_{\varepsilon,2}(r_c, s, r_c) - \mathfrak{B}_{0,2}(r_c, s, r_c)| \lesssim \varepsilon^\zeta \frac{1}{\mathfrak{m}(r_c)} \mathfrak{M}(s, r_c) \quad (\text{B.58a})$$

$$|\mathfrak{B}_{0,2}(r_c, s, r_c)| \lesssim \frac{1}{\mathfrak{m}(r_c)} \mathfrak{M}(s, r_c) \quad (\text{B.58b})$$

$$\mathbf{1}_{|s-r_c| \leq \frac{r_c}{k}} |\mathfrak{B}_{0,2}(r_c, s, r_c) - \mathfrak{B}_{0,2}(r_c, r_c, r_c)| \lesssim \frac{k^\gamma |s - r_c|^\gamma}{r_c^\gamma} \frac{1}{\mathfrak{m}(r_c)} \mathfrak{M}(s, r_c) \quad (\text{B.58c})$$

where the function $\mathfrak{M}(r, r_c)$ is defined by (B.55) above. Then we have that

$$\lim_{\varepsilon \rightarrow 0} L_\varepsilon[f](r) = L_0[f](r) \text{ as bounded operators } L^2 \rightarrow L^2,$$

where the operator L_0 is defined in (B.2). Moreover, the operator norm of $L_0[f]$ in this space is bounded by k^ζ .

Lemma B.13 is a direct consequence of Lemma B.12, which implies that (B.52) holds, and we conclude using (B.44). We omit these details. It thus remains to prove Lemma B.12.

Proof of Lemma B.12 Let $\chi_c(r, r_c)$ be defined as in (4.3). For any test function $\varphi \in L^2(dr_c)$, we may decompose

$$\begin{aligned} \langle \mathcal{M}_\varepsilon[g], \varphi \rangle &= \int_0^\infty \int_0^\infty \frac{(u(r) - u(r_c))u'(r)}{(u(r) - u(r_c))^2 + \varepsilon^2} (\mathfrak{M}_\varepsilon(r, r_c) - \mathfrak{M}_0(r, r_c)) g(r) \varphi(r_c) dr dr_c \\ &\quad + \int_0^\infty \int_0^\infty \frac{u'(r)(u(r) - u(r_c))(1 - \chi_c(r, r_c)) \mathfrak{M}_0(r, r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} g(r) \varphi(r_c) dr dr_c \\ &\quad + \int_0^\infty \int_0^\infty \frac{u'(r)(u(r) - u(r_c)) \chi_c(r, r_c) (\mathfrak{M}_0(r, r_c) - \mathfrak{M}_0(r_c, r_c))}{(u(r) - u(r_c))^2 + \varepsilon^2} \\ &\quad \times g(r) \varphi(r_c) dr dr_c \\ &\quad + \int_0^\infty \int_0^\infty \frac{u'(r)(u(r) - u(r_c)) \chi_c(r, r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} g(r) \mathfrak{M}_0(r_c, r_c) \varphi(r_c) dr dr_c \\ &=: \langle \mathcal{M}_\varepsilon^{(0)}[g], \varphi \rangle + \langle \mathcal{M}_\varepsilon^{(1)}[g], \varphi \rangle + \langle \mathcal{M}_\varepsilon^{(2)}[g], \varphi \rangle + \langle \mathcal{M}_\varepsilon^{(3)}[g], \varphi \rangle. \end{aligned} \quad (\text{B.59})$$

Next compute the limit of the first three terms on the right side of (B.59) vanish as $\varepsilon \rightarrow 0$.

By (B.12) and (B.54)–(B.55), we have

$$\begin{aligned}
 & \left| \left\langle \mathcal{M}_\varepsilon^{(0)}[g], \varphi \right\rangle \right| \\
 & \lesssim k\varepsilon^{\frac{\zeta}{2}} \int_0^\infty \int_0^\infty \frac{|u'(r)|}{|u(r) - u(r_c)|^{1-\frac{\zeta}{2}}} \mathfrak{M}(r, r_c) |g(r)\varphi(r_c)| dr dr_c \\
 & \lesssim k\varepsilon^{\frac{\zeta}{2}} \int_0^\infty \int_0^\infty \frac{r^{\frac{\zeta}{2}} r_c^{\frac{3\zeta}{2}}}{\langle r \rangle^{2\zeta} \langle r_c \rangle^{3\zeta}} \left[\frac{\mathbf{1}_{|r-r_c| \leq \frac{1}{10}}}{|r-r_c|^{1-\frac{\zeta}{2}}} + \frac{\mathbf{1}_{|r-r_c| \geq \frac{1}{10}}}{\langle r \rangle^{1-\frac{\zeta}{2}}} \right] \\
 & \quad \times \left[\mathbf{1}_{r_c \leq r} \frac{r^{\frac{1-\zeta}{2}}}{r_c^{\frac{1-\zeta}{2}}} + \mathbf{1}_{r < r_c} \frac{\langle r \rangle^{\frac{1+\zeta}{2}}}{\langle r_c \rangle^{\frac{1+\zeta}{2}}} \right] |g(r)\varphi(r_c)| dr dr_c \\
 & \lesssim k\varepsilon^{\frac{\zeta}{2}} \int_0^\infty \int_0^\infty \left[\frac{\mathbf{1}_{|r-r_c| \leq \frac{1}{10}}}{|r-r_c|^{1-\frac{\zeta}{2}}} + \frac{1}{\langle r \rangle^{\frac{1+\zeta}{2}} \langle r_c \rangle^{\frac{1+\zeta}{2}}} + \frac{\mathbf{1}_{r_c \leq 1}}{\langle r \rangle^{\frac{1+\zeta}{2}} r_c^{\frac{1-\zeta}{2}}} + \frac{\mathbf{1}_{r_c \leq 1} \mathbf{1}_{r \leq 1}}{r^{\frac{1-\zeta}{2}} r_c^{\frac{1-\zeta}{2}}} \right] \\
 & \quad \times |g(r)\varphi(r_c)| dr dr_c \\
 & \lesssim k\varepsilon^{\frac{\zeta}{2}} \|g\|_{L^2} \|\varphi\|_{L^2} \left(\left\| \frac{\mathbf{1}_{|\rho| \leq \frac{1}{10}}}{|\rho|^{1-\frac{\zeta}{2}}} \right\|_{L^1_\rho} + \left\| \frac{1}{\langle r \rangle^{\frac{1+\zeta}{2}} \langle r_c \rangle^{\frac{1+\zeta}{2}}} \right\|_{L^2_{r,r_c}} + \left\| \frac{\mathbf{1}_{r_c \leq 1}}{\langle r \rangle^{\frac{1+\zeta}{2}} r_c^{\frac{1-\zeta}{2}}} \right\|_{L^2_{r,r_c}} \right. \\
 & \quad \left. + \left\| \frac{\mathbf{1}_{r_c \leq 1} \mathbf{1}_{r \leq 1}}{r^{\frac{1-\zeta}{2}} r_c^{\frac{1-\zeta}{2}}} \right\|_{L^2_{r,r_c}} \right) \\
 & \lesssim k\varepsilon^{\frac{\zeta}{2}} \|g\|_{L^2} \|\varphi\|_{L^2} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.
 \end{aligned} \tag{B.60}$$

Similarly, using (B.12) and (B.54b)–(B.54c), which allows us to replace γ with ζ in (B.54c) since $\zeta \leq \gamma$, we get

$$\begin{aligned}
 & \left| \left\langle \mathcal{M}_\varepsilon^{(2)}[g], \varphi \right\rangle - \left\langle \mathcal{M}_0^{(2)}[g], \varphi \right\rangle \right| \\
 & \lesssim \varepsilon^{\frac{\zeta}{2}} k^\zeta \int_0^\infty \int_0^\infty \frac{|u'(r)| |r-r_c|^\zeta \mathbf{1}_{|r-r_c| \leq \frac{r_c}{k}}}{r_c^\zeta |u(r) - u(r_c)|^{1+\frac{\zeta}{2}}} \mathfrak{M}(r, r_c) |g(r)\varphi(r_c)| dr dr_c \\
 & \lesssim \varepsilon^{\frac{\zeta}{2}} k^\zeta \int_0^\infty \int_0^\infty \frac{|r-r_c|^\zeta \langle r \rangle^{2\zeta} \mathbf{1}_{|r-r_c| \leq \frac{r_c}{k}}}{r_c^\zeta r^{\frac{\zeta}{2}}} \left[\frac{\mathbf{1}_{|r-r_c| \leq \frac{1}{10}}}{|r-r_c|^{1+\frac{\zeta}{2}}} + \frac{\mathbf{1}_{|r-r_c| \geq \frac{1}{10}}}{\langle r \rangle^{1+\frac{\zeta}{2}}} \right] \\
 & \quad \times \left[\mathbf{1}_{r_c \leq r} \frac{r^{\frac{1-\zeta}{2}}}{r_c^{\frac{1-\zeta}{2}}} + \mathbf{1}_{r < r_c} \frac{\langle r \rangle^{\frac{1+\zeta}{2}}}{\langle r_c \rangle^{\frac{1+\zeta}{2}}} \right] |g(r)\varphi(r_c)| dr dr_c \\
 & \lesssim \varepsilon^{\frac{\zeta}{2}} k^\zeta \int_0^\infty \int_0^\infty \left[\frac{\mathbf{1}_{|r-r_c| \leq \frac{1}{10}}}{|r-r_c|^{1-\frac{\zeta}{2}}} + \frac{\mathbf{1}_{\frac{1}{10} \leq |r-r_c| \leq \frac{r_c}{k}}}{\langle r \rangle^{\frac{1+\zeta}{2}} \langle r_c \rangle^{\frac{1+\zeta}{2}}} \right] |g(r)\varphi(r_c)| dr dr_c \\
 & \lesssim \varepsilon^{\frac{\zeta}{2}} k^\zeta \|g\|_{L^2} \|\varphi\|_{L^2} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.
 \end{aligned} \tag{B.61}$$

Moreover, a bound similar to the above shows that

$$\left| \langle \mathcal{M}_0^{(2)}[g], \varphi \rangle \right| \lesssim k^\zeta \|g\|_{L^2} \|\varphi\|_{L^2} \quad (\text{B.62})$$

so that the limiting operator $\mathcal{M}_0^{(2)}$ is bounded on L^2 .

In a similar fashion, by using (B.12), (B.19), and (B.54b), we arrive at

$$\begin{aligned} & \left| \langle \mathcal{M}_\varepsilon^{(1)}[g], \varphi \rangle - \langle \mathcal{M}_0^{(1)}[g], \varphi \rangle \right| \\ & \lesssim \varepsilon^{\frac{\zeta}{2}} \int_0^\infty \int_0^\infty \frac{|u'(r)| \mathbf{1}_{|r-r_c| \geq \frac{r_c}{k}} \mathfrak{M}(r, r_c) |g(r) \varphi(r_c)| dr dr_c}{|u(r) - u(r_c)|^{1+\frac{\zeta}{2}}} \\ & \lesssim \varepsilon^{\frac{\zeta}{2}} \int_0^\infty \int_0^\infty \frac{\langle r \rangle^{2\zeta} \mathbf{1}_{|r-r_c| \geq \frac{r_c}{k}}}{r^{\frac{\zeta}{2}}} \left[\frac{\mathbf{1}_{|r-r_c| \leq \frac{1}{10}}}{|r-r_c|^{1+\frac{\zeta}{2}}} \left(\frac{\mathbf{1}_{r \leq 1} r^{1+\frac{\zeta}{2}}}{(r+r_c)^{1+\frac{\zeta}{2}}} + \mathbf{1}_{r \geq 1} \right) \right. \\ & \quad \left. + \mathbf{1}_{|r-r_c| \geq \frac{1}{10}} \left(\mathbf{1}_{r \leq 1} r^{1+\frac{\zeta}{2}} + \frac{\mathbf{1}_{r \geq 1}}{\langle r \rangle^{1+\frac{\zeta}{2}}} \right) \right] \\ & \quad \times \frac{r_c^{\frac{3\zeta}{2}}}{\langle r_c \rangle^{3\zeta}} \left[\mathbf{1}_{r_c \leq r} \frac{r^{\frac{1-\zeta}{2}}}{r_c^{\frac{1-\zeta}{2}}} + \mathbf{1}_{r < r_c} \frac{\langle r \rangle^{\frac{1+\zeta}{2}}}{\langle r_c \rangle^{\frac{1+\zeta}{2}}} \right] |g(r) \varphi(r_c)| dr dr_c \\ & \lesssim \varepsilon^{\frac{\zeta}{2}} \int_0^\infty \int_0^\infty \left[\frac{\mathbf{1}_{\frac{r_c}{k} \leq |r-r_c| \leq \frac{1}{10}}}{|r-r_c|^{1-\frac{\zeta}{2}}} + \frac{1}{\langle r \rangle^{\frac{1+\zeta}{2}} \langle r_c \rangle^{\frac{1+\zeta}{2}}} + \frac{\mathbf{1}_{r \leq 1} \mathbf{1}_{r_c \leq 1}}{r^{\frac{1-\zeta}{2}} r_c^{\frac{1-\zeta}{2}}} + \frac{\mathbf{1}_{r_c \leq 1}}{\langle r \rangle^{\frac{1+\zeta}{2}} r_c^{\frac{1-\zeta}{2}}} \right] \\ & \quad \times |g(r) \varphi(r_c)| dr dr_c \\ & \lesssim \varepsilon^{\frac{\zeta}{2}} \|g\|_{L^2} \|\varphi\|_{L^2} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0. \end{aligned} \quad (\text{B.63})$$

Moreover, a bound similar to the above shows that

$$\left| \langle \mathcal{M}_0^{(1)}[g], \varphi \rangle \right| \lesssim \|g\|_{L^2} \|\varphi\|_{L^2} \quad (\text{B.64})$$

so that the limiting operator $\mathcal{M}_0^{(1)}$ is bounded on L^2 .

Thus, by (B.60), (B.63)–(B.64), and (B.61)–(B.62), we know that the first three terms on the right side of (B.59) converge as $\varepsilon \rightarrow 0$ to bounded operators on $L^2(dr_c)$.

Lastly, we need to consider the fourth operator on the right side of (B.59), namely

$$\mathcal{M}_\varepsilon^{(3)}[g](r_c) = \mathfrak{M}_0(r_c, r_c) \int_0^\infty \frac{u'(r)(u(r) - u(r_c))}{(u(r) - u(r_c))^2 + \varepsilon^2} \chi_c(r, r_c) g(r) dr.$$

We note that since $u: [0, \infty) \rightarrow (0, u(0)]$ is a bijection, hence upon making a change of variables, we have that

$$\|f(r_c)\|_{L^2(dr_c)} = \left\| \frac{f(u^{-1}(c))}{\sqrt{|u'(u^{-1}(c))|}} \right\|_{L^2(dc)} \quad (\text{B.65})$$

so that we may change variables in the formula for $\mathcal{M}_\varepsilon^{(3)}[g](r_c)$ as

$$\begin{aligned}\mathcal{M}_\varepsilon^{(3)}[g](r_c) &= \mathfrak{M}_0(r_c, r_c) \sqrt{|u'(r_c)|} \int_0^\infty \frac{u'(r)(u(r) - u(r_c))}{(u(r) - u(r_c))^2 + \varepsilon^2} \\ &\quad \times \left(\frac{\sqrt{|u'(r)|} \chi_c(r, r_c)}{\sqrt{|u'(r_c)|}} \right) \frac{g(r)}{\sqrt{|u'(r)|}} dr \\ &= \sqrt{|u'(u^{-1}(c))|} \mathfrak{M}_0(u^{-1}(c), u^{-1}(c)) \\ &\quad \times \int_0^{u(0)} \frac{(y - c)}{(y - c)^2 + \varepsilon^2} \left(\frac{\sqrt{|u'(u^{-1}(y))|} \chi_c(u^{-1}(y), u^{-1}(c))}{\sqrt{|u'(u^{-1}(c))|}} \right) \\ &\quad \times \frac{g(u^{-1}(y))}{\sqrt{|u'(u^{-1}(y))|}} dy =: \sqrt{|u'(u^{-1}(c))|} \mathcal{M}_\varepsilon^{(4)} \left[\frac{g(u^{-1}(\cdot))}{\sqrt{|u'(u^{-1}(\cdot))|}} \right] (c).\end{aligned}$$

Thus, if we can show that the operator $\mathcal{M}_\varepsilon^{(4)}$ is bounded on $L^2((0, u(0)))$, uniformly in ε , and that it has a certain limit in this space, then by applying (B.65) twice, we have proven that

$$\begin{aligned}&\left\| \mathcal{M}_\varepsilon^{(3)}[g] - \mathcal{M}_0^{(3)}[g] \right\|_{L^2(dr_c)} \\ &= \left\| \mathcal{M}_\varepsilon^{(4)} \left[\frac{g(u^{-1}(\cdot))}{\sqrt{|u'(u^{-1}(\cdot))|}} \right] - \mathcal{M}_0^{(4)} \left[\frac{g(u^{-1}(\cdot))}{\sqrt{|u'(u^{-1}(\cdot))|}} \right] \right\|_{L^2(dc)} \\ &\leq \left\| \mathcal{M}_\varepsilon^{(4)} - \mathcal{M}_0^{(4)} \right\|_{L^2(dc) \rightarrow L^2(dc)} \left\| \frac{g(u^{-1}(\cdot))}{\sqrt{|u'(u^{-1}(\cdot))|}} \right\|_{L^2(dc)} \\ &= \left\| \mathcal{M}_\varepsilon^{(4)} - \mathcal{M}_0^{(4)} \right\|_{L^2(dc) \rightarrow L^2(dc)} \|g\|_{L^2(dr_c)}.\end{aligned}\tag{B.66}$$

Therefore, in order to conclude the proof of Lemma B.12, it remains to show that

$$\mathcal{M}_\varepsilon^{(4)} - \mathcal{M}_0^{(4)} \rightarrow 0 \quad \text{in } L^2(dc) \quad \text{as } \varepsilon \rightarrow 0,\tag{B.67}$$

and that the limiting operator $\mathcal{M}_0^{(4)}$, naturally defined as

$$\begin{aligned}\mathcal{M}_0^{(4)}[\psi](c) &= \mathfrak{M}_0(u^{-1}(c), u^{-1}(c)) \\ &\quad \times p.v. \int_0^{u(0)} \left(\frac{\sqrt{|u'(u^{-1}(y))|} \chi_c(u^{-1}(y), u^{-1}(c))}{\sqrt{|u'(u^{-1}(c))|}} \right) \frac{\psi(y)}{y - c} dy,\end{aligned}\tag{B.68}$$

obeys

$$\left\| \mathcal{M}_0^{(4)}[\psi] \right\|_{L^2(dc)} \lesssim \|\psi\|_{L^2(dc)}\tag{B.69}$$

Indeed, once this is achieved, we may change variables $c \mapsto r_c$, to obtain that

$$\begin{aligned}\mathcal{M}_\varepsilon^{(3)}[g](r_c) &\rightarrow \mathcal{M}_0^{(3)}[g](r_c) = \mathfrak{M}_0(r_c, r_c) p.v. \int_0^\infty \frac{u'(r)}{u(r) - u(r_c)} \chi_c(r, r_c) g(r) dr \\ &:= \mathfrak{M}_0(r_c, r_c) \lim_{\varepsilon' > 0} \int_0^\infty \frac{u'(r)}{u(r) - u(r_c)} \chi_c(r, r_c) g(r) dr\end{aligned}$$

in $L^2(dr_c)$, as $\varepsilon \rightarrow 0$. Combining the above with (B.60), (B.61)–(B.62), (B.63)–(B.64), (B.65), (B.66), and (B.67)–(B.69), we obtain that

$$\begin{aligned}\mathcal{M}_\varepsilon[g](r_c) &\rightarrow \mathcal{M}_0^{(1)}[g](r_c) + \mathcal{M}_0^{(2)}[g](r_c) + \mathcal{M}_0^{(3)}[g](r_c) \\ &= p.v. \int_0^\infty \frac{u'(r)}{u(r) - u(r_c)} \mathfrak{M}_0(r, r_c) g(r) dr\end{aligned}$$

and that the limiting operator is bounded on L^2 , as desired.

In order to prove (B.67) we note that for any $c > 0$ we have

$$\lim_{y \rightarrow c} \frac{\sqrt{|u'(u^{-1}(y))|} \chi_c(u^{-1}(y), u^{-1}(c))}{\sqrt{|u'(u^{-1}(c))|}} = 1$$

which allows us to rewrite

$$\begin{aligned}\mathcal{M}_\varepsilon^{(4)}[\psi](c) &= \mathfrak{M}_0(u^{-1}(c), u^{-1}(c)) \int_0^{u(0)} \left(\frac{\sqrt{|u'(u^{-1}(y))|} \chi_c(u^{-1}(y), u^{-1}(c))}{\sqrt{|u'(u^{-1}(c))|}} - 1 \right) \\ &\quad \times \frac{\psi(y)(y - c)}{(y - c)^2 + \varepsilon^2} dy \\ &\quad + \mathfrak{M}_0(u^{-1}(c), u^{-1}(c)) p.v. \int_0^{u(0)} \frac{\psi(y)(y - c)}{(y - c)^2 + \varepsilon^2} dy \\ &=: \mathcal{M}_\varepsilon^{(4,1)}[\psi](c) + \mathcal{M}_\varepsilon^{(4,2)}[\psi](c).\end{aligned}\tag{B.70}$$

The convergence in $L^2(dc)$ of the second part in (B.70), namely

$$\mathcal{M}_\varepsilon^{(4,2)}[\psi](c) \rightarrow \mathcal{M}_0^{(4,2)}[\psi](c) = \mathfrak{M}_0(u^{-1}(c), u^{-1}(c)) \mathcal{H}(\mathbf{1}_{(0, u(0)]} \psi)(c)$$

where \mathcal{H} is the Hilbert transform is classical. Moreover, since \mathcal{H} is unitary on L^2 , and since by (B.54b) and (B.55) we have

$$|\mathfrak{M}_0(u^{-1}(c), u^{-1}(c))| \lesssim \mathfrak{M}(r_c, r_c) \lesssim 1,$$

we have that

$$\|\mathcal{M}_0^{(4,2)}[\psi]\|_{L^2} \lesssim \|\psi\|_{L^2},$$

which is consistent with (B.69). The convergence in $L^2(dc)$ of the first part in (B.70) follows since this term is not a principle value anymore, and we have that for any test function $\varphi \in L^2(dc)$,

$$\begin{aligned}
 & \left| \langle \mathcal{M}_\varepsilon^{(4,1)}[\psi](c) - \mathcal{M}_0^{(4,1)}[\psi](c), \varphi(c) \rangle \right| \\
 & \lesssim \varepsilon^{\frac{\zeta}{2}} \int_0^{u(0)} \int_0^{u(0)} \min \left\{ u^{-1}(c)^{\frac{3\zeta}{2}}, u^{-1}(c)^{-\frac{3\zeta}{2}} \right\} \\
 & \quad \times \left| \frac{\sqrt{|u'(u^{-1}(y))|} \chi_c(u^{-1}(y), u^{-1}(c))}{\sqrt{|u'(u^{-1}(c))|}} - 1 \right| \frac{|\psi(y)| |\varphi(c)|}{|y - c|^{1+\frac{\zeta}{2}}} dy \\
 & \lesssim \varepsilon^{\frac{\zeta}{2}} \mathcal{K} \int_0^{u(0)} \int_0^{u(0)} \frac{|\psi(y)| |\varphi(c)|}{|y - c|^{1-\frac{\zeta}{4}}} dy \\
 & \lesssim \varepsilon^{\frac{\zeta}{2}} \mathcal{K} \|\psi\|_{L^2} \|\varphi\|_{L^2} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0,
 \end{aligned}$$

where

$$\begin{aligned}
 \mathcal{K} &= \sup_{y, c \in (0, u(0))} \left| \min \left\{ u^{-1}(c)^{\frac{3\zeta}{2}}, u^{-1}(c)^{-\frac{3\zeta}{2}} \right\} \frac{\frac{\sqrt{|u'(u^{-1}(y))|} \chi_c(u^{-1}(y), u^{-1}(c))}{\sqrt{|u'(u^{-1}(c))|}} - 1}{|y - c|^{\frac{3\zeta}{4}}} \right| \\
 &= \sup_{r, r_c \in [0, \infty)} \left| \frac{r_c^{\frac{3\zeta}{2}}}{\langle r_c \rangle^{3\zeta}} \frac{\frac{\sqrt{|u'(r)|} \chi_c(r, r_c)}{\sqrt{|u'(r_c)|}} - 1}{|u(r) - u(r_c)|^{\frac{3\zeta}{4}}} \right| \lesssim k^\zeta. \tag{B.71}
 \end{aligned}$$

To see that (B.71) holds, first note that

$$\frac{\mathbf{1}_{|r-r_c| \geq \frac{1}{10}}}{|u(r) - u(r_c)|} \lesssim \frac{1}{|u'(r_c)|} \min\{r_c, r_c^{-1}\} \lesssim \langle r_c \rangle^2,$$

and

$$\frac{\mathbf{1}_{|r-r_c| \leq \frac{1}{10}}}{|u(r) - u(r_c)|} \lesssim \frac{1}{|u'(r_c)|} \frac{1}{|r - r_c|} \lesssim \frac{\langle r_c \rangle^4}{r_c |r - r_c|},$$

from which it follows that

$$\frac{1 - \chi_c(r, r_c)}{|u(r) - u(r_c)|^{\frac{3\zeta}{4}}} \lesssim \langle r_c \rangle^{\frac{3\zeta}{2}} + k^{\frac{3\zeta}{4}} \frac{\langle r_c \rangle^{\frac{3\zeta}{2}}}{r_c^{\frac{3\zeta}{2}}}.$$

This proves (B.71) for the region $\chi_c(r, r_c) = 0$. Second, due to the regularity of u' , we have that

$$\begin{aligned} \chi_c(r, r_c) \frac{\frac{\sqrt{|u'(r)|}}{\sqrt{|u'(r_c)|}} - 1}{|u(r) - u(r_c)|^{\frac{3\zeta}{4}}} &\lesssim \frac{\chi_c(r, r_c) |u'(r) - u'(r_c)|}{|u'(r_c)|^{\frac{1}{2} + \frac{3\zeta}{4}} (\sqrt{|u'(r_c)|} + \sqrt{|u'(r)|})} \\ &\times \left(\frac{\mathbf{1}_{|r-r_c| \leq \frac{1}{10}}}{|r-r_c|^{\frac{3\zeta}{4}}} + \frac{\mathbf{1}_{|r-r_c| \geq \frac{1}{10}} r_c^{\frac{3\zeta}{4}}}{\langle r_c \rangle^{\frac{3\zeta}{2}}} \right) \lesssim \frac{\langle r_c \rangle^{\frac{3\zeta}{2}}}{r_c^{\frac{3\zeta}{2}}}, \end{aligned}$$

which proves (B.71) for the region $\chi_c(r, r_c) \neq 0$.

To conclude, the same argument as above shows that

$$\left| \langle \mathcal{M}_0^{(4,1)}[\psi], \varphi \rangle \right| \lesssim \mathcal{K} \|\psi\|_{L^2} \|\varphi\|_{L^2},$$

which establishes the boundedness of $\mathcal{M}_0^{(4,1)}$ on L^2 , and thus (B.69) holds. This finishes the proof of the lemma. \square

To conclude this section, we show that the conditions of Lemma B.13 hold for the specific weights we need to considering. First, the next Corollary concludes the proof of Theorem B.2.

Corollary B.14 *Let $0 \leq j \leq k-1$, and $0 \leq \ell, \ell_1, \ell_2$ be such that $\ell + \ell_1 + \ell_2 \leq j$. Let $0 < \zeta \leq \frac{\delta}{6}$. Consider the weights*

$$\begin{aligned} \mathfrak{B}_{\varepsilon,1}(r, r_c, r_c) &= \chi_1(r, r_c) \frac{\beta(r) \min(r^2, r^{-2})^j}{u'(r_c) w_{F,\delta}(r)} B_{\ell,\varepsilon}^{(1)}(r, r_c, r_c) \\ \mathfrak{B}_{\varepsilon,2}(r_c, s, r_c) &= \frac{\beta(r_c) w_{F,\frac{\delta}{4}+2\ell}(s)}{u'(s)} B_{\ell,\varepsilon}^{(2)}(r_c, s, r_c) \end{aligned}$$

where $B_{\ell,\varepsilon}^{(1)}$ is a suitable $(2\ell_1, \ell_1 + \eta/2)$ kernel of type I or II, and $B_{\ell,\varepsilon}^{(2)}$ is a suitable $(2\ell_2, \ell_2 + \eta/2)$ kernel of type I or II. Then, letting

$$\mathfrak{m}(r_c) = \frac{r_c^{\frac{3\zeta}{2}}}{\langle r_c \rangle^{3\zeta} w_{F,\frac{\delta}{2}}(r_c) (\max\{r_c^2, r_c^{-2}\})^{j-\ell_1-\eta/2}}, \quad (\text{B.72})$$

the conditions (B.57)–(B.58) of Lemma B.13, with \mathfrak{M} replaced by $k^{2\ell_1+2\ell_2}\mathfrak{M}$, and where the limiting weights are obtained by passing $\varepsilon \rightarrow 0$ in $B_{\ell,\varepsilon}^{(1)}$ and $B_{\ell,\varepsilon}^{(2)}$. Also, the operator norm of L_0 on $L^2(dr)$ is bounded as $k^{\zeta+2\ell_1+2\ell_2}$.

Proof of Corollary B.14 In view of the established pointwise (and in a Hölder class near the diagonal) convergence properties of type I and type II kernels $B_{\ell,\varepsilon}^{(1)}(r, r_c, r_c) \rightarrow$

$B_{\ell,0}^{(1)}(r, r_c, r_c)$ and $B_{\ell,\varepsilon}^{(2)}(r_c, s, r_c) \rightarrow B_{\ell,0}^{(2)}(r_c, s, r_c)$ as $\varepsilon \rightarrow 0$ (cf. Definition 6.7 and Definition 6.8), it is clear that checking conditions (B.57b) and (B.58b) is sufficient. Indeed, checking conditions (B.57a) and (B.57c), respectively (B.58a) and (B.58c), follows mutatis mutandis. For simplicity we only treat the case $\eta = 0$. The case $0 < \eta \ll 1$ is treated similarly. First, we verify (B.57b). We recall that

$$\begin{aligned} |B_{\ell,0}^{(1)}(r, r_c, r_c)| &\lesssim |u'(r_c)| \mathcal{B}(r, r_c) \mathbb{K}(r, r_c, r_c) \mathcal{L}_{2\ell_1, \ell_1}(r, r_c) \\ &= |u'(r_c)| \left(\mathbf{1}_{r_c < r} \frac{r_c^{k-\frac{1}{2}}}{r^{k-\frac{1}{2}}} + \mathbf{1}_{r < r_c} \frac{r^{k+\frac{1}{2}}}{r_c^{k+\frac{1}{2}}} \right) \langle r_c \rangle^4 \\ &\quad \times \left(\mathbf{1}_{r_c > 1} + \mathbf{1}_{r \leq r_c \leq 1} + \frac{r_c^2}{r^2} \mathbf{1}_{r_c < r < 1} + r_c^2 \mathbf{1}_{r_c < 1 < r} \right) \\ &\quad \times k^{2\ell_1} \left(\max \left\{ \frac{1}{r^2}, r^2, \frac{1}{r_c^2}, r_c^2 \right\} \right)^{\ell_1}, \end{aligned}$$

from which it follows, using the definition of $w_{F,\delta}$ in (1.10c), that

$$\begin{aligned} &\frac{r_c \langle r \rangle^4}{\langle r_c \rangle^4 \mathfrak{m}(r_c) r} |\mathfrak{B}_{0,1}(r, r_c, r_c)| \\ &\lesssim (\mathbf{1}_{\frac{r_c}{2} \leq r \leq 1} + \mathbf{1}_{r \geq 1}) \frac{r_c^{1-\frac{3\zeta}{2}} \langle r_c \rangle^{3\zeta} w_{F,\frac{\delta}{2}}(r_c)}{r \langle r \rangle^4 w_{F,\delta}(r)} \left(\mathbf{1}_{r_c < r} \frac{r_c^{k-\frac{1}{2}}}{r^{k-\frac{1}{2}}} + \mathbf{1}_{r < r_c} \frac{r^{k+\frac{1}{2}}}{r_c^{k+\frac{1}{2}}} \right) \\ &\quad \times \left(\mathbf{1}_{r_c > 1} + \mathbf{1}_{r \leq r_c \leq 1} + \frac{r_c^2}{r^2} \mathbf{1}_{r_c < r < 1} + r_c^2 \mathbf{1}_{r_c < 1 < r} \right) k^{2\ell_1} \\ &\quad \times \frac{\left(\max \left\{ \frac{1}{r^2}, r^2, \frac{1}{r_c^2}, r_c^2 \right\} \right)^{\ell_1} \left(\max \left\{ \frac{1}{r_c^2}, r_c^2 \right\} \right)^{j-\ell_1}}{\left(\max \left\{ r^2, \frac{1}{r^2} \right\} \right)^j} \\ &\lesssim k^{2\ell_1} \frac{r_c^{\frac{3\zeta}{2}}}{\langle r_c \rangle^{3\zeta}} \left[\mathbf{1}_{\frac{r_c}{2} < r < r_c < 1} + \mathbf{1}_{\frac{r_c}{2} < r < 1 < r_c} + \mathbf{1}_{1 < r < r_c} \frac{r^{2k-2j+\frac{1}{2}-\delta}}{r_c^{2k-2j+\frac{1}{2}+4-\frac{\delta}{2}-3\zeta}} \right. \\ &\quad + \mathbf{1}_{r_c < r < 1} \frac{r_c^{2k-2j+\frac{1}{2}+5-\frac{\delta}{2}-3\zeta}}{r^{2k-2j+\frac{1}{2}+5-\delta}} + \mathbf{1}_{r_c < 1 < r} r^{\frac{1}{2}-\delta} r_c^{2k-2j+\frac{1}{2}+5-\frac{\delta}{2}-3\zeta} \\ &\quad \left. + \mathbf{1}_{1 < r_c < r} \frac{r^{\frac{1}{2}-\delta}}{r_c^{\frac{1}{2}+4-\frac{\delta}{2}-3\zeta}} \right] \\ &\lesssim k^{2\ell_1} \frac{r_c^{\frac{3\zeta}{2}}}{\langle r_c \rangle^{3\zeta}} \left[\mathbf{1}_{r_c \leq r} \frac{r^{\frac{1-\zeta}{2}}}{r_c^{\frac{1-\zeta}{2}}} + \mathbf{1}_{r < r_c} \frac{\langle r \rangle^{\frac{1+\zeta}{2}}}{\langle r_c \rangle^{\frac{1+\zeta}{2}}} \right] = k^{2\ell_1} \mathfrak{M}(r, r_c) \end{aligned}$$

in view of our assumption on ζ . Here we have used that $j \leq k$. Thus, (B.57b) holds with the above definition of $\mathfrak{B}_{\varepsilon,1}(r, r_c, r_c)$, upon multiplying \mathfrak{M} by $k^{2\ell_1}$. Next, we verify (B.58b). We recall that

$$\begin{aligned}
|B_{\ell,0}^{(2)}(r_c, s, r_c)| &\lesssim |u'(s)| \mathcal{B}(r_c, s) \mathbb{K}(r_c, s, r_c) \mathcal{L}_{2\ell_2, \ell_2}(r_c, s) \\
&= |u'(s)| \left(\mathbf{1}_{s < r_c} \frac{s^{k-\frac{1}{2}}}{r_c^{k-\frac{1}{2}}} + \mathbf{1}_{r_c < s} \frac{r_c^{k+\frac{1}{2}}}{s^{k+\frac{1}{2}}} \right) \langle s \rangle^4 \\
&\quad \times \left(\mathbf{1}_{r_c > 1} + \mathbf{1}_{s \leq r_c \leq 1} + \frac{r_c^2}{s^2} \mathbf{1}_{r_c < s < 1} + r_c^2 \mathbf{1}_{r_c < 1 < s} \right) \\
&\quad \times k^{2\ell_2} \left(\max \left\{ \frac{1}{s^2}, s^2, \frac{1}{r_c^2}, r_c^2 \right\} \right)^{\ell_2}
\end{aligned}$$

from which it follows, that upon using the definition of $w_{F,\delta}$ in (1.10c) we arrive at

$$\begin{aligned}
&m(r_c) |\mathfrak{B}_{0,2}(r_c, s, r_c)| \\
&\lesssim \left(\mathbf{1}_{s < r_c} \frac{s^{k-\frac{1}{2}}}{r_c^{k-\frac{1}{2}}} + \mathbf{1}_{r_c < s} \frac{r_c^{k+\frac{1}{2}}}{s^{k+\frac{1}{2}}} \right) \frac{w_{F, \frac{\delta}{4}+2\ell}(s) \langle s \rangle^4 r_c^{\frac{3\zeta}{2}}}{\langle r_c \rangle^{6+3\zeta} w_{F, \frac{\delta}{2}}(r_c)} \\
&\quad \times \left(\mathbf{1}_{r_c > 1} + \mathbf{1}_{s \leq r_c \leq 1} + \frac{r_c^2}{s^2} \mathbf{1}_{r_c < s < 1} + r_c^2 \mathbf{1}_{r_c < 1 < s} \right) \\
&\quad \times k^{2\ell_2} \frac{\left(\max \left\{ \frac{1}{s^2}, s^2, \frac{1}{r_c^2}, r_c^2 \right\} \right)^{\ell_2}}{\left(\max \left\{ r_c^2, \frac{1}{r_c^2} \right\} \right)^{j-\ell_1}} \\
&\lesssim k^{2\ell_2} \frac{r_c^{\frac{3\zeta}{2}}}{\langle r_c \rangle^{3\zeta}} \left[\mathbf{1}_{s < r_c < 1} \frac{s^{2k-2\ell-2\ell_2+\frac{1}{2}-\frac{\delta}{4}}}{r_c^{2k-2j+2\ell_1+\frac{1}{2}-\frac{\delta}{2}}} \right. \\
&\quad + \mathbf{1}_{r_c < s < 1} \frac{s^{-2\ell+\frac{1}{2}-\frac{\delta}{4}}}{r_c^{-2j+2\ell_2+2\ell_1+\frac{1}{2}-\frac{\delta}{2}}} + \mathbf{1}_{s < 1 < r_c} \frac{s^{2k-2\ell+3-\frac{1}{2}-\frac{\delta}{4}} \max\{s^{-2\ell_2}, r_c^{2\ell_2}\}}{r_c^{2j-2\ell_1+\frac{1}{2}+\frac{\delta}{2}}} \\
&\quad + \mathbf{1}_{r_c < 1 < s} \frac{\max\{r_c^{-2\ell_2}, s^{2\ell_2}\}}{r_c^{-2j+2\ell_1+\frac{1}{2}-\frac{\delta}{2}} s^{2k-2\ell+\frac{1}{2}+1-\frac{\delta}{2}}} + \mathbf{1}_{1 < s < r_c} \frac{1}{r_c^{2j-2\ell_1-2\ell_2+1+\frac{\delta}{2}} s^{-2\ell+\frac{3}{2}-\frac{\delta}{4}}} \\
&\quad \left. + \mathbf{1}_{1 < r_c < s} \frac{r_c^{2k-2j+2\ell_1-\frac{1}{2}-\frac{\delta}{2}}}{s^{2k-2\ell-2\ell_2+\frac{3}{2}-\frac{\delta}{4}}} \right] \\
&\lesssim k^{2\ell_2} \frac{r_c^{\frac{3\zeta}{2}}}{\langle r_c \rangle^{3\zeta}} \left[\mathbf{1}_{r_c \leq s} \frac{s^{\frac{1-\zeta}{2}}}{r_c^{\frac{1-\zeta}{2}}} + \mathbf{1}_{s < r_c} \frac{\langle s \rangle^{\frac{1+\zeta}{2}}}{\langle r_c \rangle^{\frac{1+\zeta}{2}}} \right] = k^{2\ell_2} \mathfrak{M}(s, r_c).
\end{aligned}$$

Here we used that $\ell + \ell_1 + \ell_2 \leq j \leq k-1$. Therefore, (B.58b) holds with \mathfrak{M} multiplied by $k^{2\ell_2}$. \square

Similarly, the next corollary concludes the proof of Theorem B.3.

Corollary B.15 *Let $0 \leq j \leq k-1$, and $0 \leq \ell, \ell_1, \ell_2$ be such that $\ell + \ell_1 + \ell_2 \leq j$. Let $0 < \zeta \leq \frac{\delta}{6}$. Consider the weights*

$$\begin{aligned}\mathfrak{B}_{\varepsilon,1}(r, r_c, r_c) &= \mathbf{1}_{r \leq 1} \mathbf{1}_{2r \leq r_c} \frac{\beta(r) \min(r^2, r^{-2})^j}{r^{\frac{1}{2}} w_{f,\delta}(r) u'(r_c) r_c u'(r_c)} B_{\ell,\varepsilon}^{(1)}(r, r_c, r_c) \\ \mathfrak{B}_{\varepsilon,2}(r_c, s, r_c) &= \frac{\beta(r_c) w_{F,\frac{\delta}{4}+2\ell}(s)}{u'(s)} B_{\ell,\varepsilon}^{(2)}(r_c, s, r_c)\end{aligned}$$

where $B_{\ell,\varepsilon}^{(1)}$ is a suitable $(2\ell_1, \ell_1 + \eta/2)$ kernel of type I or II, and $B_{\ell,\varepsilon}^{(2)}$ is a suitable $(2\ell_2, \ell_2 + \eta/2)$ kernel of type I or II. Then, letting

$$m(r_c) = \frac{r_c^{\frac{3\zeta}{2}} \langle r_c \rangle^2}{\langle r_c \rangle^{3\zeta} w_{F,\frac{\delta}{2}}(r_c) (\max\{r_c^2, r_c^{-2}\})^{j-\ell_1}} \quad (\text{B.73})$$

the conditions (B.57)–(B.58) of Lemma B.13 are satisfied, with \mathfrak{M} replaced by $k^{2\ell_1+2\ell_2}\mathfrak{M}$, and where the limiting weights are obtained by passing $\varepsilon \rightarrow 0$ in $B_{\ell,\varepsilon}^{(1)}$ and $B_{\ell,\varepsilon}^{(2)}$. In particular, the norm on $L^2(dr)$ of the operator L_0 defined in (B.2) is bounded as $k^{\zeta+2\ell_1+2\ell_2}$.

Proof of Corollary B.15 The proof is essentially the same as the proof of Corollary B.14, hence, we only emphasize the requisite modifications. We notice that the choice of $m(r_c)$ in (B.73) is different from the one in (B.72), as we have multiplied by a factor of $\langle r_c \rangle^2$. Thus, when checking conditions (B.58a)–(B.58c) for the weight $\mathfrak{B}_{\varepsilon,2}(r_c, s, r_c)$ we need to obtain bounds which are better by an $\langle r_c \rangle^2$. This however is automatic from the decay of $\beta(r_c)$ present in the definition of $\mathfrak{B}_{\varepsilon,2}(r_c, s, r_c)$. In turn, when checking conditions (B.57a)–(B.57c) for $\mathfrak{B}_{\varepsilon,1}(r, r_c, r_c)$, we can afford estimates which are worse than those in Corollary B.14 by a factor of $\langle r_c \rangle^2$. This is natural, since recall that the $\mathfrak{B}_{\varepsilon,1}(r, r_c, r_c)$ in this corollary, differs from the one in Corollary B.14 by a factor of $\frac{\langle r_c \rangle^4 r^2}{r_c^2}$, for $r \leq 1$ and $2r \leq r_c$. The worsening of the estimate by a factor of $\langle r_c \rangle^2$ is thus compensating this extra factor when $r_c \geq 1$. On the other hand, in the case $r_c \leq 1$, we have $\frac{\langle r_c \rangle^4 r^2}{r_c^2} \leq \frac{r^2}{r_c^2} \leq \frac{1}{4}$, and thus there is nothing additional to prove. With these changes in hand, we may now follow line by the proof of Corollary B.14, and conclude the proof. \square

B.1.4 A Useful Product Formula for Weights

Lemma B.16 *With the notation of (B.24), define*

$$\begin{aligned}\mathbb{W}(r, s, s_0) &:= \left[\frac{\mathbf{1}_{r \leq 1}}{r^{k+\frac{1}{2}}} + \mathbf{1}_{r \geq 1} r^{k-\frac{1}{2}} \right] \left[\mathbf{1}_{s \leq 1} s^{k+\frac{1}{2}} + \frac{\mathbf{1}_{s \geq 1}}{s^{k-\frac{1}{2}}} \right] \frac{\mathcal{B}(r, s_0)}{\langle s_0 \rangle^4} \frac{\mathcal{B}(s_0, s)}{\langle s \rangle^4} \\ &= \left[\frac{\mathbf{1}_{r \leq 1}}{r^{k+\frac{1}{2}}} + \mathbf{1}_{r \geq 1} r^{k-\frac{1}{2}} \right] \left[\mathbf{1}_{s \leq 1} s^{k+\frac{1}{2}} + \frac{\mathbf{1}_{s \geq 1}}{s^{k-\frac{1}{2}}} \right] \\ &\quad \times \left[\mathbf{1}_{s_0 < r} \frac{s_0^{k-\frac{1}{2}}}{r^{k-\frac{1}{2}}} + \mathbf{1}_{s_0 > r} \frac{r^{k+\frac{1}{2}}}{s_0^{k+\frac{1}{2}}} \right] \left[\mathbf{1}_{s < s_0} \frac{s^{k-\frac{1}{2}}}{s_0^{k-\frac{1}{2}}} + \mathbf{1}_{s > s_0} \frac{s_0^{k+\frac{1}{2}}}{s^{k+\frac{1}{2}}} \right]. \quad (\text{B.74})\end{aligned}$$

Then we have that

$$\begin{aligned}
 \mathbb{W}(r, s, s_0) \mathbf{1}_{r < s < s_0} &= \mathbf{1}_{1 < r < s < s_0} \frac{r^{2k}}{s_0^{2k}} + \mathbf{1}_{r < 1 < s < s_0} \frac{1}{s_0^{2k}} \\
 &\quad + \mathbf{1}_{r < s < 1 < s_0} \frac{s^{2k}}{s_0^{2k}} + \mathbf{1}_{r < s < s_0 < 1} \frac{s^{2k}}{s_0^{2k}} \\
 \mathbb{W}(r, s, s_0) \mathbf{1}_{s < r < s_0} &= \mathbf{1}_{1 < s < r < s_0} \frac{r^{2k}}{s_0^{2k}} + \mathbf{1}_{s < 1 < r < s_0} \frac{r^{2k} s^{2k}}{s_0^{2k}} \\
 &\quad + \mathbf{1}_{s < r < 1 < s_0} \frac{s^{2k}}{s_0^{2k}} + \mathbf{1}_{s < r < s_0 < 1} \frac{s^{2k}}{s_0^{2k}} \\
 \mathbb{W}(r, s, s_0) \mathbf{1}_{r < s_0 < s} &= \mathbf{1}_{1 < r < s_0 < s} \frac{r^{2k}}{s^{2k}} + \mathbf{1}_{r < 1 < s_0 < s} \frac{1}{s^{2k}} + \mathbf{1}_{r < s_0 < 1 < s} \frac{1}{s^{2k}} + \mathbf{1}_{r < s_0 < s < 1} \\
 \mathbb{W}(r, s, s_0) \mathbf{1}_{s_0 < r < s} &= \mathbf{1}_{1 < s_0 < r < s} \frac{s_0^{2k}}{s^{2k}} + \mathbf{1}_{s_0 < 1 < r < s} \frac{s_0^{2k}}{s^{2k}} \\
 &\quad + \mathbf{1}_{s_0 < r < 1 < s} \frac{s_0^{2k}}{r^{2k} s^{2k}} + \mathbf{1}_{s_0 < r < s < 1} \frac{s_0^{2k}}{r^{2k}} \\
 \mathbb{W}(r, s, s_0) \mathbf{1}_{s < s_0 < r} &= \mathbf{1}_{1 < s < s_0 < r} + \mathbf{1}_{s < 1 < s_0 < r} s^{2k} + \mathbf{1}_{s < s_0 < 1 < r} s^{2k} + \mathbf{1}_{s < s_0 < r < 1} \frac{s^{2k}}{r^{2k}} \\
 \mathbb{W}(r, s, s_0) \mathbf{1}_{s_0 < s < r} &= \mathbf{1}_{1 < s_0 < s < r} \frac{s_0^{2k}}{s^{2k}} + \mathbf{1}_{s_0 < 1 < s < r} \frac{s_0^{2k}}{s^{2k}} \\
 &\quad + \mathbf{1}_{s_0 < s < 1 < r} s_0^{2k} + \mathbf{1}_{s_0 < s < r < 1} \frac{s_0^{2k}}{r^{2k}}
 \end{aligned}$$

holds. In particular, note that

$$\mathbb{W}(r, s, s_0) \leq 1.$$

Proof of Lemma B.16 The proof follows by inspection of each of the 24 possible permutations of $\{r, s, s_0, 1\}$. \square

Lemma B.17 Let $0 \leq j \leq k$, and $\ell + \ell_1 + \ell_2 \leq j$. Define

$$\mathbb{L}(r, s, s_0) := \left[\mathbf{1}_{r \leq 1} r^{2j} + \frac{\mathbf{1}_{r \geq 1}}{r^{2j}} \right] \left[\frac{\mathbf{1}_{s \leq 1}}{s^{2\ell}} + \mathbf{1}_{s \geq 1} s^{2\ell} \right] \mathcal{L}_{2\ell_1, \ell_1}(r, s_0) \mathcal{L}_{2\ell_2, \ell_2}(s_0, s).$$

Then, we have that

$$\mathbb{W}(r, s, s_0) \mathbb{L}(r, s, s_0) \lesssim 1.$$

Proof of Lemma B.17 Notice that for $j = 0$, we must have $\ell = \ell_1 = \ell_2 = 0$, and thus $\mathbb{L} = 1$. In this case the proof directly follows from Lemma B.16.

For $j \geq 1$, we use the precise formula for \mathbb{W} in Lemma B.16, and use the definition of \mathbb{L} , which is also a function of r , s , and s_0 , to obtain that

$$\begin{aligned}
 \sqrt{\mathbb{W}(r, s, s_0) \mathbb{L}(r, s, s_0)} \mathbf{1}_{r < s < s_0} &= \mathbf{1}_{1 < r < s < s_0} \frac{r^{k-j} s^\ell}{s_0^{k-\ell_1-\ell_2}} \\
 &\quad + \mathbf{1}_{r < 1 < s < s_0} \frac{s^\ell r^j}{s_0^{k-\ell_2}} \left(\frac{1}{r^{\ell_1}} + s_0^{\ell_1} \right) \\
 &\quad + \mathbf{1}_{r < s < 1 < s_0} \frac{r^j s^{k+\ell}}{s_0^k} \left(\frac{1}{r^{\ell_1}} + s_0^{\ell_1} \right) \left(\frac{1}{s^{\ell_2}} + s_0^{\ell_2} \right) \\
 &\quad + \mathbf{1}_{r < s < s_0 < 1} \frac{r^{j-\ell_1} s^{k-\ell-\ell_2}}{s_0^k} \\
 \sqrt{\mathbb{W}(r, s, s_0) \mathbb{L}(r, s, s_0)} \mathbf{1}_{s < r < s_0} &= \mathbf{1}_{1 < s < r < s_0} \frac{r^{k-j} s^\ell}{s_0^{k-\ell_1-\ell_2}} \\
 &\quad + \mathbf{1}_{s < 1 < r < s_0} \frac{r^{k-j} s^{k-\ell}}{s_0^{k-\ell_1}} \left(\frac{1}{s^{\ell_2}} + s_0^{\ell_2} \right) \\
 &\quad + \mathbf{1}_{s < r < 1 < s_0} \frac{r^j s^{k-\ell}}{s_0^k} \left(\frac{1}{r^{\ell_1}} + s_0^{\ell_1} \right) \left(\frac{1}{s^{\ell_2}} + s_0^{\ell_2} \right) \\
 &\quad + \mathbf{1}_{s < r < s_0 < 1} \frac{r^j s^{k-\ell}}{s_0^k} \\
 \sqrt{\mathbb{W}(r, s, s_0) \mathbb{L}(r, s, s_0)} \mathbf{1}_{r < s_0 < s} &= \mathbf{1}_{1 < r < s_0 < s} \frac{r^{k-j} s_0^{\ell_1}}{s^{k-\ell-\ell_2}} \\
 &\quad + \mathbf{1}_{r < 1 < s_0 < s} \frac{r^j}{s^{k-\ell-\ell_2}} \left(\frac{1}{r^{\ell_1}} + s_0^{\ell_1} \right) \\
 &\quad + \mathbf{1}_{r < s_0 < 1 < s} \frac{r^{j-\ell_1}}{s^{k-\ell}} \left(\frac{1}{s_0^{\ell_2}} + s^{\ell_2} \right) \\
 &\quad + \mathbf{1}_{r < s_0 < s < 1} \frac{r^{j-\ell_1}}{s^\ell s_0^{\ell_2}} \\
 \sqrt{\mathbb{W}(r, s, s_0) \mathbb{L}(r, s, s_0)} \mathbf{1}_{s_0 < r < s} &= \mathbf{1}_{1 < s_0 < r < s} \frac{s_0^k}{r^{j-\ell_1} s^{k-\ell-\ell_2}} \\
 &\quad + \mathbf{1}_{s_0 < 1 < r < s} \frac{s_0^k}{r^j s^{k-\ell}} \left(\frac{1}{s_0^{\ell_1}} + r^{\ell_1} \right) \left(\frac{1}{s_0^{\ell_2}} + s^{\ell_2} \right) \\
 &\quad + \mathbf{1}_{s_0 < r < 1 < s} \frac{s_0^{k-\ell_1}}{r^{k-j} s^{k-\ell}} \left(\frac{1}{s_0^{\ell_2}} + s^{\ell_2} \right) \\
 &\quad + \mathbf{1}_{s_0 < r < s < 1} \frac{s_0^{k-\ell_1-\ell_2}}{s^\ell r^{k-j}}
 \end{aligned}$$

$$\begin{aligned}
\sqrt{\mathbb{W}(r, s, s_0) \mathbb{L}(r, s, s_0) \mathbf{1}_{s < s_0 < r}} &= \mathbf{1}_{1 < s < s_0 < r} \frac{s^\ell s_0^{\ell_2}}{r^{j-\ell_1}} \\
&+ \mathbf{1}_{s < 1 < s_0 < r} \frac{s^{k-\ell}}{r^{j-\ell_1}} \left(\frac{1}{s^{\ell_2}} + s_0^{\ell_2} \right) \\
&+ \mathbf{1}_{s < s_0 < 1 < r} \frac{s^{k-\ell-\ell_2}}{r^j} \left(\frac{1}{s_0^{\ell_1}} + r^{\ell_1} \right) \\
&+ \mathbf{1}_{s < s_0 < r < 1} \frac{s^{k-\ell-\ell_2}}{r^{k-j} s_0^{\ell_1}} \\
\sqrt{\mathbb{W}(r, s, s_0) \mathbb{L}(r, s, s_0) \mathbf{1}_{s_0 < s < r}} &= \mathbf{1}_{1 < s_0 < s < r} \frac{s_0^k}{r^{j-\ell_1} s^{k-\ell-\ell_2}} \\
&+ \mathbf{1}_{s_0 < 1 < s < r} \frac{s_0^k}{r^j s^{k-\ell}} \left(\frac{1}{s_0^{\ell_1}} + r^{\ell_1} \right) \left(\frac{1}{s_0^{\ell_2}} + s^{\ell_2} \right) \\
&+ \mathbf{1}_{s_0 < s < 1 < r} \frac{s_0^{k-\ell_2}}{r^j s^{\ell}} \left(\frac{1}{s_0^{\ell_1}} + r^{\ell_1} \right) \\
&+ \mathbf{1}_{s_0 < s < r < 1} \frac{s_0^{k-\ell_1}}{r^{k-j} s^{\ell+\ell_2}}
\end{aligned}$$

Inspecting each of these 24 terms, by using the constraint $0 \leq \ell + \ell_1 + \ell_2 \leq j \leq k-1$ the proof of the lemma follows. \square

B.2 The Remaining Combinations of Iterated Operators

Similar to the results in the previous section, namely Theorems B.1, B.2, and B.3, one can prove a number of results for passing $\varepsilon \rightarrow 0$ in other combinations of three integral operators. The aforementioned theorems deal with the most difficult case, of an approximate delta function combined with two approximate singular integrals. The remaining operators all have at least one approximate delta function, and at most one approximate singular integral, which are hence easier to treat.

Theorem B.18 *Let $0 \leq j \leq k-1$, and $0 \leq \ell, \ell_1, \ell_2$ be such that $\ell + \ell_1 + \ell_2 \leq j$. Let $B_{\ell, \varepsilon}^{(1)}$ be a suitable $(2\ell_1, \ell_1 + \eta/2)$ kernel of type I or II, and $B_{\ell, \varepsilon}^{(2)}$ is a suitable $(2\ell_2, \ell_2 + \eta/2)$ kernel of type I or II. Let $0 < \zeta \leq \frac{\delta}{6}$, and consider the pairs of weights given in (B.3)–(B.4) or (B.5)–(B.6). That is, either consider the pair*

$$\begin{aligned}
\mathfrak{B}_{\varepsilon, 1}(r, s_0, r_c) &= \chi_1(r, r_c) \frac{\beta(r) \min(r^2, r^{-2})^j}{w_{F, \delta}(r) u'(s_0)} B_{\ell, \varepsilon}^{(1)}(r, s_0, r_c) \\
\mathfrak{B}_{\varepsilon, 2}(s_0, s, r_c) &= \frac{\beta(s_0) w_{F, \frac{\delta}{4} + 2\ell}(s)}{u'(s)} B_{\ell, \varepsilon}^{(2)}(s_0, s, r_c),
\end{aligned}$$

or consider the pair

$$\mathfrak{B}_{\varepsilon,1}(r, s_0, r_c) = \chi_2(r, r_c) \frac{\beta(r) \min(r^2, r^{-2})^j}{r^{\frac{1}{2}} w_{f,\delta}(r) u'(s_0) r_c u'(r_c)} B_{\ell,\varepsilon}^{(1)}(r, s_0, r_c)$$

$$\mathfrak{B}_{\varepsilon,2}(s_0, s, r_c) = \frac{\beta(s_0) w_{F,\frac{\delta}{4}+2\ell}(s)}{u'(s)} B_{\ell,\varepsilon}^{(2)}(s_0, s, r_c).$$

For each such pair of weights, we have that the following limits hold, in the sense of bounded operators on $L^2(dr)$:

$$\begin{aligned} & \int_0^\infty \int_0^\infty \int_0^\infty \frac{(u(r) - u(r_c))u'(r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} \frac{\varepsilon u'(s)}{(u(s) - u(r_c))^2 + \varepsilon^2} \frac{\varepsilon u'(s_0)}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \\ & \quad \times \mathfrak{B}_{\varepsilon,1}(r, s_0, r_c) \mathfrak{B}_{\varepsilon,2}(s_0, s, r_c) f(s) \, ds ds_0 dr_c \\ & \quad \longrightarrow \pi^2 p.v. \int_0^\infty \frac{u'(r_c)}{u(r) - u(r_c)} \mathfrak{B}_{0,1}(r, r_c, r_c) \mathfrak{B}_{0,2}(r_c, r_c, r_c) f(r_c) \, dr_c \\ & \int_0^\infty \int_0^\infty \int_0^\infty \frac{\varepsilon u'(r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} \frac{(u(s) - u(r_c))u'(s)}{(u(s) - u(r_c))^2 + \varepsilon^2} \frac{\varepsilon u'(s_0)}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \\ & \quad \times \mathfrak{B}_{0,1}(r, s_0, r_c) \mathfrak{B}_{0,2}(s_0, s, r_c) f(s) \, ds ds_0 dr_c \\ & \quad \longrightarrow \pi^2 p.v. \int_0^\infty \frac{u'(s)}{u(s) - u(r)} \mathfrak{B}_{0,1}(r, r, r) \mathfrak{B}_{0,2}(r, s, r) f(s) \, ds \\ & \int_0^\infty \int_0^\infty \int_0^\infty \frac{\varepsilon u'(r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} \frac{\varepsilon u'(s)}{(u(s) - u(r_c))^2 + \varepsilon^2} \frac{\varepsilon u'(s_0)}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \\ & \quad \times \mathfrak{B}_{\varepsilon,1}(r, s_0, r_c) \mathfrak{B}_{\varepsilon,2}(s_0, s, r_c) f(s) \, ds ds_0 dr_c \\ & \quad \longrightarrow -\pi^3 \mathfrak{B}_{0,1}(r, r, r) \mathfrak{B}_{0,2}(r, r, r) f(r) \\ & \int_0^\infty \int_0^\infty \int_0^\infty \frac{(u(r) - u(r_c))u'(r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} u'(s) \frac{\varepsilon u'(s_0)}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \\ & \quad \times \mathfrak{B}_{\varepsilon,1}(r, s_0, r_c) \mathfrak{B}_{\varepsilon,2}(s_0, s, r_c) f(s) \, ds ds_0 dr_c \\ & \quad \longrightarrow -\pi p.v. \int_0^\infty \int_0^\infty \frac{u'(r_c)}{u(r) - u(r_c)} u'(s) \mathfrak{B}_{0,1}(r, r_c, r_c) \mathfrak{B}_{0,2}(r_c, s, r_c) f(s) \, ds dr_c \\ & \int_0^\infty \int_0^\infty \int_0^\infty \frac{\varepsilon u'(r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} u'(s) \frac{\varepsilon u'(s_0)}{(u(s_0) - u(r_c))^2 + \varepsilon^2} \\ & \quad \times \mathfrak{B}_{\varepsilon,1}(r, s_0, r_c) \mathfrak{B}_{\varepsilon,2}(s_0, s, r_c) f(s) \, ds ds_0 dr_c \\ & \quad \longrightarrow \pi^2 \int_0^\infty u'(s) \mathfrak{B}_{0,1}(r, r, r) \mathfrak{B}_{0,2}(r, s, r) f(s) \, ds \\ & \int_0^\infty \int_0^\infty \frac{(u(r) - u(r_c))u'(r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} \frac{\varepsilon u'(s)}{(u(s) - u(r_c))^2 + \varepsilon^2} \mathfrak{B}_{\varepsilon,1}(r, s, r_c) f(s) \, ds dr_c \\ & \quad \longrightarrow -\pi p.v. \int_0^\infty \frac{u'(r_c)}{u(r) - u(r_c)} \mathfrak{B}_{0,1}(r, r_c, r_c) f(r_c) \, dr_c \\ & \int_0^\infty \int_0^\infty \frac{\varepsilon u'(r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} \frac{\varepsilon u'(s)}{(u(s) - u(r_c))^2 + \varepsilon^2} \mathfrak{B}_{\varepsilon,1}(r, s, r_c) f(s) \, ds dr_c \end{aligned}$$

$$\begin{aligned}
& \longrightarrow \pi^2 \mathfrak{B}_{0,1}(r, r, r) f(r) dr_c \\
& \int_0^\infty \int_0^\infty \frac{\varepsilon u'(r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} \frac{(u(s) - u(r_c)u'(s))}{(u(s) - u(r_c))^2 + \varepsilon^2} \mathfrak{B}_{\varepsilon,1}(r, s, r_c) f(s) ds ds_0 dr_c \\
& \longrightarrow -\pi p.v. \int_0^\infty \frac{u'(s)}{u(s) - u(r)} \mathfrak{B}_{0,1}(r, s, r) f(s) ds \\
& \int_0^\infty \int_0^\infty \frac{\varepsilon u'(r_c)}{(u(r) - u(r_c))^2 + \varepsilon^2} u'(s) \mathfrak{B}_{\varepsilon,1}(r, s, r_c) f(s) ds ds_0 dr_c \\
& \longrightarrow -\pi \int_0^\infty u'(s) \mathfrak{B}_{0,1}(r, s, r) f(s) ds.
\end{aligned}$$

Moreover, in each case the limiting operators are bounded on $L^2(dr)$ with norm $\lesssim k^{\zeta+2\ell_1+2\ell_2}$.

Proving each statement in Theorem B.18 amounts to following step-by-step the proof of Theorems B.1, B.2, and B.3. For every approximate delta function, first show that the contribution away from the diagonal vanishes, leaving one with the contribution close to the diagonal, in which one may pass to the limit because we have Hölder regularity of our weights in all variables not called r_c . Checking that the weights we consider are suitable for this procedure, i.e. that they give a bounded operator in the correctly weighted L^2 spaces, was already done in the proof of Theorems B.2 and B.3. This leaves us with the contribution from the joint diagonal, in which the remaining operators are either approximate singular integrals (note however that at most one singular integral may be present), or simple Hilbert–Schmidt kernels. For the approximate singular integral we again employ Lemma B.13 and Corollaries B.14 and B.15, depending on which weight we choose. For passing to the limit in the operators with Hilbert–Schmidt kernels, we simply subtract the limiting kernel, so that the difference is again Hilbert–Schmidt but of size ε^ζ for some $\zeta > 0$. Here we lose a power of k^ζ/r_c^ζ , but since $\zeta > 0$ is arbitrary, we may absorb this loss into the weights. This procedure again essentially uses the regularity of our weights in all variables not called r_c . The only non-trivial case left to discuss is the case in which we have an apparent approximate delta function in the r_c variable. We treat this operator by duality, so that the approximate delta function in r_c , becomes an approximate delta function in the r variable, acting on the L^2 test function. We pass to the limit in the dual pairing and thus obtain the formula and the boundedness of the limiting operator. Having available the estimates in the previous section, which were used to Theorems B.1, B.2, and B.3, implementing the above described program is tedious, but routine. In order to avoid this unnecessary redundancy, we omit these details.

C Properties of H_0 and H_∞

The next lemma concerns the asymptotic analysis of $R_{0,r}(z)$ and $R_{r,\infty}(z)$, which both arise in the Green's function.

Lemma C.1 For $z = c \pm i\varepsilon \in I_\alpha$ with ε sufficiently small, for $|r - r_c| \lesssim r_c/k$ there holds

$$|R_{0,r}^\varepsilon(z)| \mathbf{1}_{|r-r_c| \leq r_c/k} \lesssim \mathbf{1}_{r_c \leq 1} \max(r_c^{-3}, r_c^5) \left(k + \left| \log k \frac{|r - r_c|}{r_c} \right| \right) \quad (\text{C.1a})$$

$$|R_{0,r}^\varepsilon(z)| \mathbf{1}_{|r-r_c| \geq r_c/k} \lesssim k \mathbf{1}_{r_c \leq 1} \left(\mathbf{1}_{r < r_c} \frac{r^{2k-2}}{r_c^{2k+1}} + \mathbf{1}_{r_c < r} \frac{1}{r_c^3} \right) \quad (\text{C.1b})$$

$$+ \mathbf{1}_{r_c \geq 1} \left(\mathbf{1}_{r < 1} \frac{r^{2k-2}}{r_c^{2k-1}} + \mathbf{1}_{1 \leq r < r_c} \frac{kr^{2k+4}}{r_c^{2k-1}} + \mathbf{1}_{r > r_c} kr_c^5 \right), \quad (\text{C.1c})$$

and similarly

$$|R_{r,\infty}^\varepsilon(z)| \mathbf{1}_{|r-r_c| \lesssim r_c/k} \lesssim \mathbf{1}_{r_c \leq 1} \max(r_c^{-3}, r_c^5) \left(k + \left| \log k \frac{|r - r_c|}{r_c} \right| \right) \quad (\text{C.2a})$$

$$|R_{r,\infty}^\varepsilon(z)| \mathbf{1}_{|r-r_c| \gtrsim r_c/k} \lesssim \mathbf{1}_{r_c \leq 1} \left(\mathbf{1}_{r < r_c} \frac{k}{r_c^3} + \mathbf{1}_{r_c < r \leq 1} \frac{kr_c^{1+2k}}{r^{4+2k}} + \mathbf{1}_{r > 1} \frac{r_c^{2k+1}}{r^{2k-2}} \right) \quad (\text{C.2b})$$

$$+ k \mathbf{1}_{r_c \geq 1} \left(\mathbf{1}_{r < r_c} r_c^5 + \mathbf{1}_{r > r_c} \frac{r_c^{2k+3}}{r^{2k-2}} \right). \quad (\text{C.2c})$$

Moreover, for all η sufficiently small and $z \in I_\alpha$, we have that $\varepsilon^{-\eta}(R_{0,r}^\varepsilon(c \pm i\varepsilon) - R_{0,r}(c))$ also satisfies (C.1) and $\varepsilon^{-\eta}(R_{r,\infty}^\varepsilon(c \pm i\varepsilon) - R_{r,\infty}(c))$ also satisfies (C.2).

Proof of Lemma C.1 The proof is a straightforward variant of arguments applied in the proof of Lemma 5.2 and is hence omitted for the sake of brevity. \square

Proof of Lemma 5.4 Recall (5.25), which will be of use in the critical layer region $|r - r_c| < r_c/k$. From Theorem 4.3, followed by Lemma A.1, there holds for $|r - r_c| < r_c/k$:

$$\frac{1}{u'(r)P(r, c \pm i\varepsilon)} \approx \frac{1}{u'(r_c)} \min \left(\frac{r^{k-1/2}}{r_c^{k-1/2}}, \frac{r_c^{k+1/2}}{r^{k+1/2}} \right) \approx \max(r_c^{-1}, r_c^3).$$

Consider next (5.26). From Lemmas C.1 and A.1 we have for $|r - r_c| < r_c/k$ (for $z \in I_\alpha$ and ε sufficiently small),

$$\begin{aligned} & |\phi(r, c \pm i\varepsilon) (R_{0,r}^\varepsilon(c \pm i\varepsilon) \mp iE_{0,r}^\varepsilon(c \pm i\varepsilon))| \\ & \lesssim \mathbf{1}_{r_c \leq 1} |r - r_c| r_c^{-3} \left(k + \left| \log k \frac{|r - r_c|}{r_c} \right| \right) \\ & + \mathbf{1}_{r_c \geq 1} \frac{1}{r_c^3} |r - r_c| r_c^5 \left(k + \left| \log k \frac{|r - r_c|}{r_c} \right| \right), \end{aligned}$$

which is the desired estimate of H_0 ; the treatment of H_∞ is the same.

Turn next to the estimates (5.28a) and (5.28b). Consider just the estimates in (5.28a); the estimate (5.28b) is analogous. For $r < r_c(1 - \frac{1}{k})$, (from (2.31)),

$$|H_0(r, z)| \leq |\phi(r, z)| \int_0^r \frac{1}{|\phi^2(s, z)|} ds \lesssim |u - c| \frac{r_c^{k-1/2}}{r^{k-1/2}} \int_0^r \frac{1}{|u - c|^2} \frac{s^{2k-1}}{r_c^{2k-1}} ds.$$

Consider the case $r_c \leq 1$. Then from (A.1),

$$\begin{aligned} \mathbf{1}_{r < r_c/2} |H_0(r, z)| &\lesssim \frac{r_c^{k-1/2}}{r_c^2 r^{k-1/2}} \int_0^r \frac{s^{2k-1}}{r_c^{2k-1}} ds \approx \frac{1}{k} \frac{r^{k+1/2}}{r_c^{k+3/2}}, \\ \mathbf{1}_{r_c/2 < r < r_c(1-\frac{1}{k})} |H_0(r, z)| &\lesssim r_c |r - r_c| \frac{r_c^{k-1/2}}{r^{k-1/2}} \left(\frac{r^{2k}}{k r_c^{2k+3}} + \int_{r_c/2}^r \frac{1}{|u(s) - c|^2} \frac{s^{2k-1}}{r_c^{2k-1}} ds \right) \\ &\lesssim \frac{r^{k+1/2}}{k r_c^{k+3/2}} + \frac{\mathbf{1}_{r_c/2 < r < r_c(1-\frac{1}{k})}}{r_c |r - r_c|} \frac{1}{k} \frac{r^{k+1/2}}{r_c^{k-1/2}} \\ &\lesssim \frac{r^{k+1/2}}{r_c^{k+3/2}}. \end{aligned}$$

Once the integral crosses the critical layer, one cannot do better than the analogous upper bound on $M(z)$:

$$\begin{aligned} \mathbf{1}_{r_c < r < 1} |H_0(r, z)| &\lesssim \mathbf{1}_{r_c < r < 1} |u - c| \frac{r^{k+1/2}}{r_c^{k+1/2}} \frac{k}{r_c^3} \lesssim k \frac{r^{k+1/2}}{r_c^3} \frac{r^{k+1/2}}{r_c^{k+1/2}}, \\ \mathbf{1}_{r_c < 1} \mathbf{1}_{r \geq 1} |H_0(r, z)| &\lesssim \mathbf{1}_{r_c < 1} \mathbf{1}_{r \geq 1} \frac{k}{r_c^3} \frac{r^{k+1/2}}{r_c^{k+1/2}}. \end{aligned}$$

Analogous estimates are made for $r_c \geq 1$ and for H_∞ ; these are omitted for the sake of brevity. This completes the boundedness estimates (5.28a) and (5.28b).

Next, we consider the estimation of $H_0(r, c \pm i\varepsilon) - H_0(r, c)$. For $r < r_c(1 - 1/k)$, we write

$$\begin{aligned} H_0(r, c \pm i\varepsilon) - H_0(r, c) &= (\phi(r, c \pm i\varepsilon) - \phi(r, c)) \int_0^r \frac{1}{\phi^2(s, z)} ds \\ &\quad + \phi(r, c) \int_0^r \frac{\phi^2(s, c) - \phi^2(s, c \pm i\varepsilon)}{\phi^2(s, c \pm i\varepsilon) \phi^2(s, c)} ds. \end{aligned}$$

In this region, we apply a proof analogous to those used in Lemma 5.2. For the region $|r - r_c| < r_c/k$ we again use the complex integral expansion (5.25):

$$\begin{aligned} H_0(r, c \pm i\varepsilon) - H_0(r, c) &= \frac{1}{u'(r)P(r, c \pm i\varepsilon)} - \frac{1}{u'(r)P(r, c)} \\ &\quad + \phi(r, c \pm i\varepsilon) (R_{0,r}^\varepsilon(c \pm i\varepsilon) \mp iE_{0,r}^\varepsilon(c \pm i\varepsilon)) \\ &\quad - \phi(r, c) (R_{0,r}^\varepsilon(c) \mp iE_{0,r}^\varepsilon(c)). \end{aligned}$$

The desired estimates then follow from the convergence of $R_{0,r}^\varepsilon$ and $E_{0,r}^\varepsilon$. Convergence for $r \gtrsim r_c$ follows as in Lemma 5.2. The estimates of $\varepsilon^\eta (H_\infty(r, c \pm i\varepsilon) - H_\infty(r, c))$ follows similarly as well. \square

Proof of Lemma 5.5 The identities (5.29) follow by direct calculation from (5.25). Next, observe that $r\partial_r\phi = ru'P + (u - c)r\partial_rP$. The estimate (5.31a) then follows from Lemma C.1 and Theorem 4.3.

By direct calculation,

$$\partial_r H_0(r, z) = -\partial_r\phi \int_0^r \frac{1}{\phi(s, z)^2} ds - \frac{1}{\phi} = \left(\frac{u'}{u - z} + \frac{P'}{P} \right) H_0 - \frac{1}{\phi}; \quad (\text{C.3})$$

and analogously for H_∞ . Therefore,

$$\begin{aligned} |r\partial_r H_\infty(r, z)| \mathbf{1}_{|r-r_c| \geq r_c/k} &\lesssim \left(\frac{r|u'|}{|u-c|} + k \right) |H_\infty(r, c \pm i\varepsilon)| \\ &\quad + r \min \left(\frac{r^{k-1/2}}{r_c^{k-1/2}}, \frac{r_c^{k+1/2}}{r^{k+1/2}} \right) \end{aligned} \quad (\text{C.4})$$

$$\begin{aligned} |r\partial_r H_0(r, z)| \mathbf{1}_{|r-r_c| \geq r_c/k} &\lesssim \left(\frac{r|u'|}{|u-c|} + k \right) |H_0(r, c \pm i\varepsilon)| \\ &\quad + r \min \left(\frac{r^{k-1/2}}{r_c^{k-1/2}}, \frac{r_c^{k+1/2}}{r^{k+1/2}} \right). \end{aligned} \quad (\text{C.5})$$

Then, note from (A.1), $\left| \frac{ru'(r)}{|u-c|} \right| \mathbf{1}_{|r-r_c| > r_c/k} \lesssim k$ and moreover that

$$\begin{aligned} &r \min \left(\frac{r^{k-1/2}}{r_c^{k-1/2}}, \frac{r_c^{k+1/2}}{r^{k+1/2}} \right) \\ &\lesssim \mathbf{1}_{r_c \leq 1} \left(\mathbf{1}_{r < r_c} \frac{r^{k+1/2}}{r_c^{k+3/2}} + \mathbf{1}_{r_c < r < 1} k \frac{r^{k+2+1/2}}{r_c^{k+2+3/2}} + \mathbf{1}_{r > 1} k \frac{r^{k+1/2}}{r_c^{k+3+1/2}} \right) \\ &\quad + \mathbf{1}_{r_c > 1} \left(\mathbf{1}_{r < 1} \frac{r^{k+1/2}}{r_c^{k-1/2}} + \mathbf{1}_{1 < r < r_c} \frac{r^{k+5/2}}{r_c^{k-1/2}} + \mathbf{1}_{r > r_c} k \frac{r^{k+1/2}}{r_c^{k-5/2}} \right) \\ &r \min \left(\frac{r^{k-1/2}}{r_c^{k-1/2}}, \frac{r_c^{k+1/2}}{r^{k+1/2}} \right) \\ &\lesssim \mathbf{1}_{r_c \leq 1} \left(\mathbf{1}_{r < r_c} k \frac{r_c^{k-3/2}}{r^{k-1/2}} + \mathbf{1}_{r_c < r < 1} \frac{r_c^{k+1/2}}{r^{k+3/2}} + \mathbf{1}_{r > 1} \frac{r_c^{k+1/2}}{r^{k-1/2}} \right) \\ &\quad + \mathbf{1}_{r_c > 1} \left(\mathbf{1}_{r < 1} k \frac{r_c^{k+5-1/2}}{r^{k-1/2}} + \mathbf{1}_{1 \leq r < r_c} k \frac{r_c^{k+5-1/2}}{r^{k+3/2}} + \mathbf{1}_{r_c < r} \frac{r_c^{k+5/2}}{r^{k-1/2}} \right), \end{aligned}$$

which completes the estimates of the ∂_r derivatives. Moreover, convergence follows from (C.3) and convergence estimates already proved.

Next, turn to studying derivatives involving ∂_{r_c} . Away from the critical layer there holds

$$\begin{aligned} -\partial_{r_c} H_0(r, c) \mathbf{1}_{|r-r_c|>r_c/k} &= \partial_{r_c} ((u-z)P) \int_0^r \frac{1}{\phi^2(s, c)} ds + \phi \int_0^r \chi_{\neq} \frac{u'(r_c)}{(u-z)^3 P^2} ds \\ &\quad + \phi \int_0^r \frac{u'(r_c)u'(s)}{(u-z)^2} \partial_G \left(\frac{\chi_c}{u'P^2} \right) ds \\ &\quad + \phi \int_0^r \frac{1}{(u-z)^2} \partial_{r_c} \frac{\chi_{\neq}}{P^2} ds. \end{aligned}$$

By adapting the arguments in Lemmas 5.2, 5.3 and 5.4 the desired boundedness and convergence estimates follow (note that appropriate estimates on $\partial_{r_c} P$ follow from Theorem 4.3). Further, from Theorem 4.3, we may even take an additional $r\partial_r$ derivative and obtain similar estimates (note crucially $|r-r_c| \geq r_c/k$).

Computing ∂_G derivatives near the critical layer from Lemma 5.1:

$$\begin{aligned} \partial_G H_0 &= \frac{1}{u'(r)} \partial_r \left(\frac{1}{u'P(r, c \pm i\varepsilon)} \right) - \frac{1}{u'(r)} \partial_r \phi \left(R_{0,r}^\varepsilon(z) \mp iE_{0,r}^\varepsilon \right) - PE(r, c \pm i\varepsilon) \\ &\quad + \frac{1}{u'(r_c)} \partial_{r_c} \left(\frac{1}{u'P} \right) - \frac{1}{u'(r_c)} \partial_{r_c} \phi \left(R_{0,r}^\varepsilon(z) \mp iE_{0,r}^\varepsilon(z) \right) \\ &\quad + P\chi_c E(r, z) - \phi \int_0^r \frac{u'}{(u-z)} \partial_G (\chi_c E) ds \\ &\quad - \frac{1}{u'(r_c)} \phi \int_0^r \chi_{\neq} \frac{u'(s)u'(r_c)}{(u-z)^2} E(s, z) - \frac{u'(s)}{(u-z)} \chi_{\neq} \partial_{r_c} E(s, z) ds. \end{aligned}$$

The cancellation which ultimately removes the logarithmic singularity is the fact that $\partial_G \phi = (u-z)\partial_G P$. Taking this into account gives the identity (5.30a). The corresponding calculation on H_∞ is analogous.

Next, we prove (5.31b) and (5.31c). Consider one of the singular appearing terms in (5.30a):

$$\phi \int_0^r \frac{u'(s)}{(u-z)} \partial_G (\chi_c E) ds = \phi \int_0^r \frac{u'(s)}{(u-z)} \partial_G \left(\chi_c \frac{1}{u'(s)} \partial_s \left(\frac{1}{u'P^2} \right) \right) ds.$$

Note that ∂_G and $(u'(s))^{-1} \partial_s$ commute. Hence, this term is essentially the same as that treated in Lemma 5.3 (combining also with arguments in Lemma 5.4). Similarly, the remaining terms in (5.30a) are all easy variants of terms we have treated before in Lemmas 5.3, 5.4, and C.1. Hence, the details are omitted for brevity. The calculations involving H_∞ are analogous and are also omitted. This proves (5.31b). Similarly, convergence as $\varepsilon \rightarrow 0$ asserted in part (c) is deduced as in previous arguments.

Next, turn to (5.31c). Taking a $\partial_r \partial_G$ derivative of (5.30a) gives:

$$\begin{aligned} \partial_r \partial_G H_0 &= \partial_r \partial_G \left(\frac{1}{u' P(r, c \pm i\varepsilon)} \right) - u' (\partial_G P) (R_{0,r}^\varepsilon(z) \mp i E_{0,r}^\varepsilon) \\ &\quad - (u - z) (\partial_r \partial_G P) (R_{0,r}^\varepsilon(z) \mp i E_{0,r}^\varepsilon) - (\partial_G P) u'(r) E(r, z) \\ &\quad + (\partial_r \phi) \int_0^r \frac{u'(s)}{(u - z)} \partial_G (\chi_c E) ds + P u'(r) \partial_G (\chi_c E) \\ &\quad - \frac{1}{u'(r_c)} (\partial_r \phi) \int_0^r \chi_{\neq} \frac{u'(s) u'(r_c)}{(u - z)^2} E(s, z) - \frac{u'(s)}{(u - z)} \partial_{r_c} (\chi_{\neq} E(s, z)) ds. \end{aligned}$$

By Theorem 4.3 and the arguments used previously, we can again deduce the desired logarithmically singular upper bounds; the details are omitted as they are repetitive. \square

D Vanishing for $k \geq 2$ Outside I_α

We start by performing energy estimates on the solution to the inhomogeneous Rayleigh problem

$$\left[\partial_{rr} + \frac{1/4 - k^2}{r^2} + \frac{\beta(r)}{u(r) - c \mp i\varepsilon} \right] Y_\varepsilon(r, c) = \frac{F_\varepsilon(r, c)}{u(r) - c \mp i\varepsilon} + F_{*,\varepsilon}(r, c), \quad (\text{D.1})$$

with boundary conditions

$$Y_\varepsilon(0, c) = 0, \quad \lim_{r \rightarrow \infty} Y_\varepsilon(r, c) = 0. \quad (\text{D.2})$$

Problem (D.1) is a slight generalization of (2.13). Notice that, for every $\varepsilon > 0$, (D.1) is just a regular perturbation of Laplace's equation, and therefore

$$Y_\varepsilon(r, c) = \mathcal{O}(r^{k+1/2}) \quad \text{as } r \rightarrow 0, \quad Y_\varepsilon(r, c) = \mathcal{O}(r^{1/2-k}) \quad \text{as } r \rightarrow \infty. \quad (\text{D.3})$$

For this reason, we define the weight

$$w_Y(r) = \min\{r^{k+1}, r^{-k+1}\}, \quad w_{Y,\gamma}(r) = \min\{r^{k+1-\gamma}, r^{-k+1+\gamma}\}, \quad (\text{D.4})$$

and, using the notation as in (1.10a), prove the following theorem.

Theorem D.1 *Let Y_ε be the solution to (D.1)–(D.3), and fix any $\gamma \in (0, 2k)$. Then Y_ε satisfies the following bounds.*

- If $c \in (u(0), u(0) + 1]$, then

$$\begin{aligned} \|Y_\varepsilon(\cdot, c)\|_{L_{\psi,\gamma}^\infty}^2 + \|Y_\varepsilon(\cdot, c)\|_{L_{Y,\gamma}^2}^2 + \|r \partial_r Y_\varepsilon(\cdot, c)\|_{L_{Y,\gamma}^2}^2 &\lesssim_\gamma \left\| \langle r \rangle^2 F_\varepsilon(\cdot, c) \right\|_{L_{Y,\gamma}^2}^2 \\ &\quad + \|r^2 F_{*,\varepsilon}(\cdot, c)\|_{L_{Y,\gamma}^2}^2. \end{aligned} \quad (\text{D.5})$$

- If $c \in (0, u(0))$ is such that $r_c \leq \varepsilon^{\frac{1}{2+\alpha}}$, then

$$\begin{aligned} & \|Y_\varepsilon(\cdot, c)\|_{L_{\psi, \gamma}^\infty}^2 + \|Y_\varepsilon(\cdot, c)\|_{L_{Y, \gamma}^2}^2 + \|r\partial_r Y_\varepsilon(\cdot, c)\|_{L_{Y, \gamma}^2}^2 \\ & \lesssim_\gamma \frac{1}{r_c^{2\alpha}} \left[\|\langle r \rangle^{2+\alpha} F_\varepsilon(\cdot, c)\|_{L_{Y, \gamma}^2}^2 + \|r^2 F_{*, \varepsilon}(\cdot, c)\|_{L_{Y, \gamma}^2}^2 \right]. \end{aligned} \quad (\text{D.6})$$

- If $c > u(0) + 1$, then

$$\begin{aligned} & \|Y_\varepsilon(\cdot, c)\|_{L_{\psi, \gamma}^\infty}^2 + \|Y_\varepsilon(\cdot, c)\|_{L_{Y, \gamma}^2}^2 + \|r\partial_r Y_\varepsilon(\cdot, c)\|_{L_{Y, \gamma}^2}^2 \\ & \lesssim_\gamma \frac{1}{c^2} \|r^2 F_\varepsilon(\cdot, c)\|_{L_{Y, \gamma}^2}^2 + \|r^2 F_{*, \varepsilon}(\cdot, c)\|_{L_{Y, \gamma}^2}^2. \end{aligned} \quad (\text{D.7})$$

- If $c \leq 0$, or $c \in (0, u(0))$ is such that $r_c \geq \frac{1}{2}\varepsilon^{-\frac{1}{2+\alpha}}$, then for every $0 < \tilde{\alpha} < \gamma$ we have

$$\begin{aligned} & \|Y_\varepsilon(\cdot, c)\|_{L_{\psi, \gamma}^\infty}^2 + \|Y_\varepsilon(\cdot, c)\|_{L_{Y, \gamma}^2}^2 + \|r\partial_r Y_\varepsilon(\cdot, c)\|_{L_{Y, \gamma}^2}^2 \\ & \lesssim_\gamma \|r^2 \langle r \rangle^{2+\alpha} F_\varepsilon(\cdot, c)\|_{L_{Y, \gamma-\tilde{\alpha}}^2}^2 + \|r^2 F_{*, \varepsilon}(\cdot, c)\|_{L_{Y, \gamma-\tilde{\alpha}}^2}^2. \end{aligned} \quad (\text{D.8})$$

- There exists $R_\gamma \gg 1$ such that if $c < -R_\gamma$, then

$$\begin{aligned} & \|Y_\varepsilon(\cdot, c)\|_{L_{\psi, \gamma}^\infty}^2 + \|Y_\varepsilon(\cdot, c)\|_{L_{Y, \gamma}^2}^2 + \|r\partial_r Y_\varepsilon(\cdot, c)\|_{L_{Y, \gamma}^2}^2 \\ & \lesssim_\gamma \frac{1}{c^2} \|r^2 F_\varepsilon(\cdot, c)\|_{L_{Y, \gamma}^2}^2 + \|r^2 F_{*, \varepsilon}(\cdot, c)\|_{L_{Y, \gamma}^2}^2. \end{aligned} \quad (\text{D.9})$$

The proof of Theorem D.1 is based mostly on energy estimates, along with the Sobolev inequality

$$\|Y_\varepsilon(\cdot, c)\|_{L^\infty}^2 \leq \int_0^\infty |\partial_r Y_\varepsilon(r, c)|^2 r dr + \int_0^\infty \frac{1}{r^2} |Y_\varepsilon(r, c)|^2 r dr. \quad (\text{D.10})$$

We take the real and imaginary parts of (D.1), to obtain the system

$$\begin{aligned} & \left[\partial_{rr} + \frac{1/4 - k^2}{r^2} \right] \text{Re } Y_\varepsilon + \frac{\beta(u-c)\text{Re } Y_\varepsilon}{(u-c)^2 + \varepsilon^2} \mp \frac{\varepsilon\beta\text{Im } Y_\varepsilon}{(u-c)^2 + \varepsilon^2} = \frac{(u-c)\text{Re } F_\varepsilon}{(u-c)^2 + \varepsilon^2} \\ & \mp \frac{\varepsilon\text{Im } F_\varepsilon}{(u-c)^2 + \varepsilon^2} + \text{Re } F_{*, \varepsilon} \end{aligned} \quad (\text{D.11})$$

$$\begin{aligned} & \left[\partial_{rr} + \frac{1/4 - k^2}{r^2} \right] \text{Im } Y_\varepsilon + \frac{\beta(u-c)\text{Im } Y_\varepsilon}{(u-c)^2 + \varepsilon^2} \pm \frac{\varepsilon\beta\text{Re } Y_\varepsilon}{(u-c)^2 + \varepsilon^2} = \frac{(u-c)\text{Im } F_\varepsilon}{(u-c)^2 + \varepsilon^2} \\ & \mp \frac{\varepsilon\text{Re } F_\varepsilon}{(u-c)^2 + \varepsilon^2} + \text{Im } F_{*, \varepsilon}. \end{aligned} \quad (\text{D.12})$$

Let

$$\gamma_* \in (-k + 1/2, k + 1/2).$$

We multiply (D.11) by $\operatorname{Re} Y_\varepsilon r^{1-2\gamma_*}$ and (D.12) by $\operatorname{Im} Y_\varepsilon r^{1-2\gamma_*}$, integrate by parts using (D.3), and add the result to obtain

$$\begin{aligned} & \int_0^\infty \left| \partial_r \frac{Y_\varepsilon(r, c)}{r^{\gamma_*}} \right|^2 r dr + \left[k^2 - \frac{1}{4} - \gamma_*(\gamma_* - 1) \right] \int_0^\infty \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*+2}} r dr \\ & + \int_0^\infty \frac{\beta(r)(c - u(r))}{(u(r) - c)^2 + \varepsilon^2} \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*}} r dr \\ & \lesssim \int_0^\infty \frac{|F_\varepsilon(r, c)| |Y_\varepsilon(r, c)|}{\sqrt{(u(r) - c)^2 + \varepsilon^2}} \frac{1}{r^{2\gamma_*}} r dr + \int_0^\infty |F_{*,\varepsilon}(r, c)| |Y_\varepsilon(r, c)| \frac{1}{r^{2\gamma_*}} r dr. \end{aligned} \quad (\text{D.13})$$

Since for any small $\kappa > 0$ there holds

$$\int_0^\infty |F_{*,\varepsilon}(r, c)| |Y_\varepsilon(r, c)| \frac{1}{r^{2\gamma_*}} r dr \leq \kappa \int_0^\infty \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*+2}} r dr + \kappa^{-1} \int_0^\infty \frac{|F_{*,\varepsilon}(r, c)|^2}{r^{2\gamma_*-3}} dr, \quad (\text{D.14})$$

we obtain

$$\begin{aligned} & \int_0^\infty \left| \partial_r \frac{Y_\varepsilon(r, c)}{r^{\gamma_*}} \right|^2 r dr + \left[k^2 - \frac{1}{4} - \gamma_*(\gamma_* - 1) \right] \int_0^\infty \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*+2}} r dr \\ & + \int_0^\infty \frac{\beta(r)(c - u(r))}{(u(r) - c)^2 + \varepsilon^2} \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*}} r dr \\ & \lesssim \int_0^\infty \frac{|F_\varepsilon(r, c)| |Y_\varepsilon(r, c)|}{\sqrt{(u(r) - c)^2 + \varepsilon^2}} \frac{1}{r^{2\gamma_*}} r dr + \int_0^\infty \frac{|F_{*,\varepsilon}(r, c)|^2}{r^{2\gamma_*-3}} dr. \end{aligned} \quad (\text{D.15})$$

Cross multiplying (D.11) and (D.12) and subtracting gives

$$\begin{aligned} \varepsilon \int_0^\infty \frac{\beta(r) |Y_\varepsilon(r, c)|^2}{(u(r) - c)^2 + \varepsilon^2} \frac{1}{r^{2\gamma_*}} r dr & \lesssim \int_0^\infty \frac{|F_\varepsilon(r, c)| |Y_\varepsilon(r, c)|}{\sqrt{(u(r) - c)^2 + \varepsilon^2}} \frac{1}{r^{2\gamma_*}} r dr \\ & + \int_0^\infty \frac{|F_{*,\varepsilon}(r, c)| |Y_\varepsilon(r, c)|}{r^{2\gamma_*-1}} dr. \end{aligned} \quad (\text{D.16})$$

We use (D.15) and (D.16) in different ways, depending on the various regimes considered.

D.1 Estimates Near $c = u(0)$

We start by proving estimates (D.5)–(D.7).

D.1.1 Proof of (D.5)

When $c \in (u(0), u(0) + 1]$, then

$$c - u(r) \geq c - u(0) > 0, \quad \forall r \in [0, \infty).$$

From (D.15) and using standard arguments, we obtain

$$\begin{aligned} \int_0^\infty \frac{|F_\varepsilon(r, c)| |Y_\varepsilon(r, c)|}{\sqrt{(u(r) - c)^2 + \varepsilon^2}} \frac{1}{r^{2\gamma_*}} r dr &\leq \kappa \int_0^\infty \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*+2}} r dr \\ &\quad + \frac{1}{\kappa} \int_0^\infty \frac{|F_\varepsilon(r, c)|^2}{r^{2\gamma_*-2}(c - u(r))^2} r dr, \end{aligned} \quad (\text{D.17})$$

for any $\kappa \in (0, 1)$. Moreover, since $u(0) - u(r) \sim r^2$ near $r = 0$, we can write

$$\begin{aligned} \int_0^\infty \frac{|F_\varepsilon(r, c)|^2}{r^{2\gamma_*-2}(c - u(r))^2} r dr &\leq \int_0^1 \frac{|F_\varepsilon(r, c)|^2}{r^{2\gamma_*-2}(u(0) - u(r))^2} r dr \\ &\quad + \int_1^\infty \frac{|F_\varepsilon(r, c)|^2}{r^{2\gamma_*-2}(u(0) - u(r))^2} r dr \\ &\lesssim \int_0^1 \frac{|F_\varepsilon(r, c)|^2}{r^{2\gamma_*+2}} r dr + \int_1^\infty \frac{|F_\varepsilon(r, c)|^2}{r^{2\gamma_*-2}} r dr \\ &\lesssim \int_0^\infty \langle r \rangle^4 \frac{|F_\varepsilon(r, c)|^2}{r^{2\gamma_*+1}} dr. \end{aligned}$$

Hence, from the above estimates and (D.15), we obtain by taking $\kappa \in (0, 1)$ small enough, that

$$\begin{aligned} \int_0^\infty \left| \partial_r \frac{Y_\varepsilon(r, c)}{r^{\gamma_*}} \right|^2 r dr &+ \left[k^2 - \frac{1}{4} - \gamma_*(\gamma_* - 1) \right] \int_0^\infty \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*+2}} r dr \\ &+ \int_0^\infty \frac{\beta(r)(c - u(r))}{(u(r) - c)^2 + \varepsilon^2} \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*}} r dr \\ &\lesssim \int_0^\infty \langle r \rangle^4 \frac{|F_\varepsilon(r, c)|^2}{r^{2\gamma_*+1}} dr + \int_0^\infty \frac{|F_{*,\varepsilon}(r, c)|^2}{r^{2\gamma_*-3}} dr. \end{aligned} \quad (\text{D.18})$$

An application of (D.10) gives the estimate

$$\begin{aligned} \|r^{-\gamma_*} Y_\varepsilon(\cdot, c)\|_{L^\infty}^2 &+ \int_0^\infty \left| \partial_r \frac{Y_\varepsilon(r, c)}{r^{\gamma_*}} \right|^2 r dr + \int_0^\infty \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*+2}} r dr \\ &\lesssim_{\gamma_*} \int_0^\infty \langle r \rangle^4 \frac{|F_\varepsilon(r, c)|^2}{r^{2\gamma_*+1}} dr + \int_0^\infty \frac{|F_{*,\varepsilon}(r, c)|^2}{r^{2\gamma_*-3}} dr. \end{aligned}$$

By considering the left-hand side above restricted to $(0, 1)$ and choosing $\gamma_* = k + 1/2 - \gamma$, with $\gamma \in (0, k]$, we obtain

$$\begin{aligned} & \|r^{-k-1/2+\gamma} Y_\varepsilon(\cdot, c)\|_{L^\infty(0,1)}^2 + \|Y_\varepsilon(\cdot, c)\|_{L^2_{Y,\gamma}(0,1)}^2 + \|r \partial_r Y_\varepsilon(\cdot, c)\|_{L^2_{Y,\gamma}(0,1)}^2 \\ & \lesssim_\gamma \int_0^\infty \langle r \rangle^4 \frac{|F_\varepsilon(r, c)|^2}{r^{2\gamma_*+1}} dr + \int_0^\infty \frac{|F_{*,\varepsilon}(r, c)|^2}{r^{2\gamma_*-3}} dr \\ & \lesssim_\gamma \left\| \langle r \rangle^2 F_\varepsilon(\cdot, c) \right\|_{L^2_{Y,\gamma}}^2 + \|r^2 F_{*,\varepsilon}(\cdot, c)\|_{L^2_{Y,\gamma}}^2. \end{aligned} \quad (\text{D.19})$$

Similarly, on $(1, \infty)$ we choose $\gamma_* = -k + 1/2 + \gamma$, with $\gamma \in (0, k]$, and deduce that

$$\begin{aligned} & \|r^{k-1/2-\gamma} Y_\varepsilon(\cdot, c)\|_{L^\infty(1,\infty)}^2 + \|Y_\varepsilon(\cdot, c)\|_{L^2_{Y,\gamma}(1,\infty)}^2 + \|r \partial_r Y_\varepsilon(\cdot, c)\|_{L^2_{Y,\gamma}(1,\infty)}^2 \\ & \lesssim_\gamma \left\| \langle r \rangle^2 F_\varepsilon(\cdot, c) \right\|_{L^2_{Y,\gamma}}^2 + \|r^2 F_{*,\varepsilon}(\cdot, c)\|_{L^2_{Y,\gamma}}^2. \end{aligned} \quad (\text{D.20})$$

Adding the above two estimates together and recalling the shape of the weight (D.4), we deduce (D.5). Note that we can extend the range of γ to the interval $(0, 2k)$, as the weight $w_{Y,\gamma}$ is symmetric about $\gamma = k$.

D.1.2 Proof of (D.6)

When $c \in (0, u(0))$ complying with $r_c \leq \varepsilon^{\frac{1}{2+\alpha}}$, from (D.13) and using a similar argument to that in (D.17), we have

$$\begin{aligned} & \int_0^\infty \left| \partial_r \frac{Y_\varepsilon(r, c)}{r^{\gamma_*}} \right|^2 r dr + \left[k^2 - \frac{1}{4} - \gamma_*(\gamma_* - 1) \right] \int_0^\infty \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*+2}} r dr \\ & + \int_{r_c}^\infty \frac{\beta(r)(c - u(r))}{(u(r) - c)^2 + \varepsilon^2} \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*}} r dr \\ & \lesssim \int_0^{r_c} \frac{\beta(r)(u(r) - c)}{(u(r) - c)^2 + \varepsilon^2} \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*}} r dr + \int_0^\infty \frac{|F_\varepsilon(r, c)| |Y_\varepsilon(r, c)|}{\sqrt{(u(r) - c)^2 + \varepsilon^2}} \frac{1}{r^{2\gamma_*}} r dr \\ & + \int_0^\infty \frac{|F_{*,\varepsilon}(r, c)| |Y_\varepsilon(r, c)|}{r^{2\gamma_*-1}} dr. \end{aligned}$$

For $r \leq r_c \leq 1$ there holds $|c - u(r)| \lesssim r_c^2$ and hence by (D.16) and therefore

$$\begin{aligned} \int_0^{r_c} \frac{\beta(r)(u(r) - c)}{(u(r) - c)^2 + \varepsilon^2} \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*}} r dr & \lesssim \frac{1}{r_c^\alpha} \left[\int_0^\infty \frac{|F_\varepsilon(r, c)| |Y_\varepsilon(r, c)|}{\sqrt{(u(r) - c)^2 + \varepsilon^2}} \frac{1}{r^{2\gamma_*}} r dr \right. \\ & \left. + \int_0^\infty \frac{|F_{*,\varepsilon}(r, c)| |Y_\varepsilon(r, c)|}{r^{2\gamma_*-1}} dr \right], \end{aligned}$$

and since $r_c \ll 1$, we obtain

$$\begin{aligned} & \int_0^\infty \left| \partial_r \frac{Y_\varepsilon(r, c)}{r^{\gamma_*}} \right|^2 r dr + \left[k^2 - \frac{1}{4} - \gamma_*(\gamma_* - 1) \right] \int_0^\infty \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*+2}} r dr \\ & + \int_{r_c}^\infty \frac{\beta(r)(c - u(r))}{(u(r) - c)^2 + \varepsilon^2} \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*}} r dr \\ & \lesssim \frac{1}{r_c^\alpha} \left[\int_0^\infty \frac{|F_\varepsilon(r, c)| |Y_\varepsilon(r, c)|}{\sqrt{(u(r) - c)^2 + \varepsilon^2}} \frac{1}{r^{2\gamma_*}} r dr + \int_0^\infty |F_{*,\varepsilon}(r, c)| |Y_\varepsilon(r, c)| \frac{1}{r^{2\gamma_*}} r dr \right]. \end{aligned}$$

Now, for any $\kappa \in (0, 1)$ we have

$$\begin{aligned} \frac{1}{r_c^\alpha} \int_0^\infty \frac{|F_\varepsilon(r, c)| |Y_\varepsilon(r, c)|}{\sqrt{(u(r) - c)^2 + \varepsilon^2}} \frac{1}{r^{2\gamma_*}} r dr & \leq \kappa \int_0^\infty \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*+2}} r dr \\ & + \kappa^{-1} \frac{1}{r_c^{2\alpha}} \int_0^\infty \frac{|F_\varepsilon(r, c)|^2}{(u(r) - c)^2 + \varepsilon^2} \frac{1}{r^{2\gamma_*-2}} r dr. \end{aligned}$$

The contribution from F_ε is controlled using that $r^{2+\alpha} \lesssim \langle r \rangle^{2+\alpha} \sqrt{(u - c)^2 + \varepsilon^2}$, which implies that

$$\begin{aligned} \int_0^\infty \frac{|F_\varepsilon(r, c)|^2}{(u(r) - c)^2 + \varepsilon^2} \frac{1}{r^{2\gamma_*-2}} r dr & = \int_0^\infty \frac{r^{4+2\alpha} |F_\varepsilon(r, c)|^2}{(u(r) - c)^2 + \varepsilon^2} \frac{1}{r^{2\gamma_*+2+2\alpha}} r dr \\ & \lesssim \int_0^\infty \langle r \rangle^{4+2\alpha} \frac{|F_\varepsilon(r, c)|^2}{r^{2\gamma_*+1+2\alpha}} dr. \end{aligned}$$

On F_* , we use again (D.14). Therefore, choosing $\kappa \ll 1$ and arguing as in (D.19)–(D.20) yields the desired result.

D.1.3 Proof of (D.7)

The starting point here is again (D.15), together with (D.17), which allows us to write

$$\begin{aligned} & \int_0^\infty \left| \partial_r \frac{Y_\varepsilon(r, c)}{r^{\gamma_*}} \right|^2 r dr + \left[k^2 - \frac{1}{4} - \gamma_*(\gamma_* - 1) \right] \int_0^\infty \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*+2}} r dr \\ & + \int_0^\infty \frac{\beta(r)(c - u(r))}{(u(r) - c)^2 + \varepsilon^2} \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*}} r dr \\ & \lesssim \int_0^\infty \frac{|F_\varepsilon(r, c)|^2}{r^{2\gamma_*-2}(c - u(r))^2} r dr + \int_0^\infty \frac{|F_{*,\varepsilon}(r, c)|^2}{r^{2\gamma_*-3}} dr \lesssim \frac{1}{c^2} \int_0^\infty \frac{|F_\varepsilon(r, c)|^2}{r^{2\gamma_*-3}} dr \\ & + \int_0^\infty \frac{|F_{*,\varepsilon}(r, c)|^2}{r^{2\gamma_*-3}} dr, \end{aligned} \tag{D.21}$$

and (D.7) follows from similar arguments as in (D.19)–(D.20).

D.2 Estimates Near $c = 0$

The core of this section lies in the proof of (D.8). As we shall see below, (D.9) follows by the argument already used for (D.7).

D.2.1 Proof of (D.8)

In this case, the argument to prove uniform estimates is completely different. We aim to prove the following.

Lemma D.2 *Let Y_ε be the solution to (D.1)–(D.3), and fix $0 < \tilde{\alpha} < \gamma < 2k$. Then, for every $c \leq 0$, or any $c \in (0, u(0))$ such that*

$$r_c \geq \frac{1}{2} \varepsilon^{-\frac{1}{2+\alpha}}, \quad (\text{D.22})$$

we have that

$$\|Y_\varepsilon(\cdot, c)\|_{L^2_{Y,\gamma}} + \|r\partial_r Y_\varepsilon(\cdot, c)\|_{L^2_{Y,\gamma}} \lesssim_\gamma \|r^2 \langle r \rangle^{2+\alpha} F_\varepsilon(\cdot, c)\|_{L^2_{Y,\gamma-\tilde{\alpha}}} + \|r^2 F_{*,\varepsilon}(\cdot, c)\|_{L^2_{Y,\gamma-\tilde{\alpha}}}. \quad (\text{D.23})$$

Above, α can be take zero when $c \leq 0$. In particular, using (D.10), then (D.8) holds.

The proof of this fact is split into different lemmas. To begin with, we need an estimate with sharper weights on Y_ε in terms of slightly weaker weights.

Lemma D.3 *Let Y_ε be the solution to (D.1)–(D.3). Then, for every $c \leq 0$, and $0 < \tilde{\alpha} < \gamma < 2k$ we have that*

$$\|Y_\varepsilon(\cdot, c)\|_{L^2_{Y,\gamma}} \lesssim \left\| r^2 \langle r \rangle^2 F_\varepsilon(\cdot, c) \right\|_{L^2_{Y,\gamma-\tilde{\alpha}}} + \|r^2 F_{*,\varepsilon}(\cdot, c)\|_{L^2_{Y,\gamma-\tilde{\alpha}}} + \|Y_\varepsilon(\cdot, c)\|_{L^2_{Y,2\gamma}} \quad (\text{D.24})$$

and

$$\|r\partial_r Y_\varepsilon(\cdot, c)\|_{L^2_{Y,\gamma}} \lesssim \left\| r^2 \langle r \rangle^2 F_\varepsilon(\cdot, c) \right\|_{L^2_{Y,\gamma-\tilde{\alpha}}} + \|r^2 F_{*,\varepsilon}(\cdot, c)\|_{L^2_{Y,\gamma-\tilde{\alpha}}} + \|Y_\varepsilon(\cdot, c)\|_{L^2_{Y,2\gamma}}. \quad (\text{D.25})$$

When $c \in (0, u(0))$ is such that

$$r_c \geq \frac{1}{2} \varepsilon^{-\frac{1}{2+\alpha}},$$

then

$$\|Y_\varepsilon(\cdot, c)\|_{L^2_{Y,\gamma}} \lesssim \left\| r^2 \langle r \rangle^{2+\alpha} F_\varepsilon(\cdot, c) \right\|_{L^2_{Y,\gamma-\tilde{\alpha}}} + \|r^2 F_{*,\varepsilon}(\cdot, c)\|_{L^2_{Y,\gamma-\tilde{\alpha}}} + \|Y_\varepsilon(\cdot, c)\|_{L^2_{Y,2\gamma}} \quad (\text{D.26})$$

and

$$\|r\partial_r Y_\varepsilon(\cdot, c)\|_{L^2_{Y,\gamma}} \lesssim \|r^2\langle r \rangle^{2+\alpha} F_\varepsilon(\cdot, c)\|_{L^2_{Y,\gamma-\tilde{\alpha}}} + \|r^2 F_{*,\varepsilon}(\cdot, c)\|_{L^2_{Y,\gamma-\tilde{\alpha}}} + \|Y_\varepsilon(\cdot, c)\|_{L^2_{Y,2\gamma}}. \quad (\text{D.27})$$

Proof of Lemma D.3 By using the Green's function (4.91), we can deduce from (D.1) that

$$\begin{aligned} Y_\varepsilon(r, c) &= \int_0^\infty \mathcal{L}(r, \rho) \left[\frac{F_\varepsilon(\rho, c)}{u(\rho) - c \mp i\varepsilon} + F_{*,\varepsilon}(\rho, c) \right] d\rho \\ &\quad - \int_0^\infty \mathcal{L}(r, \rho) \frac{\beta(\rho)}{u(\rho) - c \mp i\varepsilon} Y_\varepsilon(\rho, c) d\rho. \end{aligned} \quad (\text{D.28})$$

Considering the weights in (D.4) and (1.10c), we need to prove an L^2 bound for

$$\begin{aligned} \frac{Y_\varepsilon(r, c)}{w_{Y,\gamma}(r)} &= \int_0^\infty \frac{w_{Y,\gamma-\tilde{\alpha}}(\rho)\rho^{-2}}{w_{Y,\gamma}(r)} \mathcal{L}(r, \rho) \left[\frac{1}{(u(\rho) - c \mp i\varepsilon) w_{Y,\gamma-\tilde{\alpha}}(\rho)} \frac{\rho^2 F_\varepsilon(\rho, c)}{w_{Y,\gamma-\tilde{\alpha}}(\rho)} \right. \\ &\quad \left. + \frac{\rho^2 F_{*,\varepsilon}(\rho, c)}{w_{Y,\gamma-\tilde{\alpha}}(\rho)} \right] d\rho \\ &\quad - \int_0^\infty \frac{w_{Y,2\gamma}(\rho)}{w_{Y,\gamma}(r)} \mathcal{L}(r, \rho) \frac{\beta(\rho)}{u(\rho) - c \mp i\varepsilon} \frac{Y_\varepsilon(\rho, c)}{w_{Y,2\gamma}(\rho)} d\rho. \end{aligned}$$

If we consider the case $c \leq 0$ first, we observe that since $u(\rho) \sim \langle \rho \rangle^{-2}$ for the first term we need to prove that

$$\int_0^\infty \int_0^\infty \left| \frac{w_{Y,\gamma-\tilde{\alpha}}(\rho)\rho^{-2}}{w_{Y,\gamma}(r)} \mathcal{L}(r, \rho) \right|^2 d\rho dr < \infty, \quad (\text{D.29})$$

which follows from a straightforward calculation. Similarly, we also have

$$\int_0^\infty \int_0^\infty \left| \frac{w_{Y,2\gamma}(\rho)}{w_{Y,\gamma}(r)} \mathcal{L}(r, \rho) \frac{\beta(\rho)}{u(\rho)} \right|^2 d\rho dr < \infty, \quad (\text{D.30})$$

where we need to exploit the fast decay of β at infinity. The derivative estimate follows from applying $r\partial_r$ to (D.28), and arguing in the same way as above.

When $c \in (0, u(0))$ is such that

$$r_c \geq \frac{1}{2} \varepsilon^{-\frac{1}{2+\alpha}},$$

the proof is similar. In this case, the key observation is that if $r > r_c/2$,

$$\langle r \rangle^{2+\alpha} \sqrt{(u(r) - c)^2 + \varepsilon^2} \geq \langle r \rangle^{2+\alpha} \varepsilon \gtrsim \frac{\langle r \rangle^{2+\alpha}}{r_c^{2+\alpha}} \gtrsim 1, \quad (\text{D.31})$$

while if $r \leq r_c/2$, then $|u(r) - c| \gtrsim \langle r \rangle^{-2}$, and thus again

$$\langle r \rangle^{2+\alpha} \sqrt{(u(r) - c)^2 + \varepsilon^2} \geq \langle r \rangle^{2+\alpha} |u(r) - c| \gtrsim 1. \quad (\text{D.32})$$

In view of this, the weight on F has an extra power of $\langle r \rangle^\alpha$. The proof is concluded. \square

We also need a result on the homogeneous Rayleigh problem.

Lemma D.4 Fix $0 < \gamma \ll 1$, and let $\phi \in L^2_{Y,\gamma}$ with $r\partial_r\phi \in L^2_{Y,\gamma}$ be a solution to the homogeneous Rayleigh problem (4.1), for $z = c \leq 0$. Then $\phi \equiv 0$.

Proof of Lemma D.4 If $\phi \equiv 0$ in a neighborhood of the origin, then, by unique continuation, $\phi \equiv 0$ everywhere. Hence, we may assume that $\phi(r, c) > 0$ for every $0 < r \ll 1$. Recall, the function $g(r) = r^{3/2}u(r)$ (Lemma 2.4) satisfies

$$\partial_{rr}g + \left(-\frac{3}{4r^2} + \frac{\beta(r)}{u(r)}\right)g = 0.$$

Hence,

$$g\partial_{rr}\phi - \phi\partial_{rr}g + \frac{1-k^2}{r^2}\phi g + \beta(r)\left(\frac{1}{u(r)-c} - \frac{1}{u(r)}\right)\phi g = 0. \quad (\text{D.33})$$

Let

$$\bar{r} = \sup\{r \in (0, \infty) : \phi(r', c) > 0, \quad \forall r' < r\} \in (0, \infty].$$

We integrate (D.33) on $(0, \bar{r})$. Notice that the functions involved are integrable due to the assumptions on ϕ , even if $\bar{r} = \infty$. Using that

$$g\partial_{rr}\phi - \phi\partial_{rr}g = \partial_r(g\partial_r\phi - \phi\partial_rg),$$

and the fact that since $c \leq 0$ there holds

$$\frac{1}{u(r)-c} - \frac{1}{u(r)} \leq 0,$$

we obtain

$$\begin{aligned} g(\bar{r})\partial_r\phi(\bar{r}, c) - \phi(\bar{r}, c)\partial_rg(\bar{r}) &= \int_0^{\bar{r}} \left[\frac{k^2-1}{r^2}\phi(r, c)g(r) \right. \\ &\quad \left. - \beta(r)\left(\frac{1}{u(r)-c} - \frac{1}{u(r)}\right)\phi(r, c)g(r) \right] dr \geq 0. \end{aligned}$$

Note that $\phi(\bar{r}, c) = 0$, $\partial_r\phi(\bar{r}, c) \leq 0$ and $g > 0$, so that the above implies (also in the case $\bar{r} = \infty$) that

$$\int_0^{\bar{r}} \left[\frac{k^2-1}{r^2}\phi(r, c)g(r) - \beta(r)\left(\frac{1}{u(r)-c} - \frac{1}{u(r)}\right)\phi(r, c)g(r) \right] dr = 0,$$

and therefore $\phi \equiv 0$, concluding the proof of the lemma. \square

We can now complete the proof of Lemma D.2.

Proof of Lemma D.2 Assume for contradiction that (D.23) does not hold, and let

$$M_{F,\varepsilon} = \left\| r^2 \langle r \rangle^{2+\alpha} F_\varepsilon(\cdot, c) \right\|_{L^2_{Y,\gamma-\tilde{\alpha}}} + \| r^2 F_{*,\varepsilon}(\cdot, c) \|_{L^2_{Y,\gamma-\tilde{\alpha}}}.$$

Then there exists a sequence $\varepsilon_j \rightarrow 0$ such that

$$\| Y_{\varepsilon_j}(\cdot, c) \|_{L^2_{Y,\gamma}} > j M_{F,\varepsilon_j}. \quad (\text{D.34})$$

By replacing

$$\begin{aligned} Y_{\varepsilon_j}(r, c) &\mapsto \frac{Y_{\varepsilon_j}(\cdot, c)}{\| Y_{\varepsilon_j}(\cdot, c) \|_{L^2_{Y,\gamma}}}, & F_{\varepsilon_j}(r, c) &\mapsto \frac{F_{\varepsilon_j}(r, c)}{\| Y_{\varepsilon_j}(\cdot, c) \|_{L^2_{Y,\gamma}}}, \\ F_{*,\varepsilon_j}(r, c) &\mapsto \frac{F_{*,\varepsilon_j}(r, c)}{\| Y_{\varepsilon_j}(\cdot, c) \|_{L^2_{Y,\gamma}}}, \end{aligned}$$

we may assume that $\| Y_{\varepsilon_j}(\cdot, c) \|_{L^2_{Y,\gamma}} = 1$ and $\| F_{\varepsilon_j}(\cdot, c) \|_{L^2_{Y,\gamma-\tilde{\alpha}}} + \| F_{*,\varepsilon_j}(\cdot, c) \|_{L^2_{Y,\gamma-\tilde{\alpha}}} \rightarrow 0$. Hence, thanks to standard compactness arguments, Lemma D.3 provides the existence of a subsequence (not relabeled) such that Y_{ε_j} weakly in $L^2_{Y,\gamma/2}$ and strongly in $L^2_{Y,\gamma}$, while $r \partial_r Y_{\varepsilon_j}$ converges weakly in $L^2_{Y,\gamma/2}$. Note that in that case $c \in (0, u(0))$, we have that $r_c \rightarrow \infty$ as $\varepsilon \rightarrow 0$ thanks to (D.22), which is equivalent to say that $c \rightarrow 0$. Hence, the limit $Y \in H^1_{Y,\gamma/2}$, is a weak solution to the *homogeneous* Rayleigh problem (due to $F_{\varepsilon_j} \rightarrow 0$), for some $c \leq 0$, and satisfies $\| Y \|_{L^2_{Y,\gamma}} = 1$. However, Lemma D.4 implies that $Y = 0$, which is a contradiction. With (D.23) at our disposal, the derivative estimate follows from (D.25). \square

D.2.2 Proof of (D.9)

Assume $c < -R_{\gamma_*}$, where $R_{\gamma_*} > 0$ is fixed below. Using the same ideas as in (D.21), we have

$$\begin{aligned} &\int_0^\infty \left| \partial_r \frac{Y_\varepsilon(r, c)}{r^{\gamma_*}} \right|^2 r dr + \left[k^2 - \frac{1}{4} - \gamma_*(\gamma_* - 1) \right] \int_0^\infty \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*+2}} r dr \\ &\lesssim \int_0^\infty \frac{\beta(r)(u(r) - c)}{(u(r) - c)^2 + \varepsilon^2} \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*}} r dr + \frac{1}{c^2} \int_0^\infty \frac{|F_\varepsilon(r, c)|^2}{r^{2\gamma_*-3}} dr \\ &\quad + \int_0^\infty \frac{|F_{*,\varepsilon}(r, c)|^2}{r^{2\gamma_*-3}} dr. \end{aligned} \quad (\text{D.35})$$

Now, since

$$\beta(r)r^2 \leq C_\beta$$

by standard estimates,

$$\begin{aligned} \int_0^\infty \frac{\beta(r)(u(r)-c)}{(u(r)-c)^2 + \varepsilon^2} \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*}} r dr &\leq \int_0^\infty \frac{\beta(r)r^2}{u(r)-c} \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*+2}} r dr \\ &\leq \frac{C_\beta}{R_{\gamma_*}} \int_0^\infty \frac{|Y_\varepsilon(r, c)|^2}{r^{2\gamma_*+2}} r dr. \end{aligned}$$

Thus, by taking

$$R_{\gamma_*} \gg \frac{C_\beta}{k^2 - \frac{1}{4} - \gamma_*(\gamma_* - 1)},$$

estimate (D.9) follows from (D.35).

D.3 Vanishing of f_E

The function f_E from (2.26), can be rewritten using the notation introduced in (2.20) as

$$\begin{aligned} f_E^\varepsilon(t, r) &= \frac{\beta(r)}{2\pi i \sqrt{r}} \int_{\mathbb{R}} e^{ik(u(r)-c)t} \left[\frac{u(r)-c}{(u(r)-c)^2 + \varepsilon^2} X(r, c, \varepsilon) \right. \\ &\quad \left. + \frac{i\varepsilon A(r, c, \varepsilon)}{(u(r)-c)^2 + \varepsilon^2} \right] (1 - \chi_\sigma(c)) dc, \end{aligned} \quad (\text{D.36})$$

where

$$\text{RAY}_+ X = \frac{1}{u(r)-c-i\varepsilon} \frac{2i\varepsilon[F(r) - \beta(r)Y(r, c-i\varepsilon)]}{u(r)-c+i\varepsilon}, \quad (\text{D.37})$$

and

$$\text{RAY}_+ A = \frac{1}{u(r)-c-i\varepsilon} \frac{2(u(r)-c)F(r) + 2i\varepsilon\beta(r)Y(r, c-i\varepsilon)}{u(r)-c+i\varepsilon} + 2F_*(r). \quad (\text{D.38})$$

According to (D.7) and (D.9), if $c > u(0) + 1/2$ or $c < R_\gamma$ we use Lemma 2.1 and the fact that $1 \lesssim |u(r)-c|^2$ to obtain that

$$\begin{aligned} &\left\| \frac{X(\cdot, c, \varepsilon)}{\min\{r^{k+1/2-\gamma}, r^{-k+1/2+\gamma}\}} \right\|_{L^\infty}^2 + \|X(\cdot, c, \varepsilon)\|_{L_{Y,\gamma}^2}^2 + \|r\partial_r X(\cdot, c, \varepsilon)\|_{L_{Y,\gamma}^2}^2 \\ &\lesssim_\gamma \frac{\varepsilon^2}{c^2} \|r^2 F\|_{L_{Y,\gamma}^2}^2 + \frac{\varepsilon^2}{c^2} \|r^2 \beta Y(\cdot, c-i\varepsilon)\|_{L_{Y,\gamma}^2}^2 \end{aligned}$$

$$\begin{aligned}
&\lesssim_{\mathcal{Y}} \frac{\varepsilon^2}{c^4} \|r^2 F\|_{L_{Y,\mathcal{Y}}^2}^2 + \frac{\varepsilon^2}{c^2} \|Y(\cdot, c - i\varepsilon)\|_{L_{Y,\mathcal{Y}}^2}^2 \\
&\lesssim_{\mathcal{Y}} \frac{\varepsilon^2}{c^4} \left[\|r^2 F\|_{L_{Y,\mathcal{Y}}^2}^2 + \|r^2 F_*\|_{L_{Y,\mathcal{Y}}^2}^2 \right],
\end{aligned} \tag{D.39}$$

and, similarly

$$\left\| \frac{A(\cdot, c, \varepsilon)}{\min\{r^{k+1/2-\gamma}, r^{-k+1/2+\gamma}\}} \right\|_{L^\infty}^2 \lesssim_{\mathcal{Y}} \|r^2 F\|_{L_{Y,\mathcal{Y}}^2}^2 + \|r^2 F_*\|_{L_{Y,\mathcal{Y}}^2}^2. \tag{D.40}$$

Proposition D.5 *For δ sufficiently small, there holds*

$$\lim_{\varepsilon \rightarrow 0} \|f_E^\varepsilon(t, \cdot)\|_{L_{f,\delta}^2} = 0,$$

for every $t \geq 0$.

Proof of Proposition D.5 Recalling the shape of the weight (1.10b), we split into different cases. If $r \leq 1$, we use (D.39) and (D.40) to obtain

$$\begin{aligned}
\frac{|f_E^\varepsilon(t, r)|}{r^{k+1/2-\delta}} &\lesssim \frac{\beta(r)}{r^{k+1-\delta}} \int_{\mathbb{R}} \frac{|X(r, c, \varepsilon)|}{|c|} (1 - \chi_\sigma(c)) dc \\
&\quad + \varepsilon \frac{\beta(r)}{r^{k+1-\delta}} \int_{\mathbb{R}} \frac{|A(r, c, \varepsilon)|}{|c|^2} (1 - \chi_\sigma(c)) dc \\
&\lesssim_{\delta, F, F_*} \frac{\beta(r)}{r^{1/2-\delta/2}} \varepsilon.
\end{aligned}$$

If $r \geq 1$, we get in a similar manner that

$$\begin{aligned}
\frac{|f_E^\varepsilon(t, r)|}{r^{-k+1/2-4+\delta}} &\lesssim \frac{\beta(r)}{r^{-k-3+\delta}} \int_{\mathbb{R}} \frac{|X(r, c, \varepsilon)|}{|c|} (1 - \chi_\sigma(c)) dc \\
&\quad + \varepsilon \frac{\beta(r)}{r^{-k-3+\delta}} \int_{\mathbb{R}} \frac{|A(r, c, \varepsilon)|}{|c|^2} (1 - \chi_\sigma(c)) dc \\
&\lesssim_{\delta, F, F_*} \frac{\beta(r)}{r^{-7/2+\delta/2}} \varepsilon,
\end{aligned}$$

Upon squaring and integrating over $r \in (0, \infty)$ and using Lemma 2.1, we deduce that

$$\|f_E(t, \cdot, \varepsilon)\|_{L_{f,\delta}^2} \lesssim_{\delta, F, F_*} \varepsilon,$$

concluding the proof. \square

D.4 Vanishing of f_S

We next treat f_S in (2.25), which we rewrite here

$$f_S^\varepsilon(t, r) = \frac{\beta(r)}{2\pi i \sqrt{r}} \int_{-R_\delta}^{u(0)+1} e^{ik(u(r)-c)t} \left[\frac{Y(r, c+i\varepsilon)}{u(r)-c-i\varepsilon} - \frac{Y(r, c-i\varepsilon)}{u(r)-c+i\varepsilon} \right] \chi_\sigma(c)(1-\chi_I(r_c))dc. \quad (\text{D.41})$$

Recall that

$$1 - \chi_I(r_c) \neq 0 \quad \text{if} \quad r_c \leq \varepsilon^{\frac{1}{2+\alpha}} \quad \text{or} \quad r_c \geq \frac{1}{2} \varepsilon^{-\frac{1}{2+\alpha}}. \quad (\text{D.42})$$

We prove the following result.

Proposition D.6 *For δ sufficiently small, there holds*

$$\lim_{\varepsilon \rightarrow 0} \|f_S^\varepsilon(t, \cdot)\|_{L^2_{f,\delta}} = 0,$$

for every $t \geq 0$.

Proof of Proposition D.6 Since the cut-off functions in (D.41) isolate different subsets of \mathbb{R} , we proceed case by case, using the estimates provided by Theorem D.1. We first note that

$$\frac{|f_S^\varepsilon(t, r)|}{w_{f,\delta}(r)} \lesssim J_1 + J_2 + J_3 + J_4$$

where

$$J_1^\varepsilon(t, r) = \frac{\beta(r)}{w_{f,\delta}(r)r^{1/2}} \left| \int_{r_c \leq \varepsilon^{\frac{1}{2+\alpha}}}^{u(0)} e^{ik(u(r)-c)t} \left[\frac{Y(r, c+i\varepsilon)}{u(r)-c-i\varepsilon} - \frac{Y(r, c-i\varepsilon)}{u(r)-c+i\varepsilon} \right] dc \right|, \quad (\text{D.43})$$

$$J_2^\varepsilon(t, r) = \frac{\beta(r)}{w_{f,\delta}(r)r^{1/2}} \left| \int_{u(0)}^{u(0)+1} e^{ik(u(r)-c)t} \left[\frac{u(r)-c}{(u(r)-c)^2 + \varepsilon^2} X(r, c, \varepsilon) + \frac{i\varepsilon A(r, c, \varepsilon)}{(u(r)-c)^2 + \varepsilon^2} \right] dc \right|, \quad (\text{D.44})$$

$$J_3^\varepsilon(t, r) = \frac{\beta(r)}{w_{f,\delta}(r)r^{1/2}} \left| \int_{-R_\delta}^0 e^{ik(u(r)-c)t} \left[\frac{u(r)-c}{(u(r)-c)^2 + \varepsilon^2} X(r, c, \varepsilon) + \frac{i\varepsilon}{(u(r)-c)^2 + \varepsilon^2} A(r, c, \varepsilon) \right] dc \right|, \quad (\text{D.45})$$

$$J_4^\varepsilon(t, r) = \frac{\beta(r)}{w_{f,\delta}(r)r^{1/2}} \left| \int_0^{r_c \geq \frac{1}{2}\varepsilon^{-\frac{1}{2+\alpha}}} e^{ik(u(r)-c)t} \left[\frac{Y(r, c+i\varepsilon)}{u(r)-c-i\varepsilon} - \frac{Y(r, c-i\varepsilon)}{u(r)-c+i\varepsilon} \right] dc \right|, \quad (\text{D.46})$$

and X and A satisfy (D.37) and (D.38), respectively. Let us first consider J_1 . The key observation is that on the support of the integrands in r_c , there holds for $r \leq 1$,

$$r^\delta \lesssim \left((u(r) - u(r_c))^2 + \varepsilon^2 \right)^{\frac{\delta}{2(2+\alpha)}},$$

whereas for $r > 1$ and $r_c < 1/4$, there holds

$$1 \lesssim (u(r) - u(r_c))^2 + \varepsilon^2.$$

Moreover, $u'(r_c) \approx r_c$. Hence, by (D.6) if $r \leq 1$ we have

$$\begin{aligned} & \left| \int_{r_c \leq \varepsilon^{\frac{1}{2+\alpha}}}^{u(0)} e^{ik(u(r)-c)t} \frac{Y(r, c+i\varepsilon)}{u(r)-c-i\varepsilon} (1 - \chi_\sigma(c)) dc \right| \\ & \lesssim \int_0^{\varepsilon^{\frac{1}{2+\alpha}}} |u'(r_c)| \frac{|Y(r, c+i\varepsilon)|}{\sqrt{(u(r) - u(r_c))^2 + \varepsilon^2}} dr_c \\ & \lesssim \int_0^{\varepsilon^{\frac{1}{2+\alpha}}} r_c \frac{r^{-\delta/4} |Y(r, c+i\varepsilon)|}{((u(r) - u(r_c))^2 + \varepsilon^2)^{\frac{1}{2} - \frac{\delta}{8(2+\alpha)}}} dr_c \\ & \lesssim \frac{1}{\varepsilon^{1 - \frac{\delta}{4(2+\alpha)}}} \int_0^{\varepsilon^{\frac{1}{2+\alpha}}} r_c r^{-\delta/4} |Y(r, c+i\varepsilon)| dr_c \\ & \lesssim \frac{r^{k+1/2-\delta/2}}{\varepsilon^{1 - \frac{\delta}{4(2+\alpha)}}} \int_0^{\varepsilon^{\frac{1}{2+\alpha}}} r_c \|r^{-k-1/2+\delta/4} Y(\cdot, c+i\varepsilon)\|_{L^\infty(0,1)} dr_c \\ & \lesssim_{F, F_*, \delta} r^{k+1/2-\delta/2} \varepsilon^{\frac{\delta-8\alpha}{4(2+\alpha)}}, \end{aligned}$$

while if $r > 1$ there holds

$$\begin{aligned} & \left| \int_{r_c \leq \varepsilon^{\frac{1}{2+\alpha}}}^{u(0)} e^{ik(u(r)-c)t} \frac{Y(r, c+i\varepsilon)}{u(r)-c-i\varepsilon} (1 - \chi_\sigma(c)) dc \right| \\ & \lesssim \int_0^{\varepsilon^{\frac{1}{2+\alpha}}} r_c |Y(r, c+i\varepsilon)| dr_c \\ & \lesssim r^{-k+1/2+\delta/2} \int_0^{\varepsilon^{\frac{1}{2+\alpha}}} r_c \|r^{k-1/2-\delta/2} Y(\cdot, c+i\varepsilon)\|_{L^\infty(1,\infty)} dr_c \end{aligned}$$

$$\lesssim_{F, F_*, \delta} r^{-k+1/2+\delta/2} \varepsilon^{\frac{2-\alpha}{2+\alpha}},$$

and the same holds for $Y(r, c - i\varepsilon)$. Hence we have

$$|J_1^\varepsilon(t, r)| \lesssim_{F, F_*, \delta} \begin{cases} \beta(r) r^{-1/2-\delta/2} \varepsilon^{\frac{\delta-8\alpha}{4(2+\alpha)}}, & r \in (0, 1], \\ \beta(r) r^{7/2-\delta/2} \varepsilon^{\frac{2-\alpha}{2+\alpha}}, & r > 1. \end{cases}$$

Since $\alpha < \delta/8$, thanks to Lemma 2.1 we obtain

$$\|J_1^\varepsilon(t)\|_{L^2} \lesssim_{F, F_*, \delta} \varepsilon^{\frac{\delta-8\alpha}{4(2+\alpha)}},$$

Now, J_4 is very similar. On its support, we use (D.31), (D.32), the fact that $u'(r_c) \sim r_c^{-3}$ and (D.8), to obtain for $r \leq 1$ that

$$\begin{aligned} & \left| \int_0^{r_c \geq \frac{1}{2} \varepsilon^{-\frac{1}{2+\alpha}}} e^{ik(u(r)-c)t} \frac{Y(r, c + i\varepsilon)}{u(r) - c - i\varepsilon} (1 - \chi_\sigma(c)) dc \right| \\ & \lesssim \int_{\frac{1}{2} \varepsilon^{-\frac{1}{2+\alpha}}}^\infty |u'(r_c)| \frac{|Y(r, c + i\varepsilon)|}{\sqrt{(u(r) - u(r_c))^2 + \varepsilon^2}} dr_c \\ & \lesssim \langle r \rangle^{2+\alpha} \int_{\frac{1}{2} \varepsilon^{-\frac{1}{2+\alpha}}}^\infty \frac{|Y(r, c + i\varepsilon)|}{r_c^3} dr_c \\ & \lesssim r^{k+1/2-\delta/2} \langle r \rangle^{2+\alpha} \int_{\frac{1}{2} \varepsilon^{-\frac{1}{2+\alpha}}}^\infty \frac{1}{r_c^3} \|r^{-k-1/2+\delta/2} Y(\cdot, c + i\varepsilon)\|_{L^\infty(0,1)} dr_c \\ & \lesssim_{F, F_*, \delta} r^{k+1/2-\delta/2} \langle r \rangle^{2+\alpha} \varepsilon^{\frac{2}{2+\alpha}}, \end{aligned}$$

while for $r > 1$, a similar argument implies

$$\left| \int_0^{r_c \geq \frac{1}{2} \varepsilon^{-\frac{1}{2+\alpha}}} e^{ik(u(r)-c)t} \frac{Y(r, c + i\varepsilon)}{u(r) - c - i\varepsilon} (1 - \chi_\sigma(c)) dc \right| \lesssim_{F, F_*, \delta} r^{-k+1/2+\delta/2} \langle r \rangle^{2+\alpha} \varepsilon^{\frac{2}{2+\alpha}}.$$

Similar estimates hold for $Y(r, c - i\varepsilon)$, and thus in view of Lemma 2.1 we have

$$\|J_4^\varepsilon(t)\|_{L^2} \lesssim_{F, F_*, \delta} \varepsilon^{\frac{2}{2+\alpha}}.$$

We now deal with J_2 and J_3 . For J_2 , in analogy with (D.39) and (D.40) and using (D.5), for $c \in (u(0), u(0) + 1)$ we obtain that

$$\begin{aligned} & \left\| \frac{X(\cdot, c, \varepsilon)}{\min\{r^{k+1/2-\gamma}, r^{-k+1/2+\gamma}\}} \right\|_{L^\infty}^2 + \|X(\cdot, c, \varepsilon)\|_{L_{Y,\gamma}^2}^2 + \|r \partial_r X(\cdot, c, \varepsilon)\|_{L_{Y,\gamma}^2}^2 \\ & \lesssim_\gamma \varepsilon^2 \left\| \frac{\langle r \rangle^2 F}{\sqrt{(u(r) - c)^2 + \varepsilon^2}} \right\|_{L_{Y,\gamma}^2}^2 + \varepsilon^2 \left\| \frac{\langle r \rangle^2 \beta Y(\cdot, c - i\varepsilon)}{\sqrt{(u(r) - c)^2 + \varepsilon^2}} \right\|_{L_{Y,\gamma}^2}^2 \end{aligned}$$

$$\begin{aligned}
&\lesssim_{\gamma} \frac{\varepsilon^{\delta/4}}{|u(0) - c|^{\delta/4}} \left[\|\langle r \rangle^2 F\|_{L^2_{Y,\gamma}}^2 + \|Y(\cdot, c - i\varepsilon)\|_{L^2_{Y,\gamma}}^2 \right] \\
&\lesssim_{\gamma} \frac{\varepsilon^{\delta/4}}{|u(0) - c|^{\delta/4}} \left[\|\langle r \rangle^2 F\|_{L^2_{Y,\gamma}}^2 + \|r^2 F_*\|_{L^2_{Y,\gamma}}^2 \right], \tag{D.47}
\end{aligned}$$

and, similarly

$$\left\| \frac{A(\cdot, c, \varepsilon)}{\min\{r^{k+1/2-\gamma}, r^{-k+1/2+\gamma}\}} \right\|_{L^\infty}^2 \lesssim_{\gamma} \|\langle r \rangle^2 F\|_{L^2_{Y,\gamma}}^2 + \|r^2 F_*\|_{L^2_{Y,\gamma}}^2. \tag{D.48}$$

If $r \leq 1$, taking into account that $u(r) - u(0) \sim r^2$ when $r \rightarrow 0$, we appeal to (D.47) and (D.48) to obtain

$$\begin{aligned}
J_2^\varepsilon(t, r) &\lesssim \frac{\beta(r)}{r^{k+1-\delta}} \int_{u(0)}^{u(0)+1} \frac{|X(r, c, \varepsilon)|}{|u(r) - u(0)|^{\delta/4} |u(0) - c|^{1-\delta/4}} dc \\
&\quad + \varepsilon^{\delta/8} \frac{\beta(r)}{r^{k+1-\delta}} \int_{u(0)}^{u(0)+1} \frac{|A(r, c, \varepsilon)|}{|u(r) - u(0)|^{\delta/4} |u(0) - c|^{1-\delta/8}} dc \\
&\lesssim \frac{\beta(r)}{r^{1/2-\delta/4}} \int_{u(0)}^{u(0)+1} \frac{\|r^{-k-1/2+\delta/4} X(r, c, \varepsilon)\|_{L^\infty(0,1)}}{|u(0) - c|^{1-\delta/4}} dc \\
&\quad + \varepsilon^{\delta/8} \frac{\beta(r)}{r^{1/2-\delta/4}} \int_{u(0)}^{u(0)+1} \frac{\|r^{-k-1/2+\delta/4} A(r, c, \varepsilon)\|_{L^\infty(0,1)}}{|u(0) - c|^{1-\delta/8}} dc \\
&\lesssim_{\delta, F, F_*} \frac{\beta(r)}{r^{1/2-\delta/4}} \varepsilon^{\delta/8}. \tag{D.49}
\end{aligned}$$

If $r \geq 1$, we use an even simpler version of (D.47) to get

$$\begin{aligned}
J_2^\varepsilon(t, r) &\lesssim \frac{\beta(r)}{r^{-k-3+\delta}} \int_{\mathbb{R}} |X(r, c, \varepsilon)| dc + \varepsilon \frac{\beta(r)}{r^{-k-3+\delta}} \int_{\mathbb{R}} |A(r, c, \varepsilon)| dc \\
&\lesssim_{\delta, F, F_*} \frac{\beta(r)}{r^{-7/2+\delta/2}} \varepsilon.
\end{aligned}$$

As a consequence,

$$\|J_2^\varepsilon(t)\|_{L^2} \lesssim_{F, F_*, \delta} \varepsilon^{\delta/8}.$$

The treatment of J_3 is similar. In this case, using (D.8) with $\alpha = 0$, we obtain that

$$\begin{aligned}
&\left\| \frac{X(\cdot, c, \varepsilon)}{\min\{r^{k+1/2-\gamma}, r^{-k+1/2+\gamma}\}} \right\|_{L^\infty}^2 + \|X(\cdot, c, \varepsilon)\|_{L^2_{Y,\gamma}}^2 + \|r \partial_r X(\cdot, c, \varepsilon)\|_{L^2_{Y,\gamma}}^2 \\
&\lesssim_{\gamma} \varepsilon^2 \left\| \frac{r^2 \langle r \rangle^2 F}{\sqrt{(u(r) - c)^2 + \varepsilon^2}} \right\|_{L^2_{Y,\gamma-\tilde{\alpha}}}^2 + \varepsilon^2 \left\| \frac{r^2 \langle r \rangle^2 \beta Y(\cdot, c - i\varepsilon)}{\sqrt{(u(r) - c)^2 + \varepsilon^2}} \right\|_{L^2_{Y,\gamma-\tilde{\alpha}}}^2. \tag{D.50}
\end{aligned}$$

From $u(r) \sim r^{-2}$ when $r \rightarrow \infty$ and the fact that $c \leq 0$, we have

$$\begin{aligned} \varepsilon^2 \left\| \frac{r^2 \langle r \rangle^2 F}{\sqrt{(u(r) - c)^2 + \varepsilon^2}} \right\|_{L_{Y, \gamma - \tilde{\alpha}}^2}^2 &\lesssim \varepsilon^2 \int_0^1 \frac{r^4 |F(r)|^2}{r^{2k+2-2\gamma+2\tilde{\alpha}} ((u(r) - c)^2 + \varepsilon^2)} dr \\ &\quad + \varepsilon^2 \int_1^\infty \frac{r^8 |F(r)|^2}{r^{-2k+2+2\gamma-2\tilde{\alpha}} ((u(r) - c)^2 + \varepsilon^2)} dr \\ &\lesssim_{\gamma} \varepsilon^{\delta/8} \int_0^1 \frac{r^4 |F(r)|^2}{r^{2k+2-2\gamma+2\tilde{\alpha}}} dr \\ &\quad + \varepsilon^{\delta/8} \int_1^\infty \frac{r^{8+\delta/4} |F(r)|^2}{r^{-2k+2+2\gamma-2\tilde{\alpha}}} dr \\ &\lesssim_{\gamma} \varepsilon^{\delta/8} \left\| r^2 \langle r \rangle^{2+\delta/8} F(r) \right\|_{L_{Y, \gamma - \tilde{\alpha}}^2}^2. \end{aligned}$$

Analogously, using Lemma 2.1, we have

$$\begin{aligned} \varepsilon^2 \left\| \frac{r^2 \langle r \rangle^2 \beta Y(\cdot, c - i\varepsilon)}{\sqrt{(u(r) - c)^2 + \varepsilon^2}} \right\|_{L_{Y, \gamma - \tilde{\alpha}}^2}^2 &\lesssim_{\gamma} \varepsilon^{\delta/8} \int_0^1 \frac{r^{4-2\tilde{\alpha}} |Y(r, c - i\varepsilon)|^2}{r^{2k+2-2\gamma}} dr \\ &\quad + \varepsilon^{\delta/8} \int_1^\infty \frac{\beta(r)^2 r^{8+\delta/4+2\tilde{\alpha}} |Y(r, c - i\varepsilon)|^2}{r^{-2k+2+2\gamma}} dr \\ &\lesssim_{\gamma} \varepsilon^{\delta/8} \|Y(\cdot, c - i\varepsilon)\|_{L_{Y, \gamma}^2}^2 \\ &\lesssim_{\gamma} \varepsilon^{\delta/8} \left[\left\| r^2 \langle r \rangle^2 F(r) \right\|_{L_{Y, \gamma - \tilde{\alpha}}^2}^2 + \|r^2 F_*(r)\|_{L_{Y, \gamma - \tilde{\alpha}}^2}^2 \right], \end{aligned}$$

implying that, with the choice $\tilde{\alpha} = \gamma/2 = \delta/4$, we have the control

$$\begin{aligned} &\left\| \frac{X(\cdot, c, \varepsilon)}{\min\{r^{k+1/2-\delta/2}, r^{-k+1/2+\delta/2}\}} \right\|_{L^\infty}^2 \\ &\lesssim_{\gamma} \varepsilon^{\delta/8} \left[\left\| r^2 \langle r \rangle^{2+\delta/8} F(r) \right\|_{L_{Y, \delta/4}^2}^2 + \|r^2 F_*(r)\|_{L_{Y, \delta/4}^2}^2 \right] \end{aligned} \quad (\text{D.51})$$

Regarding A , we argue in the same way and arrive at

$$\left\| \frac{A(\cdot, c, \varepsilon)}{\min\{r^{k+1/2-\delta/2}, r^{-k+1/2+\delta/2}\}} \right\|_{L^\infty}^2 \lesssim_{\gamma} \left\| r^2 \langle r \rangle^2 F(r) \right\|_{L_{Y, \delta/4}^2}^2 + \|r^2 F_*(r)\|_{L_{Y, \delta/4}^2}^2. \quad (\text{D.52})$$

In order to control J_3 , we again consider two cases. If $r \leq 1$,

$$\begin{aligned} J_3^\varepsilon(t, r) &\lesssim \frac{\beta(r)}{r^{k+1-\delta}} \int_{-R_\delta}^0 |X(r, c, \varepsilon)| dc + \varepsilon \frac{\beta(r)}{r^{k+1-\delta}} \int_{-R_\delta}^0 |A(r, c, \varepsilon)| dc \\ &\lesssim \frac{\beta(r)}{r^{1/2-\delta/2}} \int_{-R_\delta}^0 \|r^{-k-1/2+\delta/2} X(r, c, \varepsilon)\|_{L^\infty} dc \\ &\quad + \varepsilon \frac{\beta(r)}{r^{1/2-\delta/2}} \int_{-R_\delta}^0 \|r^{-k-1/2+\delta/2} A(r, c, \varepsilon)\|_{L^\infty} dc \\ &\lesssim_{\delta, F, F_*} \frac{\beta(r)}{r^{1/2-\delta/2}} \varepsilon^{\delta/16}, \end{aligned}$$

while if $r \geq 1$, since $u(r) \sim r^{-2}$ as $r \rightarrow \infty$, we use (D.51) and (D.52) to deduce that

$$\begin{aligned} J_3^\varepsilon(t, r) &\lesssim \frac{\beta(r)}{r^{-k+1+\delta-4}} \varepsilon^{-\delta/32} \int_{-R_\delta}^0 \frac{|X(r, c, \varepsilon)|}{|c|^{1-\delta/32}} dc \\ &\quad + \varepsilon^{\delta/16} \frac{\beta(r)}{r^{-k+1+\delta-4}} \int_{-R_\delta}^0 \frac{|A(r, c, \varepsilon)|}{|u(r)|^{\delta/8} |c|^{1-\delta/16}} dc \\ &\lesssim \frac{\beta(r) \varepsilon^{-\delta/32}}{r^{-7/2+\delta/2}} \int_{-R_\delta}^0 \frac{\|r^{k-1/2-\delta/2} X(r, c, \varepsilon)\|_{L^\infty}}{|c|^{1-\delta/32}} dc \\ &\quad + \frac{\beta(r) \varepsilon^{\delta/8}}{r^{-7/2+\delta/4}} \int_{-R_\delta}^0 \frac{\|r^{k-1/2-\delta/2} A(r, c, \varepsilon)\|_{L^\infty}}{|c|^{1-\delta/16}} dc \\ &\lesssim_{\delta, F, F_*} \frac{\beta(r)}{r^{-7/2+\delta/4}} \varepsilon^{\delta/32}. \end{aligned}$$

Hence,

$$\|J_3^\varepsilon(t)\|_{L^2} \lesssim_{F, F_*, \delta} \varepsilon^{\delta/32},$$

which concludes the proof. \square

D.5 Vanishing of Higher Derivatives

The aim of this section is to prove analogous results to those of Propositions D.5–D.6.

Proposition D.7 *Let δ be fixed sufficiently small, and let $j \in \{1, \dots, k\}$. Then*

$$\lim_{\varepsilon \rightarrow 0} \left[\|(r \partial_r)^j f_E^\varepsilon(t, \cdot)\|_{L_{f, \delta}^2} + \lim_{\varepsilon \rightarrow 0} \|(r \partial_r)^j f_S^\varepsilon(t, \cdot)\|_{L_{f, \delta}^2} \right] = 0,$$

for every $t \geq 0$.

The proof is based on the iteration scheme laid out in Lemma 2.8, and appropriately choosing $\gamma \in (0, 2k)$ in Theorem D.1, depending on the derivative index j and the

small parameter δ . We illustrate the proof only in one case to handle higher derivatives of f_S , for $k = 2$ and $j = 1, 2$. As in (2.28), we have

$$\begin{aligned} (r\partial_r)^j f_S^\varepsilon(t, r) = & \frac{1}{2\pi i} \int_{-R_\delta}^{u(0)+1} e^{ik(u(r)-c)t} \left[\frac{1}{u(r)-c-i\varepsilon} (ru'(r)\partial_G)^j \right. \\ & \left(\frac{\beta(r)}{\sqrt{r}} Y(r, c+i\varepsilon) \chi_\sigma(c)(1-\chi_I(r_c)) \right) \\ & - \frac{1}{u(r)-c+i\varepsilon} (ru'(r)\partial_G)^j \\ & \left. \left(\frac{\beta(r)}{\sqrt{r}} Y(r, c-i\varepsilon) \chi_\sigma(c)(1-\chi_I(r_c)) \right) \right] dc. \end{aligned} \quad (\text{D.53})$$

D.5.1 Proof for $j \in \{1, \dots, k-1\}$

Arguing as in the proof of Proposition D.6 and taking $j = 1$, an important term to bound (somewhat analogous to (D.44)) reads as

$$\begin{aligned} J_2^\varepsilon(t, r) = & \frac{\beta(r)ru'(r)}{w_{f,\delta}(r)r^{1/2}} \\ & \left| \int_{u(0)}^{u(0)+1} e^{ik(u(r)-c)t} \left[\frac{u(r)-c}{(u(r)-c)^2+\varepsilon^2} \partial_G X(r, c, \varepsilon) + \frac{i\varepsilon \partial_G A(r, c, \varepsilon)}{(u(r)-c)^2+\varepsilon^2} \right] dc \right|. \end{aligned} \quad (\text{D.54})$$

According to Lemma 2.8, we have

$$\text{RAY}_+ \partial_G X = \frac{2i\varepsilon}{(u(r)-c)^2+\varepsilon^2} \left(\frac{1}{u'(r)} \partial_r F(r) - \beta(r) \partial_G Y(r, c-i\varepsilon) - \frac{\beta'(r)}{u'(r)} Y(r, c-i\varepsilon) \right. \quad (\text{D.55})$$

$$\begin{aligned} & \left. - 2 \frac{u''(r)}{(u'(r))^2} [F(r) \right. \\ & \left. - \beta(r) Y(r, c-i\varepsilon)] \right) \end{aligned} \quad (\text{D.56})$$

$$+ \frac{1}{u-c-i\varepsilon} \left(2 \frac{u''(r)\beta(r)}{(u'(r))^2} - \frac{\beta'(r)}{u'(r)} \right) X \quad (\text{D.57})$$

$$+ \partial_{rr} \left(\frac{1}{u'(r)} \right) \partial_r X + 2 \frac{1/4-k^2}{r^3(u'(r))^2} (u'(r) + ru''(r)) X \quad (\text{D.58})$$

and

$$\begin{aligned} \text{RAY}_- \partial_G Y = & \frac{1}{u-c+i\varepsilon} \left(\left(2 \frac{u''(r)\beta(r)}{(u'(r))^2} - \frac{\beta'(r)}{u'(r)} \right) Y + \frac{1}{u'(r)} \partial_r F(r) \right. \\ & \left. - 2 \frac{u''(r)}{(u'(r))^2} F(r) \right) \end{aligned} \quad (\text{D.59})$$

$$\begin{aligned}
& + \partial_{rr} \left(\frac{1}{u'(r)} \right) \partial_r Y + 2 \frac{1/4 - k^2}{r^3 (u'(r))^2} (u'(r) + r u''(r)) Y \\
& - 2 \frac{u''(r)}{(u'(r))^2} F_*(r) + \partial_G F_*(r).
\end{aligned} \tag{D.60}$$

Next, we show that the terms in (D.58) and (D.60) are bounded in the norm $\|r^2 \cdot\|_{L^2_{Y,\gamma}}$ as they correspond to the term $F_{*,\varepsilon}$ in Theorem D.1. The terms containing F_* are harmless, since they are compactly supported away from $r = 0$. As for the other terms, notice that from Lemma 6.2, we have

$$\frac{(u'(r) + r u''(r))}{r^3 (u'(r))^2} \sim \begin{cases} r^{-4}, & r \rightarrow 0, \\ 1, & r \rightarrow \infty, \end{cases} \quad \partial_{rr} \left(\frac{1}{u'(r)} \right) \sim \begin{cases} r^{-3}, & r \rightarrow 0, \\ r, & r \rightarrow \infty. \end{cases}$$

Thus, we can choose $\gamma = 2 + \gamma' \in (0, 2k)$, deduce that

$$\begin{aligned}
& \left\| r^2 \left[\partial_{rr} \left(\frac{1}{u'(r)} \right) \partial_r Y + 2 \frac{1/4 - k^2}{r^3 (u'(r))^2} (u'(r) + r u''(r)) Y \right] \right\|_{L^2_{Y,2+\gamma'}}^2 \\
& \lesssim_\delta \|r \partial_r Y\|_{L^2_{Y,\gamma'}}^2 + \|Y\|_{L^2_{Y,\gamma'}}^2
\end{aligned} \tag{D.61}$$

and

$$\begin{aligned}
& \left\| r^2 \left[\partial_{rr} \left(\frac{1}{u'(r)} \right) \partial_r X + 2 \frac{1/4 - k^2}{r^3 (u'(r))^2} (u'(r) + r u''(r)) X \right] \right\|_{L^2_{Y,2+\gamma'}}^2 \\
& \lesssim_\delta \|r \partial_r X\|_{L^2_{Y,\gamma'}}^2 + \|X\|_{L^2_{Y,\gamma'}}^2.
\end{aligned} \tag{D.62}$$

To control the terms in (D.57) and (D.59), we note Lemma 6.2,

$$\begin{aligned}
& 2 \frac{u''(r) \beta(r)}{(u'(r))^2} - \frac{\beta'(r)}{u'(r)} \sim \begin{cases} r^{-2}, & r \rightarrow 0, \\ r^{-4}, & r \rightarrow \infty, \end{cases} \quad \frac{u''(r)}{(u'(r))^2} \sim \begin{cases} r^{-2}, & r \rightarrow 0, \\ r^2, & r \rightarrow \infty, \end{cases} \\
& \frac{1}{u'(r)} \sim \begin{cases} r^{-1}, & r \rightarrow 0, \\ r^3, & r \rightarrow \infty. \end{cases}
\end{aligned}$$

The point is that these coefficient are less singular at the origin by a power of r^{-2} , and therefore all the norms appearing in Theorem D.1 for F_ε are finite even without a gain of r^2 at the origin, upon choosing again $\gamma = 2 + \gamma'$. In the specific case when $c \in (u(0), u(0) + 1)$, we use (D.5) to have

$$\begin{aligned}
& \left\| \langle r \rangle^2 \left(\left(2 \frac{u''(r) \beta(r)}{(u'(r))^2} - \frac{\beta'(r)}{u'(r)} \right) Y + \frac{1}{u'(r)} \partial_r F(r) - 2 \frac{u''(r)}{(u'(r))^2} F(r) \right) \right\|_{L^2_{Y,2+\gamma'}}^2 \\
& \lesssim_\delta \|r \partial_r Y\|_{L^2_{Y,\gamma'}}^2 + \|Y\|_{L^2_{Y,\gamma'}}^2 + \|\langle r \rangle^2 r \partial_r F\|_{L^2_{Y,\gamma'}}^2 + \|\langle r \rangle^2 F\|_{L^2_{Y,\gamma'}}^2,
\end{aligned} \tag{D.63}$$

and

$$\left\| \langle r \rangle^2 \left(2 \frac{u''(r)\beta(r)}{(u'(r))^2} - \frac{\beta'(r)}{u'(r)} \right) X \right\|_{L^2_{Y,2+\gamma'}}^2 \lesssim_\delta \|r \partial_r X\|_{L^2_{Y,\gamma'}}^2 + \|X\|_{L^2_{Y,\gamma'}}^2. \quad (\text{D.64})$$

Lastly, the terms in (D.56) are treated as in (D.47). Accordingly, using (D.62) we obtain the bounds

$$\begin{aligned} & \left\| \frac{\partial_G Y(\cdot, c - i\varepsilon)}{\min\{r^{k+1/2-2-\gamma'}, r^{-k+1/2+2+\gamma'}\}} \right\|_{L^\infty}^2 + \|\partial_G Y(\cdot, c - i\varepsilon)\|_{L^2_{Y,2+\gamma'}}^2 \\ & \quad + \|r \partial_r \partial_G Y(\cdot, c - i\varepsilon)\|_{L^2_{Y,2+\gamma'}}^2 \\ & \lesssim \|r \partial_r Y\|_{L^2_{Y,\gamma'}}^2 + \|Y\|_{L^2_{Y,\gamma'}}^2 + \|\langle r \rangle^2 r \partial_r F\|_{L^2_{Y,\gamma'}}^2 + \|\langle r \rangle^2 F\|_{L^2_{Y,\gamma'}}^2, \end{aligned}$$

and

$$\begin{aligned} & \left\| \frac{\partial_G X(\cdot, c, \varepsilon)}{\min\{r^{k+1/2-2-\gamma'}, r^{-k+1/2+2+\gamma'}\}} \right\|_{L^\infty}^2 + \|\partial_G X(\cdot, c, \varepsilon)\|_{L^2_{Y,2+\gamma'}}^2 \\ & \quad + \|r \partial_r \partial_G X(\cdot, c, \varepsilon)\|_{L^2_{Y,2+\gamma'}}^2 \\ & \lesssim \frac{\varepsilon^{\delta/4}}{|u(0) - c|^{\delta/4}} \left[\|\langle r \rangle^2 F\|_{Y,\gamma'}^2 + \|\langle r \rangle^2 r \partial_r F\|_{Y,\gamma'}^2 + \|Y\|_{Y,\gamma'}^2 + \|\partial_G Y\|_{Y,\gamma'}^2 \right] \\ & \quad + \|r \partial_r X\|_{L^2_{Y,\gamma'}}^2 + \|X\|_{L^2_{Y,\gamma'}}^2, \end{aligned} \quad (\text{D.65})$$

which, using (D.47) and (D.48), they imply that

$$\begin{aligned} & \left\| \frac{\partial_G Y(\cdot, c - i\varepsilon)}{\min\{r^{k+1/2-2-\gamma'}, r^{-k+1/2+2+\gamma'}\}} \right\|_{L^\infty}^2 + \|\partial_G Y(\cdot, c - i\varepsilon)\|_{L^2_{Y,2+\gamma'}}^2 \\ & \quad + \|r \partial_r \partial_G Y(\cdot, c - i\varepsilon)\|_{L^2_{Y,2+\gamma'}}^2 \\ & \lesssim \|\langle r \rangle^2 r \partial_r F\|_{L^2_{Y,\gamma'}}^2 + \|\langle r \rangle^2 F\|_{L^2_{Y,\gamma'}}^2, \end{aligned}$$

and

$$\begin{aligned} & \left\| \frac{\partial_G X(\cdot, c, \varepsilon)}{\min\{r^{k+1/2-2-\gamma'}, r^{-k+1/2+2+\gamma'}\}} \right\|_{L^\infty}^2 + \|\partial_G X(\cdot, c, \varepsilon)\|_{L^2_{Y,2+\gamma'}}^2 \\ & \quad + \|r \partial_r \partial_G X(\cdot, c, \varepsilon)\|_{L^2_{Y,2+\gamma'}}^2 \\ & \lesssim \frac{\varepsilon^{\delta/4}}{|u(0) - c|^{\delta/4}} \left[\|\langle r \rangle^2 F\|_{Y,\gamma'}^2 + \|\langle r \rangle^2 r \partial_r F\|_{Y,\gamma'}^2 \right]. \end{aligned} \quad (\text{D.66})$$

Similarly, by essentially arguing that $A(\cdot, c, \varepsilon) = X(\cdot, c, \varepsilon) + 2Y(\cdot, c - i\varepsilon)$, we obtain

$$\begin{aligned} & \left\| \frac{\partial_G A(\cdot, c, \varepsilon)}{\min\{r^{k+1/2-2-\gamma'}, r^{-k+1/2+2+\gamma'}\}} \right\|_{L^\infty}^2 + \|\partial_G A(\cdot, c, \varepsilon)\|_{L_{Y,2+\gamma'}^2}^2 \\ & \quad + \|r\partial_r \partial_G A(\cdot, c, \varepsilon)\|_{L_{Y,2+\gamma'}^2}^2 \\ & \lesssim \|\langle r \rangle^2 r \partial_r F\|_{L_{Y,\gamma'}^2}^2 + \|\langle r \rangle^2 F\|_{L_{Y,\gamma'}^2}^2. \end{aligned}$$

Now, going back to (D.54) and noticing that $ru'(r) \sim r^2$ as $r \rightarrow 0$ and $ru'(r) \sim r^{-2}$ as $r \rightarrow \infty$, and collecting all the estimates above, we complete the proof of Proposition D.7 in the case $j = 1$ by choosing $\gamma' = \delta/4$.

It is worth mentioning that the proof for $k > j > 1$ is analogous: the main idea is to use Lemma 6.2, combined with the choice $\gamma = 2j + \delta < 2k$. Clearly this imposes the constraint $j \leq k - 1$, which is why the case $j = k$ is treated differently below.

D.5.2 Proof for $j = k$

We now deal with the case $j = k$. For the sake of simplicity, we consider the case $k = 2$; the others are analogous. We begin from (D.53) with $j = 1$ and take an additional $r\partial_r$ derivative, but in this case we do not exploit the ∂_G derivative. Let us only deal with the term containing X , namely

$$\begin{aligned} r\partial_r f_{S,X}^\varepsilon(t, r) &= \frac{1}{2\pi i} \int_{\mathbb{R}} e^{ik(u(r)-c)t} \frac{u(r) - c}{(u(r) - c)^2 + \varepsilon^2} r u'(r) \\ &\quad \times \partial_G \left(\frac{\beta(r)}{\sqrt{r}} X(r, c, \varepsilon) \right) (1 - \chi_\sigma(c)) dc. \end{aligned}$$

Then

$$\begin{aligned} (r\partial_r)^2 f_{S,X}^\varepsilon(t, r) &= \frac{kt}{2\pi} \int_{\mathbb{R}} e^{ik(u(r)-c)t} \frac{(u(r) - c)^2}{(u(r) - c)^2 + \varepsilon^2} r^2 u'(r) \\ &\quad \times \partial_G \left(\frac{\beta(r)}{\sqrt{r}} X(r, c, \varepsilon) \right) \chi_\sigma(c) (1 - \chi_I(r_c)) dc \\ &\quad + \frac{1}{2\pi i} \int_{\mathbb{R}} e^{ik(u(r)-c)t} \frac{\varepsilon^2 - (u(r) - c)^2}{((u(r) - c)^2 + \varepsilon^2)^2} r^2 (u'(r))^2 \\ &\quad \times \partial_G \left(\frac{\beta(r)}{\sqrt{r}} X(r, c, \varepsilon) \right) \chi_\sigma(c) (1 - \chi_I(r_c)) dc \\ &\quad + \frac{1}{2\pi i} \int_{\mathbb{R}} e^{ik(u(r)-c)t} \frac{u(r) - c}{(u(r) - c)^2 + \varepsilon^2} r (u'(r) + ru''(r)) \\ &\quad \times \partial_G \left(\frac{\beta(r)}{\sqrt{r}} X(r, c, \varepsilon) \right) \chi_\sigma(c) (1 - \chi_I(r_c)) dc \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{2\pi i} \int_{\mathbb{R}} e^{ik(u(r)-c)t} \frac{u(r)-c}{(u(r)-c)^2 + \varepsilon^2} r^2 u'(r) \partial_r \partial_G \\
& \times \left(\frac{\beta(r)}{\sqrt{r}} X(r, c, \varepsilon) \right) \chi_\sigma(c) (1 - \chi_I(r_c)) dc.
\end{aligned}$$

We show how to deal with the four terms above, when all the derivatives land on X , in the case when $c \in (u(0), u(0) + 1)$ and $r \leq 1$, when using the appropriate weight (1.10b). For the first term, we bound it using (D.66) and

$$\begin{aligned}
& \frac{kt}{r^{k+1/2-\delta}} \int_{u(0)}^{u(0)+1} r^2 u'(r) \frac{1}{\sqrt{r}} |\partial_G X(r, c, \varepsilon)| dc \\
& \lesssim \frac{kt}{r^{1/2-\delta/2}} \int_{u(0)}^{u(0)+1} r^3 \frac{|\partial_G X(r, c, \varepsilon)|}{r^{k+1/2-\delta/2}} dc \\
& \lesssim \frac{kt}{r^{1/2-\delta/2}} \int_{u(0)}^{u(0)+1} \frac{|\partial_G X(r, c, \varepsilon)|}{r^{k+1/2-2-\delta/2}} dc \\
& \lesssim kt \varepsilon^{\delta/4} \left[\|\langle r \rangle^2 F\|_{Y, \delta/2}^2 + \|\langle r \rangle^2 r \partial_r F\|_{Y, \delta/2}^2 \right],
\end{aligned}$$

The second term is similar, only slightly more delicate. It suffices to bound the following, using (D.66) once more:

$$\begin{aligned}
& \frac{1}{r^{k+1/2-\delta}} \int_{u(0)}^{u(0)+1} \frac{1}{(u(r)-c)^2 + \varepsilon^2} r^2 (u'(r))^2 \frac{1}{\sqrt{r}} |\partial_G X(r, c, \varepsilon)| dc \\
& \lesssim \frac{1}{r^{k+1/2-\delta}} \int_{u(0)}^{u(0)+1} \frac{r^4}{|u(r)-u(0)|^{1+2\delta/7}} \frac{1}{|u(0)-c|^{1-2\delta/7}} \frac{1}{\sqrt{r}} |\partial_G X(r, c, \varepsilon)| dc \\
& \lesssim \frac{1}{r^{k+1/2-\delta}} \int_{u(0)}^{u(0)+1} \frac{r^{2-4\delta/7}}{|u(0)-c|^{1-2\delta/7}} \frac{1}{\sqrt{r}} |\partial_G X(r, c, \varepsilon)| dc \\
& \lesssim \frac{1}{r^{1/2-\delta/7}} \int_{u(0)}^{u(0)+1} \frac{1}{|u(0)-c|^{1-2\delta/7}} \frac{|\partial_G X(r, c, \varepsilon)|}{r^{k+1/2-2-\delta/7}} dc \\
& \lesssim \varepsilon^{\delta/4} \left[\|\langle r \rangle^2 F\|_{Y, 2\delta/7}^2 + \|\langle r \rangle^2 r \partial_r F\|_{Y, 2\delta/7}^2 \right].
\end{aligned}$$

The next term is treated similarly, using that

$$\begin{aligned}
& \frac{1}{r^{k+1/2-\delta}} \int_{u(0)}^{u(0)+1} \frac{1}{\sqrt{(u(r)-c)^2 + \varepsilon^2}} r(u'(r) + ru''(r)) \frac{1}{\sqrt{r}} |\partial_G X(r, c, \varepsilon)| dc \\
& \lesssim \frac{1}{r^{k+1/2-\delta}} \int_{u(0)}^{u(0)+1} \frac{r^2}{|u(r)-u(0)|^{2\delta/7}} \frac{1}{|u(r)-c|^{1-2\delta/7}} \frac{1}{\sqrt{r}} |\partial_G X(r, c, \varepsilon)| dc \\
& \lesssim \frac{1}{r^{1/2-\delta/7}} \int_{u(0)}^{u(0)+1} \frac{1}{|u(r)-c|^{1-2\delta/7}} \frac{|\partial_G X(r, c, \varepsilon)|}{r^{k+1/2-2-\delta/7}} dc \\
& \lesssim \varepsilon^{\delta/4} \left[\|\langle r \rangle^2 F\|_{Y, 2\delta/7}^2 + \|\langle r \rangle^2 r \partial_r F\|_{Y, 2\delta/7}^2 \right].
\end{aligned}$$

Lastly, we have

$$\begin{aligned}
 & \frac{1}{r^{k+1/2-\delta}} \int_{u(0)}^{u(0)+1} \frac{1}{\sqrt{(u(r)-c)^2 + \varepsilon^2}} r^2 u'(r) \frac{1}{\sqrt{r}} |\partial_r \partial_G X(r, c, \varepsilon)| dc \\
 & \lesssim \frac{1}{r^{k+1/2-\delta}} \int_{u(0)}^{u(0)+1} \frac{1}{|u(r)-u(0)|^{2\delta/7}} \frac{1}{|u(0)-c|^{1-2\delta/7}} r^2 u'(r) \frac{1}{\sqrt{r}} |\partial_r \partial_G X(r, c, \varepsilon)| dc \\
 & \lesssim \frac{1}{r^{1/2-\delta/7}} \int_{u(0)}^{u(0)+1} \frac{1}{|u(0)-c|^{1-2\delta/7}} \frac{|r \partial_r \partial_G X(r, c, \varepsilon)|}{r^{k+1/2-2-\delta/7}} dc. \tag{D.67}
 \end{aligned}$$

Hence, to bound this last term we need a proper estimate L^∞ on $r \partial_r \partial_G X$. By Sobolev embeddings and the fact that $(r \partial_r)^2 = r^2 \partial_{rr} + r \partial_r$, this follows from a proper L^2 bound on $r^2 \partial_{rr}$. Referring to (D.56)–(D.58), we have that

$$r^2 \partial_{rr} \partial_G X = - \left(\frac{1}{4} - k^2 \right) \partial_G X - \frac{\beta(r)r^2}{u(r)-c-i\varepsilon} \partial_G X + r^2 [(\text{D.56}) + (\text{D.57}) + (\text{D.58})]. \tag{D.68}$$

Since for $c \in (u(0), u(0) + 1)$ we have $r^2 \lesssim \sqrt{(u(r)-c)^2 + \varepsilon^2}$ if $r \leq 1$, we use Lemma 2.1 to deduce that

$$\begin{aligned}
 \|r^2 \partial_{rr} \partial_G X\|_{L^2_{Y,2+\gamma'}}^2 & \lesssim \|\partial_G X\|_{L^2_{Y,2+\gamma'}}^2 + \left\| \frac{\beta(r)r^2}{\sqrt{(u(r)-c)^2 + \varepsilon^2}} \partial_G X \right\|_{L^2_{Y,2+\gamma'}}^2 \\
 & \quad + \left\| r^2 [(\text{D.56}) + (\text{D.57}) + (\text{D.58})] \right\|_{L^2_{Y,2+\gamma'}}^2 \\
 & \lesssim \|\partial_G X\|_{L^2_{Y,2+\gamma'}}^2 + \left\| r^2 [(\text{D.56}) + (\text{D.57}) + (\text{D.58})] \right\|_{L^2_{Y,2+\gamma'}}^2.
 \end{aligned}$$

Now, in view of (D.62), (D.64) and (D.66), we end up with the higher order estimate

$$\begin{aligned}
 & \left\| \frac{r \partial_r \partial_G X(\cdot, c, \varepsilon)}{\min\{r^{k+1/2-2-\gamma'}, r^{-k+1/2+2+\gamma'}\}} \right\|_{L^\infty}^2 + \|r \partial_r \partial_G X(\cdot, c, \varepsilon)\|_{L^2_{Y,2+\gamma'}}^2 \\
 & \quad + \left\| (r \partial_r)^2 \partial_G X(\cdot, c, \varepsilon) \right\|_{L^2_{Y,2+\gamma'}}^2 \\
 & \lesssim \frac{\varepsilon^{\delta/4}}{|u(0)-c|^{\delta/4}} \left[\|\langle r \rangle^2 F\|_{Y,\gamma'}^2 + \|\langle r \rangle^2 r \partial_r F\|_{Y,\gamma'}^2 \right]. \tag{D.69}
 \end{aligned}$$

Going back to (D.67) we now set $\gamma' = 2\delta/7$ as in the other terms. Arguing in a similar manner for $r \geq 1$, we deduce that

$$\|(r\partial_r)^2 f_{S,X}^\varepsilon(t,r)\|_{L_{f,\delta}^2} \lesssim_{\delta,t} \varepsilon^{\delta/4},$$

which is what we wanted. The treatment of all the other cases is similar, following the ideas of Propositions D.5 and D.6. The proof of Proposition D.7 is therefore concluded.

References

1. Bajer, K., Bassom, A.P., Gilbert, A.D.: Accelerated diffusion in the centre of a vortex. *J. Fluid Mech.* **437**, 395–411 (2001)
2. Balmforth, N.J., Smith, S.G.L., Young, W.R.: Disturbing vortices. *J. Fluid Mech.* **426**, 95–133 (2001)
3. Bassom, A.P., Gilbert, A.D.: The spiral wind-up of vorticity in an inviscid planar vortex. *J. Fluid Mech.* **371**, 109–140 (1998)
4. Beck, M., Wayne, C.E.: Metastability and rapid convergence to quasi-stationary bar states for the two-dimensional Navier–Stokes equations. *Proc. R. Soc. Edinb. Sect. A Math.* **143**(05), 905–927 (2013)
5. Bedrossian, J.: Nonlinear echoes and Landau damping with insufficient regularity (2016). [arXiv:1605.06841](https://arxiv.org/abs/1605.06841)
6. Bedrossian, J., Germain, P., Masmoudi, N.: Dynamics near the subcritical transition of the 3D Couette flow I: below threshold. *Mem. Am. Math. Soc.* (2015) (**To appear**). [arXiv:1506.03720](https://arxiv.org/abs/1506.03720)
7. Bedrossian, J., Germain, P., Masmoudi, N.: Dynamics near the subcritical transition of the 3D Couette flow II: above threshold (2015). [arXiv:1506.03721](https://arxiv.org/abs/1506.03721)
8. Bedrossian, J., Germain, P., Masmoudi, N.: On the stability threshold for the 3D Couette flow in Sobolev regularity. *Ann. Math.* **157**(1), 541–608 (2017)
9. Bedrossian, J., Masmoudi, N., Mouhot, C.: Landau damping in finite regularity for unconfined systems with screened interactions. *Commun. Pure Appl. Math.* **71**(3), 537–576 (2018)
10. Bedrossian, J., Masmoudi, N., Vicol, V.: Enhanced dissipation and inviscid damping in the inviscid limit of the Navier–Stokes equations near the 2D Couette flow. *Arch. Ration. Mech. Anal.* **216**(3), 1087–1159 (2016)
11. Bedrossian, J., Vicol, V., Wang, F.: The Sobolev stability threshold for 2D shear flows near Couette. *J. Nonlinear Sci.* **28**(2), 711–724 (2018)
12. Bedrossian, J., Coti Zelati, M.: Enhanced dissipation, hypoellipticity, and anomalous small noise inviscid limits in shear flows. *Arch. Ration. Mech. Anal.* **224**(3), 1161–1204 (2017)
13. Bedrossian, J., Coti Zelati, M., Glatt-Holtz, N.: Invariant measures for passive scalars in the small noise inviscid limit. *Commun. Math. Phys.* **348**(1), 101–127 (2016)
14. Bedrossian, J., Masmoudi, N.: Inviscid damping and the asymptotic stability of planar shear flows in the 2D Euler equations. *Publ. math. de l’IHÉS* **122**(1), 1–106 (2013)
15. Bedrossian, J., Masmoudi, N., Mouhot, C.: Landau damping: paraproducts and Gevrey regularity. *Ann. PDE* **2**(1), 1–71 (2016)
16. Bouchet, F., Morita, H.: Large time behavior and asymptotic stability of the 2D Euler and linearized Euler equations. *Physica D* **239**, 948–966 (2010)
17. Bracco, A., McWilliams, J.C., Murante, G., Provenzale, A., Weiss, J.B.: Revisiting freely decaying two-dimensional turbulence at millennial resolution. *Phys. Fluids* **12**(11), 2931–2941 (2000)
18. Briggs, R.J., Daugherty, J.D., Levy, R.H.: Role of Landau damping in crossed-field electron beams and inviscid shear flow. *Phys. Fluids* **13**(2), 421–432 (1970)
19. Caglioti, E., Maffei, C.: Time asymptotics for solutions of Vlasov–Poisson equation in a circle. *J. Stat. Phys.* **92**(1/2), 301–323 (1998)
20. Case, K.M.: Stability of inviscid plane Couette flow. *Phys. Fluids* **3**(2), 143–148 (1960)
21. Castro, A., Córdoba, D., Gómez-Serrano, J.: Uniformly rotating smooth solutions for the incompressible 2D Euler equations (2016). [arXiv preprint arXiv:1612.08964](https://arxiv.org/abs/1612.08964)
22. Cerfon, A.J., Freidberg, J.P., Parra, F.I., Antaya, T.A.: Analytic fluid theory of beam spiraling in high-intensity cyclotrons. *Phys. Rev. ST Accel. Beams* **16**, 024202 (2013)
23. Constantin, P., Kiselev, A., Ryzhik, L., Zlatoš, A.: Diffusion and mixing in fluid flow. *Ann. Math.* **2**(168), 643–674 (2008)
24. Coti Zelati, M., Zillinger, C.: On degenerate circular and shear flows: the point vortex and power law circular flows. Preprint

25. Degond, P.: Spectral theory of the linearized Vlasov–Poisson equation. *Trans. Am. Math. Soc.* **294**(2), 435–453 (1986)
26. Deng, W.: Resolvent estimates for a two-dimensional non-self-adjoint operator. *Commun. Pure Appl. Anal.* **12**(1), 547–596 (2013)
27. Dikii, L.A.: The stability of plane-parallel flows of an ideal fluid. *Sov. Phys. Dokl.* **135**(5), 1179 (1961)
28. Drazin, P.G., Reid, W.H.: *Hydrodynamic Stability*. Cambridge Monographs on Mechanics and Applied Mathematics. Cambridge University Press, Cambridge (1981)
29. Dritschel, D.G.: Nonlinear stability bounds for inviscid, two-dimensional, parallel or circular flows with monotonic vorticity, and the analogous three-dimensional quasi-geostrophic flows. *J. Fluid Mech.* **191**, 575–581 (1988)
30. Dubrulle, B., Nazarenko, S.: On scaling laws for the transition to turbulence in uniform-shear flows. *Eur. Phys. Lett.* **27**(2), 129 (1994)
31. Faou, E., Rousset, F.: Landau damping in Sobolev spaces for the Vlasov–HMF model. *Arch. Ration. Mech. Anal.* **219**(2), 887–902 (2016)
32. Fernandez, B., Gérard-Varet, D., Giacomini, G.: Landau damping in the Kuramoto model. *Ann. Institut Poincaré Analysis nonlinéaire (to appear)*. Preprint [arXiv:1410.6006](https://arxiv.org/abs/1410.6006)
33. Gallay, T.: Enhanced dissipation and axisymmetrization of two-dimensional viscous vortices (2017). *arXiv preprint* [arXiv:1707.05525](https://arxiv.org/abs/1707.05525)
34. Gallay, T., Wayne, C.E.: Global stability of vortex solutions of the two-dimensional Navier–Stokes equation. *Commun. Math. Phys.* **255**(1), 97–129 (2005)
35. Glassey, R., Schaeffer, J.: Time decay for solutions to the linearized Vlasov equation. *Transp. Theory Stat. Phys.* **23**(4), 411–453 (1994)
36. Hall, I.M., Bassom, A.P., Gilbert, A.D.: The effect of fine structure on the stability of planar vortices. *Eur. J. Mech. B/Fluids* **22**(2), 179–198 (2003)
37. Hwang, H.J., Velaázquez, J.J.L.: On the existence of exponentially decreasing solutions of the nonlinear Landau damping problem. *Indiana Univ. Math. J.* **58**, 2623–2660 (2009)
38. Ibrahim, S., Maekawa, Y., Masmoudi, N.: On pseudospectral bound for non-selfadjoint operators and its application to stability of Kolmogorov flows (2017). *arXiv preprint* [arXiv:1710.05132](https://arxiv.org/abs/1710.05132)
39. Kelvin, L.: Stability of fluid motion: rectilinear motion of viscous fluid between two parallel plates. *Philos. Mag.* **24**, 188 (1887)
40. Koumoutsakos, P.: Inviscid axisymmetrization of an elliptical vortex. *J. Comput. Phys.* **138**(2), 821–857 (1997)
41. Landau, L.: On the vibration of the electronic plasma. *J. Phys. USSR* **10**, 25 (1946)
42. Latini, M., Bernoff, A.J.: Transient anomalous diffusion in Poiseuille flow. *J. Fluid Mech.* **441**, 399–411 (2001)
43. Li, T., Wei, D., Zhang, Z.: Pseudospectral and spectral bounds for the Oseen vortices operator (2017). *arXiv preprint* [arXiv:1701.06269](https://arxiv.org/abs/1701.06269)
44. Lin, Z., Xu, M.: Metastability of Kolmogorov flows and inviscid damping of shear flows (2017). *arXiv preprint* [arXiv:1707.00278](https://arxiv.org/abs/1707.00278)
45. Lin, Z., Zeng, C.: Inviscid dynamical structures near Couette flow. *Arch. Ration. Mech. Anal.* **200**(3), 1075–1097 (2011)
46. Lin, Z., Zeng, C.: Small BGK waves and nonlinear Landau damping. *Commun. Math. Phys.* **306**(2), 291–331 (2011)
47. Malmberg, J., Wharton, C.: Collisionless damping of electrostatic plasma waves. *Phys. Rev. Lett.* **13**(6), 184–186 (1964)
48. Malmberg, J., Wharton, C., Gould, C., O’Neil, T.: Plasma wave echo. *Phys. Rev. Lett.* **20**(3), 95–97 (1968)
49. Melander, M.V., McWilliams, J.C., Zabusky, N.J.: Axisymmetrization and vorticity-gradient intensification of an isolated two-dimensional vortex through filamentation. *J. Fluid Mech.* **178**, 137–159 (1987)
50. Miller, P.D.: *Applied Asymptotic Analysis*, vol. 75. American Mathematical Society, Providence (2006)
51. Montgomery, M.T., Kallenbach, R.J.: A theory for vortex Rossby-waves and its application to spiral bands and intensity changes in hurricanes. *Q. J. R. Meteorol. Soc.* **123**(538), 435–465 (1997)
52. Mouhot, C., Villani, C.: On Landau damping. *Acta Math.* **207**(1), 29–201 (2011)
53. Nolan, D.S., Montgomery, M.T.: The algebraic growth of wavenumber one disturbances in hurricane-like vortices. *J. Atmos. Sci.* **57**(21), 3514–3538 (2000)

54. Orr, W.: The stability or instability of steady motions of a perfect liquid and of a viscous liquid. Part I: a perfect liquid. *Proc. R. Ir. Acad. Sect. A: Math. Phys. Sci.* **27**, 9–68 (1907)
55. Penrose, O.: Electrostatic instability of a uniform non-Maxwellian plasma. *Phys. Fluids* **3**, 258–265 (1960)
56. Rhines, P.B., Young, W.R.: How rapidly is a passive scalar mixed within closed streamlines? *J. Fluid Mech.* **133**, 133–145 (1983)
57. Scheecter, D.A., Dubin, D., Cass, A.C., Driscoll, C.F., Lansky, I.M., et al.: Inviscid damping of asymmetries on a two-dimensional vortex. *Phys. Fluids* **12**, 2397–2412 (2000)
58. Smith, G.B., Montgomery, M.T.: Vortex axisymmetrization: dependence on azimuthal wave-number or asymmetric radial structure changes. *Q. J. R. Meteorol. Soc.* **121**(527), 1615–1650 (1995)
59. Smith, R.A., Rosenbluth, M.N.: Algebraic instability of hollow electron columns and cylindrical vortices. *Phys. Rev. Lett.* **64**(6), 649 (1990)
60. Stepin, S.A.: Nonself-adjoint Friedrichs model in hydrodynamic stability. *Funct. Anal. Appl.* **29**(2), 91–101 (1995)
61. Taylor, M.E.: *Partial Differential Equations II. Qualitative Studies of Linear Equations*. Applied Mathematical Sciences, vol. 116, 2nd edn. Springer, New York (2011)
62. Thomson, W.: XXIV. Vibrations of a columnar vortex. *Lond. Edinb. Dublin Philos. Mag. J. Sci.* **10**(61), 155–168 (1880)
63. van Kampen, N.G.: On the theory of stationary waves in plasmas. *Physica* **21**, 949–963 (1955)
64. Vanneste, J.: Nonlinear dynamics of anisotropic disturbances in plane Couette flow. *SIAM J. Appl. Math.* **62**(3), 924–944 (2002)
65. Vanneste, J., Morrison, P.J., Warn, T.: Strong echo effect and nonlinear transient growth in shear flows. *Phys. Fluids* **10**, 1398 (1998)
66. Wei, D., Zhang, Z., Zhao, W.: Linear inviscid damping for a class of monotone shear flow in Sobolev spaces. *Commun. Pure Appl. Math.* **71**(4), 617–687 (2018)
67. Wei, D., Zhang, Z., Zhao, W.: Linear inviscid damping and enhanced dissipation for the Kolmogorov flow (2017). arXiv preprint [arXiv:1711.01822](https://arxiv.org/abs/1711.01822)
68. Wei, D., Zhang, Z., Zhao, W.: Linear inviscid damping and vorticity depletion for shear flows (2017). arXiv preprint [arXiv:1704.00428](https://arxiv.org/abs/1704.00428)
69. Yang, J., Lin, Z.: Linear inviscid damping for Couette flow in stratified fluid. *J. Math. Fluid Mech.* **20**, 1–28 (2016)
70. Young, B.: Landau damping in relativistic plasmas. *J. Math. Phys.* **57**(2), 021502 (2016)
71. Yu, J.H., Driscoll, C.F.: Diocotron wave echoes in a pure electron plasma. *IEEE Trans. Plasma Sci.* **30**(1), 24–25 (2002)
72. Yu, J.H., Driscoll, C.F., O’Neil, T.M.: Phase mixing and echoes in a pure electron plasma. *Phys. Plasmas* **12**, 055701 (2005)
73. Zillinger, C.: Linear inviscid damping for monotone shear flows in a finite periodic channel, boundary effects, blow-up and critical Sobolev regularity. *Arch. Ration. Mech. Anal.* **221**(3), 1449–1509 (2016)
74. Zillinger, C.: Linear inviscid damping for monotone shear flows. *Trans. Am. Math. Soc.* **369**(12), 8799–8855 (2017)
75. Zillinger, C.: On circular flows: linear stability and damping. *J. Differ. Equ.* **263**(11), 7856–7899 (2017)

Affiliations

Jacob Bedrossian¹ · Michele Coti Zelati² · Vlad Vicol^{3,4}

Jacob Bedrossian
jacob@cscamm.umd.edu

Michele Coti Zelati
m.coti-zelati@imperial.ac.uk

¹ University of Maryland, College Park, College Park, USA

² Imperial College London, London, UK

³ Princeton University, Princeton, USA

⁴ Present Address: New York University, New York, USA