The autophagy gene product BEC-1 supports normal aging and neurodevelopment in *Caenorhabditis elegans*: Integration

Nicholas Ashley and Andrea M. Holgado

St. Edward's University, Department of Biological Sciences, Austin, TX 78704, USA




Figure 1 Homozygous bec-1(ok691) mutants show a delay in the development of VD motor neurons. (A) Schematic of bec-1 gene shown in light blue and bec-1(ok691) deletion mutation in yellow (wormbase.org). (B) D-type motor neurons marked by GFP in heterozygous bec-1(ok691) mutant nematodes on day 5 of adulthood. (C) Quantification of D-type motor neuron cell bodies on days 0,1,3,5, and 7 of adulthood was recorded. Data plotted are mean \pm 1 SEM, n=60. Statistical analysis was performed using a non-parametric Kruskal-Wallis (p< .001).

Description

Aging can be defined by many different physiological pathways that emphasize changes in a variety of cells and tissues. Disease and death are often results of aging (Viña et al., 2013). However, specific mechanisms in different cell types offer multiple hypotheses regarding the definition of aging (Viña et al., 2007). Although some researchers target specific aspects of physiological processes as major contributors to aging, such as free radical theory (Harman, 2003), inflammation theory (Chung et al., 2001), mitochondrial theory (Cadenas and Davies, 2000; Sastre et al., 2000), immunologic theory (Franceschi et al., 2000), current hypotheses suggest that aging results from alterations of multiple systems working at the same time (Weinert and Timiras, 2003). Central Nervous System neural networks have been identified as particularly important for understanding cellular processes that cause aging, which is partly characterized by cell death (Yonekawa and Thorburn, 2013).

Macroautophagy (hereon referred as Autophagy) is a cellular housekeeping mechanism that uses a double membrane to target and engulf cell products for the formation of autophagosomes. These double membrane organelles then fuse



6/15/2019 – Open Access

to lysosomes where cell products are degraded and recycled (Nakamura and Yoshimori, 2018). Reports show that autophagy plays an important role in pathogen defense, development, starvation adaptations, and aging (Mizushima et al., 2008). Analysis of autophagy mutants in *C. elegans* revealed that *bec-1/Atg6/Beclin 1* is essential for dauer development, a quiescent state that survives harsh conditions such as lack of nutrients, high nematode density, and high temperatures by inducing autophagy (Meléndez et al., 2003; Meléndez and Levine 2009).

To further investigate the phenotypes associated with the bec-1(ok691) mutation, we studied nematodes possessing a 3000 base pair deletion mutation (allele ok691) in the bec-1 locus (Figure 1A). Previous reports show that the bec-1(ok691) mutation is lethal, however a small proportion of homozygous bec-1(ok691) animals reach adulthood due to maternal effect, but do not reproduce due to sterility (Melendez and Levine 2009). Our analysis of survival throughout adulthood shows that lifespan is significantly reduced in homozygous bec-1(ok691) mutants (Ashley and Holgado, 2019a). In contrast, heterozygous bec-1(ok691) mutant lifespan is not significantly different from control animals, ruling out previously discovered haploid insufficiency effects of Beclin 1 on autophagy in C. elegans (Sinha and Levine, 2008).

Previous research has found increased lifespans of daf-2 mutant C. elegans, a genetic lesion that affects the insulinlike signaling pathway and promotes constitutive dauer formation (Kimura et al. 1997). Furthermore, this daf-2dependent extension of lifespan was eliminated in bec-1(ok691); daf-2(e1370) double mutants. Together, these findings support the notion that the autophagy gene product BEC-1 is needed for the daf-2-dependent lifespan extension of C. elegans. Additionally, double mutants Atg-7 and bec-1(ok691) also show a decrease in lifespan in nematodes with inherent dietary restriction (Tóth et al., 2008).

The loss of autophagy function in the motor cortex has been associated with progression of neurodegenerative symptoms in Parkinson's disease (Kaila and Lang 2015; Fahn et al. 2004). We showed locomotion in *C. elegans* mutants for *bec-1* had significant defects (Ashley and Holgado, 2019b). These defects in locomotion suggest a role of autophagy controlling behaviors that are neuronal dependent.

Considering our findings of defective locomotion and lifespan in a bec-1(ok691) mutant background, we proceeded our research by assessing density of motor neurons in the ventral nerve cord of C. elegans. After analyzing possible effects of the bec-1(ok691) mutation on neuronal density, we followed the transgene juIs76 as it produced GFP marked D-type motor neurons (Figure 1 B). These studies showed a delay in development of ventral D-type motor neurons in bec-1(ok691) homozygous mutants (Figure 1 C). Maturation and development of VD motor neurons to the levels of controls were seen on day 5 of adulthood in bec-1(ok691) homozygous mutants (Ashley and Holgado, 2019c).

Previous research has found lineage timing of GABAergic (VD) motor neuron differentiation in *C. elegans* to occur before animals reach adulthood (Jin et al., 1994). However, our discovery in delayed lineage timing of VD motor neurons suggests a potential role of BEC-1 in neurodevelopment. This is consistent with findings in mouse models, where ortholog *Beclin 1* plays an essential role in cell differentiation during development (Cecconi and Levine, 2008). Instead of observing rapid neurodegeneration of VD motor neurons, resulting from the *bec-1(ok691)* mutation, we observed a rapid decrease in lifespan as VD motor neurons were differentiated. There is additional evidence that suggests autophagy's role in mechanisms of cell editing in early developmental stages of *C. elegans* (Di Bartolomeo et al., 2010). These conclusions should be considered as preliminary as we have not verified by an alternative line of investigation (e.g., a second allele or transgene rescue) that the observed phenotypes are specific to *bec-1(ok691)*. Specific mechanisms in which neurodegeneration act are still unknown. Future investigations observing the status of autophagy in *bec-1(ok691)* mutants may contribute novel knowledge into suggesting how autophagy may influence a delay in development of VD motor neurons or uncover pieces of the mechanism causing neurodegeneration in similar autophagy mutants.

References

Ashley, N., and Holgado, A.M. 2019a. The autophagy gene product BEC-1 supports normal aging and neurodevelopment in *Caenorhabditis elegans* (I). microPublication Biology. 10.17912/micropub.biology.000099

Ashley, N., and Holgado, A.M. 2019b. The autophagy gene product BEC-1 supports normal aging and neurodevelopment in *Caenorhabditis elegans* (II). microPublication Biology. 10.17912/micropub.biology.000100



6/15/2019 - Open Access

Ashley, N., and Hogaldo, A.M. 2019c. The autophagy gene product BEC-1 supports normal aging and neurodevelopment in *Caenorhabditis elegans* (III). microPublication Biology. 10.17912/micropub.biology.000101

Cadenas, E., and Davies, K.J. 2000. Mitochondrial Free Radical Generation, Oxidative Stress, and Aging. Free Radic. Biol. Med. **29**: 222–230. PubMed PMID: 11035250

Cecconi, F., and Levine, B. 2008. The Role of Autophagy in Mammalian Development: Cell Makeover Rather Than Cell Death. Dev. Cell 15: 344–357. PubMed PMID: 18804433

Chung, H.Y., Kim, H.J., Kim, J.W., and Yu, B.P. 2001. Molecular Inflammation Hypothesis of Aging Based on Anti-Aging Mechanism of Calorie Restriction. Microsc. Res. Tec. **59**: 327-335. PubMed PMID: 12424787

Di Bartolomeo, S., Nazio, F., Cecconi, F., and Bartolomeo, S. Di. 2010. The Role of Autophagy During Development in Higher Eukaryotes. Traffic 11: 1280–1289. PubMed PMID: 20633243

Franceschi, C., Bonafè, M., Valensin, S., Olivieri, F., De Luca, M., Ottaviani, E., and De Benedictis, g. 2000. Inflamm-aging: An Evolutionary Perspective on Immunosenescence. Ann. N. Y. Acad. Sci. **908**: 244–254. PubMed PMID: 10911963

Fahn, S., Oakes, D., Shoulson, I., Kieburtz, K., Rudolph, A., Lang, A., Olanow, CW., Tanner, C., Marek, K. 2004. Levodopa and the Progression of Parkinson's Disease. N. Engl. J. Med. 351: 2498–2508. PubMed PMID: 15590952

Harman, D. 2003. The Free Radical Theory of Aging. Antioxid. Redox Signal. 5: 557–561. PubMed PMID: 14580310

Jin, Y., Hoskins, R., Horvitz HR. 1994. Control of Type-D GABAergic Neuron Differentiation by *C. elegans* UNC-30 Homeodomain protein. Nature **372**: 780-783. PubMed PMID: 7997265

Kalia, L. V, and Lang, A.E. 2015. Parkinson's disease. Lancet 386: 896-912. PubMed PMID: 25904081

Kimura, KD, Tissenbaum HA, Liu Y, Ruvkun G. 1997. Daf-2, An Insulin Receptor-Like Gene That Regulates Longevity and Diapause in *Caenorhabditis elegans*. Science **277**: 942-946. PubMed PMID: 9252323

Meléndez, A. and Levine, B. 2009. Autophagy in C. elegans. WormBook 1–26. PubMed PMID:19705512

Meléndez, A., Tallóczy, Z., Seaman, M., Eskelinen, E.L., Hall, D.H., and Levine, B. 2003. Autophagy genes are essential for dauer development and life-span extension in *C. elegans*. Science **301**: 1387–1391. PubMed PMID: 12958363

Mizushima, N., Levine, B., Cuervo, A.M., and Klionsky, D.J. 2008. Autophagy fights disease through cellular self-digestion. Nature **451**: 1069–1075. PubMed PMID: 18305538

Nakamura, S. and Yoshimori, T. 2018. Autophagy and Longevity. Mol. Cells. 41: 66–72. PubMed PMID: 29370695

Sastre, J., Pallardo, FV., Viña, J. 2000. Mitochondrial Oxidative Stress Plays a Key Role in Aging and Apoptosis. IUBMB Life (International Union Biochem. Mol. Biol. Life) **49**: 427–435. PubMed PMID: 10902575

Sinha, S., and Levine, B. (2008). The Autophagy Effector Beclin 1: A Novel BH3-Only Protein. Oncogene 27: S137–S148. PubMed PMID: 19641499



6/15/2019 – Open Access

Tóth, M.L., Sigmond, T., Borsos, É., Barna, J., Erdélyi, P., Takács-Vellai, K., Orosz, L., Kovács, A.L., Csikós, G., Sass, M., Vellai T. 2008. Longevity Pathways Converge on Autophagy Genes to Regulate Life Span in Caenorhabditis elegans. Autophagy **4:** 330–338. PubMed PMID: 18219227

Viña, J., Borrás, C., and Miquel, J. 2007. Theories of ageing. IUBMB Life **59**: 249–254. PubMed PMID: 17505961

Viña, J., Borras, C., Abdelaziz, K.M., Garcia-Valles, R., and Gomez-Cabrera, M.C. 2013. The free radical theory of aging revisited: the cell signaling disruption theory of aging. Antioxid. Redox Signal. **19**: 779–787. PubMed PMID: 23841595

Weinert, B.T., and Timiras, P.S. 2003. Invited Review: Theories of aging. J. Appl. Physiol. **95**:1706–1716. PubMed PMID: 12970376

Yonekawa, T., and Thorburn, A. 2013. Autophagy and cell death. Essays Biochem. **55**: 105–117. PubMed PMID: 24070475

Acknowledgements

Special thanks to Dr. Stephen Steffenson for assistance with formal data analysis and Hailey Trombley for making NGM plates.

Funding

This work was supported by the National Science Foundation (award #1748523) and the Department of Biological Sciences at St. Edward's University.

Reviewed by Anonymous

Received 03/27/2019. **Accepted** 06/10/2019. **Published Online** 06/15/2019.

Copyright © 2019 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation Ashley, N; Holgado, AM (2019). The autophagy gene product BEC-1 supports normal aging and neurodevelopment in *Caenorhabditis elegans*: Integration. microPublication Biology. 10.17912/micropub.biology.000102