

Decomposing sutured monopole and instanton Floer homologies

Sudipta Ghosh and Zhenkun Li

Abstract

In this paper, we generalize the work of the second author in [19] and prove a grading shifting property, in sutured monopole and instanton Floer theories, for general balanced sutured manifolds. This result has a few consequences. First, we offer an algorithm that computes the Floer homologies of a family of sutured handle-bodies. Second, we obtain a canonical decomposition of sutured monopole and instanton Floer homologies and build polytopes for these two theories, which was initially achieved by Juhász [11] for sutured (Heegaard) Floer homology. Third, we establish a Thurston-norm detection result for monopole and instanton knot Floer homologies, which were introduced by Kronheimer and Mrowka in [16]. The same result was originally proved by Ozsváth and Szabó for link Floer homology in [23]. Last, we generalize the construction of minus versions of monopole and instanton knot Floer homology, which was initially done for knots by the second author in [19], to the case of links. Along with the construction of polytopes, we also proved that, for a balanced sutured manifold with vanishing second homology, the rank of the sutured monopole or instanton Floer homology bounds the depth of the balanced sutured manifold. As a corollary, we obtain an independent proof that monopole and instanton knot Floer homologies, as mentioned above, both detect fibred knots in S^3 . This result was originally achieved by Kronheimer and Mrowka in [16].

1 Introduction

Sutured manifold theory and Floer theory are two powerful tools in the study of 3-dimensional topology. Sutured manifolds were first introduced by Gabai in [5] and in subsequent papers. The core of sutured manifold theory is the sutured manifold hierarchy. This enables one to decompose any taut sutured manifold, in finitely many steps, into product sutured manifolds, which are the simplest possible ones. Gabai used sutured manifolds and sutured manifold hierarchies to prove some important results about 3-manifolds, including the remarkable property R conjecture.

Sutured (Heegaard) Floer homology was first introduced by Juhász in [9], while some ad hoc versions were studied by Ghiggini in [7] and Ni in [20]. In particular, Ni proved the celebrated result that the knot Floer homology, which was introduced by Ozsváth and Szabó in [22], detects fibred knots. His result is equivalent to the fact that sutured Floer homology detects product balanced sutured manifolds, which was

then generalized by Juhász in [11], where he proved that the rank of sutured Floer homology bounds the depth of a balanced sutured manifold with vanishing second homology.

The combination of sutured manifold theory with gauge theory was done by Kronheimer and Mrowka in [16], where they defined sutured monopole and instanton Floer homologies. These new Floer homologies have many significant applications in the study of knots and 3-manifolds, including a new and simpler proof of the famous property P conjecture. In [16], Kronheimer and Mrowka proved the following, in correspondence to Ni's result.

Theorem 1.1 (Kronheimer and Mrowka [16]). *Suppose (M, γ) is a balanced sutured manifold and is a homology product. Suppose further that*

$$\text{rk}(\underline{\text{SHM}}(M, \gamma)) = 1 \text{ or } \text{rk}(\underline{\text{SHI}}(M, \gamma)) = 1.$$

Then, (M, γ) is a product sutured manifold.

Theorem 1.1 has many important applications. For instance, this theorem is crucial in the proof that Khovanov homology detects unknots, by Kronheimer and Mrowka [13], and that Khovanov homology detects trefoils, by Baldwin and Sivek [4]. In this paper, we generalize Theorem 1.1 and prove the following:

Theorem 1.2. *Suppose (M, γ) is a taut balanced sutured manifold, $H_2(M) = 0$, and*

$$\text{rk}(\underline{\text{SHM}}(M, \gamma)) < 2^{k+1} \text{ or } \text{rk}(\underline{\text{SHI}}(M, \gamma)) < 2^{k+1}.$$

Then,

$$d(M, \gamma) \leq 2k.$$

Remark 1.3. If a balanced sutured manifold is a homology product, then $H_2(M) = 0$. The converse is not necessarily true. Also, $d(M, \gamma) = 0$ if and only if (M, γ) is a product sutured manifold.

As a direct corollary to Theorem 1.2, we offer a new proof to the following well-known fact.

Theorem 1.4 (Kronheimer and Mrowka [16]). *The monopole and instanton knot Floer homologies, KHM and KHI , as defined in [16], both detect fibred knots.*

In [27], Xie and Zhang constructed a version of sutured instanton Floer homology on balanced sutured manifolds with tangles. In [26], they used their construction as a tool to fully classify links whose Khovanov homologies have minimal possible ranks. One crucial step in their proofs is to show that the sutured instanton Floer homology they constructed detects product tangles inside product sutured manifolds. In this paper, with Theorem 1.2, we can prove a slightly more general result than their product-tangle-detection theorem.

Corollary 1.5. *Suppose (M, γ) is a balanced sutured manifold equipped with a vertical tangle T . Suppose further that $H_2(M \setminus T) = 0$ and $\text{SHI}(M, \gamma, T) \cong \mathbb{C}$. Then, (M, γ, T) is diffeomorphic to a product sutured manifold equipped with a product tangle, i.e.,*

$$(M, \gamma, T) \cong ([-1, 1] \times F, \{0\} \times \partial F, [-1, 1] \times \{p_1, \dots, p_n\}).$$

Here, F is a compact oriented surface-with-boundary, and $p_1, \dots, p_n$ are distinct points on F .

Remark 1.6. Note we only need the requirement that $H_2(M \setminus T) = 0$, instead of the assumption that $H_2(M) = 0$ in the hypothesis of Theorem 1.2. The later condition is stronger when tangles do exist. Our assumption is also weaker than the original assumption that (M, γ) must be a homology product in Xie and Zhang [27].

Despite many significant applications mentioned above, there are many basic aspects of sutured monopole and instanton Floer theories that remain in mystery. The usual monopole and instanton theories were defined on closed oriented 3-manifolds, while balanced sutured manifolds are compact oriented manifolds with non-trivial boundaries. So, to define the sutured monopole and instanton Floer homologies, Kronheimer and Mrowka constructed a special class of closed oriented 3-manifolds, called closures, out of the sutured data. However, the choice of closure is not unique, which lead to the following two questions:

Question 1.7. *In [14], Kronheimer and Mrowka proved that different closures give rise to isomorphic sutured monopole and instanton Floer homologies. Then, to what extent can we say that all of the essential information of sutured monopole and instanton Floer homologies is contained in the original balanced sutured manifold rather than the full closure?*

Question 1.8. *The monopole Floer homology on a closed 3-manifold decomposes along spin^c structures (see [15]). Correspondingly, the instanton Floer homology decomposes along eigenvalue functions (see [16]). Then, do we have a spin^c -type decomposition for sutured monopole or instanton Floer homology?*

Question 1.9. *How do sutured monopole and instanton Floer homologies tell us information about the Thurston norm on a balanced sutured manifold?*

Towards answering the first question, the second author proved in [19] the following proposition:

Proposition 1.10 (Li [19]). *Suppose (M, γ) is a balanced sutured manifold with a toroidal boundary, and γ consists of two components. Suppose further that Y is a closure of (M, γ) , and $\mathfrak{s}_1$ and $\mathfrak{s}_2$ are two supporting spin^c structures on Y , then there is a 1-cycle x in M so that*

$$c_1(\mathfrak{s}_1) - c_1(\mathfrak{s}_2) = P.D.[x] \in H^2(Y).$$

Similar statements hold in the instanton settings.

This theorem is central to the second author's proof of a grading shifting property for gradings associated to properly embedded surfaces inside those balanced sutured manifolds that are described in the hypothesis of Proposition 1.10. The grading shifting property has two consequences in that paper. The first is to compute the sutured monopole and instanton Floer homologies of any sutured solid tori. The second is to construct an Alexander grading on the minus versions of monopole and instanton knot Floer homologies as well as proving many fundamental properties of them.

However, in the hypothesis of Proposition 1.10, it is required that M has a toroidal boundary, and that the suture has only two components. These requirements are very restrictive. For instance, one cannot use Proposition 1.10 to construct minus versions of monopole and instanton knot Floer homologies for links. This is because the sutured manifolds arising from links may have more than one boundary component, and, thus, Proposition 1.10 does not apply. In this paper, we prove the same result, as in Proposition 1.10, for any balanced suture manifolds.

Theorem 1.11. *Suppose (M, γ) is a balanced sutured manifold. Suppose further that Y is a closure of (M, γ) , and $\mathfrak{s}_1$ and $\mathfrak{s}_2$ are two supporting spin^c structures on Y . Then, there is a 1-cycle x in M so that*

$$c_1(\mathfrak{s}_1) - c_1(\mathfrak{s}_2) = P.D.[x] \in H^2(Y).$$

Similar result holds in the instanton settings.

Thus, to answer Question 1.7, we could say that, in any closure, the difference of any two supporting spin^c structures, in terms of the Poincaré dual of their first Chern classes, is contained in the original balanced sutured manifold instead of the whole closure. Also, Theorem 1.11 leads to a generalization of the grading shifting property, which was initially discussed in Li [19], as follows.

Theorem 1.12. *Suppose (M, γ) is a balanced sutured manifold and $\alpha \in H_2(M, \partial M)$ is a non-trivial homology class. Pick two surfaces S_1 and S_2 so that*

$$[S_1, \partial S_1] = [S_2, \partial S_2] = \alpha \in H_2(M, \partial M),$$

and they are both admissible (see Definition 2.25) in (M, γ) . Then, there exist constants $l_M, l_I \in \mathbb{Z}$, so that, for any $j \in \mathbb{Z}$, we have:

$$\underline{\text{SHM}}(-M, -\gamma, S_1, j) = \underline{\text{SHM}}(-M, -\gamma, S_2, j - l_M),$$

and

$$\underline{\text{SHI}}(-M, -\gamma, S_1, j) = \underline{\text{SHI}}(-M, -\gamma, S_2, j - l_I).$$

The general grading shifting property given by Theorem 1.12 helps compute the sutured monopole and instanton Floer homology of some families of sutures on a general handle body. In section 4, we use a concrete example to present the algorithm. Theorem 1.12 also leads to a generalization of the minus version of monopole and instanton knot Floer homologies for links:

Theorem 1.13. *Suppose $L \subset Y$ is a link so that each component of L is null-homologous. Suppose further that L has r components and $\mathbf{p}$ is an r -tuple, consisting of one point on each component of L . Then, associated to the triple $(-Y, L, \mathbf{p})$, we can construct an infinite-rank module $\underline{\text{KHM}}^-(-Y, L, \mathbf{p})$ over the rings $\mathcal{R}$, the mod 2 Novikov ring. Moreover, $\underline{\text{KHM}}^-(-Y, L, \mathbf{p})$ is well defined only up to multiplication by a unit in $\mathcal{R}$ and has the following properties.*

(1) *Suppose $\{S_1, \dots, S_r\}$ is a collection of r Seifert surfaces, one for each component of L , then they together induce a $\mathbb{Z}^r$ grading on $\underline{\text{KHM}}^-(-Y, L, \mathbf{p})$ and $\underline{\text{KHI}}^-(-Y, L, \mathbf{p})$.*

(2) For each $i \in \{1, \dots, r\}$, there is a morphism

$$U_i : \underline{\text{KHM}}^-(-Y, L, \mathbf{p}) \rightarrow \underline{\text{KHM}}^-(-Y, L, \mathbf{p}),$$

which drops the grading associated to S_i by 1 and preserves all other gradings. This makes $\underline{\text{KHM}}^-(-Y, L, \mathbf{p})$ a module over $\mathcal{R}[U_1, \dots, U_r]$.

(3) There exists $N \in \mathbb{Z}$ so that if $\mathbf{j} = (j_1, \dots, j_r) \in \mathbb{Z}^r$ is a multi-grading and $j_i < N$ for some $i \in \{1, \dots, r\}$, then the morphism U_i restricts to an isomorphism

$$U_i : \underline{\text{KHM}}^-(-Y, L, \mathbf{p}, \mathbf{j}) \xrightarrow{\cong} \underline{\text{KHM}}^-(-Y, L, \mathbf{p}, \mathbf{j}^\dagger),$$

where $\mathbf{j}^\dagger$ is obtained from $\mathbf{j}$ by replacing j_i with $j_i - 1$.

Furthermore, using instanton theory, we can construct $\underline{\text{KHI}}^-(-Y, L, \mathbf{p})$, which is well defined up to a multiplication by a non-zero element in the field of complex numbers, and properties (1), (2), and (3) all hold.

To answer Question 1.8, we construct a canonical decomposition of sutured monopole and instanton Floer homologies, independent of the choices of closures. To ensure this decomposition is canonical, we need to pre-fix an element of special type inside the sutured monopole or instanton Floer homology, which we call a homogenous element.

Proposition 1.14. *Suppose (M, γ) is a balanced sutured manifold and $a \in \underline{\text{SHM}}$ is a homogenous element. Then there is a canonical decomposition*

$$\underline{\text{SHM}}(M, \gamma) = \bigoplus_{\rho \in H^2(M, \partial M; \mathbb{R})} \underline{\text{SHM}}_a(M, \gamma, \rho).$$

A similar statement holds in the instanton settings.

Thus, we could define polytopes for sutured monopole and instanton Floer theories, to be the convex hulls of sets of ρ so that $\underline{\text{SHM}}_a(M, \gamma, \rho) \neq 0$ or $\underline{\text{SHI}}_a(M, \gamma, \rho) \neq 0$. The first definition of such a polytope was introduced by Juhász in the context of sutured Floer theory. In this paper, we also proved the following:

Corollary 1.15. *Suppose (M, γ) is a taut balanced sutured manifold with $H_2(M) = 0$. Suppose further that (M, γ) is reduced, horizontally prime, and free of essential product disks. Then, the polytopes must both have maximal possible dimensions. In particular, we conclude that*

$$\text{rk}_{\mathcal{R}}(\underline{\text{SHM}}(M, \gamma)) \geq b^1(M) + 1, \text{ and } \dim_{\mathbb{C}}(\underline{\text{SHI}}(M, \gamma)) \geq b^1(M) + 1.$$

The proofs of Corollary 1.15 and Theorem 1.2 both rely on a technical result proved in section 5, which describes in detail how sutured monopole and instanton Floer homologies behave under sutured manifold decompositions. It is closely related to the decomposition theorem, Proposition 6.9, in Kronheimer and Mrowka [16].

The polytopes we defined for sutured monopole and instanton theories are closely related to the Thurston norms on the original balanced sutured manifold as well as on the closures. In particular, the canonical decomposition in Proposition 1.14 and the grading shifting property in Theorem 1.12 enable us to prove a Thurston norm detection result for monopole and instanton knot Floer homologies. The same result

was previously achieved by Ozsváth and Szábo in [23], for the link Floer homology in Heegaard Floer theory.

Suppose Y is a closed oriented 3-manifold and $L \subset Y$ is an oriented link. Let

$$L = L_1 \cup \dots \cup L_r$$

be the components of L . We require the following two conditions to hold for $L \subset Y$.

- (1) The link complement, $Y(L) = Y \setminus N(L)$, is irreducible.
- (2) The link complement $Y(L)$ is boundary-incompressible.

Let $\Gamma_\mu \subset \partial Y(L)$ be the suture consisting of a pair of oppositely oriented meridians on each boundary component of $Y(L)$. Then, by Proposition 1.14, there is a decomposition

$$\underline{\text{KHM}}(Y, L) = \underline{\text{SHM}}(Y(L), \Gamma_\mu) = \bigoplus_{\rho \in H^2(Y(L), \partial Y(L); \mathbb{Q})} \underline{\text{SHM}}_a(Y(L), \Gamma_\mu, \rho).$$

We make the following definition.

Definition 1.16. For a class $\alpha \in H_2(Y(L), \partial Y(L))$, define

$$y(\alpha) = \max_{\substack{\rho \in H^2(Y(L), \partial Y(L); \mathbb{Q}) \\ \underline{\text{SHM}}_a(Y(L), \Gamma_\mu, \rho) \neq 0}} \{\rho(\alpha)\} - \min_{\substack{\rho \in H^2(Y(L), \partial Y(L); \mathbb{Q}) \\ \underline{\text{SHM}}_a(Y(L), \Gamma_\mu, \rho) \neq 0}} \{\rho(\alpha)\}$$

Theorem 1.17. *Under the above settings,*

$$x(\alpha) + \sum_{i=1}^r |\langle \alpha, \mu_i \rangle| = y(\alpha). \quad (1)$$

Here $x(\cdot)$ is the Thurston-norm defined in Definition 2.5. $\langle, \rangle$ is to take the algebraic intersection number of a class $\alpha \in H_2(Y(L), \partial Y(L))$ with a class $[\mu_i] \in H_1(Y(L))$, where μ_i is a meridian of the link component L_i .

Remark 1.18. Theorem 1.17 offers a complete answer to Question 1.9 in the case when the boundary of the balanced sutured manifold consists of tori. Suppose M is a connected compact oriented 3-manifold so that M is irreducible and boundary-incompressible, and its boundary consists of tori, then we can perform Dehn fillings on each boundary component of M . The cores of the Dehn fillings give us a Link inside the resulting closed 3-manifold Y , which satisfies the hypothesis of Theorem 1.17. Since the Dehn surgery can be performed along any non-separating simple closed curves, Theorem 1.17 covers all the cases when the suture γ on ∂M consists of a pair of non-separating simple closed curves on each boundary component of M . For a more general suture, when it may have more than two component on some boundary component of M , we can modify the coefficients of the terms $\langle h, \mu_i \rangle$ according to the number of components, and the proof of Theorem 1.17 still applies verbatim.

Organization. In Section 2, we include the basic definitions and known results that support the proofs in this paper. In Section 3, we study the set of supporting spin^c structures on any closure of a balanced sutured manifold. This will be the basis for a generalized grading shifting result proven in Section 4. In Section 4,

we also offer an algorithm that could potentially compute the sutured monopole and instanton Floer homologies of a families of sutured handle bodies by carrying it out on a particular example. In Section 5, we use the generalized grading shifting property to construct a canonical decomposition of sutured monopole or instanton Floer homology and further construct polytopes in these two theories. Also, we prove some basic properties of the polytopes as well as the results regarding the depth of a sutured manifold. In section 6, we present some applications to knots and links: The first is to prove the Thurston-norm detection result for link complements, and the second is to construct minus versions for links.

Acknowledgement. The authors would like to thank their advisors David Shea Vela-Vick and Tomasz Mrowka, for their enormous help. The authors would like to thank András Juhász and Yi Ni for helpful comments or conversations. The first author was supported by his advisor David Shea Vela-Vick's NSF Grant 1907654 and Simons Foundation Grant 524876. The second author was supported by his advisor Tom Mrowka's NSF Grant 1808794.

2 Preliminaries

2.1 Basic definitions of Balanced sutured manifolds

Definition 2.1. A *balanced sutured manifold* is a pair (M, γ) consisting of a compact oriented 3-manifold M and a closed oriented 1-submanifold $\gamma \subset \partial M$. On ∂M , let $A(\gamma) = [-1, 1] \times \gamma$ be an annular neighborhood of $\gamma \subset \partial M$, and let

$$R(\gamma) = \partial M \setminus \text{int}(A(\gamma)).$$

They satisfy the following requirements.

- (1) Neither M nor $R(\gamma)$ has closed components.
- (2) If we orient $\partial R(\gamma) = \partial A(\gamma) = \{\pm 1\} \times \gamma$ in the same way as γ , then we require that the orientation on $\partial R(\gamma)$ induces a unique orientation on $R(\gamma)$. This orientation is called the *canonical orientation* on $R(\gamma)$. We use $R_+(\gamma)$ to denote the part of $R(\gamma)$ whose canonical orientation coincides with the boundary orientation of ∂M , and $R_-(\gamma)$ the rest.
- (3) We require that

$$\chi(R_+(\gamma)) = \chi(R_-(\gamma)).$$

Definition 2.2. A balanced sutured manifold (M, γ) is called a *product sutured manifold* if $M = [-1, 1] \times R$, $A(\gamma) = [-1, 1] \times \partial R$, $R_+(\gamma) = \{1\} \times R$ and $R_-(\gamma) = \{-1\} \times R$. Here, R is compact oriented surface with no closed components.

Definition 2.3. Suppose M is a compact oriented 3-manifold. M is called *irreducible* if every embedded 2-sphere $S^2 \subset M$ bounds an embedded 3-ball inside M .

Definition 2.4. Suppose M is a compact 3-manifold and $R \subset M$ is an embedded surface. R is called *compressible* if there is a simple closed curve $\alpha \subset R$ so that α does not bound a disk on R but bounds an embedded disk $D \subset M$ with $D \cap R = \alpha$. R is called *incompressible* if it is not compressible. A 3-manifold is called *boundary-incompressible* if its boundary is incompressible.

Definition 2.5 (Thurston norm). Suppose M is a compact 3-manifold, and suppose $U \subset \partial M$ is a submanifold of ∂M . Suppose further that S is a properly embedded surface inside M so that $\partial S \subset U$. If S is connected, then define the *norm* of S to be

$$x(S) = \max\{-\chi(S), 0\}.$$

In general, suppose the components of S are

$$S = S_1 \cup \dots \cup S_n,$$

where each S_i is connected, then define the *norm* of S to be

$$x(S) = x(S_1) + \dots + x(S_n).$$

Moreover, suppose $\alpha \in H_2(M, U)$ is a non-trivial second relative homology class, then define the *norm* of α to be

$$x(\alpha) = \min\{x(S) \mid (S, \partial S) \subset (M, U), [S, \partial S] = \alpha \in H_2(M, U)\}.$$

Definition 2.6. Suppose M is a compact 3-manifold, and $S \subset M$ is a properly embedded surface. S is called *norm-minimizing* if

$$x(S) = x(\alpha),$$

where $\alpha = [S, \partial S] \in H_2(M, N(\partial S))$. Here, $N(\partial S)$ is a neighborhood of $\partial S \subset \partial M$.

Definition 2.7 (Gabai [5]). A balanced sutured manifold (M, γ) is called *taut* if the following is true

- (1) M is irreducible.
- (2) $R_+(\gamma)$ and $R_-(\gamma)$ are both incompressible.
- (3) $R_+(\gamma)$ and $R_-(\gamma)$ are both norm-minimizing.

Definition 2.8 (Gabai [5]). Let (M, γ) be a balanced sutured manifold. A *product annulus* A in (M, γ) is an annulus properly embedded in M such that $\partial A \subset R(\gamma)$ and $\partial A \cap R_\pm(\gamma) \neq \emptyset$. A *product disk* is a disk D properly embedded in M such that $\partial D \cap A(\gamma)$ consists of two essential arcs in $A(\gamma)$.

Product annuli and product disks can detect where (M, γ) is locally a product. We have the following definition following Juhász [11].

Definition 2.9 (Juhász [11]). A balanced sutured manifold (M, γ) is called *reduced* if any product annulus $A \subset M$ either bounds a cylinder $D^2 \times I$ so that $\partial D^2 \times I = A$, or is isotopic to a component of $A(\gamma)$ inside M .

Definition 2.10 (Gabai [5]). Let (M, γ) be a taut balanced sutured manifold. A properly embedded surface $S \subset M$ is called *horizontal* if the following four properties hold.

- (1) S has no closed components and is incompressible.
- (2) $\partial S \subset A(\gamma)$, and ∂S is parallel to $\partial R_+(\gamma)$ inside $A(\gamma)$.
- (3) $[S] = [R_+(\gamma)]$ in $H_2(M, A(\gamma))$.
- (4) $\chi(S) = \chi(R_+(\gamma))$.

We say that (M, γ) is *horizontally prime* if every horizontal surface in (M, γ) is parallel to either $R_+(\gamma)$ or $R_-(\gamma)$.

Definition 2.11 (Gabai [5]). Suppose (M, γ) is a taut balanced sutured manifold. The *depth* of (M, γ) , which we write $d(M, \gamma)$, is the minimal integer n so that there exists a sequence of sutured manifold decompositions (for definitions, see Gabai [5] or Scharlemann [24])

$$(M_0, \gamma_0) \xrightarrow{S_0} (M_1, \gamma_1) \xrightarrow{S_1} \dots \xrightarrow{S_n} (M_{n+1}, \gamma_{n+1}),$$

so that each (M_i, γ_i) is taut, $(M_0, \gamma_0) = (M, \gamma)$ and (M_{n+1}, γ_{n+1}) is a product sutured manifold.

Theorem 2.12 (Gabai [5]). *For any taut balanced sutured manifold, its depth is finite.*

2.2 Monopole and instanton Floer homologies on balanced sutured manifolds

To define sutured monopole and instanton Floer homologies, one needs to construct a closed 3-manifold, together with a distinguishing surface, out of a balanced sutured manifold (M, γ) . To do this, pick T to be a connected oriented surface so that the following is true.

- (1) There is an orientation reversing diffeomorphism

$$f : \partial T \rightarrow \gamma.$$

- (2) T has genus at least 2.
- (3) There is a fixed base point $p \in T$.

Then, we can use f to glue $T \times [-1, 1]$ to M , along the annuli $A(\gamma)$, and let

$$\widetilde{M} = M \cup_{id \times f} [-1, 1] \times T.$$

The manifold $\widetilde{M}$ has two boundary components:

$$\partial \widetilde{M} = R_+ \cup R_-,$$

where

$$R_{\pm} = R_{\pm}(\gamma) \cup_f \{\pm 1\} \times T.$$

Let $h : R_+ \rightarrow R_-$ be an orientation preserving diffeomorphism so that $h(\{1\} \times \{p\}) = \{-1\} \times \{p\}$, then we can form a closed 3-manifold as follows:

$$Y = \widetilde{M} \cup_{id \cup h} [-1, 1] \times R_+.$$

Here, we use h to glue $\{1\} \times R_+$ to $R_- \subset \partial \widetilde{M}$ and use the identity map to glue $\{-1\} \times R_+$ to $R_+ \subset \partial \widetilde{M}$. Let $R = \{0\} \times R_+ \subset Y$, and we make the following definition.

Definition 2.13. The manifold $\widetilde{M}$ is called a *pre-closure* of (M, γ) . The pair (Y, R) is called a *closure* of the balanced sutured manifold (M, γ) . The choices T, f , and h are called the auxiliary data. In particular, the surface T is called an *auxiliary surface* and h a *gluing diffeomorphism*.

Definition 2.14. The *sutured monopole Floer homology* of (M, γ) is defined as

$$SHM(M, \gamma) = HM(Y|R) = \bigoplus_{\mathfrak{s} \in \mathfrak{S}^*(Y|R)} \widetilde{HM}_\bullet(Y, \mathfrak{s}; \Gamma_\eta).$$

Here, $\eta \subset R$ is a non-separating simple closed curve and

$$\mathfrak{S}^*(Y|R) = \{\mathfrak{s} \text{ spin}^c \text{ structures on } Y, c_1(\mathfrak{s})[R] = 2g(R) - 2, \widetilde{HM}_\bullet(Y, \mathfrak{s}) \neq 0\}$$

is called *the set of supporting spin^c structures*. We use the mod 2 Novikov ring $\mathcal{R}$ for coefficients.

Definition 2.15. The *sutured instanton Floer homology* of (M, γ) is defined as

$$SHI(M, \gamma) = I(Y|R) = \bigoplus_{\lambda \in \mathfrak{H}^*(Y|R)} I^\omega(Y)_\lambda.$$

Here, $\omega = S^1 \times \{p\}$, and the notation $I^\omega(Y)_\lambda$ follows from Section 6 in [18]. The set

$$\mathfrak{H}^*(Y|R) = \{\lambda \in H^2(Y; \mathbb{Q}), \lambda[R] = 2g(R) - 2, I^\omega(Y)_\lambda \neq 0\}$$

is called *the set of supporting eigenvalue functions*. We use the field of complex numbers $\mathbb{C}$ for coefficients.

Theorem 2.16. [Kronheimer and Mrowka [16]] *The isomorphism classes of $SHM(M, \gamma)$ and $SHI(M, \gamma)$ are invariants for a fixed balanced sutured manifold (M, γ) .*

Only knowing that the isomorphism class is an invariant is sometimes not enough. In [1], Baldwin and Sivek refined the definition of closures and construct canonical maps between different closures. In particular, they proved the following.

Theorem 2.17. *Suppose (M, γ) is a balanced sutured manifold. Suppose further that (Y_1, R_1) and (Y_2, R_2) are two closures of (M, γ) . Pick non-separating curves $\eta_1 \subset R_1$ and $\eta_2 \subset R_2$ to support local coefficients. Then, there exists a map*

$$\Phi_{1,2} : HM(Y_1|R_1; \Gamma_{\eta_1}) \xrightarrow{\cong} HM(Y_2|R_2; \Gamma_{\eta_2}),$$

which is well-defined up to multiplication by a unit in the base ring. Moreover, it satisfies the following properties.

(1) *If $(Y_1, R_1) = (Y_2, R_2)$, then $\Phi_{1,2} \doteq \text{id}$. Here, $\doteq$ means equal up to multiplication by a unit.*

(2) *If there is a third closure (Y_3, R_3) of (M, γ) , then*

$$\Phi_{1,3} \doteq \Phi_{2,3} \circ \Phi_{1,2}.$$

Similar results hold in the instanton settings.

Hence, for a balanced sutured manifold (M, γ) , the sutured monopole or instanton Floer homologies of closures, together with the canonical maps, form a projective transitive system, and we can derive a canonical module which we denote by

$$\underline{\text{SHM}}(M, \gamma) \text{ or } \underline{\text{SHI}}(M, \gamma).$$

They are well defined up to a unit in the corresponding base ring (field). For more details, readers are referred to Baldwin and Sivek [1].

Floer excision is a very useful tool introduced by Kronheimer and Mrowka [16] into the context of sutured monopole and instanton Floer homologies. It has several different versions, but we only present the version that is useful in later sections. Suppose (M, γ) is a balanced sutured manifold and $\widetilde{M}$ is a pre-closure of (M, γ) . Recall that

$$\partial \widetilde{M} = R_+ \cup R_-.$$

Suppose we use two gluing diffeomorphisms, h_1 and h_2 , to obtain two closures (Y_1, R_+) and (Y_2, R_+) , respectively. Let $h = h_1^{-1} \circ h_2$, and let Y_h be the mapping torus of $h : R_+ \rightarrow R_+$. Then, we can form a cobordism W , which is from $Y_1 \sqcup Y^h$ to Y_2 , as follows. Let U be a disk as depicted in Figure 1, and four parts of its boundary, μ_1, μ_2, μ_3 , and μ_4 , are each identified with the interval $[0, 1]$. Glue three pieces $[0, 1] \times \widetilde{M}$, $U \times R_+$ and $[0, 1] \times [-1, 1] \times R_+$ together, and let

$$W = ([0, 1] \times \widetilde{M}) \underset{id \cup h_1^{-1}}{\cup} (U \times R_+) \underset{id \cup h^{-1}}{\cup} ([0, 1] \times [-1, 1] \times R_+).$$

Here, we use id to glue $[0, 1] \times R_+$ to $\mu_1 \times R_+$, use h_1^{-1} to glue $[0, 1] \times R_-$ to $\mu_2 \times R_+$, use h^{-1} to glue $\mu_3 \times R_+$ to $[0, 1] \times \{1\} \times R_-$ and use id to glue $\mu_4 \times R_+$ to $[0, 1] \times \{-1\} \times R_+$. Note all the gluing maps are identity on the $[0, 1]$ direction.

Pick a non-separating curve $\eta \subset T$, and suppose the diffeomorphisms h_1 and h_2 we choose at the beginning both preserve η :

$$h_1(\{1\} \times \eta) = \{-1\} \times \eta, \text{ and } h_2(\{1\} \times \eta) = \{-1\} \times \eta.$$

Then, we can use η to support local coefficients.

Theorem 2.18 (Kronheimer and Mrowka, [16]). *The cobordism W induces an isomorphism*

$$\widetilde{HM}(W) : HM(Y_1|R_+; \Gamma_\eta) \otimes HM(Y^h|R_+; \mathbb{Z}_2) \rightarrow HM(Y_2|R_+; \Gamma_\eta).$$

There are three basic lemmas that are useful in later sections. Here we only present them in the monopole settings, but all of them have correspondences in the instanton settings.

Lemma 2.19 (Kronheimer and Mrowka [16]). *The set of supporting spin^c structures $\mathfrak{S}^*(Y^h|R_+)$ consists of a unique element $\mathfrak{s}^h$. Moreover, with $\mathbb{Z}$ coefficients,*

$$HM(Y^h|R_+) = \widetilde{HM}_\bullet(Y^h, \mathfrak{s}^h) \cong \mathbb{Z}.$$

When using local coefficients, pick a non-separating curve $\eta \subset R_+$ and suppose $\mathcal{R}$ be any suitable base ring for local coefficients, then

$$HM(Y^h|R_+; \Gamma_\eta) = \widetilde{HM}_\bullet(Y^h, \mathfrak{s}^h; \Gamma_\eta) \cong \mathcal{R}.$$




Figure 1: Gluing three parts together to get W . The middle part is $U \times R_+$, while the R_+ directions shrink to a point in the figure.

Lemma 2.20 (Kronheimer and Mrowka [15]). *Suppose Y is a closed oriented 3-manifold and $\mathfrak{s}$ is a spin^c structure on Y so that there is an embedded oriented surface $R \subset Y$ so that $g(T) \geq 1$, and $|c_1(\mathfrak{s})[R]| > 2g(R) - 2$. Then, we have*

$$\widetilde{HM}_\bullet(Y, \mathfrak{s}) = 0.$$

Similarly, for any local coefficients that could possibly be used,

$$\widetilde{HM}_\bullet(Y, \mathfrak{s}; \Gamma_\eta) = 0.$$

Lemma 2.21. *Suppose (W, ν) is a cobordism from Y to Y' . Suppose $\mathfrak{s}$ is a spin^c structure on Y and $\mathfrak{s}'$ is a spin^c structure on Y' , so that*

$$\widetilde{HM}(W)(\widetilde{HM}_\bullet(Y, \mathfrak{s})) \cap \widetilde{HM}_\bullet(Y', \mathfrak{s}') \neq \{0\},$$

then we know that

$$i_*(P.D.c_1(\mathfrak{s})) = (i')_*(P.D.c_1(\mathfrak{s}')) \in H_1(W).$$

Here $i : Y \rightarrow W$ and $i' : Y' \rightarrow W$ are the inclusions.

2.3 A grading on sutured monopole and instanton Floer homology

In [19], the second author constructed a grading on sutured monopole or instanton Floer homology, associated to a properly embedded surface with a connected boundary. We will present the construction in this subsection, while dropping the condition that ∂S is connected, using some new inputs from Kavi [12].

Definition 2.22. Suppose (M, γ) is a balanced sutured manifold, and S is a properly embedded oriented surface. A *stabilization* of S is an isotopy of S to a surface S' , so that the isotopy creates a new pair of intersection points:

$$\partial S' \cap \gamma = (\partial S \cap \gamma) \cup \{p_+, p_-\}.$$

We require that there are arcs $\alpha \subset \partial S'$ and $\beta \subset \gamma$, which are oriented in the same way as $\partial S'$ and γ , respectively, so that the following is true.

- (1) We have $\partial \alpha = \partial \beta = \{p_+, p_-\}$.
- (2) The curves α and β cobound a disk D so that $\text{int}(D) \cap (\gamma \cup \partial S') = \emptyset$. The stabilization is called *negative* if D can be oriented so that $\partial D = \alpha \cup \beta$ as oriented curves. It is called *positive* if $\partial D = (-\alpha) \cup \beta$.

Denote by $S^{\pm k}$ the result of performing k many positive or negative stabilizations of S .




Figure 2: The positive and negative stabilizations of S .

Lemma 2.23. Suppose (M, γ) is a balanced sutured manifold, and S is a properly embedded oriented surface. Suppose further that S^+ and S^- are obtained from S by performing a positive and negative stabilization, respectively. Then, we have the following.

- (1) If we decompose (M, γ) along S or S^+ , then the resulting two balanced sutured manifolds are diffeomorphic.

(2) If we decompose (M, γ) along S^- , then the resulting balanced sutured manifold, (M', γ') , is not taut, because $R_\pm(\gamma')$ are both compressible.

Remark 2.24. The positive and negative stabilization on S will be switched if we reverse the orientation of S , that is, $-(S^+)$ is the same as $(-S)^-$ and $-(S^-)$ is the same as $(-S)^+$. Accordingly, when changing the orientation of the suture, positive and negative stabilizations are also switched.

Now we present the construction of the grading. This was originally written down by Baldwin and Sivek in [4] and was then generalized one step further by the second author in [19], to fit his needs of constructing a $\mathbb{Z}$ grading in the minus version of monopole and instanton knot Floer homologies. In this subsection, we make a further generalization and introduce the most general setups of constructing such a grading.

Definition 2.25. Suppose (M, γ) is a balanced sutured manifold, and S is a properly embedded surface inside M . Suppose further that S intersects with γ transversely. S is called *admissible* inside (M, γ) if the value $(\frac{1}{2}|S \cap \gamma| - \chi(S))$ is an even integer.

Suppose (M, γ) is a balanced sutured manifold, and $S \subset M$ is an oriented admissible properly embedded surface. In case $\partial S \neq \emptyset$, suppose further that S intersects with γ transversely. Let $n = \frac{1}{2}|S \cap \gamma|$. We fix an arbitrary ordering of the boundary components of S and label all the intersection points of S with γ as follows. On each boundary component of S , index them according to the orientation of ∂S . For different components of ∂S , we first index points on the boundary component that comes first in the fixed ordering. Also, the first point to be indexed on each boundary component of S is chosen to be a positive intersection of S with γ (on ∂M). In this way, we can assume that

$$S \cap \gamma = \{p_1, \dots, p_{2n}\}.$$

In [19], when S has a connected boundary, the second author introduced the notion of balanced pairings to help construct the grading. In Kavi [12], the notion of balanced pairings was generalized to accommodate a general S . In this paper, we omit the detailed definitions of balanced pairings and will use the generalized definition from [12].

Suppose

$$\mathcal{P} = \{(i_1, j_1), \dots, (i_n, j_n)\}$$

is a balanced pairing of size n . Then, we can pick an auxiliary surface T for (M, γ) so that the following is true.

- (1) The genus of T is large enough.
- (2) The boundary of T is identified with the suture γ .
- (3) There are properly embedded arcs $\alpha_1, \dots, \alpha_n$ inside T so that the following two properties hold.
 - (a) The classes $[\alpha_1], \dots, [\alpha_n]$ are linearly independent in $H_1(T, \partial T)$.
 - (b) For $k = 1, \dots, n$, we have

$$\partial \alpha_k = \{p_{i_k}, p_{j_k}\}.$$

Then, we can form a pre-closure $\widetilde{M}$ of (M, γ) :

$$\widetilde{M} = M \cup [-1, 1] \times T.$$

The manifold $\widetilde{M}$ has two boundary components:

$$\partial\widetilde{M} = R_+ \cup R_-.$$

The surface S extends to a properly embedded surface $\widetilde{S}$ inside $\widetilde{M}$:

$$\widetilde{S} = S \cup [-1, 1] \times \alpha_1 \cup \dots \cup [-1, 1] \times \alpha_n.$$

The definition of the balanced pairing makes sure that $\widetilde{S} \cap R_+$ and $\widetilde{S} \cap R_-$ have the same number of components, and the requirement (a) for α_i makes sure that components of $\widetilde{S} \cap R_{\pm}$ represent linearly independent classes in $H_1(R_{\pm})$. Thus, there exists an orientation preserving diffeomorphism $h : R_+ \rightarrow R_-$ so that

$$h(\widetilde{S} \cap R_+) = \widetilde{S} \cap R_-.$$

We can use $\widetilde{M}$ and h to obtain a closure (Y, R_+) of (M, γ) , and, inside Y , the surface S extends to a closed surface $\bar{S}$.

Definition 2.26. Define

$$SHM(M, \gamma, S, i) = \bigoplus_{\substack{\mathfrak{s} \in \mathfrak{S}^*(Y|R) \\ c_1(\mathfrak{s})[R] = 2i}} \widetilde{HM}_{\bullet}(Y, \mathfrak{s}; \Gamma_{\eta}).$$

We say that this grading is associated to the surface $S \subset M$. The grading defined on separate closures also induces a grading on the canonical module $\underline{SHM}(M, \gamma)$, as stated in Theorem 2.27. We write this grading on the canonical module as

$$\underline{SHM}(M, \gamma, S, i).$$

Theorem 2.27 (Kavi [12] and Li [19]). *Suppose (M, γ) is a balanced sutured manifold, and $S \subset M$ is a fixed oriented admissible properly embedded surface. Then, the grading $\underline{SHM}(M, \gamma, S, i)$ is independent of all the choices made in the construction and, thus, is well-defined.*

Using the grading, we can re-formulate Kronheimer and Mrowka's decomposition theorem, Proposition 6.9, in [16], as follows.

Lemma 2.28. *Suppose (M, γ) is a balanced sutured manifold and $S \subset M$ is an oriented admissible properly embedded surface. Suppose further that S satisfies the hypothesis of Proposition 6.9 in [13], and (M', γ') is obtained from (M, γ) by a sutured manifold decomposition along S . Let*

$$g_c = \frac{1}{4}|S \cap \gamma| - \frac{1}{2}\chi(S).$$

Then, we have

$$\underline{SHM}(M, \gamma, S, g_c) \cong \underline{SHM}(M', \gamma').$$




Figure 3: The by-pass exact triangle.

2.4 Bypasses

Suppose we have three balanced sutured manifold (M, γ_1) , (M, γ_2) and (M, γ_3) so that the underlining 3-manifolds are the same, but the sutures are different. Suppose further that γ_1 , γ_2 , and γ_3 are only different within a disk $D \subset \partial M$, and, within the disk D , they are depicted as in Figure 3.

Theorem 2.29 (Baldwin and Sivek [2, 4]). *There are exact triangles relating the sutured monopole and instanton Floer homologies of three balanced sutured manifolds as follows.*

$$\begin{array}{ccc}
 \underline{\text{SHM}}(-M, -\gamma_1) & \xrightarrow{\psi_{12}} & \underline{\text{SHM}}(-M, -\gamma_2) \\
 & \swarrow \psi_{31} \quad \searrow \psi_{23} & \\
 & \underline{\text{SHM}}(-M, -\gamma_3) & \\
 \\
 \underline{\text{SHI}}(-M, -\gamma_1) & \xrightarrow{\psi_{12}} & \underline{\text{SHI}}(-M, -\gamma_2) \\
 & \swarrow \psi_{31} \quad \searrow \psi_{23} & \\
 & \underline{\text{SHI}}(-M, -\gamma_3) &
 \end{array}$$

In contact geometry, a by-pass is a half disk, which carries some special contact structure, attached along a Legendrian arc to a convex surface. For more details, readers are referred to Honda [8]. There is a description of the maps in the above by-pass exact triangle as follows. We explain how to obtain the map ψ_{12} , and the other two are the same. Let $Z = \partial M \times [0, 1]$, and we can pick the suture γ_1 on $\partial M \times \{0\}$ as well as the suture γ_2 on $\partial M \times \{1\}$. Then, there is a special contact structure ξ_{12} on Z , which corresponds to the by-pass attachment. Hence, we can attach Z to M by the identification $\partial M \times \{0\} = \partial M \subset M$. The result, $(M \cup Z, \gamma_2)$, is diffeomorphic to (M, γ_2) , and we have

$$\psi_{12} = \Phi_{\xi_{12}}.$$

Here, $\Phi_{\xi_{12}}$ is the gluing map associated to ξ_{12} , as constructed by the second author in [18].

There is a second way to interpret the maps $\psi_{\pm}$ associated to by-pass attachments based on [21]. In [21], Ozbagci proved that a by-pass attachment could be realized by attaching a contact 1-handle followed by a contact 2-handle. In sutured monopole and instanton Floer homologies, there are maps associated to the contact handle attachments, due to Baldwin and Sivek [2, 3]. So, we can compose those contact handle attaching maps to obtain $\psi_{\pm}$. This was the original way Baldwin and Sivek constructed the by-pass maps (when they define by-pass maps, there was no construction of gluing maps) and proved the existence of the exact triangle. The two interpretations are the same because of the functoriality of the gluing maps. For details, readers are referred to Li [18]. Both descriptions of the by-pass maps are useful in later sections.

3 Difference of supporting spin^c structures

3.1 A basic calculation

Suppose (M, γ) is a balanced sutured manifold, and $\widetilde{M}$ is a pre-closure of M with $\partial \widetilde{M} = R_+ \cup R_-$. Suppose further that we pick two gluing diffeomorphisms $h_1, h_2 : R_+ \rightarrow R_-$ and obtain two closures (Y_1, R_+) and (Y_2, R_+) of (M, γ) , respectively. Let $h = h_1^{-1} \circ h_2$, and let Y^h be the mapping torus of h . As in Subsection 2.2, we can construct a Floer excision cobordism W from $Y_1 \sqcup Y^h$ to Y_2 . Suppose $i : Y_2 \rightarrow W$ is the inclusion map. In section 4 of Li [18], the map

$$i_* : H_1(Y_2) \rightarrow H_1(W),$$

which is induced by the inclusion $i : Y_2 \hookrightarrow W$, has played a very important role in proving Proposition 1.10. In this subsection we compute the kernel of i_* . A first observation is that we could just work with $\mathbb{Q}$ coefficients, since for grading purpose, torsion parts have no contributions.

From the description in Subsection 2.2, we know that W is obtained by gluing three pieces together. We can compute its first homology by applying Mayer-Vietoris sequences twice and get the following result.

$$H_1(W; \mathbb{Q}) = [H_1(\widetilde{M}; \mathbb{Q}) \oplus \langle [s_1], [s^h] \rangle] / [\text{im}(h_{1,*} - 1) + \text{im}(h_* - 1)]. \quad (2)$$

Here are a few things to be explained. First, recall in section 2, we require that the gluing diffeomorphism h_1 and h_2 to fix the same base point $p \in T$. This means that there are circles $s_1 \subset Y_1$ and $s^h \subset Y^h$ of the form $\{p\} \times S^1$, respectively. Then, s_1 and s^h naturally embed into W , and the class $[s_1], [s^h] \in H_1(W)$ are represented by these two circles.

Second, recall we have a map $h_1 : R_+ \rightarrow R_-$, and, thus, there is a map

$$\begin{pmatrix} -1 & 0 \\ (h_1)_* & 0 \end{pmatrix} : H_1(R_+) \oplus H_1(R_-) \rightarrow H_1(R_+) \oplus H_1(R_-).$$

Note $H_1(R_+) \oplus H_1(R_-)$ can be viewed as $H_1(R_+ \sqcup R_-)$ and there is an inclusion $j : R_+ \sqcup R_- \rightarrow \widetilde{M}$. So, in equation (2), we use $\text{im}(h_{1,*} - 1)$ to denote the subspace

$$\text{im} \left[j_* \circ \begin{pmatrix} -1 & 0 \\ (h_1)_* & 0 \end{pmatrix} \right] \subset H_1(\widetilde{M}).$$

Note, from the maps $h : R_+ \rightarrow R_+$ and $j : R_+ \sqcup R_- \rightarrow \widetilde{M}$, we have a space $j_*[\text{im}(h_* - 1)] \subset H_1(\widetilde{M})$. Abusing the notation, in (2), we use $\text{im}(h_* - 1)$ to denote $j_*[\text{im}(h_* - 1)]$ and omit j_* from the notation. The sum $\text{im}(h_{1,*} - 1) + \text{im}(h_* - 1)$ is the sum of the two subspaces, as described above, in $H_1(\widetilde{M})$.

In a similar way, we can compute the first homology of Y_2 .

$$H_1(Y_2; \mathbb{Q}) = [H_1(\widetilde{M}; \mathbb{Q}) \oplus \langle [s_2] \rangle] / [\text{im}(h_{2,*} - 1)]. \quad (3)$$

Here, the term $\text{im}(h_{2,*} - 1)$ is similar to the term $\text{im}(h_{1,*} - 1)$ in (2), and s_2 is the circle $S^1 \times \{p\} \subset Y_2$, similar to s_1 and s^h in (2). Hence, we can deduce the following.

Lemma 3.1. *Let $i : Y_2 \rightarrow W$ be the inclusion. Then,*

$$\ker(i_*) \subset [\text{im}(h_{1,*} - 1)] / [\text{im}(h_{2,*} - 1)] \subset H_1(Y_2; \mathbb{Q}).$$

Here, the term $\text{im}(h_{1,*} - 1)$ is the same as the one appeared in (2), and the term $\text{im}(h_{2,*} - 1)$ is the same as the one appeared in (3).

Proof. It is straight forward to check that

$$i_*([s_2]) = [s_1] + [s^h] \in H_2(W; \mathbb{Q}).$$

Hence, from (3), we know that the kernel must come from the quotient of $H_1(\widetilde{M}; \mathbb{Q})$:

$$\ker(i_*) \subset H_1(\widetilde{M}; \mathbb{Q}) / [\text{im}(h_{2,*} - 1)] \subset H_1(Y_2; \mathbb{Q}).$$

Let $\alpha \in \ker(i_*)$ be any element in the kernel, then, from (3) and (2), we can find a lift $\tilde{\alpha} \in H_1(\widetilde{M}; \mathbb{Q})$ so that $\tilde{\alpha} \in \text{im}(h_{1,*} - 1) + \text{im}(h_* - 1)$. Equivalently, there are classes $\beta, \gamma \in H_1(R_+; \mathbb{Q})$ so that (recall $j : R_+ \sqcup R_- \rightarrow \widetilde{M}$ is the inclusion)

$$\tilde{\alpha} = j_*[(h_1)_*(\beta) - \beta] + j_*[h_*(\gamma) - \gamma].$$

Write

$$\tilde{\alpha}' = h_*(\gamma) - \gamma \in H_1(R_+; \mathbb{Q}) \subset H_1(R_+ \sqcup R_-; \mathbb{Q}),$$

then we know that (recall $h_2 = h_1 \circ h$)

$$\begin{aligned}\tilde{\alpha}' &= (h_1)_*(\tilde{\alpha}') + [\tilde{\alpha}' - (h_1)_*(\tilde{\alpha}')] \\ &= (h_2)_*(\gamma) - (h_1)_*(\gamma) + [\tilde{\alpha}' - (h_1)_*(\tilde{\alpha}')] \\ &= (h_2)_*(\gamma) - \gamma + [\gamma - (h_1)_*(\gamma)] + [\tilde{\alpha}' - (h_1)_*(\tilde{\alpha}')].\end{aligned}$$

Hence, we have

$$\begin{aligned}\tilde{\alpha} &= j_*[(h_2)_*(\gamma) - \gamma] \\ &\quad + j_*[\gamma - (h_1)_*(\gamma)] + j_*[\tilde{\alpha}' - (h_1)_*(\tilde{\alpha}')] + j_*[(h_1)_*(\beta) - \beta].\end{aligned}$$

The first term is in $\text{im}(h_{2,*} - 1)$ and the rest are in $\text{im}(h_{1,*} - 1)$. So, we know that

$$\alpha \in [\text{im}(h_{1,*} - 1)]/[\text{im}(h_{2,*} - 1)] \subset H_1(\widetilde{M}, \mathbb{Q})/[\text{im}(h_{2,*} - 1)] = H_1(Y_2; \mathbb{Q}),$$

and this concludes the proof of Lemma 3.1. $\square$

3.2 Adding 1-handles

Definition 3.2. Suppose (M, γ) is a balanced sutured manifold. A *product (or contact) 1-handle* is a tuple (ϕ, S, D^3, δ) , where $S \subset \partial D^3$ is the disjoint union of two embedded disks on ∂D^2 , δ is a simple closed curve on ∂D^3 , which intersects each component of S in an arc, and $\phi : S \rightarrow \partial M$ is an embedding so that $\phi(\delta \cap S) = \gamma \cap \phi(S) \subset \partial M$. Then, we can form a new balanced sutured manifold

$$(M', \gamma') = (M \cup_{\phi} D^3, \gamma' = \gamma \setminus \phi(S) \cup (\delta \setminus S)).$$

Remark 3.3. In Kronheimer and Mrowka [16], this process is called attaching a product 1-handle, while in Baldwin and Sivek [2], the same process is called attaching a contact 1-handle.

Lemma 3.4 (Kronheimer, Mrowka [16] or Baldwin, Sivek [2]). *When using auxiliary surfaces of large enough genus, any pre-closure of (M, γ) is a pre-closure of (M', γ') , and vice-versa.*

Hence, by Lemma 3.4, we can freely add 1-handles to the original (M, γ) without changing its closure. A straightforward observation is the following.

Lemma 3.5. *For any balanced sutured manifold (M, γ) , there exists a set of 1-handles, $\{h_1, \dots, h_n\}$ so that the following is true.*

(1) *If (M', γ') is the resulting balanced sutured manifold after attaching all 1-handles $h_1, \dots, h_n$, then γ' is connected.*

(2) *For $l = 1, \dots, n$, if (M_l, γ_l) is the resulting balanced sutured manifold after attaching all 1-handles $h_1, \dots, h_n$ except h_l , then $R_{\pm}(\gamma_l)$ are both connected.*

Remark 3.6. The first condition is used in the proof of Lemma 3.7, and the second is used in the proof of Lemma 3.9.

Thus, from any balanced sutured manifold (M, γ) , we can find a set of 1-handles, $\{h_1, \dots, h_n\}$, according to Lemma 3.5. Let (M', γ') be the resulting balanced sutured manifold after attaching all those 1-handles, then we have the following lemma.

Lemma 3.7. *Suppose (Y, R) is a closure of (M, γ) with $g(R)$ large enough. Then, (Y, R) can also be regarded as a closure of (M', γ') by Lemma 3.4. Suppose further that $\mathfrak{s}_1$ and $\mathfrak{s}_2$ are two supporting spin^c structures on Y , then there is a 1-cycle x in M' so that*

$$P.D.c_1(\mathfrak{s}_1) - P.D.c_1(\mathfrak{s}_2) = [x] \in H_1(Y; \mathbb{Q}).$$

Similar results hold in the instanton settings.

Proof. We start by constructing a special reference closure of (M', γ') . Pick an auxiliary surface T' , and let

$$\widetilde{M} = M' \cup [-1, 1] \times T'$$

be a pre-closure of (M', γ') that is also a pre-closure of (M, γ) . We have

$$\partial \widetilde{M} = R_+ \cup R_-, \quad R_{\pm} = R_{\pm}(\gamma') \cup \{\pm 1\} \times T'.$$

We can pick a special gluing diffeomorphism $h^r : R_+ \rightarrow R_-$ so that $h^r|_{\{1\} \times T'} = id_{T'}$. Let (Y^r, R_+) be the closure of (M', γ') arising from h^r and $\widetilde{M}$. We know that h^r can be split into two parts, $f^r = h^r|_{R_+(\gamma')}$ and $id_{T'}$.

Thus, we have an alternative interpretation for Y^r . First, we can use $f^r : R_+(\gamma') \rightarrow R_-(\gamma')$ to glue $R_+(\gamma') \subset \partial M'$ to $R_-(\gamma') \subset \partial M'$, and M' becomes a manifold M'_1 with a toroidal boundary. Note $f^r|_{\partial R_+(\gamma')} = id$, so we have a natural framing s^r and γ' on $\partial M'_1$. Here, s^r is obtained as follows. If $q \in \gamma'$ is a point, then we have an arc $[-1, 1] \times \{q\} \subset A(\gamma)$. Note f^r identifies $\{1\} \times \{q\}$ with $\{-1\} \times \{q\}$, so $[-1, 1] \times \{q\}$ becomes a circle s^r inside $\partial M'_1$. Second, we can glue M'_1 and $S^1 \times T'$ together to form Y^r :

$$Y^r = M'_1 \cup_{\phi} S^1 \times T',$$

where $\phi : \partial M'_1 \rightarrow \partial(S^1 \times T')$ maps s^1 to the S^1 direction and maps γ' to $\partial T'$ direction. Let $g(T)' = k$, and let $\{a_1, b_1, \dots, a_k, b_k\}$ be a set of generators of $H_1(T')$ as in Figure 4. Then, we can use MayerVietoris sequence to conclude the following:

$$H_1(Y^r; \mathbb{Q}) = \left(H_1(M'; \mathbb{Q}) / [\text{im}(f_*^r - 1)] \right) \oplus \langle [s^r], [a_1], \dots, [b_k] \rangle. \quad (4)$$

Here, the term $\text{im}(f_*^r - 1)$ is similar to the term $\text{im}(h_{1,*} - 1)$ in (2). Suppose $\mathfrak{s}$ is a supporting spin^c structure on Y^r , then we can write $P.D.c_1(\mathfrak{s})$ in terms of the description of $H_1(Y^r)$ in 4. The coefficient of the class $[s^r]$ can be understood to be $2g(R) - 2$ by looking at the pairing


$$c_1(\mathfrak{s})[R] = 2g(R) - 2.$$

The coefficients of $[a_1], \dots, [b_k]$ are all zero, since we can apply the adjunction inequality in Lemma 2.20 to the tori $b_1 \times S^1, \dots, a_k \times S^1 \subset Y$. Thus, we conclude the following.

Lemma 3.8. *Suppose $\mathfrak{s}$ is a supporting spin^c structure on Y^r , then there is a 1-cycle $[x]$ in M' so that*

$$P.D.c_1(\mathfrak{s}) = [x] + (2g - 2)[s^r] \in H_1(Y^r; \mathbb{Q}).$$

Similar statement holds in the instanton settings.


 Figure 4: A basis for $H_1(T')$.

Now suppose $h : R_+ \rightarrow R_-$ is any gluing diffeomorphism, and (Y, R) is the resulting closure of (M', γ') . Let $\psi = (h^r)^{-1} \circ h : R_+ \rightarrow R_+$ be a diffeomorphism, and let Y^ψ be the mapping torus of ψ . We can form an excision cobordism W from $Y^r \sqcup Y^\psi$ to Y , as in Subsection 2.2. Then, we can compute via Mayer-Vietoris sequences that

$$H_1(Y^\psi; \mathbb{Q}) = \left(H_1(R_+; \mathbb{Q}) / [\text{im}(\psi_* - 1)] \right) \oplus \langle [s^\psi] \rangle. \quad (5)$$

Here, s^ψ is the same as the term s^h in (2). Let $i : Y \rightarrow W$, $i^r : Y^r \rightarrow W$, and $i^\psi : Y^\psi \rightarrow W$ be the inclusions. Suppose $\mathfrak{s}$ is a supporting spin^c structure on Y , then, from Lemma 2.21, we know that there is a supporting spin^c structure $\mathfrak{s}^r$ on Y^r so that

$$i_*[P.D.c_1(\mathfrak{s})] = i_*^r[P.D.c_1(\mathfrak{s}^r)] + i_*^\psi[P.D.c_1(\mathfrak{s}^\psi)] \in H_1(W; \mathbb{Q}).$$

Here, $\mathfrak{s}^\psi$ is the unique spin^c structure on Y^ψ , as in Lemma 2.19. To find all possible values of $P.D.c_1(\mathfrak{s})$, we first find a class $[z] \in H_1(Y; \mathbb{Q})$ so that

$$i_*([z]) = i_*^r[P.D.c_1(\mathfrak{s}^r)] + i_*^\psi[P.D.c_1(\mathfrak{s}^\psi)] \in H_1(W; \mathbb{Q}),$$

and then

$$P.D.c_1(\mathfrak{s}) \in [z] + \ker(i_*) \subset H_1(Y; \mathbb{Q}).$$

Note $\ker(i_*)$ has been understood by Lemma 3.1.

To find the class $[z]$, by Lemma 3.8, we know that there exists a 1-cycle x in M' so that

$$P.D.c_1(\mathfrak{s}^r) = [x] + (2g - 2)[s^r] \in H_1(Y^r; \mathbb{Q}).$$

For $\mathfrak{s}^\psi$, we can check similarly (or see Subsection 4.2 in Li [19]) that there is a 1-cycle y in R_+ so that

$$P.D.c_1(\mathfrak{s}^\psi) = [y] + (2g - 2)[s^\psi] \in H_1(Y^\psi; \mathbb{Q}).$$

Inside W , there are annuli $x \times [0, 1]$ and $y \times [0, 1]$ and a pair of pants from $s^r \sqcup s^\psi \subset Y^r \sqcup Y^\psi$ to $s \subset Y$. Hence, we can take z to be the 1-cycle in Y , of the form

$$z = x + y + (2g - 2)s.$$

Note x is in M' and y is in R_+ , so there are natural ways to regard them as 1-cycles in Y . Thus, we know that

$$i_*([z]) = i_*^r[P.D.c_1(\mathfrak{s}^r)] + i_*^\psi[P.D.c_1(\mathfrak{s}^\psi)] \in H_1(W; \mathbb{Q})$$

and, hence,

$$P.D.c_1(\mathfrak{s}) \in [x] + [y] + (2g - 2)[s] + \ker(i_*) \subset H_1(Y; \mathbb{Q}). \quad (6)$$

From Lemma 3.1, we know that

$$\ker(i_*) \subset [\text{im}(h_*^r - 1)] / [\text{im}(h_* - 1)].$$

Note that $h_*^r - 1 = f_*^r - 1$ since $h|_{T' \times \{1\}} = id_{T'}$. By construction, $[\text{im}(f_*^r - 1)]$ is contained in $H_1(M')$, and, thus, we know that $\ker(i_*)$ can only contribute to the part $[x]$ in (6). As a result, we conclude that there is a 1-cycle x' in M' so that

$$P.D.c_1(\mathfrak{s}) = [x'] + [y] + (2g - 2)[s] \in H_1(Y; \mathbb{Q}).$$

Note $[y] + (2g - 2)[s]$ is independent of the choice of the supporting spin^c structure on Y , so we conclude the proof of Lemma 3.7. $\square$

3.3 Dropping 1-handles

Recall we have a balanced sutured manifold (M, γ) , and we have a set of 1-handles, $\{h_1, \dots, h_n\}$, as in Lemma 3.5. Recall further that (M', γ') is the resulting balanced sutured manifold after attaching all of those 1-handles. In Lemma 3.7, we prove that, in terms of the Poincaré dual of the first Chern classes, the difference of two supporting spin^c structures on Y is contained in M' . In this subsection, we sharpen the result and prove that the difference must lie in M instead of the whole M' , which is exactly the statement of Theorem 1.11.

Suppose, for $l = 1, \dots, n$, (M_l, γ_l) is the resulting balanced sutured manifold after attaching all 1-handles $h_1, \dots, h_n$ except h_l . Then, (M', γ') is obtained from (M_l, γ_l) by attaching h_l . Recall, from Subsection 3.2, we have an auxiliary surface T' for M' and a pre-closure $\widetilde{M}$. Then, from Lemma 3.4, $\widetilde{M}$ is also a pre-closure for (M_l, γ_l) . Thus, any closure (Y, R) arising from $\widetilde{M}$ is also a closure for (M_l, γ_l) .

Lemma 3.9. *For any fixed l , suppose (Y, R) is a closure of (M, γ) with $g(R)$ large enough, then (Y, R) is also a closure for (M_l, γ_l) by Lemma 3.4. Suppose further that $\mathfrak{s}_1, \mathfrak{s}_2$ are two supporting spin^c structures on Y , then there is a 1-cycle in M_l so that*

$$P.D.c_1(\mathfrak{s}_1) - P.D.c_1(\mathfrak{s}_2) = [x] \in H_1(Y; \mathbb{Q}).$$

Similar result holds in the instanton settings.

Proof of Theorem 1.11. Note (M', γ') and (M_l, γ_l) are obtained from (M, γ) by attaching 1-handles. So, there are injections

$$H_1(M; \mathbb{Q}) \hookrightarrow H_1(M_l; \mathbb{Q}) \hookrightarrow H_1(M'; \mathbb{Q}).$$

Also, inside $H_1(M'; \mathbb{Q})$, we have

$$H_1(M) = \bigcap_{l=1}^n H_1(M_l).$$

So, Lemma 3.9, together with Lemma 3.4, implies Theorem 1.11. $\square$

Proof of Lemma 3.9. By Lemma 3.7, we know that there is a 1-cycle x in M' which satisfies the statement of the lemma, but our goal is to show that x can be chosen inside M_l . Recall, in the proof of Lemma 3.7, we pick a special reference closure Y^r by requiring $h^r|_{T' \times \{1\}} = id_{T'}$ (and $f^r = h^r|_{R_+(\gamma')}$). To prove the current lemma, we will need an even more special closure, by making further restrictions on f^r . Though we will keep using the notations Y^r , h^r and f^r , etc.

To explain the further restriction on f^r , recall that (M', γ') is obtained from (M_l, γ_l) by attaching the 1-handle h_l . We can write

$$H_1(M'; \mathbb{Q}) = H_1(M_l; \mathbb{Q}) \oplus \langle \alpha_l \rangle,$$

where α_l consists of the core of the 1-handle h_l together with an arc inside M_l . Let D_l be the co-core of the 1-handle h_l . It is a properly embedded disk $D_l \subset M'$ that intersects α_l transversely at one point, and ∂D_l intersects $R_\pm(\gamma')$ in arcs $\beta_{l,\pm}$. See Figure 5. Note, by the condition (2) in Lemma 3.5, the arcs $\beta_{l,\pm}$ are non-separating inside $R_\pm(\gamma')$. Thus, from Lemma 3.6 in Li [18], we can find a map f^r which sends β_+ to β_- .




Figure 5: The one handle h_l .

Now we can use the new h^r to obtain the closure (Y^r, R_+) . It not only satisfies all requirements needed to conclude the proof of Lemma 3.7, but also has some new features. Suppose the co-core D_l intersects γ' at two points p_l, q_l and let $\gamma \subset T'$ be a simple arc with end points p_l, q_l . Then, we know that

$$\tilde{D}_l = D_l \cup (\gamma \times [-1, 1])$$

is a properly embedded surface in $\tilde{M}$ so that

$$\partial \tilde{D}_l \cap R_\pm = C_\pm = \beta_\pm \cup \gamma \times \{\pm 1\}.$$

From the above construction, we know that $h^r(C_+) = C_-$. Hence, $\tilde{D}_l$ becomes a torus $T_l \subset Y^r$. (It is straightforward to compute the Euler characteristic to see that it is indeed a torus.) Note this torus T_l intersects α_l transversely at one point, but is disjoint from all other generators of

$$H_1(M'; \mathbb{Q}) = H_1(M_l; \mathbb{Q}) \oplus \langle [\alpha_l] \rangle.$$

As a result, we can use the adjunction inequality in Lemma 2.20 to make a refinement of Lemma 3.8 as follows.

Lemma 3.10. *Suppose $\mathfrak{s}$ is a supporting spin^c structure on Y^r , then there is a 1-cycle $[x]$ in M_l so that*

$$P.D.c_1(\mathfrak{s}) = [x] + (2g - 2)[s^r] \in H_1(Y^r; \mathbb{Q}).$$

Similar results hold in the instanton settings.

Suppose (Y, R_+) is an arbitrary closure of (M, γ) , arising from the pre-closure $\tilde{M}$ and a gluing diffeomorphism h . We can form the diffeomorphism $\psi = (h^r)^{-1} \circ h$, the mapping torus Y^ψ , and the excision cobordism W , as in the proof of Lemma 3.7. Suppose further that $\mathfrak{s}$ is a supporting spin^c structure on Y , then we know from (6) that

$$P.D.c_1(\mathfrak{s}) \in [x] + [y] + (2g - 2)[s] + \ker(i_*) \subset H_1(Y; \mathbb{Q}),$$

where $i : Y \rightarrow W$ is the inclusion, y is a 1-cycle on R_+ , s is the curve $\{p\} \times S^1 \subset Y$, and x is a 1-cycle in M_l guaranteed by Lemma 3.10.

From Lemma 3.1, we know that

$$\ker(i_*) \subset [\text{im}(h_*^r - 1)] / [\text{im}(h_{2,*} - 1)].$$

By construction, we know that

$$\text{im}(h_*^r - 1) = \text{im}(f_*^r - 1) \subset H_1(M'; \mathbb{Q}).$$

To conclude the proof of Lemma 3.9, we need to show that

$$\text{im}(h_*^r - 1) = \text{im}(f_*^r - 1) \subset H_1(M_l; \mathbb{Q}),$$

and this is equivalent to show that, under the decomposition

$$H_1(M'; \mathbb{Q}) = H_1(M_l; \mathbb{Q}) \oplus \langle [\alpha_l] \rangle,$$

any element $[z] \in \text{im}(f_*^r - 1)$ can not have a non-zero $[\alpha_l]$ component.

To prove this final statement, suppose $[z] \in \text{im}(f_*^r - 1)$ is of the form

$$[z] = a \cdot [\alpha_l] + H_1(M_l; \mathbb{Q})$$

for some $a \in \mathbb{Q}$, then we need to show that $a = 0$. Note, from (4), we know that $[z] = 0 \in H_1(Y^r)$. Also, inside Y^r , $[z] \cdot [T_l^2] = a$, since, by construction, $[\alpha_l] \cdot [T_l^2] = 1$, and $T_l^2 \cap M_l = \emptyset$. Thus, we know that $a = 0$, and this concludes the proof of lemma 3.9. $\square$

4 General grading shifting formula

In this section, we prove the generalized grading shifting formula, Theorem 1.12, as stated in the introduction. We also use it to compute the sutured monopole and instanton Floer homologies of some particular sutured handle-bodies.

Proposition 4.1. *Suppose (M, γ) is a balanced sutured manifold, and $S \subset M$ is a properly embedded surface. Pick $i \in \mathbb{Z}$ so that the surface S^i , which is obtained from S by performing i times of stabilizations, is admissible (see Definition 2.25). Pick any $k \in \mathbb{Z}$. Then, there exist constants $l_M, l_I \in \mathbb{Z}$ so that, for any $j \in \mathbb{Z}$, we have:*

$$\underline{\text{SHM}}(-M, -\gamma, S^{i+2k}, j) = \underline{\text{SHM}}(-M, -\gamma, S^i, j - l_M),$$

and

$$\underline{\text{SHI}}(-M, -\gamma, S^{i+2k}, j) = \underline{\text{SHI}}(-M, -\gamma, S^i, j - l_I).$$

Moreover, if the sutured manifold decompositions of (M, γ) along S and $-S$ are both taut, then $l_M = l_I = k$.

Proof. We now prove the first part of the proposition, namely the existence of l and the fact that it is independent of j . As usual, we argue this in the monopole settings, but the same is true for the instanton settings. Suppose (M, γ) and S are defined the same as in the hypothesis of the proposition. We follow the idea in Subsection 2.3, to construct a closure (Y, R) of (M, γ) so that both S^i and S^{i+2k} extend to closed surfaces. To do this, put both S^i and S^{i+2k} in M so that they are transverse and $\partial S^i \cap \partial S^{i+2k} = \emptyset$ in Y . Note we can always achieve this. If ∂S intersects γ , then the positive and negative stabilizations can be performed in an arbitrarily small neighborhood of $S \cap \gamma$, so we can simply start with two parallel copies of S and perform i and $i + 2k$ stabilizations respectively. If $\partial S \cap \gamma = \emptyset$, then positive and negative stabilizations happen on different sides of S , and we can always perturb them to be distinct. See figure 6.


Now we pick a connected auxiliary surface T for (M, γ) , which is of large enough genus. For both S^i and S^{i+2k} , we can apply the construction of gradings as in Subsection 2.3. If the genus of T is chosen to be large enough, then we could arrange the arcs, which come from both ∂S^i and ∂S^{i+2k} , all represent linearly independent classes in $H_1(T, \partial T)$. We can then form the pre-closure $\widetilde{M} = M \cup [-1, 1] \times T$ and it has two boundary components $\partial \widetilde{M} = R_+ \cup R_-$. As in Subsection 2.3, we know that S^i and S^{i+2k} both extend to properly embedded surfaces $\widetilde{S}^i, \widetilde{S}^{i+2k} \subset \widetilde{M}$, and there are equal number of boundary components on R_+ and on R_- .

Thus, we can pick an orientation preserving diffeomorphism $h : R_+ \rightarrow R_-$ so that

$$h(\widetilde{S}^i \cap R_+) = \widetilde{S}^i \cap R_-, \text{ and } h(\widetilde{S}^{i+2k} \cap R_+) = \widetilde{S}^{i+2k} \cap R_-.$$

Using h and $\widetilde{M}$, we obtain a closure (Y, R_+) of $(-M, -\gamma)$ so that there are closed surfaces $\bar{S}^i \subset Y$ and $\bar{S}^{i+2k} \subset Y$. Pick a non-separating simple closed curve η which is disjoint from $\bar{S}^i \cup \bar{S}^{i+2k}$. Then, as in Subsection 2.3, we know that

$$\text{SHM}(-M, -\gamma, S^i, j) = \bigoplus_{\substack{\mathfrak{s} \in \mathfrak{G}^*(Y|_{R_+}) \\ c_1(\mathfrak{s})[\bar{S}^i] = 2j}} \widetilde{\text{HM}}_\bullet(Y, \mathfrak{s}; \Gamma_\eta),$$


 Figure 6: Pushing off S^{i+2k} .

and

$$SHM(-M, -\gamma, S^{i+2k}, j) = \bigoplus_{\substack{\mathfrak{s} \in \mathfrak{S}^*(Y|R_+) \\ c_1(\mathfrak{s})[\bar{S}^{i+2k}] = 2j}} \widetilde{HM}_\bullet(Y, \mathfrak{s}; \Gamma_\eta).$$

Now, suppose $\mathfrak{s}_1, \mathfrak{s}_2 \in \mathfrak{S}^*(Y|R_+)$ are two supporting spin^c structures, then, from Theorem 1.11, we know that there is a 1-cycle $x \subset M$ so that

$$[c_1(\mathfrak{s}_1) - c_1(\mathfrak{s}_2)] = P.D.[x] \in H^2(Y).$$

Since $x \subset M$ and S^i is isotopic to S^{i+2k} in M , we know that

$$[c_1(\mathfrak{s}_1) - c_1(\mathfrak{s}_2)][\bar{S}^i] = [c_1(\mathfrak{s}_1) - c_1(\mathfrak{s}_2)][\bar{S}^{i+2k}].$$

Thus, the number

$$l_M = \frac{1}{2}(c_1(\mathfrak{s}_1)[\bar{S}^i] - c_1(\mathfrak{s}_1)[\bar{S}^{i+2k}]) = \frac{1}{2}(c_1(\mathfrak{s}_2)[\bar{S}^i] - c_1(\mathfrak{s}_2)[\bar{S}^{i+2k}])$$

is the desired constant in the statement of the proposition.

When the decomposition of (M, γ) along S and $-S$ are both tight, then we can settle down the value of l by looking at the top or bottom non-vanishing grading and conclude $l = k$. This part of the proof is exactly the same as the proof of Proposition 4.3 in Li [19]. $\square$

Conjecture 4.2. *In general, we always have $l = k$.*

Corollary 4.3. *Suppose (M, γ) is a taut balanced sutured manifold and $D \subset M$ is a disk so that D intersects γ transversely four times. Suppose further that the decompositions of (M, γ) along D and $-D$ are (M', γ') and (M'', γ'') , respectively. If at least one of the two decompositions is taut, then*

$$\underline{SHM}(M, \gamma) \cong \underline{SHM}(M', \gamma') \oplus \underline{SHM}(M'', \gamma'').$$

Similar results holds for the instanton settings.

Proof. Without loss of generality, we can assume that (M', γ') is taut. We can perform a positive stabilization on D to make it admissible. Then, it induces a grading

$$\underline{\text{SHM}}(M, \gamma, D^+, i).$$

The adjunction inequality in Lemma 2.20 tells us that there are only three non-trivial gradings, being $i = -1, 0, 1$. Lemma 2.23 and Lemma 2.28 then imply that

$$\underline{\text{SHM}}(M, \gamma, D^+, 1) \cong \underline{\text{SHM}}(M', \gamma'),$$

$$\underline{\text{SHM}}(M, \gamma, D^+, -1) = 0.$$

Applying Proposition 4.1 and Lemma 2.28, we know that

$$\begin{aligned} \underline{\text{SHM}}(M, \gamma, D^+, 0) &= \underline{\text{SHM}}(M, \gamma, D^-, -1) \\ &= \underline{\text{SHM}}(M, \gamma, (-D)^+, 1) \\ &\cong \underline{\text{SHM}}(M'', \gamma'') \end{aligned}$$

Hence, we are done. $\square$

This gives an affirmative answer to Conjecture 4.3 in Li [17].

Corollary 4.4. *Suppose V is a solid torus, and γ^4 consists of four longitudes. When using $\mathbb{Z}$ coefficients, we have*

$$\text{SHM}(V, \gamma^4) \cong \mathbb{Z}^2.$$

With the help of Proposition 4.1, we can prove the general grading shifting property, Theorem 1.12, as stated in the introduction.

Proof of Theorem 1.12. Assume that S_1 and S_2 are transverse to each other. We need to isotope S_1 and S_2 into $S_1^{i_1, i'_1}$ and $S_2^{i_2, i'_2}$, respectively. Here, $i_1, i_2 \geq 0$ indicate the number of positive stabilizations on S_1 and S_2 , respectively. Similarly, $i'_1, i'_2 \leq 0$ correspond to the negative stabilizations. We require the following six conditions to hold.

(1) Both $S_1^{i_1, i'_1}$ and $S_2^{i_2, i'_2}$ are admissible, and no more intersection points are created during the stabilizations.

(2) Any positive intersection of $S_1^{i_1, i'_1}$ with $S_2^{i_2, i'_2}$ is contained in $R_+(\gamma)$, and any negative intersection is contained in $R_-(\gamma)$.

(3) If θ_1 is a component of $\partial S_1^{i_1, i'_1}$ and $\theta_1 \cap S_2^{i_2, i'_2} \neq \emptyset$, then $\theta_1 \cap \gamma \neq \emptyset$.

(4) If θ_2 is a component of $\partial S_2^{i_2, i'_2}$ and $\theta_2 \cap S_1^{i_1, i'_1} \neq \emptyset$, then $\theta_2 \cap \gamma \neq \emptyset$.

(5). If δ_1 is a component of $\partial S_1^{i_1, i'_1} \cap R(\gamma)$, then δ intersects $S_2^{i_2, i'_2}$ at most once.

(6). If δ_2 is a component of $\partial S_2^{i_2, i'_2} \cap R(\gamma)$, then δ intersects $S_1^{i_1, i'_1}$ at most once.

Pick a connected auxiliary surface T of large enough genus, and form the pre-closure $\widetilde{M} = M \cup [-1, 1] \times T$. It has two boundary components:

$$\partial \widetilde{M} = R_+ \cup R_-.$$

We can carry out the construction of gradings in Subsection 2.3 on both $S_1^{i_1, i'_1}$ and $S_2^{i_2, i'_2}$. Suppose

$$n_1 = \frac{1}{2}|S_1^{i_1, i'_1} \cap \gamma|, \text{ and } n_2 = \frac{1}{2}|S_2^{i_2, i'_2} \cap \gamma|.$$

Pick two balanced pairings $\mathcal{P}_1$ and $\mathcal{P}_2$, as introduced in Subsection 2.3, for the two surfaces $S_1^{i_1, i'_1}$ and $S_2^{i_2, i'_2}$, respectively. Inside T , pick a set of disjoint properly embedded arcs $\{\alpha_1, \dots, \alpha_{n_1}, \beta_1, \dots, \beta_{n_2}\}$ so that the end points of α_k are identified with the intersection points of $S_1^{i_1, i'_1} \cap \gamma$, according to the balanced pairing $\mathcal{P}_1$, and the end points of β_j are identified with the intersection points of $S_2^{i_2, i'_2} \cap \gamma$, according to the balanced pairing $\mathcal{P}_2$. There is one special requirement for $\mathcal{P}_1$:

(7) If δ_1 is a component of $\partial S_1^{i_1, i'_1} \cap R(\gamma)$, which intersects $S_2^{i_2, i'_2}$ non-trivially, then there is an arc α_{k_0} so that $\partial \alpha_{k_0}$ is identified with $\partial \delta_1$.

Strictly speaking, $\partial \delta \subset \partial R(\gamma)$, but we could regard $\partial \delta$ to be on γ , since $\partial R(\gamma)$ is parallel to γ . Similarly, we require the following for $\mathcal{P}_2$.

(8) If δ_2 is a component of $\partial S_2^{i_2, i'_2} \cap R(\gamma)$, which intersects $S_1^{i_1, i'_1}$ non-trivially, then there is an arc β_{j_0} so that $\partial \beta_{j_0}$ is identified with $\partial \delta_2$.

Note when we perform enough positive and negative stabilizations, the balanced pairing satisfying the constraints (7) and (8) always exist. When the genus of T is large enough, we can choose the arcs, $\alpha_1, \dots, \alpha_{n_1}, \beta_1, \dots, \beta_{n_2}$, to represent linearly independent classes in $H_1(T, \partial T)$. Then, inside $\widetilde{M}$, $S_1^{i_1, i'_1}$ and $S_2^{i_2, i'_2}$ extend to properly embedded surfaces $\widetilde{S}_1^{i_1, i'_1}$ and $\widetilde{S}_2^{i_2, i'_2}$, respectively. From the construction, we know that $\partial \widetilde{S}_1^{i_1, i'_1} \cap R_+$ and $\partial \widetilde{S}_1^{i_1, i'_1} \cap R_-$ have equal number of boundary components. Thus, let

$$\partial \widetilde{S}_1^{i_1, i'_1} \cap R_+ = C_{+,1} \cup \dots \cup C_{+,s}, \text{ and } \partial \widetilde{S}_1^{i_1, i'_1} \cap R_- = C_{-,1} \cup \dots \cup C_{-,s}.$$

Similarly, we can assume

$$\partial \widetilde{S}_2^{i_2, i'_2} \cap R_+ = D_{+,1} \cup \dots \cup D_{+,t}, \text{ and } \partial \widetilde{S}_2^{i_2, i'_2} \cap R_- = D_{-,1} \cup \dots \cup D_{-,t}.$$

Note the intersection points of $\widetilde{S}_1^{i_1, i'_1}$ and $\widetilde{S}_2^{i_2, i'_2}$ are in one-to-one-correspondence to the intersection points of S_1 and S_2 by requirement (1). We claim that $\partial S_1 \cap \partial S_2$ consists of an even number of positive and negative points. Indeed, $\partial S_i \cap \partial S_j = \partial S_i \cap S_j$, and it is clear that the algebraic intersection number of ∂S_i and S_j is zero. Hence, on $R_\pm$, we have a collection of circles $C_{\pm,1}, \dots, C_{\pm,s}, D_{\pm,1}, \dots, D_{\pm,t}$. They represent linearly independent classes in $H_1(R_\pm)$. There might be intersections between $C_{\pm,k}$ with $D_{\pm,j}$, but, by requirement (5), (6), (7), and (8), each $C_{\pm,k}$ intersects with at most one $D_{\pm,j}$, and each $D_{\pm,j}$ intersects with at most one $C_{\pm,k}$. Hence, the pattern of $C_{+,1}, \dots, C_{+,s}, D_{+,1}, \dots, D_{+,t}$ on R_+ is exactly the same as the pattern of $C_{-,1}, \dots, C_{-,s}, D_{-,1}, \dots, D_{-,t}$ on R_- . As a result, there exists an orientation preserving diffeomorphism $h : R_+ \rightarrow R_-$ so that

$$h(\partial \widetilde{S}_1^{i_1, i'_1} \cap R_+) = \partial \widetilde{S}_1^{i_1, i'_1} \cap R_-, \text{ and } h(\partial \widetilde{S}_2^{i_2, i'_2} \cap R_+) = \partial \widetilde{S}_2^{i_2, i'_2} \cap R_-.$$

Hence, we can obtain a closure (Y, R_+) of (M, γ) from $\widetilde{M}$ and h . Inside Y there are two closed surfaces $\bar{S}_1^{i_1, i'_1}$ and $\bar{S}_2^{i_2, i'_2}$, and they induce gradings on $\text{SHM}(M, \gamma)$ that

are associated to $S_1^{i_1, i'_1}$ and $S_2^{i_2, i'_2}$, respectively. The rest of the proof is then exactly the same as the proof of Proposition 4.1. $\square$

Next, we want to compute the sutured monopole and instanton Floer homologies of the following balanced sutured manifold (M, γ) , as depicted in Figure 7. (Strictly speaking, they are the sutured monopole and instanton Floer homologies of $(-M, -\gamma)$.)




Figure 7: The sutured manifold (M, γ) . The three curves (red, blue and green) are the sutures. The disk D is used to construct a grading.

The idea is to apply the by-pass exact triangles repeatedly. There is a graded version of by-pass exact triangles, as in Li [19], generalizing the by-pass exact triangle introduced in Subsection 2.4.

$$\begin{array}{ccc} \underline{\text{SHM}}(-M, -\gamma, D, i) & \longrightarrow & \underline{\text{SHM}}(-M, -\gamma_1, D_1^{-2}, i) \\ & \searrow & \downarrow \\ & & \underline{\text{SHM}}(-M, -\gamma_2, D_2^+, i) \end{array} \quad (7)$$

Here, the surfaces D is chosen as in Figure 7 so that it has six transverse intersection with the suture γ . The surfaces $D_1, D_2 \subset M$ are isotopic to D , but having minimal possible transverse intersection with γ_1 (two intersections) and γ_2 (four intersections) respectively. The sup-scripts in D_2^+ and D_1^{-2} imply the number of positive or negative stabilizations performed on the surfaces $D_1 \subset M$ and $D_2 \subset M$, as introduced in

Subsection 2.3. A direct check shows that (M, γ_1) is a product sutured manifold and the suture γ_2 is depicted as in Figure 8.




Figure 8: The sutured manifold (M, γ_2) . The three curves (red, blue and green) are the sutures.

Now we know that

$$\underline{\text{SHM}}(-M, -\gamma_1) = \underline{\text{SHM}}(-M, -\gamma_1, D_1, 0) = \mathcal{R}.$$

As a result, by Lemma 2.23 and Lemma 2.28, we know that

$$\underline{\text{SHM}}(-M, -\gamma_1, D_1^{-2}, i) = \begin{cases} \mathcal{R} & i = 1 \\ 0 & \text{others} \end{cases} \quad (8)$$

From the construction of gradings in Subsection 2.3 and the adjunction inequality in Lemma 2.20, we know that

$$\underline{\text{SHM}}(-M, -\gamma_2, D_2^+, i) = 0$$

for $i > 1$ or $i < -1$. From Lemma 2.23 and Lemma 2.28, we know that

$$\underline{\text{SHM}}(-M, -\gamma_2, D_2^+, 1) = 0.$$

Hence, from the graded exact triangle in (7), we conclude that

$$\underline{\text{SHM}}(-M, -\gamma) = \mathcal{R} \oplus \underline{\text{SHM}}(-M, -\gamma_2). \quad (9)$$

To compute the sutured monopole Floer homology of (M, γ_2) , we perform the same trick once more and conclude that

$$\underline{\text{SHM}}(-M, -\gamma_2) = \mathcal{R} \oplus \underline{\text{SHM}}(-M, -\gamma_3), \quad (10)$$

where (M, γ_3) is the balanced sutured manifold as depicted in Figure 9.




Figure 9: The sutured manifold (M, γ_3) . The three curves (red, blue and green) are the sutures.

To compute the sutured monopole Floer homology of (M, γ_3) , we could perform a sutured manifold decomposition along the disk D as depicted in Figure 9. Suppose (M_4, γ_4) is the resulting balanced sutured manifold, then, from Kronheimer and Mrowka [16], we know that

$$\underline{\text{SHM}}(-M, -\gamma_3) \cong \underline{\text{SHM}}(-M_4, -\gamma_4).$$

Furthermore, the balanced sutured manifold (M_4, γ_4) is a solid torus equipped with two curves of slope $\frac{1}{3}$ as the suture, as depicted in Figure 10, so from Li [19], we know that

$$\underline{\text{SHM}}(-M, -\gamma_3) \cong \underline{\text{SHM}}(-M_4, -\gamma_4) \cong \mathcal{R}^3.$$

Finally, we conclude that

Proposition 4.5. $\underline{\text{SHM}}(-M, -\gamma) \cong \mathcal{R}^5$.

The same type of arguments in instanton theory yields the following.

Corollary 4.6. $\underline{\text{SHI}}(-M, -\gamma) \cong \mathbb{C}^5$.




Figure 10: The sutured manifold (M, γ_4) . The three curves (red, blue and green) are the sutures.

5 Polytopes

5.1 Constructing the polytope

Suppose (M, γ) is a balanced sutured manifold, and $\alpha \in H_2(M, \partial M)$ is a second relative homology class. From Theorem 1.12, there is a grading on $\underline{\text{SHM}}(M, \gamma)$ or $\underline{\text{SHI}}(M, \gamma)$ associated to α , which is well-defined up to an overall grading shifting. Hence, we can define the following.

Definition 5.1. An element a in $\underline{\text{SHM}}(M, \gamma)$ or $\underline{\text{SHI}}(M, \gamma)$ is called *homogenous* if for any homology class $\alpha \in H_2(M, \partial M)$, the element a is homogenous with respect to the grading associated to α .

Lemma 5.2. For any taut balanced manifold (M, γ) , there exist non-zero homogenous elements in $\underline{\text{SHM}}(M, \gamma)$ and $\underline{\text{SHI}}(M, \gamma)$.

Proof. Pick a basis $\alpha_1, \dots, \alpha_n$ for $H_2(M, \partial M)$. Pick admissible properly embedded surfaces $S_1, \dots, S_n$ to represent $\alpha_1, \dots, \alpha_n$, respectively. We can carry out the proof of Theorem 1.12 again. This time we have n many surfaces instead of just two, but the construction is exactly the same. As a result, we obtain a special closure (Y, R) of the balanced sutured manifold (M, γ) so that, inside Y , the surfaces $S_1, \dots, S_n$ extends to closed surfaces $\tilde{S}_1, \dots, \tilde{S}_n$, respectively, and the surfaces $\tilde{S}_1, \dots, \tilde{S}_n$ are the ones used to define gradings on $\underline{\text{SHM}}(M, \gamma)$ associated to $S_1, \dots, S_n$, respectively. Also, pick some suitable non-separating simple closed curves η on R to support local

coefficients. On Y , the surfaces $\bar{S}_1, \dots, \bar{S}_n$ induces a $\mathbb{Z}^r$ -grading on $HM(Y|R; \Gamma_\eta)$ by looking at the evaluation of the first Chern classes of spin^c structures on those closed surfaces. Suppose $\mathfrak{s}_0$ is a supporting spin^c structures (see Definition 2.14) on Y , and $a \in \widetilde{HM}_\bullet(Y, \mathfrak{s}_0; \Gamma_\eta)$ is a non-zero element, then we know that the element a is homogenous with respect to all the gradings induced by $\bar{S}_1, \dots, \bar{S}_n$. We claim that it is a homogenous element as defined in Definition 5.1.

To prove the claim, suppose $\alpha \in H_2(M, \gamma)$ is any homology class. Since $\alpha_1, \dots, \alpha_n$ form a basis of $H_2(M, \gamma)$, α is a linear combination of $\alpha_1, \dots, \alpha_n$. Thus, we can perform a sequence of double curve surgeries (for definition, see Scharlemann [24]) on $S_1, \dots, S_n$ to obtain a properly embedded surface S that represents the class $\alpha \in H_2(M, \gamma)$. Correspondingly, we can perform the same set of double curve surgeries on $\bar{S}_1, \dots, \bar{S}_n$ to obtain a closed surface $\bar{S} \subset Y$, which extends S and which induces the grading associated to S . Then, we know that the element a is a homogenous element with respect to the grading associated to S , and this concludes the proof of lemma 5.2. $\square$

Lemma 5.3. *Suppose (M, γ) is a balanced sutured manifold, and $a, b \in \underline{\text{SHM}}(M, \gamma)$ are two homogenous elements. Then, there is a well defined element $\rho(a, b) \in H^2(M, \partial M; \mathbb{Q})$ associated to the (ordered) pair (a, b) .*

The same result holds in the instanton setups.

Proof. We first construct the map

$$\rho(a, b) : H_2(M, \partial M) \rightarrow \mathbb{Z}.$$

For any class $\alpha \in H_2(M, \partial M)$, we pick a surface S that represents the class α and is admissible. Define

$$\rho(a, b)(\alpha) = \text{difference between } a \text{ and } b \text{ under the grading associated to } S.$$

This is well defined by Theorem 1.12. This map is linear by essentially the same type of argument as in the proof of Lemma 5.2. Then, we can regard $\rho(a, b)$ as an element in $H^2(M, \partial M; \mathbb{Q})$. $\square$

Definition 5.4. Suppose (M, γ) is a balanced sutured manifold, and $a \in \underline{\text{SHM}}(M, \gamma)$ is a homogenous element. For an element $\rho \in H^2(M, \partial M; \mathbb{R})$, define

$$\underline{\text{SHM}}_a(M, \gamma, \rho) = \{b \in \underline{\text{SHM}}(M, \gamma), \rho(a, b) = \rho \in H^2(M, \partial M; \mathbb{R})\},$$

and

$$\underline{\text{SHI}}_a(M, \gamma, \rho) = \{b \in \underline{\text{SHI}}(M, \gamma), \rho(a, b) = \rho \in H^2(M, \partial M; \mathbb{R})\}.$$

Let

$$SM_a(M, \gamma) = \{\rho \in H^2(M, \partial M; \mathbb{R}), \underline{\text{SHM}}_a(M, \gamma, \rho) \neq \emptyset\}$$

and

$$SI_a(M, \gamma) = \{\rho \in H^2(M, \partial M; \mathbb{R}), \underline{\text{SHI}}_a(M, \gamma, \rho) \neq \emptyset\}.$$

Define the *polytopes* $PM_a(M, \gamma)$ and $PI_a(M, \gamma)$ to be the convex hull of $SM_a(M, \gamma)$ and $SI_a(M, \gamma)$, respectively.

Lemma 5.5. *Suppose a and b are two homogenous elements in $\underline{SHM}(M, \gamma)$, then the polytopes $PM_a(M, \gamma) \subset H^2(M, \partial M; \mathbb{R})$ is a translate of $PM_b(M, \gamma) \subset H^2(M, \partial M; \mathbb{R})$. The same result holds in the instanton setups.*

Proof. It is straightforward from the construction. $\square$

5.2 Dimension formula

Lemma 5.6. *Suppose (M, γ) is a taut balanced sutured manifold with $H_2(M) = 0$, and A is an incompressible product annulus. Then, we can pick an orientation of A so that the sutured manifold decomposition of (M, γ) along the oriented A yields a taut balanced sutured manifold (M', γ') , that $SHM(M', \gamma')$ is a direct summand of $SHM(M, \gamma)$, and that $SHI(M', \gamma')$ is a direct summand of $SHI(M, \gamma)$.*

Proof. Since A is incompressible, we know that $\partial A \subset R(\gamma)$ does not bound a disk. Note that Lemma 4.2 in Scharlemann [25] makes sure that no matter which orientation of A we choose, the balanced sutured manifold after the decomposition is taut. There are three cases:

Case 1. Both components of ∂A are homologically essential on $R(\gamma)$.

Case 2. Both components of ∂A are homologically trivial on $R(\gamma)$. Then, there are $V_+ \subset R_+(\gamma)$ and $V_- \subset R_-(\gamma)$ so that $\partial V_+ \cup \partial V_- = \partial A$ (as unoriented curves). Thus, we have a closed surface $V_+ \cup A \cup V_-$. The fact that $H_2(M) = 0$ implies that this closed surface is separating, or equivalently, A separates M into two parts, of which one has boundary $V_+ \cup A \cup V_-$. Thus, this part is disjoint from γ .

For the above two cases, the lemma follows from Proposition 6.7 of Kronheimer and Mrowka [16].

Case 3. One component of ∂A is homologically essential, and the other is inessential. Then we can choose a suitable orientation of A to make ∂A being boundary coherent, in the sense of Kronheimer and Mrowka [16], so that Proposition 6.9 in that paper applies, and, thus, we conclude the proof of Lemma 5.6. $\square$

Lemma 5.7. *Suppose (M, γ) is a taut balanced sutured manifold and $S \subset M$ is a properly embedded decomposing surface. Suppose $p, q \in S \cap \gamma$ are two points of different signs. Then, we can attach a product 1-handle, in the sense of Definition 3.2, to obtain a new taut balanced sutured manifold (M_1, γ_1) and a new properly embedded surface $S_1 \subset M_1$ so that the decomposition*

$$(M, \gamma) \xrightarrow{S} (M', \gamma')$$

is taut if and only if the decomposition

$$(M_1, \gamma_1) \xrightarrow{S_1} (M'_1, \gamma'_1)$$

is taut. Furthermore, there is a commutative diagram

$$\begin{array}{ccc} SHM(M', \gamma') & \hookrightarrow & SHM(M, \gamma) \\ \downarrow = & & \downarrow = \\ SHM(M'_1, \gamma'_1) & \hookrightarrow & SHM(M_1, \gamma_1). \end{array}$$

A similar statement holds in the instanton settings.

Proof. This is how Kronheimer and Mrowka proved Proposition 6.9 in [16]. S_1 is obtained from S by attaching a 2-dimensional 1-handle inside the 3-dimensional product 1-handle. See Figure 11.




Figure 11: Adding a product 1-handle.

□

Proposition 5.8. *Suppose (M, γ) is a balanced sutured manifold with $H_2(M) = 0$ and is taut, reduced, horizontally prime, and free of non-separating essential product disks. Suppose $\alpha \in H_2(M, \partial M)$ is a non-zero class. Then, we can find properly embedded surfaces S and S' in M so that*

- (1) $[S] = -[S'] = \alpha \in H_2(M, \partial M)$.
- (2) *The sutured manifold decompositions*

$$(M, \gamma) \xrightarrow{S} (M', \gamma') \text{ and } (M, \gamma) \xrightarrow{S'} (M'', \gamma'')$$

are both taut.

- (3) $SHM(M', \gamma')$ and $SHM(M'', \gamma'')$ are direct summands of $SHM(M, \gamma)$, (by Proposition 6.9 of Kronheimer and Mrowka [16]) and

$$SHM(M', \gamma') \cap SHM(M'', \gamma'') = 0$$

in $SHM(M, \gamma)$.

- (4) *The same result holds for SHI .*

Proof. From Lemma 0.7 in Gabai [6], we can pick an S so that the following holds

- (i) S represents the class $\alpha \in H_2(M, \partial M)$.
- (ii) For any component V of $R(\gamma)$, if $S \cap V$ has a closed component, then $S \cap V$ consists of parallel, parallel oriented non-separating simple closed curves.
- (iii) For any component δ of $\partial R(\gamma)$, all the intersection points of S with δ are of the same sign.

- (iv) S intersects $A(\gamma)$ in parallel and coherently oriented essential arcs.
- (v) The sutured manifold decomposition

$$(M, \gamma) \xrightarrow{S} (M', \gamma')$$

is taut.

To find S' , we proceed as follows, according to Gabai [5], Scharlemann [25] or Juhasz [11]. Pick $U = N(\partial S) \cup A(\gamma) \subset \partial M$. Pick a large enough $k \in \mathbb{Z}_+$ so that

$$x(-\alpha + (k+1)[R(\gamma)]) = x(-\alpha + k[R(\gamma)]) + x(R(\gamma)),$$

where $x(\cdot)$ is the Thurston norm for classes in $H_2(M, U)$. Pick a norm-minimizing embedded surface which represents the class $-\alpha + k[R(\gamma)]$, and disregard all components of it, which represent the zero homology class, then the remaining surface S' is the desired one. Note

$$[\partial S'] = -[\partial S] + 2k \cdot [\gamma] \in H_1(U).$$

Here, it is $2k$ rather than k , because each copy of $R(\gamma)$ contributes $2[\gamma]$. We can arrange so that $\partial S'$ is obtained from $-\partial S$ and $2k$ copies of γ by an oriented smoothing as in Figure 12. From construction of S' , we know that the sutured manifold decomposition

$$(M, \gamma) \xrightarrow{S'} (M'', \gamma'')$$

is taut.




Figure 12: The smoothing inside $A(\gamma)$.

From the above construction of S' , we know that $S \cap \gamma = S' \cap \gamma$. So, assume that

$$n = \frac{1}{2}|S \cap \gamma| = \frac{1}{2}|S' \cap \gamma|.$$

Also, assume that $\partial S \cap R(\gamma)$ (and thus $\partial S' \cap R(\gamma)$) has m closed components. Write them as

$$B_1 \cup \dots \cup B_m \subset \partial S \cap R(\gamma),$$

and orient B_i by the boundary orientation of S . Write S_0 the surface obtained from S by performing one negative stabilization on each B_i , and let $S^\dagger$ be the surface obtained by performing one positive stabilization on each B_i . By Lemma 4.5 in Juhasz [10], both S_0 and $S^\dagger$ exist. We also want to modify S' correspondingly. Note part of $\partial S'$ is coming from $-\partial S$, so positive stabilizations near this part of $\partial S'$ corresponds to negative stabilizations on S . Thus, we perform positive stabilizations on S' , in correspondence to the negative stabilizations performed on S . Since $\partial S'$ contains many copies of $\partial R(\gamma) \subset A(\gamma)$, when performing the positive stabilizations to create intersections of B_i with γ , the isotopy also creates self intersections of S' . We then perform double curve surgeries, in the sense of Scharlemann [25], to resolve all self intersections created by the positive stabilizations, and let S'_0 be the resulting surface. See Figure 13. On $A(\gamma)$, this double curve surgery behaves exactly in the same way as the oriented smoothing that is depicted in Figure 12. It is important that we choose positive stabilizations to perform on S' , so the decomposition of (M, γ) along S'_0 gives the same result as decomposing along S' :

$$(M, \gamma) \xrightarrow{S'_0} (M'', \gamma'').$$




Figure 13: A positive stabilization on S' followed by a double curve surgery.

We can attach $m + n$ copies of product 1-handle, as in Lemma 5.7, along the intersections of S_0 with γ . We require that the pair of intersection points created by a negative stabilization on S are paired together by a product 1-handle. Let (M_1, γ_1) be the resulting balanced sutured manifold. The surface S_0 extends to a properly embedded surface $S_1 \subset M_1$ as in Lemma 5.7.

The surface S'_0 can also extend, though in a slightly complicated way. As in Figure 14, in each product 1-handle, there is one (vertical) 2-dimensional 1-handle to be glued to the part of $\partial S'_0$ that comes from $-\partial S$, and $2k$ copies of (horizontal) 2-dimensional 1-handles to be glued to the part of $\partial S'_0$ that corresponds to the k copies of $\partial R(\gamma)$. We can perform a double curve surgery on those two collections of 2-dimensional 1-handles, as in Figure 14, and then glue the resulting surface to S'_0 , when gluing the product 1-handle to (M, γ) . In this way, S'_0 extends to a properly embedded surface S'_1 in (M_1, γ_1) . Let (M'_1, γ'_1) and (M''_1, γ''_1) be obtained from (M_1, γ_1) by decomposing along S_1 and S'_1 , respectively. (M''_1, γ''_1) is taut by Lemma 5.7, since it is obtained

from (M'', γ'') by attaching a few product 1-handles, as indicated in Figure 14. After decomposing along S'_1 , each product 1-handle is decomposed into $(2n + 1)$ many product 1-handles that are attached to (M'', γ'') . In that figure, $n = 3$, and the six arcs in the right-subfigure divides the original product 1-handle into 7 parts.




Figure 14: Double curve surgery on a cross section of the product 1-handle. On the right: the 2-dimensional 1-handles after the double curve surgery will cut the original product 1-handle into $2k + 2$ small ones.

From the construction, we also know that

$$S'_1 \cap R_{\pm}(\gamma_1) = -S_1 \cap R_{\pm}(\gamma_1),$$

and $S'_1 \cap A(\gamma_1)$ consists of $2k$ parallel copies of γ_1 . Now let

$$S_1 \cap R_+(\gamma_1) = B_{+,1} \cup \dots \cup B_{+,m} \cup C_{+,1} \cup \dots \cup C_{+,s}$$

and

$$S_1 \cap R_-(\gamma_1) = B_{-,1} \cup \dots \cup B_{-,m} \cup C_{-,1} \cup \dots \cup C_{-,t}$$

Here, $B_{\pm,i}$ are the boundary components of S_1 that comes from attaching a product 1-handle along the pair of intersection points created by a negative stabilization on S near B_i . Note s and t are not necessarily equal. Without loss of generality, we could assume that $s \geq t$.

Pick (V, δ) be a balanced sutured manifold where $V = S^1 \times D^2$ is a solid torus and δ consists of two longitudes. Let $D \subset V$ be a standard meridian disk in V , which has two intersections with the suture δ . Let D_0 be the surface obtained from D by performing a negative stabilization, as shown in Figure 15. It has four intersections with the suture δ . Attach two product 1-handle along two pairs of points (p_1, p_4) and (p_2, p_3) , as labeled in the figure, and let the resulting balanced sutured manifold be (V_1, γ_1) . D_0 extends to a properly embedded surface D_1 as in Lemma 5.7. Note $D_1 \cap R_+(\delta_1)$ has one components and $D_1 \cap R_-(\delta_1)$ has two components. Now we want to construct another surface D'_1 inside V_1 . First, let D' be the result of a double

curve surgery of $-D_0$ with k copies of $R(\delta)$. It is crucial that our choice of D_0 makes the sutured manifold decomposition

$$(V, \delta) \xrightarrow{D'} (V'', \gamma'')$$

taut. We can check directly that V'' is a 3-ball and γ'' is a simple closed curve on $\partial V''$, and, hence, it is a product sutured manifold. Next, when attaching two product 1-handles to (V, δ) , we repeat the procedure explained above, with which we construct S'_1 out of S'_0 , and thus construct a properly embedded surface D'_1 inside V_1 out of D'_0 . It is straightforward to check that the decomposition of (V_1, γ_1) along D'_1 yields a product sutured manifold (V''_1, γ''_1) .




Figure 15: The balanced sutured manifold (V, γ) together with the surfaces D_0 and $D^\dagger$.

So, we could form the disjoint union

$$(M_2, \gamma_2) = (M_1, \gamma_1) \sqcup (s-t)(V_1, \delta_1), \quad S_2 = S_1 \cup (s-t)D_1, \quad S'_2 = S'_1 \cup (s-t)D'_1.$$

Since the decomposition of (M_1, γ_1) along S'_1 is taut (as we have explained) and the decomposition of (V_1, γ_1) along D'_1 is also taut (by a direct check), we know that the decomposition of (M_2, γ_2) along S'_2 is taut. However, the decomposition along S_2 is not, as we will explain later.

Pick a connected auxiliary surface T for (M_2, γ_2) and form a pre-closure

$$\widetilde{M} = M_2 \cup [-1, 1] \times T, \quad \partial \widetilde{M} = R_+ \cup R_-, \quad R_\pm = R_\pm(\gamma_2) \cup \{\pm 1\} \times T.$$

Since (M_1, γ_1) is obtained from (M, γ) by attaching product 1-handles, and (V_1, δ_1) is itself a product sutured manifold, we know that $\widetilde{M}$ is also a pre-closure of (M, γ) . By construction, we know that

$$S_2 \cap R_\pm = -S'_2 \cap R_\pm,$$

and there are same number of components of $S_2 \cap R_+$ and $S_2 \cap R_-$. Moreover, since T is connected, the components of $S_2 \cap R_\pm$ represent linearly independent classes in $H_1(R_\pm)$. Thus, we could find an orientation preserving diffeomorphism $h : R_+ \rightarrow R_-$ so that

$$h(S_2 \cap R_+) = S_2 \cap R_-,$$

and we can use $\widetilde{M}$ as well as h to construct a closure (Y, R_+) for both (M, γ) and (M_2, γ_2) . Inside Y , S_2 becomes a closed surface $\bar{S}$.

Though there is no sutures on the boundary of $\widetilde{M}$, the theory of balanced sutured manifolds in Kronheimer and Mrowka [16] extends to $\widetilde{M}$ effectively. In particular, we could define

$$SHM(\widetilde{M}) = HM(Y|R_+) = SHM(M, \gamma) = SHM(M_2, \gamma_2).$$

The surface S'_2 extends to a surface $\widetilde{S}'_2 \subset \widetilde{M}$ as the union of S'_2 with $2k$ copies of T . Suppose $(\widetilde{M}'', \widetilde{\gamma}'')$ is the result of the sutured manifold decomposition of $\widetilde{M}$ along $\widetilde{S}'_2$, then $(\widetilde{M}'', \widetilde{\gamma}'')$ can be obtained from (M''_2, γ''_2) by attaching $2k+1$ copies of the product region $T \times [-1, 1]$. Recall that (M''_2, γ''_2) is obtained from (M_2, γ_2) by decomposing along S'_2 and thus is the disjoint union of (M''_1, γ''_1) with $(s-t)$ copies of product sutured manifolds (V''_1, γ''_1) . Furthermore, (M''_1, γ''_1) is obtained from (M'', γ'') by attaching a few product 1-handles, so we finally conclude that

$$SHM(\widetilde{M}'', \widetilde{\gamma}'') \cong SHM(M'', \gamma''),$$

since attaching product regions (or product 1-handles), disjoint union with product manifolds will never change the sutured monopole Floer homology.

Back to the point that $(\widetilde{M}'', \widetilde{\gamma}'')$ is the decomposition of $\widetilde{M}$ along $\widetilde{S}'_2$. The decomposition theorem, Proposition 6.9 in Kronheimer and Mrowka [16], continues to hold in this case, and we conclude that $SHM(\widetilde{M}'', \widetilde{\gamma}'')$ is a direct summand of $SHM(\widetilde{M})$. More precisely, $\widetilde{S}'_2$ becomes a closed surface $\bar{S}' \subset Y$, and we have

$$SHM(M'', \gamma'') = SHM(\widetilde{M}'', \widetilde{\gamma}'') = \bigoplus_{\substack{s \in \mathfrak{S}^*(Y|R_+) \\ c_1(s)[\bar{S}'] = 2g(\bar{S}') - 2}} \widetilde{HM}_\bullet(Y, \mathfrak{s}). \quad (11)$$

We also want to identify the summand $SHM(M', \gamma')$ inside $SHM(\widetilde{M}) = SHM(M, \gamma)$. We cannot proceed directly as we did for $SHM(M'', \gamma'')$, since the decomposition of $\widetilde{M}$ along S_2 is not taut. This is because the decomposition of (M, γ) along S_0 and (V, γ) along D_0 are both not taut, since, at the beginning, we picked S_0 and D_0 by performing negative stabilizations (see Lemma 2.23). Let $S^\dagger$ and $D^\dagger$ be obtained from S and D performing positive stabilizations instead of negative ones. We can repeat the whole construction again with S_0 and D_0 replaced by $S^\dagger$ and $D^\dagger$, respectively. Attach $m+n$ product 1-handle along the intersection points of $\partial S^\dagger$ with γ , and let $(M^\dagger_1, \gamma^\dagger_1)$ be the result. There is a properly embedded surface $S^\dagger_1 \subset M^\dagger_1$. Similarly, attach two product 1-handles to (V, γ) , and let $(V^\dagger_1, \gamma^\dagger_1)$ be the result. Then, there is a properly embedded surface $D^\dagger_1 \subset V^\dagger_1$. We can form the disjoint union

$$(M^\dagger_2, \gamma^\dagger_2) = (M^\dagger_1, \gamma^\dagger_1) \sqcup (s-t)(V^\dagger_1, \gamma^\dagger_1), \quad S^\dagger_2 = S^\dagger_1 \cup (s-t)D^\dagger_1.$$

Pick an auxiliary surface $T^\dagger$ that has the same genus as T and form a pre-closure $\widetilde{M}^\dagger$. We can pick a suitable gluing diffeomorphism $h^\dagger$ and obtain a closure $(Y^\dagger, R^\dagger_+)$. Inside $Y^\dagger$, the surface $S^\dagger_2$ becomes a closed surface $\bar{S}^\dagger$. The decomposition of $\widetilde{M}^\dagger$ along

$S_2^\dagger$ is taut. The argument that concludes (11) applies again, and we have

$$SHM(M', \gamma') = \bigoplus_{\substack{\mathfrak{s} \in \mathfrak{S}^*(Y^\dagger|_{R_+^\dagger}) \\ c_1(\mathfrak{s})[\bar{S}^\dagger] = 2g(\bar{S}^\dagger) - 2}} \widetilde{HM}_\bullet(Y^\dagger, \mathfrak{s}). \quad (12)$$

Since $g(T^\dagger) = g(T)$, we know that $g(R_+^\dagger) = g(R_+)$. Thus, as in Subsection 2.2, there is an excision cobordism W , from $Y^\dagger \cup Y_T$ to Y , which induces an isomorphism

$$\Phi : HM(Y^\dagger|_{R_+^\dagger}) \rightarrow HM(Y|_{R_+}).$$

Here, Y_T is a mapping torus of a diffeomorphism on $R_+^\dagger$, arising from $h^\dagger$, h , and a suitable identification $R_+^\dagger = R_+$. The difference between the surfaces $\bar{S}^\dagger$ and $\bar{S}$ originates from whether we performed positive or negative stabilizations. So, the proof of Proposition 4.1 in Li [19] applies to the present context, and we know that inside W ,

$$[\bar{S}^\dagger] = [\bar{S}] + [\Sigma_1] + \dots [\Sigma_m] + [\Sigma'_1] + \dots + [\Sigma'_{s-t}].$$

Here, each Σ_i or Σ'_j is a connected closed oriented surface of genus 2. Σ_i corresponds to a positive or negative stabilization on S , and Σ'_j corresponds to a positive or negative stabilization on D . As a result of the adjunction inequality in Lemma 2.20, we have

$$SHM(M', \gamma') \subset \bigoplus_{\substack{\mathfrak{s} \in \mathfrak{S}^*(Y|_{R_+}) \\ c_1(\mathfrak{s})[\bar{S}] \geq 2g(\bar{S}) - 2 - 2(s-t) - 2m}} \widetilde{HM}_\bullet(Y^\dagger, \mathfrak{s}). \quad (13)$$

Finally, we argue that

$$SHM(M', \gamma') \cap SHM(M'', \gamma'') = \{0\}.$$

Suppose not, then, from (11) and (13), there is a supporting spin^c structure $\mathfrak{s} \in S^*(Y|_{R_+})$ so that

$$c_1(\mathfrak{s})[\bar{S}] \geq 2g(\bar{S}) - 2 - 2(s-t) - 2m, \text{ and } c_1(\mathfrak{s})[\bar{S}'] = 2g(\bar{S}') - 2.$$

From the construction, we know that

$$[\bar{S}'] = -[\bar{S}] + 2k \cdot [R_+] \subset H_2(Y).$$

Hence, the above equalities and inequalities imply

$$2g(\bar{S}) - 2 - 2(s-t) - 2m + 2g(\bar{S}') - 2 \leq c_1(\mathfrak{s})[\bar{S}] + c_1(\mathfrak{s})[\bar{S}'] = 2k \cdot [2g(R_+) - 2],$$

which is equivalent to

$$\chi(\bar{S}) + \chi(\bar{S}') + 2(s-t) + 2m \geq 2k \cdot \chi(R_+). \quad (14)$$

Now let us compute each of the three terms in (14) regarding the Euler characteristics. First, $\chi(\bar{S}) = \chi(S_2)$, and, by construction, $S_2 = S_1 \sqcup (s-t)D_1$. Furthermore, we know

that S_1 is obtained from S_0 by attaching $m + n$ copies of 2-dimensional 1-handles, that S_0 is isotopic to S , and that D_1 is obtained from a disk D by attaching two copies of 2-dimensional 1-handles. Thus, we conclude that

$$\chi(S_2) = \chi(S) - (m + n) + (s - t)(-1) = \chi(S) - (m + n) - (s - t).$$

Second, we know that $\chi(\bar{S}'_2) = \chi(\tilde{S}'_2)$, and

$$\tilde{S}'_2 = S'_1 \cup D'_1 \cup (2k) \cdot T.$$

Note that there are $m + n$ product 1-handles attached to (M, γ) , and, inside each product 1-handle, there are $(2k + 1)$ copies of 2-dimensional 1-handles attached to S' . Thus, we have that

$$\chi(S'_1) = \chi(S') - (2k + 1)(m + n).$$

Similarly, we conclude that

$$\chi(D'_1) = \chi(D') - 2(2k + 1).$$

Also, D' is obtained by a double curve surgery on $-D$, which is a disk, with k copies of $R(\delta)$, which is the disjoint union of two annuli. Thus, we conclude that

$$\chi(\bar{S}') = \chi(S') - (m + n) - (s - t) + 2k \cdot \chi(T) - 2k(m + n) - 4k(s - t).$$

Third, we know that

$$R_+ = R_+(\gamma_1) \cup (s - t) \cdot R_+(\delta_1) \cup T.$$

Here, $R_+(\gamma_1)$ is obtained from $R_+(\gamma)$ by attaching $m + n$ copies of 2-dimensional 1-handles, and $R_+(\delta_1)$ is obtained from $R_+(\delta)$ (an annulus) by attaching 2 copies of 2-dimensional 1-handles. Thus, we know that

$$\chi(R_+) = \chi(R_+(\gamma)) + \chi(T) - (m + n) - 2(s - t).$$

Putting everything together, (14) is equivalent to

$$\chi(S) + \chi(S') - 2n \geq 2k \cdot \chi(R_+(\gamma)).$$

This directly contradicts Lemma 5.9, since, by definition, we have

$$n = \frac{1}{2}|S \cap \gamma|.$$

□

Lemma 5.9. *Suppose (M, γ) , S and S' are as above in Proposition 5.8, then*

$$\chi(S) + \chi(S') - |S \cap \gamma| < 2k \cdot \chi(R_+(\gamma)).$$

Proof. This is exactly the inequality

$$\chi(S) + \chi(S') + I(S) + I(S') < r(S, t) + r(S', t)$$

in the proof of Theorem 6.1 in Juhász [11]. It is by definition that

$$I(S) = I(S') = -\frac{1}{2}|S \cap \gamma|,$$

and Juhász also proved that

$$r(S, t) + r(S', t) = 2k \cdot \chi(R_+(\gamma)).$$

Note $2k$ in this paper corresponds to k in his paper. $\square$

Remark 5.10. Note the assumption in Proposition 5.8 is slightly stronger than the hypothesis of Theorem 6.1 in Juhász [11], i.e., we require both being reduced and containing no essential product disks, while Juhász only required being reduced. The difference between the two setups is some special family of balanced sutured manifolds, which are reduced but also contain essential product disks. By Lemma 2.13 in [11], there are only two such balanced sutured manifolds, namely the product sutured manifolds $(times[-1, 1] \times F, \{0\} \times \partial F)$, where F is a sphere with two or three disks removed. It is also worth mentioning that the requirement of containing no essential product disks. Clearly, the two special product sutured manifolds described above are counterexamples to Theorem 6.1 in Juhász [11]. The small error made in the proof of Theorem 6.1 in his paper is that, at some point, he used the assumption of being reduced and applied Lemma 2.13 in [11] to rule out essential product disks from the discussion, but he didn't exclude the two special product sutured manifolds from the hypothesis, where clearly Lemma 2.13 in [11] failed.

Corollary 5.11. *Suppose (M, γ) is a balanced sutured manifold with $H_2(M) = 0$. Suppose further that (M, γ) is taut, horizontally prime, reduced, and free of essential product disks. Then, the dimensions of $PM_a(M, \gamma)$ and $PI_a(M, \gamma)$ (see definition 5.4) are both $\dim_{\mathbb{Q}} H^2(M, \partial M; \mathbb{Q})$.*

Proof. We prove in the monopole settings. Pick any $\alpha \in H_2(M, \partial M)$ so that $\alpha \neq 0$. As in the proof of Proposition 5.8, we can find two properly embedded surfaces S and S' in M representing α and $-\alpha$, a closure Y of (M, γ) and suitable closed surfaces $\bar{S}$ and $\bar{S}'$ originate from S and S' , respectively. Let b be a homogenous elements in the top grading induced by $\bar{S}$ and let c be a homogenous elements in the top grading induced by $\bar{S}'$. Suppose further that they are supported by spin^c structures $\mathfrak{s}_b$ and $\mathfrak{s}_c$ on Y . We know from the definition that

$$\rho(b, c)(\alpha) = \frac{1}{2}[c_1(\mathfrak{s}_b) - c_1(\mathfrak{s}_c)](\alpha).$$

We claim that $\rho(b, c)(\alpha) \neq 0$. Suppose the contrary, then we know that

$$c_1(\mathfrak{s}_c)[\bar{S}] = c_1(\mathfrak{s}_b)[\bar{S}] \geq 2g(\bar{S}) - 2 - 2(s - t) - 2m, \quad (15)$$

and

$$c_1(\mathfrak{s}_c)[\bar{S}'] = 2g(\bar{S}') - 2. \quad (16)$$

Here, s, t, m are constants in the construction of $\bar{S}$ and $\bar{S}'$, as in the proof of Proposition 5.8. Then, formulae (15) and (16) lead to exactly the same contradiction as in the proof of Proposition 5.8.

Since $\rho(b, c)(\alpha) \neq 0$ and $\alpha \in H_2(M, \partial M)$ is chosen arbitrarily, we conclude that the polytope $PM_a(M, \gamma)$ must have maximal possible dimension. $\square$

Corollary 5.12. *Suppose (M, γ) is a taut balanced sutured manifold with $H_2(M) = 0$. Then*

$$\text{rk}(\underline{\text{SHM}}(M, \gamma)) \geq \dim_{\mathbb{R}} H^2(M, \partial M; \mathbb{R}) + 1$$

and

$$\dim_{\mathbb{C}}(\underline{\text{SHI}}(M, \gamma)) \geq \dim_{\mathbb{R}} H^2(M, \partial M; \mathbb{R}) + 1$$

Proof. To have dimension $d = \dim_{\mathbb{R}} H^2(M, \partial M; \mathbb{R})$, there must be at least $d+1$ points inside the polytope and we are done. $\square$

Now we are ready to prove Theorem 1.2 as stated in the introduction.

Proof of Theorem 1.2. This is essentially the proof of Proposition 7.6 in Juhász [11] but carried out in the monopole or the instanton settings. We will present the proof in the monopole settings, and the instanton case follows from a similar argument.

First suppose $k = 0$. By Proposition 2.16 and Proposition 2.18 in Juhász [11], we can perform a sutured manifold decomposition on (M, γ) to obtain (M', γ') that is taut, reduced, and horizontally prime. By Lemma 5.1 in Juhász [11], $H_2(M') = 0$. By Proposition 6.6 in Kronheimer and Mrowka [16] and Lemma 5.6 in this paper,

$$1 \leq \text{rk}(\text{SHM}(M', \gamma')) \leq \text{rk}(\text{SHM}(M, \gamma)) < 2.$$

Hence, $\text{rk}(\text{SHM}(M', \gamma')) = 1$. By Proposition 5.8, this implies that $H_2(M', \partial M') = 0$ and consequently, $H_1(\partial M') = 0$, which means $\partial M'$ is a sphere. ($H_2(M') = 0$ implies that $\partial M'$ is connected.) Since M' is irreducible, M' must be a 3-ball, and γ' must be connected due to tautness. So, (M', γ') is a product sutured manifold and so is (M, γ) .

Now we assume that the conclusion of the proposition holds for $k-1$, and next, we prove it for k . This part of the proof is exactly the same as in Juhász [11], so we only sketch as follows: if (M, γ) is not horizontally prime, then we could decompose along non-boundary-parallel horizontal surfaces and get a disjoint union of balanced sutured manifolds. Each component has a sutured monopole Floer homology of rank at most 2^k , and thus inductive hypothesis applies. If (M, γ) is horizontally prime we can perform a sutured manifold decomposition to make it reduced, and applying Proposition 5.8 to choose a suitable decomposition surface so that a (second) sutured manifold decomposition along the chosen surface will reduce the dimension by at least a half. Then, the inductive hypothesis applies again, and this concludes the proof of theorem 1.2. $\square$

As a corollary to the above proposition, we offer a new proof to the fact that the monopole and instanton knot Floer homology constructed by Kronheimer and Mrowka [16] detects fibred knots in S^3 .

Corollary 5.13. *Suppose $K \subset S^3$ is a knot. Then, the following three things are equivalent.*

- (1) $\text{rk}(KHM(S^3, K, g(K))) = 1$.
- (2) $\text{rk}(KHI(S^3, K, g(K))) = 1$.
- (3) K is a fibred knot.

Proof. We only prove that (1) and (3) are equivalent. From Kronheimer and Mrowka [16], we know that

$$KHM(S^3, K) = SHM(S^3(K), \Gamma_\mu)$$

where $S^3(K)$ is the knot complement and Γ_μ is the suture consisting of two meridians on $\partial S^3(K)$. Pick a minimal genus Seifert surface, $S \subset S^3$ of K . We know that the decomposition

$$(S^3(K), \Gamma_\mu) \xrightarrow{S} (M, \gamma)$$

is taut, and

$$KHM(S^3, K, g(K)) \cong SHM(M, \gamma).$$

Thus, the corollary follows from Theorem 1.2. $\square$

Proof of Corollary 1.5. By Lemma 7.10 in Xie and Zhang [27], we have an isomorphism

$$SHM(M, \gamma, T) \cong SHI(M_T, \gamma_T),$$

where (M_T, γ_T) is some balanced sutured manifold arising from the triple (M, γ, T) as explained in Section 7 in [27], and $SHI(M_T, \gamma_T)$ is the usual sutured instanton Floer homology defined by Kronheimer and Mrowka in [16]. From the description of M_T in [27], we know that $H_2(M_T) = H_2(M \setminus T)$ and hence Theorem 1.2 applies. $\square$

6 Application to knots and links

6.1 Thurston-norm detection

In this subsection, we prove Theorem 1.17. We only work in the monopole settings and the proof in the instanton settings is exactly the same. First, we need some preparations.

Lemma 6.1. *Suppose (M, γ) is a taut balanced sutured manifold so that M is boundary-incompressible and the boundary of M consists of a few tori. Suppose further that $\alpha \in H_2(M, \partial M)$ is a non-zero second relative homology class. Then, there is a properly embedded surface $S \subset M$ with the following properties.*

- (1) $[S, \partial S] = \alpha \in H_2(M, \partial M)$.
- (2) $\chi(S) = -x(\alpha)$.
- (3) For any component Σ of ∂M , $S \cap \Sigma$ consists of a disjoint union of coherently oriented non-separating simple closed curves on Σ .
- (4). S is incompressible.

Proof. Pick a surface S so that $[S, \partial S] = \alpha \in H_2(M, \partial M)$, and $x(S) = x(\alpha)$. Since M is irreducible, we can assume that there is no spherical component of S . We have assumed that M is boundary-incompressible, so we can also assume that there is no disk component of S . Thus, we have $x(\alpha) = x(S) = -\chi(S)$.

To achieve condition (3) in the conclusion of the lemma, if a component α of ∂S bounds a disk on ∂M , then we can cap off α using the disk it bounds on ∂M . Capping off by a disk does not increase the norm, so we can assume that the surface S does not have a boundary component that bounds a disk on ∂M .

Pick a component Σ of ∂M . By assumption, Σ is a torus. Since no component of $S \cap \Sigma$ bounds a disk, we know that $S \cap \Sigma$ consists of a disjoint union of parallel non-separating simple closed curves. If two components of $S \cap \Sigma$ are adjacent on Σ but are oriented reversely, we can glue the annulus, which they co-bound on Σ , to S . After possible compressions and throwing away any spherical or disk components arising from the compression, we still call the resulting surface S . Note gluing annuli, performing compressions, and throwing away spherical and disk components do not increase the norm. Thus, we conclude the proof of Lemma 6.1. $\square$

Lemma 6.2. *Suppose (M, γ) is a taut balanced sutured manifold so that M is boundary-incompressible, and the boundary of M consists of a few tori. Suppose $\alpha \in H_2(M, \gamma)$ is a non-zero second relative homology class, and S is a properly embedded surface inside M satisfying conditions (1)-(3) in Lemma 6.1. Then, the decomposition of (M, γ) along S is taut.*

Proof. Suppose the sutured manifold decomposition of (M, γ) along S yields (M', γ') , then we can regard M' as a submanifold of M . Suppose Σ is a component of ∂M , then by assumption, Σ is a torus. If $\Sigma \cap S = \emptyset$, then Σ is also a component of ∂M . Thus, we have

$$\gamma' \cap \Sigma = \gamma \cap \Sigma, \text{ and } R_{\pm}(\gamma') \cap \Sigma = R_{\pm}(\gamma) \cap V.$$

If $V \cap S \neq \emptyset$, then $\partial M' \cap V$ consists of a disjoint union of annuli, which, regardless of the orientations, are bounded by pairs of parallel curves in $\partial S \cap V$. Let $A \subset V$ be a component of $\partial M' \cap V$. There are two cases, depending on the intersection of the suture γ with the surface S . In both cases, it is straightforward to check how γ' looks like.

Case 1. $(S \cap \Sigma) \cap (\gamma \cap \Sigma) = \emptyset$. In this case, the annulus A possibly contains multiple components of γ , and they remain in γ' . Thus, A may contain either one or three components of γ' (note $\gamma \cap \Sigma$ has two components), and each component of γ' is parallel to ∂A . See Figure 16.

Case 2. $(S \cap \Sigma) \cap (\gamma \cap \Sigma) \neq \emptyset$. In this case, $\gamma \cap A$ consists of an even number of essential arcs in A , and adjacent arcs are oriented oppositely. Then, after the decomposition, A contains exactly one component of γ' , and this component is parallel to ∂A . See Figure 17.

Suppose $A_1, \dots, A_n$ are all the annular components of $\partial M' \cap \partial M$ that contain three components of γ' . Push the interiors of A_i into the interior of M' to make them properly embedded. Then, we can perform a sutured manifold decomposition




Figure 16: Left, before the decomposition. The vertical (blue) curves represent S , and the horizontal curves represent Σ . The (red) dots represent the suture γ . Right, after the decomposition. The (red) dots represent the suture γ' .

on (M', γ') , along the surface $A_1 \cup \dots \cup A_n$, after the pushing off. The resulting balanced sutured manifold (M'', γ'') is a disjoint union:

$$(M'', \gamma'') = (M''', \gamma''') \cup (V_1, \gamma_1^4) \cup \dots \cup (V_n, \gamma_n^4).$$

Here, for $i = 1, \dots, n$, V_i is a framed solid torus and γ_i^4 is the suture on ∂V consisting of four longitudes. With $\mathbb{Q}$ coefficients, we know from Li [17] that

$$SHM(V_i, \gamma_i^4) \cong \mathbb{Q}^2.$$

From Proposition 6.9 in Kronheimer and Mrowka [16], we know

$$SHM(M', \gamma') \cong SHM(M'', \gamma'') \cong SHM(M''', \gamma''') \otimes_{\mathbb{Q}} \mathbb{Q}^{2^n}.$$

Thus, (M', γ') is taut if and only if (M''', γ''') is.

It is then suffice to prove that (M''', γ''') is taut. Note we can regard $M''' = M'$, and, thus, we can assume

$$M''' = M \setminus \text{int}(N(S)).$$

Let $S_{\pm}$ be parallel copies of S in $\partial N(S)$, then $S_{\pm}$ are part of the boundary of M''' . Let Σ_{∂} be the union of components of ∂M which are disjoint from S , and let

$$F_{\pm} = \Sigma_{\partial} \cap R_{\pm}(\gamma).$$




Figure 17: Left, before the decomposition. The two (blue) circles are the boundary of A . The (red) arcs represent the suture γ . Right, after the decomposition. The two dashed circles are the boundary of A , and the (red) solid curve represent the suture γ' .

Then, we can describe $R_{\pm}(\gamma''')$ as follows:

$$R_{\pm}(\gamma''') = F_{\pm} \sqcup S_{\pm}.$$

By assumption, both $F_{\pm}$ and $S_{\pm}$ are incompressible and norm-minimizing in M , hence they are also incompressible and norm-minimizing in M''' . The fact that M is irreducible implies that M''' is the same. Thus, we conclude the proof of lemma 6.2. $\square$

Corollary 6.3. *The decomposition of (M, γ) along $-S$ is also taut.*

Proof of Theorem 1.17. For any $\alpha \in H_2(M, \partial M)$, pick a surface S as in Lemma 6.1. Then, by corollary 6.3, the decomposition of $(Y(L), \Gamma_{\mu})$ along S and $-S$ are both taut. Suppose

$$n = \frac{1}{2}|S \cap \gamma|,$$

then we know, from condition (3) in the statement of Lemma 6.1, that

$$n = \sum_{i=1}^r |\langle \alpha, \mu_i \rangle|.$$

We possibly need to perform a stabilization on S to achieve admissibility. Suppose the S^{+m} is obtained from S by perform m many positive stabilizations on S . Here, $m = 0$ or 1 . Note $m = 0$ means that the original S is admissible, and we take $S^0 = S$. From Lemma 2.20, Lemma 2.23, and Lemma 2.28, we know that

$$\underline{\text{SHM}}(Y(L), \Gamma_{\mu}, S^{+m}, g_c) \neq 0, \text{ and } \underline{\text{SHM}}(Y(L), \Gamma_{\mu}, S^{+m}, i) = 0 \text{ for } i > g_c.$$

Here, we have $g_c = \frac{1}{2}(-\chi(S) + n + m)$.

Similarly, we have

$$\underline{\text{SHM}}(Y(L), \Gamma_\mu, (-S)^{+m}, g_c) \neq 0, \text{ and } \underline{\text{SHM}}(Y(L), \Gamma_\mu, (-S)^{+m}, i) = 0 \text{ for } i > g_c.$$

Note $(-S)^{+m} = -(S^{-m})$, and, hence,

$$\underline{\text{SHM}}(Y(L), \Gamma_\mu, S^{-m}, -g_c) = \underline{\text{SHM}}(Y(L), \Gamma_\mu, (-S)^{+m}, g_c) \neq 0,$$

and

$$\underline{\text{SHM}}(Y(L), \Gamma_\mu, S^{-m}, i) = 0 \text{ for } i < -g_c.$$

Apply proposition 4.1, we know that

$$\underline{\text{SHM}}(Y(L), \Gamma_\mu, S^{+m}, m - g_c) = \underline{\text{SHM}}(Y(L), \Gamma_\mu, (-S)^{+m}, g_c) \neq 0,$$

and

$$\underline{\text{SHM}}(Y(L), \Gamma_\mu, S^{+m}, i) = 0 \text{ for } i < m - g_c.$$

From the definition of the function $y(\cdot)$ in Definition 1.16 and the construction of the canonical decomposition of sutured monopole Floer homology in 5.4, we know that

$$\begin{aligned} y(\alpha) &= \max\{i \mid \underline{\text{SHM}}(Y(L), \Gamma_\mu, S^{+m}, i) \neq 0\} - \min\{i \mid \underline{\text{SHM}}(Y(L), \Gamma_\mu, S^{+m}, i) \neq 0\} \\ &= g_c - (m - g_c) \\ &= 2g_c - m \\ &= -\chi(S) + n + m - m \\ &= x(\alpha) + \sum_{i=1}^r |\langle \alpha, \mu_i \rangle|. \end{aligned}$$

This concludes the proof of Theorem 1.17. $\square$

Suppose $L \subset S^3$ is a link with r components. Then we know that $H_2(S^3(L), \partial S^3(L)) \cong \mathbb{Z}^r$. Thus there is a $\mathbb{Z}^r$ grading on $KHM(S^3, K)$ and $KHI(S^3, K)$, according to the proof of lemma 5.2. Here KHM and KHI are the monopole and instanton knot Floer homologies introduced by Kronheimer and Mrowka in [16]. This leads to the following question.

Question 6.4. *Can we recover the multi-variable Alexander polynomial using the $\mathbb{Z}^r$ grading on KHM or KHI ?*

6.2 Minus version for links

Suppose Y is a closed oriented 3-manifold and $L \subset Y$ is an oriented link. Let $L_1, L_2, \dots, L_r$ be the components of L . We assume further that each component of L is null-homologous in Y . Thus, for $i = 1, \dots, r$, we can find (and fix) a Seifert surface $S_i \subset Y$ for L_i . Note S_i possibly intersects with L_j , for $j \neq i$. Also, for $i = 1, \dots, r$, let $p_i \in L_i$ be a fixed base point. Let $\mathbf{p} = (p_1, \dots, p_r)$. In this subsection, we construct minus versions of monopole and instanton knot Floer homologies for the triple $(-Y, L, \mathbf{p})$.

Remark 6.5. Here, we require that each component of L to be null-homologous, to fix a Seifert surface for each component of the link. It is possible to weaken this condition by simply requiring that the whole link L represents the zero class in $H_1(Y)$ and fix a Seifert surface S for it. The construction in this subsection can be easily adapted to the more general setup.

Let $Y(L) = Y \setminus N(L)$ be the knot complement and let T_i be the boundary component of $Y(L)$ corresponding to the knot L_i . The Seifert surface S_i induces a framing on T_i . We call the longitude λ_i and the meridian μ_i . For $\mathbf{n} = (n_1, \dots, n_r) \in \mathbb{Z}^r$, let $\Gamma_{\mathbf{n}}$ be the suture on $\partial Y(L)$ so that $\Gamma_{\mathbf{n}} \cap T_i$ consists of two parallel simple closed curves of class $\pm[\mu_i - n\lambda_i]$. We have the following lemma.

Remark 6.6. It seems that the choice of base points $\mathbf{p}$ does not appear in the above set up. However, $\mathbf{p}$ helps to resolve the ambiguity arising from the choice of the link complements. Since this issue is fully clarified in Baldwin and Sivek [1] and Li [19], we won't discuss on it anymore in this paper.

Lemma 6.7. *Suppose $\mathbf{n} = (n_1, \dots, n_r) \in (\mathbb{Z}_+)^r$. Let $\mathbf{n}'$ be obtained from $\mathbf{n}$ by replacing n_i with $n_i + 1$, and let $\mathbf{n}''$ be obtained from $\mathbf{n}$ by replacing n_i with $+\infty$, then there are exact triangles:*

$$\begin{array}{ccc} \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n}}) & \xrightarrow{\psi_{\pm, \mathbf{n}, i}} & \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n}'}) \\ & \swarrow \psi_{\pm, \mathbf{n}'', i} \quad \searrow \psi_{\pm, \mathbf{n}', i} & \\ & \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n}''}) & \end{array}$$

Here, $\psi_{\pm, \mathbf{n}, i}$ are the map associated to a positive or negative by-pass attached to $Y(L), \Gamma_{\mathbf{n}}$, which is performed on the boundary component T_i of $Y(L)$. (See Subsection 2.4.)

There are similar exact triangles in the instanton settings.

Proof. This is a direct application of Theorem 2.29. For more details, readers are referred to Section 2 of Li [19]. $\square$

To use a better notation, for $i = 1, \dots, r$, let $\mathbf{e}^i = (0, \dots, 1, \dots, 0) \in \mathbb{Z}^r$ be the vector whose entries are all 0 except of being 1 on the i -th place. We have the following lemma.

Lemma 6.8. *For any $\mathbf{n} \in (\mathbb{Z}_+)^r$ and $i, j \in \{1, \dots, r\}$, we have the following commutative diagram:*

$$\begin{array}{ccc} \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n}}) & \xrightarrow{\psi_{-, \mathbf{n}, i}} & \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n} + \mathbf{e}^i}) \\ \downarrow \psi_{-, \mathbf{n}, j} & & \downarrow \psi_{-, \mathbf{n} + \mathbf{e}^i, j} \\ \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n} + \mathbf{e}^j}) & \xrightarrow{\psi_{-, \mathbf{n} + \mathbf{e}^j, i}} & \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n} + \mathbf{e}^i + \mathbf{e}^j}) \end{array}$$

Proof. As explained in Subsection 2.4, the by-pass maps, $\psi_{\pm, \mathbf{n}, i}$, ultimately come from contact handle attaching maps. Since the by-passes corresponding to vertical and horizontal maps happen on different boundary components of $Y(L)$, the corresponding contact handle attachments commute, and so do the by-pass maps. $\square$

Definition 6.9. We define the minus version of monopole link Floer homology of a based link $L \subset -Y$, which is denoted by $\underline{\text{KHM}}^-(-Y, L, \mathbf{p})$, to be the direct limit of the direct system

$$\{\psi_{-, \mathbf{n}, i} : \underline{\text{SHM}}(-Y(L), \Gamma_{\mathbf{n}}) \rightarrow \underline{\text{SHM}}(-Y(L), \Gamma_{\mathbf{n} + \mathbf{e}^i}), \mathbf{n} \in (\mathbb{Z}_+)^r, i \in 1, \dots, r\}.$$

We define $\underline{\text{KHI}}^-(-Y, L, \mathbf{p})$ in a similar manner.

Lemma 6.10. For any $\mathbf{n} \in (\mathbb{Z}_+)^r$ and $i, j \in \{1, \dots, r\}$, we have the following commutative diagram:

$$\begin{array}{ccc} \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n}}) & \xrightarrow{\psi_{-, \mathbf{n}, i}} & \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n} + \mathbf{e}^i}) \\ \downarrow \psi_{+, \mathbf{n}, j} & & \downarrow \psi_{+, \mathbf{n} + \mathbf{e}^i, j} \\ \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n} + \mathbf{e}^j}) & \xrightarrow{\psi_{-, \mathbf{n} + \mathbf{e}^j, i}} & \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n} + \mathbf{e}^i + \mathbf{e}^j}) \end{array}$$

There are similar commutative diagrams in the instanton settings.

Proof. If $i \neq j$, this follows from exactly the same argument as in the proof of Lemma 6.8. If $i = j$, this follows from the proof of the same type of commutative diagram in the construction of minus versions for knots by the second author. See Corollary 2.22 in Li [19]. $\square$

Definition 6.11. For any fixed $i \in \{1, \dots, r\}$, The set of maps

$$\{\psi_{+, \mathbf{n}, i} : \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n}}) \rightarrow \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n} + \mathbf{e}^i})\}$$

induces a map

$$U_i : \underline{\text{KHM}}^-(-Y, L, \mathbf{p}) \rightarrow \underline{\text{KHM}}^-(-Y, L, \mathbf{p}),$$

which we call the i -th U map.

We define

$$U_i : \underline{\text{KHI}}^-(-Y, L, \mathbf{p}) \rightarrow \underline{\text{KHI}}^-(-Y, L, \mathbf{p})$$

in a similar manner.

Proposition 6.12. For any $i, j \in \{1, \dots, r\}$, the maps U_i and U_j commute with each other.

Proof. The proof is exactly the same as the proof of Lemma 6.8. $\square$

Next, we construct a $\mathbb{Z}^r$ grading on $\underline{\text{KHM}}^-(-Y, L, \mathbf{p})$, based on the chosen Seifert surfaces $S_1, \dots, S_r$ of $L_1, \dots, L_r$.

Recall that $L \subset Y$ has components $L_1, L_2, \dots, L_r$, and L_i has a Seifert surface S_i that could possibly intersect other components of the link. By a slight abuse of notation, let S_i also denote the intersection of the original Seifert surface with the link complement $Y(L)$. Thus, the boundary of S_i consists of a longitude on T_i and a few (possibly none) meridians on T_j , for $j \neq i$. For fixed $i \in \{1, \dots, r\}$ and $\mathbf{n} = (n_1, \dots, n_r) \in (\mathbb{Z}_+)^r$, let $S_{i,\mathbf{n}}$ be the isotopy of S_i so that $\partial S_{i,\mathbf{n}}$ has the least possible intersections with the suture $\Gamma_{\mathbf{n}} \subset \partial Y(L)$. This means that the longitudinal boundary component of $S_{i,\mathbf{n}}$ intersects $\Gamma_{\mathbf{n}}$ at $2n_i$ points, and each meridional boundary component of $S_{i,\mathbf{n}}$ intersects $\Gamma_{\mathbf{n}}$ at two points. Applying the construction of gradings in Subsection 2.3, the surface $S_{i,\mathbf{n}}$, or its stabilizations, give rise to a grading on $\underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n}})$. We then have the following proposition.

Proposition 6.13. *Fix any $i \in \{1, \dots, r\}$ and $\mathbf{n} \in (\mathbb{Z}_+)^r$. If n_i is even, then, for any $j \in \mathbb{Z}$, we have*

$$\psi_{\pm, \mathbf{n}, i}(\underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n}}, S_{i,\mathbf{n}}^{\pm}, j)) \subset \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n}+e^i}, S_{i,\mathbf{n}+e^i}, j)$$

If n_i is odd, then, for any $j \in \mathbb{Z}$,

$$\psi_{\pm, \mathbf{n}, i}(\underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n}}, S_{i,\mathbf{n}}^{\pm 2}, j)) \subset \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n}+e^i}, S_{i,\mathbf{n}+e^i}^{\pm}, j)$$

Furthermore, for any $k \neq i$, the maps $\psi_{\pm, \mathbf{n}, k}$ preserve the gradings associated to $S_{i,\mathbf{n}}$ and its stabilizations.

Similar statements hold for the instanton settings.

Proof. For $\psi_{\pm, \mathbf{n}, i}$, the proof is exactly the same as the proof of Proposition 5.5 in Li [19]. For $\psi_{\pm, \mathbf{n}, k}$ with $k \neq i$, note $S_{i,\mathbf{n}}$ has a few meridional components on the T_k , so the by-passes, which corresponds to the maps $\psi_{\pm, \mathbf{n}, k}$, can actually be made disjoint from $S_{i,\mathbf{n}}$. $\square$

Similar to the constructions in Section 5 of Li [19], if n_i is odd, let $S_{i,\mathbf{n}}^{\tau}$ be just $S_{i,\mathbf{n}}$, and if n_i is even, let $S_{i,\mathbf{n}}^{\tau}$ be a negative stabilization of $S_{i,\mathbf{n}}$ performed near T_i . We can use $S_{i,\mathbf{n}}^{\tau}$ to define a grading on $\underline{\text{SHM}}(-Y(L), \Gamma_{\mathbf{n}})$. We also need to perform a grading shift. Let

$$\underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n}}, S_{i,\mathbf{n}}^{\tau}, j)\{\sigma_i\} = \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n}}, S_{i,\mathbf{n}}^{\tau}, j + \lfloor \frac{n}{2} \rfloor),$$

where $\lfloor x \rfloor$ is to take the maximal integer which is no larger than x .

Proposition 6.14. *Using the grading $\underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n}}, S_{i,\mathbf{n}}^{\tau}, j)\{\sigma_i\}$, we can construct a $\mathbb{Z}$ -grading on $\underline{\text{KHM}}^-(-Y, L, \mathbf{p})$. The i -th U map, U_i , drops the grading by 1, and all other U maps, U_k with $k \neq i$, preserve the grading.*

Furthermore, all Seifert surfaces, $S_1, \dots, S_r$, together induce a $\mathbb{Z}^r$ grading on $\underline{\text{KHM}}^-(-Y, L, \mathbf{p})$, which we write as

$$\underline{\text{KHM}}^-(-Y, L, \mathbf{p}, \mathbf{j}).$$

Here, $\mathbf{j} \in \mathbb{Z}^r$ denote a multi-gradings. As a result, together with the commutativity of all U maps in Proposition 6.12, there is an $\mathcal{R}[U_1, \dots, U_r]$ module structure on $\underline{\text{KHM}}^-(-Y, L, \mathbf{p})$.

Similar results hold in the instanton settings.

Proof. The first half of the proposition follows from Proposition 6.13, and the second half of the proposition follows from the proof of Lemma 5.2. $\square$

The first computable example is the case of unlinks.

Proposition 6.15. *Suppose Y is a closed oriented 3-manifold and $L \subset Y$ is an unlink of r components, i.e., there exists an embedded disk $S_i \cong D^2$, for each $i \in \{1, \dots, r\}$, so that $\partial S_i = L_i$, and all S_i are disjoint from each other. Then,*

$$\underline{\text{KHM}}^-(-Y, L, \mathbf{p}) \cong \underline{\text{SHM}}(-Y(r), -\delta^r) \otimes_{\mathcal{R}} \mathcal{R}[U_1, \dots, U_r].$$

Here, $\mathbf{p}$ is a chosen set of base points, and $(Y(r), \delta^r)$ is the balanced sutured manifold obtained from Y by removing r disjoint 3-balls and picking one simple closed curve on each spherical boundary of $Y(r)$ as the suture.

Similar statements hold in the instanton settings.

Proof. For any $\mathbf{n} \in (\mathbb{Z}_+)^r$, we know that $(Y(L), \Gamma_{\mathbf{n}})$ can be obtained from the disjoint union

$$(Y(r), \delta^r) \sqcup (S^3(L_1), \Gamma_{n_1}) \sqcup \dots \sqcup (S^3(L_r), \Gamma_{n_r})$$

by attaching r many contact 1-handles (see Definition 3.2). Each 1-handle connects some $(S^3(L_i), \Gamma_{n_i})$ to $(Y(r), \delta^r)$. As in Subsection 2.4, the by-pass maps $\psi_{\pm, \mathbf{n}, i}$ can be realized as contact handle attaching maps and those contact handles are disjoint form the contact 1-handles just described above. Hence, under the isomorphism

$$\underline{\text{SHM}}(-Y(L), \Gamma_{\mathbf{n}}) \cong \underline{\text{SHM}}(Y(r), \delta^r) \otimes \underline{\text{SHM}}(S^3(L_1), \Gamma_{n_1}) \otimes \dots \otimes \underline{\text{SHM}}(S^3(L_r), \Gamma_{n_r}),$$

we have an identification

$$\psi_{\pm, \mathbf{n}, i} = id \otimes \dots \otimes \psi_{\pm, n_i} \otimes \dots \otimes id,$$

where

$$\psi_{\pm, n_i} : \underline{\text{SHM}}(-S^3(L_i), -\Gamma_{n_i}) \rightarrow \underline{\text{SHM}}(-S^3(L_i), -\Gamma_{n_i+1}).$$

Hence, we are done. $\square$

Proposition 6.16. *Under the above setups, the direct system stabilizes, that is, for any fixed $j \in \mathbb{Z}$, there exists $N \in \mathbb{Z}$, so that for all $i \in \{1, \dots, r\}$ and $\mathbf{n} = (n_1, \dots, n_r) \in (\mathbb{Z}_+)^r$ such that $n_i > N$, we have an isomorphism*

$$\psi_{-, \mathbf{n}, i} : \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n}}, S_{i, \mathbf{n}}^T, j)[\sigma_i] \cong \underline{\text{SHM}}(-Y(L), -\Gamma_{\mathbf{n}+e^i}, S_{i, \mathbf{n}+e^i}^T, j)[\sigma_i].$$

Proof. This follows from exactly the same argument as in the proof of Proposition 5.8 in Li [19]. $\square$

Proposition 6.17. *Under the above setups, there exists an integer N_0 , so that for any fixed $i \in \{1, \dots, r\}$ and any multi-grading $\mathbf{j} = (j_1, \dots, j_r) \in \mathbb{Z}^r$ with $j_i < N_0$, the map U_i restricts to an isomorphism*

$$U_i : \underline{\text{KHM}}^-(Y, L, \mathbf{p}, \mathbf{j}) \cong \underline{\text{KHM}}^-(Y, L, \mathbf{p}, \mathbf{j} - \mathbf{e}^i).$$

Proof. This follows from exactly the same argument as in the proof of Corollary 5.9 in Li [19]. $\square$

References

- [1] John A. Baldwin and Steven Sivek. Naturality in sutured monopole and instanton homology. *J. Differential Geom.*, 100(3):395–480, 2015.
- [2] John A. Baldwin and Steven Sivek. A contact invariant in sutured monopole homology. *Forum Math. Sigma*, 4:e12, 82, 2016.
- [3] John A. Baldwin and Steven Sivek. Instanton Floer homology and contact structures. *Selecta Math. (N.S.)*, 22(2):939–978, 2016.
- [4] John A. Baldwin and Steven Sivek. Khovanov homology detects the trefoils. *arXiv preprint arXiv:1801.07634*, 2018.
- [5] David Gabai. Foliations and the topology of 3-manifolds. *J. Differential Geom.*, 18(3):445–503, 1983.
- [6] David Gabai. Foliations and the topology of 3-manifolds. II. *J. Differential Geom.*, 26(3):461–478, 1987.
- [7] Paolo Ghiggini. Knot Floer homology detects genus-one fibred knots. *Amer. J. Math.*, 130(5):1151–1169, 2008.
- [8] Ko Honda. On the classification of tight contact structures. I. *Geom. Topol.*, 4:309–368, 2000.
- [9] András Juhász. Holomorphic discs and sutured manifolds. *Algebr. Geom. Topol.*, 6:1429–1457, 2006.
- [10] András Juhász. Floer homology and surface decompositions. *Geom. Topol.*, 12(1):299–350, 2008.
- [11] András Juhász. The sutured Floer homology polytope. *Geom. Topol.*, 14(3):1303–1354, 2010.
- [12] Nithin Kavi. Cutting and gluing surfaces. *arXiv preprint arXiv:1810.11954*, 2019.
- [13] P. B. Kronheimer and T. S. Mrowka. Khovanov homology is an unknot-detector. *Publ. Math. Inst. Hautes Études Sci.*, (113):97–208, 2011.
- [14] Peter Kronheimer and Tom Mrowka. Instanton Floer homology and the Alexander polynomial. *Algebr. Geom. Topol.*, 10(3):1715–1738, 2010.
- [15] Peter Kronheimer and Tomasz Mrowka. *Monopoles and three-manifolds*, volume 10 of *New Mathematical Monographs*. Cambridge University Press, Cambridge, 2007.

-
- [16] Peter Kronheimer and Tomasz Mrowka. Knots, sutures, and excision. *J. Differential Geom.*, 84(2):301–364, 2010.
 - [17] Zhenkun Li. Contact structures, excisions and sutured monopoles. *arXiv preprint arXiv:1811.11634*, 2018.
 - [18] Zhenkun Li. Gluing maps and cobordism maps for sutured monopole floor homology. *arXiv preprint arXiv:1810.13071*, 2018.
 - [19] Zhenkun Li. Direct systems and the knot monopole floor homology. *arXiv preprint arXiv:1901.06679*, 2019.
 - [20] Yi Ni. Knot Floer homology detects fibred knots. *Invent. Math.*, 170(3):577–608, 2007.
 - [21] Burak Ozbagci. Contact handle decompositions. *Topology Appl.*, 158(5):718–727, 2011.
 - [22] Peter Ozsváth and Zoltán Szabó. Holomorphic disks and knot invariants. *Adv. Math.*, 186(1):58–116, 2004.
 - [23] Peter Ozsváth and Zoltán Szabó. Link Floer homology and the Thurston norm. *J. Amer. Math. Soc.*, 21(3):671–709, 2008.
 - [24] Martin Scharlemann. Sutured manifolds and generalized Thurston norms. *J. Differential Geom.*, 29(3):557–614, 1989.
 - [25] Martin Scharlemann and Jennifer Schultens. 3-manifolds with planar presentations and the width of satellite knots. *Trans. Amer. Math. Soc.*, 358(9):3781–3805, 2006.
 - [26] Yi Xie and Boyu Zhang. Classification of links with Khovanov homology of minimal rank. *arXiv preprint arXiv:1909.10032*, 2019.
 - [27] Yi Xie and Boyu Zhang. Instanton Floer homology for sutured manifolds with tangles. *arXiv preprint arXiv:1907.00547*, 2019.