

Polygons with Prescribed Angles in 2D and 3D

Alon Efrat, Radoslav Fulek, Stephen Kobourov, Csaba D. Tóth

Abstract

We consider the construction of a polygon P with n vertices whose turning angles at the vertices are given by a sequence $A = (\alpha_0, \dots, \alpha_{n-1})$, $\alpha_i \in (-\pi, \pi)$, for $i \in \{0, \dots, n-1\}$. The problem of realizing A by a polygon can be seen as that of constructing a straight-line drawing of a graph with prescribed angles at vertices, and hence, it is a special case of the well studied problem of constructing an *angle graph*.

In 2D, we characterize sequences A for which every generic polygon $P \subset \mathbb{R}^2$ realizing A has at least c crossings, and describe an efficient algorithm that constructs, for a given sequence A , a generic polygon $P \subset \mathbb{R}^2$ that realizes A with the minimum number of crossings. In 3D, we describe an efficient algorithm that tests whether a given sequence A can be realized by a (not necessarily generic) polygon $P \subset \mathbb{R}^3$, and for every realizable sequence finds a realization.

1 Introduction

Straight-line realizations of graphs with given metric properties have been one of the earliest applications of graph theory. Rigidity theory, for example, studies realizations of graphs with prescribed edge lengths, but also considers a mixed model where the edges have prescribed lengths or directions [4, 13, 14, 15, 21]. In this paper, we extend research on the so-called *angle graphs*, introduced by in the 1980s, which are geometric graphs with prescribed angles between adjacent edges. Angle graphs found applications in mesh flattening [29], and computation of conformal transformations [8, 22] with applications in the theory of minimal surfaces and fluid dynamics.

Viyajan [27] characterized planar angle graphs under various constraints, including the case when the graph is a cycle [27, Theorem 2] and when the graph is 2-connected [27, Theorem 3]. In both cases, the characterization leads to an efficient algorithm to find a planar straight-line drawing or report that none exists. Di Battista and Vismara [6] showed that for 3-connected angle graphs (e.g., a triangulation), planarity testing reduces to solving a system of linear equations and inequalities in linear time. Garg [10] proved that planarity testing for angle graphs is NP-hard, disproving a conjecture by Viyajan. Bekos et al. [2] showed that the problem remains NP-hard even if all angles are multiples of $\pi/4$.

The problem of computing (straight-line) realizations of angle graphs can be seen as the problem of reconstructing a drawing of a graph from the given

partial information. The research problems to decide if the given data uniquely determine the realization or its parameters of interest is already interesting for cycles, where it found applications in the area of conformal transformations [22], and visibility graphs [7].

In 2D, we are concerned with realizations of angle cycles as polygons minimizing the number of crossings which, as we will see, depends only on the sum of the turning angles. It follows from the seminal work of Tutte [26] and Thomassen [25] that every positive instance of a 3-connected planar angle graph admits a crossing-free realization if the prescription of the angles implies the convexity for the faces. The convexity will also play the crucial role in our proofs.

In 3D, we test whether a given angle cycle can be realized by a (not necessarily generic) polygon. Somewhat counter-intuitively, self-intersections cannot be always avoided in a polygon realizing the given angle cycle in 3D. Di Battista et al. [5] characterized oriented polygons that can be realized in $\mathbb{R}^3$ without self-intersections with axis-parallel edges of given directions. Patrignani [20] showed that recognizing crossing-free realizability is NP-hard for graphs of maximum degree 6 in this setting.

Throughout the paper we assume modulo n arithmetic on the indices.

Angle sequences in 2-space. In the plane, an *angle sequence* A is a sequence $(\alpha_0, \dots, \alpha_{n-1})$ of real numbers such that $\alpha_i \in (-\pi, \pi)$ for all $i \in \{0, \dots, n-1\}$. Let $P \subset \mathbb{R}^2$ be an oriented polygon with n vertices $v_0, \dots, v_{n-1}$ that appear in the given order along P , which is consistent with the given orientation of P . The *turning angle* of P at v_i is the angle in $(-\pi, \pi)$ between the vector $v_i - v_{i-1}$ and $v_{i+1} - v_i$. The sign of the angle is positive if in the plane containing v_{i-1}, v_i and v_{i+1} , in which the vector $v_i - v_{i-1}$ points in the positive direction of the x -axis, the y -coordinate of $v_{i+1} - v_i$ is positive, and non-positive otherwise, see Figure 1.




Figure 1: A negative (left) and a positive (right) turning angle α_i at the vertex v_i of an oriented polygon.

The oriented polygon P *realizes* the angle sequence A if the turning angle of P at v_i is equal to α_i , for $i = 0, \dots, n-1$. A polygon P is *generic* if all its self-intersections are transversal (that is, proper crossings), vertices of P are distinct points, and no vertex of P is contained in a relative interior of an edge of P . Following the terminology of Vijayan [27], an *angle sequence* is *consistent*

if there exists a generic closed polygon P with n vertices realizing A . For a polygon P that realizes an angle sequence $A = (\alpha_0, \dots, \alpha_{n-1})$ in the plane, the *total curvature* of P is $\text{TC}(P) = \sum_{i=0}^{n-1} \alpha_i$, and the *turning number* (also known as *rotation number*) of P is $\text{tn}(P) = \text{TC}(P)/(2\pi)$; it is known that $\text{tn}(P) \in \mathbb{Z}$ in the plane [24].

The *crossing number*, denoted by $\text{cr}(P)$, of a generic polygon is the number of self-crossings of P . The *crossing number* of a consistent angle sequence A is the minimum integer k , denoted by $\text{cr}(A)$, such that there exists a generic polygon $P \in \mathbb{R}^2$ realizing A with $\text{cr}(P) = k$. Our first main result is the following theorem.

Theorem 1. *For a consistent angle sequence $A = (\alpha_0, \dots, \alpha_{n-1})$ in the plane, we have*

$$\text{cr}(A) = \begin{cases} 1 & \text{if } \sum_{i=0}^{n-1} \alpha_i = 0, \\ |j| - 1 & \text{if } \sum_{i=0}^{n-1} \alpha_i = 2j\pi \text{ and } j \neq 0. \end{cases}$$

Angle sequences in 3-space and spherical polygonal linkages. In $\mathbb{R}^d$, $d \geq 3$, the sign of a turning angle no longer plays a role: The *turning angle* of an oriented polygon P at v_i is in $(0, \pi)$, and an angle sequence $A = (\alpha_0, \dots, \alpha_{n-1})$ is in $(0, \pi)^n$. The unit-length direction vectors of the edges of P determine a spherical polygon P' . Note that the turning angles of P correspond to the spherical lengths of the segments of P' . It is not hard to see that this observation reduces the problem of realizability of A by a polygon in $\mathbb{R}^3$ to the problem of realizability of A by a spherical polygon, in the sense as defined next, that additionally contains the origin $\mathbf{0} = (0, 0, 0)$ in its convex hull.

Let $\mathbb{S}^2 \subset \mathbb{R}^3$ denote the unit 2-sphere. A *great circle* $C \subset \mathbb{S}^2$ is an intersection of $\mathbb{S}^2$ with a 2-dimensional hyperplane in $\mathbb{R}^3$ containing $\mathbf{0}$. A *spherical line segment* is a connected subset of a great circle that does not contain a pair of antipodal points of $\mathbb{S}^2$. The *length* of a spherical line segment ab equals the measure of the central angle subtended by ab . A *spherical polygon* $P \subset \mathbb{S}^2$ is a closed simple curve consisting of finitely many spherical segments; and a spherical polygon $P = (\mathbf{u}_0, \dots, \mathbf{u}_{n-1})$, $\mathbf{u}_i \in \mathbb{S}^2$, realizes an angle sequence $A = (\alpha_0, \dots, \alpha_{n-1})$ if the spherical segment $(\mathbf{u}_{i-1}, \mathbf{u}_i)$ has (spherical) length α_i , for every i . As usual, the *turning angle* of P at $\mathbf{u}_i$ is the angle in $[0, \pi]$ between the tangents to $\mathbb{S}^2$ at $\mathbf{u}_i$ that are co-planar with the great circles containing $(\mathbf{u}_i, \mathbf{u}_{i+1})$ and $(\mathbf{u}_i, \mathbf{u}_{i-1})$. Unlike for polygons in $\mathbb{R}^2$ and $\mathbb{R}^3$ we do not put any constraints on turning angles of spherical polygons in our results.

Regarding realizations of A by spherical polygons, we prove the following.

Theorem 2. *Let $A = (\alpha_0, \dots, \alpha_{n-1})$, $n \geq 3$, be an angle sequence. There exists a generic polygon $P \subset \mathbb{R}^3$ realizing A if and only if $\sum_{i=0}^{n-1} \alpha_i \geq 2\pi$ and there exists a spherical polygon $P' \subset \mathbb{S}^2$ realizing A . Furthermore, P can be constructed efficiently if P' is given.*

Theorem 3. *There exists a constructive weakly polynomial-time algorithm to test whether a given angle sequence $A = (\alpha_0, \dots, \alpha_{n-1})$ can be realized by a spherical polygon $P' \subset \mathbb{S}^2$.*

A simple exponential time algorithm for realizability of angles sequences by spherical polygons follows from a known characterization [3, Theorem 2.5], which also implies that the order of angles in A does not matter for the spherical realizability. The topology of the configuration spaces of spherical polygonal linkages have also been studied [16]. Independently, Streinu et al. [19, 23] showed that the configuration space of *noncrossing* spherical linkages is connected if $\sum_{i=0}^{n-1} \alpha_i \leq 2\pi$. However, these results do not seem to help prove Theorem 3.

The combination of Theorems 3 and 2 yields our second main result.

Theorem 4. *There exists a constructive weakly polynomial-time algorithm to test whether a given angle sequence $A = (\alpha_0, \dots, \alpha_{n-1})$ can be realized by a polygon $P \subset \mathbb{R}^3$.*

Organization. We prove Theorem 1 in Section 2 and Theorems 2, 3, and 4 in Section 3. We finish with concluding remarks in Section 4.

2 Crossing Minimization in the Plane

The first part of the following lemma gives a folklore necessary condition for the consistency of a sequence A . The condition is also sufficient except when $j = 0$. The second part follows from a result of Grünbaum and Shepard [11, Theorem 6], using a decomposition due to Wiener [28]. We provide a proof for the sake of completeness.




Figure 2: Splitting an oriented closed polygon P at a self-crossing point into 2 oriented closed polygons P' and P'' such that $\text{tn}(P) = \text{tn}(P') + \text{tn}(P'')$.

Lemma 1. *If an angle sequence $A = (\alpha_0, \dots, \alpha_{n-1})$ is consistent, then $\sum_{i=0}^{n-1} \alpha_i = 2j\pi$ for some $j \in \mathbb{Z}$. Furthermore, if $j \neq 0$ then $\text{cr}(A) \geq |j| - 1$.*

Proof. Let P be a polygon such that $\text{cr}(A) = \text{cr}(P)$. First, we prove that $\text{cr}(A) \geq |j| - 1 = |\text{tn}(P)| - 1$, by induction on $\text{cr}(P)$.

We consider the base case when $\text{cr}(P) = 0$. By Jordan-Schönflies curve theorem, P bounds a compact region homeomorphic to a disk. By a well-known fact, the internal angles at vertices of P sum up to $(n - 2)\pi$. Since A is consistent, $\sum_{i=0}^{n-1} \alpha_i = 2j\pi$, and thus, $(n - 2)\pi = \sum_{i=0}^{n-1} (\pi - \alpha_i) = (n - 2j)\pi$




Figure 3: Constructing a polygon P with $|\text{tn}(P)| - 1$ crossings.

or $(n - 2)\pi = \sum_{i=0}^{n-1} (\pi + \alpha_i) = (n + 2j)\pi$, depending on the orientation of the polygon. The claim follows since $|\text{tn}(P)| = j = 1$ in this case.

Refer to Fig. 2. In the inductive step, we have $\text{cr}(P) \geq 1$. By splitting P into two closed parts P' and P'' at a self-crossing, we obtain a pair of closed polygons such that $\text{tn}(P) = \text{tn}(P') + \text{tn}(P'')$. We have $\text{cr}(P) \geq 1 + \text{cr}(P') + \text{cr}(P'') \geq 1 + |\text{tn}(P')| - 1 + |\text{tn}(P'')| - 1 \geq |\text{tn}(P)| - 1$. Thus, the induction goes through, since both $\text{cr}(P')$ and $\text{cr}(P'')$ are smaller than $\text{cr}(P)$. $\square$

The following lemma shows that the lower bound in Lemma 1 is tight when $\alpha_i > 0$ for all $i \in \{0, \dots, n - 1\}$.

Lemma 2. *If $A = (\alpha_0, \dots, \alpha_{n-1})$ is an angle sequence such that $\sum_{i=0}^{n-1} \alpha_i = 2j\pi$, $j \neq 0$, and $\alpha_i > 0$, for all i , then $\text{cr}(A) \leq |j| - 1$.*

Proof. Refer to Figure 3. In three steps, we construct a polygon P realizing A with $|\text{tn}(P)| - 1$ self-crossings thereby proving $\text{cr}(A) \leq |j| - 1 = |\text{tn}(P)| - 1$. In the first step, we construct an oriented self-crossing-free polygonal line P' with $n + 2$ vertices, whose first and last (directed) edges are parallel to the positive x -axis, and whose internal vertices have turning angles $\alpha_0, \dots, \alpha_{n-1}$ in this order. We construct P' incrementally: The first edge has unit length starting from the origin; and every successive edge lies on a ray emanating from the endpoint of the previous edge. If the ray intersects neither the x -axis nor previous edges, then the next edge has unit length, otherwise its length is chosen to avoid any such intersection. In the second step, we prolong the last edge of P' until it creates the last self-intersection/crossing c and denote by P'' the resulting closed polygon composed of the part of P' from c to c via the prolonged part. By making the differences between the lengths of the edges of P' sufficiently large a prolongation of the last edge of P' has to eventually create at least one desired self-intersection. Hence, P'' is well-defined. Finally, we construct P realizing A from P'' by an appropriate modification of P'' in a small neighborhood of c without creating additional self-crossings. The number of self-crossings of P follows by the winding number of P w.r.t. to the point just a bit north from the end vertex of P' , which is j or $-j$. $\square$

To prove the upper bound in Theorem 1, it remains to consider the case that $A = (\alpha_0, \dots, \alpha_{n-1})$ contains both positive and negative angles. The crucial notion in the proof is that of an (essential) sign change of A which we define next. Let $A = (\alpha_0, \dots, \alpha_{n-1})$. Let $\beta_i = \sum_{j=0}^i \alpha_j \pmod{2\pi}$. Let $\mathbf{v}_i \in \mathbb{R}^2$ denote

the unit vector $(\cos \beta_i, \sin \beta_i)$. Hence, $\mathbf{v}_i$ is the direction vector of the $(i+1)$ -st edge of an oriented polygon P realizing A if the direction vector of the first edge of P is $(1, 0) \in \mathbb{R}^2$. As observed by Garg [10, Section 6], the consistency of A implies that $\mathbf{0}$ is a strictly positive convex combination of vectors $\mathbf{v}_i$, that is, there exists $\lambda_0, \dots, \lambda_{n-1} > 0$ such that $\sum_{i=0}^{n-1} \lambda \mathbf{v}_i = \mathbf{0}$ and $\sum_{i=0}^{n-1} \lambda_i = 1$.

The *sign change* of A is an index i such that $\alpha_i < 0$ and $\alpha_{i+1} > 0$, or vice versa, $\alpha_i > 0$ and $\alpha_{i+1} < 0$. Let $\text{sc}(A)$ denote the number of sign changes of A . The number of sign changes of A is even. A sign change i of a consistent angle sequence A is *essential* if $\mathbf{0}$ is not a strictly positive convex combination of $\{\mathbf{v}_0, \dots, \mathbf{v}_{i-1}, \mathbf{v}_{i+1}, \dots, \mathbf{v}_{n-1}\}$.

Lemma 3. *If $A = (\alpha_0, \dots, \alpha_{n-1})$ is a consistent angle sequence such that $\sum_{i=0}^{n-1} \alpha_i = 2j\pi$, $j \in \mathbb{Z}$, and all sign changes are essential, then $\text{cr}(A) \leq |j| - 1$.*

Proof. We distinguish between two cases depending on whether $\sum_{i=0}^{n-1} \alpha_i = 0$.

Case 1: $\sum_{i=0}^{n-1} \alpha_i = 0$. Since $\sum_{i=0}^{n-1} \alpha_i = 0$, we have $\text{sc}(A) \geq 2$. Since all sign changes are essential, for any two distinct sign changes $i \neq j$, we have $\mathbf{v}_i \neq \mathbf{v}_j$, therefore counting different vectors $\mathbf{v}_i$, where i is a sign change, is equivalent to counting sign changes. We show next that $\text{sc}(A) = 2$.

Suppose, to the contrary, that $\text{sc}(A) > 2$. Since $\text{sc}(A)$ is even, we have $\text{sc}(A) \geq 4$. Note that if $\mathbf{v}_i$ corresponds to an essential sign change i , then there is an open halfplane bounded by a line through the origin that contain only $\mathbf{v}_i$ in $\{\mathbf{v}_0, \dots, \mathbf{v}_{n-1}\}$. Thus, if i and i' are distinct essential sign changes, for any other essential sign change j we have that $\mathbf{v}_j$ is contained in a closed convex cone bounded by $-\mathbf{v}_i$ and $-\mathbf{v}_{i'}$ unless $-\mathbf{v}_i = \mathbf{v}_{i'}$. Hence, the only possibility for having 4 essential sign changes i, i', j' , and j' is if they satisfy $\mathbf{v}_i = -\mathbf{v}_{i'}$, $\mathbf{v}_j = -\mathbf{v}_{j'}$ and $\mathbf{v}_i \neq \pm \mathbf{v}_j$. Since all i, i', j , and j' are sign changes, there exists a fifth vector $\mathbf{v}_k$, which implies that one of i, i', j , and j' is not essential (contradiction).

Assume w.l.o.g. that j and $n-1$ are the only two essential sign changes. We have that $\mathbf{v}_j \neq -\mathbf{v}_{n-1}$: For otherwise, all the other $\mathbf{v}_i$'s different from $\mathbf{v}_j$ and $\mathbf{v}_{n-1}$ must be orthogonal to $\mathbf{v}_j$ and $\mathbf{v}_{n-1}$, since the sign changes j and $n-1$ are essential. Then due to the consistency of A , there exists a pair i and i' such that $\mathbf{v}_i = -\mathbf{v}_{i'}$. However, j and $n-1$ are the only sign changes, and thus, there exists k such that $\mathbf{v}_k \neq \pm \mathbf{v}_i$ (contradiction).




Figure 4: The case of exactly 2 sign changes $n-1$ and j , both of which are essential, when $\sum_{i=0}^{n-1} \alpha_i = 0$. Both missing parts of the polygon on the left are convex chains.

It follows that $\mathbf{v}_j$ and $\mathbf{v}_{n-1}$ are not collinear, and we have that the remaining $\mathbf{v}_i$'s belong to the closed convex cone bounded by $-\mathbf{v}_j$ and $-\mathbf{v}_{n-1}$. Refer to Figure 4. Thus, we may assume that (i) $\beta_{n-1} = 0$, (ii) the sign changes of A are $n-1$ and j , and (iii) $0 < \beta_0 < \dots < \beta_j$ and $\beta_j > \beta_{j+1} > \dots > \beta_{n-1} = 0$. Now, realizing A by a generic polygon with exactly 1 crossing between the line segments in the direction of $\mathbf{v}_j$ and $\mathbf{v}_{n-1}$ is a simple exercise.

Case 2: $\sum_{i=0}^{n-1} \alpha_i \neq 0$. We show that, unlike in the first case, none of the sign changes of A can be essential. Indeed, suppose j is an essential sign change, and as in Case 1, let $A' = (\alpha'_0, \dots, \alpha'_{n-2}) = (\alpha_0, \dots, \alpha_{j-1}, \alpha_j + \alpha_{j+1}, \dots, \alpha_{n-1})$ and $\beta'_i = \sum_{k=0}^i \alpha'_k \mod 2\pi$.

Furthermore, let $\mathbf{v}'_0, \dots, \mathbf{v}'_{n-2}$, where $\mathbf{v}'_i = (\cos \beta'_i, \sin \beta'_i)$. Since j is an essential sign change there exists $\mathbf{v} \neq \mathbf{0}$ such that $\langle \mathbf{v}, \mathbf{v}_j \rangle > 0$ and $\langle \mathbf{v}, \mathbf{v}'_i \rangle \leq 0$, for all i . Hence, by symmetry we assume that $0 \leq \beta'_i \leq \pi$, for all i . Then due to $-\pi < \alpha'_i < \pi$, we must have $\beta'_j = \sum_{i=0}^j \alpha'_i \mod 2\pi = \sum_{i=0}^j \alpha'_i$, which in turn implies, by Lemma 1, that $0 = \beta'_{n-2} = \sum_{i=0}^{n-2} \alpha'_i = \sum_{i=0}^{n-1} \alpha_i$ (contradiction).

We have shown that A has no sign changes. By Lemma 2, we have $\text{cr}(A) \leq |j| - 1$, which concludes the proof. $\square$

Proof of Theorem 1. The claimed lower bound $\text{cr}(A) \geq |j| - 1$ on the crossing number of A follows by Lemma 1, in the case when $j \neq 0$, and the result of Vijayan [27, Theorem 2] in the case when $j = 0$. It remains to prove the upper bound $\text{cr}(A) \leq |j| - 1$.

We proceed by induction on n . In the base case, we have $n = 3$. Then P is a triangle, $\sum_{i=0}^2 \alpha_i = \pm 2\pi$, and $\text{cr}(A) = 0$, as required. In the inductive step, assume $n \geq 4$, and that the claim holds for all shorter angle sequences. Let $A = (\alpha_0, \dots, \alpha_{n-1})$ be an angle sequence with $\sum_{i=0}^{n-1} \alpha_i = 2j\pi$.

If A has no sign changes or if all sign changes are essential, then Lemma 2 or Lemma 3 completes the proof. Otherwise, we have at least one nonessential sign change s . Let $A' = (\alpha'_0, \dots, \alpha'_{n-2}) = (\alpha_0, \dots, \alpha_{s-1}, \alpha_s + \alpha_{s+1}, \dots, \alpha_{n-1})$. Note that $\sum_{i=0}^{n-2} \alpha'_i = 2j\pi$. Since the sign change s is nonessential, $\mathbf{0}$ is a strictly positive convex combination of the β'_i 's, where $\beta'_i = \sum_{k=0}^i \alpha'_k \mod 2\pi$. Indeed, this follows from $\beta'_i = \beta_i$, for $i < k$, and $\beta'_i = \beta_{i+1}$, for $i \geq k$.




Figure 5: Re-introducing the j -th vertex to a polygon realizing A' in order to obtain a polygon realizing A .

Refer to Fig. 5. Hence, by applying the induction hypothesis we obtain a realization of A' as a generic polygon P' with $|j| - 1$ crossing. A generic

polygon realizing A is then obtained by modifying P in a small neighborhood of one of its vertices without introducing any additional crossing, similarly as in the paper by Guibas et al. [12]. $\square$

3 Realizing Angle Sequences in 3-Space

In this section, we describe a polynomial-time algorithm to decide whether an angle sequence $A = (\alpha_0, \dots, \alpha_{n-1})$ can be realized as a polygon in $\mathbb{R}^3$.

We remark that our problem can be expressed as solving a system of polynomial equations, where $3n$ variables describe the coordinates of the n vertices of P , and each of n equations is obtained by the cosine theorem applied for a vertex and two incident edges of P . However, it is not clear to us how to solve this system efficiently.

By Fenchel's theorem in differential geometry [9], the total curvature of any smooth curve in $\mathbb{R}^d$ is at least 2π . Fenchel's theorem has been adapted to closed polygons [24, Theorem 2.4], and it gives a necessary condition for an angle sequence A to have a realization in $\mathbb{R}^d$, for all $d \geq 2$.

$$\sum_{i=0}^{n-1} \alpha_i \geq 2\pi. \quad (1)$$

We show that a slightly stronger condition is both necessary and sufficient, hence it characterizes realizable angle sequences in $\mathbb{R}^3$.

Lemma 4. *Let $A = (\alpha_0, \dots, \alpha_{n-1})$, $n \geq 3$, be an angle sequence. There exists a polygon $P \subset \mathbb{R}^3$ realizing A if and only if there exists a spherical polygon $P' \subset \mathbb{S}^2$ realizing A such that $\mathbf{0} \in \text{relint}(\text{conv}(P'))$ (relative interior of $\text{conv}(P')$). Furthermore, P can be constructed efficiently if P' is given.*

Proof. Assume that an oriented polygon $P = (p_0, \dots, p_{n-1})$ realizes A in $\mathbb{R}^3$. Let $\mathbf{u}_i = (v_{i+1} - v_i) / \|v_{i+1} - v_i\| \in \mathbb{S}^2$ be the unit direction vectors of the edges of P according to its orientation. Then $P' = (\mathbf{u}_0, \dots, \mathbf{u}_{n-1})$ is a spherical polygon that realizes A . Suppose, for the sake of contradiction, that $\mathbf{0}$ is not in the relative interior of $\text{conv}(P')$. Then there is a plane H that separates $\mathbf{0}$ and P' , that is, if $\mathbf{n}$ is the normal vector of H , then $\langle \mathbf{n}, \mathbf{u}_i \rangle > 0$ for all $i \in \{0, \dots, n-1\}$. This implies $\langle \mathbf{n}, (v_{i+1} - v_i) \rangle > 0$ for all i , hence $\langle \mathbf{n}, \sum_{i=1}^{n-1} (v_{i+1} - v_i) \rangle > 0$, which contradicts the fact that $\sum_{i=1}^{n-1} (v_{i+1} - v_i) = \mathbf{0}$, and $\langle \mathbf{n}, \mathbf{0} \rangle = 0$.

Conversely, assume that there is a spherical polygon P' that realizes A , with edge lengths $\alpha_0, \dots, \alpha_{n-1}$. If all vertices of P' lie in a great circle, then $\mathbf{0} \in \text{relint}(\text{conv}(P'))$ implies $\sum_{i=0}^{n-1} \alpha_i \geq 2\pi$, and Theorem 1 completes the proof.

Otherwise we may assume that $\mathbf{0} \in \text{int}(\text{conv}(P'))$. By Carathéodory's theorem [17, Theorem 1.2.3], P' has 4 vertices whose convex combination is the origin $\mathbf{0}$. Then we can express $\mathbf{0}$ as a strictly positive convex combination of all vertices of P' . The coefficients in the convex combination encode the lengths of the edges of a polygon P realizing A , which concludes the proof in this case.

We now show how to compute strictly positive coefficients in strongly polynomial time. Let $\mathbf{c} = \frac{1}{n} \sum_{i=0}^{n-1} \mathbf{u}_i$ be the centroid of the vertices of P' . If $\mathbf{c} = \mathbf{0}$, we are done. Otherwise, we can find a tetrahedron $T = \text{conv}\{\mathbf{u}_{i_0}, \dots, \mathbf{u}_{i_3}\}$ such that $\mathbf{0} \in T$ and such that the ray from $\mathbf{0}$ in the direction $-\mathbf{c}$ intersects $\text{int}(T)$, by solving an LP feasibility problem in $\mathbb{R}^3$. By computing the intersection of the ray with the faces of T , we find the maximum $\mu > 0$ such that $-\mu\mathbf{c} \in \partial T$ (the boundary of T). We have $-\mu\mathbf{c} = \sum_{j=0}^3 \lambda_j \mathbf{u}_{i_j}$ and $\sum_{j=0}^3 \lambda_j = 1$ for suitable coefficients $\lambda_j \geq 0$. Now $\mathbf{0} = \mu\mathbf{c} - \mu\mathbf{c} = \frac{\mu}{n} \sum_{i=0}^{n-1} \mathbf{u}_i + \sum_{j=0}^3 \lambda_j \mathbf{u}_{i_j}$ is a strictly positive convex combination of the vertices of P' . $\square$

It is easy to find an angle sequence A that satisfies (1) but does not correspond to a spherical polygon P' . Consider, for example, $A = (\pi - \varepsilon, \pi - \varepsilon, \pi - \varepsilon, \varepsilon)$, for some small $\varepsilon > 0$. Points in $\mathbb{S}^2$ at (spherical) distance $\pi - \varepsilon$ are nearly antipodal. Hence, the endpoints of a polygonal chain $(\pi - \varepsilon, \pi - \varepsilon, \pi - \varepsilon)$ are nearly antipodal, as well, and cannot be connected by an edge of (spherical) length ε . Thus a spherical polygon cannot realize A .

Algorithms. In the remainder of this section, we show how to find a realization $P \subset \mathbb{R}^3$ or report that none exists, in polynomial time. Our first concern is to decide whether an angle sequence is realizable by a spherical polygon.

Theorem 3. *There exists a constructive weakly polynomial-time algorithm to test whether a given angle sequence $A = (\alpha_0, \dots, \alpha_{n-1})$ can be realized by a spherical polygon $P' \subset \mathbb{S}^2$.*

Proof of Theorem 3. Let $A = (\alpha_0, \dots, \alpha_{n-1}) \in (0, \pi)^n$ be a given angle sequence. Let $\mathbf{n} = (0, 0, 1) \in \mathbb{S}^2$ (the north pole). For $i \in \{0, 1, \dots, n-1\}$ let $U_i \subseteq \mathbb{S}^2$ be the locus of the end vertices $\mathbf{u}_i$ of all (spherical) polygonal lines $P'_i = (\mathbf{n}, \mathbf{u}_0, \dots, \mathbf{u}_i)$ with edge lengths $\alpha_0, \dots, \alpha_{i-1}$. It is clear that A is realizable by a spherical polygon P' iff $\mathbf{n} \in U_{n-1}$.

Note that for all $i \in \{0, \dots, n-1\}$, the set U_i is invariant under rotations about the z -axis, since $\mathbf{n}$ is a fixed point and rotations are isometries. We show how to compute the sets U_i , $i \in \{0, \dots, n-1\}$, efficiently.

We define a *spherical zone* as a subset of $\mathbb{S}^2$ between two horizontal planes (possibly, a circle, a spherical cap, or a pole). Recall the parameterization of $\mathbb{S}^2$ using spherical coordinates (cf. Figure 6 (left)): for every $\mathbf{v} \in \mathbb{S}^2$, $\mathbf{v}(\psi, \varphi) = (\sin \psi \sin \varphi, \cos \psi \sin \varphi, \cos \varphi)$, with longitude $\psi \in [0, 2\pi)$ and polar angle $\varphi \in [0, \pi]$, where the *polar angle* φ is the angle between $\mathbf{v}$ and $\mathbf{n}$. Using this parameterization, a spherical zone is a Cartesian product $[0, 2\pi) \times I$ for some circular arc $I \subset [0, \pi]$. In the remainder of the proof, we associate each spherical zone with such a circular arc I .

We define additions and subtraction on polar angles $\alpha, \beta \in [0, \pi]$ by

$$\alpha \oplus \beta = \min\{\alpha + \beta, 2\pi - (\alpha + \beta)\}, \quad \alpha \ominus \beta = \max\{\alpha - \beta, \beta - \alpha\};$$

see Figure 6 (right). (This may be interpreted as addition mod 2π , restricted to the quotient space defined by the equivalence relation $\varphi \sim 2\pi - \varphi$.)




Figure 6: Parametrization of the unit vectors (left). Circular arc $C_{i+1}(\varphi)$ (right).

We show that U_i is a spherical zone for all $i \in \{0, \dots, n-1\}$, and show how to compute the intervals $I_i \subset [0, \pi]$ efficiently. First note that U_0 is a circle at (spherical) distance α_0 from $\mathbf{n}$, hence U_0 is a spherical zone with $I_0 = [\alpha_0, \alpha_0]$.

Assume that U_i is a spherical zone associated with $I_i \subset [0, \pi]$. Let $\mathbf{u}_i \in U_i$, where $\mathbf{u}_i = \mathbf{v}(\psi, \varphi)$ with $\psi \in [0, 2\pi)$ and $\varphi \in I_i$. By the definition U_i , there exists a polygonal line $(\mathbf{n}, \mathbf{u}_0, \dots, \mathbf{u}_i)$ with edge lengths $\alpha_0, \dots, \alpha_i$. The locus of points in $\mathbb{S}^2$ at distance α_{i+1} from $\mathbf{u}_i$ is a circle; the polar angles of the points in the circle form an interval $C_{i+1}(\varphi)$. Specifically (see Figure 6 (right)), we have

$$C_{i+1}(\varphi) = [\min\{\varphi \ominus \alpha_{i+1}, \varphi \oplus \alpha_{i+1}\}, \max\{\varphi \ominus \alpha_{i+1}, \varphi \oplus \alpha_{i+1}\}].$$

By rotational symmetry, $U_{i+1} = [0, 2\pi) \times I_{i+1}$, where $I_{i+1} = \bigcup_{\varphi \in I_i} C_{i+1}(\varphi)$. Consequently, $I_{i+1} \subset [0, \pi]$ is connected, and hence, I_{i+1} is an interval. Therefore U_{i+1} is a spherical zone. As $\varphi \oplus \alpha_{i+1}$ and $\varphi \ominus \alpha_{i+1}$ are piecewise linear functions of φ , we can compute I_{i+1} using $O(1)$ arithmetic operations.

We can construct the intervals $I_0, \dots, I_{n-1} \subset [0, \pi]$ as described above. If $0 \notin I_{n-1}$, then $\mathbf{n} \notin U_{n-1}$ and A is not realizable. Otherwise, we can compute the vertices of a spherical realization $P' \subset \mathbb{S}^2$ by backtracking. Put $\mathbf{u}_{n-1} = \mathbf{n} = (0, 0, 1)$. Given $\mathbf{u}_i = \mathbf{v}(\psi, \varphi)$, we choose $\mathbf{u}_{i-1}$ as follows. Let $\mathbf{u}_{i-1}$ be $\mathbf{v}(\psi, \varphi \oplus \alpha_i)$ or $\mathbf{v}(\psi, \varphi \ominus \alpha_i)$ if either of them is in U_{i-1} (break ties arbitrarily). Else the spherical circle of radius α_i centered at $\mathbf{u}_i$ intersects the boundary of U_{i-1} , and then we choose $\mathbf{u}_i$ to be an arbitrary such intersection point. The decision algorithm (whether $0 \in I_{n-1}$) and the backtracking both use $O(n)$ arithmetic operations. $\square$

Enclosing the Origin. Theorem 3 provides an efficient algorithm to test whether an angle sequence can be realized by a spherical polygon, however, Lemma 4 requires a spherical polygon P' whose convex hull contains the origin. We show that this is always possible if a realization exists and $\sum_{i=0}^{n-1} \alpha_i \geq 2\pi$. The general strategy in the inductive proof of this claim is to gradually modify P' by changing the turning angle at one of its vertices to 0. This allows us to reduce the number of vertices of P' and apply induction.

Lemma 5. *Given a spherical polygon P' realizing an angle sequence $A = (\alpha_0, \dots, \alpha_{n-1})$, $n \geq 3$, with $\sum_{i=0}^{n-1} \alpha_i \geq 2\pi$, we can compute in polynomial time a spherical polygon P'' realizing A such that $\mathbf{0} \in \text{relint}(\text{conv}(P''))$.*




Figure 7: The spherical zone U_1 (or U_1^{2-}) containing $\mathbf{u}_1$ corresponding to I_1 .

We introduce some terminology for spherical polygonal linkages with one fixed endpoint. Let $P' = (\mathbf{u}_0, \dots, \mathbf{u}_{n-1})$ be a polygon in $\mathbb{S}^2$ that realizes an angle sequence $A = (\alpha_0, \dots, \alpha_{n-1})$; we do not assume $\sum_{i=1}^{n-1} \alpha_i \geq 2\pi$. Denote by U_i^{j-} the locus of the endpoints $\mathbf{u}'_i \in \mathbb{S}^2$ of all (spherical) polygonal lines $(\mathbf{u}_{i-j}, \mathbf{u}'_{i-j+1}, \dots, \mathbf{u}'_i)$, where the first vertex is fixed at $\mathbf{u}_{i-j}$, and the edge lengths are $\alpha_{i-j}, \dots, \alpha_i$. Similarly, denote by U_i^{j+} the locus of the endpoints $\mathbf{u}'_i \in \mathbb{S}^2$ of all (spherical) polygonal lines $(\mathbf{u}_{i+j}, \mathbf{u}'_{i+j-1}, \dots, \mathbf{u}'_i)$ with edge lengths $\alpha_{i+j+1}, \dots, \alpha_{i+1}$. Due to rotational symmetry about the line passing through $\mathbf{u}_{i-j}$ and $\mathbf{0}$, both U_i^{j-} and U_i^{j+} are a *spherical zone* (a subset of $\mathbb{S}^2$ bounded by two parallel circles), possibly just a circle, or a cap, or a point. In particular, the distance between $\mathbf{u}_i$ and any boundary component (circle) of U_i^{j-} or U_i^{j+} is the same; see Figure 7.

If U_i^{2+} is bounded by two circles, let T_i^{2+} and B_i^{2+} denote the two boundary circles such that $\mathbf{u}_i$ is closer to T_i^{2+} than to B_i^{2+} . If U_i^{2+} is a cap, let T_i^{2+} denote the boundary of U_i^{2+} , and let B_i^{2+} denote the center of U_i^{2+} . We define T_i^{2-} and B_i^{2-} analogously.

The vertex $\mathbf{u}_i$ of P' is a *spur* of P' if the segments $\mathbf{u}_i \mathbf{u}_{i+1}$ and $\mathbf{u}_i \mathbf{u}_{i-1}$ overlap (equivalently, the turning angle of P' at $\mathbf{u}_i$ is π). We use the following simple but crucial observation.

Observation 1. *Assume that $n \geq 4$ and U_i^{2+} is neither a circle nor a point. The turning angle of P' at $\mathbf{u}_{i+1}$ is 0 iff $\mathbf{u}_i \in B_i^{2+}$; and $\mathbf{u}_{i+1}$ is a spur of P' iff $\mathbf{u}_i \in T_i^{2+}$.*

A crucial technical tool in the proof of Lemma 5 is the following lemma based on Observation 1.

Lemma 6. *Let P' be a spherical polygon $(\mathbf{u}_0, \dots, \mathbf{u}_{n-1})$, $n \geq 4$, that realizes an angle sequence $A = (\alpha_0, \dots, \alpha_{n-1})$. Then there exists a spherical polygon $P'' = (\mathbf{u}_0, \dots, \mathbf{u}_{i-1}, \mathbf{u}'_i, \mathbf{u}'_{i+1}, \mathbf{u}_{i+2}, \dots, \mathbf{u}_{n-1})$ that also realizes A such that the turning angle at $\mathbf{u}_{i-1}$ is 0, or the turning angle at $\mathbf{u}_{i+1}$ is 0 or π .*

Proof. If $n \geq 4$, Observation 1 allows us to move vertices $\mathbf{u}_i$ and $\mathbf{u}_{i+1}$ so that the turning angle at $\mathbf{u}_{i-1}$ drops to 0, or the turning angle at $\mathbf{u}_{i+1}$ changes to 0 or π , while all other vertices of P' remain fixed. Indeed, one of the following three options holds: $U_i^{1-} \subseteq U_i^{2+}$, $U_i^{1-} \cap B_i^{2+} \neq \emptyset$, or $U_i^{1-} \cap T_i^{2+} \neq \emptyset$. If $U_i^{1-} \subseteq U_i^{2+}$, then by Observation 1 there exists $\mathbf{u}'_i \in U_i^{1-} \cap B_i^{2-} \cap U_i^{2+}$. Since $\mathbf{u}'_i \in U_i^{2+}$ there exists $\mathbf{u}'_{i+1} \in U_{i+1}^{1+}$ such that $P'' = (\mathbf{u}_0, \dots, \mathbf{u}_{i-1}, \mathbf{u}'_i, \mathbf{u}'_{i+1}, \mathbf{u}_{i+2}, \dots, \mathbf{u}_{n-1})$ realizes A and the turning angle at $\mathbf{u}_{i-1}$ equals 0. Similarly, if there exists $\mathbf{u}'_i \in U_i^{1-} \cap B_i^{2+}$ or $\mathbf{u}'_i \in U_i^{1-} \cap T_i^{2+}$, then there exists $\mathbf{u}'_{i+1} \in U_{i+1}^{1+}$ such that P'' as above realizes A with the turning angle at $\mathbf{u}_{i+1}$ equal to 0 or π respectively. $\square$

Proof of Lemma 5. We proceed by induction on the number of vertices of P' . In the basis step, we have either $n = 3$. In this case, P' is a spherical triangle. The length of every spherical triangle is at most 2π , contradicting the assumption that $\sum_{i=0}^{n-1} \alpha_i > 2\pi$. Hence the claim vacuously holds.

In the induction step, assume that $n \geq 4$ and the claim holds for smaller values of n . Assume $\mathbf{0} \notin \text{relint}(\text{conv}(P'))$, otherwise the proof is complete. We distinguish between several cases.

Case 1: a path of consecutive edges lying in a great circle contains a half-circle. We may assume w.l.o.g. that at least one endpoint of the half-circle is a vertex of P' . Since the length of each edge is less than π , the path that contains a half-circle has at least 2 edges.

Case 1.1: both endpoints of the half-circle are vertices of P' . Assume w.l.o.g., that the two endpoints of the half-circle are $\mathbf{u}_i$ and $\mathbf{u}_j$, for some $i < j$. These vertices decompose P' into two polylines, P'_1 and P'_2 . We rotate P'_2 about the line through $\mathbf{u}_i \mathbf{u}_j$ so that the turning angle at $\mathbf{u}_i$ is a suitable value in $[-\varepsilon, +\varepsilon]$ as follows. First, set the turning angle at $\mathbf{u}_i$ to be 0. If the resulting polygon P'' is contained in a great circle or $\mathbf{0} \in \text{int}(\text{conv}(P''))$ we are done. Else, P'' is contained in a hemisphere H bounded by the great circle through $\mathbf{u}_{i-1} \mathbf{u}_i \mathbf{u}_{i+1}$. In this case, we perturb the turning angle at $\mathbf{u}_i$ so that $\mathbf{u}_{i+1}$ is not contained in H thereby achieving $\mathbf{0} \in \text{int}(\text{conv}(P''))$.

Case 1.2: only one endpoint of the half-circle is a vertex of P' . Let $P'_1 = (\mathbf{u}_i, \dots, \mathbf{u}_j)$ be the longest path in P' that contains a half-circle, and lies in a great circle. Since $\mathbf{0} \notin \text{relint}(\text{conv}(P'))$, the polygon P' is contained in a hemisphere H bounded by the great circle ∂H that contains P'_1 , but P' is not contained in ∂H . By construction, $\mathbf{u}_{j+1} \notin \partial H$. In order to make the proof in this case easier, we introduce the following assumption. If a part P_0 of P' between two antipodal/identical end vertices that belong ∂H is contained in a great circle, w.l.o.g. we assume that P_0 is contained in ∂H .

W.l.o.g. $j = 0$, and we let j' be the smallest value such that $\mathbf{u}_{j'} \in \partial H$. By $\mathbf{0} \notin \text{relint}(\text{conv}(P'))$, $\mathbf{u}_0, \dots, \mathbf{u}_{j'} \in H$. We can perturb the polygon P'

into a new polygon $P'' = (\mathbf{u}'_0, \dots, \mathbf{u}'_{j'-1}, \mathbf{u}_{j'}, \dots, \mathbf{u}_{n-1})$ realizing A so that $\mathbf{0} \in \text{int}(\text{conv}(P''))$. Indeed, by Observation 1, $\mathbf{u}_0 \notin \partial U_0^{2+}$. Therefore since $(\mathbf{u}_0, \dots, \mathbf{u}_{j'})$ is not contained in a great circle by our assumption, by (a multiple use) of Observation 1, we choose $\mathbf{u}'_0, \dots, \mathbf{u}'_{j'-1}$, so that $\mathbf{u}'_0 \notin H$, and $\mathbf{u}'_1, \dots, \mathbf{u}'_{j'-1} \in \text{relint}(H)$, thereby achieving $\mathbf{0} \in \text{int}(\text{conv}(P''))$.

Case 2: the turning angle of P' is 0 at some vertex $\mathbf{u}_i$. By supressing the vertex $\mathbf{u}_i$, we obtain a spherical polygon Q' on $n - 1$ vertices that realizes the sequence $(\alpha_0, \dots, \alpha_{i-2}, \alpha_{i-1} + \alpha_i, \alpha_{i+1}, \dots, \alpha_{n-1})$ unless $\alpha_{i-1} + \alpha_i \geq \pi$, but then we are in Case 1. By induction, this sequence has a realization Q'' such that $\mathbf{0} \in \text{relint}(\text{conv}(Q''))$. Subdivision of the edge of length $\alpha_{i-1} + \alpha_i$ producers a realization P'' of A such that $\mathbf{0} \in \text{relint}(\text{conv}(Q'')) = \text{relint}(\text{conv}(P''))$.

Case 3: there is no path of consecutive edges lying in a great circle and containing a half-circle, and no turning angle is 0.

Case 3.1: $n = 4$. We claim that $U_0^{2+} \cap U_0^{2-}$ contains B_0^{2-} or B_0^{2+} . By Observation 1, this immediately implies that we can change one turning angle to 0 and proceed to Case 1.

To prove the claim, note that $U_0^{2+} \cap U_0^{2-} \neq \emptyset$ and $-2 \equiv 2 \pmod{4}$, and hence the circles T_0^{2-} , T_0^{2+} , B_0^{2-} , and B_0^{2+} are all parallel since they are all orthogonal to $\mathbf{u}_2$. Thus, by symmetry there are two cases to consider depending on whether $U_0^{2+} \subseteq U_0^{2-}$. If $U_0^{2+} \subseteq U_0^{2-}$, then $B_0^{2+} \subset U_0^{2+} \cap U_0^{2-}$. Else $U_0^{2+} \cap U_0^{2-}$ contains B_0^{2+} or B_0^{2-} , whichever is closer to $\mathbf{u}_2$, which concludes the proof of this case.

Case 3.2: $n \geq 5$. Choose $i \in \{0, \dots, n - 1\}$ so that α_{i+2} is a minimum angle in A . Note that U_i^{2+} is neither a circle nor a point since that would mean that $\mathbf{u}_{i+2}$ and $\mathbf{u}_{i+1}$, or $\mathbf{u}_i$ and $\mathbf{u}_{i+1}$ are antipodal, which is impossible.

We apply Lemma 6 and obtain a spherical polygon

$$P'' = (\mathbf{u}_0, \dots, \mathbf{u}_{i-1}, \mathbf{u}'_i, \mathbf{u}'_{i+1}, \mathbf{u}_{i+2}, \dots, \mathbf{u}_{n-1}).$$

If the turning angle of P'' at $\mathbf{u}_{i-1}$ or $\mathbf{u}'_{i+1}$ equals to 0, we proceed to Case 2. Otherwise, the turning angle of P'' at $\mathbf{u}'_{i+1}$ equals π . In other words, we introduce a spur at $\mathbf{u}'_{i+1}$. If $\alpha_{i+1} = \alpha_{i+2}$ we can make the turning angle of P'' at $\mathbf{u}_{i+2}$ equal to 0 by rotating the overlapping segments $(\mathbf{u}'_{i+1}, \mathbf{u}_{i+2})$ and $(\mathbf{u}'_{i+1}, \mathbf{u}'_i)$ around $\mathbf{u}_{i+2} = \mathbf{u}'_i$ and proceed to Case 2. Otherwise, we have $\alpha_{i+2} < \alpha_{i+1}$ by the choice of i . Let Q' denote an auxiliary polygon realizing $(\alpha_0, \dots, \alpha_i, \alpha_{i+1} - \alpha_{i+2}, \alpha_{i+3}, \dots, \alpha_{n-1})$. We construct Q' from P'' by cutting off the overlapping segments $(\mathbf{u}'_{i+1}, \mathbf{u}_{i+2})$ and $(\mathbf{u}'_{i+1}, \mathbf{u}'_i)$. We apply Lemma 6 to Q' thereby obtaining another realization

$$Q'' = (\mathbf{u}_0, \dots, \mathbf{u}_{i-1}, \mathbf{u}''_i, \mathbf{u}''_{i+1}, \mathbf{u}_{i+3}, \dots, \mathbf{u}_{n-1}).$$

We re-introduce the cut off part to Q'' at $\mathbf{u}''_{i+1}$ as an extension of length α_{i+2} of the segment $\mathbf{u}''_i \mathbf{u}''_{i+1}$, whose length in Q'' is $\alpha_{i+1} - \alpha_{i+2} > 0$, in order to recover a realization of A by the following polygon

$$R' = (\mathbf{u}_0, \dots, \mathbf{u}_{i-1}, \mathbf{u}''_i, \mathbf{u}''_{i+1}, \mathbf{u}''_{i+2}, \mathbf{u}_{i+3}, \dots, \mathbf{u}_{n-1}).$$

If the turning angle of Q'' at $\mathbf{u}_{i-1}$ equals 0, the same holds for R' and we proceed to Case 2. If the turning angle of Q'' at $\mathbf{u}''_{i+1}$ equals π , then the turning angle of

R' at $\mathbf{u}_{i+1}''$ equals 0 and we proceed to Case 2. Finally, if the turning angle of Q'' at $\mathbf{u}_{i+1}''$ equals 0, then R' has a pair of consecutive spurs at $\mathbf{u}_{i+1}''$ and $\mathbf{u}_{i+2}''$, that is, a so-called “crimp.” We may assume w.l.o.g. that $\alpha_{i+3} < \alpha_{i+1}$. Also we assume that the part $(\mathbf{u}_i'', \mathbf{u}_{i+1}'', \mathbf{u}_{i+2}'', \mathbf{u}_{i+3}'')$ of R' does not contain a pair of antipodal points, since otherwise we proceed to Case 1. Since $(\mathbf{u}_i'', \mathbf{u}_{i+1}'', \mathbf{u}_{i+2}'', \mathbf{u}_{i+3}'')$ does not contain a pair of antipodal points, $|(\mathbf{u}_i'', \mathbf{u}_{i+3}'')| = \alpha_{i+1} + \alpha_{i+3} - \alpha_{i+2}$. It follows that

$$\begin{aligned} |(\mathbf{u}_i'', \mathbf{u}_{i+3}'')| + |(\mathbf{u}_i'', \mathbf{u}_{i+1}'')| + |(\mathbf{u}_{i+1}'', \mathbf{u}_{i+2}'')| + |(\mathbf{u}_{i+2}'', \mathbf{u}_{i+3}'')| = \\ \alpha_{i+1} + \alpha_{i+3} - \alpha_{i+2} + \alpha_{i+1} + \alpha_{i+2} + \alpha_{i+3} = 2(\alpha_{i+1} + \alpha_{i+3}) \end{aligned}$$

If $\alpha_{i+3} + \alpha_{i+1} < \pi$, then the 3 angles α_{i+1} , $\alpha_{i+2} + \alpha_{i+3}$, and $|(\mathbf{u}_i'', \mathbf{u}_{i+3}'')|$ are all less than π . Moreover, their sum, which is equal to $2(\alpha_{i+3} + \alpha_{i+1})$, is less than 2π , and they satisfy the triangle inequalities. Therefore we can turn the angle at $\mathbf{u}_{i+2}''$ to 0, by replacing the path $(\mathbf{u}_i'', \mathbf{u}_{i+1}'', \mathbf{u}_{i+2}'', \mathbf{u}_{i+3}'')$ on R' by a pair of segments of lengths α_{i+1} and $\alpha_{i+2} + \alpha_{i+3}$.

Otherwise, $\alpha_{i+3} + \alpha_{i+1} \geq \pi$, and thus,

$$|(\mathbf{u}_i'', \mathbf{u}_{i+3}'')| + |(\mathbf{u}_i'', \mathbf{u}_{i+1}'')| + |(\mathbf{u}_{i+1}'', \mathbf{u}_{i+2}'')| + |(\mathbf{u}_{i+2}'', \mathbf{u}_{i+3}'')| \geq 2\pi.$$

In this case, we can apply the induction hypothesis to the closed spherical polygon $(\mathbf{u}_i'', \mathbf{u}_{i+1}'', \mathbf{u}_{i+2}'', \mathbf{u}_{i+3}'')$. In the resulting realization S' , that is w.l.o.g. fixing $\mathbf{u}_i''$ and $\mathbf{u}_{i+3}''$, we replace the segment $(\mathbf{u}_i'', \mathbf{u}_{i+3}'')$ by the remaining part of R' between $\mathbf{u}_i''$ and $\mathbf{u}_{i+3}''$. Let R'' denote the resulting realization of A . If S' is not contained in a great circle then $\mathbf{0} \in \text{int}(\text{conv}(S')) \subseteq \text{int}(\text{conv}(R''))$, and we are done. Otherwise, $S' \setminus (\mathbf{u}_{i+3}'', \mathbf{u}_i'')$ contains a pair of antipodal points on a half-circle. The same holds for R'' , and we proceed to Case 1, which concludes the proof. $\square$

The combination of Theorem 3 with Lemmas 4–5 yields Theorems 2 and 4. The inductive proof of Lemma 6 can be turned into an algorithm with a polynomial running time in n if every arithmetic operation is assumed to be carried out in $O(1)$ time. Nevertheless, we get only a weakly polynomial running time, since we are unable to guarantee a polynomial size encoding of the numerical values that are computed in the process of constructing a spherical polygon realizing A that contains $\mathbf{0}$ in its convex hull in the proof of Lemma 6.

4 Conclusion

We devised efficient algorithms to realize a consistent angle cycle with the minimum number of crossings in 2D. In 3D, we can test efficiently whether a given angle sequence is realizable, and find a realization if one exists. However, it remains an open problem to find an efficient algorithm that computes the minimum number of crossings in generic realizations. There exist sequences that are realizable, but every generic realization has crossings. It is not difficult to see that crossings are unavoidable only if every 3D realization of A is contained

in a plane, which is the case, for example, when $A = (\pi - \varepsilon, \dots, \pi - \varepsilon, (n - 1)\varepsilon)$ for $n \geq 5$ odd. Thus, an efficient algorithm for this problem would follow by Theorem 1, once one can test efficiently whether A admits a fully 3D realization.

Can our results in $\mathbb{R}^2$ or $\mathbb{R}^3$ be extended to broader interesting classes of graphs? A natural analog of our problem in $\mathbb{R}^3$ would be a construction of triangulated spheres with prescribed dihedral angles, discussed in a recent paper by Amenta and Rojas [1]. For convex polyhedra, Mazzeo and Montcouquiol [18] proved, settling Stokers' conjecture, that dihedral angles determine face angles.

Theorem 3 gave an efficient algorithm to test whether a given angle sequence A can be realized by a spherical polygon $P' \subset \mathbb{S}^2$. We wonder whether every realizable sequence A has a noncrossing realization, or possibly a noncrossing realization whose convex hull contains the origin (when $\sum_{i=0}^{n-1} \alpha_i \geq 2\pi$). If the answer is positive, can such realizations be computed efficiently? We do not know whether a realization $P \subset \mathbb{R}^3$ corresponding to a spherical realization $P' \subset \mathbb{S}^2$ (according to the method in the proof of Lemma 4) has any interesting properties when P' has no self-intersections.

References

- [1] Nina Amenta and Carlos Rojas. Dihedral deformation and rigidity. *Computational Geometry: Theory and Applications*, 2020. In press.
- [2] Michael A. Bekos, Henry Förster, and Michael Kaufmann. On smooth orthogonal and octilinear drawings: Relations, complexity and Kandinsky drawings. *Algorithmica*, 81(5):2046–2071, 2019.
- [3] R Buckman and N Schmitt. Spherical polygons and unitarization. *Preprint*, 2002. <http://www.gang.umass.edu/reu/2002/polygon.html>.
- [4] Katie Clinch, Bill Jackson, and Peter Keevash. Global rigidity of direction-length frameworks. *Journal of Combinatorial Theory, Series B*, 145:145–168, 2020.
- [5] Giuseppe Di Battista, Ethan Kim, Giuseppe Liotta, Anna Lubiw, and Sue Whitesides. The shape of orthogonal cycles in three dimensions. *Discrete Computational Geometry*, 47(3):461–491, 2012.
- [6] Giuseppe Di Battista and Luca Vismara. Angles of planar triangular graphs. *SIAM Journal on Discrete Mathematics*, 9(3):349–359, 1996.
- [7] Yann Disser, Matúš Mihalák, and Peter Widmayer. A polygon is determined by its angles. *Computational Geometry: Theory and Applications*, 44:418–426, 2011.
- [8] Tobin A Driscoll and Stephen A Vavasis. Numerical conformal mapping using cross-ratios and Delaunay triangulation. *SIAM Journal on Scientific Computing*, 19(6):1783–1803, 1998.

- [9] Werner Fenchel. On the differential geometry of closed space curves. *Bulletin of the American Mathematical Society*, 57(1):44–54, 1951.
- [10] Ashim Garg. New results on drawing angle graphs. *Computational Geometry: Theory and Applications*, 9(1-2):43–82, 1998.
- [11] Branko Grünbaum and Geoffrey Colin Shephard. Rotation and winding numbers for planar polygons and curves. *Transactions of the American Mathematical Society*, 322(1):169–187, 1990.
- [12] Leonidas Guibas, John Hersberger, and Subhash Suri. Morphing simple polygons. *Discrete & Computational Geometry*, 24(1):1–34, 2000.
- [13] Bill Jackson and Tibor Jordán. Globally rigid circuits of the direction-length rigidity matroid. *Journal of Combinatorial Theory, Series B*, 100(1):1–22, 2010.
- [14] Bill Jackson and Peter Keevash. Necessary conditions for the global rigidity of direction-length frameworks. *Discrete Computational Geometry*, 46(1):72–85, 2011.
- [15] Audrey Lee-St. John and Ileana Streinu. Angular rigidity in 3D: Combinatorial characterizations and algorithms. In *Proc. 21st Canadian Conference on Computational Geometry (CCCG)*, pages 67–70, 2009.
- [16] Michael Kapovich and John J. Millson. On the moduli space of a spherical polygonal linkage. *Canadian Mathematical Bulletin*, 42:307–320, 1999.
- [17] Jiří Matoušek. *Lectures on Discrete Geometry*, volume 212. Springer Science & Business Media, 2013.
- [18] Rafe Mazzeo and Grégoire Montcouquiol. Infinitesimal rigidity of cone-manifolds and the Stoker problem for hyperbolic and Euclidean polyhedra. *Journal of Differential Geometry*, 87(3):525–576, 2011.
- [19] Gaiane Panina and Ileana Streinu. Flattening single-vertex origami: The non-expansive case. *Computational Geometry: Theory and Applications*, 43(8):678–687, 2010.
- [20] Maurizio Patrignani. Complexity results for three-dimensional orthogonal graph drawing. *J. Discrete Algorithms*, 6(1):140–161, 2008.
- [21] Franco Saliola and Walter Whiteley. Constraining plane configurations in CAD: circles, lines, and angles in the plane. *SIAM Journal on Discrete Mathematics*, 18(2):246–271, 2004.
- [22] Jack Snoeyink. Cross-ratios and angles determine a polygon. *Discrete & Computational Geometry*, 22(4):619–631, 1999.

- [23] Ileana Streinu and Walter Whiteley. Single-vertex origami and spherical expansive motions. In *on Japanese Conference on Discrete and Computational Geometry*, volume 3742 of *LNCS*, pages 161–173. Springer, 2004.
- [24] John M. Sullivan. Curves of finite total curvature. In Alexander I. Bobenko, John M. Sullivan, Peter Schröder, and Günter M. Ziegler, editors, *Discrete Differential Geometry*, pages 137–161. Birkhäuser, Basel, 2008.
- [25] Carsten Thomassen. Planarity and duality of finite and infinite graphs. *Journal of Combinatorial Theory, Series B*, 29:244–271, 1980.
- [26] William Thomas Tutte. How to draw a graph. *Proceedings of the London Mathematical Society*, 3(1):743–767, 1963.
- [27] Gopalakrishnan Vijayan. Geometry of planar graphs with angles. In *Proc. 2nd ACM Symposium on Computational Geometry*, pages 116–124, 1986.
- [28] Christian Wiener. *Über Vielecke und Vielfläche*. Teubner, Leipzig, 1864.
- [29] Rhaleb Zayer, Christian Rössler, and Hans-Peter Seidel. Variations on angle based flattening. In N. A. Dodgson, M. S. Floater, and M. A. Sabin, editors, *Advances in Multiresolution for Geometric Modelling*, Mathematics and Visualization, pages 187–199. Springer, 2005.