

## The Impact of Cosmic Rays on the Kinematics of the Circumgalactic Medium

IRYNA S. BUTSKY,<sup>1</sup> JESSICA K. WERK,<sup>1</sup> KIRILL TCHERNYSHYOV,<sup>1</sup> DRUMMOND B. FIELDING,<sup>2</sup> JOSEPH BRENNEMAN,<sup>1</sup>  
DANIEL PIACITELLI,<sup>1</sup> THOMAS R. QUINN,<sup>1</sup> N. NICOLE SANCHEZ,<sup>1</sup> AKAXIA CRUZ,<sup>3</sup> CAMERON B. HUMMELS,<sup>4</sup>  
JOSEPH N. BURCHETT,<sup>5</sup> AND MICHAEL TREMMEL<sup>6,7</sup>

<sup>1</sup>*Astronomy Department, University of Washington, Seattle, WA 98195, USA*

<sup>2</sup>*Center for Computational Astrophysics, Flatiron Institute, 162 Fifth Avenue, New York, NY 10010, USA*

<sup>3</sup>*Physics Department, University of Washington, Seattle, WA 98195, USA*

<sup>4</sup>*TAPIR, California Institute of Technology, Pasadena, CA 91125, USA*

<sup>5</sup>*Department of Astronomy, New Mexico State University, Las Cruces, NM 88003, USA*

<sup>6</sup>*Department of Physics, Yale University, New Haven, CT 06511, USA*

<sup>7</sup>*Yale Center for Astronomy and Astrophysics, New Haven, CT 06511, USA*

Submitted to ApJ

### ABSTRACT

We use hydrodynamical simulations of two Milky Way-mass galaxies to demonstrate the impact of cosmic-ray pressure on the kinematics of cool and warm circumgalactic gas. Consistent with previous studies, we find that cosmic-ray pressure can dominate over thermal pressure in the inner 50 kpc of the circumgalactic medium (CGM), creating an overall cooler CGM than that of similar galaxy simulations run without cosmic rays. We generate synthetic sightlines of the simulated galaxies' CGM and use Voigt profile fitting methods to extract ion column densities, Doppler- $b$  parameters, and velocity centroids of individual absorbers. We directly compare these synthetic spectral line fits with HST/COS CGM absorption-line data analyses, which tend to show that metallic species with a wide range of ionization potential energies are often kinematically aligned. Compared to the Milky-Way simulation run without cosmic rays, the presence of cosmic-ray pressure in the inner CGM creates narrower O VI absorption features and broader Si III absorption features, a quality which is more consistent with observational data. Additionally, because the cool gas is buoyant due to nonthermal cosmic-ray pressure support, the velocity centroids of both cool and warm gas tend to align in the simulated Milky Way with feedback from cosmic rays. Our study demonstrates that detailed, direct comparisons between simulations and observations, focused on gas kinematics, have the potential to reveal the dominant physical mechanisms that shape the CGM.

*Keywords:* Circumgalactic medium (1879), cosmic rays (329), Galaxy evolution (594)

### 1. INTRODUCTION

Galaxies evolve embedded within a vast, gaseous halo that dwarfs the mass and spatial extent of the galactic disk. This circumgalactic medium (CGM) has a rich multiphase structure with gas temperatures, densities, and metallicities spanning several orders of magnitude (Tumlinson et al. 2017). The CGM drives galaxy evolution by controlling gas accretion, which is necessary for continued star formation, and is in turn shaped by galactic outflows, which expel the byproducts of stellar evolution. Constraints on the structure of the CGM and

the origin of its multiple phases are therefore crucial for understanding the dominant physical mechanisms driving galaxy evolution.

Because the CGM is so diffuse, much of what we know about its composition and kinematics derives from absorption-line studies, in which the sightline to a bright background source (typically a quasar) intersects a galaxy's CGM. The atoms in the CGM interact with the light from the quasar, creating absorption-line features in the resulting spectra from which we can measure properties like the ionic column density and line-of-sight velocity of the absorbing gas. Most sightlines intersecting the CGM detect absorption from high-, intermediate-, and low-ionization species at rest-frame ultraviolet (UV) wavelengths, spanning over an order of magnitude in ionization potential energies, both at

low redshift (e.g.,  $z \lesssim 1$ : Bergeron & Boissé 1991; Prochaska et al. 2011; Tripp et al. 2011; Tumlinson et al. 2013; Werk et al. 2013; Nielsen et al. 2013; Bordoloi et al. 2014; Liang & Chen 2014; Borthakur et al. 2015; Burchett et al. 2019) and high redshift (e.g.,  $2 < z < 3$ : Steidel et al. 2010; Rudie et al. 2012; Turner et al. 2015; Zahedy et al. 2019). Absorption from low-ions (e.g., Mg II, Si III), assumed to trace cool  $10^{4-5}$  K gas, is prevalent in the inner  $\sim 100$  kpc of the CGM while absorption from intermediate and high ions (e.g., C IV, O VI), assumed to trace warm  $10^{5-6}$  K gas, is detected out to, and sometimes beyond the galaxy virial radius (Johnson et al. 2015; Burchett et al. 2016; Keeney et al. 2018). Kinematic studies reveal that high ions and low ions often have similar line-of-sight velocity centroids, suggesting a common (or related) physical distribution of the warm and cool absorbers (Tripp et al. 2011; Werk et al. 2016).

Ionization modeling of the measured quantities has yielded a rich set of constraints on the gas-phase metallicity of the CGM (e.g., Lehner et al. 2013; Prochaska et al. 2017; Lehner et al. 2019), its total baryonic content (e.g., Werk et al. 2014; Stern et al. 2016), and its pressure profile (Stoche et al. 2013; Werk et al. 2016; Voit et al. 2019). In particular, around low-redshift  $L^*$  galaxies, cool CGM gas appears to have significantly lower densities than required for it to be in thermal pressure equilibrium with the hot phase (Werk et al. 2014; Stern et al. 2016).

One promising interpretation of the low densities of cool gas is that the CGM is supported by nonthermal cosmic-ray pressure. In recent years, cosmic rays have been invoked in galaxy simulations to launch winds (Ipavich 1975; Uhlig et al. 2012; Booth et al. 2013; Pakmor et al. 2016; Simpson et al. 2016; Ruszkowski et al. 2017; Farber et al. 2018; Hopkins et al. 2021a), and thereby altering the ionization structure of the CGM (Salem et al. 2016; Butsky & Quinn 2018; Ji et al. 2020b; Buck et al. 2020). Although the quantitative details of the predicted CGM structure depend on the invoked model of cosmic-ray transport (Butsky & Quinn 2018; Hopkins et al. 2021b,d), which is still poorly constrained, many of these simulations predict a cosmic-ray-pressure-supported CGM around low-redshift  $L^*$  galaxies. In these models, the cosmic-ray pressure can exceed the gas pressure by up to two orders of magnitude in the inner CGM. Significant cosmic-ray pressure support allows low-density cool gas to survive in the CGM, naturally producing high column densities of low ions. Significant cosmic-ray pressure can also qualitatively alter the kinematics of the CGM by changing the morphology of outflows (e.g., Girichidis et al. 2016; Jana et al. 2020) and preventing a virial shock Ji et al. (2020a).

In this work, we explore the implications of a cosmic-ray-pressure-supported halo on the observed kinematic alignment between multiphase ions. We start with a Milky Way mass galaxy which has successfully

reproduced O VI column densities with the help of supermassive black hole (SMBH) feedback (Sanchez et al. 2019). We then re-simulate that same galaxy, adding supernova cosmic-ray feedback. By redshift  $z = 0.25$ , this galaxy develops a cosmic-ray-pressure-supported halo, qualitatively similar to those described in the recent literature.

To compare the simulations with observations, we generate 100 synthetic spectra from sightlines that pierce the inner CGM of each simulated halo. We then analyze absorption features in these spectra with Voigt-profile fitting tools used by observers. This approach allows us to directly compare our simulated CGM against observed absorption-line properties (e.g., Doppler- $b$  parameters, line-of-sight velocity offsets) — an approach that is inaccessible to traditional simulation analysis techniques.

This paper is organized as follows. In Section 2, we describe our methods, including the simulation initial conditions, our choices of run-time physics, and the process for generating and analyzing synthetic spectra. In Section 3, we present our results. We first present a simulation-oriented view of the CGM properties and then shift the focus to properties derived from the Voigt-profile fits with an emphasis on the Doppler- $b$  parameters and velocity offsets. In Section 4, we discuss the implications of our work and future directions. In Section 5, we present a summary of this work and its conclusions. Lastly, we demonstrate the implementation of cosmic-ray physics in the CHANGA simulation code in Appendix A.

## 2. METHODS

### 2.1. Description of Galaxy Simulations

In this work, we analyze cosmological zoom-in simulations of two Milky Way-sized ( $M_{\text{vir}} = 9.9 \times 10^{11} M_{\odot}$  at  $z = 0$ ) galaxies evolved with and without cosmic-ray feedback. The base model, “Patient 0” (P0) has been studied in great detail in Sanchez et al. (2019) and Sanchez et al. (2020). “Patient 0 with cosmic rays” (P0+CR) uses the same initial conditions and physics as P0 except that it also includes cosmic-ray feedback from supernovae. We present a basic description of the simulation here for context, but encourage the reader to see the referenced works for more detail. The description of the cosmic-ray physics implementation is detailed in Appendix A.

The simulations were run with ChaNGa, a smoothed-particle hydrodynamics (SPH) astrophysical simulation library (Menon et al. 2015) with a recently updated formalism that includes a geometric density approach in the force expression (Wadsley et al. 2017). We assume a  $\Lambda$ CDM cosmology with parameters  $\Omega_m = 0.3086$ ,  $\Omega_{\Lambda} = 0.6914$ ,  $h = 0.67$ ,  $\sigma_8 = 0.77$  and an extragalactic UV background described by Haardt & Madau (2012). The mass of dark matter particles and gas particles are  $2.1 \times 10^5 M_{\odot}$  and  $1.4 \times 10^5 M_{\odot}$  respectively.

We do not include metal-line cooling for temperatures above  $10^4$  K. Although this is a limitation of the simulation, these effects are mitigated by the fact that the balance between heating and cooling at warm CGM temperatures is dominated by the extragalactic UV background. In this case, excluding metal-line cooling overestimates the cooling time by a factor of  $\sim 3 - 5$  (Shen et al. 2010). Additionally, the cooling times in  $10^{4-4.5}$  K gas are well estimated by primordial cooling (Hopkins et al. 2018).

Both simulations model star formation assuming a Kroupa IMF (Kroupa 2001) and ‘blastwave’ supernova feedback (Stinson et al. 2006). We allow star formation to happen in cold ( $T < 10^4$  K), dense ( $n > 0.2 \text{ cm}^{-3}$ ) gas with an efficiency of 15% and a characteristic timescale of  $10^6$  years. Each supernova injects  $0.75 \times 10^{51}$  erg of energy into the surrounding gas.


In P0+CR, cosmic-ray energy is injected into the simulation during supernova feedback events. The total injected supernova energy remains constant, but 10% is injected in the form of cosmic ray energy. Cosmic-ray transport is modeled as isotropic diffusion with a constant diffusion coefficient of  $\kappa = 10^{29} \text{ cm}^2 \text{ s}^{-2}$ .

We also include SMBH feedback as described in Tremmel et al. (2017). SMBH seeds form in very dense gas ( $n > 3m_H/\text{cm}^3$ ) with low metallicities ( $Z < 3 \times 10^{-4} Z_\odot$ ). SMBH accrete gas through both mergers and modified Bondi-Hoyle accretion (Tremmel et al. 2017) and redistribute 0.2% of the mass energy of their accreted material into nearby gas via thermal feedback. We do not inject additional cosmic-ray energy during SMBH feedback events.


## 2.2. Generating Synthetic Spectra

We explicitly track the evolution of hydrogen, oxygen, and iron at runtime. In the analysis phase, we use TRIDENT (Hummels et al. 2017), an extension of the widely-used simulation analysis tool, YT (Turk et al. 2012), to estimate the abundance of other chemical species as well as their ionization states. We also use TRIDENT to generate synthetic spectra that mimic the specifications of the HST COS-G130M and COS-G160M instruments.

We generate 100 randomly-oriented sightline coordinates that sample the inner CGM with impact parameters between 10 and 50 kpc. To generate the coordinates of a random sightline, we first define the position of the impact parameter – the closest point between the sightline and the galactic center – as a point on the surface of a sphere centered on the galaxy:  $p(r, \theta, \phi)$ . In this expression,  $r$  is the impact parameter,  $\theta$  is the polar angle, and  $\phi$  is the azimuthal angle. All three values are selected using a random, unweighted number generator. Once we choose the coordinates of the impact parameter, we set the orientation of the sightline to be at a random angle,  $\psi$ , in the plane



**Figure 1.** (Top) The projected paths of the 100 synthetic spectra in the x-y plane, centered on the galaxy center. The open circles show the projected position of the impact parameter – the closest point between the sightline and the galactic center. (Bottom) The random distribution of impact parameters between 10 and 50 kpc. Note that the minimum impact parameter is 10 kpc, and the appearance of impact parameters  $< 10$  kpc in the top panel is due to projection effects.



**Figure 2.** The mass-weighted projections of density, temperature and line-of-sight velocity for P0 (top) and P0+CR (bottom) at  $z = 0.25$ . The width and depth of the projection are 100 kpc. Relative to P0, the simulation with cosmic rays has a cooler CGM in the inner  $\sim 40$  kpc and a larger rotating extended disk.

tangent to the sphere. All sightlines are 500 kpc long and centered at their impact parameter.

We use the same set of galactocentric sightline coordinates to generate synthetic spectra for both, P0 and P0+CR. Figure 1 provides a visual depiction of the randomly distributed sightlines and the distribution of impact parameters (the circles).

For each sightline in each galaxy, we generate a synthetic spectrum at  $z = 0.25$ , including all transitions of the H I, Mg II, C II, C III, C IV, Si II, Si III, Si IV, N V, and O VI ions. TRIDENT deposits Voigt absorption profiles that explicitly include thermal broadening and implicitly include nonthermal broadening due to gas motions. We subtract out the bulk velocity of the 500 kpc sphere centered on the galaxy relative to the simulation frame. To make these synthetic spectra resemble real HST/COS data, we add noise with a signal-to-noise ratio of 10 per resolution element and convolve it with the line-spread functions of the G130M and G160M spectrographs. The resulting normalized synthetic spectra are characterized by a FWHM  $\approx 18$  km s $^{-1}$  when they are binned by three native spectral pixels to a dispersion of  $\Delta\lambda \approx 0.0367$  Å.

### 2.3. Analyzing Synthetic Spectra

Our spectral analysis incorporates common observational techniques for analysing medium-resolution HST/COS data (e.g., Tumlinson et al. 2011; Werk et al. 2013). Specifically, we extract information from the synthetic spectra by decomposing the absorption features into Voigt profiles. We do so without any regard to the parameters that are used by TRIDENT to generate the synthetic spectra.

The decomposition involves five steps: (1) automated identification of distinct absorption components; (2) an initial Voigt profile fit; (3) manual refinement of component definitions; (4) a final Voigt profile fit; and (5) calculation of column density upper limits for ions with no detected components. For each ion and sightline, this procedure produces either a column density upper limit or a collection of: component column densities,  $N$ ; velocity centroids relative to the halo rest frame,  $v$ ; and linewidths expressed as Doppler- $b$  parameters.

To identify absorption components for an ion along a sightline, we estimate the column density as a



function of velocity,  $N(v)$ , and then split  $N(v)$  into candidate components using a watershed-like segmentation algorithm. We use the apparent optical depth (AOD) method to estimate  $N(v)$ . The AOD method is based on the assumption that the logarithm of the degree of absorption at a velocity, the apparent optical depth, is directly proportional to  $N(v)$  (Savage & Sembach 1991). We then use the ASTRODENDRO package (Rosolowsky et al. 2008) to split  $N(v)$  up into a set of distinct and contiguous peaks; these are our candidate components. Finally, we remove candidate components whose integrated column densities are less than two times their integrated column density uncertainties.

Each candidate component has a centroid velocity, an allowed velocity range, and rough estimates of the component column density and linewidth. These values are used as initial conditions for the Voigt profile fit. We use optimization to find the likelihood-maximizing component parameter set and estimate uncertainties by calculating and inverting the Fisher information matrix (e.g., Tegmark et al. 1997) at the maximum likelihood solution.

The automated component identification procedure is imperfect. It makes two main kinds of mistakes: (1) identifying a single component where multiple components would be more appropriate and (2) identifying a single absorption feature as arising from different ions. We resolve the first kind of mistake by inspecting the initial Voigt profile fits and defining new components as needed. The second mistake happens when ions have transitions with similar rest wavelengths, such as the C II 1036Å and O VI 1037Å lines. In all such cases, at least one of the ions has more than one line, so the maximum likelihood solution for the fictitious component has a very low column density. This allows us to find and remove these components using a simultaneous cut on velocity and column density. A second round of Voigt profile fits done with the revised component structures provides our adopted set of component parameters.

If an ion has no components along a sightline, we calculate an upper limit on the ion’s column density. We use AOD to calculate  $N(v)$  and integrate the result over a 100 km s<sup>−1</sup> interval. We then convert the total column density and column density uncertainty over this interval to a 2- $\sigma$  equivalent upper limit using the formalism of Bowen et al. (2008).

### 3. RESULTS

#### 3.1. The Physical State of the CGM

Despite starting from the same initial conditions, by a redshift of 0.25, the two galaxies have distinctly different CGM properties. While some differences are naturally to be expected (even for theoretically identical cosmological simulations) due to low-level non-linear effects (Genel et al. 2019), the buildup of cosmic-ray

pressure in P0+CR produces a *qualitatively* different CGM, significantly exceeding the differences expected from chaotic processes of galaxy formation.


Figure 2 shows the mass-weighted projections of gas density, temperature, and line-of-sight velocity. The CGM of P0 is notably hotter, with temperatures in excess of 10<sup>6</sup> K, in part due to a recent burst of supernova feedback, and in part due to the thermal gas pressures required to maintain hydrostatic equilibrium. The cool, dense filament of inflowing gas traces the remnants of a recent merger with a satellite galaxy.

By contrast, the inner CGM of P0+CR is full of cool gas, supported against gravity and condensation by significant cosmic-ray pressure. This cool inner CGM extends roughly 40 kpc and has higher gas densities than the inner CGM of P0. The cool gas is pierced by pockets of hot bubbles from supernova feedback. There is a sharp transition between the cool, inner CGM and the outer CGM, marking the edge of the cosmic-ray-pressure-supported halo. The temperatures and densities in the outer CGM of P0+CR resemble those in the outer CGM of P0. The line-of-sight velocity distribution is smoother in P0+CR than in P0. The inflowing filament from the recent merger is also present in P0+CR, but is significantly diminished relative to the fast-moving filament in P0.

Figure 3 shows the projected ratio of cosmic-ray pressure to gas pressure around P0+CR. The inner  $\sim 40$  kpc of the CGM are supported by cosmic-ray pressure while the outer regions are supported by thermal pressure. The shape of the cosmic-ray-pressure-supported region traces the shape of the cool gas. The details of the cosmic-ray pressure profile and the extent of the cosmic-ray-pressure-supported region are sensitive to the choice of cosmic-ray hydrodynamics models, which are currently unconstrained. However, this simulation is qualitatively similar to the cosmic-ray-pressure-supported CGM seen in other hydrodynamics simulations with cosmic-ray diffusion (e.g., Salem et al. 2016; Butsky & Quinn 2018; Ji et al. 2020b; Buck et al. 2020).

These differences in the gas phase between the two galaxies translate into differences in their ion column densities. Figure 4 shows the column densities of O VI, Si III, and H I for the two simulated galaxies. P0 has high O VI column densities throughout, with its hot inner region exceeding column densities of 10<sup>15</sup> cm<sup>−2</sup>. While the O VI-bearing gas creates structures that span the extent of the inner CGM, the Si III and H I column density maps trace small-scale structures near the galactic disk and in the cold accretion stream. The sizes of the cloudlets that trace Si III and H I absorbers are likely limited by the resolution of the simulation (e.g., Hummels et al. 2019; Peebles et al. 2019; Suresh et al. 2019; van de Voort et al. 2019).

Relative to P0, P0+CR has lower O VI column densities with a profile that traces the shape of



**Figure 3.** The mass-weighted projection of the ratio of cosmic-ray pressure to gas pressure in P0+CR. Cosmic-ray pressure is significantly larger than thermal pressure in the inner CGM. The cosmic-ray pressure gradient counteracts gravity to maintain hydrostatic equilibrium, allowing cool, low-entropy gas to fill the inner CGM.

the cosmic-ray-pressure-supported region. The inner CGM has significant Si III and H I column densities throughout.

In Figure 5, we take a look at the differences in the gas properties of O VI and Si III absorbers between the two simulations. The images show the 2D histograms of simulated temperature and number density, colored by the ion mass probability density of O VI and Si III. The data for these plots was generated from spherical shells centered on the galaxy with radii between 10 and 50 kpc, corresponding to the region sampled with our synthetic spectra and in the previous figures.


In both galaxies, the vast majority of Si III comes from gas around  $10^{4.2}$  K, as is expected for photoionized gas. Relative to P0, the Si III in P0+CR comes from lower density gas due to nonthermal cosmic-ray pressure support. In P0, O VI traces gas with temperatures of  $T = 10^{5.3-6.2}$  K — hotter than the peak O VI ionization temperature of  $\sim 10^{5.5}$  K. In P0+CR, the majority of O VI absorption traces gas with temperatures with a peak at  $T = 10^{5.5}$  K and has a negligible fraction of photoionized O VI at low temperatures and low gas densities. Like with Si III, O VI-bearing gas in P0+CR tends to have lower densities than in P0.

### 3.2. Physical Insights from Synthetic Spectra

Next, we present the physical state of the two galaxies as seen through their synthetic spectra. Unlike real observations, synthetic spectra have the additional benefit of linking absorption features to the exact gas properties that gave rise to them. Figure 6 and Figure 7 show two examples of synthetic spectra that detect both Si III and O VI. The top panels show the ion absorption as a function of its velocity offset from the galaxy rest frame. The remaining panels show various properties of the gas particles along the one-dimensional sightline used to generate the synthetic spectra. Unlike the top panel, the x-axis of the remaining panels shows the spatial trajectory along the sightline. In all cases, the sightline trajectories are centered so that zero corresponds to the location of the impact parameter. The scattered points are colored by their line-of-sight velocity, which provides a visual aid to determine which gas particles contributed to which absorption features in the top panel.

Figure 6 highlights two example synthetic spectra from P0. The P0 spectrum on the left contains two different examples of kinematic alignment between O VI and Si III. The fast-moving O VI absorber, with a relative velocity of  $\sim 100$  km/s comes from the same physical region as the correlated Si III absorber (see the spatial alignment of the deep-red points in the bottom two panels of Figure 6). This is consistent with cold clumps being entrained in a hot background medium. By contrast, the O VI and Si III absorbers moving with a relative velocity of  $\sim 20$  km/s come from completely different regions, separated by roughly 50 kpc along the line of sight. In this case, their kinematic alignment appears to be a coincidence. Additionally, the 20 km/s Si III absorber comes from one extrapolated data point along the sightline (at  $\sim -15$  kpc), indicating that the cool cloud is not resolved and resulting in a narrow Si III absorption feature.

The second P0 spectrum, in the right panel of Figure 6, contrasts the first example in that it contains two absorption features of both O VI and Si III without any kinematic alignment. The fast-moving Si III absorber with a relative velocity of  $-200$  km/s traces the inflowing accretion of cool gas following a recent merger. Although it is spatially aligned with some O VI-absorbing gas, the line-of-sight velocity of this Si III absorber is significantly offset from any O VI features. The Si III absorber with a relative velocity of  $-120$  km/s appears to come from a single particle (medium blue point at  $\sim -5$  kpc). Although that particle is spatially aligned with an O VI cloud with similar line-of-sight velocities, the corresponding absorption features do not appear aligned. This is likely because the O VI absorption feature is broadened by gas particles moving with similar, but slightly different velocities between  $-50$  and  $-100$  km/s, whereas the unresolved Si III absorber is only composed of one particle.



**Figure 4.** The O VI, Si III, and H I column densities at  $z = 0.25$  for P0 (top) and P0+CR (bottom). The width (and depth) of the projection is 100 kpc. Compared to P0, the CGM of P0+CR has lower overall O VI column densities and higher column densities of Si III and H I in the inner 50 kpc.

The fact that the Si III structures are unresolved likely contributes to artificially narrow Si III absorbers, because the velocity structure of a fully-resolved cloud would contribute to line broadening (Peeples et al. 2019).

The P0+CR spectra in Figure 7 show a very different relationship between the kinematic and physical alignment of O VI and Si III. In both example spectra, the O VI and Si III absorbers are kinematically aligned. The first example spectrum, in the left panel of Figure 7, has two distinct absorbers of each Si III and O VI. Both sets of kinematically aligned absorbers are also spatially aligned along the line of sight.


In the second example spectrum, in the right panel of Figure 7, the cool and warm gas comes from physically distinct, yet related clouds. Although O VI and Si III absorption is spatially offset, the O VI absorption borders the Si III absorption on either side. The velocity structure of the scattered points clearly shows that together, the O VI and Si III form a coherent physical region in which the O VI and the Si III absorbers are co-rotating. The Si III absorption line is very broad because it comes from a large Si III-bearing cloud that spans a

wide range of velocity space. The two O VI absorption features, which trace the velocity at either edge of the cool cloud are nestled within the broad Si III absorber.

These four examples highlight the observed trends in and differences between the P0 and P0+CR synthetic spectra.<sup>1</sup> P0 tends to have broader O VI absorbers and more narrow Si III absorbers. A large fraction of spectra either have no Si III absorption or have Si III absorbers that are kinematically misaligned with O VI absorbers. O VI absorbers often come from hot ( $T > 10^6$  K) gas that spans tens of kpc while Si III absorbers come from small (and likely resolution-limited) clouds of  $\sim 1$  kpc scales. The O VI- and Si III- bearing gas sometimes comes from the same physical cloud, but is often physically unrelated.

In P0+CR, Si III absorbers tend to be broader and come from large, low-density clouds spanning 10 – 50 kpc. O VI absorbers tend to come from slightly cooler gas ( $T \lesssim 10^6$  K) and slightly smaller clouds (10 – 50

<sup>1</sup> The interested reader can see this plot for every spectrum in our sample at [irynabutsky.me/data](http://irynabutsky.me/data)



**Figure 5.** The ion-mass weighted 2D histograms of gas temperature and number density for P0 (left) and P0+CR (right). The data for these plots is generated from a spherical shell between 10 and 50 kpc from the galactic center. The additional pressure support from cosmic rays moves the gas to lower densities. P0 has a substantial amount of O VI absorption at temperatures above the collisional ionization equilibrium temperature of  $T \sim 10^{5.5}$  K.


kpc) than in P0. O VI and Si III absorbers are often kinematically aligned and often come from either the same physical cloud or from physically related clouds.


Note that in an attempt to highlight the trends in the kinematic and physical relationships between O VI and Si III, we omit spectra that have no measurable Si III absorption from Figure 6 and Figure 7. While these true “no-lows” (O VI absorbers without a kinematically matched Si III absorber; Werk et al. 2016) are present in both simulations, they are very common in P0 and very rare in P0+CR.

### 3.3. The Doppler- $b$ Parameter


The Doppler- $b$  parameter is a measure of both thermal and nonthermal line broadening mechanisms,  $b = \sqrt{b_{th} + b_{nt}}$ . The Voigt profile of the synthetic spectra generated with TRIDENT is a convolution a Gaussian profile that measures thermal line broadening and a Lorentzian profile that measures nonthermal broadening. Therefore, a broad absorption feature may be due to high gas temperatures, pressures, or velocity dispersions. We note that although cosmic rays are a source of nonthermal pressure support they do not explicitly affect the Doppler- $b$  parameter value. Instead, they indirectly influence the Doppler- $b$  parameter by altering the density, temperature, velocity, or spatial extent of the gas.








**Figure 7.** Similar to Figure 6, the figure above shows example synthetic spectra from P0+CR and the gas properties along their line of sight. There is an additional panel to show the ratio of cosmic-ray pressure to gas pressure,  $P_c/P_g$ , along the line of sight. The impact parameters of the sightlines in the left and right panels are 36.5 kpc and 43.0 kpc respectively. (Left) An example of kinematic alignment in P0+CR in which both Si III and O VI absorbers come from the same physical structures. In all P0+CR spectra that have both O VI and Si III absorbers, there is at least one O VI component that is kinematically aligned with Si III. P0+CR has higher incidences of kinematic alignment between Si III and O VI. (Right) In most cases in P0+CR, Si III absorption comes from large,  $\sim 10 - 50$  kpc, clouds that are enveloped by O VI absorbers. Unlike in P0, low-density Si III-bearing gas in P0+CR is resolved by many gas particles.



**Figure 8.** (Left) The column density of O VI absorbers as a function of Doppler- $b$  parameter for P0 (top) and P0+CR (bottom). Purple squares indicate O VI absorbers that are matched with a Si III absorber within 35 km/s in line-of-sight velocity space. Gray diamonds indicate O VI absorbers that have no corresponding Si III absorber within 35 km/s. We include data from [Werk et al. \(2016\)](#) for comparison in black. Upward arrows indicate upper limits on the measured column density. (Right) The histograms of the Doppler- $b$  parameters of O VI absorbers in P0 (top) and P0+CR (bottom) compared against observations. All histograms are normalized to have an area of 1. P0+CR predicts a relative abundance of narrow ( $b = 20 - 30$  km/s) O VI absorbers. Both simulations have a distinct lack of very broad ( $b > 60$  km/s) O VI absorbers.

The left panel of [Figure 8](#) shows the O VI column density as a function of Doppler- $b$  parameter for P0 and P0+CR. Each scattered point represents a single O VI absorption feature and is colored by whether or not that absorber is kinematically aligned with a Si III feature (has a matched Si III absorber within 35 km/s in line-of-sight velocity space). P0 has significantly more “no-lows” than either P0+CR or the observations. This is in part due to the fact that P0 has both more sightlines that have no detected Si III and more sightlines with detected Si III that is kinematically unrelated to O VI.

In the cosmic-ray-pressure-supported CGM of P0+CR, the distribution of O VI Doppler- $b$  values is narrower, likely because O VI-bearing gas in P0+CR tends to have cooler temperatures than O VI-bearing gas in P0 ([Figure 5](#)). P0+CR also reproduces the observed abundance of O VI Doppler- $b$  parameters that are 20 – 30 km/s and predicts a relative gap in O VI Doppler- $b$  parameters around 40 km/s. In comparison, P0 has broader O VI features, with the most abundant O VI Doppler- $b$  parameter in the 30 – 40 km/s range.



**Figure 9.** (Left) The column density of Si III absorbers as a function of Doppler- $b$  parameter for P0 (top) and P0+CR (bottom). Green squares indicate Si III absorbers that are matched with a O VI absorber within 35 km/s in line-of-sight velocity space. Gray diamonds indicate Si III absorbers that have no corresponding O VI absorber within 35 km/s. We include data from [Werk et al. \(2016\)](#) for comparison in black. Upward arrows indicate upper limits on the measured column density. (Right) The histograms of the Doppler- $b$  parameters of Si III absorbers in P0 (top) and P0+CR (bottom) compared against observations. All histograms are normalized to have an area of 1. P0+CR better reproduces the observed broad Si III absorbers but over-predicts the abundance of narrow, high-column Si III absorbers.


Neither simulated distribution reproduces the very broad O VI features seen in [Werk et al. \(2016\)](#)<sup>2</sup>. The lack of very broad O VI absorbers is likely influenced by the lack of O VI in the outskirts of the simulated CGM and the simulation resolution, which narrows the velocity distribution within an O VI cloud. The broad O VI absorbers may also be affected by the small impact parameters of our sample or potential discrepancies

in modeling the COS line-spread function in synthetic spectra.

[Figure 9](#) repeats the format of [Figure 8](#) for Si III absorbers. In the left panel, each scatter point represents individual Si III absorbers, colored by whether they have a corresponding O VI absorber within 35 km/s in line-of-sight velocity space. The CGM of P0+CR produces higher column densities of Si III, which appear consistent with the range of observed column densities. However, it is difficult to compare precisely, as many of the observed Si III absorbers are saturated. P0+CR also produces distinctly higher

<sup>2</sup> There is one additional O VI absorber with  $b = 174.7$  km/s in the [Werk et al. \(2016\)](#) data that isn't pictured in [Figure 8](#).






**Figure 10.** The histograms of the minimum velocity differences between O VI and Si III absorbers for P0 (top) and P0+CR (bottom). This excludes sightlines that have O VI but no Si III absorbers. All histograms are normalized to have an area of 1. P0+CR better reproduces the observed kinematic link between O VI and Si III absorbers.

Si III Doppler- $b$  parameters that better agree with the observed distribution. However, as with O VI absorbers, neither simulation reproduces the broad tail of Si III Doppler- $b$  parameters.


We use a two-sample Kolmogorov-Smirnov (KS) test to investigate whether either P0 or P0+CR produces ion Doppler- $b$  parameters that sample the same distribution as the COS-Halos observations. In all cases, the KS statistic is inconclusive, and neither simulation produces an obviously better fit to the data. Besides differences in physical models, there are a variety of reasons why the simulations might



**Figure 11.** The mass-weighted 2D histogram of radial velocity as a function of temperature for cool (blue) and warm (orange) CGM gas for P0 (left) and P0+CR (right). The data is taken from gas that is between 10 and 50 kpc from the galactic center. Without cosmic rays, cool gas in the inner CGM tends to be on a radially inward trajectory. With substantial cosmic-ray pressure support, low-entropy cool gas is supported against gravity and has a very similar velocity distribution to that of the warm gas, leading to kinematic alignment of multiphase absorbers.

not reproduce the observed Doppler- $b$  distributions, including resolution, the fact that the synthetic spectra only probe the inner CGM, or comparing the properties of a single galaxy, at a single point in time, against an observed variety of galaxies.

#### 3.4. The Kinematics of Cool and Warm Gas



**Figure 12.** The Wasserstein distance between O VI and Si III line-of-sight velocities and positions along the sightline. The color of the points indicates the impact parameter of the sightline. P0+CR has a clear relationship between the kinematic and spatial alignment of O VI and Si III absorbing gas. In contrast, O VI and Si III gas in P0 are less likely to be kinematically or spatially aligned.

Next, we consider the impact of cosmic rays on the kinematics of warm and cool CGM gas traced by O VI and Si III. Using synthetic spectra, we compare the velocity offsets of individual absorbers of O VI and Si III and compare them directly against the properties of the velocity offsets of the absorbers in [Werk et al. \(2016\)](#).

In [Figure 10](#), we plot the histograms of minimum difference in velocity centroids between O VI and Si III for all sightlines that have detections of both ions. P0 has significantly more spectra that have Si III absorbers that are kinematically offset from the O VI absorbers. This is because the cool, Si III absorbing, gas in P0 is often physically unrelated to the warm, O VI absorbing, gas. By comparison, P0+CR has a high incidence of kinematic alignment between O VI and Si III.

As in the previous section, we use two-sample KS tests to see whether the simulated data is drawn from the same sample as the observed data. However, we again find that the KS tests are inconclusive.

The increased kinematic alignment between gas phases in P0+CR comes from the overall effect of cosmic-ray pressure on the CGM kinematics. [Figure 11](#) highlights the radial velocity differences between cool and warm gas in the inner CGM of both simulations. In P0, the majority of cool gas is radially infalling. Even if we were to exclude the fast inflowing cool gas that originates from the accreting stream, the remaining cool gas radial velocity histogram is still offset from the warm gas radial velocity histogram. Meanwhile, the cosmic-ray pressure support in P0+CR helps support cool gas against gravity, giving it a very similar radial

velocity distribution as that of the warm gas. In this way, significant cosmic-ray pressure support naturally leads to kinematic alignment between the cool and warm gas phases.

### 3.5. Relating Kinematic Alignment to Physical Absorbers

With *synthetic* spectra, we have the unique opportunity to relate the kinematic alignment between O VI and Si III to the spatial relationship of the absorbing gas along the line of sight. To do this, we turn back to the simulated properties of each 1D sightline from which the synthetic spectra are generated.

In our spectra, we see three physical explanations of kinematic alignment. (1) Some kinematic alignment is happenstance and comes from physically unrelated gas clouds. This physically unrelated kinematic alignment accounts for 37.5% of O VI-Si III absorbers in P0 and only 5% of O VI-Si III aligned absorbers in P0+CR.

(2) Some kinematic alignment traces multiphase gas clouds – either a cool cloud embedded in a warm cloud, or an intermediate temperature cloud that produces sufficient cool and warm components. These types of clouds tend to produce small kinematic offsets and comprise 62.5% of aligned O VI-Si III absorbers in P0 and 38% percent of aligned O VI-Si III absorbers in P0+CR. It is worth mentioning that we also see instances of O VI and Si III absorbers that come from the same physical cloud but that are kinematically misaligned.

(3) The last physical explanation for kinematic alignment is when both cool and warm gas are tracing the large scale rotation of the CGM (for example, see the right panel of Figure 7). In this case, the warm gas envelops the inner cool gas, and this type of physical alignment produces slightly larger velocity offsets than clouds that are both kinematically and physically aligned. Roughly 67% of kinematic alignment in P0+CR falls into this category but is not present in P0.

In Figure 12, we quantify the offsets between line-of-sight velocities and positions of O VI- and Si III-bearing gas using the Wasserstein distance (also known as the earth mover’s distance). The Wasserstein distance,  $W_p(m, n)$ , measures the amount of “work” it takes to transform one 1D distribution ( $m$ ) into another ( $n$ ). When defining the line-of-sight velocity and position samples, we consider all particles whose O VI/Si III column density is at least  $10^{-2}$  of the total O VI/Si III column density of the sightline. We then weight each sample using the particles’ fractional column density.

Within a single sightline, if the O VI-bearing gas tends to have similar velocities as Si III-bearing gas, then the Wasserstein distance between the two velocity distributions,  $W_p(v_{\text{OVI}}, v_{\text{SiIII}})$ , will be small. Similarly, if O VI-bearing gas tends to be co-spatial with Si III-bearing gas, then the Wasserstein distance between the two line-of-sight positions,  $W_p(x_{\text{OVI}}, x_{\text{SiIII}})$ , will be small. Conversely, if cool and warm gas is physically or kinematically unrelated, then the relevant Wasserstein distance(s) will be large. Considering an ensemble of sightlines, a tightly-correlated relationship between the Wasserstein distance for the velocities and spatial offsets would suggest that kinematic alignment in spectra also implies that Si III- and O VI-absorbing gas is co-spatial. Conversely, a complete lack of correlation would imply that kinematic alignment is not related to the physical location of the absorbing gas.

The scattered points in Figure 12 show the Wasserstein distances of the line-of-sight velocities and positions of O VI- and Si III-bearing gas for all sightlines that contain detectable absorption features from both O VI and Si III. The color of each scattered point represents the impact parameter of that sightline.

In P0, there is no clear relationship between the kinematic and spatial alignment of cool and warm gas, traced by O VI and Si III. This result suggests that kinematic alignment of multiphase gas in P0 spectra is unlikely to imply a physical relationship between the gas phases. We note that the lack of scattered points with high impact parameters is due to the relative lack of Si III absorbers at large distances from the galaxy center.

In the cosmic-ray-pressure-supported halo of P0+CR, there is a positive correlation between the kinematic and spatial alignment of cool and warm gas. In particular, sightlines with impact parameters between 35-50 kpc

have the smallest kinematic and spatial offsets, implying that O VI and Si III are arising from the same physical structures. Meanwhile, sightlines that directly pierce the central region of the Si III structure have larger spatial and velocity offsets with O VI-bearing gas. This is likely because the inner CGM P0+CR is filled with a continuous Si III-absorbing cloud that is pierced by smaller O VI absorbers. The data from P0+CR predict that small kinematic offsets ( $< 10$  km/s) between O VI and Si III absorbers imply small spatial offsets ( $< 10$  kpc).

## 4. DISCUSSION

### 4.1. Implications for CGM Kinematics

In this work, we demonstrate that the dominant source of pressure support (thermal or cosmic-ray) in the CGM alters the relationship between cool and warm gas kinematics.

We first consider the case of a thermal-pressure-supported CGM, like the one present in P0. In order to maintain thermal pressure equilibrium, cool gas must have significantly higher densities than warm gas. Therefore, cool gas forms small, dense clouds that are not buoyant in the CGM. The lack of buoyancy implies that regardless of its origin, cool gas will accelerate towards the galactic disk. Even if cool gas initially shares the velocity of the warm phase (for example, cool gas that forms through thermal instability), the common velocity phase will be short-lived as cool droplets will accelerate towards the galaxy with a characteristic timescale set by the gravitational freefall time,  $t_{\text{ff}} = 2r/v_c$ . For the inner CGM ( $r = 50$  kpc) of a MW-sized galaxy, with a circular velocity of  $v_c = 200$  km/s, the freefall time is roughly  $t_{\text{ff}} = 500$  Myr. Therefore, in a thermal-pressure-supported CGM, one would expect a large fraction of cool gas absorbers to have at least some velocity offsets from warm absorbers.

In a cosmic-ray-pressure-supported CGM, the cool gas phase has significantly lower densities and is supported against gravity by the cosmic-ray pressure gradient. Rather than contracting into dense clouds, low-density cool gas fills the entire inner CGM, out to  $\sim 50$  kpc. The combination of a cosmic-ray pressure gradient contributing to maintaining hydrostatic equilibrium and the increased buoyancy of the low-density cool gas prevents cool gas from sinking as rapidly in the gravitational potential well (Butsky et al. 2020). Together, these effects result in cool gas having a similar velocity distribution to the warm gas phase. This naturally leads to a kinematic alignment between multiphase ions in absorption spectra.

In both galaxy models, the radial velocity distributions of warm gas are similar to the radial velocities of hot gas (Figure 11). Therefore, we would expect absorption from higher ionization states of oxygen (O VII, O VIII) detected in future X-ray

absorption studies to be kinematically aligned with O VI absorbers.

#### 4.2. The Relationship Between Kinematic and Spatial Alignment

In addition to altering CGM kinematics, the dominant source of pressure support (thermal or cosmic-ray) in the CGM alters the relationship between kinematic alignment and physical gas structure.

In the thermal-pressure-supported CGM of P0, gas positions and velocities are generally uncorrelated. Although in some cases, kinematic alignment does correspond with spatial alignment, we did not discover any trends that could discern such cases from coincidental kinematic alignment. In the cosmic-ray-pressure-supported CGM of P0+CR, kinematic alignment *does* correlate with spatial alignment. Absorbers with small velocity differences tend to come from the same gas structures. Spectra in which O VI absorption features were nestled inside a broad Si III absorption feature traced the rotation of the extended disk in which O VI gas enveloped Si III gas.

In both galaxy models, the kinematics of cool gas is particularly decoupled from gas positions in the case of a coherent inflowing stream. Therefore, the more we expect coherent inflows to be an important component of CGM structure, the less we should trust velocity alignment as an indicator of spatial alignment.

#### 4.3. Uncertainty in Modeling Cosmic Rays

It is now well-established that cosmic rays are produced in supernova shocks (e.g., Ackermann et al. 2013) and are an important source of energy in the galactic disk (e.g., Boulares & Cox 1990). However, despite significant recent theoretical and computational advances in cosmic-ray hydrodynamics (e.g., Jiang & Oh 2018; Thomas & Pfrommer 2019; Hopkins et al. 2021c), there is no consensus as to which existing cosmic-ray transport models (if any) realistically model the true behavior of cosmic rays. Several recent studies have demonstrated that different cosmic-ray transport models can change the predicted properties of the galactic disk, galactic outflows, and CGM structure (e.g., Ruszkowski et al. 2017; Butsky & Quinn 2018; Buck et al. 2020; Hopkins et al. 2021b).

While predictions for the degree of cosmic-ray pressure in the CGM have not yet converged, a cosmic-ray-pressure-supported CGM appears to be consistent with existing  $\gamma$ -ray observational constraints (Chan et al. 2019; Ji et al. 2020b). Nevertheless, P0+CR should be considered a limiting case, as it predicts cosmic-ray pressures that are relatively high within (yet still consistent with) the current predicted range. It is possible that advancements in cosmic-ray transport models would result in a CGM that is somewhere between that of P0 and P0+CR.

We expect that many of the quantitative details in this work, for example the exact ion column densities, the radial extent of the cosmic-ray-pressure-supported region, or the details of the gas phase distribution of our simulation will likely change with improved models for cosmic-ray transport. However, the *qualitative* predictions in this work, in particular the predictions for the kinematic link between cool and warm gas phases, are robust for any CGM model that contains significant nonthermal pressure support.

### 5. SUMMARY

In this work, we use synthetic spectroscopy to investigate the role of cosmic-ray pressure in setting the observed absorption-line properties and the kinematics of cool and warm gas in the CGM of low-redshift L\* galaxies. We simulate two Milky Way-sized galaxies, one without cosmic rays (P0), and one with cosmic-ray supernova feedback (P0+CR). By a redshift of  $z = 0.25$ , these two galaxies, which started from the same initial conditions, have evolved fundamentally different CGM properties.

P0 has a hot, thermal-pressure-supported CGM that is pierced by cool cloudlets in the inner CGM as well as an accreting stream from a recent merger event. By contrast, the inner CGM of P0+CR is supported against gravity by cosmic-ray pressure and filled with cool, low density gas.

We systematically probe the inner CGM of both galaxies with synthetic spectra and compare the extracted column densities, Doppler- $b$  parameters, and velocity centroids directly against COS-Halos data. We focus our analysis on two ions: O VI, which traces warm  $\sim 10^{5.5}$  K gas, and Si III, which traces cool  $\sim 10^{4.2}$  K gas.

Our results are summarized as follows.

(1) A cosmic-ray-pressure-supported CGM results in more narrow O VI Doppler- $b$  parameters than a thermal-pressure-supported CGM. This is due to the relatively cooler temperatures of O VI-bearing gas in the P0+CR simulation. In P0+CR, O VI absorption peaks at  $10^{5.5}$  K, while in P0, O VI traces both warm and hot gas with temperatures of  $10^{5.5-5.9}$  K (Figure 5). While P0+CR can explain the observed relative abundance of O VI Doppler- $b$  parameters between 20-30 km/s, neither simulation offers a statistically good fit to the observed data. Notably, neither simulation reproduced the observed large O VI Doppler- $b$  parameters  $> 60$  km/s (Figure 8).

(2) A cosmic-ray-pressure-supported CGM produces broader Si III Doppler- $b$  parameters. In both simulations, Si III traces gas with temperatures around  $10^{4.2}$  K (Figure 5). In P0, cool gas tends to form small cloudlets with narrow Doppler- $b$  parameters (Figure 9). In contrast, the low-density cool gas in P0+CR fills the entire inner CGM, resulting in large column densities and broader absorption features of Si III. While



the distribution of Si III Doppler- $b$  parameters in P0+CR is in better agreement with observations, it still underpredicts the observed abundance broad Si III features.

(3) A cosmic-ray-pressure-supported CGM predicts substantially more kinematic alignment of low and high ions. There is significantly more kinematic alignment between O VI and Si III in P0+CR (Figure 10). We demonstrate that in a cosmic-ray-pressure-supported halo, in which low-entropy cool gas is supported against gravity, the velocity distribution of cool and warm gas is similar, leading to a natural alignment between multiphase ions. This contrasts the kinematics of the thermally supported CGM of P0, in which cool gas is denser and is more likely to have an inward radial velocity than the warm gas (Figure 11).

(4) A cosmic-ray-pressure-supported CGM predicts a relationship between kinematic and physical alignment. We demonstrate that kinematic alignment does not necessarily imply that the absorbers trace physically related clouds. Our simulations predict that in a cosmic-ray-pressure-supported CGM, there is a correlation between the kinematic offset between O VI and Si III absorbers and their physical separation along the line of sight (Figure 12). By contrast, kinematic and spatial alignment in P0 are uncorrelated.

While the quantitative details of modeling cosmic rays in galaxy-scale simulations are still poorly-constrained, our study highlights how significant cosmic-ray pressure

support in the CGM fundamentally changes the kinematic signatures of multiphase gas. Ultimately, this work demonstrates that detailed comparisons between simulations and observations using synthetic spectroscopy are a powerful tool for constraining the dominant physical processes that govern galaxy evolution.

## ACKNOWLEDGMENTS

The authors thank Philip Hopkins, Suoqing Ji, and Peng Oh for their insightful comments. This research benefited from the KITP Program: “Fundamentals of Gaseous Halos”, and thus was supported in part by the National Science Foundation under Grant No. NSF PHY-1748958. IB was supported by HST theory grant HST-AR-15046 and by her tenure as a Blue Waters Graduate Fellow. The Blue Waters sustained-petascale computing project is supported by the National Science Foundation (Grants No. OCI-0725070 and No. ACI-1238993) and the State of Illinois. JKW acknowledges support as a Cottrell Scholar, from the Research Corporation for Science Advancement, grant ID number 26842. Additionally, JKW and KT acknowledge support for this work from NSF-AST 1812521. DBF, through the Flatiron Institute, is supported by the Simons Foundation.

## REFERENCES

- Ackermann, M., Ajello, M., Allafort, A., et al. 2013, *Science*, 339, 807, doi: [10.1126/science.1231160](https://doi.org/10.1126/science.1231160)
- Backus, I. 2017, sphglass, <https://github.com/ibackus/sphglass>, GitHub
- Bergeron, J., & Boissé, P. 1991, *A&A*, 243, 344
- Booth, C. M., Agertz, O., Kravtsov, A. V., & Gnedin, N. Y. 2013, *ApJL*, 777, L16, doi: [10.1088/2041-8205/777/1/L16](https://doi.org/10.1088/2041-8205/777/1/L16)
- Bordoloi, R., Tumlinson, J., Werk, J. K., et al. 2014, *ApJ*, 796, 136, doi: [10.1088/0004-637X/796/2/136](https://doi.org/10.1088/0004-637X/796/2/136)
- Borthakur, S., Heckman, T., Tumlinson, J., et al. 2015, *ApJ*, 813, 46, doi: [10.1088/0004-637X/813/1/46](https://doi.org/10.1088/0004-637X/813/1/46)
- Boulares, A., & Cox, D. P. 1990, *ApJ*, 365, 544, doi: [10.1086/169509](https://doi.org/10.1086/169509)
- Bowen, D. V., Jenkins, E. B., Tripp, T. M., et al. 2008, *ApJS*, 176, 59, doi: [10.1086/524773](https://doi.org/10.1086/524773)
- Buck, T., Pfrommer, C., Pakmor, R., Grand, R. J. J., & Springel, V. 2020, *MNRAS*, doi: [10.1093/mnras/staa1960](https://doi.org/10.1093/mnras/staa1960)
- Burchett, J. N., Tripp, T. M., Bordoloi, R., et al. 2016, *ApJ*, 832, 124, doi: [10.3847/0004-637X/832/2/124](https://doi.org/10.3847/0004-637X/832/2/124)
- Burchett, J. N., Tripp, T. M., Prochaska, J. X., et al. 2019, *ApJL*, 877, L20, doi: [10.3847/2041-8213/ab1f7f](https://doi.org/10.3847/2041-8213/ab1f7f)
- Butsky, I. S., Fielding, D. B., Hayward, C. C., et al. 2020, *ApJ*, 903, 77, doi: [10.3847/1538-4357/abbad2](https://doi.org/10.3847/1538-4357/abbad2)
- Butsky, I. S., & Quinn, T. R. 2018, *ApJ*, 868, 108, doi: [10.3847/1538-4357/aaeac2](https://doi.org/10.3847/1538-4357/aaeac2)
- Chan, T. K., Kereš, D., Hopkins, P. F., et al. 2019, *MNRAS*, 488, 3716, doi: [10.1093/mnras/stz1895](https://doi.org/10.1093/mnras/stz1895)
- Farber, R., Ruszkowski, M., Yang, H. Y. K., & Zweibel, E. G. 2018, *ApJ*, 856, 112, doi: [10.3847/1538-4357/aab26d](https://doi.org/10.3847/1538-4357/aab26d)
- Genel, S., Bryan, G. L., Springel, V., et al. 2019, *ApJ*, 871, 21, doi: [10.3847/1538-4357/aaf4bb](https://doi.org/10.3847/1538-4357/aaf4bb)
- Girichidis, P., Naab, T., Walch, S., et al. 2016, *ApJL*, 816, L19, doi: [10.3847/2041-8205/816/2/L19](https://doi.org/10.3847/2041-8205/816/2/L19)
- Haardt, F., & Madau, P. 2012, *ApJ*, 746, 125, doi: [10.1088/0004-637X/746/2/125](https://doi.org/10.1088/0004-637X/746/2/125)
- Hopkins, P. F., Chan, T. K., Ji, S., et al. 2021a, *MNRAS*, 501, 3640, doi: [10.1093/mnras/staa3690](https://doi.org/10.1093/mnras/staa3690)
- Hopkins, P. F., Chan, T. K., Squire, J., et al. 2021b, *MNRAS*, 501, 3663, doi: [10.1093/mnras/staa3692](https://doi.org/10.1093/mnras/staa3692)
- Hopkins, P. F., Squire, J., & Butsky, I. S. 2021c, arXiv e-prints, arXiv:2103.10443. <https://arxiv.org/abs/2103.10443>

- Hopkins, P. F., Squire, J., Chan, T. K., et al. 2021d, MNRAS, 501, 4184, doi: [10.1093/mnras/staa3691](https://doi.org/10.1093/mnras/staa3691)
- Hopkins, P. F., Wetzel, A., Kereš, D., et al. 2018, MNRAS, 477, 1578, doi: [10.1093/mnras/sty674](https://doi.org/10.1093/mnras/sty674)
- Hummels, C. B., Smith, B. D., & Silvia, D. W. 2017, ApJ, 847, 59, doi: [10.3847/1538-4357/aa7e2d](https://doi.org/10.3847/1538-4357/aa7e2d)
- Hummels, C. B., Smith, B. D., Hopkins, P. F., et al. 2019, ApJ, 882, 156, doi: [10.3847/1538-4357/ab378f](https://doi.org/10.3847/1538-4357/ab378f)
- Ipavich, F. M. 1975, ApJ, 196, 107, doi: [10.1086/153397](https://doi.org/10.1086/153397)
- Jana, R., Gupta, S., & Nath, B. B. 2020, MNRAS, 497, 2623, doi: [10.1093/mnras/staa2025](https://doi.org/10.1093/mnras/staa2025)
- Ji, S., Kereš, D., Chan, T. K., et al. 2020a, arXiv e-prints, arXiv:2011.04706. <https://arxiv.org/abs/2011.04706>
- Ji, S., Chan, T. K., Hummels, C. B., et al. 2020b, MNRAS, doi: [10.1093/mnras/staa1849](https://doi.org/10.1093/mnras/staa1849)
- Jiang, Y.-F., & Oh, S. P. 2018, ApJ, 854, 5, doi: [10.3847/1538-4357/aaa6ce](https://doi.org/10.3847/1538-4357/aaa6ce)
- Johnson, S. D., Chen, H.-W., & Mulchaey, J. S. 2015, MNRAS, 449, 3263, doi: [10.1093/mnras/stv553](https://doi.org/10.1093/mnras/stv553)
- Keeney, B. A., Stocke, J. T., Pratt, C. T., et al. 2018, ApJS, 237, 11, doi: [10.3847/1538-4365/aac727](https://doi.org/10.3847/1538-4365/aac727)
- Kroupa, P. 2001, MNRAS, 322, 231, doi: [10.1046/j.1365-8711.2001.04022.x](https://doi.org/10.1046/j.1365-8711.2001.04022.x)
- Lehner, N., Wotta, C. B., Howk, J. C., et al. 2019, ApJ, 887, 5, doi: [10.3847/1538-4357/ab41fd](https://doi.org/10.3847/1538-4357/ab41fd)
- Lehner, N., Howk, J. C., Tripp, T. M., et al. 2013, ApJ, 770, 138, doi: [10.1088/0004-637X/770/2/138](https://doi.org/10.1088/0004-637X/770/2/138)
- Liang, C. J., & Chen, H.-W. 2014, MNRAS, 445, 2061, doi: [10.1093/mnras/stu1901](https://doi.org/10.1093/mnras/stu1901)
- Menon, L., Wesolowski, F., Zheng, G., et al. 2015, in Adaptive Techniques for Clustered N-Body Cosmological Simulations. Computational Astrophysics and Cosmology, 2, 1.
- Nielsen, N. M., Churchill, C. W., & Kacprzak, G. G. 2013, ApJ, 776, 115, doi: [10.1088/0004-637X/776/2/115](https://doi.org/10.1088/0004-637X/776/2/115)
- Pakmor, R., Pfrommer, C., Simpson, C. M., & Springel, V. 2016, ApJL, 824, L30, doi: [10.3847/2041-8205/824/2/L30](https://doi.org/10.3847/2041-8205/824/2/L30)
- Peeples, M. S., Corlies, L., Tumlinson, J., et al. 2019, ApJ, 873, 129, doi: [10.3847/1538-4357/ab0654](https://doi.org/10.3847/1538-4357/ab0654)
- Pfrommer, C., Springel, V., Enßlin, T. A., & Jubelgas, M. 2006, MNRAS, 367, 113, doi: [10.1111/j.1365-2966.2005.09953.x](https://doi.org/10.1111/j.1365-2966.2005.09953.x)
- Prochaska, J. X., Weiner, B., Chen, H.-W., Mulchaey, J., & Cooksey, K. 2011, ApJ, 740, 91, doi: [10.1088/0004-637X/740/2/91](https://doi.org/10.1088/0004-637X/740/2/91)
- Prochaska, J. X., Werk, J. K., Worseck, G., et al. 2017, ApJ, 837, 169, doi: [10.3847/1538-4357/aa6007](https://doi.org/10.3847/1538-4357/aa6007)
- Rosolowsky, E. W., Pineda, J. E., Kauffmann, J., & Goodman, A. A. 2008, ApJ, 679, 1338, doi: [10.1086/587685](https://doi.org/10.1086/587685)
- Rudie, G. C., Steidel, C. C., Trainor, R. F., et al. 2012, ApJ, 750, 67, doi: [10.1088/0004-637X/750/1/67](https://doi.org/10.1088/0004-637X/750/1/67)
- Ruszkowski, M., Yang, H.-Y. K., & Zweibel, E. 2017, ApJ, 834, 208, doi: [10.3847/1538-4357/834/2/208](https://doi.org/10.3847/1538-4357/834/2/208)
- Salem, M., Bryan, G. L., & Corlies, L. 2016, MNRAS, 456, 582, doi: [10.1093/mnras/stv2641](https://doi.org/10.1093/mnras/stv2641)
- Sanchez, N. N., Tremmel, M., Werk, J. K., et al. 2020, arXiv e-prints, arXiv:2009.05581. <https://arxiv.org/abs/2009.05581>
- Sanchez, N. N., Werk, J. K., Tremmel, M., et al. 2019, ApJ, 882, 8, doi: [10.3847/1538-4357/ab3045](https://doi.org/10.3847/1538-4357/ab3045)
- Savage, B. D., & Sembach, K. R. 1991, ApJ, 379, 245, doi: [10.1086/170498](https://doi.org/10.1086/170498)
- Shen, S., Wadsley, J., & Stinson, G. 2010, MNRAS, 407, 1581, doi: [10.1111/j.1365-2966.2010.17047.x](https://doi.org/10.1111/j.1365-2966.2010.17047.x)
- Simpson, C. M., Pakmor, R., Marinacci, F., et al. 2016, ApJL, 827, L29, doi: [10.3847/2041-8205/827/2/L29](https://doi.org/10.3847/2041-8205/827/2/L29)
- Sod, G. A. 1978, Journal of Computational Physics, 27, 1, doi: [10.1016/0021-9991\(78\)90023-2](https://doi.org/10.1016/0021-9991(78)90023-2)
- Steidel, C. C., Erb, D. K., Shapley, A. E., et al. 2010, ApJ, 717, 289, doi: [10.1088/0004-637X/717/1/289](https://doi.org/10.1088/0004-637X/717/1/289)
- Stern, J., Hennawi, J. F., Prochaska, J. X., & Werk, J. K. 2016, ApJ, 830, 87, doi: [10.3847/0004-637X/830/2/87](https://doi.org/10.3847/0004-637X/830/2/87)
- Stinson, G., Seth, A., Katz, N., et al. 2006, MNRAS, 373, 1074, doi: [10.1111/j.1365-2966.2006.11097.x](https://doi.org/10.1111/j.1365-2966.2006.11097.x)
- Stocke, J. T., Keeney, B. A., Danforth, C. W., et al. 2013, ApJ, 763, 148, doi: [10.1088/0004-637X/763/2/148](https://doi.org/10.1088/0004-637X/763/2/148)
- Suresh, J., Nelson, D., Genel, S., Rubin, K. H. R., & Hernquist, L. 2019, MNRAS, 483, 4040, doi: [10.1093/mnras/sty3402](https://doi.org/10.1093/mnras/sty3402)
- Tegmark, M., Taylor, A. N., & Heavens, A. F. 1997, ApJ, 480, 22, doi: [10.1086/303939](https://doi.org/10.1086/303939)
- Thomas, T., & Pfrommer, C. 2019, MNRAS, 485, 2977, doi: [10.1093/mnras/stz263](https://doi.org/10.1093/mnras/stz263)
- Tremmel, M., Karcher, M., Governato, F., et al. 2017, MNRAS, 470, 1121, doi: [10.1093/mnras/stx1160](https://doi.org/10.1093/mnras/stx1160)
- Tripp, T. M., Meiring, J. D., Prochaska, J. X., et al. 2011, Science, 334, 952, doi: [10.1126/science.1209850](https://doi.org/10.1126/science.1209850)
- Tumlinson, J., Peeples, M. S., & Werk, J. K. 2017, ARA&A, 55, 389, doi: [10.1146/annurev-astro-091916-055240](https://doi.org/10.1146/annurev-astro-091916-055240)
- Tumlinson, J., Werk, J. K., Thom, C., et al. 2011, ApJ, 733, 111, doi: [10.1088/0004-637X/733/2/111](https://doi.org/10.1088/0004-637X/733/2/111)
- Tumlinson, J., Thom, C., Werk, J. K., et al. 2013, ApJ, 777, 59, doi: [10.1088/0004-637X/777/1/59](https://doi.org/10.1088/0004-637X/777/1/59)
- Turk, M. J., Oishi, J. S., Abel, T., & Bryan, G. L. 2012, ApJ, 745, 154, doi: [10.1088/0004-637X/745/2/154](https://doi.org/10.1088/0004-637X/745/2/154)
- Turner, M. L., Schaye, J., Steidel, C. C., Rudie, G. C., & Strom, A. L. 2015, MNRAS, 450, 2067, doi: [10.1093/mnras/stv750](https://doi.org/10.1093/mnras/stv750)

- Uhlig, M., Pfrommer, C., Sharma, M., et al. 2012, MNRAS, 423, 2374, doi: [10.1111/j.1365-2966.2012.21045.x](https://doi.org/10.1111/j.1365-2966.2012.21045.x)
- van de Voort, F., Springel, V., Mandelker, N., van den Bosch, F. C., & Pakmor, R. 2019, MNRAS, 482, L85, doi: [10.1093/mnras/sly190](https://doi.org/10.1093/mnras/sly190)
- Voit, G. M., Donahue, M., Zahedy, F., et al. 2019, ApJL, 879, L1, doi: [10.3847/2041-8213/ab2766](https://doi.org/10.3847/2041-8213/ab2766)
- Wadsley, J. W., Keller, B. W., & Quinn, T. R. 2017, MNRAS, 471, 2357, doi: [10.1093/mnras/stx1643](https://doi.org/10.1093/mnras/stx1643)
- Werk, J. K., Prochaska, J. X., Thom, C., et al. 2013, ApJS, 204, 17, doi: [10.1088/0067-0049/204/2/17](https://doi.org/10.1088/0067-0049/204/2/17)
- Werk, J. K., Prochaska, J. X., Tumlinson, J., et al. 2014, ApJ, 792, 8, doi: [10.1088/0004-637X/792/1/8](https://doi.org/10.1088/0004-637X/792/1/8)
- Werk, J. K., Prochaska, J. X., Cantalupo, S., et al. 2016, ApJ, 833, 54, doi: [10.3847/1538-4357/833/1/54](https://doi.org/10.3847/1538-4357/833/1/54)
- Zahedy, F. S., Chen, H.-W., Johnson, S. D., et al. 2019, MNRAS, 484, 2257, doi: [10.1093/mnras/sty3482](https://doi.org/10.1093/mnras/sty3482)

## APPENDIX

## A. COSMIC RAYS IN CHANGA

A.1. *Equations of Motion*

In the Lagrangian approximation, a fluid is discretized by some number of massive particles. Traditionally, each particle has a mass, velocity, and specific energy, and these quantities are evolved from their initial conditions following conservation equations. We model cosmic rays as an additional “fluid” such that each particle also contains a specific cosmic-ray energy. The evolution of these particles is governed by the equations for the conservation of mass, momentum, thermal energy, and cosmic-ray energy below.

$$\frac{D\rho}{Dt} = -\rho \nabla \cdot \mathbf{v} \quad (\text{A1})$$

$$\frac{\rho D\mathbf{v}}{Dt} + \nabla(P_{tot}) = \mathbf{g} \quad (\text{A2})$$

$$\frac{\rho Du_g}{Dt} + P_g \nabla \cdot \mathbf{v} = \Lambda_g - \Gamma_g \quad (\text{A3})$$

$$\frac{\rho Du_c}{Dt} + P_c \nabla \cdot \mathbf{v} = \nabla \cdot (\kappa_c \nabla u_c) + \Lambda_c. \quad (\text{A4})$$

We define  $\rho, \mathbf{v}, u_g$  to be the gas density, velocity, and specific internal energy (energy per mass). Similarly,  $u_c$  is the cosmic-ray specific internal energy. The gas and cosmic-ray pressures are related to the internal energy by the adiabatic index,  $P_g = (\gamma - 1)\rho u_g$  and  $P_c = (\gamma_c - 1)\rho u_c$ , where  $\gamma = 5/3$  and  $\gamma_c = 4/3$ . The total pressure of the system given by the sum of thermal and cosmic-ray pressures,  $P_{tot} = P_g + P_c$ .  $\mathbf{g}$  is the gravitational force.  $\Lambda$  and  $\Gamma$  are shorthand representations of the total source and sink terms of gas and CR energy.  $\kappa_c$  is the constant cosmic-ray diffusion coefficient. Lastly,  $t$  represents time, and  $D/Dt = \partial/\partial t + \mathbf{v} \cdot \nabla$  denotes the convective derivative.

In the SPH approach, the above equations become ordinary differential equations for the motion, internal energy, and cosmic ray energy of each particle as follows:

$$\frac{d\mathbf{v}_i}{dt} = - \sum_j m_j \left( \frac{P_{tot,i} + P_{tot,j}}{\rho_i \rho_j} + \Pi_{ij} \right) \nabla_i \bar{W}_{ij} \quad (\text{A5})$$

$$\frac{du_{g,i}}{dt} = - \sum_j m_j \left( \frac{P_{g,i} + P_{g,j}}{\rho_i \rho_j} + \Pi_{ij} \right) \nabla_i \bar{W}_{ij} \quad (\text{A6})$$

$$\frac{du_{c,i}}{dt} = - \sum_j m_j \left( \frac{P_{c,i} + P_{c,j}}{\rho_i \rho_j} + \Pi_{ij} \right) \nabla_i \bar{W}_{ij}. \quad (\text{A7})$$

In the equations above,  $m$  is the mass of an individual particle,  $\Pi$  is an artificial viscosity term and  $W$  is the general SPH kernel function. The subscript  $i$  indicates a single particle, and  $\sum_j$  is the sum of properties over that particle’s nearest neighbors. Lastly, we model the diffusion of cosmic-ray energy as

$$\frac{du_{c,i}}{dt}|_{\text{Diff}} = - \sum_j m_j \frac{\kappa_c (u_{c,i} - u_{c,j}) (\mathbf{r}_{ij} \cdot \nabla_i \bar{W}_{ij})}{\frac{1}{2}(\rho_i + \rho_j) \mathbf{r}_{ij}^2}, \quad (\text{A8})$$

where  $\mathbf{r}_{ij}$  is the distance between neighboring particles. In the following section, we test the performance of advection and diffusion of the newly added cosmic-ray fluid against analytic solutions.

## B. TESTS OF CR PERFORMANCE

B.1. *SOD Shock-tube*


We demonstrate the advection of the thermal and cosmic-ray fluids using a modified SOD shocktube test (Sod 1978; Pfrommer et al. 2006). In the initial conditions, 24,000 particles are arranged in a long, narrow three-dimensional glass (Backus 2017) with dimensions  $100 \times 100 \times 1000$  cm and periodic boundary conditions. Smoothing is done with an  $M_4$  cubic spline kernel using 64 neighboring particles. The initial discontinuity is placed at  $z = 250$  cm. To the left of the discontinuity, the gas density, pressure, and cosmic-ray pressure are initialized to  $\rho = 1 \text{ g/cm}^3$ ,  $P_g = 2/3$  kPa,  $P_c = 4/3$  kPa. To the right of the discontinuity, the gas density, pressure, and cosmic-ray pressure are initialized to  $\rho = 0.2 \text{ g/cm}^3$ ,  $P_g = 267.2$  Pa,  $P_c = 267.2$  Pa. The velocity is initialized to zero everywhere.

Figure 13 shows the evolved state of the modified SOD shocktube after  $t = 0.31$  seconds. The scattered points show the gas properties, averaged in spatial bins with  $\delta x = 5$  cm. The black dashed line shows the analytic solution for the gas density, velocity, and total pressure. The simulated shocktube follows the analytic predictions well. The slight noise to the right of the contact discontinuity ( $z \sim 400$  cm), is likely due to resolution-related particle noise.


B.2. *Diffusion*

In this work, we model cosmic-ray transport as isotropic diffusion with a constant diffusion coefficient,  $\kappa_c$ . We test the implementation of cosmic-ray diffusion by measuring the evolution of an initial overdensity of





**Figure 13.** The SOD shocktube. The state of the gas density (left), velocity (middle), and pressure (right) at  $t = 0.31$  seconds. The dashed line shows the analytic solution for the gas density, velocity, and total pressure, which is a sum of gas and cosmic-ray pressures. The shocktube was initialized with a contact discontinuity at  $x = 250$  cm at  $t = 0$ . To the left of the discontinuity, the gas density, pressure, and cosmic-ray pressure are initialized to  $\rho = 1\text{g/cm}^3$ ,  $P_g = 2/3$  kPa,  $P_c = 4/3$  kPa. To the right of the discontinuity, the gas density, pressure, and cosmic-ray pressure are initialized to  $\rho = 0.2\text{g/cm}^3$ ,  $P_g = 267.2$  Pa,  $P_c = 267.2$  Pa. The velocity is initialized to zero everywhere.



**Figure 14.** Demonstration of cosmic-ray diffusion. An initial Gaussian overdensity of cosmic-ray energy decays overtime as cosmic rays diffuse down their energy gradient with a constant diffusion coefficient. The y-axis shows radially-averaged values of the cosmic-ray energy as a function of position,  $r = x^2 + y^2$ . The black dashed lines show analytic solutions for  $t = 0$  and  $t = 0.24$  Myr.

cosmic-ray energy. We initialize 20,000 particles as an SPH glass with dimensions  $0.2 \times 0.2 \times 1$  kpc. The distribution of cosmic-ray energy density is initialized as:

$$u_c = u_{c,0} e^{-r^2/2D}, \quad (\text{B9})$$

where  $r = x^2 + y^2$  is the distance from the center. We choose constants  $u_{c,0} = 0.9$ ,  $D = 0.002$  and a diffusion coefficient of  $\kappa_c = 0.05$  kpc<sup>2</sup>/Myr. In order to only test cosmic-ray diffusion in the absence of advection, we turn off gas particle motions.

Figure 14 shows the evolution of an initial Gaussian overdensity of cosmic-ray energy assuming diffusion as the only form of cosmic-ray transport (neglecting any advection due to gas motion). Our implementation follows the expected analytic solution after  $t = 0.24$  Myr.