

2d Supersymmetric Gauge Theories, D-branes and Trialities

Sebastian Franco^{a;b;c}

^aPhysics Department, The City College of the CUNY
160 Convent Avenue, New York, NY 10031, USA

^bPhysics Program and ^cInitiative for the Theoretical Sciences
The Graduate School and University Center, The City University of New York
365 Fifth Avenue, New York NY 10016, USA

E-mail: sfranco@ccny.cuny.edu

Abstract: Engineering quantum field theories in String Theory in terms of branes is a powerful approach for understanding their dynamics. We review recent progress in the realization of 2d $\mathcal{N} = (0; 2)$ gauge theories in terms of branes. We discuss Brane Brick Models, a new class of Type IIA brane configurations which are T-dual to D1-branes over singular toric Calabi-Yau 4-folds. They fully encode the infinite class of 2d $\mathcal{N} = (0; 2)$ quiver gauge theories on the worldvolume of the D1-branes and significantly streamline their connection to the probed geometries. As an application, we explain how these constructions provide a brane realization of triality. We also comment on the realization of 2d $\mathcal{N} = (0; 1)$ theories via $\text{Spin}(7)$ orientifolds. This note is based on the author's talk at the Nankai Symposium on Mathematical Dialogues celebrating the 110th anniversary of the birth of Prof. S.-S. Chern

Contents

1	Introduction	1
2	Brane Brick Models	1
3	Triality	2
4	$\text{Spin}(7)$ Orientifolds	4

1 Introduction

Understanding the dynamics of quantum field theories in various dimensions is an active area of research. For this purpose, engineering quantum field theories in terms of branes in String and M-Theory is often a powerful tool. In this note, we will review the realization of 2d $\mathcal{N} = (0; 2)$ gauge theories on the world volume of D1-branes probing singular toric Calabi-Yau (CY) 4-folds [1, 2], as schematically shown in Figure 1.




Figure 1: D1-branes probing a CY_4 .

2 Brane Brick Models

Brane brick models are obtained from D1-branes at CY_4 singularities by T-duality. We refer the reader to [2, 5] for detailed presentations. A brane brick model is a Type IIA brane configuration consisting of D4-branes wrapping a 3-torus T^3 and suspended from an NS5-brane that wraps a holomorphic surface intersecting with T^3 . The holomorphic surface is the zero locus of the Newton polynomial defined by the toric diagram of the CY_4 . The basic ingredients of the brane setup are summarized in Table 1. The (246) directions are compactified on a T^3 . The 2d gauge theory lives on the two directions (01) common to all the branes.

Brane brick models, or equivalently their dual periodic quivers, fully encode the 2d $\mathcal{N} = (0; 2)$ quiver gauge theories on the worldvolume of D1-branes probing toric CY 4-folds. Namely, they summarize not only the quivers but also the J- and E-terms. The dictionary between brane brick models and gauge theories is summarized in Table 2.

	0	1	2	3	4	5	6	7	8	9
D4	NS5									
	{									

Table 1: Brane brick model conguration.

Brane Brick Model	Gauge Theory	Periodic Quiver
Brick	Gauge group	Node
Oriented face between bricks i and j	Bifundamental chiral eld from node i to node j	Oriented (black) arrow from node i to node j
Unoriented square face between bricks i and j	Bifundamental Fermi eld between nodes i and j	Unoriented (red) line between nodes i and j
Edge	Interaction by J - or E-term	Plaquette encoding a J - or an E-term

Table 2: Dictionary between brane brick models and 2d $N = (0; 2)$ gauge theories.

Brane brick models reduce the computation of the underlying CY_4 geometry starting from the gauge theory to a combinatorial problem, which is based on a generalization of perfect matchings [3]. Conversely, several ecient algorithms for determining the brane brick models associated to a given geometry have been developed [2, 5[8].

3 Triality

A new IR equivalence for 2d $N = (0; 2)$ gauge theory was discovered in [9]. In its simplest incarnation, which is shown in Figure 2, it relates three SQCD-like theories and was therefore dubbed triality.¹ In more general theories, triality can be locally applied to an individual gauge group, with the rest of the theory acting as a spectator. Applying triality to the same gauge group three consecutive times results in the original theory.

¹In the gure, yellow nodes indicate $SU(N_c)$ gauge groups. The theories on D1-branes have instead $U(N_c)$ gauge groups. A $U(N_c)$ version of triality was also introduced in [9]. It diers from the $SU(N_c)$ triality in Figure 2 by the presence of additional Fermi elds in the determinant representation of the gauge group for the cancellation of the Abelian anomaly. It is expected that Abelian anomalies of gauge theories on D1-branes are cancelled via a generalized Green-Schwarz mechanism. For this reason, the determinant Fermi elds are not present in those theories and triality reduces to the one considered here.




Figure 2: Triality of 2d $N = (0; 2)$ SQCD. Black arrows and red lines represent chiral and Fermi elds, respectively.

For brane brick models, triality is elegantly realized by the local transformation shown in Figure 3, which we denote cube move. More precisely, the cube move is the simplest representative of a family of local transformations implementing triality [4]. All these moves leave the underlying toric CY_4 invariant.




Figure 3: The cube move is a local transformation of a brane brick models realizing $N = (0; 2)$ triality

The correspondence between D1-brane probing toric CY_4 -folds and brane brick models can be understood systematically using mirror symmetry [10, 11]. The mirror conguration consists of D5-branes wrapping 4-spheres and the gauge theory is determined by how they intersect. In this context, triality is realized in terms of geometric transitions in the mirror geometry [11]. More generally, mirror symmetry leads to a

beautiful geometric unification of dualities in different dimensions, where the order of the duality is $n - 1$ for a CY n -fold [11, 12].

4 Spin(7) Orientifolds

As usual, it is desirable to investigate theories with less supersymmetry. The next step corresponds to $2d\ N = (0; 1)$, namely minimally supersymmetric, theories. These theories are extremely interesting, since they are barely supersymmetric and live at the borderline between non-SUSY theories and others with higher amounts of SUSY, for which powerful tools such as holomorphy become applicable. Due to the reduced SUSY, they enjoy a broad range of interesting dynamics. While there has been recent progress in their understanding, they remain relatively unexplored.

The geometric engineering of $2d\ N = (0; 1)$ gauge theories on D1-branes probing singularities was initiated in [13], where a new class of backgrounds, denoted Spin(7) orientifolds was introduced. These orientifolds are quotients of CY 4-folds by a combination of an anti-holomorphic involution leading to a Spin(7) cone and worldsheet parity. They naturally connect the view of $N = (0; 1)$ theories as real slices of $N = (0; 2)$ theories and Joyce's geometric construction of Spin(7) manifolds starting from CY 4-folds [14]. This geometric construction provides a new approach for studying $2d\ N = (0; 1)$ theories. It also gives a new perspective on the recently proposed triality of $2d\ N = (0; 1)$ gauge theories [15], in which triality translates into the fact that multiple gauge theories correspond to the same underlying orientifold [16].

Acknowledgments

I would like to thank the organizers of the Nankai Symposium on Mathematical Dialogues for putting together such an exciting meeting and for the opportunity to present my work. I am indebted to my wonderful colleagues, Cyril Closset, Dongwook Ghim, Jirui Guo, Azeem Hasan, Sangmin Lee, Alessandro Mininno, Gregg Musiker, Rak-Kyeong Song, Angel Uranga, Cumrun Vafa, Daisuke Yokoyama and Xingyang Yu, for enjoyable collaborations on the topics presented in this note. This research was supported by the U.S. National Science Foundation grants PHY-1820721, PHY-2112729 and DMS-1854179.

References

- [1] H. Garcia-Compean and A. M. Uranga, Brane box realization of chiral gauge theories in two-dimensions, *Nucl. Phys. B* **539** (1999) 329[366], [[hep-th/9806177](#)].

- [2] S. Franco, D. Ghim, S. Lee, R.-K. Seong and D. Yokoyama, 2d (0,2) Quiver Gauge Theories and D-Branes, [JHEP 09 \(2015\) 072](#), [[1506.03818](#)].
- [3] S. Franco, S. Lee and R.-K. Seong, Brane Brick Models, Toric Calabi-Yau 4-Folds and 2d (0,2) Quivers, [JHEP 02 \(2016\) 047](#), [[1510.01744](#)].
- [4] S. Franco, S. Lee and R.-K. Seong, Brane brick models and 2d (0, 2) triality, [JHEP 05 \(2016\) 020](#), [[1602.01834](#)].
- [5] S. Franco, S. Lee, R.-K. Seong and C. Vafa, Brane Brick Models in the Mirror, [JHEP 02 \(2017\) 106](#), [[1609.01723](#)].
- [6] S. Franco, S. Lee and R.-K. Seong, Orbifold Reduction and 2d (0,2) Gauge Theories, [JHEP 03 \(2017\) 016](#), [[1609.07144](#)].
- [7] S. Franco and A. Hasan, 3d printing of 2d $N = (0; 2)$ gauge theories, [JHEP 05 \(2018\) 082](#), [[1801.00799](#)].
- [8] S. Franco and A. Hasan, Calabi-Yau products: graded quivers for general toric Calabi-Yaus, [JHEP 02 \(2021\) 174](#), [[2004.13765](#)].
- [9] A. Gadde, S. Gukov and P. Putrov, (0, 2) trialities, [JHEP 03 \(2014\) 076](#), [[1310.0818](#)].
- [10] M. Futaki and K. Ueda, Tropical Coamoeba and Torus-Equivariant Homological Mirror Symmetry for the Projective Space, [Commun. Math. Phys. 332 \(2014\) 53](#)[87].
- [11] S. Franco, S. Lee, R.-K. Seong and C. Vafa, Quadrality for Supersymmetric Matrix Models, [JHEP 07 \(2017\) 053](#), [[1612.06859](#)].
- [12] S. Franco and G. Musiker, Higher Cluster Categories and QFT Dualities, [Phys. Rev. D 98 \(2018\) 046021](#), [[1711.01270](#)].
- [13] S. Franco, A. Mininno, A. M. Uranga and X. Yu, 2d $N = (0; 1)$ Gauge Theories and Spin(7) Orientifolds, [2110.03696](#).
- [14] D. Joyce, A new construction of compact 8-manifolds with holonomy Spin(7), J. Di. Geom. 53 (1999) 89{130, [[math/9910002](#)].
- [15] S. Gukov, D. Pei and P. Putrov, Trialities of minimally supersymmetric 2d gauge theories, [JHEP 04 \(2020\) 079](#), [[1910.13455](#)].
- [16] S. Franco, A. Mininno, A. M. Uranga and X. Yu, Spin(7) orientifolds and 2d $N = (0, 1)$ triality, [JHEP 01 \(2022\) 058](#), [[2112.03929](#)].