

1 **Title:** Characterization of microsatellite markers for the duckweed *Spirodela polyrhiza* and
2 *Lemna minor* tested on samples from Europe or the United States of America.

3
4 **Running title:** *Spirodela polyrhiza* and *Lemna minor* microsatellites

5
6 **Authors:** Jae E. Kerstetter^{1,2}, Andrea L. Reid³, Joshua T. Armstrong^{1,4}, Taylor A. Zallek¹, Trapper
7 T. Hobbie¹, and Martin M. Turcotte¹

8
9 ¹ Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA, USA 15260

10 ² Department of Entomology, Rutgers University, New Brunswick, NJ, USA 08901

11 ³ Department of Geodesy and Geomatics Engineering, University of New Brunswick,
12 Fredericton, NB, Canada

13 ⁴ Center for Environmental Studies, Virginia Commonwealth University, Richmond, VA,
14 USA 23284

15
16 Corresponding author: Martin Turcotte turcotte@pitt.edu

17
18 **Keywords:** Lemnaceae, simple sequence repeats, genotyping, genetic identification, molecular
19 markers

20 21 **Abstract**

22 Microsatellite primers are a valuable tool to use for both observational and experimental
23 studies in numerous taxa. Here, we develop 18 and 16 microsatellite markers for the
24 widespread duckweeds *Lemna minor* and *Spirodela polyrhiza*, respectively. All 18 *L. minor*
25 primers and 12 of the 16 *S. polyrhiza* primers amplified polymorphic loci when tested on
26 samples from Europe or Western Pennsylvania, USA.

27 28 29 **Introduction**

30 The globally distributed duckweed family (Lemnaceae) or subfamily (Lemnoideae) is 31
32 composed of 36 species (Bog *et al.* 2020) of very small floating or submerged aquatic plants
33 (Landolt, 1986; Sree *et al.* 2016). Duckweeds have a long history of scientific study given their
34 highly specialized morphology, widespread distribution, high abundance, and production of the
35 world's smallest flowers (Jacobs, 1947; Hillman 1961; Landolt, 1986; Landolt, 1992). More
36 recently, there has been an explosion in research interest given their potential applied uses
37 including for agricultural feed (Cheng and Stomp, 2009), bioremediation (Gupta and Prakash,
38 2013; Ekperusi *et al.* 2019), and biofuel production (Cui and Cheng, 2015). Furthermore, their
39 use as a model system to experimentally study numerous topics in ecology and evolutionary
40 biology is quickly expanding (Laird and Barks, 2018). This growing basic and applied interest
41 stems from their ability to reproduce clonally very quickly with population doubling times in as
42 little as 1.5 days (Ziegler *et al.* 2015). In addition, they are amenable to large scale
43 manipulative experiments in both the lab and field mesocosms (Armitage and Jones, 2019; Hart
44 *et al.* 2019; Tan *et al.* 2021; O'Brien *et al.* 2022), and have growing genomic data and tools
(Wang *et al.* 2014; Ho *et al.* 2019; Xu *et al.* 2019; Cao *et al.* 2020) and characterization of their

45 microbiome and herbivore communities (Acosta *et al.* 2020; Subramanian and Turcotte, 2020).
46 Finally, duckweed express variation in numerous traits across species and among genotypes
47 (clonal lineages) within species (Van Steveninck *et al.* 1992; Hart *et al.* 2019; Chen *et al.* 2020;
48 Hitsman and Simons, 2020; Anneberg *et al.* 2023). Therefore, being able to identify genotypes
49 may also be beneficial in many ecological studies to assess differences in traits among
50 genotypes and to determine how these genotypes may respond to different environmental
51 conditions.

52 Genetic markers, such as microsatellite markers, are important tools to study
53 population genetics. Microsatellites, also known as simple sequence repeats ‘SSRs’, are tandem
54 repeats two to 10 base pairs in length, that are flanked by conserved sequences and occur
55 ubiquitously throughout eukaryotic genomes (Tautz and Renz, 1984). They are highly
56 informative as locus-specific genetic markers due to their high abundance, high reproducibility,
57 co-dominance, and polymorphic nature (Morgante and Olivieri, 1993; Powell *et al.* 1996). The
58 length of the sequence repeats can be determined through PCR amplification using primers
59 specific to their flanking regions; variation in PCR product length is a function of the number of
60 repeated sequences. The high levels of polymorphisms observed in SSR markers (Tautz, 1989;
61 Schlotterer and Tautz, 1992) and the relative ease of detection of these polymorphisms by PCR
62 amplification has led to the wide applications of microsatellites as genetic markers (Vieira *et al.*
63 2016). Such within species markers have numerous applications including quantifying
64 biogeographic distributions, population genetic structure, evolutionary history, and mating
65 systems.

66 Moreover, a growing number of experimental evolution studies use such markers to
67 track changes in genotypic composition of asexually reproducing populations over multiple
68 generations (e.g., Turcotte *et al.* 2011; Agrawal *et al.* 2013; Hart *et al.* 2019) in large replicated
69 experiments for which genotype-by-sequencing remains too costly. These cost savings are
70 magnified when several loci can be genotyped in the same reaction (multiplexed; Markoulatos
71 *et al.* 2002). Here, we report on the development of new microsatellite markers for two
72 commonly studied and widespread duckweed species: the common duckweed *Lemna minor*
73 (L. Schleid) and the greater duckweed *Spirodela polyrhiza* (L. Schleid)

74 With the growing interest in duckweed, microsatellite markers have been developed for
75 a few duckweed species. Wani *et al.* (2014) developed nine polymorphic and 24 monomorphic
76 haplotype cpDNA-based microsatellite primers for *L. minor*. Xu *et al.* (2018) developed 60
77 microsatellite primers for *Spirodela polyrhiza*, 19 of which were polymorphic within three
78 populations of *S. polyrhiza* from China. Feng *et al.* (2017) developed three microsatellite
79 primers for the identification of *S. polyrhiza* and *Landoltia punctata* haplotypes. More recently,
80 Fu *et al.* (2020) developed 70 microsatellite primers within coding regions for *L. gibba*. It is
81 important to continue developing and reporting new microsatellite markers as populations can
82 differ in which markers function (e.g. due to null alleles) and are polymorphic (Chapuis and
83 Estoup, 2007).

84 Here we report the successful development of 18 *L. minor* and 16 *S. polyrhiza*
85 microsatellite markers. A small subset of these microsatellite primers were used to differentiate
86 genotypes in our experimental studies on evolutionary-coexistence (Hart *et al.* 2019). In
87 addition, we report genotyping results using these markers on individuals sampled in Europe
88 and the United States of America (USA).

89
90

91 **Materials & Methods:**

92 *Sample Collection*

93 Our objective when sampling was not to genetically characterize duckweed populations, but
94 instead find genotypes that differ in ecologically relevant traits to use in various experiments.
95 Thus, we genotyped few individuals from numerous bodies of water in various locations.
96 Primers were developed at ETH Zurich (Europe) and the University of Pittsburgh (USA), and thus
97 were tested on different collections of duckweeds. We collected duckweeds from numerous
98 still bodies of water (e.g. ponds, lakes, wetlands) primarily in Switzerland and Western
99 Pennsylvania (USA), however a few samples were also collected from the Netherlands and
100 Germany. In addition, some European duckweed samples were obtained from the Landolt
101 Duckweed Collection (formerly in Zurich, Switzerland) were included (see Supplemental Tables
102 S1 and S2 for collection locations). Given the two-part development of the primers, some
103 duckweed samples were only tested on the primers developed in that country (as noted in
104 Tables S1 & S2).

105 Duckweeds mostly reproduce clonally via meristematic pockets from which clonal
106 daughters emerge, creating clonal clusters of 1-8 individuals that eventually split into smaller
107 clusters (Landolt, 1986). We sampled single duckweed clusters and established isofemale
108 laboratory colonies from these clusters. We then sterilized each colony using sodium
109 hypochlorite following a method adapted from Barks *et al.* (2018). From each colony, we put
110 single individuals into individual sterile petri dishes (one individual per dish) containing sterile
111 0.5 strength Schenk and Hildebrandt growth medium (containing macro- and micronutrients as
112 described by Schenk and Hildebrandt, 1972) supplemented with sucrose (6.7 g/L), yeast extract
113 (0.067 g/L), and tryptone (0.34 g/L) for 24 hours to encourage algal and bacterial spore
114 germination. Then each individual was exposed to one of an array of concentrations of sodium
115 hypochlorite (0.3% or 0.5%) for varying amounts of time (3 or 6 minutes for *L. minor*, 4 or 7
116 minutes for *S. polyrhiza* respectively), then rinsed with autoclaved distilled water and allowed
117 to grow (Barks *et al.* 2018). Sterile colonies were maintained in sterile 0.25 strength Schenk and
118 Hildebrandt media (1972) without the additional supplements in room temperature
119 laboratories or growth chambers under plant grow lights. These collections do not reproduce
120 sexually under lab conditions.

121

122 *Microsatellite Marker Development*

123 A total of 18 *L. minor* and 16 *S. polyrhiza* microsatellite markers were developed across Europe
124 or the USA. We downloaded the whole genome shotgun sequence data for *S. polyrhiza* strain
125 7498 from the National Center for Biotechnology Information's GenBank database (accession
126 ATDW01000001.1) deposited by Wang *et al.* (2014). For *L. minor*, a draft genome (strain 8627)
127 was downloaded from www.lemna.org on October 16th 2015 (genome draft Im8627.ASMv0.1)
128 A recent study using Tubulin Based Polymorphism suggests that this lineage is in fact an
129 interspecific hybrid of *L. japonica* and *L. turionifera* both closely related to *L. minor* (Braglia *et al.*
130 *et al.* 2021). The species identity for most samples on which we report below have been
131 confirmed using morphology and/or barcoding (Fazekas *et al.* 2012; Barks *et al.* 2018). While
132 some microsatellite markers are known to amplify across more than one duckweed species (Xu

133 *et al.* 2018), we have not yet explicitly tested these markers against other species. Using
134 MSATCOMMANDER (version 1.0.8, Faircloth 2008), we identified microsatellite loci using the
135 default settings, except we avoided mononucleotide repeat motifs. We then selected loci that
136 would produce products of different lengths, had different motif lengths, and were found on
137 different contigs. The 5' end of forward primers were labeled with one of several fluorescent
138 dyes from various suppliers.

139 Primers developed at ETH Zurich were M13-tailed to reduce cost during development
140 (Boutin-Ganache *et al.* 2001). This entailed adding the full or a partial M13 sequence of
141 TGTAACGACGGCCAGT for the *S. polyrhiza* primers and GGAAACAGCTATGACCAT for *L. minor*
142 primers to the 5' end of the forward primer. The M13-labeled forward primers were used in
143 combination with a M13 primer that had the same sequence but was fluorescently dye-labeled
144 at its 5' end. Some primers amplified less polymorphic loci or did not amplify as consistently as
145 others. For these primers, we only have fragment lengths that include the M13 tail (see Tables
146 1 & 2), and we estimate that this lengthens the PCR product by 12-19 base pairs. For most
147 primers however, following initial testing with M13, we ordered new labeled primers that did
148 not include the M13 tail.

149 At least 20 duckweed samples were tested using each primer. European duckweed
150 samples were tested across 7 *L. minor* and 16 *S. polyrhiza* primers, and USA duckweed samples
151 were tested across 15 *L. minor* and 4 *S. polyrhiza* primers (see Tables S1 and S2 for details).

152
153 **Table 1:** *Lemna minor* microsatellite markers and motifs including optimized MgCl₂
154 concentrations and annealing temperatures (T_A). In addition, we report marker success rate
155 which is the number of samples successfully genotyped divided by those attempted, the
156 number of unique alleles, and number of unique genotypes for each primer. Average
157 heterozygosity (H) is the fraction of individuals that are heterozygotic for each primer. See
158 Table S1 for specific allele values. Alleles lengths with an * denote that these lengths include
159 the M13 tail sequence.

160

Primers	Forward Primer (5'-3') Reverse Primer (5'-3')	Motif	MgCl ₂ (mM)	T _A (°C)	Observed Product Length	Marker Success Rate	Unique Alleles	Unique Genotypes	Average H
LmR.1.A	F: GTTCCTAAGGATTCATCACC R: TACGAGGAGGGACACGAG	AAG	2.0	60	178-185*	75/81	2	2	0
LmR.4.A	F: AGTGGCTACGAACGGAAGAG R: AGAGGAACGTTGTGTCTGGG	AAG	0.9	63	219-234	28/28	5	5	0.036
LmR.4.B	F: CTTATTGGATCTTCGCGCCG R: AAGATATCTGACGGCGTTGG	AG	1.2	63	366-392	28/28	6	6	0.071
LmR.5.C	F: GATGCCAGTAGATCCGGC R: ACGCCTGAACACGATTGATG	AGAT	2.0	60	320-444	104/109	25	41	0.846
LmR.8.B	F: TGTACTIONCATCTGTGGGCGAG R: AACAAATTTGGCCACCGTCAG	AGAT	1.2	63	306-376	28/28	10	9	0.036
LmR.8.C	F: GACAACCTTAGGGTGACCGC R: GGAGTGAGAGCTGAGGACTG	AGG	1.2	60	435-450	28/28	3	3	0
LmR.10.A	F: TCCTTTCTCGTGTCTCCAG R: ATGCCCGACCTAGTCC	AG	2.0	60	222-254*	31/81	4	5	0.032
LmR.10.C	F: CTCTCTTTCTCTCCACGG R: ATGCAACCCTCTAGCCG	AGAT	2.0	60	179-254*	79/81	4	4	0.278
LmR.12.B	F: TCTCTGCTGACCGACTCAAG R: GCCGTTGGATCTTTCTCAG	AT	1.2	60	274-320	27/28	8	9	0.111
LmR.14.A	F: TCGCACTAGAGAGATGGGTG R: TCCATTACCAGGATGCGAG	AAT	1.2	60	261-270	24/28	3	3	0.042
LmR.14.B	F: CATGCCAGGTAAATGCCCTC R: TCGAGCTCCTTCTCAAACC	ATC	0.9	63	430-440	28/28	3	3	0
LmR.14.C	F: TTCGTTCGAGGGTATGAGCTG R: TCTCTATTTGACACGCGCG	AG	0.9	63	162-178	28/28	7	7	0.036
LmR.15.A	F: GTGACAGCGTATCCTTGTGC R: CAGCGGCAAGATCATCAAG	ATC	1.2	60	222-285	109/109	13	15	0.578
LmR.15.B	F: TCGAGCTAATCAGTGGAGCC R: GAGTGCTCGGCTTGACTTTC	AG	1.2	60	170-210	104/109	13	25	0.692
LmR.15.C	F: CATGTTCCCACTTGTGAC R: AAGGAAGAGGGAGCAAGGG	AT	1.2	60	368-400	109/109	14	26	0.743
LmR.26.B	F: GTGTCTCCGAGAGCCTACAG R: TTTAAAGCTCGGTGGTCCC	AG	1.2	63	283-329	28/28	10	7	0.964
LmR.31.A	F: GGTGATCTCAGGTAGCCGAG R: TGAGATCACCCTGTCTGCC	AAG	0.9	63	402-432	26/28	5	6	0.077
LmR.31.B	F: AGTCGGCATAGTACTTCCCG R: CTTCTCAAGACCGTTCCGC	AAG	1.2	63	155-239	28/28	7	9	0.071

162
163
164

Table 2: *Spirodela polyrhiza* microsatellite markers and sampling results as described in Table 1 with allele calls in Table S2.

Primers	Forward Primer (5'-3') Reverse Primer (5'-3')	Repeat Motif	MgCl ₂ (mM)	T _A (°C)	Observed Product Length	Marker Success Rate	Unique Alleles	Unique Genotypes	Average H
Sp.1035	F: TGCTTGGTCACTCTGTCTG R: CGATTCTAGCTCCTCTGC	AT	1.2	60	361-369	42/42	4	5	0.381
Sp.1467	F: AGTTGAGGAAGCTTCATGG R: ATTACCTCCAGCACCTCTCC	AG	2.0	58	386-411*	9/20	5	4	0.444
Sp.2597	F: TCCATTACCCACAGTCTCC R: TCATTCCACCACGTCCAC	AT	2.0	58	397-399*	14/20	2	2	0.071
Sp.5050	F: ATTAACCTTGGGCGCAGAG R: TAGCAGCAGAGTGTGAGGG	AAT	2.0	58	287*	14/20	1	1	0
Sp.5250	F: AAACGAGACCTCTACGCC R: GCCTGCGAGTAATATGTGC	ATGCCC	2.0	58	385*	19/20	1	1	0
Sp.7286	F: CGAATATGCCGAGGAATGC R: TCCTCGATCTGCCGCTTAG	CG	1.2	60	386-394	42/42	5	7	0.310
Sp.7688	F: AATGGTTGACTCGACGCTG R: TCACACCGCCATAATTCGC	AGC	2.0	58	199-211*	19/20	2	2	0.158
Sp.7814	F: AGTGTAGGGTGCAGCTGTG R: TTCGTGAAAGGCCTAGCAC	AG	1.2	60	220-228	42/42	5	6	0.095
Sp.7908	F: GAGACACATCATTGCCAGC R: TAATGCAGGCCACACAACC	AG	2.0	58	234-236	20/20	2	2	0.850
Sp.8563	F: GTATTGGGTGGGCAAATCG R: AAGGGATAGGGTCGTGTCC	AG	2.0	58	350-354*	14/20	3	4	0.071
Sp.8910	F: CCTTCCTACGTTGACTCCC R: GCGTTTCTCTGATCAGCACC	ACG -> CGT	2.0	58	358	20/20	1	1	0
Sp.9307	F: GGGAGCGAGCTGTATGAAG R: TTTCAACACCCTCACCATGC	AG	2.0	58	450-452*	9/20	2	3	0.444
Sp.9311	F: GTGAGAAAAGGAAAGGTGGC R: TGCTCAGGATTCTATGGGCC	AG	2.0	58	253-255*	10/20	2	3	0.400
Sp.Pso27	F: AAGGGTTTCAGTGGGACG R: CTCGCCTTCTCGTACATCATC	AAG	2.0	58	133*	9/20	1	1	0
Sp.Pso31	F: TCCACCGTCTCCCTGTAATG R: CCACTCCCTCGTCGTGAAG	AAG	1.2	60	240-270	32/42	7	7	0.406
Sp.Pso32	F: TGCTGGCGATGTCAATGTTG R: CTTCAGCACCAAGAGAGCTC	ATC	2.0	58	377-380*	19/20	2	3	0.895

165

166 *Microsatellite Amplification and Optimization*

167 All duckweed collections were extracted and genotyped at least twice by first sampling 4 to 10
168 individuals from each monoclonal collection and lyophilizing them for 24 hours. We then
169 extracted DNA using a modified CTAB-based method by Healey *et al.* (2014). Each
170 microsatellite primer was first optimized for annealing temperatures and Magnesium chloride
171 (MgCl₂) concentration. The most commonly used PCR recipes and thermocycler conditions are
172 presented here, and any primer specific differences are listed in Tables 1 and 2. For primers
173 developed in Europe, the conditions were the following: PCR amplification was conducted in 15
174 µL volume reactions containing 3 µL of template DNA, 3µL of 5X Colorless GoTaq Flexi buffer
175 (Promega, USA), 2.0 mM MgCl₂, 0.2 mM dNTP mix, 0.05 µM of forward primer, 0.2 µM of
176 reverse primer, 0.2 µM of M13 tagged with a fluorescent probe (e.g.: 5' 6-FAM or 5' HEX), and 1
177 unit of GoTaq G2 Flexi DNA Polymerase (Promega, USA). DNA concentrations was rarely
178 quantified as amplification was often successful across a range of values (e.g. 2- 40 ng/µL).
179 Thermocycling conditions for both *S. polyrhiza* and *L. minor* from Europe that were M13 tagged
180 were: initial denaturing at 94°C for 5 min, followed by 30 cycles of 1 min at 94°C, 1 min at 60°C,
181 1 min at 72°C, followed by eight M13 cycles consisting of 1 min at 94°C, 1 min at 53°C, 1 min at
182 72°C, followed by a final extension at 72°C for 10 minutes. For all primers developed in the USA,
183 the conditions were the following: PCR amplification was conducted in 15 µL volume reactions
184 containing 3 µL of template DNA, 3µL of 5X Colorless GoTaq Flexi buffer (Promega, USA), 1.2
185 mM MgCl₂, 0.2 mM dNTP mix, 0.08 µg/µL of Bovine Serum Albumin (BSA), 0.2 µM of each
186 forward and reverse primer, and 1 unit of GoTaq G2 Flexi DNA Polymerase (Promega, USA).
187 Thermocycling conditions for *S. polyrhiza* were: initial denaturing at 94°C for 5 min, followed by
188 34 cycles of: 1 min of denaturing at 94°C, 1 min of annealing at 60°C, and 1 min of extension at
189 72°C, followed by a final extension at 72°C for 10 minutes. For *L. minor*, touchdown PCR was
190 employed: with an initial denaturation of 94 °C for 5 min, followed by five cycles of
191 denaturation (94 °C, 1 min), annealing (67 °C, 1 min; decreasing by 1 °C per cycle), and
192 extension (72 °C, 1 min). Then 25 cycles of 1 min at 94°C, 1 min at 63°C, and 2 min at 72°C,
193 followed by a final extension at 72°C for 15 minutes.

194 Fragment length analyses for all primers were conducted on ABI 3730 Genetic Analyzers
195 (Applied Biosystems) at either the ETH Zurich Genetic Diversity Center (Switzerland), Keck DNA
196 Sequencing Lab at Yale University (USA), or the University of Pittsburgh Genomics Research
197 Core (USA), using either GeneScan™ 500 or 600 LIZ™ Dye Size Standards (Applied Biosystems).
198 Allele calls were made using either Geneious (version 9.1.6, Keaser *et al.* 2012) or GeneMarker
199 software (version 3.0.0, SoftGenetics, State College, Pennsylvania).

200

201

202 **Results and Discussion**

203 We successfully developed 18 *L. minor* and 16 *S. polyrhiza* microsatellite primers (Tables 1 & 2)
204 which were tested on samples of duckweeds from Europe or Western Pennsylvania (USA).
205 Some markers were easier to utilize than others (Tables 1 & 2). All markers amplified in some
206 samples; of these, all 18 *L. minor* primers and 12 of the 16 *S. polyrhiza* primers amplified
207 polymorphic loci, having more than one allele. Moreover, these polymorphic loci differ in
208 product length and can be used in multiplex reactions to increase efficiency and lower
209 genotyping costs. We also found that some loci were much more polymorphic than others. For

210 *L. minor*, these include loci amplified by primers LmR.5.C, LmR.8.B, LmR.15.A, LmR.15.B,
211 LmR.15.C, and LmR.26.B, some of which showed high allele richness even when tested on only
212 28 samples (Table 1). For *S. polyrhiza* these included loci amplified by primers Sp.1467, Sp.7286,
213 Sp.7814, and Sp.Pso31 (Table 2). Monomorphic loci may still be useful in different duckweed
214 populations (Chapuis and Estoup, 2007). Many microsatellite loci also showed heterogeneity
215 (Tables S1 and S2) which helps make the primers more informative to distinguish genotypes.
216 We note that some primers developed in one continent were not tested on samples from the
217 other continent (see footnotes in Tables S1 & S2); we suspect these primers will work across
218 continents given patterns observed in the others, but this remains to be tested.

219 Comparing between species, we see that although we tested less samples of *S.*
220 *polyrhiza*, it still has much lower allelic and genotypic richness across most primers. This is
221 consistent with our own recent large-scale sampling (Hobble *et al. In prep*) as well as other
222 studies using different genotyping methods, that similarly found low genetic diversity in *S.*
223 *polyrhiza* (Bog *et al.* 2015; Xu *et al.* 2015; Feng *et al.* 2017). It has been hypothesized that this
224 low genetic variation in *S. polyrhiza* is due to its low mutation rate (Xu *et al.* 2019). In addition,
225 primers differed greatly in average observed heterozygosity, but species had similar mean
226 heterozygosities (0.256 for *L. minor* and 0.283 for *S. polyrhiza*). Given that our sampling was
227 designed to find unique genotypes (shallow and widespread) and not characterize populations,
228 we limit our discussion or quantification of population genetic indices. The primers we
229 developed can help researchers address various ecological and evolutionary questions as well
230 as better identify and catalogue genotypes for applied activities.

231

232 **Supplemental Table Captions**

233

234 **Table S1:** *Lemna minor* sample collection sites and allele lengths.

235

236 **Table S2:** *Spirodela polyrhiza* sample collection sites and allele lengths.

237

238

239 **Acknowledgements**

240 We thank Walter Lämmli for sharing duckweed lineages from the Landolt Duckweed
241 Collection. We are grateful to the former Levine Plant Ecology Group at ETH Zurich and the
242 Turcotte Lab for their assistance in maintaining collections. We thank ETH Zurich Genetic
243 Diversity Center and Mary Janecka for help troubleshooting primer development. M.M.T. was
244 supported by the ETH Zurich Center for Adaptation to Changing Environments and now by an
245 NSF grant DEB-1935410.

246

247

248 **References**

249 Acosta K., Xu J., Gilbert S., Denison E., Brinkman T., Lebeis S., and Lam E. (2020). Duckweed
250 hosts a taxonomically similar bacterial assemblage as the terrestrial leaf microbiome. PLOS ONE
251 15, e0228560. doi:10.1371/journal.pone.0228560

252

- 253 Agrawal A. A., Johnson M. T. J., Hastings A. P., and Maron J. L. (2013). A field experiment
254 demonstrating plant life-history evolution and its eco-evolutionary feedback to seed predator
255 populations. *The American Naturalist* 181, S35–S45. doi:10.1086/666727
256
- 257 Anneberg T. J., O’Neill E. M., Ashman T.-L., and Turcotte M. M. (2023). Polyploidy impacts
258 population growth and competition with diploids: multigenerational experiments reveal key life
259 history tradeoffs. *New Phytologist* n/a. doi:10.1111/nph.18794
260
- 261 Armitage D. W., and Jones S. E. (2019). Negative frequency-dependent growth underlies the
262 stable coexistence of two cosmopolitan aquatic plants. *Ecology* 100, e02657.
263 doi:10.1002/ecy.2657
264
- 265 Barks P. M., Dempsey Z. W., Burg T. M., and Laird R. A. (2018). Among-strain consistency in the
266 pace and shape of senescence in duckweed. *Journal of Ecology* 106, 2132–2145.
267 doi:10.1111/1365-2745.12937
268
- 269 Bog M., Lautenschlager U., Landrock M. F., Landolt E., Fuchs J., Sowjanya Sree K., Oberprieler C.,
270 and Appenroth K.-J. (2015). Genetic characterization and barcoding of taxa in the genera
271 *Landoltia* and *Spirodela* (Lemnaceae) by three plastidic markers and amplified fragment length
272 polymorphism (AFLP). *Hydrobiologia* 749, 169–182. doi:10.1007/s10750-014-2163-3
273
- 274 Bog M., Sree K. S., Fuchs J., Hoang P. T. N., Schubert I., Kuever J., Rabenstein A., Paolacci S.,
275 Jansen M. A. K., and Appenroth K.-J. (2020). A taxonomic revision of *Lemna* sect. *Uninerves*
276 (Lemnaceae). *TAXON* 69, 56–66. doi:10.1002/tax.12188
277
- 278 Boutin-Ganache I., Raposo M., Raymond M., and Deschepper C. F. (2001). M13-tailed primers
279 improve the readability and usability of microsatellite analyses performed with two different
280 allele- sizing methods. *BioTechniques* 31, 25–28. doi:10.2144/013111bm02
281
- 282 Braglia L., Lauria M., Appenroth K. J., Bog M., Breviario D., Grasso A., Gavazzi F., and Morello L.
283 (2021). Duckweed species genotyping and interspecific hybrid discovery by tubulin-based
284 polymorphism fingerprinting. *Frontiers in Plant Science* 12, 625670.
285 doi.org/10.3389/fpls.2021.625670
286
- 287 Cao X. H., Fourounjian P., and Wang W. (2020). *The Duckweed Genomes*. (Switzerland: Springer
288 Cham), 185 p. doi:10.1007/978-3-030-11045-1
289
- 290 Chapuis M.-P., and Estoup A. (2007). Microsatellite null alleles and estimation of population
291 differentiation. *Molecular Biology and Evolution* 24, 621–631. doi:10.1093/molbev/msl191
292
- 293 Chen D., Zhang H., Wang Q., Shao M., Li X., Chen D., Zeng R., and Song Y. (2020). Intraspecific
294 variations in cadmium tolerance and phytoaccumulation in giant duckweed (*Spirodela*
295 *polyrhiza*). *J Hazard Mater* 395, 122672. doi:10.1016/j.jhazmat.2020.122672
296

- 297 Cheng J. J., and Stomp A.-M. (2009). Growing duckweed to recover nutrients from wastewaters
298 and for production of fuel ethanol and animal Feed. *CLEAN – Soil, Air, Water* 37, 17–26.
299 doi:10.1002/clen.200800210
300
- 301 Cui W., and Cheng J. J. (2015). Growing duckweed for biofuel production: a review. *Plant*
302 *Biology* 17, 16–23. doi:10.1111/plb.12216
303
- 304 Ekperusi A. O., Sikoki F. D., and Nwachukwu E. O. (2019). Application of common duckweed
305 (*Lemna minor*) in phytoremediation of chemicals in the environment: State and future
306 perspective. *Chemosphere* 223, 285–309. doi:10.1016/j.chemosphere.2019.02.025
307
- 308 Fazekas A. J., Kuzmina M. L., Newmaster S. G., and Hollingsworth P. M. (2012). DNA Barcoding
309 Methods for Land Plants. In *DNA Barcodes: Methods and Protocols*. (Totowa, NJ USA: Humana
310 Press), 223–252. doi:10.1007/978-1-61779-591-6_11
311
- 312 Feng B., Fang Y., Xu Z., Xiang C., Zhou C., Jiang F., Wang T., and Zhao H. (2017). Development of
313 a new marker system for identification of *Spirodela polyrhiza* and *Landoltia punctata*.
314 *International Journal of Genomics* 2017, e5196763. doi:10.1155/2017/5196763
315
- 316 Fu L., Ding Z., Kumpeangkeaw A., Tan D., Han B., Sun X., and Zhang J. (2020). De novo assembly,
317 transcriptome characterization, and simple sequence repeat marker development in duckweed
318 *Lemna gibba*. *Physiol Mol Biol Plants* 26, 133–142. doi:10.1007/s12298-019-00726-9
319
- 320 Gupta C., and Prakash D. (2013). Duckweed: an effective tool for phyto-remediation.
321 *Toxicological & Environmental Chemistry* 95, 1256–1266. doi:10.1080/02772248.2013.879309
322
- 323 Hart S. P., Turcotte M. M., and Levine J. M. (2019). Effects of rapid evolution on species
324 coexistence. *Proceedings of the National Academy of Sciences* 116, 2112–2117.
325 doi:10.1073/pnas.1816298116
326
- 327 Healey A., Furtado A., Cooper T., and Henry R. J. (2014). Protocol: a simple method for
328 extracting next-generation sequencing quality genomic DNA from recalcitrant plant species.
329 *Plant Methods* 10, 21. doi:10.1186/1746-4811-10-21
330
- 331 Hillman W. S. (1961). The Lemnaceae, or duckweeds. *Bot. Rev* 27, 221–287.
332 doi:10.1007/BF02860083
333
- 334 Hitsman H. W., and Simons A. M. (2020). Latitudinal variation in norms of reaction of phenology
335 in the greater duckweed *Spirodela polyrhiza*. *Journal of Evolutionary Biology* 33, 1405–1416.
336 doi:10.1111/jeb.13678
337
- 338 Ho E. K. H., Bartkowska M., Wright S. I., and Agrawal A. F. (2019). Population genomics of the
339 facultatively asexual duckweed *Spirodela polyrhiza*. *New Phytologist* 224, 1361–1371.
340 doi:10.1111/nph.16056

341
342 Jacobs D. L. (1947). An ecological life-history of *Spirodela polyrhiza* (greater duckweed) with
343 emphasis on the turion phase. *Ecological Monographs* 17, 437–469. doi:10.2307/1948596
344
345 Kearse M., Moir R., Wilson A., Stones-Havas S., Cheung M., Sturrock S., Buxton S., Cooper A.,
346 Markowitz S., Duran C., Thierer T., Ashton B., Meintjes P., and Drummond A. (2012). Geneious
347 Basic: An integrated and extendable desktop software platform for the organization and
348 analysis of sequence data. *Bioinformatics* 28, 1647–1649. doi:10.1093/bioinformatics/bts199
349
350 Laird R. A., and Barks P. M. (2018). Skimming the surface: duckweed as a model system in
351 ecology and evolution. *Am J Bot* 105, 1962–1966. doi:10.1002/ajb2.1194
352
353 Landolt E. (1986). Biosystematic investigations in the family of duckweeds (Lemnaceae). Vol. 2,
354 The family of Lemnaceae—A Monographic Study, Vol. 1 (Zürich, ETH Zürich), 566 p.
355
356 Landolt E. (1992). Lemnaceae duckweed family. *Journal of the Arizona-Nevada Academy of*
357 *Science* 26, 10–14.
358
359 Markoulatos P., Siafakas N., and Moncany M. (2002). Multiplex polymerase chain reaction: A
360 practical approach. *J Clin Lab Anal* 16, 47–51. doi:10.1002/jcla.2058
361
362 Morgante M., and Olivieri A. M. (1993). PCR-amplified microsatellites as markers in plant
363 genetics. *Plant J* 3, 175–182.
364
365 O’Brien A. M., Yu Z. H., Pencer C., Frederickson M. E., LeFevre G. H., and Passeport E. (2022).
366 Harnessing plant-microbiome interactions for bioremediation across a freshwater urbanization
367 gradient. *Water Research* 223, 118926. doi:10.1016/j.watres.2022.118926
368
369 Powell W., Machray G. C., and Provan J. (1996). Polymorphism revealed by simple sequence
370 repeats. *Trends in Plant Science* 1, 215–222. doi:10.1016/1360-1385(96)86898-1
371
372 Schenk R. U., and Hildebrandt A. C. (1972). Medium and techniques for induction and growth of
373 monocotyledonous and dicotyledonous plant cell cultures. *Can. J. Bot.* 50, 199–204.
374 doi:10.1139/b72-026
375
376 Schlötterer C., and Tautz D. (1992). Slippage synthesis of simple sequence DNA. *Nucleic Acids*
377 *Res* 20, 211–215.
378
379 Sree K. S., Bog M., and Appenroth K. J. (2016). Taxonomy of duckweeds (Lemnaceae), potential
380 new crop plants. *Emirates Journal of Food and Agriculture* 291–302. doi:10.9755/ejfa.2016-01-
381 038
382

- 383 Subramanian S. K., and Turcotte M. M. (2020). Preference, performance, and impact of the
384 water-lily aphid on multiple species of duckweed. *Ecological Entomology* 45, 1466–1475.
385 doi:10.1111/een.12932
386
- 387 Tan J., Kerstetter J. E., and Turcotte M. M. (2021). Eco-evolutionary interaction between
388 microbiome presence and rapid biofilm evolution determines plant host fitness. *Nat Ecol Evol* 5,
389 670–676. doi:10.1038/s41559-021-01406-2
390
- 391 Tautz D. (1989). Hypervariability of simple sequences as a general source for polymorphic DNA
392 markers. *Nucleic Acids Res* 17, 6463–6471.
393
- 394 Tautz D., and Renz M. (1984). Simple sequences are ubiquitous repetitive components of
395 eukaryotic genomes. *Nucleic Acids Res* 12, 4127–4138.
396
- 397 Turcotte M. M., Reznick D. N., and Hare J. D. (2011). The impact of rapid evolution on
398 population dynamics in the wild: experimental test of eco-evolutionary dynamics. *Ecology*
399 *Letters* 14, 1084–1092. doi:10.1111/j.1461-0248.2011.01676.x
400
- 401 Van Steveninck R. F. M., Van Steveninck M. E., and Fernando D. R. (1992). Heavy-metal (Zn, Cd)
402 tolerance in selected clones of duck weed (*Lemna minor*). *Plant and Soil* 146, 271–280.
403
- 404 Vieira M. L. C., Santini L., Diniz A. L., and Munhoz C. de F. (2016). Microsatellite markers: what
405 they mean and why they are so useful. *Genet Mol Biol* 39, 312–328. doi:10.1590/1678-4685-
406 GMB-2016-0027
407
- 408 Wang W., Haberer G., Gundlach H., Gläßer C., Nussbaumer T., Luo M. C., Lomsadze A.,
409 Borodovsky M., Kerstetter R. A., Shanklin J., Byrant D. W., Mockler T. C., Appenroth K. J.,
410 Grimwood J., Jenkins J., Chow J., Choi C., Adam C., Cao X.-H., Fuchs J., Schubert I., Rokhsar D.,
411 Schmutz J., Michael T. P., Mayer K. F. X., and Messing J. (2014). The *Spirodela polyrhiza* genome
412 reveals insights into its neotenus reduction fast growth and aquatic lifestyle. *Nat Commun* 5,
413 3311. doi:10.1038/ncomms4311
414
- 415 Wani G., Shah M., Reshi Z., Atangana A., and Khasa D. (2014). CPDNa microsatellite markers for
416 *lemna minor* (ARACEAE): Phylogeographic Implications. *Applications in plant sciences* 2.,
417 doi:10.3732/apps.1300099
418
- 419 Xu N., Hu F., Wu J., Zhang W., Wang M., Zhu M., and Ke J. (2018). Characterization of 19
420 polymorphic SSR markers in *Spirodela polyrhiza* (Lemnaceae) and cross-amplification in *Lemna*
421 *perpusilla*. *Applications in Plant Sciences* 6,. doi:http://dx.doi.org/10.1002/aps3.1153
422
- 423 Xu Y., Ma S., Huang M., Peng M., Bog M., Sree K. S., Appenroth K.-J., and Zhang J. (2015).
424 Species distribution, genetic diversity and barcoding in the duckweed family (Lemnaceae).
425 *Hydrobiologia* 743, 75–87. doi:10.1007/s10750-014-2014-2

426

427 Xu S., Stapley J., Gablenz S., Boyer J., Appenroth K. J., Sree K. S., Gershenzon J., Widmer A., and
428 Huber M. (2019). Low genetic variation is associated with low mutation rate in the giant
429 duckweed. *Nat Commun* 10, 1243. doi:10.1038/s41467-019-09235-5

430

431 Ziegler P., Adelman K., Zimmer S., Schmidt C., and Appenroth K.-J. (2015). Relative in vitro
432 growth rates of duckweeds (Lemnaceae) - the most rapidly growing higher plants. *Plant Biol* 17,
433 33–41. doi:10.1111/plb.12184