

On two notions of total positivity for partial flag varieties



Anthony M. Bloch ^a, Steven N. Karp ^{b,*}

^a Department of Mathematics, University of Michigan, United States

^b Department of Mathematics, University of Notre Dame, United States

ARTICLE INFO

Article history:

Received 3 July 2022

Received in revised form 12 October 2022

Accepted 23 December 2022

Available online xxxx

Communicated by L.K. Williams

MSC:

15B48

14M15

52B40

81T60

Keywords:

Total positivity

Partial flag variety

Cyclic symmetry

Bruhat interval polytope

ABSTRACT

Given integers $1 \leq k_1 < \dots < k_l \leq n-1$, let $\text{Fl}_{k_1, \dots, k_l; n}$ denote the type A partial flag variety consisting of all chains of subspaces $(V_{k_1} \subset \dots \subset V_{k_l})$ inside \mathbb{R}^n , where each V_k has dimension k . Lusztig (1994, 1998) introduced the totally positive part $\text{Fl}_{k_1, \dots, k_l; n}^{>0}$ as the subset of partial flags which can be represented by a totally positive $n \times n$ matrix, and defined the totally nonnegative part $\text{Fl}_{k_1, \dots, k_l; n}^{\geq 0}$ as the closure of $\text{Fl}_{k_1, \dots, k_l; n}^{>0}$. On the other hand, following Postnikov (2007), we define $\text{Fl}_{k_1, \dots, k_l; n}^{\Delta > 0}$ and $\text{Fl}_{k_1, \dots, k_l; n}^{\Delta \geq 0}$ as the subsets of $\text{Fl}_{k_1, \dots, k_l; n}$ where all Plücker coordinates are positive and nonnegative, respectively. It follows from the definitions that Lusztig's total positivity implies Plücker positivity, and it is natural to ask when these two notions of positivity agree. Rietsch (2009) proved that they agree in the case of the Grassmannian $\text{Fl}_{k; n}$, and Chevalier (2011) showed that the two notions are distinct for $\text{Fl}_{1, 3; 4}$. We show that in general, the two notions agree if and only if k_1, \dots, k_l are consecutive integers. We give an elementary proof of this result (including for the case of Grassmannians) based on classical results in linear algebra and the theory of total positivity. We also show that the cell decomposition of $\text{Fl}_{k_1, \dots, k_l; n}^{\geq 0}$ coincides with its matroid decomposition if and only if k_1, \dots, k_l are consecutive integers, which was previously only known for complete flag varieties, Grassmannians, and $\text{Fl}_{1, 3; 4}$. Finally, we determine

* A.M.B. was partially supported by NSF grants DMS-1613819 and DMS-2103026, and AFOSR grant FA9550-22-1-0215. S.N.K. was partially supported by an NSERC postdoctoral fellowship.

* Corresponding author.

E-mail addresses: abloch@umich.edu (A.M. Bloch), skarp2@nd.edu (S.N. Karp).

which notions of positivity are compatible with a natural action of the cyclic group of order n that rotates the index set.

© 2023 Elsevier Inc. All rights reserved.

1. Introduction

A real matrix is called *totally positive* if all of its minors (i.e. determinants of square submatrices) are positive. Totally positive matrices first appeared in the 1930's in work of Schoenberg [52] and Gantmakher and Krein [23]. In the 1990's, Lusztig [37] generalized the theory of totally positive matrices to arbitrary semisimple algebraic groups G and their partial flag varieties G/P . These spaces have since been widely studied, with connections to representation theory [37], combinatorics [45], cluster algebras [16], high-energy physics [3,4], mirror symmetry [51], topology [19], and many other topics. The purpose of this paper is to examine the definition of the totally positive and totally nonnegative parts of a partial flag variety G/P in type A .

1.1. Lusztig's total positivity vs. Plücker positivity

Given a subset $K = \{k_1 < \dots < k_l\} \subseteq \{1, \dots, n-1\}$, let $\mathrm{Fl}_{K;n}(\mathbb{R})$ denote the partial flag variety of all tuples of subspaces $(V_{k_i})_{i=1}^l$ of \mathbb{R}^n , where $V_{k_1} \subset \dots \subset V_{k_l}$ and each V_{k_i} has dimension k_i . Equivalently, we may view $\mathrm{Fl}_{K;n}(\mathbb{R})$ as a parabolic quotient of $\mathrm{GL}_n(\mathbb{R})$, where a matrix $g \in \mathrm{GL}_n(\mathbb{R})$ represents the flag $(V_{k_i})_{i=1}^l$, where V_{k_i} is the span of the first k_i columns of g . Particular cases of interest are when $K = \{1, \dots, n-1\}$, in which case $\mathrm{Fl}_{K;n}(\mathbb{R})$ is the *complete flag variety* $\mathrm{Fl}_n(\mathbb{R})$, and when $K = \{k\}$, in which case $\mathrm{Fl}_{K;n}(\mathbb{R})$ is the *Grassmannian* $\mathrm{Gr}_{k,n}(\mathbb{R})$ of all k -dimensional subspaces of \mathbb{R}^n .

Lusztig [37,38] defined the *totally positive part* $\mathrm{Fl}_{K;n}^{>0}$ of $\mathrm{Fl}_{K;n}(\mathbb{R})$ to be the subset of flags which can be represented by a totally positive matrix in $\mathrm{GL}_n(\mathbb{R})$ (or equivalently, a totally positive lower-triangular unipotent matrix; see Remark 2.6 for further discussion). He also defined the *totally nonnegative part* $\mathrm{Fl}_{K;n}^{\geq 0}$ to be the closure of $\mathrm{Fl}_{K;n}^{>0}$. There is an alternative notion of positivity which arises from the *Plücker embedding* of $\mathrm{Fl}_{K;n}(\mathbb{R})$. Namely, we say that $(V_{k_i})_{i=1}^l$ is *Plücker positive* if all its Plücker coordinates are positive, or equivalently, if it can be represented by an element of $\mathrm{GL}_n(\mathbb{R})$ whose left-justified (i.e. initial) minors of orders k_1, \dots, k_l are all positive. We similarly say that $(V_{k_i})_{i=1}^l$ is *Plücker nonnegative* if all its Plücker coordinates are nonnegative. We denote the Plücker-positive and Plücker-nonnegative parts of $\mathrm{Fl}_{K;n}(\mathbb{R})$ by $\mathrm{Fl}_{K;n}^{\Delta >0}$ and $\mathrm{Fl}_{K;n}^{\Delta \geq 0}$, respectively. The Plücker-nonnegative part $\mathrm{Gr}_{k,n}^{\Delta \geq 0}$ of the Grassmannian was studied by Postnikov [45], and the space $\mathrm{Fl}_{K;n}^{\Delta \geq 0}$ was introduced by Arkani-Hamed, Bai, and Lam [3, Section 6.3], who called it the *naive nonnegative part*.

We wish to emphasize that both total positivity and Plücker positivity are natural notions. Lusztig's total positivity is compatible with his theory of canonical bases [36]

and with the combinatorics of Coxeter groups [37]. The space $\mathrm{Fl}_{K;n}^{\geq 0}$ can be decomposed into cells [37,48], each of which admits an explicit parametrization [41], and the cell decomposition forms a regular CW complex [19]. On the other hand, Plücker positivity is more concrete, and leads to connections with matroid theory [2] and tropical geometry [53]. It also arises in the definition of loop amplituhedra [5,6], which are spaces appearing in the physics of scattering amplitudes; for example, a *1-loop amplituhedron* is a certain projection of the space $\mathrm{Fl}_{k,k+2;n}^{\Delta \geq 0}$. The construction of loop amplituhedra is incompatible with Lusztig's total positivity, due to the presence of a cyclic symmetry among the particles in the scattering amplitude, as we explain in Section 1.3.

It follows from the definitions that $\mathrm{Fl}_{K;n}^{>0} \subseteq \mathrm{Fl}_{K;n}^{\Delta >0}$ and $\mathrm{Fl}_{K;n}^{\geq 0} \subseteq \mathrm{Fl}_{K;n}^{\Delta \geq 0}$, and it is natural to ask when equality holds. This question has been explored previously in special cases, as we discuss in Section 1.4. We answer this question in general:

Theorem 1.1. *Let $n \in \mathbb{N}$ and $K \subseteq \{1, \dots, n-1\}$. Then the following are equivalent:*

- (i) $\mathrm{Fl}_{K;n}^{>0} = \mathrm{Fl}_{K;n}^{\Delta >0}$;
- (ii) $\mathrm{Fl}_{K;n}^{\geq 0} = \mathrm{Fl}_{K;n}^{\Delta \geq 0}$; and
- (iii) the set K consists of consecutive integers.

We give an elementary proof of Theorem 1.1, using classical results in linear algebra and the theory of total positivity. In particular, our proof does not rely on any previously established special cases or on the cell decomposition of $\mathrm{Fl}_{K;n}^{\geq 0}$.

1.2. Cell decomposition vs. matroid decomposition

Lusztig [37, Remark 8.15] introduced a decomposition of $\mathrm{Fl}_{K;n}^{\geq 0}$, which Rietsch [48] showed is a cell decomposition (i.e. each stratum is homeomorphic to an open ball). It is the intersection of the *projected Richardson stratification* [32] with $\mathrm{Fl}_{K;n}^{\geq 0}$. There is another natural stratification of $\mathrm{Fl}_{K;n}(\mathbb{R})$, introduced by Gelfand and Serganova [25], which is the common refinement of the vanishing and nonvanishing sets of all Plücker coordinates. We call this the *matroid decomposition*, since each stratum is labeled by a tuple of matroids on the ground set $\{1, \dots, n\}$ (or alternatively, a Coxeter matroid).

Postnikov [45] studied the matroid decomposition of the Plücker-nonnegative part of $\mathrm{Gr}_{k,n}(\mathbb{R})$, which he called the *positroid decomposition*. He showed that it forms a cell decomposition [45, Theorem 3.5], and that it coincides with the decomposition of Lusztig and Rietsch [45, Theorem 3.8] (see Section 1.4 for further discussion). The topology of this decomposition presents a stark contrast to the matroid decomposition of all of $\mathrm{Gr}_{k,n}(\mathbb{R})$, which exhibits a phenomenon known as *Mnëv universality* [42]. In general, Tsukerman and Williams [55, Section 7] showed that the cell decomposition of $\mathrm{Fl}_{K;n}^{\geq 0}$ is a refinement of the matroid decomposition. They also showed that the two decompositions coincide for complete flag varieties, but differ for $\mathrm{Fl}_{1,3;4}^{\geq 0}$. We determine in general when the two decompositions are equal:

Theorem 1.2. *Let $n \in \mathbb{N}$ and $K \subseteq \{1, \dots, n-1\}$. Then the cell decomposition of $\text{Fl}_{K;n}^{\geq 0}$ coincides with the matroid decomposition if and only if K consists of consecutive integers.*

The forward direction of Theorem 1.2 follows from Theorem 1.1, using a topological argument. However, we do not know how to deduce the reverse direction directly from Theorem 1.1. Instead, we use a result of Tsukerman and Williams [55, Theorem 7.1], which in turn builds on unpublished work of Marsh and Rietsch. It states that each cell of $\text{Fl}_{K;n}^{\geq 0}$ is contained in a single matroid stratum, which is uniquely determined by the 0-dimensional cells in its closure. These 0-dimensional cells have an explicit description in terms of the Bruhat order on the symmetric group \mathfrak{S}_n , due to Rietsch [49]. To prove the reverse direction of Theorem 1.2, we use the combinatorics of \mathfrak{S}_n to reduce the statement to the Grassmannian case, which was proved by Postnikov as described above. Our techniques also have implications in the study of the *Bruhat interval polytopes* of Kodama and Williams [34] and Tsukerman and Williams [55]; see Remark 5.17 and Remark 5.24.

1.3. Cyclic symmetry

For $\epsilon \in \mathbb{Z}/2\mathbb{Z}$, define the (*signed*) left cyclic shift map $\sigma_\epsilon \in \text{GL}_n(\mathbb{R})$ by

$$\sigma_\epsilon(v_1, \dots, v_n) := (v_2, \dots, v_n, (-1)^{\epsilon-1}v_1) \quad \text{for all } v \in \mathbb{R}^n.$$

Then σ_ϵ also acts on $\text{Fl}_{K;n}(\mathbb{R})$. If all elements of K have the same parity ϵ , then by the alternating property of the determinant, σ_ϵ acts on Plücker coordinates by rotating the index set $\{1, \dots, n\}$. Therefore σ_ϵ preserves both $\text{Fl}_{K;n}^{\Delta > 0}$ and $\text{Fl}_{K;n}^{\Delta \geq 0}$. It is natural to wonder whether σ_ϵ also preserves $\text{Fl}_{K;n}^{> 0}$ and $\text{Fl}_{K;n}^{\geq 0}$. One motivation is that the cyclic symmetry for $\text{Fl}_{k,k+2;n}^{\Delta \geq 0}$ is important in the definition of loop amplituhedra mentioned above, coming from a cyclic ordering on the n external particles.

If $K = \{k\}$ and k has parity ϵ , then σ_ϵ preserves both $\text{Gr}_{k,n}^{> 0}$ and $\text{Gr}_{k,n}^{\geq 0}$, using Theorem 1.1 for $\text{Gr}_{k,n}(\mathbb{R})$. We show, however, that in all other cases, σ_ϵ does not preserve either $\text{Fl}_{K;n}^{> 0}$ or $\text{Fl}_{K;n}^{\geq 0}$. In particular, one cannot substitute the notion of total positivity for Plücker positivity in the definition of loop amplituhedra.

Theorem 1.3. *Let $K \subseteq \{1, \dots, n-1\}$ such that $|K| \geq 2$, and let $\epsilon \in \mathbb{Z}/2\mathbb{Z}$. Then there exists $V \in \text{Fl}_{K;n}^{\geq 0}$ such that $\sigma_\epsilon(V) \notin \text{Fl}_{K;n}^{\geq 0}$. In particular, σ_ϵ does not preserve $\text{Fl}_{K;n}^{> 0}$ or $\text{Fl}_{K;n}^{\geq 0}$.*

While Theorem 1.3 does not follow directly from Theorem 1.1, we use similar techniques to prove both results.

One of our initial motivations for this work was to understand which gradient flows on $\text{Fl}_{K;n}(\mathbb{R})$ are compatible with total positivity, which we discuss in a separate paper [8]. We discovered that in certain cases, the classification of such flows differs depending

on whether one works with Lusztig's total positivity or with Plücker positivity; see Theorem 5.14, Theorem 5.18, and Remark 5.20 of [8]. Theorem 1.3 above may be regarded as an infinitesimal analogue of this phenomenon.

1.4. History

We discuss previous work related to Theorem 1.1 and Theorem 1.2. Much of this work has focused on the case when $\mathrm{Fl}_{K;n}(\mathbb{R})$ is a Grassmannian $\mathrm{Gr}_{k,n}(\mathbb{R})$, due to Postnikov's study of the matroid decomposition of $\mathrm{Gr}_{k,n}^{\Delta \geq 0}$ [45].

Rietsch [47, Section 4.5] (also announced in [38, Section 3.12]) stated that Lusztig's notion of total positivity coincides with Plücker positivity for all partial flag varieties G/P , but her proof contained an error. She later proved Theorem 1.1 for Grassmannians $\mathrm{Gr}_{k,n}(\mathbb{R})$ in unpublished notes [50]. Subsequent proofs were given by Talaska and Williams [54, Corollary 1.2], Lam [35, Remark 3.8], and Lusztig [39]. Theorem 1.2 for the complete flag variety $\mathrm{Fl}_n(\mathbb{R})$ was stated in [19, (9.15)], and proofs were given by Lusztig [40, p. 4] and Boretsky [9, Theorem 5.25]. Conversely, Chevalier [12, Example 10.1] gave an example showing that $\mathrm{Fl}_{1,3;4}^{>0} \neq \mathrm{Fl}_{1,3;4}^{\Delta >0}$; see [33, Remark 5.17] for a related discussion. In a different direction, Geiss, Leclerc, and Schröer [24, Conjecture 19.2] have conjectured an algebraic description of the totally positive part of any partial flag variety G/P .

Theorem 1.2 was proved in the case of Grassmannians $\mathrm{Gr}_{k,n}^{\geq 0}$ by Postnikov [45, Corollary 3.8]. We point out that his proof implicitly uses the fact that $\mathrm{Gr}_{k,n}^{\geq 0} = \mathrm{Gr}_{k,n}^{\Delta \geq 0}$, which was only later proved by Rietsch [50], as described above. A subsequent proof was given by Talaska and Williams [54, Corollary 1.2], and the result can also be proved using work of Tsukerman and Williams [55, Section 7]. Theorem 1.2 in the cases of $\mathrm{Fl}_n^{\geq 0}$ and $\mathrm{Fl}_{1,3;4}^{\geq 0}$ was proved by Tsukerman and Williams [55, Theorem 7.1 and Remark 7.3].

1.5. Further directions

Lusztig [37, 38] introduced the totally positive and totally nonnegative part of an arbitrary partial flag variety G/P , where G is a semisimple algebraic group G over \mathbb{R} and P is a parabolic subgroup. It would be interesting to study the problems considered in this paper for such G/P . We mention that Fomin and Zelevinsky [18] have considered analogous problems for the group G . In particular, they showed that an element of G is totally positive (respectively, totally nonnegative) in the sense of Lusztig [37] if and only if all its generalized minors are positive (respectively, nonnegative). We consider only the type A case here both for the sake of concreteness, and because our proofs of Theorem 1.1 and Theorem 1.3 are elementary. We point out that many of our arguments in Section 5 used to prove Theorem 1.2 can be applied to general Coxeter groups; however, a key tool in our proof is Postnikov's Theorem 5.16, which is only known in type A .

Lusztig [38, Theorem 3.4] showed that for any partial flag variety G/P , there exists a positive weight λ such that a partial flag is totally positive (respectively, totally nonnegative) if and only if its coordinates with respect to the canonical basis of the irre-

ducible G -module with highest weight λ are all positive (respectively, nonnegative). He also posed the problem of finding the minimal such λ . Our Theorem 1.1 gives a partial answer for type A partial flag varieties $G/P = \mathrm{Fl}_{K;n}(\mathbb{R})$: it implies that the sum of fundamental weights $\sum_{k \in K} \omega_k$ is a valid choice of λ (and hence minimal) if and only if K consists of consecutive integers.

As we described above, the combinatorics and topology of the cell decomposition of $\mathrm{Fl}_{K;n}^{\geq 0}$ have been extensively studied. For example, each cell has an explicit parametrization [41, Section 11], the closure poset is shellable [57, Theorem 1.2], and the cell decomposition forms a regular CW complex [19, Theorem 1.1]. In light of Theorem 1.1 and Theorem 1.2, it would be interesting to further study the matroid decomposition of $\mathrm{Fl}_{K;n}^{\Delta \geq 0}$. Bai, He, and Lam [6] studied the case $\mathrm{Fl}_{1,3;n}^{\Delta \geq 0}$. In [8, Corollary 6.16], we show that $\mathrm{Fl}_{K;n}^{\Delta > 0}$ is homeomorphic to an open ball and $\mathrm{Fl}_{K;n}^{\Delta \geq 0}$ is homeomorphic to a closed ball.

1.6. Outline

In Section 2, we give some background on partial flag varieties and total positivity. In Section 3, we prove Theorem 1.1. In Section 4, we prove Theorem 1.3. In Section 5, we prove Theorem 1.2.

Acknowledgments

We thank George Lusztig, Lauren Williams, and an anonymous reviewer for helpful comments.

2. Background

In this section, we define partial flag varieties and their totally positive and totally nonnegative parts, and recall some classical results in the theory of total positivity. For further details on total positivity, we refer to [14, 17, 22, 30, 37, 43].

2.1. Notation

Let $\mathbb{N} := \{0, 1, 2, \dots\}$. For $n \in \mathbb{N}$, we let $[n]$ denote $\{1, 2, \dots, n\}$, and for $i, j \in \mathbb{Z}$, we let $[i, j]$ denote the interval of integers $\{i, i+1, \dots, j\}$. Given a set S and $k \in \mathbb{N}$, we let $\binom{S}{k}$ denote the set of k -element subsets of S .

We let e_1, \dots, e_n denote the unit vectors of \mathbb{R}^n . We let $\mathbb{P}^n(\mathbb{R})$ denote n -dimensional real projective space, defined to be $\mathbb{R}^{n+1} \setminus \{0\}$ modulo multiplication by \mathbb{R}^\times . For $\lambda_1, \dots, \lambda_n \in \mathbb{R}$, we let $\mathrm{Diag}(\lambda_1, \dots, \lambda_n)$ denote the $n \times n$ diagonal matrix with diagonal entries $\lambda_1, \dots, \lambda_n$. We let $\mathrm{GL}_n(\mathbb{R})$ denote the group of invertible real $n \times n$ matrices.

Given an $m \times n$ matrix A , we let A^\top denote the transpose of A . For $1 \leq k \leq m, n$ and subsets $I \in \binom{[m]}{k}$ and $J \in \binom{[n]}{k}$, we let $\Delta_{I,J}(A)$ denote the determinant of the submatrix

of A in rows I and columns J , called a *minor* of A of *order* k . If $J = [k]$, we call $\Delta_{I,J}(A)$ a *left-justified minor* of A . We also let $\sum I$ denote the sum of the elements in I .

We will make repeated use of the *Cauchy–Binet identity* (see e.g. [20, I.(14)]): if A is an $m \times n$ matrix, B is an $n \times p$ matrix, and $1 \leq k \leq m, p$, then

$$\Delta_{I,J}(AB) = \sum_{K \in \binom{[n]}{k}} \Delta_{I,K}(A) \Delta_{K,J}(B) \quad \text{for all } I \in \binom{[m]}{k} \text{ and } J \in \binom{[p]}{k}. \quad (2.1)$$

We now introduce partial flag varieties.

Definition 2.1. Let $n \in \mathbb{N}$, and let $K = \{k_1 < \dots < k_l\} \subseteq [n-1]$. Let $\mathrm{P}_{K;n}(\mathbb{R})$ denote the parabolic subgroup of $\mathrm{GL}_n(\mathbb{R})$ of block upper-triangular matrices with diagonal blocks of sizes $k_1, k_2 - k_1, \dots, k_l - k_{l-1}, n - k_l$. We define the *partial flag variety*

$$\mathrm{Fl}_{K;n}(\mathbb{R}) := \mathrm{GL}_n(\mathbb{R}) / \mathrm{P}_{K;n}(\mathbb{R}).$$

We identify $\mathrm{Fl}_{K;n}(\mathbb{R})$ with the variety of partial flags of subspaces in \mathbb{R}^n

$$\{V = (V_{k_1}, \dots, V_{k_l}) : 0 \subset V_{k_1} \subset \dots \subset V_{k_l} \subset \mathbb{R}^n \text{ and } \dim(V_{k_i}) = k_i \text{ for } 1 \leq i \leq l\}.$$

The identification sends $g \in \mathrm{GL}_n(\mathbb{R}) / \mathrm{P}_{K;n}(\mathbb{R})$ to the tuple $(V_k)_{k \in K}$, where V_k is the span of the first k columns of g for all $k \in K$. More generally, if A is any real matrix with n rows and at least k_l columns such that V_k is the span of the first k columns of A for all $k \in K$, we say that A *represents* V . Note that $\mathrm{GL}_n(\mathbb{R})$ acts on $\mathrm{Fl}_{K;n}(\mathbb{R})$ on the left.

We have the *Plücker embedding*

$$\begin{aligned} \mathrm{Fl}_{K;n}(\mathbb{R}) &\hookrightarrow \mathbb{P}^{\binom{[n]}{k_1}-1} \times \dots \times \mathbb{P}^{\binom{[n]}{k_l}-1}, \\ V &\mapsto \left((\Delta_I(A))_{I \in \binom{[n]}{k_1}}, \dots, (\Delta_I(A))_{I \in \binom{[n]}{k_l}} \right), \end{aligned} \quad (2.2)$$

where A denotes any matrix representative of V . (We can check that the definition does not depend on the choice of A .) We call the left-justified minors $\Delta_I(A)$ appearing above the *Plücker coordinates* of $V \in \mathrm{Fl}_{K;n}(\mathbb{R})$ (also known as *flag minors*), which we denote by $\Delta_I(V)$. We point out that our Plücker coordinates differ from the *generalized Plücker coordinates* of Gelfand and Serganova [25], though each encodes essentially the same data; see Remark 5.13.

For any $K' \subseteq K$, we have a projection map

$$\mathrm{Fl}_{K;n}(\mathbb{R}) \twoheadrightarrow \mathrm{Fl}_{K';n}(\mathbb{R}), \quad (V_k)_{k \in K} \mapsto (V_k)_{k \in K'}. \quad (2.3)$$

The map (2.3) retains only the subspaces of a partial flag whose dimensions lie in K' .

We mention two instances of $\mathrm{Fl}_{K;n}(\mathbb{R})$ which are of particular interest. If $K = [n-1]$, then $\mathrm{Fl}_{K;n}(\mathbb{R})$ is the *complete flag variety* of \mathbb{R}^n , which we denote by $\mathrm{Fl}_n(\mathbb{R})$. If K is the singleton $\{k\}$, then $\mathrm{Fl}_{K;n}(\mathbb{R})$ is the *Grassmannian* of all k -dimensional subspaces of \mathbb{R}^n , which we denote by $\mathrm{Gr}_{k,n}(\mathbb{R})$. We also extend the definition of $\mathrm{Gr}_{k,n}(\mathbb{R})$ to $k = 0$ and $k = n$.

Example 2.2. Let $n := 4$ and $K := \{1, 3\}$. Then

$$\mathrm{P}_{1,3;4}(\mathbb{R}) = \left\{ \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \right\} \subseteq \mathrm{GL}_4(\mathbb{R}) \quad \text{and} \quad \mathrm{Fl}_{1,3;4}(\mathbb{R}) = \mathrm{GL}_4(\mathbb{R}) / \mathrm{P}_{1,3;4}(\mathbb{R}).$$

We identify $\mathrm{Fl}_{1,3;4}(\mathbb{R})$ with the variety of partial flags $V = (V_1, V_3)$ of subspaces of \mathbb{R}^4 , where $\dim(V_1) = 1$, $\dim(V_3) = 3$, and $V_1 \subset V_3$.

We can represent a generic partial flag $V \in \mathrm{Fl}_{1,3;4}(\mathbb{R})$ by a matrix of the form

$$A = \begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ b & 0 & 1 \\ c & d & e \end{bmatrix}, \quad \text{where } a, b, c, d, e \in \mathbb{R}.$$

That is, V_1 is spanned by the first column of A , and V_3 is spanned by all three columns of A . Then the Plücker embedding (2.2) takes V to

$$\begin{aligned} & ((\Delta_1(V) : \Delta_2(V) : \Delta_3(V) : \Delta_4(V)), (\Delta_{123}(V) : \Delta_{124}(V) : \Delta_{134}(V) : \Delta_{234}(V))) \\ & = ((1 : a : b : c), (1 : e : -d : -ad + c - be)) \in \mathbb{P}^3(\mathbb{R}) \times \mathbb{P}^3(\mathbb{R}). \quad \diamond \end{aligned}$$

Definition 2.3. We say that a real matrix is *totally positive* if all its minors are positive. For $n \in \mathbb{N}$, we let $\mathrm{GL}_n^{>0}$ denote the subset of $\mathrm{GL}_n(\mathbb{R})$ of totally positive matrices.

For example, we have $\mathrm{GL}_2^{>0} = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d, ad - bc > 0 \right\}$. We now introduce Lusztig's total positivity and Plücker positivity for partial flag varieties.

Definition 2.4. Let $n \in \mathbb{N}$ and $K \subseteq [n-1]$. Following [37, Section 8] and [38, Section 1.5], we define the *totally positive part* of $\mathrm{Fl}_{K;n}(\mathbb{R})$, denoted by $\mathrm{Fl}_{K;n}^{>0}$, as the image of $\mathrm{GL}_n^{>0}$ inside $\mathrm{Fl}_{K;n}(\mathbb{R}) = \mathrm{GL}_n(\mathbb{R}) / \mathrm{P}_{K;n}(\mathbb{R})$. Equivalently, $\mathrm{Fl}_{K;n}^{>0}$ consists of all partial flags which can be represented by a totally positive $n \times n$ matrix. We define the *totally nonnegative part* of $\mathrm{Fl}_{K;n}(\mathbb{R})$, denoted by $\mathrm{Fl}_{K;n}^{\geq 0}$, as the closure of $\mathrm{Fl}_{K;n}^{>0}$ in the Euclidean topology. Note that for any $K' \subseteq K$, the projection map (2.3) restricts to surjections

$$\mathrm{Fl}_{K;n}^{>0} \twoheadrightarrow \mathrm{Fl}_{K';n}^{>0} \quad \text{and} \quad \mathrm{Fl}_{K;n}^{\geq 0} \twoheadrightarrow \mathrm{Fl}_{K';n}^{\geq 0}. \quad (2.4)$$

We also define the *Plücker-positive part* of $\mathrm{Fl}_{K;n}(\mathbb{R})$, denoted by $\mathrm{Fl}_{K;n}^{\Delta > 0}$, as the subset of partial flags whose Plücker coordinates are all positive (up to rescaling). That is,

$\text{Fl}_{K;n}^{\Delta > 0}$ consists of all partial flags which can be represented by a matrix A such that all left-justified $k \times k$ minors of A are positive for all $k \in K$. We similarly define the *Plücker-nonnegative part* $\text{Fl}_{K;n}^{\Delta \geq 0}$ by replacing “positive” with “nonnegative” above.

Note that by definition, we have $\text{Fl}_{K;n}^{>0} \subseteq \text{Fl}_{K;n}^{\Delta > 0}$ and $\text{Fl}_{K;n}^{\geq 0} \subseteq \text{Fl}_{K;n}^{\Delta \geq 0}$. We also have that $\text{Fl}_{K;n}^{\Delta \geq 0}$ is the closure of $\text{Fl}_{K;n}^{\Delta > 0}$; see Proposition 3.3(ii).

Example 2.5. We have

$$\text{Fl}_3^{\Delta > 0} = \left\{ \begin{bmatrix} 1 & 0 & 0 \\ a+c & 1 & 0 \\ bc & b & 1 \end{bmatrix} : a, b, c > 0 \right\} \quad \text{and} \quad \text{Gr}_{2,4}^{\Delta > 0} = \left\{ \begin{bmatrix} 1 & 0 \\ a & b \\ 0 & 1 \\ -c & d \end{bmatrix} : a, b, c, d > 0 \right\}. \quad \diamond$$

Remark 2.6. Lusztig’s original definition of $\text{Fl}_{K;n}^{>0}$ is slightly different than the one we give in Definition 2.4, but is equivalent. Namely, let $\text{N}_n(\mathbb{R})$ be the subset of $\text{GL}_n(\mathbb{R})$ of all upper-triangular matrices with 1’s on the diagonal. We define $\text{N}_n^{>0}$ to be the subset of $\text{N}_n(\mathbb{R})$ of matrices whose minors are all positive, except for those which are zero due to upper triangularity. Let $(\text{N}_n^-)^{>0}$ denote the transpose of $\text{N}_n^{>0}$, and let $\text{T}_n^{>0}$ denote the subset of $\text{GL}_n(\mathbb{R})$ of diagonal matrices with positive diagonal entries. Then Lusztig [37, Section 8] defines $\text{Fl}_{K;n}^{>0}$ to be the image of $(\text{N}_n^-)^{>0}$ inside $\text{Fl}_{K;n}(\mathbb{R})$. This is equal to the image of $\text{GL}_n^{>0}$, because

$$\text{GL}_n^{>0} = (\text{N}_n^-)^{>0} \cdot \text{T}_n^{>0} \cdot \text{N}_n^{>0}, \quad (2.5)$$

and $\text{T}_n^{>0} \cdot \text{N}_n^{>0} \subseteq \text{P}_{K;n}(\mathbb{R})$. (In fact, Lusztig takes (2.5) to hold by definition; see [37, Section 2.12].) The decomposition (2.5) is a result of Cryer [13, Theorem 1.1]; we refer to [14, Chapter 2] for further discussion and references.

Remark 2.7. A real matrix is called *totally nonnegative* if all its minors are nonnegative. Every totally nonnegative matrix in $\text{GL}_n(\mathbb{R})$ represents a totally nonnegative flag in $\text{Fl}_{K;n}(\mathbb{R})$. However, not every element of $\text{Fl}_{K;n}^{\geq 0}$ is represented by a totally nonnegative matrix, unless $K = \emptyset$. For example, the element $\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \in \text{Fl}_2^{\geq 0}$ cannot be represented by a totally nonnegative matrix, since the top-left entry of a totally nonnegative matrix is positive.

We now recall two classical results from the theory of total positivity.

Lemma 2.8. *There exists a continuous function $f : \mathbb{R}_{\geq 0} \rightarrow \text{GL}_n(\mathbb{R})$ such that $f(0) = I_n$ and $f(t) \in \text{GL}_n^{>0}$ for all $t > 0$.*

Proof. By [22, II.3.(24)] or [56, (2)] (cf. [44, Problem V.76]), we may take $f(t) := (t^{(i-j)^2})_{1 \leq i,j \leq n}$. Alternatively, by [30, Theorem 3.3.4], we may take $f(t) := \exp(tA)$,

where A is any tridiagonal $n \times n$ matrix whose entries immediately above and below the diagonal are positive. \square

Theorem 2.9 (Fekete [15]; see [43, Lemma 2.1]). *Let A be an $n \times (k+1)$ matrix, where $n \geq k+1$. Suppose that all left-justified $k \times k$ minors of A are positive, and that all $(k+1) \times (k+1)$ minors of A using consecutive rows are positive. Then all $(k+1) \times (k+1)$ minors of A are positive.*

The following two results are duality and restriction statements for totally nonnegative Grassmannians. See [1, Section 3] for closely related results. We note that Lemma 2.10 follows from Jacobi's formula for the matrix inverse; we refer to [31] for further discussion and references.

Lemma 2.10 ([31, Lemma 1.11(ii)]). *Define the bilinear pairing $\langle \cdot, \cdot \rangle$ on \mathbb{R}^n by*

$$\langle v, w \rangle := v_1 w_1 - v_2 w_2 + v_3 w_3 - \cdots + (-1)^{n-1} v_n w_n.$$

Given $V \in \text{Gr}_{k,n}(\mathbb{R})$, let $V^\perp := \{w \in \mathbb{R}^n : \langle v, w \rangle = 0 \text{ for all } v \in V\} \in \text{Gr}_{n-k,n}(\mathbb{R})$. Then

$$\Delta_I(V) = \Delta_{[n] \setminus I}(V^\perp) \quad \text{for all } I \in \binom{[n]}{k}.$$

In particular, \cdot^\perp defines bijections $\text{Gr}_{k,n}^{\Delta > 0} \leftrightarrow \text{Gr}_{n-k,n}^{\Delta > 0}$ and $\text{Gr}_{k,n}^{\Delta \geq 0} \leftrightarrow \text{Gr}_{n-k,n}^{\Delta \geq 0}$.

Example 2.11. Let $V \in \text{Gr}_{2,4}(\mathbb{R})$ be represented by the matrix

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \\ a & b \\ c & d \end{bmatrix}.$$

Then $V^\perp \in \text{Gr}_{2,4}(\mathbb{R})$ is represented by the matrix

$$\begin{bmatrix} -a & c \\ b & -d \\ 1 & 0 \\ 0 & 1 \end{bmatrix}. \quad \diamond$$

Lemma 2.12. *Let $V \in \text{Gr}_{k,n}^{\Delta \geq 0}$, and let $m \leq n$. Define $W := V \cap \text{span}(e_1, \dots, e_m)$, and let $d := \dim(W)$. Then $W \in \text{Gr}_{d,m}^{\Delta \geq 0}$.*

Proof. Take an $m \times d$ matrix B representing W . Then there exists an $n \times k$ matrix A representing V of the form

$$A = \begin{bmatrix} B & * \\ 0 & C \end{bmatrix},$$

where C is an $(n - m) \times (k - d)$ matrix of rank $k - d$. We may assume that C is in reduced column echelon form, so that C restricted to some subset $J \in \binom{[n-m]}{k-d}$ of rows equals I_{k-d} . Let $J' := \{j + m : j \in J\}$. Then for all $I \in \binom{[m]}{d}$, we have

$$\Delta_I(B) = \Delta_{I \cup J'}(A), \quad \text{so} \quad \Delta_I(W) = \Delta_{I \cup J'}(V).$$

Since $V \in \text{Gr}_{k,n}^{\Delta \geq 0}$, we have $W \in \text{Gr}_{d,m}^{\Delta \geq 0}$. \square

3. Lusztig's total positivity vs. Plücker positivity

In this section we prove Theorem 1.1. Our argument will be based on the preliminary results Proposition 3.3, Lemma 3.4, and Lemma 3.5.

Lemma 3.1. *For $n \in \mathbb{N}$, we have $\text{Fl}_n^{>0} = \text{Fl}_n^{\Delta > 0}$.*

Proof. We know that $\text{Fl}_n^{>0} \subseteq \text{Fl}_n^{\Delta > 0}$. Conversely, let $V \in \text{Fl}_n^{\Delta > 0}$. Then there exists an $n \times n$ matrix A representing V whose left-justified minors are all positive. After performing left-to-right column operations, we may assume that A is lower-triangular. For $t > 0$, define

$$g := A \text{Diag}(t^{n-1}, \dots, t, 1) A^T \in \text{GL}_n(\mathbb{R}),$$

which also represents V . We claim that $g \in \text{GL}_n^{>0}$ for all t sufficiently large, whence $V \in \text{Fl}_n^{>0}$, completing the proof. (In fact, $g \in \text{GL}_n^{>0}$ for all $t > 0$, by (2.5).) To see this, note that for $1 \leq k \leq n$ and $I, J \in \binom{[n]}{k}$, by (2.1) we have that $\Delta_{I,J}(g)$ equals

$$\sum_{K \in \binom{[n]}{k}} \Delta_{I,K}(A) \Delta_{J,K}(A) t^{kn - \sum K} = \Delta_{I,[k]}(A) \Delta_{J,[k]}(A) t^{kn - \binom{k+1}{2}} + \text{lower order terms}$$

as $t \rightarrow \infty$. \square

Lemma 3.2. *Let $K \subseteq [n - 1]$, and let $g \in \text{GL}_n^{>0}$.*

- (i) *For all $V \in \text{Fl}_{K;n}^{\geq 0}$, we have $g \cdot V \in \text{Fl}_{K;n}^{>0}$.*
- (ii) *For all $V \in \text{Fl}_{K;n}^{\Delta \geq 0}$, we have $g \cdot V \in \text{Fl}_{K;n}^{\Delta > 0}$.*

Proof. Part (ii) follows from the Cauchy–Binet identity (2.1). For part (i), by (2.4), it suffices to prove the result for the complete flag variety (when $K = [n - 1]$). Since $\text{Fl}_n^{\geq 0} \subseteq \text{Fl}_n^{\Delta \geq 0}$, this case follows from part (ii) and Lemma 3.1. \square

Proposition 3.3. *Let $K \subseteq [n - 1]$.*

- (i) *$\text{Fl}_{K;n}^{\geq 0}$ is the closure of $\text{Fl}_{K;n}^{>0}$, and $\text{Fl}_{K;n}^{>0}$ is the interior of $\text{Fl}_{K;n}^{\geq 0}$.*
- (ii) *$\text{Fl}_{K;n}^{\Delta \geq 0}$ is the closure of $\text{Fl}_{K;n}^{\Delta > 0}$, and $\text{Fl}_{K;n}^{\Delta > 0}$ is the interior of $\text{Fl}_{K;n}^{\Delta \geq 0}$.*

Proof. Let $f : \mathbb{R}_{\geq 0} \rightarrow \mathrm{GL}_n(\mathbb{R})$ be as in Lemma 2.8.

(i) By definition, $\mathrm{Fl}_{K;n}^{\geq 0}$ is the closure of $\mathrm{Fl}_{K;n}^{>0}$. Also, $\mathrm{Fl}_{K;n}^{>0}$ is open, so it is contained in the interior of $\mathrm{Fl}_{K;n}^{\geq 0}$. It remains to show that the interior of $\mathrm{Fl}_{K;n}^{\geq 0}$ is contained in $\mathrm{Fl}_{K;n}^{>0}$. To see this, let $V \in \mathrm{Fl}_{K;n}^{\geq 0} \setminus \mathrm{Fl}_{K;n}^{>0}$. We claim that $f(t)^{-1} \cdot V$ is not in $\mathrm{Fl}_{K;n}^{\geq 0}$ for all $t > 0$, whence V is not in the interior of $\mathrm{Fl}_{K;n}^{\geq 0}$. Indeed, if $f(t)^{-1} \cdot V \in \mathrm{Fl}_{K;n}^{\geq 0}$ with $t > 0$, then by Lemma 3.2(i) we obtain

$$V = f(t) \cdot (f(t)^{-1} \cdot V) \in \mathrm{Fl}_{K;n}^{>0},$$

a contradiction.

(ii) Note that the closure of $\mathrm{Fl}_{K;n}^{\Delta > 0}$ is contained in $\mathrm{Fl}_{K;n}^{\Delta \geq 0}$. Conversely, given $V \in \mathrm{Fl}_{K;n}^{\Delta \geq 0}$, we have $V = \lim_{t \rightarrow 0, t > 0} f(t) \cdot V$, and $f(t) \cdot V \in \mathrm{Fl}_{K;n}^{\Delta > 0}$ for $t > 0$ by Lemma 3.2(ii). Therefore $\mathrm{Fl}_{K;n}^{\Delta \geq 0}$ is the closure of $\mathrm{Fl}_{K;n}^{\Delta > 0}$. The fact that $\mathrm{Fl}_{K;n}^{\Delta > 0}$ is the interior of $\mathrm{Fl}_{K;n}^{\Delta \geq 0}$ follows from a similar argument as in the proof of part (i). \square

Lemma 3.4. Let $V \in \mathrm{Gr}_{k,n}^{\Delta > 0}$, where $1 \leq k \leq n-1$.

- (i) There exists $W \in \mathrm{Gr}_{k+1,n}^{\Delta > 0}$ such that $V \subseteq W$.
- (ii) There exists $W \in \mathrm{Gr}_{k-1,n}^{\Delta > 0}$ such that $W \subseteq V$.

Proof. (i) Take an $n \times k$ matrix A representing V whose $k \times k$ minors are all positive. Let $B := [A \mid w]$ denote the $n \times (k+1)$ matrix formed by concatenating A and the vector $w \in \mathbb{R}^n$, where we define w as follows. We set $w_1, \dots, w_k := 0$, and for $i = k+1, \dots, n$, we take $w_i > 0$ to be sufficiently large that the minor $\Delta_{[i-k,i],[k+1]}(B)$ is positive. By Theorem 2.9, all $(k+1) \times (k+1)$ minors of B are positive. Therefore we may define W to be the column span of B .

(ii) This follows by applying part (i) to V^\perp , using Lemma 2.10. \square

Lemma 3.5. Let $V \in \mathrm{Gr}_{k,n}^{\Delta \geq 0}$ and $W \in \mathrm{Gr}_{k+1,n}^{\Delta \geq 0}$ such that $V \subseteq W$. If $e_1 + ce_n \in V$ for some $c \in \mathbb{R}$, then $e_1 \in W$.

Proof. If $e_1 \in V$, then $e_1 \in W$. Now suppose that $e_1 \notin V$ and $e_1 + ce_n \in V$ for some $c \in \mathbb{R}$, so that $c \neq 0$ and $e_n \notin V$. Let A denote the $n \times k$ matrix representing V in reduced column echelon form. Let $I \in \binom{[n]}{k}$ index the rows containing the pivot 1's of A ; equivalently, I is lexicographically minimal such that $\Delta_I(V) \neq 0$. Since $e_1 + ce_n \in V$ and $e_n \notin V$, we have $1 \in I$ and $n \notin I$, and the first column of A is $e_1 + ce_n$. Therefore $\Delta_I(A) = 1$ and $\Delta_{(I \setminus \{1\}) \cup \{n\}}(A) = (-1)^{k-1}c$. Since V is Plücker nonnegative, we get that $(-1)^{k-1}c > 0$.

Now take $w \in W \setminus V$ such that $w_i = 0$ for all $i \in I$. Let $B := [A \mid w]$ denote the $n \times (k+1)$ matrix representing W , formed by concatenating A and w . Then for $i \in [n-1] \setminus I$, there exists $\epsilon \in \{1, -1\}$ such that $\Delta_{I \cup \{i\}}(B) = \epsilon w_i$ and $\Delta_{(I \setminus \{1\}) \cup \{i,n\}}(B) =$

$\epsilon(-1)^k cw_i$. Since W is Plücker nonnegative and $(-1)^{k-1}c > 0$, we get that $w_i = 0$. Therefore w is a nonzero scalar multiple of e_n . Since $e_1 + ce_n \in W$, we obtain $e_1 \in W$. \square

Proof of Theorem 1.1. (i) \Leftrightarrow (ii): This follows from Proposition 3.3.

(iii) \Rightarrow (i): Suppose that $K = [k, l]$. Recall that $\text{Fl}_{K;n}^{>0} \subseteq \text{Fl}_{K;n}^{\Delta > 0}$. Conversely, we must show that given $V = (V_1, \dots, V_l) \in \text{Fl}_{K;n}^{\Delta > 0}$, we have $V \in \text{Fl}_{K;n}^{>0}$. By repeatedly applying Lemma 3.4, there exist $V_i \in \text{Gr}_{i,n}^{\Delta > 0}$ for $i = l+1, \dots, n$ and $i = k-1, \dots, 1$ such that $V_1 \subset \dots \subset V_{n-1}$. Let $W := (V_1, \dots, V_{n-1}) \in \text{Fl}_n(\mathbb{R})$. Then $W \in \text{Fl}_n^{\Delta > 0}$, so $W \in \text{Fl}_n^{>0}$ by Lemma 3.1. Then by (2.4), we get $V \in \text{Fl}_{K;n}^{>0}$.

(ii) \Rightarrow (iii): We prove the contrapositive. Suppose that K does not consist of consecutive integers, so that there exist consecutive elements $k < l$ of K with $l - k \geq 2$. Define the element $V = (V_i)_{i \in K}$ of $\text{Fl}_{K;n}(\mathbb{R})$ as follows:

$$V_i := \begin{cases} \text{span}(e_5, e_6, \dots, e_{i+4}), & \text{if } i < k; \\ \text{span}(e_1 + e_4, e_5, e_6, \dots, e_{k+3}), & \text{if } i = k; \\ \text{span}(e_1 + e_4, e_2, e_3, e_5, e_6, \dots, e_{i+1}), & \text{if } i \geq l. \end{cases}$$

That is, V is represented by the $n \times (n-1)$ matrix

$$A := \begin{bmatrix} 0 & (-1)^{k-1}B & 0 \\ I_{k-1} & 0 & 0 \\ 0 & 0 & I_{n-k-3} \end{bmatrix}, \quad \text{where} \quad B := \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}.$$

Note that all left-justified minors of A are nonnegative, except for a certain minor of order $k+1$. Since $k+1 \notin K$, we get that $V \in \text{Fl}_{K;n}^{\Delta \geq 0}$.

We claim that $V \notin \text{Fl}_{K;n}^{>0}$, which implies that $\text{Fl}_{K;n}^{>0} \neq \text{Fl}_{K;n}^{\Delta > 0}$. Indeed, suppose otherwise that $V \in \text{Fl}_{K;n}^{>0}$. Then by (2.4), we can extend V to a complete flag $(V_1, \dots, V_{n-1}) \in \text{Fl}_n^{>0}$. For $1 \leq i \leq n-1$, define $W_i := V_i \cap \text{span}(e_1, e_2, e_3, e_4)$. Let $d_i := \dim(W_i)$, so that $W_i \in \text{Gr}_{d_i, 4}^{\Delta > 0}$ by Lemma 2.12. Note that $W_k = \text{span}(e_1 + e_4)$ and $W_l = \text{span}(e_1 + e_4, e_2, e_3)$. Since the sequence d_k, d_{k+1}, \dots, d_l increases by 0 or 1 at each step, and $d_k = 1$ and $d_l = 3$, there exists $j \in [k, l]$ such that $d_j = 2$. Applying Lemma 3.5 to W_k and W_j , we get $e_1 \in W_j$. Since $W_j \subset W_l$, this implies $e_1 \in W_l$, a contradiction. \square

4. Cyclic symmetry

In this section, we prove Theorem 1.3.

Proof of Theorem 1.3. We claim that it suffices to construct $W \in \text{Fl}_{K;n}^{>0}$ such that $\sigma_\epsilon(W) \notin \text{Fl}_{K;n}^{>0}$. Indeed, let $f : \mathbb{R}_{\geq 0} \rightarrow \text{GL}_n(\mathbb{R})$ be as in Lemma 2.8, so that by Lemma 3.2(i), we have $f(t) \cdot W \in \text{Fl}_{K;n}^{>0}$ for all $t > 0$. If $\sigma_\epsilon(f(t) \cdot W) \in \text{Fl}_{K;n}^{>0}$ for all $t > 0$, then taking $t \rightarrow 0$ we obtain $\sigma_\epsilon(W) \in \text{Fl}_{K;n}^{>0}$, a contradiction. Therefore there exists $t > 0$ such that $\sigma_\epsilon(f(t) \cdot W) \notin \text{Fl}_{K;n}^{>0}$, whence we may take $V := f(t) \cdot W$.

Now we construct such a $W = (W_i)_{i \in K} \in \text{Fl}_{K;n}(\mathbb{R})$. Fix any two elements $k < l$ of K . We set

$$W_i := \begin{cases} \text{span}(e_3, e_4, \dots, e_{i+2}), & \text{if } i < k; \\ \text{span}(e_1 + e_2, e_3, e_4, \dots, e_{i+1}), & \text{if } i \geq k. \end{cases}$$

That is, W is represented by the $n \times (n - 1)$ matrix

$$A := \begin{bmatrix} 0 & (-1)^{k-1} & 0 \\ 0 & (-1)^{k-1} & 0 \\ I_{k-1} & 0 & 0 \\ 0 & 0 & I_{n-k-1} \end{bmatrix}.$$

Note that all left-justified minors of A are nonnegative, so A represents an element of $\text{Fl}_n^{\Delta \geq 0}$. By Theorem 1.1 we have $\text{Fl}_n^{\Delta \geq 0} = \text{Fl}_n^{\geq 0}$, so by (2.4), we get $W \in \text{Fl}_{K;n}^{\geq 0}$.

Let $X = (X_i)_{i \in K}$ denote the left cyclic shift $\sigma_\epsilon(W)$. Note that

$$X_k = \text{span}((-1)^{\epsilon-1}e_n + e_1, e_2, \dots, e_k) \quad \text{and} \quad X_l = \text{span}((-1)^{\epsilon-1}e_n + e_1, e_2, \dots, e_l).$$

Now proceed by contradiction and suppose that $X \in \text{Fl}_{K;n}^{\geq 0}$. Then by (2.4), we can extend X to a complete flag $(X_1, \dots, X_{n-1}) \in \text{Fl}_n^{\geq 0}$. Applying Lemma 3.5 to X_k and X_{k+1} , we get $e_1 \in X_{k+1}$. Since $X_{k+1} \subseteq X_l$, this implies $e_1 \in X_l$, a contradiction. \square

5. Cell decomposition vs. matroid decomposition

In this section, we prove Theorem 1.2. We begin by recalling some background in Section 5.1. We then give two proofs of the forward direction of Theorem 1.2 in Section 5.3, and prove the reverse direction in Section 5.4. Throughout this section, we fix $n \in \mathbb{N}$, and let W denote the symmetric group \mathfrak{S}_n of all permutations of $[n]$. Also, J and K will denote complementary subsets of $[n - 1]$.

5.1. Background on Coxeter combinatorics

We recall some background on the combinatorics of the Coxeter group $W = \mathfrak{S}_n$; we refer to [7] for further details.

Definition 5.1 ([7, Chapter 2]). For $1 \leq i \leq n - 1$, let $s_i := (i \ i + 1) \in W$ be the simple transposition which exchanges i and $i + 1$, and let $e \in W$ denote the identity permutation. Given $w \in W$, a *reduced word* \mathbf{w} for w is a word in s_1, \dots, s_{n-1} of minimal length whose product is w . Each reduced word for w has the same number of letters, called the *length* $\ell(w)$ of w , which is equal to the number of inversions of w . Any two reduced words for w are related by a sequence of moves of the following form:

- (M1) $s_i s_j = s_j s_i$ for $1 \leq i, j \leq n - 1$ with $|i - j| \geq 2$; and
- (M2) $s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}$ for $1 \leq i \leq n - 2$.




Fig. 1. The Hasse diagram of the Bruhat order on $W = \mathfrak{S}_3$.

In particular, if s_i appears in some reduced word for w , then it appears in every reduced word for w .

The (*strong*) *Bruhat order* \leq on W is defined as follows: $v \leq w$ if and only if for some (or equivalently, for every) reduced word \mathbf{w} for w , there exists a reduced word \mathbf{v} for v which is a subword of \mathbf{w} . The Bruhat order is graded with rank function ℓ . The Bruhat order on $W = \mathfrak{S}_3$ is shown in Fig. 1.

Example 5.2. Let $w := 5214763 \in W = \mathfrak{S}_7$. Then $\ell(w) = 9$, and a reduced word for w is $\mathbf{w} = s_1 s_3 s_4 s_3 s_2 s_1 s_5 s_6 s_5$. \diamond

We will need the following property of reduced words:

Lemma 5.3 ([7, Corollary 1.4.6(ii)]). *Let $w \in W$ and $1 \leq i \leq n - 1$. If $\ell(ws_i) < \ell(w)$, then w has a reduced word which ends in s_i .*

We define parabolic subgroups and quotients of W .

Definition 5.4 ([7, Section 2.4]). Given $J \subseteq [n - 1]$, let $W_J := \langle s_j : j \in J \rangle$ be the subgroup of W generated by the simple transpositions indexed by J , called a *parabolic subgroup*. Equivalently, W_J consists of the elements of W which setwise fix the intervals $[1, k_1], [k_1 + 1, k_2], \dots, [k_l + 1, n]$, where $[n - 1] \setminus J = \{k_1 < \dots < k_l\}$.

Let W^J denote the set of minimal-length coset representatives of the parabolic quotient W/W_J . Explicitly, we have

$$W^J = \{w \in \mathfrak{S}_n : w(j) < w(j + 1) \text{ for all } j \in J\}.$$

Each $w \in W$ has a unique factorization $w = w^J w_J$ such that $w^J \in W^J$ and $w_J \in W_J$; this factorization is length-additive. In particular, w^J is the minimal-length coset representative of w modulo W_J .

Example 5.5. Let $w := 5214763 \in W = \mathfrak{S}_7$, as in Example 5.2, and let $J := \{1, 2, 4, 6\}$. Then $w^J = 1254736 = s_3 s_4 s_3 s_6 s_5$ and $w_J = 3214576 = s_1 s_2 s_1 s_6$. \diamond

We will need the following property of the parabolic factorization:

Lemma 5.6 ([7, Proposition 2.5.1]). Let $J \subseteq [n-1]$, and let $v \leq w$ in W . Then $v^J \leq w^J$.

We now recall the *Demazure product* and *downwards Demazure product*, appearing in work of He [26, Lemma 3.3] and He and Lu [29, Appendix A]. We refer to [28, Section 2.1] for further discussion and references.

Definition 5.7 ([27, Section 1.3]). There exist binary operations $*$ and \triangleleft on W defined by

$$v * w := \max\{vx : x \leq w\} \quad \text{and} \quad v \triangleleft w := \min\{vx : x \leq w\}$$

for all $v, w \in W$. Equivalently,

$$v * (s_{i_1} \cdots s_{i_l}) = (\cdots (v * s_{i_1}) * \cdots) * s_{i_l} \quad \text{and} \quad v \triangleleft (s_{i_1} \cdots s_{i_l}) = (\cdots (v \triangleleft s_{i_1}) \triangleleft \cdots) \triangleleft s_{i_l}$$

for all $v \in W$ and reduced words $s_{i_1} \cdots s_{i_l} \in W$, where

$$v * s_i = \begin{cases} vs_i, & \text{if } \ell(vs_i) > \ell(v); \\ v, & \text{if } \ell(vs_i) < \ell(v) \end{cases} \quad \text{and} \quad v \triangleleft s_i = \begin{cases} v, & \text{if } \ell(vs_i) > \ell(v); \\ vs_i, & \text{if } \ell(vs_i) < \ell(v) \end{cases}$$

for all $1 \leq i \leq n-1$. We call $*$ the *Demazure product* and \triangleleft the *downwards Demazure product*.¹

Example 5.8. We have $s_1 s_2 s_3 * s_2 s_3 s_2 = s_1 s_2 s_3 s_2$ and $s_1 s_2 s_3 \triangleleft s_2 s_3 s_2 = s_1$. \diamond

We will need the following property of the Demazure and downwards Demazure products:

Lemma 5.9 ([27, Corollary 1 and Lemma 2]). Let $v \leq w$ in W . Then for all $x \in W$, we have $v * x \leq w * x$ and $v \triangleleft x \leq w \triangleleft x$.

5.2. Background on the cell and matroid decompositions

We recall the cell decomposition and matroid decomposition of $\mathrm{Fl}_{K;n}^{\geq 0}$, though we will mainly work with Theorem 5.15 and Theorem 5.16, rather than the definitions. We refer to [55, Sections 6–7] for further details.

Definition 5.10. Let $n \in \mathbb{N}$, and let $\mathrm{B}_n(\mathbb{R})$ and $\mathrm{B}_n^-(\mathbb{R})$ denote the subgroups of $\mathrm{GL}_n(\mathbb{R})$ of upper-triangular and lower-triangular matrices, respectively. For $w \in W$, let $\mathring{w} \in \mathrm{GL}_n(\mathbb{R})$ be any signed permutation matrix corresponding to w , i.e., $\mathring{w}_{i,j} = \pm \delta_{i,w(j)}$ for

¹ Our operation \triangleleft is the ‘mirror image’ of He’s \triangleright . We also caution that the symbol \triangleleft is used in [7] with a different meaning, namely, to denote a cover relation in the Bruhat order.

$1 \leq i, j \leq n$. Given $v, w \in W$ such that $v \leq w$, we define the (*totally nonnegative*) *Richardson cell*

$$C_{v,w} := (B_n^-(\mathbb{R}) \cdot \dot{v}) \cap (B_n(\mathbb{R}) \cdot \dot{w}) \cap \mathrm{Fl}_n^{\geq 0},$$

which is the intersection inside $\mathrm{Fl}_n^{\geq 0}$ of the opposite Schubert cell indexed by v and the Schubert cell indexed by w .

Now let J and K be complementary subsets of $[n-1]$. Given $v \in W$ and $w \in W^J$ such that $v \leq w$, we define the (*totally nonnegative*) *projected Richardson cell* $C_{v,w}^K \subseteq \mathrm{Fl}_{K;n}^{\geq 0}$ to be the image of $C_{v,w} \subseteq \mathrm{Fl}_n^{\geq 0}$ under the projection map (2.4). Rietsch [48,49] showed that $C_{v,w}^K$ is homeomorphic to an open ball of dimension $\ell(w) - \ell(v)$. We have the cell decomposition

$$\mathrm{Fl}_{K;n}^{\geq 0} = \bigsqcup_{\substack{v \in W, w \in W^J, \\ v \leq w}} C_{v,w}^K,$$

where $\mathrm{Fl}_{K;n}^{>0}$ is the unique cell of maximum dimension.

Remark 5.11. Our definition of the cell decomposition of $\mathrm{Fl}_{K;n}^{\geq 0}$ is different from, but equivalent to, the definition of Rietsch [49, Section 6]. We refer to [28, Appendix] and [19, Remark 4.9] for further discussion.

Definition 5.12. Let $K \subseteq [n-1]$. Given a tuple $M = (M_k)_{k \in K}$, where $M_k \subseteq \binom{[n]}{k}$ for $k \in K$, we define

$$S_M := \{V \in \mathrm{Fl}_{K;n}^{\geq 0} : \text{for all } k \in K \text{ and } I \in \binom{[n]}{k}, \text{ we have } \Delta_I(V) \neq 0 \Leftrightarrow I \in M_k\}.$$

If S_M is nonempty, we call it a (*totally nonnegative*) *matroid stratum*. The *matroid decomposition* (or *Gelfand–Serganova decomposition*) of $\mathrm{Fl}_{K;n}^{\geq 0}$ is its decomposition into matroid strata; equivalently, it is the common refinement of the decompositions

$$\mathrm{Fl}_{K;n}^{\geq 0} = \{V \in \mathrm{Fl}_{K;n}^{\geq 0} : \Delta_I(V) \neq 0\} \sqcup \{V \in \mathrm{Fl}_{K;n}^{\geq 0} : \Delta_I(V) = 0\}$$

for all Plücker coordinates Δ_I .

Remark 5.13. There is a different, but equivalent, way to define Plücker positivity and the matroid decomposition for partial flag varieties $\mathrm{Fl}_{K;n}(\mathbb{R})$, using the *generalized Plücker coordinates* of Gelfand and Serganova [25], rather than the Plücker coordinates of Definition 2.1. Namely, let $K = \{k_1 < \dots < k_l\} \subseteq [n-1]$. Given a tuple $I = (I_{k_1}, \dots, I_{k_l})$ such that $I_{k_1} \subset \dots \subset I_{k_l}$ and $I_k \in \binom{[n]}{k}$ for $k \in K$, define the *generalized Plücker coordinate*

$$\Delta_I := \Delta_{I_{k_1}} \Delta_{I_{k_2}} \cdots \Delta_{I_{k_l}}.$$

Then $V \in \mathrm{Fl}_{K;n}(\mathbb{R})$ is totally positive (respectively, totally nonnegative) if and only if all its generalized Plücker coordinates are positive (respectively, nonnegative). Also, the matroid decomposition of $\mathrm{Fl}_{K;n}^{\geq 0}$ is the common refinement of the decompositions

$$\mathrm{Fl}_{K;n}^{\geq 0} = \{V \in \mathrm{Fl}_{K;n}^{\geq 0} : \Delta_I(V) \neq 0\} \sqcup \{V \in \mathrm{Fl}_{K;n}^{\geq 0} : \Delta_I(V) = 0\}$$

for all generalized Plücker coordinates Δ_I . These results follow from [25, Section 9.1] (cf. [10, Chapter 1]).

Remark 5.14. We use the name *matroid decomposition* because if S_M is a matroid stratum of $\mathrm{Fl}_{K;n}^{\geq 0}$, then each M_k is a (*representable*) *matroid* of rank k on the ground set $[n]$ (in fact, M_k is a *positroid* [45, 46]). Moreover, M itself is a (*representable*) *Coxeter matroid*; see [25, Section 9.1] and [10, Section 1.7].

We also recall two results which will be key to our arguments; we refer to Section 1 for further discussion.

Theorem 5.15 (Tsukerman and Williams [55, Theorem 7.1]). *Let J and K be complementary subsets of $[n-1]$, and let $v \leq w$, where $v \in W$ and $w \in W^J$. Then the cell $C_{v,w}^K$ of $\mathrm{Fl}_{K;n}^{\geq 0}$ is contained in a single matroid stratum, which is uniquely determined by the interval $[v, w]$ modulo W_J .*

Note that Theorem 5.15 implies that the cell decomposition of $\mathrm{Fl}_{K;n}^{\geq 0}$ is a refinement of its matroid decomposition.

Theorem 5.16 (Postnikov [45, Theorem 3.8], [50]). *For $0 \leq k \leq n$, the cell decomposition of $\mathrm{Gr}_{k,n}^{\geq 0}$ coincides with the matroid decomposition.*

Remark 5.17. While it will be sufficient for our purposes to work with the combinatorial statement of Theorem 5.15, we mention that it has the following geometric interpretation; see [55, Sections 6–7] for further details. The moment polytope of $\mathrm{Fl}_{K;n}(\mathbb{C})$ is a convex polytope in \mathbb{R}^n whose vertices are indexed by W^J , or equivalently, by generalized Plücker coordinates of $\mathrm{Fl}_{K;n}(\mathbb{C})$ (see Remark 5.13). The moment polytope of $V \in \mathrm{Fl}_{K;n}(\mathbb{C})$ is contained in the moment polytope of $\mathrm{Fl}_{K;n}(\mathbb{C})$, and its vertices correspond precisely to the generalized Plücker coordinates which are nonzero at V [25, Proposition 5.1]. On the other hand, the set W^J also indexes the zero-dimensional cells of $\mathrm{Fl}_{K;n}^{\geq 0}$, i.e., the cells $C_{x,x}^K$ for $x \in W^J$. If $V \in C_{v,w}^K$, then the zero-dimensional cells in the closure of $C_{v,w}^K$ are precisely C_{x^J,x^J}^K for $x \in [v, w]$ [49, Theorem 6.1]. Theorem 5.15 can be rephrased as saying that the vertices of the moment polytope of $V \in C_{v,w}^K$ are indexed precisely by the zero-dimensional cells in the closure of $C_{v,w}^K$. Implicit in this statement is the fact that the moment polytope of V is equal to the moment polytope of $C_{v,w}^K$, even though the torus orbit of V may have dimension much less than that of $C_{v,w}^K$. This moment

polytope is called a *Bruhat interval polytope*, denoted² by $Q_{v,w}^K$. We will make a further comment about $Q_{v,w}^K$ in Remark 5.24.

5.3. Proof of the forward direction

In this subsection, we give two proofs of the forward direction of Theorem 1.2. The first proof uses Theorem 1.1, while the second proof uses Theorem 5.15.

For the first proof, we will need the following result of Rietsch [47]; see [8, Corollary 6.16] for a stronger result.

Lemma 5.18 (Rietsch [47, Lemma 5.2]). *Let $K \subseteq [n-1]$. Then $\mathrm{Fl}_{K;n}^{\Delta > 0}$ is connected.*

Proof of the forward direction of Theorem 1.2. We prove the contrapositive. Suppose that K does not consist of consecutive integers, so that by the implication (i) \Rightarrow (iii) of Theorem 1.1, $\mathrm{Fl}_{K;n}^{\Delta > 0}$ is strictly contained in $\mathrm{Fl}_{K;n}^{\Delta > 0}$. By Lemma 5.18, $\mathrm{Fl}_{K;n}^{\Delta > 0}$ is not closed in $\mathrm{Fl}_{K;n}^{\Delta > 0}$. Hence there exists a point $V \in (\mathrm{Fl}_{K;n}^{\Delta > 0} \setminus \mathrm{Fl}_{K;n}^{\Delta > 0}) \cap \mathrm{Fl}_{K;n}^{\Delta > 0}$. Then V and the cell $\mathrm{Fl}_{K;n}^{\Delta > 0}$ of $\mathrm{Fl}_{K;n}^{\Delta > 0}$ are contained in the same matroid stratum, namely, the one where all Plücker coordinates are nonzero. \square

We now proceed to the second proof of the forward direction of Theorem 1.2. It is based on the following lemma, which generalizes an example of Tsukerman and Williams [55, Remark 7.3].

Lemma 5.19. *Let $J := [2, n-2]$ and $K := \{1, n-1\}$, and let $w := (1 \ n) \in W^J$. Then for all $j \in J$, the intervals $[e, w]$ and $[s_j, w]$ are equal modulo W_J .*

Proof. Consider the reduced word

$$\mathbf{w} := s_1 s_2 \cdots s_{n-2} s_{n-1} s_{n-2} \cdots s_2 s_1$$

for w . In particular, we see that for $j \in J$, we indeed have $s_j \leq w$. Note that $[s_j, w] \subseteq [e, w]$. Conversely, we must show that given $x \in [e, w]$, there exists $y \in [s_j, w]$ such that x and y are equal modulo W_J . Take a subword \mathbf{x} of \mathbf{w} which is a reduced word for x . We will construct a reduced subword \mathbf{y} of \mathbf{w} which contains s_j , and such that the associated permutation y is equal to x modulo W_J .

If \mathbf{x} contains s_j , we set $\mathbf{y} := \mathbf{x}$. Now suppose that \mathbf{x} does not contain s_j . Note that \mathbf{w} contains two occurrences of s_{j-1} . Since \mathbf{x} does not contain s_j , it does not contain both occurrences of s_{j-1} , since otherwise we could use moves (M1) to obtain s_{j-1}^2 , contradicting the fact that \mathbf{x} is reduced. Similarly, if \mathbf{x} contains the second occurrence of s_{j-1} in \mathbf{w} , we may replace it with the first occurrence of s_{j-1} . Now let \mathbf{y} be obtained

² We caution that [55] uses the superscript J , rather than K .

from \mathbf{x} by including the second occurrence of s_j in \mathbf{w} . Since \mathbf{y} does not contain the second occurrence of s_{j-1} in \mathbf{w} , we can use (M1) to move s_j to the end of \mathbf{y} . That is, $y = xs_j$. This implies that \mathbf{y} is reduced; otherwise, by Lemma 5.3, x would have a reduced word ending in s_j , whereas \mathbf{x} (and hence every reduced word for x) does not contain s_j . Since $s_j \in W_J$, we see that y equals x modulo W_J . \square

Proof of the forward direction of Theorem 1.2. We prove the contrapositive. Suppose that K does not consist of consecutive integers, so that there exist consecutive elements $k < l$ of K with $l - k \geq 2$. Let $w := (k \ l) \in W^J$. Then by Lemma 5.19, for all $j \in [k+1, l-1]$, the intervals $[e, w]$ and $[s_j, w]$ are equal modulo W_J . Hence by Theorem 5.15, the cells $C_{e,w}^K$ and $C_{s_j,w}^K$ of $\text{Fl}_{K;n}^{\geq 0}$ are contained in the same matroid stratum. \square

5.4. Proof of the reverse direction

In this subsection, we prove the reverse direction of Theorem 1.2. We first establish two preliminary results, which will allow us to reduce the proof to Theorem 5.16.

Lemma 5.20. *Let $v \leq w$ in W , and let $J \subseteq [n-1]$. Set $v' := v \triangleleft w_J^{-1} \in W$ and $w' := w^J \in W^J$. Then $v' \leq w'$, and the intervals $[v, w]$ and $[v', w']$ are equal modulo W_J .*

Proof. Note that since the factorization $w = w^J w_J$ is length-additive, we have $w = w' * w_J$ and $w' = w \triangleleft w_J^{-1}$. In particular, $v' \leq w'$ by Lemma 5.9.

First we show that given $x' \in [v', w']$, there exists $x \in [v, w]$ such that x and x' are equal modulo W_J . We set $x := x' * w_J$. Since $w_J \in W_J$, we see that x equals x' modulo W_J . Also, by Lemma 5.9, we have

$$v \leq v' * w_J \leq x' * w_J \leq w' * w_J = w,$$

so $x \in [v, w]$.

Conversely, we show that given $x \in [v, w]$, there exists $x' \in [v', w']$ such that x and x' are equal modulo W_J . We set $x' := x \triangleleft w_J^{-1}$. Since $w_J^{-1} \in W_J$, we see that x' equals x modulo W_J . Also, $x' \in [v', w']$ by Lemma 5.9. \square

Example 5.21. As in Example 5.5, we let $w := s_1 s_3 s_4 s_3 s_2 s_1 s_5 s_6 s_5$ and $J := \{1, 2, 4, 6\}$. Take $v := s_1 s_4 s_3 s_2 s_1 s_5$, so that $v \leq w$. We set

$$v' := v \triangleleft w_J^{-1} = s_1 s_4 s_3 s_2 s_1 s_5 \triangleleft s_6 s_1 s_2 s_1 = s_4 s_3 s_5$$

and $w' := w^J = s_3 s_4 s_3 s_6 s_5$. Then Lemma 5.20 asserts that the intervals $[v, w]$ and $[v', w']$ are equal modulo W_J . Indeed, we can verify that both intervals modulo W_J are equal to

$$\{s_4 s_3 s_5, s_3 s_4 s_3 s_5, s_4 s_3 s_6 s_5, s_3 s_4 s_3 s_6 s_5\},$$

where above, we represent equivalence classes by elements of W^J . \diamond

Lemma 5.22. *Let J and K be complementary subsets of $[n-1]$, and let $v_1, v_2 \leq w$, where $v_1, v_2 \in W$ and $w \in W^J$. Suppose that the intervals $[v_1, w]$ and $[v_2, w]$ are equal modulo W_J . Then $v_1(i) = v_2(i)$ for all $i \leq \min(K)$ and all $i \geq \max(K) + 1$.*

Proof. We prove the statement for $i \leq \min(K)$; the statement for $i \geq \max(K) + 1$ follows by symmetry. Set $k := \min(K)$ and $J' := [n-1] \setminus \{k\}$, and note that $J \subseteq J'$. As in Lemma 5.20, we define $v'_1 := v_1 \triangleleft w_{J'}^{-1}$, $v'_2 := v_2 \triangleleft w_{J'}^{-1}$, and $w' := w^{J'}$, so that the intervals $[v_1, w]$, $[v_2, w]$, $[v'_1, w']$, and $[v'_2, w']$ are all equal modulo $W_{J'}$. By Theorem 5.16 (using Theorem 5.15), we obtain $v'_1 = v'_2$. Now since $w \in W^J$, we have $w(1) < \dots < w(k)$, so $w_{J'}$ is contained in the parabolic subgroup $W_{[k+1, n-1]}$. Hence $v_1(i) = v'_1(i) = v'_2(i) = v_2(i)$ for all $i \leq k$. \square

Proof of the reverse direction of Theorem 1.2. Suppose that $K = [k, l]$. Let $v_1 \leq w_1$ and $v_2 \leq w_2$, where $v_1, v_2 \in W$ and $w_1, w_2 \in W^J$. By Theorem 5.15, it suffices to show that if the intervals $[v_1, w_1]$ and $[v_2, w_2]$ are equal modulo W_J , then $v_1 = v_2$ and $w_1 = w_2$. To this end, we regard $[v_1, w_1]$ and $[v_2, w_2]$ modulo W_J as a subset of W^J ; by Lemma 5.6, this subset has minimum $v_1^J = v_2^J$ and maximum $w_1 = w_2$. In particular, $v_1(i) = v_2(i)$ for all $i \in [k+1, l]$. Therefore it remains to show that $v_1(i) = v_2(i)$ for all $i \leq k$ and all $i \geq l+1$. This follows from Lemma 5.22. \square

Example 5.23. We show how the argument above can fail when K is *not* an interval of integers. Take $n := 4$, $J := \{2\}$, $K := \{1, 3\}$, and

$$v_1 := 1234 = e, \quad v_2 := 1324 = s_2, \quad w := 4231 = s_1 s_2 s_3 s_2 s_1.$$

By Lemma 5.19 and Theorem 5.15, $C_{v_1, w}^K$ and $C_{v_2, w}^K$ are contained in the same matroid stratum. In agreement with Lemma 5.22, we have $v_1(i) = v_2(i)$ for all $i \leq 1$ and all $i \geq 4$. Also, we have $v_1^J = v_2^J = e$, but this does not imply that $v_1 = v_2$. \diamond

Remark 5.24. Recall the Bruhat interval polytopes $Q_{v,w}^K$ discussed in Remark 5.17. It follows from the theory of Coxeter matroids (namely, [10, Corollary 1.13.5]), along with a result of Tsukerman and Williams [55, Corollary 7.14] (cf. [10, Preface], [11, Theorem 6.3]), that every Bruhat interval polytope for $\text{Fl}_{K;n}(\mathbb{R})$ can be expressed as the Minkowski sum over $k \in K$ of a Bruhat interval polytope for $\text{Gr}_{k,n}(\mathbb{R})$. Lemma 5.20 allows us to write this Minkowski sum explicitly. Namely, for $v \leq w$ with $v \in W$ and $w \in W^J$, we have

$$Q_{v,w}^K = \sum_{k \in K} Q_{v \triangleleft w_{[n-1] \setminus \{k\}}^{-1}, w^{[n-1] \setminus \{k\}}}^{\{k\}}. \quad (5.1)$$

We point out that the Bruhat interval polytopes for $\text{Gr}_{k,n}(\mathbb{R})$ are known as *positroid polytopes* [1]; see [55, Proposition 2.8] for how to formulate (5.1) in terms of positroids.

As an illustration of (5.1), we adopt the setup of Example 5.23, with $v = v_1$. Then (5.1) gives

$$Q_{e, s_1 s_2 s_3 s_2 s_1}^{\{1,3\}} = Q_{e \triangleleft (s_2 s_3)^{-1}, s_3 s_2 s_1}^{\{1\}} + Q_{e \triangleleft (s_2 s_1)^{-1}, s_1 s_2 s_3}^{\{3\}},$$

or equivalently,

$$Q_{1234, 4231}^{\{1,3\}} = Q_{1234, 4123}^{\{1\}} + Q_{1234, 2341}^{\{3\}}.$$

Note that if we had instead taken $v = v_2$, we would have obtained the same Minkowski sum decomposition for $Q_{1324, 4231}^{\{1,3\}}$. Indeed, Lemma 5.19 implies that $Q_{1234, 4231}^{\{1,3\}} = Q_{1324, 4231}^{\{1,3\}}$.

References

- [1] F. Ardila, F. Rincón, L. Williams, Positroids and non-crossing partitions, *Trans. Am. Math. Soc.* 368 (1) (2016) 337–363.
- [2] F. Ardila, F. Rincón, L. Williams, Positively oriented matroids are realizable, *J. Eur. Math. Soc.* 19 (3) (2017) 815–833.
- [3] N. Arkani-Hamed, Y. Bai, T. Lam, Positive geometries and canonical forms, *J. High Energy Phys.* 2017 (2017) 039.
- [4] N. Arkani-Hamed, J. Bourjaily, F. Cachazo, A. Goncharov, A. Postnikov, J. Trnka, *Grassmannian Geometry of Scattering Amplitudes*, Cambridge University Press, Cambridge, 2016.
- [5] N. Arkani-Hamed, J. Trnka, The amplituhedron, *J. High Energy Phys.* 2014 (2014) 30.
- [6] Y. Bai, S. He, T. Lam, The amplituhedron and the one-loop Grassmannian measure, *J. High Energy Phys.* 2016 (2016) 112.
- [7] A. Björner, F. Brenti, *Combinatorics of Coxeter Groups*, Graduate Texts in Mathematics, vol. 231, Springer, New York, 2005.
- [8] A.M. Bloch, S.N. Karp, Gradient flows, adjoint orbits, and the topology of totally nonnegative flag varieties, *Commun. Math. Phys.* (2023), in press, arXiv:2109.04558.
- [9] J. Boretsky, Totally nonnegative tropical flags and the totally nonnegative flag Dressian, arXiv: 2208.09128.
- [10] A.V. Borovik, I.M. Gelfand, N. White, *Coxeter Matroids*, Progress in Mathematics, vol. 216, Birkhäuser Boston, Inc., Boston, MA, 2003.
- [11] F. Caselli, M. D’Adderio, M. Marietti, Weak generalized lifting property, Bruhat intervals, and Coxeter matroids, *Int. Math. Res. Not.* 3 (2021) 1678–1698.
- [12] N. Chevalier, Total positivity criteria for partial flag varieties, *J. Algebra* 348 (2011) 402–415.
- [13] C.W. Cryer, The LU -factorization of totally positive matrices, *Linear Algebra Appl.* 7 (1973) 83–92.
- [14] S.M. Fallat, C.R. Johnson, *Totally Nonnegative Matrices*, Princeton Series in Applied Mathematics, Princeton University Press, Princeton, NJ, 2011.
- [15] M. Fekete, G. Pólya, Über ein problem von Laguerre, *Rend. Circ. Mat. Palermo* 34 (1912) 89–120.
- [16] S. Fomin, L. Williams, A. Zelevinsky, Introduction to cluster algebras, Chapters 1–3, arXiv:1608.05735.
- [17] S. Fomin, A. Zelevinsky, Total positivity: tests and parametrizations, *Math. Intell.* 22 (1) (2000) 23–33.
- [18] S. Fomin, A. Zelevinsky, Totally nonnegative and oscillatory elements in semisimple groups, *Proc. Am. Math. Soc.* 128 (12) (2000) 3749–3759.
- [19] P. Galashin, S.N. Karp, T. Lam, Regularity theorem for totally nonnegative flag varieties, *J. Am. Math. Soc.* 35 (2) (2022) 513–579.
- [20] F.R. Gantmacher, *The Theory of Matrices*. Vols. 1, 2, Translated by K.A. Hirsch, Chelsea Publishing Co., New York, 1959.
- [21] F.R. Gantmacher, M.G. Krein, *Oscillation Matrices and Kernels and Small Vibrations of Mechanical Systems*, revised edition, AMS Chelsea Publishing, Providence, RI, 2002, Translation based on the 1950 Russian original. Edited and with a preface by Alex Eremenko.

- [22] F.R. Gantmaher, M.G. Krein, Oscillyacionye matricy i yadra i malye kolebaniya mehaničeskikh sistem, Gosudarstv. Isdat. Tehn.-Teor. Lit., Moscow-Leningrad, 1950, 2d ed. Translated into English by A. Eremenko [21].
- [23] F. Gantmakher, M. Krein, Sur les matrices complètement non négatives et oscillatoires, *Compos. Math.* 4 (1937) 445–476.
- [24] C. Geiss, B. Leclerc, J. Schröer, Preprojective algebras and cluster algebras, in: Trends in Representation Theory of Algebras and Related Topics, in: EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2008, pp. 253–283.
- [25] I.M. Gel'fand, V.V. Serganova, Combinatorial geometries and the strata of a torus on homogeneous compact manifolds, *Usp. Mat. Nauk* 42 (2) (1987) 107–134.
- [26] X. He, The G -stable pieces of the wonderful compactification, *Trans. Am. Math. Soc.* 359 (7) (2007) 3005–3024.
- [27] X. He, A subalgebra of 0-Hecke algebra, *J. Algebra* 322 (11) (2009) 4030–4039.
- [28] X. He, T. Lam, Projected Richardson varieties and affine Schubert varieties, *Ann. Inst. Fourier (Grenoble)* 65 (6) (2015) 2385–2412.
- [29] X. He, J.-H. Lu, On intersections of certain partitions of a group compactification, *Int. Math. Res. Not.* 11 (2011) 2534–2564.
- [30] S. Karlin, Total Positivity. Vol. I, Stanford University Press, Stanford, California, 1968.
- [31] S.N. Karp, Sign variation, the Grassmannian, and total positivity, *J. Comb. Theory, Ser. A* 145 (2017) 308–339.
- [32] D. Kazhdan, G. Lusztig, Representations of Coxeter groups and Hecke algebras, *Invent. Math.* 53 (2) (1979) 165–184.
- [33] A. Knutson, T. Lam, D.E. Speyer, Positroid varieties: juggling and geometry, *Compos. Math.* 149 (10) (2013) 1710–1752.
- [34] Y. Kodama, L. Williams, The full Kostant–Toda hierarchy on the positive flag variety, *Commun. Math. Phys.* 335 (1) (2015) 247–283.
- [35] T. Lam, Totally nonnegative Grassmannian and Grassmann polytopes, in: Current Developments in Mathematics 2014, Int. Press, Somerville, MA, 2016, pp. 51–152.
- [36] G. Lusztig, Canonical bases arising from quantized enveloping algebras, *J. Am. Math. Soc.* 3 (2) (1990) 447–498.
- [37] G. Lusztig, Total positivity in reductive groups, in: Lie Theory and Geometry, in: Progr. Math., vol. 123, Birkhäuser Boston, Boston, MA, 1994, pp. 531–568.
- [38] G. Lusztig, Total positivity in partial flag manifolds, *Represent. Theory* 2 (1998) 70–78.
- [39] G. Lusztig, On the totally positive grassmannian, arXiv:1905.09254.
- [40] G. Lusztig, On the totally positive grassmannian, arXiv:1905.09254, version 1.
- [41] R.J. Marsh, K. Rietsch, Parametrizations of flag varieties, *Represent. Theory* 8 (2004) 212–242.
- [42] N.E. Mnëv, The universality theorems on the classification problem of configuration varieties and convex polytopes varieties, in: Topology and Geometry—Rohlin Seminar, in: Lecture Notes in Math., vol. 1346, Springer, Berlin, 1988, pp. 527–543.
- [43] A. Pinkus, *Totally Positive Matrices*, Cambridge Tracts in Mathematics, vol. 181, Cambridge University Press, Cambridge, 2010.
- [44] G. Pólya, G. Szegö, *Aufgaben und Lehrsätze aus der Analysis. Band II*, Die Grundlehren der mathematischen Wissenschaften, vol. 20, Springer, Berlin, 1925.
- [45] A. Postnikov, Total positivity, Grassmannians, and networks, <https://math.mit.edu/~apost/papers/tpgrass.pdf>, 2007.
- [46] A. Postnikov, D. Speyer, L. Williams, Matching polytopes, toric geometry, and the totally non-negative Grassmannian, *J. Algebraic Comb.* 30 (2) (2009) 173–191.
- [47] K.C. Rietsch, Total Positivity and Real Flag Varieties, Ph.D. thesis, Massachusetts Institute of Technology, 1998.
- [48] K. Rietsch, An algebraic cell decomposition of the nonnegative part of a flag variety, *J. Algebra* 213 (1) (1999) 144–154.
- [49] K. Rietsch, Closure relations for totally nonnegative cells in G/P , *Math. Res. Lett.* 13 (5–6) (2006) 775–786.
- [50] K. Rietsch, Unpublished notes, 2009.
- [51] K. Rietsch, L. Williams, Newton–Okounkov bodies, cluster duality, and mirror symmetry for Grassmannians, *Duke Math. J.* 168 (18) (2019) 3437–3527.
- [52] I. Schoenberg, Über variationsvermindernde lineare Transformationen, *Math. Z.* 32 (1) (1930) 321–328.

- [53] D. Speyer, L. Williams, The tropical totally positive Grassmannian, *J. Algebraic Comb.* 22 (2) (2005) 189–210.
- [54] K. Talaska, L. Williams, Network parametrizations for the Grassmannian, *Algebra Number Theory* 7 (9) (2013) 2275–2311.
- [55] E. Tsukerman, L. Williams, Bruhat interval polytopes, *Adv. Math.* 285 (2015) 766–810.
- [56] A.M. Whitney, A reduction theorem for totally positive matrices, *J. Anal. Math.* 2 (1952) 88–92.
- [57] L.K. Williams, Shelling totally nonnegative flag varieties, *J. Reine Angew. Math.* 609 (2007) 1–21.