



Quasiplanar Graphs, String Graphs, and the Erdős-Gallai Problem

Jacob Fox¹, János Pach², and Andrew Suk³(✉)

¹ Stanford University, Stanford, CA, USA

jacobfox@stanford.edu

² Rényi Institute, Budapest and IST Austria, Budapest, Hungary

pach@cims.nyu.edu

³ University of California at San Diego, La Jolla, CA, USA

asuk@ucsd.edu

Abstract. An r -quasiplanar graph is a graph drawn in the plane with no r pairwise crossing edges. Let $s \geq 3$ be an integer and $r = 2^s$. We prove that there is a constant C such that every r -quasiplanar graph with $n \geq r$ vertices has at most $n(Cs^{-1} \log n)^{2s-4}$ edges.

A graph whose vertices are continuous curves in the plane, two being connected by an edge if and only if they intersect, is called a *string graph*. We show that for every $\epsilon > 0$, there exists $\delta > 0$ such that every string graph with n vertices, whose chromatic number is at least n^ϵ contains a clique of size at least n^δ . A clique of this size or a coloring using fewer than n^ϵ colors can be found by a polynomial time algorithm in terms of the size of the geometric representation of the set of strings.

In the process, we use, generalize, and strengthen previous results of Lee, Tomon, and others. All of our theorems are related to geometric variants of the following classical graph-theoretic problem of Erdős, Gallai, and Rogers. Given a K_r -free graph on n vertices and an integer $s < r$, at least how many vertices can we find such that the subgraph induced by them is K_s -free?

Keywords: Quasi-planar graphs · String graphs · Graph coloring

1 Introduction

A *topological graph* is a graph drawn in the plane with points as vertices and edges as continuous curves connecting some pairs of vertices. The curves are allowed to cross, but they may not pass through vertices other than their endpoints. If the edges are drawn as straight-line segments, then the graph is *geometric*. If no r edges in a topological graph G are pairwise crossing, then G is called *r -quasiplanar*.

J. Fox—Supported by a Packard Fellowship and by NSF award DMS-1855635.

J. Pach—Supported by NKFIH grants K-131529, Austrian Science Fund Z 342-N31, and ERC Advanced Grant “GeoScape.”

A. Suk—Supported by an NSF CAREER award and NSF award DMS-1952786.

© The Author(s), under exclusive license to Springer Nature Switzerland AG 2023
P. Angelini and R. von Hanxleden (Eds.): GD 2022, LNCS 13764, pp. 219–231, 2023.

https://doi.org/10.1007/978-3-031-22203-0_16

The following is a longstanding unsolved problem in the theory of topological graphs; see, *e.g.*, [5].

Conjecture 1. The number of edges of every r -quasiplanar graph of n vertices is $O_r(n)$.

Conjecture 1 has been proved for $r \leq 4$. See [1, 3, 4].

The *intersection graph* of a family of geometric objects, $\mathcal{S}$, is a graph with vertex set $\mathcal{S}$, in which two vertices are joined by an edge if and only if their intersection is nonempty. If $\mathcal{S}$ consists of continuous curves (or line segments) in the plane, then their intersection graph is called a *string graph* (resp., a *segment graph*).

A natural approach to prove Conjecture 1 is the following. Removing a small disc around every vertex of an r -quasiplanar graph G , we are left with a family of continuous curves $\mathcal{S}$ in the plane, no r of which are pairwise crossing. These curves define a K_r -free string graph H . Suppose that the chromatic number of H satisfies $\chi(H) \leq f(r)$. Then each color class corresponds to the edges of a planar subgraph of G . Thus, the size of each color class is at most $3n - 6$, provided that $n \geq 3$. This would immediately imply that every r -quasiplanar graph with n vertices has at most $(3n - 6)f(r) = O_r(n)$ edges, as required.

Surprisingly, this approach is not viable. In 2014, Pawlik, Kozik, Krawczyk, Lasoń, Micek, Trotter, and Walczak [28] represented a class of K_3 -free graphs originally constructed by Burling [6] as segment graphs whose chromatic numbers can be arbitrarily large. Shortly after, Walczak [35] strengthened this result by proving that there are K_3 -free segment graphs on n vertices in which every independent set is of size $O(\frac{n}{\log \log n})$.

Using the same approach, in order to prove Conjecture 1 for some r , it would be sufficient to show that there is a constant $g(r)$ with the property that the vertex set of every K_r -free string graph can be colored by $g(r)$ colors such that each (string) graph induced by one of the color classes is K_4 -free. Indeed, the result of Ackerman [1] cited above implies that the number of edges in each color class is $O(n)$. The first question to answer is the following.

Problem 1. Fix an integer $r \geq 4$. Is it true that every K_r -free segment graph on n vertices has an induced subgraph on $\Omega_r(n)$ vertices which is K_{r-1} -free?

Building upon the work of McGuinness [26], Suk [32] showed that every K_r -free *segment graph* on n vertices has a K_{r-1} -free induced subgraph with at least $\Omega_r(\frac{n}{\log n})$ vertices. (See also [29, 30].) For *string graphs*, in general, until now the best known result, due to Fox and Pach [17], was weaker: they could only guarantee the existence of an independent set and, hence, a K_{r-1} -free induced subgraph, of size at least $\frac{n}{(\log n)^{O(\log r)}}$.

In a different range, where r grows polynomially in n , Tomon [33] solved a longstanding open problem by showing that there is a constant $c' > 0$ such that every string graph on n vertices has a clique or an independent set of size $n^{c'}$.

Our next theorem slightly strengthens the result of Fox and Pach [17].

Theorem 1. *Let s be a positive integer and $r = 2^s$. Every K_r -free string graph on $n \geq r$ vertices has an independent set of size at least $n(cs/\log n)^{2s-2}$, where $c > 0$ is an absolute constant.*

At the beginning of Sect. 4, we show how to deduce from Theorem 1 the following strengthening of Tomon's above mentioned theorem [33].

Corollary 1. *For any $\epsilon > 0$, there is $\delta > 0$ such that every string graph G on n vertices has a clique of size at least n^δ or its chromatic number is at most n^ϵ . (In the latter case, obviously, G has an independent set of size at least $n^{1-\epsilon}$.)*

Theorem 1 guarantees the existence of a large *independent* set in a K_r -free string graph G . If, in the spirit of Problem 1, we want to find only a large K_{r-1} -free induced subgraph in G , we can do better.

Theorem 2. *For any $n \geq r \geq 3$, every K_r -free string graph with n vertices has a K_{r-1} -free induced subgraph with at least $c \frac{n}{\log^2 n}$ vertices, where $c > 0$ is an absolute constant.*

At the expense of another logarithmic factor, we can also find an induced subgraph with no clique of size $\lceil r/2 \rceil$.

Theorem 3. *For any $n \geq r \geq 3$, every K_r -free string graph with n vertices has a $K_{\lceil r/2 \rceil}$ -free induced subgraph with at least $c \frac{n}{\log^3 n}$ vertices, where $c > 0$ is an absolute constant.*

Now we return to the original motivation behind our present note: to estimate from above the number of edges of an r -quasiplanar *topological* graph of n vertices. As mentioned before, for $r \leq 4$, Conjecture 1 is true. For any $r \geq 5$, the best previously known upper bounds were $n(\log n)^{O(\log r)}$ and $O(n(\log n)^{4r-16})$, established in [17] and [27], respectively. For *geometric* graphs, for any fixed $r \geq 5$, Valtr [34] obtained the upper bound $O(n \log n)$. See [2] for a survey.

Using the result of Ackerman [1] as the base case of an induction argument, and exploiting several properties of string graphs established by Lee [24], Tomon [33], and Fox and Pach [16, 17], we will deduce the following improved upper bound for the number of edges of r -quasiplanar topological graphs.

Theorem 4. *Let $s \geq 3$ be an integer and $r = 2^s$. Then every r -quasiplanar graph with $n \geq r$ vertices has at most $n(\frac{c \log n}{s})^{2s-4}$ edges, where $c > 0$ is an absolute constant.*

Setting $s = 3$, for instance, we obtain that every 8-quasiplanar topological graph on n vertices has $O(n(\log n)^2)$ edges, which is better than the previously known bound of $O(n(\log n)^{16})$ [1, 27]. For $r = \delta \log n$, Theorem 4 immediately implies the following.

Corollary 2. *For any $\epsilon > 0$ there is $\delta > 0$ such that every topological graph with n vertices and at least $3n^{1+\epsilon}$ edges has n^δ pairwise crossing edges.*

The factor 3 in front of the term $n^{1+\epsilon}$ guarantees that the graph is not planar. Otherwise, we could not even guarantee that there is one crossing pair of edges.

In the special case where the strings are allowed to cross only a bounded number of times, some results very similar to Theorems 1 and 4 were established in [15].

Theorems 1, 2, 3, and Corollary 1 guarantee the existence of an independent set or a K_p -free induced subgraph for some $p > 2$ in a string graph satisfying certain conditions. All of these sets and subgraphs can be found by *efficient polynomial time algorithms* in terms of the size of a geometric representation of the underlying string graph. For example, the proof of Corollary 1 yields the following algorithmic result.

Proposition 1. *For any $\epsilon > 0$ there is $\delta > 0$ with the following property. Given a representation of a string graph on n vertices as an intersection graph of strings, there is a polynomial time algorithm which either properly colors the vertices with n^ϵ colors or finds a clique of size n^δ .*

Erdős and Gallai [11] raised the following question. Given a pair of integers, $2 \leq p < r$, how large of a K_p -free induced subgraph must be contained in every K_r -free graph of n vertices? For $p = 2$, we obtain Ramsey’s problem: how large of an independent set must be contained in every K_r -free graph of n vertices. The special case $p = r - 1$ was considered by Erdős and Rogers [12]. These problems have since been extensively studied. For many striking results, see, e.g., [9, 10, 19, 22, 23, 31, 36]. Apart from our last two results listed in the introduction, all statements in this paper can be regarded as geometric variants of the Erdős-Gallai-Rogers problem for string graphs.

The rest of this note is organized as follows. In Sect. 2, we apply the analogues of the separator theorem and the Kővári-Sós-Turán theorem for string graphs [17, 24] to establish Theorems 2 and 3. In Sect. 3, we present a simple technical lemma (Lemma 3) and some of its consequences needed for the proof of Theorems 1 and 4. The proofs of these two theorems and Corollary 1 are given in Sect. 4. The last section contains some concluding remarks.

Throughout this paper, $\log$ always stands for the binary logarithm. The letters c and C appearing in different theorems denote unrelated positive constants. Whenever they are not important, we will simply omit floor and ceiling signs.

2 Separators—Proofs of Theorems 2 and 3

In this section, we prove Theorems 2 and 3. We need the separator theorem for string graphs, due to Lee [24]. A *separator* in a graph $G = (V, E)$ is a subset S of the vertex set V such that no connected component of $G \setminus S$ has more than $\frac{2}{3}|V|$ vertices. Equivalently, S is a separator of G if there is a partition $V = S \cup V_1 \cup V_2$ with $|V_1|, |V_2| \leq \frac{2}{3}|V|$ such that no vertex in V_1 is adjacent to any vertex in V_2 .

Lemma 1 ([24]). *Every string graph with m edges has a separator of size at most $c_1\sqrt{m}$, where c_1 is an absolute constant.*

We now prove the following theorem which immediately implies Theorem 2. Let us remark that the *neighborhood* of vertex v does not include v .

Theorem 5. *There is an absolute constant $c > 0$ with the following property. Every string graph G on n vertices contains an induced subgraph G' on $c\frac{n}{\log^2 n}$ vertices whose every connected component is contained in the neighborhood of a vertex or is an isolated vertex.*

Proof. Let $c > 0$ be a sufficiently small constant to be specified later. We proceed by induction on n . The base case when $n = 1$ is trivial. For the inductive step, assume that the statement holds for all $n' < n$. Let $G = (V, E)$ be an n -vertex string graph.

If G contains a vertex v of degree at least $cn/\log^2 n$, then we are done by setting G' to be the neighborhood of v . Otherwise, we know that there are at most $cn^2/\log^2 n$ edges in G . By Lemma 1, G has a separator $S \subset V$ of size at most $c_1\sqrt{cn}/\log n$, where c_1 is the absolute constant from Lemma 1. Hence, there is a partition $V = S \cup V_1 \cup V_2$ with $|V_1|, |V_2| \leq \frac{2}{3}|V|$ such that no vertex in V_1 is adjacent to any vertex in V_2 , and $|S| \leq c_1\sqrt{cn}/\log n$. By applying induction on V_1 and V_2 and setting $c < \left(\frac{\log 3/2}{c_1}\right)^2$, we obtain an induced subgraph G' on at least

$$\begin{aligned} c \frac{|V_1|}{\log^2 |V_1|} + c \frac{|V_2|}{\log^2 |V_2|} &\geq c \frac{|V_1| + |V_2|}{\log^2 (2n/3)} \geq c \frac{n - \frac{c_1\sqrt{cn}}{\log n}}{\log^2 (2n/3)} \\ &= c \frac{n}{\log^2 n} \cdot \frac{1 - \frac{c_1\sqrt{c}}{\log n}}{\left(1 - \frac{\log(3/2)}{\log n}\right)^2} \geq c \frac{n}{\log^2 n} \end{aligned}$$

vertices such that each component of G' is contained in the neighborhood of a vertex or is an isolated vertex.

To see that Theorem 5 implies Theorem 2, it is sufficient to notice that if G' has a clique of size $r - 1$, then G has a clique of size r .

The proof of Theorem 3 is very similar to that of Theorem 2. Here, we need the following analogue of the Kővári-Sós-Turán theorem, which can also be deduced from Lemma 1 (see Conjecture 3.3 in [17]).

Lemma 2 ([17, 24]).

Every $K_{t,t}$ -free string graph on n vertices has at most $c_2 t(\log t)n$ edges, where c_2 is an absolute constant.

Proof of Theorem 3. Let $c > 0$ be a sufficiently small constant to be determined later. We proceed by induction on n . The base case $n = 1$ is trivial. For the

inductive step, assume that the statement holds for all $n' < n$. Let $G = (V, E)$ be a K_r -free string graph on n vertices, and let c_1 and c_2 be the constants from Lemmas 1 and 2, respectively.

If G has at least $cc_2 \frac{n^2}{\log^2 n}$ edges, then, by Lemma 2, G contains a complete bipartite graph $K_{t,t}$, where $t \geq c \frac{n}{\log^3 n}$. Since G is K_r -free, one of these parts must be $K_{\lceil r/2 \rceil}$ -free, and we are done.

Otherwise, if G has fewer than $cc_2 \frac{n^2}{\log^2 n}$ edges, then, by Lemma 1, there is a partition $V = S \cup V_1 \cup V_2$ with $|V_1|, |V_2| \leq \frac{2}{3}|V|$ such that no vertex in V_1 is adjacent to any vertex in V_2 , and $|S| \leq c_1 \sqrt{cc_2 n} / \log n$. Applying the induction hypothesis to V_1 and V_2 , and setting $c < \frac{\log^2(3/2)}{c_2 c_1^2}$, we obtain a $K_{\lceil r/2 \rceil}$ -free induced subgraph $G' \subseteq G$ with at least

$$\begin{aligned} c \frac{|V_1|}{\log^3 |V_1|} + c \frac{|V_2|}{\log^3 |V_2|} &\geq c \frac{|V_1| + |V_2|}{\log^3 (2n/3)} \geq c \frac{n - \frac{c_1 \sqrt{cc_2 n}}{\log n}}{\log^3 (2n/3)} \\ &= c \frac{n}{\log^3 n} \cdot \frac{1 - \frac{c_1 \sqrt{cc_2}}{\log n}}{\left(1 - \frac{\log(3/2)}{\log n}\right)^3} \geq c \frac{n}{\log^3 n} \end{aligned}$$

vertices. □

3 A Technical Lemma for String Graphs

The *average degree* in a graph $G = (V, E)$ is $d = \frac{2|E|}{|V|}$. The *edge density* of G is defined as $\frac{|E|}{\binom{|V|}{2}} = \frac{d}{|V|-1}$. We say that a graph is *dense* if its edge density is larger than some positive constant (which we will conveniently specify for our purposes).

Using Lee's separator theorem for string graphs (Lemma 1), it is easy to deduce the following technical lemma which states that every string graph G has a dense induced subgraph G' whose average degree is not much smaller than the average degree in G .

Lemma 3. *For any $\epsilon > 0$, there is $C = C(\epsilon)$ with the following property. Every string graph $G = (V, E)$ with average degree $d = 2|E|/|V|$ has an induced subgraph $G[V']$ with average degree $d' \geq (1 - \epsilon)d$ and $|V'| \leq Cd'$.*

Proof. Let $G = (V, E)$ be a string graph with average degree d . We recursively define a nested sequence of induced subgraphs $G_0 \supset G_1 \supset \dots$.

We begin with $G_0 = G$, and let $V_0 = V$, $E_0 = E$ and $d_0 = d$. After obtaining $G_i = (V_i, E_i)$ with $E_i = E(G[V_i])$ and with average degree $d_i = 2|E_i|/|V_i|$, we show that G_i is the desired induced subgraph if $d_i \geq |V_i|/C$. Otherwise if

$d_i < |V_i|/C$, we have $|E_i| \leq |V_i|^2/(2C)$, and by Lemma 1, there is a partition $V_i = U_0 \cup U_1 \cup U_2$ with $|U_1|, |U_2| \leq 2|V_i|/3$,

$$|U_0| \leq c_1 \sqrt{|E_i|} \leq c_1 \frac{|V_i|}{\sqrt{2C}} \leq |V_i|/12,$$

and there are no edges from U_1 to U_2 . The last inequality above follows from the fact that $C \geq (12c_1)^2$. We can assume this as we can choose C as large as we want.

We take G_{i+1} to be the induced subgraph on whichever of $G[U_1 \cup U_0]$ and $G[U_2 \cup U_0]$ has larger average degree. As all edges of G_i are in at least one of these two induced subgraphs and $|U_1 \cup U_0| + |U_2 \cup U_0| = |V_i| + |U_0|$, the average degree of G_{i+1} satisfies

$$\begin{aligned} d_{i+1} &\geq d_i \frac{|V_i|}{|V_i| + |U_0|} = d_i \frac{1}{1 + |U_0|/|V_i|} \\ &\geq d_i \frac{1}{1 + c_1 \sqrt{|E_i|}/|V_i|} \geq d_i \frac{1}{1 + c_1 \sqrt{d_i/(2|V_i|)}}. \end{aligned}$$

As $d_i < |V_i|/C$ and C can be chosen sufficiently large, the above inequality implies that $d_{i+1} \geq \frac{9}{10}d_i$. The inequality $|U_0| \leq |V_i|/12$ implies that $|V_{i+1}| \leq \frac{3}{4}|V_i|$. These two inequalities together imply that $d_{i+1}/|V_{i+1}| \geq \frac{6}{5}d_i/|V_i|$. It follows from the inequality above that

$$d_{i+1} = d \prod_{j=0}^i d_{j+1}/d_j \geq d \prod_{j=0}^i \frac{1}{1 + c_1 \sqrt{d_j/(2|V_j|)}} \geq de^{-\sum_{j=0}^i c_1 \sqrt{d_j/(2|V_j|)}},$$

where the last inequality uses that $\frac{1}{1+x} \geq e^{-x}$ for any $x > 0$. The sum in the exponent is dominated by a geometric series with common ratio $\sqrt{6/5} > 1$, and its largest summand is at most $c_1(1/(2C))^{1/2}$, as $d_i \leq |V_i|/C$. Hence, the sum in the exponent is $O(C^{-1/2})$. Taking C large enough, we have that $d_{i+1} \geq (1 - \epsilon)d$ for every i for which d_{i+1} is defined. (We can choose $C = O(1/\epsilon^2)$ to satisfy this.) Further, as $|V_{i+1}| \leq \frac{3}{4}|V_i| < |V_i|$ for every i for which V_{i+1} is defined, after at most $|V|$ iterations, the above process will terminate with the desired induced subgraph G_i .

The main result of [16] is that every dense string graph contains a dense spanning subgraph which is an incomparability graph. Applying Lemma 3 to this spanning subgraph with $\epsilon = 1/2$, we obtain the following corollary.

Corollary 3. *There is a constant $c > 0$ with the following property. Every string graph with n vertices and m edges has a subgraph with at least $c \frac{m}{n}$ vertices which is an incomparability graph with edge density at least c .*

Given a graph $G = (V, E)$ and two disjoint subsets of vertices $X, Y \in V$, we say that X is *complete to* Y if $xy \in E$ for all $x \in X$ and $y \in Y$.

The next lemma can be deduced by combining Corollary 3 above with Lemmas 6 and 7 of Tomon [33].

Lemma 4. *There is a constant $c > 0$ with the following property. If $G = (V, E)$ is a string graph with n vertices and at least αn^2 edges, for some $\alpha > 0$, then there are disjoint vertex subsets $X_1, \dots, X_t \subset V$ for some $t \geq 2$ such that*

1. X_i is complete to X_j for all $i \neq j$, and
2. $|X_i| \geq c\alpha \frac{n}{t^2}$ for every i .

4 Back to Quasiplanar Graphs—Proofs of Theorems 1 and 4

Before turning to the proof of Theorem 1, we show how it implies Corollary 1.

Proof of Corollary 1. The most natural technique for properly coloring a graph is by successively extracting maximum independent sets from it. Using this greedy method and the bound in Theorem 1, for $r = 2^s$, we obtain a proper coloring of any K_r -free string graph on n vertices with at most $(\frac{\log n}{cs})^{2s-2} \log n$ colors. Indeed, each time we extract a maximum independent set, the fraction of remaining vertices is at most $1 - \alpha$ with $\alpha = (\frac{cs}{\log n})^{2s-2}$. As $1 - \alpha < e^{-\alpha}$, after at most $\frac{\log n}{\alpha}$ iterations, no vertex remains.

For a given $\epsilon > 0$, choose a sufficiently small $\delta > 0$ be such that

1. $2\delta \log \frac{1}{c\delta} < \frac{\epsilon}{2}$ and
2. $\log n < n^{\epsilon/2}$ provided that $n^\delta \geq 2$.

Consider any K_{n^δ} -free string graph G on n vertices. If $n^\delta < 2$, then G has no edges and, hence, its chromatic number is $1 \leq n^\epsilon$. Otherwise, substituting $s = \delta \log n$, Theorem 1 yields that the chromatic number of G is at most

$$n^{2\delta \log \frac{1}{c\delta}} \log n < n^\epsilon.$$

□

Now we turn to the proof of Theorem 1 which gives, for $r = 2^s$, a lower bound on the independence number of a K_r -free string graph on n vertices.

Proof of Theorem 1. Our proof is by double induction on s and n . Throughout we let $r = 2^s$. The base cases are when $s = 1$ (in which case we get an independent set of size n) or $n = r$ (in which case we get an independent set of size 1) and are trivial. The induction hypothesis is that the theorem holds for all $s' < s$ and all n' , and for $s' = s$ and all $n' < n$. Note that we may assume that $r \leq n/4$, as otherwise the theorem holds. Let $\alpha = c'(\frac{s}{\log n})^2$, where $c' > 0$ is a sufficiently small absolute constant. Let G be a K_r -free string graph on n vertices.

If G has at most αn^2 edges, applying Lemma 1, there is a vertex partition $V = V_0 \cup V_1 \cup V_2$ with $|V_0| \leq c_1 \alpha^{1/2} n$, $|V_1|, |V_2| \leq 2n/3$, and there are no edges from V_1 to V_2 . Note that $|V_0| \leq n/12$ so $|V_1|, |V_2| \geq n/4$. We obtain a large independent set in G by taking the union of large independent sets in V_1 and V_2 . Using the induction hypothesis applied to $G[V_1]$ and $G[V_2]$, we obtain an independent set in G of order at least

$$|V_1| \left(\frac{cs}{\log |V_1|} \right)^{2s-2} + |V_2| \left(\frac{cs}{\log |V_2|} \right)^{2s-2} \geq (|V_1| + |V_2|) \left(\frac{cs}{\log(2n/3)} \right)^{2s-2}.$$

Note that $|V_1| + |V_2| = n - |V_0| \geq n(1 - c_1 c'^{1/2} \cdot \frac{s}{\log n})$. We also have

$$(\log(2n/3))^{2s-2} = (\log n)^{2s-2} \left(1 - \frac{\log(3/2)}{\log n} \right)^{2s-2} \leq (\log n)^{2s-2} \left(1 - \frac{s}{2 \log n} \right).$$

Substituting in these estimates and using $c' > 0$ is sufficiently small, we obtain an independent set of the desired size.

Suppose next that G has at least αn^2 edges. By Lemma 4, there is an integer $t \geq 2$ and disjoint vertex subsets $X_1, \dots, X_t$ such that X_i is complete to X_j for all $i \neq j$ and $|X_i| \geq 4c''\alpha n/t^2$ for $i = 1, \dots, t$ where $0 < 4c'' < 1$ is an absolute constant. Losing a factor at most 2 in the number of sets X_i , we may assume $t = 2^p$ for a positive integer $p < s$, which implies $|X_i| \geq c''\alpha n/t^2$. As G is K_{2s} -free, one of these sets X_i induces a subgraph which is K_{2s-p} -free. Let $n_0 = |X_i|$. Applying the induction hypothesis to $G[X_i]$, we obtain an independent set of size at least

$$\begin{aligned} n_0 \left(\frac{c(s-p)}{\log n_0} \right)^{2(s-p)-2} &\geq c''c' \left(\frac{s}{\log n} \right)^2 n 2^{-2p} \left(\frac{c(s-p)}{\log n_0} \right)^{2(s-p)-2} \\ &\geq n \left(\frac{cs}{\log n} \right)^{2s-2}. \end{aligned}$$

The last inequality holds, because after substituting $\log n_0 \leq \log n$, the ratio of the right-hand side and the expression in the middle reduces to

$$\frac{(2c)^{2p}}{c''c'} \left(\frac{s}{\log n} \right)^{2(p-1)} \left(1 + \frac{p}{s-p} \right)^{2(s-p)-2} \leq \frac{(2ec)^{2p}}{c''c'} \left(\frac{s}{\log n} \right)^{2(p-1)} \leq 1.$$

At the first inequality, we used $1 + x \leq e^x$ with $x = \frac{p}{s-p}$. As for the second inequality, we know that $s \leq \log n$, and we are free to choose the constant $c > 0$ as small as we wish (for instance, $c = c''c'/30$ will suffice). This completes the proof. $\square$

A careful inspection of the proof of Theorem 1 shows that it recursively constructs an independent set of the desired size in a K_r -free string graph in polynomial time in terms of the size of the geometric representation of the set of strings. Indeed, the proof itself is essentially algorithmic. In the first case, when the string graph is relatively sparse, we apply Lee's separator theorem for string graphs, and take the union of large independent sets from the string graph of the two remaining large vertex subsets after deleting the small separator. In the second case, when the string graph is relatively dense, we apply Lemma 4 to get in the string graph a complete multipartite subgraph with large

parts, and we can find a large independent set in one of the parts. However, this does require checking that results from several earlier papers each yield desirable structures in string graphs and incomparability graphs in polynomial time. These results include Lee's separator theorem for string graphs [24], the Fox-Pach result that every dense string graph contains a dense spanning subgraph which is an incomparability graph [16], and some extremal results of Tomon [33] for incomparability graphs.

A set of vertices $X \subseteq V$ in a graph $G = (V, E)$ is said to be r -independent if it does not induce a clique of size r , that is, if $G[X]$ is K_r -free. In particular, a 2-independent set is simply an independent set. Note that the proof of Theorem 1 carries through to the following generalization concerning the Erdős-Gallai problem for string graphs.

Theorem 6. *Let s, q be positive integers with $s > q$. Every K_{2^s} -free string graph G on $n \geq 2^s$ vertices contains a 2^q -independent set of size at least*

$$\min \left(\left(\frac{c(s+1-q)}{\log n} \right)^{2s-2q} n, \left(\frac{c(s+1-q)}{2^s \log n} \right)^2 n \right),$$

where $c > 0$ is an absolute constant.

Proof. (Sketch) We follow the proof of Theorem 1, making minor modifications. The proof is by double induction on s and n , with the base cases $s = q$ or $n = 2^s$ being trivial. We let $\alpha = c' \left(\frac{s+1-q}{\log n} \right)^2$. As in the proof of Theorem 1, if G has at most αn^2 edges, we apply the string graph separator lemma (Lemma 1). We delete the separator, use the induction hypothesis on the resulting components, and take the union of the 2^q -independent sets in the components to get a 2^q -independent set of the desired size in G .

So, we may assume G has more than αn^2 edges. By Lemma 4, there is an integer $t \geq 2$ and disjoint vertex subsets $X_1, \dots, X_t$ such that X_i is complete to X_j for all $i \neq j$ and $|X_i| \geq 4c''\alpha n/t^2$ for $i = 1, \dots, t$, where $c'' > 0$ is an absolute constant. Losing a factor at most 2 in the number of sets X_i , we may assume that $t = 2^p$ for a positive integer $p < s$, which implies $|X_i| \geq c''\alpha n/t^2$. As G is K_{2^s} -free, at least one of the sets X_i induces a subgraph which is $K_{2^{s-p}}$ -free.

The proof now splits into two cases, depending on whether $s - p > q$ or not. If $s - p > q$, the rest of the proof goes through as in the proof of Theorem 1. If $s - p \leq q$, then X_i is the desired 2^q -independent set. Indeed, we have

$$|X_i| \geq c''\alpha n/t^2 \geq c''\alpha n/2^{2s} = c''c' \left(\frac{s+1-q}{2^s \log n} \right)^2 n \geq \left(\frac{c(s+1-q)}{2^s \log n} \right)^2 n,$$

for a sufficiently small absolute constant $c > 0$, as desired.

We complete the section by proving Theorem 4, which gives an upper bound on the number of edges of a r -quasiplanar graph with n vertices for $r = 2^s$ a perfect power of two.

Proof of Theorem 4. Let $s \geq 3$ be an integer and $r = 2^s$. We have to prove that every r -quasiplanar graph on $n \geq r$ vertices has at most $n(C \frac{\log n}{s})^{2s-4}$ edges, where C is an absolute constant.

For any r -quasiplanar graph $G = (V, E)$ on n vertices, delete a small disk around each vertex and consider the string graph whose vertex set consists of the (truncated) curves in E . As G is r -quasiplanar, the resulting string graph is K_{2^s} -free.

Applying Theorem 6 with $q = 2$, we obtain a subset $E' \subset E$ with

$$|E'| \geq |E| \left(\frac{c(s-1)}{\log |E|} \right)^{2s-4} \geq |E| \left(\frac{c(s-1)}{2 \log n} \right)^{2s-4},$$

for some absolute constant $c > 0$ such that $G' = (V, E')$ is 4-quasiplanar. According to Ackerman's result [1], every 4-quasiplanar graph on n vertices has at most a linear number of edges in n , that is, we have $|E'| \leq C'n$ for a suitable constant $C' > 0$. Putting these two bounds together, we get the desired upper bound

$$|E| \leq C'n \left(\frac{2 \log n}{c(s-1)} \right)^{2s-4} \leq n \left(C \frac{\log n}{s} \right)^{2s-4},$$

provided that C is sufficiently large. □

5 Concluding Remarks

A. A family of graphs $\mathcal{G}$ is said to be *hereditary* if for any $G \in \mathcal{G}$, all induced subgraphs of G also belong to $\mathcal{G}$. Obviously, the family of string graphs is hereditary.

The proof of Lemma 3 only uses the fact that there is a separator theorem for string graphs. A careful inspection of the proof shows that the same result holds, instead of string graphs, for any hereditary family of graphs $\mathcal{G}$ such that every $G = (V, E) \in \mathcal{G}$ has a separator of size $O(|E|^\alpha |V|^{1-2\alpha})$, for a suitable constant $\alpha = \alpha(\mathcal{G}) > 0$.

Similar techniques were used in [13–15, 25]. Our Lemma 3 enables us to simplify some of the proofs in these papers.

B. Circle graphs are intersection graphs of chords of a circle. Gyárfás [18] proved that every circle graph with clique number r has chromatic number at most $O(r^2 4^r)$. Kostochka [20], and Kostochka and Kratochvíl [21] improved this bound to $O(r^2 2^r)$ and $O(2^r)$ respectively. Recently, a breakthrough was made by Davies and McCarty [7], who obtained the upper bound $O(r^2)$. Shortly after this, Davies [8] further improved this bound to $O(r \log r)$, which is asymptotically best possible due to a construction of Kostochka [20]. By taking the union of the $r - 2$ largest color classes in a proper coloring with the minimum number of colors, Davies' result implies that every circle graph on n vertices with clique number r contains an induced subgraph on $\Omega(n/\log r)$ vertices that is K_{r-1} -free. We conjecture that this “naive” bound can be improved as follows.

Conjecture 2. Every K_r -free circle graph on n vertices contains an induced subgraph on $\Omega(n)$ vertices which is K_{r-1} -free.

Acknowledgement. We are grateful to Zach Hunter for carefully reading our manuscript and pointing out several mistakes.

References

1. Ackerman, E.: On the maximum number of edges in topological graphs with no four pairwise crossing edges. *Discrete Comput. Geom.* **41**, 365–375 (2009)
2. Ackerman, E.: Quasi-planar graphs. In: Hong, S.-H., Tokuyama, T. (eds.) *Beyond Planar Graphs*, pp. 31–45. Springer, Singapore (2020). https://doi.org/10.1007/978-981-15-6533-5_3
3. Ackerman, E., Tardos, G.: On the maximum number of edges in quasi-planar graphs. *J. Comb. Theory, Ser. A* **114**(3), 563–571 (2007)
4. Agarwal, P., Aronov, B., Pach, J., Pollack, R., Sharir, M.: Quasi-planar graphs have a linear number of edges. *Combinatorica* **17**, 1–9 (1997)
5. Brass, P., Moser, W.O.J., Pach, J.: *Research problems in discrete geometry*. Springer (2005)
6. Burling, J.: On coloring problems of families of polytopes. (Ph.D. thesis), University of Colorado, Boulder 1(5) (1965)
7. Davies, J., McCarty, R.: Circle graphs are quadratically χ -bounded. *Bull. London Math. Soc.* **53**, 673–679 (2021)
8. Davis, J.: Improved bounds for colouring circle graphs. Preprint, [arXiv:2107.03585](https://arxiv.org/abs/2107.03585) (2021)
9. Dudek, A., Retter, T., Rödl, V.: On generalized ramsey numbers of erdos and rogers. *J. Combin. Theory Ser. B* **109**, 213–227 (2014)
10. Dudek, A., Rödl, V.: On k_s -free subgraphs in k_{s+k} -free graphs and vertex folkman numbers. *Combinatorica* **31**, 39–53 (2011)
11. Erdős, P., Gallai, T.: On the minimal number of vertices representing the edges of a graph. *Magyar Tud. Akad. Mat. Kutató Int. Közl.* **6**, 181–203 (1961)
12. Erdős, P., Rogers, C.A.: The construction of certain graphs. *Canad. J. Math.* **14**, 702–707 (1962)
13. Fox, J., Pach, J.: Separator theorems and turán-type results for planar intersection graphs. *Adv. Math.* **219**, 1070–1080 (2008)
14. Fox, J., Pach, J.: A separator theorem for string graphs and its applications. *Combin. Probab. Comput.* **19**, 371–390 (2010)
15. Fox, J., Pach, J.: Coloring k_k -free intersection graphs of geometric objects in the plane. *European J. Combin.* **33**, 853–866 (2012)
16. Fox, J., Pach, J.: String graphs and incomparability graphs. *Adv. Math.* **230**, 1381–1401 (2012)
17. Fox, J., Pach, J.: Applications of a new separator theorem for string graphs. *Combin. Probab. Comput.* **23**, 66–74 (2014)
18. Gyárfás, A.: On the chromatic number of multiple interval graphs and overlap graphs. *Discrete Math.* **55**, 161–166 (1985)
19. Janzer, O., Gowers, W.T.: Improved bounds for the erdos-rogers function. *Adv. Comb.* **3**, 1–27 (2020)
20. Kostochka, A.: O verkhnikh otsenkakh khromaticheskogo chisla grafov (on upper bounds for the chromatic number of graphs). In: Dement'ev, V.T. (ed.) *Modeli i metody optimizacii*, Akad. Nauk SSSR SO 10, pp. 204–226 (1988)

21. Kostochka, A., Kratochvíl, J.: Covering and coloring polygon-circle graphs. *Discrete Math.* **163**, 299–305 (1997)
22. Krivelevich, M.: k_s -free graphs without large k_r -free subgraphs. *Combin. Probab. Comput.* **3**, 349–354 (1994)
23. Krivelevich, M.: Bounding ramsey numbers through large deviation inequalities. *Random Struct. Algorithms* **7**, 145–155 (1995)
24. Lee, J.: Separators in region intersection graphs. *ITCS* **7**, 1:1–1:8 (2017). Full version: [arXiv:1608.01612](https://arxiv.org/abs/1608.01612)
25. Lipton, R., Rose, D., Tarjan, R.: Generalized nested dissections. *SIAM J. Numer. Anal.* **16**, 346–358 (1979)
26. McGuinness, S.: Colouring arcwise connected sets in the plane, i. *Graphs Combin.* **16**, 429–439 (2000)
27. Pach, J., Radoičić, R., Tóth, G.: Relaxing planarity for topological graphs. In: Akiyama, J., Kano, M. (eds.) *JCDCG 2002. LNCS*, vol. 2866, pp. 221–232. Springer, Heidelberg (2003). https://doi.org/10.1007/978-3-540-44400-8_24
28. Pawlik, A., Kozik, J., Krawczyk, T., Lasoń, M., Micek, P., Trotter, W., Walczak, B.: Triangle-free intersection graphs of line segments with large chromatic number. *J. Combin. Theory Ser. B* **105**, 6–10 (2014)
29. Rok, A., Walczak, B.: Coloring curves that cross a fixed curve. *Discrete Comput. Geom.* **61**, 830–851 (2019)
30. Rok, A., Walczak, B.: Outerstring graphs are χ -bounded. *SIAM J. Discrete Math.* **13**, 2181–2199 (2019)
31. Sudakov, B.: Large k_r -free subgraphs in k_s -free graphs and some other ramsey-type problems. *Random Struct. Algorithms* **26**, 253–265 (2005)
32. Suk, A.: Coloring intersection graphs of x -monotone curves in the plane. *Combinatorica* **34**, 487–505 (2014)
33. Tomon, I.: String graphs have the erdos-hajnal property. *arXiv preprint. arXiv:2002.10350* (2020)
34. Valtr, P.: On geometric graphs with no k pairwise parallel edges. *Discrete Comput. Geom.* **19**, 461–469 (1998)
35. Walczak, B.: Triangle-free geometric intersection graphs with no large independent sets. *Discrete Comput. Geom.* **53**, 221–225 (2015)
36. Wolfowitz, G.: k_4 -free graphs without large induced triangle-free subgraphs. *Combinatorica* **33**, 623–631 (2013)