

The Critical Locus and Rigidity of Foliations of Complex Hénon Maps

Misha Lyubich and John W. Robertson

Abstract

We study Hénon maps which are perturbations of a hyperbolic polynomial p with connected Julia set. We give a complete description of the critical locus of these maps. In particular, we show that for each critical point c of p , there is a primary component of the critical locus asymptotic to the line $y = c$. Moreover, primary components are conformally equivalent to the punctured disk, and their orbits cover the whole critical set. We also describe the holonomy maps from such a component to itself along the leaves of two natural foliations. Finally, we show that a quadratic Hénon map taken along with the natural pair of foliations, is a rigid object, in the sense that a conjugacy between two such maps respecting the foliations is a holomorphic or antiholomorphic affine map.

Contents

Introduction	3
I Foliations.	5
1 The Foliations near Degeneracy	5
1.1 The Foliations.	5
1.2 Degeneration of $\varphi_{a,-}$	6
1.3 Degeneration of $\varphi_{a,+}$	9
1.4 The Functions $\mathcal{G}_a^+$ and $\mathcal{G}_a^-$	11
II The Critical Locus.	14
2 The Critical Locus near Infinity.	14
2.1 The Foliations and the Critical Locus.	14
2.2 The Critical Locus near Infinity.	16

CONTENTS

3	Stable and Unstable Manifolds	20
3.1	Crossed Mappings	20
3.2	Recalling Context and Constructions.	21
3.3	The Stable and Unstable Manifolds as a Varies	23
4	Components of the Critical Locus.	28
4.1	Trapping and Mapping Components.	28
4.2	Classification of the Critical Components.	31
III	Rigidity	37
5	Holonomy in the Critical Locus.	37
6	Fiber Preserving Conjugacies.	40
6.1	Statement of the Theorem	40
6.2	Holomorphic circle actions	42
6.3	Rigidity Results	44
	List of Notations	49

Preamble

This paper was written in 2005–2006, but has never appeared even as a preprint. Meanwhile, the results have been developed further and have found some applications, see [Fir12, Tan16, FL17]. We are grateful to Tanya Firsova for insisting that this paper should be made available and for helping with the proof reading.

Introduction

The family of Hénon maps are a basic example of nonlinear dynamics. Both the real and the holomorphic versions of these maps have been studied extensively, and yet there is still a great deal that is not well understood about them. Some of the sources of fundamental results about Hénon maps are [FM89], [FS92], [HOV94], [HOV95], [BS91a], [BS91b], [BS92], [BLS93], [BS98a], [BS98b], and [BS99].

In this article we study holomorphic Hénon maps of $\mathbb{C}^2$. These are maps of the form

$$f_a \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} p(x) - ay \\ x \end{pmatrix}$$

where p is a monic polynomial of degree $d > 1$. Hénon maps have constant Jacobian, and the parameter a is the value of the Jacobian. In the degenerate case where $a = 0$ the map reduces to $f_0 \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} p(x) \\ x \end{pmatrix}$ and we see that the Hénon map degenerates to the polynomial map $p(x)$, acting on the copy of the complex plane given by the curve $x = p(y)$.

The Hénon maps we study here are perturbations of hyperbolic polynomial maps with connected Julia set. The Julia sets and *natural foliations* of these maps was described in great detail by Hubbard and Oberste-Vorth in [HOV94] and [HOV95]. In this paper we will describe the *tangency locus* between the natural foliations and will derive from it that the Hénon map endowed with the pair of foliations is a *rigid* object.

Let us now outline the content of the paper in more detail.

Throughout Section 1 we will recap results of Hubbard & Oberste-Vorth [HOV94] keeping careful track of what happens as the Jacobian of the Hénon map goes to zero. When the Jacobian is equal to zero, the map degenerates, but the foliations and plurisubharmonic functions associated to the map persist, and become easy to analyze.

In Section 2 we present basic facts about the critical locus and, by direct calculation, obtain a description of the tangent spaces to its primary components at infinity.

In Section 3 we recall relevant material from [HOV95] concerning the stable and unstable foliations and describe the critical locus when the Jacobian is zero.

In Section 4 we construct tubes that trap the components of the critical locus as the Jacobian varies away from zero. This allows us to prove that the primary horizontal components of the critical locus are punctured disks. We then show that every component of the critical locus is an iterate of a primary component.

In Section 5 we describe the holonomy maps on a primary horizontal component of the critical locus along the natural foliations.

The pair of natural foliations of a Hénon map can be thought of as giving natural coordinates near infinity. In Section 6 we prove that if a conjugacy between two Hénon maps in question respects these foliations then it is forced to be holomorphic or antiholomorphic near infinity. For degree two maps, this implies that it is actually affine, which is our main rigidity result.

A list of notations is provided as a reference at the end of the paper. These notations are used with the following convention.

Convention 0.1. *When we extend a certain object from $\mathbb{C}^2$ to $\mathbb{P}^1 \times \mathbb{P}^1$ we add a hat to the symbol to distinguish it, unless there is no chance for confusion. When referring to a set with the subset $\{a = 0\}$ removed we append the symbol $*$ as a superscript to the symbol denoting that set.*

Part I

Foliations.

1 The Foliations near Degeneracy

1.1 The Foliations.

In this section we define the foliations associated to a Hénon map. These foliations are not new, they were introduced and studied in [HOV94]. We give a careful development of them from scratch, following the same methods as [HOV94], in order to study what happens in the degenerate case, and to have a good handle on these foliations as the Jacobian a is allowed to vary.

In studying Hénon maps it is common to define domains V_+ and V_- such that $f_a(V_+) \subset V_+$ and $f_a^{-1}(V_-) \subset V_-$ and such that every point that has unbounded forward orbit eventually enters V_+ and every point with unbounded backward orbit eventually enters V_- . We will give precise definitions of these domains shortly.

We first recap the construction of the functions $\varphi_{a,+}: V_+ \rightarrow \mathbb{C}$ and $\varphi_{a,-}: V_- \rightarrow \mathbb{C}$, both of which are holomorphic for a in some disk such that $\varphi_{a,+} \circ f_a = \varphi_{a,+}^d$ and $\varphi_{a,+}(x, y) \sim x$ for $|x| > |y|$ as $|x| \rightarrow \infty$ and $\varphi_{a,-} \circ f_a^{-1} = \varphi_{a,-}^d/a$ holds¹ for $a \neq 0$ and $\varphi_{a,-}(x, y) \sim y$ for $|y| > |x|$ as $|y| \rightarrow \infty$.

Throughout this paper it will be convenient to consider the highest term of $p(x)$ separately, thus we write $p(x) = x^d + q(x)$ where $d \geq 2$ and $\deg q(x) < d$. We also let $d' = \deg q(x)$.

We want to construct domains V_+ and V_- where functions $\varphi_{a,+}$ and $\varphi_{a,-}$ are defined for all a in the disk $\mathbb{D}_R$ of radius R . We will need to control convergence of an infinite product to construct $\varphi_{a,+}$ and $\varphi_{a,-}$ and will choose a value r which will control the rate of convergence of this series.

Fix values $0 < r < 1$ and $R > 0$, and choose $\alpha > 0$ such that

- $\left| \frac{q(y)}{y^d} \right| + \frac{R+1}{|y^{d-1}|} < r,$
- $|p(y)| > (2R+1)|y|$

whenever $|y| \geq \alpha$.

We then define the domains V_+ and V_- to be given by

$$V_+ \equiv \{(x, y) \mid |x| > |y| \text{ and } |x| > \alpha\},$$

$$V_- \equiv \{(x, y) \mid |y| > |x| \text{ and } |y| > \alpha\},$$

We let $\begin{pmatrix} x_n \\ y_n \end{pmatrix} = f_a^{\circ n} \begin{pmatrix} x \\ y \end{pmatrix}$ for $n \in \mathbb{Z}$ so x_n and y_n are polynomial functions in x, y and $\frac{1}{a}$ for $n < 0$ and x_n and y_n are polynomial functions in x, y and a for $n > 0$.

¹The definition of φ_- given in [HOV94] has an inconsistency that is trivial to correct, but is essential to our calculations (specifically the conditions $\varphi_{a,-} \sim y$ and $\varphi_{a,-} \circ f^{-1} = \varphi_{a,-}^d$ are incompatible).

Lemma 1.1. *Let $a \in \mathbb{D}_R$. If $(x, y) \in V_-$ then $|x_{-1}| = |y| > |x|$ and $|y_{-1}| > 2|y| = 2|x_{-1}|$. Thus $f_a^{-1}(V_-) \subset V_-$.*

Proof. The statement about x_{-1} is obvious. For y_{-1} we have

$$|y_{-1}| = \left| \frac{p(y) - x}{a} \right| \geq \frac{|p(y)| - |x|}{|a|} \geq \frac{2R|y|}{|a|} > 2|y|.$$

□

Lemma 1.2. *Let $a \in \mathbb{D}_R$. If $(x, y) \in V_+$ then $|y_1| = |x| > |y|$ and $|x_1| > (R+1)|y_1| = (R+1)|x|$. Thus $f_a(V_+) \subset V_+$.*

Proof. The statement about y_1 is obvious. For x_1 we have

$$|x_1| \geq |p(x)| - |ay| \geq |p(x)| - R|x| \geq (R+1)|x| = (R+1)|y_1|.$$

□

1.2 Degeneration of $\varphi_{a,-}$.

We are chiefly interested in the degenerate case ($a = 0$) and perturbations of this case (a small). We start by working out the degree of x_{-k} and y_{-k} as polynomials in y and in $\frac{1}{a}$. In doing so it will be convenient to make the definition $\sigma_k = 1 + d + d^2 + \cdots + d^{k-1}$ for $k \geq 1$ and $\sigma_k = 0$ for $k \leq 0$.

By an easy induction we obtain

Lemma 1.3. *Given that $k \geq 1$ then the leading term of $y_{-k}(x, y, a)$ is*

$$\frac{1}{a^{\sigma_k}} (p(y) - x)^{d^{k-1}}$$

if y_{-k} is considered as a polynomial in $1/a$. The leading term of $y_{-k}(x, y, a)$ considered as a polynomial in y is just the term y^{d^k}/a^{σ_k} of it's leading term in $\frac{1}{a}$. Since $x_{-k}(x, y, a) = y_{-(k-1)}(x, y, a)$ this also gives us the leading terms of $x_{-k}(x, y, a)$ in $\frac{1}{a}$ and in y except that $x_{-1}(x, y, a) = y$.

The function $\varphi_{a,-}$ is constructed as a limit

$$\varphi_{a,-} = \lim_{n \rightarrow \infty} (y_{-n} \cdot a^{\sigma_n})^{\frac{1}{d^n}},$$

with an appropriate choice of the branch of the root. We are interested in this as a function of $V_- \times \mathbb{D}_R$ where $(x, y) \in V_-$ and $a \in \mathbb{D}_R$. Sense is made of the above limit using the telescoping formula

$$\varphi_{a,-}(x, y) = \lim_{n \rightarrow \infty} y \cdot \exp \left(\frac{1}{d} \log \frac{ay_{-1}}{y^d} + \frac{1}{d^2} \log \frac{ay_{-2}}{y^{d^2}} + \cdots \right) \quad (1.1)$$

We are most interested in this about the point $a = 0$ (where the map f_a degenerates and $\varphi_{a,-}$ can no longer be defined using its relationship with f_a).

We note that

$$\frac{ay_{-k}}{y_{-(k-1)}^d} = \frac{p(y_{-(k-1)}) - x_{-(k-1)}}{y_{-(k-1)}^d} = 1 + \frac{q(y_{-(k-1)}) - x_{-(k-1)}}{y_{-(k-1)}^d}.$$

Let

$$s_k^-(x, y, a) = \frac{q(y_{-(k-1)}) - x_{-(k-1)}}{y_{-(k-1)}^d}. \quad (1.2)$$

Lemma 1.4. $|s_k^-(x, y, a)| < r$ for $x, y \in V_-$, $a \in \mathbb{D}_R$, $k \geq 1$.

Proof. We have:

$$|s_k^-| \leq \frac{|q(y_{-(k-1)})| + |x_{-(k-1)}|}{|y_{-(k-1)}^d|} \leq \left| \frac{q(y_{-(k-1)})}{y_{-(k-1)}^d} \right| + \frac{1}{|y_{-(k-1)}^{d-1}|} < r$$

□

We evaluate $\log \frac{ay_{-k}}{y_{-(k-1)}^d} = \log(1 + s_k^-)$ using the principal branch of $\log$. By Lemma 1.4, $\log(1 + s_k^-) \leq -\log(1 - r)$. Hence the series

$$\frac{1}{d} \log \frac{ay_{-1}}{y^d} + \frac{1}{d^2} \log \frac{ay_{-2}}{y_{-1}^d} + \dots \quad (1.3)$$

converges uniformly and absolutely to a holomorphic function bounded by $\frac{-\log(1 - r)}{d - 1}$. Letting $B = (r - 1)^{-\frac{1}{d-1}}$, we conclude:

Corollary 1.5. $\varphi_{a,-}$ as defined by equation (1.1) is holomorphic as a function on $V_- \times \mathbb{D}_R$ with $(x, y) \in V_-$ and $a \in \mathbb{D}_R$. Moreover $B^{-1} < \left| \frac{\varphi_{a,-}}{y} \right| < B$.

It will be convenient at times to understand the behavior of $\varphi_{a,-}$ in a suitable compactification of $\mathbb{C}^2$. By the Riemann Extension Theorem (see e.g. [GR84] page 132),

Corollary 1.6. Let $\hat{V}_-$ denote the union of V_- and the line $y = \infty$ in $\mathbb{P}^1 \times \mathbb{P}^1$, with the point (∞, ∞) excluded. Then $\frac{\varphi_{a,-}}{y}$ extends holomorphically to $\hat{V}_- \times \mathbb{D}_R$ and the norm of the extension is bounded above by B and below by B^{-1} .

Later, when we study the extension of $\frac{\varphi_{a,-}}{y}$ described above it will be useful to understand the behavior of s_k^- near infinity. Lemma 1.3 implies:

Lemma 1.7. Letting $v = \frac{1}{y}$ then $s_k^-(x, y, a)$ vanishes to order at least d^{k-1} in v .

Corollary 1.8. $\varphi_{a,-}(x, y) \sim y$ as $|y| \rightarrow \infty$.

Proof. We know that $\frac{\varphi_{a,-}(x, y)}{y} = \exp\left(\frac{1}{d} \log(1 + s_1^-) + \frac{1}{d^2} \log(1 + s_2^-) + \dots\right)$. Since s_k^- vanishes in v for all k we see that this infinite sum is a holomorphic function in $V_- \times \mathbb{D}_R$ that vanishes in v . $\square$

Let us now study the behavior of $a \mapsto \varphi_{a,-}$ near $a = 0$.

Lemma 1.9. $s_k^-(x, y, a)$ vanishes in a precisely to order

- $(d - d') \cdot \sigma_{k-1}$ for q nonconstant;
- $d \cdot \sigma_{k-1} - \sigma_{k-2}$ for q constant.

Moreover, $s_k^-(x, 1/v, a)$ vanishes in v precisely to order

- $(d - d')d^{k-1}$ for q nonconstant.
- $d^k - d^{k-2}$ for q constant and $k \geq 2$,
- d for q constant and $k = 1$.

Proof. It follows from Lemma 1.3 that the denominator of s_k^- is a polynomial in $\frac{1}{a}$ of degree $d \cdot \sigma_{k-1}$.

The numerator of s_k^- is a polynomial in $\frac{1}{a}$ of degree

- $d' \cdot \sigma_{k-1}$ for q nonconstant;
- σ_{k-2} for q constant.

To justify that the highest degree terms in the numerator never cancel, observe that it is impossible for the degree of $q(y_{-(k-1)})$ to match the degree of $x_{-(k-1)}$ as polynomials in $1/a$ except when $k = 1$ or when q is constant and $k = 2$. It is easy to check that the Lemma still holds in these cases.

The last assertion also easily follows from Lemma 1.3. $\square$

Lemma 1.10. $\varphi_{a,-}(x, y) = (p(y) - x)^{\frac{1}{d}} + ah(x, y, a)$ for some holomorphic function h on $V_- \times \mathbb{D}_R$.

Proof. According to Lemma 1.9 all s_k^- , $k \geq 2$, vanish at $a = 0$ and hence the series (1.3) takes the form $\frac{1}{d} \log \frac{ay_1}{y^d} + ag(x, y, a)$ for some holomorphic function $g(x, y, a)$ on $V_- \times \mathbb{D}_R$. Hence by (1.1)

$$\varphi_{a,-}(x, y) = y \cdot \exp\left(\frac{1}{d} \log \frac{ay_1}{y^d}\right) \cdot \exp(ag(x, y, a)) = (p(y) - x)^{\frac{1}{d}} \exp(ag(x, y, a)),$$

and the conclusion follows. $\square$

The domain $f_a(V_-)$ swells as $a \rightarrow 0$ to include all of $\mathbb{C}^2$ except the curve $C(p)$. We make this precise:

Lemma 1.11. *Given $(x, y) \notin J_0^- = C(p)$ then $(x, y) \in f_a(V_-)$ for all sufficiently small values a . More generally if $K \subseteq \mathbb{C}^2 \setminus C(p)$ then $K \subset f_a(V_-)$ for all sufficiently small a .*

Proof. This follows because $(x, y) \in f_a(V_-)$ iff $|p(y) - x| \geq \alpha|a|$ and $|p(y) - x| \geq |y||a|$. $\square$

We let $\mathfrak{V}_{k,-}^\circ = \{(x, y, a) | (x, y) \in f_a^{\circ k}(V_-), a \in \mathbb{D}_R^*\}$ and we let $\mathfrak{V}_{k,-}$ be the union of $\mathfrak{V}_{k,-}^\circ$ and the set $\{(x, y, 0) | (x, y) \notin C(p)\}$.

Lemma 1.12. *Given $k \geq 1$ then $\varphi_{a,-}^{d^k}$ extends from a holomorphic function on V_- , to a holomorphic function on $\mathfrak{V}_{k,-}$ by defining $\varphi_{a,-}^{d^k}(x, y) \equiv a^{\sigma_k} \varphi_{a,-}(f_a^{-k}(x, y))$ for $a \neq 0$ and $\varphi_{0,-}^{d^k}(x, y) = (p(y) - x)^{d^{k-1}}$.*

Proof. If $a \neq 0$ then we can extend the function $\varphi_{a,-}^{d^k}$ to be holomorphic on $f_a^{\circ k}(V_-)$ by defining $\varphi_{a,-}^{d^k}(x, y) \equiv a^{\sigma_k} \varphi_{a,-}(f_a^{-k}(x, y))$. This agrees with $\varphi_{a,-}^{d^k}$ on V_- .

According to our definition and Lemma 1.10

$$\begin{aligned} \varphi_{a,-}^{d^k}(x, y) &= a^{\sigma_k} \varphi_{a,-}(x_{-k}, y_{-k}) \\ &= a^{\sigma_k} y_{-k} \cdot \left(1 + \frac{q(y_{-k}) - x_{-k}}{y_{-k}^d}\right)^{\frac{1}{d}} + a^{1+\sigma_k} h(x_{-k}, y_{-k}) \\ &= a^{\sigma_k} y_{-k} (1 + s_k^-(x, y))^{\frac{1}{d}} + a^{1+\sigma_k} h(x_{-k}, y_{-k}) \end{aligned} \tag{1.4}$$

for $(x, y, a) \in \mathfrak{V}_{k,-}^\circ$. Now $(x_{-k}, y_{-k}) \in V_-$ when $(x, y, a) \in \mathfrak{V}_{k,-}^\circ$ and so $y_{-k} \neq 0$. Also $s_k^-(x_{-k}, y_{-k})$ is defined and holomorphic since $y_{-k} \neq 0$. From Lemma 1.3 we see that $a^{\sigma_k} y_{-k}$ is a polynomial in x, y, a whose only term not divisible by a is $(p(y) - x)^{d^{k-1}}$. By Lemma 1.9 s_k^- vanishes in a and so $\varphi_{0,-}^{d^k}(x, y) \equiv (p(y) - x)^{d^{k-1}}$ gives a continuous, and therefore, a holomorphic, extension of $\varphi_{a,-}^{d^k}$ to $\mathfrak{V}_{k,-}$. $\square$

We let both J_0^- and K_0^- denote the curve $p(y) - x = 0$. It follows from the previous result that this is consistent with the convention that J_a^- and K_a^- will denote the sets J_- and K_- for the parameter value a .

1.3 Degeneration of $\varphi_{a,+}$

Here we include the corresponding constructions for forward iterates.

Lemma 1.13. *The leading term of $x_k(x, y, a)$ is x^{d^k} if x_k is considered as a polynomial in x . The leading term of $y_k(x, y, a)$ is $x^{d^{k-1}}$ as a polynomial in x .*

Proof. This follows from an easy induction using $\begin{pmatrix} x_1 \\ y_1 \end{pmatrix} = \begin{pmatrix} p(x) - ay \\ x \end{pmatrix}$ and $\begin{pmatrix} x_k \\ y_k \end{pmatrix} = \begin{pmatrix} p(x_{k-1}) - ay_{k-1} \\ x_{k-1} \end{pmatrix}$. $\square$

The function $\varphi_{a,+}$ is constructed as a limit $\varphi_{a,+} = \lim_{n \rightarrow \infty} x_n^{\frac{1}{d^n}}$ with an appropriate choice of root on the domain $V_+ \times \mathbb{D}_R$. Sense is made of this using the telescoping formula,

$$\varphi_{a,+}(x, y) = \lim_{n \rightarrow \infty} x \cdot \exp\left(\frac{1}{d} \log \frac{x_1}{x^d} + \frac{1}{d^2} \log \frac{x_2}{x_1^d} + \cdots\right) \quad (1.5)$$

Letting $s_k^+ = \frac{q(x_{k-1}) - ay_{k-1}}{x_{k-1}^d}$ so $\frac{x_k}{x_{k-1}^d} = 1 + s_k^+$ we see that

Lemma 1.14. $|s_k^+(x, y, a)| < r$ for $(x, y) \in V_+, a \in \mathbb{D}_R$.

Proof.

$$|s_k^+| < \frac{|q(x_{k-1})| + R|y_{k-1}|}{x_{k-1}^d} \leq \left| \frac{q(x_{k-1})}{x_{k-1}^d} \right| + \frac{R}{|x_{k-1}^{d-1}|} < r$$

□

We evaluate $\log \frac{x_k}{x_{k-1}^d} = \log(1 + s_k^+)$ using the principal branch of $\log$. Since $|s_k^+| < r$ then $|\log(1 + s_k^+)| < -\log(1 - r)$ exactly as before for s_k^- . It follows that the series

$$\frac{1}{d} \log \frac{x_1}{x^d} + \frac{1}{d^2} \log \frac{x_2}{x_1^d} + \frac{1}{d^3} \log \frac{x_3}{x_2^d} + \cdots \quad (1.6)$$

converges absolutely and uniformly.

Since $\left| \frac{1}{d^k} \log \frac{x_k}{x_{k-1}^d} \right| < \frac{-\log(1 - r)}{d^k}$ then the infinite sum (1.6) is no larger than $\log B = \frac{-\log(1 - r)}{d - 1}$. We conclude that

Corollary 1.15. *The function $\varphi_{a,+}$ defined by equation (1.5) is well defined and holomorphic for all $(x, y) \in V_+$ and all $a \in \mathbb{D}_R$. Additionally $B^{-1} < \left| \frac{\varphi_{a,+}}{x} \right| < B$.*

Proof. The final claim follows immediately from the expression (1.5) and the bounds just derived on the series (1.6). □

For the next lemma we consider $\mathbb{C}^2$ as lying in $\mathbb{P}^1 \times \mathbb{P}^1$.

Corollary 1.16. *Let $\hat{V}_+$ denote the union of V_- and the line $x = \infty$ in $\mathbb{P}^1 \times \mathbb{P}^1$, with the point (∞, ∞) excluded. Then $\frac{\varphi_{a,+}}{x}$ extends holomorphically to $\hat{V}_+ \times \mathbb{D}_R$ and $B^{-1} \leq \left| \frac{\varphi_{a,+}}{x} \right| \leq B$.*

Proof. This follows from the Riemann extension theorem (see e.g. [GR84] page 132). □

In order to better understand this extension we will need to understand s_k^+ on this extension. We extract the relevant information in the following lemma.

Lemma 1.17. *Letting $u = \frac{1}{x}$ then $s_k^+(x, y, a)$ vanishes to order at least d^{k-1} in u .*

Proof. Writing $s_k^+ = \frac{q(x_{k-1}(1/u, y, a)) - ay_{k-1}(1/u, y, a)}{x_{k-1}^d(1/u, y, a)}$ and multiplying the numerator

and denominator by u^{d^k} this follows from Lemma 1.13. □

Corollary 1.18. $\varphi_{a,+}(x, y, a) \sim x$ as $|x| \rightarrow \infty$.

Proof. We know that $\frac{\varphi_{a,+}(x, y)}{x} = \exp\left(\frac{1}{d} \log(1+s_1^+) + \frac{1}{d^2} \log(1+s_2^+) + \dots\right)$. Since s_k^+ vanishes in u for all k we see that this infinite sum is a holomorphic function on $V_+ \times \mathbb{D}_R$ that vanishes in u . $\square$

We now have an expression for $\varphi_{a,+}$ as x multiplied by the exponent of a uniformly convergent sum of functions which are holomorphic for $(x, y) \in V_+$ and $a \in \mathbb{D}_R$.

We let $\mathfrak{V}_{k,+} = \{(x, y, a) | f_a^{\circ k}(x, y) \in V_+, a \in \mathbb{D}_R\}$.

Lemma 1.19. For each $k \geq 0$ the function $\varphi_{a,+}^{d^k}$ extends to a holomorphic function on $\mathfrak{V}_{k,+}$ given by $\varphi_{a,+}^{d^k} \equiv \varphi_{a,+} \circ f_a^k$. The function $\varphi_{0,+}$ is defined and holomorphic on all of $\mathbb{C}^2 \setminus K_0^+$ and is the Böttcher coordinate $b_p(x)$ of $p(x)$.

Proof. Unlike the case of $\varphi_{a,-}$, there is no difficulty in case $a = 0$ here. It is immediate that $\varphi_{a,+}$ is holomorphic as a function on $V_+ \times \mathbb{D}_R$. It is clear from the definition that $\varphi_{0,+}$ is the Böttcher coordinate $b_p(x)$ of $p(x)$. The rest of the Lemma is obvious. $\square$

1.4 The Functions $\mathcal{G}_a^+$ and $\mathcal{G}_a^-$.

Definition 1.20. We follow the standard convention that $U_a^+ \equiv \mathbb{C}^2 \setminus K_a^+$ and $U_a^- \equiv \mathbb{C}^2 \setminus K_a^-$. We define $\tilde{U}^+ \equiv \{(x, y, a) | (x, y) \in U_a^+, a \in \mathbb{D}_R\}$ and $\tilde{U}^- \equiv \{(x, y, a) | (x, y) \in U_a^-, a \in \mathbb{D}_R\}$ with the convention that $U_0^- \equiv \mathbb{C}^2 \setminus C(p)$.

Lemma 1.21. The sets $\tilde{U}^+$ and $\tilde{U}^-$ are open in $\mathbb{C}^3$.

Proof. $\tilde{U}^+ = \bigcup_{n=0}^{\infty} f^{-n}(V_+)$ (even when $a = 0$) and therefore $\tilde{U}^+$ is open. The result follows for $\tilde{U}^-$ by a similar construction for $a \neq 0$ and by Lemma 1.11 for $a = 0$. $\square$

Another fact we will want is

Lemma 1.22. If p is hyperbolic then no point interior to K_0^+ lies in the closure of $\tilde{U}^+$.

Proof. This is a trivial consequence of the fact that any point interior to K_0^+ is attracted to an attracting cycle. $\square$

We recall² the Green's functions

$$\mathcal{G}_a^+(x, y) \equiv \begin{cases} \log |\varphi_{a,+}(x, y)| & (x, y) \in U_a^+ \\ 0 & (x, y) \in K_a^+ \end{cases}$$

and

$$\mathcal{G}_a^-(x, y) \equiv \begin{cases} \log |\varphi_{a,-}(x, y)| & (x, y) \in U_a^- \\ \frac{1}{d-1} \log |a| & (x, y) \in K_a^-. \end{cases}$$

²Again with the correction in $\mathcal{G}_a^-$, similar to the one made for $\varphi_{a,-}$. Notice that it makes the second Green function to be a *non-zero* constant on K_a^- .

We take the value of $\mathcal{G}_0^-(x, y)$ to be $-\infty$ if $(x, y) \in C(p)$, i.e. if $x = p(y)$. These satisfy

$$\mathcal{G}_a^+ \circ f_a = d \cdot \mathcal{G}_a^+$$

and

$$\mathcal{G}_a^- \circ f_a^{-1} = d \cdot \mathcal{G}_a^- - \log |a| \quad \text{for } a \neq 0.$$

This second relation is sometimes more conveniently written

$$\left(\mathcal{G}_a^- - \frac{1}{d-1} \log |a|\right) \circ f_a^{-1} = d \cdot \left(\mathcal{G}_a^- - \frac{1}{d-1} \log |a|\right).$$

Convention 1.23. We will sometimes write $\mathcal{G}^+(x, y, a)$ for $\mathcal{G}_a^+(x, y)$ and $\mathcal{G}^-(x, y, a)$ for $\mathcal{G}_a^-(x, y)$. This will be convenient, for instance, when postcomposing $\mathcal{G}^+$ with function whose output lies in $\mathbb{C}^2 \times \mathbb{D}_R$.

Hubbard & Oberste-Vorth proved that the Green's functions are continuous when f_a is nondegenerate and the same argument gives continuity in x, y and a for $\mathcal{G}_a^+$ when $a = 0$. We extend this to $\mathcal{G}_a^-$ when $a = 0$.

Theorem 1.24. The functions $\mathcal{G}_a^+(x, y)$ and $\mathcal{G}_a^-(x, y)$ are continuous in x, y and a for $a \in \mathbb{D}_R$.

Proof. This follows by the same argument as is used in [HOV94] except in the case of $\mathcal{G}^-$ when $a = 0$. For $(x', y') \notin C(p)$ the continuity of $\mathcal{G}_a^-$ at (x', y') and $a = 0$ follows from Lemma 1.12. For $(x', y') \in C(p)$ more work is required. If we restrict $\mathcal{G}_a^-$ to the slice $a = 0$ then we already have shown continuity, so we will assume for most of the rest of this proof that $a \neq 0$ (so f_a^{-1} is defined).

If $f^{-n}(x, y) \in V_-$ then $B^{-1} < \left| \frac{\varphi_{a,-}(x_{-n}, y_{-n})}{y_{-n}} \right| < B$ by Corollary 1.5. and so $B^{-1} < \left| \frac{\varphi_{a,-}^{d^n}(x, y)}{a^{\sigma_n} y_{-n}} \right| < B$. Applying log to the right hand inequality yields $\mathcal{G}_a^-(x, y) < \frac{1}{d^n} \log B + \frac{1}{d^n} \log |a^{\sigma_n} y_{-n}|$. Now $(x_{-n}, y_{-n}) = f_a^{-n}(x, y) = f_a^{-1}(x_{-(n-1)}, y_{-(n-1)}) = (y_{-(n-1)}, \frac{1}{a}(p(y_{-(n-1)}) - x_{-(n-1)}))$. Therefore $y_{-n} = \frac{1}{a}(p(y_{-(n-1)}) - y_{-(n-2)})$ for $n \geq 2$.

We let $z_{-n} \equiv a^{\sigma_n} y_{-n}$ from which it follows that

$$z_{-n} = p(z_{-(n-1)}, a^{\sigma_{n-1}}) - a^{\sigma_n - \sigma_{n-2} - 1} z_{-(n-2)} \quad \text{for } n \geq 2.$$

where $z_0 = y$ and $z_{-1} = p(y) - x$. Writing our bound for $\mathcal{G}_a^-(x, y)$ in terms of z_{-n} we get:

$$\mathcal{G}_a^-(x, y) < \frac{1}{d^n} \log B + \frac{1}{d^n} \log |z_{-n}| \tag{1.7}$$

if $f_a^{-n}(x, y) \in V_-$.

We use the convention that $p(x, y) \equiv y^d p(x/y)$. Now fix a constant $C > \frac{1}{2}$ to be greater than $(d+1)$ times the absolute value of the largest coefficient of $p(x, y)$. It is then evident that $|p(x, y)| \leq C \max\{|x|^d, |y|^d\}$. Thus from the recursion relation for z_{-n} we have

$$|z_{-n}| \leq \max\{2 \cdot |p(z_{-(n-1)}, a^{\sigma_{n-1}})|, 2 \cdot |a^{\sigma_n - \sigma_{n-2} - 1} z_{-(n-2)}|\}$$

$$\leq \max\{2C|z_{-(n-1)}|^d, 2C|a|^{d\cdot\sigma_{n-1}}, 2|a|^{d^{n-1}+d^{n-2}-1}|z_{-(n-2)}|\}.$$

Let ϵ_v be an arbitrary positive number satisfying $\epsilon_v < \frac{1}{2C}$. Given a point $(x', y') \in C(p)$ consider the neighborhood

$$\tilde{\mathcal{U}}(\epsilon_v) \equiv \left\{ (x, y, a) \mid |p(y) - x| < \epsilon_v^d, |a| < \min\{2C\epsilon_v^d, \frac{1}{2}\} \text{ and } |a|^{d-1}|y| < C\epsilon_v^d \right\}$$

of $(x', y', 0) \in \mathbb{C}^2 \times \mathbb{D}_R$. Let $\tilde{\mathcal{U}}^*(\epsilon_v) \equiv \tilde{\mathcal{U}}(\epsilon_v) \cap (\mathbb{C}^2 \times \mathbb{D}_R)$ It can be shown by induction that

$$|z_{-n}| < (2C)^{\sigma_n} \epsilon_v^{d^n} \quad \text{for } (x, y, a) \in \tilde{\mathcal{U}}^*(\epsilon_v) \text{ and } n \geq 1. \quad (1.8)$$

Combining equation (1.7) and equation (1.8) gives $\mathcal{G}_a^-(x, y) < \frac{1}{d^n} \log B + \frac{1}{d^n} \log \left| (2C)^{\sigma_n} \epsilon_v^{d^n} \right|$ if $(x, y, a) \in \tilde{\mathcal{U}}^*(\epsilon_v)$ and $f_a^{-n}(x, y) \in V_-$. If $(x, y) \notin J_a^-$ then $f_a^{-n}(x, y) \in V_-$ for all sufficiently large n so we can take the limit as $n \rightarrow \infty$ and conclude that $\mathcal{G}_a^-(x, y) < \log(|2C|^{\frac{1}{d-1}} \epsilon_v)$. On the other hand, if $(x, y) \in J_a^-$ then $\mathcal{G}_a^-(x, y) = \frac{1}{d-1} \log |a|$ by definition.

We conclude that given any $M > 0$ and any $(x', y') \in C(p)$ then there is some ϵ_v such that $\mathcal{G}_a^-(x, y) < -M$ when $(x, y, a) \in \tilde{\mathcal{U}}^*(\epsilon_v) \setminus J_a^-$, and some ϵ_a such that if $|a| < \epsilon_a$ and $(x, y) \in J_a^-$ then $\mathcal{G}_a^-(x, y) < -M$. Also there is an open subset of $a = 0$ about (x', y') on which the values of $\mathcal{G}_0^-$ are always smaller than $-M$. Combining these sets gives a neighborhood of $(x', y', 0)$ on which $\mathcal{G}_a^-(x, y) < -M$. Hence $\mathcal{G}_a^-$ is continuous in x, y and a . $\square$

Part II

The Critical Locus.

2 The Critical Locus near Infinity.

2.1 The Foliations and the Critical Locus.

The fibers of $\varphi_{a,+}$ and $\varphi_{a,-}$ form holomorphic foliations of V_+ and V_- respectively. These foliations naturally extend to much larger sets using dynamics.

Lemma 2.1. *The holomorphic foliations defined by $\varphi_{a,+}$ on V_+ and $\varphi_{a,-}$ on V_- can be extended to all of $\tilde{U}^+$ and $\tilde{U}^-$ respectively. The resulting foliations, which we denote $\mathcal{F}_a^+$ and $\mathcal{F}_a^-$ respectively, are respected by the dynamics.*

Proof. This is a consequence of Lemma 1.12 and Lemma 1.19. $\square$

Notation. We will use $\mathcal{L}_a^+(z)$ to mean the entire leaf of $\mathcal{F}_a^+$ passing through z , and similarly for $\mathcal{L}_a^-(z)$. Given a set $B \subset \mathbb{C}^2$ and a point $z \in B$, we will use $\mathcal{L}_a^+(z, B)$ (resp. $\mathcal{L}_a^-(z, B)$) to denote the connected component of $\mathcal{L}_a^+(z) \cap B$ (resp. $\mathcal{L}_a^-(z) \cap B$) containing z .

Definition 2.2. Given any holomorphic foliation $\mathcal{F}$ on a two dimensional complex manifold M and a point $z \in M$ we will say that a holomorphic function g defined in a neighborhood N of z is a *local defining function* for $\mathcal{F}$ if for each leaf $\mathcal{L}$ of $\mathcal{F}$, g is constant on each connected component of $\mathcal{L} \cap N$ and if $\mathbf{d}g$ is never zero.

Given a pair of holomorphic foliations $\mathcal{F}_1$ and $\mathcal{F}_2$ on a two-dimensional complex manifold, the critical locus $\mathcal{C}$ of $\mathcal{F}_1$ and $\mathcal{F}_2$ is the complex variety given locally as the zero set (counting multiplicity) of the holomorphic function h satisfying $h\mathbf{d}x \wedge \mathbf{d}y = \mathbf{d}g_1 \wedge \mathbf{d}g_2$ for a pair of local defining functions g_1 and g_2 of $\mathcal{F}_1$ and $\mathcal{F}_2$ respectively. Equivalently $h = \frac{\partial g_1}{\partial x} \frac{\partial g_2}{\partial y} - \frac{\partial g_2}{\partial x} \frac{\partial g_1}{\partial y}$ and hence $\mathcal{C}$ is equal to the critical set (counting multiplicity of components) of the map $(x, y) \mapsto (g_1(x, y), g_2(x, y))$.

It is straightforward to check that altering the choice of local defining functions for the foliations only results in multiplying h by a nonvanishing holomorphic function. It is clear that the critical locus of a pair of foliations is exactly the set of points on which the foliations are tangent. However, some care must be taken, as it is possible that some components of the critical locus have multiplicity higher than one. An example is given by the foliations with local defining functions x and $x + x^2y$. The critical locus is defined by x^2 , which is the y axis, but with multiplicity two. Under the circumstances we are interested in we will be able to verify that every component of the critical locus has multiplicity one. Until that is done we will need to take into account multiplicity of components when dealing with the critical locus.

Definition 2.3. Let $\mathcal{C}_a$ be the critical locus of the foliations $\mathcal{F}_a^+$ and $\mathcal{F}_a^-$. It is easy to confirm that $\mathcal{C}_a$ is a closed analytic subvariety of $U_a^+ \cap U_a^-$ invariant under f_a .

Observe that since we can define $\varphi_{a,+}$ on all of U_a^+ up to local choice of a root then we can define $\log \varphi_{a,+}$ on all of U_a^+ up to local addition of a constant, and therefore $\mathbf{d} \log \varphi_{a,+}$ is a global holomorphic form on U_a^+ . Similarly for $\mathbf{d} \log \varphi_{a,-}$ on U_a^- . If we fix a and consider $\mathbf{d} \log \varphi_{a,+} \wedge \mathbf{d} \log \varphi_{a,-} = w_a(x, y) \mathbf{d}x \wedge \mathbf{d}y$ we get the same result as if we compute $\mathbf{d}$ with a nonconstant obtaining

$$\mathbf{d} \log \varphi_+ \wedge \mathbf{d} \log \varphi_- \wedge \mathbf{d}a = \tilde{w}(x, y, a) \mathbf{d}x \wedge \mathbf{d}y \wedge \mathbf{d}a$$

in the sense that $\tilde{w}(x, y, a) = w_a(x, y)$. Since $\varphi_{a,+}$ and $\varphi_{a,-}$ do not vanish in $\mathbb{C}^2$ then $\mathbf{d} \log \varphi_{a,+} \wedge \mathbf{d} \log \varphi_{a,-} \wedge \mathbf{d}a$ and any local branch³ of $\mathbf{d}\varphi_+^{-1} \wedge \mathbf{d}\varphi_-^{-d} \wedge \mathbf{d}a$ are just multiples of each other by a nonvanishing holomorphic function. Thus on the common domain of definition $\tilde{w}$ and w_a are multiples of each other by a nonvanishing holomorphic function.

Definition 2.4. The variety defined by $\tilde{w}$ in $\tilde{U}^+ \cap \tilde{U}^-$ will also be called the critical locus and will be denoted by $\mathcal{C}$. For each a it is exactly the locus along which the foliations defined by $\varphi_{a,+}$ and $\varphi_{a,-}$ are tangent, and is equal to $\mathcal{C}_a$.

Lemma 2.5. Assume X is a one dimensional complex variety in $\mathbb{D}^2$ and Y is a smooth holomorphic curve in $\mathbb{D}^2$. Assume Y is not contained in X and that z is an intersection point of X and Y . Let $r: \mathbb{D} \rightarrow Y$ be a local parameterization of Y about z with $r(0) = z$ and let h be a local defining function for X about z . Then the order of contact, or intersection multiplicity, of X and Y at z (counting multiplicity of the components of Y at z) is the order of the zero of $h \circ r$ at $0 \in \mathbb{D}$.

Proof. Since everything is local then without loss of generality we can assume that Y is the y axis and z is the origin. Then the intersection multiplicity is

$$(X, Y) = \dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}^2, (0,0)} / (x, h(x, y)) = \dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}, 0} / (h(x, y)) =$$

$$\dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}, 0} / (h(0, y)) = \dim_{\mathbb{C}} \mathbb{C}[y] / (y^k) = k$$

where $h(0, y) = a_k y^k + a_{k+1} y^{k+1} + \dots$, $a_k \neq 0$. This is easily seen to be the order of vanishing of $h \circ r$ at zero. $\square$

Lemma 2.6. If the critical locus of a pair of holomorphic foliations in $\mathbb{D}^2$ has a singularity at a point z then the leaves of the foliations must have order of contact greater than two at z .

Proof. We can assume without loss of generality that z is the origin, that one foliation is the vertical complex lines, and that the other foliation has local defining function h in a neighborhood of the origin.

One calculates that critical locus is defined by $\frac{\partial h}{\partial y} = 0$. If either $\frac{\partial^2 h}{\partial y^2} \big|_0 \neq 0$ or $\frac{\partial^2 h}{\partial x \partial y} \big|_0 \neq 0$ then $\frac{\partial h}{\partial y} = 0$ defines a smooth variety at zero. Hence if the critical locus of the pair of foliations has a singularity at zero then $\frac{\partial^2 h}{\partial y^2} \big|_0 = 0$, and so by lemma 2.5 the leaves of the two foliations have contact of order at least three at zero. $\square$

³We use φ_+^{-1} and φ_-^{-d} because these are single valued and extend holomorphically about $(\infty, c) \in \mathbb{P}^1 \times \mathbb{P}^1$. See Section 2.2.

Lemma 2.7. *Given two holomorphic foliations $\mathcal{F}_1$ and $\mathcal{F}_2$ of a two dimensional complex manifold and their critical locus $\mathcal{C}$ then:*

1. *Given any point $z \in \mathcal{C}$, the order of contact of the leaves of $\mathcal{F}_1$ and $\mathcal{F}_2$ at z is one more than the order of contact of the leaf of either foliation with $\mathcal{C}$ at z (counting the multiplicity of components of $\mathcal{C}$).*
2. *The subset $\mathcal{K}_k \subset \mathcal{C}$ of points of $\mathcal{C}$ where the foliations have contact of order at least k is an analytic subset of $\mathcal{C}$.*

Proof. We will confirm both properties locally. Without loss of generality we can assume that $\mathcal{F}_1$ is the foliation of $\mathbb{D}^2$ by vertical complex lines and $\mathcal{F}_2$ is defined by some local defining function $h: \mathbb{D}^2 \rightarrow \mathbb{C}$. Assume $z = (z_1, z_2) \in \mathbb{D}^2$ and parameterize the leaf of $\mathcal{F}_1$ through z by $y \mapsto (z_1, y)$. Then by lemma 2.5 the order of contact between the leaves through z is the order of vanishing of $h(z_1, y) - h(z_1, z_2)$ in y at $y = z_2$. This is precisely the smallest $k \in \mathbb{N}$ such that $\frac{\partial^j h}{\partial y^j} \big|_{(z_1, z_2)} = 0$ for $j = 1, \dots, k-1$. However, since $\mathcal{C}$ is defined by $\frac{\partial h}{\partial y}$ we also conclude by lemma 2.5 that the order of contact of the vertical leaf through z with $\mathcal{C}$ at z is the vanishing order of $\frac{\partial h}{\partial y}(z_1, y)$ in y at $y = z_2$ which is exactly one less than the vanishing order of $h(z_1, y) - h(z_1, z_2)$ in y at $y = z_2$. This completes the proof of part 1.

We also conclude that for each integer ℓ , the set $\mathcal{K}_\ell \subset \mathcal{C}$ is the common zero set of $\frac{\partial^j h}{\partial y^j}$ for $j = 1, \dots, \ell-1$. This completes the proof of part 2. $\square$

Corollary 2.8. *Assume we are given two holomorphic foliations $\mathcal{F}_1$ and $\mathcal{F}_2$ defined on some complex two dimensional manifold. Let $\mathcal{C}$ be the critical locus. If the leaves of $\mathcal{F}_1$ and $\mathcal{F}_2$ have order of contact two at every point of some component X of $\mathcal{C}$ then X is smooth, X meets no other component of $\mathcal{C}$, X is a component of $\mathcal{C}$ with multiplicity one, and X is everywhere transverse to both foliations.*

Proof. Since $\mathcal{C}$ is smooth everywhere the leaves have order of contact two then X is smooth and meets no other component of $\mathcal{C}$ by lemma 2.6. It follows by part 1 of Lemma 2.7 that X must meet each leaf with order of contact one, i.e. X must be transverse to each leaf, and X must have multiplicity one as a component of $\mathcal{C}$. $\square$

2.2 The Critical Locus near Infinity.

We consider the critical locus for the extension of f_a to $\mathbb{P}^1 \times \mathbb{P}^1$. The map f_a is well defined except at (∞, ∞) . The map f_a^{-1} is well defined except at (∞, ∞) or when $a = 0$ and $(x, y) \in C(p)$.

The map f_a sends the line $y = \infty$ to the line $x = \infty$ and sends the line $x = \infty$ in turn to the point (∞, ∞) at which f_a is undefined. Similarly, the map f_a^{-1} sends the line $x = \infty$ to the line $y = \infty$ and sends the line $y = \infty$ in turn to the point (∞, ∞) at which f_a^{-1} is undefined.

In this section we will show that for each critical point c of the polynomial p , and for each $a \in \mathbb{D}_R$ the critical locus has a branch asymptotic to the curve $y = c$ as $|x| \rightarrow \infty$. We

will do this by showing that the critical locus (extended to $\mathbb{P}^1 \times \mathbb{P}^1$) contains the point (∞, c) and by computing the tangent line to the critical locus at (∞, c) .

Because we will be working in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{D}_R$ it will be convenient at times to use the coordinate systems (u, y, a) where $u = 1/x$ or (x, v, a) where $v = 1/y$ instead of using (x, y, a) . Corollary 1.16, written using (u, y, a) coordinates, states that

$$h^+(u, y, a) \equiv \frac{1}{u\varphi_{a,+}(u, y, a)} \quad (2.1)$$

is holomorphic on $\hat{V}_+ \times \mathbb{D}_R$ and $B^{-1} \leq |h^+| \leq B$. Similarly Corollary 1.6, written using (x, v, a) coordinates, states that

$$h^-(x, v, a) \equiv \frac{1}{v\varphi_{a,-}(x, v, a)} \quad (2.2)$$

is holomorphic on $\hat{V}_- \times \mathbb{D}_R$ and $B^{-1} \leq |h^-| \leq B$. We find it useful here to write f_a^{-1} with input written in the (u, y, a) coordinates, but the output is written in the coordinates (x, v, a) , i.e. $f_a^{-1}(u, y, a) = (y, \frac{au}{up(y) - 1}, a)$.

Let $\hat{\mathfrak{V}}_-^* = \{(x, y, a) | (x, y) \in f_a(\hat{V}_-), a \in \mathbb{D}_R^*\}$ and let

$$\hat{\mathfrak{V}}_- = \hat{\mathfrak{V}}_-^* \cup \left((U_0^- \cup (\{\infty\} \times \mathbb{C})) \times \{0\} \right) \subset \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{D}_R.$$

The set $\hat{\mathfrak{V}}_-$ is just the interior of the closure of $\hat{\mathfrak{V}}_-^*$ in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{D}_R$. Then from the recursion relation it follows that $\varphi_{a,-}^{-d}(u, y, a) = \frac{u}{up(y) - 1} h^-(y, \frac{au}{up(y) - 1}, a)$ on $\hat{\mathfrak{V}}_-^*$. Since both sides of the equality are holomorphic on $\hat{\mathfrak{V}}_-$ they are equal on $\hat{\mathfrak{V}}_-$.

Therefore

$$\mathbf{d}\varphi_+^{-1}(u, y, a) = \left(h^+ + u \frac{\partial h^+}{\partial u} \right) \mathbf{d}u + u \frac{\partial h^+}{\partial y} \mathbf{d}y + u \frac{\partial h^+}{\partial a} \mathbf{d}a \quad \text{on } \hat{V}_+ \times \mathbb{D}_R. \quad (2.3)$$

Also, letting $\lambda = up(y) - 1$ we obtain,

$$\begin{aligned} \mathbf{d}\varphi_-^{-d}(u, y, a) &= -\lambda^{-2} \cdot h^- \circ f_a^{-1} \mathbf{d}u \\ &\quad - \lambda^{-2} u^2 p'(y) \cdot h^- \circ f_a^{-1} \mathbf{d}y + \lambda^{-1} u \cdot \frac{\partial h^-}{\partial x} \circ f_a^{-1} \mathbf{d}y \\ &\quad + \lambda^{-1} u \cdot \frac{\partial h^-}{\partial v} \circ f_a^{-1} \cdot \left(\lambda^{-1} u \mathbf{d}a - \lambda^{-2} a \mathbf{d}u - \lambda^{-2} a u^2 p'(y) \mathbf{d}y \right) \\ &\quad + \lambda^{-1} u \cdot \frac{\partial h^-}{\partial a} \circ f_a^{-1} \mathbf{d}a \quad \text{on } \hat{\mathfrak{V}}_-. \end{aligned} \quad (2.4)$$

One then calculates

$$\begin{aligned}
 \mathbf{d}\varphi_+^{-1} \wedge \mathbf{d}\varphi_-^{-d} \wedge \mathbf{d}a = & \\
 & \left((h^+ + u \frac{\partial h^+}{\partial u}) \left(-\lambda^{-2} u^2 p'(y) \cdot h^- \circ f_a^{-1} + u \lambda^{-1} \frac{\partial h^-}{\partial x} \circ f_a^{-1} - \lambda^{-3} a u^3 p'(y) \cdot \frac{\partial h^-}{\partial v} \circ f_a^{-1} \right) \right. \\
 & \left. + u \frac{\partial h^+}{\partial y} \left(\lambda^{-2} \cdot h^- \circ f_a^{-1} + a u \lambda^{-3} \cdot \frac{\partial h^-}{\partial v} \circ f_a^{-1} \right) \right) \mathbf{d}u \wedge \mathbf{d}y \wedge \mathbf{d}a \\
 & \text{on } \hat{\mathfrak{V}}_- \cap (\hat{V}_+ \times \mathbb{D}_R). \quad (2.5)
 \end{aligned}$$

It is easy to show that the domain $\hat{\mathfrak{V}}_- \cap (\hat{V}_+ \times \mathbb{D}_R)$ contains the plane $u = 0$ in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{D}_R$.

Corollary 2.9. *The Foliations $\mathcal{F}_a^+$ and $\mathcal{F}_a^-$ extend to holomorphic foliations on $\hat{V}_+$ and $\hat{\mathfrak{V}}_-$ respectively and $\varphi_{a,+}^{-1}$ and $\varphi_{a,-}^{-d}$ are local defining functions for these foliations. In both extended foliations the leaf through $u = 0, y = c$ is the line $u = 0$ regardless of the value of a .*

Proof. Clearly, for each $a \in \mathbb{D}_R$, $\varphi_{a,+}^{-1}$ and $\varphi_{a,-}^{-d}$ define the same foliations as $\varphi_{a,+}$ and $\varphi_{a,-}$ on V_+ and on $\hat{\mathfrak{V}}_- \cap (U_a^- \times \{a\})$. Since φ_+^{-1} and φ_-^{-d} are holomorphic functions on $\hat{V}_+ \times \mathbb{D}_R$ and $\hat{\mathfrak{V}}_-$ which both vanish on $u = 0$ by equations (2.1) and (2.2) then we need only confirm that $\mathbf{d}\varphi_{a,+}^{-1}$ and $\mathbf{d}\varphi_{a,-}^{-d}$ do not vanish on $u = 0$. But this follows from equations (2.3) and (2.4). $\square$

Lemma 2.10. *$h^+|_{u=0} = 1$, and $\frac{\partial h^+}{\partial y}$ vanishes in u to order at least d . Similarly, $h^-|_{v=0} = 1$, and $\frac{\partial h^-}{\partial x}$ vanishes in v to order at least d .*

Proof. From equation (2.1), equation (1.5) and the definition of s_k^+ we know

$$h^+ = (1 + s_1^+)^{-\frac{1}{d}} \exp\left(-\frac{1}{d^2} \log(1 + s_2^+) + \dots\right) \quad (2.6)$$

We know from Lemma 1.17 that if $k \geq 2$ then s_k^+ vanishes to order at least d in u .

Since $s_1^+ = \frac{q(x) - ay}{x^d} = u^d q\left(\frac{1}{u}\right) - ay u^d$ so $\frac{\partial s_1^+}{\partial y} = -a u^d$, it follows from equation (2.6) that $\frac{\partial h^+}{\partial y}$ vanishes to order at least d in u . The claim about $h^+|_{u=0}$ follows from equation 2.6.

The results for h^- are proven similarly. $\square$

Corollary 2.11. *The critical locus of the extensions of $\mathcal{F}_a^+$ and $\mathcal{F}_a^-$ in $(\hat{V}_+ \times \mathbb{D}_R) \cap \hat{\mathfrak{V}}_-$ contains the plane $u = 0$ with multiplicity two.*

Proof. Since the v coordinate of $f_a^{-1}(u, y, a)$ is $\frac{au}{up(y) - 1}$ one concludes from Lemma 2.10 that equation (2.5) is divisible by u^2 . $\square$

Corollary 2.11 is one of the reasons these calculations are done so carefully. If we had simply approximated the critical locus at infinity using a Taylor series we could not make this conclusion.

While we need to extend $\mathcal{C}$ to a variety on $(\hat{V}_+ \times \mathbb{D}_R) \cap \hat{\mathfrak{V}}_-$, the double component $u = 0$ is spurious for our purposes. We let $\tilde{w}$ be the holomorphic function satisfying $\mathbf{d}\varphi_+^{-1} \wedge \mathbf{d}\varphi_-^d \wedge \mathbf{d}a = u^2 \tilde{w}(u, y, a) \mathbf{d}u \wedge \mathbf{d}y \wedge \mathbf{d}a$. Then the zero set of $\tilde{w}$ gives an extension of $\mathcal{C}$ to $(\hat{V}_+ \times \mathbb{D}_R) \cap \hat{\mathfrak{V}}_-$. We will abuse notation and refer to the extended variety as $\mathcal{C}$ as well. It will be obvious from the context whether we are using the extension or not. Extending $\mathcal{C}$ automatically extends $\mathcal{C}_a$ to $(\hat{V}_+ \times \mathbb{D}_R) \cap \hat{\mathfrak{V}}_-$ for each $a \in \mathbb{D}_R$ since $\mathcal{C}_a$ is the zero set of $w_a(u, y) \equiv \tilde{w}(u, y, a)$.

Lemma 2.12. *Given $a \in \mathbb{D}_R$ and a point $(0, y_0, a_0)$ (written in (u, y, a) coordinates) the defining function for $\mathcal{C}$ takes the form $\tilde{w}(u, y, a) = -p'(y) - uH$ for some holomorphic function H defined in a neighborhood of $(0, y_0, a_0)$. Thus $(0, c, a)$ is in $\mathcal{C}$ iff c is a critical point of p . Moreover if c is an order one critical point of p then $\mathcal{C}$ is smooth at $(0, c, a)$ and the tangent plane to $(0, c, a)$ is given by*

$$p''(c)\mathbf{d}y + C\mathbf{d}u = 0$$

for some C depending upon c and a .

Proof. From Lemma 2.10 we can write: $h^- \circ f_a^{-1}(u, y, a) = h^-(y, \frac{au}{up(y) - 1}, a) = 1 + uH_2$, $\frac{\partial h^-}{\partial x} \circ f_a^{-1}(u, y, a) = \frac{\partial h^-}{\partial x}(y, \frac{au}{up(y) - 1}, a) = u^d H_4$ and $\frac{\partial h^+}{\partial y} = u^d H_6$ for holomorphic functions H_2, H_4 and H_6 . From equation (2.5) one obtains $u^2 \tilde{w}(u, y, a) = (1 + uH_1) \left(\{-u^2 p'(y)(1 + uH_2)\} + uH_3 \cdot u^d H_4 - u^3 H_5 \right) - u \cdot u^d H_6 \cdot H_7 = -u^2 p'(y) - u^3 H_8$ where each H_i is a function which is holomorphic in a neighborhood of the given $y_0, a \in \mathbb{D}_R$ and for u sufficiently close to zero. Simplifying gives $\tilde{w}(u, y, a) = -p'(y) - uH_8$ for some holomorphic H_8 . $\square$

Theorem 2.13. *If c is an order one critical point of p then the critical locus $\mathcal{C}_a$ passes through the point $(0, c, a)$ (written in (u, y, a) coordinates) and is smooth at this point. Moreover $\varphi_{a,+}^{-1} \big|_{\mathcal{C}_a} \rightarrow \mathbb{C}$ is a local biholomorphism about this point.*

Proof. Smoothness follows directly from Lemma 2.12. Combining equation (2.3) with Lemma 2.10 gives $\mathbf{d}\varphi_{a,+}^{-1} \big|_{\mathcal{C}_a}(0, c, a) = \mathbf{d}u$. By Lemma 2.12 this is nondegenerate on the tangent line of $\mathcal{C}_a$ at $(0, c, a)$. $\square$

3 Stable and Unstable Manifolds

3.1 Crossed Mappings

In this section we recall the definition of and basic results about crossed mappings from [HOV95]. This is a holomorphic version of a standard construction of stable and unstable manifolds such as Theorem 6.4.9 criteria (4) in [KH95]. We will only define degree one crossed mappings, and we will not consider any crossed mappings of degree higher than one in this paper.

Let $B_1 = U_1 \times V_1$ and $B_2 = U_2 \times V_2$ be bidisks.

Definition 3.1. A (degree one) *crossed mapping* from B_1 to B_2 is a triple (W_1, W_2, f) , where

1. W_1 is an open subset of $U'_1 \times V_1$ for some open $U'_1 \subseteq U_1$.
2. W_2 is an open subset of $U_2 \times V'_2$ for some open $V'_2 \subseteq V_2$.
3. $f: W_1 \rightarrow W_2$ is a holomorphic isomorphism, such that for all $y \in V_1$ the mapping

$$\text{pr}_1 \circ f: W_1 \cap (U_1 \times \{y\}) \rightarrow U_2$$

is a biholomorphism, and for all $x \in U_2$ the mapping

$$\text{pr}_2 \circ f^{-1}: W_2 \cap (\{x\} \times V_2) \rightarrow V_1$$

is a biholomorphism.

To make the notation less cumbersome, $f: B_1 \xrightarrow{\times} B_2$ is often written leaving the precise W_1 and W_2 to be determined by the context.

Given a hyperbolic polynomial map p , Hubbard & Oberste-Vorth construct a family of bidisks in $\mathbb{C}^2$ such that if $|a|$ is sufficiently small then f_a induces crossed mappings between the bidisks of the family. They use this to get good hold on the stable and unstable manifolds of J_a .

Proposition 3.2. *If $f: B_1 \xrightarrow{\times} B_2$ is a crossed mapping of degree one and $X \subset B_1$ is the graph of an analytic map from U_1 to V_1 then the image of X in B_2 is the graph of an analytic map from U_2 to V_2 .*

Proof. This is Proposition 3.4 of [HOV95] for degree one crossed mappings. $\square$

Proposition 3.3. (a) *Let $f: B_1 \xrightarrow{\times} B_2$ be a crossed mapping of degree 1. Then $f^{-1}: B_2 \xrightarrow{\times} B_1$ is also a crossed mapping if all the coordinates are flipped.*

(b) *If B_1, B_2 and B_3 are bidisks, $W_1 \subset B_1$, $W_2 \subset B_2$, $\tilde{W}_2 \subset B_2$ and $\tilde{W}_3 \subset B_3$ and $f_1: W_1 \rightarrow \tilde{W}_2$ and $f_2: W_2 \rightarrow \tilde{W}_3$ are degree one crossed mappings, then*

$$f_2 \circ f_1: f_1^{-1}(W_2) \rightarrow f_2(\tilde{W}_2)$$

is a degree one crossed mapping from B_1 to B_3 .

Proof. This is Proposition 3.7 of [HOV95]. $\square$

Suppose that $B_1, \dots, B_{n+1}$ are bidisks such that $B_i = U_i \times V_i$. Suppose also that $W_i \subset V_i$, ($i = 1, \dots, n$) and $\tilde{W}_i \subset B_i$ ($i = 2, \dots, n+1$) are open subsets so $f_i: W_i \rightarrow \tilde{W}_{i+1}$ are crossed mappings of degree 1. Let

$$S_1^n = W_1 \cap f_1^{-1}(W_2) \cap \dots \cap (f_1^{-1} \circ \dots \circ f_{n-1}^{-1})(W_{n-1})$$

and

$$S_2^n = \tilde{W}_{n+1} \cap f_n(\tilde{W}_n) \cap \dots \cap (f_n \circ \dots \circ f_2)(\tilde{W}_2)$$

so that the restriction g of $f_n \circ \dots \circ f_1$ to S_1^n makes $g: S_1^n \rightarrow S_2^n$ a crossed mapping of degree 1 from B_1 to B_{n+1} .

Let U be a disk, and U' a relatively compact open subset. Define the *size* of U' in U to be $\frac{1}{M}$ where M is the largest modulus of an annulus $U \setminus V$ for V a compact contractible set containing U' .

Definition 3.4. Given a crossed mapping $f: U_1 \times V_1 \xrightarrow{\times} U_2 \times V_2$ we will let the *horizontal size* of f be the size of $\text{pr}_1(W_1)$ in U_1 and we will let the *vertical size* of f be the size of $\text{pr}_2(W_2)$ in V_2 .

Proposition 3.5. Let $B_0 = U_0 \times V_0$, $B_1 = U_1 \times V_1$, $\dots$ be an infinite sequence of bidisks, and $f_i: B_i \xrightarrow{\times} B_{i+1}$ be crossed mappings of degree 1, with U'_i of uniformly bounded horizontal size in U_i . Then the set

$$\left\{ \begin{bmatrix} x \\ y \end{bmatrix} \in B_0 \mid f_n \circ \dots \circ f_0 \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) \in B_n \text{ for all } n \right\}$$

is an analytic disk in B_0 , which maps by pr_2 isomorphically to V_0 , which we will call the *stable disk* of the sequence of crossed mappings.

Proof. This is Corollary 3.12 of [HOV95]. $\square$

Similarly, when we have a backward sequence of crossed mappings

$$\dots \xrightarrow[\times]{} B_{-1} \xrightarrow[\times]{} B_0$$

with uniformly bounded vertical sizes, it will have an *unstable disk*, which maps by pr_1 isomorphically to U_0 .

3.2 Recalling Context and Constructions.

In Section 2 of [HOV95] Hubbard & Oberste-Vorth an open neighborhood U of $J(p) \subset \mathbb{C}$ is constructed such that $U' \equiv p^{-1}(U) \Subset U$ and the map $p: U' \rightarrow U$ is a covering map. This is a standard construction. We can assume without loss of generality that U is chosen sufficiently small that it is a finite distance from any critical points of p . Open neighborhoods $U_z \subset U$ in $\mathbb{C}$ are attached to each point $z \in J(p)$. Later in section 4 of the same paper these

neighborhoods are used to associate an open subset B_z of $\mathbb{C}^2$ to each point $z \in J(p)$ so that $f_a: B_z \rightarrow B_{p(z)}$ is a crossed mapping for each $z \in J(p)$ and each a with $|a|$ sufficiently small. A radius $\mathbf{r}$ is chosen and U_z is defined to be the ball of radius $\mathbf{r}$ about the point z , where the distance is measured in the Kobayashi metric of U . The radius $\mathbf{r}$ is only special in that it is chosen so be so small that the neighborhoods U_z create telescopes for p of uniformly bounded modulus.

We will strengthen this requirement on $\mathbf{r}$ a small amount here by requiring that $\mathbf{r}$ is small enough that for each $z \in J(p)$ the map $p^{\circ 2}$ is a biholomorphism of U_z onto its image and the map p is a biholomorphism from the ball of radius $3\mathbf{r}$ about z onto its image.

The neighborhoods B_z are constructed in [HOV95] by first selecting a small value δ (which must satisfy various requirements). One then defines $v: \mathbb{C}^2 \rightarrow \mathbb{C}$ by

$$v(x, y) \equiv p(y) - x. \quad (3.1)$$

Taking

$$V' \equiv \text{pr}_1^{-1}(U) \cap v^{-1}(\mathbb{D}_\delta) \quad (3.2)$$

then a well-defined function $u: V' \rightarrow U'$ is constructed, which is given by $u(x, y) = p^{-1}(x)$, the inverse image always being chosen to be the one “close to y ”. We make this precise in Lemma 3.7.

We define m_p to be the minimum of $|p'|$ on $\bar{U}$. Since $\bar{U}$ is a finite distance from the critical set of p then $m_p > 0$.

Lemma 3.6. *There exists some $\beta > 0$ such that*

1. *p maps the (Euclidean) ball of radius β about an arbitrary point $y \in U$ biholomorphically onto its image.*
2. *β is smaller than the Euclidean distance from U' to ∂U .*
3. *The Euclidean ball of radius $\beta/2$ about any point $u \in U'$ is mapped biholomorphically onto its image, which contains the Euclidean disk of radius $m_p\beta/8$.*

Proof. Part 1 follows from a straightforward proof by contradiction. Part 2 is obvious. Part 3 follows from the Koebe 1/4 theorem. $\square$

We shrink δ as necessary so that $\delta < \frac{1}{8}m_p\beta$. We accordingly shrink the sets V' and the boxes B_z .

Lemma 3.7. *There is a well defined holomorphic function $u: V' \rightarrow U'$ which satisfies $p(u(x, y)) = x$ and $u(x, y)$ is the unique preimage of x within a distance $\frac{1}{2}\beta$ of y .*

Proof. That u is well defined follows using Lemma 3.6 and the definition of V' . Holomorphy of u follows from the fact that $p: U' \rightarrow U$ is a local biholomorphism. $\square$

Then for each $z \in J(p)$ the set

$$B_z \equiv \left\{ \begin{pmatrix} x \\ y \end{pmatrix} \in V' \mid u(x, y) \in U_z \right\} \quad (3.3)$$

is an open neighborhood of the point $(p(z), z)$.

Hubbard & Oberste-Vorth then prove that the mapping

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} u(x, y) \\ v(x, y) \end{pmatrix}$$

is a biholomorphic isomorphism of B_z onto the bidisk $U_z \times \mathbb{D}_\delta$, where u and v are defined in Lemma 3.7 and equation (3.1).

This can be extended to all of V' .

Lemma 3.8.

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} u(x, y) \\ v(x, y) \end{pmatrix}$$

is a biholomorphic isomorphism of V' onto $U' \times \mathbb{D}_\delta$.

Proof. Using Lemma 3.6 and Lemma 3.7 one can construct a holomorphic inverse. □

3.3 The Stable and Unstable Manifolds as a Varies

Given that p is a hyperbolic polynomial, there is some $A > 0$ such that $f_a: B_z \rightarrow B_{p(z)}$ is a crossed mapping for each $z \in J(p)$ whenever $0 < |a| < A$. It can be verified that A can be chosen so that these crossed mappings have uniformly bounded horizontal and vertical sizes. Given any $\underline{z} = (\dots, z_{-2}, z_{-1}, z_0) \in \hat{J}(p)$ the sequence of neighborhoods $\dots \rightarrow B_{z_{-2}} \rightarrow B_{z_{-1}} \rightarrow B_{z_0} \rightarrow B_{p(z_0)} \rightarrow B_{p^2(z_0)} \dots$ (all mapped by f_a) form a sequence of crossed neighborhoods. Then for each $|a| < A$ there is a natural map π_a from the natural extension $\hat{J}(p)$ of the Julia set to J_a which is a homeomorphism for each $a \neq 0$ and is the standard projection of $\hat{J}(p)$ to $J(p)$ for $a = 0$.

By definition, the point $\pi_a(\underline{z})$ is the unique point in B_{z_0} which lies in both the stable and the unstable manifold of the sequence of crossed mappings. We note that given a sequence of crossed mappings $U_0 \times V_0 \rightarrow U_1 \times V_1 \rightarrow \dots$ then the construction of the stable manifold of a crossed mapping in [HOV95] is found by first taking a point $u_n \in U_n$ and taking the preimage of $\{u_n\} \times V_n$ by $f_1 \circ \dots \circ f_n$, which is, by the hypothesis on crossed mappings, necessarily the graph of a function $g: V_1 \rightarrow U_1$. One then takes a limit of these graphs as $n \rightarrow \infty$.

We will use the notation $\tilde{f}(x, y, a) \equiv (f_a(x, y), a)$ when it is convenient to think of $f_a(x, y)$ as a self map in x, y and a .

Lemma 3.9. *Given $\underline{z} \in \hat{J}(p)$ there is a unique holomorphic map $g_+: V_0 \times \mathbb{D}_A \rightarrow U_0$ such that the local stable manifold through the point $\pi_{a'}(\underline{z}) \in J_{a'}$, $a' \neq 0$ is the graph of $g_+(\cdot, a'): V_0 \rightarrow U_0$. In the case where $a = 0$ this graph gives a vertical line through $(p(z), z)$ (the natural analogue of the stable manifold in the degenerate case).*

Proof. As the maps f_a depend holomorphically on a then for $a \neq 0$ these graphs fit together holomorphically, since they are just the preimage of $\{u_n\} \times V_n \times \mathbb{D}_A$ by $\hat{f}^{\circ n}$, which is holomorphic in both $(x, y) \in \mathbb{C}^2$ and in $a \in \mathbb{D}_A$. Thus, the stable manifold of f_a through $z \in J$ is the graph of a holomorphic map $g_{z+}: V_0 \times \mathbb{D}_A^* \rightarrow U_0$. At $a = 0$ the map is not a crossed mapping, however g_{z+} is holomorphic and clearly bounded on $V_0 \times \mathbb{D}_A^*$ and so g_{z+} has a unique holomorphic extension to a map (which we will still call g_{z+}) from $V_0 \times \mathbb{D}_A$ to U_0 .

Now the function $u - g_{z+}(v, a)$ vanishes on the stable manifold and by continuity there is a neighborhood N of $\pi_0(f^{-1}(z))$ such that $f_a(N) \subset U_0 \times V_0$ for sufficiently small a . Therefore the pullback $\tilde{f}^*(u - g_{z+}(v, a))$ by $\tilde{f}$ is defined in N for all sufficiently small a . Since the stable manifold of $\pi_a(\hat{f}^{-1}(z))$ is mapped into the stable manifold of $\pi_a(z)$ by f_a then $\tilde{f}^*(u - g_{z+}(v, a))$ vanishes on the graph $\check{g}_{\hat{f}^{-1}(z)+}: V_0 \times \mathbb{D}_A \rightarrow U_0 \times V_0 \times \mathbb{D}_A$ of $g_{\hat{f}^{-1}(z)+}$. This is true even if $a = 0$ because of continuity.

Now the image of $f_0 \circ \check{g}_{\hat{f}^{-1}(z)+}$ lies on $C(p)$ since f_0 maps all of $\mathbb{C}^2$ to $C(p)$. If the image of $\check{g}_{\hat{f}^{-1}(z)+}$ does not lie in a fiber of f_0 then it follows that $f_0 \circ \check{g}_{\hat{f}^{-1}(z)+}$ contains an open subset of $C(p)$ in its image. Therefore $u - g_{z+}(v, 0)$ would have to vanish on $C(p)$. But this is impossible since $C(p)$ is given by $v = 0$ in the (u, v) coordinates and restricting $u - g_{z+}(v, 0)$ to $v = 0$ one obtains the false statement $u - g_{z+}(0, 0) \equiv 0$. This contradiction shows that $\check{g}_{\hat{f}^{-1}(z)+}$ lies in a fiber of f_0 , that is, in a vertical line. $\square$

Observation 3.10. Recalling that for each $\underline{z} = (\dots, z_{-2}, z_{-1}, z_0) \in \hat{J}(p)$ the neighborhood B_{z_0} was equal to $U_{z_0} \times \mathbb{D}_\delta$ using (u, v) coordinates, where U_{z_0} was an open neighborhood of $z_0 \in J(p) \subset \mathbb{C}$ then we see that the local stable manifold of $\pi_a(\underline{z})$ given in Lemma 3.9 by a holomorphic function $g_{\underline{z}+}: V_0 \rightarrow U_0$ is just a holomorphic map from $\mathbb{D}_\delta$ to U_{z_0} . What is more, this stable manifold was precisely the stable manifold of the sequence $B_{z_0} \rightarrow B_{p(z_0)} \rightarrow B_{p^{\circ 2}(z_0)} \rightarrow \dots$ and was therefore dependent only on z_0 and not on any other point in the history $\underline{z}$. Thus the map $g_{\underline{z}+}$ depends only on z_0 .

Convention 3.11. In accordance with the above observation we will reduce our notation of $g_{\underline{z}+}$ to g_{z+} as g_{z+} only depends on the final term z of $\underline{z}$.

Convention 3.12. We will continue to use $\check{g}_{z\pm}$ to denote the graph in $V' \subset \mathbb{C}^2$ of $g_{z\pm}$. When we are thinking of $g_{z\pm}$ as a function of a as well, we will similarly use $\check{g}_{z\pm}$ to denote the graph in $V' \times \mathbb{D}_A \subset \mathbb{C}^3$.

Lemma 3.13. Given $z \in \hat{J}(p)$ then there is a unique holomorphic map $g_-: U_0 \times \mathbb{D}_A \rightarrow V_0$ such that the local unstable manifold through the point $\pi_a(z) \in J_a$ is the graph of $g_-(\cdot, a'): U_0 \rightarrow V_0$. For $a = 0$ it is simply the portion of the graph $x = p(y)$ about $\pi_0(z)$.

Proof. Since for all $a \neq 0$ these are just the images of some (arbitrary) $U_{-n} \times \{v_0\}$ under $f_{-n} \circ \dots \circ f_1$ in $U_0 \times V_0$ then by the hypothesis on a sequence of crossed mappings the result is a graph $g_-: U_0 \rightarrow V_0$. These graphs again all fit together to form a single holomorphic “sheet” as when put together they are simply the image of $U_{-n} \times \{v_0\} \times \mathbb{D}_A$ iterated n times by f_a (which depends holomorphically on a). For each a the limit of these graphs will be a graph. Thus the limit is a graph, which is, by [HOV95], the local unstable manifold of $\pi_a(z) \in J_a$ for each $|a| < A$. In the case $a = 0$ it is simply a portion of the graph $x = p(y)$ about the point $\pi_0(z)$ (and is therefore the appropriate version of the unstable manifold for this case). $\square$

We recall Theorem 5.9 of [BS91a].

Theorem 3.14. *Given that $a \neq 0$, if f_a is hyperbolic and $|a| \leq 1$, then $W^s(J_a) = J_a^+$. If $s_1, \dots, s_k$ are the sinks of f_a then $W^u(J_a) = J_a^- \setminus \{s_1, \dots, s_k\}$.*

Definition 3.15. Given $z \in J(p)$ let $\Delta_{z,a}$ be the image $\hat{g}_{z+}(\mathbb{D}_\delta, a) \subset \mathbb{C}^2$ which is precisely the local stable manifold in B_z corresponding to the sequence of crossed mappings $B_z \rightarrow B_{p(z)} \rightarrow B_{p^2(z)} \rightarrow \dots$. Given $r < 1$ we let $\Delta_{z,a}(r)$ be the image of $\mathbb{D}_\delta$ under $\hat{g}_{z+}$, and we let $\Sigma_{z,a}(r)$ be the image of the circle S_δ under $\hat{g}_{z+}$. Hence both $\Delta_{z,a}(r)$ and $\Sigma_{z,a}(r)$ lie in $\Delta_{z,a}$.

We will show that given $z_1, z_2 \in J(p)$, if $\Delta_{z_1,a}$ and $\Delta_{z_2,a}$ overlap then $z_1 = z_2$. First we recall the standard telescope lemma.

Lemma 3.16. *If z_1 and z_2 are points in $J(p)$ and if $U_{p^{on}(z_1)} \cap U_{p^{on}(z_2)} \neq \emptyset$ for every $n \geq 0$ then $z_1 = z_2$.*

Proof. This is the standard telescope result. □

We now show our desired result about disjointness of the sets $\Delta_{z,a}$.

Lemma 3.17. *Assume that for $z_1, z_2 \in J(p)$ one has $\Delta_{z_1,a} \cap \Delta_{z_2,a} \neq \emptyset$. Then $z_1 = z_2$.*

Proof. Assume that $\Delta_{z_1,a} \cap \Delta_{z_2,a} \neq \emptyset$. Let w be a point in the intersection. Then $f_a^{on}(w) \subset B_{p^{on}(z_1)}$ and $f_a^{on}(w) \subset B_{p^{on}(z_2)}$ for all $n \geq 0$. Recalling that $B_z = U_z \times \mathbb{D}_\delta$ (in the (u, v) coordinates defined on V') then one has $u(f_a^{on}(w)) \in U_{p^{on}(z_1)} \cap U_{p^{on}(z_2)}$ for all $n \geq 0$ and so $z_1 = z_2$ by Lemma 3.16. □

Lemma 3.18. *The maps $g_{z_0+}: \mathbb{D}_\delta \times \mathbb{D}_A \rightarrow U_{z_0}$ vary continuously (in the sense of locally uniform convergence of maps) with $z_0 \in J(p)$.*

Proof. From Lemma 3.17 the functions $g_{z_0+}: \mathbb{D}_\delta \times \mathbb{D}_A \rightarrow U'$ have disjoint graphs from which the result follows. □

Lemma 3.19. *The maps $\pi_a: \hat{J}(p) \rightarrow \mathbb{C}^2$ vary continuously with a in the sense of uniform convergence of maps for $a \in \mathbb{D}_A$.*

Proof. This is clear for nonzero a . Because $\pi_a(z)$ varies holomorphically with a it follows easily that one also has continuity at $a = 0$. □

Lemma 3.20. *If $(x, y, 0) \in \overline{\mathcal{C}}$ then either $(x, y) \in \overline{\mathcal{C}_0}$ or $x \in J(p)$ or $(x, y) \in C(p)$.*

Proof. If $z \in \overline{\mathcal{C}}$ but $z \notin \overline{\mathcal{C}_0}$ then since $\mathcal{C}$ is a closed analytic variety on its domain of definition then $z \in (K(p) \times \mathbb{C}) \cup C(p)$. Now if z lies in the interior of $K(p) \times \mathbb{C}$ then z is attracted to the cycle of an attracting periodic point α of p . It follows that z lies in the interior of K_a^+ for all sufficiently small a , which is a contradiction. Therefore $z \in J(p) \times \mathbb{C}$ or $z \in C(p)$. □

It is easy to see what the critical locus $\mathcal{C}_0$ looks like if $J(p)$ is connected. Since $\varphi_{0,+}(x, y) = b_p(x)$, the Böttcher coordinate of x , then the leaves of $\mathcal{F}_0^+$ are simply vertical. Since $\varphi_{0,-}(x, y) = (p(y) - x)^{\frac{1}{d}}$, then the leaves of $\mathcal{F}_0^-$ are translates in the x direction of the graph $C(p)$. Thus $\mathcal{F}_0^+$ and $\mathcal{F}_0^-$ will be tangent exactly along the horizontal lines $y = c$ where c is a critical point of p . Thus $\mathcal{C}_0$ is a union of horizontal lines (restricted to $U_{0+} \cap U_{0-} = \mathbb{C}^2 \setminus (C(p) \cup (K(p) \times \mathbb{C}))$), one at the level of each critical point of p . This fact will be quite important in what follows.

Lemma 3.21. *If the Julia set of p is connected then $\mathcal{C}_0$ is the union of the sets $\{(x, c) | x \notin K(p)\}$ over the critical points c of p .*

When we perturb a away from zero, we will need to be able to control the motion of the critical locus. The main difficulty is that we must control what happens at the boundary of U_{0+} and U_{0-} . We will do this by choosing a tube about each of horizontal lines in $\mathcal{C}_0$ which contains the perturbed component of $\mathcal{C}_a$ as a moves away from zero. Of course, that the different components of $\mathcal{C}_0$ remain distinct components when a is perturbed will have to be shown e.g. consider the equations $y(y - 1)x + a = 0$. When $a = 0$ this is a curve which has three components, but as a is perturbed to a nonzero value this becomes a smooth curve with only one component. Since $\mathcal{C}_0$ is not defined outside of $U_{0+} \cup U_{0-}$ it is conceivable that such a thing could happen to $\mathcal{C}_a$ as a is varied from zero, i.e. we could have a large portion of $\mathcal{C}_a$ which is “hidden” in the boundary of definition when $a = 0$. We have to demonstrate that such oddities do not occur.

We will make use of the following version of the “Inclination Lemma” about the degenerate map f_0 . Compare to [KH95] and [Rue89].⁴

Lemma 3.22 (An inclination lemma near the degenerate case.). *Given a sequence $a_k \rightarrow 0$ and a sequence of points $w_k \in \mathcal{C}_{a_k}$ converging to some point $w_\infty \in J_0^+$ then the leaves $\mathcal{L}_{a_k}^+(w_k)$ converge locally and without ramification to a vertical line through w_∞ (i.e. to the leaf of J_0^+ through w_∞).*

Sketch of Proof. We construct a neighborhood $\mathcal{N}$ of J_0 in $\mathbb{C}^3$ such that each point w of $\mathcal{N}$ lies in a box $B_{\mathfrak{z}(w)}$ and such that if w and $f_a(w)$ both lie in $\mathcal{N}$ then $f_a: B_{\mathfrak{z}(w)} \rightarrow B_{\mathfrak{z}(f_a(w))}$ is a crossed mapping.


We then apply iterates of f_a to each member of our sequence to move all the points of the sequence a definite distance away from K^+ . Then the leaves of $\mathcal{F}^+$ through the new sequence must converge to the leaves of $\mathcal{F}_0^+$, and must therefore become vertical lines in the limit.

We then pull back the leaves by iteration so that they pass through the members of the original sequence. By using the boxed mapping construction, we can guarantee that these leaves are still graphs in their respective boxes $B_{z(w_k)}$. It is then easy to show that the leaves become vertical in the limit.

Once this is established, one can apply this argument to $f_{a_k}(w_k)$ and obtain a sequence of plaques which become vertical in the limit. Taking the preimages of these plaque under f

⁴This lemma is also sometimes called the λ -lemma, but we avoid this term because the term λ -lemma is typically used in complex dynamics to refer to a result about holomorphic motions. See [MnSS83], [Lyu83].

to obtain a sequence of plaques through the points w_k , it is easy to show that the resulting plaques become vertical over arbitrarily large sets. $\square$


 Figure 1: The tube $\mathcal{T}_c^\circ$ and its environment.

4 Components of the Critical Locus.

4.1 Trapping and Mapping Components.

Standing Assumption. Throughout the rest of this paper we will make the additional assumption that the orbits of each of the critical points remain bounded, that is, that $J(p)$ is connected. We also assume that all of the critical points of p are simple.

Definition 4.1. Given a critical point c of $p(x)$ we let H_{ca} be the component of $\mathcal{C}_a$ which is asymptotic to $y = c$ as $|x| \rightarrow \infty$. We call H_{ca} the *primary horizontal component* of the critical locus corresponding to the critical point c of $p(z)$.

We will show that if a is sufficiently small then given distinct critical points c_1, c_2 then $H_{c_1,a}$ and $H_{c_2,a}$ are distinct and disjoint components of $\mathcal{C}_a$. Note that we know from Lemma 3.21 that $H_{c0} = \{(x, c) \mid x \notin K(p)\}$.

For each critical point c of $p(z)$ we select an open disk $A_c \subset \mathbb{C}$ about c . We assume these disks are chosen small enough to have disjoint closures. We let Ω_c be an open disk about $p(c)$, sufficiently small that $\overline{\Omega}_c$ lies in the basin of an attracting periodic point. We let $Q_c \equiv \mathbb{C} \setminus \overline{\Omega}_c$.

If necessary, we shrink A_c so that $Q_c \times A_c$ is a positive distance from $C(p)$. For each critical point c we let B_c be a second disk about c of half the radius of A_c . We let $\mathcal{T}_c^\circ = Q_c \times (A_c \setminus \overline{B}_c)$.

We think of $\mathcal{T}_c^\circ$ as a thin hollow tube about $\{y = c\}$ as pictured in Figure 1. We will show that for small values of a the component H_{ca} of $\mathcal{C}_a$ remains inside this tube. We let $\mathcal{T}_c^\bullet = Q_c \times B_c$ be the core of the tube $\mathcal{T}_c^\circ$ and we let $\mathcal{T}_c^\bullet = Q_c \times A_c$ be the filled tube.

Lemma 4.2. *There exists some $\delta_T > 0$ and some $\epsilon_T > 0$ such that if $|a| < \epsilon_T$ then:*

1. *The filled tube $\mathcal{T}_c^\bullet$ is distance at least δ from the set K_a^- for each critical point c of $p(z)$.*
2. *One has $\overline{\Omega}_c \times A_c \subset K_a^+$ for each critical point c .*
3. *For each critical point c , $H_{ca} \subset \mathcal{T}_c^\bullet$ and this is the only component of $\mathcal{C}_a$ which intersects $\overline{\mathcal{T}_c^\bullet}$.*

4. The foliations $\mathcal{F}_a^+$ and $\mathcal{F}_a^-$ have contact of order two at any point of H_{ca} for any critical point c . In particular, H_{ca} is smooth, has multiplicity one, and is everywhere transverse to both $\mathcal{F}_a^+$ and $\mathcal{F}_a^-$.

Proof. 1. This is a trivial consequence of the construction of $\mathcal{T}_c^\bullet$ and Lemma 1.21.

2. This follows from the fact that $\overline{\Omega_c \times A_c}$ is relatively compact in a basin of attraction of f_0 .

3. We prove first that if $|a|$ is sufficiently small then for each critical point c , any component X of $\mathcal{C}_a$ which intersects $\overline{\mathcal{T}_c^\bullet}$ lies in $\mathcal{T}_c^\bullet$. Hence, for the sake of contradiction, assume that there exists a sequence $a_i \rightarrow 0$ and a sequence z_i such that $z_i \in \mathcal{C}_{a_i} \cap \overline{\mathcal{T}_c^\bullet}$. It follows from Lemma 2.12 that the sequence z_i is bounded. Letting z_∞ be any accumulation point of z_i then either $z_\infty \in U_0^+$ or $z_\infty \in J_0^+$. One concludes from Lemma 1.12 combined with either Lemma 1.19 in the former case or with Lemma 3.22 in the latter case that $z_\infty \in \bigcup \overline{H_{c0}}$. However this is a contradiction since $\overline{\mathcal{T}_c^\bullet}$ is disjoint from each $\overline{H_{c0}}$. Thus $\mathcal{C}_a \cap \overline{\mathcal{T}_c^\bullet} = \emptyset$ for $|a|$ sufficiently small.

Now assume X is a component of $\mathcal{C}_a$ intersecting $\overline{\mathcal{T}_c^\bullet}$. Since X is disjoint from $\overline{\Omega_c \times A_c}$ by part 2 and from $\overline{\mathcal{T}_c^\bullet}$ then it follows easily from the definitions of these sets that $X \subset (\mathbb{C} \setminus \overline{\Omega_c}) \times B_c \subset \mathcal{T}_c^\bullet$. We will show that X must be H_{ca} .

Choose some sequence of points $z_k \in X$ such that

$$\lim_{k \rightarrow \infty} \mathcal{G}_a^+(z_k) = \sup_{z \in X} \mathcal{G}_a^+(z).$$

A straightforward argument by contradiction shows that $z_k \rightarrow \infty$. By Lemma 2.12, z_k converges to the point of the extension of $\mathcal{C}_a$ to $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{D}_R$ with (u, y, a) coordinates $(0, c, a)$. From Theorem 2.13, $X = H_{ca}$.

4. If there is no ϵ_T such as the lemma claims to exist then one can choose a critical point c of $p(x)$ and sequences $a_k \rightarrow 0$ and $z_k \in H_{ca_k}$ such that $\mathcal{L}_{a_k}^+(z_k)$ and $\mathcal{L}_{a_k}^-(z_k)$ have order of contact at least three at z_k for each i . We can assume without loss of generality that $|z_k| \rightarrow \infty$ or that z_k converges to some point $z_\infty \in \overline{\mathcal{T}_c^\bullet}$.

If $|z_k| \rightarrow \infty$ then we will need to use the extension of $\mathcal{C}$ to $x = \infty$, hence we change coordinates letting $u = 1/x$. Since $z_k \in \mathcal{C}_{a_k}$ by assumption and the leaves of $\mathcal{F}_{a_k}^+$ and $\mathcal{F}_{a_k}^-$ have contact of order at least three at z_k then Lemma 2.7 shows that $\mathcal{F}_{a_k}^+$ and $\mathcal{C}_{a_k}$ have intersection multiplicity at least two at z_k . Since $\mathcal{C}_{a_k}$ is defined by $w(u, y, a_k)$ for a_k and u sufficiently small and y bounded then the directional derivative of $w(u, y, a)$ along the leaf of $\mathcal{F}_{a_k}^+$ through z_k is zero by Lemma 2.5. But $\frac{\partial w}{\partial y}(0, c, 0) = 0$ since this is the directional derivative along the leaf of $\mathcal{F}_0^+$ by Corollary 2.9. But this contradicts the expression for w about $(0, c, 0)$ given in Lemma 2.12.

Having obtained a contradiction if $|z_k| \rightarrow \infty$, assume $z_k \rightarrow z_\infty = (x_\infty, y_\infty)$. By part 1 we know $z_\infty \notin K_0^-$, so $(x_\infty, y_\infty, 0) \in \mathfrak{V}_{0,-}$. By Lemma 1.22 we know $z_\infty \in \overline{\mathbb{C}^2 \setminus K_0^+}$. Hence φ_-^{dj} is well defined at $(x_\infty, y_\infty, 0)$ for some integer j . Then by either Lemma 3.22 or Lemma 1.19, depending on whether $z_\infty \in J_0^+$ or not, one concludes that one can parameterize plaques of the leaves $\mathcal{L}_{a_k}^+(z_k)$ such that they converge to a parameterization of a plaque of the leaf of J_0^+ or $\mathcal{F}_0^+$ through z_∞ (which is a vertical line in either case). Since $\mathcal{F}_{a_k}^+$ and $\mathcal{F}_{a_k}^-$ have contact of order at least three at z_k for each k then by Lemma 2.5 the directional derivative of $\varphi_{a_k,-}^{dj}$ along the leaf $\mathcal{L}_{a_k}^+(z_k)$ is zero. But by continuity of φ_-^{dj} it follows that the directional

derivative of $\varphi_{0,-}^{dj}$ in the y direction is zero. This contradicts the expression for $\varphi_{0,-}^{dj}$ given in Lemma 1.12. Hence we conclude that there is no such sequence a_k and z_k and so $\mathcal{F}_a^-$ and $\mathcal{F}_a^+$ have contact of order two at every point of H_{ca} whenever $|a|$ is sufficiently small.

The rest of the Lemma is an immediate consequence of Corollary 2.8. $\square$

Lemma 4.3. *If $|a| < \epsilon_T$ then the map $\mathcal{G}_a^+|_{H_{ca}} : H_{ca} \rightarrow (\frac{1}{d-1} \log |a|, \infty)$ is proper.*

Proof. This is easy to prove using parts 1 and 3 of Lemma 4.2. $\square$

The following is a standard fact:

Lemma 4.4. *If M is a Riemann surface and if g is harmonic on M then*

- *The zeros of ∇g are discrete*
- *The zeros of ∇g correspond to critical points of a holomorphic h such that $g = \operatorname{Re} h$ and the index of the zero is equal the negative of the order of the critical point.*

Theorem 4.5. *If $|a| < \epsilon_T$ then H_{ca} is a punctured disk and the map $\varphi_{a,+}$ extends holomorphically from $H_{ca} \cap V_+$ to a biholomorphism $\varphi_{a,+} : H_{ca} \rightarrow \mathbb{C} \setminus \overline{\mathbb{D}}$.*

Proof. Using Lemma 4.4 and part 4 of Lemma 4.2 we can conclude that $\mathcal{G}_a^+$ has no critical points on H_{ca} . Moreover, by Lemma 4.3, $\mathcal{G}_a^+$ is proper. By Theorem 2.13, Corollary 1.16 and the definition of $\mathcal{G}_a^+$ it follows that the fibers of $\mathcal{G}_a^+|_{H_{ca}}$ about the point $(\infty, c) \in H_{ca}$ are topological circles and, by Morse theory, H_{ca} is a topological annulus. Since H_{ca} contains a punctured disk about (∞, c) then H_{ca} is either $\mathbb{D}^*$ or $\mathbb{C}^*$.

The function $\varphi_{a,+} \circ f_a^{\circ k} : H_{ca} \rightarrow \mathbb{C} \setminus \overline{\mathbb{D}}$ induces the map $j \mapsto d^k \cdot j$ on $\pi_1(H_{ca}) \rightarrow \pi_1(\mathbb{C} \setminus \overline{\mathbb{D}})$ by Corollary 1.16, Theorem 2.13 and the recursion relation for $\varphi_{a,+}$. It follows that $\varphi_{a,+} \circ f_a^{\circ k}$ has a holomorphic $d^{k\text{th}}$ root which is equal to $\varphi_{a,+}$ on $H_{ca} \cap V_+$. It then follows that $\varphi_{a,+}$ has a holomorphic extension to all of H_{ca} . What is more, $\varphi_{a,+}$ is proper since $\mathcal{G}_a^+$ is. By considering $\varphi_{a,+}$ about (∞, c) it follows easily that $\varphi_{a,+}|_{H_{ca}} : H_{ca} \rightarrow \mathbb{C} \setminus \overline{\mathbb{D}}$ has degree one and is therefore a biholomorphism. $\square$

Given a, b with $|a| < \epsilon_T$ and $|b| < \epsilon_T$ we define a biholomorphism $\tau_{ab} : H_{ca} \rightarrow H_{cb}$ by $\tau_{ab} = \left(\varphi_{b,+}|_{H_{cb}} \right)^{-1} \circ \varphi_{a,+}$. Then $\varphi_{a,+} \circ \tau_{ab} = \varphi_{b,+}$.

Proposition 4.6. *The maps $\tau_{ab} : H_{ca} \rightarrow H_{cb}$ vary holomorphically in a and b .*

Proof. The precise meaning of this is that if one defines $\mathcal{H}_c \subset \mathcal{C}$ to be $\{(x, y, a) | (x, y) \in H_{ca}, a \in \mathbb{D}_{\epsilon_T}\}$ then from Lemma 2.12 and part 3 of Lemma 4.2 it is clear that $\mathcal{H}_c$ is a component of $\mathcal{C} \cap (\mathbb{C}^2 \times \mathbb{D}_{\epsilon_T})$ and this proposition states that the map $\tilde{\tau} : \mathcal{H}_c \times \mathbb{D}_{\epsilon_T} \rightarrow \mathcal{H}_c$ given by $\tilde{\tau}(x, y, a, b) \equiv \tau_{a,b}(x, y)$ is holomorphic.

The proof is elementary since from our original construction the function $\varphi_{a,+}(x, y) : V_+ \times \mathbb{D}_R \rightarrow \mathbb{C}$ is holomorphic in x, y and a . It was shown in the proof of Theorem 4.5 that $\varphi_{a,+} \circ f_a^{\circ k}|_{H_{ca}}$ has a $d^{k\text{th}}$ root which agrees with $\varphi_{a,+}$ when $|a| < \epsilon_T$ and this root gives the extension of $\varphi_{a,+}$ to $f^{-k}(V_+) \cap H_{ca}$. It follows that $\varphi_{a,+} \circ f_a^{\circ k}(x, y) : \mathcal{H}_c \cap \{(x, y, a) | f_a^{\circ k}(x, y) \in V_+, a \in \mathbb{D}_{\epsilon_T}\} \rightarrow \mathbb{C}$ has a $d^{k\text{th}}$ root which agrees with $\varphi_{a,+}(x, y)$ on $\mathcal{H}_c \cap (V_+ \times \mathbb{D}_{\epsilon_T})$. Consequently the extension $\varphi_{a,+}(x, y) : \mathcal{H}_c \rightarrow \mathbb{C} \setminus \overline{\mathbb{D}}$ is holomorphic in x, y and a . It is easy to see that the map $\widetilde{\varphi_{a,+}} \equiv (\varphi_{a,+}(x, y), a) : \mathcal{H}_c \rightarrow (\mathbb{C} \setminus \overline{\mathbb{D}}) \times \mathbb{D}_{\epsilon_T}$ is a biholomorphism. It follows that $\tilde{\tau}_{ab}$ is holomorphic in a and b from the easily verified relationship $\widetilde{\varphi_{b,+}} \circ \tilde{\tau}_{ab} = \widetilde{\varphi_{a,+}}$. $\square$

4.2 Classification of the Critical Components.

Since our strategy has been to consider the degenerate map f_0 and then to consider $\mathcal{C}_a$ as a deformation of $\mathcal{C}_0$, we need to ensure that we have accounted for every component of $\mathcal{C}_a$. It is plausible that $\mathcal{C}_a$ has a component that “escapes to infinity” as $a \rightarrow 0$, and thus this component would be invisible to us in $\mathcal{C}_0$. We will start by showing that any component of $\mathcal{C}_a$ meets either J_a^- or J_a^+ . We will be able to use this to show that any component of $\mathcal{C}_a$ is an iterate of a component of the form H_{ca} , and thus we have accounted for every component of $\mathcal{C}_a$ by accounting for the components H_{ca} and their iterates.

Lemma 4.7. *If W_a is some component of $\mathcal{C}_a$ then ∂W_a contains a point in either J_a^+ or J_a^- .*

Proof. Consider the positive plurisubharmonic function $\mathfrak{g}(x, y, a) \equiv \mathcal{G}_a^+(x, y) + \mathcal{G}_a^-(x, y) - \frac{1}{d-1} \log |a|$ on $\tilde{U}^+ \cap \tilde{U}^-$. It is easy to show that if z_n is a sequence of points of W_a such that $\lim_{n \rightarrow \infty} \mathfrak{g}(z_n) = \inf_{z \in W_a} \mathfrak{g}(z)$ then z_n has an accumulation point in $J_a^+ \cup J_a^-$. $\square$

We will need specific local stable manifolds about the points of J_a . We know that if $|a|$ is sufficiently small then given $\underline{z} = (\dots, z_{-2}, z_{-1}, z_0) \in \hat{J}(p)$ there is an associated neighborhood $B_{z_0} = V_0 \times U_{z_0}$ of $\pi_a(\underline{z})$ and the local stable manifold in B_{z_0} is the graph of a holomorphic function from $V_0 \rightarrow U_{z_0}$.

We recall that in Lemma 3.8 it was shown that (u, v) coordinates are defined on an open set V' which contains each of the sets B_z and that (u, v) coordinates provide a biholomorphic isomorphism of V' onto $U' \times \mathbb{D}_\delta$. Also, since $v(x, y) = p(y) - x$ then v is defined on all of $\mathbb{C}^2$, not just on V' .

Definition 4.8. We now fix some positive $r < 1$ such that each of the filled tubes $\mathcal{T}_c^\bullet$ is a finite distance from the set $\{(x, y) \in \mathbb{C}^2 \mid |v(x, y)| \leq r\delta\}$.

Proof that such an r exists. Each of the filled tubes lies a finite distance from $C(p)$ by construction. We let s be half the minimal distance between $C(p)$ and the nearest tube. Since $(x - v(x, y), y) \in C(p)$ for all $(x, y) \in \mathbb{C}^2$ then the set $\{(x, y) \mid |v(x, y)| \leq s\}$ is comprised of points no further than s from $C(p)$. Thus $r = s/\delta$ will do. $\square$

Lemma 4.9. *Given $\epsilon > 0$ there exists $\delta_r(\epsilon) > 0$ such that if $|a| < \delta_r(\epsilon)$ and if $(x, y) \in V' \subset \mathbb{C}^2$ then*

- $|v(x, y)| \leq r\delta$ implies that $\mathcal{G}_a^-(x, y) < \frac{1}{d} \log |r\delta| + \epsilon$
- $|v(x, y)| \geq r\delta$ implies that $\frac{1}{d} \log |r\delta| - \epsilon < \mathcal{G}_a^-(x, y)$

Proof. This is an easy consequence of Theorem 1.24. $\square$

Lemma 4.10. *There exists $\epsilon_S > 0$ such that if $|a| < \epsilon_S$ and $z \in J(p)$ then the gradient of the restriction of $\mathcal{G}_a^-$ to the local stable manifold $\check{g}_{z+}(\mathbb{D}_\delta, a)$ is defined and nonzero on the curve $\check{g}_{z+}(S_{r\delta}, a)$.*

Proof. First we recall that $\mathcal{G}_a^- - \frac{1}{d-1} \log |a|$ is pluriharmonic, and is therefore smooth, away from its zero set. From Lemma 4.9 we conclude that as long as $|a| < \delta_r(\epsilon)$ and $|a| < ((r\delta)^{1/d} e^{-\epsilon})^{d-1}$ then if $(x, y) \in \check{g}_{z+}(S_{r\delta}, a)$ then $(x, y) \in B_z \in V'$ and $|v(x, y)| = r\delta$ so

$$\mathcal{G}_a^-(x, y) - \frac{1}{d-1} \log |a| > \frac{1}{d} \log |r\delta| - \epsilon - \frac{1}{d-1} \log |a| > 0.$$

Thus $\mathcal{G}_a^-$ is smooth at such points.

Assume that no such ϵ_S existed. Then there exists a sequence $a_i \rightarrow 0$ and a sequence of points $z_i \in J(p)$ and a sequence of points $w_i \in U_{z_i} \times S_{r\delta} \subset B_{z_i}$ such that for each i the restriction of $\mathcal{G}_{a_i}^-$ to the stable manifold $\check{g}_{z_i+}(\mathbb{D}_\delta, a) \subset B_{z_i}$ has gradient zero at the point $\check{g}_{z_i+}(w_i, a_i)$. Then by compactness we can replace our sequence with a subsequence if necessary such that both z_i converges to some point $z_\infty \in J(p)$ and w_i converges to some point $w_\infty \in V'$.

Then by Lemma 3.9 and Lemma 3.18 we see that $\check{g}_{z_i+}(w_i, a_i) \rightarrow \check{g}_{z_\infty+}(w_\infty, 0)$ and $w_\infty \in U_{z_\infty} \times S_{r\delta} \subset V'$ and the gradient of the restriction of $\mathcal{G}_0^-$ to the stable manifold in B_{z_∞} is zero at the point $\check{g}_{z_\infty+}(w_\infty, 0)$ on the curve $\check{g}_{z_\infty+}(S_{r\delta}, 0)$. Since the sequence $\mathcal{G}_{a_k}^-(\check{g}_{z_k+}(\cdot, a_k))$ converges locally uniformly to $\mathcal{G}_0^-(\check{g}_{z_\infty+}(\cdot, 0))$ by Lemma 3.18, so derivatives of $\check{g}_{z_k+}(\cdot, a_k)$ converge locally uniformly to the derivatives of $\check{g}_{z_\infty+}(\cdot, 0)$, then because $\mathcal{G}_a^-$ is smooth on a neighborhood of the image of $\check{g}_{z_\infty+}(S_{r\delta}, 0)$ then the gradient of $\mathcal{G}_{a_k}^-$ projected to the tangent space of Δ_{z_k, a_k} at $\check{g}_{z_k+}(w_k, a_k)$ converges to the gradient of $\mathcal{G}_0^-$ projected to the tangent space of Δ_{z_k, a_k} at $\check{g}_{z_k+}(w_\infty, 0)$. This is a contradiction since the gradient of the restriction of $\mathcal{G}_0^- = \frac{1}{d} \log |v|$ to a vertical line does not vanish on the curve $|v| = r\delta$. $\square$

Lemma 4.11. *The index of the gradient of the restriction of $\mathcal{G}_a^-$ to $\check{g}_{z+}(\mathbb{D}_\delta)$ around the curve $\check{g}_{z+}(S_{r\delta})$ is one for all $|a| < \epsilon_S$.*

Proof. The lemma is easily seen to be true for $a = 0$ since then $\check{g}_{z+}(S_{r\delta})$ is a loop around a single singularity of $\frac{1}{d} \log |p(y) - x|$. By Lemma 4.10, the index can not change for $|a| < \epsilon_S$. $\square$

Lemma 4.12. *If a is sufficiently small then for any $z \in J(p)$ and for any $w \in p^{-1}(z)$ one has $f_a(\Delta_{w,a}(r)) \subseteq \Delta_{z,a}(r)$. Also $f_a(\Delta_{w_1,a}(r)) \cap f_a(\Delta_{w_2,a}(r)) = \emptyset$ for w_1 and w_2 distinct points of $p^{-1}(z)$.*

Proof. Using Lemma 4.9 it follows that if $(x, y) \in \overline{\Delta_{w,a}(r)}$ and $f_a(x, y) \in \overline{\Delta_{w,a}(r)} \setminus \Delta_{w,a}(r)$ then $|a| \geq \frac{(r\delta)^{1-1/d}}{\epsilon^{d+1}}$. The second statement is an immediate consequence of Lemma 3.17. $\square$

Lemma 4.13. *There exists $M > 0$ such that given $\epsilon > 0$ then for $|a| < \epsilon/M$ one has $|v(f_a(x, y))| < \epsilon$ for each $(x, y) \in \mathcal{T}_c^\bullet$ and for each critical point c of $p(x)$.*

Proof. One has $|v(f_a(x, y))| = |v(p(x) - ay, x)| = |ay|$. Choosing $M > 0$ such for each critical point c of $p(x)$ one has $\mathcal{T}_c^\bullet \subset (\mathbb{D}_M \times \mathbb{C})$ one concludes that taking $|a| < \frac{\epsilon}{M}$ is sufficient. $\square$

We will let $V'_r \equiv \{(x, y) \in V' \mid |v(x, y)| < r\}$.

Corollary 4.14. *If a is sufficiently small then for any $z \in J(p)$ the set $\overline{\Delta_{z,a}(r)} \setminus \bigcup_{w \in p^{-1}(z)} f_a(\Delta_{w,a}(r))$ contains no points of J_a .*

Proof. If a is sufficiently small then $J_a \subset V'_r$ and π_a is defined. Since $J_a \subset V'_r$ then given any $\underline{z} = (\dots, z_{-2}, z_{-1}, z_0) \in \hat{J}(p)$ then $\pi_a(\underline{z}) \in \Delta_{z,a}(r)$. From Lemma 3.17 we know that a point in J_a can lie in $\Delta_{z,a}$ iff the corresponding history in $\hat{J}(p)$ ends with the point z . Then the result is easy since the set removed from $\overline{\Delta_{z,a}(r)}$ contains all points in J_a corresponding to histories in $\hat{J}(p)$ which could end with z_0 . $\square$

Corollary 4.15. *There exists $\epsilon > 0$ such that if $|a| < \epsilon$ then the index of the vector field $\nabla(\mathcal{G}_a^-|_{\Delta_{z,a}})$ around the boundary of $\overline{\Delta_{z,a}(r)} \setminus \bigcup_{w \in p^{-1}(z)} f_a(\Delta_{w,a}(r))$ is $1 - d$.*

Proof. We note from Corollary 4.14 that $\mathcal{G}_a^- > 0$ on $\overline{\Delta_{z,a}(r)} \setminus \bigcup_{w \in p^{-1}(z)} f_a(\Delta_{w,a}(r))$ and so it will be pluriharmonic on a neighborhood of $\overline{\Delta_{z,a}(r)} \setminus \bigcup_{w \in p^{-1}(z)} f_a(\Delta_{w,a}(r))$. Thus $\nabla(\mathcal{G}_a^-|_{\Delta_{z,a}})$ will have only finitely many zeros in $\overline{\Delta_{z,a}(r)} \setminus \bigcup_{w \in p^{-1}(z)} f_a(\Delta_{w,a}(r))$. By Lemma 4.10 we know none of these zeros lie on the boundary.

The result then follows as long as ϵ is sufficiently small as a consequence of Lemma 4.11 and Lemma 4.12. $\square$

We recall [BS99] Proposition 2.7, noting that the hypothesis is satisfied for all a under consideration since f_a is hyperbolic when the crossed mapping construction of [HOV95] applies, and there is a continuous surjection from $\hat{J}(p)$ to J_a , and hence J_a is connected. Since J_a is connected and $|a| < 1$ then by Theorem 0.2 of [BS98b] it follows that f_a is unstably connected.

Proposition 4.16. *If f_a is hyperbolic and unstably connected, then the union of $\mathcal{F}_a^+$ and the stable lamination of $f_a|_{J_a}$ form a lamination of the space $U_a^+ \cup J_a^+$.*

Observation 4.17. For the maps we are studying, the union of $\mathcal{F}_a^-$ and the unstable lamination of $f_a|_{J_a}$ do not form a lamination of the space $U_a^- \cup J_a^-$. This is because critical points on the local stable manifolds are tangencies between the stable foliation and $\mathcal{F}_a^-$. Taking forward images of these tangencies gives accumulations of such tangencies near J_a . But J_a^- is transverse to J_a^+ everywhere since the map is hyperbolic, so the unstable foliation and $\mathcal{F}_a^-$ can't be part of the same foliation.

We let $K^\circ(p)$ denote the interior of the filled Julia set of $p(x)$.

Proposition 4.18. *For all sufficiently small nonzero a , given any $z \in J(p)$ then the only points of $\overline{\mathcal{C}_a}$ which lie in $f_a^{-1}(\Delta_{z,a}(r)) \setminus \bigcup_{w \in p^{-1}(z)} \Delta_{w,a}(r)$ are the points $\tau_{0a}(z, c) \in \tau_{0a}(\partial H_{c0}) = \partial H_{ca}$ where c is a critical point of $p(z)$.*

Additionally, when a and b are sufficiently small the biholomorphism $\tau_{ab}: H_{ca} \rightarrow H_{cb}$ defined in Section 4.1 extends naturally to a homeomorphism between $\overline{H_{ca}}$ and $\overline{H_{cb}}$. Since $\overline{H_{c0}}$ can be naturally identified with $\mathbb{C} \setminus K^\circ(p)$ then the same is true for $\overline{H_{ca}}$.

Proof. We choose ϵ small enough that:

1. $\epsilon \leq \epsilon_T$, so Lemma 4.2 holds for $|a| < \epsilon$,

2. $J_a \subset V'$ when $|a| < \epsilon$, which we can do by Lemma 3.19,
3. $\epsilon < r\delta/M$ for the value M in Lemma 4.13,
4. Lemma 4.12, Corollary 4.14 and Corollary 4.15 all hold for $|a| < \epsilon$.

We will show the result holds for $|a| < \epsilon$.

Given a critical point c of $p(x)$, we let $\tilde{H}_c = \{(x, y, a) | (x, y) \in H_{ca}, a \in \mathbb{D}_\epsilon\}$. Now given an arbitrary point $z \in \mathbb{C} \setminus K(p)$ we define the map $h_{cz}: \mathbb{D}_\epsilon \rightarrow \tilde{H}_c \subset \mathbb{C}^2 \times \mathbb{D}_\epsilon$ by $h_{cz}(a) \equiv (\tau_{0a}(z, c), a) \in \tilde{H}_c$. It follows from Proposition 4.6 that h_{cz} is holomorphic. One easily confirms that $\varphi_{a,+}(h_{cz}(a)) = \varphi_{0,+}(z, c)$ which is independent of a . It follows that

$$\mathcal{G}^+(h_{cz}(a)) \equiv \mathcal{G}_0^+(z, c) \text{ for } a \in \mathbb{D}_\epsilon. \quad (4.1)$$

We also note that by Corollary 1.15 there is some radius R' such that for each critical point c of $p(x)$, if $|x| \geq R'$, $(x, y) \in \overline{\mathcal{T}_c^\bullet}$ and $|a| < \epsilon$ then $\mathcal{G}_a^+(x, y) \geq 1$. Hence if $(x, y) \in H_{ca}$ and $\mathcal{G}_a^+(x, y) < 1$ then $(x, y) \in W_c \equiv \left(\overline{\mathcal{T}_c^\bullet} \cap (\mathbb{D}_{R'} \times \mathbb{C})\right) \times \mathbb{D}_\epsilon$. The set W_c is clearly a bounded set in $\mathbb{C}^3$ since the set $\mathbb{D}_{R'} \times \mathbb{C}$ has bounded x coordinates and the set $\overline{\mathcal{T}_c^\bullet}$ has bounded y coordinates.

We let $Y = \{z \in \mathbb{C} \setminus K(p) \mid \mathcal{G}_0^+(z, c) < 1\}$. If $z \in Y$ then $h_{cz}(a)$ lands in the set W_c since $\mathcal{G}^+(h_{cz}(a)) = \mathcal{G}_0^+(z, c) < 1$ for all $a \in \mathbb{D}_\epsilon$. Since W_c is bounded it follows that $\{h_{cz}|z \in Y\}$ is a normal family of maps from $\mathbb{D}_\epsilon$ into $\mathbb{C}^3$. The condition that $\mathcal{G}_0^+(z, c) < 1$ is the same as $\log|b_p(z)| < 1$ so $J(p) \times \{c\} \times \{0\} \subset \overline{Y}$.

To complete the proof of Proposition 4.18 we need two lemmas.

Lemma 4.19. *Assume z_k is a sequence of points of $\mathbb{C} \setminus K(p)$ converging to a point $z_\infty \in J(p)$. Then, for each critical point c , the limit g of any convergent subsequence of h_{cz_k} satisfies $g(a) \in f_a^{-1}(\Delta_{z_\infty, a}(r))$ for all $a \in \mathbb{D}_\epsilon$.*

Proof of Lemma 4.19. We will show that $f_a(g(a)) \in \Delta_{z_\infty, a}(r)$ for all $a \in \mathbb{D}_\epsilon$. By Lemma 4.13, since $|a| < \epsilon < r\delta/M$, then $|v(f_a(\mathcal{T}_c^\bullet))| < r\delta$. Recall that $V'_r = \{(x, y) \in V' \mid |v| < r\delta\} = U' \times \mathbb{D}_{r\delta}$. Now $\tilde{\Delta}_{z_\infty, a}(r) \equiv \{(x, y, a) | (x, y) \in \Delta_{z_\infty, a}(r), a \in \mathbb{D}_\epsilon\}$ is defined in $V'_r \times \mathbb{D}_\epsilon$ as the graph of $g_{z_\infty+}: \mathbb{D}_{r\delta} \times \mathbb{D}_\epsilon \rightarrow U'$. Thus $v - g_{z_\infty+}(u, a)$ is a holomorphic defining function for $\tilde{\Delta}_{z_\infty, a}(r) \subset V'_r \times \mathbb{D}_\epsilon$. Since $\mathcal{G}_a^+(h_{cz_k}(a)) > 0$ for each k , then $(v - g_{z_\infty+}(u, a)) \circ \tilde{f} \circ h_{cz_k}$ is nonvanishing on $\mathbb{D}_\epsilon$ for each k whenever it is defined, i.e. whenever $h_{cz_k} \in V'_r \times \mathbb{D}_\epsilon$. If g is the limit of any convergent subsequence, then $(v - g_{z_\infty+}(u, a)) \circ f_0 \circ g(0) = 0$. Since g maps $\mathbb{D}_\epsilon$ into $\{(x, y) \mid |v(x, y)| < r\delta\}$ it can be shown that the set on which $(v - g_{z_\infty+}(u, a)) \circ \tilde{f} \circ g$ vanishes is both open and closed in $\mathbb{D}_\epsilon$, so it is all of $\mathbb{D}_\epsilon$. Thus $g(a) \in f_a^{-1}(\Delta_{z_\infty, a}(r))$ for all $a \in \mathbb{D}_\epsilon$. $\square$

Lemma 4.20. *Assume z_k is a sequence of points of $\mathbb{C} \setminus K(p)$ converging to a point $z_\infty \in J(p)$. Then the limit g of any convergent subsequence of h_{cz_k} is disjoint from $\Delta_{w, a}(r)$ for each $w \in p^{-1}(z_\infty)$ for all $a \in \mathbb{D}_\epsilon$ and for each critical point c .*

Proof of Lemma 4.20. Each $g(a) \in \mathcal{T}_c^\bullet$ lies a positive distance from $|v| < r\delta$ by Definition 4.8. Since $\Delta_{w, a}(r) \subset V'_r$ this completes the proof. $\square$

Now consider an arbitrary critical point c_0 of $p(z)$. Consider also an arbitrary $a \in \mathbb{D}_\epsilon$ and a sequence of points $\{w_i\} \in H_{c_0a}$ which converge to a point $w_\infty \in \overline{H_{c_0a}} \cap (J_{a+} \cup J_{a-})$. Then let $(z_i, c_0) = \tau_{a0}(w_i) \in H_{c_00}$ and consider the sequence of maps $h_{cz_k} : \mathbb{D}_\epsilon \rightarrow \mathbb{C}^3$ for each critical point c .

This is a normal family. Choose some subsequence $h_{cz_{k_i}}$ so that $h_{cz_{k_i}}$ converges for each critical point c . For each critical point c let g_c be the limit of this subsequence. By Lemma 4.19, $g_c(a) \in f_a^{-1}(\Delta_{z_\infty, a}(r))$. By Lemma 4.2, $g_c(a) \notin J_a^-$, so $\mathcal{G}_a^-$ is smooth at $g_c(a)$. Since $g_c(a)$ is a limit of points of H_{ca_0} then by Proposition 4.16, $\mathcal{F}_a^-$ and J_a^+ are tangent at $g_c(a)$ and so $g_c(a)$ is a point where $\nabla(\mathcal{G}_a^-|_{f_a^{-1}(\Delta_{z_\infty, a}(r))})$ has nonzero index.

Now if $a \neq 0$ then since there are $d-1$ critical points of $p(x)$ then the set $\{g_c(a) | c \text{ a critical point}\}$ is a set of $d-1$ points of nonzero index in $f_a^{-1}(\Delta_{z_\infty, a}(r)) \setminus \bigcup_{w \in p^{-1}(z_\infty)} \Delta_{w, a}$.

Since the index of the gradient of the restriction of $\mathcal{G}_a^-$ is a vector field in $\Delta_{z_\infty, a}(r) \setminus \bigcup_{w \in p^{-1}(z_\infty)} f_a(\Delta_{w, a}(r))$ by Corollary 4.14, and the index around the boundary of this set is $1-d$ by Corollary 4.15, then the same is clearly true for $f_a^{-1}(\Delta_{z_\infty, a}(r)) \setminus \bigcup_{w \in p^{-1}(z_\infty)} \Delta_{w, a}(r)$. Thus each of the $d-1$ points $g_c(a)$ has index -1 and $g_c(a)$ is the unique point of nonzero index in the intersection of $f_a(\Delta_{z_\infty, a}(r)) \setminus \bigcup_{w \in p^{-1}(z_\infty)} \Delta_{w, a}(r)$ and the tube $\mathcal{T}_c^\bullet$. That the same holds for $a = 0$ is easy to verify directly.

It follows that given any critical point c of $p(z)$ then any convergent subsequence of h_{cz_k} must converge to g_c . It follows that $h_{cz_k} \rightarrow g_c$. We denote g_c by $h_{cz_\infty} : \mathbb{D}_\epsilon \rightarrow \mathbb{C}^3$. We have thus shown that, given a critical point c , if $z_k \in \mathbb{C} \setminus K(p)$ converges to $z_\infty \in J(p)$ then $h_{cz_k}(a) \equiv (\tau_{0a}(z, c), a) \in \tilde{H}_c$ converges to a holomorphic function g_c such that $g_c(a)$ is the unique point of nonzero index in the intersection of $\mathcal{T}_c^\bullet$ and $f_a^{-1}(\Delta_{z_\infty, a}(r)) \setminus \bigcup_{w \in p^{-1}(z_\infty)} \Delta_{w, a}(r)$.

We now construct the extension $\tau_{ab} : \overline{H_{ca}} \rightarrow \overline{H_{cb}}$ by defining $\tau_{ab}(w_\infty) = h_{cz_\infty}(b)$ whenever $w_\infty \in \overline{H_{ca}}$ and z_∞ is the limit of $\{\tau_{a0}(w_k)\}$ where $w_k \rightarrow w_\infty$. This is well defined because if $w_k \rightarrow w_\infty$ and $w'_k \rightarrow w_\infty$ then $w_1, w'_1, w_2, w'_2, \dots$ converges to w_∞ and by the above, the sequence of maps $h_{c\tau_{a0}(w_1)}, h_{c\tau_{a0}(w'_1)}, \dots$ converges to a single map h_{cz_∞} for $|a| < \epsilon$. This is also continuous since if $w_k \in \overline{H_{ca}}$ and w_k converges to $w_\infty \in \overline{H_{ca}}$ but $\tau_{ab}(w_k) \not\rightarrow \tau_{ab}(w_\infty)$ then there exists $\epsilon_1 > 0$ such that there are arbitrarily large values of k with $|\tau_{ab}(w_k) - \tau_{ab}(w_\infty)| > \epsilon_1$. Then replace each point w_k with a point $w'_k \in H_{ca}$ such that $|w'_k - w_k| < 1/2^k$ and $|\tau_{ab}(w'_k) - \tau_{ab}(w_k)| < \epsilon/2$ (which we can do by the definition of $\tau_{ab}(w_k)$ if $w_k \in \overline{H_{ac}} \setminus H_{ac}$ and we just take $w'_k = w_k$ otherwise). Then w'_k is a sequence in H_{ac} and $\tau_{ab}(w'_k)$ can not converge to $\tau_{ab}(w_\infty)$ because there are arbitrarily large k for which $|\tau_{ab}(w'_k) - \tau_{ab}(w_\infty)| > |\tau_{ab}(w_k) - \tau_{ab}(w_\infty)| - |\tau_{ab}(w'_k) - \tau_{ab}(w_k)| > \epsilon/2$ but $w'_k \rightarrow w_\infty$ since $|w'_k - w_k| < 1/2^k$. But this is a contradiction since $h_{c\tau_{a0}(w'_k)} \rightarrow h_{c\tau_{a0}(w_\infty)}$ by our previous work and $h_{c\tau_{a0}(w'_k)}(a) = w'_k$ and $h_{c\tau_{a0}(w'_k)}(b) = \tau_{ab}(w'_k)$ by definition. Therefore $\tau_{ab} : \overline{H_{ca}} \rightarrow \overline{H_{cb}}$ is continuous. Since τ_{ba} is clearly the inverse of τ_{ab} then τ_{ab} is a homeomorphism. This completes the proof of Proposition 4.16. $\square$

Theorem 4.21. *For all sufficiently small a every component of the critical locus is an iterate of one of the components H_c .*

Proof. The components of H_{ca} are the only components of the critical locus if $a = 0$, so assume $a \neq 0$. If W is a component of $\mathcal{C}_a$ then by Lemma 4.7 we know that ∂W contains at least one point w in either J_a^+ or J_a^- . If w lies in $J_a^+ \setminus J_a^-$ then w lies in the stable manifold of some point of J_a so there is some n' such that $f_a^{n'}(w) \in \Delta_{z', a}(r)$ for some $z' \in J(p)$.

Take n to be the smallest such n' (where n' is allowed to be negative) and take z to be the corresponding point of $J(p)$. We know a smallest such n exists since $w \in J_a^+ \setminus J_a^-$ so $\mathcal{G}_a^-(w) - \frac{1}{d-1} \log |a| > 0$ so $\mathcal{G}_a^-(f_a^{-k}(w)) - \frac{1}{d-1} \log |a| = d^n \cdot (\mathcal{G}_a^-(w) - \frac{1}{d-1} \log |a|) \rightarrow \infty$ as $k \rightarrow \infty$ but $\mathcal{G}_a^-$ is bounded on compact sets (since V_0 can certainly be assumed to be large enough to contain V').

It follows from our choice of n that $f_a^{\circ n}(w) \in \Delta_{z,a}(r) \setminus \bigcup_{y \in p^{-1}(z)} f_a(\Delta_{y,a}(r))$. But then from Proposition 4.18 and Lemma 4.2 we conclude that $f_a^{\circ n}(W)$ is an iterate of some H_c .

On the other hand if $w \in J_a^-$ then choose a sequence of points $w_i \in W$ such that $w_i \rightarrow w$. Then $\mathcal{G}_a^+(w_i) \rightarrow \mathcal{G}_a^+(w) = \mathfrak{g} \geq 0$, and $\mathcal{G}_a^-(w_i) - \frac{1}{d-1} \log |a| \rightarrow \mathcal{G}_a^-(w) - \frac{1}{d-1} \log |a| = 0$. Then for every $i \geq 1$ choose n_i such that $1 < \mathcal{G}_a^-(f_a^{-n_i}(w_i)) - \frac{1}{d-1} \log |a| \leq d$. By the recursion relation for $\mathcal{G}_a^-$ it follows that $n_i \rightarrow \infty$ as $i \rightarrow \infty$. Then $\mathcal{G}_a^+(f_a^{-n_i}(w_i)) \rightarrow 0$ as $i \rightarrow \infty$ so the sequence $f_a^{-n_i}(w_i)$ is a bounded sequence of points in iterates of $W \subset \mathcal{C}_a$ converging to J_a^+ . By Proposition 4.18 and Lemma 4.2 the members of any convergent subsequence of $f_a^{-n_i}(w_i)$ must lie in an iterate of H_c for all large i , so W must be an iterate of H_c . This completes the proof. $\square$

Part III

Rigidity

5 Holonomy in the Critical Locus.

We will now consider a single map $f = f_a$, $a \neq 0$ to which Theorem 4.21 applies and for which $|a| < \epsilon_T$ (so Lemma 4.2 and Theorem 4.5 apply). Since a is fixed we let $\psi_+ \equiv \varphi_{a,+}$ and $\psi_- \equiv \eta \cdot \varphi_{a,-} \circ f^{-1}$ where $\eta^{d-1} = 1/a$ and η is fixed. This gives the simpler relations $\psi_- \circ f^{-1} = \psi_-^d$ and $\psi_+ \circ f = \psi_+^d$. The function ψ_- is well defined and holomorphic on $f(V_-)$. We will show that ψ_- is a biholomorphism from a neighborhood of infinity in H_c to a neighborhood of infinity in $\mathbb{C}$. Because a is fixed we will omit it from the notation throughout the rest of this section.

Lemma 5.1. *There is a neighborhood H_c° of infinity in H_c which lies in $f(V_-)$ and such that ψ_- is a biholomorphism from H_c° to $\mathbb{C} \setminus \overline{\mathbb{D}}_\tau$ for some $\tau > 1$.*

Proof. First we note that

$$f(V_-) = \{(x, y) \mid |p(y) - x| > |ay| \text{ and } |p(y) - x| > |a|\alpha\}.$$

Since the y coordinate of points $(x, y) \in H_c$ remain bounded as $|x| \rightarrow \infty$, it follows that $(x, y) \in H_c$ implies $(x, y) \in f(V_-)$ whenever $|x|$ is sufficiently large. This implies the first assertion.

To show the second result we consider H_c to be lying in $\mathbb{C}^2 \subset \mathbb{P}^1 \times \mathbb{P}^1$. We then know that H_c can be completed to become a disk by adding the point (∞, c) and that its tangent space is given at (∞, c) by $p''(c)\mathbf{d}y + C\mathbf{d}u = 0$ where C is some constant and $u = \frac{1}{x}$. From Corollary 1.5 we obtain

$$B^{-1} < \left| \frac{\varphi_-(x, y)}{y} \right| < B$$

for $(x, y) \in V_-$. Now if $(x, y) \in f(V_-)$ then $f^{-1}(x, y) = \left(y, \frac{p(y) - x}{a}\right) \in V_-$ and so we obtain $B^{-1} \left| \frac{p(y) - x}{a} \right| < |\varphi_-(f^{-1}(x, y))| < B \left| \frac{p(y) - x}{a} \right|$. Since $\varphi_- \circ f^{-1} = \psi_-/\eta$ this becomes

$$B^{-1} \left| \eta \frac{p(y) - x}{a} \right| < |\psi_-(x, y)| < B \left| \eta \frac{p(y) - x}{a} \right|$$

for $(x, y) \in f(V_-)$. From Corollary 1.15 we have $B^{-1}|x| < |\psi_+(x, y)| < B|x|$ for $(x, y) \in V_+$. Dividing one equality by the other gives

$$B^{-2} \left| \eta \frac{p(y) - x}{ax} \right| < \left| \frac{\psi_-(x, y)}{\psi_+(x, y)} \right| < B^2 \left| \eta \frac{p(y) - x}{ax} \right|.$$

Since $\frac{p(y) - x}{x}$ is bounded on H_c as $|x| \rightarrow \infty$, by the Riemann extension theorem $\frac{\psi_-(x, y)}{\psi_+(x, y)}$ is holomorphic and nonzero on a neighborhood of $\infty \in H_c$. Hence by shrinking H_c° if necessary

we conclude that $\psi_-(x, y)$ is a biholomorphism from the neighborhood H_c° of ∞ onto $\mathbb{C} \setminus \overline{\mathbb{D}_\tau}$ for some large τ . $\square$

The following is shown in Proposition 6.2 of [HOV95].

Lemma 5.2. *There exists $\Re > 1$ such that for any $z \in \mathbb{C}$ with $|z| > \Re$ the fiber of ψ_+ in V_+ over z and the fiber of ψ_- in $f(V_-)$ over z are each analytic disks.*

Definition 5.3. Noting that $|\psi_+|$ and $|\psi_-|$ are well defined on all of U^+ and U^- respectively, we let

$$\mathcal{V}_+ \equiv \{z \in V_+ \mid |\psi_+(z)| > \Re\}$$

and we let

$$\mathcal{V}_- \equiv \{z \in V_- \mid |\psi_-(z)| > \Re\}.$$

We note that $f(\mathcal{V}_+) \subset \mathcal{V}_+$ and $f^{-1}(\mathcal{V}_-) \subset \mathcal{V}_-$ as follows from the recursion relations for ψ_+ and ψ_- .

The following is a consequence of the definitions:

Lemma 5.4. *Two points z_1 and z_2 are on the same leaf of $\mathcal{F}^+$ iff there exists $n \geq 0$ such that $f^{on}(z_1), f^{on}(z_2) \in \mathcal{V}_+$ and $\psi_+(f^{on}(z_1)) = \psi_+(f^{on}(z_2))$. Similarly, two points z_1 and z_2 are on the same leaf of $\mathcal{F}^-$ iff there exists $n \geq 0$ such that $f^{-n}(z_1), f^{-n}(z_2) \in \mathcal{V}_-$ and $\psi_-(f^{-n}(z_1)) = \psi_-(f^{-n}(z_2))$.*

Let $\mathcal{S} \equiv \{\omega \in \mathbb{C} \mid \omega^{d^n} = 1 \text{ for some } n \geq 0\} \cong \mathbb{Q}_d/\mathbb{Z}$. Given two points $z_1, z_2 \in U^+$, it is clear that the property $\frac{\psi_+(z_1)}{\psi_+(z_2)} \in \mathcal{S}$ is independent of the branches of ψ_+ used. Similarly for the property $\frac{\psi_-(z_1)}{\psi_-(z_2)} \in \mathcal{S}$, $z_1, z_2 \in U^-$.

Lemma 5.5. *Two points $z_1, z_2 \in U^+$ are on the same leaf of $\mathcal{F}^+$ iff $\frac{\psi_+(z_1)}{\psi_+(z_2)} \in \mathcal{S}$. Similarly, two points $z_1, z_2 \in U^-$ are on the same leaf of $\mathcal{F}^-$ iff $\frac{\psi_-(z_1)}{\psi_-(z_2)} \in \mathcal{S}$.*

Proof. If z_1 and z_2 lie in $\mathcal{V}_+$ the first assertion follows from Lemmas 5.2 and 5.4 and the recursion relationship for ψ_+ . Otherwise, choosing k such that $f^{ok}(z_1), f^{ok}(z_2) \in \mathcal{V}_+$ gives the first assertion. The second is analogous. $\square$

We will want to consider the holonomy maps of H_c determined by the foliations $\mathcal{F}^+$ and $\mathcal{F}^-$. By Theorem 4.5, H_c can be identified with $\mathbb{C} \setminus \overline{\mathbb{D}}$ using ψ_+ .

Assume that z_1 and z_2 are points of H_c for some critical point c of $p(z)$ and that z_1 and z_2 lie on the same leaf $\mathcal{L}^+(z_1)$ of $\mathcal{F}^+$. Start with $z = z_1$ and then vary $z \in H_c$. By Lemma 4.2, the leaves of $\mathcal{F}^+$ all intersect H_c transversely hence the intersection of $\mathcal{L}^+(z)$ with H_c which is near z_2 will vary holomorphically with z . By this means we get a holomorphic map h from a neighborhood $N_1 \subset H_c$ of z_1 to a neighborhood $N_2 \subset H_c$ of z_2 . Since the inverse map is given by starting with $z = z_2$, varying $z \in H_c$ and following the intersection of H_c and $\mathcal{L}^+(z)$ near z_1 then the holomorphic map from N_1 to N_2 is a biholomorphism for suitably chosen N_1 and N_2 . Let $\mathbf{m}_+ : N_1 \rightarrow N_2$ denote this biholomorphism.

Lemma 5.6. *In the coordinates on H_c given by $\psi_+ : H_c \rightarrow \mathbb{C} \setminus \overline{\mathbb{D}}$ the map $\mathbf{m}_+$ is given by $\mathbf{m}_+(z) = \omega z$ for some $\omega \in \mathcal{S}$. If we continue $\mathbf{m}_+$ by holonomy then $\mathbf{m}_+$ extends to the global automorphism of H_c given by $\mathbf{m}_+(z) = \omega z$. The same result holds on H_c° with coordinates defined by ψ_- , and using any two points $z_1, z_2 \in H_c^\circ$ on the same leaf of $\mathcal{F}^-$, along with $\mathcal{F}^-$, to construct a holonomy map $\mathbf{m}_-$.*

Proof. The function $\psi_+ : H_c \rightarrow \mathbb{C}$ is well defined and so $\frac{\psi_+(\mathbf{m}_+(z))}{\psi_+(z)}$ is a continuous function of z which takes values in $\mathcal{S}$. Hence it is constant. Thus the map $\mathbf{m}_+(z)$, which was only defined in a neighborhood of a point on H_c , takes the form $\mathbf{m}_+(z) = \omega z$ in the coordinates defined by ψ_+ , and thus such a holonomy map gives a global automorphism of H_c .

The result for H_c° is proven the same way. $\square$

One can attempt to picture this holonomy map in terms of monodromy. Assume $d = 2$. Because the Jacobian of f is very small, the set $f(H_c)$ looks approximately like the curve $C(p)$. Consequently, if $z \in \mathbb{C}$ is sufficiently large then there will be exactly two points in $f(H_c)$ which map to z under ψ_+ . The monodromy map $\mathbf{m} : z \rightarrow -z$, carried out on $f(H_c)$ instead of on H_c , interchanges such pairs. One visualizes a monodromy of a given order 2^n by looking at $f^{\circ n}(H_c) \cap V_+$ and considering the fibers of φ_+ in V_+ as we move in a large loop around K_+ along H_c .

6 Fiber Preserving Conjugacies.

6.1 Statement of the Theorem

Buzzard and Verma [BV01] proved a stability result by using the λ -lemma along the leaves of the foliation $\mathcal{F}^+$. We are interested in deforming the underlying manifold to obtain a new holomorphic self map. Unfortunately, simply deforming the leaves of $\mathcal{F}^+$ quasiconformally does not yield a well defined complex manifold structure as it destroys the complex structure transverse to the foliation. The first natural approach is to deform using both $\mathcal{F}^+$ and $\mathcal{F}^-$. However, we will show that typically Hénon maps can not be deformed in this way.

In what follows we consider Hénon maps $f(x, y) = (p(x) - ay, x)$ satisfying the following:

Condition 6.1.

- p is hyperbolic with connected Julia set and simple critical points.
- The Jacobian a is so small that the hypothesis of Theorem 4.5, Theorem 4.21, and Lemma 5.1 all hold.

Convention 6.2. We will be dealing with just two Hénon maps, f and g , in this section instead of a whole family f_a . Hence we will omit the subscript a , but will use a subscript of f or g whenever necessary, e.g. $\mathcal{F}_f^+$ and $\mathcal{F}_g^+$ instead of $\mathcal{F}_a^+$, or $\mathfrak{r}_f$ and $\mathfrak{r}_g$ instead of $\mathfrak{r}$.

Assume that we are given two different Hénon maps f and g arising from two such polynomials p_f and p_g and that f and g satisfy Condition 6.1. We will show that there are severe obstructions to the existence of a conjugacy between f and g on $U^+ \cup U^-$ which maps leaves of $\mathcal{F}_f^+$ and $\mathcal{F}_f^-$ to the leaves of $\mathcal{F}_g^+$ and $\mathcal{F}_g^-$ respectively.

The folling result can be found as Lemma 2.1 of [Buz99]. We include a proof.

Lemma 6.3. A homeomorphism from $U_f^+ \cup U_f^-$ to $U_g^+ \cup U_g^-$ which maps the leaves of $\mathcal{F}_f^+$ and $\mathcal{F}_f^-$ to the leaves of $\mathcal{F}_g^+$ and $\mathcal{F}_g^-$ respectively necessarily maps $\mathcal{C}_f$ to $\mathcal{C}_g$.

Proof. This is because two leaves $\mathcal{L}_1$ and $\mathcal{L}_2$ which are transverse in $\mathbb{C}^2$ intersect in a different manner topologically than two leaves which are not. To see this, choose convenient (x, y) coordinates so that the point of intersection is the origin, $\mathcal{L}_1$ coincides with the x axis, and $\mathcal{L}_2$ is transverse to the y axis. Then choose a biholomorphic parameterization $t \mapsto (g_1(t), g_2(t))$ of a neighborhood of the intersection in $\mathcal{L}_2$ such that $t = 0$ maps to the point of intersection.

Since the second leaf is transverse to the y axis then $g_1'(0) \neq 0$. Therefore g_1 is a local biholomorphism. It follows that we can write $g_2 = \theta g_1^d$ for some nonvanishing holomorphic function θ defined in some neighborhood of zero, where $d = 1$ iff $\mathcal{L}_1$ and $\mathcal{L}_2$ intersect transversely. Choosing ζ a holomorphic function such that $\zeta^{d-1} = \theta$, then $\zeta g_2 = (\zeta g_1)^d$ so $\mathcal{L}_2$ is parameterized by $(\zeta g_1, (\zeta g_2)^d)$. One can choose local coordinates so that that $\mathcal{L}_2$ is parameterized by (t, t^d) and $\mathcal{L}_1$ is the x axis. Then if U is any sufficiently small open neighborhood of the origin we see that the inclusion $\mathcal{L}_2 \setminus \{0\} \hookrightarrow U \setminus \mathcal{L}_1$ can induce a surjective map of fundamental groups iff $d = 1$. $\square$

Remark 6.4. We will assume that $h: U_f^+ \cup U_f^- \rightarrow U_g^+ \cup U_g^-$ is a homeomorphism which maps leaves of $\mathcal{F}_f^+$ to leaves of $\mathcal{F}_g^+$ and maps leaves of $\mathcal{F}_f^-$ to leaves of $\mathcal{F}_g^-$. Then by Lemma 6.3 h maps the critical locus $\mathcal{C}_f$ of f to the critical locus $\mathcal{C}_g$ of g . We assume that there are critical points c_f and c_g of f and g such that if we let H_f and H_g denote the components of $\mathcal{C}_f$ and $\mathcal{C}_g$ asymptotic to $y = c_f$ and $y = c_g$ respectively as $|x| \rightarrow \infty$ then h maps H_f to H_g . There is no loss of generality in assuming that h maps H_f to H_g since, by Theorem 4.21, we can always choose integers k and ℓ such that this assumption holds if we change coordinates by iterates of f and g so that in the new coordinates h becomes the map $g^{\circ k} \circ h \circ f^{\circ \ell}$.

There is one degenerate situation we wish to rule out. Since we are considering conjugacies $h: U_f^+ \cup U_f^- \rightarrow U_g^+ \cup U_g^-$ without requiring that h extends to $\mathbb{C}^2$ then it is possible that $h: H_f \rightarrow H_g$ “inverts” H_f , meaning that it maps neighborhoods of the puncture in $H_f \cong \mathbb{D}^*$ to open sets adjacent to the other boundary component of $\mathbb{D}^*$.

Condition 6.5. *The map h maps small neighborhoods of the puncture of H_f at infinity to neighborhoods of the puncture of H_g , where H_f and H_g are specified horizontal components of $\mathcal{C}_f$ and $\mathcal{C}_g$ respectively.*

Observation 6.6. Let $\mathbf{c}: \mathbb{C}^2 \rightarrow \mathbb{C}^2$ be the conjugation map $\mathbf{c}(x, y) = (\bar{x}, \bar{y})$, and let $g = \mathbf{c} \circ f \circ \mathbf{c}$. Then $\psi_{g+} = \overline{\psi_{f+} \circ \mathbf{c}}$ and $\psi_{g-} = \overline{\psi_{f-} \circ \mathbf{c}}$. Thus as a conjugacy between f and g , $\mathbf{c}$ maps the leaves of $\mathcal{F}_f^+$ and $\mathcal{F}_f^-$ to the leaves of $\mathcal{F}_g^+$ and $\mathcal{F}_g^-$ respectively.

Reduction 6.7. We can assume that $h: H_f \rightarrow H_g$ is orientation preserving by replacing g with $\mathbf{c} \circ g \circ \mathbf{c}$ and h with $\mathbf{c} \circ h$ if necessary.

Lemma 6.8. *In the coordinates on H_f and H_g given by ψ_{f+} and ψ_{g+} one has $h(\omega z) = \omega h(z)$ for any $\omega \in S^1$. The same property also holds for all $z \in H_f^\circ$ using the coordinates on H_f° and H_g° given by ψ_{f-} and ψ_{g-} ,*

Proof. We will conduct the proof for H_f and H_g with coordinates $\psi_{f+}: H_f \rightarrow \mathbb{C} \setminus \overline{D}$ and $\psi_{g+}: H_g \rightarrow \mathbb{C} \setminus \overline{D}$ given by Theorem 4.5. The proof for the other part of the lemma is the same using the coordinates given in Lemma 5.1. Given $\omega \in \mathcal{S}$ the function $\frac{h(\omega z)}{h(z)}$ only takes values in $\mathcal{S}$ so it is constant. Thus, given $\omega \in \mathcal{S}$ there exists $\theta_\omega \in \mathcal{S}$ such that $h(\omega z) = \theta_\omega h(z)$ for all $z \in H_f$. It follows that h is $\mathcal{S}$ -equivariant. Hence it maps circles (centered at the origin) to circles. Since h is a homeomorphism then the order of the points $z, \omega z, \omega^2 z, \dots, \omega^{d^n} z = z$ on the circle must be preserved or reversed. Thus $h(\omega z) = \omega h(z)$ or $h(\omega z) = \bar{\omega} h(z)$ for $\omega \in \mathcal{S}$ and hence for all $\omega \in S^1$. The latter possibility can be eliminated since h is orientation preserving. $\square$

We will write χ_{f+} , χ_{f-} , χ_{g+} and χ_{g-} for the maps $1/\psi_{f+}$, $1/\psi_{f-}$, $1/\psi_{g+}$ and $1/\psi_{g-}$. The point at infinity is a removable singularity for each of these maps, each of which sends this point to the origin. Hence $\chi_{f+}: H_f \rightarrow \mathbb{D}$, $\chi_{g+}: H_g \rightarrow \mathbb{D}$, $\chi_{f-}: H_f^\circ \rightarrow \mathbb{D}_{1/\tau_f}$ and $\chi_{g-}: H_g^\circ \rightarrow \mathbb{D}_{1/\tau_g}$ are all biholomorphisms.

Definition 6.9. We define $\mathfrak{h}: \mathbb{D} \rightarrow \mathbb{D}$ by $\mathfrak{h} = \chi_{g+} \circ h \circ \chi_{f+}^{-1}$ and we similarly define $\mathfrak{k}: \mathbb{D}_{1/\tau_f} \rightarrow \mathbb{D}_{1/\tau_g}$ by $\mathfrak{k} = \chi_{g-} \circ h \circ \chi_{f-}^{-1}$. These maps satisfy $\mathfrak{h}(\omega z) = \omega^{\pm 1} \mathfrak{h}(z)$ and $\mathfrak{k}(\omega z) = \omega^{\pm 1} \mathfrak{k}(z)$, where the $\pm$ has the same sign in both relationships.

Definition 6.10. Let $\sigma_g = \chi_{g-} \circ \chi_{g+}^{-1} : \chi_{g+}(H_g^\circ) \rightarrow \mathbb{D}_{1/\tau_g}$ and let $\sigma_f = \chi_{f-} \circ \chi_{f+}^{-1} : \chi_{f+}(H_f^\circ) \rightarrow \mathbb{D}_{1/\tau_f}$. The maps σ_g and σ_f are the biholomorphic transition maps between the coordinate systems in which $\mathfrak{h}$ and $\mathfrak{k}$ represent the map h .

It is easy to confirm that

$$\sigma_g \circ \mathfrak{h} = \mathfrak{k} \circ \sigma_f. \quad (6.1)$$

Moreover, both $\mathfrak{k} \circ \sigma_f$ and $\sigma_g \circ \mathfrak{h}$ are defined on $\Omega_f \equiv \chi_{f+}(H_f^\circ)$ and map this homeomorphically onto $\mathbb{D}_{1/\tau_g}$.

We thus have the following commutative diagram:

$$\begin{array}{ccc} \mathbb{D}_{1/\tau_f} & \xrightarrow{\mathfrak{k}} & \mathbb{D}_{1/\tau_g} \\ \uparrow \chi_{f-} & & \uparrow \chi_{g-} \\ \sigma_f \swarrow & H_f^\circ \xrightarrow{h} H_g^\circ & \searrow \sigma_g \\ \downarrow \chi_{f+} & & \downarrow \chi_{g+} \\ \Omega_f & \xrightarrow{\mathfrak{h}} & \Omega_g \end{array} \quad (6.2)$$

Proposition 6.11. *One of the following must hold:*

1. $\sigma_f(z) = \beta z$ and $\sigma_g(z) = \gamma z$ for constants $\beta, \gamma \in \mathbb{C}^*$

or

2. *there is a neighborhood of the origin about which $\mathfrak{h}(z) = \beta z$ and $\mathfrak{k}(z) = \gamma z$ for constants $\beta, \gamma \in \mathbb{C}^*$.*

In the next section we will derive Proposition 6.11 from general properties of holomorphic circle actions.

6.2 Holomorphic circle actions

Let us consider the circle $\mathbb{R}/\mathbb{Z}$ acting faithfully on a neighborhood U of 0 by biholomorphic maps fixing 0, i.e., we have a monomorphism $t \mapsto \gamma^t$ from $\mathbb{R}/\mathbb{Z}$ to the group of biholomorphic maps of U such that $\gamma^t(0) = 0$. We call it briefly a *holomorphic circle action*. The action $\gamma^t : z \mapsto e^{2\pi i t} z$ will be called *standard*.

Remark 1 (straightening). *Any holomorphic circle action is conformally conjugate to the standard one (where the conjugacy can be orientation reversing).* Indeed, take some orbit Γ of the action. It is a topological circle that bounds a topological disk $V \ni 0$. Uniformize V by the round disk, $h : V \rightarrow \mathbb{D}$. Then the maps $\tilde{\gamma}^t = h \circ \gamma^t \circ h^{-1}$ are holomorphic automorphisms of $\mathbb{D}$ fixing 0, so they are rotations. Thus, we obtain a monomorphism $\mathbb{R}/\mathbb{Z} \rightarrow \mathbb{T}$, where $\mathbb{T}$ is the group of rotations. There are only two such monomorphisms, $t \mapsto e^{\pm 2\pi i t}$, and they are conjugate by the reflection $z \mapsto \bar{z}$.

The orbits of any holomorphic circle action form an analytic foliation in the punctured neighborhood U^* of 0.

Remark 2 (tangencies) *Given a pair (γ^t, ρ^s) of holomorphic circle actions, the associated foliations either coincide (and in this case the actions either coincide or conjugate by a conformal reflection⁵) or have finitely many tangencies on any given orbit. Indeed, if the set of tangencies between the two foliations is not finite on some orbit, then they have a common leaf (since they are analytic), and hence can be simultaneously straightened.*

Given two circle actions γ^t and $\tilde{\gamma}^t$, a local homeomorphism h near 0 is called $(\gamma, \tilde{\gamma})$ -equivariant if $h(\gamma^t z) = \tilde{\gamma}^t h(z)$ or $h(\gamma^t z) = \tilde{\gamma}^{-t} h(z)$.

Proposition 6.12. *Let (γ^t, ρ^s) and $(\tilde{\gamma}^t, \tilde{\rho}^s)$ be two non-trivial pairs of holomorphic circle actions. If a local homeomorphism h is both $(\gamma, \tilde{\gamma})$ - and $(\rho, \tilde{\rho})$ -equivariant, then h is holomorphic.*

Proof. Without loss of generality, we can assume that h conjugates γ^t to $\tilde{\gamma}^t$ (otherwise, compose h with an appropriate conformal reflection).

Let us take some orbit Γ of the γ -action. By the above Remark 2, we can pick a point $z \in \Gamma$ such that Γ is transverse to the ρ -orbit at z , and the orbits of $\tilde{\gamma}$ and $\tilde{\rho}^s$ are transverse at $\tilde{z} = h(z)$. Then the map $(t, s) \mapsto (\rho^s(z), \gamma^t(\rho^s z))$ gives a smooth local chart near Γ . If h conjugates ρ^s to $\tilde{\rho}^s$, let us consider the similar local chart near $\tilde{\Gamma} = h(\Gamma)$. Otherwise, let us consider the chart $(t, s) \mapsto (\tilde{\rho}^{-s}(z), \tilde{\gamma}^t(\tilde{\rho}^{-s} z))$. In either case, the map h becomes the identity in these coordinates. Hence h is smooth near Γ , and thus, it is smooth in a punctured neighborhood of 0.

Let us now show that the joint (γ, ρ) -action is transitive on the punctured neighborhood U^* of 0, i.e., any two points in U^* can be connected by a concatenation of pieces of the γ - and ρ -orbits. Indeed, the orbits of the joint action are open since the domain of any local chart described above is contained in one orbit. Since U^* is connected, it must be a single orbit.

Let us now consider the conformal structure $\mu = h^*(\lambda)$ in U , where λ is the standard conformal structure. The structure μ is represented by a smooth family of infinitesimal ellipses in U^* . Moreover, since λ is invariant under $\tilde{\gamma}$ and $\tilde{\rho}$ and h is equivariant, μ is invariant under the joint (γ, δ) -action.

If the structure μ is standard then h is holomorphic. Otherwise, there is a non-circular ellipse $\mu(z)$. Since μ is invariant under the transitive (γ, ρ) -action, all the ellipses are non-circular on U^* . Hence the big axes $l(z) \subset T_z U^*$ of the ellipses are well defined on U^* and form a (γ, ρ) -invariant line field over U^* .

Let us take some ρ -orbit Δ_0 and consider the outermost γ -orbit Γ_0 crossing Δ . Then Γ_0 and Δ_0 have tangency of even order at some point z_0 . Rotating the line field by an appropriate angle, we can make $l(z_0)$ tangent at z_0 to both Γ_0 and Δ_0 . By invariance, $l(z)$ is then tangent to Δ_0 at any point $z \in \Delta_0$.

Let us now consider a nearby γ -orbit Γ which is closer to the origin than Γ_0 . Then Γ intersects Δ_0 transversally at two points z_+ and z_- near z_0 . Moreover, the angles $\alpha_{\pm} \in (-\pi/2, \pi/2)$ of these intersections have opposite signs. This contradicts to the invariance of the line field under the γ -action.

□

⁵We will refer to such a pair as *trivial*.

Corollary 6.13. *Under the circumstances of the above Proposition, if the γ - and $\tilde{\gamma}$ -actions are standard, then h is linear, $z \mapsto \lambda z$.*

Proof. In this case, h preserves the foliation by round circles centered at 0. But a biholomorphic map that fixes 0 and maps a circle centered at 0 to another such a circle is linear. $\square$

Proof of Proposition 6.11. We know that the maps $\mathfrak{h}$ and $\mathfrak{k}$ are equivariant with respect to the standard circle action γ^t . Let $\rho^s = \sigma_f^{-1} \circ \gamma^s \circ \sigma_f$ and $\tilde{\rho}^s = \sigma_g^{-1} \circ \gamma^s \circ \sigma_g$. By (6.1), $\mathfrak{h}$ is $(\rho, \tilde{\rho})$ -equivariant. If the maps σ_f and σ_g are not linear, then the pairs of actions, (γ, ρ) and $(\gamma, \tilde{\rho})$, are non-trivial. Then $\mathfrak{h}$ is linear by the last Corollary.

The same argument applies to $\mathfrak{k}$. $\square$

6.3 Rigidity Results

Here we translate the statement of Proposition 6.11 back into statements about the two given maps f and g which have a conjugacy h between them which maps leaves of $\mathcal{F}_f^+$ to leaves of $\mathcal{F}_g^+$ and maps leaves of $\mathcal{F}_f^-$ to leaves of $\mathcal{F}_g^-$.

Theorem 6.14. *Assume we are given two different Hénon maps f and g satisfying Condition 6.1. Assume $h: U_f^+ \cup U_f^- \rightarrow U_g^+ \cup U_g^-$ is a conjugacy between f and g such that h maps the leaves of $\mathcal{F}_f^+$ and $\mathcal{F}_f^-$ to the leaves of $\mathcal{F}_g^+$ and $\mathcal{F}_g^-$ respectively. Choose coordinates for the map $g: \mathbb{C}^2 \rightarrow \mathbb{C}^2$ so that h maps H_f to H_g for a pair of primary horizontal critical components H_f and H_g of f and g . Finally assume that $h: H_f \rightarrow H_g$ is orientation preserving and satisfies Condition 6.5. Then $h: U_f^+ \cap U_f^- \rightarrow U_g^+ \cap U_g^-$ is a biholomorphism. Also, $\psi_{g+} \circ h = \frac{1}{\beta} \psi_{f+}$ and $\psi_{g-} \circ h = \frac{1}{\gamma} \psi_{f-}$ on a neighborhood about infinity of H_f where β^{d-1} and γ^{d-1} must lie in $\mathcal{S}$.*

Corollary 6.15. *Assume we are given two different Hénon maps f and g satisfying Condition 6.1. Assume $h: U_f^+ \cup U_f^- \rightarrow U_g^+ \cup U_g^-$ is a conjugacy between f and g such that h maps the leaves of $\mathcal{F}_f^+$ and $\mathcal{F}_f^-$ to the leaves of $\mathcal{F}_g^+$ and $\mathcal{F}_g^-$ respectively. Choose coordinates for the map $g: \mathbb{C}^2 \rightarrow \mathbb{C}^2$ so that h maps H_f to H_g for a pair of primary horizontal critical components H_f and H_g of f and g . Finally assume that $h: H_f \rightarrow H_g$ is orientation reversing and satisfies Condition 6.5. Then $\mathfrak{c} \circ h: U_f^+ \cap U_f^- \rightarrow \mathfrak{c}(U_g^+ \cap U_g^-)$ is a biholomorphism. Also, $\overline{\psi_{g+} \circ h} = \frac{1}{\beta} \psi_{f+}$ and $\overline{\psi_{g-} \circ h} = \frac{1}{\gamma} \psi_{f-}$ on a neighborhood about infinity of H_f where β^{d-1} and γ^{d-1} must lie in $\mathcal{S}$.*

Proof. Replace g with $\mathfrak{c} \circ g \circ \mathfrak{c}$ and h with $\mathfrak{c} \circ h$. $\square$

Proof of Theorem. Applying Lemma 6.3 we see the critical locus of f maps by h to the critical locus of g . As in Remark 6.4 we can change coordinates using iterates of f and g so there are primary horizontal components H_f and H_g such that h maps H_f to H_g . Then Proposition 6.11 gives the following cases:

Case 1. $\sigma_f(z) = \beta z$ and $\sigma_g(z) = \gamma z$. If we write these two equations out using Definition 6.10 we obtain $\psi_{f-} = \frac{1}{\beta}\psi_{f+}$ and $\psi_{g-} = \frac{1}{\gamma}\psi_{g+}$. Now while $\psi_{f+}: U_f^+ \rightarrow \mathbb{C}^*$ is not well defined, the map to the quotient group $\psi_{f+}: U_f^+ \rightarrow \mathbb{C}^*/\mathcal{S}$ is well defined. Similarly for $\psi_{f-}: U_f^- \rightarrow \mathbb{C}^*/\mathcal{S}$.

Now there exists some $M > 0$ such that if $z \in H_f$ and $\mathcal{G}_f^+(z) > M$ then $\psi_{f-}(z) = \frac{1}{\beta}\psi_{f+}(z)$.

Now if $z \in U_f^+$ and $\mathcal{G}_f^+(z) > M$ then there is some leaf $\ell \in \mathcal{F}_f^+$ containing both z and some point $w \in H_f$. Thus $\mathcal{G}_w^+ > M$ so $\psi_{f-}(z) \sim \psi_{f-}(w) = \frac{1}{\beta}\psi_{f+}(w) \sim \frac{1}{\beta}\psi_{f+}(z)$

But then if $z \in U_f^+$ and $\mathcal{G}_z^+ > M$ then

$$\frac{1}{\beta}\psi_{f+}(z) \sim \psi_{f-}(z) \sim \psi_{f-}^d \circ f(z) \sim \frac{1}{\beta^d}\psi_{f+}^d \circ f(z) = \frac{1}{\beta^d}\psi_{f+}^{d^2}(z). \quad (6.3)$$

Hence $\beta^{d-1} \sim \psi_{f+}^{d^2-1}(z)$. However since $\psi_{f+}^{d^2-1}(z)$ has a locally continuous branch about z one concludes that this branch would have to be constant, and hence $\psi_{f+}(z)$ must be constant. This is a contradiction. Thus no Henon map f exists for which $\psi_{f-} = \frac{1}{\beta}\psi_{f+}$ for a neighborhood of infinity in H_f .


Case 2. $\mathfrak{h}(z) = \beta z$ and $\mathfrak{k}(z) = \gamma z$. If we write these two equations out using Definition 6.9 then $\psi_{g+} \circ h = \frac{1}{\beta}\psi_{f+}$ and $\psi_{g-} \circ h = \frac{1}{\gamma}\psi_{f-}$. These hold in a neighborhood of infinity in H_f . Now, as in Case 1, there exists some $M > 0$ such that if $z \in U_f^+$ and $\mathcal{G}_f^+(z) > M$ then $\psi_{g+} \circ h(z) \sim \frac{1}{\beta}\psi_{f+}(z)$ in $\mathbb{C}^*/\mathcal{S}$. Then

$$\begin{aligned} \frac{1}{\beta}\psi_{f+}^d(z) &= \frac{1}{\beta}\psi_{f+} \circ f(z) \sim \psi_{g+} \circ h \circ f(z) = \\ &\psi_{g+} \circ g \circ h(z) \sim \psi_{g+}^d \circ h(z) \sim \frac{1}{\beta^d}\psi_{f+}^d(z) \end{aligned} \quad (6.4)$$

from which it follows that $\beta^{d-1} \in \mathcal{S}$.

We will show that $h: U_f^+ \cap U_f^- \rightarrow U_g^+ \cap U_g^-$ is a biholomorphism. To do this it suffices to show that $h: U_f^+ \cap U_f^- \rightarrow U_g^+ \cap U_g^-$ is holomorphic. To accomplish this it will suffice to show that if $z \in U_f^+ \cap U_f^-$ then h is holomorphic on $\mathcal{L}_f^+(z)$ and on $\mathcal{L}_f^-(z)$. From this it will immediately follow from Osgood's theorem that h is holomorphic on $(U_f^+ \cap U_f^-) \setminus \mathcal{C}$. Since $h: U_f^+ \cap U_f^- \rightarrow U_g^+ \cap U_g^-$ is continuous then by the Riemann extension theorem it will further follow that $h: U_f^+ \cap U_f^- \rightarrow U_g^+ \cap U_g^-$ is holomorphic. In order to prove this we will need the following.

Lemma 6.16. *Given $z_0 \in (U_f^+ \cap U_f^-) \setminus \mathcal{C}_f$, assume that $\mathcal{L}_f^+(z_0)$ meets H_f at a point w_0 . Then there is a neighborhood U of z_0 in $\mathcal{L}_f^-(z_0)$ such that there is a holomorphic holonomy map $\zeta_{f,z_0,w_0}: U \rightarrow H_f$ which maps $z \in U$ to the intersection of $\mathcal{L}_f^+(z)$ and H_f near w_0 . Since $\mathcal{F}_f^+$ is transverse to $\mathcal{L}_f^-(z_0)$ and H_f at z_0 and w_0 respectively then ζ_{f,z_0,w_0} can be assumed to be a biholomorphism from U onto its image.*


 Figure 2: Holonomy using $\mathcal{F}_f^+$.

Proof. This is geometrically self evident. $\square$

Now assume $z_0 \in (U_f^+ \cap U_f^-) \setminus \mathcal{C}_f$ and $\mathcal{G}_f^+(z_0) > M$ and $\mathcal{G}_g^+(h(z_0)) > M$. Choose $w_0 \in \mathcal{L}_f^+(z_0) \cap H_f$. Then if U is a sufficiently small open neighborhood of z_0 in $\mathcal{L}_f^-(z_0)$ then one can see that

$$h|_U = \zeta_{g, h(z_0), h(w_0)}^{-1} \circ h \circ \zeta_{f, z_0, w_0} \quad (6.5)$$

because h maps H_f to H_g , maps $\mathcal{L}_f^-(z_0)$ to $\mathcal{L}_g^-(h(z_0))$ and maps leaves of $\mathcal{F}_f^+$ to leaves of $\mathcal{F}_g^+$. But because h in the right hand side of equation (6.5) is applied on H_f near infinity as long as U is sufficiently small, then it is holomorphic. Thus equation (6.5) represents $h|_U$ as a composition of three holomorphic functions. Hence $h|_U$ is holomorphic.

Now in general, if z_0 is any point in $(U_f^+ \cap U_f^-) \setminus \mathcal{C}_f$ then choosing n sufficiently large that $\mathcal{G}_f^+(f^{on}(z_0)) > M$ and $\mathcal{G}_g^+(h(f^{on}(z_0))) > M$ and writing $h = g^{-n} \circ h \circ f^{on}$ one concludes that h is holomorphic on $\mathcal{L}_f^-(z_0)$ near z_0 .

The proof that h is holomorphic on leaves of $\mathcal{F}_f^-$ on $U_f^+ \cap U_f^-$ is identical. This completes the proof. $\square$

Eliminating h from the conclusion of Theorem 6.14 we obtain

$$\gamma \psi_{g-} \circ \psi_{g+}^{-1}(z) = \psi_{f-} \circ \psi_{f+}^{-1}(\beta z) \quad (6.6)$$

for some nonzero constants β and γ .

We now assume that f has degree two, so f and g are Hénon maps of the form $f(x, y) = (x^2 + c_1 - a_1 y, x)$ and $g(x, y) = (x^2 + c_2 - a_2 y, x)$. Rewriting equation (6.6) in terms of $\chi_{f+}, \chi_{f-}, \chi_{g+}$ and χ_{g-} we obtain

$$\chi_{g-} \circ \chi_{g+}^{-1}(\beta z) = \gamma \cdot \chi_{f-} \circ \chi_{f+}^{-1}(z). \quad (6.7)$$

Our first goal will be to find all possible (nondegenerate) choices of $f(x, y) = (x^2 + c_1 - a_1 y, x)$, $g(x, y) = (x^2 + c_2 - a_2 y, x)$ and β and γ nonzero constants so that equation (6.7) holds.

The first three nonzero terms of the Taylor series expansion of

$$\chi_{g-} \circ \chi_{g+}^{-1}(\beta z) - \gamma \cdot \chi_{f-} \circ \chi_{f+}^{-1}(z) \quad (6.8)$$

are

$$(\gamma a_1^2 - \beta a_2^2)z + (\gamma a_1^2 c_1 - \beta^2 a_2^2 c_2)z^2 + (\gamma a_1^2 c_1^2 - \beta^3 a_2^2 c_2^2 + \frac{1}{2}\gamma a_1^2 c_1 - \frac{1}{2}\gamma a_1^4 c_1 + \frac{1}{2}\beta^3 a_2^4 c_2 - \frac{1}{2}\beta^3 a_2^2 c_2)z^3 \quad (6.9)$$

Since we can assume that each of a_1, a_2, β, γ are nonzero, it easily follows from the first two terms that

$$\gamma = \frac{a_2^2}{a_1^2}\beta, \quad c_1 = c_2\beta. \quad (6.10)$$

Lemma 6.17. *If $\chi_{g-} \circ \chi_{g+}^{-1}(z) = \chi_{f-} \circ \chi_{f+}^{-1}(z)$ for nonsingular*

$$f(x, y) = (x^2 + c_1 - a_1 y, x) \quad \text{and} \quad g(x, y) = (x^2 + c_2 - a_2 y, x)$$

then $f = g$.

Proof. From (6.10) it follows that $a_1 = \pm a_2$ and $c_1 = c_2$. To eliminate the remaining case assume $a_1 = -a_2$. Now (6.8) must vanish, but the coefficients of z^4, z^5 and z^7 in (6.8) generate the ideal (a_1^3) , so a_1 would have to be zero. $\square$

From the third term of (6.8) one obtains that either $a_2 = 1, a_2 = -1, c_1 = 0$, or $\beta = \frac{a_1^2 - 1}{a_2^2 - 1}$.

Each of the cases $a_2 = 1, a_2 = -1$, and $\beta = \frac{a_1^2 - 1}{a_2^2 - 1}$ can be reduced to the case $c_1 = 0$ by taking more terms of the Taylor series of (6.8) and the calculating Groebner basis of the resulting coefficients. In the case $c_1 = 0$ it can be shown that $c_2 = 0$ and that $a_1 = a_2$ (so f and g are the same Hénon map), and that $\beta = \gamma$, both of which must be the same primitive root of unity.

The following table gives the necessary details to verify these calculations using $\textcircled{R}$ Maple. Here n is the degree to which we need to calculate the Taylor series for each calculation.

Case	n	Ordering	Relevant Basis Elements	Conclusion
$\beta = \frac{a_1^2 - 1}{a_2^2 - 1}$	7	$\text{tdeg}(a_1, c_1, a_2)$	$-a_2^2 c_1 (a_1^2 - 1) \cdot (a_2^2 - 1)^5 (a_1 - a_2)$	$c_1 = 0$
$a_2 = 1$	8	$\text{plex}(\beta, c_1, a_1)$	$\beta c_1 (a_1 - 1), \beta c_1 (\beta - 1)$	$c_1 = 0$
$a_2 = -1$	8	$\text{plex}(\beta, c_1, a_1)$	$\beta c_1 (a_1 + 1), \beta c_1 (\beta - 1)$	$c_1 = 0$
$c_1 = 0$	13	$\text{tdeg}(\beta, a_2, a_1)$	$\beta a_1 a_2^2 (a_1 - a_2), \beta a_1 a_2^2 (s^3 - 1)$	$\beta^2 + \beta + 1 = 0$ $\gamma = \beta$ $a_1 = a_2$ $c_1 = c_2 = 0$

We conclude that:

Lemma 6.18. *If equation (6.6) holds with β and γ nonzero and $f(x, y) = (x^2 + c_1 - a_1y, x)$ and $g(x, y) = (x^2 + c_2 - a_2y, x)$ nondegenerate then the maps f and g must be the same Henon map and either (1) $\beta = \gamma = 1$ (the trivial solution), or else (2) $a_1 = a_2$, $c_1 = c_2 = 0$, $\beta = \gamma$ and β is a primitive cubic root of unity.*

Theorem 6.19. *Assume h is a conjugacy between quadratic Hénon maps $f(x, y) = (x^2 + c_1 - a_1y, x)$ and $g(x, y) = (x^2 + c_2 - a_2y, x)$ satisfying Condition 6.1. Assume further that h maps the leaves of $\mathcal{F}_f^+$ and $\mathcal{F}_f^-$ to the leaves of $\mathcal{F}_g^+$ and $\mathcal{F}_g^-$ respectively. Choose coordinates for the map $g: \mathbb{C}^2 \rightarrow \mathbb{C}^2$ so that h maps H_f to H_g for the primary horizontal critical components H_f and H_g of f and g . Finally assume that $h: H_f \rightarrow H_g$ is orientation preserving and satisfies Condition 6.5. Then $f = g$ and $h^{\circ 2}(x, y)$ is the identity map on $U_f^+ \cap U_f^-$.*

Proof. This is easy now, as from Theorem 6.14 $\beta \in \mathcal{S}$. By Lemma 6.18 the maps f and g must be the same Henon map, and additionally $\beta = \gamma = 1$ since primitive cubic roots of unity do not belong to $\mathcal{S}$.

To see that conclusion about $h^{\circ 2}$ one notes that since $\beta = \gamma = 1$ then $h|_{H_f}$ is the identity map. If we define $\theta(x, y) = (\psi_{f+}(x, y), \psi_{g+}(x, y))$ then about any sufficiently large point $z_0 \in H_f$ and the Jacobian of θ is a defining function for H_f about z_0 . Since H_f has multiplicity one then θ is locally a two to one map about H_f . Now $\theta(h(z)) = \theta(z)$ since $\beta = \gamma = 1$. One concludes that about such a point z_0 (which is fixed by h) h must either exchange points in the fibers of θ or must leave them fixed. Either way, $h^{\circ 2}$ must fix all points in a neighborhood of z_0 . Since $h^{\circ 2}$ is the identity map on an open set, it must be the identity map everywhere. $\square$

Corollary 6.20. *Assume h is a conjugacy between quadratic Hénon maps $f(x, y) = (x^2 + c_1 - a_1y, x)$ and $g(x, y) = (x^2 + c_2 - a_2y, x)$ satisfying Condition 6.1. Assume further that h maps the leaves of $\mathcal{F}_f^+$ and $\mathcal{F}_f^-$ to the leaves of $\mathcal{F}_g^+$ and $\mathcal{F}_g^-$ respectively. Choose coordinates for the map $g: \mathbb{C}^2 \rightarrow \mathbb{C}^2$ so that h maps H_f to H_g for the primary horizontal critical components H_f and H_g of f and g . Finally assume that $h: H_f \rightarrow H_g$ is orientation reversing and satisfies Condition 6.5. Then $f = \mathbf{c} \circ g \circ \mathbf{c}$ and $\mathbf{c} \circ h \circ \mathbf{c} \circ h$ is the identity map on $U_f^+ \cap U_f^-$.*

List of Notations

We provide a list of notations here as a reference.

Notatation	Section	Meaning
f_a		The Hénon map under consideration.
a		The Jacobian of f_a .
$p(x)$		A monic polynomial used to define f_a .
$p(x, y)$	1.4	The homogeneous version of $p(x)$.
d		The degree of $p(x)$.
$q(x)$	1.1	The polynomial $p(x)$ with its leading term removed.
d'	1.1	The degree of $q(x)$.
R	1.1	A large radius used to define V_+ and V_- .
V_+, V_-	1.1	Regions which describe the large scale behaviour of Hénon maps.
K_a^+, K_a^-		The set of points whose orbit under forward (respectively backward) iteration under f_a remain bounded.
J_a^+, J_a^-		The boundaries of K_a^+ and K_a^- respectively.
J_a		The intersection of J_a^+ and J_a^- .
U_a^+, U_a^-	1.4	The set of points whose orbit under forward (respectively backward) iteration under f_a remain bounded.
$\tilde{U}^+, \tilde{U}^-$	1.4	These are subsets of $\mathbb{C}^2 \times \mathbb{D}_R$ whose restriction to $\mathbb{C}^2 \times \{a\}$ is U_a^+ and U_a^- respectively.
$C(p)$		The curve $x = p(y)$ in $\mathbb{C}^2$. This equal to J_0^- .
$\mathbb{D}_r$		The disk of radius r in $\mathbb{C}$.
$\hat{V}_-, \hat{V}_+$	1.2, 1.3	The regions V_+ and V_- extended to $\mathbb{P}^1 \times \mathbb{P}^1$.
$\varphi_{a,+}, \varphi_{a,-}$	1.1	Holomorphic functions defined on V_+ and V_- respectively measuring that rate of escape to infinity under forward and backward iteration respectively.
$\mathcal{G}_a^+, \mathcal{G}_a^-$	1.4	Plurisubharmonic functions on $\mathbb{C}^2$ measuring the rate of escape to infinity under forward and backward iteration respectively.
s_k^-, s_k^+	1.2, 1.3	Auxilliary functions used to construct and study $\varphi_{a,+}$ and $\varphi_{a,-}$.
x_n, y_n	1.1	These are given by $(x_n, y_n) \equiv f_a^{\circ n}(x, y)$, $n \in \mathbb{Z}$.
$\mathcal{C}_a$	2.3	This is the critical locus of f_a . It tangency locus of the foliations $\mathcal{F}_a^+$ and $\mathcal{F}_a^-$.
$\mathfrak{V}_-$	1.2	This set is the forward image of $V_- \times \mathbb{D}_R$ under the map $(x, y, a) \mapsto (f_a(x, y), a)$.

Notatation	Section	Meaning
U	3.2	A neighborhood of $J(p)$ used to construct telescopes.
U'	3.2	$U' \equiv p^{-1}(U)$.
δ	3.2	A small value used to define a neighborhood of $C(p)$.
$v(x, y)$	3.2	$v(x, y) \equiv p(y) - x$
V', V'_r	3.2, 4.2	$V' = \text{pr}_1^{-1}(U) \cap v^{-1}(\mathbb{D}_\delta)$ and $V'_r = \{(x, y) \in V' \mid v(x, y) < r\delta\}$
$\mathbf{r}$	3.2	The radius of telescoping neighborhoods in the Kobayashi metric on U
β	3.6	The disk of Euclidean radius β about any point $u \in U'$ is mapped biholomorphically by p .
$u(x, y)$	3.2	A function on V' . The functions u, v give useful coordinates on V' .
A	3.3	A value choose small enough that the constructions in [HOV95] hold when $0 < a < A$.
U_z	3.2	The ball of radius $\mathbf{r}$ about $z \in J(p)$, measured in the Kobashi metric on U .
$\hat{J}(p)$	3.3	The natural extension of $p: J(p) \rightarrow J(p)$.
π_a	3.3	A homeomorphism from the natural extension $\hat{J}(p)$ to J_a for a sufficiently small.
$g_{z,\pm}, \check{g}_{z,\pm}$	3.3, 3.3	Holomorphic functions, the graph of which gives specific local stable/unstable manifolds of the point in question.
$\Delta_{z,a}, \Delta_{z,a}(r)$	3.3	$\Delta_{z,a}$ is the local stable manifold in B_z for f_a . $\Delta_{z,a}(r)$ is a smaller disk within $\Delta_{z,a}$ (not necessarily about a point of J_a).
H_{ca}	4.1	The component of the critical locus of f_a asymptotic to $y = c$.
ψ_+, ψ_-	5	Same as $\varphi_{a,+}, \varphi_{a,-}$ but with $\varphi_{a,-}$ rescaled to give the simplest iteration formula.
$\mathcal{V}_+, \mathcal{V}_-$	5	Subsets of V_+ and V_- for which the fibers of ψ_+ and ψ_- are disks.
$\tilde{f}$	3.3	$\tilde{f}(x, y, a) \equiv (f_a(x, y), a)$.
$\mathcal{S}$	5	The set of all roots of unity ω satisfying $\omega^{d^k} = 1$ for some $k \in \mathbb{N}$.

Notatation	Section	Meaning
m_+, m_-	5	The monodromy map of the critical locus determined by either $\mathcal{F}^+$ or $\mathcal{F}^-$, along with a pair of points on the critical locus on the same leaf of $\mathcal{F}^+$ or $\mathcal{F}^-$.
$H_c^\circ, \mathfrak{r}$	5	H_c° is a neighborhood of infinity in H_c which is mapped biholomorphically to $\mathbb{C} \setminus \mathbb{D}_{\mathfrak{r}}$.
$\mathfrak{c}$	6.6	The map $\mathfrak{c}(x, y) = (\bar{x}, \bar{y})$.
$\mathfrak{h}, \mathfrak{k}$	6.1	There are the restriction of a conjugacy h between two Henon maps to a component of the critical locus. $\mathfrak{h}$ and $\mathfrak{k}$ are the same conjugacy written in different coordinate systems.
σ	6.1	Change of complex coordinate functions. In each case the origin is a fixed point.
S_r		The circle of radius r about the origin in $\mathbb{C}$.

References

- [BLS93] Eric Bedford, Mikhail Lyubich, and John Smillie. Polynomial diffeomorphisms of $\mathbb{C}^2$. IV. The measure of maximal entropy and laminar currents. *Invent. Math.*, 112(1):77–125, 1993.
- [BS91a] Eric Bedford and John Smillie. Polynomial diffeomorphisms of $\mathbb{C}^2$: currents, equilibrium measure and hyperbolicity. *Invent. Math.*, 103(1):69–99, 1991.
- [BS91b] Eric Bedford and John Smillie. Polynomial diffeomorphisms of $\mathbb{C}^2$. II. Stable manifolds and recurrence. *J. Amer. Math. Soc.*, 4(4):657–679, 1991.
- [BS92] Eric Bedford and John Smillie. Polynomial diffeomorphisms of $\mathbb{C}^2$. III. Ergodicity, exponents and entropy of the equilibrium measure. *Math. Ann.*, 294(3):395–420, 1992.
- [BS98a] Eric Bedford and John Smillie. Polynomial diffeomorphisms of $\mathbb{C}^2$. V. Critical points and Lyapunov exponents. *J. Geom. Anal.*, 8(3):349–383, 1998.
- [BS98b] Eric Bedford and John Smillie. Polynomial diffeomorphisms of $\mathbb{C}^2$. VI. Connectivity of J . *Ann. of Math. (2)*, 148(2):695–735, 1998.
- [BS99] Eric Bedford and John Smillie. Polynomial diffeomorphisms of $\mathbb{C}^2$. VII. Hyperbolicity and external rays. *Ann. Sci. École Norm. Sup. (4)*, 32(4):455–497, 1999.
- [Buz99] Gregory Buzzard. Nondensity of stability for polynomial automorphisms of $\mathbb{C}^2$. *Indiana University Math Journal*, 48(3):857–865, 1999.
- [BV01] Gregory Buzzard and Kaushal Verma. Hyperbolic automorphisms and holomorphic motions in $\mathbb{C}^2$. *Mich. Math. J.*, 49(3):541–565, 2001.

- [Fir12] Tanya Firsova. The critical locus for complex Hénon maps. *Indiana Univ. Math. J.*, 61(4):1603–1641, 2012.
- [FL17] Tanya Firsova and Mikhail Lyubich. λ -lemma for families of Riemann surfaces and the critical loci of complex Hénon maps. *Conform. Geom. Dyn.*, 21:111–125, 2017.
- [FM89] Shmuel Friedland and John Milnor. Dynamical properties of plane polynomial automorphisms. *Ergodic Theory Dynam. Systems*, 9(1):67–99, 1989.
- [FS92] John Erik Fornæss and Nessim Sibony. Complex Hénon mappings in $\mathbb{C}^2$ and Fatou-Bieberbach domains. *Duke Mathematical Journal*, 65(2):345–380, 1992.
- [GR84] H. Grauert and R. Remmert. *Coherent Analytic Sheaves*. Springer-Verlag, 1984.
- [HOV94] John H. Hubbard and Ralph W. Oberste-Vorth. Hénon mappings in the complex domain. I. The global topology of dynamical space. *Inst. Hautes Études Sci. Publ. Math.*, (79):5–46, 1994.
- [HOV95] John H. Hubbard and Ralph W. Oberste-Vorth. Hénon mappings in the complex domain. II. Projective and inductive limits of polynomials. In *Real and complex dynamical systems (Hillerød, 1993)*, volume 464 of *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, pages 89–132. Kluwer Acad. Publ., Dordrecht, 1995.
- [KH95] Anatole Katok and Boris Hasselblatt. *Introduction to the modern theory of dynamical systems*, volume 54 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1995. With a supplementary chapter by Katok and Leonardo Mendoza.
- [Lyu83] M. Yu. Lyubich. Some typical properties of the dynamics of rational mappings. *Uspekhi Mat. Nauk*, 38(5(233)):197–198, 1983.
- [MnSS83] R. Mañé, P. Sad, and D. Sullivan. On the dynamics of rational maps. *Ann. Sci. École Norm. Sup. (4)*, 16(2):193–217, 1983.
- [Rue89] David Ruelle. *Elements of Differentiable Dynamics and Bifurcation Theory*. Academic Press, 1989.
- [Tan16] Raluca Tanase. Complex Hénon maps and discrete groups. *Adv. Math.*, 295:53–89, 2016.