



## Source tracking microplastics in the freshwater environment

N.L. Fahrenfeld <sup>a,\*</sup>, Georgia Arbuckle-Keil <sup>b</sup>, Nasrin Naderi Beni <sup>c</sup>,  
Shannon L. Bartelt-Hunt <sup>c</sup>



<sup>a</sup> Civil & Environmental Engineering, Rutgers, The State University of New Jersey, 500 Bartholomew Rd., Piscataway, NJ 08854, USA

<sup>b</sup> Department of Chemistry, Rutgers University-Camden, 315 Penn St, Camden, NJ 08102, USA

<sup>c</sup> Department of Civil Engineering, University of Nebraska-Lincoln, 205E Peter Kiewit Institute, 1110 South 67th Street, Omaha, NE 68182-0178, USA

### ARTICLE INFO

#### Article history:

Available online 29 November 2018

#### Keywords:

Source apportionment  
Land use  
Polymer composition  
Morphology  
Mass balance  
Sorption  
Freshwater

### ABSTRACT

Plastics are a frequently observed component of marine debris and there is growing concern about microplastic (MP) ecotoxicity, and the impacts of additives, sorbed hazardous organic contaminants, heavy metals, and biofilm on MP surfaces. The relative importance of MP from different terrestrial and freshwater sources is poorly understood and limits our ability to develop best management practices. This review focuses on evidence and methods for source apportionment of MP in freshwater environments including the use of MP characteristics, mass balance techniques, and surface characteristics. Within-study data indicated some potential for differences in polymer identity and morphology for differentiating select sources, but clear cross-study patterns were lacking. Major challenges identified include technical challenges in accurately identifying polymers, multiple classification schemes for reported MP morphologies, lack of data for several terrestrial sources, poor understanding of differential fate/transport/weathering processes for MP and surface contaminants, and methodological difficulties simultaneously confirming polymers and surface contaminants.

© 2018 Elsevier B.V. All rights reserved.

## 1. Introduction

Research conducted on occurrence of environmental microplastics (MP) has primarily focused on determining MP abundance in marine environments. Studies have been conducted to determine the prevalence of MP on the sea surface, within the water column, and in marine sediments along beaches. A review of 68 research articles conducted by Hidalgo-Ruz et al. [1] found microplastic particles ranged from 1 µm to 29 mm with concentrations varying by 7 orders of magnitude within all three of these compartments. Comparatively less research has been done on the abundance of MP in the freshwater environment; however, the growing number of studies that have been conducted offer strong evidence to support the existence of MP contamination in freshwater systems [2]. A recent review by Li et al. indicated that quantification of the MP particles in freshwater range from greater than 1 million pieces per cubic meter to less than 1 piece in 100 cubic meters [2]. A better understanding of the wide variety of

primary (manufactured in small sizes) and secondary (derived from the breakdown of other macro and micro-plastic) MP sources in freshwater is needed.

MP are generally defined as polymer particles <5 mm in size that present in a variety of morphologies (i.e., beads, fragments, fibers, films). Determining the source of MP is complicated by the fact that these particles can travel far from their point of origin, evidenced by observations of MP in remote regions of the world (e.g., Ref. [3]). There is also potential for different fate (e.g., biological uptake, photodegradation, biofouling and settling) and transport processes that may alter the profile of MP observed at a source compared to what may be observed at a sampling site that could complicate fingerprinting approaches to source tracking if not accounted for.

Mitigating MP pollution in the freshwater environment requires understanding the sources and their relative importance in terms not only of quantity but also of any potentially adverse impacts and environmental fate and transport. The implementation of voluntary phaseouts and legislation banning the addition of plastic bags and MP particles to personal care products is an example of how mitigation strategies can be implemented when sources are well defined (in addition to there being other circumstances that facilitate change) [4,5]. Multiple global reports [5,6] agree that a

\* Corresponding author.

E-mail addresses: [nfahrenf@rutgers.edu](mailto:nfahrenf@rutgers.edu) (N.L. Fahrenfeld), [arbuckle@camden.rutgers.edu](mailto:arbuckle@camden.rutgers.edu) (G. Arbuckle-Keil), [sbartelt2@unl.edu](mailto:sbartelt2@unl.edu) (S.L. Bartelt-Hunt).

## Abbreviations

ATR	attenuated total reflectance
EDS	Energy-dispersive X-ray spectroscopy
FPA	focal plane array and microscopic
FTIR	Fourier transform infrared spectroscopy
ICP-MS	inductively coupled plasma mass spectrometry
MP	microplastic
nMDS	non-metric multidimensional scaling
PE	polyethylene
PET	polyethylene terephthalate
POPs	persistent organic pollutants
PP	polypropylene
PS	polystyrene
PVC	polyvinyl chloride
PUR	polyurethane
Pyr-GC/MS	pyrolysis-gas chromatography/mass spectrometry
μ-FTIR	microscope FTIR

significant knowledge gap exists with regards to potential sources of MP. The present review compiles data from three approaches for source apportionment of MP (1) linking particle characteristics to sources, (2) source sampling and mass balance to establish inputs, and (3) using particle surface contaminants to implicate potential sources.

## 2. Linking particle characteristics to sources

Differences in polymer chemical composition and particle color, morphology [7], and size could potentially be used to help elucidate the sources of MP. Plastics Europe [8] produced a report that reported quantities of different polymer types used by different industries, suggesting some potential linkage between polymer type and application. To date, limited research has linked observations of MP characteristics to specific sources in freshwater systems.

The most commonly identified MP particles (80%) are polyethylene (PE), polypropylene (PP), polystyrene (PS), polyvinyl chloride (PVC), polyurethane (PUR), and polyethylene terephthalate (PET) with some variations including copolymer compositions depending on the collection sites, pre-treatment and analytical methods employed for identification [8,9]. Fig. 1 shows the polymers identified across four studies published over a two year period using attenuated total reflectance (ATR) Fourier transform infrared (FTIR) spectroscopy [9,10], focal plane array (FPA) FTIR [11], and microscopic FTIR ( $\mu$ -FTIR) in reflectance mode [12]. The lack of consistent methodologies for identifying polymers may be limiting the ability to use polymer identity for source tracking. MP identification techniques include optical microscope observation after sieving and/or density gradient separation, but for definitive determination of polymer composition, additional spectroscopic analysis is necessary. Analytical methodologies employed to date are FTIR spectroscopy, ideally  $\mu$ -FTIR, Raman microscopy, and pyrolysis-gas chromatography/mass spectrometry (Pyr-GC/MS) [2,13,14]. Hendrickson et al. [15] stated that the inconsistency of MP particle identification by different methods may be related to the fact that MP particles found in the environment may contain copolymers and additives. Due to the broad number of polymer/copolymer possibilities, in addition to photo-oxidation and biodegradation (primarily, presumably, of additives), accurate identification of MPs is challenging. Several combined methods have been suggested to improve polymer


identification but are time consuming and often require specific expertise. Misidentification may occur if visual microscopy is not combined with spectroscopy. Kroon et al. [16] suggests a step-by-step work flow in order to reduce bias in MP identification that relies on stereomicroscopic visual sorting, microscopic photography followed by FTIR. A validated method proposed by Danish Centre for Environment & Energy [17] identifies MP  $< 100 \mu\text{m}$  via stereomicroscopy followed by  $\mu$ -FTIR. MP particles of this size are best identified via  $\mu$ -FTIR with a FPA detector [18]. ATR FTIR spectroscopy is recommended for MP particles  $> 100 \mu\text{m}$ . Polymer molecular weight determination, such as gel permeation chromatography, may provide additional confirmation of potential MP source. These methods are time-consuming and the ability to analyze more samples within a reasonable time frame is desirable and development is on-going.

A small number of studies have inferred the source of MP contamination based on the polymer. Zhang et al. [19] determined that polystyrene identified in the Bohai Sea in China was likely due to floating foam rafts from marine agricultural areas and Styrofoam food containers, both of which were observed in a reconnaissance of beaches and shoreline near the sampling areas. Similarly Xiong et al. [20] identified the dominant polymers detected in water samples to be polystyrene and polyethylene, which was attributed by the authors to improper waste disposal of food packaging and containers along the shoreline. In contrast to observed differences in MP morphology, the polymer profile was not reported to be significantly different between influent and effluent samples in a 10 treatment plant study in the Netherlands [10]. Browne et al. [21] identified a predominance of polyester and acrylic fibers in shoreline sediments as well as wastewater effluents, suggesting that wastewater inputs were the predominant source of fibers at these locations.

Fig. 1 provides an analysis of reported polymer composition across 35 wastewater treatment plant effluent samples, one wastewater influent sample, one grit sample, one grease sample, one sludge sample, one wastewater solids sample and six compost samples. It includes data from four studies published over a two-year time period. Taken together, over 30 different polymers were identified in these samples. The composition of wastewater effluent samples was highly varied with some effluent samples containing only 2 polymers, while others contained up to twelve different polymer types. Similarly, the compost samples contained  $\geq 4$  polymers with no consistent trends observed in the percentage of a given polymer among the samples.

The morphology of MPs may also provide some evidence of their source [7]. The presence of primary MP morphologies (microbeads from personal care products and fibers from laundering of textiles) are likely associated with wastewater inputs [22]. Similarly, secondary MP such as films and fragments are associated with weathering of plastic packaging or plastic bags and may be associated with terrestrial sources such as improper solid waste disposal. Another source of MP fragments is stormwater runoff. Horton et al. [23] identified a storm drain carrying urban roadway runoff directly upgradient of a sampling location as the source of the increased abundance of fragments identified at this site.

We analyzed nineteen paired wastewater influent and effluent samples originally reported by Magnusson and Wahlberg [24] and Michielssen et al. [25] and identified the primary morphologies in each sample (Fig. 2a). The primary morphologies present in each of the paired samples were fibers, films and fragments. Morphology relative percent differences ranged from 11 to 147% for paired influent and effluent samples with an average of  $85 \pm 41\%$  relative percent difference for the two sample types. When comparing the paired influent and effluent from each location, there were decreasing percentages of fibers and films in the effluent as



**Fig. 1.** Heat map representing percent identity of total reported particles assigned a given polymer in different sources [9–12]. [MP are reported here are done so as the original authors did. Notably, authors reported Zn-sterate coated particles. Zn-sterate is a lubricant used for polymers and other materials. Cellulose-based polymers were also reported. Cellulose is a natural polymer but may be difficult to distinguish from regenerated cellulose (rayon).].


compared to the influent in 10 and 8 samples, respectively. In contrast, the fragment morphology increased from influent to effluent in 12 of the 19 samples. This indicates that fibers and films are more easily removed in wastewater treatment unit processes and may be more likely to be associated with municipal biosolids. These morphologies may enter freshwater systems predominantly through surface runoff after land application of biosolids, while fragments may be more commonly associated with wastewater effluent discharges into receiving streams. Combining morphological data across studies and matrices (water, biosolids) and levels of treatment (influent, effluent) did not indicate clustering by source (Fig. 2b). Here, we applied ordination [non-metric multidimensional scaling (nMDS)] of square root transformed percent abundance data for the different reported MP morphologies for different studies and clear clustering was not observed. However, this does not necessarily mean morphology is not a useful indicator: it should be noted that cross study comparisons are challenging given that the shape classes reported vary and may include fragments, beads, fibers, pellets, paint chips, films, and foams, and that the morphologies observed for a given sample can vary by the size class targeted. It is also possible that these patterns are not consistent across geographies.

### 3. Source sampling and mass balance to establish inputs

Direct sampling of aqueous and solid waste streams has been used to characterize the sources of MP pollution in freshwater systems. A major focus of these studies has been understanding the impact of municipal wastewater effluent as a pathway for MP entry into surface water [10–12,24–30]. Many of these studies include calculations for daily emissions of MP to the receiving water bodies (or these values are easily calculated where average daily flow rates are reported). Comparing emission mass fluxes (emission

concentrations  $\times$  flow rates) to the receiving water body mass fluxes can inform the relative importance of wastewater effluent as a point source of MP, although these dilution factors are rarely reported. Some researchers did compare the MP concentration in wastewater effluent to surrounding surface water, generally implicating the effluent as a source based on its higher concentration [26,27,29]. Studies have also sought to compare up and downstream MP concentrations to implicate wastewater effluent as a point source [31], in this case the majority of the downstream sampling sites did not have significantly higher MP concentrations compared to the paired upstream sites. In the case of sanitary and combined sewer overflows, wastewater influent would be released to freshwater sources and influent generally had an average (per study) of 63–420 times higher concentrations of MP (Fig. 3). Variation in the total MP concentration for different studies is likely due to some combination of differences in target MP size class and other analysis techniques, study site, temporal variation, and whether MP were identified visually (considered prone to false positives and negatives) [32] or with confirmation of chemical structure. The MP composition in wastewater influent would be expected to be impacted both by the inputs and activities in the sewershed (i.e., fibers from laundry, industrial inputs if present, stormwater if combined sewers, use of consumer products, etc.) and the treatment performed at the facilities themselves that can impact the effluent MP fingerprint and concentration. Differentiating these sources contributing to wastewater influent is complex and system specific. The best information available is regarding the contribution of the laundry fraction from households (as reviewed by Ref. [33]).

Runoff containing MP from a variety of sources can transport and introduce MP to surface water. Sewage solids from a variety of treatment processes have been analyzed directly [10,34,35]. Solids can serve as a sink for MP in wastewater treatment processes and




**Fig. 2.** a. Percent of total MP observed for a given morphology in a. wastewater influent and effluent [24,25]. b. nonmetric multidimensional scaling representing the square root transformed assigned morphology for a given sample. Cross matrix and cross study comparison of MP morphology (by %). Closed symbols represent wastewater influent, open symbols wastewater effluent, stars represents solids.

may be applied to land where MP can be transported to surface water in runoff. Consideration has also been given to land application of other materials to agricultural and in gardening soils including compost [36], and fertilizer [9]. Other sources of plastics exist in agricultural and home gardening (e.g., plastic mulch, landscape fabric) [37], but their impact on MP pollution has not been reported to the authors' knowledge. Attempts to estimate the contribution of MP in sludge to soil (e.g., Ref. [10]) and amount of MP transported in runoff have been made (e.g., Ref. [37]). But, controlled runoff studies estimating the MP loading in agricultural runoff have not been published, to the authors' knowledge.

Likewise, few data are available on the loading of MP in (urban) stormwater [38]. Stormwater may carry MP from tires and road markings [23,37,39], as well as other plastic litter. The potential for stormwater treatment systems to remove MP has not been reported, but removal of macroplastics has been studied (e.g., Ref. [40]).

Relatively fewer researchers have attempted to quantify atmospheric deposition of MP fibers [41], despite widespread anecdotal reports of careful measures being required to prevent contamination of field samples with fibers, likely from researchers clothing and/or air handling systems in laboratories. Comparison of up and



**Fig. 3.** Comparison of wastewater influent and effluent sample microplastic (MP) concentrations [10,12,24–26,29,30].

downstream fiber concentrations and fluxes surrounding Paris, FR indicated fibers did not increase downstream of the city [42]. Methods for polymer and natural fiber identification used in the textile industry have not been widely applied, including examination of fiber cross sections and longitudinal shapes, as well as flame tests that allow for visual and odor evaluation of the fibers on approach, within, and after removal from the flame, as well as for examination of the ashed residue [43–47]. To the authors' knowledge, comparisons between the expected loads to freshwater from clothes washing versus shedding of fibers to the atmosphere have not been reported.

Several studies have linked MP abundance to urban land uses and population densities [22,48,49]. Xiong et al. [20] compared MP abundance in an urban lake in China to the abundance in the four rivers feeding into the lake to determine whether MP contamination in the lake originated from terrestrial sources or via surface water inputs. As MP abundances in the lake were higher than those in the rivers, it was concluded that terrestrial contamination, specifically improper solid waste disposal, was the primary source to the lake; sampling was performed during the wet season when transport from land would be expected in drawing this conclusion. Ballent et al. [49] explored MP in tributaries to a lake in Canada and mapped wastewater treatment facilities and plastic industries. Limited sample size prevented mass balance approaches for implicating sources and the authors noted "assigning particular origins to the microplastics is challenging due to their small size, fragmented nature, and the unknown range of possible sources." Field observations have been used to indirectly indicate other sources than wastewater treatment plants in surface water including: the presence of MP in rivers upstream of wastewater treatment plants [31], in high concentrations in bed sediments without wastewater inputs [23], lack of mass balance in wastewater impacted surface waters [30], and comparisons of wastewater impacted versus unimpacted lakes [38]. Thus, mass balance approaches can be useful for indicating the magnitude of

commonly targeted sources (i.e., wastewater treatment plants) and for highlighting the need to study nonpoint sources. Addressing these nonpoint sources may be more challenging given that they are more diffuse and diverse. Additionally, proper study design for mass balance approaches must be considered. Particularly, to accurately calculate loads, robust approaches with appropriate sampling techniques (likely composited samples) and sufficient sample sizes to capture variability are needed, which is challenging given analytical challenges that prevent high throughput sample analysis. In lieu of the ability to perform high resolution sampling, modeling the transport of MP [50] may also help fill in gaps in knowledge.

#### 4. Can surface contaminants help to elucidate source?

While many MP analysis techniques use oxidation, enzymatic reactions, and/or acid washing to remove surface contaminants to allow for confirmation of the polymer composition, the contaminants present on the surface of MP particles may have potential utility in source tracking. One challenge with this potential approach is that surface contaminants adsorbed from the surrounding environment after MP release [51] should be considered in addition to the additives added during plastics manufacture to the polymer mixture which can be present on the surface (and within MP particles). Although challenging, identification of unique surface contaminants, including metals, organics and biofilms, could have utility for source tracking of MP pollution in freshwater systems if consistent relationships between surface contaminants and sources can be established and weathering processes understood. One significant challenge to consider is how to confirm that surface contaminants are adsorbed to MP rather than other particulate matter associated with MP particles.

Metals are commonly added as catalysts, pigments, and stabilizers during plastic manufacturing [52] therefore, targeting

specific heavy metals may help indicate the source of MP pollution if these profiles are unique in different macroplastic sources. Wang et al. [53] summarized macroplastic heavy metal profiles then attempted to match MP and macroplastic heavy metal profiles. From their field sorption tests they concluded, that most heavy metals on the surface of the MPs they studied were from production rather than environmental sources. Thus, if the plastics produced with metals had unique applications there could be potential for linking the heavy metal profile to source, assuming environmental sorption and desorption processes did not interfere. There is evidence, however, from several studies demonstrating that virgin and aged MPs can adsorb metals during different stages of their transport pathway in the environment [18]. For example, metals extracted from MPs in the San Diego Bay area [54], identified by ICP-MS (inductively coupled plasma mass spectrometry), included toxic metals (Cd, Ni, Zn and Pb). The concentrations of these metals were more uniformly found on polymers than when MP particles were exposed to organic chemicals [54], which could vary by source. For sorbed heavy metals to have utility in source tracking the rates of weathering of these metals would need to be slower than transport and the concentrations present from production understood.

Different hydrophobic organic chemicals and persistent organic pollutants (POPs) are known to bind to MPs within the environment [55,56], especially in freshwater systems where the concentrations of these chemicals are expected to be higher due to proximity to the use of these chemicals [57]. The perceived risk of plastic to serve as a vector for these toxic chemicals to accumulate via the food chain has not been linked to plastic ingestion (as recently reviewed [58]). Identification of the POPs as well as polymer degradation products involve multi-step processes, including extraction of the organic(s) from the MP particles [59] followed by chromatography before chemical determination. It has been shown that hydrophobic organic compounds have greater affinity for MPs like polyethylene, polypropylene, and polyvinyl chloride than for natural sediments [51,57]. Changes to environmental conditions, such as residence time, temperature, pH and exposure to sunlight, will influence equilibrium dynamics between chemicals and MPs, impacting their accumulation and transport [12]. However, studying organic surface contaminants found on MPs may help lead to identification of the sources. Di and Wang [60] evaluated MP particles in surface waters and sediments in China. Organics or pharmaceutical intermediates were identified via Raman spectroscopy; the authors speculated that the origin of these organic substances may be due to the MP coming from hospitals or from sorption of pharmaceuticals to MP in the environment, given local discharges from hospitals.

A third surface contaminant that may have utility for MP source tracking is the growth of biofilms on MP surfaces. Given observations of unexpected microbial community members in MP biofilm in the ocean [61], there is some evidence that terrestrially sourced biofilms can be transported long distances. MP incubated in different water sources (i.e., river water versus wastewater) had unique biofilm microbial communities that clustered by source water rather than by polymer/morphology [62]. Similarly, marine biofilm community structures grown on PET in the field varied by location and season [63]. More understanding of how robust biofilms are to weathering is needed to know if it is possible these biofilm community structures could be used to help differentiate MP sources.

## 5. Conclusions

Growing evidence suggests that MPs are present in freshwater systems and shoreline sediments worldwide and transfer of MPs

from freshwater to marine systems is likely a significant source of marine microplastic contamination. Controlling the occurrence and transport of MPs in freshwater systems requires better understanding of MP sources and pathways including wastewater effluents, urban stormwater and agricultural runoff, atmospheric transport of fibers, and improper solid waste disposal adjacent to freshwater systems. MP characteristics, including morphology and polymer type may provide clues to improve our ability to source track microplastic contamination. In addition, surface contaminants including metals, organic compounds and biofilms may provide some evidence of microplastic source, but complications arise as environmental contaminants can also associate with freshwater MPs *in situ*. Challenges associated with source tracking based on these characteristics include limitations to accurate polymer identification, poor understanding of weathering processes, and limited understanding of microplastic transport mechanisms within freshwater systems.

## Acknowledgements

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## References

- [1] V. Hidalgo-Ruz, L. Gutow, R.C. Thompson, M. Thiel, Microplastics in the marine environment: a review of the methods used for identification and quantification, *Environ. Sci. Technol.* 46 (2012) 3060–3075.
- [2] J. Li, H. Liu, J. Paul Chen, Microplastics in freshwater systems: a review on occurrence, environmental effects, and methods for microplastics detection, *Water Res.* 137 (2018) 362–374.
- [3] A.L. Lusher, V. Tirelli, I. O'Connor, R. Officer, Microplastics in Arctic polar waters: the first reported values of particles in surface and sub-surface samples, *Sci. Rep.* 5 (2015) 14947.
- [4] D. Xanthos, T.R. Walker, International policies to reduce plastic marine pollution from single-use plastics (plastic bags and microbeads): a review, *Mar. Pollut. Bull.* 118 (2017) 17–26.
- [5] A. Arroyo Schnell, N. Klein, E. Gómez Girón, J. Sousa, National Marine Plastic Litter Policies in EU Member States: an Overview, IUCN, Brussels, Belgium, 2017.
- [6] P.J. Kershaw (Editor), Joint Group of Experts on the Scientific Aspects of Marine Environmental Protection (GESAMP), Sources, Fate and Effects of Microplastics in the Marine Environment: a Global Assessment, 2015.
- [7] P.A. Helm, Improving microplastics source apportionment: a role for microplastic morphology and taxonomy? *Analytical Methods* 9 (2017) 1328–1331.
- [8] PlasticsEurope, Plastics- the Facts 2016, PlasticsEurope, Brussels, Belgium, 2016.
- [9] N. Weithmann, J.N. Möller, M.G.J. Löder, S. Piehl, C. Laforsch, R. Freitag, Organic fertilizer as a vehicle for the entry of microplastic into the environment, *Sci. Adv.* 4 (2018).
- [10] J. Vollersten, A.A. Hansen, Microplastic in Danish Wastewater: Sources, Occurrences and Fate, Danish Environmental Protection Agency, 2017.
- [11] S.M. Mintenig, I. Int-Veen, M.G.J. Löder, S. Primpke, G. Gerdts, Identification of microplastic in effluents of waste water treatment plants using focal plane array-based micro-Fourier-transform infrared imaging, *Water Res.* 108 (2017) 365–372.
- [12] F. Murphy, C. Ewins, F. Carbonnier, B. Quinn, Wastewater treatment works (WWTW) as a source of microplastics in the aquatic environment, *Environ. Sci. Technol.* 50 (2016) 5800–5808.
- [13] S. Rezania, J. Park, M.F. Md Din, S. Mat Taib, A. Talaiekhozani, K. Kumar Yadav, H. Kamyab, Microplastics pollution in different aquatic environments and biota: a review of recent studies, *Mar. Pollut. Bull.* 133 (2018) 191–208.
- [14] A.B. Silva, A.S. Bastos, C.I.L. Justino, J.P. da Costa, A.C. Duarte, T.A.P. Rocha-Santos, Microplastics in the environment: challenges in analytical chemistry - a review, *Anal. Chim. Acta* 1017 (2018) 1–19.
- [15] E. Hendrickson, E.C. Minor, K. Schreiner, Microplastic abundance and composition in western Lake Superior as determined via microscopy, Pyr-GC/MS, and FTIR, *Environ. Sci. Technol.* 52 (2018) 1787–1796.
- [16] F. Kroon, C. Motti, S. Talbot, P. Sobral, M. Puotinen, A workflow for improving estimates of microplastic contamination in marine waters: a case study from North-Western Australia, *Environ. Pollut.* 238 (2018) 26–38.
- [17] Danish Center for Environment and Energy (DCE), Proposal for a Measurement Method for Use in Investigations of Microplastics in Tap Water, 2018.
- [18] O.S. Alimi, J. Farner Budarz, L.M. Hernandez, N. Tufenkji, Microplastics and nanoplastics in aquatic environments: aggregation, deposition, and enhanced contaminant transport, *Environ. Sci. Technol.* 52 (2018) 1704–1724.

[19] W. Zhang, S. Zhang, J. Wang, Y. Wang, J. Mu, P. Wang, X. Lin, D. Ma, Microplastic pollution in the surface waters of the Bohai Sea, China, *Environ. Pollut.* 231 (2017) 541–548.

[20] X. Xiong, K. Zhang, X. Chen, H. Shi, Z. Luo, C. Wu, Sources and distribution of microplastics in China's largest inland lake – Qinghai Lake, *Environ. Pollut.* 235 (2018) 899–906.

[21] M.A. Browne, P. Crump, S.J. Niven, E. Teuten, A. Tonkin, T. Galloway, R. Thompson, Accumulation of microplastic on shorelines worldwide: sources and sinks, *Environ. Sci. Technol.* 45 (2011) 9175–9179.

[22] R. Vaughan, S.D. Turner, N.L. Rose, Microplastics in the sediments of a UK urban lake, *Environ. Pollut.* 229 (2017) 10–18.

[23] A.A. Horton, C. Svendsen, R.J. Williams, D.J. Spurgeon, E. Lahive, Large microplastic particles in sediments of tributaries of the River Thames, UK – abundance, sources and methods for effective quantification, *Mar. Pollut. Bull.* 114 (2017) 218–226.

[24] K. Magnusson, C. Wahlberg, Mikroskopiska skräppartiklar i vatten från avloppssreningsverk, Rapport NR B 2208 (2014) 33.

[25] M.R. Michielssen, E.R. Michielssen, J. Ni, M.B. Duhaime, Fate of microplastics and other small anthropogenic litter (SAL) in wastewater treatment plants depends on unit processes employed, *Environ. Sci. Water Res. Technol.* 2 (2016) 1064–1073.

[26] R. Sutton, S.A. Mason, S.K. Stanek, E. Willis-Norton, I.F. Wren, C. Box, Microplastic contamination in the San Francisco Bay, California, USA, *Mar. Pollut. Bull.* 109 (2016) 230–235.

[27] J. Talvitie, M. Heinonen, J.-P. Pääkkönen, E. Vahtera, A. Mikola, O. Setälä, R. Vaahala, Do wastewater treatment plants act as a potential point source of microplastics? Preliminary study in the coastal Gulf of Finland, Baltic Sea, *Water Sci. Technol.* 72 (2015) 1495–1504.

[28] S.A. Carr, J. Liu, A.G. Tesoro, Transport and fate of microplastic particles in wastewater treatment plants, *Water Res.* 91 (2016) 174–182.

[29] S.A. Mason, D. Garneau, R. Sutton, Y. Chu, K. Ehmann, J. Barnes, P. Fink, D. Papazissimos, D.L. Rogers, Microplastic pollution is widely detected in US municipal wastewater treatment plant effluent, *Environ. Pollut.* 218 (2016) 1045–1054.

[30] H.A. Leslie, S.H. Brandsma, M.J.M. van Velzen, A.D. Vethaak, Microplastics en route: field measurements in the Dutch river delta and Amsterdam canals, wastewater treatment plants, North Sea sediments and biota, *Environ. Int.* 101 (2017) 133–142.

[31] S. Estahbanati, N.L. Fahrenfeld, Influence of wastewater treatment plants on microplastics in surface waters, *Chemosphere* 162 (2016), 277–184.

[32] S. Ziajahromi, P.A. Neale, L. Rintoul, F.D.L. Leusch, Wastewater treatment plants as a pathway for microplastics: development of a new approach to sample wastewater-based microplastics, *Water Res.* 112 (2017) 93–99.

[33] F. Salvador Cesa, A. Turra, J. Baroque-Ramos, Synthetic fibers as microplastics in the marine environment: a review from textile perspective with a focus on domestic washings, *Sci. Total Environ.* 598 (2017) 1116–1129.

[34] K.A.V. Zubris, B.K. Richards, Synthetic fibers as an indicator of land application of sludge, *Environ. Pollut.* 138 (2005) 201–211.

[35] A.M. Mahon, B. O'Connell, M.G. Healy, I. O'Connor, R. Officer, R. Nash, L. Morrison, Microplastics in sewage sludge: effects of treatment, *Environ. Sci. Technol.* 51 (2017) 810–818.

[36] M. Bläsing, W. Amelung, Plastics in soil: analytical methods and possible sources, *Sci. Total Environ.* 612 (2018) 422–435.

[37] K. Magnusson, K. Eliasson, A. Fräne, K. Haikonen, J. Hultén, M. Olshammar, J. Stadmark, A. Voisin, Swedish Sources and Pathways for Microplastics to the Marine Environment, IVL Svenska miljöinstitutet, Stockholm, 2016.

[38] S. Lasee, J. Mauricio, W.A. Thompson, A. Karnjanapiboonwong, J. Kasumba, S. Subbiah, A.N. Morse, T.A. Anderson, Microplastics in a freshwater environment receiving treated wastewater effluent, *Integrated Environ. Assess. Manag.* 13 (2017) 528–532.

[39] P.J. Kole, A.J. Löhr, F. Van Belleghem, A. Ragas, Wear and tear of tyres: a stealthy source of microplastics in the environment, *Int. J. Environ. Res. Publ. Health* 14 (2017) 1265.

[40] Q. Guo, J. Kim, Quantity and Quality of Stormwater Solids Trapped by Hydrodynamic Separators at Highway Sites, World Environmental and Water Resources Congress 2010: Challenges of Change, 2010, pp. 3291–3300.

[41] R. Dris, J. Gasperi, M. Saad, C. Mirande, B. Tassin, Synthetic fibers in atmospheric fallout: a source of microplastics in the environment? *Mar. Pollut. Bull.* 104 (2016) 290–293.

[42] R. Dris, J. Gasperi, V. Rocher, B. Tassin, Synthetic and non-synthetic anthropogenic fibers in a river under the impact of Paris Megacity: sampling methodological aspects and flux estimations, *Sci. Total Environ.* 618 (2018) 157–164.

[43] A.N. Khan, N. Abir, M.A.N. Rakib, E.M. Saberin Bhuiyan, M.R. Howlader, A review paper on textile fiber identification, *IOSR J. Polym. Textil. Eng.* 04 (2017) 14–20.

[44] D.R. Perry, Identification of Textile Materials, seventh ed., The Textile Institute, Manchester, UK, 1985.

[45] D. Braun, Simple Methods for Identification of Plastics, fourth ed., Hanser Gardner Publications, Cincinnati, OH, 1999.

[46] International Organization for Standardization, ISO/TR 11827 Textiles – Composition Testing – Identification of Fibers, 2012.

[47] M. Lewin, Handbook of Fiber Chemistry, CRC Press, Boca Raton, FL, 2006.

[48] S. Sruthy, E.V. Ramasamy, Microplastic pollution in Vembanad Lake, Kerala, India: the first report of microplastics in lake and estuarine sediments in India, *Environ. Pollut.* 222 (2017) 315–322.

[49] A. Ballent, P.L. Corcoran, O. Madden, P.A. Helm, F.J. Longstaffe, Sources and sinks of microplastics in Canadian Lake Ontario nearshore, tributary and beach sediments, *Mar. Pollut. Bull.* 110 (2016) 383–395.

[50] R.N. Cable, D. Beletsky, K. Wigginton, B.W. Locke, M.B. Duhaime, Distribution and modeled transport of plastic pollution in the Great Lakes, the world's largest freshwater resource, *Front. Environ. Sci.* 5 (2017) 45.

[51] B. Graca, K. Szewc, D. Zakrzewska, A. Dołęga, M. Szczerbowska-Boruchowska, Sources and fate of microplastics in marine and beach sediments of the Southern Baltic Sea – a preliminary study, *Environ. Sci. Pollut. Control Ser.* 24 (2017) 7650–7661.

[52] E. Nakashima, A. Isobe, S.I. Kako, T. Itai, S. Takahashi, Quantification of toxic metals derived from macroplastic litter on Ookushi Beach, Japan, *Environ. Sci. Technol.* 46 (2012) 10099–10105.

[53] J. Wang, J. Peng, Z. Tan, Y. Gao, Z. Zhan, Q. Chen, L. Cai, Microplastics in the surface sediments from the Beijiang River littoral zone: composition, abundance, surface textures and interaction with heavy metals, *Chemosphere* 171 (2017) 248–258.

[54] C.M. Rochman, B.T. Hentschel, S.J. Teh, Long-term sorption of metals is similar among plastic types: implications for plastic debris in aquatic environments, *PLoS One* 9 (2014), e85433.

[55] C.M. Rochman, E. Hoh, B.T. Hentschel, S. Kaye, Long-term field measurement of sorption of organic contaminants to five types of plastic pellets: implications for plastic marine debris, *Environ. Sci. Technol.* 47 (2013) 1646–1654.

[56] H. Lee, W.J. Shim, J.-H. Kwon, Sorption capacity of plastic debris for hydrophobic organic chemicals, *Sci. Total Environ.* 470 (2014) 1545–1552.

[57] A.A. Horton, A. Walton, D.J. Spurgeon, E. Lahive, C. Svendsen, Microplastics in freshwater and terrestrial environments: evaluating the current understanding to identify the knowledge gaps and future research priorities, *Sci. Total Environ.* 586 (2017) 127–141.

[58] A.A. Koelmans, E. Besseling, E. Fockema, M. Kooi, S. Mintenig, B.C. Ossendorp, P.E. Redondo-Hasselerharm, A. Verschoor, A.P. van Wezel, M. Scheffer, Risks of plastic debris: unravelling fact, opinion, perception, and belief, *Environ. Sci. Technol.* 51 (2017) 11513–11519.

[59] A. Ceccarini, A. Corti, F. Erba, F. Modugno, J. La Nasa, S. Bianchi, V. Castelvetro, The hidden microplastics: new insights and figures from the thorough separation and characterization of microplastics and of their degradation byproducts in coastal sediments, *Environ. Sci. Technol.* 52 (2018) 5634–5643.

[60] M. Di, J. Wang, Microplastics in surface waters and sediments of the three Gorges Reservoir, China, *Sci. Total Environ.* 616–617 (2018) 1620–1627.

[61] J.P. Harrison, T.J. Hoellein, M. Sapp, A.S. Tagg, Y. Ju-Nam, J.J. Ojeda, Microplastic-associated Biofilms: a Comparison of Freshwater and Marine Environments, *Freshwater Microplastics*, Springer, 2018, pp. 181–201.

[62] K. Parrish, N.L. Fahrenfeld, Microbial Ecology of Microplastic Biofilm in Fresh- and Wastewater as a Function Material and Size Class, DOI (in review).

[63] S. Oberbeckmann, A.M. Osborn, M.B. Duhaime, Microbes on a bottle: substrate, season and geography influence community composition of microbes colonizing marine plastic debris, *PLoS One* 11 (2016), e0159289.