

NUCLEUS I: ADJUNCTION SPECTRA IN RECOMMENDER SYSTEMS AND DESCENT

DUSKO PAVLOVIC ^a AND DOMINIC J. D. HUGHES ^{b,c}

^a University of Hawaii, Honolulu, HI

^b Apple Inc., Cupertino, CA

^c UC Berkeley, Berkeley, CA

ABSTRACT. Recommender systems build user profiles using concept analysis of usage matrices. The concepts are mined as spectra and form Galois connections. Descent is a general method for spectral decomposition in algebraic geometry and topology which also leads to generalized Galois connections. Both recommender systems and descent theory are vast research areas, separated by a technical gap so large that trying to establish a link would seem foolish. Yet a formal link emerged, all on its own, bottom-up, against authors' intentions and better judgment. Familiar problems of data analysis led to a novel solution in category theory. The present paper arose from a series of earlier efforts to provide a top-down account of these developments.

CONTENTS

1. Introduction	2
1.1. Idea and result	2
1.2. Background	3
1.3. Overview	4
2. Posetal nuclei: from contexts to concepts	5
2.1. Idea of Formal Concept Analysis	5
2.2. Formalizing concept analysis	6
2.3. Posetal matrices	6
2.4. Completions make matrices representable as adjunctions	7
2.5. Posetal adjunctions induce closure and interior operators	8
2.6. Closure and interior operators have adjunction resolutions	8
2.7. Concept lattices as nuclei	8
3. From categorical contexts to adjunctions, monads, and comonads	9
3.1. Lifting concept analysis to categories — and not further	9
3.2. Matrices (distributors, profunctors, bimodules)	10
3.3. Completions make matrices representable as adjunctions again	11

Key words and phrases: adjunction, monad, concept analysis, spectral decomposition.

^aSupported by NSF and AFOSR..

^cVisiting scholar at Logic Group. Thanks to Wes Holliday and Dana Scott.

3.4.	From adjunctions to monads and comonads	12
3.5.	From monads and comonads to adjunctions	14
3.6.	Monads and comonads are reflective in adjunctions	15
4.	The nucleus functor	16
4.1.	Nucleus decomposes descent monads and comonads	17
4.2.	Nucleus is an adjunction	18
5.	The nucleus monad is idempotent	21
5.1.	From weak to strong equivalences through absolute completions	22
5.2.	The Nucleus Theorem	23
5.3.	Consequences of the Nucleus Theorem	28
6.	Simple nucleus	29
7.	New frameworks for old big pictures	32
7.1.	Descent to nucleus	32
7.2.	Kan's adjunction	35
8.	Closing	38
	References	40
Appendix A.	Comprehending presheaves and matrices as discrete fibrations	44
Appendix B.	Idempotents, absolute (Cauchy) completions, and weak (Morita) equivalences	45

1. INTRODUCTION

1.1. Idea and result.

Definition 1.1. We say that an adjunction $F = (F^* \dashv F_* : \mathbb{B} \rightarrow \mathbb{A})$ is nuclear when the right adjoint F_* is monadic and the left adjoint F^* is comonadic.

Explanation. For an adjunction $F = (F^* \dashv F_* : \mathbb{B} \rightarrow \mathbb{A})$, we say that

- F_* is monadic when $\mathbb{B}$ is equivalent to the category $\mathbb{A}^{\overleftarrow{F}}$ of algebras for the monad $\overleftarrow{F} = F_* F^* : \mathbb{A} \rightarrow \mathbb{A}$, and that
- F^* is comonadic when $\mathbb{A}$ is equivalent to the category $\mathbb{B}^{\overrightarrow{F}}$ of coalgebras for the comonad $\overrightarrow{F} = F^* F_* : \mathbb{B} \rightarrow \mathbb{B}$.

The adjunction F is thus nuclear when it displays how the categories $\mathbb{A}$ and $\mathbb{B}$ determine each other along it: $\mathbb{A}$ as $\mathbb{B}^{\overrightarrow{F}}$ and $\mathbb{B}$ as $\mathbb{A}^{\overleftarrow{F}}$. The situation is reminiscent of Maurits Escher's “*Drawing hands*” in Fig.1. In a formal sense, nuclear adjunctions are the lax version of equivalences.

Claim. Any adjunction F induces a nuclear adjunction $\mathfrak{N}F$

$$\frac{F = (F^* \dashv F_* : \mathbb{B} \rightarrow \mathbb{A})}{\mathfrak{N}F = \left(F^\# \dashv F_\# : \mathbb{A}^{\overleftarrow{F}} \rightarrow \mathbb{B}^{\overrightarrow{F}} \right)} \quad (1.1)$$

where $F_\#$ is formed by composing the forgetful functor $\mathbb{A}^{\overleftarrow{F}} \rightarrow \mathbb{A}$ with the comparison functor $\mathbb{A} \rightarrow \mathbb{B}^{\overrightarrow{F}}$, and $F^\#$ by composing $\mathbb{B}^{\overrightarrow{F}} \rightarrow \mathbb{B}$ with $\mathbb{B} \rightarrow \mathbb{A}^{\overleftarrow{F}}$. Hence the left-hand square in Fig. 2. We prove that the right-hand square is an equivalence of adjunctions.

Upshot. The equivalences in Fig. 2 present $\overleftarrow{F}$ -algebras as algebras over $\overrightarrow{F}$ -coalgebras and $\overrightarrow{F}$ -coalgebras as coalgebras over $\overleftarrow{F}$ -algebras. Simplifying these presentations furnishes the *simple* nucleus construction in





Figure 1: An adjunction $(F^* \dashv F_*)$ is nuclear when $A \simeq B^{\overleftarrow{F}}$ and $B \simeq A^{\overrightarrow{F}}$.




Figure 2: The nucleus construction induces an idempotent monad on adjunctions.

Sec. 6, which opens up a new view of $\overleftarrow{F}$ -algebras and $\overrightarrow{F}$ -coalgebras, complementing the familiar Eilenberg-Moore construction [EM⁺65]. This new view was used as a programming tool in [PS17] and as a mathematical tool in [PH22]. The resulting reconstruction of monadicity and comonadicity in terms of idempotent splittings echoes Paré's explanations in terms of absolute colimits [Par69, Par71], and contrasts with Beck's fascinating but less transparent treatment in terms of split coequalizers [BW85, Bec67]. Applications branch in many directions, spanning a wide gamut from descent theory in algebraic geometry to concept analysis on the web. Curiously, the presented results did not evolve from pure mathematics to its applications, but the other way around. Conceptual advances were driven by the ideas evolving in response to practical problems.

1.2. Background. Nuclear adjunctions have been studied since the early days of category theory, albeit without a name. The problem of characterizing situations when the left adjoint of a monadic functor is comonadic is the topic of Michael Barr's paper in the proceedings of the legendary Battelle conference [Bar69]. Ever since, the coalgebras over algebras and the algebras over coalgebras have been emerging in the wide range of applications, from semantics of computation [BK11, Jac94, Jac13, and the references therein] on one end, to algebraic geometry [CDGV07, Mes06b] and homotopy theory on the other [Bal12, Hes10]. The monad-comonad couplings on two sides of an adjunction could be gleaned already on the primordial

examples of adjunctions [Kan58a] and the efforts to grasp them propelled the early research relating them with monads and comonads [AT70, Bec67, Hub61]. There are, however, different ways in which monad-comonad couplings arise from adjunctions, and from other monads and comonads. In [AT69], Applegate and Tierney considered monads and comonads formed on the two sides of the comparison functors from adjunctions to their final resolutions in the induced categories of coalgebras. They found that such couplings in general induce further such couplings and form transfinite towers. We describe this process in Sec. 7. Confusingly, Applegate-Tierney’s towers of adjunctions of comparison functors left a false impression that the adjunctions between algebras over coalgebras and coalgebras over algebras also lead to transfinite towers. This impression blended into folklore and spread through literature.¹ We show that the latter towers settle in one step, displayed in Fig. 2.

Name. Although they appeared in many avatars, nuclear adjunctions don’t seem to have been named. We call them nuclear in reference to the nuclear operators on Banach spaces, which generalize spectral decompositions of hermitians and the singular value decompositions of linear operators. The terminology was introduced in Grothendieck’s thesis [Gro55].




Figure 3: The paths to the nucleus

1.3. Overview. The paths to the nucleus are mapped in Fig. 3. They lead from observations tabulated in data Matrices to concepts collected in Nuclei. They lead through Adjunctions and then branch through Monads and Comonads. The Nuclei arise at the intersection of the two branches. Sec. 2 presents the posetal case, which is intuitive and well-known². The categorical setting is in Sec. 3. The nucleus functor is defined in Sec. 4. The fact that it is an idempotent monad is proved in Sec. 5. The *simple nucleus* construction, offering an alternative to the familiar Eilenberg-Moore presentation of algebras and coalgebras, is in Sec. 6. Sec. 7 touches upon the ways in which the nucleus concept feeds back into the ideas of descent and of

¹There is an interesting exception outside the literature. In a fax message sent to Paul Taylor on 9/9/99 [Lac], a copy of which was kindly provided after the present paper appeared on arxiv, Steve Lack set out to determine the conditions under which the tower of coalgebras over algebras, which "a priori continues indefinitely", settles to equivalence at a finite stage. Within 7 pages of diagrams, the question was reduced to splitting a certain idempotent. While the argument is succinct, it does seem to prove a claim which, together with its dual, seems equivalent to our Prop. 4.1. The claim was not pursued any further. This episode from the early life of the nucleus underscores its message: that a concept is within reach whenever there is an adjunction but it does need to be spelled out to be recognized.

²In the theory of locales, the word “nucleus” is sometimes used to refer to the meet-preserving closure operators and to the sublocales that they induce [Sim78, Joh82, §II.2.2–4]. This usage is unrelated to the present concept or its background.

homotopy-invariant space decompositions, from which it indirectly originates. The closing section returns to the big picture and comments about the remaining work.

2. POSETAL NUCLEI: FROM CONTEXTS TO CONCEPTS

2.1. Idea of Formal Concept Analysis. Consider a market with A sellers and B buyers. Their interactions are recorded in an adjacency matrix $A \times B \xrightarrow{\Phi} 2$, where 2 is the set $\{0, 1\}$, and the entry Φ_{ab} is 1 if the seller $a \in A$ at some point sold goods to the buyer $b \in B$; otherwise it is 0. It is often convenient to move between the adjacency matrices and the induced binary relations along the correspondence

$$\left(A \times B \xrightarrow{\Phi} 2 \right) \leftrightarrow \widehat{\Phi} = \{ \langle a, b \rangle \in A \times B \mid \Phi_{ab} = 1 \} \quad (2.1)$$

which allows writing $a\widehat{\Phi}b$ instead of $\Phi_{ab} = 1$. In the literature on Formal Concept Analysis (FCA) [CR04, GW99, GSW05], the adjacency matrices Φ and the corresponding relations $\widehat{\Phi}$ are called *contexts*. The latent *concepts* that they carry are extracted as pairs $\langle U, V \rangle \in \wp A \times \wp B$ where every seller in U is $\widehat{\Phi}$ -related to every buyer in V , and vice versa. Fig. 4 displays an example. The binary relation $\widehat{\Phi} \subseteq A \times B$ is presented as a




Figure 4: A context Φ , its four concepts, and their concept lattice

bipartite graph, and the pairs $\langle U, V \rangle$ are complete subgraphs. E.g., if buyers a_0 and a_4 have farms, and sellers b_1, b_2 and b_3 sell farming equipment, but seller b_0 does not, then the sets $U = \{a_0, a_4\}$ and $V = \{b_1, b_2, b_3\}$

span a complete subgraph of the bipartite graph Φ , which corresponds to the concept "farming". If the buyers from the set $U' = \{a_0, a_1, a_2, a_3\}$ own cars but the buyer a_4 does not, and the sellers $V' = \{b_0, b_1, b_2\}$ sell car accessories but the seller b_3 does not, then the sets $\langle U', V' \rangle$ span another complete subgraph, this time corresponding to the concept "car". The latent concepts are extracted from a context, viewed as a bipartite graph, as its complete subgraphs.

2.2. Formalizing concept analysis. The subgraph of a bipartite graph $\widehat{\Phi} \subseteq A \times B$ spanned by a pair of sets of nodes $\langle U, V \rangle \in \wp A \times \wp B$ is complete if

$$U = \bigcap_{v \in V} \{x \in A \mid x \widehat{\Phi} v\} \quad V = \bigcap_{u \in U} \{y \in B \mid u \widehat{\Phi} y\}$$

Such pairs are ordered by the relation

$$\langle U, V \rangle \leq \langle U', V' \rangle \iff U \subseteq U' \wedge V \supseteq V' \quad (2.2)$$

They are obviously a subposet of the lattice $\wp A \times \wp^o B$, where $\wp A$ is the set of subsets of A ordered by the inclusion $\subseteq$ and $\wp^o B$ is the set of subsets of B ordered by reverse inclusion $\supseteq$. We will see in a moment that they are also a quotient of this lattice, and thus form a lattice $\overleftarrow{\mathfrak{N}}_0 \Phi$. This is the *concept lattice* induced by the *context matrix* Φ .

2.3. Posetal matrices. The posetal case of Fig. 3 starts from the category

$$|\text{Mat}_0| = \coprod_{A, B \in \text{Pos}} \text{Pos}(A^o \times B, \mathbb{2}) \quad (2.3)$$

$$\text{Mat}_0(\Phi, \Psi) = \{\langle h, k \rangle \in \text{Pos}(A, C) \times \text{Pos}(B, D) \mid \Phi(a, b) = \Psi(ha, kb)\}$$

A, B, C, D are posets and $\Phi \in \text{Pos}(A^o \times B, \mathbb{2})$ and $\Psi \in \text{Pos}(C^o \times D, \mathbb{2})$ are matrices with entries from the poset $\mathbb{2} = \{0 < 1\}$. Note that the sets A of sellers and B of buyers, discussed above, generally come with partial orders, e.g. from previous concept analyses.

Comprehensive view. Generalizing (2.1), context matrices can also be viewed as subposets of product posets using the isomorphisms

$$\begin{aligned} \text{Pos}(A^o \times B, \mathbb{2}) & \xrightleftharpoons[\chi]{(-)} \text{Sub } A \times B^o \\ \Phi & \mapsto \widehat{\Phi} = \{\langle x, y \rangle \in A \times B^o \mid \Phi(x, y) = 1\} \\ \chi_S(x, y) = \begin{cases} 1 & \text{if } \langle x, y \rangle \in S \\ 0 & \text{otherwise} \end{cases} & \leftarrow (S \subseteq A \times B^o) \end{aligned} \quad (2.4)$$

The poset $\widehat{\Phi}$ can be thought of as the "comprehension" of Φ viewed as a "predicate" [Law70, Pav90, Sec. III.3]. The poset $\widehat{\Phi}$ refines the set-theoretic comprehension schema by imposing the order-closure requirement:

$$a \leq a' \wedge a' \widehat{\Phi} b' \wedge b' \leq b \implies a \widehat{\Phi} b \quad (2.5)$$

2.4. Completions make matrices representable as adjunctions. The concepts are extracted from a context $\widehat{\Phi} \subseteq A \times B$ by finding the pairs of a lower-closed subset of A and an upper-closed subset of B coupled by the context. The lower-closed and the upper-closed subsets of a poset form the complete semilattices $(\Downarrow A, \bigvee)$ and $(\Uparrow B, \bigwedge)$, where

$$\Downarrow A = \{X \subseteq A \mid a \leq a' \in X \implies a \in X\} \quad \Uparrow B = \{Y \subseteq B \mid Y \ni b' \leq b \implies Y \ni b\} \quad (2.6)$$

and $\bigvee$ in $\Downarrow A$ and $\bigwedge$ in $\Uparrow B$ are the unions. The embeddings

$$\begin{array}{ll} A \xrightarrow{\nabla} \Downarrow A & B \xrightarrow{\blacktriangle} \Uparrow B \\ a \mapsto \{x \leq a\} & b \mapsto \{y \geq b\} \end{array}$$

are respectively the $\bigvee$ -completion of A and the $\bigwedge$ -completion of B . These semilattice completions allow extending the context matrix $\widehat{\Phi} \subseteq A \times B$ from (2.1) along (2.4) to $\overline{\Phi} \subseteq \Downarrow A \times \Uparrow B$, setting

$$X \overline{\Phi} Y \iff \forall a \in X \forall b \in Y. a \widehat{\Phi} b \quad (2.7)$$

Since its domain and codomain are complete semilattices $\Downarrow A$ and $\Uparrow B$, the matrix $\overline{\Phi}$ is representable in the form

$$\Phi^* X \subseteq Y \iff X \overline{\Phi} Y \iff X \supseteq \Phi_* Y \quad (2.8)$$

The adjoints $\Phi^* \dashv \Phi_*$ extract the complete bipartite subgraphs, illustrated in Fig. 4, as intersections:

$$\begin{array}{ccccc} X & & \Downarrow A & & \bigcap_{y \in Y} \bullet \Phi y \\ \downarrow & & \updownarrow \begin{array}{c} \Phi^* \dashv \Phi_* \end{array} & & \uparrow \\ \bigcap_{x \in X} x \Phi \bullet & & \Uparrow B & & Y \end{array} \quad (2.9)$$

where $\bullet \Phi y = \{x \in A \mid x \Phi y\}$ and $x \Phi \bullet = \{y \in B \mid x \widehat{\Phi} y\}$ are the images along the transpositions

$$\frac{\Phi : A^o \times B \rightarrow \mathbb{2}}{\bullet \Phi : B \rightarrow \mathbb{2}^{A^o} \cong \Downarrow A \quad \Phi \bullet : A \rightarrow (\mathbb{2}^B)^o \cong \Uparrow B} \quad (2.10)$$

The poset adjunctions like (2.9) are often called *Galois connections*. They form the category

$$|\text{Adj}_0| = \coprod_{A, B \in \text{Pos}} \{ \langle \Phi^*, \Phi_* \rangle \in \text{Pos}(A, B) \times \text{Pos}(B, A) \mid \Phi^* x \leq y \iff x \leq \Phi_* y \} \quad (2.11)$$

$$\text{Adj}_0(\Phi, \Psi) = \{ \langle H, K \rangle \in \text{Pos}(A, C) \times \text{Pos}(B, D) \mid K \Phi^* = \Psi^* H \wedge H \Phi_* = \Psi_* K \}$$

The matrix extension at the first step of the posetal case of Fig. 3 is thus

$$\begin{array}{ll} \text{MA}_0 : \text{Mat}_0 & \longrightarrow \text{Adj}_0 \\ \Phi & \longmapsto (\Phi^* \dashv \Phi_* : \Uparrow B \rightarrow \Downarrow A) \text{ defined in (2.9)} \end{array} \quad (2.12)$$

2.5. Posetal adjunctions induce closure and interior operators. The further steps through the posetal Fig. 3 go to the categories of closure and interior operators, respectively:

$$\begin{aligned} |\mathbf{Mnd}_0| &= \coprod_{A \in \mathbf{Pos}} \left\{ \overleftarrow{\Phi} \in \mathbf{Pos}(A, A) \mid x \leq \overleftarrow{\Phi}x \geq \overleftarrow{\Phi}\overleftarrow{\Phi}(x) \right\} \\ \mathbf{Mnd}_0(\overleftarrow{\Phi}_A, \overleftarrow{\Psi}_C) &= \left\{ H \in \mathbf{Pos}(A, C) \mid H\overleftarrow{\Phi} = \overleftarrow{\Psi}H \right\} \end{aligned} \quad (2.13)$$

$$\begin{aligned} |\mathbf{Cmn}_0| &= \coprod_{B \in \mathbf{Pos}} \left\{ \overrightarrow{\Phi} \in \mathbf{Pos}(B, B) \mid x \geq \overrightarrow{\Phi}x \leq \overrightarrow{\Phi}\overrightarrow{\Phi}(x) \right\} \\ \mathbf{Cmn}_0(\overrightarrow{\Phi}_B, \overrightarrow{\Psi}_D) &= \left\{ K \in \mathbf{Pos}(B, D) \mid K\overrightarrow{\Phi} = \overrightarrow{\Psi}K \right\} \end{aligned} \quad (2.14)$$

along the functors

$$\begin{aligned} \mathbf{Cmn}_0 &\xleftarrow{\mathbf{AC}_0} \mathbf{Adj}_0 \xrightarrow{\mathbf{AM}_0} \mathbf{Mnd}_0 \\ \Phi^* \Phi_* &\longleftarrow (\Phi^* \dashv \Phi_*) \longrightarrow \Phi_* \Phi^* \end{aligned} \quad (2.15)$$

2.6. Closure and interior operators have adjunction resolutions. The functors from adjunctions to closure and interior operators have full and faithful left adjoints:

$$\begin{aligned} \mathbf{Cmn}_0 &\xrightarrow{\mathbf{EC}_0} \mathbf{Adj}_0 \xleftarrow{\mathbf{EM}_0} \mathbf{Mnd}_0 \\ (A \xrightleftharpoons[\top]{\top} A^{\overleftarrow{\Phi}}) &\longleftarrow \overleftarrow{\Phi}_A \\ \overrightarrow{\Phi}_B &\longmapsto (B^{\overrightarrow{\Phi}} \xleftarrow[\top]{\top} B) \end{aligned} \quad (2.16)$$

where the posets

$$A^{\overleftarrow{\Phi}} = \{x \in A \mid \overleftarrow{\Phi}x = x\} \quad B^{\overrightarrow{\Phi}} = \{y \in B \mid \overrightarrow{\Phi}y = y\} \quad (2.17)$$

inherit the order from A and B . The claimed adjunctions $\mathbf{AM}_0 \dashv \mathbf{EM}_0$ and $\mathbf{AC}_0 \dashv \mathbf{EC}_0$ are easily checked, as are the natural isomorphisms $\mathbf{AM}_0\mathbf{EM}_0 \cong \text{id}$ and $\mathbf{AC}_0\mathbf{EC}_0 \cong \text{id}$, making $\mathbf{Mnd}_0$ and $\mathbf{Cmn}_0$ into reflective subcategories of $\mathbf{Adj}_0$.

2.7. Concept lattices as nuclei. The posetal version of Fig. 3 is comprised of the following steps

$$\begin{aligned} &\Phi : A^o \times B \rightarrow \mathbb{2} \\ &\overline{\mathbf{MA}_0\Phi} = \Phi_*^* = \left(\Downarrow A \xrightleftharpoons[\Phi^*]{\Phi_*} \Uparrow B \right) \\ &\overleftarrow{\mathbf{EM}_0\Phi_*^*} = \left(\Downarrow A \xrightleftharpoons[\top]{\top} \Downarrow A^{\overleftarrow{\Phi}} \right) \quad \overleftarrow{\mathbf{EC}_0\Phi_*^*} = \left(\Uparrow B^{\overrightarrow{\Phi}} \xleftarrow[\top]{\top} \Uparrow B \right) \\ &\overleftarrow{\mathfrak{R}_0\Phi} = \left(\Uparrow B^{\overrightarrow{\Phi}} \xleftarrow[\Phi^\#]{\Phi^\#} \Downarrow A^{\overleftarrow{\Phi}} \right) \end{aligned} \quad (2.18)$$

where $\overleftarrow{\mathbf{EM}}_0 = \mathbf{EM}_0 \circ \mathbf{AM}_0$, and $\overleftarrow{\mathbf{EC}}_0 = \mathbf{EC}_0 \circ \mathbf{AC}_0$ are the idempotent monads displaying $\mathbf{Mnd}_0$ and $\mathbf{Cmn}_0$ as reflective subcategories of $\mathbf{Adj}_0$. A peculiarity of the posetal case is that they are also idempotent comonads, arising from the adjunctions $\mathbf{EM}_0 \dashv \mathbf{AM}_0$ and $\mathbf{EC}_0 \dashv \mathbf{AC}_0$, from which it follows not only that $\mathbf{Mnd}_0$ and

Cmn_0 are also coreflective in Adj_0 , but also that they are equivalent. A concept lattice can thus be obtained using $\overleftarrow{\text{EM}}_0$ and presented in terms of a closure operator, or using $\overleftarrow{\text{EC}}_0$ and presented in terms of an interior operator, or using $\overleftarrow{\mathfrak{N}}_0 \cong \overleftarrow{\text{EM}}_0 \overleftarrow{\text{EC}}_0 \cong \overleftarrow{\text{EC}}_0 \overleftarrow{\text{EM}}_0$ and presented in terms of an adjunction, reducing the context $\Phi: A^\circ \times B \xrightarrow{\Phi} \mathbb{2}$ to the concept lattice

$$\overleftarrow{\mathfrak{N}}_0 \Phi = \{ \langle X, Y \rangle \in \Downarrow A \times \Uparrow B \mid X = \Phi_* Y \wedge \Phi^* X = Y \} \quad (2.19)$$

The elements of this lattice generalize Dedekind cuts of rational numbers, and the nucleus construction generalizes the Dedekind-MacNeille completion to matrices [BB67, Ded01, Mac36]. All of this is well-known and valid not only for categories of posets but also for posetally enriched categories. Our present concern is the general case, where the categories Mnd and Cmn are not equivalent.

3. FROM CATEGORICAL CONTEXTS TO ADJUNCTIONS, MONADS, AND COMONADS

The need for categorical concept analysis. The posetal nuclei from the preceding section lift to poset-enriched categories with no major surprises, though with significant new applications [BTV22, Pav12, Pav13, Will13]. Large-scale concept analyses in recommender systems [RRS22], however, require going beyond the posetal and numeric contexts, to proper categories. The reason is that the posetal contexts do not capture out-of-band dependencies or hidden variables. The vector space models preclude hidden variables by imposing a tacit but very consequential assumption that all analyzed sources are statistically independent. This assumption is implied by the linearity of the operations on context matrices which aggregate correlations as sums of *products* of the observed frequencies. While the source interference is normally measured as the deviation from the product distribution, in the vector space model, the product distribution is built-in. This conceptual shortcoming, briefly described at the outset of our work [KP15], in the meantime grew into a practical problem of tremendous impact, leading to wholesale correlation amplifications, information cascades, and echo chambers. On the theoretical side, the task could be stated in terms of Grothendieck’s Galois descent [Gro60, Gro71, JT04, Vis05] but required substantial reinterpretation and narrowing. The basic ideas, sketched in [KP15, PS17], evolved bottom-up, from implementations to constructions. Here we attempt to provide a top-down account.

3.1. Lifting concept analysis to categories — and not further. Spanned over categories $\mathbb{A}$ and $\mathbb{B}$, data matrices become functors $\mathbb{A}^\circ \times \mathbb{B} \xrightarrow{\Phi} \text{Set}$, assigning to each pair of objects $\langle a, b \rangle$ the set Φ_{ab} of observations linking them. Whether the categories $\mathbb{A}$ and $\mathbb{B}$ represent sellers and buyers like in Sec. 2, or structural components and functional modules of a machine like in [KP15], or some other pair of types, the matrix entries record the interactions between the entities of type $\mathbb{A}$ and the entities of type $\mathbb{B}$ and the arrows in each of the categories capture the dependencies within each of the types. The arrow part of the functor Φ thus tracks how the dependencies propagate through the interactions. In category theory, the functors generalizing matrices in this way are variably called distributors, profunctors, or bimodules [Bor94, Vol. 1, Sec. 7.8]. We continue to call them matrices not so much to expand the terminological choices but to remember the application at hand.

The 2-dimensional structure. Given the matrices $\mathbb{A}^\circ \times \mathbb{B} \xrightarrow{\Phi} \text{Set}$ and $\mathbb{B}^\circ \times \mathbb{C} \xrightarrow{\Psi} \text{Set}$, the usual matrix composition is generalized using the *coend* operation

$$(\Phi; \Psi)_{ac} = \lim_{y \in \mathbb{B}} (\Phi_{ay} \times \Psi_{yc}) \quad (3.1)$$

See [Lor21] for details. The compositions accumulate and propagate data dependencies between $\mathbb{A}$ and $\mathbb{C}$ across the intermediary $\mathbb{B}$. The main point is that, for each $\mathbb{A}$ and $\mathbb{B}$, the $\mathbb{A}^\circ \times \mathbb{B}$ -matrices form a category recording their correlations, and that the correlations can be composed, functorially up to natural isomorphism.

The categorical matrices (i.e. distributors, profunctors, bimodules) thus form a *bicategory* [Bén67, Bor94, Vol. 1, Sec. 7.7]. The categories of adjunctions, monads, and comonads form 2-categories, meaning that the morphisms between any pair of adjunctions (resp. monads, comonads) also form a category, and the representable data correlations captured there also compose functorially, this time not up to isomorphism but strictly [Bor94, Vol. 1, Sec. 7.1]. The 2-categorical structure of adjunctions, monads, and comonads goes back to [Aud74, LS02, SS86, Str72], and could be gleaned already in [Mar66].

The nucleus construction is 1-dimensional. The nucleus construction combines monadic and comonadic adjunctions. To lift a monad morphism to a morphism between the induced monadic adjunctions, we use the 2-cell component of the monad morphism, which stipulates how the 1-cell distributes over the two monads. Ditto for lifting a comonad morphism to a morphism between the induced comonadic adjunctions. The problem that arises for the nucleus construction is that the two distributivities, one needed to map monad morphisms to adjunction morphisms between final resolutions, the other one to map the comonad morphisms similarly, are required to run in opposite directions. To compose the two 2-monads on the 2-category of adjunctions, one making them monadic, the other making them comonadic, the 2-dimensional structure must be restricted to invertible distributivity 2-cells. But then the invertible distributivities can be absorbed into 1-cells and the 2-cells can be left out. This will turn out to be the feature making the nucleus monad strongly idempotent. In 2-dimensional category theory, a monad is usually weakly idempotent, in the lax sense that its algebras are adjoint to the unit [Koc95, Str80, Zöb76]. The nucleus monad is strongly idempotent, in the sense that its algebras are equivalences. It is a categorical construction, not 2-categorical. The following definitions and the results that they enable provide the evidence.

3.2. Matrices (distributors, profunctors, bimodules). The category of matrices over categories with the morphisms up to invertible 2-cells is defined in Fig. 5.

$$\begin{aligned}
 |\mathbf{Mat}| &= \coprod_{\mathbf{A}, \mathbf{B} \in \mathbf{CAT}} \mathbf{CAT}(\mathbf{A}^o \times \mathbf{B}, \mathbf{Set}) \\
 \mathbf{Mat}(\Phi, \Psi) &= \coprod_{\substack{H \in \mathbf{CAT}(\mathbf{A}, \mathbf{C}) \\ K \in \mathbf{CAT}(\mathbf{B}, \mathbf{D})}} \left\{ \begin{array}{ccc} \mathbf{A}^o \times \mathbf{B} & \xrightarrow{H^o \times K} & \mathbf{C}^o \times \mathbf{D} \\ & \searrow \Phi \quad \swarrow \Psi & \\ & \mathbf{Set} & \end{array} \right\}
 \end{aligned}$$

Figure 5: $\gamma \in \text{Iso}(\Phi, \Psi(H^o \times K))$ are natural isomorphisms

Comprehensive view. The posetal comprehension (2.4) lifts in categories to the equivalence of presheaves and discrete fibrations

$$\mathbf{Cat}(\mathbf{A}^o \times \mathbf{B}, \mathbf{Set}) \begin{array}{c} \xrightarrow{(-)} \\ \simeq \\ \xleftarrow{\Xi} \end{array} \mathbf{Dfib} / \mathbf{A} \times \mathbf{B}^o \quad (3.2)$$

The details are similar to the posetal case and also widely known, as the categorical fibrations, generalizing the posetal lower sets, go back to Grothendieck [Gro60, Gro71] and have been reviewed many times [Jac01, Pav90, Str18]. For completeness, we reproduce the details of (3.2) in Appendix A. Since switching between the $\mathbf{Set}$ -functors and their comprehensions is as routinely in category theory as the switching between the predicates as $\mathbb{2}$ -functions and their comprehensions is in logic, we often use $\Downarrow \mathbf{A}$ and $\Uparrow \mathbf{B}$ to refer to both sides of the equivalences

$$\mathbf{Set}^{\mathbf{A}^o} \simeq \Downarrow \mathbf{A} = \mathbf{Dfib} / \mathbf{A} \quad (\mathbf{Set}^{\mathbf{B}})^o \simeq \Uparrow \mathbf{B} = (\mathbf{Dfib} / \mathbf{B}^o)^o \quad (3.3)$$

where $\mathbf{Set}^{\mathbb{X}}$ abbreviates the category $\mathbf{Cat}(\mathbb{X}, \mathbf{Set})$. The embeddings

$$\begin{array}{ccc} \mathbb{A} & \xrightarrow{\blacktriangledown} & \Downarrow \mathbb{A} \\ a & \mapsto & \left(\mathbb{A}/a \xrightarrow{\text{Dom}} \mathbb{A} \right) \end{array} \qquad \begin{array}{ccc} \mathbb{B} & \xrightarrow{\blacktriangle} & \Uparrow \mathbb{B} \\ b & \mapsto & \left(b/\mathbb{B} \xrightarrow{\text{Cod}} \mathbb{B} \right) \end{array}$$

are this time, respectively, the $\varinjlim$ -completion of $\mathbb{A}$ and the $\varprojlim$ -completion of $\mathbb{B}$. As usually, $\mathbb{A}/a$ denotes the category of $\mathbb{A}$ -arrows into a , whereas $b/\mathbb{B}$ denotes the category of $\mathbb{B}$ arrows from b . The fact that $\Downarrow \mathbb{A}$ is a $\varinjlim$ -completion and that $\Uparrow \mathbb{B}$ is a $\varprojlim$ -completion means that every $(\mathbb{X} \xrightarrow{X} \mathbb{A}) \in \Downarrow \mathbb{A}$ and $(\mathbb{Y} \xrightarrow{Y} \mathbb{B}) \in \Uparrow \mathbb{B}$ satisfy

$$X = \varinjlim \left(\mathbb{X} \xrightarrow{X} \mathbb{A} \xrightarrow{\blacktriangledown} \Downarrow \mathbb{A} \right) \qquad Y = \varprojlim \left(\mathbb{Y} \xrightarrow{Y} \mathbb{B} \xrightarrow{\blacktriangle} \Uparrow \mathbb{B} \right) \quad (3.4)$$

3.3. Completions make matrices representable as adjunctions again. A matrix $\Phi: \mathbb{A}^o \times \mathbb{B} \rightarrow \mathbf{Set}$ can be represented as the following adjunction between the completions

$$\begin{array}{ccccc} \mathbb{X} \xrightarrow{X} \mathbb{A} & & \Downarrow \mathbb{A} & & \varprojlim \left(\mathbb{Y} \xrightarrow{Y} \mathbb{B} \xrightarrow{\bullet\Phi} \Downarrow \mathbb{A} \right) \\ \downarrow & & \Phi^* \left(\begin{array}{c} \uparrow \\ \downarrow \end{array} \right) \Phi_* & & \uparrow \\ \varinjlim \left(\mathbb{X} \xrightarrow{X} \mathbb{A} \xrightarrow{\Phi_\bullet} \Uparrow \mathbb{B} \right) & & \Uparrow \mathbb{B} & & \mathbb{Y} \xrightarrow{Y} \mathbb{B} \end{array} \quad (3.5)$$

where $\Phi_\bullet$ and $\bullet\Phi$ are the two transposes of Φ

$$\begin{array}{c} \Phi: \mathbb{A}^o \times \mathbb{B} \rightarrow \mathbf{Set} \\ \hline \Phi_\bullet: \mathbb{A} \rightarrow (\mathbf{Set}^{\mathbb{B}})^o \simeq \Uparrow \mathbb{B} \qquad \bullet\Phi: \mathbb{B} \rightarrow \mathbf{Set}^{\mathbb{A}^o} \simeq \Downarrow \mathbb{A} \\ \hline \Phi^*: \Downarrow \mathbb{A} \longrightarrow \Uparrow \mathbb{B} \qquad \Phi_*: \Uparrow \mathbb{B} \longrightarrow \Downarrow \mathbb{A} \end{array} \quad (3.6)$$

and the definitions of Φ^* and Φ_* are based on the fact that any left adjoint must preserve $\varinjlim$, and that any right adjoint must preserve $\varprojlim$. In other words, Φ^* is the left Kan extension of $\Phi_\bullet$ along $\blacktriangledown: \mathbb{A} \rightarrow \Downarrow \mathbb{A}$, whereas Φ_* is the right Kan extension of $\bullet\Phi$ along $\blacktriangle: \mathbb{B} \rightarrow \Uparrow \mathbb{B}$. It is useful to notice that both are computed pointwise, just like both extensions in (2.9) are computed as set intersections. The reason is that dualizing along the equivalence $\Uparrow \mathbb{B} \simeq (\mathbf{Set}^{\mathbb{B}})^o$ gives

$$\varinjlim \left(\mathbb{X} \xrightarrow{X} \mathbb{A} \xrightarrow{\Phi_\bullet} \Uparrow \mathbb{B} \right) = \varprojlim \left(\mathbb{X}^o \xrightarrow{X^o} \mathbb{A}^o \xrightarrow{\Phi_\bullet^o} \mathbf{Set}^{\mathbb{B}} \right)$$

But the limits in $\mathbf{Set}^{\mathbb{B}}$ are pointwise, since the Yoneda lemma gives for any diagram $D: \mathbb{D} \rightarrow \mathbf{Set}^{\mathbb{B}}$

$$\left(\varprojlim D \right) b = \mathbf{Set}^{\mathbb{B}} \left(\blacktriangle b, \varprojlim D \right) = \text{Cones}(b, D)$$

In words, the limit of D is the functor $\mathbb{B} \rightarrow \mathbf{Set}$ whose value at b is the set of commutative cones from b to D . The matrix extension

$$\begin{array}{ccc} \mathbf{MA}: \mathbf{Mat} & \longrightarrow & \mathbf{Adj} \\ \Phi & \longmapsto & (\Phi^* \dashv \Phi_*: \Uparrow \mathbb{B} \rightarrow \Downarrow \mathbb{A}) \text{ defined in (3.5)} \end{array} \quad (3.7)$$

takes us to the category of adjunctions in Fig. 6.

$$\begin{aligned}
|\text{Adj}| &= \coprod_{\mathbb{A}, \mathbb{B} \in \text{CAT}} \coprod_{\substack{F^* \in \text{CAT}(\mathbb{A}, \mathbb{B}) \\ F_* \in \text{CAT}(\mathbb{B}, \mathbb{A})}} \left\{ \left\langle \begin{array}{c} \eta \\ \varepsilon \end{array} \right\rangle \in \begin{array}{c} \text{Nat}(\text{Id}, F_* F^*) \\ \times \\ \text{Nat}(F^* F_*, \text{Id}) \end{array} \mid \begin{array}{c} F_* F^* F_* \\ \begin{array}{ccc} \nearrow \eta F_* & & \nwarrow F_* \varepsilon \\ F_* & \xlongequal{\quad} & F_* \\ \nwarrow F^* \eta & & \nearrow \varepsilon F^* \\ F^* F_* F^* & & \end{array} \end{array} \right\} \\
\text{Adj}(F, G) &= \coprod_{\substack{H \in \text{CAT}(\mathbb{A}, \mathbb{C}) \\ K \in \text{CAT}(\mathbb{B}, \mathbb{D})}} \left\{ \left\langle \begin{array}{c} v^* \\ v_* \end{array} \right\rangle \in \begin{array}{c} \text{Iso}(KF^*, G^* H) \\ \times \\ \text{Iso}(HF_*, G_* K) \end{array} \mid \begin{array}{ccccc} HF_* F^* & G^* G_* K & \xrightarrow{\varepsilon^G K} & K & \\ \uparrow \eta^F & \uparrow v_* F^* & \uparrow G^* v_* & \uparrow K \varepsilon^F & \\ H \eta^F & G_* K F^* & G^* H F_* & K \varepsilon^F & \\ \downarrow & \downarrow G_* v^* & \downarrow v^* F_* & \downarrow & \\ H & \xrightarrow{\eta^G H} & G_* G^* H & K F^* F_* & \end{array} \right\}
\end{aligned}$$

Figure 6: $\text{Nat}(X, Y)$ are the natural transformations, $\text{Iso}(X, Y)$ the natural isomorphisms from X to Y

$$\begin{aligned}
|\text{Mnd}| &= \coprod_{\mathbb{A} \in \text{CAT}} \coprod_{\overleftarrow{T} \in \text{CAT}(\mathbb{A}, \mathbb{A})} \left\{ \left\langle \begin{array}{c} \eta \\ \mu \end{array} \right\rangle \in \begin{array}{c} \text{Nat}(\text{Id}, \overleftarrow{T}) \\ \times \\ \text{Nat}(\overleftarrow{T} \overleftarrow{T}, \overleftarrow{T}) \end{array} \mid \begin{array}{c} \overleftarrow{T} \overleftarrow{T} \overleftarrow{T} \\ \begin{array}{ccc} \nwarrow \overleftarrow{T} \mu & & \swarrow \mu \overleftarrow{T} \\ \overleftarrow{T} \overleftarrow{T} & \xrightarrow{\mu} & \overleftarrow{T} \\ \nwarrow \eta \overleftarrow{T} & \parallel & \swarrow \overleftarrow{T} \eta \\ \overleftarrow{T} & & \overleftarrow{T} \end{array} \end{array} \right\} \\
\text{Mnd}(\overleftarrow{T}_{\mathbb{A}}, \overleftarrow{S}_{\mathbb{C}}) &= \coprod_{H \in \text{CAT}(\mathbb{A}, \mathbb{C})} \left\{ \chi \in \text{Iso}(H \overleftarrow{T}, \overleftarrow{S} H) \mid \begin{array}{ccccc} & & H \overleftarrow{T} & \leftarrow H \mu^T - & H \overleftarrow{T} \overleftarrow{T} \\ & \nearrow H \eta^T & \uparrow \chi & \uparrow \chi^T & \uparrow \overleftarrow{S} H \overleftarrow{T} \\ H & & \downarrow \chi & \downarrow \overleftarrow{S} \chi & \downarrow \overleftarrow{S} \chi \\ & \searrow \eta^S H & \downarrow \overleftarrow{S} H & \leftarrow \mu^S H - & \overleftarrow{S} \overleftarrow{S} H \end{array} \right\}
\end{aligned}$$

Figure 7: Monads generalize closure operators (2.13).

3.4. From adjunctions to monads and comonads. Lifting definitions (2.13–2.14) from posets to categories requires imposing the commutativity conditions in Fig. 7, which were automatic in posets. The object parts of the functors generalizing (2.15) are now

$$\begin{array}{ccccc}
\text{Cmn} & \xleftarrow{\text{AC}} & \text{Adj} & \xrightarrow{\text{AM}} & \text{Mnd} \\
\langle F^* F_*, \varepsilon, \nu \rangle & \longleftarrow & \langle F^*, F_*, \eta, \varepsilon \rangle & \longrightarrow & \langle F_* F^*, \eta, \mu \rangle
\end{array} \tag{3.8}$$

$$\begin{aligned}
|\text{Cmn}| &= \coprod_{\mathbb{B} \in \text{CAT}} \coprod_{\vec{T} \in \text{CAT}(\mathbb{B}, \mathbb{B})} \left\{ \left\langle \begin{array}{c} \varepsilon \\ \nu \end{array} \right\rangle \in \begin{array}{c} \text{Nat}(\vec{T}, \text{Id}) \\ \times \\ \text{Nat}(\vec{T}, \vec{T}\vec{T}) \end{array} \mid \begin{array}{c} \vec{T}\vec{T}\vec{T} \\ \nearrow \quad \nwarrow \\ \vec{T}\vec{T} \xleftarrow{\nu} \vec{T} \xrightarrow{\nu} \vec{T}\vec{T} \\ \nwarrow \quad \nearrow \\ \vec{T} \end{array} \right\} \\
\text{Cmn}(\vec{T}_{\mathbb{B}}, \vec{S}_{\mathbb{D}}) &= \coprod_{K \in \text{CAT}(\mathbb{B}, \mathbb{D})} \left\{ \kappa \in \text{Iso}(K\vec{T}, \vec{S}K) \mid \begin{array}{c} K\vec{T} \xrightarrow{K\varepsilon^T} K\vec{T} \xrightarrow{K\nu^T} K\vec{T}\vec{T} \\ \nwarrow \quad \uparrow \quad \nearrow \\ \vec{S}K \xrightarrow{\varepsilon^S K} \vec{S}K \xrightarrow{K\nu^S} \vec{S}\vec{S}K \\ \nwarrow \quad \downarrow \quad \nearrow \\ K \end{array} \right\}
\end{aligned}$$

Figure 8: Comonads generalize interior operators (2.14).

where the the comonad (chain) evaluation ν and the monad (cochain) evaluation μ are defined

$$\mu = \left(F_* F^* F_* F^* \xrightarrow{F_* \varepsilon F^*} F_* F^* \right) \quad \nu = \left(F^* F_* \xrightarrow{F^* \eta F_*} F^* F_* F^* F_* \right) \quad (3.9)$$

The arrow part of (3.8), whereby the adjunction morphisms induce the monad and comonad morphisms, is displayed in Fig. 9.




Figure 9: $\kappa = \left(KF_* F^* \xleftrightarrow{\nu^* F_*} G^* HF_* \xleftrightarrow{G^* \nu_*} G^* G_* K \right)$ and $\chi = \left(HF_* F^* \xleftrightarrow{\nu_* F^*} G_* KF^* \xleftrightarrow{G_* \nu^*} G_* G^* H \right)$

3.5. From monads and comonads to adjunctions. To lift (2.16) from posets to categories, we first replace the fixpoint posets in (2.17) with the categories of algebras $\mathbb{A}^{\overleftarrow{T}}$ and coalgebras $\mathbb{B}^{\overrightarrow{T}}$, as specified in Fig. 10. The object parts of EC and EM are displayed in Figures 11 and 13. The arrow parts are in Fig. 12, where

$$|\mathbb{A}^{\overleftarrow{T}}| = \coprod_{a \in |\mathbb{A}|} \left\{ \alpha \in \mathbb{A}(\overleftarrow{T}a, a) \mid \begin{array}{ccc} \overleftarrow{T}\overleftarrow{T}a & \xrightarrow{\mu} & \overleftarrow{T}a \\ \downarrow \overleftarrow{T}\alpha & \nearrow \eta & \downarrow \alpha \\ \overleftarrow{T}a & \xrightarrow{\alpha} & a \end{array} \right\} \quad |\mathbb{B}^{\overrightarrow{T}}| = \coprod_{b \in |\mathbb{B}|} \left\{ \beta \in \mathbb{B}(b, \overrightarrow{T}b) \mid \begin{array}{ccc} \overrightarrow{T}\overrightarrow{T}b & \xleftarrow{\nu} & \overrightarrow{T}b \\ \uparrow \overrightarrow{T}\beta & \nwarrow \varepsilon & \uparrow \beta \\ \overrightarrow{T}b & \xleftarrow{\beta} & b \end{array} \right\}$$

$$\mathbb{A}^{\overleftarrow{T}}(\alpha, \gamma) = \left\{ h \in \mathbb{A}(a, c) \mid \begin{array}{ccc} \overleftarrow{T}a & \xrightarrow{\overleftarrow{T}h} & \overleftarrow{T}c \\ \downarrow \alpha & & \downarrow \gamma \\ a & \xrightarrow{h} & c \end{array} \right\} \quad \mathbb{B}^{\overrightarrow{T}}(\beta, \delta) = \left\{ k \in \mathbb{B}(b, d) \mid \begin{array}{ccc} \overrightarrow{T}b & \xleftarrow{\overrightarrow{T}k} & \overrightarrow{T}c \\ \uparrow \beta & & \uparrow \delta \\ b & \xleftarrow{k} & d \end{array} \right\}$$

Figure 10: The final resolutions of the monad $\overleftarrow{T}$ and the comonad $\overrightarrow{T}$




Figure 11: The object part of EM and EC. The U -functors are specified in Fig. 13.

where K^κ and H^χ are

$$\frac{b \xrightarrow{\beta} \overrightarrow{T}b}{K^\kappa \beta = \left(Kb \xrightarrow{K\beta} K\overrightarrow{T}b \xleftarrow{\kappa} \overrightarrow{T}Kb \right)} \quad \frac{\overleftarrow{T}a \xrightarrow{\alpha} a}{H^\chi \alpha = \left(\overleftarrow{T}Ha \xleftarrow{\chi} H\overleftarrow{T}a \xrightarrow{H\alpha} Ha \right)} \quad (3.10)$$

The functors EC and EM require opposite distributivities. Note that the functor components K^κ and H^χ of $\text{EC}(K, \kappa)$ and $\text{EM}(H, \chi)$, displayed in Fig. 12 and defined in (3.10), require different directions of κ and χ , because

- $\langle K, \kappa \rangle \in \text{Cmn}(\overrightarrow{S}, \overrightarrow{T})$ lifts $K: \mathbb{B} \rightarrow \mathbb{D}$ to $K^\kappa: \mathbb{B}^{\overrightarrow{S}} \rightarrow \mathbb{D}^{\overrightarrow{T}}$ using $\kappa: K\overrightarrow{S} \rightarrow \overrightarrow{T}K$, whereas
- $\langle H, \chi \rangle \in \text{Mnd}(\overleftarrow{S}, \overleftarrow{T})$ lifts $H: \mathbb{A} \rightarrow \mathbb{C}$ to $H^\chi: \mathbb{A}^{\overleftarrow{S}} \rightarrow \mathbb{C}^{\overleftarrow{T}}$ using $\chi: \overleftarrow{T}H \rightarrow H\overleftarrow{S}$.




Figure 12: The arrow part of the functors in Fig. 11

Although κ and χ are 2-cell components of morphisms in different categories, $\mathbf{Cmn}$ and $\mathbf{Mnd}$ respectively, their images $\mathbf{EC}(K, \kappa)$ and $\mathbf{EM}(H, \chi)$ land in the same category $\mathbf{Adj}$. To accommodate both $\mathbf{EC}$ and $\mathbf{EM}$, the 2-cell components of the morphisms in $\mathbf{Adj}$, and consequently in $\mathbf{Mnd}$, and $\mathbf{Cmn}$, are required to be invertible. The functors $\mathbf{KC}$ and $\mathbf{KM}$, mentioned in the following section and described in detail later, also require opposite distributivities, as do $\mathbf{KC}$ and $\mathbf{EC}$ on one hand, and $\mathbf{KM}$ and $\mathbf{EM}$ on the other.




Figure 13: The adjunction units $\mathbf{EC} \circ \mathbf{AC}(F) \leftarrow F \rightarrow \mathbf{EM} \circ \mathbf{AM}(F)$ of $\mathbf{AC} \dashv \mathbf{EC}$ and $\mathbf{AM} \dashv \mathbf{EM}$

3.6. Monads and comonads are reflective in adjunctions. The adjunction equipment of

$$\mathbf{AC} \dashv \mathbf{EC}: \mathbf{Cmn} \rightarrow \mathbf{Adj} \quad \text{and} \quad \mathbf{AM} \dashv \mathbf{EM}: \mathbf{Mnd} \rightarrow \mathbf{Adj} \quad (3.11)$$

is displayed in Fig. 13, with the *comparison* functions defined

$$H^0 a = \left(F^* a \xrightarrow{F^* \eta} F^* F_* F^* a \right) \quad \text{and} \quad H_1 b = \left(F_* F^* F_* b \xrightarrow{F_* \varepsilon} F_* b \right) \quad (3.12)$$

The constructions underlying the functors $\mathbf{AC}$ and $\mathbf{AM}$ go back to Huber [Hub62]. Soon after Huber, Kleisli in [Kle65], and Eilenberg and Moore in [EM⁺65] provided two different inverses of Huber's constructions, decomposing arbitrary (co)monads into adjunctions. The construction by Eilenberg and Moore induces the

functors EC and EM above, whereas Kleisli's construction induces functors KC and KM, which turned out to be adjoint to AC and AM on the other side:

$$\text{KM} \dashv \text{AM} \dashv \text{EM} \quad \text{and} \quad \text{KC} \dashv \text{AC} \dashv \text{EC} \quad (3.13)$$

The (co)monad morphisms and the adjunction morphisms with respect to which Kleisli's and Eilenberg-Moore's constructions are functorial were introduced in [Mar66] and the induced 2-categories were studied in [Aud74, Str72]. The details are still being studied [BVMV11, VT10, e.g.]. Lifting ideas from homological algebra, the multiple adjunctions that induce the same (co)monad have been called its *resolutions* [LS86, Sec. 0.6]. It was established already in [Mar66] that Kleisli's resolution was initial, whereas Eilenberg-Moore's was final, meaning that the two constructions provide the adjoint reflections displayed in Fig. 14. Note that Fig. 3 only displays half of the monad and comonad resolution picture³. Since both Kleisli's and




Figure 14: Both comonads and monads are reflective and coreflective in adjunctions

Eilenberg-Moore's constructions invert Huber's

$$\text{AC} \circ \text{EC} = \text{AC} \circ \text{KC} = \text{Id}_{\text{Cmn}} \quad \text{and} \quad \text{AM} \circ \text{EM} = \text{AM} \circ \text{KM} = \text{Id}_{\text{Mnd}} \quad (3.14)$$

they induce idempotent monads

$$\begin{aligned} \overleftarrow{\text{EC}} &= \text{EC} \circ \text{AC} & \overleftarrow{\text{EM}} &= \text{EM} \circ \text{AM} \\ \overrightarrow{\text{KC}} &= \text{KC} \circ \text{AC} & \overrightarrow{\text{KM}} &= \text{KM} \circ \text{AM} \end{aligned}$$

The first two are studied next, in Thm. 5.1. For the record, note that both monads and comonads are localized as subcategories of adjunctions in *two* extremal ways and provide a typical example of Lawvere's *unity-and-identity of the opposites* [KL89, Law89, Law96].

4. THE NUCLEUS FUNCTOR

Fig. 15 defines the object part of the nucleus construction $\overleftarrow{\mathfrak{N}}$. Fig. 16 defines the arrow part, which is




Figure 15: The object part of $\overleftarrow{\mathfrak{N}}$ is comprised of $F^\sharp = H_1 \circ V_{\overleftarrow{F}}$ and $F_\sharp = H^0 \circ U_{\overleftarrow{F}}$ as defined in Fig. 11

³This is mostly to avoid clutter, and because the initial resolutions only really concern us towards the end.

$$\Upsilon = \langle H, K, v^*, v_* \rangle : F \rightarrow G$$

$$\overleftarrow{\mathfrak{N}}(\Upsilon) = \langle K^\nu, H^\nu, v^\sharp, v_\sharp \rangle : \overleftarrow{\mathfrak{N}}(F) \rightarrow \overleftarrow{\mathfrak{N}}(G)$$

$$\frac{b \xrightarrow{\beta} F^* F_* b}{K^\nu \beta = \left(Kb \xrightarrow{K\beta} KF^* F_* b \xleftarrow{v^* F_*} G^* HF_* b \xleftarrow{G^* v_*} G^* G_* Kb \right)}$$

$$v^\sharp = G^* (G_* v^* \circ v_* F^*)$$

$$\frac{F_* F^* a \xrightarrow{\alpha} a}{H^\nu \alpha = \left(G_* G^* Ha \xleftarrow{G_* v^*} G_* KF^* a \xleftarrow{v_* F^*} HF_* F^* a \xrightarrow{H\alpha} Ha \right)}$$

$$v_\sharp = G_* (G^* v_* \circ v F_*)$$




Figure 16: The arrow part of $\overleftarrow{\mathfrak{N}}$ maps $\langle H, K, v^*, v_* \rangle$ to $\langle K^\nu, H^\nu, v^\sharp, v_\sharp \rangle$ as defined in (3.10)

Prop. 4.1 below claims that Figures 15 and 16 together define a functor $\overleftarrow{\mathfrak{N}} : \mathbf{Adj} \rightarrow \mathbf{Adj}$. Before proving that, we discharge a useful lemma that follows from the definition of $F^\sharp$ and $F_\sharp$ alone.

4.1. Nucleus decomposes descent monads and comonads. Descent monads and comonads emerge as a common denominator of a variety of structures echoing the monad-comonad couplings in descent theory [Bal12, CDGV07, Hes10, Mes06a]. They will come in handy in Sections 5.2 and 7.1.

Definition 4.1. For an adjunction $F = (F^* \dashv F_* : \mathbb{B} \rightarrow \mathbb{A})$, the monad $\overleftarrow{F} = F_* F^*$ on $\mathbb{A}$ and $\overrightarrow{F} = F^* F_*$ on $\mathbb{B}$ with the final (Eilenberg-Moore) resolutions

$$U(F) = \left(U^{\overleftarrow{F}} \dashv U_{\overleftarrow{F}} : \mathbb{A}^{\overleftarrow{F}} \rightarrow \mathbb{A} \right) \quad V(F) = \left(V^{\overrightarrow{F}} \dashv V_{\overrightarrow{F}} : \mathbb{B} \rightarrow \mathbb{B}^{\overrightarrow{F}} \right)$$

the induced comonad and monad of F -descent are defined

$$\overrightarrow{\overleftarrow{F}} = U^{\overleftarrow{F}} U_{\overleftarrow{F}} : \mathbb{A}^{\overleftarrow{F}} \rightarrow \mathbb{A}^{\overleftarrow{F}} \quad \overleftarrow{\overrightarrow{F}} = V_{\overrightarrow{F}} V^{\overrightarrow{F}} : \mathbb{B}^{\overrightarrow{F}} \rightarrow \mathbb{B}^{\overrightarrow{F}}$$

Lemma 4.2. The descent monad $\overrightarrow{\overleftarrow{F}}$ and the descent comonad $\overleftarrow{\overrightarrow{F}}$ also arise as composites of the nucleus components $F^\sharp = H_1 \circ V^{\overrightarrow{F}}$ and $F_\sharp = H^0 \circ U^{\overleftarrow{F}}$.

$$U^{\overleftarrow{F}} U_{\overleftarrow{F}} = \overrightarrow{\overleftarrow{F}} = F^\sharp F_\sharp \quad V_{\overrightarrow{F}} V^{\overrightarrow{F}} = \overleftarrow{\overrightarrow{F}} = F_\sharp F^\sharp \quad (4.1)$$

Proof. Fig. 17 and the definitions of $F^\sharp$ and $F_\sharp$ give

$$F^\sharp F_\sharp = H_1 V^{\vec{F}} H^0 U^{\overleftarrow{F}} = U^{\overleftarrow{F}} U_{\overleftarrow{F}} \quad F_\sharp F^\sharp = H^0 U^{\overleftarrow{F}} H_1 V^{\vec{F}} = V_{\vec{F}} V^{\vec{F}}$$

where $H_1 V^{\vec{F}} H^0 = U^{\overleftarrow{F}}$ and $H^0 U^{\overleftarrow{F}} H_1 = V_{\vec{F}}$ follow from (3.12). $\square$




Figure 17: The monad $\overleftarrow{\overleftarrow{F}} = V_{\vec{F}} V^{\vec{F}}$ and the comonad $U^{\overleftarrow{F}} U_{\overleftarrow{F}} = \overrightarrow{\overrightarrow{F}}$ also decompose by $F_\sharp$ and $F^\sharp$

4.2. Nucleus is an adjunction.

Proposition 4.1. *The nucleus functor $\overleftarrow{\mathfrak{N}}: \text{Adj} \rightarrow \text{Adj}$ is well-defined:*

- $\overleftarrow{\mathfrak{N}}(F)$, specified in Fig. 15, is an adjunction; and
- $\overleftarrow{\mathfrak{N}}(H, K, v^*, v_*)$ in Fig. 16 is an adjunction morphism.

Proof. Checking that $\langle K^\nu, H^\nu, v^\sharp, v_\sharp \rangle$ satisfies the two commutativity requirements of adjunction morphisms in Fig. 6, as soon as $\langle H, K, v^*, v_* \rangle$ satisfies them, is straightforward. Checking the adjunction equipment of $F^\sharp \dashv F_\sharp$ requires work. The object parts of the functors $F_\sharp$ and $F^\sharp$ from Fig. 15 are unfolded in Fig. 18 again. The arrow part of $F_\sharp$ is the same as F^* and the arrow part of $F^\sharp$ is the same as F_* . For the functors $F^\sharp$ and $F_\sharp$, we now prove that the correspondence

$$\begin{aligned} \mathbb{A}^{\overleftarrow{F}}(F^\sharp \beta, \alpha) &\cong \mathbb{B}^{\vec{F}}(\beta, F_\sharp \alpha) \\ f &\mapsto \bar{f} = F^* f \circ \beta \end{aligned} \quad (4.2)$$

is a natural bijection. More precisely, the claim is that

- a) f is an algebra homomorphism if and only if $\bar{f}$ is a coalgebra homomorphism: each of the following squares commutes if and only if the other one commutes

$$\begin{array}{ccc} F_* F^* F_* y & \xrightarrow{F_* F^* f} & F_* F^* x \\ \downarrow F_* \varepsilon = F^\sharp \beta & & \downarrow \alpha \\ F_* y & \xrightarrow{f} & x \end{array} \quad \Longleftrightarrow \quad \begin{array}{ccc} F^* F_* y & \xrightarrow{F^* F_* \bar{f}} & F^* F_* F^* x \\ \uparrow \beta & & \uparrow F_\sharp \alpha = F^* \eta \\ y & \xrightarrow{\bar{f}} & F^* x \end{array} \quad (4.3)$$

$$\begin{array}{c}
x \vdash \dashrightarrow \left(\begin{array}{c} F^* x \\ \downarrow F^* \eta \\ F^* F_* F^* x \end{array} \right) \longleftarrow \left(\begin{array}{c} F_* F^* x \\ \downarrow \alpha \\ x \end{array} \right) \\
\\
A \overset{H^0}{\dashrightarrow} B \overset{\vec{F}}{\longrightarrow} A \overset{\overleftarrow{F}}{\longleftarrow} A \\
\quad \quad U_* \curvearrowright \\
\\
B \overset{V^*}{\dashrightarrow} A \overset{\overleftarrow{F}}{\longleftarrow} B \overset{\vec{F}}{\longrightarrow} B \\
\quad \quad H_1 \curvearrowright \\
\\
y \vdash \dashrightarrow \left(\begin{array}{c} F_* F^* F_* y \\ \downarrow F_* \varepsilon \\ F_* y \end{array} \right) \longleftarrow \left(\begin{array}{c} y \\ \downarrow \beta \\ F^* F_* y \end{array} \right)
\end{array}$$

Figure 18: The definitions of $F_{\#}$ and $F^{\#}$

b) the map $f \mapsto \bar{f}$ is a bijection, natural along the coalgebra homomorphisms on the left and along the algebra homomorphisms on the right.

Claim (a) is proved as Lemma 4.4. The bijection part of claim (b) is proved as Lemma 4.3. The naturality of the correspondence is straightforward. $\square$

Lemma 4.3. *For an arbitrary adjunction $F = (F^* \dashv F_* : \mathbb{B} \rightarrow \mathbb{A})$, any algebra $F_*F^*x \xrightarrow{\alpha} x$, and any coalgebra $y \xrightarrow{\beta} F^*F_*y$ in $\mathbb{B}$, the mappings*

$$\begin{array}{ccc} & \xrightarrow{\quad \overline{(-)} \quad} & \\ \mathbb{A}(F_*y, x) & & \mathbb{B}(y, F^*x) \\ & \xleftarrow{\quad \overline{(-)} \quad} & \end{array}$$

defined by

$$\overline{f} = F^* f \circ \beta \qquad \underline{g} = \alpha \circ F_* g$$

induce a bijection between the subsets

$$\{f \in \mathbb{A}(F_*y, x) \mid f = \alpha \circ F_*F^*f \circ F_*\beta\} \cong \{g \in \mathbb{B}(y, F^*x) \mid g = F^*\alpha \circ F^*F_*g \circ \beta\}$$

illustrated in the following diagram.

$$\begin{array}{ccc}
 F_* F^* F_* y & \xrightarrow{F_* F^* f} & F_* F^* x \\
 \uparrow F_* \beta & \nearrow F_* g & \downarrow \alpha \\
 F_* y & \xrightarrow{f} & x
 \end{array}
 \quad \Leftrightarrow \quad
 \begin{array}{ccc}
 F^* F_* y & \xrightarrow{F^* F_* g} & F^* F_* F^* x \\
 \uparrow \beta & \searrow F^* f & \downarrow F_* \alpha \\
 y & \xrightarrow{g} & F^* x
 \end{array}$$

Proof. Following each of the mappings "there and back" gives

$$\begin{array}{ccccc} f & \longmapsto & \bar{f} = F^* f \circ \beta & \longmapsto & \underline{\bar{f}} = \alpha \circ F_* F^* f \circ F_* \beta = f \\ g & \longmapsto & g = \alpha \circ F_* g & \longmapsto & \bar{g} = F^* \alpha \circ F^* F_* g \circ \beta = g \end{array}$$

□

Lemma 4.4. For any adjunction $F = (F^* \dashv F_* : \mathbb{B} \rightarrow \mathbb{A})$, algebra $F_*F^*x \xrightarrow{\alpha} x$ in $\mathbb{A}$, coalgebra $y \xrightarrow{\beta} F^*F_*y$ in $\mathbb{B}$, arrow $f \in \mathbb{A}(F_*y, x)$ and $\bar{f} = F^*f \circ \beta \in \mathbb{B}(x, F^*y)$, if any of the squares (1-4) in Fig. 19 commutes, then they all commute. In particular, a square on one side of any of the equivalences (a–c) commutes if and only if

$$\begin{array}{ccc}
 \begin{array}{ccc}
 F_*F^*F_*y & \xrightarrow{F_*F^*f} & F_*F^*x \\
 \downarrow F_*\varepsilon & & \downarrow \alpha \\
 F_*y & \xrightarrow{f} & x
 \end{array} & (1) & \\
 \Downarrow (a) & & \\
 \begin{array}{ccc}
 F_*F^*F_*y & \xrightarrow{F_*F^*f} & F_*F^*x \\
 \uparrow F_*\beta & & \downarrow \alpha \\
 F_*y & \xrightarrow{f} & x
 \end{array} & (2) &
 \end{array}
 \quad (b) \Leftrightarrow \quad
 \begin{array}{ccc}
 \begin{array}{ccc}
 F^*F_*y & \xrightarrow{F^*F_*\bar{f}} & F^*F_*F^*x \\
 \uparrow \beta & & \uparrow F^*\eta \\
 y & \xrightarrow{\bar{f}} & F^*x
 \end{array} & (4) & \\
 \Downarrow (c) & & \\
 \begin{array}{ccc}
 F^*F_*y & \xrightarrow{F^*F_*\bar{f}} & F^*F_*F^*x \\
 \uparrow \beta & & \downarrow F_*\alpha \\
 y & \xrightarrow{\bar{f}} & F^*x
 \end{array} & (3) &
 \end{array}$$

Figure 19: Proof schema for (4.3)

the square on the other side of the equivalence commutes.

Proof. The claims are established as follows.

(1) $\xRightarrow{(a)}$ (2): Using the commutativity of (1) and (*) the counit equation $\varepsilon \circ \beta = \text{id}$ for the coalgebra β , we derive (2) as

$$\alpha \circ F_*F^*f \circ F_*\beta \stackrel{(1)}{=} f \circ F_*\varepsilon \circ F_*\beta \stackrel{(*)}{=} f$$

(2) $\stackrel{(a)}{\Rightarrow}$ (1). is proved by chasing the following diagram:

$$\begin{array}{ccccc}
F_*F^*F_*F^*F_*y & \xrightarrow{F_*F^*F_*F^*f} & F_*F^*F_*F^*x & & \\
\downarrow F_*\varepsilon & \swarrow F_*F^*F_*\beta & \searrow F_*F^*\alpha & & \downarrow F_*\varepsilon \\
& F_*F^*F_*y & \xrightarrow{F_*F^*f} & F_*F^*x & \\
& \downarrow F_*\varepsilon & (1) & \downarrow \alpha & \\
& F_*y & \xrightarrow{f} & x & \\
& \swarrow F_*\beta & (2) & \nwarrow \alpha & \\
F_*F^*F_*y & \xrightarrow{F_*F^*f} & F_*F^*x & &
\end{array}$$

The top and the bottom trapezoids commute by assumption (2), whereas the left-hand trapezoid (denoted $(\dagger)$) and the outer square (denoted $(\square)$) commute by the naturality of ε . The right-hand trapezoid (denoted $(\ddagger)$) commutes by the cochain condition for the algebra α . It follows that the inner square (denoted (1)) must also commute:

$$\begin{aligned}
f \circ F_*\varepsilon &\stackrel{(2)}{=} \alpha \circ F_*F^*f \circ F_*\beta \circ F_*\varepsilon \\
&\stackrel{(\dagger)}{=} \alpha \circ F_*F^*f \circ F_*\varepsilon \circ F_*F^*F_*\beta \\
&\stackrel{(\square)}{=} \alpha \circ F_*\varepsilon \circ F_*F^*F_*F^*f \circ F_*F^*F_*\beta \\
&\stackrel{(\ddagger)}{=} \alpha \circ F_*\alpha^* \circ F_*F^*F_*F^*f \circ F_*F^*F_*\beta \\
&\stackrel{(2)}{=} \alpha \circ F_*F^*f
\end{aligned}$$

(4) $\stackrel{(c)}{\Leftrightarrow}$ (3). is proven dually to (1) $\stackrel{(a)}{\Leftrightarrow}$ (2) above. The duality consists of reversing the arrows, switching F_* and F^* , and also α and β , and replacing ε with η .

(2) $\stackrel{(b)}{\Leftrightarrow}$ (3). follows from Lemma 4.3. □

5. THE NUCLEUS MONAD IS IDEMPOTENT

The claim is that the nucleus of a nucleus is equivalent to it. An easy way to prove this is to show that the construction of the nucleus $\overleftarrow{\mathfrak{N}}(F)$ can be decomposed into the final (Eilenberg-Moore) resolutions of the monad and the comonad induced by the adjunction $F = (F^* \dashv F_*)$. More precisely, first resolving the monad $\overleftarrow{F}$ induced by F into $\overleftarrow{\text{EM}}(F)$ and then resolving the comonad $\overrightarrow{F}$ induced by $\overleftarrow{\text{EM}}(F)$ to $\overleftarrow{\text{EC}}(\overleftarrow{\text{EM}}(F))$ is the

same as first resolving F to $\overleftarrow{\text{EC}}(F)$ and then to $\overleftarrow{\text{EM}}(\overleftarrow{\text{EC}}(F))$. The two pairs of resolutions are displayed in Fig. 20. Now we claim that the adjunctions $\overleftarrow{\text{EM}}(\overleftarrow{\text{EC}}(F))$ and $\overleftarrow{\text{EC}}(\overleftarrow{\text{EM}}(F))$ on the two ends are equivalent,




Figure 20: Alternating the monad and comonad localizations of an adjunction F

and that they are equivalent to $\overleftarrow{\mathfrak{N}}(F)$. Since $\overleftarrow{\text{EC}}$ and $\overleftarrow{\text{EM}}$ are idempotent monads, the claim that they commute implies that their composite is also an idempotent monad, which is also their intersection in the lattice of idempotent monads. This composite is the nucleus monad $\overleftarrow{\mathfrak{N}}$.

5.1. From weak to strong equivalences through absolute completions. The claim that the resolutions $\overleftarrow{\text{EC}} \circ \overleftarrow{\text{EM}}(F)$ and $\overleftarrow{\text{EM}} \circ \overleftarrow{\text{EC}}(F)$ are equivalent is true either

- a) for arbitrary categories and weak (Morita-)equivalences, or
- b) for absolutely (Cauchy-)complete categories and strong equivalences.

See [Bor94, Vol. 1, Sec. 7.9] for definitions of weak equivalences and absolute completions. In the setting from (a), nuclei could be studied entirely in terms of matrices, without getting into adjunctions, monads, or comonads⁴. But the technical overhead of working with weak equivalences is significant and many simple ideas get obscured by irrelevant implementation details. Approach (b) is easier, since the absolute completions of categories are simple, succinct, always available, and make weak equivalences strong.⁵ Appendix B provides an overview. To get a simple strong equivalence between a nucleus and its nucleus, we pursue approach (b) and construct them over absolutely complete categories. This, however, does not mean that the result needs to be restricted to absolutely complete categories. Since the absolute completion functor $\underline{(-)}: \text{Cat} \rightarrow \text{Cat}$ is an idempotent monad itself (as explained in Appendix B), which readily lifts to adjunctions (and also to monads and comonads), we can precompose all functors in sight with the absolute completion monad and work with absolutely complete categories with no loss of generality. In particular, we precompose the monads $\overleftarrow{\text{EC}}, \overleftarrow{\text{EM}}, \overleftarrow{\mathfrak{N}}: \text{Adj} \rightarrow \text{Adj}$ with the absolute completion monad $\underline{(-)}: \text{Adj} \rightarrow \text{Adj}$, lifted from categories to adjunctions, and define functors $\underline{\overleftarrow{\text{EC}}}, \underline{\overleftarrow{\text{EM}}}, \underline{\overleftarrow{\mathfrak{N}}}: \text{Adj} \rightarrow \text{Adj}$ by

$$\underline{\overleftarrow{\text{EC}}}(F) = \overleftarrow{\text{EC}}(\underline{F}) \quad \underline{\overleftarrow{\mathfrak{N}}}(F) = \overleftarrow{\mathfrak{N}}(\underline{F}) \quad \underline{\overleftarrow{\text{EM}}}(F) = \overleftarrow{\text{EM}}(\underline{F}) \quad (5.1)$$

⁴Grothendieck's generalization of Galois theory [Gro60, Gro71, Exposé VI] was originally stated in that way. The theory of monadicity was developed as a more salient version [BB69, Bec67, BR70]. Turning the tables, descent could be viewed as monadicity without monads [Pav91a].

⁵The tradeoffs like (a) vs (b) often arise in mathematics. Scientists could work with the sequences of rational numbers that arise from their measurements and use weak equality to compare them. Or they can construct the Cauchy-completion of the rationals, the real numbers, and work with the strong equality of those. The latter approach is usually preferred.

where $\underline{\mathbb{C}}$ denotes the absolute completion of a category $\mathbb{C}$ and $\underline{F}: \underline{\mathbb{C}} \rightarrow \underline{\mathbb{D}}$ is the lifting of the functor $F: \mathbb{C} \rightarrow \mathbb{D}$. Checking that $\underline{\underline{\mathbb{E}\mathbb{M}}}$ and $\underline{\underline{\mathbb{E}\mathbb{C}}}$ are idempotent monads is straightforward, and checking that $\underline{\underline{\mathfrak{N}}}$ is strongly equivalent to both of their composites and thus the idempotent monad at their intersection, will be the task of the theorem below. Since the liftings are unique, the underlining is omitted whenever the confusion is unlikely. We reiterate that this is inessential for the main constructions and could be avoided at the cost of more involved arguments.

5.2. The Nucleus Theorem.

Theorem 5.1. *The idempotent monads*

$$\underline{\underline{\mathbb{E}\mathbb{C}}} = \left(\text{Adj} \xrightarrow{\text{AC}} \text{Cmn} \xrightarrow{\text{EC}} \text{Adj} \right) \quad \text{and} \quad \underline{\underline{\mathbb{E}\mathbb{M}}} = \left(\text{Adj} \xrightarrow{\text{AM}} \text{Mnd} \xrightarrow{\text{EM}} \text{Adj} \right) \quad (5.2)$$

commute and their composites are isomorphic to the nucleus

$$\underline{\underline{\mathbb{E}\mathbb{C}}} \circ \underline{\underline{\mathbb{E}\mathbb{M}}}(F) \cong \underline{\underline{\mathfrak{N}}}(F) \cong \underline{\underline{\mathbb{E}\mathbb{M}}} \circ \underline{\underline{\mathbb{E}\mathbb{C}}}(F) \quad (5.3)$$

naturally in F . With the leftmost and the rightmost adjunctions in Fig. 20 isomorphic, the monad units η displayed there can be folded into the commutative square in Adj , displayed in Fig. 21, presenting the unit of the monad $\underline{\underline{\mathfrak{N}}}$.




Figure 21: $\underline{\underline{\mathfrak{N}}}(F)$ factorized into $\underline{\underline{\mathbb{E}\mathbb{M}}} \circ \underline{\underline{\mathbb{E}\mathbb{C}}}(F)$ and $\underline{\underline{\mathbb{E}\mathbb{C}}} \circ \underline{\underline{\mathbb{E}\mathbb{M}}}(F)$

Proof. The claim is that the leftmost resolution $\underline{\underline{\mathbb{E}\mathbb{M}}} \circ \underline{\underline{\mathbb{E}\mathbb{C}}}(F)$ and the rightmost resolution $\underline{\underline{\mathbb{E}\mathbb{C}}} \circ \underline{\underline{\mathbb{E}\mathbb{M}}}(F)$ in Fig. 20 are isomorphic, and that they are both isomorphic to $\underline{\underline{\mathfrak{N}}}(F)$, as displayed in Fig. 22. Recall from Lemma 4.2 that $\underline{\underline{\mathfrak{N}}}(F) = (F^\# \dashv F_\# : \underline{\underline{A}}^{\underline{\underline{F}}} \rightarrow \underline{\underline{B}}^{\underline{\underline{F}}})$ as a resolution of the descent comonad $\overrightarrow{F}$ on $\underline{\underline{A}}^{\underline{\underline{F}}}$. We show

Figure 22: The equivalences claimed in Thm. 5.1

that for the induced comparison functor $\Xi^0: \underline{\mathbb{B}}^{\vec{F}} \rightarrow (\underline{\mathbb{A}}^{\overleftarrow{F}})^{\overrightarrow{F}}$ is an equivalence. Ξ^0 is defined in Fig. 23 by instantiating the standard comparison functor construction from (3.12) to $\overrightarrow{F}$. The right adjoint Ξ_0 is defined

Figure 23: The comparison functor from $\underline{\mathfrak{M}}(F)$ to $\underline{\mathbb{E}\mathbb{C}} \circ \underline{\mathbb{E}\mathbb{M}}(F)$

in Fig. 24, where d is the structure map of the coalgebra $\alpha \xrightarrow{d} F^\#F_\#\alpha$ in $\underline{\mathbb{A}}^{\overleftarrow{F}}$ and y is defined by splitting the idempotent $\varepsilon \circ F^*d$. **This is where the absolute completeness of $\mathbb{B}$ is used.** To show that Ξ^0 and Ξ_0 make $(\underline{\mathbb{A}}^{\overleftarrow{F}})^{\overrightarrow{F}}$ and $\underline{\mathbb{B}}^{\vec{F}}$ equivalent, we construct natural isomorphisms $\Xi_0\Xi^0 \cong \text{Id}$ and $\Xi^0\Xi_0 \cong \text{Id}$. Towards $\Xi_0\Xi^0 \cong \text{Id}$, note that instantiating $\Xi^0\beta: F^\#\beta \rightarrow F^\#F_\#F^\#\beta$ (the right-hand square in Fig. 23) as $\delta: \alpha \rightarrow F^\#F_\#\alpha$ (the left-hand square in Fig. 24) reduces the right-hand equalizer in Fig. 24 to Fig. 25. The commutativity of Fig. 25 is equivalent to the fact that β is a $\overrightarrow{F}$ -coalgebra [BW85, Sec. 3.6]. To construct the isomorphism $\Xi^0\Xi_0 \cong \text{Id}$, take an arbitrary coalgebra $\alpha \xrightarrow{\delta} F^\#F_\#\alpha$ from $(\underline{\mathbb{A}}^{\overleftarrow{F}})^{\overrightarrow{F}}$ and consider the functor Ξ^0 in Fig. 23 instantiated to $\beta = \Xi_0\delta$. By extending the right-hand side of this instance of Fig. 23 by the F_* -image of the right-hand side of Fig. 24, we get Fig. 26. The claim is now that $x \xrightarrow{d} F_*F^*x$ equalizes the parallel

$$\begin{array}{ccc}
\left(\mathbb{A}^{\overleftarrow{F}}\right)^{\overrightarrow{F}} & \xrightarrow{\Xi_0} & \mathbb{B}^{\overrightarrow{F}} \\
\\
\begin{array}{ccc}
F_*F^*x & \xrightarrow{\alpha} & x \\
\downarrow F_*F^*d & \downarrow \delta & \downarrow d \\
F_*F^*F_*F^*x & \xrightarrow[=F_*\varepsilon_{F^*x}]{F^\sharp F_\sharp \alpha} & F_*F^*x
\end{array} & \mapsto & \begin{array}{ccccc}
& & \varepsilon & & \\
& & \swarrow & & \searrow \\
y & \xrightarrow{e} & F^*x & \xrightarrow[F^*\eta]{F^*d} & F^*F_*F^*x \\
& \swarrow r & \downarrow F_\sharp \alpha = F^*\eta & & \downarrow F^*\eta \\
& & F^*F_*y & \xrightarrow[F^*F_*\eta]{F^*F_*e} & F^*F_*F^*x & \xrightarrow[F^*F_*\eta]{F^*F_*F^*d} & F^*F_*F_*F^*x \\
& & \downarrow \Xi_0\delta & & \downarrow F^*\eta & & \downarrow F^*\eta \\
& & F^*F_*y & \xrightarrow[F^*F_*\eta]{F^*F_*e} & F^*F_*F^*x & \xrightarrow[F^*F_*\eta]{F^*F_*F^*d} & F^*F_*F_*F^*x \\
& & \swarrow F^*F_*r & & \swarrow F^*F_*\varepsilon & & \swarrow F^*F_*\varepsilon
\end{array}
\end{array}$$

Figure 24: The right adjoint of the comparison from $\overleftarrow{\mathfrak{H}}(F)$ to $\overleftarrow{\mathbf{EC}} \circ \overleftarrow{\mathbf{EM}}(F)$

$$\begin{array}{ccccc}
& & \varepsilon & & \\
& & \swarrow & & \searrow \\
y & \xrightarrow{\beta} & F^*F_*y & \xrightarrow[F^*\eta]{F^*F_*\beta} & F^*F_*F^*F_*y \\
& \swarrow \varepsilon & \downarrow F^*\eta & & \downarrow F^*\eta \\
& & F^*F_*y & \xrightarrow[F^*F_*\eta]{F^*F_*\beta} & F^*F_*F^*F_*y \\
& & \downarrow \Xi_0\Xi^0\beta = \beta & & \downarrow F^*\eta \\
& & F^*F_*y & \xrightarrow[F^*F_*\eta]{F^*F_*\beta} & F^*F_*F^*F_*y \\
& & \swarrow F^*F_*\varepsilon & & \swarrow F^*F_*\varepsilon
\end{array}$$

Figure 25: $\Xi_0\Xi^0 \cong \text{id}$

pair $\langle F_*F^*\eta, F_*F^*d \rangle$ in the first row. Since $y \xrightarrow{e} F^*x$ was defined in Fig. 24 as a split equalizer of the pair $\langle F^*\eta, F^*d \rangle$, and all functors preserve split equalizers, it follows that $F_*y \xrightarrow{F_*e} F_*F^*x$ is also an equalizer of the same pair $\langle F_*F^*\eta, F_*F^*d \rangle$. Hence the isomorphism $x \cong F_*y$, which gives $\Xi_0\Xi^0\delta \cong \delta$. To prove the claim that $x \xrightarrow{d} F_*F^*x$ equalizes the first row, note that, just like the coalgebra $y \xrightarrow{\beta} F^*F_*y$ in $\mathbb{B}^{\overrightarrow{F}}$ was determined up to isomorphism by the split equalizer in $\mathbb{B}$, shown in Fig. 25, the coalgebra $\alpha \xrightarrow{\delta} F^\sharp F_\sharp \alpha$ in $\left(\mathbb{A}^{\overleftarrow{F}}\right)^{\overrightarrow{F}}$ is determined




Figure 26: Construction of $\Xi^0 \Xi_0 \delta$

up to isomorphism by the following split equalizer in $\mathbb{A}^{\overleftarrow{F}}$

$$\begin{array}{ccccc}
 & & \xleftarrow{\epsilon} & & \\
 & & F^\# F_\# \delta & & \\
 \alpha & \xrightarrow{\delta} & F^\# F_\# \alpha & \xrightarrow{F^\# \eta} & F^\# F_\# F^\# F_\# \alpha \\
 & \xleftarrow{\epsilon} & & &
 \end{array} \tag{5.4}$$

In $\mathbb{A}$, (5.4) unfolds to the lower squares of the Fig. 27. Since the upper right-hand squares also commute




Figure 27: The unfolding of (5.4) and its splitting in $\mathbb{A}$

(by the naturality of η), they also induce the factoring of the split equalizers in the upper left-hand square.

both $F_*y \xrightarrow{F_*e} F_*F^*x$ and $x \xrightarrow{d_*} F_*F^*x$ are split equalizers of the same pair yields the isomorphism $F_*y \xrightarrow{\iota} x$ in $\mathbb{A}$, which turns out to be a coalgebra isomorphism $\Xi^0\Xi_0\delta \xrightarrow[\sim]{\iota} \delta$ in $\left(\mathbb{A}^{\overleftarrow{F}}\right)^{\overrightarrow{F}}$, as shown in Fig. 28. Here the

$$\begin{array}{ccccc}
F_*F^*x & \xrightarrow{\alpha} & & & x \\
\downarrow F_*F^*d & \swarrow F_*F^*\iota & F_*F^*F_*y & \xrightarrow{F_*\varepsilon} & F_*y & \searrow \iota & \downarrow d \\
& & \downarrow F_*F^*F_*\Xi_0\delta & & \downarrow F_*\Xi_0\delta & & \\
& & F_*F^*F_*F^*F_*y & \xrightarrow{F_*\varepsilon} & F_*F^*F_*y & \searrow F_*F^*\iota & \\
& \swarrow F_*F^*F_*F^*\iota & & & & & \\
F_*F^*F_*F^*x & \xrightarrow{F_*\varepsilon} & & & F_*F^*x
\end{array}$$

Figure 28: $\Xi^0 \Xi_0 \cong \text{Id}$

outer square is δ , as in Fig. 24 on the left, whereas the inner square is $\Xi^0 \Xi_0 \delta$, as in Fig. 26 on the left. The right-hand trapezoid commutes because the middle square in Fig. 26 commutes, and can be chased down to Fig. 29 using the fact that ι is defined by $F_* e = d \circ \iota$. The commutativity of the left-hand trapezoid in

Figure 29: Commutativity of the right-hand trapezoid in Fig. 28

Fig. 28 follows, because it is an F_*F^* -image of the right-hand trapezoid. The bottom trapezoid commutes by the naturality of ε . The top trapezoid commutes because everything else commutes, and d is a monic. The commutative diagram in Fig. 28 thus displays the claimed isomorphism $\Xi^0 \Xi_0 \delta \xrightarrow{\iota} \delta$. This completes the proof that $\Xi^0 \Xi_0 \cong \text{Id}$. Together with the proof that $\Xi_0 \Xi^0 \cong \text{Id}$ in Fig. 25, it completes the proof that $\Xi = (\Xi^0 \dashv \Xi_0)$ provides the first of the equivalences

$$\mathbb{B}^{\overrightarrow{F}} \simeq \left(\mathbb{A}^{\overleftarrow{F}} \right)^{\overrightarrow{F}} \quad \text{and} \quad \left(\mathbb{B}^{\overrightarrow{F}} \right)^{\overleftarrow{F}} \simeq \mathbb{A}^{\overleftarrow{F}}$$

claimed in Fig. 22. The second one is dual. $\square$

The adjunction $\overleftarrow{\mathfrak{N}}(F) = (F^\# \dashv F_\# : \mathbb{B}^{\overrightarrow{F}} \rightarrow \mathbb{A}^{\overleftarrow{F}})$ is monadic and comonadic. By definition, an adjunction $G^* \dashv G_* : \mathbb{D} \rightarrow \mathbb{C}$ is monadic if the comparison functor $H^1 : \mathbb{D} \rightarrow \mathbb{C}^{\overleftarrow{G}}$ in (3.12) on the right is an equivalence. It is comonadic if the comparison functor $H_0 : \mathbb{C} \rightarrow \mathbb{D}^{\overrightarrow{G}}$ in (3.12) on the right is an equivalence. Thm. 5.1 thus implies that the nucleus $\overleftarrow{\mathfrak{N}}(F)$ of any adjunction F is monadic and comonadic, as anticipated in Fig. 1.

5.3. Consequences of the Nucleus Theorem. The equivalence $\Xi = (\Xi^* \dashv \Xi_*)$ proves the claim in the title of this section.

Corollary 5.1. $\overleftarrow{\mathfrak{N}} : \text{Adj} \rightarrow \text{Adj}$ is an idempotent monad, with the natural isomorphism $\overleftarrow{\mathfrak{N}}(F) \cong \overleftarrow{\mathfrak{N}}\overleftarrow{\mathfrak{N}}(F)$ realized by the unit in Fig. 30.

$$\begin{array}{ccc}
 \begin{array}{c} \overleftarrow{F} \\ \curvearrowright \\ \mathbb{B}^{\overrightarrow{F}} \end{array} & \xrightarrow{\simeq} \xrightarrow{\Xi^0} & \begin{array}{c} \overleftarrow{F} \\ \left(\mathbb{A}^{\overleftarrow{F}} \right)^{\overrightarrow{F}} \end{array} \\
 \uparrow & \eta & \uparrow \\
 \overleftarrow{\mathfrak{N}}(F) & \xrightarrow{\eta} & \overleftarrow{\mathfrak{N}}\overleftarrow{\mathfrak{N}}(F) \\
 \downarrow & & \downarrow \\
 \mathbb{A}^{\overleftarrow{F}} & \xrightarrow{\simeq} \xrightarrow{\Xi_1} & \begin{array}{c} \overleftarrow{F} \\ \left(\mathbb{B}^{\overrightarrow{F}} \right)^{\overleftarrow{F}} \end{array} \\
 \begin{array}{c} \overrightarrow{F} \\ \curvearrowright \end{array} & &
 \end{array}$$

Figure 30: The unit $\eta : \overleftarrow{\mathfrak{N}}(F) \rightarrow \overleftarrow{\mathfrak{N}}\overleftarrow{\mathfrak{N}}(F)$

Corollary 5.2. If $F = (F^* \dashv F_* : \mathbb{B} \rightarrow \mathbb{A})$ and $G = (G^* \dashv G_* : \mathbb{D} \rightarrow \mathbb{A})$ induce the same monad, then they have the same nucleus:

$$F_*F^* = \overleftarrow{T} = G_*G^* \implies \overleftarrow{\mathfrak{N}}(F) \cong \overleftarrow{\mathfrak{N}}(G) \quad (5.5)$$

In particular, $F_*F^* = G_*G^*$ on $\mathbb{A}$ implies $\mathbb{B}^{\overrightarrow{F}} \simeq \mathbb{D}^{\overrightarrow{G}}$.

Proof. The assumption $F_*F^* = \overleftarrow{T} = G_*G^*$ means that F and G have the same final resolution $\overleftarrow{\text{EM}}(F) = (U_{\overleftarrow{T}} \dashv U^{\overleftarrow{T}} : \mathbb{A}^{\overleftarrow{T}} \rightarrow \mathbb{A}) = \overleftarrow{\text{EM}}(G)$. It follows that they also induce the same descent comonad $\overrightarrow{T} = U^{\overleftarrow{T}}U_{\overleftarrow{T}}$ on $\mathbb{A}^{\overleftarrow{T}}$. Lemma 4.2 now gives

$$F^\#F_\# = \overrightarrow{T} = G^\#G_\#$$

whereas Thm. 5.1 proves

$$\mathbb{B}^{\overrightarrow{F}} \simeq \left(\mathbb{A}^{\overleftarrow{T}} \right)^{\overrightarrow{T}} \simeq \mathbb{D}^{\overrightarrow{G}}$$

Since the comparison functors commute with the nucleus functors, we have proved $\overleftarrow{\mathfrak{N}}(F) \cong \overleftarrow{\mathfrak{N}}(G)$. $\square$

Remark. Notice that the proof and its dual in essence prove

$$\overleftarrow{\mathbf{EM}}(F) \cong \overleftarrow{\mathbf{EM}}(G) \vee \overleftarrow{\mathbf{EC}}(F) \cong \overleftarrow{\mathbf{EC}}(G) \implies \overleftarrow{\mathbf{N}}(F) \cong \overleftarrow{\mathbf{N}}(G) \quad (5.6)$$

Stated in this form, the claim is an obvious consequence of Thm. 5.1. However, unpacking the implications from the two disjuncts provides useful statements. The first disjunct says that any equivalences $\underline{\mathbb{A}} \simeq \underline{\mathbb{C}}$ and $\underline{\mathbb{A}}^{\overleftarrow{F}} \simeq \underline{\mathbb{C}}^{\overleftarrow{G}}$, coherent with the forgetful and the free functors, imply $\underline{\mathbb{B}}^{\overrightarrow{F}} \simeq \underline{\mathbb{D}}^{\overrightarrow{G}}$. Dually, the second disjunct says that coherent equivalences $\underline{\mathbb{B}} \simeq \underline{\mathbb{D}}$ and $\underline{\mathbb{B}}^{\overrightarrow{F}} \simeq \underline{\mathbb{D}}^{\overrightarrow{G}}$ imply $\underline{\mathbb{A}}^{\overleftarrow{F}} \simeq \underline{\mathbb{C}}^{\overleftarrow{G}}$.

6. SIMPLE NUCLEUS

The equivalences $\underline{\mathbb{A}}^{\overleftarrow{F}} \simeq (\underline{\mathbb{B}}^{\overrightarrow{F}})^{\overleftarrow{F}}$ and $\underline{\mathbb{B}}^{\overrightarrow{F}} \simeq (\underline{\mathbb{A}}^{\overleftarrow{F}})^{\overrightarrow{F}}$ from Thm. 5.1 present $\overleftarrow{F}$ -algebras as $\overleftarrow{\overleftarrow{F}}$ -algebras over $\overrightarrow{F}$ -coalgebras, and $\overrightarrow{F}$ -coalgebras as $\overrightarrow{\overrightarrow{F}}$ -coalgebras over $\overleftarrow{F}$ -algebras. But these presentations are redundant, and discharging the redundancies yields interesting and useful alternatives to the familiar Eilenberg-Moore views of algebras and coalgebras — in terms of idempotents. In a subtly different form, they were used in [PS17] for a particular security application. Here we prepare them for mathematical applications.

Proposition 6.1. *Given an adjunction $F = (F^* \dashv F_* : \underline{\mathbb{B}} \rightarrow \underline{\mathbb{A}})$, consider the categories*

$$|\underline{\mathbb{A}}^{\overrightarrow{\overleftarrow{F}}}| = \coprod_{x \in |\underline{\mathbb{A}}|} \left\{ \alpha_x \in \underline{\mathbb{B}}(F^*x, F^*x) \mid \begin{array}{ccccc} F^*x & & F_*F^*x & \xrightarrow{\quad} & x \\ \downarrow \alpha_x & \searrow & \downarrow & \searrow & \downarrow \tilde{\alpha}_x \\ F^*x & \xrightarrow{\alpha_x} & F^*x & \xrightarrow{\quad} & F_*F^*x \end{array} \right\} \quad (6.1)$$

$$\underline{\mathbb{A}}^{\overrightarrow{\overleftarrow{F}}}(\alpha_x, \gamma_z) = \left\{ f \in \underline{\mathbb{A}}(x, z) \mid \begin{array}{ccc} F^*x & \xrightarrow{F^*f} & F^*z \\ \downarrow \alpha_x & & \downarrow \gamma_z \\ F^*x & \xrightarrow{F^*f} & F^*z \end{array} \right\}$$

$$|\underline{\mathbb{B}}^{\overleftarrow{\overrightarrow{F}}}| = \coprod_{u \in |\underline{\mathbb{B}}|} \left\{ \beta^u \in \underline{\mathbb{A}}(F_*u, F_*u) \mid \begin{array}{ccccc} F_*u & & F^*F_*u & \xrightarrow{\tilde{\beta}^u} & u \\ \downarrow \beta^u & \searrow & \downarrow & \searrow & \downarrow \\ F_*u & \xrightarrow{\beta^u} & F_*u & \xrightarrow{\quad} & F^*F_*u \end{array} \right\} \quad (6.2)$$

$$\underline{\mathbb{B}}^{\overleftarrow{\overrightarrow{F}}}(\beta^u, \delta^w) = \left\{ g \in \underline{\mathbb{B}}(u, w) \mid \begin{array}{ccc} F_*u & \xrightarrow{F_*g} & F_*w \\ \downarrow \beta^u & & \downarrow \delta^w \\ F_*u & \xrightarrow{F_*g} & F_*w \end{array} \right\}$$

where $x \xrightarrow{\tilde{\alpha}_x} F_*F^*x$ is the transpose of $F^*x \xrightarrow{\alpha_x} F^*x$, and $F^*F_*u \xrightarrow{\tilde{\beta}^u} u$ is the transpose of $F_*u \xrightarrow{\beta} F_*u$. The adjunction $F^\natural \dashv F_\natural : \underline{\mathbb{B}}^{\overleftarrow{\overrightarrow{F}}} \rightarrow \underline{\mathbb{A}}^{\overrightarrow{\overleftarrow{F}}}$ defined in Fig. 31 with the comparison functors




Figure 31: The simple nucleus $F^\sharp \dashv F_\sharp$ of $F^* \dashv F_*$

$$\begin{array}{ccc}
 K^0 : \mathbb{A} \longrightarrow \underline{\mathbb{A}}^{\overline{F}} & & K_1 : \mathbb{B} \longrightarrow \underline{\mathbb{B}}^{\overleftarrow{F}} \\
 x \mapsto \left\langle F_* F^* x, \begin{array}{c} F^* F_* F^* x \\ \varepsilon F^* \downarrow \\ F^* x \\ F^* \eta \downarrow \\ F^* F_* F^* x \end{array} \right\rangle & & u \mapsto \left\langle F^* F_* u, \begin{array}{c} F_* F^* F_* u \\ F_* \varepsilon \downarrow \\ F_* u \\ \eta F_* \downarrow \\ F_* F^* F_* u \end{array} \right\rangle
 \end{array}$$

is equivalent to the nucleus, i.e.

$$\overleftarrow{\mathfrak{N}}(F^* \dashv F_*) \simeq (F^\sharp \dashv F_\sharp)$$

Lemma 6.2. If $(F_* F^* x \xrightarrow{F_* \alpha_x} F_* F^* x) = (F_* F^* x \xrightarrow{\alpha^x} x \xrightarrow{\tilde{\alpha}_x} F_* F^* x)$, where $\tilde{\alpha}_x$ is the adjoint transpose of α_x , then $\alpha^x \circ \tilde{\alpha}_x = \text{id}$ implies $\alpha^x \circ \eta_x = \text{id}$.

Proof. The assumption that $\tilde{\alpha}_x$ is the adjoint transpose implies $\tilde{\alpha}_x \circ \alpha^x = F_* \alpha_x \circ \eta_x \circ \alpha^x = \tilde{\alpha}_x \circ \alpha^x \circ \eta_x \circ \alpha^x$. Pre- and postcomposing with $\alpha^x \circ \tilde{\alpha}_x = \text{id}$ gives $\alpha^x \circ \eta_x = \alpha^x \circ (\tilde{\alpha}_x \circ \alpha^x \circ \eta_x \circ \alpha^x) \circ \tilde{\alpha}_x = \alpha^x \circ \tilde{\alpha}_x \circ \alpha^x \circ \tilde{\alpha}_x = \text{id}$. $\square$

Proof of Prop. 6.1. Splitting the idempotent $F_* F^* x \xrightarrow{F_* \alpha_x} F_* F^* x$ into $F_* F^* x \xrightarrow{\alpha^x} x \xrightarrow{\tilde{\alpha}_x} F_* F^* x$ as above gives

$$\alpha^x \circ \tilde{\alpha}_x = \text{id}_x \quad \text{and} \quad \alpha^x \circ \eta_x = \text{id}_x \quad (6.3)$$

Analogous reasoning proves

$$\alpha^x \circ F_* F^* \alpha^x = \alpha^x \circ F_* \varepsilon_{F_* x} \quad \text{and} \quad \tilde{\alpha}_x \circ \alpha^x = F_* \varepsilon_{F_* x} \circ F_* F^* \tilde{\alpha}_x \quad (6.4)$$

Together, (6.3–6.4) say that $F_* F^* x \xrightarrow{\alpha^x} x$ is an algebra in $\underline{\mathbb{A}}^{\overline{F}}$ and that $\tilde{\alpha}_x \in \underline{\mathbb{A}}^{\overline{F}}(\alpha^x, \mu_x)$ is a coalgebra over α^x in $(\underline{\mathbb{A}}^{\overline{F}})^{\overline{F}}$. Hence a functor $\underline{\mathbb{A}}^{\overline{F}} \rightarrow (\underline{\mathbb{A}}^{\overline{F}})^{\overline{F}}$. Similar construction yields a similar functor $\underline{\mathbb{B}}^{\overleftarrow{F}} \rightarrow (\underline{\mathbb{B}}^{\overleftarrow{F}})^{\overleftarrow{F}}$. Hence the equivalences

$$\underline{\mathbb{A}}^{\overline{F}} \simeq (\underline{\mathbb{A}}^{\overline{F}})^{\overline{F}} \quad \underline{\mathbb{B}}^{\overleftarrow{F}} \simeq (\underline{\mathbb{B}}^{\overleftarrow{F}})^{\overleftarrow{F}} \quad (6.5)$$

The equivalences

$$\underline{\mathbb{A}}^{\overline{F}} \simeq \underline{\mathbb{B}}^{\overrightarrow{F}} \quad \underline{\mathbb{B}}^{\overleftarrow{F}} \simeq \underline{\mathbb{A}}^{\overleftarrow{F}} \quad (6.6)$$

follow from Thm 5.1 and have also been verified directly in [PS17]. Every object $\langle x, F^*x \xrightarrow{\alpha_x} F^*x \rangle$ of $\underline{\mathbb{A}}^{\overline{\overline{F}}}$ is thus isomorphic to one in the form $\langle F_*y, F^*F_* \xrightarrow{\varepsilon} y \xrightarrow{\beta} F^*F_*y \rangle$, where β is a coalgebra in $\underline{\mathbb{B}}^{\overline{F}}$. It follows that the inner and the outer squares in the following diagram commute

$$\begin{array}{ccccc}
 F^*x & \xrightarrow{F^*\eta} & F^*F_*F^*x & & \\
 \downarrow \alpha_x & \swarrow F^*\iota & \downarrow F^*F_*F^*\iota & & \downarrow \alpha_x \\
 & F^*F_*y & \xrightarrow{F^*\eta F_*} & F^*F_*F^*F_*y & \\
 & \downarrow \varepsilon & & \downarrow F^*F_*\varepsilon & \\
 & y & \xrightarrow{\beta} & F^*F_*y & \\
 & \downarrow \beta & & \downarrow F^*F_*\beta & \\
 & F^*F_*y & \xrightarrow{F^*\eta F_*} & F^*F_*F^*F_*y & \\
 \downarrow \alpha_x & \swarrow F^*\iota & \downarrow F^*F_*F^*\iota & & \downarrow \alpha_x \\
 F^*x & \xrightarrow{F^*\eta} & F^*F_*F^*x & &
 \end{array} \tag{6.7}$$

and that they are linked by an isomorphism $x \leftarrow \iota \rightarrow F_*y$ in $\underline{\mathbb{A}}$. Transferring the nuclear adjunction $F^\sharp \dashv F_\sharp : \underline{\mathbb{A}}^{\overline{\overline{F}}} \rightarrow \underline{\mathbb{B}}^{\overline{F}}$ along the equivalences yields the nuclear adjunction $F^\natural \dashv F_\natural : \underline{\mathbb{B}}^{\overline{F}} \rightarrow \underline{\mathbb{A}}^{\overline{\overline{F}}}$, with the natural correspondence

$$\begin{aligned}
 \underline{\mathbb{B}}^{\overline{F}}(F^\natural \alpha_x, \beta^u) &\cong \underline{\mathbb{A}}^{\overline{\overline{F}}}(\alpha_x, F_\natural \beta^u) \\
 (F^*x \xrightarrow{f} u) &\mapsto \tilde{f} = (x \xrightarrow{\eta} F_*F^*x \xrightarrow{F_*f} F_*u)
 \end{aligned}$$

The transpositions along $F^* \dashv F_* : \underline{\mathbb{B}} \rightarrow \underline{\mathbb{A}}$ lift to $F^\natural \dashv F_\natural : \underline{\mathbb{B}}^{\overline{F}} \rightarrow \underline{\mathbb{A}}^{\overline{\overline{F}}}$ because each of the following squares commutes if and only if the other one does:

$$\begin{array}{ccc}
 F_*F^*x & \xrightarrow{F_*f} & F_*u \\
 \downarrow F_*\alpha_x & & \downarrow \beta^u \\
 F_*F^*x & \xrightarrow{F_*f} & F_*u
 \end{array}
 \iff
 \begin{array}{ccc}
 F^*x & \xrightarrow{F^*(\tilde{f})} & F^*F_*u \\
 \downarrow \alpha_x & & \downarrow F^*\beta^u \\
 F^*x & \xrightarrow{F^*(\tilde{f})} & F^*F_*u
 \end{array} \tag{6.8}$$

To see this, suppose that the left-hand side square commutes. Take the F^* -image of the right-hand side square and precompose it with the outer square from (6.7), as in the following diagram.

$$\begin{array}{ccccc}
 & & F^*(\tilde{f}) & & \\
 & \nearrow & & \searrow & \\
 F^*x & \xrightarrow{F^*\eta} & F^*F_*F^*x & \xrightarrow{F^*F_*f} & F^*F_*u \\
 \downarrow \alpha_x & & \downarrow F^*F_*\alpha_x & & \downarrow F^*\beta^u \\
 F^*x & \xrightarrow{F^*\eta} & F^*F_*F^*x & \xrightarrow{F^*F_*f} & F^*F_*u \\
 & \nwarrow & & \swarrow & \\
 & & F^*(\tilde{f}) & &
 \end{array} \tag{6.9}$$

The two outer paths around this diagram are the paths around the right-hand square in (6.8). The right-to-left implication is analogous. $\square$

7. NEW FRAMEWORKS FOR OLD BIG PICTURES

7.1. Descent to nucleus. In an (over)simplified form, the idea of descent is to descend to a complex object by covering it with simpler objects. Sheaves can be viewed as a special case. A sheaf is a set ς , continuously varying over a topological space X . The continuous variation means that it is indexed over the open neighborhoods of X and the elements of ςB , for an open neighborhood $B \subseteq X$, can be described by descending from the simpler elements of ςA_i down a covering $B = \bigcup_{i \in I} A_i$. The sets varying over X are formalized as functors $OX^o \rightarrow \mathbf{Set}$, the presheaves, where OX is the lattice of open neighborhoods of X . A presheaf $\varsigma: OX^o \rightarrow \mathbf{Set}$ is a sheaf if every covering $B = \bigcup_{i \in I} A_i$ is mapped to a decomposition

$$\varsigma B \cong \left\{ \langle si \rangle_{i \in I} \in \prod_{i \in I} \varsigma A_i \mid \forall i, j \in I. si \upharpoonright_{A_i \cap A_j} = sj \upharpoonright_{A_i \cap A_j} \right\} \tag{7.1}$$

where $sU \upharpoonright_V$ is the restriction of $sU \in \varsigma U$, mapped by ς -image $\varsigma U \xrightarrow{\upharpoonright} \varsigma V$ of the morphism $U \supseteq V$ in OX^o . This *sheaf condition* formalizes the intuition that the set ςB descends from the family of sets $\langle \varsigma A_i \rangle_{i \in I}$ just like B descends from its cover $\{A_i\}_{i \in I}$: that is, just like B is obtained by gluing A_i s and A_j s along the intersections $A_i \cap A_j$, the set ςB is obtained by gluing ςA_i s and ςA_j s along the intersections $\varsigma(A_i \cap A_j)$.

The (over)simplified idea of Grothendieck descent [Gro60, Gro71] is to lift the concept of gluing from sheaves as continuously variable sets $OX^o \rightarrow \mathbf{Set}$ to continuously variable categories $\mathbb{S}^o \rightarrow \mathbf{Cat}$, where the continuity with respect to $\mathbb{S}$ is expressed in terms of coverings of objects by *sieves* in the slices [Gir64]. An indexed category $\mathbb{S}^o \rightarrow \mathbf{Cat}$ satisfying the suitable gluing condition, usually comprehended as a fibration, is called a *stack*. The concept originated in algebraic geometry [Gir71, Vis05], expanded into homotopy theory [Gro18, Jar19] and many areas of mathematics in-between and beyond [JT04]. The breadth of applications makes the presentations wildly divergent and implementation-dependent, and it is not easy to see the forest for the trees. The first author proposed a logical interpretation of descent as an interpolation condition [Pav91a]. The nucleus seems to suggest a further simplification.

For intuition, consider an indexed category $\mathcal{P}: \mathbb{S}^o \rightarrow \mathbf{Cat}$ mapping types (or sets) from a universe $\mathbb{S}$ to categories $\mathcal{P}X$ whose objects, written $\alpha(x), \gamma(u), \dots$, can be thought of as observable properties or

predicates over X .⁶ The morphisms $p \in \mathcal{P}X(\alpha, \gamma)$ are then thought of as proofs of $\alpha(x) \Rightarrow \gamma(x)$, along the lines of the Brouwer-Heyting-Kolmogorov interpretation of constructivist logic, echoed in the propositions-as-types interpretation of type theory. See [CH88, Jac01, ML84, Pav90] for formal developments of such interpretations. The starting point of the corresponding categorical developments was Lawvere's definition of *hyperdoctrines* [Law69, Law70]. We call $\mathcal{P}$ a hyperdoctrine even if the closed structure, central in the original treatments, presently plays no role. A hyperdoctrine $\mathcal{P}$ is thus a functor assigning to every function f in the universe $\mathbb{S}$ an adjoint pair of functors $f_! \dashv f^*$, where f^* is interpreted as the substitution operation and $f_!$ provides the corresponding existential quantifiers. The notation in Fig. 32 was formalized and used in [Pav90, Pav91b, Pav96]. In a high-level logical view, the idea of descent is that the predicates over B

$$\frac{f: A \rightarrow B}{f_! \dashv f^*: \mathcal{P}B \rightarrow \mathcal{P}A} \quad f^*(\beta(y)) = \beta(f(x)) \quad f_!(\alpha(x)) = \exists x: A. f(x) \stackrel{B}{=} y \wedge \alpha(x)$$

Figure 32: The arrow part of the functor $\mathcal{P}: C^o \rightarrow \text{Cat}$

can be glued from predicates over A along a covering morphism $f: A \rightarrow B$. The requirement in (7.1), that a sheaf ς must map every covering $\bigcup_{i \in I} A_i = B$ to a corresponding decomposition of the elements of ςB is generalized to the requirement that $\mathcal{P}$ must map every cover by morphisms $\{f_i: A_i \rightarrow B\}_{i \in I}$, with the intersections annotated as in Fig. 33, to a decomposition of the predicates in the form

$$\mathcal{P}B \simeq \left\{ \langle d_{ij} \rangle_{i,j \in I} \in \prod_{i,j \in I} \coprod_{\substack{x_i \in \mathcal{P}A_i \\ x_j \in \mathcal{P}A_j}} \mathcal{P}E_{ij}(\delta_{ij}^* x_i, \rho_{ij}^* x_j) \mid \right. \\ \left. \forall i \in I. \eta_i^* d_{ii} = \text{id}_{x_i} \wedge \forall ijk \in I. p_{ij}^* d_{ij} \circ p_{jk}^* d_{jk} = p_{ik}^* d_{ik} \right\} \quad (7.2)$$

A hyperdoctrine $\mathcal{P}$ satisfying (7.2) is called a *stack* [Gir71, Ch. II]⁷. Note that (7.2) only specifies the object




Figure 33: The intersections of the elements of cover $\{f_i: A_i \rightarrow B\}_{i \in I}$ are their pullbacks

⁶If $\mathbb{S} = \text{Esp}$ is the category of topological spaces and $\mathcal{P}X = \mathcal{O}X$ is the lattice of open neighborhoods of a space X , then a predicate is a continuous map $\alpha: X \rightarrow \mathbb{2}$ into the Sierpiński space $\mathbb{2} = \{0, 1\}$ with $\{1\}$ open. The comprehension of the indexed category $\mathcal{O}: \text{Esp}^o \rightarrow \text{Cat}$ is then the fibration $\text{Esp}/\mathbb{2} \xrightarrow{\text{Dom}} \text{Esp}$. Here we restrict to the discrete case since we just need to build logical intuition.

part of the *stack condition*. To lift the sheaf condition from sets to categories, the stack condition also has a morphism part, which is not hard to reconstruct. Here, though, we are interested in a special case of the stack condition, and in special stack morphisms. When $\mathbb{S}$ has stable coproducts $A = \coprod_{i \in I} A_i$, or when we freely adjoin them, then B can be covered by a single morphism $f: A \rightarrow B$, where $f = [f_i]_{i \in I}$, and the stack requirement in (7.2) boils down to the *descent condition*

$$\mathcal{P}B \simeq \coprod_{x \in \mathcal{P}A} \left\{ d \in \mathcal{P}E(\delta^*x, \rho^*x) \mid \eta^*d = \text{id}_x \wedge p_0^*d \circ p_1^*d = p_2^*d \right\} \quad (7.3)$$

The notations are obtained from Fig. 33 again, renaming $p_{ij} \mapsto p_0$, $p_{jk} \mapsto p_1$, and $p_{ik} \mapsto p_2$ and erasing all other subscripts. Note that the right-hand side still only specifies the objects of the category that the descent condition requires to be equivalent to $\mathcal{P}B$. These objects are the *descent data*. The sheaf condition in (7.1) evolved to the stack condition in (7.2), and to the descent condition in (7.3). The category of descent data along f is usually denoted by Desc_f . Its objects, described in (7.3) on the right, are conveniently written in the form $x = (\delta^*x \xrightarrow{d_x} \rho^*x)$. The morphisms between descent data are:

$$\text{Desc}_f(x, y) = \left\{ \varphi \in \mathcal{P}A(x, y) \mid \begin{array}{ccc} \delta^*x & \xrightarrow{\delta^*\varphi} & \delta^*y \\ \downarrow d_x & & \downarrow d_y \\ \rho^*x & \xrightarrow{\rho^*\varphi} & \rho^*y \end{array} \right\} \quad (7.4)$$

Note the similarity with the simple nucleus morphisms in (6.1–6.2). This is not an accident. Fig. 34 shows

$$\frac{\delta^*x \xrightarrow{d} \rho^*x}{\rho_! \delta^*x \xrightarrow{d'} x} \quad \frac{\rho_! \delta^*x \xrightarrow{d'} x}{f^* f_! x \xrightarrow{d''} x}$$

Figure 34: The descent data down the kernel $\langle \delta, \rho \rangle$ of f correspond to $f^* f_!$ -algebras

the transpositions of the f -descent data using the adjunction $\rho_! \dashv \rho^*: \mathcal{P}A \rightarrow \mathcal{P}E$ and the *Beck-Chevalley* isomorphism⁸

$$\rho_! \delta^*x \cong f^* f_! \quad (7.5)$$

As explained in [Pav91a], the Beck-Chevalley isomorphism assures that the predicates of a hyperdoctrine $\mathcal{P}$ recognize the pullbacks in $\mathbb{S}$. In the logical notation of Fig. 32, (7.5) becomes

$$\exists e : E. \delta(e) = x \wedge \rho(e) = x' \iff f(x) = f(x') \quad (7.6)$$

If the subobject $E \xrightarrow{\langle \delta, \rho \rangle} A \times A$ is viewed as a binary relation, and $\exists e : E. \delta(e) = x \wedge \rho(e) = x'$ is abbreviated to xEx' , then (7.5) becomes $xEx' \iff f(x) = f(x')$. E is thus the kernel of the function f . The crucial observation of [BR70] was that the transpositions in Fig. 34 map the conditions in (7.3), which make $\delta^*x \xrightarrow{d} \rho^*x$ into descent data, precisely correspond to the conditions in (10), which make $f^* f_! x \xrightarrow{d''} x$ into an $\overleftarrow{f}$ -algebra for the monad $\overleftarrow{f} = f^* f_!$. The category Desc_f of descent data is equivalent to the category $\mathcal{P}A^{\overleftarrow{f}}$

⁷The book is in French, so stacks are called *champs*. The stack condition goes back to a much earlier seminar on Grothendieck's reconstruction of Teichmüller spaces. Although Giraud did not work with indexed categories but with their fibered comprehensions, and hyperdoctrines were not yet defined, the idea of stack condition was there.

⁸This isomorphism was attributed to unpublished work of Jon Beck by Lawvere [Law70] and to unpublished work of Claude Chevalley by Bénabou and Roubaud [BR70].

of algebras. The descent condition (7.3) thus implies that $\mathcal{P}B$ must be equivalent to $\mathcal{P}A^{\overleftarrow{f}}$. But Prop. 6.1 says that $\mathcal{P}A^{\overleftarrow{f}}$ is equivalent to $\mathcal{P}B^{\overleftarrow{f}}$. In summary, a morphism $f: A \rightarrow B$ in $\mathbb{S}$ satisfies the descent requirement if and only if

$$\mathcal{P}B \simeq \text{Desc}_f \simeq \mathcal{P}A^{\overleftarrow{f}} \simeq \mathcal{P}B^{\overleftarrow{f}} \quad (7.7)$$

where $\mathcal{P}A$ and $\mathcal{P}B$ are assumed to be absolutely complete.⁹

Simple descent monads. In words, (7.7) says that $\mathcal{P}A$ covers $\mathcal{P}B$ along a descent morphism $f: A \rightarrow B$ precisely when the objects of $\mathcal{P}B$ are retracts of the objects of $\mathcal{P}A$ along f^* , as specified in $\mathcal{P}B^{\overleftarrow{f}}$. Lifting the f^* -retractions to $\overleftarrow{f}$ -algebras in $\mathcal{P}A$, and then transposing them to $\mathcal{P}E$ along the Beck-Chevalley and $(\rho! \dashv \rho^*)$ -adjunction yields the f -descent data in Desc_f . On the other hand, the equivalence (6.5) of the retractions in $\mathcal{P}B^{\overleftarrow{f}}$ and the $\overleftarrow{F}$ -algebras in $\mathcal{P}B^{\overleftarrow{f}}$ explain the utility of descent monads and comonads [Bal12, CDGV07, Hes10, Mes06a].

7.2. Kan’s adjunction. In the final section, we consider the initial example of adjunctions. The idea of adjunction goes back at least to Évariste Galois, or, depending on how you think of it, much further back to Heraclitus [Lam81], and still further into the roots of logic [Law69]. Yet the categorical definition of an adjunction as two categories, two functors, and two natural transformations goes back to the late 1950s, to Daniel Kan’s work in homotopy theory [Kan58a]. Kan defined the Kan extensions to capture this particular adjunction, like Eilenberg and MacLane defined categories and functors to define certain natural transformations.

Simplices and the simplex category. Eilenberg simplified computing homology groups by decomposing topological spaces into simplices [Eil44]. An m -simplex is the set

$$\Delta_{[m]} = \left\{ \vec{x} \in [0, 1]^{m+1} \mid \sum_{i=0}^m x_i = 1 \right\} \quad (7.8)$$

with the product topology induced by the open intervals on $[0, 1]$. The relevant structure of a topological space X is captured by the families of continuous maps $\Delta_{[m]} \rightarrow X$, for all $m \in \mathbb{N}$. Some of these maps do not *embed* some simplices into the space, but contain degeneracies or singularities; yet considering all such maps assures that any simplices that can be embedded into X will be embedded by some of the maps. Since the simplicial structure is captured by each $\Delta_{[m]}$ ’s projections onto all $\Delta_{[\ell]}$ s for $\ell < m$, and by $\Delta_{[m]}$ ’s embeddings into all $\Delta_{[n]}$ s for $n > m$, a simplicial structure of a space corresponds to a functor of the form $\Delta_{[-]} : \Delta \rightarrow \text{Esp}$, where Esp is the category of topological spaces and continuous maps¹⁰, and Δ is the simplex category. Its objects are finite ordinals

$$[m] = \{0 < 1 < 2 < \dots < m\}$$

while its morphisms are the order-preserving functions [EZ50]. All information about the simplicial structure of topological spaces is thus captured by the matrix

$$\begin{aligned} \widehat{\Delta}: \Delta^o \times \text{Esp} &\rightarrow \text{Set} \\ [m] \times X &\mapsto \text{Esp}(\Delta_{[m]}, X) \end{aligned} \quad (7.9)$$

In the language of Sec. 2, this is the “*context matrix*” of homotopy theory. Its “*concept analysis*” starts by making this matrix representable by adjoint functors along the lines of Sec. 3.3, instantiating (3.6–3.5) to

⁹Without this assumption, $\mathcal{P}B$ and $\mathcal{P}B^{\overleftarrow{f}}$ must be weakly (Morita) equivalent.

¹⁰We denote the category of topological spaces by the abbreviation Esp of the French word *espace*, not just because there are other things called Top in the same contexts, but also as authors’ reminder-to-self of the tacit sources of the approach [Gro71, AGV64].

$\Phi = \Delta$. But the matrix $\widehat{\Delta}$ is already representable by the functor $\Delta_{[-]} : \Delta \rightarrow \mathbf{Esp}$, and $\mathbf{Esp}$ is already complete. Leaving out of (3.6) the completion of $\mathbb{B} = \mathbf{Esp}$, the adjunction in (3.5) boils down to

$$\begin{array}{ccc}
 \mathbb{K} \xrightarrow{K} \Delta & \mathbf{sSet} & (\Delta_{[-]}/X \xrightarrow{\text{Dom}} \Delta) \\
 \downarrow & \Delta^* \begin{array}{c} \uparrow \\ \downarrow \end{array} \Delta_* & \uparrow \\
 \varinjlim (\mathbb{K} \xrightarrow{K} \Delta \xrightarrow{\Delta_{[-]}} \mathbf{Esp}) & \mathbf{Esp} & X
 \end{array} \tag{7.10}$$

where the extension Φ_* from (3.5) becomes Δ_* in (7.10) because

$$\varprojlim (1 \xrightarrow{X} \mathbf{Esp} \xrightarrow{\Delta_*} \mathbf{Dfib}/\Delta) = (\Delta_{[-]}/X \xrightarrow{\text{Dom}} \Delta)$$

The adjunction $(\Delta^* \dashv \Delta_* : \mathbf{Esp} \rightarrow \mathbf{sSet})$ has been extensively studied. $\Delta^* : \mathbf{sSet} \rightarrow \mathbf{Esp}$ is the *geometric realization* of simplicial sets [Mil57], whereas $\Delta_* : \mathbf{Esp} \rightarrow \mathbf{sSet}$ is the *singular decomposition* of spaces. Eilenberg's singular homology was also based on it [Eil44]. Kan spelled out the concept of adjunction as the abstract structure underlying the concrete relationship between these two functors [Kan58a, Kan58c]. The idea was that the adjunction $\Delta^* \dashv \Delta_*$ should provide a general method for transferring the invariants of interest between the geometric universe of spaces and an algebraic or combinatorial framework for computations. For a geometric realization $\Delta^* K \in \mathbf{Esp}$ of a complex $K \in \mathbf{sSet}$, the homotopy groups can indeed be computed combinatorially, from the structure of K alone [Kan58b], since the spaces in the form $\Delta^* K$ boil down to Whitehead's CW-complexes [Mil57, Whi49]. But not all spaces are in this form.

Trouble with localizations. The upshot of Kan's adjunction $\Delta^* \dashv \Delta_* : \mathbf{Esp} \rightarrow \mathbf{sSet}$ is that for any space X , we can construct a CW-complex $\overrightarrow{\Delta}X = \Delta^* \Delta_* X$, with a continuous map $\overrightarrow{\Delta}X \xrightarrow{\varepsilon} X$, the counit of Kan's adjunction. This is the best approximation of X by a CW-complex. When do such approximations preserve the invariants of interest? By the late 1950s, it was clear that combinatorial approximations work in many cases, certainly whenever ε is invertible. In general, though, even $\overrightarrow{\Delta}\overrightarrow{\Delta}X \xrightarrow{\varepsilon} \overrightarrow{\Delta}X$ is not always invertible. The idea of approximating topological spaces by combinatorial complexes thus grew into a quest for making the units or the counits of adjunctions invertible. Which spaces have the same invariants as the geometric realizations of their singular¹¹ decompositions? For particular invariants, there are direct answers [EM45, EM50]. In general, though, the localizations at suitable spaces along suitable reflections or coreflections is the abstract form of spectral analysis and can be construed as an extension of concept analysis, as presented in Sec. 2. Some of the most influential methods of algebraic topology can be thought of in these terms. Grossly oversimplifying, we mention three approaches.

The direct approach [GZ12, Bor94, Vol. I, Ch. 5] was to enlarge the given category by formal inverses of a family of arrows, usually called weak equivalences $\mathcal{W}$. They are made invertible in a category of fractions just like the non-zero integers are made invertible in the field of rationals. But adjoining formal inverses for a large class $\mathcal{W}$ in a large category like $\mathbf{Esp}$ usually leads to a category with large classes of morphisms between pairs of small objects.

Another approach [DHKS05, Qui67] was to factor out the $\mathcal{W}$ -arrows using two families of arrows, called fibrations $\mathcal{F}$ and cofibrations $\mathcal{C}$, such that the pairs of families $(\mathcal{C} \cap \mathcal{W}, \mathcal{F})$ and $(\mathcal{C}, \mathcal{W} \cap \mathcal{F})$ form weak factorization systems [Rie14, Part III]. It follows that every morphism f can be factored as in Fig. 35. Localizing every f at the corresponding $\mathcal{C} \cap \mathcal{F}$ -component $f^\#$ makes the weak equivalences $\mathcal{W}$ invertible. The triples $(\mathcal{C}, \mathcal{W}, \mathcal{F})$ are called *model structures* since they make categories into abstract models of homotopy theories. The general idea originates from the model structures in $\mathbf{Esp}$ and $\mathbf{sSet}$, making the adjunction in

¹¹The word "singular" here means that the simplices, into which space may be decomposed, do not have to be embedded into it, which would make the decomposition *regular*, but that the continuous maps from their geometric realizations may have *singularities*.




Figure 35: A decomposition of f by a model structure $(C, \mathcal{W}, \mathcal{F})$

(7.10) into an equivalence. This was Dan Quillen's approach [Qui69, Qui67]. Here we worked out a model structure on the category of adjunctions.

The third approach [AT69, AT70] tackles the task of making the arrows $\vec{\Delta}X \xrightarrow{\varepsilon} X$ invertible by modifying the comonad $\vec{\Delta}$ until it becomes idempotent and then localizing at the coalgebras of this idempotent comonad. Note that this approach does not tamper with the continuous maps in $\mathbf{Esp}$, be it to make some of them formally invertible, or to factor them out. The idea is that a comonad $\vec{\Delta}_\infty : \mathbf{Esp} \rightarrow \mathbf{Esp}$, should localize any space X at a space $\vec{\Delta}_\infty X$ such that $\vec{\Delta} \vec{\Delta}_\infty X \xrightarrow{\varepsilon} \vec{\Delta}_\infty X$. It follows that $\vec{\Delta}_\infty \vec{\Delta}_\infty X \xrightarrow{\varepsilon} \vec{\Delta}_\infty X$ must hold and the comonad $\vec{\Delta}_\infty$ must be idempotent. The quest for such a comonad is illustrated in Fig. 36 where $\mathbf{Esp}^{\vec{\Delta}}$




Figure 36: Iterating the comonad resolutions for $\vec{\Delta}$

is the category of $\vec{\Delta}$ -coalgebras, $V^* \dashv V_* : \mathbf{Esp} \rightarrow \mathbf{Esp}^{\vec{\Delta}}$ is its final resolution, and Δ^0 is the comparison functor, mapping a complex K to the coalgebra $\Delta^* K \xrightarrow{\eta^*} \Delta^* \Delta_* \Delta^* K$. Since $\mathbf{sSet}$ is a complete category, Δ^0 has a right adjoint Δ_0 , and they induce the comonad $\vec{\Delta}_0 = \Delta^0 \Delta_0$ on $\mathbf{Esp}^{\vec{\Delta}}$. If $\vec{\Delta}$ was idempotent, the final resolution $V^* \dashv V_*$ would be a coreflection, and the comonad $\vec{\Delta}_0$ would be isomorphic to the identity. But $\vec{\Delta}$ is not idempotent, $\vec{\Delta}_0$ is not the identity, and its final resolution, generically denoted $V^* \dashv V_*$ again, induces a comparison functor Δ_1 , which induces the comonad $\vec{\Delta}_1 = \Delta^1 \Delta_1$, which induces the comonad $\vec{\Delta}_2$, and so on. The tower of final resolutions of these monads, displayed in Fig. 36 $V^* \dashv V_*$, continues infinitely and produces coalgebras over coalgebras over coalgebras... , which span inductive towers in $\mathbf{Esp}$. At each step, the parts of a space that are not approximable by geometric realizations are projected away, but because of that, some previously approximable parts cease to be approximable. This keeps occurring after infinitely many steps and pushes the inductive tower into ordinal lengths, echoing Cantor's quest for accumulation points of the convergence domains of the Fourier series, which led him to discover ordinals. Since $\mathbf{Esp}$ is

cocomplete, the process eventually stabilizes, and the resulting idempotent comonad $\overrightarrow{\Delta}_\infty$ localizes at the spaces that can be analyzed combinatorially, in terms of simplicial sets alone, using Kan's adjunction $\Delta^* \dashv \Delta_*$.




Figure 37: The nucleus of the Kan adjunction

The nuclear solution. The nucleus of Kan's adjunction, displayed in Fig. 37, allows analyzing a much larger family of spaces combinatorially, also in terms of simplicial sets, but equipped with $\overleftarrow{\Delta}$ -algebra structures for the monad $\overleftarrow{\Delta} = \Delta_* \Delta^*$. This larger family of spaces arises already at the first step in Fig. 36, without going down the tower. They are the spaces that can be recovered as retracts of geometric realizations of their singular decompositions. The embedding parts of the retractions make them into $\overrightarrow{\Delta}$ -coalgebras in $\text{Esp}^{\overrightarrow{\Delta}}$. Thm. 5.1 allows presenting such spaces as coalgebras over the simplicial algebras in $\text{sSet}^{\overleftarrow{\Delta}}$. If that presentation, albeit combinatorial, seems complicated, note that Prop. 6.1 allows simplifying them, and provides an effective equivalence between the spaces with coalgebras in $\text{Esp}^{\overrightarrow{\Delta}}$ and the simplicial sets with idempotents in $\text{sSet}^{\overleftarrow{\Delta}}$.

8. CLOSING

Categorical dimension reduction. The task of dimension reduction often arises in mathematics: manifolds are reduced to their boundaries, curves are approximated by points, linear operators are diagonalized using spectra. Spectral methods are the tenet of concept mining and principal component analysis of data [AFK⁺01]. In category theory, on the other hand, the way up into higher categories has been more attractive than the way down, for a variety of reasons. One is that categories themselves expand into the higher dimensions [BM09, Lei04, Lur09, Sim11]. Another is that Grothendieck's *homotopy hypothesis* [Gro18, Mal05] made higher category theory into a geometric pursuit, by some accounts subsuming homotopy theory. The nucleus results seem to suggest that there might be a way down even in category theory. On the object level, the nucleus construction ostensibly follows the familiar procedures of spectral decomposition. The nuclear adjunction induced by a categorical matrix can be construed as a diagonal matrix, spanned by the eigenvectors of the original matrix. On the meta-level, the fact that the categories of categories needed for concept analysis do not form higher categories but interact with each other within their own dimensionality seems unusual. It may signal a shortcoming of the current approach. Or it may signal the robustness and stability of categories

as a mathematical tool for the application areas where the way up is the way down, like statistics, machine learning, and data analysis; the areas where the problems get simpler as we go, and not more complicated.

Summary. We saw adjunctions in action in recommender systems in Sec. 2, and in algebraic geometry and homotopy theory in Sec. 7. In Sections 3–6, we derived a common denominator: the nuclear adjunctions. It is a powerful tool of mathematical conceptualizations; in a certain sense universal. In the meantime, it proved its worth by untying (as inadvertently as it came about) a fundamental problem, posed by Jim Lambek in 1966: *to complete any given category under limits and colimits in such a way that any limits and colimits that may already exist are preserved* [Lam66]. We call such completions *tight*. Dedekind constructed the reals as a tight completion of the rationals. MacNeille generalized Dedekind’s construction to posets. The task was thus to lift the Dedekind-MacNeille completions from posets to categories. It lifts readily to poset-enriched categories, with no major surprises, yet with significant and varied applications [Pav12, Pav13, PS17, Wil13]. Yet the ordinary, Set-enriched categories offer resistance to being tightly completed under all limits and colimits, for fundamental reasons spelled out by John Isbell [Isb68, Thm. 3.1]. On the other hand, the concept analysis requires nothing less than tight completions: completing the data contexts from above and from below at once, while preserving the previous completions. And in recommender systems, avoiding blind amplifications and echoing everything requires going beyond the posetal and numeric models. So be it as it may, Isbell notwithstanding, we cannot avoid adding some limits and colimits tightly. Which ones? This is where the nucleus kicked in. The idea was sketched in [KP15]. It has been worked out in several forms, and an attempt to present it was made in [PH22].

Back to Escher. The nucleus of an adjunction, illustrated in the Introduction by Escher’s “Drawing hands”, is a descendant of Dedekind’s construction of the real numbers as *cuts* of the rationals. A cut is a pair $c = \langle \overleftarrow{c}, \overrightarrow{c} \rangle \in \Downarrow \mathbb{Q} \times \Uparrow \mathbb{Q}$ where $\overleftarrow{c}$ is the set of lower bounds of $\overrightarrow{c}$, and $\overrightarrow{c}$ is the set of upper bounds of $\overleftarrow{c}$. The “Drawing hands” of Dedekind’s cuts are displayed in Fig. 38 on the left. The “Drawing hands” of a simple




Figure 38: The nucleus of an adjunction echoes the idea of Dedekind cuts

nucleus are displayed on the right. Since the components $\overleftarrow{c}$ and $\overrightarrow{c}$ of a cut determine each other, they both carry the same information, and either can be used alone to represent a real number. The real numbers have three isomorphic representations:

$$\{\overleftarrow{c} \in \Downarrow \mathbb{Q} \mid \overleftarrow{c} = \nabla \Delta \overleftarrow{c}\} \cong \{ \langle \overleftarrow{c}, \overrightarrow{c} \rangle \in \Downarrow \mathbb{Q} \times \Uparrow \mathbb{Q} \mid \overleftarrow{c} = \nabla \overrightarrow{c} \wedge \Delta \overleftarrow{c} = \overrightarrow{c} \} \cong \{ \overrightarrow{c} \in \Uparrow \mathbb{Q} \mid \Delta \nabla \overrightarrow{c} = \overrightarrow{c} \}$$

This just says that the same real number can be approached from below, from above, or tightly, from both directions. The nucleus construction shows that concepts are like real numbers: a concept can be presented as an algebra, or as a coalgebra, and they determine each other. And it shows that concepts are unlike real numbers since the algebra and the coalgebra are not each other’s images, but mere retracts. As a nucleus of an adjunction, an instance of itself, the nucleus construction seems easier to use than to understand. The path

forward might be to conceptualize it in its own terms. Part II of this work will describe the nucleus monad on monads and comonads, covering itself. Part III will spell out the induced model structures and localizations.

Acknowledgements and apology. While working on the results presented in this paper, the first author was assisted by Liang-Ting Chen, Tobias Heindel, Toshiki Kataoka, Peter-Michael Seidel, and Christina Vassilakopoulou. Toshiki coauthored [KP15], where an early version of Prop. 4.1 was proved. Peter-Michael coauthored [PS17], where parts of the proof of Thm. 5.1 were constructed. A longer version of the present paper, covering a wider family of examples and the results now left for parts II and III, has been circulated since 2019 and posted on arxiv in 2020. It also contained an account of the nucleus monads on monads and comonads, induced by the nucleus monad on adjunctions presented here. The dualities between monads and comonads in abstract 2-categories, studied by Ross Street in the final sections of [Str72], induce closely related 2-monads directly, without a detour into adjunctions. Ross presented the result concerning the 2-monad on monads in his seminar, provided helpful comments, and considered the mentioned previous version for publication in a journal that he recommended. After a couple of years, the anonymous referees returned very succinct reviews, dispensing pedagogic style advice as a conditional acceptance recommendation. In view of the absence of substantive comments, suggesting an apparent lack of interest, we withdrew the submission, but realized that a part of the problem was our weaving and knotting of several narratives into one, each of the coauthors trusting that the other one would assure a tidy presentation. We apologize for the confusion and thank the readers and the users seeing us through it.

REFERENCES

- [AFK⁺01] Yossi Azar, Amos Fiat, Anna Karlin, Frank McSherry, and Jared Saia. Spectral analysis of data. In *Proceedings of the thirty-third annual ACM Symposium on Theory of Computing*, STOC '01, pages 619–626, New York, NY, USA, 2001. ACM. URL: <http://doi.acm.org/10.1145/380752.380859>, doi:<http://doi.acm.org/10.1145/380752.380859>.
- [AGV64] Michael Artin, Alexander Grothendieck, and Jean-Louis Verdier, editors. *Séminaire de Géométrie Algébrique: Théorie des Topos et Cohomologie Étale des Schemas (SGA 4)*, volume 269,270,305 of *Lecture Notes in Mathematics*. Springer-Verlag, 1964. Second edition, 1972.
- [AT69] H. Applegate and Myles Tierney. Categories with models. In Eckmann and Tierney [ET69], pages 156–244. Reprinted in *Theory and Applications of Categories* 18(2008).
- [AT70] H. Applegate and Myles Tierney. Iterated cotriples. In S. et al MacLane, editor, *Reports of the Midwest Category Seminar IV*, pages 56–99, Berlin, Heidelberg, 1970. Springer.
- [Aud74] Claude Auderset. Adjonctions et monades au niveau des 2-catégories. *Cahiers de Topologie et Géométrie Différentielle Catégoriques*, 15(1):3–20, 1974.
- [Bal12] Paul Balmer. Descent in triangulated categories. *Mathematische Annalen*, 353:109–125, 2012. URL: <http://doi.org/10.1007/s00208-011-0674-z>, doi:10.1007/s00208-011-0674-z.
- [Bar69] Michael Barr. Coalgebras in a category of algebras. In *Category Theory, Homology Theory and their Applications I*, volume 86 of *Lecture Notes in Mathematics*, pages 1–12. Springer, 1969. URL: <http://dx.doi.org/10.1007/BFb0079381>, doi:10.1007/BFb0079381.
- [BB67] B. Banaschewski and G. Bruns. Categorical characterization of the MacNeille completion. *Archiv der Mathematik*, 18(4):369–377, September 1967.
- [BB69] Michael Barr and Jon Beck. Homology and standard constructions. In Eckmann and Tierney [ET69], pages 245–335. Reprinted in *Theory and Applications of Categories* 18(2008).
- [Bec67] Jonathan Mock Beck. *Triples, Algebras and Cohomology*. PhD thesis, Columbia University, 1967. Reprinted in *Theory and Applications of Categories*, No. 2, 2003, pp. 1–59.
- [Bén67] Jean Bénabou. Introduction to bicategories. In *Reports of the Midwest Category Seminar*, number 47 in *Lecture Notes in Mathematics*, pages 1–77. Springer, 1967. doi:<https://doi.org/10.1007/BFb0074299>.
- [BK11] Adriana Balan and Alexander Kurz. On coalgebras over algebras. *Theoretical Computer Science*, 412(38):4989 – 5005, 2011. CMCS Tenth Anniversary Meeting. URL: <http://www.sciencedirect.com/science/article/pii/S0304397511002519>, doi:<https://doi.org/10.1016/j.tcs.2011.03.021>.
- [BM09] John C. Baez and J. Peter May. *Towards Higher Categories*, volume 152 of *IMA Volumes in Mathematics and its Applications*. Springer, 2009.

- [Bor94] Francis Borceux. *Handbook of Categorical Algebra*. Number 50 in Encyclopedia of Mathematics and its Applications. Cambridge University Press, 1994. Three volumes.
- [BR70] Jean Bénabou and Jacques Roubaud. Monades et descente. *CR Acad. Sci. Paris Sér. AB*, 270(A96-A98):2, 1970.
- [BTv22] Tai-Danae Bradley, John Terilla, and Yiannis Vlassopoulos. An enriched category theory of language: from syntax to semantics. *La Matematica*, 1(2):551–580, 2022.
- [Bun13] Marta Bunge. Tightly bounded completions. *Theory and Applications of Categories*, 28(8):213–240, 2013.
- [BVMV11] Tomasz Brzeziński, Adrian Vazquez Marquez, and Joost Vercruysse. The Eilenberg-Moore category and a Beck-type theorem for a Morita context. *Applied Categorical Structures*, 19:821–858, 2011.
- [BW85] Michael Barr and Charles Wells. *Toposes, Triples, and Theories*. Number 278 in Grundlehren der mathematischen Wissenschaften. Springer-Verlag, 1985. Republished in: Reprints in Theory and Applications of Categories, No. 12 (2005).
- [CDGV07] Stefaan Caenepeel, Erwin De Groot, and Joost Vercruysse. Galois theory for comatrix corings: descent theory, morita theory, frobenius and separability properties. *Transactions of the American Mathematical Society*, 359(1):185–226, 2007.
- [CH88] Thierry Coquand and Gérard P. Huet. The calculus of constructions. *Inf. Comput.*, 76:95–120, 1988.
- [CR04] Claudio Carpineto and Giovanni Romano. *Concept Data Analysis: Theory and Applications*. John Wiley & Sons, 2004.
- [Ded01] Richard Dedekind. *Essays on the Theory of Numbers*. The Religion of Science Library. Open Court Publishing Company, 1901. reprinted by Dover in 1963.
- [DHKS05] William G. Dwyer, Philip S. Hirschhorn, Daniel M. Kan, and Jeffrey H. Smith. *Homotopy Limit Functors on Model Categories and Homotopical Categories*, volume 113 of *Mathematical surveys and monographs*. American Mathematical Society, 2005.
- [Eil44] Samuel Eilenberg. Singular homology theory. *Annals of Mathematics*, 45(3):407–447, 1944.
- [EM45] Samuel Eilenberg and Saunders MacLane. Relations between homology and homotopy groups of spaces. *Annals of Mathematics*, 46(3):480–509, 1945.
- [EM50] Samuel Eilenberg and Saunders MacLane. Relations between homology and homotopy groups of spaces. II. *Annals of Mathematics*, 51(3):514–533, 1950.
- [EM⁺65] Samuel Eilenberg, John C Moore, et al. Adjoint functors and triples. *Illinois Journal of Mathematics*, 9(3):381–398, 1965.
- [ET69] Beno Eckmann and Myles Tierney, editors. *Seminar on Triples and Categorical Homology Theory*, number 80 in Lecture Notes in Mathematics. Springer-Verlag, 1969. Reprinted in *Theory and Applications of Categories* 18(2008).
- [EZ50] Samuel Eilenberg and Joseph A Zilber. Semi-simplicial complexes and singular homology. *Annals of Mathematics*, 51(3):499–513, 1950.
- [Fre64] Peter Freyd. *Abelian Categories: an Introduction to the Theory of Functors*. Harper and Row, 1964. Republished in: Reprints in Theory and Applications of Categories, No. 3 (2003).
- [Gir64] Jean Giraud. *Méthode de la descente*. Number 2 in Mémoires de la Société Mathématique de France. Société mathématique de France, 1964. URL: http://www.numdam.org/item/MSMF_1964__2__R3_0/, doi:10.24033/msmf.2.
- [Gir71] Jean Giraud. *Cohomologie non-abélienne*. Number 179 in Grundlehren der mathematischen Wissenschaften. Springer-Verlag, 1971.
- [Gro55] Alexander Grothendieck. *Produits tensoriels topologiques et espaces nucléaires*, volume 16 of *Memoirs of the American Mathematical Soc.* AMS, 1955.
- [Gro60] Alexander Grothendieck. Technique de descente et théorèmes d’existence en géométrie algébrique. i. généralités. descente par morphismes fidèlement plats. In *Séminaire Bourbaki : années 1958/59 - 1959/60, exposés 169-204*, number 5 in Séminaire Bourbaki, pages 299–327. Société mathématique de France, 1960. talk:190. URL: http://www.numdam.org/item/SB_1958-1960__5__299_0.
- [Gro71] Alexander Grothendieck. *Revêtement étales et groupe fondamental (SGA1)*, volume 224 of *Lecture Notes in Mathematics*. Springer, 1971.
- [Gro18] Alexander Grothendieck. Pursuing stacks. *CoRR*, abs/2111.01000, 2018. URL: <http://arxiv.org/abs/2111.01000>, arXiv:2111.01000.
- [GSW05] Bernhard Ganter, Gerd Stumme, and Rudolf Wille, editors. *Formal Concept Analysis, Foundations and Applications*, volume 3626 of *Lecture Notes in Computer Science*. Springer, 2005.
- [GW99] Bernhard Ganter and Rudolf Wille. *Formal Concept Analysis: Mathematical Foundations*. Springer, Berlin/Heidelberg, 1999.
- [GZ12] Peter Gabriel and Michel Zisman. *Calculus of fractions and homotopy theory*, volume 35 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, 2012.
- [Hes10] Kathryn Hess. A general framework for homotopic descent and codescent. *CoRR*, abs/1001.1556, 2010. URL: <http://arxiv.org/abs/1001.1556>, arXiv:1001.1556.

- [Hub61] Peter J. Huber. Homotopy theory in general categories. *Mathematische Annalen*, 144:361–385, 1961. URL: <http://doi.org/10.1007/bf01396534>, doi:10.1007/bf01396534.
- [Hub62] Peter J. Huber. Standard constructions in abelian categories. *Mathematische Annalen*, 146(4):321–325, 1962.
- [Isb68] John R. Isbell. Small categories and completeness. *Mathematical Systems Theory*, 2(1):27–50, 1968.
- [Jac94] Bart Jacobs. Coalgebras and approximation. In A. Nerode and Yu. V. Matiyasevich, editors, *International Symposium on Logical Foundations of Computer Science*, volume 813 of *Lecture Notes in Computer Science*, pages 173–183. Springer, 1994.
- [Jac01] Bart Jacobs. *Categorical Logic and Type Theory*. Studies in logic and the foundations of mathematics. Elsevier, 2001. URL: <https://books.google.com/books?id=0hhyL4WG3ngC>.
- [Jac13] Bart Jacobs. Bases as coalgebras. *Logical Methods in Computer Science*, 9(3), 2013. doi:10.2168/LMCS-9(3:23)2013.
- [Jar19] J.F. Jardine. Galois descent criteria. In D.G. Davis et al, editor, *Homotopy Theory: Tools and Applications*, volume 729 of *Contemporary Mathematics*, pages 129–156. American Mathematical Society, 2019.
- [Joh82] Peter Johnstone. *Stone Spaces*. Number 3 in Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1982.
- [JT04] George Janelidze and Walter Tholen. Facets of descent, I–III. *Applied Categorical Structures*, 2,5,12:245–281,229–248,461–477, 1994,1997,2004.
- [Kan58a] Daniel M. Kan. Adjoint functors. *Transactions of the American Mathematical Society*, 87(2):294–329, 1958.
- [Kan58b] Daniel M Kan. A combinatorial definition of homotopy groups. *Annals of Mathematics*, 67(2):282–312, 1958.
- [Kan58c] Daniel M. Kan. Functors involving c.s.s. complexes. *Transactions of the American Mathematical Society*, 87(2):330–346, 1958.
- [KL89] G. Max Kelly and F. William Lawvere. On the complete lattice of essential localizations. *Bull. Soc. Math. de Belgique*, 41:289–319, 1989.
- [Kle65] Heinrich Kleisli. Every standard construction is induced by a pair of adjoint functors. *Proceedings of the American Mathematical Society*, pages 544–546, 1965.
- [Koc95] Anders Kock. Monads for which structures are adjoint to units. *J. Pure Appl. Algebra*, 104:41–59, 1995. doi:[https://doi.org/10.1016/0022-4049\(94\)00111-U](https://doi.org/10.1016/0022-4049(94)00111-U).
- [KP15] Toshiki Kataoka and Dusko Pavlovic. Towards Concept Analysis in Categories: Limit Inferior as Algebra, Limit Superior as Coalgebra. In L.S. Moss and P. Sobocinski, editors, *Proceedings of CALCO 2015*, volume 35 of *LIPICs*, pages 130–155, Dagstuhl, Germany, 2015. Leibniz-Zentrum für Informatik. arxiv:1505.01098. doi:10.4230/LIPICs.CALCO.2015.130.
- [Lac] Steve Lack. Facsimile of 9/9/99, sent to Paul Taylor. 8 pages; provided by Ross Street.
- [Lam66] Joachim Lambek. *Completions of categories: seminar lectures given 1966 in Zurich*. Number 24 in Springer Lecture Notes in Mathematics. Springer-Verlag, 1966.
- [Lam81] Jim Lambek. The influence of Heraclitus on modern mathematics. In J. Agassi et al., editor, *Scientific Philosophy Today: Essays in Honor of Mario Bunge*, pages 111–121. Springer, 1981.
- [Law69] F. William Lawvere. Adjointness in foundations. *Dialectica*, 23:281–296, 1969. reprint in *Theory and Applications of Categories*, No. 16, 2006, pp.1–16.
- [Law70] F. William Lawvere. Equality in hyperdoctrines and the comprehension schema as an adjoint functor. In Alex Heller, editor, *Applications of Categorical Algebra*, number 17 in *Proceedings of Symposia in Pure Mathematics*, pages 1–14. American Mathematical Society, 1970.
- [Law73] F. William Lawvere. Metric spaces, generalised logic, and closed categories. In *Rendiconti del Seminario Matematico e Fisico di Milano*, volume 43. Tipografia Fusi, Pavia, 1973. Reprinted in *Theory and Applications of Categories*, No. 1, 2002, pp. 1–37.
- [Law89] F. William Lawvere. Display of graphics and their applications, as exemplified by 2-categories and the hegelian 'taco'. In Teodor Rus, editor, *Proceedings of AMAST 1989*, pages 51–74. University of Iowa, 1989.
- [Law96] F William Lawvere. Unity and identity of opposites in calculus and physics. *Applied categorical structures*, 4(2):167–174, 1996.
- [Lei04] Tom Leinster. *Higher Operads, Higher Categories*, volume 298 of *London Mathematical Society Lecture Note*. Cambridge University Press, 2004.
- [Lor21] Fosco Loregian. *(Co)end Calculus*. London Mathematical Society Lecture Note Series. Cambridge University Press, 2021. doi:10.1017/9781108778657.
- [LS86] Joachim Lambek and Philip Scott. *Introduction to Higher Order Categorical Logic*. Number 7 in Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1986.
- [LS02] Stephen Lack and Ross Street. The formal theory of monads II. *Journal of Pure and Applied Algebra*, 175(1):243–265, 2002.
- [Lur09] Jacob Lurie. *Higher Topos Theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, 2009.

- [Mac36] Holbrook M. MacNeille. Extensions of partially ordered sets. *Proc. Nat. Acad. Sci. USA*, 22:45–50, 1936.
- [Mal05] Georges Maltsiniotis. *La théorie de l’homotopie de Grothendieck*, volume 301 of *Astérisque*. Société mathématique de France, 2005.
- [Mar66] Jean-Marie Maranda. On fundamental constructions and adjoint functors. *Canadian Mathematical Bulletin*, 9(5):581–591, 1966. doi:10.4153/CMB-1966-072-9.
- [Mes06a] Bachuki Mesablishvili. Monads of effective descent type and comonadicity. *Theory Appl. Categ*, 16(1):1–45, 2006.
- [Mes06b] Bachuki Mesablishvili. Monads of effective descent type and comonadicity. *Theory Appl. Categ*, 16(1):1–45, 2006.
- [Mil57] John Milnor. The geometric realization of a semi-simplicial complex. *Annals of Mathematics*, 65(2):357–362, 1957.
- [ML84] Per Martin-Löf. *Intuitionistic Type Theory*. Bibliopolis, Naples, 1984.
- [Par69] Robert Paré. Absolute coequalizers. In *Category Theory, Homology Theory and their Applications I*, volume 86 of *Lecture Notes in Mathematics*, pages 132–145. Springer, 1969. URL: <http://dx.doi.org/10.1007/BFb0079381>, doi:10.1007/BFb0079381.
- [Par71] Robert Paré. On absolute colimits. *J. Alg.*, 19:80–95, 1971.
- [Pav90] Dusko Pavlovic. *Predicates and Fibrations*. PhD thesis, Rijksuniversiteit Utrecht, 1990. URL: <http://dusko.org/math-and-computation/>.
- [Pav91a] Dusko Pavlovic. Categorical interpolation: descent and the Beck-Chevalley condition without direct images. In A. Carboni et al., editor, *Category Theory, Como 1990*, volume 1488 of *Lecture Notes in Mathematics*, pages 306–326. Springer Verlag, 1991.
- [Pav91b] Dusko Pavlovic. Constructions and predicates. In D. Pitt et al., editor, *Category Theory and Computer Science ’91*, volume 530 of *Lecture Notes in Computer Science*, pages 173–197. Springer Verlag, 1991.
- [Pav96] Dusko Pavlovic. Maps II: Chasing diagrams in categorical proof theory. *J. of the IGPL*, 4(2):1–36, 1996. doi:10.1093/jigpal/4.2.159.
- [Pav12] Dusko Pavlovic. Quantitative Concept Analysis. In Florent Domenach, Dmitry I. Ignatov, and Jonas Poelmans, editors, *Proceedings of ICFC 2012*, volume 7278 of *Lecture Notes in Artificial Intelligence*, pages 260–277. Springer Verlag, 2012. arXiv:1204.5802. doi:10.1007/978-3-642-29892-9_24.
- [Pav13] Dusko Pavlovic. Bicompletions of distance matrices. In Bob Coecke, Luke Ong, and Prakash Panangaden, editors, *Computation, Logic, Games and Quantum Foundations. The Many Facets of Samson Abramsky*, volume 7860 of *Lecture Notes in Computer Science*, pages 291–310. Springer Verlag, 2013. doi:10.1007/978-3-642-38164-5_20.
- [PH22] Dusko Pavlovic and Dominic J. D. Hughes. Tight limits and completions from dedekind-macneille to lambek-isbell. *CoRR*, abs/2204.09285, 2022. arxiv:2204.09285. arXiv:2204.09285, doi:10.48550/arXiv.2204.09285.
- [PS17] Dusko Pavlovic and Peter-Michael Seidel. Quotients in monadic programming: Projective algebras are equivalent to coalgebras. In *32nd Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2017, Reykjavik, Iceland, June 20-23, 2017*, pages 1–12. IEEE Computer Society, 2017. arxiv:1701.07601. URL: <http://ieeexplore.ieee.org/xpl/mostRecentIssue.jsp?punumber=7999337>, doi:10.1109/LICS.2017.8005110.
- [Qui67] Daniel G. Quillen. *Homotopical algebra*, volume 43 of *Lecture Notes in Mathematics*. Springer, 1967.
- [Qui69] Daniel Quillen. Rational homotopy theory. *Annals of Mathematics*, pages 205–295, 1969.
- [Rie14] Emily Riehl. *Categorical Homotopy Theory*. Categorical Homotopy Theory. Cambridge University Press, 2014. URL: <https://books.google.com/books?id=6xpvAwAAQBAJ>.
- [RRS22] F. Ricci, L. Rokach, and B. Shapira. *Recommender Systems Handbook*. Springer US, 2022.
- [Sim78] Harold Simmons. A framework for topology. In A. et al. Macintyre, editor, *Logic Colloquium ’77*, Studies in Logic and the Foundations of Mathematics, pages 239–251. North-Holland, 1978.
- [Sim11] Carlos Simpson. *Homotopy Theory of Higher Categories*, volume 19 of *New Mathematical Monographs*. Cambridge University Press, 2011.
- [SS86] Stephen Schanuel and Ross Street. The free adjunction. *Cahiers de topologie et géométrie différentielle catégoriques*, 27(1):81–83, 1986.
- [Str72] Ross Street. The formal theory of monads. *Journal of Pure and Applied Algebra*, 2(2):149–168, 1972.
- [Str80] Ross Street. Fibrations in bicategories. *Cahiers de Topologie et Géométrie Différentielle Catégoriques*, 21(2):111–160, 1980.
- [Str18] Thomas Streicher. Fibered categories à la Jean Benabou. *CoRR*, abs/1801.02927, 2018. URL: <http://arxiv.org/abs/1801.02927>, arXiv:1801.02927.
- [Vis05] Angelo Vistoli. Grothendieck topologies, fibered categories and descent theory. In *Fundamental Algebraic Geometry: Grothendieck’s FGA Explained*, Mathematical surveys and monographs, pages 1–137. American Mathematical Society, 2005.
- [VT10] Joan B. Climent Vidal and Juan Carlos Soliveres Tur. Kleisli and Eilenberg-Moore constructions as parts of biadjoint situations. *Extracta Mathematicae*, 25(1):1–61, 2010.
- [Whi49] John H.C. Whitehead. Combinatorial homotopy. I and II. *Bulletin of the American Mathematical Society*, 55(3, 5):213–245, 453–496, 1949.

- [Wil13] Simon Willerton. Tight spans, Isbell completions and semi-tropical modules. *Theory and Applications of Categories*, 28(22):696–732, August 2013.
- [Zöb76] Volker Zöberlein. Doctrines on 2-categories. *Math. Zeitschrift*, 148:267–279, 1976.

APPENDIX A. COMPREHENDING PRESHEAVES AND MATRICES AS DISCRETE FIBRATIONS

Following the step from (2.3) to (2.4), the comprehension correspondence (2.4) now lifts to

$$\begin{array}{ccc}
 \text{Cat}(\mathbb{A}^o \times \mathbb{B}, \text{Set}) & \xrightleftharpoons[\Xi]{\widehat{(-)}} & \text{Dfib} / \mathbb{A} \times \mathbb{B}^o \\
 \left(\mathbb{A}^o \times \mathbb{B} \xrightarrow{\Phi} \text{Set} \right) & \mapsto & \left(\int \Phi \xrightarrow{\widehat{\Phi}} \mathbb{A} \times \mathbb{B}^o \right) \\
 \left(\mathbb{A}^o \times \mathbb{B} \xrightarrow{\Xi_E} \text{Set} \right) & \leftarrow & \left(\mathbb{E} \xrightarrow{E} \mathbb{A} \times \mathbb{B}^o \right)
 \end{array} \tag{A.1}$$

Transposing the arrow part of Φ , which maps every pair $f \in \mathbb{A}(a, a')$ and $g \in \mathbb{B}(b', b)$ into $\Phi(a', b') \xrightarrow{\Phi_{fg}} \Phi(a, b)$, the closure property expressed by the implication in (2.5) becomes the mapping

$$\mathbb{A}(a, a') \times \Phi(a', b') \times \mathbb{B}(b', b) \longrightarrow \Phi(a, b) \tag{A.2}$$

The *lower-upper* closure property expressed by (2.5) is now captured as the structure of the total category $\int \Phi$, defined as follows:

$$\begin{aligned}
 \left| \int \Phi \right| &= \coprod_{\substack{a \in \mathbb{A} \\ b \in \mathbb{B}}} \Phi(a, b) \\
 \int \Phi(x_{ab}, x'_{a'b'}) &= \{ \langle f, g \rangle \in \mathbb{A}(a, a') \times \mathbb{B}(b', b) \mid x = \Phi_{fg}(x') \}
 \end{aligned} \tag{A.3}$$

It is easy to see that the obvious projection

$$\begin{array}{ccc}
 \int \Phi & \xrightarrow{\widehat{\Phi}} & \mathbb{A} \times \mathbb{B}^o \\
 x_{ab} & \mapsto & \langle a, b \rangle
 \end{array} \tag{A.4}$$

is a discrete fibration, i.e., an object of $\text{Dfib} / \mathbb{A} \times \mathbb{B}^o$. In general, a functor $\mathbb{F} \xrightarrow{F} \mathbb{C}$ is a discrete fibration over $\mathbb{C}$ when for all $x \in \mathbb{F}$ the obvious induced functors $\mathbb{F}/x \xrightarrow{F_x} \mathbb{C}/Fx$ are isomorphisms. In other words, for every $x \in \mathbb{F}$ and every morphism $c \xrightarrow{t} Fx$ in $\mathbb{C}$, there is a unique lifting $t^! x \xrightarrow{\theta^!} x$ of t to $\mathbb{F}$, i.e., a unique $\mathbb{F}$ -morphism into x such that $F(\theta^!) = t$. For a discrete fibration $\mathbb{E} \xrightarrow{E} \mathbb{A} \times \mathbb{B}^o$, such liftings induce the arrow part of the corresponding presheaf

$$\begin{array}{ccc}
 \Xi_E: \mathbb{A}^o \times \mathbb{B} & \rightarrow & \text{Set} \\
 \langle a, b \rangle & \mapsto & \{ x \in \mathbb{E} \mid Ex = \langle a, b \rangle \}
 \end{array}$$

because any pair of morphisms $\langle f, g \rangle \in \mathbb{A}(a, a') \times \mathbb{B}^o(b, b')$ lifts to a function $\Xi_E(f, g) = \langle f, g \rangle^! : \Xi_E(a', b') \rightarrow \Xi_E(a, b)$. The equivalences in (A.1) thus yield an equivalent version of the category Mat of matrices:

$$\begin{aligned}
 |\text{Mat}| &= \coprod_{\mathbb{A}, \mathbb{B} \in \text{CAT}} \text{Dfib} / \mathbb{A} \times \mathbb{B}^o \\
 \text{Mat}(\Phi, \Psi) &= \coprod_{\substack{H \in \text{CAT}(\mathbb{A}, \mathbb{C}) \\ K \in \text{CAT}(\mathbb{B}, \mathbb{D})}} \left(\text{Dfib} / \mathbb{A} \times \mathbb{B}^o \right) \left(\Phi, (H \times K^o)^* \Psi \right)
 \end{aligned} \tag{A.5}$$

where $\Phi \in \mathbf{Dfib} / \mathbb{A} \times \mathbb{B}^o$, $\Psi \in \mathbf{Dfib} / \mathbb{C} \times \mathbb{D}^o$, and $(H \times K^o)^* \Psi$ is a pullback of Ψ along

$$(H \times K^o) : \mathbb{A} \times \mathbb{B}^o \longrightarrow \mathbb{C} \times \mathbb{D}^o$$

The notational abuse of $\mathbf{Mat}$ to denote both (2.4) and (A.5) is not just technically harmless, but its tacit identification of the two sides of (A.1) is conceptually justified by the recurring theme of categorical comprehension, used already in the next section.

APPENDIX B. IDEMPOTENTS, ABSOLUTE (CAUCHY) COMPLETIONS, AND WEAK (MORITA) EQUIVALENCES

Idempotent splitting. An endomorphism $\varphi : A \rightarrow A$ is *idempotent* if $\varphi \circ \varphi = \varphi$. A *splitting* of an idempotent φ is its epi-mono (quotient-injective) factorization

$$\begin{array}{ccc} A & \xrightarrow{\varphi} & A \\ & \searrow q & \nearrow i \\ & B & \end{array}$$

It is easy to prove (and goes back at least as far as [AGV64, Sec. IV.7.5]) that the following statements are equivalent:

- (a) $\varphi \circ \varphi = \varphi$
- (b) $q \circ i = \text{id}$
- (c) i is an equalizer and q is a coequalizer of φ and the identity

$$\begin{array}{ccc} & B & \\ i \swarrow & & \nwarrow q \\ A & \xrightarrow{\varphi} & A \\ \text{id} \xrightarrow{\quad} & & \end{array}$$

The idempotent splittings are often drawn in the form $A \begin{smallmatrix} \xrightarrow{q} \\ \xleftarrow{i} \end{smallmatrix} B$, suggesting $i \circ q = \varphi$ and $q \circ i = \text{id}$.

Since idempotent splittings are equationally characterized, and functors preserve equations, the idempotent splittings are preserved by all functors. A categorical property that is preserved by all functors is called *absolute*. By (c) above, the idempotent splittings are absolute equalizers and coequalizers. It was shown in [Par71] that these are the only absolute limits or colimits. Completing a category under all absolute limits and colimits boils down to adjoining the idempotent splittings.

Absolute and Cauchy completions. For an arbitrary category $\mathbb{C}$, the absolute completion $\underline{\mathbb{C}}$ consists of the $\mathbb{C}$ -idempotents as the objects, and the idempotent-preserving homomorphisms:

$$|\underline{\mathbb{C}}| = \coprod_{A \in |\mathbb{C}|} \{A \xrightarrow{\varphi} A \mid \varphi \circ \varphi = \varphi\} \quad (\text{B.1})$$

$$\underline{\mathbb{C}}(A \xrightarrow{\varphi} A, B \xrightarrow{\psi} B) = \left\{ f \in \mathbb{C}(A, B) \mid \begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow \varphi & & \uparrow \psi \\ A & \xrightarrow{f} & B \end{array} \right\} \quad (\text{B.2})$$

Note that the homomorphism condition $\psi \circ f \circ \varphi = f$ is equivalent to the conjunction of $f \circ \varphi = f$ and $\psi \circ f = f$. In [AGV64, Sec. IV.7.5], the construction of the category $\underline{\mathbb{C}}$ was attributed to Max Karoubi, so it came to be called the *Karoubi envelope*. It also appeared as exercise 2-B in [Fre64, p. 61].

A category $\mathbb{C}$ is said to be *Cauchy complete* if every matrix $\mathbb{C}^o \xrightarrow{\Gamma} \mathbf{Set}$ with a right adjoint $\mathbb{C} \xrightarrow{\Gamma_*} \mathbf{Set}$, where $\Gamma_*(x)$ is the set of cocones from Γ to x , is representable by some $c \in \mathbb{C}$ as $\Gamma(x) = \mathbb{C}(x, c)$. The name is motivated by the observation that the corresponding property in metric spaces, viewed as enriched categories,

characterizes the convergence of Cauchy sequences [Law73]. See also [Bor94, Vol. 1, Sec. 7.9] and [Bun13]. The following statements are equivalent for any category $\mathbb{C}$:

- (a) $\mathbb{C}$ is Cauchy complete,
- (b) $\mathbb{C}$ is absolutely complete,
- (c) all idempotents split in $\mathbb{C}$,
- (d) the embedding $\mathbb{C} \hookrightarrow \underline{\mathbb{C}}$ is an equivalence of categories.

Weak (Morita) equivalences. Categories $\mathbb{C}$ and $\mathbb{D}$ are said to be *weakly equivalent* when their absolute completions are strongly equivalent; i.e., $\mathbb{C} \sim \mathbb{D}$ means that $\underline{\mathbb{C}} \simeq \underline{\mathbb{D}}$. The weak equivalence is often named the *Morita equivalence of categories* because it is also characterized by the strong equivalence of the categories $\text{Set}^{\mathbb{C}} \simeq \text{Set}^{\mathbb{D}}$. A proof can be found in [Bor94, Vol. 1, Sec. 7.9]. This terminology is justified by the analogy of the categories $\text{Set}^{\mathbb{C}}$ and $\text{Set}^{\mathbb{D}}$ with the abelian categories Ab^R and Ab^S of R -modules and S -modules, whose equivalence was studied by Morita.¹² All of our results are valid up to weak equivalence of categories or can be construed as speaking of absolute completions. To shorten proofs, we take the latter approach, introduce absolute completions in the statements, and prove strong equivalences. Omitting this would lead to shorter statements, longer proofs, and essentially equivalent theory.

Absolute reflections. Let $\underline{\text{Cat}}$ denote the full subcategory of Cat spanned by absolutely complete categories $\underline{\mathbb{A}}, \underline{\mathbb{B}}$, etc. Checking that the idempotent splitting construction in (B.1) induces a left adjoint to the inclusion $\underline{\text{Cat}} \hookrightarrow \text{Cat}$ is a standard exercise. Any functor $F : \mathbb{A} \rightarrow \mathbb{B}$ lifts to $\underline{F} : \underline{\mathbb{A}} \rightarrow \underline{\mathbb{B}}$, where $\underline{F}x = Fx$, whether x is an idempotent or a homomorphism of idempotents. Any natural transformation $\tau : F \rightarrow G$ between any pair of functors $F, G : \mathbb{A} \rightarrow \mathbb{B}$ lifts to a natural family $\underline{\tau} : \underline{F} \rightarrow \underline{G}$ comprised of

$$(\underline{F}\varphi \xrightarrow{\underline{\tau}_\varphi} \underline{G}\varphi) = (FA \xrightarrow{F\varphi} FA \xrightarrow{\tau_A} GA \xrightarrow{G\varphi} GA) \quad (\text{B.3})$$

The naturality of τ implies not only the naturality of the family $\underline{\tau}$ but also the functoriality of the induced mappings $\text{Cat}(\mathbb{A}, \mathbb{B}) \rightarrow \text{Cat}(\underline{\mathbb{A}}, \underline{\mathbb{B}})$. The idempotent splitting in (B.1) thus induces a 2-functor $\text{Cat} \rightarrow \underline{\text{Cat}}$. This means that any monad $(\overline{T}, \eta, \mu)$ on $\mathbb{A}$ lifts to a monad $(\underline{\overline{T}}, \underline{\eta}, \underline{\mu})$ on $\underline{\mathbb{A}}$. Ditto for any comonad, adjunction, matrix, etc. Hence the corresponding 2-categories $\underline{\text{Mnd}}, \underline{\text{Cmn}}, \underline{\text{Adj}}$, etc., of the various categorical structures lifted to absolute completions, in all cases fully embedded into the general categorical structures.

Notation. Since the embedding $\underline{\text{Cat}} \hookrightarrow \text{Cat}$ is conservative, we write $F^* \dashv F_* : \underline{\mathbb{B}} \rightarrow \underline{\mathbb{A}}$ instead of $\underline{F}^* \dashv \underline{F}_* : \underline{\mathbb{B}} \rightarrow \underline{\mathbb{A}}$, since the latter provides no additional information.

¹²Neither Morita nor Cauchy were involved with the categorical liftings of the concepts that carry their names. History is often used for mnemonic purposes.