1	On the Role of the Meridional Jet and Horizontal Potential Vorticity Dipole
2	
3	in the Iowa Derecho of 10 August 2020
4	
5	
6	
7	
8	Matthew H. Hitchman and Shellie M. Rowe
9	
10	Department of Atmospheric and Oceanic Sciences
11	University of Wisconsin – Madison
12	
13	
14	
15	
16	
17	
18	
19	
20	Submitted: May 5, 2022
21	Revised: August 3, 2023; November 30, 2023; March 11, 2024
22	
23	Corresponding Author:
24	Shellie M. Rowe
25	rowe1@wisc.edu
26	
27	
28	
29	
30	

Abstract. On 10 August 2020, a derecho caused widespread damage across Iowa and Illinois. Des Moines station data show that the arrival of the gust front was characterized by an abrupt shift to northerly flow, exceeding 22 m/s for ~ 20 min. To test the hypothesis that this northerly jet is associated with a horizontal potential vorticity (PV) dipole in the lower troposphere, we investigated the structure of PV in the University of Wisconsin Nonhydrostratic Modeling System (UWNMS) and of absolute vorticity in High Resolution Rapid Refresh (HRRR) forecast analyses.

This structure is described here for the first time. The negative PV member coincides with the downdraft, while the positive PV member coincides with the updraft, with a northerly jet between. The westerly inflow jet descends anticyclonically in the downdraft, joining with northerly flow from the surface anticyclone. The resulting northerly outflow jet creates the trailing comma-shaped radar echo.

The speed of propagation of the derecho is similar to the westerly wind maximum in the 3-5 km layer associated with the approaching synoptic cyclone, which acts as a steering level for resonant amplification. Idealized diagrams and 3D isosurfaces illustrate the commonality of the PV dipole / northerly jet structure. Differences in this structure among three model states are related to low-level wind shear theory. The PV dipole coincides with the pattern of diabatic stretching tendency, which shifts westward and downward relative to the updraft/downdraft with increasing tilt. The PV dipole can contribute toward dynamical stability in a derecho.

Significance Statement. The purpose of this work is to investigate the structure of potential vorticity (PV) in the lower troposphere in a derecho. It is found that a northerly outflow jet occurs between an east-west oriented horizontal PV dipole, which is described here for the first time. The negative PV member coincides with the downdraft and is inertially unstable, while the positive PV member coincides with the updraft. This work contributes toward the theory of resonant structures and longevity. The 3-5 km westerly inflow layer constitutes a steering level, which controls propagation speed despite differences in structure. The degree of westward tilt with height is related to the pattern of forcing by diabatic stretching in producing the PV dipole.

1. Introduction

Deep convective updrafts and downdrafts play a crucial role in determining the growth and structure of mesoscale convective systems (MCSs). Davies-Jones (1984), Montgomery et al. (2006), Chagnon and Gray (2009), Davis and Galarneau (2009), and others have shown that a horizontal potential vorticity (PV) dipole can be formed by an updraft in shear, where horizontally-oriented vorticity can be bent into a horseshoe shape by an updraft caused by diabatic heating. Hitchman and Rowe (2017, 2019) studied horizontal PV dipoles in the upper troposphere / lower stratosphere (UTLS) over deep convection in tropical and midlatitude cyclones, and found amplitudes of $\sim +/-10 - 20$ PVU, where 1 PVU = 10^{-6} m² K kg⁻¹ s⁻¹. They suggested that UTLS PV dipole formation by the diabatic tilting term can be interpreted in terms of momentum transport by an updraft in shear. Air transported into an environment moving at a different speed and direction, will create a mesoscale jet, and therefore an associated horizontal PV dipole. Since vertical motion is limited by both the ground and the discontinuity in static stability at the tropopause, it was hypothesized that a downdraft in shear might create a mesoscale jet and horizontal PV dipole in the lower troposphere. Since a downdraft in shear is an essential aspect of a derecho, it was decided to investigate the PV structure of a derecho in the lower troposphere.

It is of interest to study PV dipoles since they can have a longer lifetime than the convection that initiates them (Chagnon and Gray 2009, Weijenborg et al. 2017). Emanation of gravity wave activity has been shown to occur simultaneously with the creation of horizontal vortex dipoles (e.g., Snyder et al. 2007, 2009, Muraki and Snyder 2007, Wang et al. 2009, Hernandez-Duenas et al. 2014, Plougonven and Zhang 2014, and Rocha et al. 2018). The formation of horizontal PV dipoles and emission of gravity waves can play a significant role in the dynamical adjustment process by which a derecho evolves into a balanced structure.

The derecho is a particularly damaging and long-lasting type of MCS, with a characteristic westerly inflow layer and downdraft. Corfidi et al. (2016) define a derecho as "a family of damaging downburst clusters associated with a forward-propagating MCS that, during part of its existence, displays evidence of one or more sustained bow echoes with mesoscale vortices and/or rear-inflow jets. The damage swath must be nearly continuous, at least 100 km wide along most of its extent, and 650 km long".

On 10 August 2020, a derecho formed in northeastern Nebraska at \sim 1300 UTC (8 am Central Daylight Time; CDT = UTC – 5 hr) and travelled eastward across Iowa, Illinois, Indiana, and western Ohio. Wind damage over 90,000 square miles resulted in \$11.5B loss in property. Peak wind gusts topped 140 mph in Cedar Rapids, Iowa, and at least 26 tornadoes were spawned. More than 1,000,000 customers lost power, numerous injuries were counted, and four people lost their lives (Chicago Weather Forecast Office 2021).

The Iowa derecho of 10 August 2020 was a high-end event (three or more gusts exceeding 32 m/s), which occur every 2-3 years, in a climatological maximum extending from Minnesota southeastward into Indiana, Ohio and West Virginia (Johns and Hirt 1987, Coniglio and Stensrud 2004, Guastini and Bosart 2016). Summertime configurations favorable to the formation of derechos include an upper-level trough approaching a mid-tropospheric ridge from the west (Coniglio et al. 2004). As will be shown, the synoptic setting for the 10 August 2020 derecho includes an approaching upper-level cyclone, and a lower-tropospheric anticyclone.

To investigate the hypothesis that a downdraft in a derecho is associated with a lower-tropospheric PV dipole, we carried out simulations of the Iowa derecho of 10 August 2020 with the University of Wisconsin Nonhydrostatic Modeling System (UWNMS), which has Ertel's PV as a standard output variable. An idealized diagram of the meridional jet / PV dipole found in UWNMS simulations (Fig. 1) illustrates the basic structure. The westerly inflow jet turns clockwise as it sinks in the downdraft, becoming a northerly jet. The updraft and positive PV anomaly coincide, while the downdraft and negative PV anomaly coincide, with the northerly outflow jet between.




Figure 1. Idealized 3D representation in the lower troposphere of the relationship between a) the westerly inflow jet (yellow), northerly surface outflow jet (green arrow), lower-tropospheric PV dipole (blue/negative and red/positive), downdraft (brown arrow), and updraft (black arrow), and b) perspective view showing the air mass pathways, with horizontal reference planes near the surface and 4 km altitude.

We then investigated the dynamical structure depicted in High-Resolution Rapid Refresh (HRRR) model forecast analyses, which includes absolute vorticity as an output variable. The radar reflectivity pattern of a derecho often evolves from a bow-echo to a comma echo (Przybylinski 1995, Weisman and Trapp 2003, Wakimoto et al. 2006, Atkins and St. Laurent 2009). This paper documents the relationship between the updraft / downdraft pair, lower-tropospheric PV dipole, and northerly jet in the 10 August 2020 derecho, and its relationship to the characteristic equatorward bulge in the trailing comma radar echo pattern.

The development of a rear inflow jet is crucial to the formation of a derecho (Weisman 1992, 1993) and the westward tilt with height of the downdraft / updraft pair (Rotunno et al. 1988, Weisman and Rotunno 2004). The gust front updraft is forced by the cold outflow from the downdraft, where lifting of warm, moist air with high potential temperature by the density current leads to instability in the updraft. In an MCS the degree of westward tilt of the updraft and the shape of the anvil outflow in the UTLS depends on the environmental wind shear profile. Shallow shear tends to promote a strongly-sloping front-to-rear flow with an extensive anvil developing to the west and trailing stratiform precipitation, while deep shear tends to promote an extensive anvil developing to the east and leading stratiform precipitation (Lin, 2007).

The existence of a critical level at which the ambient wind speed, U, is equal to the phase speed of the derecho, c, implies that there is a steering level for the storm (Lin 2007). This also implies that latent heating in the updraft and downdraft can effectively amplify the interaction between the updraft and downdraft. In an idealized modeling study, Weisman and Rotunno (2004) found that, when shear in the lowest 5 km is increased from 10 m s⁻¹/5 km to 20 m s⁻¹/5 km, the updraft becomes more upright, with a stronger forward anvil outflow. At this level of shear there is still significant front-to rear outflow. However, with a low-level shear of 30 m s⁻¹/5 km the front-to-rear flow disappears and the updraft is nearly upright. This suggests that a low-level shear of ~ 20 -25 m s⁻¹/5 km might be optimal for a derecho, fostering longevity through structural stabilization, with positive feedback between forced condensation and hydrometeor cooling. The inflow jet experiences diabatic cooling and drag by hydrometeors in the downdraft (Markowski and Richardson 2010, Seigel and van den Heever 2013), which transports mid-tropospheric air with high momentum to near the surface and creates a cold pool.

Since nearly straight-line motion is an essential aspect of a derecho, most studies have focussed on motions in longitude-altitude. However, Evans et al. (2014) describe a family of

trajectories in a derecho which "descend from the north and northwest". In this study we explore the role of the meridional wind component and its relationship to PV structures, updrafts, downdrafts, and shape of the trailing comma echo.

Lin and Smith (1986) investigated the response to a travelling region of cooling associated with falling hydrometeors in the downdraft of a squall line, which included a westward-tilted updraft/downdraft pair, and source of gravity wave activity. The simultaneous formation of gravity waves and updraft/downdraft pair has been studied by Bretherton and Smolarkiewicz (1989), Pandya et al. (2000), Shige and Satomura (2000), Fovell et al. (2006), Lane and Zhang (2011), and Adams-Selin (2020).

Raymond and Jiang (1990) proposed that an MCS can maintain itself over a long period of time through interaction between the diabatic effects of moist convection and the vertical motion field. They explored the contribution of diabatic heating in producing mesoscale PV anomalies by transporting mass across isentropic surfaces (see also Haynes and McIntyre 1990). Fritsch et al. (1994), Jiang and Raymond (1995), Wernli and Davies (1997), Lin (2007), and Chagnon and Gray (2009) found that diabatic vortex stretching in an updraft will generate a positive PV anomaly below the heating maximum. In a negative buoyancy-driven downdraft, diabatic vortex shrinking will occur below the level of maximum cooling (Lin and Smith 1986), producing a negative PV anomaly in the lower troposphere. In examining PV tendency in a convective situations, other authors have shown that the diabatic tilting term is important when there is significant vertical shear (Chagnon and Gray 2009, Weijenborg et al. 2017, Oertel et al. 2020, Harvey et al. 2020, Prince and Evans 2022).

It is helpful for discussion to introduce the conservation equations for PV and absolute vorticity, since UWNMS output fields include Ertel's PV at constant altitude, while HRRR output fields include absolute vorticity at constant pressure. The vertical component of Ertel's PV is $P = \frac{1}{\rho_o} \frac{\partial \theta}{\partial z} (f + \zeta) - \frac{1}{\rho_o} \frac{\partial \theta}{\partial x} \frac{\partial v}{\partial z} + \frac{1}{\rho_o} \frac{\partial \theta}{\partial y} \frac{\partial u}{\partial z}$, where relative vorticity is $\zeta = \frac{\partial v}{\partial x} - \frac{1}{\cos \phi} \frac{\partial (\cos \phi u)}{\partial y}$,

 θ is potential temperature, f is the Coriolis parameter, ρ_o is the basic state density, $\partial x = a\cos\phi \,\partial\lambda$, and $\partial y = a\partial\phi$ (eqn. 3.1.4 Andrews et al. 1987). Neglecting the net viscous force,

the effects of diabatic stretching and diabating tilting on PV following the motion are given by

178
$$\frac{DP}{Dt} = \frac{1}{\rho_o} \left[(f + \zeta) \frac{\partial Q}{\partial z} + \frac{\partial Q}{\partial y} \frac{\partial u}{\partial z} - \frac{\partial Q}{\partial x} \frac{\partial v}{\partial z} \right], \quad (1)$$

(derived from eqn. 3.1.5 in Andrews et al. 1987). Although *Q* is not available as an output field for the UWNMS, we can use scale analysis to estimate the likely relevance of the diabatic stretching and tilting terms. Neglecting friction and solenoidal terms, the tendency equation for absolute vorticity following the motion is

183
$$\frac{d}{dt}(\zeta + f) = (\zeta + f)\frac{\partial w}{\partial z} + \frac{\partial w}{\partial y}\frac{\partial u}{\partial z} - \frac{\partial w}{\partial x}\frac{\partial v}{\partial z}$$
 (2)

- 184 (Holton and Hakim 2013). Since $w \frac{\partial \theta}{\partial z} \approx Q$ in convection, similar scaling arguments can be made for PV and absolute vorticity.
- Considering the diabatic stretching term $\frac{1}{\rho_o}(f+\zeta)\frac{\partial Q}{\partial z}$, below a heating maximum in an
- updraft, $\frac{\partial Q}{\partial z} > 0$ will induce a region of P > 0. Below a cooling maximum in a downdraft, $\frac{\partial Q}{\partial z} < 0$
- 188 0 will induce a region of P < 0 (Lin and Smith 1986). Typical magnitudes may be estimated
- near 700 hPa, where $\rho \sim$ 0.8 kg m⁻³. If we assume that $f + \zeta \sim$ 10⁻⁴ s⁻¹, $Q \sim$ +/- 10 K/hr, and $\delta z \sim$
- 3 km, the resulting PV tendency would be \sim +/- 0.4 PVU/hr.
- To estimate the diabatic tilting term $-\frac{1}{\rho_0} \frac{\partial Q}{\partial x} \frac{\partial v}{\partial z}$ near 700 hPa, if $Q \sim 10$ K/hr, $\delta x \sim 50$ km,
- and $\frac{\partial v}{\partial z} \sim 20 \text{ m s}^{-1} / 5 \text{ km}$, and if the two gradient fields are in phase, the diabatic tilting tendency
- would be \sim +/- 1 PVU per hour. Near the tropopause (150 hPa), $\rho \sim 0.2$ kg m⁻³. If Q \sim 1 K/hr,
- 194 $\delta x \sim 50$ km, and $\frac{\partial v}{\partial z} \sim 30$ m s⁻¹ / 2 km in the UTLS, the diabatic tilting tendency would be $\sim +/-1.5$
- 195 PVU per hour. Once PV anomalies are formed by diabatic stretching and diabatic tilting they
- can be advected upward or downward along isentropes.
- Evans et al. (2014) carried out a PV analysis for the formation of the cyclonic meso-betavortex at the north end of the bow echo in the Super Derecho of 8 May 2009 (Weisman et al. 2013). Xu et al. (2015) studied the genesis of mesovortices in the bow echo system of this derecho using a model with a 4 km outer grid and 0.8 km inner grid. They found a long-lived
- derecho using a model with a 4 km outer grid and 0.8 km inner grid. They found a long-live cyclonic meso-vortex near the cusp of the bow echo which is somewhat similar to the east
- 202 (cyclonic) half of the PV dipole to be described below. In the present study, an east-west
- oriented PV dipole is diagnosed, with a northerly jet between. To our knowledge, this structure
- 204 has not been previously described in the literature.
- The observations, UWNMS simulations, and HRRR analyses provide complementary information. The UWNMS, with explicit moist convection, includes Ertel's PV, equivalent

potential temperature, θ_e , and streamfunction as output variables, and evolves without updating information in the interior. HRRR ingests radar and satellite data in a rapid update cycle, and includes absolute vorticity as an output variable. This provides the opportunity for comparison of the location, rate of progression of the derecho, and dynamical structure.

Can a derecho can be produced in a model internally from "first principles" and a suitable initial state, or does one need to ingest detailed observations during integration to achieve a reasonable derecho structure? Requisite elements include a sufficiently moist environment, forcing which provides uplifting, and environmental shear which helps to organize the storm. We investigated this by running the UWNMS using relatively coarse initialization data and updating only on the side boundaries. The mesoscale detail that evolves in the UWNMS is a dynamically distinct solution relative to HRRR, which incorporates high resolution 3D information into the interior of the grid throughout integration. If similar PV structures are found in the HRRR and UWNMS, this would support the robustness of the result.

The observational data sets, HRRR analyses, and UWNMS simulations used in this study are described in section 2. A depiction of the derecho as seen in ground-based and satellite observations is given in section 3, including the temporal signature of the northerly jet at Des Moines, and the shape of the outflow cirrus. HRRR synoptic-scale wind analyses at tropospheric pressure levels show the relationship between the approaching upper-level cyclone, westerly inflow jet, and surface anticyclone. The evolution of three simulations is described in section 4. The PV dipole / meridional jet structure, relationship with updraft / downdraft pair and westerly inflow jet are shown for UWNMS simulations in section 5 and for HRRR in section 6. Dynamical aspects discussed in section 7 include the role of the inflow jet in determining the speed and strength of the derecho, critical level theory, and the generation mechanism for the lower-tropospheric PV dipole. Results are summarized in section 8.

2. Observations, HRRR analysis, and UWNMS simulations

a) Radiosonde, surface, radar, and satellite observations

Hourly radar plots were used to estimate the rate of eastward progression of the gust front. A surface station map was generated for 1600 UTC (11 am CDT) on 10 August 2020, when the gust front reached Des Moines, IA. A special sounding was taken at Davenport, IA (DVN) at 1700 UTC (noon CDT), about 2 hours before the arrival of the gust front.

238 Surface observations at 1-min interval were obtained for Des Moines, IA (DSM) for the 239 12-hr period 1200 – 2400 UTC (7 am – 7 pm CDT) on 10 August 2020. Both 1-min average 240 wind speed ("wind speed") and "gust speed" were obtained, along with 1-min average wind 241 direction. The National Weather Service defines a 2-min average gust speed as "the values of 242 the 3-sec peak from the 5-sec sample containing the highest 3-sec peak wind speed". The Iowa 243 Mesonet provided a 1-min average gust speed. 244 Lightning strikes detected by the GOES-16 Geostationary Lightning Mapper (GLM) 245 during a 20 second period were superimposed on GOES-16 ABI L2 images of cloud top 246 temperature and height at \sim 1800 UTC (\sim 1 pm CDT). 247 b) HRRR forecast analyses 248 249 High-Resolution Rapid Refresh (HRRR) model forecast analyses were obtained from NOAA. The HRRR model has 3 km grid spacing, with radar data assimilated every 15 minutes 250 251 over a 1-hr period (Dowell et al. 2022). The HRRRv4 CONUS domain, with 1800 x 1060 grid 252 points, spans the contiguous U.S. We analyzed the HRRR forecast starting at 0000 UTC 10 253 August 2020. Three-dimensional isosurfaces were created using MATLAB. 254 255 c) The University of Wisconsin Nonhydrostatic Modeling System (UWNMS) The UWNMS (Tripoli 1992a, b) was derived from the CSU model of Tripoli and Cotton 256 257 (1982) and was designed to study the interaction between convection and its environment. We 258 have used it to study inertial instability and midlatitude jets (Rowe and Hitchman 2015, 2016), 259 UTLS PV dipoles (Hitchman and Rowe 2017, 2019), summertime flooding (Rowe and Hitchman 260 2020), and tropopause folds (Hitchman and Rowe 2021). 261 For the derecho simulations, the UWNMS was initialized with 0.5° data from the 262 National Centers for Environmental Prediction (NCEP) Global Forecast System (GFS) at 0000 263 UTC 10 August and run for 24 hours, updating boundary conditions every 15 minutes (Table 1). 264 265 266 267 268

Model	UWNMS version 8i
Time	0000 UTC 10 August 2020 initialization, run 24 hr, output every 20 min
Grid	Domain 1200 km x 1200 km, spherical grid center point 41°N, 91°W 300 m vertical grid spacing, 95 vertical levels, 5 soil grid points 26.7 km model top with 1.5 km Rayleigh friction sponge layer 85 vertical grid points in visualized domain (25.2 km top)
Physics	explicit convection, turbulent kinetic energy prediction Hashino and Tripoli (2007, 2011) microphysics lateral and upper boundaries - Hack/Schubert normal mode method
Initialization	0.5° NCEP GFS data, boundary values updated every 15 min
Runs	4 km horizontal grid spacing (UWNMS-4) 8 km horizontal grid spacing (UWNMS-8)

Table 1. Specifications for UWNMS simulations.

We used a domain of 1200 x 1200 km centered in eastern Iowa (41°N, 91°W), covering 36°N-46°N and 84°W-98°W. Simulations were carried out with 4, 6, 8, 10, and 12 km grid spacing, keeping all other parameters constant. The coherence of a derecho-like structure generally improved with smaller grid spacing (not shown). Each simulation exhibited a different evolution history, with significant variation in amplitude and structure with time within each simulation. In section 5 results are shown for simulations with grid spacings of 4 km (UWNMS-4) and 8 km (UWNMS-8).

3. Observational depiction of the Iowa Derecho of 10 August 2020

a) Evolution of the gust front

An hourly sequence of radar reflectivity images during 1250 – 2350 UTC 10 August 2020 (Fig. 2) shows that the derecho reached Des Moines at 1600 UTC (Fig. 2e) and Davenport, IA at 1900 UTC (Fig. 2h). Gusts in the range 25-40 m/s were observed across Iowa during 1500 - 1900 UTC (Iowa Environmental Mesonet). This derecho developed a comma echo, with an initially linear convective cluster extending to the northeast (Fig. 2). An equatorward bulge of the echo pattern developed as the derecho propagated into Illinois and Indiana (Figs. 2h-m). The leading edge of the comma traveled toward the east, but the surface outflow behind the gust front was northerly.




Figure 2. a) Progression of the arc of high reflectivity from 1250 UTC 10 August to 0050 UTC 11 August 2020 (7:50 am to 8:50 pm CDT 10 August). Panels b) - m) show hourly snapshots from 1250 to 2350 UTC. The locations of Des Moines, IA (DSM), Davenport, IA (DVN), and Lincoln, IL (ILX) are indicated in panel a). Values of dBZ less than zero are not plotted.

Campbell et al. (2017) used "the approximate center of the leading edge of the feature of interest" to define derecho speed. Here we track the easternmost point of the echo pattern (Fig. 2a). The eastward speed was ~ 18 m/s at 1400 UTC, increasing to ~ 30 m/s at 1900 UTC, then decreasing to ~ 21 m/s by 2300 UTC, with a mean of 24.5 m/s (Fig. 3a).




Figure 3. a) Speed of propagation during 1400 UTC 10 August - 0000 UTC 11 August (9 am - 7 pm CDT 10 August) 2020. Approximate arrival times at DSM, DVN, and ILX are indicated. b) Surface station plot for 1600 UTC (11 am CDT) 10 August 2020. The position of the cold front is shown. Dew point temperatures exceeding 75°F are shaded light green, less than 65°F light brown. The locations of Iowa (IA), Wisconsin (WI), and Illinois (IL) are indicated, along with stations DSM, DVN, and ILX (Lincoln, IL). c) National Weather Service special sounding from Davenport, IA (DVN) at 1700 UTC (Noon CDT) on 10 August 10 (about 2 hours before the

arrival of the derecho). Profiles of temperature (°C, dark blue), dew point temperature (°C, light blue), and horizontal wind vectors (knots) are shown. The level of free convection (LFC), equilibrium level (EL) and CAPE are indicated.

Mahoney et al. (2009) compared the speed of propagation of MCSs with maximum zonal wind speed in the cold pool and mean 900-200 hPa wind speed. They found speeds of 10-22 m/s, similar to the maximum zonal wind speed in the cold pool. Corfidi et al. (1996) defined the advective component of an MCS to be proportional to the mean wind in the 850-300 hPa layer. In section 4 we will compare the speed of the gust front with the speed of the 850-700 hPa westerly inflow jet. The 850-700 hPa (~ 3-5 km) layer was chosen because it is the layer of maximum westerly flow in this case. It can be related to critical surface theory for low-level shear, where the outflow pattern in the upper troposphere depends on how low-level shear influences storm structure.

b) Surface station chart

At 1600 UTC 10 August 2020 a cold front extended from Kansas to Wisconsin (Fig. 3b). Dew point temperatures, T_d , exceeded 75°F (24°C) over Iowa and most of Illinois, with values near 64°F (18°C) along the cold front, and less than 60°F (16°C) to the north and west of Iowa (Fig. 3b).

c) Davenport, IA sounding at 1700 UTC

In the 1700 UTC DVN sounding (Fig. 3c) the level of free convection was \sim 2200 m and equilibrium level \sim 14,760 m, for a moderate CAPE value of \sim 1700 J/kg. However, Evans and Doswell (2001) found that when synoptic-scale forcing is strong, derechos can develop and persist within environments with almost no CAPE evident on nearby soundings. From the surface to 3 km, wind speed increased upward by \sim 15 m/s (wind shear \sim 5x10⁻³ s⁻¹), shifting from southwesterly to westerly. Dry air with southwesterly flow is seen in the 6.5-8 km layer, above which the flow becomes increasingly westerly, reaching \sim 35 m/s near the tropopause (\sim 12 km). In the 300-100 hPa layer there are alternating layers of stronger and weaker westerlies (Fig. 3c).

d) Satellite cloud-top images

At 1800 UTC the outflow gust front was located near 91.5°W, about an hour west of Davenport, IA (Fig. 2g). A region of cloud top temperatures less than 210 K (Fig. 4a) and cloud top heights exceeding 16 km (Fig. 4b) is seen over east-central Iowa, with coincident lightning (Figs. 4a, b), indicating the presence of deep convection. The cirrus outflow boundary coincides approximately with the 240 K contour. The cirrus shield extends westward over Iowa, with the trailing stratiform region overlying the mid-level inflow jet (Smull and Houze 1987, Adams-Selin and Johnson 2013), as well as northward over Minnesota, and northeastward over Wisconsin. The shape of the cirrus shield is consistent with a vertical wind profile which allows for significant outflow both upstream and downstream in the UTLS.

In HRRR similar features are indicated, but the region of lowest cloud top pressure (Fig. 4c) and highest cloud top echo height (Fig. 4d) is located ~ 80 km to the west and ~ 25 km to the south. In addition, the line of convective activity extending eastward over northern Illinois in satellite observations (Figs. 4a, c) and radar reflectivity (Fig. 2g) instead extends northeastward across Wisconsin in the HRRR analyses (Figs. 4b, d).




Figure 4. a) GOES-16 ABI-L2 cloud top temperature (K), b) GOES-16 ABI-L2 cloud top altitude (km), c) HRRR cloud top pressure (hPa), and d) HRRR cloud top echo altitude (km) at 1800 UTC 10 August 2020. The locations of lightning strikes detected by the GOES-16 GLM mapper during the 20 second period 17:59:40 – 18:00:00 UTC are shown with black dots.

The coldest cloud top satellite temperatures are less than 200 K and the highest altitudes occur at 18.0 km ($\sim 55,000'$) (Figs. 4a, b). In HRRR the lowest cloud-top pressure is 90 hPa and highest cloud top echo heights are 15.3 km (Figs. 4c, d).

e) Surface observation time series at Des Moines

During 1200-1530 UTC maximum gusts at Des Moines were ~ 5 m/s or less (Figs. 5a, b), with the surface wind direction shifting gradually from southerly to westerly (Fig. 5b). During 1400-1530 UTC surface pressure rose gradually, with a series of oscillations of amplitude ~ 0.5 hPa and period ~ 15 -30 min (Fig. 5a).

Around 1600 UTC gust speed at DSM increased from westerly at ~ 5 m/s to northerly at ~ 33 m/s in less than 10 min (Fig. 5b). Typical gust speeds are $\sim 30\text{-}50\%$ higher than 1-min average wind speeds (not shown). After 1600 UTC the wind speed decreased exponentially to ~ 12 m/s in an hour (Fig. 5a). Northerly flow exceeded 22 m/s for ~ 20 minutes. With an eastward translation speed of 28 m/s for the derecho past Des Moines, this duration implies a ~ 35 km zonal width for the meridional jet exceeding 22 m/s. The duration of northerly flow exceeding 15 m/s implies a zonal width of ~ 90 km. During 1600 - 1800 UTC the wind direction shifted gradually from north-northwesterly ($\sim 345^{\circ}$) to northeasterly ($\sim 60^{\circ}$), held steady during 1800 - 1930 UTC, then abruptly switched to southerly upon arrival of the west end of the cold pool (Figs. 5b, c). Over the subsequent 5 hr, winds gradually shifted to northwesterly (Fig. 5b).




Figure 5. Iowa Environmental Mesonet surface meteorological observations at Des Moines, Iowa (DSM) during 1200 – 2400 UTC (7 am – 7 pm CDT) 10 August 2020 of a) surface pressure (red line) and gust speed (black), b) gust speed (black) and wind direction (blue), and c) temperature (yellow) and dew point temperature (green). Data are shown at 1-min intervals. Vertical dashed lines indicate the duration of the cold pool at DSM.

An increase in surface pressure of ~ 5 hPa occurred over ~ 15 min (~ 0.5 Pa/s) near 1600 UTC (Fig. 5a). The peak in pressure occurred ~ 5 min after the peak in wind speed. This is consistent with an outflow boundary "pressure head", with a sharp eastward decrease in pressure accelerating the flow ahead of the pressure maximum (e.g., Fig. 6.8 of Lin 2007). The period of elevated pressure lasted ~ 2.5 hours (Fig. 5a). The 3.5-hr period 1600 – 1930 UTC is characterized by northerly surface flow (Fig. 5b).

During 1200 –1530 UTC the temperature rose from 73°F to 81°F (23°C to 27°C), while T_d held steady near 71°F (22°C) (Fig. 5c). With the arrival of the gust front at 1600 UTC,

temperatures dropped to \sim 62°F (17°C) in \sim 20 minutes, and T_d dropped to \sim 60°F (16°C). In the wake of the initial outflow boundary and downburst, temperatures gradually returned to 79°F (26°C) over the next 7 hours (Fig. 5c). The dew point stayed low during the \sim 3.5 hr duration of the cold pool, then rose slightly, but remained below 65°F (18°C).

This sharp northerly jet (Fig. 5b) implies the existence of a relative vorticity dipole in the lower troposphere, with a positive leading anomaly to the east and a negative anomaly to the west (Fig. 1).

f) HRRR wind charts

At 1800 UTC on 10 August 2020 a synoptic-scale trough is seen at 150 hPa over the upper Midwest, with southwesterly flow over eastern Iowa and Wisconsin (Fig. 6a). This flow is interrupted by a mesoscale region of low-speed air at 42°N, 92.5°W (Fig. 6a). At 300 hPa, stronger winds are evident in the trough and the mesoscale region of low-speed air is larger, with noticeable outflow toward the north (Fig. 6b).




Figure 6. HRRR horizontal wind speed (color bar, interval 2.5 m/s) and wind vectors (reference length 10 m/s) at 1800 UTC on 10 August 2020 for a) 150 hPa, b) 300 hPa, c) 500 hPa, d) 700 hPa, e) 850 hPa, and f) 950 hPa.

The wind pattern at 500 hPa shows an approaching jet streak, extending from the northwest into western Iowa, in a meridionally-concentrated band about ~ 200 km wide (Fig. 6c). A minimum in wind speed is again seen near 42°N, 92.5°W (Fig. 6c). At 700 hPa, flow converges into the westerly inflow jet, exceeding 25 m/s over central Iowa (Fig. 6d). An anticyclonic flow pattern is found to the west at 850 hPa (Fig. 6e), with northerly flow entering northwestern Iowa. Note also the mesoscale region of westerlies at 850 hPa exceeding 35 m/s over eastern Iowa (Fig. 6e). At 950 hPa, continuous northerly flow extends from Minnesota to southern Iowa (Fig. 6f), with a comma-shaped speed maximum consistent with the abrupt switch to northerly flow seen at Des Moines (Figs. 5b, 6f). From 700 to 950 hPa the speed maximum turns from westerly to northerly (Figs. 6d-f). The mesoscale northerly jet at 950 hPa is continuous with the northerly ouflow from the anticyclone to the north (Fig. 6f).

4. Comparison of storm evolution

The evolution of hourly samples in updraft speed during the life of the derecho for UWNMS-4 and UWNMS-8 at 3 km altitude, and for HRRR at 700 hPa, are shown in Fig. 7a. In HRRR analyses, starting at 1400 UTC, updraft speed increased from ~ 0.3 m/s to ~ 3.5 m/s by 1700 UTC, then gradually declined to ~ 1 m/s at 2000 UTC, with a second peak of ~ 3.5 m/s at 2100 UTC (Fig. 7a). In the UWNMS-4 simulation, updraft speed was negligible at first, increasing from ~ 0 to ~ 3 m/s during 1500-1700 UTC. It then generated an unrealistically strong updraft (> 14 m/s) at 1800 UTC, followed by a pronounced minimum in speed at 1900 UTC, then a resurgence to ~ 8 m/s at 2000 UTC (Fig. 7a). In the UWNMS-8 simulation, development of the updraft was delayed by 2-3 hr, finally strengthening to 1 m/s at 1800 UTC, then ~ 4 m/s at 1900 UTC, and remaining strong for four hours (Fig. 7a).




Figure 7. Hourly time series during the period 1400 - 2300 UTC 10 August 2020 of a) maximum updraft velocity (m/s) along 42°N at 700 hPa for HRRR (thick black), and at 3.18 km altitude for UWNMS-4 (dashed blue), and UWNMS-8 (blue) simulations and b) radar gust front speed from Fig. 3a (m/s, thick blue), HRRR zonal wind speed 100 km upstream of the gust front averaged in the layer 500-700 hPa (black), gust front speed estimated from HRRR surface wind pattern (dashed black), and UWNMS-8 zonal wind speed at 4.5 km altitude 50 km upstream and 50 km poleward of the gust front (blue).

The average eastward speed of the gust front was observed to be 24.5 m/s during 1400 – 2300 UTC and 28.0 m/s during 1700 – 2100 UTC (Figs. 3a, 7b). Hourly sections of zonal wind

along 42°N in HRRR (Fig. S1) were used to make a separate estimate of the speed of propagation, which is ~ 5 m/s less than deduced from radar echos (Fig. 7b). The average speed of the westerly inflow layer in the 750-500 hPa layer 100 km west of the gust front in HRRR grows and then weakens, similar to gust front speed, but it is ~ 5 -10 m/s slower (Fig. 7b). The westerly wind in UWNMS-8 at 4.5 km altitude sampled 50 km upstream and 50 km poleward of the gust front shows better agreement with the observed speed of the gust front (Fig. 7b). These results may be compared with Fig. 7 of Mahoney et al. (2007).

The evolution history in each of the three simulations was quite distinct. The UWNMS-4 simulation initially showed similar development to HRRR, but then underwent two large amplification/diminution cycles. The UWNMS-8 simulation developed too slowly, but then achieved an updraft strength comparable to HRRR, with a coherent structure lasting for ~ 5 hr.

5. Depiction of structure in UWNMS simulations

The center of the updraft/downdraft pair in both UWNMS simulations propagated nearly eastward along 42°N, with the timing of primary dynamical features similar to the gust front seen in radar reflectivity (Fig. 2) and Des Moines surface data (Fig. 5). The two simulations differ in the timing of amplification and in the degree of westward tilt with height.

During 1800 – 2000 UTC both UWNMS simulations show an east-west PV dipole, with the leading positive PV member of the dipole coinciding with the updraft and the trailing negative PV member coinciding with the downdraft (Fig. 8). During 1800 – 2000 UTC there is a transition in the UWNMS-4 from a strong updraft burst subsiding into a dispersed set of mesoscale convective vortices (MCVs; Figs. 7a, 8a-c), but in the UWNMS-8 a derecho-like structure grows to moderate, steady strength (Figs. 7a, 8d-f).

The robustness of the lower-tropospheric PV dipole structure is dependent on a sufficiently strong and vertically-continuous updraft / downdraft pair. For example, at 1800 UTC a PV dipole in UWNMS-4 is apparent (Fig. 8a), with an updraft speed exceeding 14 m/s (Fig. 7a), but at 1900 and 2000 UTC there is instead a set of smaller-scale MCVs arranged along the outflow boundary (Figs. 8b, c), with updraft speeds of less than 2 m/s at 1900 UTC (Fig. 7a).




Figure 8. Plan views of PV (color bar) and vertical velocity (m/s, contours, negative dashed) at 3 km altitude for UWNMS-4 (left) and UWNMS-8 (right) for a), d) 1800 UTC, b), e) 1900 UTC, and c), f) 2000 UTC 10 August 2020. In UWNMS-4 vertical velocity is contoured every 2 m/s and PV every 2 PVU, while in UWNMS-8 vertical velocity is contoured every 0.8 m/s and PV every 0.5 PVU.

In contrast, UWNMS-8 showed a steady derecho-like structure during 1900-2100 UTC (Figs. 8d-f), with an updraft speed of ~ 5 m/s (Fig. 7a) and a lower-tropospheric PV dipole coincident with the downdraft/updraft pair. In UWNMS-8, during 1800-2000 UTC the negative PV anomaly grew from ~ -1 PVU to -2 PVU while the downdraft grew from ~ -1 to -2 m/s, and the positive PV anomaly grew from ~ 0.5 PVU to 2 PVU while the updraft grew from ~ 0.6 to 4 m/s (Figs. 8 d-f). It will be shown that these values of vertical motion and PV anomalies are similar to those in HRRR, with similar horizontal scales of ~ 50 km.

In order to illustrate the range in structure of PV and 3D winds in the UWNMS, and for comparison with HRRR structure and with shear theory, we chose a high amplitude state of the

UWNMS-4 at 1800 UTC (Fig. 9) and a representative state of the UWNMS-8 at 1900 UTC (Figs. 10-12).

Longitude-altitude sections of UWNMS-4 variables along 42°N (Fig. 9) are shown with an aspect ratio of 2:1 in the domain 0-21 km altitude, 95.0°W - 90.5°W, to facilitate comparison with HRRR. The location of this section was chosen to fit the locus of updraft/downdraft and PV dipoles in the UWNMS (inset, Fig. 9b). The meridional deviation from 42°N of the track of the center of the PV dipole was less than 5 km as it progressed across the domain.

The location of the gust front can be determined from where the northerly flow maximum abruptly increases, near 91.5°W at 1800 UTC (Fig. 9b). A westerly inflow jet exceeding 25 m/s is seen in the 4-6 km layer about 50 km to the west of the gust front. This pattern extends down to near the surface, with zonal winds exceeding 30 m/s (Fig. 9a) and meridional winds exceeding -25 m/s (Fig. 9b). This descent of the westerly inflow layer in UWNMS-4 occurs fairly abruptly and the updraft is nearly upright (Fig. 9c). The surface westerly and northerly jet maxima coincide with the downdraft (Figs. 9a-c). The northerly jet extends upward to 10 km altitude. It lies between a PV dipole of -10 / + 20 PVU maximizing near 7 km (Figs. 8b, d).

The updraft in UWNMS-4 exceeds 25 m/s and reaches 14 km altitude (Fig. 9c). The westerly inflow jet, with $\theta_e < 330$ K (Figs. 9a, c), splits near 8 km altitude, with a weak rising branch joining the main updraft (Fig. 9c). In the UTLS, a westward outflow layer is found near 12 km altitude and a strong eastward outflow layer exceeding 70 m/s is found near 14 km (Fig. 9a). The zonal flow in the updraft is markedly easterly (Fig. 9a), with flow reaching -20 to -30 m/s in the upper troposphere, extending to 16 km and coinciding with southerly flow (Fig. 9b).

An east-west PV dipole is seen in the 14-16 km layer at the top of the updraft (Fig. 9d). The wavy pattern in potential temperature and PV which tilts westward with height into the stratosphere (Figs. 9c, d) represents upward propagation of gravity wave activity and downward transport of westerly momentum.

The updraft/downdraft pair in UWNMS-4 at 1800 UTC exhibits weak westward tilt with height (Fig. 9c), consistent with a weak and variable cold pool below 800 hPa near 92°-94°W (Fig. 9c). The influence of convection can be seen up to ~18 km (Fig. 9), which is a higher altitude than the maximum HRRR cloud top echo height (Fig. 4d), but is similar to satellite observations (Fig. 4b).




Figure 9. Longitude-altitude sections in UWNMS-4 at 1800 UTC 10 August 2020 for a) zonal wind (contour interval 5 m/s, negative dashed), b) meridional wind (contour interval 5 m/s, negative dashed), c) equivalent potential temperature (blue contours, interval 10 K) and vertical wind (contour interval 3 m/s, negative dashed), and d) PV (contour interval 10 PVU, negative dashed). The vertical line in each panel indicates the longitude of maximum northerly flow in b). The location of this section is shown in the inset of panel b).

In UWNMS-8, the updraft/downdraft pair at 1900 UTC exhibits a stronger westward tilt with height (Fig. 10). At 1900 UTC the gust front was located near 90.5°W (Figs. 10a, b). A westerly inflow jet of ~ 25 m/s is seen in the 4-6 km layer about 50 km to the west of the gust front, which extends continuously down to near the surface, and coincides with a northerly jet exceeding -18 m/s near 2 km altitude (Fig. 10b). The surface westerly and northerly jet maxima occur in the vicinity of the downdraft (Figs. 10a-c).

In UWNMS-8, the northerly jet lies between a PV dipole of -1 / + 4 PVU which maximizes near 6 km (Figs. 10b, d). The axis of maximum northerly wind (Fig. 10b) lies between the updraft/downdraft pair (Fig. 10c) and between the PV dipole (Fig. 10d), separating the westerly inflow layer from the westward-tilting updraft (Fig. 10a).

In UWNMS-8 at 1900 UTC the northerly jet extends upward to \sim 7 km and the updraft extends to \sim 12 km altitude (Fig. 10c), exhibiting less vertical development compared with

UWNMS-4 at 1800 UTC (Fig. 9c). The updraft (Fig. 10c), axis of maximum northerly flow (Fig. 10b), separation between strong and weak westerlies (Fig. 10a), and the PV dipole all tilt westward with height by \sim - 30 km in 10 km altitude, with an elevation angle of \sim 17° (Fig. 10a), It will be shown that this is comparable to the tilt of the updraft in HRRR (\sim 15°), but the elevation angle of the downdraft is much smaller in HRRR (\sim 1.7°).




Figure 10. Longitude-altitude sections in UWNMS-8 at 1900 UTC 10 August 2020 for a) zonal wind (contour interval 3 m/s, negative dashed), b) meridional wind (contour interval 3 m/s, negative dashed), c) equivalent potential temperature (blue contours, interval 10 K) and vertical wind (contour interval 2 m/s, negative dashed), and d) PV (contour interval 1 PVU, negative dashed). The tilted vertical line in each panel indicates the axis of maximum northerly wind.

The horizontal extent of the updraft and downdraft in UWNMS-8 near 3 km at 1900 UTC (Fig. 10c) is \sim 30 km, similar to scales depicted in Smull and Houze (1987). In UWNMS-8 a cold pool with $\theta_e < 340$ K is seen to the west of the gust front (91.2-90.8°W) below 800 hPa, which is somewhat more realistic than in UWNMS-4 (Fig. 9c). Note also values of $\theta_e > 350$ K in the boundary layer ahead of the gust front (Fig. 10c).

In UWNMS-8 at 1900 UTC an easterly front-to-rear outflow layer occurs near 9 km altitude, with a westerly outflow layer exceeding 70 m/s near 12 km (Fig. 10a). The easterly

outflow layer occurs at a lower altitude than in the UWNMS-4 at 1800 UTC. The zonal flow in the updraft (Fig. 10a) is much less easterly than in UWNMS-4 (Fig. 9a), reaching -3 m/s near 12 km (Fig. 10a). The axis of northerly flow in the lower troposphere lies between the westerly flow in the downdraft and weak westerly flow in the updraft (Figs. 10a-c), and between the PV dipole (Fig. 10d). The negative PV anomaly coincides with the downdraft and the positive PV anomaly coincides with the updraft (Figs. 10c, d). Over the next two hours, the downdraft/updraft pair and PV dipole remained coherent in the UWNMS-8. Eastward movement of the PV dipole / northerly jet can explain the development of the trailing equatorward bulge of the gust front into southern Illinois (Fig. 2).

UWNMS-8 exhibited a development phase (1700 and 1800 UTC), mature phase (1900 - 2000 UTC), and decaying phase (2100 UTC) (Fig. 7a). This evolution is similar to the observed changes in the speed of propagation of the derecho, which was fastest during 1800-2100 UTC (Fig. 7a). The basic structure of a westerly mid-level inflow jet of ~ 25 m/s and a surface westerly maximum coinciding with the leading edge of the downdraft supports the idea that the speed of the westerly inflow jet is fundamentally related to the rate of eastward progression of the derecho.

The existence of a northerly jet is in agreement with the observations at DSM (Fig. 5b). Since the the gust front arrived at DVN (90.58°W) at \sim 1900 UTC, we explore wind structure in the meridional plane at 90.58°W at 1900 and 1920 UTC (Fig. 11). A region of near-surface westerlies, \sim 50 km in meridional extent, is seen near 42°N, coinciding with the western half of the downdraft (Fig. 11a). This layer extends upward and poleward and is continuous with a layer of westerly flow to the north centered at \sim 6 km and exceeding \sim 20 m/s (Fig. 11a). A northerly jet maximizes at -18 m/s near 2 km and coincides with the center and eastern half of the downdraft (Fig. 11b). This region of northerly flow extends northward and upward from the gust front (Fig. 11b), lying between the westerlies to the north and weak zonal flow in the poleward-tilted updraft (Figs. 11a). In the UTLS, outflow jets extend primarily northward near 12 km from the updraft, and also southward at 14 km (Fig. 11c).

Twenty minutes later, the region of southward flow exhibited a similar pattern (Figs. 11c, d), but the region of zonal flow exceeding 20 m/s has filled in the whole 1-5 km layer immediately behind the gust front (Fig. 11b). The axis of northerly flow associated with the gust front tilts northward with height (Fig. 11b, d). It appears that air in the surface northerly jet

is a blending of air streams associated with the cyclonic member, the anticyclonic member, and the surface anticyclone to the north (Fig. 6f). Note the offset in altitude for outflow to the north (~12 km) and to the south (~14 km) (Fig. 11d).




Figure 11. Meridional sections along 90.58°W of UWNMS-8 (left) zonal wind and (right) meridional wind (black contours every 3 m/s, negative dashed), with vertical wind (blue contours every 2 m/s, negative dashed) at (top) 1900 and (bottom) 1920 UTC on 10 August 2020. The view is toward the east, with north to the left. The locations of the section ends (dots) are shown in the inset.

Perspective views of speed isosurfaces show that the westerly inflow layer descends anticyclonically into the northerly outflow jet (Figs. 12a, b). The location of the northerly jet between the PV dipole is shown in Fig. 12c. Since the negative PV dipole member is inertially unstable, this can contribute toward acceleration of the northerly flow.




Figure 12. Spatial relationship between the westerly inflow jet, lower-tropospheric northerly jet, and the downdraft in UWNMS-8 at 1900 UTC 10 August 2020. Perspective views from a) the southwest and b) south showing isosurfaces of zonal wind (U, 22 m/s, yellow), meridional wind (V, -5 m/s, green), and vertical wind (W, -0.8 m/s, orange), and c) from the southwest showing an isosurface of meridional wind (V, -5 m/s, green), together with the -1 PVU (blue) and +2 PVU (red) isosurfaces. Selected altitudes are indicated in b) and c).

6. Depiction of derecho in HRRR analyses

a) Evolution of absolute vorticity at 150 hPa and 700 hPa

During 8-10 August 2020 an upper tropospheric ridge over the southeastern U.S. gradually weakened as a sequence of eastward-travelling troughs approached from the Gulf of Alaska (not shown). These troughs interacted with a ridge over the North Pacific in anticyclonic Rossby wave breaking events, producing a filament of stratospheric air with high 150 hPa absolute vorticity extending southwestward across Colorado at 1600 UTC 10 August (Fig. 13a). Interrupting this filament of high absolute vorticity is a mesoscale absolute vorticity dipole, with positive member lying to the east, and amplitude +/- $10x10^{-4}$ s⁻¹ (Fig. 13a). Since isentropic density $\sigma = \rho \frac{\partial z}{\partial \theta} \sim 460$ kg K⁻¹ m⁻² near 150 hPa, from values in Fig. 3c, this corresponds to a PV dipole of amplitude \sim +/- 2 PVU. This UTLS PV dipole is observed to progress eastward with the derecho over the next three hours (Figs. 2, 13a-d).

A similar eastward-progressing absolute vorticity dipole is seen at 700 hPa, with the positive member lying to the east (Figs. 13e-g). The dipole at 150 hPa lies \sim 50 km poleward of the dipole at 700 hPa, indicating a poleward tilt of the updraft. The zonal scale of each anomaly at 700 hPa is \sim 50 km, with magnitude \sim +/- $10x10^{-4}$ s⁻¹. Since $\sigma \sim 200$ kg K⁻¹ m⁻² near 700 hPa, this corresponds to an east-west PV dipole with strength \sim +/- 5 PVU.




Figure 13. HRRR absolute vorticity (color bar) at 150 hPa for a) 1600 UTC, b) 1700 UTC, c) 1800 UTC, and d) 1900 UTC 10 August 2020, and at 700 hPa for e) 1700 UTC, f) 1800 UTC, and g) 1900 UTC. The color bar interval is 0.5×10^{-4} s⁻¹ in panel a) and 2×10^{-4} s⁻¹ in b)-g).

The updraft at 150 hPa (Fig. 14a) coincides with the ~ 50 km scale zonal flow minimum (Fig. 6a) and with the negative absolute vorticity maximum (Fig. 13c). The updraft near 700 hPa occurs in an arc shape at the eastern end of the downdraft (Fig. 14b). The zonal scale of the anomalies at 700 hPa is ~ 50 km (Fig. 6f, 14b). The downdraft and negative PV anomaly at 700

hPa have a wider zonal scale (\sim 100-150 km) than the updraft and positive PV anomaly (\sim 50 km) (Figs. 13f, 14b). Maximum vertical velocities in the updraft are \sim - 40 Pa/s (\sim 6 m/s) and \sim 10 Pa/s (\sim -1 m/s) in the downdraft (Fig. 14b).




Figure 14. HRRR pressure velocity (color bar, interval 0.5 Pa/s) at 1800 UTC on 10 August 2020 for a) 150 hPa and b) 700 hPa.

The updraft in HRRR weakened from 1700 to 2000 UTC (Fig. 7a), whereas observations show more steady intensity (Fig. 2). The absolute vorticity dipole at 700 hPa is oriented eastwest at 1700 UTC (Fig. 13e) but by 1800 UTC (Fig. 13f) the downdraft has become more diffuse and spread out to the south. This degradation of dipole structure in HRRR, compared with its peristence in the UWNMS-8, is consistent with the hypothesis that a stable dipole helps to maintain the strength of the derecho.

c) HRRR zonal and meridional sections

A movie of the longitude-altitude structure of zonal wind (Fig. S1) shows the evolution of the derecho in HRRR. In the early stages the updraft was associated with moderate easterlies. The structure at 1800 UTC is now diagnosed in detail. Westerlies in the 700-400 hPa layer near 94° W extend eastward and downward to the surface near 92° W, with values exceeding ~ 35 m/s near 850 hPa, 92.3° W (Fig. 15a). A northerly jet occurs near the gust front, exceeding ~ 20 m/s between 92.5° W and 92° W in the 800-900 hPa layer (Fig. 15b). The mid-tropospheric westerly inflow jet curves equatorward as it descends in the downdraft (Figs. 15a-c). West of 92.5° W, a layer of easterly flow near 200 hPa (Fig. 15a) lies between the westerlies at 150 and 300 hPa (Fig. 6a, b).




Figure 15. Longitude-altitude sections at 42°N (looking north) of HRRR a) zonal wind (color bar range, interval 5 m/s), b) meridional wind (color bar, interval 5 m/s), c) pressure velocity (color bar, interval 2.5 Pa/s), and d) absolute vorticity (color bar, interval 0.00025 s⁻¹) at 1800 UTC 10 August 2020. The zero vertical velocity line (heavy dashed) and locations of updrafts and downdraft (arrows) in c) are transposed onto a), b), and d). The locus of southward flow in b) is indicated with outward arrow symbols (⊙) and transposed onto a), c), and d). Distance east of the center of the HRRR domain (km) is shown along the x-axis. Pressure (hPa) and altitude (km, gray tick marks) are shown on the y-axis.

Air in the westerly inflow layer splits, with one branch descending and curving southward, and the other rising into the upper troposphere and curving northward, where

westerly flow exceeds 10 m/s (Figs. 15a, b). The downdraft coincides with the maximum in eastward flow (Figs. 15a, c), with values of ~ 5 Pa/s (-0.5 m/s) near 850 hPa. The downdraft (Fig. 15c), axis of maximum northerly flow (Fig. 15b), and westerly inflow layer (Fig. 15a) all tilt westward with height, with slope ~ -170 km / 5 km (elevation angle $\sim 1.7^{\circ}$). The updraft ahead of the gust front (Fig. 15c) tilts westward with altitude, with slope ~ -30 km / 8 km (elevation angle $\sim 15^{\circ}$). Updraft values exceed ~ -15 Pa/s (1.5 m/s) near 850 hPa and ~ 3 m/s near 400 hPa. Above ~ 650 hPa the updraft coincides with weak westerly flow (Fig. 15a, c). Below 650 hPa the updraft lies just to the east of the northerly jet maximum (Figs. 15b, c).

An east-west absolute vorticity dipole is found below 700 hPa, with values exceeding - $5 \times 10^{-4} \, \text{s}^{-1}$ and $20 \times 10^{-4} \, \text{s}^{-1}$ (\sim - 2.5 and 10 PVU) (Fig. 15d). The positive absolute vorticity region is located in the updraft while the negative region is located in the downdraft (Figs. 15c, d). The maximum in northerly flow (Fig. 15b) tilts westward with altitude and lies along the zero contour separating the two absolute vorticity anomalies (Fig. 15d). This meridional jet is kinematically consistent with the east-west dipole in absolute vorticity. This meso-beta scale PV dipole structure is distinct from the meso-beta cyclonic book-end vortex and meso-gamma scale MCVs. (Since the updraft also tilts poleward with altitude, the PV dipole in the UTLS maximizes \sim 50 km poleward of 42°N (Fig. 13c), such that this feature is not apparent in Fig. 15d.)

The pattern of strong westerlies which enters the downdraft descends from \sim 700 hPa (\sim 3 km) to the surface (Fig. 15a). The updraft and downdraft near 800 hPa are \sim 50 km wide (Fig. 15c). The meridional breadth of the region of strong zonal winds is \sim 50 km at 92°W (Fig. 6e), but is \sim 150 km in the 700 - 850 hPa layer at 92.5°W (Fig. 16a). The meridional breadth of the updraft and downdraft (Figs. 16c, g) and of the absolute vorticity anomaly (Figs. 16d, h) in the lower troposphere is \sim 50 km.

The updraft and downdraft in a MCS are part of a pattern of gravity wave phase axes which tilts westward with altitude (Lin 2007), with characteristic horizontal and vertical wavelengths $L_x \sim 200$ km and $L_z \sim 12$ km (Fig. 15c). These gravity waves may be related to the oscillations in pressure and wind speed seen at Des Moines after the passage of the gust front (Fig. 5a), which have a period of ~ 2 hours. For a gravity wave to travel eastward 200 km in 2 hours requires a zonal phase speed c ~ 100 km/hr, or 28 m/s. Since this is the speed of propagation of the derecho, this suggests that there is a critical surface, or steering level in the

mid-troposphere. In linear gravity wave theory, the relationship between pressure amplitude and wind amplitude is $P_o \rho c \sim V_o$. A pressure amplitude of 1.5 hPa (Fig. 5a) implies $V_o \sim 5$ m/s, which is consistent with the variation in Fig. 16b.




Figure 16. Latitude-altitude sections (eastward view) at 92.5°W (left) and 92°W (right) of HRRR a) zonal wind (color bar range, interval 5 m/s), b) meridional wind (color bar, interval 5 m/s), c) pressure velocity (color bar, interval 2.5 Pa/s), and d) absolute vorticity (color bar, interval .00025 s⁻¹) at 1800 UTC 10 August 2020. The dashed vertical line shows 42°N, the latitude of sections in Fig. 8. Distance north of the center of the HRRR domain is shown in km. Pressure (hPa) and altitude (km, gray tick marks) are shown on the vertical axis.

A meridional slice at 92.5° W (Fig. 16a) shows that the westerly inflow jet below 650 hPa has a maximum of ~ 35 m/s near 42° N, which is continuous with westerlies in the 800-700 hPa

layer to the north (Figs. 6d, 16a). At 92°W the "nose" of westerly flow is meridionally ~ 100 km wide (Fig. 16b). The layer of westerlies at 92.5°W coincides with northerly flow, with the layer descending as it moves southward (Figs. 16a-c). The "nose" of the derecho at 92°W contains a prominent northerly flow component (Figs. 16e, f).

At 92.5°W, the 600 hPa level separates the portion of the westerly inflow jet which rises poleward into the upper troposphere from that which descends equatorward in the downdraft (Figs. 16a-c). The meridional section of pressure velocity at 92°W (Fig. 16g) slices through the horseshoe-shaped, westward-tilting updraft, with a portion of the downdraft appearing in the middle near the surface, indicating that this is close to the gust front. At 92.5°W the downdraft coincides with negative absolute vorticity in the lower troposphere (Figs. 16c, d), while near 92°W, positive absolute vorticity coincides with the updraft, which extends into the upper troposphere (Figs. 16g, h).

d) Reynolds stresses

Products of velocity components are useful for highlighting fundamental flow structures, estimating momentum fluxes, and relating them to the synoptic flow. Below the level of zero vertical wind, westerly momentum is transported southward (Figs. 17a, d). In the downdraft, westerly momentum (Figs. 17b, e), and northerly momentum (Figs. 17 c, f) are transported downward within the derecho.

In the 500-700 hPa westerly inflow layer, uv > 0 (Fig. 17a), consistent with northward transport of westerly momentum associated with the approaching trough (Fig. 6c). In the UTLS, uv exhibits a quadrupole pattern in longitude-altitude associated with updraft outflow (Fig. 17a). The negative region near 200 hPa is consistent with southeasterly front-to-rear outflow and trailing stratiform precipitation (Houze et al. 1989). The positive region near 150 hPa is consistent with southwesterly outflow creating a layer of leading stratiform precipitation (Figs. 17a, 3a). At 92.5°W (Fig. 17d) one may see that this region of uv > 0 extends well to the north of the derecho.

Above ~ 600 hPa there is upward transport of westerly momentum, where the upper half of the westerly inflow jet merges with the gust front updraft (Figs. 17b, e). This pattern of momentum flux divergence transports momentum away from the 700-500 westerly inflow jet, consistent with the deceleration observed in Fig. 15a. Near the top of the updraft at 150 hPa, the

vertical momentum flux convergence near 92.2°W (Fig. 17b) is compatible with westerly acceleration in the outflow jet.




Figure 17. Reynolds stresses u v (top, color bar, interval 100 m² s⁻²), $u \omega$ (middle, color bar, interval 50 m Pa s⁻²), and $v \omega$ (bottom, color bar, interval 50 m Pa s⁻²) in HRRR for longitude-altitude sections at 42°N looking north (a, b, c) and for a latitude-altitude section at 92.5°W looking east (d, e, f). The zero vertical wind line in Fig. 15c) is transposed onto panels a), b), and c). At left, both longitude and distance (km) east of the center of the HRRR domain are shown. At right, both latitude and distance north of the center of the HRRR domain are shown. Pressure (hPa) and altitude (km, gray tick marks) are shown on the vertical axis.

Below the zero vertical wind line near ~ 600 hPa, northerly and westerly momentum are transported downward in the downdraft, while above ~ 600 hPa, southerly and westerly momentum are transported upward in the updraft (Figs. 17c, e). These results are consistent with down-gradient transport from the mid-tropospheric westerly jet streak associated with the cyclone and from the northerly winds in the surface anticyclone.

e) 3D isosurfaces

The westerly inflow jet curves anticyclonically as it descends and becomes the northerly outflow jet, as indicated by their spatial co-location viewed from the southwest and south (Figs. 18a, b). This relationship is similar to that shows for UWNMS-8 in Figs. 12a, b, except the downdraft slopes more gently in HRRR. The pattern in the v = -18 m/s isosurface (Fig. 18b) shows the core outflow, plus the trailing speed maximum at the south edge of the gust front, as in Fig. 6f. This outflow jet lies between the absolute vorticity dipole in HRRR (Fig. 18c), similar to UWNMS-8 (Fig. 12).




Figure 18. HRRR 3D isosurfaces for 1800 UTC 10 August 2020 of a) u = 23 m/s (yellow), v = -18 m/s (green), and $\omega = 6$ Pa/s (orange) viewed obliquely from the southwest, b) as in a) except viewed from the south, and c) v = -18 m/s (green), $\zeta + f = 10x10^{-4}$ s⁻¹ (red), and $\zeta + f = -7x10^{-4}$ s⁻¹ (blue) viewed from the south.

A view from the north reinforces these spatial relationships (Fig. 19). The region of westerlies associated with the upper-level cyclone is continuous with the westerly inflow jet and northerly outflow jet, which occurs to the east of the positive absolute vorticity anomaly (Fig. 19a). The meridional jet lies between the absolute vorticity dipole (Fig. 19b). The westerly inflow jet and northerly jet blend, wrapping around the negative absolute vorticity anomaly (Fig. 19c).




Figure 19. HRRR 3D isosurfaces for 1800 UTC 10 August 2020 viewed from the north of a) u = 23 m/s (yellow), v = -18 m/s (green), and $\zeta + f = 10x10^{-4}$ s⁻¹ (red), b) v = -18 m/s (green), $\zeta + f = 10x10^{-4}$ s⁻¹ (red), and $\zeta + f = -7x10^{-4}$ s⁻¹ (blue), and c) u = 23 m/s (yellow), v = -18 m/s (green), and $\zeta + f = -7x10^{-4}$ s⁻¹ (blue).

7. Discussion

a) Summary comparison

All three numerical depictions show an east-west PV dipole structure in the mid- to lower troposphere, coincident with the updraft/downdraft pair, with a northerly jet between. HRRR maintained a continuous derecho-like structure in time, but lagged observed locations by ~ 80 km, with reduced zonal translation speed and weakening during 1700 – 2000 UTC. UWNMS-4 generated a derecho-like structure 2 hours earlier than UWNMS-8, but the tilt of the updraft/downdraft was too weak, and rapid amplification was followed by a breakdown into MCVs. UWNMS-8 was slow to amplify but then maintained a coherent derecho-like structure for ~ 2 hours, with amplitudes similar to HRRR, and exhibited a somewhat more realistic westward tilt with height. The gentle slope of the downdraft in HRRR was more realistic.

b) Role of steering level in the eastward progression of the derecho

To remain coherent for several hours, the gust front cannot significantly outrun the downdraft. Under the steering level hypothesis, the speed of the gust front, c, should be related to a spatial average of the fastest portions of the westerly inflow layer, U. The existence of such a steering level implies that there is a critical level where $c \sim U$. This may imply that the existence of a derecho depends on the existence of a mid-tropospheric westerly wind maximum, with moderate westerlies in the upper troposphere. This zonal wind profile, with shear ~ 25 m/s km⁻¹, is consistent with the proper combination of westward tilt with height and similar degree of outflow toward the northwest and northeast in the UTLS, as in the idealized modeling results of Weisman and Rotunno (1994).

Since the derecho speed was similar in UWNMS-8 and UWNMS-4, features such as a realistic tilt of the updraft/downdraft and a well-developed cold pool may be less import in determining storm motion than the zonal wind speed upstream in the 750-500 hPa layer. Note the existence of strong westerly flow in the 3-5 km layer to the north of the gust front (Figs. 11a, b, 16a). The movement of this "arm" of high speed air associated with the upper tropospheric cyclone guides the steering level of the derecho eastward.

c) Role of steering level and comparison with breaking orographic gravity waves

Peltier and Clark (1979), Lilly and Klemp (1979), and Durran and Klemp (1983) studied the saturation process of orographic gravity wave amplification to the point of turbulent breakdown, in which the downdraft brings high westerly wind speed air to the surface, followed

by a hydraulic jump. The resulting region of stalled zonal flow in the updraft is similar to zonal wind sections in the UWNMS (Figs. 9a, 10a) and HRRR (Figs. 15a, S1). The zero wind surface is a critical surface for orographic gravity waves. In a derecho, the ~ 25 m/s zonal wind surface can be regarded as a critical surface for waves with $c \sim 25$ m/s, which grow by resonant amplification with the diabatic heating pattern travelling at the same rate. Critical surface theory can help explain both the observed emission of gravity waves from the storm and nonlinear saturation, or reaching a finite-amplitude quasi-steady state.

A westward tilt with height is required to achieve this configuration. In the case of the derecho, the forcing pattern is moving at the speed of the flow (Figs. 8 and 9 of Lin and Smith 1986). The speed of a phenomenon arising from a resonant instability is most likely to travel at the speed of the forcing agent. The derecho is somewhat like an orographic gravity wave propagating at the speed of the inflow layer. Instead of a hydraulic jump, however, there is vertical shear of the meridional wind, with a northerly flow maximum at the leading edge of the downdraft and a southerly component to the updraft.

d) Mechanism of PV dipole formation

In the mature phase, an east-west PV dipole forms in association with the tilted updraft/downdraft pair, with a northerly flow maximum between. Kinematically, a northerly jet implies that positive relative vorticity exists to its east and negative vorticity to its west. Formation of the horizontal PV dipole in the lower troposphere occurred in conjunction with the formation of an updraft and downdraft, which are caused by diabatic heating/cooling.

Evans et al. (2014) and Xu et al. (2015) emphasized the role of surface friction and tilting of near-surface horizontal vorticity as a generation mechanism for mesovortices in a derecho. However, the amplitude of the PV dipole diagnosed in the 10 August 2020 derecho increases upward from the surface and the PV dipole coincides with the westward-tilting updraft/downdraft pair (Figs. 9, 10, 15d).

It is of interest to explore the effect of tilt on the diabatic stretching and tilting terms by considering the three progressively tilted structures shown in snapshots from the UWNMS-8 (Fig. 9), UWNMS-4 (Fig. 10), and HRRR (Fig. 15). The updraft and downdraft are most vertically-oriented in the case of UWNMS-4 at 1800 UTC. The diabatic stretching tendency of P in (1) may be estimated as follows. Assuming that $w \frac{\partial \theta}{\partial z} \sim Q$ and $\frac{\partial \theta}{\partial z} = \frac{\theta}{T} (\frac{\partial T}{\partial z} + \Gamma_d)$, where Γ_d is

the dry adiabatic lapse rate, an ambient moist adiabatic lapse rate $\frac{\partial T}{\partial z} \sim -6.5$ K/km, $\frac{\partial w}{\partial z} \sim 3$ m s⁻¹/2 km in the downdraft near 6 km (Fig. 9c), and $\zeta + f \sim -10^{-4}$ s⁻¹, the diabatic stretching term in (1) is estimated to be \sim - 3 PVU/hr, perhaps sufficient to generate the negative PV anomaly in Fig. 9d. As an example from UWNMS-8, the updraft in Fig. 10c exhibits $\frac{\partial w}{\partial z} \sim 4$ m s⁻¹/2 km near 4 km, with $\zeta + f \sim 2 \times 10^{-4}$ s⁻¹, giving a diabatic stretching tendency of \sim 3 PVU/hr, which may be adequate to generate the modeled positive PV anomaly in Fig. 10d.

As the tilt increases, the axes of maximum diabatic stretching tendency shift westward and underneath relative to the updraft and downdraft axes. This may be seen by estimating the diabatic stretching tendency in (2) for HRRR at 800 hPa, 92.5°W (just below the downdraft maximum), in the updraft near 850 hPa, 91.8°W (Fig. 15c), and near 650 hPa, 92.3°W (below the updraft maximum). Taking $\frac{\partial \omega}{\partial p} \sim \frac{5 Pa/s}{-100 \ hPa}$ and $\zeta + f \sim 10^{-4} \ s^{-1}$ in the westerly inflow jet below the downdraft maximum, vortex stretching gives $\sim -2x10^{-4} \ s^{-1}$ per hr. Taking $\frac{\partial \omega}{\partial p} \sim \frac{-10 \ Pa/s}{-100 \ hPa}$ in the updraft, vortex stretching gives $\sim 4x10^{-4} \ s^{-1}$ per hr, while taking $\frac{\partial \omega}{\partial p} \sim \frac{-15 \ Pa/s}{-200 \ hPa}$ below the updraft maximum, vortex stretching gives $\sim 3x10^{-4} \ s^{-1}$ per hr, comparable with absolute vorticity magnitudes in Fig. 15d. Note that the positive absolute vorticity maximum tilts westward such that it coincides with the zero wind line (Fig. 15d), where the vertical gradient in pressure velocity (Fig. 15c), hence diabatic stretching, is large. This westward and downward displacement in diabatic tilting tendency is a direct result of increasing tilt (Fig. 20).




Figure 20. Idealized diagram showing the maxima in diabatic stretching tendency for a) weak tilt and b) strong tilt. The downdraft (convergent arrows) coincides with negative PV anomaly (blue), the updraft (divergent arrows) coincides with positive PV anomaly (red), with a northerly jet between (dotted blue axis). With increasing tilt, the maxima in diabatic stretching tendency,

as indicated by the curve below each diagram, are displaced westward and downward relative to the updraft and downdraft.

Since the gradients in the diabating tilting terms maximize between the PV dipole they cannot strengthen the dipole. In the case of weak tilt, both diabatic tilting terms contribute toward negative PV tendency, maximizing along the axis of northerly flow, in quadrature with the PV anomalies. In the case of stronger tilt, products of gradients yield fine-scale patterns between the PV anomalies, with $\frac{\partial w}{\partial y} \frac{\partial u}{\partial z}$ in (2) causing positive tendency to the east of the negative PV anomaly and negative tendency to the west of the positive PV anomaly, while $-\frac{\partial w}{\partial x} \frac{\partial v}{\partial z}$ causes the opposite pattern, tending to cancel the other out. At the three locations described above (Fig. 15), $\frac{\partial \omega}{\partial x} \sim 0 \sim \frac{\partial \omega}{\partial y}$, so the diabatic tilting term is expected to be small. This analysis suggests that the diabatic stretching term is primarily responsible for generating the PV dipole. The generation and maintenance of the PV dipole may facilitate the coherence and longevity of a derecho.

The westerly inflow layer which curves anticyclonically in the downdraft and gust front inflow layer which curves cyclonically in the updraft can be understood in terms of the horizontal momentum equations following the motion

904 (a)
$$\frac{du}{dt} = fv - \frac{1}{\rho} \frac{\partial p}{\partial x}$$
, (b) $\frac{dv}{dt} = -fu - \frac{1}{\rho} \frac{\partial p}{\partial y}$. (3)

As air in the westerly inflow layer travels eastward, the northward pressure gradient force diminishes as it descends, such that northerly acceleration occurs (3b). An increase in the westward pressure gradient force and the Coriolis torque on northerly flow causes the zonal flow to decelerate (3a), yielding anticyclonic vorticity in the downdraft. As air in the gust front inflow layer ascends, an increasing northward pressure gradient force and Coriolis torque on easterly flow cause southerly acceleration (3b), yielding cyclonic vorticity in the updraft. The existence of a stagnation point at the east end of the westerly inflow jet in the mid-troposphere implies the existence of a meso-high, which steers the westerly inflow layer southward as it descends, and the front-to-rear flow northward as it ascends. As the westerly inflow jet curves around the anticyclonic member, angular momentum decreases radially outward, consistent with inertial instability, where air with high angular momentum diverges outward, creating the northerly outflow jet.

8. Conclusions

The Iowa derecho of 10 August 2020 was characterized by westerly winds exceeding 30 m/s in a ~100 km wide swath through central Iowa, developing an equatorward expansion as it traveled eastward, forming a comma echo. When the gust front reached DSM at 1600 UTC, the temperature dropped 11 K in 10 minutes, due to advection of low values of θ_e by the downdraft, replacing the previous boundary layer air. The pressure rose initially by ~ 5 hPa and remained elevated for several hours during the passage of the meso-anticyclone. The arrival of the gust front was characterized by an abrupt shift to northerly flow and a rapid increase in speed (Fig. 5b). The speed of the flow in the 500-750 hPa westerly inflow layer upstream of the gust front is similar to the speed of propagation of the derecho. The westerly inflow layer is a southward extension of a region of high wind speed (jet streak) associated with the approaching synoptic trough. As pointed out by Hinrichs (1888), there is a significant northerly aspect to the surface outflow. Below 850 hPa, the northerly jet between the updraft/downdraft and PV dipole exists proximal to the southeastern flank of the lower-tropospheric anticyclone.

When the updraft/downdraft pair strengthened, an east-west PV dipole formed primarily by diabatic stretching, and propagated eastward with the derecho, with the northerly outflow jet creating a comma radar echo pattern. These structures in HRRR and UWNMS confirm that the observed northerly jet at Des Moines implies an east-west PV dipole in the lower troposphere.

In a derecho the positive PV anomaly extends to higher altitudes than the negative PV anomaly, with westerly inflow above 5 km contributing to the updraft and the negative PV anomaly beginning near the top of the downdraft, which is confined below 5 km. The inertially unstable property of the negative PV member may help facilitate northerly acceleration, promoting the equatorward bulge of the radar echo.

The fundamental structure of a westward-tilted PV dipole may be on a continuum, which includes the more upright structure in a mature MCS studied by Fritsch et al. (1994) and Jiang and Raymond (1995), where a mesoscale warm-core vortex (positive PV anomaly) in the latent heat-driven updraft maximizes in the mid-troposphere, with a shallow anticyclone (negative PV anomaly) near the surface. With a westerly inflow layer, a tilted structure forms with the near-surface PV anomaly maximizing in the mid-troposphere. If the cold pool is poorly developed, such as in UWNMS-4 at 1800 UTC (Fig. 9), the PV dipole extends through most of the depth of

the troposphere, with the negative member maximizing at the level of westerly inflow where the vertical wind diverges (Figs. 9a, c, d). With a more developed cold pool, such as in UWNMS-8 at 1900 UTC (Fig. 10), the PV dipole maximizes at a lower altitude. Finally, in HRRR (Fig. 15), the slope of the downdraft is much more gentle and the negative member of the PV dipole tilts more and maximizes at a still lower altitude.

These results can be interpreted in the context of the idealized study by Weisman and Rotunno (2004). The mid-tropospheric westerly inflow layer was shown to be associated with a region of westerlies extending southward from the approaching cyclone. This jet streak provides a steering level or critical level for the derecho which can help amplify the structure. In this present case, the wind profile is most similar to their case with shallow shear at 20 m s⁻¹ / 5 km and constant westerlies above. The formation of a horizontal PV dipole may contribute to strengthening and stabilizing the self-perpetuating structure of a derecho. Since the speed of propagation of the derecho was very similar in HRRR, UWNMS-4, and UWNMS-8, despite differing strength and structure, this supports the hypothesis that the speed of the derecho is controlled by a steering level flow which is related to the jet streak in the synoptic-scale cyclone.

Acknowledgements. The authors would like to acknowledge support from NSF Grant AGS-1947658. We would like to thank Pete Pokrandt for help with converting the HRRR grid to local latitude / longitude. We are grateful to the reviewers who provided many helpful comments, and especially Editor Russ Schumacher's suggestion of using HRRR data.

- 969 Data Availability. Composite radar files are available from
- 970 https://mesonet.agron.iastate.edu/current/mcview.phtml?prod=comprad&java=script&mode=arc
- 971 <u>hive&frames</u>. Surface meteorological observations are available from
- 972 https://mesonet.agron.iastate.edu/request/asos/1min.phtml#.
- 973 The Davenport, IA sounding is available from
- 974 https://www.spc.noaa.gov/exper/archive/event.php?date=20200810.
- Observations for creating the surface station plot are available from https://rda.ucar.edu/.
- 976 NOAA NEXRAD-II GOES-16 images and lightning data are available from
- 977 https://www.ncdc.noaa.gov/wct/.

- NOAA High-Resolution Rapid Refresh (HRRR) Model was last accessed on July 15, 2023 from
- 979 https://registry.opendata.aws/noaa-hrrr-pds.
- 980 UWNMS images may be reproduced from the VIS5D files of the simulations, which are stored
- on a dedicated computer in the AOS Department at the UW-Madison, available from the authors.

- 983 Literature Cited
- Adams-Selin, R. D., and R. H. Johnson, 2013: Examination of gravity waves associated with the
- 985 13 March 2003 bow echo. *Mon. Wea. Rev.*, **141**, 3735–3756.
- Adams-Selin, R. D., 2020: Impact of convectively generated low-frequency gravity waves on
- evolution of mesoscale convective systems. *J. Atmos. Sci.*, **77**, 3341-3460.
- 988 Andrews, D. G., J. R. Holton, and C. B. Leovy, 1987: Middle Atmosphere Dynamics, Academic
- 989 Press, 489 pp.
- 990 Atkins, N. T., and M. St. Laurent, 2009: Bow echo mesovortices. Part I: Processes that
- influence their damaging potential. *Mon. Wea. Rev.*, **137**, 1497-1513.
- 992 Bretherton, C. S., and Smolarkiewicz, P., 1989: Gravity waves, compensating subsidence and
- detrainment around cumulus clouds. *J. Atmos. Sci.*, **46**, 740-759.
- Campbell, M. A., A. E. Cohen, M. C. Coniglio, A. R. Dean, S. F. Corfidi, S. J. Corfidi, and C.
- 995 M. Mead, 2017: Structure and motion of severe-wind-producing mesoscale convective
- systems and derechos in relation to the mean wind. Wea. Forecasting, **32**, 423-439.
- 997 Chagnon, J. M. and S. L. Gray, 2009: Horizontal potential vorticity dipoles on the convective
- 998 storm scale. *Quart. J. Roy. Meteorol. Soc.*, **135**, 1392–1408.
- 999 Chicago Weather Forecast Office, 2021: "August 10, 2020: Corn Belt Derecho",
- https://www.weather.gov/lot/2020aug10.
- 1001 Coniglio, M. C., and D. J. Stensrud, 2004: Interpreting the climatology of derechos. Weather
- *and Forecasting*, **19**, 595-605.
- 1003 Coniglio, M. C., D. J. Stensrud, and M. B. Richman, 2004: An observational study of derecho-
- producing convective systems. Weather and Forecasting, 19, 320-337.
- 1005 Corfidi. S. F., J. H. Merritt, and J. M. Fritsch, 1996: Predicting the movement of mesoscale
- convective complexes. Wea. Forecasting, 11, 41–46.
- 1007 Corfidi, S. F., M. C. Coniglio, A. E. Cohen, and C. M. Mead, 2016: A proposed revision to the
- definition of "derecho." Bull. Amer. Meteor. Soc., 97, 935–949.

- Davies-Jones, R., 1984: Streamwise vorticity: The origin of updraft rotation in supercell storms.
- 1010 J. Atmos. Sci., 30, 2991-3006.
- Davis, C. A., and T. J. Galarneau, 2009: The vertical structure of mesoscale convective vortices.
- 1012 *J. Atmos. Sci.*, **66**, 686–704.
- Dowell, D. C., et al., 2022: The High-Resolution Rapid Refresh (HRRR): An hourly updating
- convection-allowing forecast model. Part I: Motivation and system description. Wea.
- 1015 *Forecasting*, **37**, 1371-1395.
- Durran, D.R., and J.B. Klemp, 1983: A compressible model for the simulation of moist mountain
- 1017 waves. Mon. Wea. Rev., 111, 2341-2361.
- 1018 Evans, J. S., and C. A. Doswell III, 2001: Examination of derecho environments using proximity
- 1019 soundings. *Wea. Forecasting*, **16**, 329-342.
- 1020 Evans, C., M. L. Weisman, and L. F. Bosart, 2014: Development of an intense, warm-core
- mesoscale vortex associated with the 8 May 2009 "Super Derecho" convective event. J.
- 1022 Atmos. Sci., 71, 1218-1240.
- Fovell, R. G., G. L. Mullendore, and S.-H. Kim, 2006: Discrete propagation in numerically
- simulated nocturnal squall lines. *Mon. Wea. Rev.*, **134**, 3735–3752.
- Fritsch, J. M., J. D. Murphy, and J. S. Kain, 1994: Warm core vortex amplification over land. J.
- 1026 Atmos. Sci., **51**, 1780-1807.
- 1027 Guastini, C. T., and L. F. Bosart, 2016: Analysis of a progressive derecho climatology and
- associated formation environments. *Mon. Wea. Rev.*, **144**, 1363-1382.
- Harvey, B., J. Methven, C. Sanchez, and A. Schäfler, 2020: Diabatic generation of negative
- potential vorticity and its impact on the North Atlantic jet stream. Q. J. Roy. Meteorol.
- 1031 *Soc.*, **146(17)**, DOI: 10.1002/qj.3747.
- Hashino, T. and G. J. Tripoli, 2007: The Spectral Ice Habit Prediction System (SHIPS), part 1:
- Model description and simulation of the vapor deposition process. J. Atmos. Sci., 64,
- 1034 2210-2237.
- Hashino, T. and Tripoli, G.J., 2011: The Spectral Ice Habit Prediction System (SHIPS), part 4:
- Box model simulations of the habit-dependent aggregation process. J. Atmos. Sci., 68,
- 1037 1142-1161.
- Haynes, P. H., and M. E. McIntyre, 1990: On the conservation and impermeability theorems for
- 1039 potential vorticity. *J. Atmos. Sci.*, **47**, 2021-2031.

- Hinrichs, G., 1888: Tornados and derechos. *American Meteorological Journal*, 306-349.
- Hernandez-Duenas, G., L. M. Smith, and S. N. Stechmann, 2014: Investigation of Boussinesq
- dynamics using intermediate models based on wave–vertical interactions. *J. Fluid*
- 1043 *Mech.*, **747**, 247–287.
- Hitchman, M. H., and S. M. Rowe, 2017: On the similarity of lower stratospheric potential
- vorticity dipoles above tropical and midlatitude deep convection. J. Atmos. Sci., 74, 2593-
- 1046 2613.
- Hitchman, M. H., and S. M. Rowe, 2019: On the 3D structure and formation of UTLS jetlets
- associated with potential vorticity dipoles in tropical cyclones. *Mon. Wea. Rev.*, **147**,
- 1049 4107-4125.
- Hitchman, M. H., and S. M. Rowe, 2021: On the formation of tropopause folds and constituent
- gradient enhancement near westerly jets. *J. Atmos. Sci.*, **78**, 2057-2074.
- Holton, J. R., and G. J. Hakim, 2013: "An Introduction to Dynamic Meteorology", Fifth Ed.,
- Academic Press, NY, 532 pp.
- Jiang, H., and D. J. Raymond, 1995: Simulation of a mature mesoscale convective system using
- a nonlinear baance model. *J. Atmos. Sci.*, **52**, 161-175.
- Johns, R. H., and W. D. Hirt, 1987: Derechos: Widespread convectively induced windstorms.
- 1057 *Wea. Forecasting*, **2**, 32–49.
- Lane, T. P., and F. Zhang, 2011: Coupling between gravity waves and tropical convection at
- mesoscales. J. Atmos. Sci., **68**, 2582-2598.
- Lilly, D. K., and J. B. Klemp, 1979: The effects of terrain shape on nonlinear hydrostatic
- 1061 mountain waves. *J. Fluid Mech.*, **95**, 241-262.
- Lin, Y.-L., and R. B. Smith, 1986: Transient dynamics of airflow near a local heat source. J.
- 1063 Atmos. Sci., 43, 40-49.
- Lin, Y.-L., 2007: *Mesoscale Dynamics*. Cambridge University Press, New York, 630 pp.
- Mahoney, K. M., G. M. Lackmann, and M. D. Parker, 2009: The role of momentum transport in
- the motion of a quasi-idealized mesoscale convective system. Mon. Wea. Rev., 137,
- 1067 3316-3338.
- Markowski, P., and Y. Richardson, 2010: Mesoscale Meteorology in Midlatitudes. John Wiley
- and Sons, Ltd., Chichester, United Kingdom, 407 pp.

- 1070 Montgomery, M. T., M. E. Nicholls, T A. Cram, and A. B. Saunders, 2006: A vortical hot tower
- route to tropical cyclogenesis. *J. Atmos. Sci.*, **63**, 355-386.
- 1072 Muraki, D., and C. Snyder, 2007: Vortex dipoles for surface quasigeostrophic models. *J. Atmos*.
- 1073 *Sci.*, **64**, 2961–2967.
- Oertel, A., M. Boettcher, H. Joos, M. Sprenger, M., and H. Wernli, 2020: Potential vorticity
- structure of embedded convection in a warm conveyor belt and its relevance for large-
- scale dynamics. Weather Clim. Dynam., 1, 127–153.
- Pandya, R. E., D. R. Durran, and M. L. Weisman, 2000: The influence of convective thermal
- forcing on the three-dimensional circulation around squall lines. J. Atmos. Sci., 57, 29–
- 1079 45.
- Peltier, W. R., and T. L. Clark, 1979: The evolution and stability of finite-amplitude mountain
- waves. Part II. Surface wave drag and severe downslope windstorms. J. Atmos. Sci., 36,
- 1082 1498-1529.
- Plougonven, R., and F. Zhang, 2014: Internal gravity waves from atmospheric jets and fronts.
- 1084 *Rev. Geophys.*, **52**, 33–76.
- Prince, K. C., and C. Evans, 2022: Convectively generated negative potential vorticity
- enhancing the jet stream through an inverse energy cascade during the extratropical
- transition of Hurricane Irma. J. Atmos. Sci., 79, 2901-2918.
- 1088 Przybylinski, R., 1995: The bow echo: Observations, numerical siulations, and severe weather
- detection methods. *Weather and Forecasting*, **10**, 2013-218.
- Rappin, E. D., M. C. Morgan, and G. J. Tripoli, 2011: The impact of outflow environment on
- tropical cyclone intensification and structure. J. Atmos. Sci., 68, 3224-3242.
- Raymond, D. J., and H. Jiang, 1990: A theory for long-lived mesoscale convective systems. J.
- 1093 *Atmos. Sci.*, **47**, 3067-3077.
- Rocha, C. B., G. L. Wagner, and W. R. Young, 2018: Stimulated generation: extraction of
- energy from balanced flow by near-inertial waves. *J. Fluid Mech.*, **847**, 417-451.
- Rotunno, R., J. B. Klemp, and M. L. Weisman, 1988: A theory for strong, long-lived squall lines.
- 1097 *J. Atmos. Sci.*, **45**, 463–485.
- Rowe, S. M., and M. H. Hitchman, 2015: On the role of inertial instability in stratosphere
- troposphere exchange near midlatitude cyclones. J. Atmos. Sci., 72, 2131-2151.

- Rowe, S. M., and M. H. Hitchman, 2016: On the relationship between inertial instability,
- poleward momentum surges, and jet intensifications near midlatitude cyclones. *J. Atmos.*
- 1102 *Sci.*, **73**, 2299-2315.
- Rowe, S. M., and M. H. Hitchman, 2020: The destruction of a stratospheric potential vorticity
- intrusion due to inertial instability in the UTLS as it relates to the 20 August 2018
- Southern Wisconsin extreme flooding event. *Mon. Wea. Rev.*, **148**, 4397–4414.
- Shige, S., and T. Satomura, 2000: The gravity wave response in the troposphere around deep
- 1107 convection. *J. Meteor. Soc. Japan*, **78**, 789–801.
- Seigel, R. B., and S. C. van den Heever, 2013: Squall-line intensification via hydrometeor
- 1109 recirculation. *J. Atmos. Sci.*, **70**, 2012-2031.
- Smull, B. F., and R. A. Houze, Jr., 1987: Rear inflow in squall lines with trailing stratiform
- 1111 precipitation. *Mon. Wea. Rev.*, **115**, 2869-2889.
- 1112 Snyder, C. D., D. J. Muraki, R. Plougonven, and F. Zhang, 2007: Inertia–gravity waves
- generated within a dipole vortex. J. Atmos. Sci., 64, 4417–4431.
- 1114 Snyder, C. D., R. Plougonven, and D. J. Muiraki, 2009: Mechanisms for spontaneous gravity
- wave generation within a dipole vortex. J. Atmos. Sci., 66, 3464–3478.
- 1116 Tripoli, G. J., and W. R. Cotton, 1982: The Colorado State University three-dimensional
- cloud/mesoscale model-1982, Part I: General theoretical framework and sensitivity
- 1118 experiments. J. de Rech. Atmos., 16(3), 185-219.
- 1119 Tripoli, G. J., 1992a: A non-hydrostatic mesoscale model designed to simulate scale interaction.
- 1120 *Mon. Wea. Rev.*, **120**, 1342-1359.
- 1121 Tripoli, G.J., 1992b: An explicit three-dimensional nonhydrostatic numerical simulation of a
- tropical cyclone. *Meteorol. Atmos. Phys.*, **49**, 229-254.
- Wakimoto, R. M., J. V. Murphey, C. A. Davis, and N. T. Atkins, 2006: High winds generated
- by bow echoes. Part II: The relationship between the mesovortices and damaging
- straight-line winds. *Mon. Wea. Rev.*, **134**, 2813-2829.
- Wang, S., F. Zhang, and C. Snyder, 2009: Generation and propagation of inertia-gravity waves
- from vortex dipoles and jets. *J. Atmos. Sci.*, **66**, 1294–1314.
- Weijenborg, C., J. M. Chagnon, P. Friederichs, S. L. Gray, and A. Hense, 2017: Coherent
- evolution of potential vorticity anomalies associated with deep moist convection. Q. J. R.
- 1130 *Meteorol. Soc.*, **143**, 1254–1267.

1131 Weisman, M. L., 1992: The role of convectively generated rear inflow jets in the evolution of long-lived mesoconvective systems. J. Atmos. Sci., 49, 1826–1847. 1132 1133 Weisman, M. L., 1993: The genesis of severe, long-lived bow echoes. J. Atmos. Sci., 50, 645– 1134 670. Weisman, M. L., and R. F. Trapp, 2003: Low-level mesovortices within squall lines and bow 1135 1136 echoes. Part I: Overview and dependence on environmental shear. Mon. Wea. Rev., **131**, 2779-2803. 1137 Weisman, M. L, and R. Rotunno, 2004: "A theory for strong long-lived squall lines" revisited. J. 1138 1139 Atmos. Sci., **61**, 361-382. 1140 Weisman, M. L., C. Evans, and L. Bosart, 2013: The 8 May 2009 Superderecho: Analysis of a real-time explicit convective forecast. Wea. Forecasting, 28, 863-892. 1141 Wernli, H., and H. Davies, 1997: A Lagrangian-based analysis of extratropical cyclones. I: The 1142 method and some applications. Quart. J. Roy. Meteorol. Soc., 123, 467-489. 1143 Xu, X, M. Xue, and Y. Wang, 2015: The genesis of mesovortices within a real-data simulation 1144 of a bow echo system. J. Atmos. Sci., 72, 1963-1986. 1145 1146 1147