

TORUS FIXED POINT SETS OF HESSENBERG SCHUBERT VARIETIES IN REGULAR SEMISIMPLE HESSENBERG VARIETIES

MEGUMI HARADA and MARTHA PRECUP

(Received February 10, 2022, revised June 10, 2022)

Abstract

It is well-known that the T -fixed points of a Schubert variety in the flag variety $GL_n(\mathbb{C})/B$ can be characterized purely combinatorially in terms of Bruhat order on the symmetric group \mathfrak{S}_n . In a recent preprint, Cho, Hong, and Lee give a combinatorial description of the T -fixed points of Hessenberg analogues of Schubert varieties (which we call Hessenberg Schubert varieties) in a regular semisimple Hessenberg variety. This note gives an interpretation of their result in terms of Bruhat order by making use of a partition of the symmetric group defined using so-called subsets of Weyl type. The Appendix, written by Michael Zeng, proves a lemma concerning subsets of Weyl type which is required in our arguments.

1. Introduction

The main result of this note is a characterization of the T -fixed point sets of (opposite) Hessenberg Schubert varieties in terms of Bruhat order. We achieve this by giving an interpretation in terms of Bruhat order of the notion of *reachability* – a concept introduced by Cho, Hong, and Lee in [3]. Hessenberg varieties have recently garnered great interest in different research communities due to their connections (which have come to light in the past decade) to many areas, not the least of which is the famously unsolved Stanley–Stembridge conjecture in algebraic combinatorics [12]; see [1] for an overview. In [3], the authors analyze the Białynicki–Birula decomposition of a regular semisimple Hessenberg variety, and they introduce and use the notion of reachability to give an explicit description of the T -fixed points of the closure of a given Białynicki–Birula cell. Such a closure is called a Hessenberg Schubert variety, since these are analogues of classical Schubert varieties in the flag variety.

To describe our results, we briefly recall some terminology. Let $w \in \mathfrak{S}_n$ and $\Omega_w^\circ := B_w B / B$ denote the corresponding opposite Schubert cell in the type A flag variety $GL_n(\mathbb{C})/B$. Here B and B_- denote the Borel subgroups in $GL_n(\mathbb{C})$ of upper-triangular and lower-triangular matrices, respectively. Given $u \in \mathfrak{S}_n$, the partial order on \mathfrak{S}_n defined by

$$(1.1) \quad w \leq u \text{ whenever } uB \in \Omega_w^\circ := \overline{\Omega_w^\circ}$$

is called **Bruhat order**. This order is fundamental in the study of the symmetric group and the geometry of flag varieties and related spaces. Now let $\mathbf{S} \in \mathfrak{gl}_n(\mathbb{C})$ be a diagonal matrix with distinct eigenvalues, $h : [n] \rightarrow [n]$ a Hessenberg function, and $\mathcal{Hess}(\mathbf{S}, h)$ the corresponding regular semisimple Hessenberg variety in $GL_n(\mathbb{C})/B$ (for definitions see

Section 2). We call the intersection $\Omega_{w,h}^\circ = \Omega_w^\circ \cap \mathcal{Hess}(\mathbf{S}, h)$ the (opposite) Hessenberg Schubert cell indexed by $w \in \mathfrak{S}_n$, and its closure $\Omega_{w,h} := \overline{\Omega_{w,h}^\circ}$ is the **opposite Hessenberg Schubert variety indexed by w** . The torus $T := B \cap B_-$ of diagonal matrices in $GL_n(\mathbb{C})$ acts on $\mathcal{Hess}(\mathbf{S}, h)$ by left multiplication. Cho, Hong, and Lee gave an explicit characterization of the T -fixed points in $\Omega_{w,h}$ in terms of reachability [3]. Our main contribution in this note is a reinterpretation, using subsets of Weyl type and the corresponding partition of \mathfrak{S}_n , of reachability in terms of Bruhat order (Proposition 3.13), which then allows us to give a concrete description of the T -fixed points in $\Omega_{w,h}$ in terms of Bruhat order (Theorem 4.8 and Corollary 4.12).

As is implied above, our main tool is a partition of \mathfrak{S}_n into sets $\mathcal{W}(\mathcal{S}, h)$ defined by particular subsets \mathcal{S} in the type A root system, called subsets of Weyl type (with respect to h); cf. Definition 2.8 below. This partition was introduced by Sommers–Tymoczko in [11] and subsequently used by the second author in [10] to prove that the Betti numbers of regular Hessenberg varieties are palindromic in all Lie types. Lemma 2.11, proven in Appendix A below by Michael Zeng, proves that each set $\mathcal{W}(\mathcal{S}, h)$ is a weak Bruhat interval. Let $w_{\mathcal{S}} \in \mathcal{W}(\mathcal{S}, h)$ denote the maximal element in this interval. Theorem 4.8 below shows that $\Omega_{w_{\mathcal{S}}, h}^T = \Omega_{w_{\mathcal{S}}}^T$. Thus the notion of reachability, which defines a Hessenberg variation of the partial order in (1.1), can be characterized using Bruhat order.

Example 4.16 below shows that $\Omega_{w_{\mathcal{S}}, h}$ is not simply a union of the Hessenberg Schubert cells indexed by $u \geq w_{\mathcal{S}}$. In other words, our Theorem 4.8 does not yield a complete description of the Hessenberg Schubert variety $\Omega_{w_{\mathcal{S}}, h}$.

We expect that our interpretation of the results of Cho–Hong–Lee in the language of subsets of Weyl type and Bruhat order will yield further insights into the geometry and combinatorics of Hessenberg varieties. We hope that this interpretation will shed light on the still-unsolved problem of fully characterizing the closures of (opposite) Hessenberg Schubert cells. We leave this problem for future work.

2. Background

2.1. Hessenberg Varieties and Hessenberg Schubert cells. Hessenberg varieties in Lie type A are subvarieties of the (full) flag variety $GL_n(\mathbb{C})/B$ where B is the Borel subgroup of upper triangular matrices in $GL_n(\mathbb{C})$. Let $G = GL_n(\mathbb{C})$ and let B_- denote the Borel subgroup of lower triangular matrices. The following two cell decompositions of G/B (both called a Bruhat decomposition of G/B) are well-studied:

$$(2.1) \quad G/B = \bigsqcup_{w \in \mathfrak{S}_n} X_w^\circ = \bigsqcup_{w \in \mathfrak{S}_n} \Omega_w^\circ$$

where $X_w^\circ := BwB/B$ is the **Schubert cell** and $\Omega_w^\circ = B_-wB/B$ is the **opposite Schubert cell**. The closure $X_w := \overline{X_w^\circ}$ (respectively $\Omega_w := \overline{\Omega_w^\circ}$) is called the **Schubert variety** (respectively **opposite Schubert variety**) for $w \in \mathfrak{S}_n$. It is an important and well-known fact that

$$(2.2) \quad X_w = \bigsqcup_{u \leq w} X_u^\circ \quad \text{and} \quad \Omega_w = \bigsqcup_{u \geq w} \Omega_u^\circ$$

where \leq denotes the Bruhat order on \mathfrak{S}_n defined in (1.1).

We denote the root system of $gl_n(\mathbb{C})$ by $\Phi = \{t_i - t_j \mid 1 \leq i \neq j \leq n\}$ with positive roots

$\Phi^+ = \{t_i - t_j \in \Phi \mid i < j\}$, negative roots $\Phi^- = \{t_i - t_j \in \Phi \mid i > j\}$, and simple positive roots $\Delta = \{t_i - t_{i+1} \mid 1 \leq i \leq n-1\}$. Given $w \in \mathfrak{S}_n$ the **inversion set** of w is

$$N(w) := \{t_i - t_j \in \Phi^+ \mid w(t_i - t_j) \in \Phi^-\} = \Phi^+ \cap w^{-1}(\Phi^-).$$

We set $\ell(w) := |N(w)| = |\{i < j \mid w(i) > w(j)\}|$. It is known that $X_w^\circ \simeq \mathbb{C}^{\ell(w)}$ and $\Omega_w^\circ \simeq \mathbb{C}^{N-\ell(w)}$ where $N = \sum_{i=1}^{n-1} (n-i) = \dim_{\mathbb{C}} GL_n(\mathbb{C})/B$.

A Hessenberg variety in G/B is specified by two pieces of data: a **Hessenberg function**, that is, a nondecreasing function $h : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$ such that $h(i) \geq i$ for all i , and a choice of an element \mathbf{X} in $gl(n, \mathbb{C})$. We frequently write a Hessenberg function by listing its values in sequence, i.e., $h = (h(1), h(2), \dots, h(n))$. The **Hessenberg variety** associated to the linear operator \mathbf{X} and Hessenberg function h is defined as

$$(2.3) \quad \mathcal{Hess}(\mathbf{X}, h) = \{gB \mid \mathbf{X}g_i \in \text{span}_{\mathbb{C}}\{g_1, \dots, g_{h(i)}\} \text{ for all } i \in \{1, 2, \dots, n\}\}$$

where g_1, \dots, g_n denote the columns of $g \in GL_n(\mathbb{C})$. In this paper, we focus on the case when \mathbf{X} is a regular semisimple operator \mathbf{S} (i.e., diagonalizable with distinct eigenvalues); more specifically, we fix \mathbf{S} to be a diagonal matrix with distinct eigenvalues. We refer to the corresponding Hessenberg variety $\mathcal{Hess}(\mathbf{S}, h)$ as a **regular semisimple Hessenberg variety**. It was established in [5] that $\mathcal{Hess}(\mathbf{S}, h)$ is a smooth, irreducible variety of dimension $N_h = \sum_{i=1}^{n-1} (h(i) - i)$.

For each $w \in \mathfrak{S}_n$ we consider the **Hessenberg Schubert cell**, defined as

$$X_{w,h}^\circ = X_w^\circ \cap \mathcal{Hess}(\mathbf{S}, h) = BwB/B \cap \mathcal{Hess}(\mathbf{S}, h)$$

and also the **opposite Hessenberg Schubert cell**, defined as

$$(2.4) \quad \Omega_{w,h}^\circ = \Omega_w^\circ \cap \mathcal{Hess}(\mathbf{S}, h) = B_-wB/B \cap \mathcal{Hess}(\mathbf{S}, h).$$

From this, we obtain decompositions of the regular semisimple Hessenberg variety

$$(2.5) \quad \mathcal{Hess}(\mathbf{S}, h) = \bigsqcup_{w \in \mathfrak{S}_n} X_{w,h}^\circ = \bigsqcup_{w \in \mathfrak{S}_n} \Omega_{w,h}^\circ$$

where $X_{w,h}^\circ \simeq \mathbb{C}^{\ell_h(w)}$ and $\Omega_{w,h}^\circ \simeq \mathbb{C}^{N_h - \ell_h(w)}$ for

$$(2.6) \quad \ell_h(w) := |\{i < j \mid w(i) > w(j) \text{ and } j \leq h(i)\}|.$$

When $h = (n, n, \dots, n)$ then $\mathcal{Hess}(\mathbf{S}, h) = G/B$ and we recover the Bruhat decomposition of $GL_n(\mathbb{C})/B$ from (2.1). We now define the **Hessenberg Schubert variety** for $w \in \mathfrak{S}_n$ to be $X_{w,h} := \overline{X_{w,h}^\circ}$ and the **opposite Hessenberg Schubert variety** to be $\Omega_{w,h} := \overline{\Omega_{w,h}^\circ}$. Our main result gives a combinatorial characterization of $\Omega_{w,h}$ for certain permutations w .

REMARK 2.7. One may define the Hessenberg Schubert cells and opposite Hessenberg Schubert cells in the language of Białyński–Birula strata, as in [3]. However, our definition is equivalent, and the Morse-theoretic point of view is not necessary for our purposes.

2.2. Subsets of Weyl type and acyclic orientations. As mentioned above, one of the contributions of this note is to introduce the theory of subsets of Weyl type into the study of Hessenberg Schubert closure relations. We briefly recall the relevant terminology and results.

Let $h : [n] \rightarrow [n]$ be a Hessenberg function. Then h determines a subset of Φ^+ defined by

$$\Phi_h^+ = \{t_i - t_j \in \Phi \mid i < j \text{ and } j \leq h(i)\}$$

and similarly, we let $\Phi_h^- = \{t_j - t_i \mid i < j \text{ and } j \leq h(i)\} \subseteq \Phi^-$. Note that $\ell_h(w) = |N(w) \cap \Phi_h^+|$ where ℓ_h is the Hessenberg length function defined in (2.6) above.

DEFINITION 2.8. Given a subset $S \subseteq \Phi_h^+$ we say that S is **Φ_h^+ -closed** if for all $\alpha, \beta \in S$ such that $\alpha + \beta \in \Phi_h^+$, then $\alpha + \beta \in S$ as well. Given such a subset $S \subseteq \Phi_h^+$, we say that S is a **subset of Weyl type (with respect to h)** if both S and its complement $\Phi_h^+ \setminus S$ are Φ_h^+ -closed. Denote the set of all subsets $S \subseteq \Phi_h^+$ of Weyl type (with respect to h) by \mathcal{W}_h .

It is a well known theorem of Kostant [9, Prop. 5.10] that $S \subseteq \Phi^+$ is a subset of Weyl type if and only if $S = N(w)$ for some $w \in \mathfrak{S}_n$. Sommers and Tymoczko generalized that result to the setting of subsets of Weyl type in Φ_h^+ . The following summarizes their results from [11] in the form most useful for our purposes.

Theorem 2.9 (Sommers–Tymoczko [11]). *Let $h : [n] \rightarrow [n]$ be a Hessenberg function and $S \in \mathcal{W}_h$.*

- (1) *There exists $w \in \mathfrak{S}_n$ such that $S = N(w) \cap \Phi_h^+$, and S is a subset of Weyl type with respect to h if and only if it is of this form.*
- (2) *There exists a unique element $z_S \in \mathfrak{S}_n$ satisfying both $S = N(z_S) \cap \Phi_h^+$ and $z_S^{-1}(-\Delta) \cap \Phi^+ \subseteq \Phi_h^+$.*
- (3) *$N(z_S) \subseteq N(y)$ for any $y \in \mathfrak{S}_n$ with $S \subseteq N(y)$.*

Given a fixed $S \in \mathcal{W}_h$, we now consider

$$\mathcal{W}(S, h) := \{w \in \mathfrak{S}_n \mid N(w) \cap \Phi_h^+ = S\} \subseteq \mathfrak{S}_n$$

i.e., $\mathcal{W}(S, h)$ is the set of permutations whose associated subset of Weyl type is exactly S . Note that $\mathcal{W}(S, h)$ is always non-empty for any $S \in \mathcal{W}_h$ by Theorem 2.9, and we obtain a partition $\mathfrak{S}_n = \bigsqcup_{S \in \mathcal{W}_h} \mathcal{W}(S, h)$. Recall that (left) **weak Bruhat order** is the partial order on \mathfrak{S}_n defined by

$$(2.10) \quad u \leq_L v \text{ if } v = s_{i_1} \cdots s_{i_k} u \text{ for simple reflections } s_{i_1}, \dots, s_{i_k} \text{ such that } \ell(v) = \ell(u) + k.$$

The weak Bruhat order is stronger than Bruhat order in the sense that $u \leq_L v$ implies $u \leq v$ for all $u, v \in \mathfrak{S}_n$. Note that $u \leq_L v$ if and only if $N(u) \subseteq N(v)$ [2, Prop. 3.1.3]. The following lemma tells us that $\mathcal{W}(S, h)$ is a weak Bruhat interval. A proof can be found in Appendix A.

Lemma 2.11. *Let $h : [n] \rightarrow [n]$ be a Hessenberg function and $S \in \mathcal{W}_h$. There exist elements $z_S, w_S \in \mathcal{W}(S, h)$ such that $\mathcal{W}(S, h)$ is precisely the weak (left) Bruhat interval*

$$[z_S, w_S]_L := \{v \in \mathfrak{S}_n \mid z_S \leq_L v \leq_L w_S\}.$$

In order to apply the results of [3] needed below, we now introduce a graph Γ_h uniquely determined by a Hessenberg function h .

DEFINITION 2.12. Let $h : [n] \rightarrow [n]$ be a Hessenberg function. The **incomparability graph** $\Gamma_h = (V(\Gamma_h), E(\Gamma_h))$ is the graph on vertex set $V(\Gamma_h) = [n]$ with edges $E(\Gamma_h) := \{(i, j) \mid j < i \text{ and } i \leq h(j)\}$.

EXAMPLE 2.13. The incomparability graph Γ_h for $h = (2, 4, 4, 4)$ and $h = (3, 4, 5, 5, 5)$ are given below.



The incomparability graph for h plays a key role in the results of [6] and also appears in [3, 4] (with the notation $G_{e,h}$). An **acyclic orientation** o of Γ_h is an assignment of a direction (i.e. orientation) to each edge $e \in E(\Gamma_h)$ such that the resulting oriented graph contains no directed cycles. Given $S \in \mathcal{W}_h$, we obtain an orientation $o_h(S)$ of Γ_h defined by the rule

$$(2.14) \quad j \longleftarrow i \quad \text{for } \{i, j\} \in E(\Gamma_h) \text{ with } j < i \text{ if and only if } t_j - t_i \in S.$$

In other words, $o_h(S)$ is obtained by orienting edges corresponding to the roots in S to the left, and oriented all other edges to the right.

EXAMPLE 2.15. Let $n = 4$. Consider the following acyclic orientations, the first of Γ_h for $h = (3, 4, 4, 4)$, and the second of Γ_h for $h = (2, 3, 4, 4)$.



The acyclic orientation on the LHS is $o_{(3,4,4,4)}(\{t_2 - t_3, t_1 - t_3\})$ and the one on the RHS is $o_{(2,3,4,4)}(\{t_1 - t_2\})$.

The following observation yields a bijection between acyclic orientations and subsets of Weyl type.

Lemma 2.16. *The set of all acyclic orientations of Γ_h is precisely $\{o_h(S) \mid S \in \mathcal{W}_h\}$.*

Sketch of Proof. Consider first the case in which $h = (n, n, \dots, n)$ so $\Gamma_h = K_n$ is the complete graph on n vertices. Since it is the complete graph, the set of edges is the set of all pairs $\{i, j\}$ with $j < i$. As in (2.14) an orientation of $\Gamma_h = K_n$ uniquely corresponds to the subset of edges which point to the left, and we may view this set as a subset S of the positive roots Φ^+ . Since Γ_h is the complete graph, it is straightforward to see that such an orientation will be acyclic if and only if both S and its complement $\Phi^+ \setminus S$ is closed under addition, i.e., S is a subset of Φ^+ of Weyl type. By the above-mentioned theorem of Kostant, subsets of Φ^+ correspond uniquely to permutations via the identification $w \mapsto N(w)$, so there are $n!$ acyclic orientations, as claimed.

Returning to the case of an arbitrary Hessenberg function, note that Γ_h is a subgraph of K_n . Let $S \in \mathcal{W}_h$. By Theorem 2.9 there exists $w \in \mathfrak{S}_n$ such that $S = N(w) \cap \Phi_h^+$. The orientation $o_h(S)$ defined in (2.14) is the orientation induced by the that of w on K_n as in the previous paragraph. This shows that $o_h(S)$ is an acyclic orientation. Since every acyclic orientation of Γ_h is the restriction of an acyclic orientation on K_n it follows that every acyclic orientation is of this form. \square

REMARK 2.17. In [4], the authors define and study an equivalence class of permutations. In their notation, the equivalence class $[w]_h$ from [4, Definition 3.4] is precisely the set

$\mathcal{W}(\mathcal{S}, h)$ above for $\mathcal{S} = N(w) \cap \Phi_h^+$. We also note that in [3, 4] the authors use the language of subgraphs $G_{w,h}$ of Γ_h for different permutations w , but this data can be equivalently characterized by acyclic orientations, which is what we choose to do in this manuscript.

The relation between $\mathcal{W}(\mathcal{S}, h)$ (and specifically the maximal element $w_{\mathcal{S}}$ of $\mathcal{W}(\mathcal{S}, h)$) to the acyclic orientation $o_h(\mathcal{S})$ is developed further in the next section.

3. Reachability

A key contribution of this manuscript is to connect the work of Cho, Hong, and Lee in [3]—in which they use the combinatorial notion of reachability to study Hessenberg Schubert cells—to the theory of subsets of Weyl type. In this section, we make this connection precise. The essential result is Proposition 3.13, which is the technical engine driving the proof of our main theorem (Theorem 4.8).

We begin with the definition of reachability, taken from [3], relating two vertices i and j on the incompatibility graph Γ_h .

DEFINITION 3.1. Let $h : [n] \rightarrow [n]$ be a Hessenberg function and Γ_h its associated incomparability graph, equipped with an acyclic orientation $o_h(\mathcal{S})$. Suppose $i > j$. We say that i is **reachable from j with respect to \mathcal{S} (or $o_h(\mathcal{S})$)** if $i = v_m$ and $j = v_0$ and there exists a sequence of vertices $j = v_0 < v_1 < \dots < i = v_m$ of Γ_h such that there is an oriented edge from each v_ℓ to $v_{\ell+1}$, i.e., there is a sequence of oriented edges of the form $j = v_0 \longrightarrow v_1 \longrightarrow \dots \longrightarrow v_{m-1} \longrightarrow v_m = i$ in Γ_h (equipped with the orientation $o_h(\mathcal{S})$). We allow m to be 0, that is, j is always reachable from j .

We say that a vertex k in an oriented graph is a **source** (with respect to the given orientation) if all edges adjacent to k point “out” of k , i.e., each such edge is of the form $k \longrightarrow i$ for all i adjacent to k . Since our incomparability graphs have vertices labelled by sets of positive integers $\{1, 2, \dots, n\}$, we say that a vertex k is the **largest source** if k is a source and moreover, if j is another source, then $k > j$. The next lemma shows that any vertex larger than the largest source k is reachable from k .

Lemma 3.2. *Let $h : [n] \rightarrow [n]$ be a Hessenberg function, Γ_h its associated incomparability graph, and $\mathcal{S} \in \mathcal{W}_h$. Suppose that k is the largest source of Γ_h with respect to the acyclic orientation $o_h(\mathcal{S})$ and i is a vertex with $i > k$. Then i is reachable from k .*

Proof. To prove the claim of the lemma, it suffices to show that the set


$$(3.3) \quad \{i > k \mid i \text{ is not reachable from } k\}$$

is empty. For the sake of obtaining a contradiction, suppose not, and consider the oriented subgraph Γ' induced by the vertices in set (3.3). Since the original oriented graph Γ_h is acyclic, so is the subgraph. Any acyclic orientation must have a source, so there exists a vertex i' which is a source of Γ' . We claim that the vertex i' must also be a source of the original oriented graph Γ_h . To see this, we must show that any edge $\{j, i'\} \in E(\Gamma_h)$ has orientation $i' \longrightarrow j$. To argue this, we first observe that $i' > h(k)$, because if $i' \leq h(k)$ then by definition of Γ_h there would be an edge from k to i' , and since k is a source by assumption, the edge between k and i' would be oriented as $k \longrightarrow i'$, making i' reachable from k . This

contradicts the assumption that i' is in the set (3.3). This implies that there is no edge in Γ_h between i' and any vertex j with $j \leq k$. Thus we may now assume without loss of generality that $j > k$. We take cases.

Suppose that $k < j < i'$. If there is no edge between j and i' in Γ_h then there is nothing to prove, so suppose there is an edge. If j is not reachable by k , then j is in the set (3.3) and we have assumed that i' is a source of the subgraph, so we must have $i' \rightarrow j$. On the other hand, if j is reachable by k , then by reasoning similar to the above, we must also have $j \leftarrow i'$ since otherwise i' would be reachable by k .

Next suppose $j > i'$. Again, if there is no edge between j and i' then there is nothing to prove, so we may suppose $\{i', j\}$ is an edge of Γ_h . If j is not reachable by k then by the same reasoning as above we already know the edge is oriented as $i' \rightarrow j$, so suppose j is reachable by k . Recall that we wish to show $i' \rightarrow j$ in $o_h(\mathcal{S})$. Suppose for the sake of contradiction that $i' \leftarrow j$ instead. Since j is reachable from k , there exists a sequence $v_0 < v_1 < \dots < v_m$ of vertices such that $k = v_0 \rightarrow v_1 \rightarrow \dots \rightarrow v_{m-1} \rightarrow v_m = j$ in $o_h(\mathcal{S})$. Let v_ℓ be the largest vertex in that list such that $v_\ell < i'$. Since $\{v_\ell, v_{\ell+1}\}$ is an edge in Γ_h and $i' < v_{\ell+1}$ we conclude that $\{v_\ell, i'\} \in E(\Gamma_h)$. Since v_ℓ is reachable from k , by the same reasoning as above we know that $v_\ell \leftarrow i'$ and we may visualize the graph as



in $o_h(\mathcal{S})$, which shows that there is a cycle starting and ending at i' , contradicting the fact that $o_h(\mathcal{S})$ is an acyclic orientation. Thus we must have $i' \rightarrow j$.

We have now shown that i' is a source in Γ_h , contradicting that k is the largest source. Therefore (3.3) is indeed empty as was to be shown. \square

The next lemma shows that reachability implies an inequality among entries in the one-line notation of w for $w \in \mathcal{W}(\mathcal{S}, h)$.

Lemma 3.4. *Let $j \leq i$ be vertices in Γ_h and $\mathcal{S} \in \mathcal{W}_h$. If i is reachable from j with respect to $o_h(\mathcal{S})$, then $w(j) \leq w(i)$ for all $w \in \mathcal{W}(\mathcal{S}, h)$.*

Proof. Let $w \in \mathcal{W}(\mathcal{S}, h)$. Suppose i is reachable from j so we have a sequence of vertices $v_0 < v_1 < \dots < v_m$ such that $j = v_0 \rightarrow v_1 \rightarrow \dots \rightarrow v_{m-1} \rightarrow v_m = i$ in $o_h(\mathcal{S})$. Thus for each ℓ such that $1 \leq \ell \leq m$, we have $t_{v_{\ell-1}} - t_{v_\ell} \in \Phi_h^+ \setminus \mathcal{S}$ and since $\mathcal{S} = N(w) \cap \Phi_h^+$, we get $t_{v_{\ell-1}} - t_{v_\ell} \notin N(w)$. This means $w(v_{\ell-1}) \leq w(v_\ell)$ for all $1 \leq \ell \leq m$, and by putting the inequalities together we obtain $w(j) \leq w(i)$, as desired. \square

It also turns out that the location of 1 in the one-line notation of $w \in \mathcal{W}(\mathcal{S}, h)$ is significant.

Lemma 3.5. *Let $h : [n] \rightarrow [n]$ be a Hessenberg function and $\mathcal{S} \in \mathcal{W}_h$. If $w \in \mathcal{W}(\mathcal{S}, h)$, then $w^{-1}(1)$ is a source of Γ_h equipped with the orientation $o_h(\mathcal{S})$.*

Proof. Suppose $j = w^{-1}(1)$. It follows directly from the definition of the inversion set that

$$\{t_1 - t_j, t_2 - t_j, \dots, t_{j-1} - t_j\} \subseteq N(w)$$

since $w(j) = 1$ is strictly smaller than any $w(1), w(2), \dots, w(j-1)$. By similar reasoning, since $w(j) = 1$ is also smaller than $w(j+1), \dots, w(n)$, we have

$$\{t_j - t_{j+1}, t_j - t_{j+2}, \dots, t_j - t_n\} \subseteq \Phi^+ \setminus N(w).$$

As $w \in \mathcal{W}(\mathcal{S}, h)$ by hypothesis, we have that $\mathcal{S} = N(w) \cap \Phi_h^+$. From the definition of $o_h(\mathcal{S})$ in (2.14) we see that the edges $\{k, j\}$ with $k < j$ must be oriented to the left and the edges for $k > j$ are oriented toward the right. Thus j is a source, as desired. \square

We now give an inductive construction using the Hessenberg function h and graph Γ_h that will be useful in what follows. Let $k \in [n]$. Consider the smaller graph on $n-1$ vertices which is obtained from Γ_h by deleting vertex k and all adjacent edges to k , and for convenience in our arguments below, relabeling the vertex set to be $\{2, 3, \dots, n\}$ (so $\{1, 2, \dots, k-1\}$ gets relabelled as $\{2, 3, \dots, k\}$ respectively, and the labels of $\{k+1, \dots, n\}$ are unchanged). We let $h^{(k)} : \{2, 3, \dots, n\} \rightarrow \{2, 3, \dots, n\}$ denote the Hessenberg function whose associated graph $\Gamma_{h^{(k)}}$ is precisely the graph just described. Alternatively, suppose the Hessenberg function h is visualized as a collection of boxes in an $n \times n$ array where the (i, j) -th box (in row i and column j) is said to be in the collection if $i \leq h(j)$. Then the collection of boxes in the $(n-1) \times (n-1)$ array corresponding to $h^{(k)}$ is obtained from that of h by deleting the k -th row and column.

Set $\Psi := \{t_i - t_j \mid i \neq j \text{ and } i, j \in \{2, \dots, n\}\}$. Then Ψ is a root system of type A_{n-2} , and the only difference between Ψ and the standard root system is that we have shifted the index set to be $\{2, 3, \dots, n\}$ instead of $\{1, 2, \dots, n-1\}$. Fix $k \in [n]$. In analogy with how we defined Φ_h^+ above, let us define $\Psi_{h^{(k)}}^+ := \{t_i - t_j \in \Psi \mid i < j \text{ and } j \leq h^{(k)}(i)\}$. From the description of the graph $\Gamma_{h^{(k)}}$ it is not hard to check that

$$(3.6) \quad \Psi_{h^{(k)}}^+ = u_k(\Phi_h^+) \cap \Psi$$

where $u_k := s_1 s_2 \cdots s_{k-1} \in \mathfrak{S}_n$ (in one-line notation we have $u_k = [2, 3, \dots, k, 1, k+1, k+2, \dots, n]$). Here we take $u_1 := e \in \mathfrak{S}_n$. Given $\mathcal{S} \in \mathcal{W}_h$, we can consider the orientation or $\Gamma_{h^{(k)}}$ induced by $o_h(\mathcal{S})$, which must necessarily be acyclic. Let $\mathcal{S}' \subseteq \Psi_{h^{(k)}}^+$ denote the corresponding subset of Weyl type. By (3.6) we have

$$\mathcal{S}' = u_k(\mathcal{S}) \cap \Psi.$$

The next lemma relates certain elements in $\mathcal{W}(\mathcal{S}, h)$ with those in $\mathcal{W}(\mathcal{S}', h^{(k)})$.

Lemma 3.7. *Let $y \in \langle s_2, s_3, \dots, s_{n-1} \rangle$ and k be a source of Γ_h with respect to $o_h(\mathcal{S})$ for $\mathcal{S} \in \mathcal{W}_h$. Let u_k be as above and $w = yu_k \in \mathfrak{S}_n$. Then $y \in \mathcal{W}(\mathcal{S}', h^{(k)})$ if and only if $w \in \mathcal{W}(\mathcal{S}, h)$.*

Proof. First suppose $w \in \mathcal{W}(\mathcal{S}, h)$, so $N(w) \cap \Phi_h^+ = \mathcal{S}$. On the other hand, since the decomposition $w = yu_k$ satisfies $\ell(w) = \ell(y) + \ell(u_k)$ (note u_k is a minimal coset representative of $\mathfrak{S}_{n-1} \setminus \mathfrak{S}_n$) we have $N(w) = N(u_k) \sqcup u_k^{-1}N(y)$ (cf. for instance [7, Section 1.7]). Combining these facts we obtain

$$(3.8) \quad \mathcal{S} = (N(u_k) \sqcup u_k^{-1}N(y)) \cap \Phi_h^+ \Leftrightarrow u_k(\mathcal{S}) = (u_kN(u_k) \sqcup N(y)) \cap u_k(\Phi_h^+)$$

and thus

$$\mathcal{S}' = u_k(\mathcal{S}) \cap \Psi = N(y) \cap \Psi \cap u_k\Phi_h^+ = N(y) \cap \Psi_{h^{(k)}}^+$$

where the first and third equalities follow from (3.6) and the second equality follows from

an explicit computation of $u_k N(u_k)$ which shows that $u_k N(u_k) \cap \Psi = \emptyset$. This proves $y \in \mathcal{W}(\mathcal{S}', h^{(k)})$.

To see the other direction, suppose $y \in \mathcal{W}(\mathcal{S}', h^{(k)})$. We want to show $w \in \mathcal{W}(\mathcal{S}, h)$, for which it suffices to prove (3.8). Since k is a source we have

$$\mathcal{S} = \{t_j - t_i \in \mathcal{S} \mid i, j \in [n] \setminus \{k\}\} \sqcup \{t_j - t_k \mid j < k, k \leq h(j)\}.$$

Since $u_k([n] \setminus \{k\}) = \{2, \dots, n\}$ we have

$$(3.9) \quad \{t_j - t_i \in \mathcal{S} \mid i, j \in [n] \setminus \{k\}\} = u_k^{-1}(\mathcal{S}')$$

and now the claim follows from the observation that (3.6) implies

$$u_k^{-1}(\mathcal{S}') = u_k^{-1}\left(N(y) \cap \Psi_{h^{(k)}}^+\right) = u_k^{-1}N(y) \cap \Phi_h^+$$

and that $N(u_k) \cap \Phi_h^+ = \{t_j - t_k \mid j < k, k \leq h(j)\}$. This completes the proof. \square

The next lemma is a kind of converse to Lemma 3.5.

Lemma 3.10. *Let $\mathcal{S} \in \mathcal{W}_h$. If k is a source of Γ_h with respect to the orientation $o_h(\mathcal{S})$, then there exists $w \in \mathcal{W}(\mathcal{S}, h)$ such that $w^{-1}(1) = k$.*

Proof. Since $\mathcal{W}(\mathcal{S}', h^{(k)})$ is non-empty, there exists $y \in \mathcal{W}(\mathcal{S}', h^{(k)}) \subseteq \langle s_2, \dots, s_{n-1} \rangle$. Consider $w = yu_k$. By Lemma 3.7, $w \in \mathcal{W}(\mathcal{S}, h)$, and by construction $w(k) = 1$. \square

The relationship is even tighter between the largest source and the maximal element $w_{\mathcal{S}}$ of $\mathcal{W}(\mathcal{S}, h)$.

Lemma 3.11. *Let $h : [n] \rightarrow [n]$ be a Hessenberg function and $\mathcal{S} \in \mathcal{W}_h$. Then the vertex k is the largest source of Γ_h with respect to $o_h(\mathcal{S})$ if and only if $w_{\mathcal{S}}^{-1}(1) = k$.*

Proof. Suppose k is the largest source of Γ_h with respect to $o_h(\mathcal{S})$. By Lemma 3.10 there exists $w \in \mathcal{W}(\mathcal{S}, h)$ such that $w^{-1}(1) = k$ so we may write $w = yu_k$ for some $y \in \langle s_2, \dots, s_{n-1} \rangle$. Let $j := w_{\mathcal{S}}^{-1}(1)$ or equivalently, $w_{\mathcal{S}}(j) = 1$. Then by Lemma 3.5 we know j must be a source, and since k is the largest source, we conclude $j \leq k$. To complete the argument that $j = w_{\mathcal{S}}^{-1}(1) = k$ we show that $k \leq j$. To see this, write $w_{\mathcal{S}} = y_{\mathcal{S}}u_j$ for some $y_{\mathcal{S}} \in \langle s_2, \dots, s_n \rangle$, which is possible since $j = w_{\mathcal{S}}^{-1}(1)$. We know that $w_{\mathcal{S}}$ is the maximal element of $\mathcal{W}(\mathcal{S}, h)$, so


$$w \leq w_{\mathcal{S}} \Rightarrow yu_k \leq y_{\mathcal{S}}u_j \Rightarrow u_k \leq u_j \Rightarrow k \leq j$$

where the second implication follows from the fact that the map from \mathfrak{S}_n to the minimal coset representatives $\{u_1 := e, u_2, \dots, u_n\}$ of the subgroup $\langle s_2, s_3, \dots, s_n \rangle$ is order-preserving [2, Prop. 2.5.1]. Thus $j = w_{\mathcal{S}}^{-1}(1) = k$ as desired. The converse follows by similar reasoning. \square

The lemma and other inductive structure established above yield a simple way to compute $w_{\mathcal{S}}$ given the corresponding acyclic orientation.

EXAMPLE 3.12. Let $n = 4$. We consider as in Example 2.15 the orientation $o_{(3,4,4,4)}(\{t_2 - t_3, t_1 - t_3\})$ of graph Γ_h for $h = (3, 4, 4, 4)$. Using the graph pictured in that example,

Lemma 3.11 tells us that $w_S(3) = 1$. Deleting 3 and all adjacent edges tells us that $w_S(1) = 2$. Continuing in this way gives the sequence of graphs:



and $w_S = [2, 3, 1, 4]$.

We can now prove one of our important technical results, which characterizes reachability in terms of the maximal elements w_S for $S \in \mathcal{W}_h$. This reformulation is the tool which allows us to prove our characterization of T -fixed points in the opposite Hessenberg Schubert variety in the next section.

Proposition 3.13. *Let $h : [n] \rightarrow [n]$ be a Hessenberg function and $S \in \mathcal{W}_h$. Suppose $j \leq i$. Then i is reachable from j with respect to S if and only if $w_S(j) \leq w_S(i)$.*

Proof. First suppose that $j \leq i$ and i is reachable from j . We wish to show that $w_S(j) \leq w_S(i)$. This follows immediately from Lemma 3.4.

So now suppose $j \leq i$ and that $w_S(j) \leq w_S(i)$. We need to show that i is reachable from j . We proceed to prove the contrapositive statement by an induction on n . When $n = 1$, we have $j = i = 1$ and $w_S = e$ and the claim is trivial. Now suppose $n \geq 2$ and that the claim is true for $n - 1$. Note that we may assume $j \neq i$ since if $j = i$ then i is reachable from j by convention and the claim is immediate. So suppose $j < i$ and additionally suppose that i is not reachable from j . Let $k := w_S^{-1}(1)$ and write $w_S = y_S u_k$ for some $y_S \in \langle s_2, \dots, s_{n-1} \rangle$. If $i = k$, then our claim follows immediately since $w_S(k) = w_S(i) = 1 < w_S(j)$. Next, notice that by Lemma 3.11, k is the largest source of Γ_h with respect to $o_h(S)$. If $j = k$, then by Lemma 3.2 we would have that i is reachable from j , but we have assumed that i is not reachable from j , so we conclude $j \neq k$.

We now have that $i \neq k$, $j \neq k$ and $j < i$. Consider the graph $\Gamma_{h^{(k)}}$ obtained from Γ_h with acyclic orientation $o_{h^{(k)}}(S')$ induced by $o_h(S)$. By Lemma 3.7, since $w_S \in \mathcal{W}(S, h)$ we know $y_S \in \mathcal{W}(S', h^{(k)})$. We wish to show y_S is the maximal element in $\mathcal{W}(S', h^{(k)})$. To see this, suppose \tilde{y}_S is the maximal element. First observe $y_S \leq \tilde{y}_S$ since \tilde{y}_S is maximal. On the other hand, $\tilde{y}_S u_k \in \mathcal{W}(S, h)$ by Lemma 3.7 and since w_S is maximal, we conclude $\tilde{y}_S u_k \leq w_S = y_S u_k$. Now the properties of Bruhat order imply $\tilde{y}_S \leq y_S$. Hence $y_S = \tilde{y}_S$ and y_S is maximal in $\mathcal{W}(S', h^{(k)})$. Our assumptions imply that $i' = u_k(i)$ is not reachable from $j' = u_k(j)$. As $\Gamma_{h^{(k)}}$ is a Hessenberg graph on $n - 1$ vertices and y_S is maximal, we know $y_S(i') < y_S(j')$ by induction. Since $w_S = y_S u_k$, we conclude $w_S(i) < w_S(j)$ as desired. \square

As a first geometric application of the results in this section, we prove the following. Recall that from Lemma 2.11 we know that $\mathcal{W}(S, h)$ is a weak Bruhat interval of the form $[z_S, w_S]_L$. Moreover, from the definition (2.10) of weak Bruhat order, it follows that if $v \in \mathcal{W}(S, h)$ then $v \leq_L w_S$ so there exists an element $u \in \mathfrak{S}_n$ such that $w_S = u^{-1}v$ with $\ell(w_S) = \ell(v) + \ell(u)$.

Proposition 3.14. *Suppose $S \in \mathcal{W}_h$ and $v \in \mathcal{W}(S, h)$ and let $w_S = u^{-1}v$ for $u \in \mathfrak{S}_n$ where $\ell(w_S) = \ell(v) + \ell(u)$. Then*

$$\Omega_{v,h}^\circ = u \left(\Omega_{w_S}^\circ \cap \mathcal{Hess}(u^{-1} \mathfrak{S} u, h) \right).$$

Proof. Let b denote a lower triangular matrix with 1's on the diagonal. Recall that $wbB \in \Omega_w^\circ$ if and only if $b_{ij} = 0$ for all $i > j$ such that $w(i) < w(j)$, or equivalently, $b_{ij} = 0$ for all $t_j - t_i \in N(w)$. Our assumptions imply that $v \leq_L w_S$ so $N(v) \subseteq N(w_S)$.

Suppose $w_S bB \in \Omega_{w_S}^\circ$, so $b_{ij} = 0$ for all $t_j - t_i \in N(w_S)$ implying that $b_{ij} = 0$ for all $t_j - t_i \in N(v)$ also. Thus $uw_S bB = vbB \in \Omega_v^\circ$. We now have

$$\begin{aligned} u\Omega_{w_S}^\circ \subseteq \Omega_v^\circ &\Rightarrow u\Omega_{w_S}^\circ \cap \text{Hess}(\mathbf{S}, h) \subseteq \Omega_v^\circ \cap \text{Hess}(\mathbf{S}, h) \\ &\Rightarrow u\left(\Omega_{w_S}^\circ \cap \text{Hess}(u^{-1}\mathbf{S}u, h)\right) \subseteq \Omega_{v,h}^\circ. \end{aligned}$$

Here the second implication uses the fact that $\text{Hess}(\mathbf{S}, h) = u\text{Hess}(u^{-1}\mathbf{S}u, h)$.

To prove the opposite inclusion, suppose $vbB \in \Omega_{v,h}^\circ$. By [3, Corollary 3.9], $b_{ij} = 0$ for all $i > j$ such that i is not reachable from j . By Proposition 3.13, this implies $b_{ij} = 0$ for all $i > j$ such that $w_S(i) < w_S(j)$ so $w_S bB \in \Omega_{w_S}^\circ$. Furthermore, since $vbB \in \text{Hess}(\mathbf{S}, h)$ it follows immediately that $u^{-1}vbB = w_S bB \in \text{Hess}(u^{-1}\mathbf{S}u, h)$. Thus $u^{-1}vbB \in \Omega_{w_S}^\circ \cap \text{Hess}(u^{-1}\mathbf{S}u, h)$, as desired. \square

4. Connection to Bruhat Order

In this section we state and prove our main result, Theorem 4.8. To do so, we need some terminology and notation from the work of Cho, Hong, and Lee [3].

Let n be a positive integer and $1 \leq k \leq n$. Denote by $I_{k,n}$ the set of k -tuples of positive integers $\underline{i} = (i_1, i_2, \dots, i_k) \in \mathbb{Z}^k$ satisfying $1 \leq i_1 < i_2 < \dots < i_k \leq n$. Suppose we are given a permutation $w \in S_n$ and $\underline{i} \in I_{k,n}$. We can consider the k -tuple of not-necessarily-strictly-increasing integers $(w(i_1), w(i_2), \dots, w(i_k)) \in \mathbb{Z}^k$ and then re-order the entries in such a way that they are strictly increasing; we denote the result as $w \cdot \underline{i}$, and by construction, we have $w \cdot \underline{i} \in I_{k,n}$.

We also need a certain subset $J_{w,h,k}$ of $I_{k,n}$ associated to a permutation $w \in S_n$, a Hessenberg function $h : [n] \rightarrow [n]$, and an integer k such that $1 \leq k \leq n$. To define it, we need the following terminology from [3], which extends the notion of reachability (described in Section 3) to two subsets of $[n]$.

DEFINITION 4.1. Let $A = \{1 \leq a_1 < a_2 < \dots < a_r \leq n\}$ and $B = \{1 \leq b_1 < b_2 < \dots < b_r \leq n\}$ be two subsets of $[n]$ of the same cardinality r . We say A is **reachable** from B if there exists a permutation $\sigma \in S_r$ such that $a_{\sigma(i)}$ is reachable from b_i for all $1 \leq i \leq r$.

We now define $J_{w,h,k}$ as in [3, Equation (3.2)] in terms of reachability with respect to the acyclic orientation $o_h(\mathcal{S})$ determined by $\mathcal{S} = N(w) \cap \Phi_h^+$:

$$(4.2) \quad J_{w,h,k} := \{\underline{i} = (i_1, i_2, \dots, i_k) \in I_{k,n} \mid \{i_1, i_2, \dots, i_k\} \text{ is reachable from } \{1, 2, \dots, k\}\}.$$

With this notation in place, we can state the following result of Cho, Hong, and Lee [3, Theorem 3.5].

Theorem 4.3 (Cho–Hong–Lee). *Let $h : [n] \rightarrow [n]$ be a Hessenberg function and $w \in \mathfrak{S}_n$. Then*

$$(\Omega_{w,h})^T = \{u \in \mathfrak{S}_n \mid u \cdot (1, \dots, k) \in \{w \cdot (i_1, \dots, i_k) \mid (i_1, \dots, i_k) \in J_{w,h,k}\} \text{ for all } 1 \leq k \leq n-1\}.$$

Here and below we define and use the following partial order \leq on $I_{k,n}$:

$$(4.4) \quad (i_1, i_2, \dots, i_k) \geq (j_1, j_2, \dots, j_k) \text{ if and only if } i_\ell \geq j_\ell \text{ for all } \ell, 1 \leq \ell \leq k.$$

In order to obtain a description of $(\Omega_{w,h})^T$ in terms of Bruhat order using Theorem 4.3, we need the following lemma, which characterizes the set $J_{w,h,k}$ defined in (4.2) in terms of the maximal element w_S of $\mathcal{W}(S, h)$. This connects the discussion to that of Section 3 and allows us to use the results therein.

Lemma 4.5. *Suppose $S \in \mathcal{W}_h$. For all $w \in \mathcal{W}(S, h)$ and all k such that $1 \leq k \leq n-1$, we have*

$$J_{w,h,k} = \{(i_1, \dots, i_k) \in I_{k,n} \mid w_S \cdot (i_1, \dots, i_k) \geq w_S \cdot (1, \dots, k)\}$$

where $J_{w,h,k}$ is the set defined by (4.2).

Proof. By its definition in (4.2), and from the definition of reachability, $J_{w,h,k}$ consists of $(i_1, \dots, i_k) \in I_{k,h}$ such that there exists a permutation $\sigma \in \mathfrak{S}_k$ with the property that $i_{\sigma(\ell)}$ is reachable from ℓ for all $1 \leq \ell \leq k$ with respect to $o_h(S)$. On the other hand, by Proposition 3.13 this is equivalent to $w_S(i_{\sigma(\ell)}) \geq w_S(\ell)$ for all $1 \leq \ell \leq k$. The claim now follows from our definition of the notation $w_S \cdot (i_1, \dots, i_k) \geq w_S \cdot (1, \dots, k)$. \square

We also need the following description of the Bruhat order in \mathfrak{S}_n , which we now briefly recall (cf. [2, Propositions 2.4.8, 2.5.1, and Theorem 2.6.3]).

Lemma 4.6. *Let $w, v \in \mathfrak{S}_n$. Then $w \leq v$ in Bruhat order if and only if*

$$(4.7) \quad w \cdot (1, \dots, k) \leq v \cdot (1, \dots, k) \text{ for all } 1 \leq k \leq n-1.$$

We are now ready to state and prove the main result of this note.

Theorem 4.8. *Let $h : [n] \rightarrow [n]$ be a Hessenberg function and fix $S \in \mathcal{W}_h$. Then*

$$(4.9) \quad \Omega_{w_S, h}^T = [w_S, w_0],$$

where $[w_S, w_0]$ denotes the Bruhat interval of $w \in \mathfrak{S}_n$ such that $w_S \leq w$.

Proof. First, as $\Omega_{w_S, h}^{\circ} \subseteq \Omega_w \cap \text{Hess}(S, h)$ and the intersection $\Omega_w \cap \text{Hess}(S, h)$ is closed, we have $\Omega_{w,h} \subseteq \Omega_w \cap \text{Hess}(S, h)$. It follows immediately that $\Omega_{w_S, h}^T \subseteq [w_S, w_0]$.

To prove the opposite inclusion, suppose $u \in [w_S, w_0]$. By Lemma 4.6 we obtain

$$(4.10) \quad w_S \cdot (1, \dots, k) \leq u \cdot (1, \dots, k) \text{ for all } 1 \leq k \leq n-1.$$

As w_S is a permutation there exists $i_1, \dots, i_{n-1} \in [n]$ such that $w_S(i_\ell) = u(\ell)$ for all $1 \leq \ell \leq n-1$. Equation (4.10) now yields

$$w_S \cdot (i_1, \dots, i_k) = u \cdot (1, \dots, k) \geq w_S \cdot (1, \dots, k) \text{ for all } 1 \leq k \leq n-1.$$

Now $(i_1, \dots, i_k) \in J_{w_S, h, k}$ by Lemma 4.5 and therefore $u \in \Omega_{w_S, h}^T$ by Theorem 4.3. \square

Let $w_0 = [n, n-1, \dots, 2, 1]$ denote the longest element in \mathfrak{S}_n . By translating by w_0 , we easily obtain the analogous result for Hessenberg Schubert cells corresponding to the minimal element z_S of $\mathcal{W}(S, h)$.

Corollary 4.11. *Let $h : [n] \rightarrow [n]$ be a Hessenberg function and $\mathcal{S} \in \mathcal{W}_h$. Then*

$$X_{z_{\mathcal{S}}, h}^T = [e, z_{\mathcal{S}}],$$

where $[e, z_{\mathcal{S}}]$ denotes the Bruhat interval of $w \in \mathfrak{S}_n$ such that $w \leq z_{\mathcal{S}}$.

Proof. Let $\bar{\mathcal{S}} = \Phi_h^+ \setminus \mathcal{S} \in \mathcal{W}_h$. Since $w_0 B_- w_0^{-1} = B$ we have $w_0 \Omega_w^\circ = X_{w_0 w}^\circ$. Using the fact that $w_0 w_{\bar{\mathcal{S}}} = z_{\mathcal{S}}$ as in the proof of Corollary A.3 we obtain

$$w_0 \left(\Omega_{w_{\bar{\mathcal{S}}}}^\circ \cap \text{Hess}(w_0 \mathcal{S} w_0, h) \right) = X_{z_{\mathcal{S}}}^\circ \cap \text{Hess}(\mathcal{S}, h) = X_{z_{\mathcal{S}}, h}^\circ.$$

Since left multiplication respects closure relations in the flag variety, the corollary now follows by application of Theorem 4.8 to the Hessenberg Schubert variety $\Omega_{w_{\bar{\mathcal{S}}}}^\circ \cap \text{Hess}(w_0 \mathcal{S} w_0, h)$ and the fact that $w_{\bar{\mathcal{S}}} \leq w \Leftrightarrow z_{\mathcal{S}} \geq w_0 w$ since left multiplication by w_0 is an order reversing involution of \mathfrak{S}_n . \square

We can also give a reformulation of Theorem 4.3 in the language of Bruhat order.

Corollary 4.12. *Let $h : [n] \rightarrow [n]$ be a Hessenberg function and $\mathcal{S} \in \mathcal{W}_h$. Suppose $v \in \mathcal{W}(\mathcal{S}, h)$ and let $w_{\mathcal{S}} = u^{-1} v$ for $u \in \mathfrak{S}_n$ such that $\ell(w_{\mathcal{S}}) = \ell(v) + \ell(u)$. Then*

$$\Omega_{v, h}^T = u[w_{\mathcal{S}}, w_0].$$

Proof. Left multiplication respects closure relations in the flag variety, so by Proposition 3.14 we have $\Omega_{v, h} = u \left(\overline{\Omega_{w_{\mathcal{S}}}^\circ \cap \text{Hess}(u^{-1} \mathcal{S} u, h)} \right)$. Theorem 4.8 now yields

$$\left(\overline{\Omega_{w_{\mathcal{S}}}^\circ \cap \text{Hess}(u^{-1} \mathcal{S} u, h)} \right)^T = [w_{\mathcal{S}}, w_0]$$

and the corollary follows. \square

We conclude this section with a discussion of the geometry of the Hessenberg Schubert variety $\Omega_{w_{\mathcal{S}}, h}$. The results of Theorem 4.8 tell us that

$$(4.13) \quad \Omega_{w_{\mathcal{S}}, h}^T = \Omega_{w_{\mathcal{S}}}^T = (\text{Hess}(\mathcal{S}, h) \cap \Omega_{w_{\mathcal{S}}})^T$$

where the second equality follows from [5, Proposition 3]. Since

$$(4.14) \quad \dim \Omega_{w_{\mathcal{S}}, h} = \dim \Omega_{w_{\mathcal{S}}}^\circ = \dim (\Omega_{w_{\mathcal{S}}} \cap \text{Hess}(\mathcal{S}, h))$$

by definition, the equality (4.13) might lead one to hope that $\Omega_{w_{\mathcal{S}}, h} = \Omega_{w_{\mathcal{S}}} \cap \text{Hess}(\mathcal{S}, h)$. This is not true in general. More specifically, although $\Omega_{w_{\mathcal{S}}, h} = \Omega_{w_{\mathcal{S}}} \cap \text{Hess}(\mathcal{S}, h)$ holds in many cases (see Remark 4.17), we can conclude in general only that $\Omega_{w_{\mathcal{S}}, h}$ is an irreducible component of $\Omega_{w_{\mathcal{S}}} \cap \text{Hess}(\mathcal{S}, h) = \bigcup_{w \geq w_{\mathcal{S}}} \Omega_{w, h}^\circ$. We give more details below.

Let $w \in \mathfrak{S}_n$. As $\Omega_{w, h}$ is the closure of the affine cell $\Omega_{w, h}^\circ$, it is irreducible. Now (4.14) and fact that $\Omega_{w, h} \subseteq \Omega_w \cap \text{Hess}(\mathcal{S}, h)$ together imply that $\Omega_{w, h}$ is indeed an irreducible component of $\Omega_w \cap \text{Hess}(\mathcal{S}, h)$. It now follows that the equality

$$(4.15) \quad \Omega_{w, h} = \Omega_w \cap \text{Hess}(\mathcal{S}, h)$$

holds if and only if $\Omega_w \cap \text{Hess}(\mathcal{S}, h)$ is irreducible. We consider cases. If w is not the maximal element of $\mathcal{W}(\mathcal{S}, h)$ for some $\mathcal{S} \in \mathcal{W}_h$, then $\Omega_w \cap \text{Hess}(\mathcal{S}, h)$ is reducible, since $\Omega_{w, h}^T \subseteq [w, w_0] = (\Omega_w \cap \text{Hess}(\mathcal{S}, h))^T$ by Corollary 4.12. On the other hand, if $w = w_{\mathcal{S}}$, the

following example shows that $\Omega_{w_S} \cap \text{Hess}(\mathbf{S}, h)$ is, in general, still reducible.

EXAMPLE 4.16. Let $n = 4$ and $h = (3, 4, 4, 4)$. In this case, $\Phi_h^+ = \Phi^+ \setminus \{t_1 - t_4\}$ and $\dim \text{Hess}(\mathbf{S}, h) = 5$. Consider $\mathcal{S} = \{t_2 - t_3, t_1 - t_3\} \in \mathcal{W}_h$. In Example 3.12 above, it was shown that $w_{\mathcal{S}} = [2, 3, 1, 4]$. We also have

$$\dim(\Omega_{w_S} \cap \text{Hess}(\mathbf{S}, h)) = \dim(\Omega_{w_S, h}^\circ) = 5 - |\mathcal{S}| = 3.$$

Consider $w = [2, 3, 4, 1]$, which satisfies $w_{\mathcal{S}} < w$ so $\Omega_{w, h} \subseteq \Omega_{w_S} \cap \text{Hess}(\mathbf{S}, h)$. As

$$N(w) = \{t_3 - t_4, t_2 - t_4, t_1 - t_4\} \Rightarrow \ell_h(w) = |N(w) \cap \Phi_h^+| = 2$$

we have $\dim \Omega_{w, h}^\circ = 3$. Now the fact that $\Omega_{w_S, h}^\circ \cap \Omega_{w, h}^\circ = \emptyset$ and $\dim \Omega_{w, h} = \dim(\Omega_{w_S} \cap \text{Hess}(\mathbf{S}, h))$ implies $\Omega_{w, h}$ is an irreducible component of $\Omega_{w_S} \cap \text{Hess}(\mathbf{S}, h)$ with $\Omega_{w, h} \neq \Omega_{w_S, h}$. This shows $\Omega_{w_S} \cap \text{Hess}(\mathbf{S}, h)$ is reducible.

REMARK 4.17. When $h = (2, 3, 4, \dots, n, n)$, the corresponding regular semisimple Hessenberg variety is the toric variety associated to the Weyl chambers, also called the permutohedral variety. In this well-studied case, the equality from (4.15) holds for all w_S with $\mathcal{S} \in \mathcal{W}_h$. This follows from an application of [8, Theorem 3.10].

Appendix A Proof of Lemma 2.11 by Michael Zeng

We now present a proof of Lemma 2.11, written by Michael Zeng as part of an undergraduate research project with the second author. Let $w_0 = [n, n-1, \dots, 2, 1]$ denote the longest element of \mathfrak{S}_n . Consider the map

$$\varphi : \mathfrak{S}_n \rightarrow \mathfrak{S}_n, \varphi(w) = w_0 w.$$

Since

$$(A.1) \quad N(w_0 w) = \Phi^+ \setminus N(w)$$

for all $w \in \mathfrak{S}_n$ it follows that φ defines an order-reversing involution with respect to the weak order, that is, $u \leq_L v \Leftrightarrow \varphi(u) \geq_L \varphi(v)$ [2, Prop. 3.1.5].

Lemma A.2. *Let $\mathcal{S} \in \mathcal{W}_h$. The restriction of φ to $\mathcal{W}(\mathcal{S}, h)$ is a bijection between $\mathcal{W}(\mathcal{S}, h)$ and $\mathcal{W}(\Phi_h^+ \setminus \mathcal{S}, h)$.*

Proof. Since φ is an involution, it suffices to show that $\varphi(w) \in \mathcal{W}(\Phi_h^+ \setminus \mathcal{S}, h)$ for all $w \in \mathcal{W}(\mathcal{S}, h)$. Intersecting both sides of equation (A.1) with Φ_h^+ we obtain $N(w_0 w) \cap \Phi_h^+ = \Phi_h^+ \setminus (N(w) \cap \Phi_h^+)$, which is equivalent to $N(w_0 w) \cap \Phi_h^+ = \Phi_h^+ \setminus \mathcal{S}$. Thus $\varphi(w) = w_0 w \in \mathcal{W}(\Phi_h^+ \setminus \mathcal{S}, h)$, as desired. \square

Using the previous lemma and Theorem 2.9, we obtain the following corollary.

Corollary A.3. *For each $\mathcal{S} \in \mathcal{W}_h$, $\mathcal{W}(\mathcal{S}, h)$ has a unique maximal element with respect to weak Bruhat order.*

Proof. Since \mathcal{S} is a subset of Weyl type with respect to h , the complement $\bar{\mathcal{S}} = \Phi_h^+ \setminus \mathcal{S}$ is also a subset of Weyl type with respect to h . By Theorem 2.9(3), the set $\mathcal{W}(\Phi_h^+ \setminus \mathcal{S}, h)$ has a unique minimal element with respect to the weak Bruhat order, denoted $z_{\bar{\mathcal{S}}}$. By Lemma A.2,

$\varphi_S(z_S) \in \mathcal{W}(S, h)$. Since φ is an order reversing involution and z_S is the unique minimal element in $\mathcal{W}(\Phi_h^+ \setminus S, h)$, we conclude that $\varphi(z_S)$ is the unique maximal element in $W(S, h)$. \square

Finally, we prove Lemma 2.11: $\mathcal{W}(S, h)$ is an interval in the weak Bruhat order.

Proof of Lemma 2.11. Let z_S and w_S be the unique minimum and maximum elements, respectively, of $\mathcal{W}(S, h)$ with respect to the weak (left) Bruhat order. Such elements must exist by Theorem 2.9(3) and Corollary A.3. The inclusion $\mathcal{W}(S, h) \subseteq [z_S, w_S]_L$ is immediate. To show the converse, let $w \in [z_S, w_S]_L$. This means $z_S \leq_L w \leq_L w_S$, or equivalently, $N(z_S) \subseteq N(w) \subseteq N(w_S)$. Taking the intersection with Φ_h^+ at each term in this chain, we obtain $N(z_S) \cap \Phi_h^+ \subseteq N(w) \cap \Phi_h^+ \subseteq N(w_S) \cap \Phi_h^+$. Since $z_S, w_S \in \mathcal{W}(S, h)$ we have $N(z_S) \cap \Phi_h^+ = N(w_S) \cap \Phi_h^+ = S$, implying $N(w) \cap \Phi_h^+ = S$. This proves $w \in \mathcal{W}(S, h)$, as desired. \square

ACKNOWLEDGEMENTS. This work was supported in part by the National Security Agency under Grant No. H98230-19-1-0119, The Lyda Hill Foundation, The McGovern Foundation, and Microsoft Research, as part of the Mathematical Sciences Research Institute Summer Research for Women program. The first author is supported by a Natural Science and Engineering Research Council Discovery Grant and a Canada Research Chair (Tier 2) from the Government of Canada. The second author is supported in part by NSF DMS-1954001. The Appendix is by Michael Zeng, and is part of an undergraduate research project conducted under the second author's supervision in 2020.

References

- [1] H. Abe and T. Horiguchi: *A survey of recent developments on Hessenberg varieties*; in Schubert calculus and its applications in combinatorics and representation theory, Springer Proc. Math. Stat. **332**, Springer, Singapore, 2020, 251–279.
- [2] A. Björner and F. Brenti: Combinatorics of Coxeter groups, Graduate Texts in Mathematics, **231**, Springer, New York, 2005.
- [3] S. Cho, J. Hong and E. Lee: *Bases of the equivariant cohomologies of regular semisimple Hessenberg varieties*, ArXiv: 2008.12500, 2020.
- [4] S. Cho, J. Hong and E. Lee: *Permutation module decomposition of the second cohomology of a regular semisimple Hessenberg variety*, ArXiv: 2107.00863, 2021.
- [5] F. De Mari, C. Procesi and M.A. Shayman: *Hessenberg varieties*, Trans. Amer. Math. Soc. **332** (1992), 529–534.
- [6] M. Harada and M. Precup: *The cohomology of abelian Hessenberg varieties and the Stanley-Stembridge conjecture*, Algebr. Comb. **2** (2019), 1059–1108.
- [7] J. Humphreys: Reflection groups and Coxeter groups, Cambridge Studies in Advanced Mathematics **29**, Cambridge University Press, Cambridge, 1990.
- [8] E. Insko and M. Precup: *The singular locus of semisimple Hessenberg varieties*, J. Algebra **521** (2019), 55–96.
- [9] B. Kostant: *Lie algebra cohomology and the generalized Borel-Weil theorem*, Ann. of Math. (2) **74** (1961), 329–387.

- [10] M. Precup: *The Betti numbers of regular Hessenberg varieties are palindromic*, Transf. Groups **23** (2018), 491–499.
- [11] E. Sommers and J. Tymoczko: *Exponents for B-stable ideals*, Trans. Amer. Math. Soc. **358** (2006), 3493–3509.
- [12] J. Shareshian and M. Wachs: *Chromatic quasisymmetric functions*, Adv. Math., **295** (2016), 497–551.

Megumi Harada
Dept. of Mathematics and Statistics, McMaster University
1280 Main Street West
Hamilton, Ontario L8S 4K1
Canada
e-mail: haradam@mcmaster.ca

Martha Precup
Department of Mathematics and Statistics
Washington University in St. Louis
One Brookings Drive
St. Louis, Missouri 63130
U.S.A.
e-mail: martha.precup@wustl.edu