

Developing Nursing Students' Practice Readiness with Shadow Health Digital Clinical Experiences: A Transmodal Analysis

Yeyu Wang¹, Mamta Shah², Francisco A. Jimenez², Cheryl Wilson², Muhammad Ashiq¹, Brendan Eagan¹, David Williamson Shaffer¹
10.1007/978-3-031-47014-1_25

¹University of Wisconsin-Madison
ywang2466, ashig, beagan[@wisc.edu], dws@education.wisc.edu
²Elsevier
m.shah, f.jimenez, c.l.wilson [@elsevier.com]

Abstract. This study applied transmodal analysis, a quantitative ethnographic approach, to examine whether and how virtual patient simulations can aid in educating undergraduate nursing students with competencies that exemplify practice-ready nurses. Multimodal transcripts capturing patient interactions, exam actions and documentation were obtained from two students who used Elsevier's Shadow Health® Digital Clinical Experiences™ (DCE) from Fall 2022 to Spring 2023. Patient scenarios were situated in three content areas (Gerontology, Mental Health, Community Health) and two assignment types (focused assessment and contact tracing). In each scenario, DCE engaged both students in learning activities such as collecting patient data and establishing a caring relationship in a similar manner. These activities indicated how students practiced recognizing and analyzing cues, subjective assessment, diagnosing and prioritizing hypotheses, generating solutions, evaluating outcomes, therapeutic communication, and care coordination and management in relation to each patient's needs and conditions. A statistically significant difference was observed between competencies practiced while completing focused assessments and contact tracing assignments. This study provides evidence for using simulations to facilitate competency-based education in nursing. It also provides motivation for using transmodal and ordered network analysis to advance quantitative ethnography research in health care and health professions education.

Keywords: quantitative ethnography, transmodal analysis, nursing education, competency-based education, virtual patient simulations

1. Introduction

New graduates' insufficient practice readiness persists even as the demand for nurses is growing in the United States [1]. This is a multifaceted challenge since nursing educators are faced with several paradigm shifts and competing gaps in preparing pre-licensure students for the profession. For instance, with the recent release of *The*

Essentials: Core Competencies for Professional Nursing Education [2] and NCSBN Clinical Judgment Measurement Model [3], nursing regulatory bodies have placed competency-based education at the forefront, prompting programs to transform teaching, learning and assessment practices. In addition, the Future of Nursing 2020-2030 report [4] has underscored the need for new nurses to be prepared to (a) treat patients that reflect diversity in social determinants of health and (b) promote health equity across communities. Furthermore, U.S nursing schools are having to turn away thousands of qualified applicants due to shortages of clinical sites, faculty and resource constraints [5]. Lastly, nursing leaders are foreseeing a continued trend towards online/remote education. This pedagogical movement along with rapid digital transformation is likely to create new opportunities and challenges for nursing programs and regulations [6].

We believe that virtual screen based simulations have the potential to cultivate students' practice readiness and aid nursing educators in addressing the aforementioned shifts and gaps in the discipline. [7] concluded that utilizing virtual patient simulations (VPS)-a type of screen-based simulation- had a positive effect on multiple learning outcomes for nursing students. Eighty-six percent of studies in their review demonstrated that VPS were efficient at enhancing nursing students' knowledge acquisition, skill development, critical thinking, self-assurance, and satisfaction with learning. Recently, [8] urged researchers to go investigate learner performance as a direction for advancing simulation use for competency-based education. Quantitative ethnography (QE) has enabled researchers and practitioners to investigate and illustrate complex patterns in human behavior in several domains.

We build upon extant QE research, especially in nursing education, and apply TransModal Analysis (TMA) for investigating students' engagement in multimodal learning activities in Shadow Health® Digital Clinical Experiences™ (DCE). DCE is a type of VPS designed to cultivate nursing students' knowledge, skills and attitudes for providing comprehensive and compassionate care to digital standardized patients.

In what follows, we describe learning activities typically afforded by DCE, and provide an overview of simulation scenarios designed for gerontology, mental health and community health content areas. Next, we describe The Essentials [2] and its application in this study. This is followed by a justification for using TMA to advance QE research on simulations in nursing education. Thereafter, we describe our methodological procedures and modeling decisions. This is followed by a report of findings for Rose and Roshni (pseudonyms) who completed two focused assessments (End of Life scenario in Gerontology, Bipolar Disorder scenario in Mental Health) and one contact tracing assignment (HIV Diagnosis and with Contact Tracing scenario in Community Health) in DCE from 2022-2023. We conclude this paper by discussing our findings and outlining implications for future research.

2. Shadow Health Digital® Clinical Experiences™ (DCE)

The DCE provides an array of standardized clinical scenarios across each course in undergraduate nursing education to guide learners in developing the knowledge, skills, and attitudes needed to care for diverse patients in a safe environment. Typically in a DCE scenario, learners can interact with virtual patients and ask questions in an open environment to explore their medical and sociocultural background. Learners can also perform physical exams and interpret their findings in the simulated electronic health

record (EHR), which mimics the clinical decision making process based on the patient information gathered. During the virtual exam, learners can also express empathy when the virtual patient shares emotional, physical, or experiential difficulties, and offer educational statements when the patient reveals gaps in their understanding of relevant medical topics. The virtual patients are programmed to recognize and respond to thousands of questions and statements related to the learning objectives covered in each scenario, making the conversation feel natural and realistic. As such, learners engage in a clinical reasoning process by completing patient care activities such as collection of history and physical examination data, therapeutic communication skills, and creation of care plans.

DCE simulations have been successful at increasing critical thinking, confidence, and satisfaction among undergraduate nursing students [9]. Students as early as in their first year of nursing education have demonstrated significant efficiency gains when it comes to gathering patient data, applying therapeutic communication, and creating care plans using DCE [10]. In addition, recent studies have shown that DCE scenarios can be an effective means of teaching nursing students about patient care issues that they may not encounter as part of their clinical education, such as fostering cultural competence and sensitivity when caring for transgender patients [11].

For this study, we focus on DCE scenarios from the following content areas in undergraduate nursing education: Gerontology, Mental Health and Community Health. In Gerontology scenarios, students interact with a diverse range of older adult patients, gathering data to assess risk for geriatric syndromes and medication contraindications using Beers Criteria. They take complete health histories, perform problem-focused physical assessments and construct care plans. In Mental Health scenarios, students engage with a set of patients who are experiencing a variety of mental health conditions. They take complete health histories, perform mental status assessments, conduct problem-focused physical exams and complete either care plans or Situation Background Assessment Recommendation (SBAR) handoffs. In Community Health scenarios, students explore a systems-view approach to healthcare—assessing community strengths and weaknesses, tracing the spread of disease, advocating for vulnerable populations, treating individual patients, and creating a care plan for the community.

3. Theoretical Framework

The Essentials [2] provides a competency-based education framework for guiding the development and revision of nursing curricula to prepare entry-level and advanced-level nurses. In addition, it outlines programmatic expectations for teaching, learning and assessment at both levels. Competencies and sub-competencies are organized within 10 domains. These are applicable across all health care areas and diversity of patient populations. At the entry-level, learners should demonstrate attainment and integration of level 1 sub-competencies. Eight concepts (Clinical Judgment; Communication; Compassionate Care; Diversity, Equity, and Inclusion; Ethics; Evidence-based Practice; Health Policy; Social Determinants of Health) that are central to professional nursing practice integrate across the domains and competencies are also included in *The Essentials*.

In this study, we examined competencies characterizing Domains 1 (Knowledge of Nursing Practice), 2 (Person-Centered Care), and 9 (Professionalism) described for entry-level programs. These included: Demonstrating clinical judgment founded on a broad knowledge base (1.3), Engaging with the individual in establishing a caring relationship (2.1), Communicating effectively with individuals (2.2), Integrating assessment skills in practice (2.3), Diagnosing actual or potential health problems and needs (2.4), Promoting self-care management (2.8), Providing care coordination (2.9), and Employing participatory approach to nursing care (9.2). For each of these competencies, we used select level 1 sub-competencies to guide our operationalization of theoretical constructs we examined in this study (see Table 1 for codebook). We also used NCSBN Clinical Judgment Measurement Model Layer 3 [3] to guide our examination of learner performance for competency 1.3 in DCE.

At the entry-level, it is important for nursing students to be exposed to varied experiences in four spheres of care (Disease Prevention/Promotion of Health and Well-Being, Chronic Disease Care, Regenerative / Hospice / Restorative Care and Hospice / Restorative Care Palliative Care) with diverse populations and ages (AACN, 2021). These guidelines along with the core concepts informed our choice of specific scenarios from the three content areas and two types of assignments (focused exam and contact tracing). We examined transcripts of students' interactions with three virtual patients; namely, Regina Walker from gerontology, Lucas Callahan from mental health and Quan Tran from community health. Regina is a 69-year-old Black/African American, cisgender and heterosexual woman. She is a retired family coach and program director at a non-profit. Regina is experiencing increased pain and decreased activity due to metastatic cancer. She needs recommendations for and discussion on hospice care. Lucas is a 25-year-old White cisgender and heteroflexible man who is currently unemployed. He is at a risk of intentions to harm himself or others. Lucas needs education on symptoms common with hypomanic state including lack of sleep. He also needs a care plan and recommendation for outpatient therapy. Quan Tran is a 52-year-old Vietnamese American man. He chooses not to disclose his sexual orientation and is employed as a manager at a trading company. Quan was recently diagnosed with HIV at a community center. The possible contacts of this communicable disease need to be traced using evidence-based guidelines. Quan and the contacts also need to be educated and cared for with empathy. Through these scenarios, students were exposed to patients needing Hospice/Palliative Care (i.e., Regina-End of Life), Chronic Disease Care (i.e., Lucas-Bipolar Disorder), and Prevention/promotion of Health and Wellbeing (i.e., Quan-HIV Diagnosis and Contact Tracing).

4. Quantitative Ethnography in Nursing Education Research

Quantitative Ethnography (QE) is an emerging field for understanding complex processes and discovering meaningful patterns in various disciplines such as education [12], and policy [13]. As a unified approach of qualitative and quantitative analyses, QE provides both thick descriptions and statistical warrants on a given analytic claim [14]. Recent studies in nursing education have applied QE methods to examine alignment of curricular content [15], trace student learning trajectories [16], model instructor facilitation and classroom interaction across pre-briefing, simulation, and debriefing phases for scenarios in fundamentals of nursing [16, 17].

However, nursing education by nature involves multiple modalities such as dialogue and physical examinations. Despite this, existing work in QE mainly relies on unimodal data. For example, [18] adopted epistemic network analysis (ENA) to investigate student learning trajectories based on discourse data collected from virtual reality simulation sessions. According to this study, ENA represented connections made among constructs derived from frameworks such as NCJMM and Quality and Safety Education for Nurses (QSEN). However, this study had two limitations: (1) ENA did not provide ordered relationships for the connections between any pair of constructs or self-references and, (2) ENA was not initially designed to analyze multimodal data and often requires a great deal of additional model parameterization. In ENA, it is not possible to assign different window sizes for different modalities and as a result researchers often need to account for differences in temporal influence of different datastreams by manual adjustment in their model.

In contrast, TransModal Analysis (TMA) models multimodal data by specifying a function or functions that describes, for each data modality, how events interact and using those functions to include multiple modalities in the same model [19]. As such, in this study, we implemented TMA using Ordered Network Analysis [20] by specifying a different window size for each data modality. Described in further detail below, students engaged in DCE through conversations, virtual exams, and documentations. These modalities (dialog, click, documentation) are highly interactive and interwoven during the learning processes in DCE. Hence, we chose to use TMA to model the cross-modality interactions in complex thinking and activity in virtual patient simulations.

5. Methods

5.1 Participants and Settings

Purposive sampling was applied to identify two Bachelor of Science in Nursing (BSN) students from the same cohort at a public university in south central United States. The students were selected because they had completed the three scenarios at the time of the study. Rose and Roshni (pseudonyms) were enrolled in the Adaptation in Aging and Psychiatric-Mental Health Nursing courses in Fall 2022 where they completed the scenarios for Regina Walker and Lucas Callahan respectively. They completed Quan Tran's scenario in Spring 2023 as part of their Community Health Nursing course.

In this study, the interview guide was fully enabled by the instructor for all three assignments in DCE. The interview guide is meant to scaffold students' DCE as they engage in subjective data collection. The interview guide shows students the high-level data within each section of the patient interview they will need to collect. Faculty may choose to (partially or fully) enable or disable the interview guide when assigning assignments to students. The full option allowed Rose and Roshni to see explicitly what subjective findings were scored in the interview (e.g. asked about chief complaint) before they uncovered them. This was the typical preference for faculty at the institution where the two students were enrolled.

5.2 Transcripts, Codebook and Coding

We organized and examined a total of 1760 lines of timestamped utterances for the two students' (Rose and Roshni) transcripts from three scenarios (Regina, Lucas, Quan). The utterances included a variety of interactions (answer, clarification, exam action, feedback, greet, prompt, response, statement) logged between participants (student, patient, the system, and other virtual characters). These interactions characterized the nature of specific learning activities (e.g., objective data collection, subjective data collection, education and empathy) students typically engage in DCE across three modalities (click, dialog, documentation) and distinct phases (assessment, care plan, contact tracing) in a scenario.

The nested nature of each utterance provided insight into the overall pedagogical structure of the simulation experience. For instance, the dialogue data involved the student conversing with the virtual patient for subjective data collection related to but not limited to the history of the patient's present illness, past medical history, review of systems, social history. Dialog data also captured students' empathizing and educating their patients. The click data involved the student examining the patient, performing physical assessments, interpreting observations for any abnormalities, and practicing contact tracing. The documentation data involved the student summarizing and/or interpreting the state of the patient throughout the scenario. The temporal structure of the utterances was meaningful to understand the sequence each student followed in a specific scenario and the amount of time they spent in each learning activity.

Table 1. Codebook

Code name	Definition	Example (from Lucas Callahan's Scenario)
Recognize Cues (RC)	Determining what client findings are significant, most important, and of immediate concern to the nurse (relevant cues)	Inspected right-left forearm, right-left wrist
Analyze Cues (AC)	Organizing and linking the relevant cues with client conditions/problems	• Observations-Evidence of self-harm • Wounds or Scars-Scar visible • Location Of Visible Abnormality-Right and left wrist
Diagnosis + Prioritizing Hypothesis (DPH)	Ranking client conditions/problems according to urgency, complexity, and time. Diagnosing actual or potential health problems and needs	• Diagnosis: Risk for injury • Signs and Symptoms: Poor impulse control, Sleeping 3 hours or fewer in the past few days, Prior suicide attempt

Generate Solutions (GS)	Identifying interventions that meet desired outcomes for the client; can include collecting additional assessment data	Short-term Goal-The patient will remain injury free until he can be evaluated by a psychiatric provider.
Evaluate Outcomes (EO)	Comparing actual client outcomes with desired client outcomes to determine effectiveness of care	"[Patient] is writing poetry and reciting. That is not injurious to him. He is safe and free from injury."
Therapeutic Communication (TC)	Engaging with the individual in establishing a caring relationship. Communicating effectively with individuals. Employing a participatory approach to nursing care	"I am so sorry you feel that way. For your safety we will frequently check on you. We are moving you to a private room."
Subjective Assessment (SA)	Integrating assessment skills in practice	• Have any other people noticed your mood or energy shifts? • Do you ever have excessive irritability or anger? • Have you been in fights or arguments?
Care-Management and Coordination (CMC)	Promoting self-care management and providing care coordination	Interventions- • Coordinate with staff for frequent checks • Provide the patient with structured, solitary activities that do not present a risk for injury • Request order for Prozac (fluoxetine) from a healthcare provider

We applied a combination of manual and automated approaches to code the dataset using the codebook above (See Table 1). Automated coding allows researchers to operate fast and minimize human efforts; however, it is challenging to use this approach for coding (1) affect-intensive complex constructs and (2) domain-specific jargons and terms. In a survey on irony detection, [21] showed that automated coding and classification remained in a relatively low model performance using Natural

Language Processing tools (NLP). Similarly, [22] highlighted challenges in automated detection of humor using a language model. Another challenge of applying automation is in coding jargon and terms in specialized domains, such as medical studies. [23] illustrated this further while applying NLP to detect cancer from medical records.

Considering the affordances and constraints of each coding method and our grounded understanding of the dataset used in this study, we manually coded the constructs of Therapeutic Communication (TC), Subjective Assessment (SA), and Care-Management and Coordination (CMC) using social moderation [24]. These codes manifested in unique ways in students' transcripts based on the context of each scenario. Automation was applied to code students' actions for recognizing cues (RC), analyzing cues (AC), diagnosing and prioritizing hypothesis, generating solutions (GS) and evaluating outcomes (EO) because the conceptual definitions of these codes were consistent with procedural definitions (keyword matching) as captured by the DCE system. For example, logs were coded for an occurrence of RC when the student performed physical assessments using the exam action (e.g., assessed vitals). If the student correctly interpreted the result of their exam action (e.g., normothermic), this was coded as an occurrence of AC.

5.3 Model Construction and Research Questions

We took a TMA approach and constructed a model with ONA to represent patterns of student performance across the three scenarios in DCE. In particular, we specified different window lengths for learning events in different modalities. This is important to note because specifying different window lengths is an attempt to account for these differences in multimodal learning data. For instance, in a hypothetical context of collaborative problem solving where a student is involved in peer discussion and has access to resources in a system, chats may have a shorter window of impact to future learning events. That is, topics can rapidly change in a discussion; but engaging with a resource may have a longer influence in a student's connection making compared to chatting.

In this study, we defined the smallest unit of analysis as students within learning activities. For each unit, ONA calculated and accumulated connections across eight codes (See Table 1) within recent temporal contexts. With learning events collected from three modalities, we operationalize recent temporal contexts for clicks, dialogue, and documents respectively. That is, we configured unique window sizes for different modalities. Both patient interactions (dialog data) and exams actions (click data) have a strong temporal dependency and continuity due to interactivity between learners and the system. According to our qualitative analysis, we specified a window of five learning events for these two modalities. However, we selected a shorter window for the documentation modality, the length of two learning events, because each documentation is usually connected with the next action due to system design.

Using the methodological procedures and coding decisions described above, we developed and interpreted a combination of TMA and ONA (T/ONA) graphs to answer the two research questions (RQ): *How do students engage in DCE scenarios?* We examined the alignment between learning patterns and guided instructional design across the three scenarios. *Is there a difference in the connections students make to clinical competencies based on simulation assignment types in DCE?* We compared the patterns of connection making across the focused exam and contact tracing scenarios.

6. Results

To answer each research, we describe the connection patterns visually and statistically and interpret the dimensions based on the node positions in respective T/ONA graphs. These are supported by qualitative examples from students' performance in DCE scenarios.

6.1 RQ1: How do students engage in DCE scenarios?

Figure 1 represents the grand mean connections (strength, self-referencing, direction) of learning patterns for Rose and Roshni across all scenarios. The darker and thicker edges indicate strength of connections; arrows on edges indicate the main direction of connections; and the radius of the outer contour of a node indicates the total receiving degree, while the radius of the inner contour of a node indicates degree of self-references.

In the two-dimensional space of Figure 1, competencies related to Domain 1 (RC, AC, DPH, GS, EO) are clustered on the most positive side of the y-axis, while competencies related to Domains 2 and 9 (CMC, TC, SA) are located in the middle or negative side of the dimension. Students made self-references within SA. That is, students spent a prolonged period performing a holistic assessment and obtaining a complete history of the patient. Self-referencing was also observed within AC. Students spent a bulk of their time in the scenarios interpreting patients' cues to relevant medical conditions/health problems. These activities preceded students' decision making to foster patients' well-being. TC was central to Rose and Roshni's participation in the three scenarios. This allowed them to establish relationship-centered care, demonstrate empathy, practice humility and cultural sensitivity especially while engaging in subjective assessment and generating solutions for three different patients. Another pattern worth highlighting is students' practice of CMC as a response to subjective assessment and generating solutions. Students were observed educating patients, promoting self-care and facilitating continuity of care through coordination with family members and the healthcare team. Across the three scenarios, there was no significant difference between ONA scores for both students ($t(74.424) = 1.692$, $p = 0.095$, Cohen's $d = 0.378$). That is, both students practiced competencies related to clinical judgment, person centered-care, and professionalism in a similar manner as a result of using DCE. Below, we illustrate these connections to clinical competencies by drawing examples from Rose and Roshni's engagement in the gerontology scenario.

Students engaged in two phases in the end of life focused exam for virtual patient Regina Walker: assessment and care plan. During assessment, Rose and Roshni inquired about major health deviations. This included interacting with Regina about her chief complaint (increased pain), history of present illness (onset, duration, location, characteristics, aggravating factors, relieving factors, and severity of pain), past medical history (existing health conditions, general and medication allergies), social history (substance use), review of systems (head, eyes, ears, nose, throat, respiratory, cardiovascular, gastrointestinal, neurological, psychological). A big portion of their dialog centered around Regina's functional status and geriatric syndromes for which older adults may be at risk. This included questions about depression, living environment, feeling safe at home, bathing, dressing, toileting, transferring, continence, ability to eat independently, sleeping habits, confusion, evidence of falls, gait and

ambulatory aids, skin breakdowns, weight changes, weight loss, oral or dental problems, appetite changes, health and social activity, perception of health, fatigue, ER visits or hospitalizations). Both students complemented this subjective data collection by performing exam actions, and noting their observations for any abnormalities. This included (a) assessing vitals, IV bag, IV pump, IV site, urine quality; (b) inspecting eyes, mouth, skin; (c) auscultating carotids, breath sounds, heart sounds, bowel sounds; (d) palpating abdomen, bladder; (e) testing cognition, skin turgor and capillary refill. In a relatively brief but important part of this phase, Rose and Roshni discussed Regina's comfort and preference for hospice care, the family's need for health care services, and implications for health policy, financing and service availability.

Their care plan focused on symptom management and advocacy for appropriate palliative/hospice care. Specifically, Rose and Roshni indicated their diagnosis for Regina (Readiness for effective coping), identified signs and symptoms (Interest in hospice care, increased home care needs outlined short-term goals) and recommended interventions (e.g., Evaluate the patient's current understanding of coping strategies supporting their transition to hospice care), discuss the plan with the patient, and evaluate their own understanding of hospice care as a result of their simulated experience with Regina.




Fig 1. Grand mean plot illustrating Rose and Roshni's patterns of engagement across the gerontology, mental health and community health scenarios

There were subtle differences in students' participation patterns. For instance, Rose engaged in a dialogue with Regina before performing exam actions. Roshni chose the opposite approach. Rose was also more compassionate in her communication during both phases; she (a) offered information about what to expect during assessment, (b) provided reassurance during sensitive discussions, (c) expressed consideration for

family and Regina while discussing an end of life transition plan. The following statements are an example:

Rose: Ms. Walker, we are going to work on ways for you to cope with the upcoming changes you are about to face. This will help the transition be less demanding on you as well as your family. I know this is going to be hard on you all, so I am here for any questions or concerns. I want to know how much education you have on coping strategies, give you some supplemental strategies, and also give you referrals.

Roshni: "I will provide you and your family members a referral to the funeral service, spiritual support and financial assistance if that is ok with you.

6.2 RQ2: Is there a difference in the connections students make to clinical competencies based on simulation assignment types in DCE?

Figure 2 depicts the differences in connections made by students in the two types of simulation assignment type in DCE. In this subtracted plot, edges and nodes are colored based on the stronger connections made by a certain scenario type (i.e., purple for focused exams and yellow for contact tracing). In the T/ONA space depicted in Figure 2, patterns of node positions are similar to the grand mean plot in Figure 1—competencies related to Domain 1 cluster on the negative side of the x-axis, while codes depicting Domains 2 and 9 cluster on the positive side. However, according to t-test of ONA scores, there is a significant difference for connections made in the focused exams and contact tracing tasks ($t(76.471) = 11.3$, $p < .001^{**}$, Cohen's $d = 1.985$). That is, the type of assignment in DCE influenced the clinical competencies students applied during learning activities in a scenario.

For connections made in focused exam scenarios, both CMC and Diagnosis and Prioritize Hypothesis (DPH) were observed as common responses to AC. That is, after analyzing cues from the virtual patients, students tended to rank signs and symptoms, diagnose conditions, and provide care advice. Additionally, students made more connections from GS to CMC in focused exams. In other words, after generating solutions related to the virtual patients' conditions, Rose and Roshni offered corresponding care-management advice and coordinations such as scheduling follow-up appointments. For the contact tracing scenario, students made more (1) self-references within SA, (2) self-references with TC and (3) connections from SA to TC and no connections to the clinical judgment codes (RC, AC, DPH, GS, EO) because there was no objective data collection and care planning in this scenario. Self-references and connections were influenced by a focus on conducting a comprehensive patient assessment, facilitating health literacy, preventing disease, and promoting well-being. Below, we illustrate Rose and Roshni's engagement in the community health scenario.




Fig 2. Subtraction plot illustrating differences in connections made by Rose and Roshni in focused exam (purple) vs contact tracing assignment (yellow)

Students engaged in two phases in the contact tracing assignment for virtual patient Quan Tran: assessment and contact tracing. During assessment, Rose and Roshni inquired about Quan's chief complaint (i.e., HIV diagnosis), history of present illness (testing history, prodrome), past medical history (vaccinations, allergies, past hospitalizations), social history (home life, support system, substance use, typical diet), medication (herbal supplements, antiretroviral prescription), sexual history (sexual partners), review of relevant systems (constitutional and mental health; integumentary, respiratory, and cardiovascular system), patient needs (goals and priorities), and social determinants of health (employment, health insurance, education). Once again there were subtle differences in Rose and Roshni's participation. For instance, Rose followed up with Quan when he reported not understanding how HIV infection is transmitted, taking an herbal supplement consisting of echinacea and goldenseal, and wanting to keep his diagnosis hidden from coworkers and family members. Roshni did not respond to Quan's lack of understanding of HIV transmission; she followed up on his use of herbal supplements, reaction of others to HIV status. She also educated him when Quan reported unfamiliarity with his antiretroviral medication, and feeling anxious about his diagnosis.

Students were prompted and guided by the system during the second phase of the scenario to model the process of tracing contacts and notifying partners. This included asking contacts about the results of their sexually transmitted infection (STI) panel or offering assistance to identify a clinic that performs STI testing, underscoring the importance of treatment from an HIV specialist and an appropriate medication regimen, encouraging regular tracking of viral load, maintaining confidentiality, and promoting safe sex practices.

7. Discussion and Implications

Professional organizations and regulatory bodies such as the American Association of Colleges of Nursing (AACN) and National Council of State Boards of Nursing (NCSBN) are keen on helping nursing programs address the practice-readiness gap. An emphasis on competency-based education provides the opportunity to enhance interprofessional education, increase the use of simulation, and improve clinical judgment in new graduate and advanced practice nurses [25]. Virtual patient simulations provide a beneficial modality in which learners apply and practice their clinical reasoning and critical thinking abilities before interacting with real patients [26]. However, current research on simulations heavily relies on self-evaluations [8]. In this study, we examined two undergraduate students' performance in DCE and provided evidence for their practice of clinical competencies characterized in Domains 1 (Knowledge of Nursing Practice), 2 (Person-Centered Care), and 9 (Professionalism) for entry-level programs in The Essentials [2] and Layer 3 of NCSBN Clinical Judgment Measurement Model [3].

There is a growing body of research in health care and health professions education that applies Quantitative Ethnography to investigate complex questions about professional enculturation and practice [27]; [28]. However, few studies have used multimodal data and fewer exist in the nursing context [29]. This paper applied TMA to examine data obtained from three types of modalities (click, dialog, documentation) to make sense of students' engagement in collecting and interpreting patient data, synthesizing evidence, and promoting care that is suitable for each patient's condition (i.e., Disease Prevention/Promotion of Health and Well-Being, Chronic Disease Care, Hospice/Palliative Care). TMA allowed us to set and account for the impact of different window sizes for a variety of data types. For a complete discussion of both the mechanisms of TMA, its conceptual and theoretical underpinnings, please refer to a forthcoming paper [19].

In a previous study on DCE, researchers observed students practicing recognizing cues for a prolonged period and demonstrated the ways in which this clinical skill manifested in a pediatric and a geriatric patient scenario in the health assessment content area [30]. However, this self-referencing phenomena could not be modeled using Epistemic Network Analysis (ENA). In this study, Ordered Network Analysis allowed us to not only capture the strength of connections among multiple clinical competencies that ENA typically allows, but also illustrate directionality and self-referencing (Figures 1-4) across two assignment types (in three content areas (gerontology, mental health, community health)).

Nursing programs are likely to continue their expanded use of virtual simulations even after the peak of COVID-19 pandemic [31]. At the same time extant QE research is providing valuable insight into the design and enactment of simulation-based learning in undergraduate nursing education (manikin-based, virtual reality, digital standardized patients) [15]; [18]; [30]. This study provides additional impetus for continuing the application of research in this discipline. In the future, researchers should broaden examinations using virtual patient simulations like DCE to include additional (a) foundational and specialty content areas in nursing, (b) clinical competencies, (c) no. of participants. Researchers should also consider deepening their examinations by investigating (a) students' performance in DCE scenarios where the

interview guide is partially enabled or turned off, and (b) the impact of interaction types (e.g., questions, clarification) between participants (e.g., student, patient, system). Findings from these studies can yield recommendations to inform regulations on how nursing faculty and administrators can use virtual simulations for fostering students' practice readiness.

References

1. National Council of State Boards of Nursing: "The NCSBN 2023 Environmental Scan: Nursing at a Crossroads—An Opportunity for Action: Nursing at a Crossroads: An Opportunity for Action". *Journal of Nursing Regulation*, no. 13.4, (2023), S1–S48.
2. American Association of Colleges of Nursing: *The Essentials: Core Competencies for Professional Nursing Education*. (2021).
3. Dickinson, P. et al.: "NCSBN Clinical Judgment Measurement model Clarification". *Journal of Nursing Education*, no. 59.7, (2020), pp. 365–65.
4. Wakefield, M. et al.: *The future of nursing 2020-2030: Charting a path to achieve health equity*. (2021).
5. American Association of Colleges of Nursing: "Data Spotlight: The Impact of Insufficient Clinical Sites on Baccalaureate Program Admissions". (2022).
6. Alexander, M.: "Nursing Challenges Continue Into 2022". *Journal of Nursing Regulation*, no. 12.4, (2022).
7. Foronda, C. L. et al.: "Virtual Simulation in nursing education: A systematic review spanning 1996 to 2018". *Simulation in Healthcare*, 15(1), (2020), pp. 46–54.
8. Cole, H. S.: "Competency-based Evaluations in Undergraduate Nursing Simulation: A State of The Literature". *Clinical Simulation in Nursing*, no. 76, (2023), pp. 1–16.
9. Turrise, S. L. et al.: "Virtual simulation: Comparing critical thinking and satisfaction in RN-BSN students". *Clinical Simulation in Nursing* 46, (2020), pp. 57–61.
10. Santarelli, T. et al.: "Enhancing Nursing Student Efficiency and Efficacy in Patient Care through Virtual Patient Simulation". *MODSIM World 2022*, (2022).
11. Altmiller, G. et al.: "Impact of a Virtual Patient Simulation on Nursing Students' Attitudes of Transgender Care". *Nurse Educator*, (2022), pp. 10–1097.
12. Carmona, G. et al.: "Exploring Interactions Between Computational and Critical Thinking in Model-Eliciting Activities Through Epistemic Network Analysis". (2022).
13. Siebert-Evenstone, A. et al.: "Cause and Because: Using Epistemic Network Analysis to Model Causality in the Next Generation Science Standards". (2019).
14. Shaffer, D.W.: *Quantitative Ethnography*. (2017).
15. Shah, M. et al.: "Alignment of Content In Simulation Learning System to Clinical Judgment Competencies". White paper. Elsevier Education, (2023).
https://evolve.elsevier.com/education/wp-content/uploads/sites/2/2023/02/23-NHPgap-0065-SLS_QE_Alignment_Whitepaper_HR.pdf
16. Shah, M. et al.: "Student Learning in Simulation Learning System with Virtual Reality: A Quantitative Ethnographic Examination of a Fundamentals of Nursing Scenario". White Paper. Elsevier Education, (2022).
https://evolve.elsevier.com/education/wp-content/uploads/sites/2/22-NHPep-0060_WhitePaper-StudentLearning-SLSwVR_Final-1.pdf
17. Shah, M. et al.: "Modeling Educator Use of Virtual Reality Simulations in Nursing Education Using Epistemic Network Analysis". 7th Conference of the Immersive Learning Research Network (iLRN), IEEE, (2021), pp. 1–8.
18. Shah, M. et al.: "Quality and Safety Education for Nursing (QSEN) in Virtual Reality Simulations: A Quantitative Ethnographic Examination". In: *Advances in Quantitative Ethnography: Third International Conference, ICQE 2021*, Online. November 8-11, 2021, Proceedings, (2021).

19. Shaffer, D.W. et al.: "Transmodal Analysis". (2023).
20. Tan, Y. et al.: "Ordered Network Analysis". In: *Advances in Quantitative Ethnography: Fourth International Conference, ICQE 2022, Copenhagen, Denmark, October 15–19, 2022, Proceedings*, (2022).
21. Wallace, B.C. et al.: "Computational irony: A survey and new perspectives". *Artificial Intelligence Review* 43, (2015), pp. 467–83.
22. Gultchin, L. et al.: "Humor in word embeddings: Cockamamie gobbledegook for nincompoops". PMLR, (2019).
23. Savova, G. et al.: "Use of Natural Language Processing to Extract Clinical Cancer Phenotypes from Electronic Medical Records; Natural Language Processing for Cancer Phenotypes from EMRs". *Cancer Research*, (79)21, (2019), pp. 5463–70.
24. Herrenkohl, L.R. et al.: "Investigating elementary students' scientific and historical Argumentation". *Journal of the Learning Sciences*, no. 22.3, (2013), pp. 413–61.
25. Lewis, L.S. et al.: "Nursing Education Practice Update 2022: Competency-Based Education in Nursing". *Sage Open Nursing*, no. 8, (2022).
26. Kononowicz, A. et al.: "Virtual patient simulations in health professions education: Systematic review and meta-analysis by the digital health education collaboration". *Journal of Medical Internet Research*, 21(7), (2019).
27. Shah, M. et al.: "Communicating QE: A Two-Part Resource for Quantitative Ethnographers in Health Education and Health Care Contexts (Part 1 of 2)". In: *Fourth International Conference on Quantitative Ethnography: Conference Proceedings Supplement*, (2022).
38. Ruis, A. et al.: "Communicating QE: A Two-Part Resource for Quantitative Ethnographers in Health Education and Health Care Contexts (Part 2 of 2)". In: *Fourth International Conference on Quantitative Ethnography: Conference Proceedings Supplement*, (2022).
29. Buckingham-Shum, S. et al.: "The multimodal matrix as a quantitative ethnography methodology". In: *Advances in Quantitative Ethnography: First International Conference, ICQE 2019, Madison WI, USA, October 20-22, 2019, Proceedings*, (2019).
30. Shah, M. et al.: "Understanding How Undergraduate Nursing Students (Learn to) Recognize Cues in Digital Clinical Experiences: A Transmodal Analysis". In: *Fourth International Conference on Quantitative Ethnography: Conference Proceedings Supplement*, (2022).
31. Harder, N. et al.: "How the Pandemic Impacted Simulation: Where Do We Go From Here?" *Clinical Simulation In Nursing*, no. 74, (2023), pp. 1–2.