

Lack of testosterone-sensitive TRPM8 Ca²⁺ channel disrupts measures of anxiety and depression in mice: Implications for dopamine



Erdmier, A.¹, Saldes, E.¹, Shepler, B.³, Palmisano, H.³, & Koeltzow, T.E.^{1,3}, Asuthkar, S.^{1,2}
¹Department of Psychology, Bradley University, Peoria, IL 61625, ²Pediatrics, University of Illinois College of Medicine at Peoria
³Department of Cancer Biology and Pharmacology, University of Illinois College of Medicine at Peoria



Introduction

- Transient Receptor Potential Melastatin-8 (TRPM8) cation channel being a high affinity target for testosterone suggests a mechanism by which androgens can rapidly influence central nervous system activity (Asuthkar et al., 2015).
- Sex differences are a signature feature of anxiety, depression and substance use, yet the identification of neurobiological mechanism to account for this dimorphism has not been meaningfully identified (Altemus et al., 2014; Bangasser & Cuarenta, 2021; Santos-Toscano et al., 2023).
- In addition to its well-established role in peripheral cold sensation (Bautista et al., 2007), TRPM8 is highly expressed in the central nervous system, including the hypothalamus, and limbic and cortical regions relevant to motivated behavior.
- Male TRPM8-KO mice exhibit disrupted sexual behavior, including increased mounting, indiscriminate approach, and a prolonged latency to satiety. These behaviors were linked to altered dopamine (DA) neuron activity in the ventral tegmental area (Mohandass et al., 2020). DA exerts an essential role in a variety of motivated behaviors, including response to novelty and reinforcement learning, and is critically implicated in substance use disorders and depression.

Hypothesis

It is possible that the effects of TRPM8 deletion on sexual reward will translate to more global reward processes. Therefore, we subjected TRPM8 null mutant mice to a battery of tests that are sensitive to DA and/or are established rodent models of anxiety, depression, and substance abuse. We hypothesize that male mice lacking the TRPM8 gene will exhibit impairment on these measures compared to wildtype mice.

Method

Animals: Female and male C57BL/6J WT and TRPM8-KO (#008198) were obtained from Jackson Laboratories (Bar Harbor, ME). Mice were group housed by sex at the University of Illinois College of Medicine in Peoria with their conspecifics on a 12:12 light/dark cycle under constant temperature and humidity and with food and water available *ad libitum*.

Behavioral Approaches


Locomotion in Activity Frame (AF), Sucrose 2-Bottle Preference, Light/Dark Test, & Forced Swim Test

Pharmacological Approach

Amphetamine Administration: Mice were placed in AF and allowed to habituate for 20 min prior to an injection of physiological saline (3ml/kg body weight, s.c.). After 20 minutes, mice were injected with either 1.0 mg/kg (Day 1) or 3.0 mg/kg (Days 2-6), and activity was recorded for an additional 70 minutes.


All procedures were performed in compliance with an approved IACUC protocol. Statistical analyses were performed using SPSS using either ANOVA or Student's t tests.

TRPM8 is primarily activated by testosterone




TRPM8 channel activity was assessed in planar lipid bilayers in the presence of varying concentrations of testosterone and membrane impermeable BSA-testosterone. Single channel currents at 100 mV of TRPM8 activated in the presence of testosterone and membrane impermeable BSA-testosterone. However, BSA-testosterone induced TRPM8 openings only into the small conductance. Suggesting that TRPM8 is primary a testosterone receptor.

Results




Spontaneous Locomotor Activity: As expected, female mice were more active than male mice across the 48 h. test ($F(1,26) = 16.35, p < 0.001$; Left). There were no differences between TRPM8-KO and WT mice. Inset represents cumulative activity totals. Shaded area denotes dark cycle. DA plays a role in habituating to a novel environment. Male TRPM8-KO mice exhibited a trend towards increased latency to habituate ($F(1,12) = 1.60, p = 0.2$; Right). Asterisks denote statistically significant differences among planned comparisons. Error bars depict SEM.



Sucrose Preference: Male TRPM8 deficient mice exhibited a preference for a low concentration of sucrose (0.1%) compared to male wildtype mice ($t(10) = 2.01, p < 0.05$; $d = 1.00$). In contrast, deletion of the TRPM8 gene had no impact on sucrose preference among females across three different concentrations.

Light Dark Test: During the 10 min test, male mice lacking the TRPM8 gene made fewer crosses between the light and dark side ($t(10) = 3.34, p < 0.005$), and spent statistically significantly less time in the light side of the chamber ($t(10) = 2.44, p < 0.05$). **Forced Swim Test:** TRPM8-KO mice exhibited a statistically significant decrease in the latency to immobility, a measure of behavioral despair, compared to wildtype mice ($t(10) = 3.70, p < 0.005$; right). Error bars depict SEM.



Repeated Amphetamine: On Day 1 of testing, mice were allowed 40 minutes to acclimate to the test environment (Habituation), prior to an injection of physiological Saline (3 ml/kg body weight, i.p.) prior to an injection of a low dose of Amphetamine (1.0 mg/kg). Procedures were identical for 5 subsequent days, except the dose of amphetamine was increased (3.0 mg/kg). Data analysis indicate no differences between groups during habituation or in response to saline across 6 days of testing. However, RMANOVA revealed a statistically significant main effect of test day 3 ($F(5,65) = 157, p < 0.001$). Error bars depict SEM.

Discussion

- Deletion of the TRPM8 receptor did not reliably influence spontaneous locomotor activity among either male or female mice. However, there was a trend for male TRPM8 null mutant mice to habituate to the test environment faster than wildtype mice. **These findings suggest that DA is not altered under basal conditions.**
- Male TRPM8^{-/-} mice exhibited a preference for a low concentration of sucrose; this effect was not observed among female mice. In contrast, deletion of the TRPM8 receptor resulted in decreased activity in the light-dark test, and reduced time in the anxiogenic light side of the chamber. These mice also exhibited a decreased latency to immobility in the forced swim test. Together, these findings suggest that **TRPM8 may play a protective role in the etiology of anxiety and depression.**
- Male TRPM8-KO mice exhibited a diminished response to amphetamine compared to wildtype, suggesting that **TRPM8 may be involved in substance use disorders.**
- Future research seeks to clarify the relationship of testosterone, TRPM8, and DA.

References

- Altemus, Margaret, Nilofar Sarvaiya, and C. Neill Epperson. "Sex differences in anxiety and depression clinical perspectives." *Frontiers in neuroendocrinology* 35.3 (2014): 320-330.
- Asuthkar, Swapna, et al. "The TRPM8 protein is a testosterone receptor: I. Biochemical evidence for direct TRPM8-testosterone interactions." *Journal of Biological Chemistry* 290.5 (2015): 2659-2669.
- Bangasser, Debra A., and Amelia Cuarenta. "Sex differences in anxiety and depression: circuits and mechanisms." *Nature Reviews Neuroscience* 22.11 (2021): 674-684.
- Bautista, Diana M., et al. "The menthol receptor TRPM8 is the principal detector of environmental cold." *Nature* 448.7150 (2007): 204-208.
- Mohandass, Adithya, et al. "TRPM8 as the rapid testosterone signaling receptor: Implications in the regulation of dimorphic sexual and social behaviors." *The FASEB Journal* 34.8 (2020): 10887-10906.

Acknowledgements

This work is supported by the National Science Foundation: (IOS-1922428)