
Multi-branch Spatio-Temporal Graph Neural Network For Efficient Ice Layer Thickness Prediction

Zesheng Liu

Department of Computer Science and Engineering
Lehigh University
Bethlehem, PA 18015
zel220@lehigh.edu

Maryam Rahnemoonfar*

Department of Computer Science and Engineering
Department of Civil and Environmental Engineering
Lehigh University Bethlehem, PA 18015
maryam@lehigh.edu

Abstract

Understanding spatio-temporal patterns in polar ice layers is essential for tracking changes in ice sheet balance and assessing ice dynamics. While convolutional neural networks are widely used in learning ice layer patterns from raw echogram images captured by airborne snow radar sensors, noise in the echogram images prevents researchers from getting high-quality results. Instead, we focus on geometric deep learning using graph neural networks, aiming to build a spatio-temporal graph neural network that learns from thickness information of the top ice layers and predicts for deeper layers. In this paper, we developed a novel multi-branch spatio-temporal graph neural network that used the GraphSAGE framework for spatio features learning and a temporal convolution operation to capture temporal changes, enabling different branches of the network to be more specialized and focusing on a single learning task. We found that our proposed multi-branch network can consistently outperform the current fused spatio-temporal graph neural network in both accuracy and efficiency.

1 Introduction

As global temperatures continue to rise, research has shown that the accelerated mass loss of polar ice sheets is increasingly contributing to climate change [2, 8, 11, 21]. Polar ice sheets comprise several internal ice layers formed in different years. A better understanding of the status of internal ice layers can provide valuable information on snowfall melting and accumulation and enable a comprehensive understanding of the global climate system and future climate change.

Traditional methods to study the internal ice layer are through onsite ice core [9]. However, the limited and discrete coverage makes it impossible to study the continuous change of the ice layer. Additionally, onsite ice cores are expensive to obtain and will cause destructive damage to the ice sheet. In recent years, airborne snow radar sensors have proven to be a more effective way to study ice layers. Internal layers with different depths are captured continuously as radargrams by measuring the reflected signal strength [1], shown in Figure 1(a).

*Corresponding Author

With the development of deep learning techniques, various convolution-based neural networks[10, 13-15] have been proposed to extract ice layer boundaries from radargrams. However, noise in the radargrams is shown to be a major obstacle to achieve high-quality results. Instead of the convolution-based neural networks, Zalatan et al.[17-19] focus on learning the relationship between ice layers formed in different years and graph neural network. They utilized a fused spatio-temporal graph neural network, AGCN-LSTM, and aimed to make predictions on the thickness of deeper ice layers based on the thickness information of shallow layers. In their network, graph convolutional network(GCN) for spatio features is fused into the long short-term memory(LSTM) structure for temporal changes. Although their proposed methods have decent performance in predicting ice thickness, training the fused spatio-temporal graph neural network usually takes a few hours due to high computational costs.

In this paper, we aim to build upon the work of Zalatan et al.[17-19] by improving the network performance in both accuracy and efficiency. Our major contributions are: 1) We developed a novel multi-branch spatio-temporal graph neural network that uses the GraphSAGE framework for spatio feature learning and uses temporal convolution operation to learn temporal changes, making different parts of the network more specialized to one single learning task. 2) We conducted extensive experiments on comparison with fused spatio-temporal graph neural networks, and results show that our proposed network can consistently outperform other methods in efficiency and accuracy.

2 From Radargram To Graph Dataset

In this work, we will start with radargrams captured over the Greenland Region in 2012 via airborne snow radar sensor operated by CReSIS[4]. Each radargram varies from 1200 pixels to 1700 pixels in depth, with a fixed 256 pixels in width. The value of each pixel is determined by the reflected signal strength, where brighter pixels in the radargram mean a stronger reflection[1], shown in Figure 1(a). Based on radargrams, corresponding labeled images are generated manually by NASA scientists by tracking out each layer, as illustrated in Figure 1(b). Based on the position of ice layer boundary pixels, layer thickness can be calculated as the difference between its upper and lower boundaries. Moreover, while capturing the radargrams, airborne snow radar sensors will also keep records of current latitude and longitude simultaneously.

Our graph dataset is then generated on the thickness and geographical location information from the radargrams and labeled images with additional pre-processing steps. As shown in Figure 1(b), we will use the top five layers (formed in 2007-2011) to predict the thickness of the deeper ice layers (formed in 1992-2006). To ensure the high quality of our graph dataset, we will eliminate images with less than 20 complete layers, dividing the remaining images into training, testing, and validation sets with a ratio of 3 : 1 : 1. This pre-processing step reduces the total number of valid images to 1660. Each valid labeled image will be represented as two temporal sequences of spatial graphs: one sequence of five graphs as input and another sequence of fifteen as the ground truth. Each spatial graph represents one internal ice layer formed in a specific year, with 256 nodes that correspond to the 256 pixels in the labeled images. Nodes will be fully connected and undirected. Edge weights between two nodes i, j are defined as

$$w_{i,j} = \frac{1}{2 \arcsin(hav(\phi_j - \phi_i) + \cos \phi_i \cos \phi_j \hav(\lambda_j - \lambda_i))} \quad (1)$$

where $\hav(\theta) = \sin^2(\frac{\theta}{2})$. Each node will contain three node features: latitude, longitude, and the thickness of the current ice layer.

3 Key Designs

The key idea of our proposed multi-branch network is to make different parts of the network more specialized. Unlike previous AGCN-LSTM and GraphSAGE-LSTM, in which the network learns spatio and temporal features in one block, we will have individual branches to learn spatio patterns or temporal changes. Figure 1(c) shows the architecture of our proposed network. For each branch, we will have a dimension reduction block to let the GraphSAGE framework or temporal convolution focus more on the relevant features. Outputs of each branch are concatenated together and passed into three linear layers for final prediction.

3.1 Dimension Reduction

In order to let each branch focus only on relevant features, we will have dimension reduction blocks for different branches. Based on signal transmission properties, pixels in the same pixel column but different ice layers will have the same latitude and longitude. Therefore, for the spatial branch, we will condense the overall node feature matrix of the input graph sequence from $5 \times (256, 3)$ to $(256, 7)$, where the seven node features are latitude, longitude and concatenated thickness information of the previous five ice layers. For the temporal branch, we will modify the dimension of the node feature matrix to $5 \times (256, 1)$ by removing the latitude and longitude.

3.2 GraphSAGE Inductive Framework

GraphSAGE [5] is an inductive framework that generates node embedding based on a sampling and aggregating process on its local neighbors [5, 20]. For a known node i with node feature \mathbf{x}_i , it is defined as follows:

$$\mathbf{x}'_i = \mathbf{W}_1 \mathbf{x}_i + \mathbf{W}_2 \cdot \text{mean}_{j \in \mathcal{N}(i)} \mathbf{x}_j \quad (2)$$

where \mathbf{x}'_i is the learned node embedding via GraphSAGE, $\mathbf{W}_1, \mathbf{W}_2$ are layer weights, $\mathcal{N}(i)$ is the neighbor list of node i that may include neighbors with different depth, \mathbf{x}_j is the neighbor node features and mean is the aggregator function. GraphSAGE can be understood as a linear approximation of localized spectral convolution [5], and the sampling and aggregating process can reduce the effect of possible outliers and noise and increase the model's generalization ability.

3.3 Temporal Convolution

Recurrent neural networks, like long short-term memory (LSTM), can effectively learn temporal changes with high computational cost and long training time. In this work, we will replace the LSTM structure with a gated temporal convolution block proposed by Yu et al. [16] to improve its efficiency.

Inspired by Gehring et al. [3], the temporal convolution block will learn the temporal features from the original node features. An input graph sequence with node feature matrix X will pass through three two-dimensional convolution operation to achieve P, Q, R . P, Q are then passed into a gated linear unit (GLU) and R is added to the output as a skip connection. Therefore, the gated temporal convolution is defined as follows:

$$X' = \text{ReLU}(P \times \sigma(Q) + R) \quad (3)$$

where X' is the learned temporal features, \times is an element-wise Hadamard product, and σ is the sigmoid function that serves as the gate part of the GLU.




Figure 1: Diagram of our graph dataset and network structure. (a) Radargram captured by airborne snow radar sensor. (b) Labeled image where each ice layer is labeled out from the radargram. (c) Architecture of our proposed multi-branch network.

Table 1: Experiment results of GCN-LSTM, GraphSAGE-LSTM, and our proposed Multi-branch model. Results are reported as the mean and standard deviation of the RMSE on the test dataset over five individual trials. Train time is reported as the average train time over five individual trials

Model	RMSE Results	Average Train Time
GCN-LSTM	3.2106 ± 0.1188	1:58:56
GraphSAGE-LSTM	3.1949 ± 0.0332	1:16:14
Proposed Multi-branch(SAGE+TempConv)	3.1236 ± 0.0548	0:16:18

4 Experiment and Results

To verify the design of our proposed multi-branch spatio-temporal graph neural network and the choice of branches, we compare its performance and efficiency with several different graph neural networks, including GCN-LSTM[12] and SAGE-LSTM[7]. All the networks are trained on 8 NVIDIA A5000 GPUs and Inter(R) Xeon(R) Gold 6430 CPU. Mean-squared error loss is used as the loss function for all the networks. Adam optimizer[6] with 0.01 as an initial learning rate and 0.0001 as a weight decay coefficient. A step learning rate scheduler halves the learning rate every 75 epochs.

The mean of the training time and the mean and standard deviations of prediction error over the five trials are reported as the model efficiency and accuracy, shown in Table 1. Our proposed multi-branch spatio-temporal graph neural network can significantly outperform previous fused spatio-temporal networks in both efficiency and accuracy. Figure 2 shows the qualitative results of our trained model. Compared with the previously fused spatio-temporal graph neural networks, our proposed multi-branch method significantly improves the accuracy for bottom layers and pixels around the image boundaries.

5 Conclusion

In this work, we develop a novel multi-branch spatio-temporal graph neural network for learning deeper ice layer thickness. This network utilizes GraphSAGE as the spatial branch and temporal convolution to learn temporal changes over time. Experiments show that compared with previous fused spatio-temporal networks, our proposed method consistently performs better in both accuracy and efficiency.




Figure 2: Qualitative results of model predictions. The green line is the groundtruth (manually-labeled ice layers) and the red line is the model prediction.

Acknowledgments and Disclosure of Funding

This work is supported by NSF BIGDATA awards (IIS- 1838230, IIS-1838024), IBM, and Amazon. We acknowledge data and data products from CReSIS generated with support from the University of Kansas and NASA Operation IceBridge.

References

- [1] Emily Arnold, Carl Leuschen, Fernando Rodriguez-Morales, Jilu Li, John Paden, Richard Hale, and Shawn Keshmiri. Cresis airborne radars and platforms for ice and snow sounding. *Annals of Glaciology*, 61(81):58–67, 2020. doi: 10.1017/aog.2019.37.
- [2] Rene Forsberg, Louise Sørensen, and Sebastian Simonsen. *Greenland and Antarctica Ice Sheet Mass Changes and Effects on Global Sea Level*, pages 91–106. Springer International Publishing, Cham, 2017. ISBN 978-3-319-56490-6. doi: 10.1007/978-3-319-56490-6_5. URL https://doi.org/10.1007/978-3-319-56490-6_5
- [3] Jonas Gehring, Michael Auli, David Grangier, Denis Yarats, and Yann N. Dauphin. Convolutional sequence to sequence learning, 2017.
- [4] S. Gogineni, J. B. Yan, D. Gomez, F. Rodriguez-Morales, J. Paden, and C. Leuschen. Ultra-wideband radars for remote sensing of snow and ice. In *IEEE MTT-S International Microwave and RF Conference*, pages 1–4, 2013. doi: 10.1109/IMaRC.2013.6777743.
- [5] William L. Hamilton, Rex Ying, and Jure Leskovec. Inductive representation learning on large graphs, 2018.
- [6] Diederik P. Kingma and Jimmy Ba. Adam: A method for stochastic optimization, 2017.
- [7] Zesheng Liu and Maryam Rahnemoonfar. Learning spatio-temporal patterns of polar ice layers with physics-informed graph neural network, 2024. URL <https://arxiv.org/abs/2406.15299>
- [8] Jérémie Mouginot, Eric Rignot, Anders A. Bjørk, Michiel van den Broeke, Romain Millan, Mathieu Morlighem, Brice Noël, Bernd Scheuchl, and Michael Wood. Forty-six years of greenland ice sheet mass balance from 1972 to 2018. *Proceedings of the National Academy of Sciences*, 116(19):9239–9244, 2019. doi: 10.1073/pnas.1904242116. URL <https://www.pnas.org/doi/abs/10.1073/pnas.1904242116>
- [9] W.S.B. Paterson. 15 - ice core studies. In W.S.B. PATERSON, editor, *The Physics of Glaciers (Third Edition)*, pages 378–409. Pergamon, Amsterdam, third edition edition, 1994. ISBN 978-0-08-037944-9. doi: <https://doi.org/10.1016/B978-0-08-037944-9.50021-2>. URL <https://www.sciencedirect.com/science/article/pii/B9780080379449500212>
- [10] Maryam Rahnemoonfar, Masoud Yari, John Paden, Lora Koenig, and Oluwanisola Ibikunle. Deep multi-scale learning for automatic tracking of internal layers of ice in radar data. *Journal of Glaciology*, 67(261):39–48, 2021. doi: 10.1017/jog.2020.80.
- [11] E. Rignot, I. Velicogna, M. R. van den Broeke, A. Monaghan, and J. T. M. Lenaerts. Acceleration of the contribution of the greenland and antarctic ice sheets to sea level rise. *Geophysical Research Letters*, 38(5), 2011. doi: <https://doi.org/10.1029/2011GL046583>. URL <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2011GL046583>
- [12] Youngjoo Seo, Michaël Defferrard, Pierre Vandergheynst, and Xavier Bresson. Structured sequence modeling with graph convolutional recurrent networks, 2016.
- [13] Debjrat Varshney, Maryam Rahnemoonfar, Masoud Yari, and John Paden. Deep ice layer tracking and thickness estimation using fully convolutional networks. In *2020 IEEE International Conference on Big Data (Big Data)*, pages 3943–3952, 2020. doi: 10.1109/BigData50022.2020.9378070.

- [14] Debrat Varshney, Maryam Rahnemoonfar, Masoud Yari, John Paden, Oluwanisola Ibikunle, and Jilu Li. Deep learning on airborne radar echograms for tracing snow accumulation layers of the greenland ice sheet. *Remote Sensing*, 13(14), 2021. ISSN 2072-4292. doi: 10.3390/rs13142707. URL <https://www.mdpi.com/2072-4292/13/14/2707>
- [15] Masoud Yari, Oluwanisola Ibikunle, Debrat Varshney, Tashnim Chowdhury, Argho Sarkar, John Paden, Jilu Li, and Maryam Rahnemoonfar. Airborne snow radar data simulation with deep learning and physics-driven methods. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 14:12035–12047, 2021. doi: 10.1109/JSTARS.2021.3126547.
- [16] Bing Yu, Haoteng Yin, and Zhanxing Zhu. Spatio-temporal graph convolutional networks: A deep learning framework for traffic forecasting. In *Proceedings of the Twenty-Seventh International Joint Conference on Artificial Intelligence*, IJCAI-2018. International Joint Conferences on Artificial Intelligence Organization, july 2018. doi: 10.24963/ijcai.2018/505. URL <http://dx.doi.org/10.24963/ijcai.2018/505>
- [17] Benjamin Zalatan and Maryam Rahnemoonfar. Recurrent graph convolutional networks for spatiotemporal prediction of snow accumulation using airborne radar. In *2023 IEEE Radar Conference (RadarConf23)*, pages 1–6, 2023. doi: 10.1109/RadarConf2351548.2023.10149562.
- [18] Benjamin Zalatan and Maryam Rahnemoonfar. Prediction of annual snow accumulation using a recurrent graph convolutional approach. In *IGARSS 2023 - 2023 IEEE International Geoscience and Remote Sensing Symposium*, pages 5344–5347, 2023. doi: 10.1109/IGARSS52108.2023.10283236.
- [19] Benjamin Zalatan and Maryam Rahnemoonfar. Prediction of deep ice layer thickness using adaptive recurrent graph neural networks. In *2023 IEEE International Conference on Image Processing (ICIP)*, pages 2835–2839, 2023. doi: 10.1109/ICIP49359.2023.10222391.
- [20] Jie Zhou, Ganqu Cui, Shengding Hu, Zhengyan Zhang, Cheng Yang, Zhiyuan Liu, Lifeng Wang, Changcheng Li, and Maosong Sun. Graph neural networks: A review of methods and applications. *AI Open*, 1:57–81, 2020. ISSN 2666-6510. doi: <https://doi.org/10.1016/j.aiopen.2021.01.001>. URL <https://www.sciencedirect.com/science/article/pii/S2666651021000012>
- [21] H. Jay Zwally, Jun Li, Anita C. Brenner, Matthew Beckley, Helen G. Cornejo, John DiMarzio, Mario B. Giovinetto, Thomas A. Neumann, John Robbins, Jack L. Saba, and et al. Greenland ice sheet mass balance: distribution of increased mass loss with climate warming; 2003–07 versus 1992–2002. *Journal of Glaciology*, 57(201):88–102, 2011. doi: 10.3189/002214311795306682.