



Reply to: Limitations of the iterative electron density reconstruction algorithm from solution scattering data

Thomas D. Grant

REPLYING TO P. V. Konarev & D. I. Svergun *Nature Methods* <https://doi.org/10.1038/s41592-021-01082-x> (2021)

In their Matters Arising¹, Konarev and Svergun claim that several statements made in the original article² are not accurate, namely that DENSS (DENSity from Solution Scattering) cannot reconstruct multiple-density particles from one-dimensional solution scattering data and that the averaging procedure results in a type of low-pass filter effect that negates the accuracy of the averaged density map. Here I demonstrate that, while three-dimensional (3D) *ab initio* reconstructions from DENSS must not be overinterpreted, DENSS can indeed reconstruct multi-density particles provided that sufficient contrast differences exist between components, and that the averaging procedure is an effective way to improve the accuracy of the reconstructions.

The authors raise some valid points highlighting constraints on DENSS that users of the algorithm should take into account. Small-angle X-ray scattering (SAXS) is not only a low-resolution technique, but also a low-information one, leading to ambiguity in reconstructions of 3D objects *ab initio*. Conventional shape restoration algorithms cannot generate higher-resolution reconstructions for two primary reasons: low information content and implicit assumptions of uniform density. The iterative structure factor retrieval algorithm implemented in DENSS removes the assumption of uniform density. However, it does not remove the more fundamental barrier of limited information content. Thus, it is important for researchers to understand that DENSS is still subject to the same ambiguity and low resolution limiting conventional algorithms, which correspond to a fundamental limitation of SAXS.

DENSS is an entirely new approach to the problem of 3D reconstruction, using iterative structure factor retrieval, and this fact should not be lost on the reader. DENSS is capable of creating reconstructions of similar quality to those of traditional shape reconstruction methods while removing one of the primary restraints, demonstrating the power of the approach and its utility for researchers. It is of course always advantageous to achieve similar or better solutions with fewer restraints. The added advantage of DENSS over conventional shape reconstruction is its ability to visualize density within the object envelope. However, owing to the low information content in SAXS, the density difference between two regions must be sufficiently large to be distinguishable at low resolution.

Konarev and Svergun claim that the protein–micelle complex is an unsuitable example of density difference, as bead modeling can also reconstruct hollow spheres in a dumbbell shape. It is true that the coenzyme A (CoA) micelles are reasonably well approximated by

hollow spheres at low resolution, so it is valuable to also use an example where this approximation is no longer valid. Some common use cases include SAXS or small-angle neutron scattering (SANS) experiments on membrane proteins in micelles, nanodiscs, detergents, etc., as well as non-biological targets such as block copolymers, nanoparticles and other soft matter^{3–6}. In the case of some micelles, such as those with DDM detergent, the density of the hydrophobic core is often so low as to exhibit substantial negative contrast in the interior^{3,7}. In such cases, the default restraint of positivity can be disabled when running DENSS to allow for negative contrast; the algorithm is then able to successfully reconstruct the multi-density particle showing large negative density in the interior (Fig. 1a), which is not possible with bead modeling, where uniform density is assumed. Multi-phase bead modeling algorithms can theoretically produce a viable bead model with a contrast series of multiple scattering curves, although this does not appear to be possible with only one scattering curve⁷, as can be done with DENSS. The ability to model negative contrast is indeed a unique advantage of DENSS.

Konarev and Svergun next claim that protein–nucleic acid complexes would be more appropriate examples to demonstrate that DENSS can distinguish the nucleic acid in the reconstruction, as nucleic acid has twice the contrast of protein. However, this assumption is an oversimplification of the actual problem. It is true that nucleic acid has higher contrast than protein owing to the high proportion of phosphates. However, the density of a biological molecule is not a single value, but rather is a broad distribution of values, and the distribution of nucleic acid densities largely overlaps that of protein⁸. Even for intermediate-resolution cryo-electron microscopy (cryo-EM) reconstructions, nucleic acid cannot be distinguished from protein using density alone; rather, advanced algorithms are required to discern it⁸. The assumption of higher density for nucleic acid does not take atomic packing into consideration. Proteins are often much more compact than nucleic acids, as RNA and DNA mostly form helical or double-helical structures containing large voids in the major and minor grooves. At low resolution, such packing must be accounted for when estimating density. Figure 1b shows low-resolution density calculated from atomic coordinates using EMAN2 (ref. 9) for the given examples. In each case, it is clear that the nucleic acid cannot be distinguished from the protein on the basis of higher density, and thus the nucleic acid is also not expected to be distinguishable in DENSS reconstructions.

However, if the density of the nucleic acid were substantially greater, DENSS should be able to distinguish the two components in

Department of Structural Biology, Jacobs School of Medicine and Biomedical Sciences, SUNY University at Buffalo, Buffalo, NY, USA.

✉e-mail: tdgrant@buffalo.edu




Fig. 1 | Calculated or reconstructed low-resolution density of DDM micelles and protein-nucleic acid complexes. **a**, Reconstructions from SAXS data of DDM micelles with large interior negative contrast. The average of 20 DENSS reconstructions is shown as a 3D transparent volume (left) and a cross-section (right). Density is colored in green (positive density) and red (negative density). **b**, Calculated density from atomic coordinates at low resolution using EMAN2 (ref. ⁹) for the protein-nucleic acid complexes shown by Konarev and Svergun. Maps were calculated at the resolution shown in the figure as estimated by DENSS reconstructions. Maps are colored from low density (blue) to high density (red). **c**, Calculated and reconstructed electron density maps of the modified protein-DNA complex for PDB ID 5Y0D with the same coloring as in **b**.

such a scenario. To test this with one example, I have modified the structure for DNA-wrapped histone (PDB ID 5Y0D), duplicating the DNA molecule to artificially double its contribution to the scattering. This increases the contrast to the point where low-resolution density calculated from the atomic coordinates shows an expected maximum where the DNA is located (Fig. 1c). DENSS reconstructions demonstrate that the final averaged density does indeed reflect the higher expected density at the outer edge of the histone disc where the DNA is located (Fig. 1c). Thus, if given sufficient contrast, DENSS is capable of reconstructing multi-density systems. As noted by Konarev and Svergun, in some circumstances where the density difference between two regions is sufficient to be distinguished by DENSS, bead modeling can approximate the shape as having hollow regions where there is low relative density. However, in such cases, DENSS reproduces actual density where it exists, even if it is low (for example, the protein in this simulation and the hydrophobic core in the DDM micelle), and is thus a more accurate representation of the true object.

Konarev and Svergun also simulate an ellipsoid that is cut in half in different orientations, assigning different contrasts to each half. DENSS reconstructions from simulated data do not discriminate between the different orientations in this case. This is a valid criticism and shows opportunities for future improvements to DENSS. This is likely due to the sharp cutoffs of uniform density in the simulated system, as such unnatural, sharp cutoffs are known to result in aliasing and striping artifacts in image reconstruction, resulting in stagnation of the phase retrieval procedure and falling into local minima. Various advanced algorithms for dealing with such artifacts have been developed^{10,11} and could be incorporated into DENSS. In such cases, default parameters may not be appropriate to yield reliable reconstructions and may need to be adjusted for adequate performance. It is noteworthy that the three distinct ellipsoid models have highly similar scattering profiles, as seen in Supplementary Fig. 2 of the critique, and thus likely represent an example of a fundamental inability of SAXS to distinguish between such similar objects.

Konarev and Svergun's final criticism is that the averaging procedure used for DENSS reconstructions results in a type of

low-pass filter on the maps, producing averages that do not fit the data and thus are unreliable reconstructions. It is true that the averaged maps do not fit the data. However, the assumption that an averaged map is unreliable because it does not fit the data is not correct. Supplementary Fig. 1 shows a plot of the Fourier shell correlation (FSC) between 20 individual reconstructions and the known structure along with the FSC between the averaged map and the known structure (using PDB ID 4FE9 as the example¹², chosen for its highly ambiguous shape as estimated by AMBIMETER¹³). One can clearly see that the averaged density has a better correlation to the actual structure than any of the individual reconstructions, despite the poorer fit of the scattering profile. Additionally, the poor fit of the average to the data is a common occurrence in *ab initio* SAXS reconstructions and is similarly seen when generating an average of multiple bead models. Similarly to how bead modeling programs offer refinement for such purposes (for example, the DAMSTART program¹⁴), DENSS also provides a refinement script for a similar reason (denss.refine.py), although in my tests this does not appear to improve the FSC resolution. Future improvements to DENSS may be realized by incorporating goodness of fit into the averaging procedure, to aid generation of averaged maps that simultaneously fit the data.

The primary aim of the original DENSS publication was to describe a new approach to solving the phase problem for experiments that do not yield full 3D Fourier amplitudes. This approach is not limited to SAXS but can theoretically be used for other experiments where the information content is substantially greater, for example, in fiber diffraction. I used one-dimensional solution scattering as the example as it provided a low-information limiting case showing the power of the technique. I have provided the DENSS software open source so that others may take advantage of it and investigate the algorithm themselves. There are likely cases where DENSS and traditional shape reconstruction methods perform equivalently, cases where DENSS excels and cases where traditional methods excel. Researchers should use the tool most appropriate for their needs.

Online content

Any methods, additional references, Nature Research reporting summaries, source data, extended data, supplementary information, acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41592-021-01083-w>.

Received: 12 March 2019; Accepted: 1 February 2021;

Published online: 1 March 2021

References

- Konarev, P. V. & Svergun, D. I. Limitations of the iterative electron density reconstruction algorithm from solution scattering data. *Nat. Methods* <https://doi.org/10.1038/s41592-021-01082-x> (2021).
- Grant, T. D. Ab initio electron density determination directly from solution scattering data. *Nat. Methods* **15**, 191–193 (2018).
- Garavito, R. M. & Ferguson-Miller, S. Detergents as tools in membrane biochemistry. *J. Biol. Chem.* **276**, 32403–32406 (2001).
- Hyland, L. L., Taraban, M. B. & Yu, Y. B. Using small-angle scattering techniques to understand mechanical properties of biopolymer-based biomaterials. *Soft Matter* <https://doi.org/10.1039/C3SM51209F> (2018).
- Manet, S. et al. Structure of micelles of a nonionic block copolymer determined by SANS and SAXS. *J. Phys. Chem. B* **115**, 11318–11329 (2011).
- Li, T., Senesi, A. J. & Lee, B. Small angle X-ray scattering for nanoparticle research. *Chem. Rev.* **116**, 11128–11180 (2016).
- Gabel, F., Engilberge, S., Perez, J. & Girard, E. Medical contrast media as possible tools for SAXS contrast variation. *IUCr* **6**, 521–525 (2019).
- Spahn, C. M., Penczek, P. A., Leith, A. & Frank, J. A method for differentiating proteins from nucleic acids in intermediate-resolution density maps: cryo-electron microscopy defines the quaternary structure of the *Escherichia coli* 70S ribosome. *Structure* **8**, 937–948 (2000).
- Tang, G. et al. EMAN2: an extensible image processing suite for electron microscopy. *J. Struct. Biol.* **157**, 38–46 (2007).
- Fienup, J. R. & Wackerman, C. C. Phase-retrieval stagnation problems and solutions. *J. Opt. Soc. Am. A* **3**, 1897–1907 (1986).
- Wackerman, C. C. & Yagle, A. E. Use of Fourier domain real-plane zeros to overcome a phase retrieval stagnation. *J. Opt. Soc. Am. A* **8**, 1898–1904 (1991).
- Cameron, E. A. et al. Multidomain carbohydrate-binding proteins involved in *Bacteroides thetaiotaomicron* starch metabolism. *J. Biol. Chem.* **287**, 34614–34625 (2012).
- Petoukhov, M. V. & Svergun, D. I. Ambiguity assessment of small-angle scattering curves from monodisperse systems. *Acta Crystallogr. D Biol. Crystallogr.* **71**, 1051–1058 (2015).
- Volkov, V. V. & Svergun, D. I. Uniqueness of ab-initio shape determination in small-angle scattering. *J. Appl. Crystallogr.* **36**, 860–864 (2003).

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© Springer Nature America, Inc. 2021

Methods

The DDM micelle shown in Fig. 1a was reconstructed using the MEMBRANE mode of DENSS (v1.6.0) to allow for negative contrast and using the default 20 reconstructions for averaging. The simulated maps in Fig. 1b were calculated using the EMAN2 script `e2pdb2mrc.py`, and the resolution was set to the resolution of the ab initio reconstruction calculated by DENSS. The scattering profile of the modified DNA-wrapped histone structure was calculated using FoXS¹⁵, and the reconstruction shown in Fig. 1c was performed using the default SLOW mode of DENSS, with 100 reconstructions for averaging. The scattering profile for PDB ID 4FE9 was calculated using FoXS and reconstructed using the default SLOW mode of DENSS with 20 reconstructions for averaging. EMAN2 was used to align the maps to the atomic coordinates and to calculate the FSC curves shown in Supplementary Fig. 1.

Reporting Summary. Further information on research design is available in the Nature Research Reporting Summary linked to this article.

Data availability

The experimental DDM micelle data were kindly provided by F. Gabel and are available as described in ref. ⁷.

Code availability

DENSS is available open source on GitHub at <https://github.com/tdgrant1/denss>, [git](#), and instructions for downloading, installing and running DENSS can be found on the GitHub page and at <https://www.tdgrant.com/denss/>.

References

15. Schneidman-Duhovny, D., Hammel, M., Tainer, J. A. & Sali, A. Accurate SAXS profile computation and its assessment by contrast variation experiments. *Biophys. J.* **105**, 962–974 (2013).

Acknowledgements

The author thanks F. Gabel for providing SAXS data of the DDM micelle⁷. The author acknowledges support from the BioXFEL NSF Science and Technology Center (NSF 1231306) and from NIH award R01GM133998.

Competing interests

The author declares no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41592-021-01083-w>.

Correspondence and requests for materials should be addressed to T.D.G.

Reprints and permissions information is available at www.nature.com/reprints.

Reporting Summary

Nature Research wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Research policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

n/a Confirmed

- ☒ ☐ The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
- ☒ ☐ A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- ☒ ☐ The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
- ☒ ☐ A description of all covariates tested
- ☒ ☐ A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- ☒ ☐ A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- ☒ ☐ For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted
Give P values as exact values whenever suitable.
- ☒ ☐ For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- ☒ ☐ For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- ☒ ☐ Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

Data collection SAXS data for the DDM micelle was provided by Dr. Frank Gabel and previously published in Reference 5.

Data analysis DENSS v1.6.0 was used for generating the DDM micelle density, using the "MEMBRANE" mode option of DENSS to disable the positivity restraint. 20 reconstructions were averaged together using denss.all.py.

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Research [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A list of figures that have associated raw data
- A description of any restrictions on data availability

SAXS data for the DDM micelle was provided by Dr. Frank Gabel and previously published in Reference 5.

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

☒ Life sciences ☐ Behavioural & social sciences ☐ Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see [nature.com/documents/nr-reporting-summary-flat.pdf](https://www.nature.com/documents/nr-reporting-summary-flat.pdf)

Life sciences study design

All studies must disclose on these points even when the disclosure is negative.

Sample size	N/A
Data exclusions	N/A
Replication	N/A
Randomization	N/A
Blinding	N/A

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems

n/a	Involved in the study
☒	☐ Antibodies
☒	☐ Eukaryotic cell lines
☒	☐ Palaeontology and archaeology
☒	☐ Animals and other organisms
☒	☐ Human research participants
☒	☐ Clinical data
☒	☐ Dual use research of concern

Methods

n/a	Involved in the study
☒	☐ ChIP-seq
☒	☐ Flow cytometry
☒	☐ MRI-based neuroimaging