

ON THE EMERGENCE OF ALMOST-HONEYCOMB STRUCTURES IN LOW-ENERGY PLANAR CLUSTERS

M. CAROCCIA, K. DEMASON, AND F. MAGGI

ABSTRACT. Several commonly observed physical and biological systems are arranged in shapes that closely resemble an honeycomb cluster, that is, a tessellation of the plane by regular hexagons. Although these shapes are not always the direct product of energy minimization, they can still be understood, at least phenomenologically, as low-energy configurations. In this paper, explicit quantitative estimates on the geometry of such low-energy configurations are provided, showing in particular that the vast majority of the chambers must be generalized polygons with six edges, and be closely resembling regular hexagons. Part of our arguments is a detailed revision of the estimates behind the global isoperimetric principle for honeycomb clusters due to Hales [Hal01].

CONTENTS

1. Introduction	1
2. From the quantitative Hales' inequality to the main theorem	9
3. The arc function	11
4. Proof of the quantitative Hales' hexagonal inequality, part one	12
5. Proof of the quantitative Hales' hexagonal inequality, part two	17
Appendix A. The isoperimetric inequality for immersed polygons	28
References	32

1. INTRODUCTION

1.1. Overview. Honeycomb-like structures are commonly observed in physical and biological systems. While in some cases (e.g., in foams) their emergence is consequence of an *energy minimization* property, in many other systems, like those resulting from geological mechanisms (e.g., basalt columns), animal behavior (e.g., beehives) or animal morphology (e.g., compound eyes of arthropods), what is observed results from complex processes leading to the formation of *low-energy* (rather than energy minimizing) configurations. This fact provides the motivation to extend and adapt Hales' isoperimetric theorem [Hal01], which characterizes **honeycomb clusters** (i.e., tessellations of the plane by unit-area, regular hexagons) as the only *energy minimizing* configurations *under periodic boundary conditions*, to the case of low-energy configurations (and without the periodicity restriction). In this paper, partly by a comprehensive review of Hales' methods, we undertake the analysis of such low-energy configurations. In rough terms, our main result (Theorem 1.2 below) states that *every low-energy planar cluster with N -many unit-area, simply connected chambers must be close, in a quantitative way as $N \rightarrow \infty$, to an ideal honeycomb, with a controlled number of defects and with a vast preponderance of almost-hexagonal cells*; and this, without said cluster being required to possess any energy minimizing property.

1.2. Isoperimetric clusters and Hales' theorem. We frame our work in the setting of Almgren's theory of isoperimetric clusters [Alm76] as presented in [Mag12, Part IV]. Since we only consider *planar* clusters, we do not need to work with Borel sets and sets of finite perimeter, and work directly with open sets with Lipschitz boundary $E \subset \mathbb{R}^2$. We

denote by $|E|$ the area (Lebesgue measure) of any such E , and by $P(E) = \mathcal{H}^1(\partial E)$ its perimeter (one-dimensional Hausdorff measure of its topological boundary).

Given $N \in \mathbb{N}$, $N \geq 2$, a planar N -**cluster** is a family $\mathcal{E} = \{\mathcal{E}(h)\}_{h=1}^N$ of mutually disjoint, non-empty open sets $\mathcal{E}(h) \subset \mathbb{R}^2$ (the **chambers** of $\mathcal{E}$) with finite area and Lipschitz regular boundary. The **area** and **perimeter** of an N -cluster $\mathcal{E}$ are then

$$|\mathcal{E}| = (|\mathcal{E}(1)|, \dots, |\mathcal{E}(N)|) \in \mathbb{R}^N, \quad P(\mathcal{E}) = \frac{1}{2} \sum_{h=0}^N P(\mathcal{E}(h)) = \mathcal{H}^1(\partial \mathcal{E}),$$

where $\mathcal{E}(0) = \mathbb{R}^2 \setminus \bigcup_{h=1}^N \overline{\mathcal{E}(h)}$ denotes the **exterior chamber** of $\mathcal{E}$ (so that $|\mathcal{E}(0)| = +\infty$), and where $\partial \mathcal{E} = \bigcup_{h=1}^N \partial \mathcal{E}(h)$ is the **boundary** of $\mathcal{E}$. We say that $\mathcal{E}$ is an **isoperimetric cluster** if

$$P(\mathcal{E}) \leq P(\mathcal{F}), \quad \text{for all clusters } \mathcal{F} \text{ with } |\mathcal{F}| = |\mathcal{E}|;$$

and a **unit-area isoperimetric cluster** if, in addition, $|\mathcal{E}(h)| = 1$ for all $h = 1, \dots, N$.

Notice that the chambers of an N -cluster are not assumed to be connected. A disjoint family of N' -many open, connected subsets of $\mathbb{R}^2$ with Lipschitz boundaries could thus be regarded (in more than one way) as an N -cluster for different values of $N \leq N'$. From a conceptual viewpoint, this is clearly a somehow flawed feature of the notion of N -cluster used here. However, this feature is also crucially important for proving Almgren's existence theorem [Alm76] for isoperimetric clusters: *for every $v \in \mathbb{R}^N$ with positive coordinates, there exist isoperimetric N -clusters $\mathcal{E}$ with $|\mathcal{E}| = v$; see [Mag12, Chapter 29].*

This said, we have the following natural **connectedness conjecture** about isoperimetric clusters:

$$\textit{isoperimetric clusters have simply connected chambers.} \quad (\text{CC})$$

This conjecture is widely open. Its validity has been confirmed in the few cases ($N = 2$ [FAB⁺93], $N = 3$ [Wic04], and $N = 4$ with equal areas [PT18, PT20]) where a complete classification of isoperimetric clusters is known. One does not expect to carry this approach much further, as obtaining characterizations of isoperimetric clusters with arbitrary values of N and v seems out of question. The two basic facts about isoperimetric clusters that are known for every N and v are the validity of **Plateau laws** (boundaries of isoperimetric clusters consist of finitely many circular arcs/segments meeting in threes at 120-degrees; see, e.g. [Mag12, Theorem 30.7]), and the local finiteness and constancy in v of the possible diffeomorphic types of $\partial \mathcal{E}$; see [CLM16, Theorem 1.9].

In his celebrated paper [Hal01], Hales presents an argument that serves to prove two isoperimetric principles concerning “honeycombs”. The first one states that, if $a, b > 0$ are such that the flat torus $\mathbb{R}^2[a, b]$ of width a and height b admits a tiling $\mathcal{H}$ by N unit-area, regular hexagons, then $\mathcal{H}$ is the unique unit-area isoperimetric N -cluster (modulo translations); see [CM16] for a quantitative analysis of this isoperimetric principle. The second one states that if $\mathcal{E}$ is a unit-area isoperimetric N -cluster in $\mathbb{R}^2$, then¹

$$\psi(N) > (12)^{1/4} N, \quad \forall N \geq 2, \quad (1.1)$$

where we have set

$$\begin{aligned} \psi(N) &= P(\mathcal{E}_N), \\ (\mathcal{E}_N \text{ a generic unit-area isoperimetric } N\text{-cluster}). \end{aligned}$$

¹By a slight refinement of Hales' argument, (1.1) can actually be improved to $\psi(N) \geq (12)^{1/4} N + K_0 \sqrt{N}$ with $K_0 = \sqrt{\pi} - \sqrt[4]{3}/\sqrt{2} \approx 0.84$. Such refinement simply consists in applying “Dido's inequality” to quantify the size of some non-negative terms that were just discarded in Hales' original presentation; see the proof of [HM04, Theorem 2.1].




FIGURE 1.1. The idea behind the low-energy condition (1.3) is that it identifies unit-area N -clusters whose “internal perimeter” is comparable to that of an $\sqrt{N} \times \sqrt{N}$ -chunk of ideal honeycomb, and whose “external perimeter” is comparable to $\sqrt{N}$ (i.e., the square root of the area of the bulk of the cluster). Unit-area locally minimizing clusters may fail to satisfy this condition. For example, the N -cluster depicted here satisfies, for some $r_0 > 0$, the local isoperimetric property $P(\mathcal{E}) \leq P(\mathcal{F})$ for every $\mathcal{F}$ with $|\mathcal{F}| = |\mathcal{E}|$ with $\text{diam}(\mathcal{F}(h) \Delta \mathcal{E}(h)) \leq r_0$, $h = 1, \dots, N$; but it does not satisfy (1.3) – as it can be seen, for example, by looking at its external perimeter, which is $O(N)$, compare with (1.13) in Theorem 1.2.

The energy bound (1.1) is the only evidence towards a second fascinating, challenging, and largely unexplored **honeycomb conjecture**:

$$\begin{aligned} &\text{unit-area isoperimetric } N\text{-clusters with } N \text{ large} \\ &\text{should locally coincide with honeycombs.} \end{aligned} \tag{HC}$$

To understand the connection between (1.1) and the honeycomb geometry, we notice that, starting from a unit-area regular hexagon H , one can add a first complete layer of unit-area regular hexagons around H (which, when complete, results in a cluster of $N = 7$ unit-area regular hexagons), then a second complete layer (which, when complete, results in a cluster of $N = 19$ unit-area regular hexagons), and so on. For every $N \geq 2$ one can smooth out the exterior edges of such clusters into circular arcs. The resulting construction, detailed in [HM04, Theorem 2.1], gives

$$\psi(N) \leq (12)^{1/4} N + M_0 \sqrt{N} + 3, \quad \forall N \geq 2, \tag{1.2}$$

(where $M_0 = \pi/A_0^{1/2} \approx 1.95$ and $A_0 = 3\sqrt{3}/2$ is the area of a regular hexagon of unit side length). The upper bound (1.2) on $\psi(N)$ implies the sharpness of (1.1), and clarifies its connection with the honeycomb geometry.

1.3. Low-energy clusters and main result. Motivated by (1.2), we say that a planar, unit-area N -cluster $\mathcal{E}$ is a **low-energy cluster with exterior energy density M** , if

$$P(\mathcal{E}) \leq (12)^{1/4} N + M \sqrt{N}. \tag{1.3}$$

This condition amounts in asking that that the “bulk” of the cluster $\mathcal{E}$, that is, the set $E = \mathbb{R}^2 \setminus \mathcal{E}(0)$, has perimeter $P(E) = O(\sqrt{|E|})$, while the “internal perimeter” of $\mathcal{E}$, that is, the length of $\partial\mathcal{E}$ due to interfaces $\partial\mathcal{E}(h) \cap \partial\mathcal{E}(k)$ with $h \neq k$, $h, k \neq 0$, is approximately that of an $\sqrt{N} \times \sqrt{N}$ -chunk of the ideal honeycomb $\mathcal{H}$.

By (1.2), unit-area isoperimetric clusters satisfy (1.3). At the same time, (1.3) rules out many N -clusters that are only *locally*, but not globally, isoperimetric, and that may fail to look like honeycombs; see Figure 1.1.

In summary, (1.3) identifies a large class of N -clusters which, although not possessing any isoperimetric property, may still be expected, for N large, to be close to honeycombs. Our main result confirms this expectation in a class $\mathcal{C}(N, M)$ of planar unit-area N -clusters that *contains every unit-area isoperimetric N -cluster with simply connected chambers* (i.e., every unit-area isoperimetric cluster, should conjecture (CC) hold true); see Remark 1.1 below.

We say that a unit-area N -cluster $\mathcal{E}$ in $\mathbb{R}^2$ belongs to $\mathcal{C}(N, M)$ if:

- (C1) each $\mathcal{E}(h)$ has finitely many connected components $\{E_j^h\}_{j=1}^{N_h}$, called **cells** (of $\mathcal{E}(h)$ or of $\mathcal{E}$, depending on the context); each cell of $\mathcal{E}$ is simply connected;
- (C2) the boundary of each cell E_j^h consists of finitely many bounded Lipschitz injective curves, called **edges** (of E_j^h or of $\mathcal{E}$, depending on the context); endpoints of edges are called **vertexes**;
- (C3) each vertex is the endpoint of *exactly three* edges, and each cell has *at least two* edges;
- (C4) the boundary $\partial\mathcal{E}$ of $\mathcal{E}$ is connected and satisfies (1.3), that is

$$P(\mathcal{E}) \leq (12)^{1/4} N + M \sqrt{N};$$

- (C5) denoting by $k_j^h = k_j^h(\mathcal{E})$ the number of edges of the cell E_j^h of $\mathcal{E}$, we require that

$$|E_j^h| \geq \frac{1}{100}, \quad \text{if } 2 \leq k_j^h \leq 6. \quad (1.4)$$

Remark 1.1 (The class $\mathcal{C}(N, M)$ and isoperimetric clusters). If $\mathcal{E}$ is a unit-area N -isoperimetric cluster, then: (i) each chamber of $\mathcal{E}$ has finitely many connected components, so that the first part of (C1) holds true; (ii) each cell of $\mathcal{E}$ is bounded by finitely many circular arcs/segments, therefore (C2) holds true; (iii) the validity of (C3) is a simple consequence of the Plateau laws; (iv) a sliding argument shows that $\partial\mathcal{E}$ is connected, so that $\mathcal{E}$ satisfies (C4) thanks to (1.2). As a consequence, if each chamber $\mathcal{E}(h)$ of $\mathcal{E}$ has just one single (simply connected) component E_1^h , then (C1) and (C5) hold true (with $|E_1^h| = 1$), and $\mathcal{E} \in \mathcal{C}(N, M)$ for some M . In summary, *every unit-area N -isoperimetric cluster with simply connected chambers* is an element of $\mathcal{C}(N, M)$ for some M .

As a last step towards the statement of our main result we introduce the following notation. For $\mathcal{E} \in \mathcal{C}(N, M)$ we introduce the family $\text{Hex}(\mathcal{E})$ of those (indexes of) chambers of $\mathcal{E}$ that **are connected and have six edges**,

$$\text{Hex}(\mathcal{E}) := \left\{ h : N_h = 1, k_1^h = 6 \right\}, \quad (1.5)$$

and, for each $k \geq 2$, the family $\text{Ch}_k(\mathcal{E})$ of those (indexes of) chambers of $\mathcal{E}$ **having at least one cell with k -sides**

$$\text{Ch}_k(\mathcal{E}) := \left\{ h : \exists j \in \{1, \dots, N_h\} \text{ s.t. } k_j^h = k \right\}. \quad (1.6)$$

We also define the **exterior perimeter of $\mathcal{E}$** by setting

$$P_{\text{ext}}(\mathcal{E}) := P(E_1^0), \quad (1.7)$$

where, we recall, E_1^0 is the unique connected component of the exterior chamber $\mathcal{E}(0)$ with infinite area; the **number of exterior edges of $\mathcal{E}$** is defined by setting

$$\text{edge}_{\text{ext}}(\mathcal{E}) := k_1^0; \quad (1.8)$$

and the **interior void of $\mathcal{E}$** as

$$\mathcal{E}_{\text{void}} := \mathcal{E}(0) \setminus E_1^0. \quad (1.9)$$

(Notice that this set must have finite area). Finally, we denote by

$$d_{\text{hex}}(E) = \inf \left\{ |E\Delta(x + re^{i\theta}[H])| : x \in \mathbb{R}^2, \theta \in [0, 2\pi], r^2 = |E| \right\}, \quad (1.10)$$

the distance in area of $E \subset \mathbb{R}^2$ from a regular hexagon with area $|E|$ (here H denotes a reference unit-area regular hexagon and $e^{i\theta}$ the angle θ counter-clockwise rotation of $\mathbb{R}^2$ around the origin). With this terminology in place, we state our main theorem.

Theorem 1.2 (Honeycomb-like structure of low-energy clusters). *There exists a computable constant C_0 with the following property. If $N \geq 2$ and $\mathcal{E} \in \mathcal{C}(N, M)$, then*

$$\# \text{Hex}(\mathcal{E}) \geq N - C_0 M \sqrt{N}, \quad (1.11)$$

with

$$\frac{1}{\# \text{Hex}(\mathcal{E})} \sum_{h \in \text{Hex}(\mathcal{E})} d_{\text{hex}}(\mathcal{E}(h))^2 \leq \frac{C_0 M}{\sqrt{N}}, \quad (1.12)$$

and

$$P_{\text{ext}}(\mathcal{E}) \leq C_0 M \sqrt{N}, \quad (1.13)$$

$$\text{edge}_{\text{ext}}(\mathcal{E}) \leq C_0 M \sqrt{N}, \quad (1.14)$$

$$|\mathcal{E}_{\text{void}}| \leq C_0 M \sqrt{N}, \quad (1.15)$$

$$\# \text{Ch}_k(\mathcal{E}) \leq \frac{C_0 M}{|k - 6|} \sqrt{N}. \quad (1.16)$$

Finally, there is at least one $k \leq 5$ such that $\text{Ch}_k(\mathcal{E}) \neq \emptyset$.

Remark 1.3. Conclusions (1.11) and (1.12) state that the vast majority of the chambers of $\mathcal{E}$ are (simply) connected, possess six curvilinear edges, and are close to being regular hexagons. Conclusion (1.13) implies that only a very small fraction of the total cluster perimeter $P(\mathcal{E}) = O(N)$ is used to compound the bulk of the chambers of $\mathcal{E}$. Given conclusion (1.13), conclusion (1.14) indicates that the generic external edge of $\mathcal{E}$ must have length of order one. Simple examples show that *all these conclusions are sharp* in the class $\mathcal{C}(N, M)$.

Remark 1.4 (Connection with the connectedness conjecture). Thanks to Remark 1.1, should one be able to prove the validity of the area lower bound (1.4) for the cells of isoperimetric clusters, then conclusion (1.11) in Theorem 1.2 would imply, in particular, a partial answer to the connectedness conjecture (CC), namely, that for a unit-area isoperimetric N -cluster, $(N - O(\sqrt{N}))$ -many chambers are connected.

1.4. Hales' hexagonal isoperimetric inequality and strategy of proof. The proof of Theorem 1.2 is based on a careful extension of the methods developed by Hales in [Hal01]. The key result in Hales' paper is an inequality for immersed planar curves, called here *Hales' hexagonal isoperimetric inequality*, see (1.27) below. The key step in proving Theorem 1.2 is obtaining a quantitative improvement of Hales' hexagonal isoperimetric inequality. Thus, in order to illustrate our strategy of proof, we need to introduce Hales' hexagonal isoperimetric inequality.

Hales' hexagonal isoperimetric inequality is a direct improvement of the hexagonal isoperimetric inequality. Denoting by Pol_k the family of planar polygons with k -sides (k -polygons) and by $p(k)$ the perimeter of a reference unit-area regular k -polygon, the **k -polygonal isoperimetric inequality** states that

$$P(\Pi) \geq p(k) \sqrt{|\Pi|}, \quad \forall \Pi \in \text{Pol}_k, \quad (1.17)$$

with equality if and only if Π is a regular k -polygon. Here $p(k)$ is explicitly given by

$$p(k) = 2 \sqrt{k \tan(\pi/k)}. \quad (1.18)$$

The case $k = 6$ of (1.17) is of course the **hexagonal isoperimetric inequality**,

$$P(\Pi) \geq 2(12)^{1/4} \sqrt{|\Pi|}, \quad \forall \Pi \in \text{Pol}_6, \quad (1.19)$$

with equality if and only if Π is a regular hexagon. Now, on noticing that $t \mapsto 2\sqrt{t \tan(\pi/t)}$ is strictly decreasing and convex, and setting

$$a(k) = \frac{p(k) - p(6)}{6 - k}, \quad \forall k \neq 6,$$

(so that $a(k) > 0$), we can deduce from (1.17) that if $a \in (a(7), a(5))$, then

$$P(\Pi) + a(k - 6) \geq 2(12)^{1/4} \sqrt{|\Pi|}, \quad \forall \Pi \in \bigcup_{k=3}^{\infty} \text{Pol}_k, \quad (1.20)$$

with equality if and only if Π is a regular hexagon. This is of course a rather trivial generalization of (1.19), but it is already sufficient to prove the lower bound $P(\mathcal{E}) > (12)^{1/4} N$ on the special class of **unit-area polygonal Plateau-type N -clusters $\mathcal{E}$** whose chambers have unit area and consist of finitely many connected polygonal cells, and whose vertexes are the endpoints of *exactly three segments*. Indeed, since all the cells E_j^h of $\mathcal{E}$ with $(h, j) \neq (0, 1)$ are polygons with finite area, we can apply (1.20) on them. Denoting by k_j^h the number of sides of E_j^h , we thus find that

$$2P(\mathcal{E}) \geq \sum_{(h,j) \neq (0,1)} P(E_j^h) \geq 2(12)^{1/4} \sum_{(h,j) \neq (0,1)} \sqrt{|E_j^h|} + a \sum_{(h,j) \neq (0,1)} (6 - k_j^h). \quad (1.21)$$

On the one hand, since $|E_j^h| \leq |\mathcal{E}(h)| = 1$ for $h = 1, \dots, N$ implies $|E_j^h| \leq \sqrt{|E_j^h|}$, we get

$$\sum_{(h,j) \neq (0,1)} \sqrt{|E_j^h|} = \sum_{j=2}^{N_0} \sqrt{|E_j^0|} + \sum_{(h,j), h \neq 0} \sqrt{|E_j^h|} \geq \sum_{j=2}^{N_0} \sqrt{|E_j^0|} + \sum_{h=1}^N |\mathcal{E}(h)| \geq N. \quad (1.22)$$

On the other hand, counting that each vertex of $\partial\mathcal{E}$ is the endpoint of three segments, we can associate to $\partial\mathcal{E}$ a regular graph on $\mathbb{S}^2$ with numbers of faces F , edges E , and vertexes V given by

$$F = \sum_{h=0}^N N_h, \quad E = \frac{1}{2} \sum_{h=0}^N \sum_{j=0}^{N_h} k_j^h, \quad V = \frac{2}{3} E,$$

so that, by Euler's formula $2 = V - E + F$,

$$2 = -\frac{1}{6} \sum_{h=0}^N \sum_{j=0}^{N_h} k_j^h + \sum_{h=0}^N N_h, \quad \text{that is} \quad 12 = \sum_{h=0}^N \sum_{j=1}^{N_h} (6 - k_j^h),$$

and thus

$$\sum_{(h,j) \neq (0,1)} (6 - k_j^h) = 6 + k_1^0. \quad (1.23)$$

The combination of $a > 0$, (1.21), (1.22), and (1.23) gives $P(\mathcal{E}) > (12)^{1/4} N$ for every unit-area polygonal Plateau-type N -cluster $\mathcal{E}$ as defined above.

The above argument, due to Fejes Tóth [FT43], has been presented in detail since the class of unit-area isoperimetric N -clusters $\mathcal{E}$ considered in Hales' isoperimetric principle (1.1) is actually not that far from the class of unit-area polygonal Plateau-type N -clusters: the only difference is that, in the former case, cells may be bounded by circular arcs (with possible non-zero curvature) rather than just by segments (as in the latter). Hales' hexagonal isoperimetric inequality consists of an extension of (1.20) that suffices to repeat Fejes Tóth's argument on isoperimetric clusters.

The following notation will be needed in stating Hales' hexagonal isoperimetric inequality. Given $s, t \in \mathbb{S}^1 = \{z \in \mathbb{C} : |z| = 1\}$, $s \neq t$, we denote by $[s, t]$ the set of points in $\mathbb{S}^1$ obtained by moving from s to t in the orientation of $\mathbb{S}^1$ induced by $e^{i\theta}$. We say that I is




FIGURE 1.2. In the example in the picture, the quantity $\alpha(\gamma, s, t)$ defined in (1.24) is obtained by subtracting the areas depicted in light grey from the areas depicted in dark grey.

an **interval** of $\mathbb{S}^1$, if $I = \mathbb{S}^1$ or $I = [s, t]$ for some $s \neq t$. If $\gamma \in \text{Lip}(I; \mathbb{R}^2)$, then we denote by $L(\gamma)$ and $A(\gamma)$ the **length** and **oriented area** of γ , defined by setting

$$L(\gamma) = \int_I |\gamma'(t)| dt, \quad A(\gamma) = \int_\gamma x dy = \int_I \gamma^{(1)}(t) (\gamma^{(2)})'(t) dt.$$

(Here and in the following $x^{(i)}$ is the i -th component of $x \in \mathbb{R}^2$. Also, with this convention, if $\gamma(\theta) = e^{i\theta}$ ($\theta \in \mathbb{S}^1$), then $A(\gamma) = \pi$, and if $\gamma(\theta) = e^{-i\theta}$ ($\theta \in \mathbb{S}^1$), then $A(\gamma) = -\pi$.) Given $x, y \in \mathbb{R}^2$ we denote by $\llbracket x, y \rrbracket$ a constant speed parametrization (over a closed interval in $\mathbb{S}^1$) of the segment $\{tx + (1-t)y : t \in [0, 1]\}$. With this notation, given $\gamma \in \text{Lip}(\mathbb{S}^1; \mathbb{R}^2)$ and $[s, t] \subset \mathbb{S}^1$, we introduce the **signed secant area of γ over $[s, t]$** by setting

$$\alpha(\gamma, s, t) = A(\gamma|_{[s,t]}) + A(\llbracket \gamma(t), \gamma(s) \rrbracket); \quad (1.24)$$

see Figure 1.4. Finally, given $k \geq 2$, we say that $\mathbf{t} : \{1, \dots, k\} \rightarrow \mathbb{S}^1$ defines an **ordered k -partition** of $\mathbb{S}^1$ if, for every $i = 1, \dots, k-1$, $[\mathbf{t}(i), \mathbf{t}(i+1)]$ does not contain any $\mathbf{t}(j)$ with $j \neq i, i+1$. Denoting by $\mathcal{P}_k$ the set of ordered k -partitions of $\mathbb{S}^1$, and setting $\mathbf{t}(k+1) = \mathbf{t}(1)$ if $\mathbf{t} \in \mathcal{P}_k$, we define a functional

$$\sigma : \text{Lip}(\mathbb{S}^1; \mathbb{R}^2) \times \mathcal{P}_k \rightarrow \mathbb{R},$$

by taking

$$\sigma(\gamma, \mathbf{t}) = \sum_{i=1}^k \min \left\{ \frac{1}{2}, \max \left\{ -\frac{1}{2}, \alpha(\gamma, \mathbf{t}(i), \mathbf{t}(i+1)) \right\} \right\}. \quad (1.25)$$

Thus $\sigma(\gamma, \mathbf{t})$ is obtained by adding up the signed secant areas defined by γ and the nodes of the ordered k -partition $\mathbf{t}$, truncated in the interval $[-1/2, 1/2]$.

Theorem A (Hales' hexagonal isoperimetric inequality, [Hal01]). *There exists a constant $a > 0$ with the following property. If $k \geq 2$, $(\gamma, \mathbf{t}) \in \text{Lip}(\mathbb{S}^1; \mathbb{R}^2) \times \mathcal{P}_k$, and*

$$A(\gamma) \geq \frac{2\pi}{\sqrt{3}k^2}, \quad (1.26)$$

then

$$L(\gamma) + a(k-6) + (12)^{1/4} \sigma(\gamma, \mathbf{t}) \geq 2(12)^{1/4} \min\{1, A(\gamma)\} \quad (1.27)$$

with equality if and only if $(\gamma, \mathbf{t})$ corresponds² to a unit-area regular hexagon.

Remark 1.5. (i): Hales proves (1.27) with $a = 0.0505$. Notice that this choice satisfies $a \in (a(7), a(5))$, and is thus admissible in (1.20). In particular, when γ parameterizes the boundary of a unit-area polygon and $\mathbf{t}$ ranges through the vertexes of that polygon, then (1.27) boils down to (1.20); **(ii):** One cannot take a different value than $(12)^{1/4}$ for the

²That is, γ is an injective parametrization of the boundary of a unit-area regular hexagon and $\{\gamma(\mathbf{t}(i))\}_{i=1}^6$ are the vertexes of said hexagon.

constant in front of $\sigma(\gamma, \mathbf{t})$: this can be verified by taking $k = 6$, γ a parametrization of a unit-area regular hexagon H , $\mathbf{t}$ ranging through the vertexes of H , and then modifying γ by taking small circular variations of the edges of H and subsequent rescalings; **(iii)**: The inequality cannot hold with $A(\gamma)$ in place of $\min\{1, A(\gamma)\}$ because of the different scaling of $L(\gamma)$ and $A(\gamma)$ – of course, a variant of (1.27) with $\sqrt{A(\gamma)}$ in place of $\min\{1, A(\gamma)\}$ would have looked more natural; **(iv)**: taking $k < 6$, and testing (1.27) with $t\gamma$ in place of γ , we eventually obtain a contradiction as $t \rightarrow 0^+$; this is why a lower bound on $A(\gamma)$ must be assumed.

Remark 1.6 (Proof of (1.1) and assumption (1.26)). It seems useful to recall how Theorem A is used in proving (1.1), as this point allows us to illustrate the reason for the particular form (1.26) of the lower bound on $A(\gamma)$ assumed in Theorem A. To this end, let us introduce the functional

$$\delta(\mathcal{F}) = P(\mathcal{F}) - (12)^{1/4} \sum_{h=1}^N \min\{1, |\mathcal{F}(h)|\}, \quad \mathcal{F} \text{ an } N\text{-cluster}.$$

When $\mathcal{E}$ is a unit-area isoperimetric N -cluster, then $\delta(\mathcal{E}) = P(\mathcal{E}) - (12)^{1/4} N$, and (1.1) is equivalent to $\delta(\mathcal{E}) > 0$. As Hales argues in [Hal01, Remark 2.7], if a cell E_j^h of $\mathcal{E}$ is such that $|E_j^h| < 2\pi/(\sqrt{3}(k_j^h)^2)$, then a new N -cluster $\mathcal{F}$ can be defined so that $\delta(\mathcal{F}) \leq \delta(\mathcal{E})$ and $\mathcal{F}(h)$ has one less cell than $\mathcal{E}(h)$ (one simply has to add E_j^h to a cell E_i^m , $m \neq h$, sharing with E_j^h its longest edge). In finitely many steps, one constructs an N -cluster Γ such that $\delta(\Gamma) \leq \delta(\mathcal{E})$, each cell of Γ is simply connected and satisfies $|G_j^h| \geq 2\pi/(\sqrt{3}(k_j^h)^2)$. Theorem A can then be applied to each cell of Γ , and, adding up over the number of cells, Fejes Tóth's argument can be repeated with minor modifications to conclude that $\delta(\Gamma)$, thus $\delta(\mathcal{E})$, is positive (compare with the proof of Theorem 1.2).

Coming back to the presentation of the proof of our main result, Theorem 1.2, the key ingredient will be obtaining the following *quantitative improvement* of Hales' hexagonal isoperimetric inequality (1.27).

Theorem 1.7 (A quantitative Hales' hexagonal isoperimetric inequality). *There exist positive constants a_1 , a_2 , and a_3 such that, setting $a_3(k) = 0$ if $k \neq 6$ and $a_3(6) = a_3$, then the following holds. If $k \geq 2$, $(\gamma, \mathbf{t}) \in \text{Lip}(\mathbb{S}^1; \mathbb{R}^2) \times \mathcal{P}_k$, γ is injective, $A(\gamma) \leq 1$, and*

$$A(\gamma) \geq \frac{1}{100}, \quad \text{if } 2 \leq k \leq 6, \quad (1.28)$$

then

$$\begin{aligned} L(\gamma) + a_1(k-6) + (12)^{1/4} \sigma(\gamma, \mathbf{t}) &\geq 2(12)^{1/4} A(\gamma) \\ &\quad + a_2|k-6| \\ &\quad + a_3(k) \left\{ d_{\text{hex}}(E_\gamma)^2 + (1 - A(\gamma)) \right\}, \end{aligned} \quad (1.29)$$

where E_γ denotes the bounded connected component of $\mathbb{R}^2 \setminus \gamma(\mathbb{S}^1)$ defined by γ according to Jordan's theorem.

Remark 1.8. **(i)**: With respect to assumption (1.26) of Theorem A, we assume no lower bound on $A(\gamma)$ when $k \geq 7$, and we weaken the lower bound on $A(\gamma) \geq 2\pi/(\sqrt{3}k^2)$ when $2 \leq k \leq 6$ (indeed $2\pi/(\sqrt{3}(6^2)) > 1/10$); **(ii)**: A more cosmetic than substantial change is the replacement of the term $2(12)^{1/4} \min\{1, A(\gamma)\}$ in (1.27) with the term $2(12)^{1/4} A(\gamma)$ in (1.29). By Remark 1.5-(iii), this change calls for adding the assumption $A(\gamma) \leq 1$; **(iii)**: The second and third term on the right-hand side of (1.29) quantify the distance of $(\gamma, \mathbf{t})$ from corresponding to a unit-area regular hexagon (unique equality cases of (1.27)).

1.5. Organization of the paper. In Section 2, Theorem (1.2) is deduced from Theorem 1.7. After a brief introduction of the arc function in Section 3, the rest of the paper is thus concerned with the proof of Theorem 1.7, which is broken down in two sections. In Section 4 we present the part of the argument making use of quantitative isoperimetry methods, while in Section 5 we present the part of the argument which follows more closely [Hal01].

Acknowledgement: MC has been partially supported by the fund “Analisi di problemi variazionali e differenziali, teoria degli operatori (F.S.R Politecnico di Milano)”. MC thanks the financial support of PRIN 2022R537CS “Nodal optimization, nonlinear elliptic equations, nonlocal geometric problems, with a focus on regularity” funded by the European Union under Next Generation EU. KDM is supported by NSF-DMS RTG 1840314 and the NSF Graduate Research Fellowship Program under Grant DGE 2137420. FM is supported by NSF-DMS RTG 1840314 and NSF-DMS 2247544. This work was partially completed while KDM was hosted at the math department of Politecnico di Milano.

2. FROM THE QUANTITATIVE HALES’ INEQUALITY TO THE MAIN THEOREM

In this section we show how to deduce Theorem 1.2 from Theorem 1.7. We shall use “Dido’s inequality”: *if a curve of length ℓ bounds an area a with its chord, then $\ell \geq \sqrt{2\pi a}$.* This is easily proved by reflecting the circular arc with respect to the chord and by applying the isoperimetric inequality to find $2\ell \geq 2\sqrt{\pi}\sqrt{2a}$.

Proof of Theorem 1.2. Let $\mathcal{E} \in \mathcal{C}(N, M)$, let $\{E_j^h\}_{j=1}^{N_h}$ be the cells of the chamber $\mathcal{E}(h)$, and let k_j^h be the number of vertexes of E_j^h (see the defining properties (C1)–(C5) of $\mathcal{C}(N, M)$ for the notation and terminology used here). For each cell E_j^h we can find an injective $\gamma_j^h \in \text{Lip}(\mathbb{S}^1; \mathbb{R}^2)$ and $\mathbf{t}_j^h \in \mathcal{P}_{k_j^h}$ such that:

- (i): $\gamma_j^h(\mathbf{t}_j^h(i))$ is the i -th vertex of E_j^h ;
- (ii): if $(h, j) \neq (0, 1)$ (and thus E_j^h is not the cell E_1^0 of $\mathcal{E}$ with infinite area), then the orientation of γ_j^h can be chosen so that $A(\gamma_j^h) = |E_j^h|$ and $L(\gamma_j^h) = P(E_j^h)$;
- (iii): the orientation of γ_1^0 is such that

$$\sum_{(h,j) \neq (0,1)} \sigma(\gamma_j^h, \mathbf{t}_j^h) = -\sigma(\gamma_1^0, \mathbf{t}_1^0), \quad (2.1)$$

where we are using the cancelations due to opposite orientations along each internal edge of $\mathcal{E}$.

By (ii), since $P(E_j^h) = L(\gamma_j^h)$, we notice that

$$2P(\mathcal{E}) = P(E_1^0) + \sum_{(h,j) \neq (0,1)} L(\gamma_j^h), \quad (2.2)$$

and bound from below the two terms on the right-hand side as follows. Concerning $P(E_1^0)$, setting for brevity

$$A_i = \alpha(\gamma_1^0, \mathbf{t}_1^0(i), \mathbf{t}_1^0(i+1)),$$

(see (1.24)) and applying Dido’s inequality, we find that

$$\frac{P(E_1^0)}{\sqrt{2\pi}} \geq \sum_{i=1}^{k_1^0} \sqrt{|A_i|} \geq \sum_{i=1}^{k_1^0} \sqrt{\min\{1/2, |A_i|\}} \geq \sum_{i=1}^{k_1^0} \min\{1/2, |A_i|\} \geq |\sigma(\gamma_1^0, \mathbf{t}_1^0)|. \quad (2.3)$$

Assumption (1.28) of Theorem 1.7 holds for each γ_j^h with $(h, j) \neq (0, 1)$ thanks to property (C5). We can thus apply (1.29) to each γ_j^h and adding up the results we find

$$\begin{aligned} \sum_{(h,j) \neq (0,1)} L(\gamma_j^h) &\geq \sum_{(h,j) \neq (0,1)} \left\{ a_1 (6 - k_j^h) - (12)^{1/4} \sigma(\gamma_j^h, \mathbf{t}_j^h) \right\} \\ &+ \sum_{(h,j) \neq (0,1)} \left\{ 2(12)^{1/4} |E_j^h| + a_2 |k_j^h - 6| + a_3(k_j^h) \left(d_{\text{hex}}(E_j^h)^2 + (1 - |E_j^h|) \right) \right\}. \end{aligned} \quad (2.4)$$

Property (C3) of $\mathcal{C}(N, M)$ ensures that we can repeat Fejes Tóth's argument to deduce (1.23), that is $\sum_{(h,j) \neq (0,1)} (6 - k_j^h) = 6 + k_1^0$. Combining this last identity,

$$\sum_{(h,j) \neq (0,1)} |E_j^h| = N + |\mathcal{E}_{\text{void}}|$$

(recall (1.9)), and (2.1) into (2.4), and then recalling (2.2), we find that

$$\begin{aligned} 2P(\mathcal{E}) &\geq 2(12)^{1/4} N + 2(12)^{1/4} |\mathcal{E}_{\text{void}}| + P(E_1^0) + (12)^{1/4} \sigma(\gamma_1^0, \mathbf{t}_1^0) + a_1 (6 + k_1^0) \\ &+ \sum_{(h,j) \neq (0,1)} \left\{ a_2 |k_j^h - 6| + a_3(k_j^h) \left(d_{\text{hex}}(E_j^h)^2 + (1 - |E_j^h|) \right) \right\}. \end{aligned} \quad (2.5)$$

Now, taking into account that $\sqrt{2\pi} - (12)^{1/4} > 0$, thanks to (2.3) we find that

$$P(E_1^0) + (12)^{1/4} \sigma(\gamma_1^0, \mathbf{t}_1^0) \geq 2c_0 P(E_1^0), \quad \text{if} \quad c_0 = \frac{\sqrt{2\pi} - (12)^{1/4}}{2\sqrt{2\pi}}. \quad (2.6)$$

By combining (2.5) and (2.6) with the low-energy condition $P(\mathcal{E}) \leq (12)^{1/4} N + M\sqrt{N}$, we immediately deduce

$$2(12)^{1/4} |\mathcal{E}_{\text{void}}| + a_1 (6 + k_1^0) + c_0 P(E_1^0) + a_2 \sum_{(h,j) \neq (0,1)} |k_j^h - 6| \leq 2M\sqrt{N}, \quad (2.7)$$

$$\sum_{(h,j) \neq (0,1)} a_3(k_j^h) \left(d_{\text{hex}}(E_j^h)^2 + (1 - |E_j^h|) \right) \leq 2M\sqrt{N}. \quad (2.8)$$

Conclusions (1.13), (1.14), (1.15), and (1.16) follow immediately from (2.7) and

$$|k - 6| \#(\text{Ch}_k(\mathcal{E})) \leq \sum_{(h,j) \neq (0,1)} |k_j^h - 6|.$$

To prove (1.11) and (1.12), we begin by noticing that $\text{Hex}(\mathcal{E}) = I_1 \cap I_2$ where

$$\begin{aligned} I_1 &= \left\{ h : k_j^h = 6, \forall j \in \{1, \dots, N_h\} \right\}, \\ I_2 &= \left\{ h : N_h = 1 \right\}. \end{aligned}$$

Now, by (2.7), setting $I = \{1, \dots, N\}$, we find

$$\#(I \setminus I_1) \leq \sum_{\{(h,j): k_j^h \neq 6\}} |k_j^h - 6| \leq \sum_{(h,j) \neq (0,1)} |k_j^h - 6| \leq C M \sqrt{N},$$

and, similarly, by (2.8), we obtain

$$\begin{aligned} C M \sqrt{N} &\geq \sum_{\{(h,j) \neq (0,1): k_j^h = 6\}} (1 - |E_j^h|) \geq \sum_{h \in I_1} \sum_{j=1}^{N_h} (1 - |E_j^h|) \\ &= \sum_{h \in I_1} (N_h - 1) \geq \#(I_1 \setminus I_2). \end{aligned}$$




FIGURE 3.1. (a) $\text{arc}(\ell, x)$ is defined as the length of a circular arc (depicted in bold) subtending a chord of length ℓ and including a secant area x ; (b) An implicit formula for arc_1 on the interval $[0, \pi/2]$ can be obtained by referring to this picture. The second argument of arc in (3.5) is obtained by subtracting the area of a rectangle with sidelengths $R \sin \theta$ and $R \cos \theta$ from the area θR^2 of an angular sector whose amplitude equals 2θ radians.

We thus conclude that

$$\#\text{Hex}(\mathcal{E}) = \#(I_1) - \#(I_1 \setminus I_2) = \#I - \#(I \setminus I_1) - \#(I_1 \setminus I_2) \geq N - C M \sqrt{N},$$

that is (1.11). Of course, (1.12) follows immediately from (1.11) and (2.8).

Finally, let us assume, by way of contradiction that $k_j^h \geq 6$ for all $(h, j) \neq (0, 1)$. In this case, $k_j^h - 6$ being non-negative for every $(h, j) \neq (0, 1)$, we can go back to (2.4) and apply a version of (1.29) where, in place of a_1 , an arbitrarily large constant L appears. Correspondingly, in place of (2.7) we now deduce an inequality that implies, in particular, $6L \leq M \sqrt{N}$. By taking L large enough in terms of M and N we obtain a contradiction. $\square$

3. THE ARC FUNCTION

Starting from the next section we will make repeated use of the function

$$\text{arc}(\ell, x), \quad \ell \geq 0, x \geq 0,$$

defined as the length of a circular arc subtending a segment of length ℓ and bounding a region of area x . Clearly $\text{arc}(0, x) = 2\sqrt{\pi}\sqrt{x}$ is the isoperimetric profile of $\mathbb{R}^2$. By scaling,

$$\text{arc}(\ell, x) = \ell \text{arc}\left(1, \frac{x}{\ell^2}\right), \quad \forall \ell > 0, x \geq 0, \quad (3.1)$$

so that we can directly focus on $\text{arc}_1 = \text{arc}(1, \cdot)$. We claim that

$$\text{arc}'_1 > 0 \text{ on } (0, \infty) \text{ with } \text{arc}_1(0) = 1, \text{arc}_1(+\infty) = +\infty, \quad (3.2)$$

$$\text{arc}'_1(0) = 0, \text{arc}''_1(0) = 12, \quad (3.3)$$

$$\text{arc}''_1 > 0 \text{ on } [0, \pi/8) \text{ and } \text{arc}''_1 < 0 \text{ on } (\pi/8, \infty). \quad (3.4)$$

We can obtain implicit formulas for arc_1 that can be used in proving (3.2), (3.3), and (3.4). For example, by combining (3.1) with the identity

$$2\theta R = \text{arc}\left(2R \sin \theta, R^2(\theta - \sin \theta \cos \theta)\right), \quad \theta \in [0, \pi/2], R \geq 0, \quad (3.5)$$

see Figure 3, we obtain the following implicit representation³ of arc_1 on the interval $[0, \pi/8]$:

$$\frac{\theta}{\sin \theta} = \text{arc}_1(p(\theta)), \quad \text{for } p(\theta) = \frac{\theta - \sin \theta \cos \theta}{4 \sin^2 \theta}, \quad \theta \in [0, \pi/2]. \quad (3.6)$$

³Notice that p is strictly increasing from $[0, \pi/2]$ to $[0, \pi/8]$.

(A similar construction can be used to represent arc_1 on $[\pi/8, \infty)$.) By combining (3.6) with a Taylor expansion we easily prove (3.3). Similarly, we can use (3.6) to prove (3.2) and (3.4) on $[0, \pi/8]$, and a representation of arc_1 on $[\pi/8, \infty)$ can be used to complete the proof of (3.2) and (3.4). Since this approach (although formally correct) is a bit obscure, we prove (3.2) and (3.4) with the following argument, which seems more transparent. The key remark is that by the variational definition of (mean) curvature (see, e.g., [Mag12, Remark 17.6]), we have

$$\frac{\partial \text{arc}}{\partial x}(\ell, x) = \text{curvature of a circular arc} \quad (3.7)$$

enclosing an area x above a segment of length ℓ .

By (3.7) (with $\ell = 1$) we see that $\text{arc}'_1 > 0$ on $(0, \infty)$, thus deducing (3.2). Moreover, as x increases from 0 to $\pi/8$ (with $x = \pi/8$ corresponding to $\theta = \pi/2$ in Figure 3-(b)), we see from (3.7) that arc'_1 strictly increases from 0 to 2, thus establishing that $\text{arc}''_1 > 0$ on $(0, \pi/8)$; similarly, as x increases from $\pi/8$ to the limit value $+\infty$, we see from (3.7) that arc'_1 strictly decreases from 2 back to the limit value 0, so that $\text{arc}''_1 < 0$ on $(\pi/8, \infty)$.

4. PROOF OF THE QUANTITATIVE HALES' HEXAGONAL INEQUALITY, PART ONE

In this section we begin the proof of Theorem 1.7. More precisely, we prove Theorem 1.7 *conditionally* to the validity of an improvement of Theorem A (Hales' hexagonal isoperimetric inequality). This improvement of Theorem A will be established in Section 5 as Theorem 5.1, and is based on a refinement of the arguments employed in [Hal01]. The argument presented in this section, instead, makes use of a quantitative hexagonal isoperimetric inequality proved in [CM16, Corollary 2.2] (as an elaboration on [IN15, Corollary 1.3]), as stated in Theorem 4.1 below. In the following, given two closed sets C and K in $\mathbb{R}^2$, we denote by

$$\text{hd}(C, K) := \max \left\{ \sup_{x \in C} \text{dist}(x, K), \sup_{x \in K} \text{dist}(x, C) \right\},$$

the Hausdorff distance between C and K .

Theorem 4.1. *There exist positive constants c and η such that if $\Pi \subset \mathbb{R}^2$ is a convex hexagon with $\text{hd}(\Pi, H) \leq \eta$ for some regular hexagon H , then*

$$P(\Pi) - 2(12)^{1/4} \sqrt{|\Pi|} \geq c |\Pi \Delta H^*|^2,$$

for a regular hexagon H^ with $|H^*| = |\Pi|$.*

Proof of Theorem 4.1. By [CM16, Corollary 2.2] (which is stated on the two-dimensional flat torus, but obviously holds on the plane too) there exist positive constants c' and η such that if $\Pi \subset \mathbb{R}^2$ is a convex hexagon with $\text{hd}(\partial\Pi, \partial H) \leq \eta$ for some regular hexagon H , then, for a regular hexagon H^* with $|H^*| = |\Pi|$,

$$P(\Pi) - 2(12)^{1/4} \sqrt{|\Pi|} \geq c' \text{hd}(\partial\Pi, \partial H^*)^2.$$

Since $\text{hd}(\partial\Pi, \partial H) = \text{hd}(\Pi, H)$, we conclude the proof by observing that, for a positive constant C (independent of Π), it holds $|\Pi \Delta H^*| \leq C \text{hd}(\partial\Pi, \partial H^*)$. $\square$

Proof of Theorem 1.7, part one. We prove Theorem 1.7 conditionally to the validity of Theorem 5.1 (that will be proved in Section 5). Theorem 5.1 asserts the following:

There exist positive constants a_1 and a_2 with the following property. If $k \geq 2$, $(\gamma, \mathbf{t}) \in \text{Lip}(\mathbb{S}^1; \mathbb{R}^2) \times \mathcal{P}_k$, and

$$A(\gamma) > 0 \quad \text{if } k = 6, \quad A(\gamma) \geq \frac{1}{100} \quad \text{if } 2 \leq k \leq 5, \quad (4.1)$$

then

$$L(\gamma) + a_1(k-6) + (12)^{1/4} \sigma(\gamma, \mathbf{t}) \geq 2(12)^{1/4} \min\{1, A(\gamma)\} + a_2|k-6|, \quad (4.2)$$

with equality if and only if $(\gamma, \mathbf{t})$ corresponds to a unit-area regular hexagon.

This statement implies Theorem 1.7 when $k \neq 6$. Therefore, for the remaining part of the proof, we shall assume that $k = 6$. We thus want to prove the existence of a positive constant a_3 such that if $(\gamma, \mathbf{t}) \in \text{Lip}(\mathbb{S}^1, \mathbb{R}^2) \times \mathcal{P}_6$, γ is injective, and

$$1 \geq A(\gamma) \geq \frac{1}{100}, \quad (4.3)$$

then

$$L(\gamma) + (12)^{1/4} \sigma(\gamma, \mathbf{t}) \geq 2(12)^{1/4} A(\gamma) + a_3 \left\{ d_{\text{hex}}(E_\gamma)^2 + (1 - A(\gamma)) \right\}, \quad (4.4)$$

where E_γ denotes the bounded connected component of $\mathbb{R}^2 \setminus \gamma(\mathbb{S}^1)$ identified by γ (thanks to Jordan's theorem).

Step one: In this step we set, for $(\gamma, \mathbf{t}) \in \text{Lip}(\mathbb{S}^1; \mathbb{R}^2) \times \mathcal{P}_6$,

$$\delta(\gamma, \mathbf{t}) = L(\gamma) + (12)^{1/4} \sigma(\gamma, \mathbf{t}) - 2(12)^{1/4} A(\gamma), \quad (4.5)$$

and prove that for every $\varepsilon > 0$ there is $\delta > 0$ such that, if $\delta(\gamma, \mathbf{t}) \leq \delta$, then there is a unit-area, regular hexagon H_0 with

$$\max \left\{ \text{hd}(E_\gamma, H_0), \text{hd}(\Pi_{\gamma, \mathbf{t}}, H_0), |E_\gamma \Delta \Pi_{\gamma, \mathbf{t}}| \right\} \leq \varepsilon, \quad (4.6)$$

where $\Pi_{\gamma, \mathbf{t}}$ is the convex envelope of $\{\gamma(\mathbf{t}(i))\}_{i=1}^6$. Notice, in particular, that if ε is small enough, then (4.6) implies that $\Pi_{\gamma, \mathbf{t}}$ is a convex hexagon.

We prove this arguing by contradiction, aiming at using our assumption that, under (4.1), (4.2) holds, and thus $\delta(\gamma, \mathbf{t}) \geq 0$ with equality if and only if $k = 6$ and $(\gamma, \mathbf{t})$ corresponds to a unit-area, regular hexagon.

We thus consider $\varepsilon_* > 0$ and a sequence $\{(\gamma_j, \mathbf{t}_j)\}_j$ in $\text{Lip}(\mathbb{S}^1; \mathbb{R}^2) \times \mathcal{P}_6$ with $1 \geq A(\gamma_j) \geq 1/100$, $\delta(\gamma_j, \mathbf{t}_j) \rightarrow 0^+$ as $j \rightarrow \infty$, and

$$\max \left\{ \inf_H \left\{ \max\{\text{hd}(E_{\gamma_j}, H), \text{hd}(\Pi_{\gamma_j, \mathbf{t}_j}, H)\} \right\}, |E_{\gamma_j} \Delta \Pi_{\gamma_j, \mathbf{t}_j}| \right\} \geq \varepsilon_*, \quad \forall j. \quad (4.7)$$

(Here H denotes a generic unit-area, regular hexagon.) Up to a reparametrization we can assume that

$$|\gamma'_j| = \frac{L(\gamma_j)}{2\pi} \quad \text{on } \mathbb{S}^1. \quad (4.8)$$

By $|\sigma(\gamma_j, \mathbf{t}_j)| \leq 1/2$, $A(\gamma_j) \leq 1$, and $\delta(\gamma_j, \mathbf{t}_j) \rightarrow 0$ we find that $\ell = \sup_j L(\gamma_j) < \infty$. By (4.8), up to extracting subsequences and up to translations, there is $\gamma \in \text{Lip}(\mathbb{S}^1; \mathbb{R}^2)$ such that $\gamma_j \rightarrow \gamma$ uniformly on $\mathbb{S}^1$ and $\gamma'_j \xrightarrow{*} \gamma'$ in $L^\infty(\mathbb{S}^1; \mathbb{R}^2)$: in particular,

$$A(\gamma_j) = \int_{\mathbb{S}^1} \gamma_j^{(1)}(\gamma_j')^{(2)} \rightarrow \int_{\mathbb{S}^1} \gamma^{(1)}(\gamma')^{(2)} = A(\gamma) \quad (4.9)$$

as $j \rightarrow \infty$ (here we are denoting by $x^{(i)}$ the components of $x \in \mathbb{R}^2$), and from the corresponding bounds on $A(\gamma_j)$ we find that

$$1 \geq A(\gamma) \geq \frac{1}{100}.$$

Now, up to extracting subsequences, there is $\{s_i : 1 \leq i \leq 6\} \subset \mathbb{S}^1$ such that, as $j \rightarrow \infty$, $\mathbf{t}_j(i) \rightarrow s_i$ for each $i = 1, \dots, 6$. In particular, for each $i = 1, \dots, 6$, and setting $s_7 = s_1$, we

have $s_i \leq s_{i+1}$ in the ordering of $\mathbb{S}^1$. To compute the limit of $\sigma(\gamma_j, \mathbf{t}_j)$ we start noticing that

$$\alpha(\gamma_j, \mathbf{t}_j(i), \mathbf{t}_j(i+1)) = \int_{[\mathbf{t}_j(i), \mathbf{t}_j(i+1)]} \gamma_j^{(1)} (\gamma_j')^{(2)} + A(\llbracket \gamma_j(\mathbf{t}_j(i+1)), \gamma_j(\mathbf{t}_j(i)) \rrbracket) . \quad (4.10)$$

Starting from (4.10) and thanks to the fact that $\gamma_j \rightarrow \gamma$ uniformly on $\mathbb{S}^1$ and $\gamma_j' \xrightarrow{*} \gamma'$ in $L^\infty(\mathbb{S}^1; \mathbb{R}^2)$, we see that if $s_i < s_{i+1}$, then

$$\lim_{j \rightarrow \infty} \alpha(\gamma_j, \mathbf{t}_j(i), \mathbf{t}_j(i+1)) = \alpha(\gamma, s_i, s_{i+1}) ;$$

while if $s_i = s_{i+1}$ but there is at least one $\ell \neq i$ such that $s_\ell \neq s_{\ell+1}$, then

$$\lim_{j \rightarrow \infty} \alpha(\gamma_j, \mathbf{t}_j(i), \mathbf{t}_j(i+1)) = 0 ; \quad (4.11)$$

and, finally, if $s_i = s_{i+1}$ for all i , then there is i_0 such that (4.11) holds for all $i \neq i_0$, and

$$\lim_{j \rightarrow \infty} \alpha(\gamma_j, \mathbf{t}_j(i_0), \mathbf{t}_j(i_0+1)) = A(\gamma) .$$

By using these three facts we conclude that, either $s_i = s_{i+1}$ for all i and

$$\lim_{j \rightarrow \infty} \sigma(\gamma_j, \mathbf{t}_j) = \min \left\{ \frac{1}{2}, A(\gamma) \right\} , \quad (4.12)$$

or there exists $2 \leq k \leq 6$ and $\mathbf{t} \in \mathcal{P}_k$ such that

$$\lim_{j \rightarrow \infty} \sigma(\gamma_j, \mathbf{t}_j) = \sigma(\gamma, \mathbf{t}) . \quad (4.13)$$

If we are in the case when (4.12) holds, then we deduce from the isoperimetric inequality, $\delta(\gamma_j, \mathbf{t}_j) \geq 0$, (4.9), and (4.12) that

$$\begin{aligned} 2\sqrt{\pi} \sqrt{A(\gamma)} &\leq L(\gamma) \leq \liminf_{j \rightarrow \infty} L(\gamma_j) \leq \lim_{j \rightarrow \infty} 2(12)^{1/4} A(\gamma_j) - (12)^{1/4} \sigma(\gamma_j, \mathbf{t}_j) \\ &= 2(12)^{1/4} A(\gamma) - (12)^{1/4} \min \left\{ \frac{1}{2}, A(\gamma) \right\} . \end{aligned} \quad (4.14)$$

Now, the function

$$F(x) = 2\sqrt{\pi}x - 2(12)^{1/4}x + (12)^{1/4} \min\{x, 1/2\} ,$$

is concave on $[0, 1]$, with $F(0) = 0$ and $F(1) = 2\sqrt{\pi} - (12)^{1/4} > 0$, so that (4.14) implies $A(\gamma) = 0$ (a contradiction, as $A(\gamma) \geq 1/100$). Therefore (4.12) never occurs, and by combining (4.9) with (4.13), $\delta(\gamma_j, \mathbf{t}_j) \geq 0$, and $\delta(\gamma_j, \mathbf{t}_j) \rightarrow 0$ as $j \rightarrow \infty$ we conclude that $\delta(\gamma, \mathbf{t}) = 0$ for $\mathbf{t}$ as in (4.13). By our assumption, $\delta(\gamma, \mathbf{t}) = 0$ implies that $k = 6$, γ is a monotone parametrization of the boundary of a regular, unit-area hexagon, $\{\gamma(\mathbf{t}(i))\}_{i=1}^6$ are the vertexes of such hexagon, and $\mathbf{t}_j(i) \rightarrow \mathbf{t}(i)$ for $1 \leq i \leq 6$ as $j \rightarrow \infty$. As $\gamma_j \rightarrow \gamma$ uniformly on $\mathbb{S}^1$ we conclude that

$$\max \left\{ \inf_H \left\{ \max\{\text{hd}(E_{\gamma_j}, H), \text{hd}(\Pi_{\gamma_j, \mathbf{t}_j}, H)\} \right\}, |E_{\gamma_j} \Delta \Pi_{\gamma_j, \mathbf{t}_j}| \right\} \rightarrow 0 ,$$

as $j \rightarrow \infty$, in contradiction with (4.7).

Step two: We conclude the proof of Theorem 1.7. Let $(\gamma, \mathbf{t}) \in \text{Lip}(\mathbb{S}^1, \mathbb{R}^2) \times \mathcal{P}_6$ be such that γ is injective and $1 \geq A(\gamma) \geq 1/100$, and let E_γ be the connected component of $\mathbb{R}^2 \setminus \gamma(\mathbb{S}^1)$ identified by Jordan's theorem (so that $|E_\gamma| = A(\gamma)$). We want to prove the existence of $a_3 > 0$ such that

$$\delta(\gamma, \mathbf{t}) \geq a_3 \left\{ d_{\text{hex}}(E_\gamma)^2 + (1 - A(\gamma)) \right\} . \quad (4.15)$$

Since $d_{\text{hex}}(E_\gamma) < 2|E_\gamma| \leq 2$, if $\delta(\gamma, \mathbf{t}) \geq \delta$ for some constant $\delta > 0$, then (4.15) holds for every $a_3 \leq \delta/5$. For $\varepsilon > 0$ to be chosen later, we select $\delta > 0$ depending on ε as determined

in step one, and reduce to the situation when there is a unit-area, regular hexagon H_0 such that

$$\max \left\{ \text{hd}(E_\gamma, H_0), \text{hd}(\Pi_{\gamma, \mathbf{t}}, H_0), |E_\gamma \Delta \Pi_{\gamma, \mathbf{t}}| \right\} < \varepsilon, \quad \Pi_{\gamma, \mathbf{t}} \text{ is a convex hexagon.} \quad (4.16)$$

If we denote by x_i the total area enclosed between the i -th (curvilinear) edge of E_γ and the corresponding i -th edge of $\Pi_{\gamma, \mathbf{t}}$, and by ℓ_i the length $|\gamma(\mathbf{t}(i+1)) - \gamma(\mathbf{t}(i))|$ of the i -th edge of $\Pi_{\gamma, \mathbf{t}}$, then by (4.16) we have that, for some positive constant C ,

$$\sum_{i=1}^6 x_i + \max_{1 \leq i \leq 6} \left| \ell_i - \frac{(12)^{1/4}}{3} \right| < C \varepsilon, \quad (4.17)$$

where of course $(12)^{1/4}/3$ is the length of one edge of a unit-area, regular hexagon. Since $(12)^{1/4}/3 < 1$, provided ε is small enough and by (4.17), we deduce that $\max_i \ell_i < 1$. In particular, using the scaling property (3.1) of arc_1 , and the fact that arc_1 is increasing on $[0, \infty)$ (recall (3.2)), we find that

$$L(\gamma) \geq \sum_{i=1}^6 \text{arc}(\ell_i, x_i) = \sum_{i=1}^6 \text{arc}_1\left(\frac{x_i}{\ell_i^2}\right) \geq \sum_{i=1}^6 \ell_i \text{arc}_1\left(\frac{x_i}{\ell_i}\right).$$

By (4.17), up to further decreasing the value of ε , we can ensure that $\max_i x_i/\ell_i \leq \pi/8$, and then combine the convexity of arc_1 on $[0, \pi/8]$ (recall (3.4)) with Jensen's inequality to conclude that

$$L(\gamma) \geq \sum_{i=1}^6 \ell_i \text{arc}_1\left(\frac{x_i}{\ell_i}\right) \geq \left(\sum_{i=1}^6 \ell_i\right) \text{arc}_1\left(\frac{\sum_{i=1}^6 x_i}{\sum_{i=1}^6 \ell_i}\right) = P(\Pi_{\gamma, \mathbf{t}}) \text{arc}_1\left(\frac{|E_\gamma \Delta \Pi_{\gamma, \mathbf{t}}|}{P(\Pi_{\gamma, \mathbf{t}})}\right). \quad (4.18)$$

(Inequality (4.18) is strictly related to the *chordal isoperimetric inequality* found in [Hal01, Proposition 6.1-A] and [Mor09, 15.5], although it does not seem to exactly fit in those statements.) Now, by $\text{arc}_1(0) = 1$, $\text{arc}'_1(0) = 0$, and $\text{arc}''_1(0) = 12$ (recall (3.2) and (3.3)), we can find a positive constant C such that

$$\text{arc}_1(x) \geq 1 + 6x^2 - C|x|^3, \quad \forall x \in [0, 1/2]. \quad (4.19)$$

Since we have $|E_\gamma \Delta \Pi_{\gamma, \mathbf{t}}| \rightarrow 0$ and $P(\Pi_{\gamma, \mathbf{t}}) \rightarrow P(H_0) = 2(12)^{1/4}$ in the limit $\varepsilon \rightarrow 0^+$, by further decreasing the value of ε , thanks to (4.17), we can ensure that $|E_\gamma \Delta \Pi_{\gamma, \mathbf{t}}| < P(\Pi_{\gamma, \mathbf{t}})/2$, and thus deduce from (4.18), (4.19), and (4.16) that

$$L(\gamma) \geq P(\Pi_{\gamma, \mathbf{t}}) + \left(\frac{6}{2(12)^{1/4}} - C\varepsilon\right) |E_\gamma \Delta \Pi_{\gamma, \mathbf{t}}|^2. \quad (4.20)$$

For $\eta > 0$ as in Theorem 4.1, up to further decreasing ε so to entail that $\text{hd}(\Pi_{\gamma, \mathbf{t}}, H_0) < \eta$, and thanks to the convexity of $\Pi_{\gamma, \mathbf{t}}$, we can combine Theorem 4.1 with (4.20) to find that, for some positive constant C ,

$$L(\gamma) \geq 2(12)^{1/4} \sqrt{|\Pi_{\gamma, \mathbf{t}}|} + \frac{|\Pi_{\gamma, \mathbf{t}} \Delta H_*|^2}{C} + \left(\frac{3}{(12)^{1/4}} - C\varepsilon\right) |E_\gamma \Delta \Pi_{\gamma, \mathbf{t}}|^2, \quad (4.21)$$

where H_* is a regular hexagon with $|H_*| = |\Pi_{\gamma, \mathbf{t}}|$. Now, let us recall that, by (4.3), $A(\gamma) > 0$, so that, in particular, the orientation of γ is such that $A(\gamma) = |E_\gamma|$. Denoting by $\circ$ the operation of concatenating curves, we see that

$$\pi_{\gamma, \mathbf{t}} = [\gamma(\mathbf{t}(1)), \gamma(\mathbf{t}(2))] \circ [\gamma(\mathbf{t}(2)), \gamma(\mathbf{t}(3))] \circ \cdots \circ [\gamma(\mathbf{t}(6)), \gamma(\mathbf{t}(1))],$$

defines a Lipschitz map $\pi_{\gamma, \mathbf{t}} \in \text{Lip}(\mathbb{S}^1; \mathbb{R}^2)$ that, thanks to (4.16), maps injectively $\mathbb{S}^1$ into $\partial \Pi_{\gamma, \mathbf{t}}$, and in such a way that $A(\pi_{\gamma, \mathbf{t}}) = |\Pi_{\gamma, \mathbf{t}}|$. By definition of A , $\pi_{\gamma, \mathbf{t}}$, and of the secant

oriented area functional α (recall (1.24)), we thus find that

$$\begin{aligned} |E_\gamma| - |\Pi_{\gamma, \mathbf{t}}| &= \sum_{i=1}^k A(\gamma|_{[\mathbf{t}(i), \mathbf{t}(i+1)]}) - \sum_{i=1}^k A([\gamma(\mathbf{t}(i)), \gamma(\mathbf{t}(i+1))]) \\ &= \sum_{k=1}^6 \alpha(\gamma, \mathbf{t}(i), \mathbf{t}(i+1)), \end{aligned}$$

so that, up to decreasing the value of ε , so to enforce from (4.16) that $|\alpha(\gamma, \mathbf{t}(i), \mathbf{t}(i+1))| \leq |E_\gamma \Delta \Pi_{\gamma, \mathbf{t}}| < 1/2$, we conclude that

$$|\Pi_{\gamma, \mathbf{t}}| = |E_\gamma| - \sigma(\gamma, \mathbf{t}), \quad (4.22)$$

Aiming at providing an efficient lower bound for the first term on the right-hand side of (4.21), we use (4.22) to find that

$$\begin{aligned} \sqrt{|\Pi_{\gamma, \mathbf{t}}|} &= \sqrt{|E_\gamma|} \sqrt{\frac{|\Pi_{\gamma, \mathbf{t}}|}{|E_\gamma|}} = |E_\gamma| \sqrt{1 - \frac{\sigma(\gamma, \mathbf{t})}{|E_\gamma|}} + \left(\sqrt{|E_\gamma|} - |E_\gamma| \right) \sqrt{\frac{|\Pi_{\gamma, \mathbf{t}}|}{|E_\gamma|}} \\ &= |E_\gamma| \sqrt{1 - \frac{\sigma(\gamma, \mathbf{t})}{|E_\gamma|}} + \frac{1 - |E_\gamma|}{1 + \sqrt{|E_\gamma|}} \sqrt{|\Pi_{\gamma, \mathbf{t}}|} \end{aligned} \quad (4.23)$$

$$\geq |E_\gamma| \sqrt{1 - \frac{\sigma(\gamma, \mathbf{t})}{|E_\gamma|}} + \frac{1 - |E_\gamma|}{C} \quad (4.24)$$

where we have decreased ε to ensure $|\Pi_{\gamma, \mathbf{t}}| \geq 1/2$, and where C is a positive constant. By means of the Taylor expansion we find that

$$\sqrt{1-s} = 1 - \frac{s}{2} - \frac{s^2}{8} + O(s^3), \quad \text{as } s \rightarrow 0^+, \quad (4.25)$$

and use (4.25) with (4.24) to bound from below the first and the third term on the right hand side of (4.21) as

$$\begin{aligned} &2(12)^{1/4} \sqrt{|\Pi_{\gamma, \mathbf{t}}|} + \left(\frac{3}{(12)^{1/4}} - C\varepsilon \right) |E_\gamma \Delta \Pi_{\gamma, \mathbf{t}}|^2 \\ &\geq 2(12)^{1/4} |E_\gamma| - (12)^{1/4} \sigma(\gamma, \mathbf{t}) + \frac{1 - |E_\gamma|}{C} \\ &\quad - 2(12)^{1/4} \left(\frac{1}{8} - C\varepsilon \right) \sigma(\gamma, \mathbf{t})^2 + \left(\frac{3}{(12)^{1/4}} - C\varepsilon \right) |E_\gamma \Delta \Pi_{\gamma, \mathbf{t}}|^2. \end{aligned} \quad (4.26)$$

Therefore, up to further decreasing the value of ε , we find that

$$|\sigma(\gamma, \mathbf{t})| \leq |E_\gamma \Delta \Pi_{\gamma, \mathbf{t}}|, \quad \text{and} \quad \frac{3}{(12)^{1/4}} > \frac{2(12)^{1/4}}{8},$$

we finally conclude from (4.21) and (4.26) that

$$L(\gamma) \geq 2(12)^{1/4} |E_\gamma| + \frac{|\Pi_{\gamma, \mathbf{t}} \Delta H_*|^2}{C} - (12)^{1/4} \sigma(\gamma, \mathbf{t}) + \frac{|E_\gamma \Delta \Pi_{\gamma, \mathbf{t}}|^2 + (1 - |E_\gamma|)}{C},$$

that is

$$C \delta(\gamma, \mathbf{t}) \geq |\Pi_{\gamma, \mathbf{t}} \Delta H_*|^2 + |E_\gamma \Delta \Pi_{\gamma, \mathbf{t}}|^2 + (1 - |E_\gamma|). \quad (4.27)$$

Since $d_{\text{hex}}(E_\gamma) \leq C(|\Pi_{\gamma, \mathbf{t}} \Delta H_*| + |E_\gamma \Delta \Pi_{\gamma, \mathbf{t}}|)$, we easily see that (4.27) implies (4.4). $\square$

5. PROOF OF THE QUANTITATIVE HALES' HEXAGONAL INEQUALITY, PART TWO

Thanks to the argument presented in Section 4, in order to complete the proof of Theorem 1.7, and thus of Theorem 1.2, we are left to prove the following theorem.

Theorem 5.1. *There exist positive constants a_1 and a_2 with the following property. If $k \geq 2$, $(\gamma, \mathbf{t}) \in \text{Lip}(\mathbb{S}^1; \mathbb{R}^2) \times \mathcal{P}_k$, and*

$$A(\gamma) > 0 \quad \text{if } k = 6, \quad A(\gamma) \geq \frac{1}{100} \quad \text{if } 2 \leq k \leq 5, \quad (5.1)$$

then

$$L(\gamma) + a_1(k-6) + (12)^{1/4} \sigma(\gamma, \mathbf{t}) \geq 2(12)^{1/4} \min\{1, A(\gamma)\} + a_2|k-6|, \quad (5.2)$$

with equality if and only if $(\gamma, \mathbf{t})$ corresponds to a unit-area regular hexagon.

Theorem 5.1 improves on Hales' hexagonal isoperimetric inequality (Theorem A) since it requires no lower bound on $A(\gamma)$ when $k \geq 6$ and weaker ones when $2 \leq k \leq 5$, and since, when $k \neq 7$, it provides the additional lower bound $a_2|k-6|$. These improvements are obtained by carefully revisiting Hales' original argument, which is why the methods of this section overlap with those of [Hal01].

Proof of Theorem 5.1. We make the following claim:

Claim: There are positive constants a and c such that, if $k \geq 2$, $k \neq 6$, and $(\gamma, \mathbf{t}) \in \text{Lip}(\mathbb{S}^1; \mathbb{R}^2) \times \mathcal{P}_k$ satisfies (5.1), then

$$L(\gamma) + a(k-6) + (12)^{1/4} \sigma(\gamma, \mathbf{t}) \geq 2(12)^{1/4} \min\{1, A(\gamma)\} + c; \quad (5.3)$$

and, if $(\gamma, \mathbf{t}) \in \text{Lip}(\mathbb{S}^1; \mathbb{R}^2) \times \mathcal{P}_6$ with $1/5 \geq A(\gamma) > 0$, then

$$L(\gamma) + (12)^{1/4} \sigma(\gamma, \mathbf{t}) > 2(12)^{1/4} \min\{1, A(\gamma)\}. \quad (5.4)$$

The claim implies the theorem: Indeed, when $k = 6$, Theorem A implies that

$$L(\gamma) + (12)^{1/4} \sigma(\gamma, \mathbf{t}) \geq 2(12)^{1/4} \min\{1, A(\gamma)\}$$

whenever $(\gamma, \mathbf{t}) \in \text{Lip}(\mathbb{S}^1; \mathbb{R}^2) \times \mathcal{P}_6$ and $A(\gamma) \geq 2\pi/(\sqrt{3}(6^2))$, and with equality if and only if $(\gamma, \mathbf{t})$ corresponds to a regular unit-area hexagon. Since $2\pi/(\sqrt{3}(6^2)) < 1/5$, the combination of (5.4) with Theorem A proves the case $k = 6$ of Theorem 5.1. When $k \geq 2$, $k \neq 6$, we set

$$a_1 = a_2 + a, \quad a_2 = \frac{c}{8},$$

and deduce (5.2) from (5.3) as follows: (5.2) with $k > 6$ is equivalent to

$$L(\gamma) + (a_1 - a_2)(k-6) + (12)^{1/4} \sigma(\gamma, \mathbf{t}) \geq 2(12)^{1/4} \min\{1, A(\gamma)\},$$

which is implied by (5.3) thanks to $a_1 - a_2 = a$; if, instead, $2 \leq k \leq 5$, then by (5.3) we get

$$\begin{aligned} L(\gamma) + (12)^{1/4} \sigma(\gamma, \mathbf{t}) &\geq 2(12)^{1/4} \min\{1, A(\gamma)\} + c - a(k-6) \\ &= 2(12)^{1/4} \min\{1, A(\gamma)\} + 8a_2 + (a_2 - a_1)(k-6), \\ &\geq 2(12)^{1/4} \min\{1, A(\gamma)\} - (a_1 + a_2)(k-6), \end{aligned}$$

that is (5.2).

We can thus focus on the proof of the above claim. It will be convenient to set some notation. For every $(\gamma, \mathbf{t}) \in \text{Lip}(\mathbb{S}^1; \mathbb{R}^2) \times \mathcal{P}_k$ we set for brevity

$$\varepsilon(\gamma, \mathbf{t}) = a(k-6) + (12)^{1/4} \sigma(\gamma, \mathbf{t}) - 2(12)^{1/4} \min\{1, A(\gamma)\},$$

where $a > 0$ is a constant to be determined. Moreover, dropping the dependency on $\mathbf{t}$ in the interest of brevity, we set, for each $i = 1, \dots, k+1$,

$$\gamma_i = \gamma|_{[\mathbf{t}(i), \mathbf{t}(i+1)]}, \quad (5.5)$$

$$A_i = A(\gamma_i) + A(\llbracket \gamma(\mathbf{t}(i+1)), \gamma(\mathbf{t}(i)) \rrbracket), \quad (5.6)$$

so that

$$L(\gamma) = \sum_{i=1}^k L(\gamma_i), \quad \sigma(\gamma, \mathbf{t}) = \sum_{i=1}^k \max \left\{ -\frac{1}{2}, \min \left\{ A_i, \frac{1}{2} \right\} \right\}.$$

We now present a series of reduction steps.

Step one: For each $(\gamma, \mathbf{t}) \in \text{Lip}(\mathbb{S}^1; \mathbb{R}^2) \times \mathcal{P}_k$, there is $\bar{\gamma} \in \text{Lip}(\mathbb{S}^1; \mathbb{R}^2)$ such that, for each $i = 1, \dots, k$, the restriction of $\bar{\gamma}$ to the interval $[\mathbf{t}(i), \mathbf{t}(i+1)]$ is the unit speed parametrization of a circular arc with $\bar{\gamma}(\mathbf{t}(i)) = \gamma(\mathbf{t}(i))$, and moreover

$$L(\gamma) + \varepsilon(\gamma, \mathbf{t}) \geq L(\bar{\gamma}) + \varepsilon(\bar{\gamma}, \mathbf{t}), \quad (5.7)$$

for every choice of $a > 0$. (Notice carefully that neither γ or $\bar{\gamma}$ are assumed or required to be injective.)

To prove this, it is enough to define $\bar{\gamma}$ on $[\mathbf{t}(i), \mathbf{t}(i+1)]$, $1 \leq i \leq k$, as the unit speed parametrization of a circular arc with endpoints $\bar{\gamma}(\mathbf{t}(i)) = \gamma(\mathbf{t}(i))$ and $\bar{\gamma}(\mathbf{t}(i+1)) = \gamma(\mathbf{t}(i+1))$ and such that

$$A(\bar{\gamma}_i) = A(\gamma_i),$$

(where $\bar{\gamma}_i$ is the restriction of $\bar{\gamma}$ to $[\mathbf{t}(i), \mathbf{t}(i+1)]$, see (5.5)). Then, by construction, $A(\bar{\gamma}) = A(\gamma)$ and $\sigma(\bar{\gamma}, \mathbf{t}) = \sigma(\gamma, \mathbf{t})$, while, by Dido's inequality, $L(\bar{\gamma}) \leq L(\gamma)$, so that (5.7) holds.

Having proved step one, for the rest of the proof we will be able to work directly with pairs

$$(\gamma, \mathbf{t}) \in \text{Cir}_k,$$

as defined by the condition that, for each $i = 1, \dots, k$, the restriction of γ to the interval $[\mathbf{t}(i), \mathbf{t}(i+1)]$ is the unit speed parametrization of a circular arc.

Step two: We prove that, for each $(\gamma, \mathbf{t}) \in \text{Cir}_k$, there is $\bar{\gamma}$ such that $(\bar{\gamma}, \mathbf{t}) \in \text{Cir}_k$,

$$\bar{A}_i \geq -\frac{1}{2}, \quad \forall i = 1, \dots, k,$$

and (5.7) holds (for any choice of $a > 0$). Here $\bar{A}_i$ is defined from $\bar{\gamma}$ as A_i was defined from γ in (5.6).

To prove this, we set $\bar{\gamma} = \gamma$ on those intervals $[\mathbf{t}(i), \mathbf{t}(i+1)]$ corresponding to $i = 1, \dots, k$ such that $A_i \geq -1/2$; if, instead, $A_i < -1/2$, then we define $\bar{\gamma}$ on $[\mathbf{t}(i), \mathbf{t}(i+1)]$ as a unit-speed parametrization of a circular arc with endpoints $\gamma(\mathbf{t}(i))$ and $\gamma(\mathbf{t}(i+1))$ such that $\bar{A}_i = -1/2$: since $\bar{\gamma}$ and γ have the same endpoints and $A_i < -1/2$, we see that this implies

$$A(\bar{\gamma}_i) > A(\gamma_i), \quad L(\bar{\gamma}_i) \leq L(\gamma_i).$$

In particular, $L(\gamma) \geq L(\bar{\gamma})$, $A(\bar{\gamma}) \geq A(\gamma)$, and $\sigma(\bar{\gamma}, \mathbf{t}) = \sigma(\gamma, \mathbf{t})$, which implies the validity of (5.7).

Thanks to step one and step two we have reduced to the case when

$$(\gamma, \mathbf{t}) \in \text{Cir}_k, \quad A_i \geq -\frac{1}{2}, \quad \forall i = 1, \dots, k. \quad (5.8)$$

Step three: We prove that given $(\gamma, \mathbf{t})$ as in (5.8) there is $(\bar{\gamma}, \mathbf{t}) \in \text{Cir}_k$ such that (5.7) holds (for any value of a), and such that all the $\bar{A}_i$'s in $[-1/2, 1/2]$ have the same sign, that is, setting

$$\bar{I} = \left\{ i = 1, \dots, k : |\bar{A}_i| \leq \frac{1}{2} \right\},$$

we have that either $\bar{A}_i \geq 0$ for all $i \in \bar{I}$ or $\bar{A}_i \leq 0$ for all $i \in \bar{I}$.

To prove this, let us consider the set K of those pairs (i, j) with $1 \leq i, j \leq k$, $i \neq j$, and

$$\frac{1}{2} \geq A_i > 0 > A_j \geq -\frac{1}{2},$$

which we can assume to be non-empty, otherwise, we have nothing to prove. Let $(i, j) \in K$ and let $\beta = A_i + A_j$. Let us assume for a moment that $\beta \geq 0$. In this case, we define γ^* by taking $\gamma^* = \gamma$ everywhere on $\mathbb{S}^1$ except on: $[\mathbf{t}(j), \mathbf{t}(j+1)]$, on which we define γ^* as the unit speed parametrization of the segment from $\gamma(\mathbf{t}(j))$ to $\gamma(\mathbf{t}(j+1))$ (so that $A_j^* = 0$ provided A_j^* is defined from γ^* in analogy to (5.6)); and $[\mathbf{t}(i), \mathbf{t}(i+1)]$, on which we define γ^* as the unit speed parametrization of a circular arc from $\gamma(\mathbf{t}(i))$ to $\gamma(\mathbf{t}(i+1))$ such that $A_i^* = \beta \geq 0$. In this way $A(\gamma^*) = A(\gamma)$ and $\sigma(\gamma^*, \mathbf{t}) = \sigma(\gamma, \mathbf{t})$ (recall the restriction $1/2 \geq |A_i|, |A_j|$). Moreover, $L(\gamma_j^*) \leq L(\gamma_j)$ (as on $[\mathbf{t}(j), \mathbf{t}(j+1)]$ we have replaced a circular arc with its chord segment), and $L(\gamma_i^*) \leq L(\gamma_i)$ (since $A_i^* = \beta \leq A_i$, on $[\mathbf{t}(i), \mathbf{t}(i+1)]$ we have modified the curvature of a circular arc with fixed endpoints so to decrease the amount of enclosed area), so that, in total, $L(\gamma^*) \leq L(\gamma)$.

In summary, when $\beta \geq 0$ we have constructed a curve γ^* such that $L(\gamma^*) + \varepsilon(\gamma^*, \mathbf{t}) \leq L(\gamma) + \varepsilon(\gamma, \mathbf{t})$, $A_i^* = \beta$, $A_j^* = 0$, and $\gamma^* = \gamma$ on $[\mathbf{t}(h), \mathbf{t}(h+1)]$ for all $h \neq i, j$. In the case $\beta < 0$, an analogous construction produces γ^* such that $L(\gamma^*) + \varepsilon(\gamma^*, \mathbf{t}) \leq L(\gamma) + \varepsilon(\gamma, \mathbf{t})$, $A_i^* = 0$, $A_j^* = \beta$, and $\gamma^* = \gamma$ on $[\mathbf{t}(h), \mathbf{t}(h+1)]$ for all $h \neq i, j$. By iterating this construction finitely many times we end up constructing a curve $\bar{\gamma}$ satisfying (5.7) and such that, with $I = \{i : |\bar{A}_i| \leq 1/2\}$, we have that either $\bar{A}_i \geq 0$ or $\bar{A}_i \leq 0$ for all $i \in I$.

Therefore, we have so far reduced to prove that (5.3) holds when $k \geq 2$, $k \neq 6$,

$$(\gamma, \mathbf{t}) \in \text{Cir}_k, \quad A_i \geq -\frac{1}{2}, \quad \forall i = 1, \dots, k, \quad (5.9)$$

and, setting $I = \{i : |A_i| \leq 1/2\}$, we have that

$$\text{either } A_i \geq 0 \text{ for all } i \in I, \text{ or } A_i \leq 0 \text{ for all } i \in I. \quad (5.10)$$

Step four: In this step we impose the first restriction on the constant $a > 0$ appearing in the definition of $\varepsilon(\gamma, \mathbf{t})$, that is, we impose

$$a < \frac{\sqrt{\pi}}{2} - \frac{3}{8}(12)^{1/4}. \quad (5.11)$$

With this restriction on a , we prove that if $(\gamma, \mathbf{t}) \in \text{Cir}_k$ satisfies (5.9), (5.10), and is such that $A_i > 1/2$ for some $1 \leq i \leq k$, then

$$L(\gamma) + \varepsilon(\gamma, \mathbf{t}) \geq \min \left\{ \frac{7}{10}, c(a) \right\}, \quad (5.12)$$

where

$$c(a) = 2\sqrt{\pi} - (3/2)(12)^{1/4} - 4a, \quad (5.13)$$

is strictly positive thanks to the strict sign in (5.11). We notice that this step, combined with the previous ones, effectively reduces the proof of (5.3) (thus of the theorem) to the case when $k \geq 2$, $k \neq 6$, $(\gamma, \mathbf{t}) \in \text{Cir}_k$ and

$$\text{either } \frac{1}{2} \geq A_i \geq 0 \quad \forall i, \quad \text{or} \quad 0 \geq A_i \geq -\frac{1}{2} \quad \forall i. \quad (5.14)$$

We now turn to the proof of (5.12):

Let $(\gamma, \mathbf{t}) \in \text{Cir}_k$ satisfy (5.8), (5.10) and be such that $A_{i_0} > 1/2$ for some i_0 . Set

$$I = \left\{ i : A_i > \frac{1}{2} \right\}, \quad J = \left\{ i : |A_i| \leq \frac{1}{2} \right\},$$

so that $\{1, \dots, k\} = I \cup J$, $I \cap J = \emptyset$, $I \neq \emptyset$, and, setting

$$J^+ = \{i \in J : A_i \geq 0\}, \quad J^- = \{i \in J : A_i \leq 0\},$$

and recalling (5.10), either $J = J^+$ or $J = J^-$. We address the two cases by different arguments.

Case one, $J = J^+$: In this case by $I \neq \emptyset$ we have

$$\sigma(\gamma, \mathbf{t}) = \frac{\#I}{2} + \sum_{i \in J} A_i \geq \frac{1}{2}.$$

Hence, by the trivial bound $a(k-6) \geq -4a$ (recall, $k \geq 2$), and by the isoperimetric inequality $L(\gamma) \geq 2\sqrt{\pi} \sqrt{A(\gamma)}$ (recall that $A(\gamma) \geq 0$ by (5.1)), we find

$$\begin{aligned} L(\gamma) + \varepsilon(\gamma, \mathbf{t}) &\geq 2\sqrt{\pi} \sqrt{A(\gamma)} - 4a + \frac{(12)^{1/4}}{2} - 2(12)^{1/4} \min\{1, A(\gamma)\} \\ &\geq 2\sqrt{\pi} - 4a + \frac{(12)^{1/4}}{2} - 2(12)^{1/4} = \left(2\sqrt{\pi} - \frac{3}{2}(12)^{1/4}\right) - 4a, \end{aligned}$$

where in the last step we have used the fact that $x \geq 0 \mapsto 2\sqrt{\pi} \sqrt{x} - 2(12)^{1/4} \min\{1, x\}$ achieves its minimum at $x = 1$. This proves (5.12) when $J = J^+$.

Case two, $J = J^-$: Setting for brevity

$$x = A(\gamma), \quad z = \sum_{i \in J} (-A_i),$$

we have $x, z \geq 0$, and, thanks to $I \neq \emptyset$,

$$\sigma(\gamma, \mathbf{t}) = \frac{\#I}{2} + \sum_{j \in J} A_j \geq \frac{1}{2} - z. \quad (5.15)$$

Now, by Dido's inequality, we have that

$$L(\gamma_i) \geq \sqrt{2\pi} |A_i|, \quad \forall i = 1, \dots, k, \quad (5.16)$$

so that $A_i > 1/2$ for $i \in I$ implies that

$$L(\gamma) = \sum_{i=1}^k L(\gamma_i) \geq \sqrt{2\pi} \sum_{i=1}^k \sqrt{|A_i|} \geq \sqrt{\pi} \#I + \sqrt{2\pi} \sum_{i \in J} \sqrt{|A_i|}.$$

Since $|A_i| \leq 1/2$ for $i \in J$ and $\sqrt{2}|t| \leq \sqrt{|t|}$ for $|t| \leq 1/2$ we conclude that

$$L(\gamma) \geq \sqrt{\pi} \#I + 2\sqrt{\pi} \sum_{i \in J} |A_i| \geq \sqrt{\pi} + 2\sqrt{\pi} z, \quad (5.17)$$

taking into account $I \neq \emptyset$ and the definition of z . By using as done in the other case that $a(k-6) \geq -4a$, we thus obtain a first lower bound on $L(\gamma) + \varepsilon(\gamma, \mathbf{t})$, namely

$$\begin{aligned} L(\gamma) + \varepsilon(\gamma, \mathbf{t}) &= L(\gamma) + a(k-6) + (12)^{1/4} \sigma(\gamma, \mathbf{t}) - 2(12)^{1/4} \min\{1, A(\gamma)\} \\ &\geq \sqrt{\pi} + 2\sqrt{\pi} z - 4a + (12)^{1/4} \left(\frac{1}{2} - z\right) - 2(12)^{1/4} \\ &\geq \left(2\sqrt{\pi} - (12)^{1/4}\right) z + \left(\sqrt{\pi} - \frac{3}{2}(12)^{1/4}\right) - 4a. \end{aligned} \quad (5.18)$$

This lower bound will be sufficient to prove (5.12) only for certain values of z . For this reason, before discussing the latter point, we obtain a second, complementary, lower bound, that combined with (5.18) will allow us to deduce (5.12). To obtain this second lower

bound, consider $\bar{\gamma}$ obtained from γ by “reflecting” with respect to their chords all the circular arcs γ_i corresponding to $i \in J$. In this way $L(\bar{\gamma}) = L(\gamma)$, while $A(\bar{\gamma}_i) = -A(\gamma_i)$ for all $i \in J$, gives

$$A(\bar{\gamma}) = \sum_{i \in I} A(\gamma_i) - \sum_{i \in J} A(\gamma_i) = A(\gamma) - 2 \sum_{i \in J} A_i = x + 2z. \quad (5.19)$$

By applying the isoperimetric inequality to $\bar{\gamma}$ (notice that $A(\bar{\gamma}) \geq A(\gamma) \geq 0$) and using (5.19) we thus find

$$L(\gamma) = L(\bar{\gamma}) \geq 2\sqrt{\pi} \sqrt{A(\bar{\gamma})} \geq 2\sqrt{\pi} \sqrt{x + 2z}, \quad (5.20)$$

which, combined with (5.15), gives

$$\begin{aligned} L(\gamma) + \varepsilon(\gamma, \mathbf{t}) &= L(\gamma) + a(k-6) + (12)^{1/4} \sigma(\gamma, \mathbf{t}) - 2(12)^{1/4} \min\{1, A(\gamma)\} \\ &\geq 2\sqrt{\pi} \sqrt{x + 2z} - 4a + (12)^{1/4} \left(\frac{1}{2} - z\right) - 2(12)^{1/4} \min\{1, x\}, \\ &\geq 2\sqrt{\pi} \sqrt{1 + 2z} - 4a + (12)^{1/4} \left(\frac{1}{2} - z\right) - 2(12)^{1/4}, \end{aligned} \quad (5.21)$$

where we have used $x \geq 0 \mapsto 2\sqrt{\pi} \sqrt{x + 2z} - 2(12)^{1/4} \min\{1, x\}$ has a minimum at $x = 1$. In summary, setting

$$\begin{aligned} f(z) &= \left(2\sqrt{\pi} - (12)^{1/4}\right)z + \left(\sqrt{\pi} - \frac{3}{2}(12)^{1/4}\right) - 4a, \\ g(z) &= 2\sqrt{\pi} \sqrt{1 + 2z} - (12)^{1/4}z - \frac{3}{2}(12)^{1/4} - 4a, \end{aligned}$$

and $h = \max\{f, g\}$, we are left to prove that

$$\inf_{z \geq 0} h(z) = \min\left\{\frac{7}{10}, c(a)\right\}.$$

Since f is an affine, increasing function, g is concave on $[0, \infty)$, and $f(3/2) = g(3/2)$, we have that $h = g$ on $[0, 3/2]$ and $h = f$ on $[3/2, \infty)$. In particular, if $z \geq 3/2$, then, recalling (5.11), we find

$$h(z) = f(z) \geq f(3/2) = 4\sqrt{\pi} - 3(12)^{1/4} - 4a \geq 2\sqrt{\pi} - \frac{3}{2}(12)^{1/4} \geq \frac{7}{10}.$$

If, instead, $z \in [0, 3/2]$, then $h(z) \geq g(z) \geq g(0) = c(a)$ by definition (5.13) of $c(a)$.

Step five: We conclude the proof of the theorem. Based on the previous four steps, we have to prove the following reduced version of our opening claim:

Reduced Claim: There are positive constants a and c such that the following holds. Let $k \geq 2$ and $(\gamma, \mathbf{t}) \in \text{Cir}_k$ satisfy

$$A(\gamma) \geq \frac{1}{100}, \quad \text{if } 2 \leq k \leq 5, \quad (5.22)$$

and let either $0 \leq A_i \leq 1/2$ for all $i = 1, \dots, k$ or $0 \geq A_i \geq -1/2$ for all $i = 1, \dots, k$. Then, when $k \neq 6$,

$$L(\gamma) + a(k-6) + (12)^{1/4} \sigma(\gamma, \mathbf{t}) \geq 2(12)^{1/4} \min\{1, A(\gamma)\} + c; \quad (5.23)$$

and, when $k = 6$ and $1/5 \geq A(\gamma) > 0$,

$$L(\gamma) + (12)^{1/4} \sigma(\gamma, \mathbf{t}) > 2(12)^{1/4} \min\{1, A(\gamma)\}. \quad (5.24)$$

To begin the proof of this reduced claim, let us recall that, so far, we have only imposed on a the constraint (5.11). Since $(\sqrt{\pi}/2) - (3/8)(12)^{1/4} > 1/10$, we can work with any $a \leq 1/10$. The choice

$$a = \frac{3}{50}, \quad (5.25)$$

is enforced from now on, for the sake of definiteness.

Next, we notice that the case $k = 2$ is easily dealt with. Indeed, in this case, $|A_i| \leq 1/2$ implies $\sigma(\gamma, \mathbf{t}) = A_1 + A_2 = A(\gamma)$ as well as $\min\{A(\gamma), 1\} = A(\gamma)$: therefore, by applying the isoperimetric inequality we find that

$$L(\gamma) + \varepsilon(\gamma, \mathbf{t}) \geq 2\sqrt{\pi A(\gamma)} - 4a - (12)^{1/4} A(\gamma) \geq \min\left\{h(1), h\left(\frac{1}{100}\right)\right\} > \frac{9}{100},$$

where we have used that $h(x) = 2\sqrt{\pi x} - (12)^{1/4}x - (6/25)$ is concave on $[0, 1]$ and that $1 \geq A(\gamma) \geq 1/100$.

When $k \geq 3$ there is no immediate relation between $x = A(\gamma)$ and $y = \sigma(\gamma, \mathbf{t}) = \sum_{i=1}^k A_i$ that we can use. To discuss this case it is convenient to collect the following four lower bounds:

$$L(\gamma) \geq 2\sqrt{\pi x}, \quad (5.26)$$

$$L(\gamma) \geq 2\sqrt{\pi(x + 2y^-)}, \quad (5.27)$$

$$L(\gamma) \geq 2\sqrt{\pi}|y|, \quad (5.28)$$

$$L(\gamma) \geq 2\sqrt{k \tan\left(\frac{\pi}{k}\right) \sqrt{(x - y)^+}}, \quad (5.29)$$

where $z^+ = \max\{z, 0\}$ and $z^- = \max\{-z, 0\}$. Of course (5.26) is just the isoperimetric inequality, as already used repeatedly, while (5.27) follows from the isoperimetric inequality by the argument used in proving (5.20). Concerning (5.28), by Dido's inequality $L(\gamma_i) \geq \sqrt{2\pi|A_i|}$ with $|A_i| \leq 1/2$ for each i we obtain

$$L(\gamma) \geq \sqrt{2\pi} \sum_{i=1}^k \sqrt{|A_i|} \geq 2\sqrt{\pi} \sum_{i=1}^k |A_i| = 2\sqrt{\pi} |\sigma(\gamma, \mathbf{t})|,$$

where in the last identity we have taken into account that, under our assumptions on γ , all the A_i 's have the same sign. Finally, let us define $\pi_\gamma \in \text{Lip}(\mathbb{S}^1; \mathbb{R}^2)$ by taking

$$\pi_\gamma = [\gamma(\mathbf{t}(1)), \gamma(\mathbf{t}(2))] \circ [\gamma(\mathbf{t}(2)), \gamma(\mathbf{t}(3))] \circ \cdots \circ [\gamma(\mathbf{t}(k-1)), \gamma(\mathbf{t}(k))] \circ [\gamma(\mathbf{t}(k)), \gamma(\mathbf{t}(1))].$$

Clearly $L(\gamma) \geq L(\pi_\gamma)$ and π_γ is a polygonal curve with k vertexes. By the polygonal isoperimetric inequality *for immersed curves*⁴ we have

$$L(\pi_\gamma)^2 \geq 4k \tan\left(\frac{\pi}{k}\right) A(\pi_\gamma) = 4k \tan\left(\frac{\pi}{k}\right) (A(\gamma) - \sigma(\gamma, \mathbf{t})), \quad (5.30)$$

where in asserting $A(\pi_\gamma) = A(\gamma) - \sigma(\gamma, \mathbf{t})$ we have used $|A_i| \leq 1/2$ for all $i = 1, \dots, k$.

Based on (5.26), (5.27), (5.28), and (5.29), we now set

$$\begin{aligned} g_k(x, y) &= a(k-6) + (12)^{1/4} \left(y - 2 \min\{1, x\}\right), \\ f_{k,1}(x, y) &= g_k(x, y) + 2\sqrt{\pi x}, \\ f_{k,2}(x, y) &= g_k(x, y) + 2\sqrt{\pi(x + 2y^-)}, \\ f_{k,3}(x, y) &= g_k(x, y) + 2\sqrt{\pi}|y|, \\ f_{k,4}(x, y) &= g_k(x, y) + 2\sqrt{k \tan\left(\frac{\pi}{k}\right) \sqrt{(x - y)^+}}. \end{aligned}$$

Taking into account that each $f_{k,i}(\cdot, y)$ is increasing on $x \geq 1$, and setting

$$f_k = \max\{f_{k,1}, f_{k,2}, f_{k,3}, f_{k,4}\},$$

⁴The point here is that π_γ may not be injective, therefore we are not able to bound $L(\pi_\gamma)$ from below by directly using the polygonal isoperimetric inequality *for sets*, namely (1.17)!

we can conclude the proof of the reduced claim, and thus of the theorem, by showing that

$$\inf_{k \geq 7} \inf_{[0,1] \times \mathbb{R}} f_k > 0, \quad (5.31)$$

$$\inf_{\mathbb{R}} f_6(x, \cdot) > 0, \quad \forall x \in \left(0, \frac{1}{5}\right], \quad (5.32)$$

$$\inf_{3 \leq k \leq 5} \inf_{[1/100, 1] \times \mathbb{R}} f_k > 0. \quad (5.33)$$

To prove this we will combine different lower bounds on each the $f_{k,i}$'s.

First lower bound: We prove the existence of a positive (computable) constant c_0 such that, for all $k \geq 3$,

$$f_k(x, y) \geq c_0, \quad \forall (x, y) \in [0, 1] \times \left\{ \left(-\infty, -\frac{5}{2}\right] \cup \left[\frac{4}{5}, \infty\right) \right\}. \quad (5.34)$$

Indeed, if $k \geq 3$, $x \in [0, 1]$, and $y \in \mathbb{R}$, then

$$\begin{aligned} f_k(x, y) &\geq f_{k,3}(x, y) = 2\sqrt{\pi}|y| + a(k-6) + (12)^{1/4}y - 2(12)^{1/4}x \\ &\geq 2\sqrt{\pi}|y| + (12)^{1/4}y - 2(12)^{1/4} - 3a \end{aligned}$$

This last function is non-negative if and only if (recall that $a = 3/50$)

$$\text{either } y \geq \frac{2(12)^{1/4} + (9/50)}{2\sqrt{\pi} + (12)^{1/4}}, \quad \text{or } y \leq -\frac{2(12)^{1/4} + (9/50)}{2\sqrt{\pi} - (12)^{1/4}}.$$

Taking into account that

$$\frac{2(12)^{1/4} + (9/50)}{2\sqrt{\pi} + (12)^{1/4}} < \frac{4}{5}, \quad \frac{2(12)^{1/4} + (9/50)}{2\sqrt{\pi} - (12)^{1/4}} < \frac{5}{2},$$

we conclude that $2\sqrt{\pi}|y| + (12)^{1/4}y - 2(12)^{1/4} - 3a$ is uniformly positive when either $y \leq -5/2$ or $y \geq 4/5$.

Second lower bound: We prove that, for all $k \geq 3$,

$$f_k(x, y) \geq f_{k,1}(x, 0) \quad \forall x \in [0, 1], y \geq -\frac{5}{2}. \quad (5.35)$$

Since $f_k \geq f_{k,1}$, this is obvious from the definition of $f_{k,1}$ when $y \geq 0$ (one just drops the term $(12)^{1/4}y$). Assuming now that $y \in [-5/2, 0]$ we notice that

$$\begin{aligned} f_k(x, y) &\geq f_{k,2}(x, y) = 2\sqrt{\pi(x+2y^-)} + a(k-6) + (12)^{1/4}y - 2(12)^{1/4}x \\ &= 2\sqrt{\pi(x+2|y|)} - (12)^{1/4}|y| + a(k-6) - 2(12)^{1/4}x \\ &\geq 2\sqrt{\pi x} + a(k-6) - 2(12)^{1/4}x = f_{k,1}(x, 0), \end{aligned}$$

where in the last step we have used that

$$F(x, b) = 2\sqrt{\pi(x+2b)} - (12)^{1/4}b - 2\sqrt{\pi x} \geq 0, \quad \forall (x, b) \in [0, 1] \times \left[0, \frac{5}{2}\right]. \quad (5.36)$$

To prove (5.36) we just notice that, for every $x \geq 0$, $F(x, \cdot)$ is concave on $[0, \infty)$, so that

$$\inf_{[0, 5/2]} F(x, \cdot) \geq \min \left\{ F(x, 0), F\left(x, \frac{5}{2}\right) \right\} = \min \left\{ 0, F\left(x, \frac{5}{2}\right) \right\}.$$

Now, for every $x \in (0, 1]$ we have

$$\begin{aligned} F\left(x, \frac{5}{2}\right) &= 2\sqrt{\pi}\sqrt{x+5} - \frac{5(12)^{1/4}}{2} - 2\sqrt{\pi x}, \\ \frac{\partial F}{\partial x}\left(x, \frac{5}{2}\right) &= \sqrt{\frac{\pi}{x+5}} - \sqrt{\frac{\pi}{x}} < 0, \end{aligned}$$

so that

$$\inf_{0 \leq x \leq 1} F\left(x, \frac{5}{2}\right) \geq F\left(1, \frac{5}{2}\right) = 2\sqrt{6\pi} - \frac{5(12)^{1/4}}{2} - 2\sqrt{\pi} > \frac{2}{5}.$$

This proves (5.36), and thus (5.35).

Third lower bound: Motivated by (5.35), we prove the existence of a (computable) positive constant c_1 such that

$$f_{k,1}(x, 0) \geq c_1, \quad \forall x \in [0, 1], \quad \text{if } k \geq 9, \quad (5.37)$$

$$f_{k,1}(x, 0) \geq c_1, \quad \forall x \in \left[0, \frac{9}{10}\right], \quad \text{if } 7 \leq k \leq 8, \quad (5.38)$$

$$f_{k,1}(x, 0) \geq c_1, \quad \forall x \in \left[\frac{1}{100}, \frac{4}{5}\right], \quad \text{if } 3 \leq k \leq 5, \quad (5.39)$$

and also show that

$$f_{6,1}(x, 0) > 0, \quad \forall x \in \left(0, \frac{9}{10}\right]. \quad (5.40)$$

To prove this, let us notice that

$$f_{k,1}(x, 0) = 2\sqrt{\pi x} + a(k-6) - 2(12)^{1/4}x = -q_k(\sqrt{x}),$$

where

$$q_k(t) = 2(12)^{1/4}t^2 - 2\sqrt{\pi}t - a(k-6).$$

The roots of q_k are given by

$$t_k^\pm = \frac{2\sqrt{\pi} \pm \sqrt{4\pi + 4a(k-6)2(12)^{1/4}}}{4(12)^{1/4}}.$$

Thanks to $a = 3/50$ and $k \geq 3$ we have

$$4\pi + 4a(k-6)2(12)^{1/4} \geq 4\pi - \frac{36}{25}(12)^{1/4} > 9.$$

Therefore both $t_k^\pm$ are real, with

$$\begin{aligned} t_k^- &< 0 < t_k^+, & \text{if } k \geq 7, \\ t_6^- &= 0 < t_6^+, \\ 0 &< t_k^- < t_k^+, & \text{if } 3 \leq k \leq 5. \end{aligned}$$

In particular, if $k \geq 6$, then we have

$$\{x \geq 0 : f_{k,1}(x, 0) \geq 0\} = [0, z_k], \quad (5.41)$$

$$\text{where } z_k := (t_k^+)^2 = \frac{\pi + a(k-6)(12)^{1/4} + \sqrt{\pi} \sqrt{\pi + a(k-6)2(12)^{1/4}}}{2(12)^{1/2}},$$

while if $3 \leq k \leq 5$, then (with z_k as in (5.41))

$$\{x \geq 0 : f_{k,1}(x, 0) \geq 0\} = [y_k, z_k], \quad (5.42)$$

$$\text{where } y_k := (t_k^-)^2 = \frac{\pi + a(k-6)(12)^{1/4} - \sqrt{\pi} \sqrt{\pi + a(k-6)2(12)^{1/4}}}{2(12)^{1/2}},$$

Exploiting $a = 3/50$ we find that if $k \geq 9$, then

$$z_k \geq z_9 = \frac{\pi + (9/50)(12)^{1/4} + \sqrt{\pi} \sqrt{\pi + (9/25)(12)^{1/4}}}{2(12)^{1/2}} > 1,$$

and since the last sign is strict (and $t_k^- < 0$ for $k \geq 7$) we conclude that

$$\inf_{[0,1]} f_{k,1}(\cdot, 0) \geq \inf_{[0,1]} f_{9,1}(\cdot, 0) > 0, \quad \forall k \geq 9. \quad (5.43)$$

For $8 \geq k \geq 6$ we have

$$z_k \geq z_6 = \frac{\pi}{(12)^{1/2}} > \frac{9}{10}, \quad (5.44)$$

and the strict sign in (5.44), $t_k^- < 0$ if $k = 7, 8$, and $t_6^- = 0$, gives

$$\min_{k=7,8} \inf_{[0,9/10]} f_{k,1}(\cdot, 0) \geq \inf_{[0,9/10]} f_{7,1}(\cdot, 0) > 0, \quad (5.45)$$

and prove (5.40), respectively. Finally, for the remaining cases $3 \leq k \leq 5$, we see that $k \geq 3$ gives

$$z_k \geq z_3 = \frac{\pi - (9/50)(12)^{1/4} + \sqrt{\pi} \sqrt{\pi - (9/25)(12)^{1/4}}}{2(12)^{1/2}} > \frac{4}{5}; \quad (5.46)$$

moreover, by differentiation, $t \in [2, 6] \mapsto y_t$ is decreasing on $[2, 6]$ and takes its minimum value at $k = 6$ ($y_6 = 0$), so that, for $3 \leq k \leq 5$, we have

$$y_k \leq y_3 = \frac{\pi - (9/50)(12)^{1/4} - \sqrt{\pi} \sqrt{\pi - (9/25)(12)^{1/4}}}{2(12)^{1/2}} < \frac{1}{500} < \frac{1}{100}. \quad (5.47)$$

Thanks to the strict signs in (5.46) and (5.47) we find that

$$\inf_{3 \leq k \leq 8} \inf_{[1/100, 4/5]} f_{k,1}(x, 0) > 0. \quad (5.48)$$

We conclude the proof of (5.37), (5.38), and (5.39) by setting

$$c_1 = \min \left\{ \inf_{[0,1]} f_{9,1}(\cdot, 0), \inf_{[0,9/10]} f_{7,1}(\cdot, 0), \min_{3 \leq k \leq 5} \inf_{[1/100, 4/5]} f_{k,1}(\cdot, 0) \right\},$$

where $c_1 > 0$ by (5.43), (5.45), and (5.48).

Conclusion of the proof: We conclude the proof of the theorem by proving (5.31), (5.32) and (5.33). To begin with, we notice that (5.34), (5.35) and (5.40) imply (5.32).

Next, we prove (5.31). First of all, by (5.34), (5.35), and (5.37), we find that

$$\inf_{[0,1] \times \mathbb{R}} f_k \geq \min\{c_0, c_1\} > 0, \quad \forall k \geq 9. \quad (5.49)$$

that is (5.31) restricted to $k \geq 9$. Combining (5.49) with (5.34), (5.35), and (5.38), we see that to complete the proof of (5.31) we are left to show that

$$\min_{k=7,8} \inf_{[9/10,1] \times [-5/2, 4/5]} f_k > 0. \quad (5.50)$$

We begin the proof of (5.50) by showing that

$$\min_{k=7,8} \inf_{[9/10,1] \times [-5/2, -1/10]} f_k > 0. \quad (5.51)$$

The idea is obtaining a positive lower bound on $f_{k,2}$ all the way up to $y \leq -1/10$ under the restriction $x \in [9/10, 1]$ (recall that (5.35) holds only up to $y \leq -5/2$, but on the wider range $x \in [0, 1]$). To this end, let us recall that, if $y \leq 0$, then

$$f_k(x, y) \geq f_{k,2}(x, y) = 2\sqrt{\pi(x + 2|y|)} - (12)^{1/4}|y| + a(k - 6) - 2(12)^{1/4}x;$$

now, $f_{k,2}(\cdot, y)$ ($y \leq 0$) is decreasing on $[1/2, 1]$ (and, thus, on $[9/10, 1]$) since

$$\frac{\partial f_{k,2}}{\partial x}(x, y) = \sqrt{\frac{\pi}{x + 2|y|}} - 2(12)^{1/4} \leq \sqrt{\frac{\pi}{1/2}} - 2(12)^{1/4} < -1,$$

therefore, for all $y \leq 0$ and $k \geq 7$,

$$\inf_{[9/10,1]} f_k(\cdot, y) \geq f_{k,2}(1, y) \geq f_{7,2}(1, y) = 2\sqrt{\pi(1 + 2|y|)} + \frac{3}{50} - (12)^{1/4}|y| - 2(12)^{1/4}.$$

Since $f_{7,2}(1, \cdot)$ is concave on $(-\infty, 0]$ we find

$$\begin{aligned} \min_{k=7,8} \inf_{[9/10,1] \times [-5/2,-1/10]} f_k &\geq \inf_{[-5/2,-1/10]} f_{7,2}(1, \cdot) \\ &= \min \left\{ f_{7,2}\left(1, -\frac{5}{2}\right), f_{7,2}\left(1, -\frac{1}{10}\right) \right\} \geq \min \left\{ \frac{3}{10}, \frac{3}{100} \right\}. \end{aligned}$$

(Notice that we cannot extend this lower bound to $[-5/2, 0]$ since $f_{7,2}(1, 0) < 0$.) Having proved (5.51), we next show

$$\min_{k=7,8} \inf_{[9/10,1] \times [1/10,4/5]} f_k > 0. \quad (5.52)$$

Working with $f_{k,1}$, and using the fact that $f_{k,1}(x, y)$ is increasing in y and concave in x , we find that

$$\begin{aligned} \min_{k=7,8} \inf_{[9/10,1] \times [1/10,4/5]} f_k &\geq \inf_{[9/10,1]} f_{7,1}\left(\cdot, \frac{1}{10}\right) \\ &\geq \min \left\{ f_{7,1}\left(\frac{9}{10}, \frac{1}{10}\right), f_{7,1}\left(1, \frac{1}{10}\right) \right\} \geq \min \left\{ \frac{1}{4}, \frac{3}{50} \right\}, \end{aligned}$$

thus proving (5.52). Thanks to (5.51) and (5.52), in order to prove (5.50), and thus to conclude the proof of (5.31), we are left to show that

$$\min_{k=7,8} \inf_{[9/10,1] \times [-1/10,1/10]} f_k > 0. \quad (5.53)$$

To this end we shall bound f_k from below by using $f_{k,4}$. We first notice that since $x \geq 9/10$, $y \leq 1/10$, $k \mapsto k \tan(\pi/k)$ is decreasing, and $k \geq 7$, we have

$$\frac{\partial f_{k,4}}{\partial x}(x, y) = \sqrt{\frac{k \tan(\pi/k)}{x-y}} - 2(12)^{1/4} \leq \sqrt{\frac{7 \tan(\pi/7)}{(8/10)}} - 2(12)^{1/4} < -1.$$

Hence

$$\min_{k=7,8} \inf_{[9/10,1] \times [-1/10,1/10]} f_k \geq \min_{k=7,8} \inf_{[-1/10,1/10]} f_{k,4}(1, \cdot). \quad (5.54)$$

Next we notice that

$$f_{k,4}(1, y) = 2 \sqrt{k \tan\left(\frac{\pi}{k}\right)} \sqrt{1-y} + a(k-6) + (12)^{1/4}(y-2),$$

is concave on $y \in [-1/10, 1/10]$, with

$$\begin{aligned} f_{7,4}\left(1, -\frac{1}{10}\right) &\geq \frac{1}{500}, & f_{7,4}\left(1, \frac{1}{10}\right) &\geq \frac{7}{1000}, \\ f_{8,4}\left(1, -\frac{1}{10}\right) &\geq \frac{1}{50}, & f_{8,4}\left(1, \frac{1}{10}\right) &\geq \frac{3}{100}, \end{aligned}$$

which, combined with (5.54), leads to (5.53). This completes the proof of (5.31).

We finally prove (5.33). Thanks to (5.34), (5.35), and (5.39), it is enough to show that

$$\min_{3 \leq k \leq 5} \inf_{[7/10,1] \times [-5/2,4/5]} f_k > 0. \quad (5.55)$$

To this end, we first show that

$$\min_{3 \leq k \leq 5} \inf_{[7/10,1] \times [-5/2,-3/10]} f_k > 0. \quad (5.56)$$

Indeed, as already noticed in the proof of (5.51), for every $y \leq 0$, $f_{k,2}(\cdot, y)$ is decreasing on $[1/2, 1]$. In particular,

$$\min_{3 \leq k \leq 5} \inf_{[7/10,1] \times [-5/2,-3/10]} f_k \geq \inf_{[-5/2,-3/10]} f_{3,2}(1, \cdot),$$

where we have also used that $k \mapsto f_{k,2}(x, y)$ is increasing. Now,

$$f_{3,2}(1, y) = 2 \sqrt{\pi(1+2|y|)} - (12)^{1/4}|y| - 3a - 2(12)^{1/4},$$

is concave on $y \in [-5/2, -3/10]$, with

$$f_{3,2}\left(1, -\frac{5}{2}\right) > \frac{1}{10}, \quad f_{3,2}\left(1, -\frac{3}{10}\right) > \frac{1}{50},$$

so that (5.56) follows. We next prove that

$$\min_{k=3,4} \inf_{[4/5,1] \times [-3/10,1/5]} f_k > 0, \quad (5.57)$$

$$\inf_{[4/5,1] \times [-3/10,19/100]} f_5 > 0, \quad (5.58)$$

To this end, we start noticing that when $3 \leq k \leq 5$, $x \geq 4/5$, and $y \leq 1/5$, we have

$$\frac{\partial f_{k,4}}{\partial x}(x, y) = \sqrt{\frac{k \tan(\pi/k)}{x-y}} - 2(12)^{1/4} \leq \sqrt{\frac{3 \tan(\pi/3)}{3/5}} - 2(12)^{1/4} \leq -\frac{4}{5}.$$

In particular, using the concavity in y of $f_{k,4}(1, y)$, we find

$$\begin{aligned} \min_{k=3,4} \inf_{[4/5,1] \times [-3/10,1/5]} f_k &\geq \min_{k=3,4} \inf_{[-3/10,1/5]} f_{k,4}(1, \cdot) \\ &= \min_{k=3,4} \min \left\{ f_{k,4}\left(1, -\frac{3}{10}\right), f_{k,4}\left(1, \frac{1}{5}\right) \right\}, \end{aligned}$$

where

$$\begin{aligned} f_{3,4}\left(1, -\frac{3}{10}\right) &\geq \frac{7}{10}, & f_{3,4}\left(1, \frac{1}{5}\right) &\geq \frac{1}{2}, \\ f_{4,4}\left(1, -\frac{3}{10}\right) &\geq \frac{1}{10}, & f_{4,4}\left(1, \frac{1}{5}\right) &\geq \frac{1}{10}, \end{aligned}$$

thus proving (5.57). In the case $k = 5$ we would like to repeat the same argument but, unfortunately, $f_{5,4}(1, 1/5) < 0$. This is why, when $k = 5$, we need to stop using the $f_{k,4}$ bound a bit below $y = 1/5$. Stopping at $y = 19/100$ works, and we find

$$\inf_{[4/5,1] \times [-3/10,19/100]} f_5 \geq \inf_{[-3/10,19/100]} f_{5,4}(1, y) \geq \min \left\{ f_{5,4}\left(1, -\frac{3}{10}\right), f_{5,4}\left(1, \frac{19}{100}\right) \right\},$$

where

$$f_{5,4}\left(1, -\frac{3}{10}\right) \geq \frac{1}{200}, \quad f_{5,4}\left(1, \frac{19}{100}\right) \geq \frac{1}{1000},$$

thus proving (5.58). Thanks to (5.56), (5.57), and (5.58), in order to deduce (5.55) (and thus complete the proof of the theorem), we are left to show that

$$\min_{k=3,4} \inf_{[4/5,1] \times [1/5,4/5]} f_k > 0, \quad (5.59)$$

$$\inf_{[4/5,1] \times [19/100,4/5]} f_5 > 0. \quad (5.60)$$

Using that $f_{k,1}$ is increasing in y and concave in x we find that

$$\min_{k=3,4} \inf_{[4/5,1] \times [1/5,4/5]} f_k \geq \min_{k=3,4} \inf_{[4/5,1]} f_{k,1}\left(\cdot, \frac{1}{5}\right) \geq \min_{k=3,4} \min \left\{ f_{k,1}\left(\frac{4}{5}, \frac{1}{5}\right), f_{k,1}\left(1, \frac{1}{5}\right) \right\},$$

where

$$\begin{aligned} f_{3,1}\left(\frac{4}{5}, \frac{1}{5}\right) &\geq \frac{3}{10}, & f_{3,1}\left(1, \frac{1}{5}\right) &\geq \frac{1}{100}, \\ f_{4,1}\left(\frac{4}{5}, \frac{1}{5}\right) &\geq \frac{2}{5}, & f_{4,1}\left(1, \frac{1}{5}\right) &\geq \frac{7}{100}, \end{aligned}$$

thus proving (5.59). Similarly,

$$\inf_{[4/5,1] \times [19/100,4/5]} f_5 \geq \inf_{[4/5,1]} f_{5,1}\left(\cdot, \frac{19}{100}\right) \geq \min \left\{ f_{5,1}\left(\frac{4}{5}, \frac{19}{100}\right), f_{5,1}\left(1, \frac{19}{100}\right) \right\},$$

where

$$f_{5,1}\left(\frac{4}{5}, \frac{19}{100}\right) \geq \frac{2}{5}, \quad f_{5,1}\left(1, \frac{19}{100}\right) \geq \frac{1}{10},$$

thus leading to (5.60). This completes the proof of the theorem. $\square$

APPENDIX A. THE ISOPERIMETRIC INEQUALITY FOR IMMERSSED POLYGONS

In the proof of Theorem 1.2, and specifically in the proof of Theorem 5.1, Section 5, we have claimed that the isoperimetric inequality for k -gon (1.17) is valid for *immersed* polygons (with the notion of oriented area used in place of the standard notion of area); see (5.30). Since we have not been able to find a proof of this more general inequality in the literature, we include one in this appendix.

We identify a generic immersed polygon with k -edges with an ordered collection Π_k of points $p_j \in \mathbb{R}^2$, $j = 1, \dots, k$. Setting $p_j = (x_j, y_j)$ for the coordinates of these points, with $p_{k+1} = p_1$ the perimeter and oriented area of Π_k are given by

$$P(\Pi_k) = \sum_{j=1}^k \sqrt{(x_j - x_{j+1})^2 + (y_j - y_{j+1})^2},$$

$$A(\Pi_k) = \frac{1}{2} \sum_{j=1}^k (x_j y_{j+1} - x_{j+1} y_j).$$

Theorem A.1 (Isoperimetry for immersed k -gons). *For every Π_k as above we have*

$$P(\Pi_k)^2 \geq p(k)^2 A(\Pi_k),$$

where $p(k)$ is the perimeter of a unit-area regular k -gon as defined in (1.18).

Proof. After scaling, we can prove Theorem A.1 by showing that the maximum of $A(\Pi_k)$ under the constraint $P(\Pi_k) = 1$ is achieved when Π_k is a regular k -gon with unit-perimeter. Clearly a maximum point Π_k exists⁵ and satisfies $A(\Pi_k) > 0$. As shown in [KP08, Leg18], the fact that Π_k is a critical point of A at P fixed implies that all the vertexes of Π_k belong to a circle or to a line, and since $A(\Pi_k) > 0$, the first case holds. For some $R > 0$ we can thus set $p_j = (R \cos \theta_j, R \sin \theta_j)$ for $1 \leq j \leq k$. Setting

$$x \equiv y \quad \text{if and only if} \quad x = y + 2\pi h \quad \text{for some } h \in \mathbb{Z},$$

extending the definition of θ_j to every $j \in \mathbb{Z}$ by setting $\theta_{j+hk} = \theta_j$ for all $j = 1, \dots, k$ and $h \in \mathbb{Z}$, and noticing that $\theta_{j+1} \not\equiv \theta_j$ for all j (for, otherwise, Π_k would have less than k edges), we see that

$$A(\Pi_k) = \frac{R^2}{2} \sum_{j=1}^k \sin(\theta_{j+1} - \theta_j), \quad P(\Pi_k) = \sqrt{2} R \sum_{j=1}^k \sqrt{1 - \cos(\theta_{j+1} - \theta_j)}.$$

Exploiting $P(\Pi_k) = 1$ to solve for R , we come to define

$$a(\theta_1, \dots, \theta_k) := 4 A(\Pi_k) = \frac{N}{D^2}, \quad \text{where}$$

$$\omega_j = \theta_{j+1} - \theta_j, \quad N = \sum_{j=1}^k \sin(\omega_j), \quad D = \sum_{j=1}^k \sqrt{1 - \cos(\omega_j)}.$$

We prove the theorem by showing that, at a maximum point of a , it holds $\omega_j \equiv 2\pi/k$ for all j .

⁵A compactness argument shows the existence of a maximizer with at least 3 and at most k edges. To see that a maximizer has exactly k edges one notices that, should this not be the case, area could be increased by first taking a “triangular variation” of one edge (thus adding one additional edge), and then by rescaling so to preserve the perimeter constraint.

Step one: We prove that

$$\omega_j \equiv \omega_1, \quad \text{if } j \text{ is odd}, \quad (\text{A.1})$$

$$\omega_j \equiv \omega_2, \quad \text{if } j \text{ is even}. \quad (\text{A.2})$$

To this end, it will suffice to show

$$\theta_j - \theta_{j-1} \equiv \theta_{j+2} - \theta_{j+1}, \quad \forall j. \quad (\text{A.3})$$

We start by computing (recall that $\omega_j \neq 0$ for all j)

$$\frac{\partial a}{\partial \theta_j} = \frac{\cos(\omega_{j-1}) - \cos(\omega_j)}{D^2} + \frac{N}{D^3} \left\{ \frac{\sin(\omega_{j-1})}{\sqrt{1 - \cos(\omega_{j-1})}} - \frac{\sin(\omega_j)}{\sqrt{1 - \cos(\omega_j)}} \right\}.$$

Since $\cos(x) = \cos^2(x/2) - \sin^2(x/2)$ implies $\sqrt{1 - \cos(x)} = \sqrt{2} |\sin(x/2)|$, using also $\sin(x) = 2 \sin(x/2) \cos(x/2)$, we find that

$$\frac{\sin(\omega_j)}{\sqrt{1 - \cos(\omega_j)}} = \frac{2 \sin(\omega_j/2) \cos(\omega_j/2)}{\sqrt{2} |\sin(\omega_j/2)|} = \sqrt{2} \sigma_j \cos\left(\frac{\omega_j}{2}\right)$$

where we have set $\sigma_j = \text{sign}(\sin(\omega_j/2))$. Hence,

$$D^3 \frac{\partial a}{\partial \theta_j} = D \left\{ \cos(\omega_{j-1}) - \cos(\omega_j) \right\} + \sqrt{2} N \left\{ \sigma_j \cos\left(\frac{\omega_j}{2}\right) - \sigma_{j-1} \cos\left(\frac{\omega_{j-1}}{2}\right) \right\}.$$

The condition that a achieves its maximum at $(\theta_1, \dots, \theta_k)$ thus implies that

$$\frac{\cos(\omega_j) - \cos(\omega_{j-1})}{\sigma_j \cos(\omega_j/2) - \sigma_{j-1} \cos(\omega_{j-1}/2)} = \frac{\sqrt{2} N}{D}, \quad \forall j = 1, \dots, k, \quad (\text{A.4})$$

where, notably, the right-hand side is independent of j . By $\cos(x) = \cos^2(x/2) - \sin^2(x/2) = 2 \cos^2(x/2) - 1$ and thanks to $\sigma_j^2 = 1$, we find that

$$\begin{aligned} \cos(\omega_j) - \cos(\omega_{j-1}) &= 2 \left(\cos^2(\omega_j/2) - \cos^2(\omega_{j-1}/2) \right) \\ &= 2 \left(\sigma_j \cos(\omega_j/2) - \sigma_{j-1} \cos(\omega_{j-1}/2) \right) \left(\sigma_j \cos(\omega_j/2) + \sigma_{j-1} \cos(\omega_{j-1}/2) \right), \end{aligned}$$

which combined into (A.4) gives

$$\sigma_j \cos\left(\frac{\omega_j}{2}\right) + \sigma_{j-1} \cos\left(\frac{\omega_{j-1}}{2}\right) = \frac{N}{\sqrt{2} D}, \quad \forall j = 1, \dots, k.$$

In particular, for all $j = 1, \dots, k$,

$$\sigma_{j+1} \cos\left(\frac{\omega_{j+1}}{2}\right) + \sigma_j \cos\left(\frac{\omega_j}{2}\right) = \sigma_j \cos\left(\frac{\omega_j}{2}\right) + \sigma_{j-1} \cos\left(\frac{\omega_{j-1}}{2}\right),$$

that is (setting $\theta_0 = \theta_k$, $\omega_0 = \omega_k$, and $\sigma_0 = \sigma_k$)

$$\sigma_{j+1} \cos\left(\frac{\omega_{j+1}}{2}\right) = \sigma_{j-1} \cos\left(\frac{\omega_{j-1}}{2}\right), \quad \forall j = 1, \dots, k. \quad (\text{A.5})$$

Let us now denote by t_j the unique element of $(-\pi, 0) \cup (0, \pi)$ such that $t_j \equiv \omega_j/2$, so that $\sigma_j = \text{sign}(\sin(t_j))$ and (A.5) takes the form

$$\sigma_{j+1} \cos(t_{j+1}) = \sigma_{j-1} \cos(t_{j-1}), \quad \forall j = 1, \dots, k. \quad (\text{A.6})$$

If $t_{j+1}, t_{j-1} \in (0, \pi)$, then $\sigma_{j+1} = \sigma_{j-1} = 1$, and $\cos(t_{j+1}) = \cos(t_{j-1})$ implies $t_{j+1} = t_{j-1}$; similarly, if $t_{j+1}, t_{j-1} \in (-\pi, 0)$, then $\sigma_{j+1} = \sigma_{j-1} = -1$, and (A.6) gives again $t_{j+1} = t_{j-1}$; finally, if, without loss of generality, $t_{j+1} \in (0, \pi)$ and $t_{j-1} \in (-\pi, 0)$, then (A.6) gives $\cos(t_{j+1}) = -\cos(t_{j-1})$, which in turn implies $t_{j+1} = t_{j-1} + \pi$. Looking back at the definition of t_j , this proves (A.3).

Step two: We prove that there is $n \in \{1, \dots, k-1\}$ such that, for all j ,

$$\omega_j \equiv 2\pi \frac{n}{k}, \quad \forall j. \quad (\text{A.7})$$

We have separate arguments depending on whether k is odd or even.

When k is odd, using first that $k+1$ is even in combination with (A.2), and then $\theta_{j+k} = \theta_j$ for all j , we find that

$$\omega_2 \equiv \omega_{k+1} = \theta_{k+2} - \theta_{k+1} = \theta_2 - \theta_1 = \omega_1,$$

that is, $\omega_2 \equiv \omega_1$ and thus, by (A.2) and (A.1), $\omega_j \equiv \omega_1$ for all j . By repeatedly using this fact we find

$$\theta_1 = \theta_{k+1} \equiv \omega_1 + \theta_k = \omega_1 + \sum_{j=2}^k (\theta_j - \theta_{j-1}) + \theta_1 \equiv k\omega_1 + \theta_1, \quad (\text{A.8})$$

that is, $\omega_1 \equiv 2\pi(n/k)$ for some $n \in \{1, \dots, k\}$. Since $\omega_j \not\equiv 0$ for all j , it cannot be that $n = k$, and the conclusion follows.

When k is even, using (A.2) and (A.1), in place of (A.8) we find

$$\theta_1 = \theta_{k+1} = \theta_1 + \sum_{j=2}^{k+1} (\theta_j - \theta_{j-1}) \equiv \theta_1 + \frac{k}{2}\omega_2 + \frac{k}{2}\omega_1. \quad (\text{A.9})$$

In particular, there exists $n \in \mathbb{N}$ such that

$$\omega_1 + \omega_2 = 4\pi \frac{n}{k}. \quad (\text{A.10})$$

Notice that n cannot be an integer multiple of k or of $k/2$ (for, otherwise, $\omega_2 \equiv -\omega_1$ would imply that all the points p_j lie on a same line, thus implying $A(\Pi_k) = 0$). Without loss of generality, we can thus assume that

$$\omega_1 + \omega_2 = 4\tau, \quad \tau := \pi \frac{n}{k} \quad n = 1, \dots, k-1, \quad n \neq \frac{k}{2}. \quad (\text{A.11})$$

and use (A.2), (A.1), and (A.11) to find that

$$\begin{aligned} a(\theta_1, \dots, \theta_k) &= \frac{\sum_{j=1}^k \sin(\omega_j)}{(\sum_{j=1}^k \sqrt{1 - \cos(\omega_j)})^2} \\ &= \frac{(k/2)(\sin \omega_1 + \sin \omega_2)}{(k/2)^2 (\sqrt{1 - \cos \omega_1} + \sqrt{1 - \cos \omega_2})^2} \\ &= \frac{2}{k} \frac{\sin(\omega_1) + \sin(4\tau - \omega_1)}{(\sqrt{1 - \cos(\omega_1)} + \sqrt{1 - \cos(4\tau - \omega_1)})^2}. \end{aligned}$$

Notice that, up to this point, we know that $\omega_1 \not\equiv 0$ (because, otherwise, we would have $p_1 = p_2$) and that $\omega_1 \not\equiv 4\tau$ (because, otherwise, (A.11) would give $\omega_2 \equiv 0$ and thus $p_3 = p_2$). Setting $f(\omega) = g(\omega)/h(\omega)^2$ for

$$g(\omega) = \sin(\omega) + \sin(4\tau - \omega), \quad h(\omega) = \sqrt{1 - \cos(\omega)} + \sqrt{1 - \cos(4\tau - \omega)},$$

the fact that Π_k maximizes A under the constraint that $P = 1$ implies that $\omega = \omega_1$ is a maximum point of the function $f(\omega)$ over the open set $\Omega = \{\omega \in \mathbb{R} : \omega \not\equiv 0, 4\tau\}$. Now, f is differentiable in Ω , with

$$\begin{aligned} f'(\omega) &= \frac{\cos(\omega) - \cos(4\tau - \omega)}{h(\omega)^2} - \frac{2g(\omega)}{h(\omega)^3} \left\{ \frac{\sin(\omega)}{2\sqrt{1 - \cos(\omega)}} + \frac{-\sin(4\tau - \omega)}{2\sqrt{1 - \cos(4\tau - \omega)}} \right\} \\ &= \frac{\cos(\omega) - \cos(4\tau - \omega)}{h(\omega)^2} - \frac{\sqrt{2}g(\omega)}{h(\omega)^3} \left\{ \sigma\left(\frac{\omega}{2}\right) \cos\left(\frac{\omega}{2}\right) - \sigma\left(2\tau - \frac{\omega}{2}\right) \cos\left(2\tau - \frac{\omega}{2}\right) \right\}, \end{aligned}$$

where we have used again $\sqrt{1 - \cos(x)} = \sqrt{2} |\sin(x/2)|$ and $\sin(x) = 2 \sin(x/2) \cos(x/2)$, and where we have set $\sigma(\theta) = \text{sign}(\sin \theta)$. By $|\sin \theta| = \sigma(\theta) \sin \theta$ we find

$$\begin{aligned} \frac{h(\omega)^3}{\sqrt{2}} f'(\omega) &= \left(\cos(\omega) - \cos(4\tau - \omega) \right) \left\{ \sigma\left(\frac{\omega}{2}\right) \sin\left(\frac{\omega}{2}\right) + \sigma\left(2\tau - \frac{\omega}{2}\right) \sin\left(2\tau - \frac{\omega}{2}\right) \right\} \\ &\quad - \left(\sin(\omega) + \sin(4\tau - \omega) \right) \left\{ \sigma\left(\frac{\omega}{2}\right) \cos\left(\frac{\omega}{2}\right) - \sigma\left(2\tau - \frac{\omega}{2}\right) \cos\left(2\tau - \frac{\omega}{2}\right) \right\} \\ &= -2 \sin(2\tau) \sin(\omega - 2\tau) \left\{ \sigma\left(\frac{\omega}{2}\right) \sin\left(\frac{\omega}{2}\right) + \sigma\left(2\tau - \frac{\omega}{2}\right) \sin\left(2\tau - \frac{\omega}{2}\right) \right\} \\ &\quad - 2 \sin(2\tau) \cos(\omega - 2\tau) \left\{ \sigma\left(\frac{\omega}{2}\right) \cos\left(\frac{\omega}{2}\right) - \sigma\left(2\tau - \frac{\omega}{2}\right) \cos\left(2\tau - \frac{\omega}{2}\right) \right\}. \end{aligned}$$

The conditions on τ in (A.11) guarantee that $\sin(2\tau) \neq 0$, and thus allow us to infer the following identity from the fact that ω_1 is a maximum point of f on the open set Ω ,

$$\begin{aligned} &\sigma\left(\frac{\omega_1}{2}\right) \left\{ \cos(\omega_1 - 2\tau) \cos\left(\frac{\omega_1}{2}\right) + \sin(\omega_1 - 2\tau) \sin\left(\frac{\omega_1}{2}\right) \right\} \\ &= \sigma\left(2\tau - \frac{\omega_1}{2}\right) \left\{ \cos(\omega_1 - 2\tau) \cos\left(2\tau - \frac{\omega_1}{2}\right) - \sin(\omega_1 - 2\tau) \sin\left(2\tau - \frac{\omega_1}{2}\right) \right\}, \end{aligned}$$

that is, by $\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$,

$$\sigma\left(\frac{\omega_1}{2}\right) \cos\left(2\tau - \frac{1}{2}\omega_1\right) = \sigma\left(2\tau - \frac{\omega_1}{2}\right) \cos\left(\frac{1}{2}\omega_1\right). \quad (\text{A.12})$$

If both σ 's in (A.12) have the same sign, then (by $\cos(x) = \cos(y)$ if and only if $x \equiv y$ or $x \equiv -y$) we either have

$$2\tau - \frac{1}{2}\omega_1 \equiv \frac{1}{2}\omega_1, \quad \text{i.e.} \quad \omega_1 \equiv 2\tau,$$

which implies by (A.11) that $\omega_2 \equiv 2\tau$, and hence $\omega_1 \equiv \omega_2$; or

$$2\tau - \frac{1}{2}\omega_1 \equiv -\frac{1}{2}\omega_1,$$

which leads to $2\tau \equiv 0$, in contradiction with (A.11). If, instead, the σ 's in (A.12) have opposite signs, recalling that $\cos(x) = -\cos(y) = \cos(y + \pi)$ if and only if $x \equiv y + \pi$ or $x \equiv -y + \pi$, we either have

$$2\tau - \frac{1}{2}\omega_1 \equiv \frac{1}{2}\omega_1 + \pi, \quad \text{i.e.} \quad \omega_1 \equiv 2\tau - \pi,$$

which implies by (A.11) that $\omega_2 \equiv 2\tau + \pi$, and hence also $\omega_1 \equiv \omega_2$; or

$$2\tau - \frac{1}{2}\omega_1 \equiv -\frac{1}{2}\omega_1 + \pi,$$

which leads to $2\tau \equiv \pi$, again in contradiction with (A.11). We have thus proved that $\omega_1 \equiv \omega_2$. Combining this fact with (A.11) gives (A.7).

Conclusion: We have proved so far that there is $n \in \{1, \dots, k-1\}$ such that $\omega_j = \theta_{j+1} - \theta_j = 2\pi n/k$ for all j . We are thus left to prove that $n = 1$. To this end we notice that

$$a(\theta_1, \dots, \theta_k) = \frac{\sum_{j=1}^k \sin(\omega_j)}{(\sum_{j=1}^k \sqrt{1 - \cos(\omega_j)})^2} = \frac{1}{k} \frac{\sin(2\pi n/k)}{1 - \cos(2\pi n/k)}.$$

Since $t \mapsto \sin(t)/(1 - \cos(t))$ is decreasing for $t \in (0, 2\pi)$, and since Π_k is a maximum point of A under $P = 1$, we conclude that $n = 1$. $\square$

REFERENCES

- [Alm76] F. J. Jr. Almgren. Existence and regularity almost everywhere of solutions to elliptic variational problems with constraints. *Mem. Amer. Math. Soc.*, 4(165):viii+199 pp, 1976.
- [CLM16] Marco Cicalese, Gian Paolo Leonardi, and Francesco Maggi. Improved convergence theorems for bubble clusters I. The planar case. *Indiana Univ. Math. J.*, 65(6):1979–2050, 2016.
- [CM16] M. Caroccia and F. Maggi. A sharp quantitative version of Hales’ isoperimetric honeycomb theorem. *J. Math. Pures Appl. (9)*, 106(5):935–956, 2016.
- [FAB⁺93] Joel Foisy, Manuel Alfaro, Jeffrey Brock, Nickalous Hodges, and Jason Zimba. The standard double soap bubble in $\mathbf{R}^2$ uniquely minimizes perimeter. *Pacific J. Math.*, 159(1):47–59, 1993.
- [FT43] László Fejes Toth. Über das kürzeste Kurvennetz, das eine Kugeloberfläche in flächengleiche konvexe Teile zerlegt. *Math. Naturwiss. Anz. Ungar. Akad. Wiss.*, 62:349–354, 1943.
- [Hal01] T. C. Hales. The honeycomb conjecture. *Discrete Comput. Geom.*, 25(1):1–22, 2001.
- [HM04] Aladar Heppes and Frank Morgan. Planar clusters. 2004.
- [IN15] E. Indrei and L. Nurbekyan. On the stability of the polygonal isoperimetric inequality. *Adv. Math.*, 276:62786, 2015.
- [KP08] G. Khimshiashvili and G. Panina. Cyclic polygons are critical points of area. *Zapiski Nauchnykh Seminarov POMI*, 360:238–245, 2008.
- [Leg18] J.C. Leger. Aire, périmètre et polygones cocycliques. 2018. arXiv: 1805.05423.
- [Mag12] F. Maggi. *Sets of finite perimeter and geometric variational problems*, volume 135 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2012. ISBN 978-1-107-02103-7. xx+454 pp. An introduction to Geometric Measure Theory.
- [Mor09] F. Morgan. *Geometric measure theory. A beginner’s guide. Fourth edition*. Elsevier/Academic Press, Amsterdam, 2009. viii+249 pp.
- [PT18] Emanuele Paolini and Andrea Tamagnini. Minimal clusters of four planar regions with the same area. *ESAIM Control Optim. Calc. Var.*, 24(3):1303–1331, 2018.
- [PT20] E. Paolini and V. M. Tortorelli. The quadruple planar bubble enclosing equal areas is symmetric. *Calc. Var. Partial Differential Equations*, 59(1):Paper No. 20, 9, 2020.
- [Wic04] W. Wichiramala. Proof of the planar triple bubble conjecture. *J. Reine Angew. Math.*, 567:1–49, 2004.

POLITECNICO DI MILANO, DIPARTIMENTO DI MATEMATICA, VIA EDOARDO BONARDI 9 - 20133 MILANO, ITALY

Email address: marco.caroccia@polimi.it

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF TEXAS AT AUSTIN, 2515 SPEEDWAY STOP C1200, AUSTIN, TEXAS 78712-1202, USA

Email address: kdemason@utexas.edu

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF TEXAS AT AUSTIN, 2515 SPEEDWAY STOP C1200, AUSTIN, TEXAS 78712-1202, USA

Email address: maggi@math.utexas.edu