

DEBIASING FEDERATED LEARNING WITH CORRELATED CLIENT PARTICIPATION

Zhenyu Sun*, Ziyang Zhang*, Zheng Xu†, Gauri Joshi‡, Pranay Sharma‡, Ermin Wei*

*Northwestern University, †Google Research, ‡Carnegie Mellon University

{zhenyusun2026, ziyangzhang2024}@u.northwestern.edu, xuzheng@google.com, {pranaysh, gaurij}@andrew.cmu.edu, ermin.wei@northwestern.edu

ABSTRACT

In cross-device federated learning (FL) with millions of mobile clients, only a small subset of clients participate in training in every communication round, and Federated Averaging (FedAvg) is the most popular algorithm in practice. Existing analyses of FedAvg usually assume the participating clients are independently sampled in each round from a uniform distribution, which does not reflect real-world scenarios. This paper introduces a theoretical framework that models client participation in FL as a Markov chain to study optimization convergence when clients have non-uniform and correlated participation across rounds. We apply this framework to analyze a more practical pattern: every client must wait a minimum number of R rounds (minimum separation) before re-participating. We theoretically prove and empirically observe that increasing minimum separation reduces the bias induced by intrinsic non-uniformity of client availability in cross-device FL systems. Furthermore, we develop an effective debiasing algorithm for FedAvg that provably converges to the unbiased optimal solution under arbitrary minimum separation and unknown client availability distribution.

1 INTRODUCTION

The massive amounts of data generated on edge devices such as cellphones or sensors offers an opportunity to train machine learning (ML) models for various applications. However, communication and privacy constraints of edge devices preclude the transfer of raw data to the cloud. Federated learning (FL) (McMahan et al., 2017; Kairouz et al., 2019; Li et al., 2020a; Yang et al., 2019) has emerged as a powerful framework to operate within these constraints by keeping decentralized data on the edge devices and instead moving model training to the edge. Federated model training operates in communication rounds. In each round, the current model is sent by the central server to edge clients, which perform model updates using their own local data, and the resulting models are then averaged by the central server. A typical cross-device FL framework consists of millions of intermittently connected edge clients, in each round only a small subset of them participate in training (Bonawitz et al., 2019). The subset of participating clients is affected by devices’ intrinsic properties such as battery status and network connectivity, and also system induced constraints for efficiency and privacy. In this paper, we seek understand the effect of such client participation patterns on convergence of federated training.

The federated averaging (FedAvg) algorithm and its variants are widely used in practice (Kairouz et al., 2019; Wang et al., 2021; Hard et al., 2018; Xu et al., 2023), and the convergence has been extensively analyzed in literature (Li et al., 2020b; Woodworth et al., 2020; Wang et al., 2022; Karimireddy et al., 2019; Wang & Joshi, 2021; Wang et al., 2020). However, most works assume uniform client participation which ensures that the model update applied to the global model is an unbiased estimate of the model update in the full client participation setting. This enables convergence results for the full-participation setting to be extended to the partial participation setting resulting in an additional variance term appearing in the convergence bound (Jhunjhunwala et al., 2022; Karimireddy et al., 2019; Wang et al., 2020). A generalization of the uniform client participation model is to consider that each client has an intrinsic availability probability p_i that is either known or unknown

to the central server. The set of participating clients is chosen according to this probability. Such non-uniform client participation introduces a bias in the model updates received by the server, with more frequently participating clients dominating the average update. To counter the bias, the central server can normalize the updates by the corresponding availability probabilities (Wang & Ji, 2022; Cho et al., 2022) or their estimates (Wang & Ji, 2023; Ríbero et al., 2022). We consider the setting of unknown client availability and analyze the convergence.

Both the uniform and non-uniform client participation models described above assume that client participation follows a Bernoulli process that is independent across clients and rounds. This assumption fails to capture practical settings where the client participation are correlated across rounds due to memory or time-dependence constraints. In cross-device FL systems, a device can only be available for training when it is plugged in for charging, connected to unmetered network and not being actively used by the owner (Hard et al., 2018; Paulik et al., 2021; Huba et al., 2022). These criterion, which typically occurs during the night of the devices' local time, not only results in the client availability probability for non-uniform client participation, but also correlated client participation of a periodic pattern due to user preference and time zone (Kairouz et al., 2019; Eichner et al., 2019; Zhu et al., 2021). More recently, a new criteria is introduced on devices in a FL system to impose a minimum separation constraint on successive participation instances of a client (McMahan & Thakurta, 2022; Xu et al., 2023). Specifically, once a client participates in training, it cannot become available to participate for at least R more rounds (R specified by the central aggregating server). The minimum separation is introduced to effectively combine differential privacy (DP) and FL (Kairouz et al., 2021; Choquette-Choo et al., 2023; McMahan et al., 2024) as advanced privacy-preserving methods, and quickly becomes the default criterion in many FL applications (Xu et al., 2023; Xu & Zhang, 2024). The client participation across rounds are correlated under the minimum separation criterion, and the extreme case of very large R will force cyclic client participation as studied in (Cho et al., 2023; Malinovsky et al., 2023). However, setting R to be the exact value for cyclic client participation can be challenging and may cause system slowdown, and these recent work did not study non-uniform client participation or the large spectrum of minimum separation R in practice. Other existing convergence analyses of federated training with generalized client participation (Wang & Ji, 2022; Rodio et al., 2023) do not fully explain the effect of such correlated client participation patterns, calling for new theoretical advances. See Appendix A for more related work discussions.

In this paper we bridge the gap of algorithms in practical FL system and the theoretical guarantees on their convergence with correlated client participation and unknown client availability. Our paper makes the following key contributions: (1) To the best of our knowledge we are the first to analyze the convergence of FedAvg with a minimum separation constraint on successive participation instances of each client, which is a general setting widely used in practical FL systems. We show that such correlated participation patterns can be captured by a Markov chain model. (2) We show that as the minimum separation R increases, the effective client participation probabilities become more uniform and reduces the asymptotic bias in the solution attained by the FedAvg algorithm. (3) We propose a debiased FedAvg algorithm that estimates the unknown client participation probabilities and incorporates them in the local updates. We prove that this algorithm achieves an unbiased solution that is consistent with the global FL objective under arbitrary minimum separation R .

Notations: For any positive integer N , we denote $[N] = \{1, \dots, N\}$. Let $\|\cdot\|$, $\|\cdot\|_1$ and $\|\cdot\|_\infty$ denote l_2 -norm, l_1 -norm and l_∞ -norm, respectively. For an ordered sequence $\{i_1, \dots, i_k\}$, it is represented by (i_1, \dots, i_k) and we use the same notation for a vector when the context causes no confusion. Unless otherwise specified, $\mathbb{E}(\cdot)$ means the total expectation taken on all randomness. We use \mathbf{c} to denote the vector where all entries are c . The d -dimensional Euclidean space is denoted by \mathbb{R}^d , and \mathbb{R}_+^d is the space formed vectors where every entry is strictly positive.

2 PROBLEM FORMULATION

We consider the federated learning setting where N clients cooperate to minimize the following global objective:

$$\min_x F(x) := \frac{1}{N} \sum_{i=1}^N f_i(x) \quad (1)$$

where f_i is the local objective function of client i . We aim to solve problem equation 1 in the federated learning setting, i.e., the system implements the some federated learning algorithm which

operates in rounds. In each round, a subset of the clients participate in training, and each of the clients in the subset performs multiple local updates based on the local gradients and then communicates with the server.

Non-uniform and correlated client participation. In this paper, we consider the scenario where each client requires some resting periods between participation and hence the participation pattern is correlated over time. Specifically, once participating in the system, a client has to wait as least R rounds until its next participation, where R is called *the minimum separation*. In other words, suppose client i 's last participation is in round t_i . It may join again at any round t with $t \geq t_i + 1 + R$ and not before then. Moreover, when a client is available to be sampled, instead of assuming uniform sampling, we consider that each client is associated with some *unknown* strictly positive scalar $p_i > 0$ to characterize its intrinsic willingness to be sampled at every round. Without loss of generality, we assume $\sum_{i=1}^N p_i = 1$ and hence refer to p_i as *the availability probability* of client i . Therefore, the client participation pattern is as follows: at each communication round, client i is sampled to participate in the training process with probability proportional to p_i if it has waited for R rounds after its last participation; otherwise client i cannot be sampled.

The above setting encompasses many of those in existing literature as special cases. For instance, note that $R = 0$ means each client is sampled at every round with probability p_i independently, which is consistent with (Wang & Ji, 2023). And the cyclic participation corresponds to the case $R = \frac{N}{B} - 1$ where B number of clients are sampled in each round (Cho et al., 2023), assuming the total number of clients in the FL population N is divisible by B . We investigate the potential bias introduced by the non-uniform and correlated client participation on federated algorithm performance and propose debiasing scheme to mitigate it.

3 MARKOV CHAIN MODEL AND ITS PROPERTIES

In this section, we propose a Markov chain model to capture the correlated participation scenario described above. Intuitively, the fact that every client cannot be sampled again within R rounds motivates us to maintain a memory window with length R to track which clients have not waited for R rounds. In other words, clients that are possible to be sampled in the current round only depend on which clients appearing in the memory window. This calls for a Markov chain with R -memory, also known as R -order Markov chain, defined as below.

Definition 1. Let $\{X_t\}_{t=0}^\infty$ be a stochastic process where $X_t \in \mathcal{X}, \forall t \geq 0$. It is said to be an R -order Markov chain if

$$P(X_t | X_{t-1}, X_{t-2}, \dots, X_0) = P(X_t | X_{t-1}, \dots, X_{t-R}), \forall t \geq R.$$

\mathcal{X} is called the state space.

If $R = 1$ it reduces to conventional Markov chain; if $R = 0$, then the clients can be sampled at each round with probability p_i , independent of the history. In a conventional Markov chain (with $R = 1$) with finite state space \mathcal{X} , we can use the transition probability matrix P to represent the Markov chain, where the (i, j) -th entry of P is $[P]_{i,j} = P(X_t = j | X_{t-1} = i)$, i.e., the probability of transitioning from state i to state j .

Recall that each client i is associated with a strictly positive availability probability $p_i > 0, \forall i \in [N]$. At each round t , the server samples a size- B subset of clients S_t , where $|S_t| = B$, with probability for each client proportional to p_i to join the training system. Note that only clients that have waited for R rounds are available. In other words, set S_t is sampled with probability proportional to $\sum_{i \in S_t} p_i$ from all subsets of size B formed by the available clients. We assume $N = MB$ for some $M > 0$ and note that the minimum separation R ranges from 0 to $M - 1$, where $R = M - 1$ corresponds to a cyclic participation pattern where subsets of clients participate in training in a fixed order.¹

Denote \mathcal{X} as the collection of all possible ordered subsets of $[N]$ with exactly B elements. Then, $|\mathcal{X}| = \sigma(N, B)$ where $\sigma(N, B) = \frac{N!}{(N-B)!}$ represents the total number of B -permutations of $[N]$.

¹Any $R > M - 1$ would result in periods with insufficient available clients. We do not consider those cases here.

Considering the stochastic process $\{X_t\}_{t=0}^\infty$ where $X_t \in \mathcal{X}$, the participation pattern in Section 2 can be precisely described by an R -order Markov chain defined in Definition 1. Formally,

$$P(X_t = \mathcal{I}_0 \mid X_{t-1} = \mathcal{I}_1, X_{t-2} = \mathcal{I}_2, \dots, X_0 = \mathcal{I}_t) = P(X_t = \mathcal{I}_0 \mid X_{t-1} = \mathcal{I}_1, \dots, X_{t-R} = \mathcal{I}_R) \quad (2)$$

where each state $\mathcal{I}_k \in \mathcal{X}$ represents which ordered subset of size B has been sampled at round k . For example, suppose clients 1 to B are sampled during the current round. $(1, 2, \dots, B)$ and $(2, 1, 3, \dots, B)$ are two different states, although the probability of these two states to appear is the same. The reason we consider this ordered case is that it allows us to cleanly define the probability of client i to be sampled (which is the marginal distribution of $P(X_t)$) by noting that $P(i \text{ to be sampled at round } t) = \sum_{i_2, \dots, i_B} P(X_t = (i, i_2, \dots, i_B))$. Here we calculate the probability of client i appearing as the first element in the ordered set X_t . The probability of i being sampled in any position would need an additional scaling factor of B . Since the scaling factor B is the same for all clients and only the relative frequency across clients contribute towards any bias effect, ignoring this factor of B would not affect the debiasing calculation.

The above high-order Markov chain equation 2 has some nice properties as summarized below (see Appendix C for proofs). The insights are important for deriving Theorem 2, which is the reason we formally present them here.

Proposition 1. *The R -th order Markov chain equation 2 maintains the following properties:*

- (1). *The ordered sequence $(\mathcal{I}_0, \mathcal{I}_1, \dots, \mathcal{I}_R)$ is non-repeated, meaning $\mathcal{I}_l \cap \mathcal{I}_k = \emptyset, \forall l \neq k$.*
- (2). *For any non-repeated $(\mathcal{I}_0, \dots, \mathcal{I}_R)$,*

$$P(X_t = \mathcal{I}_0 \mid X_{t-1} = \mathcal{I}_1, \dots, X_{t-R} = \mathcal{I}_R) = \frac{p_{\mathcal{I}_0}}{\sum_{\mathcal{J} \in S_{\mathcal{I}_{1:R}}^c} p_{\mathcal{J}}} =: p_{(\mathcal{I}_1, \dots, \mathcal{I}_R) \rightarrow \mathcal{I}_0}. \quad (3)$$

Otherwise $P(X_t = \mathcal{I}_0 \mid X_{t-1} = \mathcal{I}_1, \dots, X_{t-R} = \mathcal{I}_R) = 0$. Since \mathcal{I}_k is a set with B unique elements, we define $p_{\mathcal{I}_k} := \prod_{e \in \mathcal{I}_k} p_e, \forall \mathcal{I}_k$. $S_{\mathcal{I}_{1:R}}^c$ is the collection containing all B -permutations of $[N] \setminus \bigcup_{k=1}^R \mathcal{I}_k$.

(3). For $t \geq R-1$, define $Y_t = (X_t, \dots, X_{t-R+1}) \in \mathbb{R}^R$. Then $\{Y_t\}_{t=R-1}^\infty$ is a conventional Markov chain with its cardinality of the state space being $d(M, R)$, where $d(M, R) = \prod_{k=0}^{R-1} \sigma(B(M-k), B)$. Moreover its transition probability is

$$P(Y_t = (\mathcal{I}_0, \mathcal{J}_1, \dots, \mathcal{J}_{R-1}) \mid Y_{t-1} = (\mathcal{I}_1, \dots, \mathcal{I}_R)) = \begin{cases} p_{(\mathcal{I}_1, \dots, \mathcal{I}_R) \rightarrow \mathcal{I}_0} & , \quad \mathcal{J}_k = \mathcal{I}_k, k \in [R-1] \\ 0 & , \quad \text{otherwise} \end{cases} \quad (4)$$

for any non-repeated $(\mathcal{I}_0, \dots, \mathcal{I}_R)$.

(4). Define vector $u_{(\mathcal{I}_1, \dots, \mathcal{I}_R)} \in \mathbb{R}^{d(M, R)}$ with $(\mathcal{I}_0, \mathcal{I}_1, \dots, \mathcal{I}_{R-1})$ -th entry as $P(Y_t = (\mathcal{I}_0, \mathcal{I}_1, \dots, \mathcal{I}_{R-1}) \mid Y_{t-1} = (\mathcal{I}_1, \dots, \mathcal{I}_R))$. Then, $u_{(\mathcal{I}_1, \dots, \mathcal{I}_R)} \in \mathbb{R}_+^{\sigma(B(M-R), B)} \subset \mathbb{R}^{d(M, R)}$ and $u_{(\mathcal{I}_1, \dots, \mathcal{I}_R)}[(\mathcal{I}_0, \dots, \mathcal{I}_{R-1})] = p_{\mathcal{I}_0}(\sum_{\mathcal{J} \in S_{\mathcal{I}_{1:R}}^c} p_{\mathcal{J}})^{-1} > 0, \forall \mathcal{I}_0 \in S_{\mathcal{I}_{1:R}}^c$.

(5). Denote $v_{(\mathcal{J}_0, \dots, \mathcal{J}_{R-1})} \in \mathbb{R}^{d(M, R)}$ with $(\mathcal{J}_1, \mathcal{J}_2, \dots, \mathcal{J}_R)$ -th entry as $P(Y_t = (\mathcal{J}_0, \dots, \mathcal{J}_{R-1}) \mid Y_{t-1} = (\mathcal{J}_1, \mathcal{J}_2, \dots, \mathcal{J}_R))$. Then, $v_{(\mathcal{J}_1, \dots, \mathcal{J}_R)} \in \mathbb{R}_+^{\sigma(B(M-R), B)}$ and $v_{(\mathcal{J}_0, \dots, \mathcal{J}_{R-1})}[(\mathcal{J}_1, \dots, \mathcal{J}_R)] = p_{\mathcal{J}_0}(\sum_{\mathcal{J} \in S_{\mathcal{J}_{1:R}}^c} p_{\mathcal{J}})^{-1} > 0$ for any $\mathcal{J}_R \in S_{\mathcal{J}_{0:R-1}}^c$.

Properties (1),(2) essentially state that clients to be sampled in the current round cannot be those who have not waited for R rounds, which establish the equivalence of our Markov-chain modeling equation 2 and the participation pattern in Section 2. Property (3) means that we can augment our state space by considering R -length history to formulate an equivalent Markov chain $\{Y_t\}_{t=R}^\infty$ with order 1. The last two properties explicitly shows what entries are for each row and column of the transition probability matrix of the new Markov chain $\{Y_t\}_{t=R}^\infty$. Also since there are only $\sigma(B(M-R), B) \ll d(M, R)$ non-zero entries in every row and column, the transition matrix is sparse.

A main benefit of this Markov-chain modeling is allowing us to look into the probability of each client to be sampled as t goes on. Specifically, given any R , denote $P_R \in \mathbb{R}^{d(M, R) \times d(M, R)}$ as the

transition probability matrix of the Markov chain $\{Y_t\}_{t=R}^\infty$ where its entry is given by equation 4. Let $\phi_R(t) \in \mathbb{R}^{d(M,R)}$ be the state distribution at round t of the Markov chain $\{Y_t\}_{t=R}^\infty$ and $\eta_R(t) \in \mathbb{R}^N$ be the distribution of clients to be sampled at round t . We have the following evolution of distributions with respect to t :

$$\eta_R(t) = Q_R^T \phi_R(t), \quad \phi_R(t+1) = P_R^T \phi_R(t) \quad (5)$$

for any initial distribution $\eta_R(0)$ and corresponding $\phi_R(0)$ such that $\eta_R(0) = Q_R^T \phi_R(0)$, where $Q_R = Q_{R,1} Q_{R,2}$ and $Q_{R,1} \in \mathbb{R}^{d(M,R) \times \sigma(N,B)}$ is defined by

$$[Q_{R,1}]_{(\mathcal{I}_1, \dots, \mathcal{I}_R), \mathcal{J}} = \begin{cases} p_{(\mathcal{I}_1, \dots, \mathcal{I}_R) \rightarrow \mathcal{J}} & , \quad \{\mathcal{J}, \mathcal{I}_1, \dots, \mathcal{I}_R\} \text{ non-repeated} \\ 0 & , \quad \text{otherwise.} \end{cases}$$

and $Q_{R,2} \in \mathbb{R}^{\sigma(N,B) \times N}$ is defined by

$$[Q_{R,2}]_{\mathcal{J}, j} = \begin{cases} 1 & , \quad \mathcal{J} = (j, *) \\ 0 & , \quad \text{otherwise,} \end{cases}$$

where $\mathcal{J} = (j, *)$ denotes that the first entry of \mathcal{I} is j . We are particularly interested in the distribution of $\eta_R(t)$ as $t \rightarrow \infty$ because it helps us characterize the asymptotic performance of existing FL algorithms. From classical Markov chain literature, we know that if a Markov chain is irreducible and aperiodic (see formal definitions in Appendix B), it has a stationary distribution which is unique and strictly positive. We denote $\zeta_R = \lim_{t \rightarrow \infty} \phi_R(t)$ as the stationary distribution of Markov chain P_R and we have

$$\zeta_R^T = \zeta_R^T P_R, \quad \pi_R^T = \zeta_R^T Q_R. \quad (6)$$

where $\pi_R \in \mathbb{R}^N$ is marginal stationary distribution of clients to be sampled, i.e., the i -th entry of π_R is given by $\pi_R^i = \lim_{t \rightarrow \infty} \eta_R^i(t)$. On the other hand, if the Markov chain is irreducible and peroidic, we let ζ_R be the Perron vector², which is also strictly positive. We now show our Markov chain is irreducible and (a)periodic to justify the definitions of ζ_R and π_R in Lemma 1. The proof is in Appendix C.

Lemma 1. *The Markov chain $\{Y_t\}_{t=R}^\infty$ with transition matrix P_R defined by equation 4 is irreducible for all $M \geq 1$ and $0 \leq R \leq M-1$. Further, when $R \leq M-2$, it is also aperiodic.*

We provide an example to illustrate the intuition of our Markov-chain model above, considering the case of $N = 4, B = 1, R = 2$, i.e., every round one client is sampled, then it has to wait for two rounds. For instance, if client 1 and client 2 are consecutively selected in the first two rounds, in the third round only client 3 or 4 can be selected with probabilities of $p_3/(p_3 + p_4)$ or $p_4/(p_3 + p_4)$ respectively. Then, the state $(2, 1)$ can only transition to $(3, 2)$ or $(4, 2)$, where the second index is sampled before the first one as is in equation 2. Similarly, if we are currently at state $(1, 4)$, the previous state has to be $(4, 3)$ or $(4, 2)$. One can easily check that Proposition 1 holds. To see how π is calculated, we take the first entry of π_R as an example:

$$\pi_R^1 = \zeta^{(2,3)} p_{(2,3) \rightarrow 1} + \zeta^{(2,4)} p_{(2,4) \rightarrow 1} + \zeta^{(3,2)} p_{(3,2) \rightarrow 1} + \zeta^{(3,4)} p_{(3,4) \rightarrow 1} + \zeta^{(4,2)} p_{(4,2) \rightarrow 1} + \zeta^{(4,3)} p_{(4,3) \rightarrow 1}$$

by noting that the remaining $p_{(i,j) \rightarrow 1} = 0$, if i or $j = 1$.

The vectors in equation 6 characterize the final distribution according to which clients will be sampled when the communication round t becomes infinitely large. In other words, each client i is sampled with probability π_R^i given some fixed R . Although π_{M-1} is the uniform distribution no matter what p_i 's are (by observing that all clients follow a cyclic participation), we note that π_R for $R < M-1$ does not necessarily follow the uniform distribution, because $\{p_1, \dots, p_N\}$ are arbitrary. This will be problematic in the sense that existing federated learning algorithms may no longer guarantee convergence to the correct and optimal solution of equation 1 no matter how many rounds of training are implemented. We call this phenomenon *the asymptotic bias induced by π_R* . We will characterize both empirically and theoretically this phenomenon in the next section.

²We say v is the Perron vector of the transition matrix P if $v^T = v^T P$, i.e., v is right eigenvector of P corresponding to eigenvalue 1 and $v^T \mathbf{1} = 1$.

4 ASYMPTOTIC BIAS UNDER NON-UNIFORM CORRELATED PARTICIPATION

In this section, we utilize the Markov chain model in the previous section to analyze asymptotic bias of existing federated learning algorithms caused by $R < M - 2$ and arbitrary p_i 's. In particular, we consider FedAvg with local gradient descent updates, i.e., at each round, a set S_t with $|S_t| = B$ clients are sampled and after being selected client i updates its model as

$$x_{t,0}^i = x_t, \quad x_{t,k+1}^i = x_{t,k}^i - \alpha \nabla f_i(x_{t,k}^i), \quad k = 0, \dots, K-1 \quad (7)$$

where x_t denotes the server's model at round t and $x_{t,k}^i$ is the local model maintained by client i at k -th iteration. The server then updates $x_{t+1} = \frac{1}{B} \sum_{i \in S_t} x_{t,K}^i$. We next show in the following that FedAvg may not converge to the desired optimal solutions of equation 1. Instead there may exist some error neighborhood, i.e., the asymptotic bias, that is related to π_R , even as t goes to infinity. Before we formally deliver the result, two standard assumptions are needed.

Assumption 1. *There exists $G > 0$ such that $\|\nabla f_i(x) - \nabla F(x)\|^2 \leq G^2, \forall x$ and $\forall i \in [N]$.*

Assumption 2. *Each f_i is L -smooth, i.e., $\|\nabla f_i(x) - \nabla f_i(y)\| \leq L\|x - y\|, \forall x, y$ and $\forall i \in [N]$.*

Then we are ready to state the convergence of FedAvg under correlated client participation (see Appendix F for the proof).

Theorem 1. *Suppose Assumptions 1,2 hold and assume $\|\nabla F(x)\| \leq D, \forall x$ with some finite $D > 0$. Then for any $T > 2\tau_{mix} \log \tau_{mix}$ choosing $\alpha = \tilde{\mathcal{O}}(1/(\tau_{mix} K \sqrt{T}))$, FedAvg with local updates equation 7 generates the trajectory $\{x_t\}_{t=0}^{T-1}$ satisfying*

$$\mathbb{E} \|\nabla F(\tilde{x}_T)\|^2 \leq \tilde{\mathcal{O}}\left(\frac{\tau_{mix}}{\sqrt{T}}\right) + \mathcal{O}\left(\frac{1}{T}\right) + \mathcal{O}\left(\|\pi_R - \frac{1}{N} \mathbf{1}_N\|_1^2\right), \quad (8)$$

for any $0 \leq R < M - 1$, where \tilde{x}_T is drawn uniformly from x_0, \dots, x_{T-1} , $\tilde{\mathcal{O}}(\cdot)$ hides logarithmic factors, and τ_{mix} denotes the mixing time³ of Markov chain equation 5. Moreover, the bias term $\mathcal{O}(\|\pi_R - \frac{1}{N} \mathbf{1}_N\|_1^2)$ shown in equation 8 is unavoidable.

Theorem 1 implies that without any debiasing technique, FedAvg can only converge to a solution with unavoidable asymptotic bias which is measured by the distance between π_R (defined in equation 6) and the uniform distribution. Except for $R = M - 1$, where π_{M-1} is the uniform distribution, for $R \leq M - 2$, there is generally some gap between π_R and $(1/N) \mathbf{1}_N$, which shows that FedAvg may fail to perform under correlated client participation. However, if π_R is not too far away from the uniform distribution, we expect FedAvg to converge to a solution reasonably close to the optimal solution of equation 1. We next investigate what factors influence the distance from π_R to the uniform distribution. We find that one factor is the spread among p_i 's. Stated by the following proposition, if all p_i 's are equal, no gap between π_R and $(1/N) \mathbf{1}_N$ exists (see Appendix D for the proof).

Proposition 2. *Suppose $p_1 = p_2 = \dots = p_N = \frac{1}{N}$. Then for any $0 \leq R \leq M - 1$, $\pi_R = \frac{1}{N} \mathbf{1}_N$.*

When p_i 's are not equal to each other, we turn to understand how R affect π_R . In fact, we empirically observe that π_R approaches the uniform distribution as R increases. This key observation is illustrated in Figure 1. We consider the case where $N = 500, B = 1$ and assign each client a random $p_i > 0$. We then calculate π_R for each R ranging from 0 to $N - 1$ and measure its distance from the uniform distribution. As shown in the figure, increasing R causes π_R moving towards the uniform distribution. One explanation for this observation is that when R becomes larger, fewer clients are ready to be sampled in the current round, because many clients have not waited for enough rounds and hence are not available. Rather than dictated by the availability probability p_i 's, which is the case for a small R and many available clients, here the sampling process is mostly determined by the waiting requirement. In the extreme case, when $R = M - 1$, at each round, only B clients are available, hence all clients are sampled with equal frequency. Another point suggested by this observation is that we can choose a large minimum separation R in the practical scenario to reduce the asymptotic bias for existing FL algorithms, even with unknown p_i 's.

³Please refer to Appendix B for the formal definition of the mixing time.

Algorithm 1 Debiasing FedAvg for correlated client participation

```

1: Input: initial point  $x_0$ , stepsizes  $\{\alpha\}$ , some  $\tau > 0$ ,  $\lambda_0 = \mathbf{0}_N$ ,  $t_i = 0, \forall i \in [N]$  for each client
2: for  $t = 0, 1, \dots, T$  do
3:   A batch of clients  $S_t$  with size  $|S_t| = B$  is selected. The server sends current  $t$  and model  $x_t$  to clients in  $S_t$ .
4:   for  $i \in S_t$  in parallel do
5:     Each client sets  $t_i \leftarrow t_i + 1$  and calculates  $\lambda_t^i = \frac{t_i}{(t+1)B}$  and  $\nu_t^i = \frac{1}{\lambda_t^i N}$ .
6:     for  $k = 0, 1, \dots, K - 1$  do
7:       Client  $i$  updates its local model by
           
$$x_{t,k+1}^i = x_{t,k}^i - \alpha \nu_t^i \nabla f_i(x_{t,k}^i). \quad (9)$$

8:     end for
9:   end for
10:  The server updates its model  $x_{t+1} = \frac{1}{B} \sum_{i \in S_t} x_{t,K}^i$ .
11: end for
12: Output:  $\tilde{x}_T$  sampled uniformly from  $\{x_t\}_{t=0}^{T-1}$ 

```

The above empirical observation verifies the formal theorem that characterizes the debiasing effect of increasing minimum separation R in Theorem 2. (see Appendix D for the proof).

Theorem 2. *Given a set of p_i 's, with at least one element $p_i \neq \frac{1}{N}$. Without loss of generality, let p_1, \dots, p_B be the B smallest values among all p_i 's. Define $q_B := \sum_{j=1}^B p_j$, then $q_B < 1/M$. There exists a $\bar{\delta} > 0$, such that if any size- B batch of clients \mathcal{B}_j picking from $[N] \setminus [B]$, $\delta_j := \left| \sum_{i \in \mathcal{B}_j} p_i - \frac{1-q_B}{M-1} \right| \leq \bar{\delta}$, then π_R converges to a neighborhood of $\frac{1}{N} \mathbf{1}_N$ characterized by $\{\pi \mid \|\pi - \frac{1}{N} \mathbf{1}_N\|_1 = \mathcal{O}(N^{-1})\}$ as R ranging from 0 to $M-1$. When $R = M-1$, π_{M-1} is the uniform distribution supported on $[N]$.*

Theorem 2 states that when the availability probabilities p_i 's of clients are not too far away from each other or when B is relatively large (i.e., δ_j 's are small for all j), and when the total number of clients N is large, π_R approaches the uniform distribution as R increases. It is worth noting that practically when the requirements in Theorem 2 are not strictly satisfied, the effect of increasing R on π_R can be still observed as shown in Figure 1.

5 DEBIASING FEDAVG AND ITS CONVERGENCE

As we discuss in the previous section, existing federated learning algorithms like FedAvg cannot guarantee convergence to the correct optimal solution if $R \leq M-2$ and p_i 's are arbitrary. Although we can reduce the asymptotic bias caused by π_R by increasing R , it may still be problematic under some particular circumstances. Clients have intermittent and non-uniform availability, and forcing a large minimum separation R in practice may cause significant slowdown of the training in the FL system due to the small number of available clients. The minimum separation R can be relatively small and the p_i 's can be very different from each other, which then suggests by Figure 1 and Theorem 2, π_R can be far from the uniform distribution, making the asymptotic bias non-negligible. We next design a debiasing process that can be easily integrated into the existing federated learning algorithms to address asymptotic bias. Our proposed algorithm based on FedAvg is given by Algorithm 1.

The main difference between our algorithm and vanilla FedAvg lies in the stage of local updates (Lines 5 and 7). Specifically, we require each client to maintain an estimator of its corresponding




Figure 1: $\|\pi_R - \mathbf{1}_N/N\|_1$ as R increases ($N = 500, B = 1$)

component of π_R , which is only updated when the client is sampled. This estimator is later used to scale the gradient step during the local update. The estimator is designed by counting the times the client has been sampled and then used to compute the running empirical frequency of the client’s participation. Recall that π_R^i represents the frequency of client i to be selected when t is large enough (i.e., when the Markov chain equation 5 becomes steady, meaning $\phi_R(\infty) = \zeta_R$). If we reweigh the local objective function f_i by $\frac{1}{\pi_R^i}$ (corresponding to $\nu_t^i = \frac{1}{\pi_R^i N}$ in equation 9), this weighting cancels the asymptotic bias introduced by unbalanced sampling, which drives the trajectory of the server’s models towards the correct solution of equation 1. If we know π_R^i for every client in prior, the above-mentioned reweighting method provides us with unbiased solutions. Then, λ_t^i serves as a role to iteratively approximate π_R^i round by round, which yields Algorithm 1. Also note that Algorithm 1 reduces to FedAvg if fixing $\lambda_t^i = 1/N, \forall i \in [N]$. This shows the advantage of our algorithm: it is computationally cheap in the sense that each client only maintains two additional scalars (λ_t^i and ν_t^i) and can be easily embedded with existing algorithms by just multiplying the learning rates by ν_t^i . We note that other federated algorithms suffering from asymptotic bias due to non-uniform sampling could also benefit from our debiasing technique based on simple counting.

However, formally characterizing the convergence of ν_t^i to $\frac{1}{\pi_R^i N}$ remains challenging due to the samples of clients are not independent across different rounds. In particular, the clients sampled in the current round may affect those in the future, which makes the conventional concentration tools and law of large numbers not applicable. To address this challenge, we carefully analyze the transition of the Markov chain equation 5 and its influences on the marginal distribution of clients to be sampled to conclude that λ_t^i is an unbiased estimate of π_R^i asymptotically. Then, we further leverage the fact that the Markov chain is irreducible as stated in Lemma 1 to show that λ_t^i is almost surely strictly positive even t is infinite, concluding the convergence of ν_t^i to $\frac{1}{\pi_R^i N}$, as summarized in Lemma 2 (see Corollary 2 in Appendix G for the proof).

Lemma 2. *Given $\lambda_0 = \mathbf{0}_N$, then $\nu_t^i, \forall i \in [N]$ in Algorithm 1 satisfies*

$$\mathbb{E}\|\tilde{\nu}_t\|_\infty^2 \leq \mathcal{O}\left(\frac{\tau_{mix}}{t}\right)$$

for any $t > 0$, where $\tilde{\nu}_t^i = \nu_t^i - \frac{1}{\pi_R^i N}$ and $\tilde{\nu}_t = (\tilde{\nu}_t^1, \dots, \tilde{\nu}_t^N)$; τ_{mix} is the mixing time of Markov chain equation 5.

Based on the above, we can achieve the following convergence result of Algorithm 1 (see Appendix G for the proof).

Theorem 3. *Suppose Assumptions 1 and 2 hold. For any $0 \leq R < M - 1$ and $T > c^\dagger \tau_{mix} \log \tau_{mix}$ (with c^\dagger being some constant), choosing $\alpha = \tilde{\mathcal{O}}(1/(\tau_{mix} K \sqrt{T}))$, the output of Algorithm 1 satisfies*

$$\mathbb{E}\|\nabla F(\tilde{x}_T)\|^2 = \tilde{\mathcal{O}}\left(\frac{\tau_{mix}}{\sqrt{T}}\right) + \mathcal{O}\left(\frac{1}{T}\right)$$

where \tilde{x}_T is defined as that in Theorem 1; τ_{mix} is the mixing time of Markov chain equation 5.

Comparing to Theorem 1, no bounded gradient assumption is needed to reach the convergence of our algorithm. Unlike the result in (Cho et al., 2023) where clients are forced to participate in the system cyclically, our bound shown in Theorem 3 does not grow as the number of clients increases. Particularly, for the bounds in (Cho et al., 2023) to be non-vacuous, the total number of communication round T should be proportional to the number of clients, which could be hard to satisfy in practice especially when client number is super large. To prove Theorem 3 we critically rely on the fact that the Markov chain equation 5 is aperiodic to make analysis go through. That is to say our bound does not suit for $R = M - 1$, which is the limitation of our analysis. However, since $R = M - 1$ is the cyclic case, where the Markov chain follows much nicer structure, one may be able to get a better bound (Cho et al., 2023).

We remark that our convergence result achieves nearly the same order of rate as Markov-sampling SGD literature (Beznosikov et al., 2024; Even, 2023) (where rates of $\mathcal{O}(\sqrt{\tau_{mix}}/\sqrt{T}) + \tau_{mix}/T$) are obtained). However, their analysis only suits for the first-order Markov chain and no debiasing results are presented, while our results generalize to high-order Markov chain and guarantee approaching unbiased solutions. We note that utilizing variance-reduced techniques may accelerate the

convergence rate for Markov-sampling SGD (Even, 2023). Then whether variance reduction can be used in our problem to design faster algorithms would be an interesting future direction.

It is worth noting that although a uniform minimum separation R for all clients is placed throughout the paper, we also allow each client maintains its own specific $R_i, \forall i \in [N]$. In this more general case, we could still use the same modeling technique as in Section 3 where the order of the Markov chain is chosen to be an upper bound of all R_i 's (e.g. $\max_i R_i$). Then Theorems 1 and 3 can be obtained without any modification as the analysis stays valid for any irreducible and aperiodic Markov chain. However, Theorem 2 becomes tricky in this case as our proof highly relies on nice properties of the Markov chain summarized by Proposition 1 which now cease to hold. Therefore, more advanced mathematical tools might be needed in order to obtain similar statements as Theorem 2 when clients have various R_i 's.

6 NUMERICAL RESULTS

In this section, we provide numerical experiments to illustrate our theoretical results. In particular, we compare vanilla FedAvg with our proposed algorithm (Algorithm 1) under non-uniform and correlated client participation described in Section 2. For simplicity, we partition the N clients into M groups and exactly one group of clients are selected at each round to fully participate in the system. Here we choose $N = 100, M = 20$. Since all clients in the same group participate in the system together once being sampled, we only need to associate availability probabilities to each group, where $p_i \propto i^{-1.5}, i \in [M]$ is a long-tailed distribution.

Synthetic dataset. We test Vanilla FedAvg and Debiasing FedAvg (Algorithm 1) under a synthetic dataset constructed following (Sun & Wei, 2022): for each client i , $A_i \in \mathbb{R}^{n_i \times d}$ is the feature matrix, where n_i is the number of local samples and d is the feature dimension. Every entry of A_i is generated by a Gaussian distribution $\mathcal{N}(0, (0.5i)^{-2})$. We then generate $b_i \in \mathbb{R}^{n_i}$, the labels of client i , by first generating a reference point $\theta_i \in \mathbb{R}^d$, where $\theta_i \sim \mathcal{N}(\mu_i, I_d)$. And μ_i is drawn from $\mathcal{N}(\alpha, 1)$ with $\alpha \sim \mathcal{N}(0, 100)$. Then $b_i = A_i \theta_i + \epsilon_i$ with $\epsilon_i \sim \mathcal{N}(0, 0.25I_{n_i})$. We set $d = 20, n_i = 100, \forall i \in [N]$. And we define $f_i(x) = \frac{1}{n_i} \sum_{j=1}^{n_i} \log(\frac{1}{2}(\langle A_i[j, :], x \rangle + b_i[j])^2 + 1)$ where $A_i[j, :]$ represents the j -th row of A_i and $b_i[j]$ is the j -th entry of b_i . The outcomes are shown in Figures 2a,2b, where Figure 2a shows that Vanilla FedAvg suffers from bias which can be mitigated by increasing R , and Figure 2b shows that Debiasing FedAvg effectively reduces bias no matter what value of R is set.

MNIST dataset. We also test our proposed algorithm under the MNIST dataset. Each client maintains a three-layer fully-connected neural network for training. All learning rates are chosen to be with the order of $\mathcal{O}(10^{-3})$. In Figure 3c, we compare Debiasing FedAvg with Vanilla FedAvg and FedVARP (Jhunjhunwala et al., 2022), and Debiasing FedAvg can effectively mitigate the bias effect. Another interesting empirical observation is that increasing R can possibly fasten the speed of both Debiasing and Vanilla FedAvg (as shown by Figures 3a,3b). This is yet not characterized by our theoretical demonstration. Here we conjecture that larger R corresponds to smaller mixing time τ_{mix} and hence faster rate as the bounds in Theorems 1,3 scale with respect to τ_{mix} . We provide more detailed and intuitive discussions in Appendix I.

7 LIMITATIONS

Our Markov-chain framework works for that all clients share the same static minimum separation R and in the last paragraph of Section 5 we further allow static client-specific R_i 's. However, more practical scenarios call for even time-varying R , which lies outside the scope of this paper. In Theorem 2, we force the availability probabilities p_i 's not too far away from each other to make the theory hold, while this assumption is not required in practice. Moreover, Theorems 1,3 do not enjoy speedup in the number of clients as FL literature when clients are uniformly sampled. We believe it is mainly due to the non-uniformity and time-correlation of the client sampling process and more advanced mathematical tools are needed to show speedup in the Markov setting.

Many of the challenges in cross-device federated learning can also be addressed by system design in addition to algorithm design. For example, incorporating trusted execution environment (TEEs) in




Figure 2: Experiments on synthetic dataset. (a) The training loss of Vanilla FedAvg (after convergence) with different R is shown. Larger R leads to smaller bias. (b) Debiasing FedAvg is tested under different values of R , where the red line represents Vanilla FedAvg when clients are sampled under an oracle uniform distribution. The subfigure on the right shows that all curves reach unbiased objective after convergence, indicating that the asymptotic bias is effectively canceled.




Figure 3: Experiments on MNIST. (a) The convergence of our Debiasing FedAvg under different client minimum separation R configurations. The red horizontal line is the convergence value of the objective function by vanilla FedAvg when clients are sampled under an oracle uniform distribution. Our Debiasing FedAvg converges to the unbiased objective with larger R converges faster. (b) For Vanilla FedAvg, increasing R causes smaller bias. (c) When $R = 8$, Vanilla FedAvg, FedVARP and Debiasing FedAvg are compared. Note that both Vanilla FedAvg and FedVARP are designed only for uniform client sampling and hence are significantly affected by bias from client participation.

the FL system (Huba et al., 2022; Daly et al., 2024) can potentially provide more control on client sampling.

8 CONCLUSION

In this paper, we consider FL with non-uniform and correlated client participation, where every client must wait as least R rounds (minimum separation) before participating again, and each client has their own availability probability. A high-order Markov chain is introduced to model this practical scenario. Based on this Markov-chain modeling, we are able to study the convergence performances of existing FL algorithms. Due to the effect of non-uniformity and time correlation, FL algorithms can only converge with asymptotic bias, which can be reduced by increasing minimum separation R as shown by our empirical and theoretical results. Finally, we propose a debiasing algorithm for FedAvg that guarantee convergence to unbiased solutions given arbitrary non-uniformity and minimum separation R .

ACKNOWLEDGMENTS

This work was supported in part by NSF grants CCF-2045694, ECCS-2216970, CMMI-2024774, CNS-2112471, CPS-2111751, ONR N00014-23-1-2149, and a Google Research Scholar Award to GJ.

REFERENCES

Zeyuan Allen-Zhu. Katyusha: The first direct acceleration of stochastic gradient methods. In *Proceedings of Symposium on Theory of Computing (STOC)*, pp. 1200–1205, 2017.

Zeyuan Allen-Zhu and Elad Hazan. Variance reduction for faster non-convex optimization. In *International conference on machine learning*, pp. 699–707. PMLR, 2016.

Aleksandr Beznosikov, Sergey Samsonov, Marina Sheshukova, Alexander Gasnikov, Alexey Naumov, and Eric Moulines. First order methods with markovian noise: from acceleration to variational inequalities. *Advances in Neural Information Processing Systems*, 36, 2024.

Shalabh Bhatnagar, Mohammad Ghavamzadeh, Mark Lee, and Richard S Sutton. Incremental natural actor-critic algorithms. *Advances in neural information processing systems*, 20, 2007.

Jieming Bian, Lei Wang, Kun Yang, Cong Shen, and Jie Xu. Accelerating hybrid federated learning convergence under partial participation. *IEEE Transactions on Signal Processing*, 2024.

Keith Bonawitz, Hubert Eichner, Wolfgang Grieskamp, Dzmitry Huba, Alex Ingerman, Vladimir Ivanov, Chloe Kiddon, Jakub Konecny, Stefano Mazzocchi, H. Brendan McMahan, Timon Van Overveldt, David Petrou, Daniel Ramage, and Jason Roselander. Towards Federated Learning at Scale: System Design. *SysML*, April 2019. URL <https://www.sysml.cc/doc/2019/193.pdf>.

Wenlin Chen, Samuel Horvath, and Peter Richtarik. Optimal client sampling for federated learning. *arXiv preprint arXiv:2010.13723*, 2020.

Yae Jee Cho, Jianyu Wang, and Gauri Joshi. Client selection in federated learning: Convergence analysis and power-of-choice selection strategies. In *Proceedings of the International Conference on Artificial Intelligence and Statistics (AISTATS)*, 2022. URL "<https://arxiv.org/abs/2010.01243>".

Yae Jee Cho, Pranay Sharma, Gauri Joshi, Zheng Xu, Satyen Kale, and Tong Zhang. On the convergence of federated averaging with cyclic client participation. In *International Conference on Machine Learning*, pp. 5677–5721. PMLR, 2023.

Christopher A Choquette-Choo, Arun Ganesh, Ryan McKenna, H Brendan McMahan, John Rush, Abhradeep Guha Thakurta, and Zheng Xu. (amplified) banded matrix factorization: A unified approach to private training. *Advances in Neural Information Processing Systems*, 36, 2023.

Katharine Daly, Hubert Eichner, Peter Kairouz, H Brendan McMahan, Daniel Ramage, and Zheng Xu. Federated learning in practice: reflections and projections. In *2024 IEEE 6th International Conference on Trust, Privacy and Security in Intelligent Systems, and Applications (TPS-ISA)*, pp. 148–156. IEEE, 2024.

Aaron Defazio, Francis Bach, and Simon Lacoste-Julien. SAGA: A fast incremental gradient method with support for non-strongly convex composite objectives. In *Advances in Neural Information Processing Systems 27*, pp. 1646–1654, 2014.

Hubert Eichner, Tomer Koren, Brendan McMahan, Nathan Srebro, and Kunal Talwar. Semi-cyclic stochastic gradient descent. In *International Conference on Machine Learning*, pp. 1764–1773. PMLR, 2019.

Mathieu Even. Stochastic gradient descent under markovian sampling schemes. In *International Conference on Machine Learning*, pp. 9412–9439. PMLR, 2023.

Yann Fraboni, Richard Vidal, Laetitia Kameni, and Marco Lorenzi. Clustered sampling: Low-variance and improved representativity for clients selection in federated learning. In *International Conference on Machine Learning*, pp. 3407–3416. PMLR, 2021.

Andrew Hard, Kanishka Rao, Rajiv Mathews, Swaroop Ramaswamy, Françoise Beaufays, Sean Augenstein, Hubert Eichner, Chloé Kiddon, and Daniel Ramage. Federated learning for mobile keyboard prediction. *arXiv preprint arXiv:1811.03604*, 2018.

Dzmitry Huba, John Nguyen, Kshitiz Malik, Ruiyu Zhu, Mike Rabbat, Ashkan Yousefpour, Carole-Jean Wu, Hongyuan Zhan, Pavel Ustinov, Harish Srinivas, et al. Papaya: Practical, private, and scalable federated learning. *Proceedings of Machine Learning and Systems*, 4:814–832, 2022.

Divyansh Jhunjhunwala, Pranay Sharma, Aushim Nagarkatti, and Gauri Joshi. FedVARP: Tackling the variance due to partial client participation in federated learning. In *Proceedings of the Conference on Uncertainty in Artificial Intelligence (UAI)*, aug 2022. URL <https://arxiv.org/pdf/2207.14130.pdf>.

Rie Johnson and Tong Zhang. Accelerating stochastic gradient descent using predictive variance reduction. *Advances in neural information processing systems*, 26, 2013.

Peter Kairouz, H. Brendan McMahan, Brendan Avent, Aurelien Bellet, Mehdi Bennis, Arjun Nitin Bhagoji, Keith Bonawitz, Zachary Charles, Graham Cormode, Rachel Cummings, Rafael G. L. D’Oliveira, Salim El Rouayheb, David Evans, Josh Gardner, Zachary Garrett, Adria Gascon, Badih Ghazi, Phillip B. Gibbons, Marco Gruteser, Zaid Harchaoui, Chaoyang He, Lie He, Zhouyuan Huo, Ben Hutchinson, Justin Hsu, Martin Jaggi, Tara Javidi, Gauri Joshi, Mikhail Khodak, Jakub Konecny, Aleksandra Korolova, Farinaz Koushanfar, Sanmi Koyejo, Tancrede Lepoint, Yang Liu, Prateek Mittal, Mehryar Mohri, Richard Nock, Ayfer Ozgur, Rasmus Pagh, Mariana Raykova, Hang Qi, Daniel Ramage, Ramesh Raskar, Dawn Song, Weikang Song, Sebastian U. Stich, Ziteng Sun, Ananda Theertha Suresh, Florian Tramer, Praneeth Vepakomma, Jianyu Wang, Li Xiong, Zheng Xu, Qiang Yang, Felix X. Yu, Han Yu, and Sen Zhao. Advances and open problems in federated learning. *arXiv preprint arXiv:1912.04977*, 2019.

Peter Kairouz, Brendan McMahan, Shuang Song, Om Thakkar, Abhradeep Thakurta, and Zheng Xu. Practical and private (deep) learning without sampling or shuffling. In *International Conference on Machine Learning*, pp. 5213–5225. PMLR, 2021.

Sai Praneeth Karimireddy, Satyen Kale, Mehryar Mohri, Sashank J Reddi, Sebastian U Stich, and Ananda Theertha Suresh. SCAFFOLD: Stochastic controlled averaging for on-device federated learning. *arXiv preprint arXiv:1910.06378*, 2019.

David A Levin and Yuval Peres. *Markov chains and mixing times*, volume 107. American Mathematical Soc., 2017.

Tian Li, Anit Kumar Sahu, Ameet Talwalkar, and Virginia Smith. Federated learning: Challenges, methods, and future directions. *IEEE Signal Processing Magazine*, 37(3):50–60, 2020a.

Xiang Li, Kaixuan Huang, Wenhao Yang, Shusen Wang, and Zhihua Zhang. On the convergence of fedavg on non-iid data. In *International Conference on Learning Representations (ICLR)*, July 2020b. URL <https://arxiv.org/abs/1907.02189>.

Grigory Malinovsky, Samuel Horváth, Konstantin Burlachenko, and Peter Richtárik. Federated learning with regularized client participation, 2023.

Brendan McMahan and Abhradeep Thakurta. Federated learning with formal differential privacy guarantees. *Google AI Blog*, 2022.

H. Brendan McMahan, Eider Moore, Daniel Ramage, Seth Hampson, and Blaise Agøura y Arcas. Communication-Efficient Learning of Deep Networks from Decentralized Data. *International Conference on Artificial Intelligence and Statistics (AISTATS)*, April 2017. URL <https://arxiv.org/abs/1602.05629>.

H Brendan McMahan, Zheng Xu, and Yanxiang Zhang. A hassle-free algorithm for private learning in practice: Don’t use tree aggregation, use blts. *arXiv preprint arXiv:2408.08868*, 2024.

Kumar Kshitij Patel, Lingxiao Wang, Blake E Woodworth, Brian Bullins, and Nati Srebro. Towards optimal communication complexity in distributed non-convex optimization. *Advances in Neural Information Processing Systems*, 35:13316–13328, 2022.

Matthias Paulik, Matt Seigel, Henry Mason, Dominic Telaar, Joris Kluivers, Rogier van Dalen, Chi Wai Lau, Luke Carlson, Filip Granqvist, Chris Vandevelde, et al. Federated evaluation and tuning for on-device personalization: System design & applications. *arXiv preprint arXiv:2102.08503*, 2021.

S Sundhar Ram, A Nedić, and Venugopal V Veeravalli. Asynchronous gossip algorithms for stochastic optimization. In *Proceedings of the 48h IEEE Conference on Decision and Control (CDC) held jointly with 2009 28th Chinese Control Conference*, pp. 3581–3586. IEEE, 2009.

Mónica Ríbero, Haris Vikalo, and Gustavo De Veciana. Federated learning under intermittent client availability and time-varying communication constraints. *IEEE Journal of Selected Topics in Signal Processing*, 17(1):98–111, 2022.

Angelo Rodio, Francescomaria Faticanti, Othmane Marfoq, Giovanni Neglia, and Emilio Leonardi. Federated learning under heterogeneous and correlated client availability. In *IEEE INFOCOM 2023 - IEEE Conference on Computer Communications*, pp. 1–10, 2023. doi: 10.1109/INFOCOM53939.2023.10228876.

Zhenyu Sun and Ermin Wei. A communication-efficient algorithm with linear convergence for federated minimax learning. *Advances in Neural Information Processing Systems*, 35:6060–6073, 2022.

Richard S Sutton, David McAllester, Satinder Singh, and Yishay Mansour. Policy gradient methods for reinforcement learning with function approximation. *Advances in neural information processing systems*, 12, 1999.

John Tsitsiklis and Benjamin Van Roy. Analysis of temporal-difference learning with function approximation. *Advances in neural information processing systems*, 9, 1996.

John N Tsitsiklis and Benjamin Van Roy. Average cost temporal-difference learning. *Automatica*, 35(11):1799–1808, 1999.

Jianyu Wang and Gauri Joshi. Cooperative SGD: A unified framework for the design and analysis of communication-efficient SGD algorithms. *Journal of Machine Learning Research (JMLR)*, 2021. URL <https://arxiv.org/abs/1808.07576>.

Jianyu Wang, Qinghua Liu, Hao Liang, Gauri Joshi, and H. Vincent Poor. Tackling the Objective Inconsistency Problem in Heterogeneous Federated Optimization. *preprint*, May 2020. URL <https://arxiv.org/abs/2007.07481>.

Jianyu Wang, Zachary Charles, Zheng Xu, Gauri Joshi, H Brendan McMahan, Maruan Al-Shedivat, Galen Andrew, Salman Avestimehr, Katharine Daly, Deepesh Data, et al. A field guide to federated optimization. *arXiv preprint arXiv:2107.06917*, 2021.

Jianyu Wang, Rudrajit Das, Gauri Joshi, Satyen Kale, Zheng Xu, and Tong Zhang. On the unreasonable effectiveness of federated averaging with heterogeneous data. *arXiv preprint arXiv:2206.04723*, 2022.

Shiqiang Wang and Mingyue Ji. A unified analysis of federated learning with arbitrary client participation. In *Advances in Neural Information Processing Systems*, 2022.

Shiqiang Wang and Mingyue Ji. A lightweight method for tackling unknown participation probabilities in federated averaging. *arXiv preprint arXiv:2306.03401*, 2023.

Blake E Woodworth, Kumar Kshitij Patel, and Nati Srebro. Minibatch vs local sgd for heterogeneous distributed learning. *Advances in Neural Information Processing Systems*, 33:6281–6292, 2020.

Ming Xiang, Stratis Ioannidis, Edmund Yeh, Carlee Joe-Wong, and Lili Su. Efficient federated learning against heterogeneous and non-stationary client unavailability. *arXiv preprint arXiv:2409.17446*, 2024.

Zheng Xu and Yanxiang Zhang. Advances in private training for production on-device language models. *Goolge Research Blog*, 2024.

Zheng Xu, Yanxiang Zhang, Galen Andrew, Christopher Choquette, Peter Kairouz, Brendan McMahan, Jesse Rosenstock, and Yuanbo Zhang. Federated learning of gboard language models with differential privacy. *ACL Industry*, 2023.

Haibo Yang, Minghong Fang, and Jia Liu. Achieving linear speedup with partial worker participation in non-iid federated learning. *arXiv preprint arXiv:2101.11203*, 2021.

Qiang Yang, Yang Liu, Tianjian Chen, and Yongxin Tong. Federated machine learning: Concept and applications. *ACM Transactions on Intelligent Systems and Technology (TIST)*, 10(2):1–19, 2019.

Chen Zhu, Zheng Xu, Mingqing Chen, Jakub Konečný, Andrew Hard, and Tom Goldstein. Diurnal or nocturnal? federated learning of multi-branch networks from periodically shifting distributions. In *International Conference on Learning Representations*, 2021.

A RELATED WORK

Non-uniform & correlated client participation. There is a recent surge of efforts to investigate FL with non-uniform client participation both from theoretical and empirical perspectives. Earlier work presumes that clients are sampled by the server uniformly, which guarantees the global model held by the server is an unbiased estimate as that in the full participation setting and hence allows extension of convergence results for the full-participation setting to the partial-participation setting (Jhunjhunwala et al., 2022; Karimireddy et al., 2019; Yang et al., 2021; Bian et al., 2024). The above-mentioned uniform participation is, however, far from the reality as clients may have their intrinsic sampling probabilities p_i 's that are non-uniform due to, for example, intermittent availability resulting from practical constraints. Recent works analyzed the convergence behaviors of FL algorithms when such p_i 's are known as a prior or controllable (Wang & Ji, 2022; Karimireddy et al., 2019; Chen et al., 2020; Fraboni et al., 2021). However, pointed out by (Bonawitz et al., 2019; Wang et al., 2021), client participation pattern can highly depend on the underlying system characteristics, which is thus hard to know or control. As characterized by (Wang & Ji, 2022; Xiang et al., 2024), such unknown and non-uniform participation statistics causes a bias in the model updates as more frequently participating clients dominate the average update. In order to mitigate the effect of bias, (Patel et al., 2022; Ribero et al., 2022; Wang & Ji, 2023) introduced reweighting mechanisms combined with dynamically estimating client participation distributions. Such idea is also introduced in asynchronous distributed learning literature (Ram et al., 2009). Most works aiming at analyzing non-uniform participation, however, rely on the unrealistic assumption that every client participates in the system independently, which fails to capture practical scenarios where each client's participation is influenced by others across rounds (Kairouz et al., 2019; Eichner et al., 2019; Zhu et al., 2021). One interesting time-correlated participation pattern is that clients have to wait for at least R (called minimum separation) rounds between consecutive participation (McMahan & Thakurta, 2022; Xu et al., 2023). In particular, imposing a minimum separation constraint has been empirically shown to benefit privacy preservation in FL applications (Kairouz et al., 2021; Choquette-Choo et al., 2023; Xu et al., 2023; Xu & Zhang, 2024). Instead, such time-correlated participation has not been fully investigated theoretically. The only work that partially captures the above case is (Cho et al., 2023) where the clients are forced to follow a cyclic participation, which is an extreme case of very large R . Therefore, in this paper we study convergence performances of FL algorithms under non-uniform and correlated client participation, which provides theoretical explanations for their empirical counterparts in practice.

Stochastic optimization with Markov-sampling. Another line of related works is stochastic gradient-based optimization under Markov-sampling. Unlike classical stochastic optimization literature where i.i.d. samples are drawn during the training process (Allen-Zhu & Hazan, 2016; Allen-Zhu, 2017; Johnson & Zhang, 2013; Defazio et al., 2014), many contexts, including TD-learning and reinforcement learning (RL), require to optimize the objective function by utilizing samples generated by a Markov chain (Tsitsiklis & Van Roy, 1996; 1999; Bhatnagar et al., 2007; Sutton et al., 1999). Recently, the work (Even, 2023) provided convergence guarantees for SGD under Markov-sampling when the objectives are convex, strongly convex and non-convex. Then (Beznosikov et al., 2024) further proposed an accelerated method and generalized the analysis to variational inequalities. Both of them restrict on the first-order Markov chains. It has been shown by literature that gradient-based methods converge to the optimal solution of the objective induced by the stationary distribution of the underlying Markov chain (Even, 2023; Beznosikov et al., 2024). This indicates that the final solution is biased if the stationary distribution is non-uniform and existing literature cannot deal with such bias problem. In contrast, in this paper our analysis suits for higher-order Markov chains and the proposed algorithm enables the convergence to an unbiased solution without any information and constraint on the Markov chain and stationary distribution.

B PRELIMINARIES OF MARKOV CHAINS

In this section, we summarize several notions and properties of the conventional Markov chain (i.e., first-order Markov chain). We only focus on finite Markov chains, meaning the state space is finite. Note that for a finite Markov chain, we can use its transition matrix to uniquely represent it.

Definition 2. *Given a finite Markov chain with transition matrix P , we say it is irreducible if its induced graph is strongly connected, i.e., every state can be reached from every other state.*

Note that $[P^k]_{i,j}$ is the probability transiting from state i to state j with exactly k steps, based on which we introduce the definition of aperiodic and periodic Markov chains.

Definition 3. *The period of state i is the greatest common divisor (g.c.d.) of the set $\{k \in \mathbb{N} \mid [P^k]_{i,i} > 0\}$. If every state has period 1 then the Markov chain is aperiodic, otherwise it is periodic.*

In other words, the period of state i can be achieved by calculating the g.c.d. of the number of steps starting from i and returning back. If the Markov chain is also irreducible, we have the following.

Lemma 3. *If the Markov chain is irreducible, every state has the same period.*

Next important result states the convergence of the Markov chain.

Lemma 4. *Suppose a finite Markov chain with transition matrix P is irreducible and aperiodic. Then, there exist some $\rho \in (0, 1)$ and $C > 0$ such that*

$$\max_x \|P^k(x, \cdot) - \pi\|_{TV} \leq C\rho^k$$

where π is the unique, strictly positive stationary distribution; $\|\cdot\|_{TV}$ denotes the total variation.

Lemma 4 implies that starting from any initial distribution, the Markov chain converges to the stationary distribution at linear rate. Without confusion, we denote $d_{TV}(P^k, \mathbf{1}\pi^T) = \max_x \|P^k(x, \cdot) - \pi\|_{TV}$. Note that $d_{TV}(P^k, \mathbf{1}\pi^T) = \frac{1}{2}\|P^k - \mathbf{1}\pi^T\|_\infty$. Then, we define the mixing time of the chain.

Definition 4. *Given any $\epsilon > 0$, the mixing time $t_{mix}(\epsilon)$ is defined as $t_{mix}(\epsilon) := \inf\{l \geq 1 \mid d_{TV}(P^l, \mathbf{1}\pi^T) \leq \epsilon\}$. Conventionally, we denote $\tau_{mix} = t_{mix}(1/4)$.*

Lemma 5. *We have the following statements:*

$$(1). d_{TV}(P^{t+1}, \mathbf{1}\pi^T) \leq d_{TV}(P^t, \mathbf{1}\pi^T), \forall t \geq 0.$$

$$(2). \text{For } k \geq 2, t_{mix}(2^{-k}) \leq (k-1)\tau_{mix}.$$

$$(3). \text{Moreover,}$$

$$\sum_{k=0}^T d_{TV}(P^k, \mathbf{1}\pi^T) \leq c_0\tau_{mix}, \forall T \geq 0$$

for some constant $c_0 > 0$.

Proof. The first two claims are shown in (Levin & Peres, 2017). To see the third claim, we note that

$$\begin{aligned} \sum_{k=0}^T d_{TV}(P^k, \mathbf{1}\pi^T) &\leq \sum_{k=0}^{\infty} d_{TV}(P^k, \mathbf{1}\pi^T) \\ &\leq \sum_{l=0}^{\tau_{mix}} d_{TV}(P^l, \mathbf{1}\pi^T) + \sum_{k=2}^{\infty} \sum_{l=t_{mix}(2^{-k})+1}^{t_{mix}(2^{-(k+1)})} d_{TV}(P^l, \mathbf{1}\pi^T) \\ &\leq d_{TV}(P, \mathbf{1}\pi^T)\tau_{mix} + \sum_{k=2}^{\infty} (t_{mix}(2^{-(k+1)}) - t_{mix}(2^{-k}))2^{-k} \\ &\leq d_{TV}(P, \mathbf{1}\pi^T)\tau_{mix} + \sum_{k=2}^{\infty} k2^{-k}\tau_{mix} \\ &\leq d_{TV}(P, \mathbf{1}\pi^T)\tau_{mix} + 2\tau_{mix} \end{aligned}$$

which completes the proof with $c_0 = d_{TV}(P, \mathbf{1}\pi^T) + 2$. \square

C PROOFS OF PROPOSITION 1 AND LEMMA 1

C.1 PROOF OF PROPOSITION 1

Property (1) follows from equation 2, where by definition within $R + 1$ rounds, the clients cannot participate in the system twice or more. Property (2) follows from the fact that every client is

sampled with probability proportional to its availability probability p_i if it has waited for R rounds. Property (3) then directly follows from the definition of the first-order Markov chain and Properties (1), (2). Properties (4) and (5) are due to the observation that every row and column has exactly $\sigma(B(M - R), B)$ non-zero entries due to Property (3).

C.2 PROOF OF LEMMA 1

It is obvious that the Markov chain is irreducible in the sense that all ordered sequences $(\mathcal{I}_1, \dots, \mathcal{I}_R)$ can be observed due to every client has strictly positive probability to be selected. To see that it is aperiodic for $R \leq M - 2$, we only need to show that starting from the state $(\mathcal{I}_1, \dots, \mathcal{I}_R)$ where $\mathcal{I}_k = ((k-1)B+1, \dots, kB)$, $k = 1, \dots, R$, both $R+1$ steps and $R+2$ steps can be possibly taken such that the first return happens, which implies aperiodicity. This is because if a Markov chain is irreducible, all the states have the same period by Lemma 3.

Then, we consider the following two constructed sequence. Let $h_1 = (\mathcal{I}_1, \dots, \mathcal{I}_R, \mathcal{I}_{R+1}, \mathcal{I}_1, \dots, \mathcal{I}_R)$ for state $\mathcal{I}_{R+1} = (RB+1, \dots, (R+1)B)$, where the length of h_1 is $2R+1$. Denote $h_1[k]$ as the entry at the k -th position. We construct the sequence $\{Y_t, Y_{t+1}, \dots, Y_{t+R}\}$ as $Y_{t+k-1} = (h_1[k \bmod (2R+1)], \dots, h_1[(k+R-1) \bmod (2R+1)])$, $k = 1, \dots, 2R+1$, i.e., starting from $(\mathcal{I}_1, \dots, \mathcal{I}_R)$ exactly $R+1$ steps are taken to firstly return. Similar to the definition of h_1 , let $h_2 = (\mathcal{I}_1, \dots, \mathcal{I}_R, \mathcal{I}_{R+1}, \mathcal{I}_{R+2}, \mathcal{I}_1, \dots, \mathcal{I}_R)$ with its length $2R+2$ and state $\mathcal{I}_{R+2} = ((R+1)B+1, (R+2)B)$. We then construct the sequence $\{Y_t, \dots, Y_{t+R+1}\}$ as $Y_{t+k-1} = (h_2[k \bmod (2R+2)], \dots, h_2[(k+R-1) \bmod (2R+2)])$, $k = 1, \dots, 2R+2$, which then suggests exactly $R+2$ steps are required to return back to $(\mathcal{I}_1, \dots, \mathcal{I}_R)$. Combining these two cases leads to the Markov chain is aperiodic for any $R \leq M - 2$.

D PROOFS OF PROPOSITION 2 AND THEOREM 2

D.1 PROOF OF THEOREM 2

Let us first consider the case when $B = 1$ and given $p_1 > 0$, $p_i = \frac{1-p_1}{N-1}$, $\forall i = 2, \dots, N$. Then, for any $0 < R \leq N - 1$ and any (j_0, \dots, j_{R-1}) , pick an arbitrary $j_R \in \{j_0, \dots, j_{R-1}\}^c$. By denoting $b_R = b(P_R[\cdot, (j_0, \dots, j_{R-1})])$, $b_{R+1} = b(P_{R+1}[\cdot, (j_0, \dots, j_R)])$ (which are the column sums for each column of P_R and P_{R+1} , respectively) and letting $S_R := \{j_0, \dots, j_{R-1}\}$, $S_{R+1} := \{j_0, \dots, j_R\}$ for notation simplicity. By observing that when π_R is exactly the uniform distribution, the sum of P_R for each column is exactly one, we then tend to prove that the column sum of P_R asymptotically approaches one as R increases. We have four cases.

Case I: $j_0 = \{1\}$. Then, for any $0 \leq R \leq N - 2$, utilizing last two properties in Proposition 1,

$$\begin{aligned} b_{R+1} - b_R &= p_1 \sum_{k \in S_{R+1}^c} (p_1 + \sum_{i \in S_{R+1}^c} p_i - p_k)^{-1} - p_1 \sum_{k \in S_R^c} (p_1 + \sum_{i \in S_R^c} p_i - p_k)^{-1} \\ &= p_1 \sum_{k \in S_{R+1}^c} (p_1 + \frac{1-p_1}{N-1}(N-R-2))^{-1} - p_1 \sum_{k \in S_R^c} (p_1 + \frac{1-p_1}{N-1}(N-R-1))^{-1} \\ &= p_1(N-R-1)(p_1 + \frac{1-p_1}{N-1}(N-R-2))^{-1} - p_1(N-R)(p_1 + \frac{1-p_1}{N-1}(N-R-1))^{-1} \end{aligned}$$

Let $r = N - R - 1$. We simply b_R as

$$b_R = \frac{p_1 r}{p_1 + \frac{1-p_1}{N-1}(r-1)} = \frac{p_1(N-1)}{1-p_1} + \frac{p_1(1 - \frac{p_1(N-1)}{1-p_1})}{p_1 + \frac{1-p_1}{N-1}(r-1)}$$

Then,

$$\begin{aligned} b_{R+1} - b_R &= p_1(1 - \frac{p_1(N-1)}{1-p_1}) \left(\frac{1}{p_1 + \frac{1-p_1}{N-1}(r-1)} - \frac{1}{p_1 + \frac{1-p_1}{N-1}r} \right) \\ &= \frac{p_1(1-p_1)}{N-1} \left(1 - \frac{p_1(N-1)}{1-p_1} \right) (p_1 + \frac{1-p_1}{N-1}(r-1))^{-1} (p_1 + \frac{1-p_1}{N-1}r)^{-1} \end{aligned}$$

which is strictly positive for $p_1 < 1/N$ for all $0 \leq R \leq N - 2$.

Case II: $\{1\} \in S_{R+1}^c$. Then, we obtain $p_{j_0} = \frac{1-p_1}{N-1}$ and hence

$$\begin{aligned} b_R/p_{j_0} &= (p_{j_0} + \frac{1-p_1}{N-1}(N-R-1))^{-1} + (N-R-1)(p_1 + \frac{1-p_1}{N-1}(N-R-1))^{-1} \\ &= \frac{N-1}{(1-p_1)(r+1)} + r(p_1 + \frac{1-p_1}{N-1}r)^{-1} \\ &= \frac{N-1}{(1-p_1)(r+1)} + \frac{N-1}{1-p_1} - \frac{p_1(N-1)}{1-p_1} \frac{1}{p_1 + \frac{1-p_1}{N-1}r} \end{aligned}$$

where we let $r = N - R - 1$. Then, denoting $\bar{p} = \frac{1-p_1}{N-1}$ and $\alpha = p_1/\bar{p}$ yields

$$\begin{aligned} (b_{R+1} - b_R)/p_{j_0} &= \frac{1}{\bar{p}} \frac{1}{r(r+1)} - \frac{p_1}{\bar{p}^2(r+\alpha-1)(r+\alpha)} \\ &= \frac{(r+\alpha)(r+\alpha-1) - \alpha r(r+1)}{\bar{p}r(r+1)(r+\alpha)(r+\alpha-1)} \\ &= \frac{(1-\alpha)r^2 + (\alpha-1)r + \alpha(\alpha-1)}{\bar{p}r(r+1)(r+\alpha)(r+\alpha-1)} \\ &= \frac{(1-\alpha)(r^2 - r - \alpha)}{\bar{p}r(r+1)(r+\alpha)(r+\alpha-1)}. \end{aligned}$$

Note that when $p_1 < 1/N$, $\alpha < 1$, which indicates $b_{R+1} - b_R > 0$, $\forall 0 \leq R \leq N - 3$ by observing $r^2 - r - \alpha \geq 0$. Moreover, note that $b_R > 1$, $\forall R \leq N - 2$ in this case by

$$b_R = \frac{1}{r+1} + 1 - \frac{\alpha}{r+\alpha} = \frac{(1-\alpha)r}{(r+1)(r+\alpha)} + 1 > 1$$

for $\alpha < 1$. And a straightforward calculation gives $b_{N-2} < \frac{3}{2}$, which then indicates $|b_R - 1| < \frac{1}{2}$, $\forall R \leq N - 1$.

Case III: $\{1\} \in S_R^c$ and $\{1\} \notin S_{R+1}^c$. In this case, $p_{j_0} = \frac{1-p_1}{N-1} = \bar{p}$. Then, a simple calculation gives

$$(b_{R+1} - b_R)/p_{j_0} = \frac{1}{\bar{p}} \frac{(\alpha-1)r}{(r+1)(r+\alpha)} < 0$$

when $p_1 < 1/N$.

Case IV: $\{1\} \notin S_R^c$. Then, all the clients are available in both S_R^c and S_{R+1}^c have availability probability \bar{p} . Then, it is obvious that $b_R = 1$, $\forall 0 \leq R \leq N - 1$.

For Cases I, III and IV, we conclude that when $p_1 < 1/N$ and $p_i = \frac{1-p_1}{N-1}$, $i = 2, \dots, N$, $|b_{R+1} - 1| < |b_R - 1|$, $\forall 0 \leq R \leq N - 2$ by further noting that $b_{N-1} = 1$. By Case II, we then have all $|b_R - 1|$ converges to $[0, 0.5]$ as R increases. Observe $b_{N-1} = 1$ corresponds to the case that ζ_{N-1} is exactly the uniform distribution and so is π_{N-1} . This indicates that π_R converges to some neighborhood of the uniform distribution $\frac{1}{N}\mathbf{1}_N$. In order to characterize this neighborhood, we turn to carefully analyze Case II, i.e., $|b_R - 1| < 0.5$. Noting that Case II corresponds to at most $1 - R/N$ portion of columns in P_R and so does π_R , therefore the neighborhood is characterized by $\{\pi \mid \|\pi - \frac{1}{N}\mathbf{1}_N\|_1 = \mathcal{O}(1/N)\}$.

Next, in order to prove the statement, we perturb each $p_i = \frac{1-p_1}{N-1}$, $i = 2, \dots, N$ by some scalar ϵ_i such that $\sum_{i=2}^N \epsilon_i = 0$. Note that $b_{R+1} - b_R$ is continuous in $(\epsilon_2, \dots, \epsilon_N)$ and so is π_R , which then implies that there exists some positive $\Delta > 0$ such that $b_{R+1} - b_R$ preserves the original properties as before the perturbation is added for all $|\epsilon_i| \leq \Delta$. Therefore, we achieve the statement that π_R converges to the neighborhood $\{\pi \mid \|\pi - \frac{1}{N}\mathbf{1}_N\|_1 = \mathcal{O}(1/N)\}$ when $B = 1$. Obtaining the statement for $B > 1$ follows the same technique by noting that we can always calculate the equivalent \tilde{p}_i for each batch with size B . Specifically, given a batch of clients, say \mathcal{B}_i , then $\tilde{p}_i = \prod_{j \in \mathcal{B}_i} p_j / C$ with suitable normalization constant C and we can then obtain the convergence of π_R to a neighborhood of the uniform distribution by similar development.

D.2 PROOF OF PROPOSITION 2

The proof of Proposition 2 is straightforward by observing that $b_R = 1, \forall R$ when $p_i = 1/N, \forall i \in [N]$. Then $\mathbf{1}^T P_R = \mathbf{1}^T, \forall R$ which indicates π_R is always the uniform distribution.

E INTERMEDIATE LEMMAS

In this section, we present some useful intermediate results under the following generalized setting: we consider a general global objective function defined as $F_w(x) := \sum_{i=1}^N w_i f_i(x)$ where $\sum_{i=1}^N w_i = 1$ and $w_i \geq 0, \forall i \in [N]$. And we consider the following local update

$$x_{t,k+1}^i = x_{t,k}^i - \alpha q_t^i \nabla f_i(x_{t,k}^i) \quad (10)$$

where $q_t^i = \frac{w_i}{y_t^i}$ for some positive sequence y_t^i . Note that the above update equation 10 is a generalized version of equation 9 in Algorithm 1. Then we have the following useful lemmas when forcing the update equation 10.

Lemma 6. *Under Assumption 1, we have for any x*

$$\begin{aligned} \|\nabla F_w(x) - \nabla F(x)\| &\leq G \\ \|\nabla f_i(x) - \nabla F_w(x)\| &\leq 2G, \forall i \in [N]. \end{aligned}$$

Proof. Note that Assumption 1 implies

$$\begin{aligned} \|\nabla F_w(x) - \nabla F(x)\| &= \left\| \sum_{i=1}^N w_i (\nabla f_i(x) - \nabla F(x)) \right\| \\ &\leq \sum_{i=1}^N w_i \|\nabla f_i(x) - \nabla F(x)\| \\ &\leq G \sum_{i=1}^N w_i = G. \end{aligned}$$

Then, for any $i \in [N]$

$$\|\nabla f_i(x) - \nabla F_w(x)\| \leq \|\nabla f_i(x) - \nabla F(x)\| + \|\nabla F_w(x) - \nabla F(x)\| \leq 2G.$$

□

Lemma 7. *Given any t and i , we have $\|x_{t,k}^i - x_t\|^2 \leq \gamma^2 L^{-2} \|\nabla F_w(x_t)\|^2 + 4\gamma^2 L^{-2} G^2$, $\forall k = 0, \dots, K$, when $\alpha \leq \min \left\{ \frac{\gamma}{8KL}, \frac{\gamma}{8KLq_t^i} \right\}$ and $\gamma \leq 1/3$.*

Proof. During the t -th communication round, S_t and q_t^i are fixed. Then, for any $\beta > 0$ and $\alpha \leq \min \left\{ \frac{\gamma}{\beta L}, \frac{\gamma}{\beta L q_t^i} \right\}$, using Lemma 6 gives

$$\begin{aligned} \|x_{k+1}^i - x_t\|^2 &\leq (1 + \beta^{-1}) \|x_k^i - x_t\|^2 + (1 + \beta)(\alpha)^2 (q_t^i)^2 \|\nabla f_i(x_k^i)\|^2 \\ &\leq (1 + \beta^{-1}) \|x_k^i - x_t\|^2 + 3(1 + \beta)(\alpha)^2 (q_t^i)^2 (\|\nabla f_i(x_k^i) - \nabla f_i(x_t)\|^2 \\ &\quad + \|\nabla f_i(x_t) - \nabla F_w(x_t)\|^2 + \|\nabla F_w(x_t)\|^2) \\ &\leq (1 + \beta^{-1}) \|x_k^i - x_t\|^2 + 3(1 + \beta)(\alpha q_t^{s_t})^2 (L^2 \|x_k^i - x_t\|^2 + 4G^2 + \|\nabla F_w(x_t)\|^2) \\ &\leq (1 + \beta^{-1}) \|x_k^i - x_t\|^2 + \frac{3(1 + \beta)\gamma^2}{\beta^2 L^2} (L^2 \|x_k^i - x_t\|^2 + 4G^2 + \|\nabla F_w(x_t)\|^2) \\ &= (1 + (1 + 3\gamma^2)\beta^{-1} + 3\gamma^2\beta^{-2}) \|x_k^i - x_t\|^2 + \frac{3(1 + \beta)\gamma^2}{\beta^2 L^2} (4G^2 + \|\nabla F_w(x_t)\|^2) \\ &\leq \exp \left(\frac{1 + 6\gamma^2}{\beta} \right) \|x_k^i - x_t\|^2 + \frac{3(1 + \beta)\gamma^2}{\beta^2 L^2} (4G^2 + \|\nabla F_w(x_t)\|^2) \end{aligned}$$

for any $\beta \geq 1$. Unrolling the above gives for any $k = 0, \dots, K-1$

$$\|x_k^i - x_t\|^2 \leq \sum_{k=0}^{K-1} \exp\left(\frac{1+6\gamma^2}{\beta} k\right) \frac{3(1+\beta)\gamma^2}{\beta^2 L^2} (G^2 + \|\nabla F_w(x_t)\|^2)$$

which further indicates by choosing $\gamma \leq 1/3$

$$\begin{aligned} \|x_k^i - x_t\|^2 &\leq \sum_{k=0}^{K-1} e^{2k\beta^{-1}} \frac{3(1+\beta)\gamma^2}{\beta^2 L^2} (4G^2 + \|\nabla F_w(x_t)\|^2) \\ &= \frac{1-e^{2K/\beta}}{1-e^{2/\beta}} \cdot \frac{3(1+\beta)\gamma^2}{\beta^2 L^2} (4G^2 + \|\nabla F_w(x_t)\|^2) \\ &\leq \frac{(e^{2K/\beta}-1)3\gamma^2}{L^2} (4G^2 + \|\nabla F_w(x_t)\|^2) \\ &\leq \frac{\gamma^2}{L^2} (4G^2 + \|\nabla F_w(x_t)\|^2) \end{aligned}$$

when choosing $\beta = 8K$.

□

Lemma 8. For any $t \geq \tau$, we have $\|x_t - x_{t-\tau}\|^2 \leq 4\gamma^2 L^{-2} \tau^2 G^2 + \gamma^2 L^{-2} \tau \sum_{l=t-\tau}^{t-1} \|\nabla F_w(x_l)\|^2$ when $\alpha \leq \min\{\frac{\gamma}{8KLq}, \frac{\gamma}{8KLq_t^i}\}$ and $\gamma \leq 1/3$.

Proof. Note that

$$\begin{aligned} \|x_{t+1} - x_t\|^2 &= \left\| \frac{1}{|S_t|} \sum_{i \in S_t} x_{t,K}^i - x_t \right\|^2 \\ &\leq \frac{1}{|S_t|} \sum_{i \in S_t} \|x_{t,K}^i - x_t\|^2 \\ &\leq \frac{\gamma^2}{L^2} (4G^2 + \|\nabla F_w(x_t)\|^2). \end{aligned}$$

Then,

$$\begin{aligned} \|x_t - x_{t-\tau}\|^2 &= \left\| \sum_{l=t-\tau}^{t-1} x_{l+1} - x_l \right\|^2 \\ &\leq \tau \sum_{l=t-\tau}^{t-1} \|x_{l+1} - x_l\|^2 \\ &\leq \frac{4\gamma^2}{L^2} \tau^2 G^2 + \frac{\gamma^2}{L^2} \tau \sum_{l=t-\tau}^{t-1} \|\nabla F_w(x_l)\|^2. \end{aligned}$$

□

Lemma 9. For any $l \in [t-\tau, t]$ with $t \geq \tau \geq 1$ and $\alpha \leq \min\{\frac{\gamma}{8KLq}, \frac{\gamma}{8KLq_t^i}\}$ with $\gamma \leq \min\{\frac{1}{2\tau}, \frac{1}{3}\}$ we have

$$\max_{t-\tau \leq l \leq t} \mathbb{E} \|\nabla F_w(x_l)\|^2 \leq 4\mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 + 16\tau^2 \gamma^2 G^2.$$

Proof. For any $t-\tau \leq l \leq t$, we have

$$\begin{aligned} \mathbb{E} \|\nabla F_w(x_l)\|^2 &\leq 2\mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 + 2\mathbb{E} \|\nabla F_w(x_l) - \nabla F_w(x_{t-\tau})\|^2 \\ &\leq 2\tau\gamma^2 \sum_{l=t-\tau}^{t-1} \mathbb{E} \|\nabla F_w(x_l)\|^2 + 8\tau^2 \gamma^2 G^2 + 2\mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 \\ &\leq 2\tau^2 \gamma^2 \max_{t-\tau \leq l \leq t} \mathbb{E} \|\nabla F_w(x_l)\|^2 + 8\tau^2 \gamma^2 G^2 + 2\mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 \\ &\leq \frac{1}{2} \max_{t-\tau \leq l \leq t} \mathbb{E} \|\nabla F_w(x_l)\|^2 + 8\tau^2 \gamma^2 G^2 + 2\mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 \end{aligned}$$

where the second inequality follows Lemma 8 and we use $\gamma \leq 1/(2\tau)$ in the last inequality. Finally, taking the maximum over l on the left-hand side completes the proof. \square

Lemma 10. Define $F_w := \sum_{i=1}^N w_i f_i$ for $\sum_{i=1}^N w_i = 1, w_i \geq 0$. Suppose Assumptions 1,2 hold. Considering any sequence y_t^i that satisfies $\sum_{i=1}^N y_t^i = 1, y_t^i \geq a^{-1} > 0, \forall i \in [N], t \geq 0$ and letting $q_t^i = \frac{w_i}{y_t^i}, \forall i \in [N]$, then, given $\tau \geq \tau_{\text{mix}} \log(1/\delta)$ with $0 < \delta < 1$, for $\alpha \leq \frac{\gamma}{8\bar{a}KL}$ with $\gamma \leq \min\{\frac{1}{384\tau L}, \frac{L}{384\tau}, \frac{1}{3}\}$, we have $\forall T > \tau$,

$$\begin{aligned} \frac{1}{T-\tau} \sum_{t=\tau}^{T-1} \mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 &\leq \frac{32\bar{a}L\Delta_\tau}{\gamma(T-\tau)} + \frac{8}{T-\tau} \sum_{t=\tau}^{T-1} \mathbb{E} [\|\tilde{q}_t\|_\infty^2 \|\nabla F_w(x_{t-\tau})\|^2] \\ &\quad + 32\bar{a}LG^2 \left(3\gamma + 6\gamma\tau^2 + \frac{2\gamma}{L} + \frac{3\gamma}{16L^2} + \frac{\gamma^2}{16aL} \right) \\ &\quad + \frac{32G^2}{T-\tau} \sum_{t=\tau}^{T-1} \mathbb{E} \|\tilde{q}_t\|_\infty^2 + 8c_1^2 \delta^2 G^2 \end{aligned}$$

where $\bar{a} = a \max_i \{w_i\}$, $\tilde{q}_t = (\tilde{q}_t^1, \dots, \tilde{q}_t^N)$ with $\tilde{q}_t^i = q_t^i - \frac{w_i}{\pi_i}$, and c_1 is some constant. Moreover,

$$\Delta_\tau := \mathbb{E}[F_w(x_\tau) - \min_x F_w(x)] \leq \frac{\gamma\tau}{2\bar{a}L} G^2 + \mathbb{E}[F_w(x_0) - F_w^*].$$

Proof. For notation simplicity, we drop subscript t for $x_{t,k}^i$. Define $q_t^i = \frac{w_i}{y_t^i}$. Note that

$$\begin{aligned} x_K^i &= x_t - \sum_{k=0}^{K-1} \alpha q_t^i \nabla f_i(x_k^i) \\ x_{t+1} &= x_t - \frac{1}{B} \sum_{i \in S_t} \sum_{k=0}^{K-1} \alpha q_t^i \nabla f_i(x_k^i) \end{aligned}$$

where S_t denotes the subset of clients drawn in the t -th round. Due to the smoothness of every f_i , we have

$$\mathbb{E}[F_w(x_{t+1}) - F_w(x_t)] \leq \mathbb{E}\langle \nabla F_w(x_t), x_{t+1} - x_t \rangle + \frac{L}{2} \mathbb{E}\|x_{t+1} - x_t\|^2.$$

Considering $t \geq \tau$ for any $\tau \geq 0$,

$$\begin{aligned} \mathbb{E}\langle \nabla F_w(x_t), x_{t+1} - x_t \rangle &= -\mathbb{E}\langle \nabla F_w(x_t), \frac{1}{B} \sum_{i \in S_t} \sum_{k=0}^{K-1} \alpha q_t^i \nabla f_i(x_k^i) \rangle \\ &= \underbrace{\mathbb{E}\langle \nabla F_w(x_{t-\tau}) - \nabla F_w(x_t), \frac{1}{B} \sum_{i \in S_t} \sum_{k=0}^{K-1} \alpha q_t^i \nabla f_i(x_k^i) \rangle}_{e_1} \\ &\quad + \underbrace{\mathbb{E}\langle -\nabla F_w(x_{t-\tau}), \frac{1}{B} \sum_{i \in S_t} \sum_{k=0}^{K-1} \alpha q_t^i \nabla f_i(x_{t-\tau}) \rangle}_{e_2} \\ &\quad + \underbrace{\mathbb{E}\langle -\nabla F_w(x_{t-\tau}), \frac{1}{B} \sum_{i \in S_t} \sum_{k=0}^{K-1} \alpha q_t^i (\nabla f_i(x_k^i) - \nabla f_i(x_t)) \rangle}_{e_3} \\ &\quad + \underbrace{\mathbb{E}\langle -\nabla F_w(x_{t-\tau}), \frac{1}{B} \sum_{i \in S_t} \sum_{k=0}^{K-1} \alpha q_t^i (\nabla f_i(x_t) - \nabla f_i(x_{t-\tau})) \rangle}_{e_4}. \end{aligned}$$

We first note that according to the conditions on y_t^i , $w_i \leq q_t^i \leq aw_i$ with some positive constant $a < \infty$ for every $i \in [N]$ and $\forall t \geq 0$. Then by choosing $\alpha \leq \frac{\gamma}{8aKLw_m} \leq \min\{\frac{\gamma}{8KL}, \frac{\gamma}{8KL \max_i \{q_t^i\}}\}$ with $\gamma \leq 1/3$ and $w_m = \max_i w_i$.

$$\begin{aligned}
e_1 &\leq \frac{1}{2} \mathbb{E} \|\nabla F_w(x_t) - \nabla F(x_{t-\tau})\|^2 + \frac{1}{2} \mathbb{E} \left\| \frac{1}{B} \sum_{i \in S_t} \sum_{k=0}^{K-1} \alpha q_t^i \nabla f_i(x_k^i) \right\|^2 \\
&\leq \frac{L^2}{2} \mathbb{E} \|x_t - x_{t-\tau}\|^2 + \mathbb{E} \left\| \frac{1}{B} \sum_{i \in S_t} \sum_{k=0}^{K-1} \alpha q_t^i (\nabla f_i(x_k^i) - \nabla f_i(x_t)) \right\|^2 \\
&\quad + \mathbb{E} \left\| \frac{1}{B} \sum_{i \in S_t} \sum_{k=0}^{K-1} \alpha q_t^i \nabla f_i(x_t) \right\|^2 \\
&\leq \frac{L^2}{2} \mathbb{E} \|x_t - x_{t-\tau}\|^2 + K \mathbb{E} \left[\frac{1}{B} \sum_{i \in S_t} \sum_{k=0}^{K-1} (\alpha)^2 L^2 (q_t^i)^2 \|x_k^i - x_t\|^2 \right] \\
&\quad + \mathbb{E} \left\| \frac{1}{B} \sum_{i \in S_t} \sum_{k=0}^{K-1} \alpha q_t^i \nabla f_i(x_t) \right\|^2 \\
&\leq \frac{\tau \gamma^2}{2} \sum_{l=t-\tau}^{t-1} \mathbb{E} \|\nabla F_w(x_l)\|^2 + 2\tau^2 \gamma^2 G^2 + \frac{\gamma^2}{64L^2} \mathbb{E} \|\nabla F_w(x_t)\|^2 + \frac{\gamma^2 G^2}{16L^2} \\
&\quad + \frac{\gamma^2}{64L^2} \mathbb{E} \left\| \frac{1}{B} \sum_{i \in S_t} \nabla f_i(x_t) \right\|^2 \\
&\leq \frac{\tau \gamma^2}{2} \sum_{l=t-\tau}^{t-1} \mathbb{E} \|\nabla F_w(x_l)\|^2 + \left(2\tau^2 + \frac{1}{16L^2} + \frac{1}{8BL^2} \right) \gamma^2 G^2 + \frac{3\gamma^2}{64L^2} \mathbb{E} \|\nabla F(x_t)\|^2
\end{aligned}$$

where we use Lemmas 7 and 8 in the fourth inequality; we use the fact $\mathbb{E} \|\frac{1}{B} \sum_{i \in S_t} \nabla f_i(x_t)\|^2 \leq 2\mathbb{E} \|\nabla F_w(x_t)\|^2 + 8G^2/B$ in the last inequality. Next we turn to bound e_2 . Note that

$$\begin{aligned}
e_2 &= -\alpha K \mathbb{E} \left[\mathbb{E}(\langle \nabla F_w(x_{t-\tau}), \frac{1}{B} \sum_{i \in S_t} q_t^i \nabla f_i(x_{t-\tau}) \rangle \mid \mathcal{F}_{t-\tau}) \right] \\
&= \frac{\alpha K}{2} \mathbb{E} \|\nabla F_w(x_{t-\tau}) - \mathbb{E}(\frac{1}{B} \sum_{i \in S_t} q_t^i \nabla f_i(x_{t-\tau}) \mid \mathcal{F}_{t-\tau})\|^2 - \frac{\alpha K}{2} \mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 \\
&\quad - \frac{\alpha K}{2} \mathbb{E} \|\mathbb{E}(\frac{1}{B} \sum_{i \in S_t} q_t^i \nabla f_i(x_{t-\tau}) \mid \mathcal{F}_{t-\tau})\|^2 \\
&\leq \frac{\alpha K}{2} \mathbb{E} \|\nabla F_w(x_{t-\tau}) - \mathbb{E}(\frac{1}{B} \sum_{i \in S_t} q_t^i \nabla f_i(x_{t-\tau}) \mid \mathcal{F}_{t-\tau})\|^2 - \frac{\alpha K}{2} \mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 \\
&\leq \alpha K \mathbb{E} \|\nabla F_w(x_{t-\tau}) - \mathbb{E}(\frac{1}{B} \sum_{i \in S_t} q_*^i \nabla f_i(x_{t-\tau}) \mid \mathcal{F}_{t-\tau})\|^2 - \frac{\alpha K}{2} \mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 \\
&\quad + \alpha K \mathbb{E} \left\| \frac{1}{B} \sum_{i \in S_t} (q_t^i - q_*^i) \nabla f_i(x_{t-\tau}) \right\|^2
\end{aligned}$$

where $q_*^i = \frac{w_i}{\pi_i}$ and $\mathcal{F}_{t-\tau}$ is the filtration up to $t-\tau$. Next, we provide the bound for $\mathbb{E} \|\nabla F_w(x_{t-\tau}) - \mathbb{E}(\frac{1}{B} \sum_{i \in S_t} q_*^i \nabla f_i(x_{t-\tau}) \mid \mathcal{F}_{t-\tau})\|^2$. Since we are focusing on the case for a particular R , without causing confusion, we drop R for notation simplicity in the following analysis.

Denoting $\psi_S := \lim_{t \rightarrow \infty} P(S_t = S)$, we have

$$\pi^i = \frac{\sum_{\hat{S}_i} \psi_{\hat{S}_i}}{\sum_{i=1}^N \sum_{\hat{S}_i} \psi_{\hat{S}_i}} = \frac{\sum_{\hat{S}_i} \psi_{\hat{S}_i}}{B}$$

where \hat{S}_i denotes any set with size B containing i . Then, for any vectors $\{v_i\}_{i=1}^N$, we have

$$\sum_{S \in \mathcal{S}} \sum_{i \in S} \frac{\psi_S}{\pi^i} v_i = \sum_{i=1}^N \sum_{\hat{S}_i} \frac{\psi_{\hat{S}_i}}{\pi^i} v_i = B \sum_{i=1}^N v_i.$$

where \mathcal{S} is the collection of all sets with size B . Thus, by letting $v_i = w_i \nabla f_i(x_{t-\tau})$ in the above, we obtain

$$\begin{aligned} & \mathbb{E} \|\nabla F_w(x_{t-\tau}) - \mathbb{E} \left(\frac{1}{B} \sum_{i \in S_t} q_*^i \nabla f_i(x_{t-\tau}) \mid \mathcal{F}_{t-\tau} \right) \|^2 \\ &= \mathbb{E} \|\nabla F_w(x_{t-\tau}) - \frac{1}{B} \sum_{S \in \mathcal{S}} \sum_{i \in S} P(S_t = S \mid \mathcal{F}_{t-\tau}) q_*^i \nabla f_i(x_{t-\tau}) \|^2 \\ &= \mathbb{E} \left\| \frac{1}{B} \sum_{S \in \mathcal{S}} \sum_{i \in S} (P(S_t = S \mid \mathcal{F}_{t-\tau}) - \psi_S) q_*^i \nabla f_i(x_{t-\tau}) \right\|^2 \end{aligned}$$

by noting $q_*^i = w_i / \pi^i$. Moreover, $P(S_t = \cdot)$ can be uniquely induced by $\phi_R(t)$ defined by equation 6 under proper linear transformations, which also indicates that $P(S_t = \cdot \mid \mathcal{F}_{t-\tau}) = P(S_t = \cdot \mid S_{t-\tau})$. Thus, Lemma 4 implies $|P(S_t = S \mid \mathcal{F}_{t-\tau}) - \psi_S| \leq c_1 \delta \pi_{\min} / \sqrt{C_N^B}$ for some $c_1 > 0$, $\forall S$ when $\tau \geq \tau_{\text{mix}} \log(1/\delta)$ with $C_N^B = \binom{N}{B}$. Then,

$$\begin{aligned} & \mathbb{E} \left\| \nabla F_w(x_{t-\tau}) - \mathbb{E} \left(\frac{1}{B} \sum_{i \in S_t} q_*^i \nabla f_i(x_{t-\tau}) \mid \mathcal{F}_{t-\tau} \right) \right\|^2 \\ &= \mathbb{E} \left\| \frac{1}{B} \sum_{S \in \mathcal{S}} \sum_{i \in S} (P(S_t = S \mid \mathcal{F}_{t-\tau}) - \psi_S) q_*^i \nabla f_i(x_{t-\tau}) \right\|^2 \\ &\leq \frac{1}{B} \mathbb{E} \left[\sum_{i \in S} \left\| \sum_{S \in \mathcal{S}} (P(S_t = S \mid \mathcal{F}_{t-\tau}) - \psi_S) q_*^i \nabla f_i(x_{t-\tau}) \right\|^2 \right] \\ &\leq c_1^2 \pi_{\min}^2 \delta^2 \mathbb{E} \left[\frac{1}{B} \sum_{i \in S} \left\| q_*^i \nabla f_i(x_{t-\tau}) \right\|^2 \right] \\ &\leq c_1^2 \delta^2 (\mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 + 4G^2) \end{aligned}$$

where we use the fact that

$$\|\nabla f_i(x_{t-\tau})\|^2 \leq 2 \|\nabla F_w(x_{t-\tau})\|^2 + 8G^2.$$

Utilizing the following

$$\begin{aligned} \mathbb{E} \left\| \frac{1}{B} \sum_{i \in S_t} (q_t^i - q_*^i) \nabla f_i(x_{t-\tau}) \right\|^2 &= \mathbb{E} \left\| \frac{1}{B} \sum_{i \in S_t} \tilde{q}_t^i (\nabla f_i(x_{t-\tau}) - \nabla F(x_{t-\tau}) + \nabla F(x_{t-\tau})) \right\|^2 \\ &\leq 8G^2 \mathbb{E} \|\tilde{q}_t\|_\infty^2 + 2\mathbb{E} [\|\tilde{q}_t\|_\infty^2 \|\nabla F(x_{t-\tau})\|^2] \end{aligned}$$

where we denote $\tilde{q}_t^i = q_t^i - q_*^i$. Then we bound e_2 as

$$e_2 \leq \frac{\alpha K}{2} (2c_1^2 \delta^2 - 1) \mathbb{E} \|\nabla F(x_{t-\tau})\|^2 + 2\alpha K G^2 (\delta^2 + 4\mathbb{E} \|\tilde{q}_t\|_\infty^2) + 2\alpha K \mathbb{E} [\|\tilde{q}_t\|_\infty^2 \|\nabla F(x_{t-\tau})\|^2].$$

In order to bound e_3 , note that according to Lemma 7 for $\alpha \leq \frac{\gamma}{8KLa} \leq \frac{\gamma}{8KLq_t^i}$

$$\begin{aligned}
e_3 &\leq \mathbb{E} \left[\frac{1}{B} \sum_{i \in S_t} \sum_{k=0}^{K-1} \alpha q_t^i \|\nabla F(x_{t-\tau})\| \|\nabla f_i(x_k^i) - \nabla f_i(x_t)\| \right] \\
&\leq \mathbb{E} \left[\frac{1}{B} \sum_{i \in S_t} \sum_{k=0}^{K-1} \left(\frac{(\alpha q_t^i)^2 K}{2} \|\nabla F_w(x_{t-\tau})\|^2 + \frac{L^2}{2K} \|x_k^i - x_t\|^2 \right) \right] \\
&\leq \mathbb{E} \left[\frac{1}{B} \sum_{i \in S_t} \sum_{k=0}^{K-1} \left(\frac{\gamma^2}{128L^2K} \|\nabla F_w(x_{t-\tau})\|^2 + \frac{\gamma^2}{2K} (\|\nabla F_w(x_t)\|^2 + 4G^2) \right) \right] \\
&\leq \frac{\gamma^2}{128L^2} \mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 + \frac{\gamma^2}{2} \mathbb{E} \|\nabla F_w(x_t)\|^2 + 2\gamma^2 G^2.
\end{aligned}$$

Finally, based on Lemma 8, similarly we obtain

$$\begin{aligned}
e_4 &\leq \mathbb{E} \left[\frac{1}{B} \sum_{i \in S_t} \sum_{k=0}^{K-1} \alpha q_t^i \|\nabla F(x_{t-\tau})\| \|\nabla f_i(x_t) - \nabla f_i(x_{t-\tau})\| \right] \\
&\leq \mathbb{E} \left[\frac{1}{B} \sum_{i \in S_t} \sum_{k=0}^{K-1} \left(\frac{(\alpha q_t^i)^2 K}{2} \|\nabla F(x_{t-\tau})\|^2 + \frac{L^2}{2K} \|x_t - x_{t-\tau}\|^2 \right) \right] \\
&\leq \frac{\gamma^2}{128L^2} \mathbb{E} \|\nabla F(x_{t-\tau})\|^2 + \frac{\gamma^4 \tau}{2} \left(\sum_{l=t-\tau}^{t-1} \mathbb{E} \|\nabla F_w(x_l)\|^2 + 4\tau G^2 \right).
\end{aligned}$$

Thus, denoting $\bar{a} = a \max_i \{w_i\}$

$$\begin{aligned}
\frac{\gamma}{16\bar{a}L} \mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 &\leq \mathbb{E} [F_w(x_t) - F_w(x_{t+1})] + \frac{\tau\gamma^2(1+\gamma^2)}{2} \sum_{l=t-\tau}^{t-1} \mathbb{E} \|\nabla F_w(x_l)\|^2 \\
&\quad + \left(\frac{\gamma^2}{2} + \frac{\gamma^2}{2L} + \frac{3\gamma^2}{64L^2} \right) \mathbb{E} \|\nabla F_w(x_t)\|^2 + \frac{\gamma}{aL} G^2 \mathbb{E} \|\tilde{q}_t\|_\infty^2 \\
&\quad + \frac{\gamma}{4\bar{a}L} \mathbb{E} [\|\tilde{q}_t\|_\infty^2 \|\nabla F_w(x_{t-\tau})\|^2] + \frac{\gamma c_1^2 \delta^2}{4\bar{a}L} G^2 \\
&\quad + \gamma^2 G^2 \left(2 + 2\tau^2 + 2\gamma^2 \tau^2 + \frac{2}{L} + \frac{3}{16L^2} \right) \\
&\quad + \left(\frac{\gamma \delta^2}{8\bar{a}L} + \frac{\gamma^2}{64L^2} \right) \mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2
\end{aligned}$$

which implies that

$$\begin{aligned}
\gamma \mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 &\leq 16\bar{a}L\mathbb{E}[F_w(x_t) - F_w(x_{t+1})] + \frac{16\bar{a}L\tau\gamma^2(1+\gamma^2)}{2} \sum_{l=t-\tau}^{t-1} \mathbb{E} \|\nabla F_w(x_l)\|^2 \\
&\quad + \gamma \left(16\bar{a}L\gamma + 8\bar{a}\gamma + \frac{3\bar{a}\gamma}{4L} \right) \mathbb{E} \|\nabla F_w(x_t)\|^2 + 4\gamma c_1^2 \delta^2 G^2 \\
&\quad + 4\gamma \mathbb{E} [\|\tilde{q}_t\|_\infty^2 \|\nabla F_w(x_{t-\tau})\|^2] + 16\gamma G^2 \mathbb{E} \|\tilde{q}_t\|_\infty^2 \\
&\quad + 16\bar{a}L\gamma^2 G^2 \left(2 + 2\tau^2 + 2\gamma^2\tau^2 + \frac{2}{L} + \frac{3}{16L^2} \right) \\
&\quad + \gamma \left(2c_1^2 \delta^2 + \frac{\bar{a}\gamma}{4L} \right) \mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 \\
&\leq 16\bar{a}L\mathbb{E}[F_w(x_t) - F_w(x_{t+1})] + \gamma \left(16\bar{a}L\gamma + 8\bar{a}\gamma + \frac{3\bar{a}\gamma}{4L} \right) \mathbb{E} \|\nabla F_w(x_t)\|^2 \\
&\quad + \gamma \left(2c_1^2 \delta^2 + \frac{\bar{a}\gamma}{4L} + 32\bar{a}L\tau\gamma(1+\gamma^2) \right) \mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 \\
&\quad + 4\gamma \mathbb{E} [\|\tilde{q}_t\|_\infty^2 \|\nabla F_w(x_{t-\tau})\|^2] + 16\gamma G^2 \mathbb{E} \|\tilde{q}_t\|_\infty^2 + 4\gamma c_1^2 \delta^2 G^2 \\
&\quad + 16\bar{a}L\gamma^2 G^2 \left(2 + 2\tau^2 + 2\gamma^2\tau^2 + 8\tau^4\gamma^2(1+\gamma^2) + \frac{2}{L} + \frac{3}{16L^2} \right)
\end{aligned}$$

where we make use of

$$\sum_{l=t-\tau}^{t-1} \mathbb{E} \|\nabla F_w(x_l)\|^2 \leq 4\tau \mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 + 16\tau^3 \gamma^2 G^2$$

by Lemma 9. Under the following conditions

$$\begin{aligned}
2c_1^2 \delta^2 &\leq \frac{1}{6}, \quad \frac{\bar{a}\gamma}{4L} \leq \frac{1}{36}, \quad \gamma \leq \min\left\{\frac{1}{2\tau}, \frac{1}{384\bar{a}}\right\}, \\
64\bar{a}L\tau\gamma &\leq \frac{1}{12},
\end{aligned}$$

which implies $32\bar{a}L\tau\gamma(1+\gamma^2) \leq \frac{1}{6}$ and hence $2c_1^2 \delta^2 + \frac{\bar{a}\gamma}{4L} + 32\bar{a}L\tau\gamma(1+\gamma^2) \leq \frac{1}{2}$, then we obtain

$$\begin{aligned}
\gamma \mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 &\leq 32\bar{a}L\mathbb{E}[F_w(x_t) - F_w(x_{t+1})] + 2\gamma \left(16\bar{a}L\gamma + 8\bar{a}\gamma + \frac{3\bar{a}\gamma}{4L} \right) \mathbb{E} \|\nabla F_w(x_t)\|^2 \\
&\quad + 8\gamma \mathbb{E} [\|\tilde{q}_t\|_\infty^2 \|\nabla F_w(x_{t-\tau})\|^2] + 32\gamma G^2 \mathbb{E} \|\tilde{q}_t\|_\infty^2 + 8\gamma c_1^2 \delta^2 G^2 \\
&\quad + 32\bar{a}L\gamma^2 G^2 \left(2 + 2\tau^2 + 2\gamma^2\tau^2 + 8\tau^4\gamma^2(1+\gamma^2) + \frac{2}{L} + \frac{3}{16L^2} \right).
\end{aligned}$$

Summing over $\tau \leq t \leq T-1$ gives

$$\begin{aligned}
\gamma \sum_{t=\tau}^{T-1} \mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 &\leq 32\bar{a}L\Delta_\tau + 2\gamma \left(16\bar{a}L\gamma + 8\bar{a}\gamma + \frac{3\bar{a}\gamma}{4L} \right) \sum_{t=\tau}^{T-1} \mathbb{E} \|\nabla F_w(x_t)\|^2 \\
&\quad + 8\gamma \sum_{t=\tau}^{T-1} \mathbb{E} [\|\tilde{q}_t\|_\infty^2 \|\nabla F_w(x_{t-\tau})\|^2] + 32\gamma G^2 \sum_{t=\tau}^{T-1} \mathbb{E} \|\tilde{q}_t\|_\infty^2 \\
&\quad + 32\bar{a}L\gamma^2 G^2 \left(3 + 6\tau^2 + \frac{2}{L} + \frac{3}{16L^2} \right) (T-\tau) + 8\gamma c_1^2 \delta^2 G^2 (T-\tau).
\end{aligned}$$

where $\Delta_\tau = \mathbb{E}[F_w(x_\tau) - F^*]$ and we use $\gamma^2\tau^2 \leq 1/4$. Again leveraging Lemma 9, we observe

$$\sum_{t=\tau}^{T-1} \mathbb{E} \|\nabla F_w(x_t)\|^2 \leq 4 \sum_{t=\tau}^{T-1} \mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 + 16\tau^2 \gamma^2 G^2 (T-\tau)$$

which thus renders

$$\begin{aligned} \frac{1}{T-\tau} \sum_{t=\tau}^{T-1} \mathbb{E} \|\nabla F_w(x_{t-\tau})\|^2 &\leq \frac{32\bar{a}L\Delta_\tau}{\gamma(T-\tau)} + \frac{8}{T-\tau} \sum_{t=\tau}^{T-1} \mathbb{E} [\|\tilde{q}_t\|_\infty^2 \|\nabla F_w(x_{t-\tau})\|^2] \\ &\quad + 32\bar{a}LG^2 \left(3\gamma + 6\gamma\tau^2 + \frac{2\gamma}{L} + \frac{3\gamma}{16L^2} + \frac{\gamma^2}{16aL} \right) + 8c_1^2\delta^2G^2 \\ &\quad + \frac{32G^2}{T-\tau} \sum_{t=\tau}^{T-1} \mathbb{E} \|\tilde{q}_t\|_\infty^2 \end{aligned}$$

by noting that $16\bar{a}L\gamma + 8a\gamma + \frac{3\bar{a}\gamma}{4L} \leq \frac{1}{16}$.

In the following, we turn to bound Δ_τ . Noting that

$$\begin{aligned} F_w(x_{t+1}) - F_w(x_t) &\leq -\alpha K \langle \nabla F_w(x_t), \frac{1}{BK} \sum_{i \in S_t} \sum_{k=0}^{K-1} \nabla f_i(x_k^i) \rangle + \frac{\alpha^2 L}{2B^2} \left\| \sum_{i \in S_t} \sum_{k=0}^{K-1} \nabla f_i(x_k^i) \right\|^2 \\ &\leq \frac{\alpha K}{2} \left\| \frac{1}{BK} \sum_{i \in S_t} \sum_{k=0}^{K-1} (\nabla f_i(x_k^i) - \nabla F_w(x_t)) \right\|^2 - \frac{\alpha K}{2} \|\nabla F_w(x_t)\|^2 \end{aligned}$$

by $\alpha \leq \frac{\gamma}{8aLK} \leq \frac{1}{2LK}$. Moreover, since

$$\begin{aligned} \left\| \frac{1}{BK} \sum_{i \in S_t} \sum_{k=0}^{K-1} (\nabla f_i(x_k^i) - \nabla F_w(x_t)) \right\|^2 &\leq \frac{2}{BK} \sum_{i \in S_t} \sum_{k=0}^K (L^2 \|x_k^i - x_t\|^2 + 4G^2) \\ &\leq 2\gamma^2 \|\nabla F_w(x_t)\|^2 + 8G^2 \end{aligned}$$

we conclude that

$$F_w(x_{t+1}) - F_w(x_t) \leq -\frac{\alpha K}{2} (1 - 2\gamma^2) \|\nabla F_w(x_t)\|^2 + 4\alpha K G^2 \leq \frac{\gamma}{2\bar{a}L} G^2$$

which implies

$$\Delta_\tau = \mathbb{E}[F_w(x_\tau) - F^*] \leq \frac{\gamma\tau}{2\bar{a}L} G^2 + F_w(x_0) - F_w^*.$$

□

F CONVERGENCE ANALYSIS OF FEDAVG UNDER CORRELATED CLIENT PARTICIPATION

In this section, we provide the convergence analysis of Vanilla FedAvg for correlated client participation. We first show FedAvg suffers from unavoidable bias (stated in Theorem 1), summarized by the following proposition.

Proposition 3. *There exists a problem case such that FedAvg converges with unavoidable asymptotic bias.*

Proof. We consider a problem case with $N = 3, B = 1, R = 1$. We set $p_1 = 0.25, p_2 = 0.25, p_3 = 0.5$ and $f_i(x) = \frac{1}{2}(x - i)^2, i = 1, 2, 3$ and $x \in \mathbb{R}$. In this case, we have the Markov chain induced by the problem denoted by $P \in \mathbb{R}^{3 \times 3}$. Letting $\pi \in \mathbb{R}^3$ be the stationary distribution of P , a straightforward calculation gives $\pi_1 = \pi_2 = 0.3, \pi_3 = 0.4$. Then we obtain the server's update of FedAvg given by

$$x_{t+1} = \beta x_t + (1 - \beta) i_t$$

where $\beta = (1 - \alpha)^K < 1$ with α being the stepsize of local updates; i_t is the index of the sampled client at round t which is a random variable. Taking the expectation on both sides yields

$$\begin{aligned} \mathbb{E}[x_{t+1}] &= \beta \mathbb{E}[x_t] + (1 - \beta) \mu^T P^t I \\ &= \beta \mathbb{E}[x_t] + (1 - \beta) (\mu^T P^t - \pi^T) I + (1 - \beta) \pi^T I \end{aligned}$$

where $\mu = (p_1, p_2, p_3)$, and $I = (1, 2, 3)$ is the vector formed by clients' indices. Noting that the third term vanishes as $t \rightarrow \infty$ due to the convergence of the Markov chain (shown by Lemma 4), we conclude that $\lim_{t \rightarrow \infty} \mathbb{E}[x_t] = \sum_{i=1}^3 \pi_i i$ which is the minimizer of $F_\pi(x) := \sum_{i=1}^3 \pi_i f_i(x)$ but not $F(x) = \frac{1}{3} \sum_{i=1}^3 f_i(x)$. And $|F'(\pi^T I)| = |I^T(\pi - \frac{1}{3} \mathbf{1}_3)|$. Therefore, the bias in Theorem 1 is unavoidable. \square

Then we show the convergence result of FedAvg.

Theorem 4. *Suppose Assumptions 1,2 hold and assume $\|\nabla F(x)\| \leq D, \forall x$. Then, by choosing $\alpha = \mathcal{O}(\frac{\gamma}{K})$ and $T \geq 2\tau_{mix} \log \tau_{mix}$, the output \tilde{x}_T generated by FedAvg satisfies*

$$\begin{aligned} \mathbb{E}\|\nabla F(\tilde{x}_T)\|^2 &= \mathcal{O}\left(\frac{\Delta_0}{\gamma T}\right) + \mathcal{O}\left(\frac{\tau_{mix} \log T G^2}{T}\right) + \mathcal{O}((\gamma \tau_{mix}^2 \log^2 T + \gamma^2) G^2) \\ &\quad + \mathcal{O}\left((G^2 + D^2)\|\pi - \frac{1}{N} \mathbf{1}_N\|_1^2\right) \end{aligned}$$

where $\Delta_0 := \mathbb{E}[F(x_0) - \min_x F(x)]$ and τ_{mix} is the mixing time.

Proof. For FedAvg, we have $y_t^i = 1/N$. Utilizing Lemma 10 and setting $w_i = \frac{1}{N}$, it yields

$$\begin{aligned} \frac{1}{T-\tau} \sum_{t=0}^{T-\tau-1} \mathbb{E}\|\nabla F(x_t)\|^2 &\leq \frac{32L\Delta_0}{\gamma(T-\tau)} + \frac{8D^2}{T-\tau} \sum_{t=\tau}^{T-1} \mathbb{E}\|\tilde{q}_t\|_\infty^2 + \frac{36G^2}{T-\tau} \sum_{t=\tau}^{T-1} \mathbb{E}\|\tilde{q}_t\|_\infty^2 + \frac{16\tau G^2}{T-\tau} \\ &\quad + 32LG^2 \left(3\gamma + 6\gamma\tau^2 + \frac{2\gamma}{L} + \frac{3\gamma}{16L^2} + \frac{\gamma^2}{16L}\right) + 8c_1^2 \delta^2 G^2. \end{aligned}$$

Then noting that $\|\tilde{q}_t\|_\infty^2 \leq \pi_{min}^{-2} \|\pi - \frac{1}{N} \mathbf{1}_N\|_1^2$, we conclude

$$\mathbb{E}\|\nabla F(\tilde{x}_T)\|^2 = \mathcal{O}\left(\frac{\Delta_0}{\gamma T}\right) + \mathcal{O}\left(\frac{\tau G^2}{T}\right) + \mathcal{O}((\gamma \tau^2 + \gamma^2) G^2) + \mathcal{O}\left((G^2 + D^2)\|\pi - \frac{1}{N} \mathbf{1}_N\|_1^2\right)$$

by setting $\delta = 1/\sqrt{T}$. For the above to be true, we need $T \geq \tau = \tau_{mix} \log T$, which is actually always satisfied for $T \geq 2\tau_{mix} \log \tau_{mix}$. To see this, we observe that if $T \leq \tau_{mix}^2$, $\tau_{mix} \log T \leq 2\tau_{mix} \log \tau_{mix}$; if $T \geq \tau_{mix}^2$, $\tau_{mix} \log T \leq \sqrt{T} \log T \leq T$. This completes the proof. \square

The following corollary restates the convergence result of Theorem 1.

Corollary 1. *Suppose all conditions in Theorem 4 hold. Then, choosing $\alpha = \tilde{\mathcal{O}}(1/(K\tau_{mix}\sqrt{T}))$, the output \tilde{x}_T of FedAvg satisfies*

$$\mathbb{E}\|\nabla F(\tilde{x}_T)\|^2 \leq \tilde{\mathcal{O}}\left(\frac{\tau_{mix}}{\sqrt{T}}\right) + \mathcal{O}\left((D^2 + G^2)\|\pi_R - \frac{1}{N} \mathbf{1}_N\|_1^2\right).$$

Proof. The proof is straightforward by simply plugging in $\gamma = \mathcal{O}(1/(\tau\sqrt{T}))$ and $\tau = \tau_{mix} \log T$ to Theorem 4. \square

G CONVERGENCE ANALYSIS OF ALGORITHM 1

We first provide the following theorem showing that y_t^i serves as a reasonable estimation of π^i .

Theorem 5. *For any real-valued function $f \in \mathbb{R}^N$ and any initial distribution $\mu \in \mathbb{R}^N$, we have the following:*

$$\begin{aligned} \mathbb{E}_\mu \left(\frac{1}{T} \sum_{t=0}^{T-1} f(X_t) - \pi_R^T f \right) &= \frac{1}{T} \sum_{t=0}^{T-1} \mu^T Q_\mu^\dagger (P_R^t - \mathbf{1} \zeta_R^T) Q_R f \\ T \mathbb{E}_{\pi_R} \left(\frac{1}{T} \sum_{t=0}^{T-1} f(X_t) - \pi_R^T f \right)^2 &\leq f^T \Pi_R (I - \mathbf{1}_N \pi_R^T) f + c_0 \pi_{\max} \|f\|_\infty^2 N \tau_{mix} \\ T \mathbb{E}_\mu \left(\frac{1}{T} \sum_{t=0}^{T-1} f(X_t) - \pi_R^T f \right)^2 &\leq T \mathbb{E}_{\pi_R} \left(\frac{1}{T} \sum_{t=0}^{T-1} f(X_t) - \pi_R^T f \right)^2 + 3c_0 N^2 \|g\|_\infty^2 \tau_{mix} \end{aligned}$$

where $\mathbb{E}_\mu(\cdot)$ means the initial state X_0 follows μ ; $\Pi_R = \text{diag}(\pi_R[i])$ and Q_μ^\dagger is defined such that $\mu^T Q_\mu^\dagger = \zeta_\mu$ and $\zeta_\mu^T Q_R = \mu$; $g = f - \pi_R^T f \mathbf{1}_N$; τ_{mix} is the mixing time of P_R .

Proof. We firstly show the first equality. Note that

$$\begin{aligned}\mathbb{E}_\mu \left(\frac{1}{T} \sum_{t=0}^{T-1} f(X_t) - \pi_R^T f \right) &= \frac{1}{T} \sum_{k=0}^{T-1} (\mu^T P_R^k Q_R f - \zeta_R^T Q_R f) \\ &= \frac{1}{T} \sum_{k=0}^{T-1} \mu^T (P_R^k - \mathbf{1} \zeta_R^T) Q_R f\end{aligned}$$

where we observe that $\mu^T \mathbf{1} = 1$.

Then we turn to show the second inequality. By the definition, we have

$$T \mathbb{E}_{\pi_R} \left(\frac{1}{T} \sum_{t=0}^{T-1} f(X_t) - \pi_R^T f \right)^2 = \text{Var}_{\pi_R}(f(X_0)) + \frac{2}{T} \sum_{k=1}^{T-1} (T-k) \text{Cov}_{\pi_R}(f(X_0), f(X_k)). \quad (11)$$

For any k , let ζ_k and π_k be the distributions after the Markov chain evolves k steps. Then, we have $\zeta_{k+1}^T = \zeta_k^T P_R$ and $\pi_k^T = \zeta_k^T Q_R$. Defining $\hat{Q}_k \in \mathbb{R}^{N \times d(M,R)}$ as an inverse mapping from π_k to ζ_k , i.e., $\zeta_k^T = \pi_k^T \hat{Q}_k$, it is straightforward to verify that we can always pick a nonnegative \hat{Q}_k such that $\hat{Q}_k \mathbf{1} = \mathbf{1}_N$ in the sense that the freedom of \hat{Q}_k is $(N-1) \times d(M,R) - N$ when forcing both $\zeta_k^T = \pi_k^T \hat{Q}_k$ and $\hat{Q}_k \mathbf{1} = \mathbf{1}_N$ to hold. Moreover,

$$\begin{aligned}\text{Cov}_{\pi_R}(f(X_0), f(X_k)) &= \sum_i \pi_R[i] f(i) \sum_j [\hat{Q}_k P_R^k Q_R]_{i,j} f(j) - \sum_{i,j} \pi_R[i] \pi_R[j] f(i) f(j) \\ &= f^T \Pi_R \hat{Q}_k P_R^k Q_R f - f^T \Pi_R \mathbf{1}_N \pi_R^T f \\ &= f^T \Pi_R \hat{Q}_k (P_R^k - \mathbf{1} \zeta_R^T) Q_R f\end{aligned}$$

where we utilize $\hat{Q}_k \mathbf{1} = \mathbf{1}_N$. Further, $\|\hat{Q}_k\|_\infty = 1, \forall k \geq 0$ since \hat{Q}_k is nonnegative. Then,

$$\text{Cov}_{\pi_R}(f(X_0), f(X_k)) \leq \pi_{\max} \|f\|_\infty^2 \|Q_R\|_\infty \|P_R^k - \mathbf{1} \zeta_R^T\|_\infty.$$

Substituting it into equation 11 yields

$$\begin{aligned}T \mathbb{E}_{\pi_R} \left(\frac{1}{T} \sum_{t=0}^{T-1} f(X_t) - \pi_R^T f \right)^2 &\leq \text{Var}_{\pi_R}(f(X_0)) + 2 \sum_{k=1}^{\infty} \text{Cov}_{\pi_R}(f(X_0), f(X_k)) \\ &\leq \text{Var}_{\pi_R}(f(X_0)) + 2 \pi_{\max} \|f\|_\infty^2 \|Q_R\|_\infty \sum_{k=0}^T \|P_R^k - \mathbf{1} \zeta_R^T\|_\infty \\ &\leq f^T \Pi_R (I - \mathbf{1}_N \pi_R^T) f + c_0 \pi_{\max} \|f\|_\infty^2 \|Q_R\|_\infty \tau_{mix}\end{aligned}$$

where we make use of Lemma 5. Finally noting that $\|Q_R\|_\infty \leq \|Q_{R,1}\|_\infty \|Q_{R,2}\|_\infty \leq N$ completes the proof of the second inequality.

To obtain the third inequality, defining $g(i) = f(i) - \pi_R^T f$ we aim to bound

$$\begin{aligned}T \left| \mathbb{E}_\mu \left(\frac{1}{T} \sum_{k=0}^{T-1} g(X_k) \right)^2 - \mathbb{E}_{\pi_R} \left(\frac{1}{T} \sum_{k=0}^{T-1} g(X_k) \right)^2 \right| \\ \leq \left| \frac{1}{T} \sum_{k=0}^{T-1} \mathbb{E}_\mu g^2(X_k) - \mathbb{E}_{\pi_R} g^2(X_k) \right| + \frac{2}{T} \sum_{k=0}^{T-1} \sum_{l=k+1}^{T-1} \left| \mathbb{E}_\mu(g(X_k)g(X_l)) - \mathbb{E}_{\pi_R}(g(X_k)g(X_l)) \right|.\end{aligned}$$

For notation simplicity, we drop the subscript R without confusion to get

$$\begin{aligned}
& |\mathbb{E}_\mu(g(X_k)g(X_l)) - \mathbb{E}_{\pi_R}(g(X_k)g(X_l))| \\
&= \left| \sum_{i,j} \mu_i g(j) ((\hat{Q}_k P^k Q)_{i,j} - \pi_j) \sum_r ((\hat{Q}_l P^{l-k} Q)_{j,r} - \pi_r) g(r) \right| \\
&= \left| \sum_{i,j} \mu_i g(j) (\hat{Q}_k (P^k - \mathbf{1}\zeta^T) Q)_{i,j} \sum_r (\hat{Q}_l (P^{l-k} - \mathbf{1}\zeta^T) Q)_{j,r} g(r) \right| \\
&\leq \|g\|_\infty^2 N^2 \|P^l - \mathbf{1}\zeta^T\|_\infty.
\end{aligned}$$

Thus, by Lemma 5,

$$\begin{aligned}
& T \left| \mathbb{E}_\mu \left(\frac{1}{T} \sum_{k=0}^{T-1} g(X_k) \right)^2 - \mathbb{E}_{\pi_R} \left(\frac{1}{T} \sum_{k=0}^{T-1} g(X_k) \right)^2 \right| \\
&\leq \frac{1}{T} \sum_{k=0}^{T-1} \mu^T Q_\mu^\dagger (P^k - \mathbf{1}\zeta^T) Q g^2 + \frac{2}{T} c_0 N^2 \|g\|_\infty^2 \sum_{k=0}^{T-1} \tau_{mix} \\
&\leq \frac{1}{T} \sum_{k=0}^{T-1} \mu^T Q_\mu^\dagger (P^k - \mathbf{1}\zeta^T) Q g^2 + 2c_0 N^2 \|g\|_\infty^2 \tau_{mix} \\
&\leq \frac{1}{T} c_0 \|g\|_\infty^2 N \tau_{mix} + 2c_0 N^2 \|g\|_\infty^2 \tau_{mix} \\
&\leq 3c_0 N^2 \|g\|_\infty^2 \tau_{mix}
\end{aligned}$$

where g^2 denotes the elementwise square of g . Combining all the above completes the proof. \square

Then, the following corollary induced by Theorem 5 is exactly Lemma 2.

Corollary 2. *Given initial $\lambda_0 = \mathbf{0}_N$ and let $\nu_t^i = \frac{1}{\lambda_t^i N}$ as in Algorithm 1, we have*

$$\mathbb{E} \|\tilde{\nu}_t\|_\infty^2 \leq \mathcal{O} \left(\frac{\tau_{mix}}{t} \right)$$

where $\tilde{\nu}_t^i = \nu_t^i - \frac{1}{\pi_i N}$ and $\tilde{\nu}_t = (\tilde{\nu}_t^1, \dots, \tilde{\nu}_t^N)$.

Proof. By Theorem 5, setting $f = \mathbf{e}_i$ for any i , we have

$$\mathbb{E}(\lambda_t^i - \pi_i)^2 = \mathcal{O} \left(\frac{N^2 \tau_{mix}}{t} \right) \quad (12)$$

Note that

$$\begin{aligned}
\mathbb{E}(\tilde{\nu}_t^i)^2 &= \frac{1}{N^2} \mathbb{E} \left(\frac{\lambda_t^i - \pi_i}{\lambda_t^i \pi_i} \right)^2 = \frac{1}{N^2} \mathbb{E} \left[\left(\frac{\lambda_t^i - \pi_i}{\lambda_t^i \pi_i} \right)^2 \mid \lambda_t^i \geq a \right] P(\lambda_t^i \geq a) \\
&\quad + \frac{1}{N^2} \mathbb{E} \left[\left(\frac{\lambda_t^i - \pi_i}{\lambda_t^i \pi_i} \right)^2 \mid \lambda_t^i < a \right] P(\lambda_t^i < a) \quad (13)
\end{aligned}$$

for any positive a . Moreover, due to the Markov chain in Section 3 is irreducible by Lemma 1, every client will be visited infinitely as t goes to infinite, which then implies there always exists some strictly positive constant a_0 independent of t such that $\lambda_t^i \geq a_0 > 0$ almost surely for any $i \in [N]$. Combining equation 12, equation 13 we conclude

$$\mathbb{E} \|\tilde{\nu}_t^i\|_\infty^2 = \mathcal{O} \left(\frac{\tau_{mix}}{t} \right).$$

\square

G.1 CONVERGENCE PROOF OF ALGORITHM 1

The following lemma is useful to derive the convergence proof of Algorithm 1.

Lemma 11. *Supposing that the stochastic scalar sequence $\mathbb{E}[U_1(t)^2] \leq u(t)$ with u being a monotonically decreasing positive function w.r.t. t and assuming that $U_1(t) \leq \bar{u} < \infty$ almost surely, then given any $\delta, \epsilon > 0$, for all $t \geq \inf\{t_0 \mid u(t_0)/\delta^2 \leq \epsilon/\bar{u}^2\}$ and stochastic scalar sequence $U_2(t)$,*

$$\mathbb{E}[U_1(t)^2 U_2(t)] \leq (\epsilon + \delta^2) \mathbb{E}[U_2(t)].$$

Proof. For any $\delta > 0$, we have for all $t \geq \inf\{t_0 \mid u(t_0)/\delta^2 \leq \epsilon/\bar{u}^2\}$

$$\begin{aligned} \mathbb{E}[U_1(t)^2 U_2(t)] &= P(U_1(t) > \delta) \mathbb{E}[U_1(t)^2 U_2(t) \mid U_1(t) > \delta] + P(U_1(t) \leq \delta) \mathbb{E}[U_1(t)^2 U_2(t) \mid U_1(t) \leq \delta] \\ &\leq P(U_1(t) > \delta) \bar{u}^2 \mathbb{E}[U_2(t)] + \delta^2 \mathbb{E}[U_2(t)] \\ &\leq (\epsilon + \delta^2) \mathbb{E}[U_2(t)] \end{aligned}$$

where we use the Markov inequality in the last step, i.e.,

$$P(U_1(t) > \delta) \leq P(U_1(t)^2 > \delta^2) \leq \frac{u(t)}{\delta^2}.$$

□

Then we are ready to provide the proof for Theorem 3.

Proof of Theorem 3: As discussed in the proof of Corollary 2, we know that there exists a positive a^{-1} which lower bounds each λ_t^i for all t almost surely, implying that $\tilde{\nu}_t^i \leq \frac{1}{N}(a + \pi_{\min}^{-1})$. Then for any $t > \tau > c'\tau_{\text{mix}}$ (with c' being some constant), we have

$$\mathbb{E}[\|\tilde{\nu}_t\|_\infty^2 \|\nabla F(x_{t-\tau})\|^2] \leq \frac{1}{16} \mathbb{E}\|\nabla F(x_{t-\tau})\|^2$$

by Lemmas 2 and 11. Further Utilizing Lemma 10 with setting $w_i = \frac{1}{N}$, we obtain

$$\begin{aligned} \frac{1}{T-\tau} \sum_{t=\tau}^{T-1} \mathbb{E}\|\nabla F(x_{t-\tau})\|^2 &\leq \frac{64\bar{a}L\Delta_0}{\gamma(T-\tau)} + \frac{64G^2}{T-\tau} \sum_{t=\tau}^{T-1} \mathbb{E}\|\tilde{\nu}_t\|_\infty^2 + \frac{32\tau G^2}{T-\tau} + 16c_1^2\delta^2G^2 \\ &\quad + 64\bar{a}LG^2 \left(3\gamma + 6\gamma\tau^2 + \frac{2\gamma}{L} + \frac{3\gamma}{16L^2} + \frac{\gamma^2}{16aL} \right) \end{aligned}$$

for $\tau \geq \tau_{\text{mix}} \max\{c', \log(1/\delta)\}$. Similar to the proofs of Theorem 4, setting $\delta = 1/\sqrt{T}$, with $T \geq c'\tau_{\text{mix}} \log \tau_{\text{mix}}$ for some constant c' , we finally conclude that

$$\mathbb{E}\|\nabla F(\tilde{x}_T)\|^2 = \tilde{\mathcal{O}}\left(\frac{\tau_{\text{mix}}}{\sqrt{T}}\right) + \mathcal{O}\left(\frac{1}{T}\right)$$

by choosing $\gamma = \mathcal{O}(1/(\tau\sqrt{T}))$ with $\tau = \Omega(\tau_{\text{mix}} \log T)$ and by leveraging the fact that $\sum_{t=\tau}^{T-1} \mathbb{E}\|\tilde{\nu}_t\|_\infty^2 = \mathcal{O}(\tau_{\text{mix}} \log T)$ implied by Lemma 2. This completes the proof.

H ADDITIONAL EXPERIMENTS

In this section, we compare Vanilla FedAvg, Debiasing FedAvg (ours) and FedVARP under the CIFAR10 dataset given the same participation pattern as in Section 6. Each client maintains a CNN with three convolution layers. Learning rates for three algorithms are selected to be with the order of $\mathcal{O}(10^{-3})$. In Figure 4, Debiasing FedAvg achieves the highest training accuracy due to its debiasing nature as shown in Theorem 3, while Vanilla FedAvg and FedVARP suffer from bias. Moreover, in Table 1, the training and test accuracies for different R are presented, where one can see that Debiasing FedAvg achieves the best performance.




Figure 4: Experiments on CIFAR10. The training accuracies of Vanilla FedAvg, Debiasing FedAvg (ours) and FedVARP are compared given different values of R . The results show that Debiasing FedAvg achieves the highest accuracy and outperforms the other two, since FedAvg and FedVARP suffer from bias.

Algorithms	$R = 5$		$R = 10$		$R = 15$	
	Train acc	Test acc	Train acc	Test acc	Train acc	Test acc
FedAvg	74.9%	67.6%	75.3%	68.8%	75.6%	70.1%
FedVARP	76.7%	68.0%	77.5%	69.4%	79.5%	72.5%
Debiasing FedAvg	79.3%	73.8%	81.5%	74.9%	82.9%	74.1%

Table 1: Training and test accuracies for different R under CIFAR10

I THE INFLUENCE OF R ON CONVERGENCE RATES

In this section, we discuss the effect of different values of R on the convergence rates of Debiasing FedAvg and Vanilla FedAvg as observed empirically in Figure 3. We simulate the "effective" client sampling distribution (i.e., $\eta_R(t)$) as time evolves for different minimum separation R , where we set $N = 100, B = 1$. The code for all experiments can be found through https://github.com/Starrskyy/debias_f1. Figure 5 shows the total variation distance of the evolution of client sampling distributions to their corresponding stationary π_R 's. Clearly increasing R , the convergence rate of "effective" client sampling distribution to the stationary distribution also increases, implying the decrease of mixing time τ_{mix} (see Appendix B for details). Combining this observation together with Theorems 1 and 3 leads to that larger R implies faster convergence rate, which then consistently explains the observation in Figure 3. However, the above explanation is only from an empirical perspective. More rigorous explanations need theoretical advance in the convergence results to reveal explicitly the relation between the rates and values of R .




Figure 5: Convergence of client sampling distribution to π_R for different R ($N = 100, B = 1$).