

# Scalable Second-order Riemannian Optimization for $K$ -means Clustering

Peng Xu<sup>\*1</sup>, Chun-Ying Hou<sup>\*2</sup>, Xiaohui Chen<sup>3</sup>, and Richard Y. Zhang<sup>2</sup>

<sup>1</sup>Department of Statistics, University of Illinois at Urbana-Champaign  
`pengxu1@illinois.edu`

<sup>2</sup>Department of Electrical and Computer Engineering, University of Illinois at Urbana-Champaign  
`{cyhou2,ryz}@illinois.edu`

<sup>3</sup>Department of Mathematics, University of Southern California  
`xiaohuic@usc.edu`

September 29, 2025

## Abstract

Clustering is a hard discrete optimization problem. Nonconvex approaches such as low-rank semidefinite programming (SDP) have recently demonstrated promising statistical and local algorithmic guarantees for cluster recovery. Due to the combinatorial structure of the  $K$ -means clustering problem, current relaxation algorithms struggle to balance their constraint feasibility and objective optimality, presenting tremendous challenges in computing the second-order critical points with rigorous guarantees. In this paper, we provide a new formulation of the  $K$ -means problem as a smooth unconstrained optimization over a submanifold and characterize its Riemannian structures to allow it to be solved using a second-order cubic-regularized Riemannian Newton algorithm. By factorizing the  $K$ -means manifold into a product manifold, we show how each Newton subproblem can be solved in linear time. Our numerical experiments show that the proposed method converges significantly faster than the state-of-the-art first-order nonnegative low-rank factorization method, while achieving similarly optimal statistical accuracy.

## 1 Introduction

Clustering is a cornerstone of modern unsupervised learning, where the goal is to group similar observations into meaningful clusters. The problem is commonly approached through the  $K$ -means formulation, which seeks to partition  $n$  data points  $X_1, X_2, \dots, X_n \in \mathbb{R}^d$  into  $K$  disjoint groups  $G_1, \dots, G_K$  by maximizing the total intra-cluster similarity:

$$\max_{G_1, \dots, G_K} \left\{ \sum_{k=1}^K \frac{1}{|G_k|} \sum_{i,j \in G_k} \langle X_i, X_j \rangle : \bigsqcup_{k=1}^K G_k = [n] \right\}. \quad (1)$$

Here, the inner product  $\langle X_i, X_j \rangle = X_i^\top X_j$  is used to measure pairwise similarity,  $|G_k|$  denotes the cardinality of  $G_k$ , and  $\sqcup$  denotes disjoint union. Most common algorithms for  $K$ -means clustering, including Lloyd's algorithm [31] and spectral clustering [36, 32], can be understood as heuristics for finding “good enough” solutions to the discrete optimization (1). These methods do not come with any guarantees of local optimality,

---

<sup>\*</sup>These authors contributed equally.

let alone global optimality. Indeed, it is commonly argued that globally solving (1) is NP-hard in the worst-case [19, 3], and would lead to statistically meaningless clustering that overfits the data.

Yet in average-case regimes, globally solving the  $K$ -means optimization problem (1) can be both computationally tractable as well as statistically optimal. In particular, when the data  $X_1, \dots, X_n$  arise from a Gaussian mixture model with sufficiently well-separated components, Chen and Yang [17] showed that a well-known semidefinite programming (SDP) relaxation of Peng and Wei [38], written

$$\max_{Z \in \mathbb{R}^{n \times n}} \left\{ \langle XX^\top, Z \rangle + \mu \sum_{i,j} \log(Z_{i,j})_+ : Z\mathbf{1}_n = \mathbf{1}_n, \text{tr}(Z) = K, Z \succeq 0 \right\}, \quad (2)$$

where  $X = [X_1, \dots, X_n]^\top$  and  $\log(Z_{i,j})_+ := \log(\max\{Z_{i,j}, 0\})$ , is guaranteed to compute the globally optimal clusters  $G_1^*, \dots, G_K^*$  for (1) in the limit  $\mu \rightarrow 0^+$  in polynomial time, that in turn recover the ground truth partitions. Note that the formulation (2) is equivalent to the standard  $K$ -means SDP formulation with the elementwise nonnegativity constraint  $Z_{i,j} \geq 0$  in Peng and Wei [38] and Chen and Yang [17] (see Appendix A for more discussions). Moreover, this recovery occurs as soon as the separation between the clusters is large enough for it to be possible. Put in another way, if solving (2) does not recover the ground truth partitions, then the clusters are too closely spaced in a way that makes recovery inherently impossible in an information-theoretic limit sense, see Section 2.1 for more details.

Unfortunately, the SDP (2) is not a practical means of solving (1) to global optimality, due to its need to optimize over an  $n \times n$  matrix to cluster  $n$  samples. Following Burer and Monteiro [13] and Boumal, Voroninski, and Bandeira [11], a natural alternative is to factor  $Z = UU^\top$  into its  $n \times r$  factor matrix  $U$  for rank parameter  $r \geq K$ , impose the logarithmic penalty over  $U$  instead of  $Z$ , and then directly optimize over  $U$ :

$$\max_{U \in \mathbb{R}^{n \times r}} \left\{ \langle XX^\top, UU^\top \rangle + \mu \sum_{i,j} \log(U_{i,j})_+ : UU^\top \mathbf{1}_n = \mathbf{1}_n, \text{tr}(UU^\top) = K \right\}. \quad (3)$$

This reduces the number of variables and constraints from  $O(n^2)$  down to  $O(n)$ , but at the cost of giving up the convexity of the SDP. In general, we can at best hope to compute critical points, which may be spurious local minima or saddle points. The core motivation for our approach, and the impetus for this paper, is the surprising empirical observation that all second-order critical points are global optima in this setting; this is formalized as the following assumption.

**Assumption 1** (Benign nonconvexity). *In the average-case regime when (2) globally solves (1), all approximate second-order critical points in (3) are within a neighborhood of a global optimum.*

The phenomenon of benign nonconvexity is well-documented in the *unconstrained* version—optimizing over semidefinite  $Z \succeq 0$  by factorizing  $Z = UU^\top$ —dating back to the early works of Burer and Monteiro [13]. In contrast, it is rarely seen in our *nonnegative* variant, which adds the elementwise constraint  $U \geq 0$  to enforce doubly nonnegativity in  $Z = UU^\top$ . Despite a superficial similarity, the two formulations differ in fundamental ways, with the nonnegative case known to admit numerous spurious critical points; see Section 1.2 for some classic and recent examples. Nevertheless, we consistently observe that all second-order critical points correspond to global optima, that in turn successfully recover the optimal clusters.

## 1.1 Contributions: Cheap and fast convergence to second-order critical points

Under Assumption 1, globally solving the  $K$ -means optimization problem (1) reduces to that of computing a second-order critical point for (3). Unfortunately, in the constrained nonconvex setting, there is no general-purpose algorithm that is rigorously guaranteed to compute a second-order critical point. The core issue is the need to maintain *feasibility*, i.e. for each iterate  $U$  to satisfy the nonconvex constraints  $UU^\top \mathbf{1}_n = \mathbf{1}_n$  and  $\text{tr}(UU^\top) = K$ , while making progress towards optimality. General-purpose solvers like `fmincon` [14] and `knitr` [15] promise convergence only to critical points of an underlying merit function, which may be infeasible for the original problem. Augmented Lagrangian methods guarantee convergence only to first-order critical points, and only when starting within a local neighborhood [47]. This is a significant

departure from the unconstrained nonconvex setting, where a diverse range of algorithms—both cheap first-order algorithms like gradient descent, as well as rapidly-converging second-order methods like trust-region Newton’s method—globally converge to a second-order critical point starting from any initial point.

Our first contribution is to present an interpretation of (3) as a *smooth unconstrained optimization over a Riemannian manifold*. This allows the immediate benefit of extending the wide array of unconstrained optimization algorithms to the constrained setting, as well as their accompanying guarantees for first- and second-order optimality. For the first time in the context of  $K$ -means, we open the possibility to guarantee global convergence to first- and second-order optimality.

Our second contribution is to show that *second-order Riemannian algorithms can be implemented with linear per-iteration costs* with respect to the number of samples  $n$ . In other words, of all practical algorithms to compute second-order critical points, we show that the one with the best iteration complexity (second-order methods) can be improved to have the same per-iteration costs as first-order methods. Our final algorithm computes  $\epsilon$  second-order points in  $n \cdot \epsilon^{-3/2} \cdot \text{poly}(r, d)$  time.

## 1.2 Related work

Benign nonconvexity in the unconstrained Burer–Monteiro factorization  $Z = UU^\top$  has been empirically observed since the early 2000s [13], and widely exploited in nonconvex low-rank algorithms in machine learning. In the past decade, theory has been developed to explain this phenomenon under some specialized settings [6, 23, 5, 10, 22]. Unfortunately, these guarantees tend to be conservative in the number of samples or the level of noise; they capture the general phenomenon but cannot rigorously explain what is broadly observed in practice.

In contrast, the *nonnegative* Burer–Monteiro factorization  $Z = UU^\top$  with  $U \geq 0$  is widely understood *not* to exhibit benign nonconvexity. To give two simple examples, the functions  $f(U) = \langle SU, U \rangle$  and  $f(U) = \|UU^\top - U_*U_*^\top\|_F^2$  are easily confirmed to exhibit benign nonconvexity over  $U \in \mathbb{R}^{n \times r}$ . But imposing  $U \geq 0$  causes spurious local minima to proliferate; this is unsurprising because both problems, namely copositive testing [35] and complete positive testing [20], are well-known to be NP-hard. For a more sophisticated example, the function  $f(U) = \|\mathcal{A}(UU^\top - U_*U_*^\top)\|^2$  is well known to exhibit benign nonconvexity when the linear operator  $\mathcal{A} : S^n(\mathbb{R}) \rightarrow \mathbb{R}^m$  satisfies the restricted isometry property (RIP) [6]. In this context, a recent arXiv preprint [45] gave a strong counterexample for the equivalent statement over  $U \geq 0$ .

Therefore, even though  $K$ -means is widely known to admit a nonnegative Burer–Monteiro reformulation [38], there have been only two prior works that actually follow this approach, to the best of our knowledge. Neither of these can rigorously guarantee global optimality under Assumption 1. The first is the first-order Riemannian method introduced by Carson et al. [16]. It solves the following:

$$\min_{U \in \mathcal{M}'} \left\{ -\langle XX^\top, UU^\top \rangle + \lambda \|U_-\|_F^2 \right\} \quad (4)$$

where  $\mathcal{M}' := \{U \in \mathbb{R}^{n \times K} : U^\top U = I_K, UU^\top \mathbf{1}_n = \mathbf{1}_n\}$ ,  $U_- = \max\{-U, 0\}$  is the (entrywise) negative part of  $U$  and  $\lambda \geq 0$  is the penalty parameter for  $U \geq 0$ . Although superficially similar, their approach fundamentally lacks a convergence guarantee to a second-order critical point, due to: (i) their nonsmooth objective; (ii) their use of a smooth penalty, which cannot truly enforce feasibility  $U \geq 0$ ; (iii) their use of a first-order method, which can get trapped at a saddle point. Moreover, their manifold is geometrically complicated, necessitating an expensive retraction that costs  $O(n^2)$  time, which prevents their method from scaling to large datasets.

The second work is the nonnegative low-rank (NLR) method of Zhuang et al. [47]. This is a simple projected gradient descent that directly projects  $U$  onto the nonnegative spherical constraint and deals the row sum constraint  $UU^\top \mathbf{1}_n = \mathbf{1}_n$  via the augmented Lagrangian method. It is a first-order primal-dual method that can only achieve local linear convergence in a neighborhood of its global solution. Like Carson et al. [16], it is unclear whether there is a pathway that this algorithm can lead to a global optimality guarantee, or even to second-order optimality.

## 2 Background

### 2.1 SDP relaxation of $K$ -means

Despite the worst-case NP-hardness of the  $K$ -means clustering optimization problem (1), common practical heuristics and relaxed formulations like Lloyd's algorithm [31], spectral clustering [36, 32], nonnegative matrix factorization (NMF) [24, 28, 42] and SDPs [38, 40, 21, 18] work surprisingly well at solving it for real-world data. To explain this discrepancy between theory and practice, suppose that the data  $X_1, \dots, X_n \in \mathbb{R}^d$  are generated from a standard Gaussian mixture model (GMM)

$$X_i = \mu_k + \varepsilon_i, \quad \varepsilon_i \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, \sigma^2 I_d), \quad \text{for } i \in G_k^*, \quad (5)$$

where  $G_k^*$  denotes the ground truth clusters. Chen and Yang [17] proved that the SDP (2) of Peng and Wei [38] (as  $\mu \rightarrow 0^+$ ) achieves a *sharp phase transition* on the separation of centroids for the clustering problem, in any dimension  $d$  and sample size  $n$ . Let

$$\bar{\Theta}^2 := 4\sigma^2 \left( 1 + \sqrt{1 + \frac{Kd}{n \log n}} \right) \log n, \quad (6)$$

and  $\Theta_{\min} := \min_{1 \leq j < k \leq K} \|\mu_j - \mu_k\|$  be the minimum centroid separation. Assume that  $m = n/K$  is an integer without loss of generality and consider any  $\alpha > 0$ . As soon as the exact recovery becomes possible in the regime  $\Theta_{\min} \geq (1 + \alpha)\bar{\Theta}$ , the SDP approach (2) solves the  $K$ -means problem without clustering error with high probability. For precise statements on the information-theoretic threshold, please refer to Theorem 3 in Appendix B. As an immediate consequence of the global optimality guarantee of the  $K$ -means SDP in (2), we deduce that the global solution of the nonconvex low-rank SDP in (3) solves the  $K$ -means clustering problem in (1) in the exact recovery regime.

Next, from the membership matrix  $Z$ , we would like to convert it to the cluster label.

**Lemma 1.** *Let  $Z = Z^\top \in \mathbb{R}^{n \times n}$  be the symmetric block diagonal matrix defined by  $Z_{ij} = 1/|G_k|$  if  $i, j \in G_k$ , and  $Z_{ij} = 0$  otherwise. Then for any integer  $r \in [K, n]$ , there is a unique (up to column permutation)  $U \in \mathbb{R}_+^{n \times K}$  such that  $Z = UU^\top$ . Moreover,  $U$  can be recovered from any  $\hat{U} \in \mathbb{R}^{n \times r}$  satisfying  $Z = \hat{U}\hat{U}^\top$  in  $n \cdot \text{poly}(r)$  time.*

For each block diagonal membership matrix  $Z$ , the unique  $U \in \mathbb{R}_+^{n \times K}$  in Lemma 1 is the associated group assignment matrix, i.e. the  $k$ -th column of  $U$  provides a one-hot encoding of membership in the  $k$ -th cluster.

### 2.2 Critical points in constrained optimization

The problems considered in this paper are instances of the following

$$\begin{aligned} \min_U \quad & f(U), \\ \text{s.t.} \quad & U \in \mathcal{M} := \{U \in \mathbb{R}^{n \times r} : \mathcal{A}(UU^\top) + \mathcal{B}(U) = c\}, \end{aligned} \quad (7)$$

where the linear operators  $\mathcal{A} : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^m$  and  $\mathcal{B} : \mathbb{R}^{n \times r} \rightarrow \mathbb{R}^m$  and right-hand side  $c \in \mathbb{R}^m$  together are assumed to satisfy the independence constraint qualification (LICQ)

$$2[\mathcal{A}^\top(y)]U + \mathcal{B}^\top(y) = \mathbf{0} \iff y = \mathbf{0} \quad \forall U \in \mathcal{M}. \quad (8)$$

In this context,  $U \in \mathbb{R}^{n \times r}$  is said to be *feasible* if it satisfies  $U \in \mathcal{M}$ . The feasible point  $U$  is an  $\epsilon$ -first-order critical point if it satisfies

$$\text{exists } y \in \mathbb{R}^m \quad \text{s.t.} \quad \|\nabla f(U) + 2[\mathcal{A}^\top(y)]U + \mathcal{B}^\top(y)\| \leq \epsilon, \quad (9)$$

and an  $\epsilon$ -second-order critical point if it additionally satisfies

$$\langle \nabla^2 f(U) + 2[\mathcal{A}^\top(y)], \dot{U} \dot{U}^\top \rangle \geq \sqrt{\epsilon} \|\dot{U}\|^2 \quad \forall \dot{U} \in T_U \mathcal{M} \quad (10)$$

over the *tangent space* of  $\mathcal{M}$  at the point  $U$ , given by  $T_U \mathcal{M} = \{\dot{U} \in \mathbb{R}^{n \times r} : \mathcal{A}(U \dot{U}^\top + \dot{U} U^\top) + \mathcal{B}(\dot{U}) = 0\}$ . Under LICQ (8), every local minimum (and hence the global minimum) is guaranteed to be an  $\epsilon$ -second-order critical point (for any  $\epsilon \geq 0$ ). Unfortunately, there is no general-purpose algorithm that is guaranteed to converge to a critical point, due to the need to achieve and maintain feasibility across all iterates.

### 2.3 Second-order Riemannian optimization

Riemannian algorithms are special algorithms that maintain feasible iterates through a problem-specific *retraction* operator, and are hence able to rigorously guarantee convergence to critical points. The basic idea is to improve a feasible iterate  $U \in \mathcal{M}$  by tracing a smooth curve on the feasible set  $\gamma : [0, \epsilon) \rightarrow \mathcal{M}$  that begins at  $\gamma(0) = U$  and proceeds in a direction of descent  $\dot{\gamma}(0) = \dot{U} \in T_U \mathcal{M}$ . In analogy with unconstrained algorithms, a good choice of  $\dot{U} \in T_U \mathcal{M}$  is found through a local Taylor expansion

$$f(\gamma(t)) = f(U) + t \langle \text{grad } f(U), \dot{U} \rangle + t^2 \langle \text{Hess } f(U)[\dot{U}], \dot{U} \rangle + O(t^3), \quad (11)$$

where  $\text{grad } f$  and  $\text{Hess } f$  are respectively the *Riemannian gradient* and *Riemannian Hessian* of  $f$  on the manifold  $\mathcal{M}$ . Afterwards, we trace the curve  $\gamma(t) = \text{Retr}_U(t\dot{U})$  using a *second-order retraction* operator  $\text{Retr}_U : T_U \mathcal{M} \rightarrow \mathcal{M}$  satisfying

$$\text{Retr}_U(0) = U, \quad \frac{d}{dt} \text{Retr}_U(t\dot{U}) \Big|_{t=0} = \dot{U}, \quad \frac{d^2}{dt^2} \text{Retr}_U(t\dot{U}) \Big|_{t=0} \perp T_U \mathcal{M},$$

for all  $U \in \mathcal{M}$  and all  $\dot{U} \in T_U \mathcal{M}$ . After choosing step-size  $t$  so that  $U_{\text{new}} = \gamma(t)$  makes a sufficient improvement over  $U$ , we repeat the algorithm until it reaches an  $\epsilon$ -second-order critical point satisfying  $\|\text{grad } f(U)\| \leq \epsilon$  and  $\lambda_{\min}(\text{Hess } f(U)) \geq -\sqrt{\epsilon}$ , which incidentally corresponds exactly to (9) and (10). Proofs for the following convergence result can be found in Zhang and Zhang [44], Boumal, Absil, and Cartis [9], and Agarwal et al. [2]; we have chosen the simplest but most restrictive settings to ease the exposition.

**Theorem 1** (Riemannian cubic-regularized Newton). *Suppose that  $\min_{U \in \mathcal{M}} f(U) > -\infty$ , and that the pullback  $\hat{f} = f \circ \text{Retr}_U$  has Lipschitz continuous Hessian for all  $U \in \mathcal{M}$ . Then, there exists a sufficiently large regularizer  $L$  such that  $U_{k+1} = \text{Retr}_{U_k}(\dot{U}_k)$  where*

$$\dot{U}_k = \underset{\dot{U} \in T_{U_k} \mathcal{M}}{\text{argmin}} f(U) + \langle \text{grad } f(U), \dot{U} \rangle + \frac{1}{2} \langle \text{Hess } f(U)[\dot{U}], \dot{U} \rangle + \frac{L}{6} \|\dot{U}\|^3$$

converges to an  $\epsilon$ -second order critical point in  $O(\epsilon^{-3/2})$  iterations, independent of dimension.

Each iteration of Riemannian cubic-regularized Newton solves an expensive Newton subproblem. Although it converges in far fewer iterations compared to gradient methods, it is practically competitive only when the added cost of solving the Newton subproblem can be offset by the corresponding reduction in iteration count.

## 3 Formulation and solution of K-means as manifold optimization

We now explain how we solve (3) using a Riemannian optimization approach. As a first attempt, we can indeed verify that the the constraint set in (3), written

$$\mathcal{M} := \mathcal{M}_r = \{U \in \mathbb{R}^{n \times r} : UU^\top \mathbf{1}_n = \mathbf{1}_n, \text{tr}(UU^\top) = K\}, \quad (12)$$

is a manifold by checking that (8) holds (cf. Lemma 4 in the appendix). In fact, directly applying Riemannian optimization techniques results in a  $K$ -means algorithm very similar to the one proposed in Carson et al.

[16]. The immediate and critical difficulty with this approach is the lack of an efficient retraction operator, which must be called at every iteration to keep iterates feasible  $U \in \mathcal{M}$ . For example, Carson et al. [16] used a complicated exponential retraction that costs  $O(n^2)$  time, hence bottlenecking the entire algorithm and preventing it from scaling to large  $n$ .

Instead, our first contribution in this paper is to reformulate (3) by establishing a submersion from the product manifold  $\widetilde{\mathcal{M}} = \mathcal{V} \times \text{Orth}(r)$  to  $\mathcal{M}$ , where  $\mathcal{V} = \{V \in \mathbb{R}^{n \times (r-1)} : \mathbf{1}_n^\top V = 0, \text{tr}(VV^\top) = K - 1\}$  and  $\text{Orth}(r) = \{Q \in \mathbb{R}^{r \times r} : QQ^\top = I_r\}$ . In words,  $\mathcal{V}$  is a projected hypersphere and  $\text{Orth}(r)$  is the set of  $r \times r$  orthonormal matrices.

**Theorem 2.** *We have  $\mathcal{M} = \varphi(\widetilde{\mathcal{M}})$  for  $\varphi(V, Q) = [\hat{\mathbf{1}}_n \quad V] Q$ , where  $\hat{\mathbf{1}}_n := (1/\sqrt{n})\mathbf{1}_n$ . Moreover, the Jacobian  $D\varphi : T\widetilde{\mathcal{M}} \rightarrow T\mathcal{M}$  is surjective for all  $(V, Q) \in \widetilde{\mathcal{M}}$ , i.e.,  $\varphi$  is a submersion.*




Figure 1: **Local convergence to second-order critical points yields global optimality.** In the case of GMMs, where ground truths can be planted, we consistently observe local convergence to global optimality, yielding zero clustering error. This provides strong numerical evidence that near-second-order critical points are near-globally optimal, as hypothesized in Assumption 1.

Having established the submersion property of  $\varphi$ , it is a standard result that every  $\epsilon$ -second order point of  $\widetilde{\mathcal{M}}$  is also an  $c\epsilon$ -second order point on  $\mathcal{M}$  for some constant rescaling factor  $c$ ; see e.g. Example 3.14 and the surrounding text in Levin, Kileel, and Boumal [29]. Therefore, to solve (3), we equivalently solve

$$\min_{(V, Q) \in \widetilde{\mathcal{M}}} \langle C, VV^\top \rangle - \mu \sum_{i,j} \log(\varphi_{i,j}(V, Q))_+, \quad (13)$$

where  $C = -XX^\top$  is the (negative) data Gram matrix, and  $\varphi_{i,j}$  is the  $(i, j)$ -th element of the operator  $\varphi$  in Theorem 2. A basic but critical benefit of the reformulation (13) is that the product manifold  $\widetilde{\mathcal{M}}$  admits a simple second-order retraction via its Euclidean projection [8, Sec. 5.12]

$$\text{Retr}_{(V, Q)}(\dot{V}, \dot{Q}) = [\Pi_{\mathcal{V}}(V + \dot{V}) \quad \Pi_{\text{Orth}(r)}(Q + \dot{Q})],$$

where  $\Pi_{\mathcal{V}}(V) = \sqrt{K-1}(V - n^{-1}\mathbf{1}_n\mathbf{1}_n^\top V)/\|V - n^{-1}\mathbf{1}_n\mathbf{1}_n^\top V\|$  and  $\Pi_{\text{Orth}(r)}(Q) = (QQ^\top)^{-1/2}Q$ . It is easy to check that the retraction above costs just  $O(nr + r^3)$  time to evaluate. In Section C.2, we give explicit expressions for the Riemannian gradient and Hessian and explain how they can be computed in  $O(nr + r^3)$  time.

The appearance of the logarithmic penalty in (13) presents two difficulties. First, as a practical concern, any algorithm for (13) must begin at a *strictly feasible* point  $(V_0, Q_0) \in \widetilde{\mathcal{M}}$  that additionally satisfies  $\varphi(V_0, Q_0) > 0$ . In Section C.4, we provide a good strictly feasible initial point, and prove that points exist only if the search rank is over-parameterized as  $r > K$ . Second, some special care is needed to rigorously

apply the guarantees from Section 2.3, given that the penalty  $\varphi(V, Q)$  is Lipschitz only when restricted to a closed and strictly feasible subset; see Section C.5 for details.

Together, these ingredients allow us to apply Riemannian gradient descent [9] to (13) to compute an  $\epsilon$ -first-order critical point in  $(n/\epsilon) \cdot \text{poly}(r, d, K)$  time. In practice, the algorithm often converges to an  $\epsilon$ -second-order critical point, though this is not rigorously guaranteed without a carefully-tuned noise perturbation. Alternatively, we can apply the conjugate-gradients (CG) variant of the Riemannian trust-region algorithm (RTR), a general-purpose solver available in packages like MANOPT [7] or PYMANOPT [41], to guarantee convergence to an  $\epsilon$ -second-order critical point. Unfortunately, in our experiments, we observed that all of these algorithms experience unsatisfactorily slow convergence, due to the severe ill-conditioning introduced by the logarithmic penalty.

Instead, our best numerical results were obtained by the Riemannian cubic-regularized Newton (Theorem 1). Our key insight is that the algorithm can be implemented with just  $O(nr^3)$  time per-iteration, by exploiting the underlying block-diagonal-plus-low-rank structure of the Riemannian Hessian. To explain, our core difficulty is to efficiently solve the Newton subproblem

$$\min_{Ap=0} g^\top p + \frac{1}{2} p^\top H p + \frac{L}{6} \|p\|^3,$$

where  $g$  and  $H$  denote the vectorized Riemannian gradient and Hessian respectively, and  $A$  implements the tangent space constraint  $(\dot{V}, \dot{Q}) \in T_{(V, Q)} \mathcal{M}$ . We can verify that the subproblem contains  $n(r-1) + r^2 = O(nr)$  variables and is subject to  $m = r + r(r+1)/2 = O(r^2)$  constraints. Given that the subproblem has only linear constraints, its local minima must always satisfy the first- and second-optimality conditions (9) and (10), which read

$$\begin{bmatrix} H + \lambda I & A^\top \\ A & 0 \end{bmatrix} \begin{bmatrix} p \\ q \end{bmatrix} = \begin{bmatrix} -g \\ 0 \end{bmatrix}, \quad \lambda = \frac{L}{2} \|p\|, \quad \xi^\top (H + \lambda I) \xi \geq 0 \text{ for all } \xi \text{ satisfying } A\xi = 0.$$

The following standard result shows that, with sufficient regularization  $L$ , the global minimum corresponds to the unique second-order critical point.

**Lemma 2.** *Let  $A$  have full row-rank (i.e.  $AA^\top \succ 0$ ) and let  $\lambda_{\min} = \min_{\|\xi\|=1, A\xi=0} \xi^\top H \xi$ . For  $\lambda > -\lambda_{\min}$ , the parameterized solution  $p(\lambda) = \begin{bmatrix} I \\ 0 \end{bmatrix}^\top \begin{bmatrix} H + \lambda I & A^\top \\ A & 0 \end{bmatrix}^{-1} \begin{bmatrix} -g \\ 0 \end{bmatrix}$  is well-defined and  $\|p(\lambda)\|$  is monotonously decreasing with respect to  $\lambda$ .*

The same lemma also suggests solving the Newton subproblem by simple bisection search. Indeed, the solution is just  $p(\lambda_{\text{opt}})$ , where  $\lambda_{\text{opt}}$  is the solution to the *monotone* equation  $2\lambda = L\|p(\lambda)\|$  (via Lemma 2). Thus, we pick a very small  $\lambda_{\text{lb}} \approx -\lambda_{\min}$  such that  $\|p(\lambda_{\text{lb}})\| > 2\lambda_{\text{lb}}/L$ , a very large  $\lambda_{\text{ub}}$  such that  $2\lambda_{\text{ub}}/L > \|p(\lambda_{\text{ub}})\|$ , and then perform bisection until  $2\lambda_{\text{opt}} = L\|p(\lambda_{\text{opt}})\|$  is approximately found. For each  $\lambda$ , if  $2\lambda < L\|p(\lambda)\|$ , then we increase  $\lambda$ ; otherwise, we decrease  $\lambda$ .

The main cost of the bisection search is the computation of  $p(\lambda)$ , which naively costs  $O(n^3 r^3)$  time. For our specific problem, we explain in Section E how a block-diagonal-plus-low-rank structure in the Hessian  $H$  reduces the computation cost to just  $n \cdot \text{poly}(r, d)$  time. Applying Theorem 1 shows that the overall method computes an  $\epsilon$ -second-order critical point in  $(n/\epsilon^{1.5}) \cdot \text{poly}(r, d, K)$  time.

## 4 Numerical results

In this section, we showcase the superior performance of our proposed Riemannian second-order method for clustering on both synthetic Gaussian mixture models (GMM) and real-world mass cytometry (CyTOF) datasets. Compared to existing state-of-the-art methods, such as the nonnegative low-rank (NLR) factorization [47] and prior Riemannian  $K$ -means algorithms [16], our approach achieves faster convergence, higher clustering accuracy, and more reliable recovery of ground-truth cluster memberships. These results highlight the convergence and accuracy advantages of second-order methods when they can be implemented with per-iteration costs of just  $O(n)$  time. The implementation details are deferred to Section G.




Figure 2: **Real-world benchmark on CyTOF data.** We compared our method to NLR, the previous state-of-the-art, as well as classical benchmarks SC, NMF, and  $KM++$ . Our method and NLR achieve the most consistently accurate clustering, with the smallest variance and the fewest outliers (left), but we outperform NLR in ground truth recovery (right).

**Datasets.** We conducted experiments on both synthetic and real datasets. The synthetic data was generated from a standard  $K$ -component,  $d$ -dimensional Gaussian mixture model (GMM), with centroids placed at simplex vertices such that their separation equals  $\gamma\bar{\Theta}^2$ , where  $\bar{\Theta}$  is the information-theoretic threshold for exact recovery in (6), and  $\gamma$  controls separation. The real dataset came from mass cytometry (CyTOF) [30, 43]. It consists of 265,627 cell protein expression profiles across 32 markers, labeled into 14 gated cell populations. Following Zhuang et al. [47], we uniformly sample 1,800 cells from  $K = 4$  unbalanced clusters (labels 2, 7, 8, 9) from individual 1 for our experiment.

**Global optimality at second-order critical points (validation of Assumption 1).** Figure 1 shows the convergence behaviors of loss function (13) for GMMs ( $n = 500$ ,  $\gamma = 1.2$ ,  $\mu = 0.01$ ) with 50 randomized initializations. We consistently observe that: (i) the loss value steadily decreases over iterations and converges rapidly near the globally optimal point; (ii) the Riemannian gradient norm dynamics suggest that our algorithm initially attempts to escape saddle points (with increased gradient norm) and eventually converges to second-order local optimality, where zero-loss is achieved, indicating global optimality. To verify second-order local optimality, we also plot the minimum eigenvalue of the Riemannian Hessian. This provides strong numerical evidence that near-second-order critical points are near-globally optimal, as posited by Assumption 1.

**Benchmark on real world data.** Prior studies on mass cytometry (CyTOF) and computer vision (CIFAR-10) datasets identified the nonnegative low-rank (NLR) factorization [47] as the most reliable clustering solver, attaining the lowest average mis-clustering error and the tightest variance compared to classical baselines such spectral clustering (SC), nonnegative matrix factorization (NMF), and  $K$ -means++ [4] ( $KM++$ ). Our algorithm optimizes the same nonnegative low-rank model, so it inherits this reliability. Because it applies second-order Hessian updates rather than first-order gradients, it refines each iterate more thoroughly and therefore recovers the ground-truth membership matrix more accurately. Figure 2 illustrates this on CyTOF: both methods keep mis-clustering near zero, yet our solver achieves a smaller Frobenius gap to the oracle solution. The experiment was repeated 50 times on random subsamples of size  $n = 1800$ .

**Comparison with NLR.** Next, we compare our method directly to the nonnegative low-rank (NLR) factorization. Figure 3 shows experimental results for GMM with  $n = 100$ ,  $\gamma = 0.8$ ,  $\mu = 0.1$ , and with varying  $n$ . Main observations: (i) Each Newton step is solved in  $O(n)$  time, matching the theory in Section 3. (ii)




Figure 3: **Comparison with previous state-of-the-art NLR on GMM.** Our second-order method reaches optimality in 152 iterations, while NLR needs 80k. Even though each second-order iteration costs  $\approx 30\text{--}100$  NLR steps, the total runtime is still two to four times shorter. (Left and middle) clustering accuracy vs log iterations and linear time; (right) per-iteration time vs sample size  $n$ .

A Newton step is about 30–100 times costlier than a single NLR update. This is to be expected because the Newton step solves several linear systems, while NLR performs only a single matrix–vector product. (iii) Our solver reaches the optimum in hundreds of iterations, whereas NLR needs tens of thousands. The orders-of-magnitude reduction in iterations more than offsets the costlier step, so wall-clock time drops by a factor of two to four.




Figure 4: **Comparison with prior Riemannian  $K$ -means method of Carson et al. [16] on real-world data.** Each run is warm-started from the previous and the penalty is stepped through  $\lambda_i = 0, 10^4, 10^6, 10^7$ . However: (Left) average mis-clustering exceeds 30%; (Middle) the recovery error  $\|Z - Z^*\|_F$  remains large; (Right) the infeasibility  $\|U_-\|$  never vanishes. Our Riemannian method, shown for reference, enforces  $U_- = 0$  by design and achieves near-zero error in both metrics.

**Comparison with prior Riemannian  $K$ -means methods.** We evaluate the clustering performance of the algorithm proposed by Carson et al. [16] to solve the penalized formulation (4) on the CyTOF data. Figure 4 presents the performance of this first-order manifold method. Unfortunately, we were unable to identify a sequence of  $\lambda$  that would produce acceptable clustering results. While increasing the penalty parameter  $\lambda$  improves feasibility, it also degrades clustering performance. This highlights the difficulty in solving (4) using the method of Carson et al. [16], as it struggles to balance strict constraint satisfaction with objective optimality in the  $K$ -means problem.




Figure 5: **Comparison with classical Riemannian Trust Region (RTR) on GMM.** Our method drives both loss and gradient norm to machine precision in around 360 iterations. In contrast, RTR stagnates for over 21k iterations due to the extreme ill-conditioning induced by the log penalty.




Figure 6: **Dependence on hyperparameters.** (Left) Error exhibits phase transition as penalty parameter  $\mu$  gets too large. (Right): Error seems insensitive to the search rank  $r = K + 1, \dots, 3K$ .

**Comparison with classical Riemannian algorithms.** As discussed in Section 3, Problem (13) can be solved with CG or RTR as the gradient, Hessian, and retraction are all available. Nevertheless, both CG and RTR perform poorly because the log-barrier induces an extremely ill-conditioned landscape. To illustrate this, we solve (13) on GMM data using PYMANOPT’s implementation of CG and RTR. For CG, we were unable to tune the method to produce a meaningful solution, as its updates frequently lead to infeasible points. While RTR can converge to a solution comparable to ours, it requires significantly more iterations and time, as shown in Figure 5.

**Effect of hyperparameters.** Our method displays a sharp phase transition with respect to the regularization parameter  $\mu$ . Below a critical threshold, the algorithm consistently converges to optimal solutions and remains robust to variations in  $\mu$ . However, once  $\mu$  exceeds this threshold, the method fails to yield meaningful results. This behavior is illustrated in Figure 6 where we used GMM synthetic data with four clusters and separation  $\gamma = 0.8$ . Figure 6 also shows the clustering errors evaluated across different search ranks  $r$ . Clustering performance appears insensitive to  $r$ , so the smallest feasible  $r = K + 1$  should be used in practice. A brief discussion on the choice of hyperparameters is provided in Section F.

## References

- [1] P-A Absil et al. *All roads lead to Newton: Feasible second-order methods for equality-constrained optimization*. Technical Report. 2009.
- [2] Naman Agarwal et al. “Adaptive regularization with cubics on manifolds”. In: *Mathematical Programming* 188 (2021), pp. 85–134.
- [3] Daniel Aloise et al. “NP-hardness of Euclidean sum-of-squares clustering”. In: *Machine learning* 75.2 (2009), pp. 245–248.
- [4] David Arthur and Sergei Vassilvitskii. “**k-means++**: the advantages of careful seeding”. In: *Proceedings of the Eighteenth Annual ACM-SIAM Symposium on Discrete Algorithms*. SODA '07. New Orleans, Louisiana: Society for Industrial and Applied Mathematics, 2007, pp. 1027–1035. ISBN: 9780898716245.
- [5] Afonso S Bandeira, Nicolas Boumal, and Vladislav Voroninski. “On the low-rank approach for semidefinite programs arising in synchronization and community detection”. In: *Conference on learning theory*. PMLR. 2016, pp. 361–382.
- [6] Srinadh Bhojanapalli, Behnam Neyshabur, and Nati Srebro. “Global optimality of local search for low rank matrix recovery”. In: *Advances in Neural Information Processing Systems* 29 (2016).
- [7] N. Boumal et al. “Manopt, a Matlab Toolbox for Optimization on Manifolds”. In: *Journal of Machine Learning Research* 15.42 (2014), pp. 1455–1459. URL: <https://www.manopt.org>.
- [8] Nicolas Boumal. *An Introduction to Optimization on Smooth Manifolds*. Cambridge University Press, 2023. DOI: 10.1017/9781009166164. URL: <https://www.nicolasboumal.net/book>.
- [9] Nicolas Boumal, Pierre-Antoine Absil, and Coralia Cartis. “Global rates of convergence for nonconvex optimization on manifolds”. In: *IMA Journal of Numerical Analysis* 39.1 (2019), pp. 1–33.
- [10] Nicolas Boumal, Vlad Voroninski, and Afonso Bandeira. “The non-convex Burer-Monteiro approach works on smooth semidefinite programs”. In: *Advances in Neural Information Processing Systems* 29 (2016).
- [11] Nicolas Boumal, Vladislav Voroninski, and Afonso S Bandeira. “Deterministic Guarantees for Burer-Monteiro Factorizations of Smooth Semidefinite Programs”. In: *Communications on Pure and Applied Mathematics* 73.3 (2020), pp. 581–608.
- [12] S.P. Boyd and L. Vandenberghe. *Convex Optimization*. Cambridge University Press, 2004. ISBN: 9780521833783.
- [13] Samuel Burer and Renato D. C. Monteiro. “A nonlinear programming algorithm for solving semidefinite programs via low-rank factorization”. In: *Mathematical Programming* 95.2 (Feb. 1, 2003), pp. 329–357. ISSN: 1436-4646. DOI: 10.1007/s10107-002-0352-8.
- [14] Richard H Byrd, Jean Charles Gilbert, and Jorge Nocedal. “A trust region method based on interior point techniques for nonlinear programming”. In: *Mathematical programming* 89 (2000), pp. 149–185.
- [15] Richard H Byrd, Jorge Nocedal, and Richard A Waltz. “Knitro: An integrated package for nonlinear optimization”. In: *Large-scale nonlinear optimization* (2006), pp. 35–59.
- [16] Timothy Carson et al. “Manifold optimization for  $k$ -means clustering”. In: *2017 International Conference on Sampling Theory and Applications (SampTA)*. 2017, pp. 73–77. DOI: 10.1109/SAMPTA.2017.8024388.
- [17] Xiaohui Chen and Yun Yang. “Cutoff for exact recovery of Gaussian mixture models”. In: *IEEE Transactions on Information Theory* 67.6 (2021), pp. 4223–4238.
- [18] Xiaohui Chen and Yun Yang. “Hanson–Wright inequality in Hilbert spaces with application to  $K$ -means clustering for non-Euclidean data”. In: *Bernoulli* 27.1 (2021), pp. 586–614. DOI: 10.3150/20-BEJ1251.
- [19] Sanjoy Dasgupta. “The hardness of  $k$ -means clustering”. In: *Technical Report CS2007-0890, University of California, San Diego* (2007).

[20] Peter JC Dickinson and Luuk Gijben. “On the computational complexity of membership problems for the completely positive cone and its dual”. In: *Computational optimization and applications* 57 (2014), pp. 403–415.

[21] Yingjie Fei and Yudong Chen. “Hidden Integrality of SDP Relaxations for Sub-Gaussian Mixture Models”. In: *Proceedings of the 31st Conference On Learning Theory*. Ed. by Sébastien Bubeck, Vianney Perchet, and Philippe Rigollet. Vol. 75. Proceedings of Machine Learning Research. PMLR, July 6, 2018, pp. 1931–1965. URL: <https://proceedings.mlr.press/v75/fei18a.html>.

[22] Rong Ge, Chi Jin, and Yi Zheng. “No spurious local minima in nonconvex low rank problems: A unified geometric analysis”. In: *International Conference on Machine Learning*. PMLR. 2017, pp. 1233–1242.

[23] Rong Ge, Jason D Lee, and Tengyu Ma. “Matrix completion has no spurious local minimum”. In: *Advances in neural information processing systems* 29 (2016).

[24] Zhaoshui He et al. “Symmetric Nonnegative Matrix Factorization: Algorithms and Applications to Probabilistic Clustering”. In: *IEEE Transactions on Neural Networks* 22.12 (2011), pp. 2117–2131. DOI: [10.1109/TNN.2011.2172457](https://doi.org/10.1109/TNN.2011.2172457).

[25] Wen Huang et al. “A Riemannian Optimization Approach to Clustering Problems”. In: *Journal of Scientific Computing* 103.1 (Feb. 2025), p. 8. ISSN: 1573-7691. DOI: [10.1007/s10915-025-02806-3](https://doi.org/10.1007/s10915-025-02806-3).

[26] V. Kalofolias and E. Gallopolous. “Computing symmetric nonnegative rank factorizations”. In: *Linear Algebra and its Applications* 436.2 (2012). Special Issue devoted to the Applied Linear Algebra Conference (Novi Sad 2010), pp. 421–435. ISSN: 0024-3795. DOI: [10.1016/j.laa.2011.03.016](https://doi.org/10.1016/j.laa.2011.03.016). URL: <https://www.sciencedirect.com/science/article/pii/S0024379511002199>.

[27] Etienne Klerk. *Aspects of Semidefinite Programming*. Springer New York, NY, 2006. ISBN: 978-0-306-47819-2. DOI: [10.1007/b105286](https://doi.org/10.1007/b105286).

[28] Da Kuang, Sangwoon Yun, and Haesun Park. “SymNMF: nonnegative low-rank approximation of a similarity matrix for graph clustering”. In: *Journal of Global Optimization* 62 (2015), pp. 545–574.

[29] Eitan Levin, Joe Kileel, and Nicolas Boumal. “The effect of smooth parametrizations on nonconvex optimization landscapes”. In: *Mathematical Programming* 209.1 (Jan. 2025), pp. 63–111. ISSN: 1436-4646. DOI: [10.1007/s10107-024-02058-3](https://doi.org/10.1007/s10107-024-02058-3).

[30] Jacob H. Levine et al. “Data-Driven Phenotypic Dissection of AML Reveals Progenitor-like Cells that Correlate with Prognosis”. In: *Cell* 162.1 (2015), pp. 184–197. ISSN: 0092-8674. DOI: [10.1016/j.cell.2015.05.047](https://doi.org/10.1016/j.cell.2015.05.047). URL: <https://www.sciencedirect.com/science/article/pii/S0092867415006376>.

[31] Stuart Lloyd. “Least squares quantization in PCM”. In: *IEEE Transactions on Information Theory* 28 (1982), pp. 129–137.

[32] Ulrike von Luxburg. “A tutorial on spectral clustering”. In: *Statistics and Computing* 17.4 (2007), pp. 395–416.

[33] Jan R. Magnus and Heinz Neudecker. *Matrix Differential Calculus with Applications in Statistics and Econometrics*. Third. Wiley Series in Probability and Statistics. John Wiley, 2019. ISBN: 9781119541202. DOI: [10.1002/9781119541219](https://doi.org/10.1002/9781119541219).

[34] Bamdev Mishra and Rodolphe Sepulchre. “Riemannian preconditioning”. In: *SIAM Journal on Optimization* 26.1 (2016), pp. 635–660.

[35] Katta G Murty and Santosh N Kabadi. “Some NP-complete problems in quadratic and nonlinear programming”. In: *Mathematical Programming* 39 (1987), pp. 117–129.

[36] Andrew Y. Ng, Michael I. Jordan, and Yair Weiss. “On Spectral Clustering: Analysis and an algorithm”. In: *Advances in Neural Information Processing Systems*. MIT Press, 2001, pp. 849–856.

[37] Jorge Nocedal and Stephen J. Wright. *Numerical Optimization*. 2nd. Springer Series in Operations Research and Financial Engineering. Springer New York, NY, 2006. ISBN: 978-0-387-40065-5. DOI: [10.1007/978-0-387-40065-5](https://doi.org/10.1007/978-0-387-40065-5).

[38] Jiming Peng and Yu Wei. “Approximating  $K$ -means-type clustering via semidefinite programming”. In: *SIAM J. OPTIM* 18.1 (2007), pp. 186–205.

[39] Wei Qian, Yuqian Zhang, and Yudong Chen. “Structures of Spurious Local Minima in k-Means”. In: *IEEE Transactions on Information Theory* 68.1 (2022), pp. 395–422. DOI: 10.1109/TIT.2021.3122465.

[40] Martin Royer. “Adaptive Clustering through Semidefinite Programming”. In: *Advances in Neural Information Processing Systems 30*. Ed. by I. Guyon et al. Curran Associates, Inc., 2017, pp. 1795–1803.

[41] James Townsend, Niklas Koep, and Sebastian Weichwald. “Pymanopt: A Python Toolbox for Optimization on Manifolds using Automatic Differentiation”. In: *Journal of Machine Learning Research* 17.137 (2016), pp. 1–5. URL: <http://jmlr.org/papers/v17/16-177.html>.

[42] Yu-Xiong Wang and Yu-Jin Zhang. “Nonnegative matrix factorization: A comprehensive review”. In: *IEEE Transactions on knowledge and data engineering* 25.6 (2012), pp. 1336–1353.

[43] Lukas Weber. *Clustering benchmark data: 32-dimensional data set from Levine et al.* GitHub Repository. 2015. URL: <https://github.com/lmweber/benchmark-data-Levine-32-dim>.

[44] Junyu Zhang and Shuzhong Zhang. *A cubic regularized Newton’s method over Riemannian manifolds*. 2018. arXiv: 1805.05565 [math.OC].

[45] Richard Y Zhang. *Nonnegative Low-rank Matrix Recovery Can Have Spurious Local Minima*. 2025. arXiv: 2505.03717 [math.OC].

[46] Yubo Zhuang, Xiaohui Chen, and Yun Yang. “Wasserstein  $K$ -means for clustering probability distributions”. In: *Advances in Neural Information Processing Systems*. 2022.

[47] Yubo Zhuang et al. “Statistically Optimal  $K$ -means Clustering via Nonnegative Low-rank Semidefinite Programming”. In: *The Twelfth International Conference on Learning Representations*. 2024.

# Appendices

## A Relationship between SDP Formulations of $K$ -means Clustering: Standard and Interior Point Versions (2)

The standard  $K$ -means clustering for data in  $\mathbb{R}^d$  has two formulations in the literature: (i) *centroid-based* optimization

$$\min_{\beta_1, \dots, \beta_K \in \mathbb{R}^d} \sum_{i=1}^n \min_{k \in [K]} \|X_i - \beta_k\|_2^2;$$

and (ii) *partition-based* optimization

$$\bigsqcup_{k=1}^K \min_{G_k=[n]} \sum_{k=1}^K \sum_{i \in G_k} \|X_i - \bar{X}_k\|_2^2,$$

where  $\bar{X}_k = |G_k|^{-1} \sum_{j \in G_k} X_j$  is the empirical centroid of cluster  $G_k$ . Formulations (i) and (ii) are known to be equivalent, cf. Zhuang, Chen, and Yang [46, Eqn. (1)] or Qian, Zhang, and Chen [39, Appx. A]. Using the parallelogram law in Zhuang, Chen, and Yang [46, Eqn. (5)]

$$\sum_{i,j \in G_k} \|X_i - X_j\|_2^2 = 2|G_k| \sum_{i \in G_k} \|X_i - \bar{X}_k\|_2^2,$$

we may write the partition-based objective function as

$$\bigsqcup_{k=1}^K \min_{G_k=[n]} \sum_{k=1}^K \frac{1}{2|G_k|} \sum_{i,j \in G_k} \|X_i - X_j\|_2^2.$$

Next, expanding the pairwise squared Euclidean distance and dropping  $\sum_{i=1}^n \|X_i\|_2^2$  (no longer depending on any partition  $G_1, \dots, G_K$ ), we arrive at (1), in the form of maximizing the total intra-cluster similarity in terms of the Gram matrix  $\{\langle X_i, X_j \rangle\}_{i,j=1}^n$ .

For general data without a likelihood derivation as in Chen and Yang [17], we can replace the  $\mathbb{R}^d$ -inner product with any (positive semidefinite) kernel  $k: \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  and consider the kernelized version of  $K$ -means clustering that involves data  $X_1, \dots, X_n$  only via their Gram matrix  $\{k(X_i, X_j)\}_{i,j=1}^n$ . Thus, our manifold formulation of this paper carries over to the general kernel method setting with possibly nonlinear boundary structure.

Next, we convexify the  $K$ -means problem (1) into an SDP. Note that each partition  $G_1, \dots, G_K$  via one-hot encoding is equivalent to an assignment matrix  $H_{n \times K}$  (up to cluster relabel) where the latter is a binary matrix with exactly one non-zero entry in each row, i.e.  $H_{ik} = 1$  if  $i \in G_k$ . With this reparameterization, one can write (1) as a mixed zero-one integer program:

$$\max_{H \in \{0,1\}^{n \times K}} \{\langle XX^\top, H B H^\top \rangle : H \mathbf{1}_K = \mathbf{1}_n\}.$$

Now, applying the change of variables  $Z_{n \times n} = H B H^\top$  and noting that the membership matrix  $Z$  and assignment matrix  $H$  are not one-to-one, we relax the  $K$ -means problem by preserving the key properties of  $Z$  as the following constraints:

$$Z \succeq 0, \quad \text{tr}(Z) = K, \quad Z \mathbf{1}_n = \mathbf{1}_n, \quad Z \geq 0,$$

which no longer depend on the assignment matrix  $H$ . Then, we arrive at the standard SDP relaxation for  $K$ -means clustering, cf. Peng and Wei [38, Eqn. (13)] or Chen and Yang [17, Eqn. (11)]:

$$\max_{Z \in \mathbb{R}^{n \times n}} \{\langle XX^\top, Z \rangle : Z \succeq 0, \text{tr}(Z) = K, Z \mathbf{1}_n = \mathbf{1}_n, Z \geq 0\}. \quad (14)$$

In practice, the elementwise nonnegativity constraint  $Z \geq 0$  is almost always enforced by a logarithmic barrier. This means that we can make (14) more explicitly in the form

$$\max_{Z \in \mathbb{R}^{n \times n}} \left\{ \langle XX^\top, Z \rangle + \mu \sum_{i,j=1}^n \log(Z_{i,j})_+ : Z \succeq 0, \text{tr}(Z) = K, Z\mathbf{1}_n = \mathbf{1}_n \right\}, \quad (15)$$

which is precisely how any practical interior-point solver would solve the original SDP in (14). The barrier cost is the actual objective used internally, with  $\mu$  set to reflect the solver's target accuracy, cf. Boyd and Vandenberghe [12, Chapter 11] or Nocedal and Wright [37, Chapters 14 & 19]. In other words, we take the standard SDP in (14), and make explicit the logarithmic penalty that is already implicit in how such an SDP is actually solved.

## B Information-theoretic Threshold for Exact Recovery

The following theorem is a precise statement of the information-theoretic threshold of Chen and Yang [17].

**Theorem 3** (Average-case phase transition for exact recovery). *Let  $\alpha > 0$  and*

$$\Theta_{\min} := \min_{1 \leq j < k \leq K} \|\mu_j - \mu_k\|$$

*be the minimum centroid separation. Suppose that data  $X_1, \dots, X_n$  are generated from the Gaussian mixture model (5) with equal cluster size  $|G_1^*| = \dots = |G_K^*| = m$ . Then we have the following dichotomy.*

1. *If  $K \leq \log n / \log \log(n)$  and  $\Theta_{\min} \geq (1 + \alpha)\bar{\Theta}$ , then there exist constants  $C_1, C_2 > 0$  depending only on  $\alpha$  such that, with probability at least  $1 - C_1(\log n)^{-C_2}$ , the SDP (2) as  $\mu \rightarrow 0^+$  (cf. Appendix A for the equivalence of two SDP formulations) has a unique solution that exactly recovers the true partition  $G_1^*, \dots, G_K^*$ .*
2. *If  $K \leq \log n$  and  $\Theta_{\min} \leq (1 - \alpha)\bar{\Theta}$ , then there exists a constant  $C_3 > 0$  depending only on  $\alpha$  such that*

$$\inf_{\hat{G}_1, \dots, \hat{G}_K} \sup_{\Xi(n, K, \Theta_{\min})} \mathbb{P}(\exists k : \hat{G}_k \neq G_k^*) \geq 1 - \frac{C_3 K}{n},$$

*where the infimum is taken over all possible estimators  $(\hat{G}_1, \dots, \hat{G}_K)$  for  $(G_1^*, \dots, G_K^*)$  and the parameter space is defined as*

$$\Xi(n, K, \Theta) := \left\{ (G_1, \dots, G_K, \mu_1, \dots, \mu_K) \left| \begin{array}{l} \|\mu_j - \mu_k\| \geq \Theta, \forall j, k \in [K], j \neq k \\ (1 - \delta_n)m \leq |G_k| \leq (1 + \delta_n)m \end{array} \right. \right\}$$

*with  $\delta_n = C\sqrt{K \log(n)/n}$  for some large enough constant  $C > 0$ .*

## C Additional Details on our Riemannian Formulation

### C.1 Equivalence between Riemannian Optimization and SQP

Let us consider a constraint optimization problem,

$$\begin{aligned} \min_{u \in \mathbb{R}^n} \quad & f(u), \\ \text{s.t.} \quad & g_i(u) = 0 \quad \forall i. \end{aligned} \quad (16)$$

where LICQ:

$$\sum_i y_i \nabla g_i(u) = 0 \iff y_i = 0 \quad \forall i.$$

holds for every  $u$  in the constraint set. Then  $\mathcal{M} := \{u \in \mathbb{R}^n : g_i(u) = 0 \forall i\}$  is a smooth embedded manifold [8], and we may employ Riemannian methods to solve (16).

The Riemannian method solves  $\min_{u \in \mathcal{M}} f(u)$  by solving

$$\min_{\dot{u} \in T_u \mathcal{M}} f(u) + \langle \text{grad } f(u), \dot{u} \rangle + \frac{1}{2} \langle \text{Hess } f(u)[\dot{u}], \dot{u} \rangle + \frac{L}{6} \|\dot{u}\|^3 \quad (17)$$

at each iteration. One can show that this is equivalent to the SQP method that solve (16) by minimizing the Lagrangian

$$\begin{aligned} \min_{\dot{u}} \quad & \mathcal{L}(u, y^{(u)}) + \langle \nabla_u \mathcal{L}(u, y^{(u)}), \dot{u} \rangle + \frac{1}{2} \langle \nabla_{uu}^2 \mathcal{L}(u, y^{(u)})[\dot{u}], \dot{u} \rangle + \frac{L}{6} \|\dot{u}\|^3, \\ \text{s.t.} \quad & \langle \nabla g_i(u), \dot{u} \rangle = 0 \forall i. \end{aligned} \quad (18)$$

where

$$\mathcal{L}(u, y^{(u)}) = f(u) + \sum_i y_i^{(u)} g_i(u) \quad \text{and} \quad y^{(u)} = \operatorname{argmin}_y \|\nabla_u \mathcal{L}(u, y)\|.$$

Hence, our contribution is to efficiently solve the SQP subproblem by exploiting a block-diagonal-plus-low-rank structure in the Hessian, and the fact that there are only  $r + r(r+1)/2 \ll n$  constraints. We provide more details in Appendix C.2 and Section C.3.

To establish the equivalence, we first observe that the search space of (17) and (18) are the same. Indeed, the tangent space  $T_u \mathcal{M} = \{\dot{u} : \langle \nabla g_i(u), \dot{u} \rangle = 0 \forall i\}$ . Next, we write the expressions for the Riemannian gradient and Hessian (Proposition 3.61 and Corollary 5.16 [8]):

$$\text{grad } f(u) := \text{Proj}_u(\nabla f(u)), \quad \text{Hess } f(u)[\dot{u}] := \text{Proj}_u(D_u \text{grad } f(u)[\dot{u}]), \quad (19)$$

where  $\text{Proj}_u(v) := \operatorname{argmin}_{\dot{u} \in T_u \mathcal{M}} \|v - \dot{u}\|$ , and  $D_u$  denotes the usual differential operator. We then obtain

$$\text{grad } f(u) := \text{Proj}_u(\nabla f(u)) = \nabla f(u) + \sum_i y_i^{(u)} \nabla g_i(u) = \nabla_u \mathcal{L}(u, y^{(u)}). \quad (20)$$

For the second equality, see Equation (7.75) in Boumal [8].

For the second order terms, we have  $\langle \text{Hess } f(u)[\dot{u}], \dot{u} \rangle = \langle \nabla_{uu}^2 \mathcal{L}(u)[\dot{u}], \dot{u} \rangle$  for all  $\dot{u}$  in  $T_u \mathcal{M}$  by the facts that

- (i)  $\text{Hess } f(u)[\dot{u}] = \text{Proj}_u(\nabla_{uu}^2 \mathcal{L}(u, y^{(u)})[\dot{u}])$ .
- (ii)  $\langle \text{Proj}_u(v), \dot{u} \rangle = \langle v, \dot{u} \rangle$ .

We obtain fact (i) by (19) and (20):

$$\begin{aligned} \text{Hess } f(u)[\dot{u}] &:= \text{Proj}_u(D \text{grad } f(u)[\dot{u}]) \\ &= \text{Proj}_u(\nabla^2 f(u)[\dot{u}] + \sum_i (D_u y_i^{(u)}) \nabla g_i(u)[\dot{u}] + \sum_i y_i^{(u)} \nabla^2 g_i(u)[\dot{u}]) \\ &= \text{Proj}_u(\nabla^2 f(u)[\dot{u}] + \sum_i y_i^{(u)} \nabla^2 g_i(u)[\dot{u}]) \\ &= \text{Proj}_u(\nabla_{uu}^2 \mathcal{L}(u, y^{(u)})[\dot{u}]). \end{aligned}$$

Inside the projection operator, the term  $\sum_i (D y_i^{(u)}) \nabla g_i(u)$  vanishes because  $T_u \mathcal{M}$  is a linear subspace, and  $(T_u \mathcal{M})^\perp = \text{span}(\nabla g_i(u))$ . Fact (ii) is due to that the projection operator is self-adjoint and that the projection of any tangent vector is itself. For more on the connection between SQP and Riemannian Newton method, we refer to Absil et al. [1] and Mishra and Sepulchre [34].

## C.2 General form of the Riemannian gradient and Hessian

Throughout this section, we use a bar over a function defined on the manifold  $\mathcal{M}$  to denote its smooth extension defined on a neighborhood of  $\mathcal{M}$  so that the Euclidean gradient and Hessian can be defined on  $\mathcal{M}$ . Namely, for  $f: \mathcal{M} \rightarrow \mathbb{R}^m$ , we use  $\bar{f}: N(\mathcal{M}) \rightarrow \mathbb{R}^m$  to denote the smooth extension. The notations  $\text{grad } f$  and  $\text{Hess } f$  denote the Riemannian gradient and Hessian;  $\nabla \bar{f}$  and  $\nabla^2 \bar{f}$  denote the Euclidean gradient and Hessian.

For manifolds that can be defined by

$$\min_{U \in \mathcal{M}} f(U), \quad \mathcal{M} = \{U \in \mathbb{R}^{n \times r} : \mathcal{A}(UU^\top) + \mathcal{B}(U) = c\},$$

where  $\mathcal{A}: \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^m$  and  $\mathcal{B}: \mathbb{R}^{n \times r} \rightarrow \mathbb{R}^m$  are linear operators, and  $c \in \mathbb{R}^m$ , its tangent space can be written as:

$$T_U \mathcal{M} = \{\dot{U} \in \mathbb{R}^{n \times r} : \mathcal{A}(\dot{U}U^\top + U\dot{U}^\top) + \mathcal{B}(\dot{U}) = 0\}.$$

We call the function  $\mathcal{A}(UU^\top) + \mathcal{B}(U) = c$  as the *defining function* of  $\mathcal{M}$ . Let us denote

$$L(\dot{U}) := \mathcal{A}(\dot{U}U^\top + U\dot{U}^\top) + \mathcal{B}(\dot{U}). \quad (21)$$

Immediately, we see that  $T_U \mathcal{M} = \ker(L)$ , and the adjoint operator

$$L^*(y) = 2\mathcal{A}^\top(y)U + \mathcal{B}^\top(y). \quad (22)$$

The projection operator onto the tangent space is defined to be

$$\text{Proj}_U(W) := \underset{\dot{U} \in T_U \mathcal{M}}{\text{argmin}} \|W - \dot{U}\|.$$

We also know that  $\ker(L)^\perp = \text{image}(L^*)$ . Thus, by the orthogonal projection theorem, we may see that

$$\text{Proj}_U(W) = W - W^\perp = W - L^*(\tilde{y}), \quad (23)$$

where  $\tilde{y} = \underset{y \in \mathbb{R}^m}{\text{argmin}} \|W - L^*(y)\|$  is the solution to the linear system  $W - L^*(y) \in \ker(L) = T_U \mathcal{M}$ , and both  $\text{Proj}_U(W)$  and  $\tilde{y}$  are unique.

Consequently, the Riemannian gradient and the Hessian matrix-vector product have the following form:

$$\begin{cases} \text{grad } f(U) := \text{Proj}_U(\nabla \bar{f}(U)) = \nabla \bar{f}(U) + 2[\mathcal{A}^\top(y_U)]U + \mathcal{B}^\top(y_U), \\ \text{Hess } f(U)[\dot{U}] := \text{Proj}_U(\text{D grad } f(U)[\dot{U}]) = \text{Proj}_U(\nabla^2 \bar{f}(U)[\dot{U}] + 2[\mathcal{A}^\top(y_U)]\dot{U}), \end{cases} \quad (24)$$

where  $y_U = -\tilde{y}$  in (23) is the unique Lagrange multipliers

$$y_U = \underset{y \in \mathbb{R}^m}{\text{argmin}} \|\nabla f(U) + 2[\mathcal{A}^\top(y)]U + \mathcal{B}^\top(y)\|. \quad (25)$$

For a detailed proof, see Boumal, Voroninski, and Bandeira [11].

## C.3 Efficient computation of the Riemannian gradient and Hessian

We first write down the Euclidean gradient and Hessian for our objective function, and then explain how to compute the Riemannian counterparts efficiently. Specifically, we show that  $y_U$  can be computed in  $O(nr + r^3)$  time.

We decompose the objective function as  $f = f_1 + \mu f_2$ , where

$$f_1(V, Q) := \langle C, VV^\top \rangle, \quad f_2(V, Q) := -\sum_{i,j} \log \varphi_{i,j}(V, Q).$$

The Euclidean gradients are respectively:  $\nabla_V f_1(V, Q) = 2CV$ ,  $\nabla_Q f_1(V, Q) = 0$ , and

$$\nabla_V f_2(V, Q) = -U^{(-1)} \begin{bmatrix} \mathbf{0}_{r-1} \\ I_{r-1} \end{bmatrix} Q^\top, \quad \nabla_Q f_2(V, Q) = - \begin{bmatrix} \hat{\mathbf{1}}_n^\top \\ V^\top \end{bmatrix} U^{(-1)},$$

where  $U := \varphi(V, Q)$  and  $U^{(-1)}$  is its element-wise inverse, i.e.,  $[U^{(-1)}]_{i,j} = (U_{i,j})^{-1}$ . The Euclidean Hessians are given in vectorized form as

$$\nabla^2 f_1(V, Q) = \begin{bmatrix} 2I_{r-1} \otimes C & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix}, \quad \nabla^2 f_2(V, Q) = \begin{bmatrix} J_V^\top \\ J_Q^\top \end{bmatrix} D^2 \begin{bmatrix} J_V & J_Q \end{bmatrix} - \begin{bmatrix} \mathbf{0} & H_{VQ} \\ H_{QV} & \mathbf{0} \end{bmatrix},$$

where

$$D = \text{dvec}(U^{(-1)}), \quad J_V = \left( Q^\top \begin{bmatrix} \mathbf{0} \\ I_{r-1} \end{bmatrix} \otimes I_n \right), \quad J_Q = I_r \otimes Q U^\top,$$

and

$$H_{VQ} = \left( [\mathbf{0} \quad I_{r-1}] \otimes U^{(-1)} \right) K^{(r,r)} = H_{QV}^\top,$$

with  $\text{dvec}(U^{(-1)}) := \text{diag}[\text{vec}(U^{(-1)})]$ , and  $K^{(n,r-1)}, K^{(r,r)}$  denoting the commutation matrices [33, Sec. 3.7]. We can then compute the Riemannian gradient and Hessian-vector-product according to (24) and (25). In the remainder of this section, we show how to efficiently solve (25).

The manifold we consider,  $\widetilde{\mathcal{M}} := \mathcal{V} \times \text{Orth}(r)$ , can be written as

$$\widetilde{\mathcal{M}} = \left\{ (V, Q) \in \mathbb{R}^{n \times (r-1)} \times \mathbb{R}^{r \times r} : \mathbf{1}_n^\top V = \mathbf{0}_{r-1}, \text{tr}(VV^\top) = K - 1, \text{svec}(QQ^\top - I_r) = \mathbf{0} \right\},$$

where  $\text{svec}$  denotes the symmetric vectorization operator [27, Appx. E]. Note that there are no cross terms ( $VQ^\top$  or  $QV^\top$ ) in the defining functions of  $\mathcal{M}$ . Thus, we can treat the defining functions with respect to  $V$  and  $Q$  separately. The corresponding terms are

$$\mathcal{A}_V(VV^\top) := \begin{bmatrix} \mathbf{0}_{r-1} \\ \text{tr}(VV^\top) \end{bmatrix}, \quad \mathcal{B}_V(V) := \begin{bmatrix} \mathbf{1}_n^\top V \\ 0 \end{bmatrix}, \quad \mathcal{A}_Q(QQ^\top) := \text{svec}(QQ^\top)$$

and

$$c_V := \begin{bmatrix} \mathbf{0}_{r-1} \\ K - 1 \end{bmatrix} \quad \text{and} \quad c_Q := \text{svec}(I_r).$$

Mimicking (21) and (22), we use the notation  $L_V$ ,  $L_Q$ ,  $L_V^*$ , and  $L_Q^*$  respectively. For any  $(y_1, y_2) \in \mathbb{R}^{r-1} \times \mathbb{R}$  and  $y_3 \in \mathbb{R}^{r(r+1)/2}$ , we have

$$L_V^*(y_1, y_2) = \mathbf{1}_n y_1^\top + 2y_2 V \quad \text{and} \quad L_Q^*(y_3) = 2 \text{smat}(y_3) Q,$$

where  $\text{smat}$  is the inverse of  $\text{svec}$ , that is,  $\text{smat}(\text{svec}(M)) = M$  for all symmetric matrices  $M$ . Denote  $G_V := \nabla_V f(V, Q)$  and  $G_Q := \nabla_Q f(V, Q)$ . By solving the linear systems

$$G_V - L_V^*(y_1, y_2) \in \ker(L_V) \quad \text{and} \quad G_Q - L_Q^*(y_3) \in \ker(L_Q),$$

we obtain the following closed form solutions:

$$\tilde{y}_1 = \frac{1}{n} G_V^\top \mathbf{1}_n, \quad \tilde{y}_2 = \frac{1}{2(K-1)} \langle G_V, V \rangle, \quad \tilde{y}_3 = \frac{1}{4} \text{svec}(G_Q Q^\top + Q G_Q^\top). \quad (26)$$

The computation of  $\tilde{y}_1$ ,  $\tilde{y}_2$ , and  $\tilde{y}_3$  in total requires  $O(nr + r^3)$  time. Therefore, we can compute  $y_U = -(\tilde{y}_1, \tilde{y}_2, \tilde{y}_3)$  with the same cost. Given a Euclidean gradient and a Euclidean Hessian-vector product, we may write out explicitly the Riemannian gradient:

$$\text{grad } f(V, Q) = \left[ \left( I - \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^\top \right) G_V - \frac{1}{K-1} \langle G_V, V \rangle V \quad \frac{G_Q Q^\top - Q G_Q^\top}{2} Q \right]$$

and the Riemannian Hessian-vector product:

$$\text{Hess } f(V, Q)[\dot{V}, \dot{Q}] = \begin{bmatrix} \text{Proj}_V \left( \nabla^2 f(V, Q)[\dot{V}, \dot{Q}]_V - \frac{1}{K-1} \langle G_V, V \rangle \dot{V} \right) \\ \text{Proj}_Q \left( \nabla^2 f(V, Q)[\dot{V}, \dot{Q}]_Q - \frac{G_Q Q^\top + Q G_Q^\top}{2} \dot{Q} \right) \end{bmatrix}^\top.$$

#### C.4 Feasible initial point

In this section, we show that  $r > K$  is necessary and sufficient for the existence of an interior point of  $\mathcal{M}$ . The following Lemma 3 shows the necessity of  $r > K$ . When  $r = K$ , the structure of the unique  $U \in \mathbb{R}_+^{n \times K}$  in Lemma 1 can be explicitly written as

$$U = \left[ \frac{1}{\sqrt{|G_1|}} \mathbf{1}_{G_1}, \frac{1}{\sqrt{|G_2|}} \mathbf{1}_{G_2}, \dots, \frac{1}{\sqrt{|G_K|}} \mathbf{1}_{G_K} \right], \quad (27)$$

where  $\mathbf{1}_{G_k} \in \{0, 1\}^n$  denotes the binary vector with its support being  $G_k$ .

**Lemma 3** (Isolated feasibility when  $r = K$ ). *Let  $\mathcal{M}_+ = \mathcal{M} \cap \mathbb{R}_+^{n \times K}$  and  $\mathcal{M}'_+ = \mathcal{M}' \cap \mathbb{R}_+^{n \times K}$ , where  $\mathbb{R}_+^{n \times K} = \{U \in \mathbb{R}^{n \times K} : U \geq 0\}$ . Then, we have: (i)  $\mathcal{M}_+ = \mathcal{M}'_+$ ; (ii)  $U \in \mathcal{M}_+$  if and only if  $U$  is a group assignment matrix defined in (27); (iii) if  $U$  is a group assignment matrix, then the intersection of the tangent space  $T_U \mathcal{M}$  and the cone  $\mathcal{C}_U := \{V \in \mathbb{R}^{n \times K} : v_{ij} \geq 0, \forall u_{ij} = 0\}$  is trivial, i.e.,  $T_U \mathcal{M} \cap \mathcal{C}_U = \{0\}$ .*

In Lemma 5, we moreover provide a complete analytical construction for an interior point of  $\mathcal{M}$  when  $r > K$ . Here, we present the construction when  $n = qr$  is an integer multiple of  $r$ . Let  $U_0 = (x - y)I + y\mathbf{1}_n\mathbf{1}_n^\top$ , where  $x = r^{-1}(1 + \sqrt{(r-1)(K-1)})$  and  $y = r^{-1}(\sqrt{r-1} - \sqrt{K-1})$ .

Then  $U = \hat{\mathbf{1}}_q \otimes U_0$  is an interior point of  $\mathcal{M}$ . Next, we show how to compute the pair  $V, Q$  corresponding to the interior point  $U$  by SVD. For a given  $U \in \mathcal{M}_r$ , let  $U = P_U \Sigma Q_U^\top$  be the SVD of  $U$ . We can find  $(V, Q)$  such that  $U = [\hat{\mathbf{1}}_n \ V] Q$  by  $V = \text{sgn}(P_{U(1,1)})[P_U \Sigma]_{(:,2:r)}$  and  $Q = \text{sgn}(P_{U(1,1)})Q_U^\top$ .

#### C.5 Lipschitz continuity of penalty

To apply the guarantees in Section 2.3, we need to take care of the logarithmic penalty in (13) since it does not have Lipschitz gradients nor Hessians over its whole domain. The standard workaround, widely used in the analysis of nonlinear interior-point methods, is to observe that all iterates  $U_k = \varphi(V_k, Q_k)$  remain strictly feasible. Consequently, the penalty could be modified by a Huber-style smoothing, where  $\delta = \min_{i,j,k} (U_k)_{i,j} > 0$ :

$$r(x) = \begin{cases} \log x & x \geq \delta \\ \log \delta + \frac{(x - \delta)}{\delta} - \frac{(x - \delta)^2}{2\delta^2} + \frac{(x - \delta)^3}{2\delta^3} & x < \delta \end{cases}$$

The function  $r(x)$  is both concave and has Lipschitz Hessians. Therefore, the guarantees in Section 2.3 apply. The smoothing is only needed for theoretical purposes. In practice, we apply the Riemannian algorithms directly to  $\log x$ , and not to  $r(x)$ . Since we have assumed that all queries satisfy  $x \geq \delta$ , the actual behavior remains consistent with the smoothed model.

## D Proofs

*Proof of Lemma 1.* Note that the membership matrix  $Z$  associated to a partition  $G_1, \dots, G_K$  contains a diagonal principal submatrix of rank  $K$ . The lemma follows from Theorem 4 in Kalofolias and Gallopolous [26].  $\square$

*Proof of Lemma 2.* Let  $N$  denote the null space basis of  $A$ , such that  $AN = 0$  and  $N^\top N = I$ . Then, we have  $\lambda_{\min} = \lambda_{\min}(N^\top HN)$  and  $p(\lambda) = N\hat{p}(\lambda)$  where  $(N^\top HN + \lambda I)\hat{p}(\lambda) = -N^\top g$ . Then,  $\lambda > -\lambda_{\min}$  implies that  $N^\top HN + \lambda I \succ 0$ , so  $p(\lambda) = -N(N^\top HN + \lambda I)^{-1}N^\top g$  is always well-defined. Moreover,  $\|p(\lambda)\| = \|\hat{p}(\lambda)\|$  is monotonously decreasing because all the eigenvalues of  $N^\top HN + \lambda I$  are strictly positive and increasing with  $\lambda$ .  $\square$

*Proof of Lemma 3.* To prove the first statement, note that the implication

$$U \in \mathbb{R}_{\geq 0}^{n \times K} \wedge UU^\top \mathbf{1}_n = \mathbf{1}_n \wedge U^\top U = I_K \implies U \in \mathbb{R}_{\geq 0}^{n \times K} \wedge UU^\top \mathbf{1}_n = \mathbf{1}_n \wedge \|U\|_F^2 = K$$

is straightforward. To see the converse, note that  $UU^\top$  is a (doubly) stochastic matrix,  $\text{tr}(UU^\top) = K$ , and  $\text{rank}(UU^\top) = K$ , thus all the  $K$  eigenvalues of  $UU^\top$  are 1, *i.e.*  $U \in \text{St}(n, K)$ .

For the second statement, it is trivial that

$$\begin{aligned} U &= \left[ \frac{1}{\sqrt{|G_1|}} \mathbf{1}_{G_1} \quad \cdots \quad \frac{1}{\sqrt{|G_K|}} \mathbf{1}_{G_K} \right] \wedge \bigsqcup_{k=1}^K G_k = [n] \\ &\implies U \in \mathbb{R}_{\geq 0}^{n \times K} \wedge UU^\top \mathbf{1}_n = \mathbf{1}_n \wedge U^\top U = I_K. \end{aligned}$$

The converse is also true. Let us denote  $G_k := \text{supp}(u_k)$ . Observe that  $G_i \cap G_j = \emptyset$  for all  $i \neq j$  since  $U \geq 0$  and  $u_i^\top u_j = 0$  for all  $i \neq j$ . Then  $UU^\top \mathbf{1}_n = \mathbf{1}_n$  implies  $u_k u_k^\top \mathbf{1}_{G_k} = \mathbf{1}_{G_k}$  and  $\bigsqcup_{k=1}^K G_k = [n]$ . Since  $\|u_k\| = 1$ , we have that  $u_k = \mathbf{1}_{G_k} / \sqrt{|G_k|}$ .

Finally, we will prove the third statement. Let  $U$  be a group assignment matrix as defined by (27). Note that for all  $V \in T_U \mathcal{M} \cap \mathcal{C}_U$ ,  $V$  must satisfy  $(UV^\top + VU^\top)\mathbf{1}_n = \mathbf{0}_n$ ,  $\langle U, V \rangle = 0$ , and  $v_{i,j} \geq 0, \forall u_{i,j} = 0$ . Define  $A : [n] \rightarrow [K]$  to be the group assigning function, *i.e.*  $A(i) = \sum_{k=1}^K k \mathbb{I}\{u_{i,k} \neq 0\}$ , then

$$\begin{aligned} UV^\top &= \left[ \frac{1}{\sqrt{|G_{A(i)}|}} v_{j,A(i)} \right] \\ &= \begin{bmatrix} \frac{1}{\sqrt{|G_{A(1)}|}} v_{1,A(1)} & \frac{1}{\sqrt{|G_{A(1)}|}} v_{2,A(1)} & \cdots & \frac{1}{\sqrt{|G_{A(1)}|}} v_{n,A(1)} \\ \vdots & \ddots & & \vdots \\ \frac{1}{\sqrt{|G_{A(2)}|}} v_{1,A(2)} & & & \vdots \\ \vdots & & & \vdots \\ \frac{1}{\sqrt{|G_{A(n)}|}} v_{1,A(n)} & \cdots & \cdots & \frac{1}{\sqrt{|G_{A(n)}|}} v_{n,A(n)} \end{bmatrix}. \end{aligned}$$

Observe that

- (i)  $A(i) = A(j) \iff i \in G_{A(j)} \iff j \in G_{A(i)}$
- (ii)  $i \notin G_{A(j)} \iff (u_{i,A(j)} = 0 \wedge u_{j,A(i)} = 0) \implies (v_{i,A(j)} \geq 0 \wedge v_{j,A(i)} \geq 0)$
- (iii)  $\langle U, V \rangle = 0 \iff \forall j \in [n], \sum_{i \in G_{A(j)}} v_{i,A(j)} = 0$ .

Denote  $\mathbf{w} := (UV^\top + VU^\top)\mathbf{1}$ , then

$$\begin{aligned} w_j &= \sum_{i \in [n]} \left( \frac{1}{\sqrt{|G_{A(j)}|}} v_{i,A(j)} + \frac{1}{\sqrt{|G_{A(i)}|}} v_{j,A(i)} \right) \\ &= \sum_{i \in G_{A(j)}} \left( \frac{1}{\sqrt{|G_{A(j)}|}} v_{i,A(j)} + \frac{1}{\sqrt{|G_{A(i)}|}} v_{j,A(i)} \right) \\ &\quad + \sum_{i \notin G_{A(j)}} \left( \frac{1}{\sqrt{|G_{A(j)}|}} v_{i,A(j)} + \frac{1}{\sqrt{|G_{A(i)}|}} v_{j,A(i)} \right). \end{aligned}$$

By (i) and (iii), we know that

$$\sum_{i \in G_{A(j)}} \left( \frac{1}{\sqrt{|G_{A(j)}|}} v_{i,A(j)} + \frac{1}{\sqrt{|G_{A(i)}|}} v_{j,A(i)} \right) = \sqrt{|G_{A(j)}|} v_{j,A(j)}.$$

By (ii), we know that

$$\sum_{i \notin G_{A(j)}} \left( \frac{1}{\sqrt{|G_{A(j)}|}} v_{i,A(j)} + \frac{1}{\sqrt{|G_{A(i)}|}} v_{j,A(i)} \right) \geq 0.$$

Thus we can write

$$w_j = \sqrt{|G_{A(j)}|} v_{j,A(j)} + R_j \quad \text{for some } R_j \geq 0.$$

Next, we use proof by contradiction. Suppose there exists  $V \in T_U \mathcal{M} \cap \mathcal{C}_U$  such that  $V \neq 0$ , then there must be both positive and negative entries in  $V$  to satisfy  $\langle U, V \rangle = 0$ . This implies that there exists  $j_1 \in [n]$  such that  $v_{j_1,A(j_1)} < 0$  since  $v_{j,A(j)}$ 's are the only entries that can take negative value. To satisfy  $\langle U, V \rangle = 0$ , there must exist  $j_2 \in G_{A(j_1)}$  such that  $v_{j_2,A(j_1)} > 0$ . Then for such  $j_2$ ,

$$w_{j_2} = \sqrt{|G_{A(j_2)}|} v_{j_2,A(j_2)} + R_{j_2} = \sqrt{|G_{A(j_2)}|} v_{j_2,A(j_1)} + R_{j_2} > 0.$$

This contradicts  $(UV^\top + VU^\top)\mathbf{1} = \mathbf{0}$ . Therefore,  $T_U \mathcal{M} \cap \mathcal{C}_U = \{0\}$ .  $\square$

**Lemma 4.** *If  $K \geq 2$ , then the set  $\mathcal{M}$  defined in (12) satisfies linear independence constraint qualification (LICQ) for all  $U \in \mathbb{R}^{n \times r}$ , and is therefore a smooth submanifold of  $\mathbb{R}^{n \times r}$ .*

*Proof of Lemma 4.* Following Boumal, Voroninski, and Bandeira [11], the set  $\mathcal{M} = \{U \in \mathbb{R}^{n \times r} : \langle A_i, UU^\top \rangle = b_i \text{ for all } i\}$  is a smooth submanifold of  $\mathbb{R}^{n \times r}$  if LICQ holds for all  $U \in \mathcal{M}$ , i.e., that  $\sum y_i A_i U = 0$  if and only  $y = 0$ . For the definition in (12), we can verify that

$$\begin{aligned} \left\| \sum y_i A_i U \right\|_F^2 &= \|(\mathbf{1}y^\top + y\mathbf{1}^\top + y_0 I)U\|_F^2 \\ &= \|(\mathbf{1}y^\top + y\mathbf{1}^\top)U\|_F^2 + 2\langle (\mathbf{1}y^\top + y\mathbf{1}^\top)U, y_0 U \rangle + y_0^2 \|U\|_F^2 \\ &= [ny^\top(I + UU^\top)y + 2(\mathbf{1}^\top y)^2] + 4y_0(\mathbf{1}^\top y) + y_0^2 K \\ &\geq [2(\mathbf{1}^\top y)^2 + 4y_0(\mathbf{1}^\top y) + y_0^2] + n\|y\|^2 + (K-1)y_0^2 \\ &= (2\mathbf{1}^\top y + y_0)^2 + n\|y\|^2 + (K-1)y_0^2 \\ &\geq \|y\|^2 + y_0^2. \end{aligned}$$

The third line is because  $\langle (\mathbf{1}y^\top + y\mathbf{1}^\top)U, U \rangle = 2\langle y, UU^\top \mathbf{1} \rangle = 2(y^\top \mathbf{1})$  and

$$\begin{aligned} \|(\mathbf{1}y^\top + y\mathbf{1}^\top)U\|_F^2 &= \text{tr}[(\mathbf{1}y^\top + y\mathbf{1}^\top)UU^\top(\mathbf{1}y^\top + y\mathbf{1}^\top)] \\ &= \text{tr}[\mathbf{1}y^\top UU^\top y\mathbf{1}^\top + y\mathbf{1}^\top UU^\top \mathbf{1}y^\top + 2 \cdot \mathbf{1}y^\top UU^\top \mathbf{1}y^\top] \\ &= \text{tr}[\mathbf{1}y^\top UU^\top y\mathbf{1}^\top + y\mathbf{1}^\top \mathbf{1}y^\top + 2 \cdot \mathbf{1}y^\top \mathbf{1}y^\top] \\ &= ny^\top UUy + ny^\top y + 2(\mathbf{1}^\top y)^2. \end{aligned}$$

$\square$

**Lemma 5** (Interior point construction for  $\mathcal{M}_r$ ). *Given a  $K \in \mathbb{N}$ , for any  $r$  such that  $r > K$ , for large enough  $n$ , we have the following two cases:*

**Case 1:**  $n \equiv 0 \pmod{r}$

Denote  $q := n/r$ , let  $U_0 = (x - y)I + y\mathbf{1}\mathbf{1}^\top$ , where

$$x = \frac{1}{r} \left( 1 + \sqrt{(r-1)(K-1)} \right), \quad y = \frac{1}{r} \left( \sqrt{r-1} - \sqrt{K-1} \right).$$

Then  $U = (1/\sqrt{q})\mathbf{1}_q \otimes U_0$  is an interior point of  $\mathcal{M}_r$ .

**Case 2:**  $n \not\equiv 0 \pmod{r}$

Let us denote  $q := \lfloor n/r \rfloor$  and  $p := n \pmod{r}$ . Construct the block matrix  $B \in \mathbb{R}^{r \times r}$ :

$$B = \begin{bmatrix} B_{1,1} & B_{1,2} \\ B_{2,1} & B_{2,2} \end{bmatrix},$$

where

$$\begin{aligned} B_{1,1} &= (x - y)I + y\mathbf{1}_p\mathbf{1}_p^\top, & B_{1,2} &= z\mathbf{1}_p\mathbf{1}_{r-p}^\top, \\ B_{2,2} &= (w - z)I + z\mathbf{1}_{r-p}\mathbf{1}_{r-p}^\top, & B_{2,1} &= y\mathbf{1}_{r-p}\mathbf{1}_p^\top, \end{aligned}$$

The coefficients  $x, y, z$  and  $w$  depends on  $n, K$  and  $r$ . They will be specified in the proof. Then

$$U = [I_n \quad \mathbf{0}] (\mathbf{1}_q \otimes B)$$

is an interior point of  $\mathcal{M}_r$ .

*Proof of Lemma 5.* For a general large enough  $n$ ,  $n$  is either divisible or nondivisible by  $r$ . We present two different constructions of an interior point of  $\mathcal{M}_r$  corresponding to the two cases.

**Case 1:**  $n \equiv 0 \pmod{r}$

We first construct a  $U_0 \in \mathbb{R}^{r \times r}$  such that  $U_0 U_0^\top \mathbf{1}_r = \mathbf{1}_r$ , and  $\|U_0\|_F^2 = K$ . Using the ansatz  $U_0 = (x - y)I + y\mathbf{1}\mathbf{1}^\top$ , where  $x, y > 0$ , we can find  $x$  and  $y$  by solving the system:

$$\begin{cases} x + (r-1)y = 1 & (U_0 U_0^\top \mathbf{1}_r = \mathbf{1}_r) \\ x^2 + (r-1)y^2 = K/r & (\|U_0\|_F^2 = K) \end{cases}.$$

The first equation gives  $x = 1 - (r-1)y$ . By substituting into the second equation, we obtain the following quadratic equation of  $y$ :

$$r(r-1)y^2 - 2(r-1)y + 1 - \frac{K}{r} = 0.$$

By the quadratic formula and  $x, y > 0$ , we have the following solution

$$\begin{cases} x = 1 - (r-1)y = \frac{1}{r}(1 + \sqrt{(r-1)(K-1)}), \\ y = \frac{r-1 - \sqrt{(r-1)(K-1)}}{r(r-1)} = \frac{1}{r}(\sqrt{r-1} - \sqrt{K-1}). \end{cases}$$

Note that if  $r = K$ , we can still solve the quadratic equation, but without an all positive solution. Denote  $q := n/r$ , then  $U = (1/\sqrt{q})\mathbf{1}_q \otimes U_0$  is an interior point of  $\mathcal{M}_r$ .

**Case 2:**  $n \not\equiv 0 \pmod{r}$

Let us denote  $q := \lfloor n/r \rfloor$  and  $p := n \pmod{r}$ . We consider the ansatz

$$U = [I_n \quad \mathbf{0}] (\mathbf{1}_q \otimes B) \quad \text{for some block matrix } B = \begin{bmatrix} B_{1,1} & B_{1,2} \\ B_{2,1} & B_{2,2} \end{bmatrix} \in \mathbb{R}^{r \times r},$$

where

$$\begin{aligned} B_{1,1} &= (x - y)I + y\mathbf{1}_p\mathbf{1}_p^\top, & B_{1,2} &= z\mathbf{1}_p\mathbf{1}_{r-p}^\top, \\ B_{2,2} &= (w - z)I + z\mathbf{1}_{r-p}\mathbf{1}_{r-p}^\top, & B_{2,1} &= y\mathbf{1}_{r-p}\mathbf{1}_p^\top. \end{aligned}$$

Additional to the constraints  $\|U\|_F^2 = K$ , and  $UU^\top = \mathbf{1}_n$ , we assume that  $U\mathbf{1}_r = c_r\mathbf{1}_n$ ,  $U^\top\mathbf{1}_n = c_c\mathbf{1}_r$ , and  $c_r c_c = 1$  for some  $c_r$  and  $c_c$ , which are sufficient for  $UU^\top = \mathbf{1}_n$ . Then we can find  $x, y, z$ , and  $w$  by solving

the system:

$$\begin{cases} x + (p-1)y + (r-p)z = py + w + (r-p-1)z, \\ (q+1)x + q(r-1)y + (p-1)y = qw + q(r-1)z + pz, \\ (py + w + (r-p-1)z)(qw + q(r-1)z + pz) = 1, \\ (q+1)p(x^2 + (p-1)y^2 + (r-p)z^2) + q(r-p)(py^2 + w^2 + (r-p-1)z^2) = K. \end{cases} \quad (28)$$

The four equations correspond to the following constraints, respectively:  $U\mathbf{1}_r = c_r\mathbf{1}_n$ ,  $U^\top\mathbf{1}_n = c_c\mathbf{1}_r$ ,  $c_r c_c = 1$ , and  $\|U_0\|_F^2 = K$ . From the first two equations, we can express  $x$  and  $y$  in terms of  $z$  and  $w$  (note that  $n = qr + p$ ):

$$x = \left(1 - \frac{1}{n}\right)w + \frac{1}{n}z, \quad y = \left(1 + \frac{1}{n}\right)z - \frac{1}{n}w. \quad (29)$$

By substituting (29) to the third and fourth equations of (28), we are left with a system of quadratic equations of two variables:

$$\begin{cases} a_1z^2 + a_2zw + a_3w^2 + c_1 = 0, \\ b_1z^2 + b_2zw + b_3w^2 + c_2 = 0, \end{cases} \quad (30)$$

where

$$\begin{aligned} a_1 &= nr - qr + p - p(2q + 1)/n, & a_2 &= 2p(1 + 2q - n)/n, \\ a_3 &= n - p(2q + 1)/n, & b_1 &= (r + p/n - 1)(n - q), \\ b_2 &= (1 - p/n)(n - q) + q(r + p/n - 1), & b_3 &= q(1 - p/n), \\ c_1 &= -K, & c_2 &= -1. \end{aligned}$$

Now our goal is to solve (30). By multiplying the first equation with  $b_1$  and the second one with  $a_1$  and subtraction, we can express  $z$  in terms of  $w$ :

$$z = aw + \frac{b}{w} \quad \text{with } a = \frac{a_3b_1 - a_1b_3}{a_1b_2 - a_2b_1}, b = \frac{c_1b_1 - a_1c_2}{a_1b_2 - a_2b_1}. \quad (31)$$

Suppose that  $w \neq 0$ , we substitute (31) into the second equation and multiply by  $w^2$ . The result is a quintic equation:

$$(b_1a^2 + b_2a + b_3)w^4 + (2abb_1 + bb_2 + c_2)w^2 + b^2b_1 = 0. \quad (32)$$

By solving (32) with  $z = aw + b/w > 0$ , we obtain

$$w = \sqrt{\frac{-(2abb_1 + bb_2 + c_2) + \sqrt{(2abb_1 + bb_2 + c_2)^2 - 4(a^2b_1 + ab_2 + b_3)b^2b_1}}{2(a^2b_1 + ab_2 + b_3)}}. \quad (33)$$

The proof is completed by combining (29), (31), and (33).  $\square$

*Proof of Theorem 2.* For “ $\supseteq$ ,” if  $U = [\hat{\mathbf{1}}_n \ V] Q$ , then  $UU^\top = (1/n)\mathbf{1}_n\mathbf{1}_n^\top + VV^\top$  and hence  $UU^\top\mathbf{1} = \mathbf{1}$  and  $\text{tr}(UU^\top) = K$  respectively, because  $\mathbf{1}_n^\top V = 0$  and  $\text{tr}(VV^\top) = K - 1$ . For “ $\subseteq$ ,” let  $U = P\Sigma Q$  denote the singular value decomposition with  $P^\top P = QQ^\top = I_r$ . Since  $UU^\top\mathbf{1}_n = \mathbf{1}_n$ , the decomposition can be chosen so that  $P\Sigma e_1 = (1/\sqrt{n})\mathbf{1}_n = \hat{\mathbf{1}}_n$ . So if  $V = P\Sigma[e_2, \dots, e_r]$ , then  $\mathbf{1}_n^\top V = e_1^\top [e_2, \dots, e_r] = 0$  and  $\|V\|^2 = \|U\|^2 - \|P\Sigma e_1\|^2 = K - 1$ .

In the final part, we first construct the inner approximation  $S$  of the tangent space

$$\begin{aligned} T_{(V,Q)}\widetilde{\mathcal{M}} &= \{(\dot{V}, \dot{Q}) : \mathbf{1}_n^\top \dot{V} = 0, \langle V, \dot{V} \rangle = 0, Q\dot{Q}^\top + \dot{Q}Q^\top = 0\} \\ &\supseteq \left\{(\dot{V}, \dot{Q}) : \mathbf{1}_n^\top \dot{V} = 0, \langle V, \dot{V} \rangle = 0, \dot{Q}Q^\top = \begin{bmatrix} \mathbf{0} & -h^\top \\ h & \mathbf{0} \end{bmatrix} \right\} = S. \end{aligned}$$

We observe that

$$[\hat{\mathbf{1}}_n \quad V] \dot{Q} = [\hat{\mathbf{1}}_n \quad V] \begin{bmatrix} \mathbf{0} & -h^\top \\ h & \mathbf{0} \end{bmatrix} Q = [Vh \quad \hat{\mathbf{1}}_n h^\top] Q$$

and therefore the Jacobian operator is injective for all  $(\dot{V}, \dot{Q}) \in S$ :

$$\begin{aligned} \|\mathbf{D}\varphi(V, Q)[\dot{V}, \dot{Q}]\|^2 &= \|[\mathbf{0} \quad \dot{V}] Q + [\hat{\mathbf{1}}_n \quad V] \dot{Q}\|^2 \\ &= \|[\mathbf{0} \quad \dot{V}]\|^2 + \|[Vh \quad \hat{\mathbf{1}}_n h^\top]\|^2 \geq \|\dot{V}\|^2 + \|h\|^2 \geq \frac{1}{\sqrt{2}} \|(\dot{V}, \dot{Q})\|^2 \end{aligned}$$

where we used the fact that  $\mathbf{1}_n^\top \dot{V} = 0$ . Hence, the Jacobian operator is surjective, as claimed:

$$\dim(\text{image}(\mathbf{D}\varphi(V, Q))) \geq \dim(S) = n(r-1) - 1 = \dim(\mathbf{T}_U \mathcal{M}).$$

□

## E Efficient Implementation and Cost of Bisection Search

To implement the proposed method, we vectorize the input as  $u = [\text{vec}(V^\top) \quad \text{vec}(Q)]^\top$ . The cost function we used is

$$f(u) = -\|X^\top V\|^2 - \mu \mathbf{1}_n^\top \log(\varphi(V, Q)) \mathbf{1}_n,$$

and the constraint functions are defined as

$$g_1(u) = \|V\|^2 - (K-1), \quad g_2(u) = \mathbf{1}_n^\top V, \quad g_3(u) = \text{svec}(Q^\top Q - I_r).$$

The Jacobian  $J$  and Hessian  $H$  are computed analytically, as in Section C.3, in order to exploit their sparsity.

For convenience, we list some of the derivatives in this section. The (Euclidean) Jacobian of the constraints can be written in block form as

$$J = \begin{bmatrix} J_{1v} & \mathbf{0} \\ J_{2v} & \mathbf{0} \\ \mathbf{0} & J_{3q} \end{bmatrix},$$

where

$$J_{1v} = 2 \text{vec}(V^\top), \quad J_{2v} = (I_{r-1} \otimes \mathbf{1}_n^\top) K^{(r-1,n)}, \quad J_{3q} = 2\Psi_r^\top (I_r \otimes Q^\top),$$

with  $\Psi_r$  being the orthogonal basis matrix of size- $r$  symmetric matrices.

Computing the second-order derivatives of  $g_1, g_2$  are trivial, since  $J_{1v}$  is linear in  $V$  and  $J_{2v}$  is constant. The second-order derivatives of  $g_3$  can be computed explicitly by noticing that the entries of  $\text{svec}(Q^\top Q)$  can be expressed as inner products with bases outer products, i.e.

$$\langle E_{i,j}, Q^\top Q \rangle = \text{vec}(Q)^\top (E_{i,j} \otimes I_r) \text{vec}(Q),$$

where

$$E_{i,j} = \begin{cases} e_i e_j^\top & i = j, \\ \frac{e_i e_j^\top + e_j e_i^\top}{\sqrt{2}} & i \neq j. \end{cases}$$

It follows then the second-order derivative w.r.t. each entry is given by  $2(E_{i,j} \otimes I_r)$ .

Collecting results, we have

$$H = \begin{pmatrix} H_{vv} - BB^\top & H_{vq} \\ H_{qv} & H_{qq} \end{pmatrix},$$

with

$$B = \sqrt{2}(X \otimes I_{r-1}) \quad (34)$$

$$H_{vv} = \mu K^{(n,r-1)}(\hat{Q} \otimes I_n) \text{dvec}(U^{(-2)})(\hat{Q}^\top \otimes I_n)K^{(r-1,n)} + 2\tilde{y}_2 I, \quad (35)$$

$$H_{qq} = \mu(I_r \otimes \hat{V}^\top) \text{dvec}(U^{(-2)})(I_r \otimes \hat{V}) + 2(I_r \otimes \text{smat}(\tilde{y}_3)), \quad (36)$$

and

$$H_{vq} = -\mu(U^{(-1)} \otimes [\mathbf{0}_{r-1} \quad I_{r-1}]) + \mu(I_n \otimes \hat{Q}) \text{dvec}(U^{(-2)})(I_r \otimes \hat{V}) = H_{qv}^\top, \quad (37)$$

where  $\hat{Q} := [\mathbf{0}_{r-1} \quad I_{r-1}] Q$ ,  $\hat{V} := [\hat{\mathbf{1}}_n \quad V]$ , and  $\tilde{y}_2, \tilde{y}_3$  follow (26).

To solve the saddle point problem with bisection search, we solve the linear system

$$\left[ \begin{array}{c|cc} H_{vv} + \lambda I - BB^\top & H_{vq} & J_v^\top \\ \hline H_{qv} & H_{qq} + \lambda I & J_q^\top \\ J_v & J_q & \mathbf{0} \end{array} \right] \left[ \begin{array}{c} p_v \\ p_q \\ r \end{array} \right] = \left[ \begin{array}{c} -g_v \\ -g_q \\ \mathbf{0} \end{array} \right].$$

Repartition along the lines shown to yield the following

$$\left[ \begin{array}{c|c} K_{11} & K_{12} \\ \hline K_{21} & K_{22} \end{array} \right] \left[ \begin{array}{c} x_1 \\ x_2 \end{array} \right] = \left[ \begin{array}{c} b_1 \\ b_2 \end{array} \right]$$

with dimensions  $x_1, b_1 \in \mathbb{R}^{n(r-1)}$  and  $x_2, b_2 \in \mathbb{R}^{r^2+m}$ . Critically, we observe that the block  $K_{11}$ , which corresponds to the Riemannian Hessian with respect to the  $v$  variable, has the form  $K_{11} = D_{11} - BB^\top$ , where  $D_{11} = H_{vv} + \lambda I$  is block-diagonal, with  $n$  blocks of  $r-1$ , and  $B$  has at most  $dr$  columns. Therefore, we instead solve

$$\left[ \begin{array}{c|cc} D_{11} & K_{12} & B \\ \hline K_{21} & K_{22} & \mathbf{0} \\ B^\top & \mathbf{0} & I \end{array} \right] \left[ \begin{array}{c} \dot{v} \\ \dot{q} \\ z \end{array} \right] = \left[ \begin{array}{c} b_1 \\ b_2 \\ \mathbf{0} \end{array} \right].$$

First, it costs  $n(r-1)^3 = O(nr^3)$  time to invert  $D_{11}$ . Afterwards, forming and solving the size  $m + r^2 + rd$  Schur complement problem:

$$(L_{22} - L_{12}^\top D_{11}^{-1} L_{12}) \begin{bmatrix} \dot{q} \\ z \end{bmatrix} = \begin{bmatrix} b_2 \\ \mathbf{0} \end{bmatrix} - L_{12}^\top D_{11}^{-1} b_1, \quad (38)$$

where

$$L_{12} := [K_{21} \quad B] \quad L_{22} := \begin{bmatrix} K_{22} & \mathbf{0} \\ \mathbf{0} & I \end{bmatrix},$$

cost  $O(nr^3(d+r) + r^6 + r^3d^3)$  time. In the end, we substitute to recover

$$\dot{v} = D_{11}^{-1} \left( b_1 - L_{12} \begin{bmatrix} \dot{q} \\ z \end{bmatrix} \right)$$

in  $O(nr^3(d+r))$  time, and apply retractions to  $\dot{v}$  and  $\dot{q}$ . In total, it takes  $O(nr^3(d+r) + r^6 + r^3d^3)$  time to solve the system, which is indeed  $n \cdot \text{poly}(r, d)$ . Putting pieces together, a pseudo-code of our Riemannian method is shown in Algorithm 1.

## F Additional Numerical Details

We collect additional numerical results in this section.

**Hardware information.** All experiments in this work were conducted on a machine equipped with a single Intel Core i9-14900K CPU and 32 GB of RAM.

**License of assets.** The `MANOPT` solver is distributed under the terms of the GPLv3 license; the `PYMANOPT` solver is released under the 3-Clause BSD license; the CyTOF dataset is the work of Levine et al. [30], cleaned and distributed by Weber [43] under the MIT license.

**Dataset visualization.** Figure 7 and Figure 8 display the first two principal components of the GMM dataset and CyTOF dataset, respectively.




Figure 7: Visualizing the effect of the separation parameter in GMMs. As  $\gamma$  decreases, the clusters become increasingly difficult to distinguish.

**Hyperparameters tuning.** In the various numerical experiments, we observed that a smaller value of  $\mu$  led to more accurate solutions but at the cost of slower convergence. Therefore, we recommend selecting the largest possible  $\mu$  that does not trigger the phase transition. A good heuristic we found is to choose such that the initial penalty term  $\mu f_2$  remains less than 20 times the main term  $f_1$  in the loss function. The onset of phase transition is also easy to notice, as the algorithm will quickly stagnate and terminate in just a few iterations. If higher accuracy is desired, one can reduce  $\mu$  gradually, using the solution obtained with a larger  $\mu$  as initialization. This warm-start strategy significantly speeds up convergence compared to using a small  $\mu$  from the start.

The other hyperparameters in Algorithm 1 primarily influence the speed of the inner optimization. The initial multiplier  $\lambda$  affects only the number of inner steps required during the first iteration. We recommend doing a simple trial run with only two iterations; the resulting optimization history typically offers a reliable guide for choosing an appropriate initial scale for  $\lambda$ . For the other two parameters, we suggest setting  $\kappa_-$  slightly smaller than  $\kappa_+$ . Empirically, we found  $\kappa_- = 1.1$  and  $\kappa_+ = 1.3$  work well.

**Robustness to initialization.** As illustrated by Figure 1, our method is robust to initialization, all 50 trials successfully converged to second-order optimal solutions. Although the solutions differ (Figure 9), their corresponding membership matrices  $Z$  are close to each other (Figure 10), and yield identical clustering result. Moreover, the minimum eigenvalues upon convergence form distinct clusters that align with clusters in the recovered membership matrix  $Z$ , as shown in Figure 10. These local critical points consistently produce perfect clustering, indicating that they remain close to the global optimum.




Figure 8: Visualization of the CyTOF dataset. Two clusters exhibit significant overlap, implying the difficulty of clustering.




Figure 9: **Difference between the solutions.** First 25 rows of selected solution  $U$  obtained from the global optimality experiment described in Section 4.

**Comparison with another Riemannian clustering method.** Inexact Accelerated Manifold Proximal Gradient Method (I-AManPG) by Huang et al. [25] is a recent first-order Riemannian method for solving general problems of the form

$$\begin{aligned} \min_X \quad & f(X) + \lambda \|X\|_1, \\ \text{s.t.} \quad & X \in \mathcal{F}_v := \{X : X^\top X = I, v \in \text{span}(X)\}. \end{aligned} \tag{39}$$

We evaluated its performance on the clustering problem using the on the CyTOF dataset with 50 repetitions (same settings as in Figure 2). The results are shown in Figure 11. While I-AManPG is generally fast and




Figure 10: **Similarities of the membership matrices.** Frobenius distances between the membership matrices  $Z$  obtained from the global optimality experiment in Section 4, sorted according to their corresponding minimum Hessian eigenvalues.

accurate, its median error is higher than that of our methods. In particular, several runs of I-AManPG exhibited large errors, indicating convergence failures.




Figure 11: **Comparison with I-AManPG using CyTOF.** Performance of I-AManPG is comparable to that of other clustering methods; however, it suffers from convergence failures occasionally and requires careful parameter tuning. In contrast, our method consistently demonstrates both accuracy and stability.

**Additional convergence plots.** Figure 12, Figure 13, and Figure 14 illustrate the convergence of our method on GMM with different parameters and on the CyTOF dataset, demonstrating its stability across different datasets.




Figure 12: Convergence of our method on synthetic Gaussian mixture data with perfect separation ( $n = 500, \gamma = 1.0$ ). The loss value steadily decreases over iterations and converges rapidly near the optimal point. This example achieved a perfect final clustering result in the end.




Figure 13: Convergence of our method on synthetic Gaussian mixture data with low separation ( $n = 500, \gamma = 0.25$ ).




Figure 14: Convergence behavior on real world CyTOF dataset.

## G Pseudocode

This section presents the pseudocode of our implementation. For the derivation, see Section E.

---

**Algorithm 1** Riemannian Second-order Method

---

**Require:** Data  $X$ ,

Initial point  $(V_0, Q_0)$ ,  
 Initial multiplier  $\lambda$ ,  
 Increment/decrement factors  $(\kappa_+, \kappa_-)$ ,  
 log-barrier penalty  $\mu$ ,  
 Max number of outer/inner iterations  $T$  and  $B$ .

```

1:  $(V, Q) \leftarrow (V_0, Q_0)$ 
2: for  $i = 1, \dots, T$  do
3:   Vectorize input  $u \leftarrow [\text{vec}(V^\top), \text{vec}(Q)]^\top$ .
4:   Compute current loss  $\mathcal{L} = f(u)$ .
5:   Compute Riemannian gradient  $d \leftarrow \nabla f(u)$  and Jacobian  $J \leftarrow Dg(u)$ .
6:   Compute Riemannian Hessians  $H_f \leftarrow \nabla^2 f(u)$  and  $H_i \leftarrow D^2 g_i(u)$ .
7:   Solve  $y_u = \min_y \|d + Jy\|$  by least-squares.
8:   Compute  $\nabla_u \mathcal{L}(u, y_u)$ :  $G \leftarrow d + Jy_x$ 
9:   Compute  $\nabla_u^2 \mathcal{L}(u, y_u)$ :  $H \leftarrow H_f + \sum_{i=1}^3 (y_u)_i H_i$ 
10:  for  $j = 1, \dots, B$  do
11:     $\dot{v}, \dot{q} \leftarrow \text{SOLVEINNER}(H, G, J, \lambda)$ 
12:    Reconstruct  $\dot{V}, \dot{Q}$  from vector  $\dot{v}, \dot{q}$ 
13:     $(V, Q) \leftarrow \text{Retr}_{(V, Q)}(\dot{V}, \dot{Q})$ 
14:    Compute new loss  $\mathcal{L}'$  from  $(V, Q)$ 
15:    if  $\mathcal{L} > \mathcal{L}'$  then
16:       $\lambda \leftarrow \lambda / \kappa_-$ 
17:      break
18:    else
19:       $\lambda \leftarrow \lambda \cdot \kappa_+$ 
20:  output  $(V, Q)$ 
21:
22: function  $\text{SOLVEINNER}(H, G, J, \lambda)$ 
23:   Adding  $\lambda I$  to the  $V$  and  $Q$  blocks of  $H$ 
24:   Forming block matrices as in (38)
25:    $S \leftarrow L_{22} - L_{12}^\top D_{11}^{-1} L_{12}$ 
26:    $\begin{bmatrix} \dot{q} \\ z \end{bmatrix} \leftarrow \begin{bmatrix} b_2 \\ 0 \end{bmatrix} - L_{12}^\top D_{11}^{-1} b_1$ 
27:    $\dot{v} \leftarrow D_{11}^{-1} \left( b_1 - L^{12} \begin{bmatrix} \dot{q} \\ z \end{bmatrix} \right)$ 
28:   return  $\dot{v}, \dot{q}$ 

```

---