

FIG. 1. Star-shaped oscillation patterns of Leidenfrost drops on a curved, aluminum surface ( $T = 350$  °C). Oscillation modes with  $n = 2$  to  $n = 13$  are shown when the lobes are at their maximum displacement. The scale bar (2 cm) applies to all images. Source: APS-DFD (<http://gfm.aps.org/meetings/dfd-2014/541347fc69702d585c400000>)

## The many faces of a Leidenfrost drop

Xiaolei Ma,<sup>a)</sup> Juan-José Liétor-Santos, and Justin C. Burton

*Department of Physics, Emory University, Atlanta, Georgia 30322, USA*

<sup>a)</sup> Electronic mail: [xiaolei.ma@emory.edu](mailto:xiaolei.ma@emory.edu)

When deposited on a hot surface well above its boiling point, a liquid drop can survive for minutes by levitating on an insulating cushion of evaporated vapor. This phenomenon, known as the Leidenfrost effect,<sup>1</sup> can be easily observed on a hot pan over a cook stove. The vapor layer is maintained by a sustained evaporation of the liquid beneath the drop,<sup>2,3</sup> resulting in near-frictionless motion on the

surface. These properties can lead to a variety of interesting dynamics,<sup>4, 5</sup> such as self-propulsion on textured surfaces,<sup>6</sup> and self-organized “star-shaped” oscillations of liquid puddles.<sup>7</sup> The latter depends on a sensitive coupling between deformations of the liquid/vapor interface and lubrication flow in the thin ( $\approx 100 \mu\text{m}$ ) vapor layer.

We investigated star-shaped oscillation modes of water drops on a hot aluminum surface with mode numbers from  $n = 2$  to  $n = 13$ . We used a high-speed camera (Phantom V7.11) to image the drops from above, as shown in Fig. 1. The aluminum surface was curved in order to keep drops stationary and suppress the buoyancy-driven Rayleigh-Taylor instability in the vapor layer.<sup>4</sup> Surprisingly, we found that all of the oscillation modes in Fig. 1 share the same azimuthal wavelength ( $\lambda \approx 1.2 \text{ cm}$ ). The frequency of the oscillations is nearly constant ( $f \approx 13 \text{ Hz}$ ) as can be seen in Fig. 2a, and is consistent with a quasi-2D dispersion relation.<sup>8</sup>




FIG. 2. (a) Star-shaped oscillation frequency and peak pressure frequency for different mode numbers. Each data point corresponds to an individual drop. The error bar is defined as the half width of the highest peak in (c). (b) and (c) represent the variations and corresponding Fast Fourier Transform of the pressure in the vapor layer beneath a Leidenfrost drop during an  $n = 8$  oscillation within 1 s and 20 s, respectively.

In order to gain insight into the nature of the star-shaped oscillations, we used a pressure sensor (GEMS Sensors) connected to a small, 1 mm hole at the center of the curved aluminum surface to detect pressure variations in the vapor layer directly beneath the drop. Fig. 2b shows the measured pressure during a 1 s interval of an  $n = 8$  mode, which contains multiple frequency components. Fig. 2c shows the Fast Fourier Transform (FFT) of the pressure data over a 20 s interval, indicating that the highest peak is located at approximately 26 Hz, which is twice the oscillation frequency of the drop. This relationship

holds true for all the modes we probed as shown in Fig. 2a. Note that higher harmonics are also visible, suggesting a nonlinear mechanism for generating the star-shaped oscillations.

Similar star-shaped oscillations have also been observed in driven systems, which is due to a parametric coupling mechanism between the azimuthal oscillations of drops and vertical external forces.<sup>7</sup> Our results exhibit the same characteristics of the parametric coupling in the absence of external forcing fields. However, the origin of the pressure oscillations observed in our experiment remains unclear. We hypothesize that capillary waves excited underneath the drop lead to the pressure variations thereby generating the star-shaped oscillations. Although we do not currently have a model for these capillary waves, they may be related to “brim waves” which exist in viscous drops levitated by an air flow through a porous mold.<sup>9</sup>

We gratefully acknowledge financial support from NSF DMR-1455086.

<sup>1</sup> J. G. Leidenfrost, *De aquae communis nonnullis qualitatibus tractatus* (Ovenius, 1756).

<sup>2</sup> J. Burton, A. Sharpe, R. van der Veen, A. Franco, and S. Nagel, "Geometry of the vapor layer under a Leidenfrost drop," *Phys. Rev. Lett.* **109**, 074301 (2012).

<sup>3</sup> J. H. Snoeijer, P. Brunet, and J. Eggers, "Maximum size of drops levitated by an air cushion," *Phys. Rev. E* **79**, 036307 (2009).

<sup>4</sup> D. Quéré, "Leidenfrost dynamics," *Annu. Rev. Fluid Mech.* **45**, 197 (2013).

<sup>5</sup> T. A. Caswell, "Dynamics of the vapor layer below a Leidenfrost drop," *Phys. Rev. E* **90**, 013014 (2014).

<sup>6</sup> H. Linke, B. Alemán, L. Melling, M. Taormina, M. Francis, C. Dow-Hygelund, V. Narayanan, R. Taylor, and A. Stout, "Self-propelled Leidenfrost droplets," *Phys. Rev. Lett.* **96**, 154502 (2006).

<sup>7</sup> P. Brunet, and J. H. Snoeijer, "Star-drops formed by periodic excitation and on an air cushion—A short review," *Eur. Phys. J. Spec. Top.* **192**, 207 (2011).

<sup>8</sup> R. Takaki, and K. Adachi, "Vibration of a flattened drop. ii. normal mode analysis," *J. Phys. Soc. Jpn.* **54**, 2462 (1985).

<sup>9</sup> L. Duchemin, J. Lister, and U. Lange, "Static shapes of levitated viscous drops," *J. Fluid Mech.* **533**, 161 (2005).