

Scaling Theory of Entanglement at the Many-Body Localization Transition

Philipp T. Dumitrescu,^{1,*} Romain Vasseur,^{2,3,4} and Andrew C. Potter¹

¹*Department of Physics, University of Texas at Austin, Austin, Texas 78712, USA*

²*Department of Physics, University of California, Berkeley, California 94720, USA*

³*Materials Science Division, Lawrence Berkeley National Laboratories, Berkeley, California 94720, USA*

⁴*Department of Physics, University of Massachusetts, Amherst, Massachusetts 01003, USA*

(Received 23 January 2017; published 14 September 2017)

We study the universal properties of eigenstate entanglement entropy across the transition between many-body localized (MBL) and thermal phases. We develop an improved real space renormalization group approach that enables numerical simulation of large system sizes and systematic extrapolation to the infinite system size limit. For systems smaller than the correlation length, the average entanglement follows a subthermal volume law, whose coefficient is a universal scaling function. The full distribution of entanglement follows a universal scaling form, and exhibits a bimodal structure that produces universal subleading power-law corrections to the leading volume law. For systems larger than the correlation length, the short interval entanglement exhibits a discontinuous jump at the transition from fully thermal volume law on the thermal side, to pure area law on the MBL side.

DOI: 10.1103/PhysRevLett.119.110604

Recent experimental advances in synthesizing isolated quantum many-body systems, such as cold atoms [1–4], trapped ions [5,6], or impurity spins in solids [7,8], have raised fundamental questions about the nature of statistical mechanics. Even when decoupled from external sources of dissipation, large interacting quantum systems tend to act as their own heat baths and reach thermal equilibrium. This behavior is formalized in the eigenstate thermalization hypothesis (ETH) [9,10]. Generic excited eigenstates of such thermal systems are highly entangled, with the entanglement of a subregion scaling as the volume of that region (volume law). This results in incoherent, classical dynamics at long times. In contrast, strong disorder can dramatically alter this picture by pinning excitations that would otherwise propagate heat and entanglement [11–17]. In such many-body localized (MBL) systems [18–20], generic eigenstates have properties akin to those of ground states. They exhibit short-range entanglement that scales like the perimeter of the subregion [17] (area law), and have quantum coherent dynamics up to arbitrarily long time scales [21–27], even at high energy densities [17,26,28–31].

A transition between MBL and thermal regimes requires a singular rearrangement of eigenstates from area-law to volume-law entanglement. This many-body (de)localization transition (MBLT) represents an entirely new class of critical phenomena, outside the conventional framework of equilibrium thermal or quantum phase transitions. Developing a systematic theory of this transition promises not only to expand our understanding of possible critical phenomena, but also to yield universal insights into the nature of the proximate MBL and thermal phases.

The eigenstate entanglement entropy can be viewed as a nonequilibrium analog of the thermodynamic free energy

for a conventional thermal phase transition, and plays a central role in our conceptual understanding of the MBL and ETH phases. Describing the entanglement across the MBLT requires addressing the challenging combination of disorder, interactions, and dynamics. Consequently, most studies have resorted to fully microscopic simulation methods like exact diagonalization (ED) [15,16,32–34]. The exponential complexity of such methods fundamentally limits them to small systems ($\lesssim 30$ sites), preventing them from accurately capturing universal scaling properties. For example, critical exponents computed from ED violate rigorous scaling bounds [35,36].

A promising alternative is to eschew a microscopic description, which is not required to compute universal scaling properties, and instead develop a coarse grained renormalization group (RG) description. Two related RG approaches [37,38] have produced a consistent picture of the MBLT (see also [39]). Nonetheless, both approaches rest on *ad hoc* albeit plausible heuristics for computing many-body matrix elements. In this paper, we develop a RG scheme building upon [38], but whose steps are rooted in well-established properties of matrix elements in MBL and thermal systems. Using this modified RG scheme, we compute the full scaling structure of entanglement across the transition, by simulating large systems sizes with many (10^5 – 10^6) disorder realizations that allow systematic extrapolation to the infinite size limit. The resulting scaling properties depart dramatically from those of conventional equilibrium critical points, highlighting the unusual nature of the MBLT.

RG approach.—Our RG approach builds a coarse-grained picture of eigenstates by identifying collective many-body resonances that destabilize the MBL phase. Although this approach is not tied to a particular microscopic

model, we picture a chain of spinless fermions with Hamiltonian $H = \sum_x (-c_x^\dagger c_{x+1} + \text{H.c.} - \mu_x \rho_x + V \rho_x \rho_{x+1})$; here $\rho_x = c_x^\dagger c_x$ is the fermion density on site x , and μ_x is a random chemical potential drawn from a uniform distribution on $[0, W]$. The noninteracting system ($V = 0$) is Anderson localized with localization length $x_0 \approx 2/\log(1 + W^2)$ [38]. Interactions ($|V| > 0$) can drive multi-particle collective resonances. For weak interactions, $V \ll W$, the system remains MBL and these resonances restructure the local integrals of motion (LIOM) from weakly dressed single-particle orbitals to few-body LIOM [40–43]. For sufficiently strong interactions, MBL breaks down as all degrees of freedom resonate.

While finding the true resonances is tantamount to solving the many-body Hamiltonian, close to the continuous MBLT, one expects a scale-invariant structure in which resonances are organized hierarchically and can be constructed iteratively [38]. Since large many-fermion resonances will drive the MBLT, it is natural to consider an effective model in terms of resonant clusters, i.e., groups of inter-resonating single-particle orbitals, characterized only by coarse grained information: the effective bandwidth Λ_i and the typical level spacing δ_i .

To characterize cluster interactions, we retain only the typical amplitude Γ_{ij} of matrix elements for transitions changing the states of clusters i and j , and compare this to the corresponding typical energy mismatch ΔE_{ij} between those states. For $\Gamma_{ij} \gg \Delta E_{ij}$, states of i and j will resonantly admix, whereas for $\Gamma_{ij} \ll \Delta E_{ij}$, the clusters will remain decoupled apart from weak virtual dressing. We divide these regimes sharply and define a resonant coupling if $\Gamma_{ij} > \Delta E_{ij}$. The ambiguity of this partition becomes unimportant for the large clusters determining the transition, since both Γ_{ij} and ΔE_{ij} depend exponentially on fluctuating extensive quantities, and are rarely comparable.

The RG procedure for a chain of L sites with periodic boundary conditions proceeds as follows. Initially, each cluster corresponds to a localized single-particle orbital with bandwidth $\Lambda_i = \epsilon_i \approx \mu_i$ (ϵ_i being the noninteracting single-particle energy), $\Delta E_{ij} = |\mu_i - \mu_j|$, and $\Gamma_{ij} = V(e^{-|i-j|/x_0} + e^{-|i-j-L|/x_0})$. We set $V = 0.3$ throughout. During a RG step, all clusters connected by a path of resonating bonds are merged into a new cluster $\{i\} \rightarrow i'$. The coarse grained parameters of the newly formed cluster are chosen as [44]: $\Lambda_{i'} = [\sum_i \Lambda_i^2 + \sum_{ij} \Gamma_{ij}^2]^{1/2}$, $\delta_{i'} = \Lambda_{i'}/(2^{n_{i'}} - 1)$, and $\Delta E_{i'j'} = \delta_{i'} \delta_{j'}/\min(\Lambda_{i'}, \Lambda_{j'})$ where $n_{i'}$ is the number of sites in cluster i' .

The effective intercluster couplings are changed according to two distinct rules, locally mirroring MBL or ETH behavior (Fig. 1). First, consider two clusters not modified during a RG step. In isolation, these clusters would form a small MBL region, with decoupled LIOM that project onto the separate states of each cluster. Any further resonance




FIG. 1. Schematic of a RG step. Eight initial clusters (dashed squares) interact with each other; those connected by a resonant path ($\Gamma_{ij} > \Delta E_{ij}$) merge into bigger, new clusters (colored rectangles). The coupling between new clusters is turned off or renormalized from the previous step (see the text).

between these two clusters must be mediated by other clusters; we can therefore neglect the direct coupling between them and set $\Gamma_{i'j'} = 0$. Second, if at least one of the clusters is modified during the RG step, the new coupling between two clusters is [45]

$$\Gamma_{i'j'} = \left[\max_{i_1 \in \{i\}, i_2 \in \{j\}} \Gamma_{ij} \right] e^{-(n_{i'} + n_{j'} - n_{i_1} - n_{i_2})s_{\text{th}}/2}. \quad (1)$$

Here, $\max \Gamma$ selects the strongest resonating pathway. The exponential factor approximates the resonating clusters as small locally thermal subsystems with entropy $s_{\text{th}} = \log 2$ per site. This form holds for matrix elements of local operators in a finite-size, ETH system [9,45].

The renormalization of intercluster couplings is different from those of [37,38], but has similarities to those of [46]. The coupling Γ_{ij} sets the time scale over which clusters can resonate to change each other's state. Early in the RG, resonances are fast and occur directly between a few strongly coupled sites. Later in the RG, resonances are more collective and involve many sites. Although the direct coupling Γ_{ij} is set to zero if two clusters cannot resonate at a given time scale, they can still resonate later, if mediated via coupling to other clusters [45].

Approximating Γ_{ij} by the limiting MBL and ETH forms becomes self-consistently justified since the width of the distribution of resonance parameters $g_{ij} = \Gamma_{ij}/\Delta E_{ij}$ [37,38,47] increases with each RG step. In an infinite critical system, the width of the distribution of g increases without bound along the RG flow so that one asymptotically encounters only the cases $g \ll 1$ (MBL) or $g \gg 1$ (ETH) and almost never faces marginal cases where $g \approx 1$. This flow to infinite randomness of g justifies the RG approximations in an analogous fashion to other microscopic RG approaches for quantum phase transitions in disordered spin chains [30,48–51].

Rooting the Γ_{ij} renormalization in well-established asymptotic properties more accurately captures the competition between locally MBL regions being thermalized by nearby locally thermal clusters, or isolating them. These rules cleanly prevent unphysical “avalanche” instabilities of the MBL phase [46,52] in which an atypically large resonant cluster becomes increasingly thermal as it grows, enabling it to thermalize an arbitrarily large MBL region [53].




FIG. 2. Universal scaling of bipartite entanglement. Normalized bipartite entanglement $\hat{s}(L/2)$ as a function of disorder bandwidth W for different system sizes L . Inset: scaling collapse of $\hat{s}(L/2)$ (upper) and fluctuations $\sigma_{\hat{s}(L/2)}$ (lower), with $W_c = 2.05$ and $\nu = 3.2$. Data with $L \leq 1000$ have 2.5×10^5 or 10^6 disorder realizations; those with $L \geq 1500$ have 10^5 . Error bars were calculated using jackknife resampling, but are not shown when smaller than marker sizes.

The RG terminates if no resonant bonds remain or the system fully thermalizes. Like [38], our approach allows for a distribution of various cluster sizes in the final configuration. This feature is important, as typical configurations at criticality are predominantly MBL with few large clusters [38]—a picture supported by recent ED numerics [34]. In contrast, the approach of [37] allowed both MBL and thermal blocks (clusters) to grow until the system is one large block that is either thermal or MBL.

Half-system entanglement at criticality.—For each disorder realization, the RG produces a configuration of decoupled locally thermal clusters. We calculate the entanglement of a subinterval by summing the thermal volume law contribution from each cluster spanning the interval boundaries. A cluster partitioned into m and n sites contributes $S_{m,n} = \min(m, n)s_{\text{th}}$.

Figure 2 depicts the normalized entanglement entropy, $\hat{s}(x) = \overline{S(x, L)}/xs_{\text{th}}$, for $x = L/2$, where $\overline{(\dots)}$ denotes averaging over disorder realizations and interval location. It shows the transition from a fully thermal system consisting of a single large cluster to the localized system made from many small clusters, indicated by curves of different L crossing at critical disorder $W_c = 2.05 \pm 0.01$. The curves satisfy a scaling form $\hat{s} = f([W - W_c]L^{1/\nu})$, with critical exponent $\nu = 3.2 \pm 0.3$ (Fig. 2 upper inset). This indicates the presence of a single diverging correlation length $\xi \approx |W - W_c|^{-\nu}$. A variety of observables give the same estimates of W_c and ν and our extracted ν lies within error bars of those obtained in [37,38]. Notably, we find two distinct values of ν for average and typical correlation length exponents $\nu_{\text{typ}} \approx 2.1 \pm 0.2$ [45], consistent with a flow to infinite randomness. Together with the small value \hat{s} at the




FIG. 3. Bipartite entanglement at criticality. Normalized histogram over disorder realizations of the bipartite entanglement entropy near criticality ($W = 2.04$), using 100 linearly spaced bins. Inset: scaling collapse of the fraction of fully thermalized configurations $N_{\text{th}}/N_{\text{tot}}$; error bars are 95% confidence intervals expected for binomial distribution.

crossing, this demonstrates that the transition is driven by rare thermal clusters separated by large MBL regions.

Figure 3 shows the full histogram of entanglement over disorder realizations at W_c . The distribution has a bimodal structure consisting of a power-law tail, $P(s) \approx s^{-\alpha}$ with $\alpha = 1.4 \pm 0.2$, fit over the interval $s \in [0.1, 0.8]$, and a distinct sharp peak near the fully thermal value $s = 1$. Away from criticality, the weight of the thermal peak scales like a universal function of L/ξ (Fig. 3 inset). Indications of a bimodal structure were observed in small-scale ED simulations [54]. Our RG approach allows an extensive exploration of this structure.

At criticality, the thermal peak gives a volume-law contribution to the bipartite entanglement with a coefficient $a = (0.8 \pm 0.3) \times 10^{-2}$ far below the thermal value. The power-law component gives a universal subleading power-law contribution intermediate between area and volume law,

$$\overline{S(x = L/2, L, W = W_c)} \approx ax + bx^{1-\alpha} + \dots \quad (2)$$

These results differ from those of [37], whose proxy for half-system entanglement showed a smaller power law [$P(s) \approx s^{-0.9}$] and lacked a thermal peak.

Nonlocal influence of system size.—Consider next an infinite system slightly away from the critical point. Near a conventional continuous phase transition, observables (including entanglement) measured over distance x exhibit critical behavior over an extended “critical fan” $x \ll \xi$ extending across both sides of the transition. Moreover, they become independent of system size as $L \rightarrow \infty$, since critical fluctuations are determined by local physics. Entanglement at the MBLT departs dramatically from this conventional behavior, and instead shows a strong nonlocal dependence on system size, since an infinite thermal

system can act as a bath for any finite subsystem, no matter its local properties. Hence, all subintervals of an infinite system must exhibit fully thermal entanglement $\hat{s}(x, L = \infty) = 1$ for $W < W_c$ and $L \gg \xi$ [55]. The conventional scaling picture would then suggest full thermal entanglement also on the MBL side ($W > W_c$) for $x \ll \xi$ [55]. Instead, ED simulations in [34] give evidence that this region actually has subthermal entanglement, consistent with the picture of [38] that the critical regime mainly contains large MBL regions. Together with [55], this implies that the entanglement jumps discontinuously from fully thermal to subthermal across the MBLT for $L = \infty$ [34].

Our RG approach can directly demonstrate this predicted discontinuity by systematically extrapolating to the limit $L \rightarrow \infty$ with $x \ll \xi \ll L$. Figure 4 shows the normalized entanglement for a fixed interval $x = 10$ and various system sizes L . While one can never observe a true discontinuity in a finite-size system, we observe a clear finite-size flow towards a nonanalytic jump with increasing L . Similar $L \rightarrow \infty$ extrapolations are obtained for all x .

This discontinuous jump establishes that the entanglement on the MBL side is subthermal for all x . However, many functional forms are consistent with this requirement. Unlike the thermal behavior for $L = \infty$ and $W < W_c$, which follows from analytic constraints [55], determining the entanglement scaling for $W > W_c$ requires a threefold hierarchy of scales $1 \ll x \ll \xi \ll L$ (Fig. 5). This necessitates large systems with at least $\mathcal{O}(10^3)$ sites, making our RG approach uniquely suited to address this question.

Having an objective measure of the correlation length ξ is vital to identify the desired scaling regime and separate it from the distinct crossover behavior when $\xi \approx L$. To this end, we examine the distribution of cluster sizes, which




FIG. 4. Infinite system entanglement. The normalized entanglement entropy for an interval $x = 10$ develops a nonanalytic step on the thermal side of the MBLT as $L \rightarrow \infty$. Points labeled $+\infty$ are extrapolations in L , assuming the leading scaling form $\propto L^{1/\nu}$ along fixed \hat{s} . Cubic spline interpolation was used between data points. The error bars reflect the uncertainty in $\nu = 3.2 \pm 0.3$. The transition $W_c = 2.05 \pm 0.1$ is indicated by the dashed line and gray shaded region.

exhibit power-law decay up to a scale that we identify as ξ , beyond which they decay exponentially [45]. For $L \gg \xi$, the entanglement curves show a small nonuniversal rise over $x \lesssim 1-10$ and then remain perfectly flat as x crosses through ξ , indicating that the entanglement follows a pure area law everywhere on the MBL side of the transition, even for $x \ll \xi$.

The absence of scaling on the MBL side is particular to the disorder averaged entanglement, for which critical fluctuations affect only subleading terms that vanish for large L . Other observables, like higher moments of entanglement, can exhibit universal power-law singularities as $W \rightarrow W_c^+$. We also note that the discontinuous behavior of entanglement for $L \rightarrow \infty$ is special to static eigenstate properties (equivalently, infinite time averaged quantities). In contrast, due to the logarithmic causal cone for dynamics at the MBLT [37,38], dynamical measurements on time scales $\log t \ll L$ are insensitive to the system size, and exhibit a more conventional critical scaling fan.

Full scaling form of $S(x, L, W)$.—For infinite systems, we have seen that the entanglement jumps discontinuously at the MBLT. For finite L , this jump becomes a smooth crossover. What universal data can we extract from this crossover? The entanglement is itself generically not a scaling variable. In addition to nonuniversal, subleading terms, different parts of the entanglement may be universal




FIG. 5. Entanglement finite-size crossover. (a) Entanglement entropy as a function of interval size x for system size $L = 1000$ and various $W > W_c$. The error bars correspond to the vertical thickness of the curves. The black points are lower bounds on the estimate of ξ taken from the cluster size histograms. (b) Normalized entanglement entropy for $L = 4000$ for different fractions of system size $f = x/L$. (c) Plot as in (a) zoomed out for disorder values $W = 1.4$ (yellow, linear volume law) and $1.96 \leq W \leq 2.3$ in steps of 0.02.

for different critical points; identifying an appropriate scaling form is not straightforward. For example, in one-dimensional conformal field theories one needs to consider $\partial S/\partial \log x$ in the limit $x, L \gg 1$ [56,57].

By performing scaling collapses of $S(x, L)$ for fixed x/L and various W [45], we find evidence that the volume law coefficient is a universal scaling function

$$\hat{s}(x, L) = \frac{\overline{S(x, L)}}{xs_{\text{th}}} = \mathcal{A}\left(\frac{x}{\xi}, \frac{L}{\xi}, \text{sgn}\delta W\right) + (\dots). \quad (3)$$

Here (\dots) indicates subleading corrections in x and L that vanish in the scaling limit $x, L \gg 1$. The scaled form as the function of the variables $x/L, L/\xi \text{sgn } \delta W$ is shown in Fig. 5(b). At finite L , the above scaling form with a single universal exponent ν is relatively conventional. The large L limit, however, is different from the scaling of conventional correlation functions. The nonlocal system size dependence shown above implies that in the limit $L/\xi \rightarrow \infty$, \mathcal{A} depends only on $\text{sgn}\delta W$; there is absolutely no dependence on x/ξ . The striking discrepancy in scaling highlights the unusual and asymmetric nature of thermalization and the MBL transition.

We thank D. Huse and T. Grover for insightful discussions and especially S. A. Parameswaran for collaboration on previous work. We acknowledge the Texas Advanced Computing Center (TACC) at the University of Texas at Austin for computational resources and LBNL Quantum Materials program for support (R. V.).

*philippd@utexas.edu

- [1] I. Bloch, J. Dalibard, and W. Zwerger, *Rev. Mod. Phys.* **80**, 885 (2008).
- [2] M. Schreiber, S. S. Hodgman, P. Bordia, H. P. Lüschen, M. H. Fischer, R. Vosk, E. Altman, U. Schneider, and I. Bloch, *Science* **349**, 842 (2015).
- [3] J.-y. Choi, S. Hild, J. Zeiher, P. Schauß, A. Rubio-Abadal, T. Yefsah, V. Khemani, D. A. Huse, I. Bloch, and C. Gross, *Science* **352**, 1547 (2016).
- [4] H. P. Lüschen, P. Bordia, S. Scherg, F. Alet, E. Altman, U. Schneider, and I. Bloch, [arXiv:1612.07173](https://arxiv.org/abs/1612.07173).
- [5] J. Smith, A. Lee, P. Richerme, B. Neyenhuis, P. W. Hess, P. Hauke, M. Heyl, D. A. Huse, and C. Monroe, *Nat. Phys.* **12**, 907 (2016).
- [6] J. Zhang, P. W. Hess, A. Kyprianidis, P. Becker, A. Lee, J. Smith, G. Pagano, I. D. Potirniche, A. C. Potter, A. Vishwanath, N. Y. Yao, and C. Monroe, *Nature (London)* **543**, 217 (2017).
- [7] S. Choi, J. Choi, R. Landig, G. Kucska, H. Zhou, J. Isoya, F. Jelezko, S. Onoda, H. Sumiya, V. Khemani, C. von Keyserlingk, N. Y. Yao, E. Demler, and M. D. Lukin, *Nature (London)* **543**, 221 (2017).
- [8] K. X. Wei, C. Ramanathan, and P. Cappellaro, [arXiv: 1612.05249](https://arxiv.org/abs/1612.05249).
- [9] M. Srednicki, *Phys. Rev. E* **50**, 888 (1994).
- [10] J. M. Deutsch, *Phys. Rev. A* **43**, 2046 (1991).
- [11] P. W. Anderson, *Phys. Rev.* **109**, 1492 (1958).
- [12] L. Fleishman and P. W. Anderson, *Phys. Rev. B* **21**, 2366 (1980).
- [13] I. V. Gornyi, A. D. Mirlin, and D. G. Polyakov, *Phys. Rev. Lett.* **95**, 206603 (2005).
- [14] D. Basko, I. Aleiner, and B. Altshuler, *Ann. Phys. (N.Y.)* **321**, 1126 (2006).
- [15] V. Oganesyan and D. A. Huse, *Phys. Rev. B* **75**, 155111 (2007).
- [16] A. Pal and D. A. Huse, *Phys. Rev. B* **82**, 174411 (2010).
- [17] B. Bauer and C. Nayak, *J. Stat. Mech.* 2013, P09005 (2013).
- [18] R. Nandkishore and D. A. Huse, *Annu. Rev. Condens. Matter Phys.* **6**, 15 (2015).
- [19] E. Altman and R. Vosk, *Annu. Rev. Condens. Matter Phys.* **6**, 383 (2015).
- [20] R. Vasseur and J. E. Moore, *J. Stat. Mech.* 2016, 064010 (2016).
- [21] M. Znidaric, T. Prosen, and P. Prelovsek, *Phys. Rev. B* **77**, 064426 (2008).
- [22] J. H. Bardarson, F. Pollmann, and J. E. Moore, *Phys. Rev. Lett.* **109**, 017202 (2012).
- [23] M. Serbyn, Z. Papić, and D. A. Abanin, *Phys. Rev. Lett.* **110**, 260601 (2013).
- [24] M. Serbyn, M. Knap, S. Gopalakrishnan, Z. Papić, N. Y. Yao, C. R. Laumann, D. A. Abanin, M. D. Lukin, and E. A. Demler, *Phys. Rev. Lett.* **113**, 147204 (2014).
- [25] R. Vasseur, S. A. Parameswaran, and J. E. Moore, *Phys. Rev. B* **91**, 140202 (2015).
- [26] Y. Bahri, R. Vosk, E. Altman, and A. Vishwanath, *Nat. Commun.* **6**, 7341 (2015).
- [27] M. Serbyn, Z. Papić, and D. A. Abanin, *Phys. Rev. B* **90**, 174302 (2014).
- [28] D. A. Huse, R. Nandkishore, V. Oganesyan, A. Pal, and S. L. Sondhi, *Phys. Rev. B* **88**, 014206 (2013).
- [29] A. Chandran, V. Khemani, C. R. Laumann, and S. L. Sondhi, *Phys. Rev. B* **89**, 144201 (2014).
- [30] D. Pekker, G. Refael, E. Altman, E. Demler, and V. Oganesyan, *Phys. Rev. X* **4**, 011052 (2014).
- [31] J. A. Kjäll, J. H. Bardarson, and F. Pollmann, *Phys. Rev. Lett.* **113**, 107204 (2014).
- [32] D. J. Luitz, N. Laflorencie, and F. Alet, *Phys. Rev. B* **91**, 081103 (2015).
- [33] D. J. Luitz, N. Laflorencie, and F. Alet, *Phys. Rev. B* **93**, 060201 (2016).
- [34] V. Khemani, S. P. Lim, D. N. Sheng, and D. A. Huse, *Phys. Rev. X* **7**, 021013 (2017).
- [35] J. T. Chayes, L. Chayes, D. S. Fisher, and T. Spencer, *Phys. Rev. Lett.* **57**, 2999 (1986).
- [36] A. Chandran, C. Laumann, and V. Oganesyan, [arXiv: 1509.04285](https://arxiv.org/abs/1509.04285).
- [37] R. Vosk, D. A. Huse, and E. Altman, *Phys. Rev. X* **5**, 031032 (2015).
- [38] A. C. Potter, R. Vasseur, and S. A. Parameswaran, *Phys. Rev. X* **5**, 031033 (2015).
- [39] L. Zhang, B. Zhao, T. Devakul, and D. A. Huse, *Phys. Rev. B* **93**, 224201 (2016).
- [40] M. Serbyn, Z. Papić, and D. A. Abanin, *Phys. Rev. Lett.* **111**, 127201 (2013).

[41] D. A. Huse, R. Nandkishore, and V. Oganesyan, *Phys. Rev. B* **90**, 174202 (2014).

[42] V. Ros, M. Müller, and A. Scardicchio, *Nucl. Phys. B* **891**, 420 (2015).

[43] J. Z. Imbrie, *Phys. Rev. Lett.* **117**, 027201 (2016).

[44] Unless the level spacing of one cluster exceeds the bandwidth of the other; then $\Lambda_{i'} \geq \delta_{i'} \geq \Lambda_{j'} \geq \delta_{j'}$ and $\Delta E_{i'j'} = \max(\delta_{i'} - \Lambda_{j'}, \delta_{j'})$.

[45] See Supplemental Material <http://link.aps.org/supplemental/10.1103/PhysRevLett.119.110604> for details regarding the matrix element renormalization procedure as well as additional plots of the cluster size scaling, universal entanglement crossover and cluster size histograms.

[46] W. De Roeck and F. Huveneers, *Phys. Rev. B* **95**, 155129 (2017).

[47] M. Serbyn, Z. Papić, and D. A. Abanin, *Phys. Rev. X* **5**, 041047 (2015).

[48] D. S. Fisher, *Phys. Rev. Lett.* **69**, 534 (1992).

[49] D. S. Fisher, *Phys. Rev. B* **50**, 3799 (1994).

[50] R. Vosk and E. Altman, *Phys. Rev. Lett.* **112**, 217204 (2014).

[51] R. Vasseur, A. C. Potter, and S. A. Parameswaran, *Phys. Rev. Lett.* **114**, 217201 (2015).

[52] K. Agarwal, E. Altman, E. Demler, S. Gopalakrishnan, D. A. Huse, and M. Knap, *Ann. Phys. (Berlin)* **529**, 1600326 (2017).

[53] Reexamining [38] showed that those RG rules allowed rare avalanches; this does not quantitatively alter the reported critical properties.

[54] X. Yu, D. J. Luitz, and B. K. Clark, *Phys. Rev. B* **94**, 184202 (2016).

[55] T. Grover, [arXiv:1405.1471](https://arxiv.org/abs/1405.1471).

[56] R. C. Myers and A. Singh, *J. High Energy Phys.* 04 (2012), 122.

[57] H. Liu and M. Mezei, *J. High Energy Phys.* 04 (2013), 162.