

15 **Abstract**

16 Earthworm species with different feeding, burrowing, and/or casting behaviors can
17 lead to distinct microbial communities through complex direct and indirect processes.
18 European earthworm invasion into temperate deciduous forests in North America has
19 been shown to alter microbial biomass in the soil and reduce the fungi-to-bacteria ratios.
20 It is unclear how changes in earthworm species composition due to interspecific
21 competition may alter this dynamics, especially under the ongoing invasion by Asian
22 *Amyntas* and *Metaphire* species. Furthermore, it is also poorly understood how
23 interspecific interactions involving different species may have species-dependent, non-
24 additive effects on different groups of soil microorganisms. By conducting a two-year
25 field experiment in a deciduous forest in the Mid-Atlantic, we examined how the Asian
26 species *Amyntas agrestis* and *Amyntas corticis*, the European species *Lumbricus*
27 *rubellus* and *Octolasion lacteum*, and their interactions affect soil microbial communities,
28 and tested the hypotheses that the Asian species and the interaction between the two
29 European species negatively affect bacterial biomass. We showed that while *A. corticis*
30 generally had no effect on soil microorganisms, *A. agrestis* had a positive effect on
31 microbial biomass, primarily by increasing the biomass of Gram-positive and Gram-
32 negative bacteria. Consistent with our hypothesis, the interaction between *L. rubellus* and
33 *O. lacteum* had a negative effect on bacteria biomass. We concluded that the ongoing
34 invasion of the Asian earthworm *A. agrestis* in forest soils and potential displacement of
35 *L. rubellus* due to interspecific competition may lead to increased bacterial biomass
36 through increasing resource availability and disrupting the interaction between *L.*

37 *rubellus* and *O. lacteum*, potentially causing increased carbon mineralization and
38 reducing soil carbon storage.

39

40 **Keywords:** Earthworm invasion; *Amyntas agrestis*; Microbial community; Phospholipid
41 fatty acid; Neutral lipid fatty acid; Interspecific interaction

1. Introduction

Since European settlement, soils in temperate deciduous forests in North America have gone through major changes as a result of past land use, climate change and species gains and losses (Compton and Boone, 2000; Foster and Aber, 2003; Yesilonis et al., 2016). European earthworm invasion into habitats with low native earthworm abundance or previously devoid of earthworms in this region has resulted in major changes in soil carbon cycles. Through their feeding, burrowing, and casting behaviors, invasive earthworms increase litter decomposition rate, incorporate organic matter into aggregates, change the pool size and distribution of carbon and nitrogen in the soil profile, and affect carbon mineralization rates (Hale et al., 2005a; Eisenhauer et al., 2007; Szlavecz et al., 2011; Fahey et al., 2013; Ma et al., 2013, 2014; Yavitt et al., 2015). The altered forms and distribution of soil carbon and nitrogen can change resource availability to soil microbial communities, which in turn affect carbon transformation and mineralization (Schimel and Weintraub, 2003; Waring et al., 2013; Cotrufo et al., 2015). While the effects of invasive earthworms on soil carbon biogeochemistry in temperate deciduous forests have been the focus of many recent studies (e.g. Crumsey et al., 2015 and the studies cited above), the effects of earthworms on the structure of microbial communities in these forests have received less attention (but see Groffman et al., 2004).

It is generally assumed that the burrowing behaviors of earthworms disrupt fungal hyphae in the soil and therefore would shift soil food webs in temperate forests from a fungi-dominated system to a bacteria-dominated system (Bohlen et al., 2004b). This hypothesis has recently gained support from comparing forest stands invaded by European earthworms with those that are earthworm-free. Field observations in

65 hardwood forests in Northeastern USA showed that earthworms decreased the fungi-to-
66 bacteria ratio in the soil profile primarily by eliminating the organic horizon, where
67 fungal biomass dominates (Dempsey et al., 2011, 2013). Consistent with the original
68 hypothesis, the fungi-to-bacteria ratio also decreased in the surface organic soil, even
69 though the impact was much weaker.

70 However, field observations focusing on total microbial biomass have reported
71 contradictory results. Some deciduous forests in North America showed a net increase of
72 microbial biomass in the overall soil profile (Groffman et al., 2004, 2015), while others
73 showed significant net decrease (Eisenhauer et al., 2007, 2011). All of these studies
74 documented consistent decrease in microbial biomass in the O horizon, but the
75 inconsistency appeared to take place in the surface of the mineral soil, where microbial
76 biomass decreased in some studies (Eisenhauer et al., 2007, 2011) while increased in
77 others (Groffman et al., 2004, 2015). The relatively limited number of studies and the
78 inconclusive results therein highlight our lack of understanding on the factors
79 contributing to the contradictory impacts.

80 One potential explanation is the ecology of earthworm species involved. Earthworms
81 affect soil microorganisms and activities through complex direct and indirect processes,
82 and species with different feeding and/or burrowing behaviors can have potentially
83 distinct effects (McLean et al., 2006). Earthworms change resource availability to soil
84 microorganisms by consuming leaf litter and soil organic matter (SOM), by vertical
85 translocation of C and N in the soil profile, and by promoting aggregate formation
86 through casting (Eisenhauer et al., 2007; Dempsey et al., 2011). Fragmentation and
87 vertical mixing of leaf litter by litter-feeding species into surface soil increase the

88 concentration of readily mineralizable C and tend to increase microbial abundance in the
89 bulk soil. In contrast, soil-feeding species may decrease microbial biomass in the bulk
90 soil as they feed primarily on soil organic matter and reduce resource availability to
91 microbes (McLean et al., 2006). The activity of soil living earthworms can disrupt fungal
92 hyphae (Butenschoen et al., 2007), leading to the reduction of soil fungi. Some
93 earthworm species selectively feed on bacteria (Jayasinghe and Parkinson, 2009; Larsen
94 et al., 2016), reducing bacteria abundance. Newly produced earthworm casts, middens,
95 and burrows are rich in soluble and readily accessible carbon and thus generally
96 considered hotspots of microbial activity (Aira et al., 2009; Stromberger et al., 2012).
97 However, the casting behavior also incorporates organic matter into aggregates, making it
98 inaccessible to microorganisms. Altogether, these different processes highlight that
99 earthworm communities with different species compositions can have contrasting overall
100 effects on soil microbial communities, depending on the ecology of the involved species.

101 In North America, the majority of studies on invasive earthworm effects reported on
102 less than 10 species, all European lumbricids. Invasion of several Asian species,
103 including *Amyntas corticis*, *A. gracilis*, *A. agrestis*, *A. tokioensis*, and *Metaphire*
104 *hilgendorfi* (also known as *A. hilgendorfi*), have received little attention until recent years.
105 The last three species, *A. agrestis*, *A. tokioensis*, and *M. hilgendorfi*, although being
106 present in most East Coast states for at least 80 years (Gates, 1982; Chang et al., 2016c),
107 have recently been reported spreading within those states, as well as into Midwest and
108 West Coast states, including Wisconsin in 2013 (Qiu and Turner, 2016) and Oregon in
109 2016 (Bruce Snyder, personal communication). Unlike most common European
110 earthworms, which have a lifespan of more than a year, the three Asian species have an

annual life cycle. They overwinter exclusively as cocoons, hatch in spring, grow rapidly to 5-15 cm long, reproduce at the end of summer, and die by the end of fall (Callaham et al., 2003; Görres et al., 2014). Their ongoing rapid spreading, which is likely facilitated by nursery plants and commercial mulch (Bellitürk et al., 2015), especially in the form of cocoons, has caused increasing concern. The Asian species can reach extremely high abundance and biomass (Görres et al., 2014), are able to rapidly eliminate the leaf litter layer and change the structure of surface soil (Snyder et al, 2011; Greiner et al., 2012), and have significant impact on soil carbon pools and mineralization (Greiner et al., 2012; Chang et al., 2016a, b) and nutrient concentrations (Qiu et al., 2016). Moreover, they have been shown to negatively affect other organisms, such as millipedes and salamanders (Snyder et al., 2013; Ziemba et al., 2016). The Asian species are also superior competitors for leaf litter against one of the most common European earthworm, *Lumbricus rubellus*, (Zhang et al., 2010; Chang et al., 2016a) and, when present, are usually the dominant earthworm group (Dávalos et al., 2015).

Several recent studies suggested that the common Asian and European invasive earthworms have distinct effects on soil microbial communities (Zhang et al., 2010; Chang et al., 2016a, b). The European species *Lumbricus rubellus* and *Octolasion lacteum* have been shown to have no effect on bacterial biomass, while the Asian species *A. agrestis* and *M. hilgendorfi* reduce the biomass of both Gram-positive and Gram-negative bacteria (Zhang et al., 2010; Chang et al., 2016a). Moreover, the interaction between *L. rubellus* and *O. lacteum* has a positive effect on soil carbon content, and a negative effect on soil respiration, suggesting reduction of carbon mineralization rates (Chang et al., 2016b), while the interaction between *M. hilgendorfi* and *L. rubellus* or *O.*

134 *lacteum* has no effect. All of these contrasting findings between Asian and European
135 earthworms were observed in short-term (<30 days), laboratory mesocosm experiments
136 under well-controlled soil temperature and moisture with nearly no roots and mycorrhizal
137 fungi. It is unclear if any of the changes in microbial communities observed in the
138 laboratory will hold true in the field. Clearly, a sizable gap exists in our understanding
139 between what has been learned from laboratory mesocosm experiments and those from
140 field observational studies.

141 To fill this gap, and to understand species-specific effects and their interactions on
142 soil microbial communities in temperate deciduous forest in Eastern US under the context
143 of European and Asian earthworm invasion, we conducted a two-year, field experiment
144 focusing on two Asian (*A. agrestis* and *A. corticis*) and two European (*L. rubellus* and *O.*
145 *lacteum*) species in a temperate deciduous forest in the Mid-Atlantic region. We
146 hypothesized that (1) earthworms reduce the fungi-to-bacteria ratio in the surface soil, (2)
147 the effects of earthworms on soil microbial community are species-specific and non-
148 additive, (3) the two Asian earthworms, *A. agrestis* and *A. corticis*, have negative effects
149 on soil bacterial biomass, while the two European species, *L. rubellus* and *O. lacteum*,
150 have no effect, and (4) the interaction between *L. rubellus* and *O. lacteum* has a negative
151 effect on soil bacterial biomass.

153 2. Methods

155 2.1. Field site

A two-year field experiment was conducted in a forest stand called Treefall at the Smithsonian Environmental Research Center (SERC), Edgewater, Maryland, USA (38°53'17.0"N, 76°33'14.3"W; www.serc.si.edu). The upland secondary forests at SERC, ranging from 50 to 150 years old (Yesilonis et al., 2016), are dominated by tulip poplar (*Liriodendron tulipifera*), sweet gum (*Liquidambar styraciflua*), red maple (*Acer rubrum*), black cherry (*Prunus serotina*), box elder maple (*Acer negundo*), American beech (*Fagus grandifolia*), oaks (e.g. *Quercus falcata*, *Q. alba*) and hickories (e.g. *Carya tomentosa*, *C. glabra*) (Higman 1968). Soils at our plot locations have been classified as Collington sandy loam (fine-loamy mixed, active, mesic Typic Hapludult) (Szlavecz et al. 2011), with an average pH of 5.1, 32% silt, 20% clay, and 5.6% organic matter content (Yesilonis et al., 2016). The forests have an established community of European earthworms, with *Lumbricus friendi*, *L. rubellus*, *Aporrectodea caliginosa*, *Octolasion lacteum*, *O. cyaneum*, and *Eisenoides lonnbergi* being the most common species, and are currently been invaded by four Asian earthworm species: *Amyntas agrestis*, *Metaphire hilgendorfi*, *A. tokioensis*, and *A. corticis* (Szlavecz and Csuzdi, 2007; Chang et al., 2016a; C.-H. Chang, personal observation).

2.2. Experimental design

Four species of earthworms were selected for the experiment: *Amyntas agrestis*, *A. corticis*, *Lumbricus rubellus*, and *Octolasion lacteum*. *A. agrestis* and *A. corticis* are both originally from Asia and epi-endogeic, but they have contrasting life history: *A. agrestis* is an annual species, reproduces at the end of summer, and dies by the end of fall, while *A. corticis* lives for more than a year (Chang et al. 2016c). The other two species, *L. rubellus*

and *O. lacteum*, are among the most common European earthworms found in temperate deciduous forests in the US. *L. rubellus* is also epi-endogeic, while *O. lacteum* is usually considered polyhumic endogeic. All four species are found at SERC. Individuals of the two *Amyntas* species were collected from forests along the Stony Run Trail, Baltimore, MD; individuals of *L. rubellus* and *O. lacteum* were collected from SERC. All earthworm individuals collected for the experiment were mature (with a clitellum).

Six plots were established in the Treefall forest stand (Fig. 1). Each plot was 3 m x 4 m, at least 1 m away from the nearest trees, and at least 50 m away from the other plots. Each plot was equally divided into a 3 x 4 grid (Fig. 1). Field mesocosms were constructed by burying 23-liter (30 cm in diameter, 35 cm in height) mesh metal wastebaskets 25 cm into the soil at each grid point. Since there was a total of eleven treatments (see below), one randomly chosen grid point in row 4 received no mesocosm. To improve drainage several holes were drilled in the bottom of the basket, which was then covered by a single layer of gravel. The interior of the baskets was then covered by a polyester mesh bag with a 0.8 mm pore size. The pore size allowed fine roots and fungal hyphae to grow into the mesocosms but kept earthworms from going through. Within one plot, the soil retrieved when installing the baskets was divided into the upper 10 cm and the lower 15 cm, which roughly equal the organic and mineral soils, respectively, sieved through a 5 mm sieve, and well mixed. Roots, earthworms, and other soil macrofauna were hand-sorted and removed during the process. Mesocosms were reconstructed by sequentially adding 7.4 kg mineral soil and 5.1 kg organic soils into the basket, and 30 gram of tulip poplar leaf litter on the soil surface.

Earthworms were added into the field mesocosms in June 2012, two weeks after the mesocosms were constructed. A total of 10 treatments and one control were established in each of the six plots: four were single-species treatments with each of the four earthworm species and six were two-species treatments with all combinations of the four species. The control contained no earthworm. The treatments were randomly assigned to the grid points. Eight *A. agrestis*, six *A. corticis*, twelve *L. rubellus*, and twenty-four *O. lacteum* individuals were added into the single-species treatments with the respective species. For the two-species treatments, four *A. agrestis*, three *A. corticis*, six *L. rubellus*, and twelve *O. lacteum* were added into the respective treatments. The numbers of individuals were chosen to take into account both density in the field and individual biomass differences to mimic potential co-occurrence conditions in the field. The mean fresh biomass of individuals (including gut content) was 0.88 g for *A. agrestis*, 0.92 g for *A. corticis*, 0.58 g for *L. rubellus*, and 0.21 g for *O. lacteum*. After earthworm addition the mesh bags were closed, and the plot area was covered by bird nettings (Fig. 1). To supply food for earthworms, additional 60 g Tulip poplar litter was added in two increments, 30 g in October 2012 and 30 g in June 2013. Before the first litter addition, earthworm abundance in the mesocosms was monitored using electroshocking (Szlavec et al., 2013). Before the second litter addition, electroshocking was used to remove all earthworms in each mesocosm, including all individuals without a clitellum, and the abundance of mature earthworms was reestablished to the initial conditions.

The mesocosms were destructively sampled in October 2013. The timing was chosen so that the sampling took place when the soil was moist, and before mature *A. agrestis* reached the end of its life cycle (Görres et al., 2014), and was also based on high

microbial biomass previously reported in temperate deciduous forests in Eastern US (Dempsey et al., 2011, 2013). First, the remaining leaf litter was collected from the soil surface. Then, three soil cores were collected from each mesocosm using a plastic corer with a diameter of 3.5 cm and then divided into 0-2 cm, 2-5 cm, and 5-10 cm increments. Finally, all earthworms from the mesocosms were hand-sorted and preserved in 70% ethanol. The soil samples were sieved through a 2-mm sieve, homogenized, and stored at -20 °C for microbial analysis. Soils from 2-5 cm and 5-10 cm from one of the plot were lost during sample processing and were not analyzed. Soil pH was measured using a 1:2, soil:water slurry in 0.01 M CaCl₂.

2.3. Microbial community analysis

Phospholipid lipid fatty acid (PLFA) analysis was used to characterize soil microbial communities, including Gram-positive bacteria, Gram-negative bacteria, actinomycetes, fungi, and protozoa. The PLFA commonly used as a biomarker for arbuscular mycorrhizal (AM) fungi, 16:1 ω 5, is also known to be found in Gram-negative bacteria, and is a poor indicator of AM fungi in soils with high bacterial biomass (Frostegård et al., 2011). Therefore, to better estimate changes in AM fungi, instead of using PLFA, we used the neutral lipid fatty acid (NLFA) 16:1 ω 5cis as the biomarker for AM fungi (Olsson, 1999), as suggested by Frostegård et al. (2011) and Sharma and Buyer (2015).

Samples were prepared and analyzed as described (Buyer and Sasser, 2012; Sharma and Buyer, 2015) using 19:0 phosphatidylcholine (Avanti Polar Lipids, Alabaster, Alabama, USA) and trionadecanoin glyceride (catalog # T-165, Nu-Chek Prep, Elysian,

MN, USA) as internal standards for quantitative analysis. Gas chromatography was conducted on an Agilent 6890 gas chromatograph (Agilent Technologies, Wilmington, Delaware, USA) coupled with autosampler, split-splitless injector, and flame ionization detector. The system was controlled with MIS Sherlock (Microbial ID, Inc., Newark, DE, USA) and Agilent ChemStation software. Fatty acids were identified using the PLFAD1 calibration mix and PLFAD1 peak library (Microbial ID). Random samples were run on a Clarus 500 GC-MS (Perkin-Elmer, Waltham, MA, USA) to confirm fatty acid identifications.

2.4. Statistical analysis

Fatty acids were combined into groups of biomarkers for Gram-negative bacteria (monounsaturated fatty acids and cyclopropyl 17:0 and 19:0), Gram-positive bacteria (iso and anteiso saturated branched fatty acids), actinomycetes (10-methyl fatty acids), fungi (18:2 ω 6 cis), AM fungi (NLFA 16:1 ω 5cis), and protozoa (20:3 and 20:4 fatty acids) following Buyer and Sasser (2012) and Sharma and Buyer (2015). These biomarkers are not completely universal for all members in each organism group, and are not entirely specific to each group, either. Therefore, they need to be interpreted with caution (Frostegård et al., 2011).

All statistical tests were conducted in R v3.1.2 (R Core Team 2014). PLFA and NLFA were combined into biomarker groups (Buyer and Sasser, 2012) and analyzed using two approaches. First, the effects of earthworm species and species interactions on total PLFA, bacteria PLFA, AM fungi NLFA, fungi-to-bacteria ratios, and PLFA

biomarkers for Gram-negative bacteria, Gram-positive bacteria, actinomycetes, fungi, and protozoa were analyzed using mixed effect models as implemented in the package lme4 (Bates et al., 2015). In the mixed effect models, ‘plot’ was the random effects and the biomass of each earthworm species was considered as fixed effects. Second, the data were Hellinger-transformed (square root of proportion), and microbial community compositions were analyzed using redundancy analyses as implemented in the package vegan (Oksanen et al., 2016) with soil moisture, soil pH, and the biomass of individual earthworm species as constrained variables. Permutation tests with 999 permutations were used to test the significances of the overall models, the constrained axes, soil pH, and the biomass of earthworm species. Linear regressions were used to investigate the correlations between the PLFA 16:1 ω 5cis (AM fungi biomarkers) and the NLFA biomarkers for AM fungi or the PLFA biomarkers for Gram-negative bacteria. The purpose of this regression analysis was to evaluate whether the PLFA 16:1 ω 5cis is an acceptable biomarker for AM fungi in our system, or if it has more contribution from Gram-negative bacteria.

3. Results

The overall survival rates were 12% for *A. agrestis*, 14% for *A. corticis*, 19% for *L. rubellus*, and 32% for *O. lacteum* (Table A1, Appendix). The mortality was most likely caused by extreme heat and drought during summer and was expected for an experiment conducted in field conditions without any control for temperature and soil moisture. Leaf litter remaining on the soil surface was highest in the control treatment (17.9 \pm 5.6 g;

mean \pm SE), and lowest in the *A. corticis* + *O. lacteum* treatment (7.2 \pm 2.1 g) (Table A2, Appendix).

Amyntas agrestis significantly increased total PLFA ($P = 0.003$) and Gram-negative ($P = 0.010$), Gram-positive ($P = 0.004$), and total bacteria ($P = 0.004$) PLFA biomarkers in the 0-2 cm soil layer (Fig. 2; Table 1), but not in the 2-5 cm and 5-10 cm soils. In contrast, *L. rubellus* x *O. lacteum* interaction had significant negative effects on total PLFA ($P = 0.012$) and Gram-negative ($P = 0.016$), Gram-positive ($P = 0.025$), and total bacteria ($P = 0.012$) PLFA biomarkers in the 0-2 cm soil layer (Table 1). In the 2-5 cm layer, *A. agrestis* x *O. lacteum* interaction had significant positive effects on total PLFA ($P = 0.047$) and Gram-negative ($P = 0.032$), Gram-positive ($P = 0.019$), and total bacteria ($P = 0.018$) PLFA biomarkers (Tables A3-A5, Appendix).

In general, earthworms had non-significant effects on fungi PLFA and AM fungi NLFA (Fig. 3; Table 1; Tables A3-A5, Appendix). The only significant effect was found in *A. corticis* x *O. lacteum* interaction, which had a positive effect on fungi PLFA ($P = 0.015$). Similarly, the fungi-to-bacteria ratios were not significantly affected either by earthworm species or their interactions (Fig. 3; Table 1) with only two exceptions. *A. agrestis* x *A. corticis* interaction and *A. corticis* x *O. lacteum* interaction had significant positive effects on the fungi-to-bacteria ratios in the 2-5 cm and 5-10 cm soils, respectively (Table 1). The PLFA 16:1 ω 5cis, frequently attributed to AM fungi, correlated relatively poorly to the NLFA biomarkers for AM fungi ($R^2 = 0.20$, 0.15 and 0.08 for 0-2 cm, 2-5 cm and 5-10 cm soils, respectively); in contrast, the PLFA 16:1 ω 5cis were highly correlated to the PLFA biomarkers for Gram-negative bacteria ($R^2 = 0.74$, 0.82 and 0.73 for 0-2 cm, 2-5 cm and 5-10 cm soils, respectively). These correlations also

had lower AIC (Akaike information criterion) values when compared to the respective linear regressions with the NLFA biomarkers for AM fungi ($\Delta AIC = 68, 84$ and 66 for 0-2 cm, 2-5 cm and 5-10 cm soils, respectively) (Fig. 4), confirming our choice of including NLFA for AM fungi analysis.

In addition to the biomarkers for bacteria, fungi, and protozoa, our samples also contained compounds that matched entries in the PLFAD1 library as dimethyl acetals, which are biomarkers for anaerobes. These compounds are produced from plasmalogens, the class of phospholipids found in anaerobes, by acid-catalyzed transesterification. However, we used base-catalyzed transesterification in the PLFA procedure, and dimethyl acetals would not be produced under basic conditions (Christie 2003). These identifications are therefore incorrect. The concentrations of these unknown compounds were positively affected in the 0-2 cm soils by *A. agrestis* x *A. corticis* interaction ($P = 0.003$) and in the 5-10 cm soils by *L. rubellus* x *O. lacteum* interaction ($P < 0.001$) (Fig. 5; Table 1).

The redundancy analysis (RDA) of microbial community composition showed that in 0-2 cm soil the constrained axes (soil pH and the biomass of individual earthworm species) explained 20.2% of the variance. The overall model was significant ($P = 0.016$), and so was axis 1 ($P = 0.001$). Among the constrained variables, soil pH ($P = 0.004$) and *A. agrestis* had significant effects ($P = 0.034$), while the effects of the other three earthworm species were not significant (Fig. 6). Soil pH was positively associated with the proportion of AM fungi, and negatively associated with the proportion of Gram-positive bacteria. The biomass of *A. agrestis* was positively associated with AM fungi, and negatively associated with the proportion of Gram-negative bacteria. RDA of

microbial community composition in 2-5 and 5-10 cm soils showed that the constrained axes explained 10.0% and 12.0% of the variance, respectively, but the overall models were not significant ($P = 0.366$ for 2-5 cm; $P = 0.247$ for 5-10 cm).

4. Discussion

By conducting a two-year field manipulation, we showed that the Asian species *Amyntas agrestis* had a consistent positive effect on total microbial and bacterial biomass in the surface soil, including the biomass of Gram-positive bacteria, Gram-negative bacteria, and Actinomycetes, as indicated by PLFA biomarkers, while the other Asian species and the two European species had no significant effect. In contrast, the interspecific interaction between *Lumbricus rubellus* and *Octolasion lacteum* had negative effects on total microbial biomass and the biomass of both Gram-positive and Gram-negative bacteria. These results partially support our second hypothesis that the effects of earthworms are species-specific and non-additive, but contradict our third hypothesis regarding the negative relationship between microbial biomass and the presence of *A. agrestis*.

4.1 The impact of *Amyntas agrestis* invasion

Our finding that *A. agrestis* increased microbial and bacterial biomass in the surface soil is consistent with that reported on non-native European earthworms in northern hardwood forests in Northeastern US (Dempsey et al., 2011, 2013; Groffman et al., 2004, 2015), but is contradictory to a previous laboratory mesocosm study documenting that *A. agrestis* reduces total microbial biomass and the biomass of Gram-negative and Gram-

positive bacteria (Zhang et al., 2010). These contradictory results highlighted that data from short-term, laboratory mesocosm experiments need to be interpreted with caution before they are confirmed in the field. *A. agrestis* is an epi-endogeic species that is primarily active and casting in the surface soil. Its activity has been known to increase extracellular enzyme activity (Bellitürk et al., 2015) and the concentrations of nitrate, inorganic nitrogen, and dissolved organic carbon in the soil (Qiu and Turner, 2016). These observations suggest that the activity of *A. agrestis* increases carbon and nitrogen readily available to microorganisms, leading to our observed increase in bacterial biomass. Altogether, the increase in microbial biomass and resource availability to microorganisms can potentially increase carbon mineralization and lead to higher carbon losses (Schimel and Weintraub, 2003). Our result further suggests that previous field observations showing no apparent correlation between microbial biomass and *A. agrestis* presence (Snyder et al., 2011) in the Great Smoky Mountains might be caused by confounding factors that are common in heterogeneous field environments, such as soil texture, soil moisture, and topography.

The fact that *A. agrestis* increases, as opposed to decreases, bacterial biomass in the field lead us to reevaluate how it may interact with another epi-endogeic species, the European *Lumbricus rubellus*, which is common in temperate deciduous forests in North America. Zhang et al. (2010) suggested that *A. agrestis* competes with *L. rubellus* by negatively affecting soil bacteria, which were thought to be important for *L. rubellus* nutrition or for conditioning leaf litter (Zhang et al., 2010). The fact that both Gram-negative and Gram-positive bacterial biomass increased in the field does not support that hypothesis. An alternative hypothesis regarding interspecific competition in earthworms

has been proposed for another Asian invasive species, *Metaphire hilgendorfi*, often referred to as *Amyntas hilgendorfi* in North American literature (Chang et al., 2016c). *Metaphire hilgendorfi* is a species closely related and ecologically similar to *A. agrestis* (Chang et al., 2016c). It has been shown to directly compete with *L. rubellus* for leaf litter, and as the superior competitor, it has the potential to outcompete the latter (Chang et al., 2016a). Considering the similarities between *A. agrestis* and *M. hilgendorfi*, including their body size, annual life cycle, reproduction, and their moving, feeding, burrowing and casting behaviors (Callaham et al., 2003; Richardson et al., 2009; Snyder et al., 2011, 2013; Greiner et al., 2012; Görres et al., 2012, 2014; Chang et al., 2016a, b, c; Qiu and Turner, 2016), we believe that the mechanisms of competition between *A. agrestis* and *L. rubellus* are similar to those reported for *M. hilgendorfi* (Chang et al., 2016a), with *A. agrestis* being the superior competitor.

In a short-term, laboratory mesocosm experiment, Chang et al. (2016b) reported that the interaction between *L. rubellus* and *O. lacteum* has a positive effect on soil carbon content and a negative effect on soil respiration. Consistent with these reports, our data from the field experiment supported our last hypothesis that the interaction between the two European species has a negative effect on the biomass of both Gram-negative and Gram-positive bacteria, and further suggested that decreases in microbial biomass may be responsible for the previously observed changes in soil respiration and soil carbon content. The fact that the same type of interaction did not happen between *O. lacteum* and the other two epi-endogeic species in our experiment, *A. agrestis* and *A. corticis*, further strengthens the idea that when soil carbon biogeochemistry is concerned, earthworm

species composition, and more importantly, species identities, need to be taken into account (Neilson et al., 2000; Zicsi et al., 2011; Chang et al., 2016a).

When taking into account both the effects of *A. agrestis* and the interspecific interaction between epi-endogeic species and *O. lacteum* (endogeic), the ongoing invasion of *A. agrestis* has an unexpected consequence. Most of the forests at SERC, as well as many temperate deciduous forests in Eastern and Northeastern US, are currently dominated by the epi-endogeic species *L. rubellus*, the anecic species *Lumbricus friendi* or *L. terrestris*, and one or two of the common endogeic species, including *Eisenoides lonnbergi*, *Aporrectodea caliginosa*, *Octolasion cyaneum*, and *O. lacteum* (Bohlen et al., 2004a; Hale et al., 2005b; Szlavecz and Csuzdi, 2007; Filley et al., 2008; Nuzzo et al., 2009). Recent studies have collectively indicated displacement of *L. rubellus* by *A. agrestis* as a result of interspecific competition (Zhang et al., 2010; Greiner et al., 2012; Dávalos et al., 2015). Adding *A. agrestis* into the system may result in replacing *L. rubellus* by *A. agrestis* as a result of competition, and thus disrupting the existing interaction between *L. rubellus* and *O. lacteum*. This shift in community composition may then increase carbon mineralization rates and lead to future carbon loss through increasing soil microbial biomass. This conclusion, albeit unexpected, would not have been possible without knowing the interaction between *L. rubellus* and *O. lacteum*, or conducting field experiments.

4.2 The effect of earthworms on fungi-to-bacteria ratio

In general, our data showed little effect of earthworms on the fungi-to-bacteria ratio in the soil, and did not support our first hypothesis. These results are inconsistent with

field observations reported by Dempsey et al. (2011, 2013), which is most likely due to the different initial conditions at the onset of the experiment. Dempsey et al (2011, 2013) documented how earthworms changed soil microbial communities in a previously earthworm-free ecosystem with a thick O horizon rich in fungal biomass. Our experiment captured the dynamics of microorganisms under the presence or absence of different earthworm species in forest soils with a long history of land use and earthworm legacy (Szlavec and Csuzdi, 2007; Szlavec et al., 2011). The soil surface in our forest has not redeveloped a thick O horizon even after 150 years of agricultural abandonment (Yesilonis et al., 2016). As soil sieving at the beginning of our experiment destroyed fungal hyphae and bacteria were probably feeding on the dead fungal biomass, the system was in a state of recovery for the duration of the experiment. The lack of the O horizon may have limited fungi growth into the mesocosms, including the control.

Positive interspecific interactions between *A. agrestis* and *A. corticis* (in 2-5 cm soil) and between *A. corticis* and *O. lacteum* (in 5-10 cm soil) on fungi-to-bacteria ratios are intriguing, and it appears that the effects under these species interactions were mainly caused by positive effects on fungal biomass. The treatment with both *A. corticis* and *O. lacteum* had the highest fungal biomass in the 5-10 cm soil among all treatments, but its AM fungal biomass is the lowest, suggesting that the effects we observed were primarily contributed by increasing biomass of saprotrophic and/or ectomycorrhizal fungi, as opposed to AM fungi. This result is consistent with our observations that fine roots from the surrounding trees and saplings, primarily *Fagus grandifolia* (American beech, associated with ectomycorrhizal fungi), were able to successfully grow back into most of

our mesocosms after the initial disruption of soil structure, presumably providing the source of ectomycorrhizal fungi.

4.3 Technical consideration for PLFA

The present study also documented the importance of choosing the appropriate biomarkers for AM fungi. The PLFA 16:1 ω 5cis, commonly used as a biomarker for AM fungi, has also been found repeatedly in Gram-negative bacteria (Frostegård et al., 2011), and may be appropriate as a biomarker in some systems (e.g. Dempsey et al., 2013) while not in others (Frostegård et al., 2011; Sharma and Buyer, 2015). Our data clearly showed that the assumed AM fungi biomarker PLFA 16:1 ω 5cis had a stronger correlation with Gram-negative bacteria PLFA biomarkers than with the reliable AM fungi biomarker NLFA 16:1 ω 5cis (Olsson, 1999; Frostegård et al., 2011; Sharma and Buyer, 2015), suggesting that the assumed biomarker reflects more of the biomass of Gram-negative bacteria rather than that of AM fungi. Accordingly, in our forest plots at SERC, where the dominant tree species are *Liriodendron tulipifera* (associated with AM fungi) and *Fagus grandifolia* (associated with ectomycorrhizal fungi), choosing a reliable AM fungi fatty acid biomarker is crucial in estimating the relative biomass change among different treatments, and the PLFA 16:1 ω 5cis should never be used.

Chang et al. (2016b) claimed that the interaction between *L. rubellus* and *O. lacteum* has a positive effect on the PLFA biomarkers of anaerobes. However, we recently discovered that the assertion is unfounded. As noted earlier, the biomarkers for anaerobes, dimethyl acetals, are not produced under the base-catalyzed transesterification procedure we adopted (Christie 2003). Therefore, the compounds in our samples are not dimethyl

acetals. We have not been able to identify the compounds due to their low concentrations in the soil. However, the concentration of these unknown compounds is consistently increased in both laboratory (Chang et al. 2016b) and field (this study) conditions when both *L. rubellus* and *O. lacteum* are present. Further study is needed to understand the ecological implication of this increase.

5. Conclusions

Our study documented microbial communities in a field experiment with different earthworm species and species combinations in a temperate deciduous forest in Eastern US. Our results highlighted that three different epi-endogeic species and their interactions with the endogeic earthworm *O. lacteum* have distinct effects on bacterial biomass. The ongoing invasion of the Asian earthworm *A. agrestis* in forest soils will lead to increased microbial and bacterial biomass. Moreover, through outcompeting the common European species *L. rubellus* and changing interspecific interactions with *O. lacteum*, *A. agrestis* can further exacerbate the condition, potentially causing increased carbon mineralization and reducing soil carbon storage.

Acknowledgements

We are grateful to those who helped in the field, especially Jason Hu, Samuel Keamy-Minor, Beverly Lasan, and Michael Bernard. We also thank Stanley Tesch for helping with PLFA and NLFA. Funding for this study was partially provided by the EPS Field

493 Funds and the National Science Foundation (EEC-0540832, ACI-1244820, and EF-
494 1550795).

495

496 **References**

497 Aira, M., McNamara, N.P., Pearce, T.G., Dominguez, J., 2009. Microbial communities
498 of *Lumbricus terrestris* L. middens: structure, activity, and changes through time in
499 relation to earthworm presence. *Journal of Soils and Sediments* 9, 54–61.

500 Bates, D., Maechler, M., Bolker, B.M., Walker, S.C., 2015. Fitting Linear Mixed-Effects
501 Models Using lme4. *Journal of Statistical Software* 67, 1–48.

502 Bellitürk, K., Goerres, J.H., Kunkle, J., Melnichuk, R.D.S., 2015. Can commercial
503 mulches be reservoirs of invasive earthworms? Promotion of ligninolytic enzyme
504 activity and survival of *Amyntas agrestis* (Goto and Hatai, 1899). *Applied Soil*
505 *Ecology* 87, 27–31.

506 Bohlen, P.J., Groffman, P.M., Fahey, T.J., Fisk, M.C., Suarez, E., Pelletier, D.M., Fahey,
507 R.T., 2004a. Ecosystem consequences of exotic earthworm invasion of north
508 temperate forests. *Ecosystems* 7, 1–12.

509 Bohlen, P.J., Scheu, S., Hale, C.M., McLean, M.A., Migge, S., Groffman, P.M.,
510 Parkinson, D., 2004b. Non-native invasive earthworms as agents of change in
511 northern temperate forests. *Frontiers in Ecology and the Environment* 2, 427–435.

512 Butenschoen, O., Poll, C., Langel, R., Kandeler, E., Marhan, S., Scheu, S., 2007.
513 Endogeic earthworms alter carbon translocation by fungi at the soil-litter interface.
514 *Soil Biology & Biochemistry* 39, 2854–2864.

515 Buyer, J.S., Sasser, M., 2012. High throughput phospholipid fatty acid analysis of soils.

516 Applied Soil Ecology 61, 127–130.

517 Callaham, M.A., Hendrix, P.F., Phillips, R.J., 2003. Occurrence of an exotic earthworm

518 (*Amyntas agrestis*) in undisturbed soils of the southern Appalachian Mountains,

519 USA. *Pedobiologia* 47, 466–470.

520 Chang, C.-H., Szlavecz, K., Filley, T., Buyer, J.S., Bernard, M.J., Pitz, S.L., 2016a.

521 Belowground competition among invading detritivores. *Ecology* 97, 160–170.

522 Chang, C.-H., Szlavecz, K., Buyer, J.S., 2016b. Species-specific effects of earthworms on

523 microbial communities and the fate of litter-derived carbon. *Soil Biology &*

524 *Biochemistry* 100, 129–139.

525 Chang, C.-H., Snyder, B.A., Szlavecz, K., 2016c. Asian pheretimoid earthworms in

526 North America north of Mexico: An illustrated key to the genera *Amyntas*,

527 *Metaphire*, *Pithemera*, and *Polypheretima* (Clitellata: Megascolecidae). *Zootaxa*

528 4179 (3), 495–529.

529 Christie, W.W., 2003. *Lipid Analysis*, Third ed. PJ Barnes & Associates, Bridgwater,

530 England, 416 pp.

531 Compton, J.E., Boone, R.D., 2000. Long-term impacts of agriculture on soil carbon and

532 nitrogen in New England forests. *Ecology* 81, 2314–2330.

533 Cotrufo, M.F., Soong, J.L., Horton, A.J., Campbell, E.E., Haddix, M.L., Wall, D.H.,

534 Parton, A.J., 2015. Formation of soil organic matter via biochemical and physical

535 pathways of litter mass loss. *Nature Geoscience* 8, 776–779.

536 Crumsey, J.M., Capowiez, Y., Goodsitt, M.M., Larson, S., Le Moine, J.M., Bird, J.A.,

537 Kling, G.W., Nadelhoffer, K.J., 2015. Exotic earthworm community composition

538 interacts with soil texture to affect redistribution and retention of litter-derived C

539 and N in northern temperate forest soils. *Biogeochemistry* 126, 379–395.

540 Davalos, A., Simpson, E., Nuzzo, V., Blossey, B., 2015. Non-consumptive Effects of
 541 Native Deer on Introduced Earthworm Abundance. *Ecosystems* 18, 1029–1042.

542 Dempsey, M.A., Fisk, M.C., Fahey, T.J., 2011. Earthworms increase the ratio of bacteria
 543 to fungi in northern hardwood forest soils, primarily by eliminating the organic
 544 horizon. *Soil Biology & Biochemistry* 43, 2135–2141.

545 Dempsey, M.A., Fisk, M.C., Yavitt, J.B., Fahey, T.J., Balser, T.C., 2013. Exotic
 546 earthworms alter soil microbial community composition and function. *Soil Biology
 547 & Biochemistry* 67, 263–270.

548 Eisenhauer, N., Partsch, S., Parkinson, D., Scheu, S., 2007. Invasion of a deciduous forest
 549 by earthworms: Changes in soil chemistry, microflora, microarthropods and
 550 vegetation. *Soil Biology & Biochemistry* 39, 1099–1110.

551 Eisenhauer, N., Schlaghamersky, J., Reich, P.B., Frelich, L.E., 2011. The wave towards a
 552 new steady state: effects of earthworm invasion on soil microbial functions.
 553 *Biological Invasions* 13, 2191–2196.

554 Fahey, T.J., Yavitt, J.B., Sherman, R.E., Maerz, J.C., Groffman, P.M., Fisk, M.C.,
 555 Bohlen, P.J., 2013. Earthworm effects on the incorporation of litter C and N into
 556 soil organic matter in a sugar maple forest. *Ecological Applications* 23, 1185–1201.

557 Filley, T.R., McCormick, M.K., Crow, S.E., Szlavecz, K., Whigham, D.F., Johnston,
 558 C.T., van den Heuvel, R.N., 2008. Comparison of the chemical alteration trajectory
 559 of *Liriodendron tulipifera* L. leaf litter among forests with different earthworm
 560 abundance. *Journal of Geophysical Research-Biogeosciences* 113.

561 Foster, D., Aber, J., 2003. Forest in Time, the environmental consequences of 1000 years

562 of change in New England. Yale University Press, New Haven.
 563 Frostegard, A., Tunlid, A., Baath, E., 2011. Use and misuse of PLFA measurements in
 564 soils. *Soil Biology & Biochemistry* 43, 1621–1625.
 565 Gates, G.E., 1982. Farewell to North American megadriles. *Megadrilogica* 4, 12–77.
 566 Görres, J.H., Melnichuk, R.D.S., 2012. Asian invasive earthworms of the genus
 567 *Amyntas* Kinberg in Vermont. *Northeastern Naturalist* 19, 313–322.
 568 Görres, J.H., Melnichuk, R.D.S., Bellitürk, K., 2014. Mortality pattern relative to size
 569 variation within *Amyntas agrestis* (Goto & Hatai 1899) (Oligochaeta:
 570 Megascolecidae) populations in the Champlain Valley of Vermont, USA.
 571 *Megadrilogica* 16, 9–14.
 572 Greiner, H.G., Kashian, D.R., Tiegs, S.D., 2012. Impacts of invasive Asian (*Amyntas*
 573 *hilgendorfi*) and European (*Lumbricus rubellus*) earthworms in a North American
 574 temperate deciduous forest. *Biological Invasions* 14, 2017–2027.
 575 Groffman, P.M., Bohlen, P.J., Fisk, M.C., Fahey, T.J., 2004. Exotic earthworm invasion
 576 and microbial biomass in temperate forest soils. *Ecosystems* 7, 45–54.
 577 Groffman, P.M., Fahey, T.J., Fisk, M.C., Yavitt, J.B., Sherman, R.E., Bohlen, P.J.,
 578 Maerz, J.C., 2015. Earthworms increase soil microbial biomass carrying capacity
 579 and nitrogen retention in northern hardwood forests. *Soil Biology & Biochemistry*
 580 87, 51–58.
 581 Hale, C.M., Frelich, L.E., Reich, P.B., 2005a. Exotic European earthworm invasion
 582 dynamics in northern hardwood forests of Minnesota, USA. *Ecological*
 583 *Applications* 15, 848–860.
 584 Hale, C.M., Frelich, L.E., Reich, P.B., Pastor, J., 2005b. Effects of European earthworm

585 invasion on soil characteristics in northern hardwood forests of Minnesota, USA.
 586 Ecosystems 8, 911–927.

587 Higman, D., 1968. An ecologically annotated checklist of the vascular flora at the
 588 Chesapeake Bay Center for field biology, with keys. Smithsonian Institution,
 589 Washington, DC, USA.

590 Jayasinghe, B.A.T.D., Parkinson, D., 2009. Earthworms as the vectors of actinomycetes
 591 antagonistic to litter decomposer fungi. Applied Soil Ecology 43, 1–10.

592 Larsen, T., Pollierer, M.M., Holmstrup, M., D'Annibale, A., Maraldo, K., Andersen, N.,
 593 Eriksen, J., 2016. Substantial nutritional contribution of bacterial amino acids to
 594 earthworms and enchytraeids: A case study from organic grasslands. Soil Biology
 595 & Biochemistry 99, 21–27.

596 Ma, Y., Filley, T.R., Johnston, C.T., Crow, S.E., Szlavecz, K., McCormick, M.K., 2013.
 597 The combined controls of land use legacy and earthworm activity on soil organic
 598 matter chemistry and particle association during afforestation. Organic
 599 Geochemistry 58, 56–68.

600 Ma, Y., Filley, T.R., Szlavecz, K., McCormick, M.K., 2014. Controls on wood and leaf
 601 litter incorporation into soil fractions in forests at different successional stages. Soil
 602 Biology & Biochemistry 69, 212–222.

603 McLean, M.A., Migge-Kleian, S., Parkinson, D., 2006. Earthworm invasions of
 604 ecosystems devoid of earthworms: effects on soil microbes. Biological Invasions 8,
 605 1257–1273.

606 Neilson, R., Boag, B., Smith, M., 2000. Earthworm $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analyses suggest
 607 that putative functional classifications of earthworms are site-specific and may also

608 indicate habitat diversity. *Soil Biology & Biochemistry* 32, 1053–1061.

609 Nuzzo, V.A., Maerz, J.C., Blossey, B., 2009. Earthworm Invasion as the Driving Force
610 Behind Plant Invasion and Community Change in Northeastern North American
611 Forests. *Conservation Biology* 23, 966–974.

612 Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'Hara, R.B.,
613 Simpson, G.L., Solymos, P., Stevens, M.H.H., Wagner, H., 2016. *vegan*:
614 community ecology package. R package version 2.3-5. [http://CRAN.R-](http://CRAN.R-project.org/package=vegan)
615 [project.org/package=vegan](http://CRAN.R-project.org/package=vegan)

616 Olsson, P.A., 1999. Signature fatty acids provide tools for determination of the
617 distribution and interactions of mycorrhizal fungi in soil. *FEMS Microbiology*
618 *Ecology* 29, 303–310.

619 R Core Team, 2014. R: A language and environment for statistical computing. R
620 Foundation for Statistical Computing, Vienna, Austria.

621 Richardson, D.R., Snyder, B.A., Hendrix, P.F., 2009. Soil moisture and temperature:
622 tolerances and optima for a non-native earthworm species, *Amyntas agrestis*
623 (Oligochaeta: Opisthopora: Megascolecidae). *Southeastern Naturalist* 8, 325–334.

624 Schimel, J.P., Weintraub, M.N., 2003. The implications of exoenzyme activity on
625 microbial carbon and nitrogen limitation in soil: a theoretical model. *Soil Biology &*
626 *Biochemistry* 35, 549–563.

627 Sharma, M.P., Buyer, J.S., 2015. Comparison of biochemical and microscopic methods
628 for quantification of arbuscular mycorrhizal fungi in soil and roots. *Applied Soil*
629 *Ecology* 95, 86–89.

630 Snyder, B.A., Callahan, M.A., Jr., Hendrix, P.F., 2011. Spatial variability of an invasive

631 earthworm (*Amyntas agrestis*) population and potential impacts on soil
 632 characteristics and millipedes in the Great Smoky Mountains National Park, USA.
 633 Biological Invasions 13, 349–358.

634 Snyder, B.A., Callahan, M.A., Jr., Lowe, C.N., Hendrix, P.F., 2013. Earthworm invasion
 635 in North America: Food resource competition affects native millipede survival and
 636 invasive earthworm reproduction. Soil Biology & Biochemistry 57, 212–216.

637 Stromberger, M.E., Keith, A.M., Schmidt, O., 2012. Distinct microbial and faunal
 638 communities and translocated carbon in *Lumbricus terrestris* drilospheres. Soil
 639 Biology & Biochemistry 46, 155–162.

640 Szlavecz, K., Csuzdi, C., 2007. Land use change affects earthworm communities in
 641 Eastern Maryland, USA. European Journal of Soil Biology 43, S79–S85.

642 Szlavecz, K., McCormick, M., Xia, L., Saunders, J., Morcol, T., Whigham, D., Filley, T.,
 643 Csuzdi, C., 2011. Ecosystem effects of non-native earthworms in Mid-Atlantic
 644 deciduous forests. Biological Invasions 13, 1165–1182.

645 Waring, B.G., Averill, C., Hawkes, C.V., 2013. Differences in fungal and bacterial
 646 physiology alter soil carbon and nitrogen cycling: insights from meta-analysis and
 647 theoretical models. Ecology Letters 16, 887–894.

648 Yavitt, J.B., Fahey, T.J., Sherman, R.E., Groffman, P.M., 2015. Lumbricid earthworm
 649 effects on incorporation of root and leaf litter into aggregates in a forest soil, New
 650 York State. Biogeochemistry 125, 261–273.

651 Yesilonis, I., Szlavecz, K., Pouyat, R., Whigham, D., Xia, L., 2016. Historical land use
 652 and stand age effects on forest soil properties in the Mid-Atlantic US. Forest
 653 Ecology and Management 370, 83–92.

654 Zhang, W., Hendrix, P.F., Snyder, B.A., Molina, M., Li, J., Rao, X., Siemann, E., Fu, S.,
655 2010. Dietary flexibility aids Asian earthworm invasion in North American forests.
656 Ecology 91, 2070–2079.

657 Zicsi, A., Szlavecz, K., Csuzdi, C., 2011. Leaf litter acceptance and cast deposition by
658 peregrine and endemic European lumbricids (Oligochaeta: Lumbricidae).
659 Pedobiologia 54, S145–S152.

660 Ziemba, J.L., Hickerson, C.-A.M., Anthony, C.D., 2016. Invasive Asian earthworms
661 negatively impact keystone terrestrial salamanders. PLoS ONE 11(5), e0151591.
662





Fig. 1. One of the plots containing 11 mesocosms in the Treefall stand at the Smithsonian Environmental Research Center, Maryland, USA. (A) Photo of Plot 3. (B) Diagram of Plot 3 with dimension information and treatment assignment. Ctr: control; Aa: *A. agrestis*; Ac: *A. corticis*; Lr: *L. rubellus*; Ol: *O. lacteum*.



669

670 **Fig. 2.** Bulk soil (0-2 cm) phospholipid fatty acid (PLFA) biomarker concentrations in
 671 the control (Ctr) and in the single-species treatments containing *A. agrestis* (Aa), *A.*
 672 *corticis* (Ac), *L. rubellus* (Lr), or *O. lacteum* (Ol), showing increased PLFA biomarker
 673 concentration in the *A. agrestis* treatment for (A) total PLFA, (B) total bacteria PLFA,
 674 (C) Gram-positive bacteria, and (D) Gram-negative bacteria. The effects of *A. agrestis*
 675 were significant (positive; $P < 0.05$) in all four groups of PLFA biomarkers, while all of
 676 the other three species were not significant. Error bars show SE.

677




Fig. 3. Bulk soil (0-2 cm) fungi biomarkers and fungi-to-bacteria ratio in the control (Ctr) and in the single-species treatments containing *A. agrestis* (Aa), *A. corticis* (Ac), *L. rubellus* (Lr), or *O. lacteum* (Ol). The effects of all four earthworm species were not significant ($P > 0.05$) for fungi PLFA (A), AM fungi NLFA (B), and fungi-to-bacteria ratio (C). Error bars show SE.




Fig. 4. Correlations between the PLFA 16:1w5cis and the NLFA AM fungi biomarker (A-C) or the PLFA biomarkers for Gram-negative bacteria (D-F) in 0-2 cm (A and D), 2-5 cm (B and E), and 5-10 cm (C and F) soils. All correlations were significant, but the correlations were stronger (higher R^2 values) with the Gram-negative bacteria than with the AM fungi biomarkers.




Fig. 5. Concentrations of unknown compounds matching the entries of dimethyl acetals in the PLFAD1 library in the 5-10 cm soil, showing concentrations in the control (Ctr), in the single-species treatments containing *A. agrestis* (Aa), *A. corticis* (Ac), *L. rubellus* (Lr), or *O. lacteum* (Ol), and in the two-species treatments. The effect of *L. rubellus* x *O. lacteum* interaction was significant (positive; $P < 0.05$), while all of the other factors (species and their interactions) were not significant. Error bars show SE.




Fig. 6. Redundancy analysis of soil microbial community structures in 0-2 cm soil based on phospholipid fatty acid (PLFA) and neutral lipid fatty acid (NLFA) biomarkers. GN: Gram-negative bacteria; GP: Gram-positive bacteria; AT: Actinomycetes; FG: fungi; AM: arbuscular mycorrhizal fungi; PT: protozoa; pH: soil pH; *agrestis*: *Amyntas agrestis*; *corticis*: *A. corticis*; *rubellus*: *Lumbricus rubellus*; *lacteum*: *Octolasion lacteum*. Among the five constrained variables, only soil pH and *A. agrestis* were significant. Vectors for microbes were not shown for clarity.

708 **Table 1**

709 Results of mixed models testing the effects of species and their interactions on PLFA or NLFA biomarkers for bacteria, fungi, and protozoa in
 710 the soil. *P*-values from likelihood ratio tests with a χ^2 distribution and a *df*=1.

Factor*	Gram-negative bacteria			Gram-positive bacteria			Actinomycetes			Total bacteria			Total PLFA		
	0-2 cm	2-5 cm	5-10 cm	0-2 cm	2-5 cm	5-10 cm	0-2 cm	2-5 cm	5-10 cm	0-2 cm	2-5 cm	5-10 cm	0-2 cm	2-5 cm	5-10 cm
Aa	↑0.010	0.349	0.117	↑0.004	0.937	0.137	↑0.005	0.817	0.145	↑0.004	0.633	0.115	↑0.003	0.528	0.166
Ac	0.184	0.952	0.842	0.946	0.398	0.512	0.871	0.427	0.327	0.427	0.647	0.877	0.374	0.860	0.993
Lr	0.445	0.301	0.407	0.216	0.746	0.995	0.397	0.772	0.897	0.303	0.464	0.611	0.320	0.481	0.610
Ol	0.200	0.432	0.699	0.854	0.140	0.892	0.535	0.153	0.906	0.474	0.241	0.770	0.495	0.285	0.808
Aa x Ac	0.302	0.530	0.418	0.684	0.235	0.455	0.765	0.598	0.638	0.410	0.347	0.419	0.521	0.609	0.425
Aa x Lr	0.869	0.272	0.966	0.279	0.840	0.944	0.415	0.979	0.681	0.555	0.617	0.995	0.451	0.775	0.927
Aa x Ol	0.264	↑0.032	0.999	0.582	↑0.019	0.952	0.413	↑0.089	0.984	0.348	↑0.018	0.980	0.380	↑0.047	0.814
Ac x Lr	0.295	0.973	0.229	0.326	0.255	↑0.053	0.188	0.593	↑0.056	0.275	0.583	0.123	0.276	0.613	0.234
Ac x Ol	0.722	0.434	0.550	0.601	0.406	0.962	0.652	0.309	0.375	0.647	0.397	0.700	0.679	0.489	0.462
Lr x Ol	↓0.016	0.323	0.441	↓0.025	↓0.016	0.947	↓0.022	↓0.040	0.540	↓0.012	↓0.082	0.659	↓0.012	0.106	0.684
Factor*	Fungi			AM fungi			Protozoa			Fungi-to-bacteria ratio			Unknown compounds [#]		
	0-2 cm	2-5 cm	5-10 cm	0-2 cm	2-5 cm	5-10 cm	0-2 cm	2-5 cm	5-10 cm	0-2 cm	2-5 cm	5-10 cm	0-2 cm	2-5 cm	5-10 cm
Aa	0.894	0.291	0.764	↑0.059	0.198	0.212	0.189	0.603	0.379	0.125	0.344	0.992	0.213	↑0.097	0.473
Ac	0.191	0.329	0.555	0.319	0.890	0.479	↓0.048	0.813	0.395	0.252	0.249	0.591	0.493	0.613	0.498
Lr	0.700	0.533	0.733	0.280	0.961	0.434	0.902	0.824	0.298	0.996	0.695	0.749	0.661	0.754	0.474
Ol	0.288	0.693	0.662	0.207	0.559	0.228	↓0.014	0.332	0.549	0.443	0.844	0.754	0.669	0.161	0.704
Aa x Ac	0.856	↑0.069	0.752	0.472	0.215	↓0.079	0.404	↑0.091	0.389	0.597	↑0.026	0.945	↑0.003	0.439	0.631
Aa x Lr	↑0.092	0.778	0.820	0.800	0.135	0.226	↑0.035	0.797	0.469	0.311	0.810	0.862	0.381	0.579	0.503
Aa x Ol	0.497	0.577	0.383	0.907	0.405	0.819	0.406	0.637	0.781	0.787	0.259	0.321	0.386	0.128	↓0.066
Ac x Lr	0.914	0.495	0.449	0.442	0.268	0.808	0.578	0.691	0.232	0.541	0.323	0.211	0.613	0.330	0.926
Ac x Ol	0.672	0.473	↑0.015	0.439	0.369	0.770	0.317	0.282	0.808	0.538	0.306	↑0.012	0.152	0.983	0.536
Lr x Ol	0.187	0.871	0.965	0.597	0.363	0.283	0.321	0.411	0.722	0.940	0.575	0.874	0.909	0.850	↑<0.001

711 Notes: ↑, positive effect; ↓, negative effect; significant effects (*P* < 0.05) are given in bold.

712 * Biomass of *Amyntas agrestis* (Aa), *Amyntas corticis* (Ac), *Lumbricus rubellus* (Lr), *Octolasion lacteum* (Ol) and their interactions.

713 # Unknown compounds matching entries for anaerobic biomarkers in the PLFAD1 library.