

DAVIES' METHOD FOR ANOMALOUS DIFFUSIONS

MATHAV MURUGAN AND LAURENT SALOFF-COSTE

(Communicated by Mark M. Meerschaert)

ABSTRACT. Davies' method of perturbed semigroups is a classical technique to obtain off-diagonal upper bounds on the heat kernel. However Davies' method does not apply to anomalous diffusions due to the singularity of energy measures. In this note, we overcome the difficulty by modifying the Davies' perturbation method to obtain sub-Gaussian upper bounds on the heat kernel. Our computations closely follow the seminal work of Carlen, Kusuoka and Stroock (1987). However, a cutoff Sobolev inequality due to Andres and Barlow (2015) is used to bound the energy measure.

1. INTRODUCTION

Davies' method of perturbed semigroups is a well-known method to obtain off-diagonal upper bounds on the heat kernel. It was introduced by E. B. Davies to obtain the explicit constants in the exponential term for Gaussian upper bounds [11] using the logarithmic-Sobolev inequality. Davies' method was extended by Carlen, Kusuoka and Stroock to a non-local setting [9, Section 3] using Nash inequality. Moreover, Davies extended this technique to higher order elliptic operators on $\mathbb{R}^n$ [12, Section 6 and 7]. More recently Barlow, Grigor'yan and Kumagai applied Davies' method as presented in [9] to obtain off-diagonal upper bounds for the heat kernel of heavy tailed jump processes [7, Section 3].

Despite these triumphs, Davies' perturbation method has not yet been made to work in the following contexts:

- (a) Anomalous diffusions (see [4, Section 4.2]).
- (b) Jump processes with jump index greater than or equal to 2 (see [22, Remark 1(d)] and [14, Section 1]).

The goal of this work is to extend Davies' method to anomalous diffusions in order to obtain sub-Gaussian upper bounds. In the anomalous diffusion setting, we use cutoff functions satisfying a cutoff Sobolev inequality to perturb the corresponding heat semigroup. We use a recent work of Andres and Barlow [1] to construct these cutoff functions. We extend the techniques developed here in a sequel to a non-local setting for the jump processes mentioned in (b) above [23]. In [23], we consider the analogue of symmetric stable processes on fractals, while in this work we are motivated by Brownian motion on fractals.

Before we proceed, we briefly outline Davies' method as presented in [9] and point out the main difficulty in extending it to the anomalous diffusion setting.

Received by the editors February 3, 2016 and, in revised form, May 26, 2016 and June 6, 2016.
 2010 *Mathematics Subject Classification.* Primary 60J60, 60J35.
 Both authors were partially supported by NSF grant DMS 1404435.

Consider a metric measure space (M, d, μ) and a Markov semigroup $(P_t)_{t \geq 0}$ symmetric with respect to μ . The most classical case is that of the heat semigroup in $\mathbb{R}^n$ (corresponding to Brownian motion in $\mathbb{R}^n$) associated with the Dirichlet form $\mathcal{E}(f, f) = \int_{\mathbb{R}^n} |\nabla f|^2 d\mu$, where μ is the Lebesgue measure.

Instead of considering the original Markov semigroup $(P_t)_{t \geq 0}$, we consider the perturbed semigroup

$$(1.1) \quad (P_t^\psi f)(x) = e^{\psi(x)} (P_t(e^{-\psi} f))(x)$$

where ψ is a ‘sufficiently nice function’. Given an ultracontractive estimate

$$(1.2) \quad \|P_t\|_{1 \rightarrow \infty} \leq m(t)$$

for the diffusion semigroup, Davies’ method yields an ultracontractive estimate for the perturbed semigroup

$$(1.3) \quad \|P_t^\psi\|_{1 \rightarrow \infty} \leq m_\psi(t).$$

If $p_t(x, y)$ is the kernel of P_t , then the kernel of P_t^ψ is $p_t^\psi(x, y) = e^{-\psi(x)} p_t(x, y) e^{\psi(y)}$. Therefore by (1.3), we obtain the off-diagonal estimate

$$(1.4) \quad p_t(x, y) \leq m_\psi(t) \exp(\psi(y) - \psi(x)).$$

By varying ψ over a class of ‘nice functions’ to minimize the right hand side of (1.4), Davies obtained off-diagonal upper bounds. In Davies’ method as presented in [9, 11], it is crucial that the function ψ satisfies

$$e^{-2\psi} \Gamma(e^\psi, e^\psi) \ll \mu \quad \text{and} \quad e^{2\psi} \Gamma(e^{-\psi}, e^{-\psi}) \ll \mu,$$

where $\Gamma(\cdot, \cdot)$ denotes the corresponding energy measure (cf. Definition 1.1). For the classical example of heat semigroup in $\mathbb{R}^n$ described above, the energy measure $\Gamma(f, g)$ is $\nabla f \cdot \nabla g d\mu$, where μ is the Lebesgue measure.

In fact the expression of m_ψ in (1.3) depends on the uniform bound on the Radon-Nikodym derivatives of the energy measure given by (see [9, Theorem 3.25])

$$\Gamma(\psi) := \left\| \frac{de^{-2\psi} \Gamma(e^\psi, e^\psi)}{d\mu} \right\|_\infty \vee \left\| \frac{de^{2\psi} \Gamma(e^{-\psi}, e^{-\psi})}{d\mu} \right\|_\infty.$$

The main difficulty in extending Davies’ method to anomalous diffusions is that for many ‘typical fractals’ that satisfy a sub-Gaussian estimate, the energy measure $\Gamma(\cdot, \cdot)$ is singular with respect to the underlying symmetric measure μ [8, 16, 20]. This difficulty is well-known to experts (for instance, [5, p. 1507] or [19, p. 86]). In this context, the condition $e^{-2\psi} \Gamma(e^\psi, e^\psi) \ll \mu$ implies that ψ is necessarily a constant, in which case the off-diagonal estimate of (1.4) is not an improvement over the diagonal estimate (1.3).

We briefly recall some fundamental notions regarding Dirichlet form and refer the reader to [13] for details. Let (M, d, μ) be a locally compact metric measure space where μ is a positive Radon measure on M with $\text{supp}(\mu) = M$. We denote by $\langle \cdot, \cdot \rangle$ the inner product on $L^2(M, \mu)$. Let $\mathbf{X} = (\Omega, \mathcal{F}_\infty, \mathcal{F}_t, X_t, \mathbb{P}_x)$ denote the diffusion corresponding to a strongly local regular Dirichlet form. Here Ω denotes the totality of right continuous paths with left-limits from $[0, \infty)$ to M and $\mathbb{P}_x$ denotes the law of the process conditioned to start at $X_0 = x$. The corresponding Markov semigroup $\{P_t : t \geq 0\}$ of $\mathbf{X}$ is defined by $P_t f(x) := \mathbb{E}_x[f(X_t)]$, where $\mathbb{E}_x$ denotes the expectation with respect to the measure $\mathbb{P}_x$. These operators $\{P_t : t \geq 0\}$

form a strongly continuous semigroup of self-adjoint contractions. The Dirichlet form $(\mathcal{E}, \mathcal{D})$ associated with $\mathbf{X}$ is the symmetric, bilinear form

$$\mathcal{E}(u, v) := \lim_{t \downarrow 0} \frac{1}{t} \langle u - P_t u, v \rangle$$

defined on the domain

$$\mathcal{D} := \left\{ u \in L^2(M, \mu) : \sup_{t > 0} \frac{1}{t} \langle u - P_t u, u \rangle < \infty \right\}.$$

Recall that a Dirichlet form $(\mathcal{E}, \mathcal{D})$ on $L^2(M, \mu)$ is said to be *regular* if $C_c(M) \cap \mathcal{D}$ is dense in both $(C_c(M), \|\cdot\|_\infty)$ and the Hilbert space $(\mathcal{D}, \mathcal{E}_1)$. Here $C_c(M)$ is the space of continuous functions with compact support in M and $\mathcal{E}_1(\cdot, \cdot) := \mathcal{E}(\cdot, \cdot) + \langle \cdot, \cdot \rangle$ denotes the inner product on $\mathcal{D}$. For a μ -measurable function u let $\text{Supp}[u]$ denote the support of the measure $u d\mu$. We say that a Dirichlet form $(\mathcal{E}, \mathcal{D})$ on $L^2(M, \mu)$ is *strongly local* if it satisfies the following property: For all functions $u, v \in \mathcal{D}$ such that $\text{Supp}[u], \text{Supp}[v]$ are compact and v is constant on a neighborhood of $\text{Supp}[v]$, we have $\mathcal{E}(u, v) = 0$. For example, the form corresponding to the heat semigroup on $\mathbb{R}^n$ defined by $(f \mapsto \int_{\mathbb{R}^n} |\nabla f|^2 d\mu, W^{1,2}(\mathbb{R}^n))$ is a regular, strongly local Dirichlet form on $L^2(\mathbb{R}^n, \mu)$, where μ is the Lebesgue measure and $W^{1,2}$ denotes the Sobolev space.

We denote by $B(x, r) := \{y \in M : d(x, y) < r\}$ the ball centered at x with radius r and by

$$V(x, r) := \mu(B(x, r))$$

the corresponding volume. We assume that the metric measure space is Ahlfors-regular, meaning that there exist $C_1 > 0$ and $d_f > 0$ such that

$$V(d_f) \quad C_1^{-1} r^{d_f} \leq V(x, r) \leq C_1 r^{d_f}$$

for all $x \in M$ and for all $r \geq 0$. The quantity $d_f > 0$ is called the *volume growth exponent* or *fractal dimension*. Let $p_t(\cdot, \cdot)$ be the (regularized) kernel of P_t with respect to μ [1, eq. (1.10)]. We are interested in obtaining sub-Gaussian upper bounds of the form

$$\text{USG}(d_f, d_w) \quad p_t(x, y) \leq \frac{C_1}{t^{d_f/d_w}} \exp \left(-C_2 \left(\frac{d(x, y)^{d_w}}{t} \right)^{1/(d_w-1)} \right)$$

where $d_w \geq 2$ is the *escape time exponent* or *walk dimension*. It is known that if the heat kernel p_t satisfies $\text{USG}(d_f, d_w)$, then $d_w \geq 2$ (cf. [17, p. 252]). The corresponding diffusion X_t then has a diffusive speed of at least t^{1/d_w} (up to constants). This means that a process starting at x first exits a ball $B(x, r)$ at the time $\tau_{B(x, r)} \gtrsim r^{d_w}$ (cf. [1, Lemma 5.3]). Moreover, if the process satisfies a matching sub-Gaussian lower bound for p_t with different constants, then $\tau_{B(x, r)} \asymp r^{d_w}$. For comparison, recall that the Brownian motion on Euclidean space has a Gaussian heat kernel and satisfies $\tau_{B(x, r)} \asymp r^2$.

Such sub-Gaussian estimates are typical of many fractals (cf. [3, Theorem 8.18]). We assume the on-diagonal bound corresponding to the sub-Gaussian estimate of $\text{USG}(d_f, d_w)$. That is, we assume that there exists $C_1 > 0$ such that

$$(1.5) \quad p_t(x, x) \leq \frac{C_1}{t^{d_f/d_w}}$$

for all $x \in M$ and for all $t > 0$. The on-diagonal estimate of (1.5) is equivalent to the following Nash inequality ([9, Theorem 2.1]): there exists $C_N > 0$ such that

$$N(d_f, d_w) \quad \|f\|_2^{2(1+d_w/d_f)} \leq C_N \mathcal{E}(f, f) \|f\|_1^{2d_w/d_f}$$

for all $f \in \mathcal{D} \cap L^1(M, \mu)$. The Nash inequality $N(d_f, d_w)$ may be replaced by an equivalent Sobolev inequality, a logarithmic Sobolev inequality or a Faber-Krahn inequality (see [2]). However, we will follow the approach of [9] and use the Nash inequality. Such a Nash inequality can be obtained from geometric assumptions like a Poincaré inequality and a volume growth assumption like $V(d_f)$.

Since $\mathcal{E}$ is regular, it follows that $\mathcal{E}(f, g)$ can be written in terms of a signed measure $\Gamma(f, g)$ as

$$\mathcal{E}(f, g) = \int_M \Gamma(f, g),$$

where the energy measure Γ is defined as follows.

Definition 1.1. For any essentially bounded $f \in \mathcal{D}$, $\Gamma(f, f)$ is the unique Borel measure on M (called the energy measure) on M satisfying

$$\int_M g d\Gamma(f, f) = \mathcal{E}(f, fg) - \frac{1}{2} \mathcal{E}(f^2, g)$$

for all essentially bounded $g \in \mathcal{D} \cap C_c(M)$; $\Gamma(f, g)$ is then defined by polarization.

We shall use the following properties of the energy measure.

- (i) *Locality*: For all functions $f, g \in \mathcal{D}$ and all measurable sets $G \subset M$ on which f is constant

$$\mathbf{1}_G d\Gamma(f, g) = 0.$$

- (ii) *Leibniz and chain rules*: For $f, g \in \mathcal{D}$ essentially bounded and $\phi \in C^1(\mathbb{R})$,

$$(1.6) \quad d\Gamma(fg, h) = f d\Gamma(g, h) + g d\Gamma(f, h),$$

$$(1.7) \quad f\Gamma(\phi(f), g) = \phi'(f) d\Gamma(f, g).$$

We wish to obtain an off-diagonal estimate using Davies' perturbation method. The main difference from the previous implementations of the method is that, in addition to an on-diagonal upper bound (or equivalently Nash inequality), we also require a cutoff Sobolev inequality. Spaces satisfying the sub-Gaussian upper bound given in $USG(d_f, d_w)$ necessarily satisfy the cutoff Sobolev annulus inequality $CSA(d_w)$, a condition introduced by Andres and Barlow [1]. The condition CSA simplifies the cutoff Sobolev inequalities CS which were originally introduced by Barlow and Bass [5] for weighted graphs. The significance of the cutoff Sobolev inequalities CS and CSA is that they are stable under bounded perturbations of the Dirichlet form (cf. [1, Corollary 5.2]). Moreover, the condition CS is stable under quasi-isometries (rough isometries) of the underlying space [6, Theorem 2.21(b)]. Therefore cutoff Sobolev inequalities provide a robust method to obtain heat kernel estimates with anomalous time-space scaling. We now define the cutoff Sobolev inequality $CSA(d_w)$.

Definition 1.2. Let $U \subset V$ be open sets in M with $U \subset \bar{U} \subset V$. We say that a continuous function ϕ is a cutoff function for $U \subset V$ if $\phi \equiv 1$ on U and $\phi \equiv 0$ on V^c .

Definition 1.3 ([1, Definition 1.10]). We say $\text{CSA}(d_w)$ holds if there exist $C_1, C_2 > 0$ such that for every $x \in M$, $R > 0$, $r > 0$, there exists a cutoff function ϕ for $B(x, R) \subset B(x, R + r)$ such that if $f \in \mathcal{D}$, then

$$\text{CSA}(d_w) \quad \int_U f^2 d\Gamma(\phi, \phi) \leq C_1 \int_U \phi^2 d\Gamma(f, f) + \frac{C_2}{r^{d_w}} \int_U f^2 d\mu,$$

where $U = B(x, R + r) \setminus \overline{B(x, r)}$.

It is clear that the condition $\text{CSA}(d_w)$ is preserved by bounded perturbations of the Dirichlet form. The above definition is slightly different from the one introduced in [1, Definition 1.10], where the constant C_1 is taken to be $1/8$. However both definitions are equivalent due to a ‘self-improving’ property of $\text{CSA}(d_w)$ [1, Lemma 5.1].

Our main result is that the Nash inequality $N(d_f, d_w)$ and the cutoff Sobolev inequality $\text{CSA}(d_w)$ imply the desired sub-Gaussian estimate $\text{USG}(d_f, d_w)$. By [1, Theorem 1.12], it is known that both $N(d_f, d_w)$ and $\text{CSA}(d_w)$ are also necessary for the sub-Gaussian estimate $\text{USG}(d_f, d_w)$ to hold. More precisely,

Theorem 1.4. *Let (M, d, μ) be a locally compact metric measure space that satisfies $V(d_f)$ with volume growth exponent d_f . Let $(\mathcal{E}, \mathcal{D})$ be a strongly local, regular Dirichlet form whose energy measure Γ satisfies the cutoff Sobolev inequality $\text{CSA}(d_w)$ for some $d_w \geq 2$. Then the Nash inequality $N(d_f, d_w)$ implies the sub-Gaussian upper bound $\text{USG}(d_f, d_w)$.*

Remark 1.5. The above properties given by $V(d_f)$ and $\text{USG}(d_f, d_w)$ are a special case of the more general assumptions of volume doubling and heat kernel upper bounds with a general time-space scaling of [1]. In fact, Theorem 1.4 is subsumed by [1, Theorem 1.12]. A recent work of Lierl provides an alternate proof of the sub-Gaussian estimates in [1] using Moser’s iteration method and extends the results to certain time-dependent, non-symmetric local bilinear forms [21]. Like earlier work by Andres and Barlow and the present work, Lierl’s arguments involve improved control on some cutoff functions.

Our methods give an alternate proof to [1, Theorem 1.12] in a restricted setting. Moreover we show in [23] that this technique can be adapted to the non-local setting to provide new results and resolve the conjecture posed in [22, Remark 1(d)].

2. OFF-DIAGONAL ESTIMATES USING DAVIES’ METHOD

Spaces satisfying $\text{CSA}(d_w)$ have a rich class of cutoff functions with low energy. We start by studying energy estimates of these cutoff functions.

2.1. Self-improving property of CSA. The cutoff Sobolev inequality $\text{CSA}(d_w)$ has a self-improving property which states that the constants C_1, C_2 in $\text{CSA}(d_w)$ are flexible. For example, we can decrease the value of C_1 in $\text{CSA}(d_w)$ by increasing C_2 appropriately. This is quantified in Lemma 2.1. Lemma 2.1 is essentially contained in [1]; we simplify the proof and obtain a slightly stronger result.

Lemma 2.1. *Let (M, d, μ) satisfy $V(d_f)$. Let $(\mathcal{E}, \mathcal{D})$ denote a strongly local, regular Dirichlet form with energy measure Γ that satisfies $\text{CSA}(d_w)$. There exist $C_1, C_2 > 0$ such that for all $\rho \in (0, 1]$, for all $R, r > 0$ and for all $x \in M$, there*

exists a cutoff function $\phi_\rho = \phi_\rho(f)$ for $B(x, R) \subset B(x, R+r)$ that satisfies

$$(2.1) \quad \int_U f^2 d\Gamma(\phi_\rho, \phi_\rho) \leq 4C_1\rho^2 \int_U d\Gamma(f, f) + \frac{C_2\rho^{2-d_w}}{r^{d_w}} \int_U f^2 d\mu$$

for all $f \in \mathcal{D}$, where C_1, C_2 are the constants in $\text{CSA}(d_w)$. Further the cutoff function ϕ_ρ above satisfies

$$(2.2) \quad \left| \phi_\rho(y) - \left(\frac{R+r-d(x, y)}{r} \right) \right| \leq 2\rho$$

for all $y \in B(x, R+r) \setminus B(x, R)$.

Remark 2.2. Lemma 2.1 is essentially contained in work of Andres and Barlow [1, Lemma 5.1]. More recently, following [1, Lemma 5.1], Lierl obtained a cutoff Sobolev inequality [21, Lemma 2.3] that is similar to Lemma 2.1. However, the estimate (2.2) is new and it shows that the cutoff functions converge in L^∞ norm as $\rho \rightarrow 0$ to the ‘linear cutoff function’. The constructions in [1, Lemma 5.1] and [21, Lemma 2.3] converge in L^∞ norm as $\rho \rightarrow 0$ to a somewhat more complicated cutoff function that depends on d_w . The proof below was suggested to us by Martin Barlow.

Proof of Lemma 2.1. Let $x \in M$, $r > 0$, $R > 0$, $\rho > 0$, $f \in \mathcal{D}$. Define $n := \lfloor \rho^{-1} \rfloor \in [\rho^{-1}/2, \rho^{-1}]$. We divide the annulus $U = B(x, R+r) \setminus B(x, R)$ into n -annuli $U_1, U_2, \dots, U_n$ of equal width, where

$$U_i := B(x, R+ir/n) \setminus B(x, R+(i-1)r/n), \quad i = 1, 2, \dots, n.$$

By $\text{CSA}(d_w)$, there exists a cutoff function ϕ_i for $B(x, R+(i-1)r/n) \subset B(x, R+ir/n)$ satisfying

$$(2.3) \quad \int_{U_i} f^2 d\Gamma(\phi_i, \phi_i) \leq C_1 \int_{U_i} d\Gamma(f, f) + \frac{C_2}{(r/n)^{d_w}} \int_{U_i} f^2 d\mu$$

for $i = 1, 2, \dots, n$. We define $\phi = n^{-1} \sum_{i=1}^n \phi_i$. By locality, we have

$$(2.4) \quad d\Gamma(\phi, \phi) = \frac{1}{n^2} \sum_{i=1}^n d\Gamma(\phi_i, \phi_i).$$

Therefore by (2.4), (2.3) and $\rho^{-1}/2 \leq n = \lfloor \rho^{-1} \rfloor \leq \rho^{-1}$, we obtain

$$\begin{aligned} \int_U f^2 d\Gamma(\phi, \phi) &= n^{-2} \sum_{i=1}^n \int_U f^2 d\Gamma(\phi, \phi) \\ &\leq n^{-2} \sum_{i=1}^n \left(C_1 \int_{U_i} d\Gamma(f, f) + \frac{C_2}{(r/n)^{d_w}} \int_{U_i} f^2 d\mu \right) \\ &\leq C_1 n^{-2} \int_U d\Gamma(f, f) + \frac{C_2 n^{d_w-2}}{r^{d_w}} \int_{U_i} f^2 d\mu \\ &\leq 4C_1 \rho^2 \int_U d\Gamma(f, f) + \frac{C_2 \rho^{2-d_w}}{r^{d_w}} \int_{U_i} f^2 d\mu. \end{aligned}$$

This completes the proof of (2.1).

Note that if $y \in U_i$, then $1-i/n \leq \phi(y) \leq 1-(i-1)/n$ and $R+(i-1)r/n \leq d(x, y) < R+ir/n$, for each $1 \leq i \leq n$. This along with $n^{-1} \leq 2\rho$ implies (2.2). $\square$

Observe that by (2.2), the cutoff function ϕ_ρ for $B(x, r) \subset B(x, R + r)$ satisfies

$$\lim_{\rho \downarrow 0} \phi_\rho(y) = 1 \wedge \left(\frac{(R + r - d(x, y))_+}{r} \right).$$

2.2. Estimates on perturbed forms. The key to carrying out Davies' method is the following elementary inequality.

Lemma 2.3. *Let $(\mathcal{E}, \mathcal{D})$ be a strongly local, regular Dirichlet form. Then*

$$(2.5) \quad \mathcal{E}(e^\psi f^{2p-1}, e^{-\psi} f) \geq \frac{1}{p} \mathcal{E}(f^p, f^p) - p \int_M f^{2p} d\Gamma(\psi, \psi)$$

for all $f \in \mathcal{D}$, $\psi \in \mathcal{D}$ and $p \in [1, \infty)$.

Proof. Using Leibniz rule (1.6) and chain rule (1.7), we obtain

$$(2.6) \quad \begin{aligned} & \Gamma(e^\psi f^{2p-1}, e^{-\psi} f) - \frac{1}{p} \Gamma(f^p, f^p) - p f^{2p} \Gamma(\psi, \psi) \\ &= (p-1) \left(f^{2(p-1)} \Gamma(f, f) + f^{2p} \Gamma(\psi, \psi) - 2 f^{2p-1} \Gamma(f, \psi) \right). \end{aligned}$$

By [9, Theorem 3.7] and the Cauchy-Schwarz inequality, we have

$$\int_M f^{2p-1} d\Gamma(f, \psi) \leq \left(\int_M f^{2(p-1)} d\Gamma(f, f) \cdot \int_M f^{2p} d\Gamma(\psi, \psi) \right)^{1/2}.$$

Therefore

$$(2.7) \quad 2 \int_M f^{2p-1} d\Gamma(f, \psi) \leq \int_M f^{2(p-1)} d\Gamma(f, f) + \int_M f^{2p} d\Gamma(\psi, \psi).$$

By integrating (2.6) and using (2.7), we obtain (2.5). $\square$

Davies used the bound

$$\int_M f^{2p} d\Gamma(\psi, \psi) \leq \left\| \frac{d\Gamma(\psi, \psi)}{d\mu} \right\|_\infty \|f\|_{2p}^{2p}$$

to control a term in (2.5). However, for anomalous diffusions, the energy measure is singular to μ . We will instead use $\text{CSA}(d_w)$ to bound $\int_M f^{2p} d\Gamma(\psi, \psi)$ by choosing ψ to be a multiple of the cutoff function satisfying $\text{CSA}(d_w)$. The following estimate is analogous to [9, Theorem 3.9], but unlike in [9], the cutoff functions depend on both p and λ . This raises new difficulties in the implementation of Davies' method.

Proposition 2.4. *Let (M, d, μ) be a metric measure space. Let $(\mathcal{E}, \mathcal{D})$ be a strongly local, regular Dirichlet form on M satisfying $\text{CSA}(d_w)$. There exists $C > 0$ such that, for all $\lambda \geq 1$, for all $r > 0$, for all $x \in M$, and for all $p \in [1, \infty)$, there exists a cutoff function $\phi = \phi_{p,\lambda}$ on $B(x, r) \subset B(x, 2r)$ such that*

$$(2.8) \quad \mathcal{E}(e^{\lambda\phi} f^{2p-1}, e^{-\lambda\phi} f) \geq \frac{1}{2p} \mathcal{E}(f^p, f^p) - C \frac{\lambda^{d_w} p^{d_w-1}}{r^{d_w}} \|f\|_{2p}^{2p},$$

for all $f \in \mathcal{D}$. There exists $C' > 0$ such that the cutoff functions $\phi_{p,\lambda}$ above satisfy

$$(2.9) \quad \|\exp(\lambda(\phi_{p,\lambda} - \phi_{2p,\lambda}))\|_\infty \vee \|\exp(-\lambda(\phi_{p,\lambda} - \phi_{2p,\lambda}))\|_\infty \leq \exp(C'/p)$$

for all $\lambda \geq 1$ and for all $p \geq 1$.

Proof. This theorem follows from Lemma 2.3 and Lemma 2.1. Let $x \in M$ and $r > 0$ be arbitrary. Using (2.5), we obtain

$$(2.10) \quad \mathcal{E}(e^{\lambda\phi} f^{2p-1}, e^{-\lambda\phi} f) \geq \frac{1}{p} \left(\mathcal{E}(f^p, f^p) - (p\lambda)^2 \int_M f^{2p} d\Gamma(\phi, \phi) \right).$$

By Lemma 2.1 and fixing $\rho^2 = (p\lambda)^{-2}/(8C_1)$ in (2.1), we obtain a cutoff function $\phi = \phi_{p,\lambda}$ for $B(x, r) \subset B(x, 2r)$ and $C > 0$ such that

$$(2.11) \quad (p\lambda)^2 \int_M f^{2p} d\Gamma(\phi, \phi) \leq \frac{1}{2} \mathcal{E}(f^p, f^p) + C \frac{(\lambda p)^{d_w}}{r^{d_w}} \int_M f^{2p} d\mu.$$

By (2.10) and (2.11), we obtain (2.8).

By (2.2) and the above calculations, there exists $C' > 0$ such that the cutoff functions $\phi_{p,\lambda} = \phi_{p,\lambda}(f)$ satisfy

$$\|\phi_{p,\lambda} - \phi_{2p,\lambda}\|_\infty \leq \frac{C'}{p\lambda}$$

for all $p \geq 1$, for all $\lambda \geq 1$, for all $f \in \mathcal{D}$, for all $x \in M$ and for all $r > 0$. This immediately implies (2.9). $\square$

Remark 2.5. Estimates similar to (2.8) were introduced by Davies in [12, equation (3)] to obtain off-diagonal estimates for higher order (order greater than 2) elliptic operators. Roughly speaking, the generator $\mathcal{L}$ for anomalous diffusion with walk dimension d_w behaves like an ‘elliptic operator of order d_w ’. However the theory presented in [12] is complete only when the ‘order’ d_w is bigger than the volume growth exponent d_f , i.e. in the strongly recurrent case. This is because the method in [12] relies on a Gagliardo-Nirenberg inequality which is true only in the strongly recurrent setting. We believe that one can adapt the methods of [12] to obtain an easier proof for the strongly recurrent case. However, we will not impose any such restrictions and our proof will closely follow the one in [9].

2.3. Proof of Theorem 1.4. Let $\lambda \geq 1$ and $x \in M$ and $r > 0$. Let $p_k = 2^k$ and let $\psi_k = \lambda\phi_{p_k,\lambda}$, where $\phi_{p_k,\lambda}$ is a cutoff function on $B(x, r) \subset B(x, 2r)$ given by Proposition 2.4. We write

$$(2.12) \quad f_{t,k} := P_t^{\psi_k} f$$

for all $k \in \mathbb{N}$, where $f \in \mathcal{D}$ and $P_t^{\psi_k}$ denotes the perturbed semigroup as in (1.1).

Using (2.8), there exists $C_0 > 0$ such that

$$(2.13) \quad \begin{aligned} \frac{d}{dt} \|f_{t,0}\|_2^2 &= -2\mathcal{E}(e^{\psi_1} f_{t,0}, e^{-\psi_1} f_{t,0}) \\ &\leq 2C_0 \frac{\lambda^{d_w}}{r^{d_w}} \|f_{t,0}\|_2^2 \end{aligned}$$

and

$$(2.14) \quad \begin{aligned} \frac{d}{dt} \|f_{t,k}\|_{2p_k}^{2p_k} &= -2p_k \mathcal{E}(e^{\psi_k} f_{t,k}^{2p_k-1}, e^{-\psi_k} f_{t,k}) \\ &\leq -\mathcal{E}(f_{t,k}^{p_k}, f_{t,k}^{p_k}) + 2C_0 \left(\frac{\lambda p_k}{r} \right)^{d_w} \|f_{t,k}\|_{2p_k}^{2p_k} \end{aligned}$$

for all $k \in \mathbb{N}^*$. By (2.13), we obtain

$$(2.15) \quad \|f_{t,0}\|_{p_1} = \|f_{t,0}\|_2 \leq \exp(C_0 \lambda^{d_w} t / r^{d_w}) \|f\|_2.$$

Using (2.14) and the Nash inequality $N(d_f, d_w)$, we obtain
(2.16)

$$\frac{d}{dt} \|f_{t,k}\|_{2p_k} \leq -\frac{1}{2C_N p_k} \|f_{t,k}\|_{2p_k}^{1+2d_w p_k/d_f} \|f_{t,k}\|_{p_k}^{-2d_w p_k/d_f} + C_0 p_k^{d_w-1} \left(\frac{\lambda}{r}\right)^{d_w} \|f_{t,k}\|_{2p_k}$$

for all $k \in \mathbb{N}^*$. By (2.9) and the fact that P_t is a contraction on L^∞ , we have

$$(2.17) \quad \exp(-2C_1/p_k) f_{t,k+1} \leq f_{t,k} \leq \exp(2C_1/p_k) f_{t,k+1}$$

for all $k \in \mathbb{N}_{\geq 0}$. Combining (2.16) and (2.17), we obtain

$$(2.18) \quad \frac{d}{dt} \|f_{t,k}\|_{2p_k} \leq -\frac{1}{C_A p_k} \|f_{t,k}\|_{2p_k}^{1+2d_w p_k/d_f} \|f_{t,k-1}\|_{p_k}^{-2d_w p_k/d_f} + C_0 p_k^{d_w-1} \left(\frac{\lambda}{r}\right)^{d_w} \|f_{t,k}\|_{2p_k}$$

for all $k \in \mathbb{N}^*$, where $C_A = 2C_N \exp(8d_w C_1/d_f)$.

To obtain off-diagonal estimates using the differential inequalities (2.18) we use the following lemma. The following lemma is analogous to [9, Lemma 3.21], but the statement and its proof are slightly modified to suit our anomalous diffusion context with walk dimension d_w .

Lemma 2.6. *Let $w : [0, \infty) \rightarrow (0, \infty)$ be a non-decreasing function and suppose that $u \in C^1([0, \infty); (0, \infty))$ satisfies*

$$(2.19) \quad u'(t) \leq -\frac{\epsilon}{p} \left(\frac{t^{(p-2)/\theta p}}{w(t)} \right)^{\theta p} u^{1+\theta p}(t) + \delta p^{d_w-1} u(t)$$

for some positive ϵ, θ and δ , $d_w \in [2, \infty)$ and $p = 2^k$ for some $k \in \mathbb{N}^*$. Then u satisfies

$$(2.20) \quad u(t) \leq \left(\frac{2p^{d_w}}{\epsilon \theta} \right)^{1/\theta p} t^{(1-p)/\theta p} w(t) e^{\delta t/p}.$$

Proof. Set $v(t) = e^{-\delta p^{d_w-1} t} u(t)$. By (2.19), we have

$$v'(t) = e^{-\delta p^{d_w-1} t} (u'(t) - \delta p^{d_w-1} u(t)) \leq -\frac{\epsilon t^{p-2}}{p w(t)^{\theta p}} e^{\theta \delta p^{d_w} t} v(t)^{1+\theta p}.$$

Hence

$$\frac{d}{dt} (v(t))^{-\theta p} \geq \epsilon \theta t^{p-2} w(t)^{-\theta p} e^{\theta \delta p^{d_w} t},$$

and so, since w is non-decreasing,

$$(2.21) \quad e^{\delta \theta p^{d_w} t} u(t)^{-\theta p} \geq \epsilon \theta w(t)^{-\theta p} \int_0^t s^{(p-2)} e^{\theta \delta p^{d_w} s} ds.$$

Note that

$$\begin{aligned} \int_0^t s^{(p-2)} e^{\theta \delta p^{d_w} s} ds &\geq (t/\delta \theta p^{d_w})^{p-1} \int_{\delta \theta p^{d_w} (1-1/p^{d_w})}^{\delta \theta p^{d_w}} y^{(p-2)} e^{ty} dy \\ &\geq \frac{t^{p-1}}{p-1} \exp(\delta \theta p^{d_w} t - \delta \theta t) [1 - (1 - p^{-d_w})^{p-1}] \\ (2.22) \quad &\geq \frac{t^{p-1}}{2p^{d_w}} \exp(\delta \theta p^{d_w} t - \delta \theta t). \end{aligned}$$

In the last line above, we used the bound $(1 - p^{-d_w})^{p-1} \geq 1 - p^{-d_w}(p-1)$ for all $p, d_w \geq 2$. Combining (2.21) and (2.22) yields (2.20). $\square$

We now pick $f \in L^2(M, \mu)$ and $f \geq 0$ with $\|f\|_2 = 1$. Let $u_k(t) = \|f_{t,k-1}\|_{p_k}$ and let

$$w_k(t) = \sup\{s^{d_f(p_k-2)/(2d_w p_k)} u_k(s) : s \in (0, t]\}.$$

By (2.15), $w_1(t) \leq \exp(C_0 \lambda^{d_w} t / r^{d_w})$. Further by (2.18), u_{k+1} satisfies (2.19) with $\epsilon = 1/C_A$, $\theta = 2d_w/d_f$, $\delta = C_0(\lambda/r)^{d_w}$, $w = w_k$ and $p = p_k$. Hence by (2.20),

$$u_{k+1}(t) \leq (2^{d_w k+1}/\epsilon\theta)^{1/(\theta p_k)} t^{(1-p_k)/\theta p_k} e^{\delta t/p_k} w_k(t).$$

Therefore

$$w_{k+1}(t)/w_k(t) \leq (2^{d_w k+1}/\epsilon\theta)^{1/(\theta 2^k)} t^{-1/(2^k \theta)} e^{\delta t/2^k}$$

for $k \in \mathbb{N}^*$. Hence, we obtain

$$\lim_{k \rightarrow \infty} w_k(t) \leq C_2 t^{-1/\theta} e^{\delta t} w_1(t) \leq C_2 t^{-1/\theta} \exp(C_0 \lambda^{d_w} t / r^{d_w})$$

where $C_2 = C_2(d_w, \epsilon, \theta)$. Since P_t is a contraction on all $L^p(M, \mu)$ for $1 \leq p \leq \infty$, we obtain

$$\lim_{k \rightarrow \infty} u_k(t) = \|P_t^{\psi_\infty} f\|_\infty \leq \frac{C_2}{t^{d_f/2d_w}} \exp(C_0 \lambda^{d_w} t / r^{d_w})$$

where $\psi_\infty = \lim_{k \rightarrow \infty} \psi_k$. Since the above bound holds for all $f \in L^2(M, \mu)$ with $\|f\|_2 = 1$, we have

$$\|P_t^{\psi_\infty}\|_{2 \rightarrow \infty} \leq \frac{C_2}{t^{d_f/2d_w}} \exp(C_0 \lambda^{d_w} t / r^{d_w}).$$

The estimate is unchanged if we replace ψ_k by $-\psi_k$. Since $P_t^{-\psi}$ is the adjoint of P_t^ψ , by duality we have that

$$\|P_t^{\psi_\infty}\|_{1 \rightarrow 2} \leq \frac{C_2}{t^{d_f/2d_w}} \exp(C_0 \lambda^{d_w} t / r^{d_w}).$$

Combining the above, we have

$$(2.23) \quad \|P_t^{\psi_\infty}\|_{1 \rightarrow \infty} \leq \|P_{t/2}^{\psi_\infty}\|_{1 \rightarrow 2} \|P_{t/2}^{\psi_\infty}\|_{2 \rightarrow \infty} \leq \frac{C_2 2^{d_f/d_w}}{t^{d_f/d_w}} \exp(C_0 \lambda^{d_w} t / r^{d_w}).$$

Therefore

$$p_t(x, y) \leq \frac{C_2 2^{d_f/d_w}}{t^{d_f/d_w}} \exp(C_0 \lambda^{d_w} t / r^{d_w} + \psi_\infty(y) - \psi_\infty(x))$$

for all $x, y \in M$ and for all $r, t > 0$ and $\lambda \geq 1$. If we choose $r = d(x, y)/2$, we have $\psi_\infty(y) - \psi_\infty(x) = -\lambda$. This yields

$$p_t(x, y) \leq \frac{C_3}{t^{d_f/d_w}} \exp(C_4 \lambda^{d_w} t / d(x, y)^{d_w} - \lambda)$$

where $C_3, C_4 > 1$. Assuming $\lambda = C_4^{-1/(d_w-1)} (d(x, y)^{d_w} / t)^{1/(d_w-1)} \geq 1$ in the above equation, we obtain

$$p_t(x, y) \leq \frac{C_3}{t^{d_f/d_w}} \exp\left(-\left(\frac{d(x, y)^{d_w}}{C_5 t}\right)^{1/(d_w-1)}\right)$$

for all $x, y \in M$ and for all $t > 0$ such that $d(x, y)^{d_w} \geq C_4 t$.

If $d(x, y)^{d_w} < C_4 t$, the on-diagonal estimate (1.5) suffices to obtain the desired sub-Gaussian upper bound. $\square$

Remark 2.7. The following generalized capacity estimate is a weaker form of the cutoff Sobolev inequality $\text{CSA}(d_w)$, where the cutoff function ϕ is allowed to depend on the function f . This generalized capacity estimate was introduced by Grigor'yan, Hu and Lau, and they obtain a sub-Gaussian estimate under this weaker assumption [15, Theorem 1.2].

Definition 2.8. We say $\text{Gcap}(d_w)$ holds if there exists $C_1, C_2 > 0$ such that for every $x \in M$, $R > 0$, $r > 0$, $f \in \mathcal{D}$, there exists a cutoff function $\phi = \phi(f)$ for $B(x, R) \subset B(x, R + r)$ such that

$$\text{Gcap}(d_w) \quad \int_U f^2 d\Gamma(\phi, \phi) \leq C_1 \int_U \phi^2 d\Gamma(f, f) + \frac{C_2}{r^{d_w}} \int_U f^2 d\mu,$$

where $U = B(x, R + r) \setminus \overline{B(x, r)}$.

We refer the reader to [10, Definition 1.5] for an analogous generalized capacity estimate in a non-local setting.

It is an interesting open problem to modify the proof of Theorem 1.4 under the above weaker assumption. The main difficulty for carrying out Davies' method under the weaker generalized capacity estimate assumption is that we require the inequalities (2.13) and (2.14) as $t > 0$ varies. This would require the cutoff function ψ_k to depend on $f_{t,k}$ for each t . Therefore the derivatives computed in (2.13) and (2.14) will have additional terms, since ψ_k varies with time t .

We were informed of the reference [18] by the referee during the revision stage. In [18], the techniques developed here and in the companion paper [23] are extended to Dirichlet forms on metric measure spaces (possibly non-local) with jumps satisfying a polynomial type upper bound.

ACKNOWLEDGEMENTS

The authors thank Martin Barlow for his enlightening remarks on the cutoff Sobolev inequalities and for suggesting the proof of Lemma 2.1 which replaced a more complicated proof in an earlier draft. The authors also thank Evan Randles for clarifying various aspects of Davies' work [12] on higher order operators and Tom Hutchcroft for proofreading part of the manuscript. The authors thank the referee for useful remarks and suggestions that led to an improved presentation.

REFERENCES

- [1] Sebastian Andres and Martin T. Barlow, *Energy inequalities for cutoff functions and some applications*, J. Reine Angew. Math. **699** (2015), 183–215, DOI 10.1515/crelle-2013-0009. MR3305925
- [2] D. Bakry, T. Coulhon, M. Ledoux, and L. Saloff-Coste, *Sobolev inequalities in disguise*, Indiana Univ. Math. J. **44** (1995), no. 4, 1033–1074, DOI 10.1512/iumj.1995.44.2019. MR1386760
- [3] Martin T. Barlow, *Diffusions on fractals*, Lectures on probability theory and statistics (Saint-Flour, 1995), Lecture Notes in Math., vol. 1690, Springer, Berlin, 1998, pp. 1–121, DOI 10.1007/BFb0092537. MR1668115
- [4] Martin T. Barlow, *Heat kernels and sets with fractal structure*, Heat kernels and analysis on manifolds, graphs, and metric spaces (Paris, 2002), Contemp. Math., vol. 338, Amer. Math. Soc., Providence, RI, 2003, pp. 11–40, DOI 10.1090/conm/338/06069. MR2039950
- [5] Martin T. Barlow and Richard F. Bass, *Stability of parabolic Harnack inequalities*, Trans. Amer. Math. Soc. **356** (2004), no. 4, 1501–1533 (electronic), DOI 10.1090/S0002-9947-03-03414-7. MR2034316

- [6] Martin T. Barlow, Richard F. Bass, and Takashi Kumagai, *Stability of parabolic Harnack inequalities on metric measure spaces*, J. Math. Soc. Japan **58** (2006), no. 2, 485–519. MR2228569
- [7] Martin T. Barlow, Alexander Grigor'yan, and Takashi Kumagai, *Heat kernel upper bounds for jump processes and the first exit time*, J. Reine Angew. Math. **626** (2009), 135–157, DOI 10.1515/CRELLE.2009.005. MR2492992
- [8] Oren Ben-Bassat, Robert S. Strichartz, and Alexander Teplyaev, *What is not in the domain of the Laplacian on Sierpinski gasket type fractals*, J. Funct. Anal. **166** (1999), no. 2, 197–217, DOI 10.1006/jfan.1999.3431. MR1707752
- [9] E. A. Carlen, S. Kusuoka, and D. W. Stroock, *Upper bounds for symmetric Markov transition functions* (English, with French summary), Ann. Inst. H. Poincaré Probab. Statist. **23** (1987), no. 2, suppl., 245–287. MR898496
- [10] Z.-Q. Chen, T. Kumagai, and J. Wang, *Stability of heat kernel estimates for symmetric jump processes on metric measure spaces*, [arxiv:1604.04035](#)
- [11] E. B. Davies, *Explicit constants for Gaussian upper bounds on heat kernels*, Amer. J. Math. **109** (1987), no. 2, 319–333, DOI 10.2307/2374577. MR882426
- [12] E. B. Davies, *Uniformly elliptic operators with measurable coefficients*, J. Funct. Anal. **132** (1995), no. 1, 141–169, DOI 10.1006/jfan.1995.1103. MR1346221
- [13] Masatoshi Fukushima, Yōichi Ōshima, and Masayoshi Takeda, *Dirichlet forms and symmetric Markov processes*, de Gruyter Studies in Mathematics, vol. 19, Walter de Gruyter & Co., Berlin, 1994. MR1303354
- [14] Alexander Grigor'yan, Jiaxin Hu, and Ka-Sing Lau, *Estimates of heat kernels for non-local regular Dirichlet forms*, Trans. Amer. Math. Soc. **366** (2014), no. 12, 6397–6441, DOI 10.1090/S0002-9947-2014-06034-0. MR3267014
- [15] Alexander Grigor'yan, Jiaxin Hu, and Ka-Sing Lau, *Generalized capacity, Harnack inequality and heat kernels of Dirichlet forms on metric measure spaces*, J. Math. Soc. Japan **67** (2015), no. 4, 1485–1549, DOI 10.2969/jmsj/06741485. MR3417504
- [16] Masanori Hino, *On singularity of energy measures on self-similar sets*, Probab. Theory Related Fields **132** (2005), no. 2, 265–290, DOI 10.1007/s00440-004-0396-1. MR2199293
- [17] Masanori Hino, *On short time asymptotic behavior of some symmetric diffusions on general state spaces*, Potential Anal. **16** (2002), no. 3, 249–264, DOI 10.1023/A:1014033208581. MR1885762
- [18] J. Hu and X. Li, *The Davies method revisited for heat kernel upper bounds of regular Dirichlet forms on metric measure spaces*, [arxiv:1605.05548](#)
- [19] Takashi Kumagai, *Recent developments of analysis on fractals [MR 2105988]*, Selected papers on analysis and related topics, Amer. Math. Soc. Transl. Ser. 2, vol. 223, Amer. Math. Soc., Providence, RI, 2008, pp. 81–95. MR2441420
- [20] Shigeo Kusuoka, *Dirichlet forms on fractals and products of random matrices*, Publ. Res. Inst. Math. Sci. **25** (1989), no. 4, 659–680, DOI 10.2977/prims/1195173187. MR1025071
- [21] J. Lierl, *Parabolic Harnack inequality on fractal-like metric measure Dirichlet spaces*, [arxiv:1509.04804v2](#)
- [22] Mathav Murugan and Laurent Saloff-Coste, *Anomalous threshold behavior of long range random walks*, Electron. J. Probab. **20** (2015), no. 74, 21, DOI 10.1214/EJP.v20-3989. MR3371433
- [23] M. Murugan and L. Saloff-Coste, *Heat kernel estimates for anomalous heavy-tailed random walks*, [arxiv:1512.02361](#)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA AND PACIFIC INSTITUTE
FOR THE MATHEMATICAL SCIENCES, VANCOUVER, BRITISH COLUMBIA V6T 1Z2, CANADA
E-mail address: mathav@math.ubc.ca

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, ITHACA, NEW YORK 14853
E-mail address: lsc@math.cornell.edu