



Search for heavy resonances decaying into WW in the $e\nu\mu\nu$ final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector

The ATLAS Collaboration

A search for neutral heavy resonances is performed in the $WW \rightarrow e\nu\mu\nu$ decay channel using pp collision data corresponding to an integrated luminosity of 36.1 fb^{-1} , collected at a centre-of-mass energy of 13 TeV by the ATLAS detector at the Large Hadron Collider. No evidence of such heavy resonances is found. In the search for production via the quark–antiquark annihilation or gluon–gluon fusion process, upper limits on $\sigma_X \times B(X \rightarrow WW)$ as a function of the resonance mass are obtained in the mass range between 200 GeV and up to 5 TeV for various benchmark models: a Higgs-like scalar in different width scenarios, a two-Higgs-doublet model, a heavy vector triplet model, and a warped extra dimensions model. In the vector-boson fusion process, constraints are also obtained on these resonances, as well as on a Higgs boson in the Georgi–Machacek model and a heavy tensor particle coupling only to gauge bosons.

1 Introduction

The measured properties [1–4] of the Higgs boson discovered in 2012 by the ATLAS [5] and CMS [6] collaborations at the Large Hadron Collider (LHC) are, within experimental uncertainties, consistent with those predicted for the Standard Model (SM) Higgs boson, h . Nevertheless, the SM is thought to be an incomplete theory and many scenarios beyond the SM (BSM) predict an extended Higgs sector [7, 8]. Diboson vector and tensor resonances are also predicted in several other extensions to the SM, such as in composite Higgs models [9, 10] and models with warped extra dimensions [11–14].

This article reports on the results of a search for heavy neutral resonances decaying into two W bosons, which then decay into the $e\nu\mu\nu$ final state, either directly or via leptonic tau decays with additional neutrinos. The analysis is based on the full pp collision dataset collected by the ATLAS detector in 2015 and 2016 at the centre-of-mass energy of $\sqrt{s} = 13$ TeV, corresponding to an integrated luminosity of 36.1 fb^{-1} .

The results are interpreted in terms of different benchmark models. For the case of a scalar resonance produced by gluon–gluon fusion (ggF) or vector-boson fusion (VBF), two scenarios with different intrinsic widths are considered. Constraints on the heavy neutral scalar in two-Higgs-doublet models (2HDM) are also obtained. The neutral member of the fiveplet in the Georgi–Machacek (GM) model [15, 16] also serves as a reference model in the VBF production mode. The parameterisation of heavy vector triplet (HVT) Lagrangians [17, 18] permits the interpretation of searches for spin-1 resonances in a generic way. The bulk Randall–Sundrum (RS) model [11, 19] features a spin-2 Kaluza–Klein (KK) graviton excitation (G_{KK}) decaying into WW , while a tensor resonance signal in the VBF production mode is based on an effective Lagrangian model (ELM) [20].

A previous search for a heavy Higgs boson in the $e\nu\mu\nu$ final state was performed by ATLAS [21] based on a data sample with an integrated luminosity of 20.3 fb^{-1} at $\sqrt{s} = 8$ TeV. The CMS Collaboration also published a search for a high-mass scalar decaying into two W bosons in the fully leptonic final state [22], using datasets at $\sqrt{s} = 7$ and 8 TeV with integrated luminosities of 5.1 fb^{-1} and 19.5 fb^{-1} , respectively. A search for heavy resonances in the RS models in the leptonic decays of the WW channel, using a dataset of 4.7 fb^{-1} at 7 TeV [23], was reported by the ATLAS Collaboration. The ATLAS and CMS collaborations have obtained constraints on the HVT and bulk RS models, based on other decay modes of the VV channels, with V being either a W or a Z boson [24–36]. The search in the $e\nu\mu\nu$ decay mode is complementary to searches performed in other decay modes. In particular, the sensitivity to low mass resonances is higher in the fully leptonic final state than in final states that include jets due to background from jet production.

The article is organised as follows. Section 2 presents the various models used in this analysis. Section 3 describes the ATLAS detector. The data and simulated event samples are discussed in Section 4. The event reconstruction and selection are described in Sections 5 and 6, respectively, followed by the background estimation techniques in Section 7. Systematic uncertainties are discussed in Section 8 and the results are presented in Section 9. Finally, the conclusions are given in Section 10.

2 Theoretical models

The different signal models studied are presented in Table 1. One scenario for the heavy scalar assumes that the scalar has a width much smaller than the detector resolution. This is referred to as the narrow-

Table 1: Summary of the different signal models and resonances considered in the analysis. The resonance spin and production mode are also specified with ggF for gluon–gluon fusion, qqA for quark–antiquark annihilation and VBF for vector-boson fusion.

Model	Resonance spin	Production mode		
		ggF	qqA	VBF
NWA	Spin-0	x		x
2HDM		x		x
LWA		x		x
GM				x
HVT	Spin-1		x	x
Bulk RS	Spin-2	x		
ELM				x

width approximation (NWA). Larger widths (large-width assumption, LWA) of 5%, 10% and 15% of the heavy Higgs boson mass, are also considered. The choice of the width range for the heavy Higgs boson is motivated by the fact that, for several of the most relevant BSM models, widths above 15% are already excluded by indirect limits [37].

The 2HDM comes in different types [38], defined by assumptions about the couplings of each of the Higgs doublets and the discrete symmetries imposed. This analysis considers Type I, where one Higgs doublet couples to vector bosons while the other couples to fermions, and Type II of the minimal supersymmetric (SUSY)-like model in which one Higgs doublet couples to up-type quarks and the other one to down-type quarks and charged leptons. This analysis uses a generic charge-conjugation- and parity-conserving (CP-conserving) 2HDM with a softly broken Z_2 symmetry [38] which has several free parameters: (i) four masses m_h , m_H , m_A and $m_{H^\pm}$ for the two CP-even neutral states, the pseudo-scalar and the charged Higgs boson pair, respectively, (ii) a mixing angle α between the CP-even neutral Higgs fields, and (iii) the ratio of the vacuum expectation values of the two Higgs doublets $\tan \beta = v_2/v_1$. The benchmark is defined by setting $m_h = 125$ GeV and the masses of the supersymmetric particles heavy enough so that Higgs boson decays into SUSY particles are kinematically forbidden. The cross sections and branching fractions are calculated with SusHi and 2HDMC [39, 40].

The GM model extends the Higgs sector with the addition of a real and a complex triplet of $SU(2)_L$ in a way which preserves the SM value of $\rho = M_W^2/(M_Z^2 \cos^2 \theta_W) = 1$ at tree level, with m_W , m_Z and θ_W being the W and Z boson mass and the weak mixing angle, respectively. The physical states include a fermiophobic fiveplet, H_5^0 , $H_5^\pm$, and $H_5^{\pm\pm}$, of custodial $SU(2)$ symmetry which couples preferentially to vector bosons [41]. For that reason, the GM model is less constrained [42], when produced by the VBF process, than other standard benchmark models of a triplet Higgs field, such as the little Higgs model [43] or the left–right symmetric model [44]. The model has many parameters [45, 46], but, if the other new Higgs bosons are heavier than those of the H_5 multiplet, the only production mode is via the VBF process. The cross section and decay width into VV are then proportional to a single parameter, $\sin^2 \theta_H$, which characterises the fraction of the gauge boson masses generated by the triplet Higgs fields.

The HVT Lagrangian [18] parameterises the couplings of the new spin-1 heavy bosons to SM particles in a generic manner and allows their mixing with SM gauge bosons. The s -channel production mechanism of the heavy gauge bosons is primarily via $q\bar{q}$ annihilation (qqA). The HVT bosons couple to the Higgs boson and SM gauge bosons with coupling strength $c_h g_V$ and to the fermions with coupling strength $g^2 c_F/g_V$, where g is the SM $SU(2)_L$ gauge coupling, c_h and c_F are multiplicative factors that modify the

couplings to the Higgs boson and to the fermions, and g_V represents its coupling strength to the W and Z bosons. For the case of vector-boson fusion, it is assumed that there is no coupling to fermions so that non-VBF production processes are suppressed.

The spin-2 G_{KK} is the first Kaluza–Klein excitation of the graviton in the RS model with a warped extra dimension [11, 19], where the SM fields are localised in the bulk [12–14]. This model is characterised by the dimensionless coupling constant $k/\bar{M}_{Pl} \sim \mathcal{O}(1)$ where k determines the curvature of the space, and where $\bar{M}_{Pl} = M_{Pl}/\sqrt{8\pi}$ is the reduced Planck scale.

For the VBF production mode, the spin-2 signal is based on an effective Lagrangian approach with Λ as a characteristic energy scale of the underlying new physics [20],

$$\mathcal{L} = \frac{1}{\Lambda} T_{\mu\nu} \left(f_1 B^{\alpha\nu} B_\alpha^\mu + f_2 W_i^{\alpha\nu} W_\alpha^{i,\mu} + 2f_5 (D^\mu \Phi)^\dagger (D^\nu \Phi) \right).$$

Here, f_i are variable coupling parameters, $T_{\mu\nu}$ is the spin-2 singlet field, $B^{\alpha\nu}$ and $W_i^{\alpha\nu}$ are the electroweak field strength tensors, and Φ is the scalar Higgs field. The covariant derivative D^μ is $D^\mu = \partial^\mu - igW_i^\mu \sigma^i/2 - ig'YB^\mu$, where σ^i are the Pauli matrices, Y the weak hypercharge, and g and g' the corresponding gauge coupling constants. The model differs from the RS model in that the couplings to fermions or gluons are not included in the Lagrangian. Also, the BSM amplitude is multiplied by a form factor which is a function of a cut-off scale Λ_{ff} and a suppression power n_{ff} in order to preserve unitarity at high energies:

$$f(p_1^2, p_2^2, k_{sp2}^2) = \left(\frac{\Lambda_{ff}^2}{|p_1^2| + \Lambda_{ff}^2} \cdot \frac{\Lambda_{ff}^2}{|p_2^2| + \Lambda_{ff}^2} \cdot \frac{\Lambda_{ff}^2}{|k_{sp2}^2| + \Lambda_{ff}^2} \right)^{n_{ff}},$$

where p_1^2 and p_2^2 are the squared invariant masses of the incoming electroweak bosons and k_{sp2}^2 is the squared invariant mass of the sum of the initial boson momenta, equivalent to that of an s -channel spin-2 particle. The specific parameter settings for the signal models used are given in Section 4.

3 ATLAS detector

The ATLAS detector [47, 48] is a general-purpose particle detector used to investigate a broad range of physics processes. It includes an inner tracking detector (ID) surrounded by a thin superconducting solenoid, electromagnetic and hadronic calorimeters and a muon spectrometer (MS) incorporating three large superconducting toroidal magnets with eight coils each. The ID consists of fine-granularity silicon pixel and microstrip detectors, and a straw-tube tracker. It is immersed in a 2 tesla axial magnetic field produced by the solenoid and provides precision tracking for charged particles in the range $|\eta| < 2.5$, where η is the pseudorapidity of the particle.¹ The straw-tube detector also provides transition radiation measurements for electron identification. The calorimeter system covers the pseudorapidity range $|\eta| < 4.9$. It is composed of sampling calorimeters with either liquid argon (LAr) or scintillator tiles as the active medium, and lead, steel, copper, or tungsten as the absorber material. The MS provides muon identification and momentum measurements for $|\eta| < 2.7$. The ATLAS detector has a two-level trigger system [49] to select events for further analysis.

¹ ATLAS uses a right-handed coordinate system with its origin at the nominal interaction point (IP) in the centre of the detector and the z -axis along the beam pipe. The x -axis points from the IP to the centre of the LHC ring, and the y -axis points upwards. Cylindrical coordinates (r, ϕ) are used in the transverse plane, ϕ being the azimuthal angle around the beam pipe. The pseudorapidity is defined in terms of the polar angle θ as $\eta = -\ln \tan(\theta/2)$. Angular distance is measured in units of $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$.

4 Data and simulation samples

The data used in this analysis were collected with a single-electron or single-muon trigger. These triggers have a transverse energy or momentum threshold, E_T or p_T , that depends on the data-taking period, with the lowest threshold varying between 20 GeV and 26 GeV. The trigger efficiency for WW events passing the offline event selection (Section 6) is greater than 99%. Data quality criteria are applied to ensure that events are recorded with stable beam conditions and with all relevant subdetector systems operational.

Samples of simulated signal and background events are used to optimise the event selection and to estimate the signal acceptance and the background yields from various SM processes.

The sample for the NWA heavy Higgs boson signal was produced with POWHEG-Box 2.0 [50–52] which calculates separately the ggF [53] and VBF [54] production mechanisms with matrix elements up to next-to-leading order (NLO) in quantum chromodynamics (QCD). It uses the CT10 NLO parton distribution function (PDF) set [55] and is interfaced with PYTHIA 8.186 [56] for the $H \rightarrow WW$ decays, for parton showering and hadronisation. A set of tuned parameters called the AZNLO tune [57] is used to describe the underlying event. The NWA Higgs boson is generated with a width of 4 MeV. This event sample is also used to constrain the 2HDM. The LWA heavy Higgs boson signal was simulated at NLO using the MADGRAPH5_AMC@NLO 2.3.2 event generator [58] with the NNPDF23LO PDF set [59]. The generated particles at matrix element level are showered by PYTHIA 8.186 with the A14 tune [60] for the underlying event. The mass of the heavy Higgs boson signals considered in this analysis spans the range between 200 GeV and 4 (3) TeV for the ggF-induced (VBF-induced) signals. Both NWA and LWA samples were generated in steps of 100 GeV up to 1 TeV, and in steps of 200 GeV thereafter.

The POWHEG-Box samples describe the production of a ggF-induced heavy Higgs boson in association with one jet at leading-order (LO) precision, while further jets are emulated by the parton shower generator, PYTHIA. A more precise calculation of higher jet multiplicities is provided by using MADGRAPH5_AMC@NLO 2.3.2 to simulate $gg \rightarrow H$ events in association with up to two jets at NLO precision. Here, the overlap between identical final states generated at the matrix element (ME) and the parton shower (PS) stage is removed using FxFx merging [61]. The fraction of ggF events passing the event selection requirements of the $N_{\text{jet}} = 1$ and $N_{\text{jet}} \geq 2$ VBF categories (defined later in Section 6) predicted by the POWHEG-Box event generator is reweighted to match that of the MADGRAPH5_AMC@NLO FxFx samples. The corresponding scale factors are calculated for several hypothetical heavy Higgs boson masses. It is the largest, 1.14, for the 200 GeV mass point, and decreases with increasing resonance mass to a value of 0.85 for the 4 TeV mass point, for the $N_{\text{jet}} = 1$ VBF category. The corresponding numbers are 0.91 and 0.73 for the $N_{\text{jet}} \geq 2$ VBF category.

Benchmark samples for the GM, HVT and bulk RS models were generated at LO using MADGRAPH5_AMC@NLO interfaced to PYTHIA 8.186 with the NNPDF23LO PDF set. A value of $\sin \theta_H = 0.4$ is chosen for the GM benchmark model. For the HVT interpretation in the $q\bar{q}$ annihilation mode, samples were generated according to the extended gauge symmetry model A [18] with $g_V = 1$. In the VBF mode, samples were generated using the same g_V value but setting the couplings to the fermions to zero so that the new vector boson couples only to the SM vector and Higgs bosons. For the bulk RS model, a curvature scale parameter $k/\bar{M}_{\text{Pl}}$ of either 0.5 or 1 is considered. The ELM VBF spin-2 signals were generated at LO with VBFNLO 3.0.0 beta 2 [62] with the NNPDF30LO PDF set [63] and using the following parameter setting [20]: $\Lambda_{ff} = 3$ TeV, $n_{ff} = 4$, $\Lambda = 1.5$ TeV and $f_1 = f_2 = f_5 = 1$. The mass range considered is between 200 GeV and 5 TeV for the KK graviton signal, between 250 GeV and 5 TeV for the HVT qqA

signal, between 200 GeV and 1 TeV for the GM and ELM VBF signals, and between 300 GeV and 1 TeV for the HVT VBF signal.

The main sources of SM background include events from the production of single top quarks, $t\bar{t}$, dibosons (WW , WZ and ZZ), $Z/\gamma^* + \text{jets}$ and $W + \text{jets}$. Single-top-quark simulated events were generated with POWHEG-Box 2.0 [64, 65] using the CT10 NLO PDF set interfaced to PYTHIA 6.428 [66] for parton showering and hadronisation, with the PERUGIA2012 tune [67] and CTEQ6L1 PDF [68] to describe the underlying event. The $t\bar{t}$ events were generated with POWHEG-Box 2.0 [69] using the NNPDF30NLO PDF set [63] interfaced to PYTHIA 8.186 for parton showering and hadronisation, with the A14 tune and CTEQ6L1 PDF to describe the underlying event. The sample was generated by setting the resummation damping parameter h_{damp} to 1.5 times the top-quark mass, m_{top} , which was set to 172.5 GeV. The h_{damp} parameter controls the ME/PS matching and effectively regulates the high- p_T radiation. The EVT-GEN 1.2.0 [70] package was used to model the properties of the bottom and charm hadron decays. Diboson samples were generated with SHERPA 2.1.1 [71–75] for the gg production processes and SHERPA 2.2.1 for the $q\bar{q}$ production processes, using the CT10 NLO and NNPDF30NNLO PDF sets, respectively. The SHERPA event generator for the latter processes produces up to one additional parton at NLO and up to three additional partons at LO. Production of W and Z bosons in association with jets was also simulated using SHERPA 2.1.1 with the CT10 NLO PDF set, where b - and c -quarks are treated as massive particles. The $gg \rightarrow WW$ production also includes the contribution of the SM Higgs boson at 125 GeV and the interference effects between the continuum and Higgs resonance processes. The VBF part of SM Higgs boson production was generated with POWHEG-Box [54] interfaced to PYTHIA 8.186 for parton showering and hadronisation.

The effect of multiple pp interactions in the same and neighbouring bunch crossings (pile-up) was included by overlaying minimum-bias collisions, simulated with PYTHIA 8.186, on each generated signal and background event. The number of overlaid collisions is such that the distribution of the average number of interactions per pp bunch crossing in the simulation matches the pile-up conditions observed in the data, which is about 25 interactions per bunch crossing on average. The generated samples were processed through a GEANT4-based detector simulation [76, 77], followed by the standard ATLAS reconstruction software used for collision data.

5 Event reconstruction

Events used in this analysis are required to have at least one primary vertex with a minimum of two associated tracks, each with transverse momentum $p_T > 400$ MeV. If there is more than one vertex reconstructed in an event that meets these conditions, the one with the highest sum of track p_T^2 is chosen as the primary vertex.

Electrons are reconstructed from clusters of energy deposits in the electromagnetic calorimeter that match a track reconstructed in the ID. They are identified using the likelihood identification criteria described in Ref. [78]. The electrons used in this analysis are required to pass the “MediumLH” selection for $p_T > 25$ GeV ² or the “TightLH” selection for $p_T < 25$ GeV and be within $|\eta| < 2.47$, excluding the transition region between the barrel and endcaps in the LAr calorimeter ($1.37 < |\eta| < 1.52$). These “MediumLH” and “TightLH” selection categories have identification efficiencies of 84% and 74%,

² For electrons, p_T is defined as the magnitude of the transverse component of the electron momentum as measured using the electromagnetic calorimeter.

respectively, for electrons with p_T of 25 GeV. The corresponding probabilities to misidentify hadrons as electrons are approximately 0.5% and 0.3%, respectively.

Muons are reconstructed by combining ID and MS tracks that have consistent trajectories and curvatures [79]. The muon candidates used in this analysis are required to have $|\eta| < 2.5$ and pass the “Medium” selection for $p_T > 25$ GeV or the “Tight” selection for $p_T < 25$ GeV, defined on the basis of the quality of the reconstruction and identification. These selections have a reconstruction efficiency of approximately 96% and 92%, respectively, for muons originating from the decay of W bosons [80]. The corresponding probabilities to misidentify hadrons as muons are approximately 0.2% and 0.1%, respectively.

To ensure that leptons originate from the interaction point, a requirement of $|d_0|/\sigma_{d_0} < 5$ (3) is imposed on the electrons (muons) and $|z_0 \sin \theta| < 0.5$ mm is applied to both lepton types. Here d_0 and z_0 are the transverse and longitudinal impact parameters of the lepton with respect to the primary vertex, respectively, and σ_{d_0} is the uncertainty in the measured value of d_0 . In addition, electrons and muons are required to be isolated from other tracks and calorimetric activities by applying p_T - and η -dependent isolation criteria. For muons, the calorimeter isolation is based on energy deposits in the calorimeter within a cone ΔR of 0.2 around the muons. The muon track isolation uses a variable cone size starting at $\Delta R = 0.3$ and shrinking with increasing p_T of the muon [81]. The same calorimeter isolation is used for electrons, and the electron track isolation uses a variable cone size starting at $\Delta R = 0.2$. The efficiency of these isolation requirements is 90% for both lepton types with p_T of 25 GeV, increasing to 99% at 60 GeV.

Jets are reconstructed from three-dimensional clusters of energy deposits in the calorimeters using the anti- k_t algorithm [82] with a radius parameter of $R = 0.4$ implemented in the FastJet package [83]. The four-momenta of the jets are calculated as the sum of the four-momenta of their constituents, which are assumed to be massless. Jets are corrected for energy from pile-up using the pile-up subtraction based on jet areas [84]. The jet energy scale is estimated in Ref. [85]. Jets are required to have $p_T > 30$ GeV and $|\eta| < 4.5$.

For jets with $p_T < 60$ GeV and $|\eta| < 2.5$, the multivariate “jet vertex tagger” algorithm [86] is used to suppress jets from pile-up interactions. To avoid double counting, jets of any transverse momentum are discarded if they are within a cone of size $\Delta R = 0.2$ around an electron candidate or if they have fewer than three associated tracks and are within a cone of size $\Delta R = 0.2$ around a muon candidate. However, if a jet with three or more associated tracks is within a cone of size $\Delta R < 0.4$ of a muon candidate, or the separation between an electron and any jet is within $0.2 < \Delta R < 0.4$, the corresponding muon or electron candidate is rejected.

To estimate the number of b -tags in the event, jets with $p_T > 20$ GeV and within $|\eta| < 2.5$ are considered to contain a b -hadron if they yield a b -tagging algorithm discriminant value exceeding a reference value. The MV2c10 algorithm [87, 88] is chosen at the 85% b -tagging efficiency benchmark point, estimated from b -jets in simulated $t\bar{t}$ events. The misidentification rate for jets which originate from a light quark or gluon is less than 1%, while it is approximately 17% for c -jets.

The missing transverse momentum, with magnitude E_T^{miss} , is calculated as the negative vectorial sum of the transverse momenta of calibrated electrons, muons, and jets originating from the primary vertex, as well as tracks with $p_T > 500$ MeV compatible with the primary vertex and not associated with any of these [89].

Table 2: Selection conditions and phase space definitions used in the ggF and VBF signal regions.

SR _{ggF}	SR _{VBF1J}	SR _{VBF2J}
Common selections		
$N_{b\text{-tag}} = 0$ $ \Delta\eta_{\ell\ell} < 1.8$ $m_{\ell\ell} > 55 \text{ GeV}$ $p_T^{\ell, \text{lead}} > 45 \text{ GeV}$ $p_T^{\ell, \text{sublead}} > 30 \text{ GeV}$ veto if $p_T^{\ell, \text{other}} > 15 \text{ GeV}$ $\max(m_T^W) > 50 \text{ GeV}$		
ggF phase space	VBF1J phase space	VBF2J phase space
Inclusive in N_{jet} but excluding VBF1J and VBF2J phase space	$N_{\text{jet}} = 1$ and $ \eta_j > 2.4, \min(\Delta\eta_{j\ell}) > 1.75$	$N_{\text{jet}} \geq 2$ and $m_{jj} > 500 \text{ GeV}, \Delta y_{jj} > 4$

6 Event selection

As a first step, WW candidate events are selected by requiring two oppositely charged, different-flavour leptons (e or μ). Both leptons must satisfy the minimal quality criteria discussed in Section 5. When ordered in p_T , these leptons are called the leading and subleading ones, $p_T^{\ell, (\text{sub})\text{lead}}$. In order to suppress the background from diboson processes, a veto is imposed on events with an additional lepton with $p_T^{\ell, \text{other}} > 15 \text{ GeV}$.

Table 2 summaries the selections and the definition of signal regions (SRs). The variables used in the selections are the most discriminating ones chosen by a boosted decision tree (BDT) [90], based on the NWA signal samples. These are $p_T^{\ell, \text{lead}}$, the invariant mass of the leading and subleading leptons, $m_{\ell\ell}$, and the pseudorapidity difference between the two leptons, $\Delta\eta_{\ell\ell}$. The first two variables provide good separation between a heavy resonance signal and the WW and top-quark background. The separation of signal from background based on the $\Delta\eta_{\ell\ell}$ distribution is found to have a reasonable efficiency and allows, at the same time, a control region to be defined for the WW background (Section 7.2). For each selected variable, the selection criterion is set by maximising the signal significance in the presence of background. The optimised selection is checked to be applicable to the LWA signals.

In order to further suppress the top-quark background, events with at least one b -tagged jet ($N_{b\text{-tag}} \geq 1$) are rejected from the signal regions. To reduce the Z +jets and W +jets contributions, two other variables are used: $p_T^{\ell, \text{sublead}}$ and the maximum value of the transverse mass calculated with either of the two leptons and the missing transverse momentum, m_T^W . The latter variable is defined as:

$$m_T^W = \sqrt{2p_T^\ell E_T^{\text{miss}} (1 - \cos(\phi^\ell - \phi_T^{E^{\text{miss}}}))},$$

where p_T^ℓ and ϕ^ℓ are the transverse momentum and azimuthal angle of a given lepton and $\phi_T^{E^{\text{miss}}}$ is the azimuthal angle of the missing transverse momentum vector.

Three event categories are defined: two disjoint categories optimised for the VBF production, VBF $N_{\text{jet}} = 1$ and VBF $N_{\text{jet}} \geq 2$ (SR_{VBF1J} and SR_{VBF2J}), and one quasi-inclusive category (excluding the




Figure 1: Acceptance times efficiency as a function of signal mass for the ggF or qqA (left) and VBF (right) productions. All three signal event categories are combined. The hatched band around the NWA signal curve shows the typical size of the total statistical and systematic uncertainties.

VBF phase space) dedicated to the ggF or qqA signal (SR_{ggF}). For the VBF $N_{jet} = 1$ category, two discriminating variables are used to minimise the contribution of the ggF signal: the pseudorapidity of the jet, η_j , and the minimum value of the pseudorapidity difference between the jet and either of the leptons, $\min(|\Delta\eta_{j\ell}|)$. For the VBF $N_{jet} \geq 2$ category, the invariant mass, m_{jj} , and the rapidity difference, Δy_{jj} , of the two leading jets are used to select the VBF signal.

The NWA and LWA signal acceptance times the efficiency, after all selection requirements for a 700 GeV ggF signal, is approximately 50% in the quasi-inclusive ggF category and 5% or less in the VBF $N_{jet} = 1$ and $N_{jet} \geq 2$ categories. For a 700 GeV VBF signal, it is between 15% and 25% for the three event categories. The acceptance times efficiency for the three event categories combined, as a function of resonance mass, is shown in Figure 1 for the different signals. For the spin-1 and spin-2 signals, the range up to 1 TeV is considered in the case of VBF model processes. For samples with lower resonance masses, the acceptance times efficiency is lower because the leptons are softer. This is also the reason why the search is limited to signal mass values greater than about 200 GeV. The same selection is applied to all models and the different selection efficiencies between the models are mainly due to different $\Delta\eta_{\ell\ell}$ distributions for the different spin states.

The discriminating variable used for the statistical analysis (Section 9) in this search is the transverse mass defined as

$$m_T = \sqrt{(E_T^{\ell\ell} + E_T^{\text{miss}})^2 - |\mathbf{p}_T^{\ell\ell} + \mathbf{E}_T^{\text{miss}}|^2},$$

where

$$E_T^{\ell\ell} = \sqrt{|\mathbf{p}_T^{\ell\ell}|^2 + m_{\ell\ell}^2},$$

and $\mathbf{p}_T^{\ell\ell}$ is the transverse momentum vector of the leading and subleading leptons.

7 Background estimation

The dominant background for the $e\nu\mu\nu$ final state is due to events with top quarks and due to SM WW events. Additional contributions to the background arise from V +jets and the diboson processes VZ , $V\gamma$ and $V\gamma^*$. Since the discriminating variable used for this search is the transverse mass, m_T , both the

normalisation and the shape of the background m_T distribution must be estimated. The shape of the background is modelled using simulated events while the top-quark and WW background normalisations are determined by a simultaneous fit (Section 9) to the data in m_T -binned distributions in the signal regions and the total event yields in control regions. The normalisation factors of the fit, named “post-fit” normalisation factors ³ hereafter, provide the best overall matching between the number of observed data events and the corresponding SM background expectations in all the signal and control regions. The control regions are defined by criteria similar to those used for the signal regions, but with some requirements loosened or reversed to obtain signal-depleted samples, enriched in the relevant background. These criteria are summarised in Table 3.

The following subsections describe the methods used to estimate the most important background processes, namely top quark, WW , and W +jets. The Z/γ^* +jets and non- WW diboson background contributions are small. The Z/γ^* +jets Monte Carlo (MC) samples are normalised using NNLO cross sections [91] and the non- WW ones with NLO cross sections from the SHERPA event generator. The small background from the $m_h \simeq 125$ GeV Higgs boson resonance and its off-shell component is included and its interference with the continuum WW background is taken into account.

7.1 Top-quark background

Events containing top quarks can be produced as a $t\bar{t}$ pair or as a single top quark in association with either a W boson or a quark of another flavour. In this analysis, contributions from $t\bar{t}$ and single-top-quark events are estimated together, with their relative contributions determined by their predicted cross sections and by their relative acceptances obtained from MC simulation. The single-top-quark contribution varies from about 10% to 30% depending on the signal event category.

The normalisation of the top-quark background for the quasi-inclusive ggF category is determined in a control region (Top CR_{ggF}) where one jet is required to be b -tagged in addition to the signal region selection. The purity of the top-quark background in this CR is high (97%) and thus allows the modelling of the MC simulation to be validated. The distribution of the simulated leading lepton p_T in the Top CR_{ggF} is found to disagree with the data and the ratio between the data and the simulation decreases with increasing $p_T^{\ell, \text{lead}}$. The simulated distribution is corrected in the SR_{ggF} and corresponding CRs with factors obtained by fitting the ratio with a linear function. The correction varies between +4% and −10% as $p_T^{\ell, \text{lead}}$ increases from 50 GeV to 200 GeV.

The top-quark background control regions for the VBF categories (Top CR_{VBF}) have a small number of data events and are therefore merged. At least one jet is required to be b -tagged. In addition, the selection thresholds imposed on $m_{\ell\ell}$ and $p_T^{\ell, (\text{sub})\text{lead}}$ are relaxed to 10 GeV and 25 GeV, respectively, and the selection on $|\Delta\eta_{\ell\ell}|$ and $\max(m_T^W)$ is removed. The threshold value on $m_{\ell\ell}$ of 10 GeV is used to suppress background contributions from low-mass resonances decaying into different-flavour final states via $\tau^+\tau^-$. In this control region, the purity of the top-quark background is 96%, and no mis-modelling of the $p_T^{\ell, \text{lead}}$ distribution is observed.

The post-fit normalisation factors from the simultaneous fit are 0.96 ± 0.05 and $1.12^{+0.13}_{-0.12}$ in the ggF and the VBF control regions, respectively, where the uncertainty quoted corresponds to the combined statistical and systematic uncertainties.

³ The post-fit normalisation factors are checked to be consistent within the quoted uncertainties with the pre-fit ones obtained using the control regions only.

Table 3: Summary of all the selections used in the ggF and VBF WW and top-quark control regions. The common selection “veto if $p_T^{\ell, \text{other}} > 15 \text{ GeV}$ ” applied to all the regions is not explicitly shown.

WW CR _{ggF}	Top CR _{ggF}	WW CR _{VBF1J}	Top CR _{VBF}
$N_{b\text{-tag}} = 0$ $ \Delta\eta_{\ell\ell} > 1.8$ $m_{\ell\ell} > 55 \text{ GeV}$ $p_T^{\ell, \text{lead}} > 45 \text{ GeV}$ $p_T^{\ell, \text{sublead}} > 30 \text{ GeV}$ $\max(m_T^W) > 50 \text{ GeV}$	$N_{b\text{-tag}} = 1$ $ \Delta\eta_{\ell\ell} < 1.8$	$N_{b\text{-tag}} = 0$ $(\Delta\eta_{\ell\ell} > 1.8 \text{ or } 10 \text{ GeV} < m_{\ell\ell} < 55 \text{ GeV})$ $p_T^{\ell, \text{lead}} > 25 \text{ GeV}$ $p_T^{\ell, \text{sublead}} > 25 \text{ GeV}$ —	$N_{b\text{-tag}} \geq 1$ — $m_{\ell\ell} > 10 \text{ GeV}$
Excluding VBF1J and VBF2J phase space		VBF1J phase space	VBF1J and VBF2J phase space




Figure 2: Transverse mass distribution in the ggF (left) and VBF (right) top-quark control regions. In each plot, the last bin contains the overflow. The hatched band in the upper and lower panels shows the combined statistical, experimental and theoretical uncertainties in the predictions. The arrow in the lower right panel indicates that an entry is outside of the vertical scale. The top-quark and WW background event yields are scaled using the indicated normalisation factors obtained from the simultaneous fit to all signal and control regions. The heavy Higgs boson signal event yield, normalised to the expected limits on $\sigma_H \times B(H \rightarrow WW)$, is shown for masses of 700 GeV and 2 TeV in the NWA scenario.

Figure 2 shows the m_T distributions in the ggF and VBF top-quark CRs. The different background components are scaled according to the event yields obtained from the simultaneous fit. In the control regions the fit uses only the integrated event yields. The shape of the distributions is compared between data and MC predictions and found to be in good agreement after the application of the $p_T^{\ell, \text{lead}}$ correction described above for the ggF top-quark CR. The shapes of the m_T distribution for 700 GeV and 2 TeV NWA Higgs boson signals are also shown, normalised to the expected limits on $\sigma_H \times B(H \rightarrow WW)$ from this analysis. The ggF contribution from the SM Higgs boson is included in the WW component. The SM Higgs boson VBF contribution is negligibly small and is not shown in this and following figures.

7.2 WW background

The WW CR for the quasi-inclusive ggF category (WW CR_{ggF}) uses the same selection as for the SR except for $|\Delta\eta_{\ell\ell}|$ which is reversed so that the CR and SR are orthogonal. The selection conditions are shown in Table 3. The m_T distributions of the $q\bar{q} \rightarrow WW$ SHERPA MC sample in the SR_{ggF} and WW CR_{ggF} are compared at MC generator level with corresponding predictions combining NNLO QCD calculations [92] with NLO electroweak (EW) corrections [93]. While the integrated yields of the distributions agree within 3% in both the SR_{ggF} and the WW CR_{ggF}, a small m_T shape difference is observed, particularly in the SR. The m_T distributions of the SHERPA samples are thus reweighted to the combined NNLO QCD and NLO EW predictions. The post-fit normalisation factor obtained from the simultaneous fit for the WW contributions in the quasi-inclusive ggF categories is 1.14 ± 0.09 , where the uncertainty quoted corresponds to the combined statistical and systematic uncertainties. The post-fit purity of the WW background in the control region is 51%.

In order to select more data events, the WW CR for the $N_{\text{jet}} = 1$ VBF category (WW CR_{VBF1J}) uses a slightly different selection (shown in Table 3) from the one in the SR, but still disjoint from the SR. The normalisation factor obtained from the same simultaneous fit for the WW contribution in the WW CR_{VBF1J} is 1.0 ± 0.2 , where the uncertainty quoted corresponds to the combined statistical and systematic uncertainties. The post-fit purity of the WW background in the control region is 44%.

The WW contribution in the $N_{\text{jet}} \geq 2$ VBF category is about 20%, and its prediction is taken from simulation because it is difficult to isolate a kinematic region with a sufficient number of WW events and with a small contamination from the top-quark background.

Figure 3 shows the m_T distributions in the WW CR_{ggF} and CR_{VBF1J}. The different background contributions are scaled according to the event yields obtained from the simultaneous fit. For the WW control regions only integrated event yields are used in the fit, like in the fits of the top control regions.

7.3 W +jets background

Events with W bosons produced in association with jets may enter the SR when a jet is misidentified as a lepton. Due to the difficulties in accurately modelling the misidentification process in the simulation, the W +jets background contribution is estimated using the data-driven method developed for the SM $h \rightarrow WW$ analysis [94]. A sample of events is used which satisfies all event selection criteria, except that one of the two lepton candidates fails to meet the quality criteria for being an identified lepton but satisfies a less restrictive selection, referred to as “anti-identified”. Anti-identified muons (electrons) have loosened isolation and impact parameter (likelihood identification) selection criteria as compared to the identified selection. From this data sample the non- W +jets contribution, dominated by top-quark and WW background processes, is subtracted on the basis of MC predictions. The W +jets purity of the samples is 46%, 59% and 22% for the quasi-inclusive ggF, $N_{\text{jet}} = 1$ and $N_{\text{jet}} \geq 2$ VBF categories, respectively.

The W +jets contamination in the signal region is then determined by scaling the number of events in the background-subtracted data sample by an extrapolation factor, which is the ratio of the number of identified leptons to the number of anti-identified leptons in a data sample of dijet events in bins of lepton p_T and η . The dijet sample is collected using prescaled low- p_T single-lepton triggers with thresholds of 12 GeV for electrons and 14 GeV for muons. Events are selected with exactly one candidate lepton, back-to-back with the leading jet. Electroweak processes in the dijet event sample, dominated by W +jets and Z/γ^* background contributions, are subtracted. The dominant systematic uncertainty in the estimation




Figure 3: Transverse mass distribution in the quasi-inclusive ggF (left) and $N_{\text{jet}} = 1$ VBF WW (right) control regions. In each plot, the last bin contains the overflow. The hatched band in the upper and lower panels shows the combined statistical, experimental and theoretical uncertainties in the predictions. The top-quark and WW background events are scaled using the indicated normalisation factors obtained from the simultaneous fit to all signal and control regions. The heavy Higgs boson signal event yield, normalised to the expected limits on $\sigma_H \times B(H \rightarrow WW)$, is shown for masses of 700 GeV and 2 TeV in the NWA scenario.

of the W +jets background is due to the differences between dijet and W +jets sample characteristics. All systematic uncertainties associated with this background estimate are listed in Section 8.1.

8 Systematic uncertainties

In this section, experimental and theoretical uncertainties in the normalisation and shape of the m_T distributions of the background and the signal are described. Except for those explicitly mentioned here, the shape uncertainties are small and thus neglected. Overall, the systematic uncertainty dominates, except in the tails of the m_T distributions where the statistical uncertainty is larger.

8.1 Experimental uncertainties

The dominant sources of experimental uncertainty in the signal and background yields are the jet energy scale and resolution (Jet) [85], the b -tagging efficiency (b -tag) [87], and the pile-up modelling [86]. Other systematic uncertainties such as those associated with trigger efficiencies, lepton reconstruction and identification efficiencies, lepton momentum scales and resolutions [78, 80], missing transverse momentum reconstruction [89] and the jet vertex tagger [86] are also considered when evaluating systematic effects on the shape and normalisation of the background, or the shape and efficiency of the signal yield. The uncertainty associated with the pile-up modelling is assessed by performing a variation of $\pm 9\%$ in the number of simulated pile-up interactions to cover the uncertainty in the ratio of the predicted and measured cross sections of non-diffractive inelastic events producing a hadronic system of mass $m_{X,\text{had}} > 13$ GeV [95].

For the main background from top-quark and WW processes, the impact of the most important experimental systematic uncertainties is summarised in Tables 4 and 5 together with dominant theoretical uncertainties. The maximum changes in yield for the up and down variations are shown in the various signal and control regions. The correlation between the SRs and CRs is taken into account in the simultaneous fit.

Systematic effects due to lepton identification efficiencies, momentum and scale resolutions, are found to be approximately 1%. They are not shown in the tables. The last column in the tables shows the total uncertainty, including these small uncertainty sources.

The data-driven W +jets background estimate is subject to several sources of systematic uncertainty. The subtraction of the subdominant electroweak processes (Section 7.3) has a significant impact on the extrapolation factor calculation at high lepton p_T . The subtraction is varied, as described in Ref. [94], and the variation of the event yield in the signal region is taken as the uncertainty. The method assumes that the extrapolation factors of the dijet and W +jets samples are equal. Differences in the jet flavour composition between dijet and W +jets events introduce an additional systematic uncertainty. This is evaluated as the sum in quadrature of two contributions: differences between the extrapolation factors calculated with dijet samples and Z +jets samples in data, and differences between the extrapolation factors evaluated with W +jets and Z +jets MC samples. Finally, the statistical uncertainties of the different data and MC samples used to evaluate the extrapolation factors are taken as an additional source of systematic uncertainty. The overall relative systematic uncertainty of the W +jets background is found to be approximately 35% for each of the three signal event categories, with the dominant uncertainty being associated with the jet flavour composition.

The uncertainty in the total 2015 and 2016 integrated luminosity is 2.1%. It is derived, following a methodology similar to that detailed in Ref. [96], from van der Meer scans performed in August 2015 and May 2016, calibrated at high luminosity by various luminosity detectors.

8.2 Theoretical uncertainties of the background

For background sources which are normalised using control regions, theoretical uncertainties are evaluated for the extrapolation from the control region to the signal region.

For the top-quark and WW background, theoretical uncertainties in the extrapolation are evaluated according to the prescription from the LHC Higgs Cross Section Working Group [97]. The uncertainties include the impact of missing higher-order corrections, PDF variations and other MC modelling. The dominant theoretical uncertainties are shown in Tables 4 and 5.

For the top-quark background, the uncertainty from the event generator and parton shower modelling (ME+PS) is estimated by comparing the nominal POWHEG-Box+PYHTIA8 generated samples with those from an alternative event generator, SHERPA 2.2.1. The uncertainty named “Scale” corresponds to variations of the renormalisation μ_R and factorisation μ_F scales as well as h_{damp} . The variations for μ_R and μ_F are between 0.5 and 2 from their nominal scale of $\sqrt{m_{\text{top}}^2 + p_T^2}$, with p_T being the top-quark transverse momentum. The parameter h_{damp} is varied between m_{top} and $2 \cdot m_{\text{top}}$ from its nominal scale $h_{\text{damp}} = 1.5 \cdot m_{\text{top}}$. In the analysis the single-top-quark and $t\bar{t}$ processes are studied together. An uncertainty of 20% [98, 99] is assigned to the relative contribution of the single-top-quark processes, corresponding to the source “Single top” in Table 4. The PDF uncertainty is obtained by taking the envelope of the uncertainty of the NNPDF30NLO PDF set and its differences in central value with the CT14 [100] and MMHT 2014 [101] PDF sets, following the recommendations of Ref. [55]. The PDF uncertainties are m_T dependent and

Table 4: Relative impact (in %) of dominant experimental and theoretical uncertainties in the event yields for the top-quark background processes in the three signal regions (SR_{ggF} , SR_{VBF1J} and SR_{VBF2J}) and the top-quark and WW control regions (Top $CR_{ggF/VBF}$ and the WW $CR_{ggF/VBF1J}$). Jet and b -tag sources dominate the experimental uncertainty while ME+PS, Scale, Single top and PDF are the dominant theoretical uncertainties. The last column shows the total uncertainty including those not listed here.

Source	Jet	b -tag	ME+PS	Scale	Single top	PDF	Total
SR_{ggF}	5.2	17	1.3	3.0	4.2	2.5	19
SR_{VBF1J}	9.6	7.8	1.0	1.6	5.9	2.6	15
SR_{VBF2J}	9.7	14	9.5	5.0	2.1	3.4	21
Top CR_{ggF}	2.2	4.8	0.34	0.21	2.6	3.0	6.6
WW CR_{ggF}	5.3	18	1.1	6.3	4.0	3.2	20
Top CR_{VBF}	8.2	3.5	10	1.5	1.3	3.7	14
WW CR_{VBF1J}	9.9	8.3	9.4	3.9	5.3	2.7	18

Table 5: Relative impact (in %) of dominant experimental and theoretical uncertainties in the event yields for the WW background processes in the three signal regions (SR_{ggF} , SR_{VBF1J} and SR_{VBF2J}) and the WW control regions (WW $CR_{ggF/VBF1J}$). Jet and Pile-up sources dominate the experimental uncertainty while ME+PS, μ_R , Resummation and PDF are the dominant theoretical uncertainties. The last column shows the total uncertainty including those not listed here.

Source	Jet	Pile-up	ME+PS	μ_R	Resummation	PDF	Total
SR_{ggF}	1.2	1.8	2.4	1.7	3.1	2.7	5.5
SR_{VBF1J}	17	2.8	11	7.3	5.0	2.3	23
SR_{VBF2J}	18	3.1	38	18	1.4	2.1	47
WW CR_{ggF}	1.1	1.8	2.6	0.95	2.9	3.6	5.9
WW CR_{VBF1J}	16	4.5	12	11	2.3	2.8	23

increase from 2% to 10% with m_T . This m_T dependence is taken into account in the signal regions. In the ggF quasi-inclusive category, two additional shape systematic uncertainties associated with the scale variations and the p_T reweighting for the leading lepton in the top-quark background are applied, the latter corresponding to $\pm 50\%$ of the reweighting correction. These two uncertainties are comparable and vary from a few percent at low m_T to about 10% at $m_T \simeq 1$ TeV, without affecting the integrated event yield of the top-quark background in the category.

For the WW background, the ME+PS modelling uncertainty is obtained by comparing the nominal SHERPA 2.2.1 sample with an alternative sample generated with POWHEG-Box+PYTHIA8. The renormalisation, factorisation, and resummation scales are varied separately by factors of 0.5 and 2. The uncertainty corresponding to the factorisation scale variation is smaller than the other uncertainties and is not shown. The PDF uncertainty for the WW background is obtained and treated in the same way as for the top-quark background. In the ggF quasi-inclusive category, an additional shape uncertainty from ME+PS is applied. It varies from a few percent at low m_T to about 20% at $m_T \simeq 1$ TeV. There are no significant shape uncertainties in the m_T distributions in the VBF categories.

In addition to the scale uncertainties described above, a relative uncertainty of $\pm 50\%$ is assigned to the reweighting corrections of the $q\bar{q} \rightarrow WW$ SHERPA sample to the combined NNLO QCD and NLO EW predictions in the ggF SR and WW CR.

The $gg \rightarrow (h^*) \rightarrow WW$ process, where the SM 125 GeV Higgs boson is off-shell, is modelled at leading order with the SHERPA event generator with a K -factor of 1.7 that is used to account for higher-order cross-section corrections with an uncertainty of 60%, following the studies in Refs. [102–105].

Other small background processes, such as WZ , ZZ , $Z/\gamma^* + \text{jets}$ and WW in the $N_{\text{jet}} \geq 2$ VBF category, do not have their own control regions. They are normalised to the theoretical predictions. The uncertainties in their yields due to the uncertainties in the predictions are evaluated with the same prescription as described above. The impact of these uncertainties is small (see Tables 6 and 7 in Section 9).

8.3 Theoretical uncertainties in the signal predictions

Theoretical uncertainties in the signal acceptance include effects due to the choice of QCD renormalisation and factorisation scales, the PDF set as well as the underlying-event modelling, the parton shower model and the parton shower tune. These uncertainties are evaluated separately in each of the three event categories as a function of the resonance mass and independently for ggF- and VBF-induced resonances.

The effect of missing higher-order corrections in QCD on the signal acceptance is estimated by varying the renormalisation and factorisation scales independently by factors of 0.5 and 2 from the nominal scale of $\sqrt{m_H^2 + p_{T,H}^2}$, with m_H and $p_{T,H}$ being the mass and the transverse momentum of the heavy Higgs boson, respectively. The acceptance values obtained with these modified MC samples are compared to the signal acceptance of the nominal sample. For resonances produced via ggF, these uncertainties are found to be negligible in the quasi-inclusive ggF and $N_{\text{jet}} = 1$ VBF categories, while in the $N_{\text{jet}} \geq 2$ VBF category they range between 2.5% and 0.2% for a resonance mass varying from 200 GeV to 4 TeV (unless stated otherwise, the following uncertainties are quoted for the same mass range). For resonances produced via vector-boson fusion, these uncertainties range from 0.9% to 2.8% in the quasi-inclusive ggF category, from 1.9% to 3.6% in the $N_{\text{jet}} = 1$ VBF category and from 1.0% to 7.3% in the $N_{\text{jet}} \geq 2$ VBF category.

The PDF-induced uncertainties in the signal acceptance are determined in the same way as for the top-quark and WW background processes. For the ggF-induced (VBF-induced) signal, these uncertainties reach 0.4% (1.7%), 1.5% (1.2%) and 1.6% (1.5%) for the quasi-inclusive ggF, $N_{\text{jet}} = 1$ and $N_{\text{jet}} \geq 2$ VBF event categories, respectively.

The uncertainties corresponding to the parton shower tune and the underlying event are derived by moving independently, up or down, the PYTHIA internal parameters that are associated with final-state radiation or the multiple parton interactions to study separately their influence on the signal acceptance of the various signal mass points. These uncertainties are compared for each event category and mass point to the uncertainties from the choice of parton shower model, which are estimated by comparing the results obtained for the nominal parton shower generator to those obtained using HERWIG++ [106, 107]. The tune uncertainties are found to be smaller than the shower uncertainties for all mass points. Thus only the latter uncertainties are considered in the final results. The corresponding uncertainties for ggF-induced signals increase from 1.3% to 3.1%, from 13% to 28%, and from 2.3% to 15% for increasing resonance masses in the quasi-inclusive ggF, $N_{\text{jet}} = 1$ and $N_{\text{jet}} \geq 2$ VBF categories, respectively. The uncertainties for VBF-induced signals increase from 4.3% to 19%, from 5.1% to 9.0%, and from 3.3% to 8.0% in the three categories.

Table 6: Event yields in the signal and control regions for the quasi-inclusive ggF category. The predicted background yields and uncertainties are calculated after the simultaneous fit to the data in all the SRs and the CRs including those from Table 7. The statistical and systematic uncertainties are combined. The notation “VV” represents non- WW diboson background.

	SR _{ggF}	Top CR _{ggF}	WW CR _{ggF}
WW	11 500 $\pm$ 800	820 $\pm$ 120	3 360 $\pm$ 220
Top quark	11 800 $\pm$ 600	52 550 $\pm$ 330	2 610 $\pm$ 180
Z/γ^*	1 420 $\pm$ 110	111 $\pm$ 20	20.9 $\pm$ 2.0
W+jets	1 180 $\pm$ 320	710 $\pm$ 190	280 $\pm$ 70
VV	866 $\pm$ 34	101 $\pm$ 12	250 $\pm$ 11
Background	26 740 $\pm$ 170	54 290 $\pm$ 250	6 510 $\pm$ 80
Data	26 739	54 295	6 515

In addition, uncertainties due to missing higher-order corrections in QCD are evaluated for ggF-induced processes for each event category, considering also event migration effects between different event categories. This follows the method proposed by Stewart and Tackmann [108]. The corresponding uncertainties range from 3% to 10% for the quasi-inclusive ggF category and from 4% to 30% (30% – 60%) for the $N_{\text{jet}} = 1$ ($N_{\text{jet}} \geq 2$) VBF event categories.

9 Results

The statistical method used to interpret the results of the search is described in Ref. [109]. A likelihood function $\mathcal{L}$ is defined as the product of Poisson probabilities associated with the number of events in bins of the m_T distributions in the signal regions and of the total yields in the control regions. Each source of systematic uncertainty is parameterised by a corresponding nuisance parameter θ constrained by a Gaussian function.

The m_T distributions in the signal regions are divided into 18 (8) bins for the ggF quasi-inclusive ($N_{\text{jet}} = 1$ and ≥ 2 VBF) categories. The bins are of variable size to reflect the increasing width of the m_T distribution of the expected signal with increasing mass, while keeping the statistical precision of the background contributions in each bin sufficiently high.

The numbers of events predicted and observed in the signal and control regions are shown for the quasi-inclusive ggF categories in Table 6 and for the VBF $N_{\text{jet}} = 1$ and ≥ 2 categories in Table 7. These yields are obtained from a simultaneous fit to the data in all the SRs and the CRs. The fitted signal event yield is consistent with zero. The background compositions depend strongly on the event categories: the top-quark and WW processes are comparable in SR_{ggF} and SR_{VBF1J} while the top-quark events dominate in SR_{VBF2J}. The large reduction of the total background uncertainty is due to strong anti-correlations between some of the uncertainty sources of the top-quark and WW background. The m_T distributions in SR_{ggF}, SR_{VBF1J} and SR_{VBF2J} are shown in Figure 4. As no excess over the background prediction is observed, upper limits at 95% confidence level (CL) are set on the production cross section times the branching fraction, $\sigma_X \times B(X \rightarrow WW)$, for signals in each benchmark model.

The 95% CL upper limits are computed using the modified frequentist method known as CL_s [110], using the asymptotic approximation of the distribution of a test statistic [111], q_μ , a function of the signal




Figure 4: Post-fit distributions of the transverse mass m_T in the SR_{ggF} (top left), SR_{VBF1J} (top right) and SR_{VBF2J} (bottom) categories. In each plot, the last bin contains the overflow. The hatched band in the upper and lower panels shows the total uncertainty of the fit. The top-quark and WW background event yields are scaled using the indicated normalisation factors obtained from the simultaneous fit to all signal and control regions. The heavy Higgs boson signal event yield is normalised to the expected limits on $\sigma_H \times B(H \rightarrow WW)$ and is shown for masses of 700 GeV and 2 TeV in the NWA scenario.

Table 7: Event yields in the signal and control regions for the $N_{\text{jet}} = 1$ and ≥ 2 VBF categories. The predicted background yields and uncertainties are calculated after the same simultaneous fit to the data in all the event categories as in Table 6. The statistical and systematic uncertainties are combined. The notation “VV” represents non- WW diboson background.

	SR _{VBF1J}	SR _{VBF2J}	Top CR _{VBF}	WW CR _{VBF1J}
WW	390 ± 50	120 ± 26	61 ± 11	265 ± 32
Top quark	450 ± 50	391 ± 24	5 650 ± 90	167 ± 18
Z/γ*	45 ± 11	24 ± 6	68 ± 19	74 ± 12
W+jets	52 ± 13	8.9 ± 2.5	91 ± 24	43 ± 11
VV	32 ± 7	16.6 ± 1.9	20 ± 9	38 ± 4
Background	972 ± 29	563 ± 22	5 890 ± 80	596 ± 22
Data	978	560	5 889	594

strength μ , defined as the ratio of the measured $\sigma_X \times B(X \rightarrow WW)$ to that of the prediction:

$$q_\mu = -2 \ln \left(\frac{\mathcal{L}(\mu; \hat{\theta}_\mu)}{\mathcal{L}(\hat{\mu}; \hat{\theta})} \right).$$

The quantities $\hat{\mu}$ and $\hat{\theta}$ are those values of μ and θ , respectively, that unconditionally maximise $\mathcal{L}$. The numerator depends on the values $\hat{\theta}_\mu$ that maximise $\mathcal{L}$ for a given value of μ .

Limits are obtained separately for ggF and VBF production for the NWA and LWA signal hypotheses. To derive the expected limits on the ggF (VBF) production modes, the VBF (ggF) production cross section is set to zero so that the expected limits correspond to the background-only hypothesis. To derive the observed limits on the ggF (VBF) production mode, the VBF (ggF) production cross section is treated as a nuisance parameter in the fit and profiled, in the same way as dealing with the normalisation factors of the different background processes. This approach avoids making any assumption about the presence or absence of the signal in any of these production modes.

Figure 5 shows the 95% CL upper limits on $\sigma_H \times B(H \rightarrow WW)$ as a function of m_H for a Higgs boson in the NWA scenario in the mass range $200 \text{ GeV} \leq m_H \leq 4(3) \text{ TeV}$ for the ggF (VBF) production. Values above 6.4 pb (1.3 pb) at $m_H = 200 \text{ GeV}$ and above 0.008 pb (0.006 pb) at 4 (3) TeV are excluded at 95% CL by the quasi-inclusive ggF (VBF) NWA analysis. The main systematic uncertainties affecting the limits are the p_T correction for the leading lepton in the top-quark background, scale variations for the top-quark background, the parton shower modelling of the WW MC generator, and the jet energy scale and resolution uncertainties. Limits are consistent with those expected in the absence of a signal over the investigated mass range. The fact that the observed limits are more stringent than the expected ones for mass values beyond 2 TeV is explained by the deficit in data at the high m_T tail in Figure 4. These limits are extracted using the asymptotic approximation and their accuracy is verified to be consistent within about 5% at 800 GeV and better than 20% at 2 TeV and beyond using pseudo-experiments.

The analysis can be extended to a more general case where the relative fraction of the ggF production cross section varies over the total ggF and VBF production cross section. The corresponding 95% CL upper exclusion limits for a signal at 800 GeV are shown in Figure 6. The dependence of the limits on the ggF fraction for other masses is similar but becomes slightly stronger (weaker) for lower (higher) mass values. The limit values for a ggF fraction of 0 and 1 are comparable with the VBF and ggF limits shown




Figure 5: Upper limits at 95% CL on the Higgs boson production cross section times branching fraction $\sigma_H \times B(H \rightarrow WW)$ in the $e\nu\mu\nu$ channel, for ggF (left) and VBF (right) signals with narrow-width lineshape as a function of the signal mass. The inner and outer bands show the $\pm 1\sigma$ and $\pm 2\sigma$ ranges around the expected limit.




Figure 6: Upper limits at 95% CL on the total ggF and VBF Higgs boson production cross section times branching fraction $\sigma_H \times B(H \rightarrow WW)$ in the $e\nu\mu\nu$ channel, for a signal at 800 GeV as a function of the ggF cross section divided by the combined ggF and VBF production cross section. The inner and outer bands show the $\pm 1\sigma$ and $\pm 2\sigma$ ranges around the expected limit.

in Figure 5 at the same mass value. The VBF limits are tighter than the ggF limits since the VBF $N_{\text{jet}} \geq 2$ signal region has the smallest background contribution and thus is the most sensitive.

The NWA exclusion limit shown above can be further translated to exclusion contours in the 2HDM for the phase space where the narrow-width approximation is valid. The 95% CL exclusion contours for Type I and Type II in the plane of $\tan \beta$ and $\cos(\beta - \alpha)$ for three mass values of 200 GeV, 300 GeV and 500 GeV are shown in Figure 7. For a fixed value of $\cos(\beta - \alpha) = -0.1$, 95% CL exclusion limits on $\tan \beta$ as a function of the heavy Higgs boson mass are shown in Figure 8. The coupling of the heaviest CP-even

Higgs boson to vector bosons is proportional to $\cos(\beta - \alpha)$ and in the decoupling limit $\cos(\beta - \alpha) \rightarrow 0$, the light CP-even Higgs boson is indistinguishable from a SM Higgs boson with the same mass. The range of $\cos(\beta - \alpha)$ and $\tan \beta$ explored is limited to the region where the assumption of a heavy narrow-width Higgs boson with negligible interference is valid. When calculating the limits at a given choice of $\cos(\beta - \alpha)$ and $\tan \beta$, the relative rate of ggF and VBF production in the fit is set to the prediction of the 2HDM for that parameter choice. The white regions in the exclusion plots indicate regions of parameter space which are not excluded by the present analysis.

For the LWA scenario, the interference effects among the heavy boson, the light Higgs boson at 125 GeV and the SM WW continuum background were studied and found to have negligible impact on the exclusion limits. The 95% CL upper limits are shown in Figure 9. The limits for signal widths of 5%, 10% and 15% are comparable with those from the NWA scenario for the VBF signals while for the ggF signals, the limits weaken slightly at high masses as the width increases. For the LWA 15% case, the upper exclusion limit ranges between 5.2 pb (1.3 pb) at $m_H = 200$ GeV and 0.02 pb (0.006 pb) at 4 (3) TeV for the ggF (VBF) signals.

Figure 10 shows the limits on the resonance production cross section times branching fraction $\sigma_X \times B(X \rightarrow WW)$ and $\sin \theta_H$ for a scalar GM signal with masses between 200 GeV and 1 TeV. At the observed limit, the width is narrower than the experimental resolution [46]. The current sensitivity is not sufficient to exclude the benchmark model with $\sin \theta_H = 0.4$.

Limits are derived in the mass range from 250 GeV to 5 TeV and from 300 GeV to 1 TeV for a qqA and VBF HVT signal, respectively, as shown in Figure 11. For the qqA production, signals below about 1.3 TeV are excluded at 95% CL. No limit can be set for the VBF production in the benchmark model that assumes a coupling strength to gauge bosons $g_V = 1$ and a coupling to fermions $c_F = 0$. The model has an intrinsic width much narrower than the detector resolution.

Figure 12 shows the limits on a $G_{KK} \rightarrow WW$ signal for two different couplings: $k/\bar{M}_{Pl} = 1$ and $k/\bar{M}_{Pl} = 0.5$, for masses between 200 GeV and 5 TeV, and for an ELM spin-2 VBF signal for masses between 200 GeV and 1 TeV. The observed limits exclude a KK graviton signal lighter than 1.1 TeV (750 GeV) with the higher (lower) coupling, while the current sensitivity is not sufficient to exclude the ELM spin-2 VBF signal.




Figure 7: Exclusion contours at 95% CL in the plane of $\tan\beta$ and $\cos(\beta - \alpha)$ for Type I (left) and Type II (right) 2HDM signals with three mass values of 200 GeV (top), 300 GeV (middle) and 500 GeV (bottom). The inner and outer bands show the $\pm 1\sigma$ and $\pm 2\sigma$ ranges around the expected limit and the hatched regions are excluded.




Figure 8: Exclusion contours at 95% CL in the plane of $\tan\beta$ and m_H for Type I (left) and Type II (right) 2HDM signals with $\cos(\beta - \alpha) = -0.1$. The inner and outer bands show the $\pm 1\sigma$ and $\pm 2\sigma$ ranges around the expected limit and the hatched regions are excluded. The other heavy Higgs boson states A and $H^\pm$ are assumed to have the same mass as H .




Figure 9: Upper limits at 95% CL on the Higgs boson production cross section times branching fraction $\sigma_H \times B(H \rightarrow WW)$ in the $\nu_e \nu_\mu$ channel, for a signal with a width of 15% of the mass (top) and the comparison of three different widths (bottom) for the ggF (left) and VBF (right) production. The inner and outer bands show the $\pm 1\sigma$ and $\pm 2\sigma$ ranges around the expected limit.




Figure 10: Upper limits at 95% CL on the resonance production cross section times branching fraction $\sigma_X \times B(X \rightarrow WW)$ (left) and on $\sin\theta_H$ (right) in the $e\nu\mu\nu$ channel, for a GM signal. The inner and outer bands show the $\pm 1\sigma$ and $\pm 2\sigma$ ranges around the expected limit. The full curves without dots correspond to the predicted theoretical cross section and the model parameter used in the benchmark model, respectively.




Figure 11: Upper limits at 95% CL on the resonance production cross section times branching fraction $\sigma_X \times B(X \rightarrow WW)$ in the $e\nu\mu\nu$ channel, for HVT qqA (left) and VBF (right) signals. The inner and outer bands show the $\pm 1\sigma$ and $\pm 2\sigma$ ranges around the expected limit. The full curves without dots correspond to the predicted theoretical cross sections.




Figure 12: Upper limits at 95% CL on the resonance production cross section times branching fraction $\sigma_X \times B(X \rightarrow WW)$ in the $e\nu\mu\nu$ channel, for a graviton signal with two different couplings of $k/\bar{M}_{Pl} = 1$ (left) and $k/\bar{M}_{Pl} = 0.5$ (right), and for an ELM spin-2 VBF signal (bottom). The inner and outer bands show the $\pm 1\sigma$ and $\pm 2\sigma$ ranges around the expected limit. The full curves without dots correspond to the predicted theoretical cross sections.

10 Conclusion

A search for heavy neutral resonances decaying into a WW boson pair in the $e\nu\mu\nu$ channel performed by the ATLAS Collaboration at the LHC is presented. The search uses proton–proton collision data collected at a centre-of-mass energy of 13 TeV corresponding to an integrated luminosity of 36.1 fb^{-1} . No significant excess of events beyond the Standard Model background prediction is found in the mass range between 200 GeV and up to 5 TeV. Upper limits are set on the product of the production cross section and the $X \rightarrow WW$ branching fraction in several scenarios: a high-mass Higgs boson with a narrow width or with intermediate widths (of 5%, 10% and 15% of the heavy Higgs boson mass), as well as other spin-0, spin-1, and spin-2 signals. For the narrow-width heavy Higgs boson signals, values above 6.4 pb at $m_H = 200 \text{ GeV}$ and above 0.008 pb at 4 TeV are excluded at 95% confidence level for the gluon–gluon fusion production mode. The corresponding values for the vector-boson fusion production modes are 1.3 pb and 0.006 pb at 200 GeV and 3 TeV, respectively. For the signals of the heavy vector triplet model A produced by quark–antiquark annihilation and of the Randall–Sundrum graviton model with $k/\bar{M}_{\text{Pl}} = 1$ and 0.5, mass values below 1.3 TeV, 1.1 TeV and 750 GeV are excluded, respectively.

Acknowledgements

We thank CERN for the very successful operation of the LHC, as well as the support staff from our institutions without whom ATLAS could not be operated efficiently.

We acknowledge the support of ANPCyT, Argentina; YerPhI, Armenia; ARC, Australia; BMWFW and FWF, Austria; ANAS, Azerbaijan; SSTC, Belarus; CNPq and FAPESP, Brazil; NSERC, NRC and CFI, Canada; CERN; CONICYT, Chile; CAS, MOST and NSFC, China; COLCIENCIAS, Colombia; MSMT CR, MPO CR and VSC CR, Czech Republic; DNRF and DNSRC, Denmark; IN2P3-CNRS, CEA-DRF/IRFU, France; SRNSF, Georgia; BMBF, HGF, and MPG, Germany; GSRT, Greece; RGC, Hong Kong SAR, China; ISF, I-CORE and Benoziyo Center, Israel; INFN, Italy; MEXT and JSPS, Japan; CNRST, Morocco; NWO, Netherlands; RCN, Norway; MNiSW and NCN, Poland; FCT, Portugal; MNE/IFA, Romania; MES of Russia and NRC KI, Russian Federation; JINR; MESTD, Serbia; MSSR, Slovakia; ARRS and MIZŠ, Slovenia; DST/NRF, South Africa; MINECO, Spain; SRC and Wallenberg Foundation, Sweden; SERI, SNSF and Cantons of Bern and Geneva, Switzerland; MOST, Taiwan; TAEK, Turkey; STFC, United Kingdom; DOE and NSF, United States of America. In addition, individual groups and members have received support from BCKDF, the Canada Council, CANARIE, CRC, Compute Canada, FQRNT, and the Ontario Innovation Trust, Canada; EPLANET, ERC, ERDF, FP7, Horizon 2020 and Marie Skłodowska-Curie Actions, European Union; Investissements d’Avenir Labex and Idex, ANR, Région Auvergne and Fondation Partager le Savoir, France; DFG and AvH Foundation, Germany; Herakleitos, Thales and Aristeia programmes co-financed by EU-ESF and the Greek NSRF; BSF, GIF and Minerva, Israel; BRF, Norway; CERCA Programme Generalitat de Catalunya, Generalitat Valenciana, Spain; the Royal Society and Leverhulme Trust, United Kingdom.

The crucial computing support from all WLCG partners is acknowledged gratefully, in particular from CERN, the ATLAS Tier-1 facilities at TRIUMF (Canada), NDGF (Denmark, Norway, Sweden), CC-IN2P3 (France), KIT/GridKA (Germany), INFN-CNAF (Italy), NL-T1 (Netherlands), PIC (Spain), ASGC (Taiwan), RAL (UK) and BNL (USA), the Tier-2 facilities worldwide and large non-WLCG resource providers. Major contributors of computing resources are listed in Ref. [112].

References

- [1] ATLAS Collaboration, *Measurements of the Higgs boson production and decay rates and coupling strengths using pp collision data at $\sqrt{s} = 7$ and 8 TeV in the ATLAS experiment*, [Eur. Phys. J. C **76** \(2016\) 6](#), arXiv: [1507.04548 \[hep-ex\]](#).
- [2] ATLAS Collaboration, *Study of the spin and parity of the Higgs boson in diboson decays with the ATLAS detector*, [Eur. Phys. J. C **75** \(2015\) 476](#), arXiv: [1506.05669 \[hep-ex\]](#).
- [3] CMS Collaboration, *Precise determination of the mass of the Higgs boson and tests of compatibility of its couplings with the standard model predictions using proton collisions at 7 and 8 TeV*, [Eur. Phys. J. C **75** \(2015\) 212](#), arXiv: [1412.8662 \[hep-ex\]](#).
- [4] ATLAS and CMS Collaborations, *Measurements of the Higgs boson production and decay rates and constraints on its couplings from a combined ATLAS and CMS analysis of the LHC pp collision data at $\sqrt{s} = 7$ and 8 TeV*, [JHEP **08** \(2016\) 045](#), arXiv: [1606.02266 \[hep-ex\]](#).
- [5] ATLAS Collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, [Phys. Lett. B **716** \(2012\) 1](#), arXiv: [1207.7214 \[hep-ex\]](#).
- [6] CMS Collaboration, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, [Phys. Lett. B **716** \(2012\) 30](#), arXiv: [1207.7235 \[hep-ex\]](#).
- [7] D. de Florian et al., *Handbook of LHC Higgs cross sections: 4. Deciphering the nature of the Higgs sector*, 2016, arXiv: [1610.07922 \[hep-ph\]](#).
- [8] I. P. Ivanov, *Building and testing models with extended Higgs sectors*, [Prog. Part. Nucl. Phys. **95** \(2017\) 160](#), arXiv: [1702.03776 \[hep-ph\]](#).
- [9] K. Agashe, R. Contino and A. Pomarol, *The minimal composite Higgs model*, [Nucl. Phys. B **719** \(2005\) 165](#), arXiv: [hep-ph/0412089](#).
- [10] G. Giudice et al., *The strongly-interacting light Higgs*, [JHEP **06** \(2007\) 045](#), arXiv: [hep-ph/0703164](#).
- [11] L. Randall and R. Sundrum, *A large mass hierarchy from a small extra dimension*, [Phys. Rev. Lett. **83** \(1999\) 3370](#), arXiv: [hep-ph/9905221](#).
- [12] K. Agashe et al., *Warped gravitons at the LHC and beyond*, [Phys. Rev. D **76** \(2007\) 036006](#), arXiv: [hep-ph/0701186](#).
- [13] O. Antipin, D. Atwood and A. Soni, *Search for RS gravitons via $W_L W_L$ decays*, [Phys. Lett. B **666** \(2008\) 155](#), arXiv: [0711.3175 \[hep-ph\]](#).
- [14] O. Antipin and A. Soni, *Towards establishing the spin of warped gravitons*, [JHEP **10** \(2008\) 018](#), arXiv: [0806.3427 \[hep-ph\]](#).
- [15] H. Georgi and M. Machacek, *Doubly charged Higgs bosons*, [Nucl. Phys. B **262** \(1985\) 463](#).
- [16] M. S. Chanowitz and M. Golden, *Higgs boson triplets with $M_W = M_Z \cos \theta_W$* , [Phys. Lett. B **165** \(1985\) 105](#).
- [17] J. de Blas, J. M. Lizana and M. Perez-Victoria, *Combining searches of Z' and W' bosons*, [JHEP **01** \(2013\) 166](#), arXiv: [1211.2229 \[hep-ph\]](#).
- [18] D. Pappadopulo et al., *Heavy vector triplets: bridging theory and data*, [JHEP **09** \(2014\) 060](#), arXiv: [1402.4431 \[hep-ph\]](#).

- [19] T. Han, J. D. Lykken and R.-J. Zhang, *On Kaluza-Klein states from large extra dimensions*, [Phys. Rev. D **59** \(1999\) 105006](#), arXiv: [hep-ph/9811350](#).
- [20] J. Frank, M. Rauch and D. Zeppenfeld, *Spin-2 resonances in vector-boson-fusion processes at next-to-leading order QCD*, [Phys. Rev. D **87** \(2013\) 055020](#), arXiv: [1211.3658 \[hep-ph\]](#).
- [21] ATLAS Collaboration, *Search for a high-mass Higgs boson decaying to a W boson pair in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, [JHEP **01** \(2016\) 032](#), arXiv: [1509.00389 \[hep-ex\]](#).
- [22] CMS Collaboration, *Search for a Higgs boson in the mass range from 145 to 1000 GeV decaying to a pair of W or Z bosons*, [JHEP **10** \(2015\) 144](#), arXiv: [1504.00936 \[hep-ex\]](#).
- [23] ATLAS Collaboration, *Search for new phenomena in the $WW \rightarrow \ell\nu\ell'\nu'$ final state in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector*, [Phys. Lett. B **718** \(2013\) 860](#), arXiv: [1208.2880 \[hep-ex\]](#).
- [24] CMS Collaboration, *Search for massive resonances in dijet systems containing jets tagged as W or Z boson decays in pp collisions at $\sqrt{s} = 8$ TeV*, [JHEP **08** \(2014\) 173](#), arXiv: [1405.1994 \[hep-ex\]](#).
- [25] CMS Collaboration, *Search for massive resonances decaying into pairs of boosted bosons in semi-leptonic final states at $\sqrt{s} = 8$ TeV*, [JHEP **08** \(2014\) 174](#), arXiv: [1405.3447 \[hep-ex\]](#).
- [26] ATLAS Collaboration, *Search for WZ resonances in the fully leptonic channel using pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, [Phys. Lett. B **737** \(2014\) 223](#), arXiv: [1406.4456 \[hep-ex\]](#).
- [27] CMS Collaboration, *Search for new resonances decaying via WZ to leptons in proton-proton collisions at $\sqrt{s} = 8$ TeV*, [Phys. Lett. B **740** \(2015\) 83](#), arXiv: [1407.3476 \[hep-ex\]](#).
- [28] ATLAS Collaboration, *Search for production of WW/WZ resonances decaying to a lepton, neutrino and jets in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, [Eur. Phys. J. C **75** \(2015\) 209](#), Erratum: [Eur. Phys. J. C **75** \(2015\) 370](#), arXiv: [1503.04677 \[hep-ex\]](#).
- [29] ATLAS Collaboration, *Search for high-mass diboson resonances with boson-tagged jets in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, [JHEP **12** \(2015\) 055](#), arXiv: [1506.00962 \[hep-ex\]](#).
- [30] ATLAS Collaboration, *Combination of searches for WW, WZ, and ZZ resonances in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, [Phys. Lett. B **755** \(2016\) 285](#), arXiv: [1512.05099 \[hep-ex\]](#).
- [31] ATLAS Collaboration, *Searches for heavy diboson resonances in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, [JHEP **09** \(2016\) 173](#), arXiv: [1606.04833 \[hep-ex\]](#).
- [32] CMS Collaboration, *Search for massive resonances decaying into WW, WZ or ZZ bosons in proton-proton collisions at $\sqrt{s} = 13$ TeV*, [JHEP **03** \(2017\) 162](#), arXiv: [1612.09159 \[hep-ex\]](#).
- [33] CMS Collaboration, *Combination of searches for heavy resonances decaying to WW, WZ, ZZ, WH, and ZH boson pairs in proton-proton collisions at $\sqrt{s} = 8$ and 13 TeV*, [Phys. Lett. B **774** \(2017\) 533](#), arXiv: [1705.09171 \[hep-ex\]](#).
- [34] ATLAS Collaboration, *Search for diboson resonances with boson-tagged jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, 2017, arXiv: [1708.04445 \[hep-ex\]](#).
- [35] CMS Collaboration, *Search for massive resonances decaying into WW, WZ, ZZ, qW, and qZ with dijet final states at $\sqrt{s} = 13$ TeV*, 2017, arXiv: [1708.05379 \[hep-ex\]](#).

- [36] ATLAS Collaboration, *Searches for heavy ZZ and ZW resonances in the $\ell\ell qq$ and $\nu\nu qq$ final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, 2017, arXiv: [1708.09638 \[hep-ex\]](#).
- [37] ATLAS Collaboration, *Constraints on new phenomena via Higgs boson couplings and invisible decays with the ATLAS detector*, *JHEP* **11** (2015) 206, arXiv: [1509.00672 \[hep-ex\]](#).
- [38] G. Branco et al., *Theory and phenomenology of two-Higgs-doublet models*, *Phys. Rept.* **516** (2012) 1, arXiv: [1106.0034 \[hep-ph\]](#).
- [39] R. V. Harlander, S. Liebler and H. Mantler, *SusHi: a program for the calculation of Higgs production in gluon fusion and bottom-quark annihilation in the Standard Model and the MSSM*, *Comput. Phys. Commun.* **184** (2013) 1605, arXiv: [1212.3249 \[hep-ph\]](#).
- [40] D. Eriksson, J. Rathsmann and O. Stal, *2HDMC: Two-Higgs-doublet model calculator*, *Comput. Phys. Commun.* **181** (2010) 189, Erratum: *Comput. Phys. Commun.* **181** (2010) 985, arXiv: [0902.0851 \[hep-ph\]](#).
- [41] C. Degrande et al., *Automatic predictions in the Georgi-Machacek model at next-to-leading order accuracy*, *Phys. Rev. D* **93** (2016) 035004, arXiv: [1512.01243 \[hep-ph\]](#).
- [42] S. Godfrey and K. Moats, *Exploring Higgs triplet models via vector boson scattering at the LHC*, *Phys. Rev. D* **81** (2010) 075026, arXiv: [1003.3033 \[hep-ph\]](#).
- [43] N. Arkani-Hamed et al., *The littlest Higgs*, *JHEP* **07** (2002) 034, arXiv: [hep-ph/0206021](#).
- [44] J. F. Gunion et al., *Higgs bosons in left-right symmetric models*, *Phys. Rev. D* **40** (1989) 1546.
- [45] K. Hartling, K. Kumar and H. E. Logan, *GMCALC: a calculator for the Georgi-Machacek model*, 2014, arXiv: [1412.7387 \[hep-ph\]](#).
- [46] M. Zaro and H. Logan, *Recommendations for the interpretation of LHC searches for H_5^0 , $H_5^\pm$, and $H_5^{\pm\pm}$ in vector boson fusion with decays to vector boson pairs*, LHC Higgs Cross Section Working Group, LHCHXSWG-2015-001, URL: <https://cds.cern.ch/record/2002500>.
- [47] ATLAS Collaboration, *The ATLAS experiment at the CERN large hadron collider*, *JINST* **3** (2008) S08003.
- [48] ATLAS Collaboration, *ATLAS insertable B-layer technical design report*, ATLAS-TDR-19, 2010, URL: <https://cds.cern.ch/record/1291633>, *ATLAS insertable B-layer technical design report addendum*, ATLAS-TDR-19-ADD-1, 2012, URL: <https://cds.cern.ch/record/1451888>.
- [49] ATLAS Collaboration, *Performance of the ATLAS trigger system in 2015*, *Eur. Phys. J. C* **77** (2017) 317, arXiv: [1611.09661 \[hep-ex\]](#).
- [50] P. Nason, *A new method for combining NLO QCD with shower Monte Carlo algorithms*, *JHEP* **11** (2004) 040, arXiv: [hep-ph/0409146](#).
- [51] S. Frixione, P. Nason and C. Oleari, *Matching NLO QCD computations with parton shower simulations: the POWHEG method*, *JHEP* **11** (2007) 070, arXiv: [0709.2092 \[hep-ph\]](#).
- [52] S. Alioli et al., *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, *JHEP* **06** (2010) 043, arXiv: [1002.2581 \[hep-ph\]](#).
- [53] S. Alioli et al., *NLO Higgs boson production via gluon fusion matched with shower in POWHEG*, *JHEP* **04** (2009) 002, arXiv: [0812.0578 \[hep-ph\]](#).

- [54] P. Nason and C. Oleari, *NLO Higgs boson production via vector-boson fusion matched with shower in POWHEG*, *JHEP* **02** (2010) 037, arXiv: [0911.5299 \[hep-ph\]](#).
- [55] H.-L. Lai et al., *New parton distributions for collider physics*, *Phys. Rev. D* **82** (2010) 074024, arXiv: [1007.2241 \[hep-ph\]](#).
- [56] T. Sjöstrand, S. Mrenna and P. Z. Skands, *A brief Introduction to PYTHIA 8.1*, *Comput. Phys. Commun.* **178** (2008) 852, arXiv: [0710.3820 \[hep-ph\]](#).
- [57] ATLAS Collaboration, *Measurement of the Z/γ^* boson transverse momentum distribution in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector*, *JHEP* **09** (2014) 145, arXiv: [1406.3660 \[hep-ex\]](#).
- [58] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP* **07** (2014) 079, arXiv: [1405.0301 \[hep-ph\]](#).
- [59] S. Carrazza, S. Forte and J. Rojo, *Parton distributions and event generators*, 2013, arXiv: [1311.5887 \[hep-ph\]](#).
- [60] ATLAS Collaboration, *ATLAS PYTHIA 8 tunes to 7 TeV data*, ATL-PHYS-PUB-2014-021, 2014, URL: <https://cds.cern.ch/record/1966419>.
- [61] R. Frederix and S. Frixione, *Merging meets matching in MC@NLO*, *JHEP* **12** (2012) 061, arXiv: [1209.6215 \[hep-ph\]](#).
- [62] J. Baglio et al., *Release note - VBFNLO 2.7.0*, 2014, arXiv: [1404.3940 \[hep-ph\]](#).
- [63] R. D. Ball et al., *Parton distributions for the LHC Run II*, *JHEP* **04** (2015) 040, arXiv: [1410.8849 \[hep-ph\]](#).
- [64] S. Alioli et al., *NLO single-top production matched with shower in POWHEG: s- and t-channel contributions*, *JHEP* **09** (2009) 111, Erratum: *JHEP* **02** (2010) 011, arXiv: [0907.4076 \[hep-ph\]](#).
- [65] E. Re, *Single-top Wt-channel production matched with parton showers using the POWHEG method*, *Eur. Phys. J. C* **71** (2011) 1547, arXiv: [1009.2450 \[hep-ph\]](#).
- [66] T. Sjöstrand, S. Mrenna and P. Z. Skands, *PYTHIA 6.4 physics and manual*, *JHEP* **05** (2006) 026, arXiv: [hep-ph/0603175](#).
- [67] P. Z. Skands, *Tuning Monte Carlo generators: The Perugia tunes*, *Phys. Rev. D* **82** (2010) 074018, arXiv: [1005.3457 \[hep-ph\]](#).
- [68] J. Pumplin et al., *New generation of parton distributions with uncertainties from global QCD analysis*, *JHEP* **07** (2002) 012, arXiv: [hep-ph/0201195](#).
- [69] S. Alioli, S.-O. Moch and P. Uwer, *Hadronic top-quark pair-production with one jet and parton showering*, *JHEP* **01** (2012) 137, arXiv: [1110.5251 \[hep-ph\]](#).
- [70] D. J. Lange, *The EVTGEN particle decay simulation package*, *Nucl. Instrum. Meth. A* **462** (2001) 152.
- [71] T. Gleisberg et al., *Event generation with SHERPA 1.1*, *JHEP* **02** (2009) 007, arXiv: [0811.4622 \[hep-ph\]](#).
- [72] T. Gleisberg and S. Höche, *Comix, a new matrix element generator*, *JHEP* **12** (2008) 039, arXiv: [0808.3674 \[hep-ph\]](#).
- [73] F. Cascioli, P. Maierhofer and S. Pozzorini, *Scattering amplitudes with open loops*, *Phys. Rev. Lett.* **108** (2012) 111601, arXiv: [1111.5206 \[hep-ph\]](#).

- [74] S. Schumann and F. Krauss, *A Parton shower algorithm based on Catani-Seymour dipole factorisation*, *JHEP* **03** (2008) 038, arXiv: [0709.1027 \[hep-ph\]](#).
- [75] S. Höche et al., *QCD matrix elements + parton showers: The NLO case*, *JHEP* **04** (2013) 027, arXiv: [1207.5030 \[hep-ph\]](#).
- [76] S. Agostinelli et al., *GEANT4: a simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250.
- [77] ATLAS Collaboration, *The ATLAS simulation infrastructure*, *Eur. Phys. J. C* **70** (2010) 823, arXiv: [1005.4568 \[physics.ins-det\]](#).
- [78] ATLAS Collaboration, *Electron efficiency measurements with the ATLAS detector using the 2015 LHC proton–proton collision data*, ATLAS-CONF-2016-024, 2016, URL: <https://cds.cern.ch/record/2157687>.
- [79] ATLAS Collaboration, *Measurement of the muon reconstruction performance of the ATLAS detector using 2011 and 2012 LHC proton–proton collision data*, *Eur. Phys. J. C* **74** (2014) 3130, arXiv: [1407.3935 \[hep-ex\]](#).
- [80] ATLAS Collaboration, *Muon reconstruction performance of the ATLAS detector in proton-proton collision data at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **76** (2016) 292, arXiv: [1603.05598 \[hep-ex\]](#).
- [81] K. Rehermann and B. Tweedie, *Efficient identification of boosted semileptonic top quarks at the LHC*, *JHEP* **03** (2011) 059, arXiv: [1007.2221 \[hep-ph\]](#).
- [82] M. Cacciari, G. P. Salam and G. Soyez, *The anti- k_t jet clustering algorithm*, *JHEP* **04** (2008) 063, arXiv: [0802.1189 \[hep-ph\]](#).
- [83] M. Cacciari, G. P. Salam and G. Soyez, *FastJet user manual*, *Eur. Phys. J. C* **72** (2012) 1896, arXiv: [1111.6097 \[hep-ph\]](#).
- [84] M. Cacciari and G. P. Salam, *Pileup subtraction using jet areas*, *Phys. Lett. B* **659** (2008) 119.
- [85] ATLAS Collaboration, *Jet energy scale measurements and their systematic uncertainties in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Rev. D* **96** (2017), arXiv: [1703.09665 \[hep-ex\]](#).
- [86] ATLAS Collaboration, *Tagging and suppression of pileup jets with the ATLAS detector*, ATLAS-CONF-2014-018, 2014, URL: <https://cds.cern.ch/record/1700870>.
- [87] ATLAS Collaboration, *Performance of b-jet identification in the ATLAS experiment*, *JINST* **11** (2016) P04008, arXiv: [1512.01094 \[hep-ex\]](#).
- [88] ATLAS Collaboration, *Optimisation of the ATLAS b-tagging performance for the 2016 LHC run*, ATL-PHYS-PUB-2016-012, 2016, URL: <https://cds.cern.ch/record/2160731>.
- [89] ATLAS Collaboration, *Performance of missing transverse momentum reconstruction with the ATLAS detector in the first proton-proton collisions at $\sqrt{s} = 13$ TeV*, ATL-PHYS-PUB-2015-027, 2015, URL: <https://cds.cern.ch/record/2037904>.
- [90] A. Hoecker et al., *TMVA: Toolkit for multivariate data analysis*, 2007, arXiv: [physics/0703039](#).
- [91] K. Melnikov and F. Petriello, *Electroweak gauge boson production at hadron colliders through $O(\alpha_s^2)$* , *Phys. Rev. D* **74** (2006) 114017, arXiv: [hep-ph/0609070](#).
- [92] M. Grazzini et al., *W^+W^- production at the LHC: fiducial cross sections and distributions in NNLO QCD*, *JHEP* **08** (2016) 140, arXiv: [1605.02716 \[hep-ph\]](#).
- [93] B. Biedermann et al., *Next-to-leading-order electroweak corrections to $pp \rightarrow W^+W^- \rightarrow 4$ leptons at the LHC*, *JHEP* **06** (2016) 065, arXiv: [1605.03419 \[hep-ph\]](#).

- [94] ATLAS Collaboration, *Observation and measurement of Higgs boson decays to WW^* with the ATLAS detector*, *Phys. Rev. D* **92** (2015) 012006, arXiv: [1412.2641 \[hep-ex\]](#).
- [95] ATLAS Collaboration, *Measurement of the inelastic proton-proton cross section at $\sqrt{s} = 13$ TeV with the ATLAS detector at the LHC*, *Phys. Rev. Lett.* **117** (2016) 182002, arXiv: [1606.02625 \[hep-ex\]](#).
- [96] ATLAS Collaboration, *Luminosity determination in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector at the LHC*, *Eur. Phys. J. C* **76** (2016) 653, arXiv: [1608.03953 \[hep-ex\]](#).
- [97] S. Dittmaier et al., *Handbook of LHC Higgs cross sections: 2. differential distributions*, 2012, arXiv: [1201.3084 \[hep-ph\]](#).
- [98] ATLAS Collaboration, *Measurement of the production cross-section of a single top quark in association with a W boson at 8 TeV with the ATLAS experiment*, *JHEP* **01** (2016) 064, arXiv: [1510.03752 \[hep-ex\]](#).
- [99] ATLAS Collaboration, *Measurement of the cross-section for producing a W boson in association with a single top quark in pp collisions at $\sqrt{s} = 13$ TeV with ATLAS*, (2016), arXiv: [1612.07231 \[hep-ex\]](#).
- [100] S. Dulat et al., *New parton distribution functions from a global analysis of quantum chromodynamics*, *Phys. Rev. D* **93** (2016) 033006, arXiv: [1506.07443 \[hep-ph\]](#).
- [101] L. A. Harland-Lang et al., *Parton distributions in the LHC era: MMHT 2014 PDFs*, *Eur. Phys. J. C* **75** (2015) 204, arXiv: [1412.3989 \[hep-ph\]](#).
- [102] K. Melnikov and M. Dowling, *Production of two Z -bosons in gluon fusion in the heavy top quark approximation*, *Phys. Lett. B* **744** (2015) 43, arXiv: [1503.01274 \[hep-ph\]](#).
- [103] F. Caola et al., *QCD corrections to W^+W^- production through gluon fusion*, *Phys. Lett. B* **754** (2016) 275, arXiv: [1511.08617 \[hep-ph\]](#).
- [104] M. Bonvini et al., *Signal-background interference effects for $gg \rightarrow H \rightarrow W^+W^-$ beyond leading order*, *Phys. Rev. D* **88** (2013) 034032, arXiv: [1304.3053 \[hep-ph\]](#).
- [105] C. S. Li et al., *Soft gluon resummation in the signal-background interference process of $gg(\rightarrow h^*) \rightarrow ZZ$* , *JHEP* **08** (2015) 065, arXiv: [1504.02388 \[hep-ph\]](#).
- [106] M. Bahr et al., *HERWIG++ physics and manual*, *Eur. Phys. J. C* **58** (2008) 639, arXiv: [0803.0883 \[hep-ph\]](#).
- [107] J. Bellm et al., *HERWIG++ 2.7 release note*, 2013, arXiv: [1310.6877 \[hep-ph\]](#).
- [108] I. Stewart and F. Tackmann, *Theory uncertainties for Higgs mass and other searches using jet bins*, *Phys. Rev. D* **85** (2012) 034011, arXiv: [1107.2117 \[hep-ph\]](#).
- [109] ATLAS Collaboration, *Combined search for the Standard Model Higgs boson in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector*, *Phys. Rev. D* **86** (2012) 032003, arXiv: [1207.0319 \[hep-ex\]](#).
- [110] A.L. Read, *Presentation of search results: the CL_s technique*, *J. Phys. G* **28** (2002) 2693.
- [111] G. Cowan et al., *Asymptotic formulae for likelihood-based tests of new physics*, *Eur. Phys. J. C* **71** (2011) 1554, Erratum: *Eur. Phys. J. C* **73** (2013) 2501, arXiv: [1007.1727 \[physics.data-an\]](#).
- [112] ATLAS Collaboration, *ATLAS computing acknowledgements 2016–2017*, ATL-GEN-PUB-2016-002, URL: <https://cds.cern.ch/record/2202407>.

The ATLAS Collaboration

M. Aaboud^{137d}, G. Aad⁸⁸, B. Abbott¹¹⁵, O. Abidinov^{12,*}, B. Abeloos¹¹⁹, S.H. Abidi¹⁶¹, O.S. AbouZeid¹³⁹, N.L. Abraham¹⁵¹, H. Abramowicz¹⁵⁵, H. Abreu¹⁵⁴, R. Abreu¹¹⁸, Y. Abulaiti^{148a,148b}, B.S. Acharya^{167a,167b,a}, S. Adachi¹⁵⁷, L. Adamczyk^{41a}, J. Adelman¹¹⁰, M. Adersberger¹⁰², T. Adye¹³³, A.A. Affolder¹³⁹, Y. Afik¹⁵⁴, C. Agheorghiesei^{28c}, J.A. Aguilar-Saavedra^{128a,128f}, S.P. Ahlen²⁴, F. Ahmadov^{68,b}, G. Aielli^{135a,135b}, S. Akatsuka⁷¹, H. Akerstedt^{148a,148b}, T.P.A. Åkesson⁸⁴, E. Akilli⁵², A.V. Akimov⁹⁸, G.L. Alberghi^{22a,22b}, J. Albert¹⁷², P. Albicocco⁵⁰, M.J. Alconada Verzini⁷⁴, S.C. Alderweireldt¹⁰⁸, M. Aleksa³², I.N. Aleksandrov⁶⁸, C. Alexa^{28b}, G. Alexander¹⁵⁵, T. Alexopoulos¹⁰, M. Alhroob¹¹⁵, B. Ali¹³⁰, M. Aliev^{76a,76b}, G. Alimonti^{94a}, J. Alison³³, S.P. Alkire³⁸, B.M.M. Allbrooke¹⁵¹, B.W. Allen¹¹⁸, P.P. Allport¹⁹, A. Aloisio^{106a,106b}, A. Alonso³⁹, F. Alonso⁷⁴, C. Alpigiani¹⁴⁰, A.A. Alshehri⁵⁶, M.I. Alstady⁸⁸, B. Alvarez Gonzalez³², D. Álvarez Piqueras¹⁷⁰, M.G. Alvigi^{106a,106b}, B.T. Amadio¹⁶, Y. Amaral Coutinho^{26a}, C. Amelung²⁵, D. Amidei⁹², S.P. Amor Dos Santos^{128a,128c}, S. Amoroso³², C. Anastopoulos¹⁴¹, L.S. Ancu⁵², N. Andari¹⁹, T. Andeen¹¹, C.F. Anders^{60b}, J.K. Anders⁷⁷, K.J. Anderson³³, A. Andreazza^{94a,94b}, V. Andrei^{60a}, S. Angelidakis³⁷, I. Angelozzi¹⁰⁹, A. Angerami³⁸, A.V. Anisenkov^{111,c}, N. Anjos¹³, A. Annovi^{126a}, C. Antel^{60a}, M. Antonelli⁵⁰, A. Antonov^{100,*}, D.J. Antrim¹⁶⁶, F. Anulli^{134a}, M. Aoki⁶⁹, L. Aperio Bella³², G. Arabidze⁹³, Y. Arai⁶⁹, J.P. Araque^{128a}, V. Araujo Ferraz^{26a}, A.T.H. Arce⁴⁸, R.E. Ardell⁸⁰, F.A. Arduh⁷⁴, J-F. Arguin⁹⁷, S. Argyropoulos⁶⁶, M. Arik^{20a}, A.J. Armbruster³², L.J. Armitage⁷⁹, O. Arnaez¹⁶¹, H. Arnold⁵¹, M. Arratia³⁰, O. Arslan²³, A. Artamonov^{99,*}, G. Artoni¹²², S. Artz⁸⁶, S. Asai¹⁵⁷, N. Asbah⁴⁵, A. Ashkenazi¹⁵⁵, L. Asquith¹⁵¹, K. Assamagan²⁷, R. Astalos^{146a}, M. Atkinson¹⁶⁹, N.B. Atlay¹⁴³, K. Augsten¹³⁰, G. Avolio³², B. Axen¹⁶, M.K. Ayoub^{35a}, G. Azuelos^{97,d}, A.E. Baas^{60a}, M.J. Baca¹⁹, H. Bachacou¹³⁸, K. Bachas^{76a,76b}, M. Backes¹²², P. Bagnaia^{134a,134b}, M. Bahmani⁴², H. Bahrasemani¹⁴⁴, J.T. Baines¹³³, M. Bajic³⁹, O.K. Baker¹⁷⁹, P.J. Bakker¹⁰⁹, E.M. Baldin^{111,c}, P. Balek¹⁷⁵, F. Balli¹³⁸, W.K. Balunas¹²⁴, E. Banas⁴², A. Bandyopadhyay²³, Sw. Banerjee^{176,e}, A.A.E. Bannoura¹⁷⁸, L. Barak¹⁵⁵, E.L. Barberio⁹¹, D. Barberis^{53a,53b}, M. Barbero⁸⁸, T. Barillari¹⁰³, M-S Barisits⁶⁵, J.T. Barkeloo¹¹⁸, T. Barklow¹⁴⁵, N. Barlow³⁰, S.L. Barnes^{36c}, B.M. Barnett¹³³, R.M. Barnett¹⁶, Z. Barnovska-Blenessy^{36a}, A. Baroncelli^{136a}, G. Barone²⁵, A.J. Barr¹²², L. Barranco Navarro¹⁷⁰, F. Barreiro⁸⁵, J. Barreiro Guimarães da Costa^{35a}, R. Bartoldus¹⁴⁵, A.E. Barton⁷⁵, P. Bartos^{146a}, A. Basalaev¹²⁵, A. Bassalat^{119,f}, R.L. Bates⁵⁶, S.J. Batista¹⁶¹, J.R. Batley³⁰, M. Battaglia¹³⁹, M. Bause^{134a,134b}, F. Bauer¹³⁸, K.T. Bauer¹⁶⁶, H.S. Bawa^{145,g}, J.B. Beacham¹¹³, M.D. Beattie⁷⁵, T. Beau⁸³, P.H. Beauchemin¹⁶⁵, P. Bechtel²³, H.P. Beck^{18,h}, H.C. Beck⁵⁷, K. Becker¹²², M. Becker⁸⁶, C. Becot¹¹², A.J. Beddall^{20e}, A. Beddall^{20b}, V.A. Bednyakov⁶⁸, M. Bedognetti¹⁰⁹, C.P. Bee¹⁵⁰, T.A. Beermann³², M. Begalli^{26a}, M. Begel²⁷, J.K. Behr⁴⁵, A.S. Bell⁸¹, G. Bella¹⁵⁵, L. Bellagamba^{22a}, A. Bellerive³¹, M. Bellomo¹⁵⁴, K. Belotskiy¹⁰⁰, O. Beltramello³², N.L. Belyaev¹⁰⁰, O. Benary^{155,*}, D. Benckekroun^{137a}, M. Bender¹⁰², N. Benekos¹⁰, Y. Benhammou¹⁵⁵, E. Benhar Noccioli¹⁷⁹, J. Benitez⁶⁶, D.P. Benjamin⁴⁸, M. Benoit⁵², J.R. Bensinger²⁵, S. Bentvelsen¹⁰⁹, L. Beresford¹²², M. Beretta⁵⁰, D. Berge¹⁰⁹, E. Bergeaas Kuutmann¹⁶⁸, N. Berger⁵, L.J. Bergsten²⁵, J. Beringer¹⁶, S. Berlendis⁵⁸, N.R. Bernard⁸⁹, G. Bernardi⁸³, C. Bernius¹⁴⁵, F.U. Bernlochner²³, T. Berry⁸⁰, P. Berta⁸⁶, C. Bertella^{35a}, G. Bertoli^{148a,148b}, I.A. Bertram⁷⁵, C. Bertsche⁴⁵, G.J. Besjes³⁹, O. Bessidskaia Bylund^{148a,148b}, M. Bessner⁴⁵, N. Besson¹³⁸, A. Bethani⁸⁷, S. Bethke¹⁰³, A. Betti²³, A.J. Bevan⁷⁹, J. Beyer¹⁰³, R.M. Bianchi¹²⁷, O. Biebel¹⁰², D. Biedermann¹⁷, R. Bielski⁸⁷, K. Bierwagen⁸⁶, N.V. Biesuz^{126a,126b}, M. Biglietti^{136a}, T.R.V. Billoud⁹⁷, H. Bilokon⁵⁰, M. Bindi⁵⁷, A. Bingul^{20b}, C. Bini^{134a,134b}, S. Biondi^{22a,22b}, T. Bisanz⁵⁷, C. Bittrich⁴⁷, D.M. Bjergaard⁴⁸, J.E. Black¹⁴⁵, K.M. Black²⁴, R.E. Blair⁶, T. Blazek^{146a}, I. Bloch⁴⁵, C. Blocker²⁵, A. Blue⁵⁶, U. Blumenschein⁷⁹, Dr. Blunier^{34a}, G.J. Bobbink¹⁰⁹, V.S. Bobrovnikov^{111,c}, S.S. Bocchetta⁸⁴,

A. Bocci⁴⁸, C. Bock¹⁰², M. Boehler⁵¹, D. Boerner¹⁷⁸, D. Bogavac¹⁰², A.G. Bogdanchikov¹¹¹,
 C. Bohm^{148a}, V. Boisvert⁸⁰, P. Bokan^{168,i}, T. Bold^{41a}, A.S. Boldyrev¹⁰¹, A.E. Bolz^{60b}, M. Bomben⁸³,
 M. Bona⁷⁹, M. Boonekamp¹³⁸, A. Borisov¹³², G. Borissov⁷⁵, J. Bortfeldt³², D. Bortoletto¹²²,
 V. Bortolotto^{62a}, D. Boscherini^{22a}, M. Bosman¹³, J.D. Bossio Sola²⁹, J. Boudreau¹²⁷,
 E.V. Bouhova-Thacker⁷⁵, D. Boumediene³⁷, C. Bourdarios¹¹⁹, S.K. Boutle⁵⁶, A. Boveia¹¹³, J. Boyd³²,
 I.R. Boyko⁶⁸, A.J. Bozson⁸⁰, J. Bracinik¹⁹, A. Brandt⁸, G. Brandt⁵⁷, O. Brandt^{60a}, F. Braren⁴⁵,
 U. Bratzler¹⁵⁸, B. Brau⁸⁹, J.E. Brau¹¹⁸, W.D. Breaden Madden⁵⁶, K. Brendlinger⁴⁵, A.J. Brennan⁹¹,
 L. Brenner¹⁰⁹, R. Brenner¹⁶⁸, S. Bressler¹⁷⁵, D.L. Briglin¹⁹, T.M. Bristow⁴⁹, D. Britton⁵⁶, D. Britzger⁴⁵,
 F.M. Brochu³⁰, I. Brock²³, R. Brock⁹³, G. Brooijmans³⁸, T. Brooks⁸⁰, W.K. Brooks^{34b}, E. Brost¹¹⁰,
 J.H. Broughton¹⁹, P.A. Bruckman de Renstrom⁴², D. Bruncko^{146b}, A. Bruni^{22a}, G. Bruni^{22a},
 L.S. Bruni¹⁰⁹, S. Bruno^{135a,135b}, B.H. Brunt³⁰, M. Bruschi^{22a}, N. Bruscino¹²⁷, P. Bryant³³,
 L. Bryngemark⁴⁵, T. Buanes¹⁵, Q. Buat¹⁴⁴, P. Buchholz¹⁴³, A.G. Buckley⁵⁶, I.A. Budagov⁶⁸,
 F. Buehrer⁵¹, M.K. Bugge¹²¹, O. Bulekov¹⁰⁰, D. Bullock⁸, T.J. Burch¹¹⁰, S. Burdin⁷⁷, C.D. Burgard¹⁰⁹,
 A.M. Burger⁵, B. Burghgrave¹¹⁰, K. Burka⁴², S. Burke¹³³, I. Burmeister⁴⁶, J.T.P. Burr¹²², D. Büscher⁵¹,
 V. Büscher⁸⁶, P. Bussey⁵⁶, J.M. Butler²⁴, C.M. Buttar⁵⁶, J.M. Butterworth⁸¹, P. Butti³², W. Buttinger²⁷,
 A. Buzatu¹⁵³, A.R. Buzykaev^{111,c}, Changqiao C.-Q.^{36a}, S. Cabrera Urbán¹⁷⁰, D. Caforio¹³⁰, H. Cai¹⁶⁹,
 V.M. Cairo^{40a,40b}, O. Cakir^{4a}, N. Calace⁵², P. Calafiura¹⁶, A. Calandri⁸⁸, G. Calderini⁸³, P. Calfayan⁶⁴,
 G. Callea^{40a,40b}, L.P. Caloba^{26a}, S. Calvente Lopez⁸⁵, D. Calvet³⁷, S. Calvet³⁷, T.P. Calvet⁸⁸,
 R. Camacho Toro³³, S. Camarda³², P. Camarri^{135a,135b}, D. Cameron¹²¹, R. Caminal Armadans¹⁶⁹,
 C. Camincher⁵⁸, S. Campana³², M. Campanelli⁸¹, A. Camplani^{94a,94b}, A. Campoverde¹⁴³,
 V. Canale^{106a,106b}, M. Cano Bret^{36c}, J. Cantero¹¹⁶, T. Cao¹⁵⁵, M.D.M. Capeans Garrido³², I. Caprini^{28b},
 M. Caprini^{28b}, M. Capua^{40a,40b}, R.M. Carbone³⁸, R. Cardarelli^{135a}, F. Cardillo⁵¹, I. Carli¹³¹, T. Carli³²,
 G. Carlino^{106a}, B.T. Carlson¹²⁷, L. Carminati^{94a,94b}, R.M.D. Carney^{148a,148b}, S. Caron¹⁰⁸, E. Carquin^{34b},
 S. Carrá^{94a,94b}, G.D. Carrillo-Montoya³², D. Casadei¹⁹, M.P. Casado^{13,j}, A.F. Casha¹⁶¹, M. Casolino¹³,
 D.W. Casper¹⁶⁶, R. Castelijin¹⁰⁹, V. Castillo Gimenez¹⁷⁰, N.F. Castro^{128a,k}, A. Catinaccio³²,
 J.R. Catmore¹²¹, A. Cattai³², J. Caudron²³, V. Cavaliere¹⁶⁹, E. Cavallaro¹³, D. Cavalli^{94a},
 M. Cavalli-Sforza¹³, V. Cavasinni^{126a,126b}, E. Celebi^{20d}, F. Ceradini^{136a,136b}, L. Cerda Alberich¹⁷⁰,
 A.S. Cerqueira^{26b}, A. Cerri¹⁵¹, L. Cerrito^{135a,135b}, F. Cerutti¹⁶, A. Cervelli^{22a,22b}, S.A. Cetin^{20d},
 A. Chafaq^{137a}, D. Chakraborty¹¹⁰, S.K. Chan⁵⁹, W.S. Chan¹⁰⁹, Y.L. Chan^{62a}, P. Chang¹⁶⁹,
 J.D. Chapman³⁰, D.G. Charlton¹⁹, C.C. Chau³¹, C.A. Chavez Barajas¹⁵¹, S. Che¹¹³,
 S. Cheatham^{167a,167c}, A. Chegwidan⁹³, S. Chekanov⁶, S.V. Chekulaev^{163a}, G.A. Chelkov^{68,l},
 M.A. Chelstowska³², C. Chen^{36a}, C. Chen⁶⁷, H. Chen²⁷, J. Chen^{36a}, S. Chen^{35b}, S. Chen¹⁵⁷,
 X. Chen^{35c,m}, Y. Chen⁷⁰, H.C. Cheng⁹², H.J. Cheng^{35a,35d}, A. Cheplakov⁶⁸, E. Cheremushkina¹³²,
 R. Cherkaoui El Moursli^{137e}, E. Cheu⁷, K. Cheung⁶³, L. Chevalier¹³⁸, V. Chiarella⁵⁰, G. Chiarelli^{126a},
 G. Chiodini^{76a}, A.S. Chisholm³², A. Chitan^{28b}, Y.H. Chiu¹⁷², M.V. Chizhov⁶⁸, K. Choi⁶⁴,
 A.R. Chomont³⁷, S. Chouridou¹⁵⁶, Y.S. Chow^{62a}, V. Christodoulou⁸¹, M.C. Chu^{62a}, J. Chudoba¹²⁹,
 A.J. Chuinard⁹⁰, J.J. Chwastowski⁴², L. Chytka¹¹⁷, A.K. Ciftci^{4a}, D. Cinca⁴⁶, V. Cindro⁷⁸, I.A. Cioara²³,
 A. Ciocio¹⁶, F. Ciotto^{106a,106b}, Z.H. Citron¹⁷⁵, M. Citterio^{94a}, M. Ciubancan^{28b}, A. Clark⁵²,
 M.R. Clark³⁸, P.J. Clark⁴⁹, R.N. Clarke¹⁶, C. Clement^{148a,148b}, Y. Coadou⁸⁸, M. Cobal^{167a,167c},
 A. Coccaro⁵², J. Cochran⁶⁷, L. Colasurdo¹⁰⁸, B. Cole³⁸, A.P. Colijn¹⁰⁹, J. Collot⁵⁸, T. Colombo¹⁶⁶,
 P. Conde Muiño^{128a,128b}, E. Coniavitis⁵¹, S.H. Connell^{147b}, I.A. Connelly⁸⁷, S. Constantinescu^{28b},
 G. Conti³², F. Conventi^{106a,n}, M. Cooke¹⁶, A.M. Cooper-Sarkar¹²², F. Cormier¹⁷¹, K.J.R. Cormier¹⁶¹,
 M. Corradi^{134a,134b}, E.E. Corrigan⁸⁴, F. Corriveau^{90,o}, A. Cortes-Gonzalez³², G. Costa^{94a}, M.J. Costa¹⁷⁰,
 D. Costanzo¹⁴¹, G. Cottin³⁰, G. Cowan⁸⁰, B.E. Cox⁸⁷, K. Cranmer¹¹², S.J. Crawley⁵⁶, R.A. Creager¹²⁴,
 G. Cree³¹, S. Crépe-Renaudin⁵⁸, F. Crescioli⁸³, W.A. Cribbs^{148a,148b}, M. Cristinziani²³, V. Croft¹¹²,
 G. Crosetti^{40a,40b}, A. Cueto⁸⁵, T. Cuhadar Donszelmann¹⁴¹, A.R. Cukierman¹⁴⁵, J. Cummings¹⁷⁹,
 M. Curatolo⁵⁰, J. Cúth⁸⁶, S. Czekierda⁴², P. Czodrowski³², G. D'amen^{22a,22b}, S. D'Auria⁵⁶,

L. D'era⁸³, M. D'Onofrio⁷⁷, M.J. Da Cunha Sargedas De Sousa^{128a,128b}, C. Da Via⁸⁷, W. Dabrowski^{41a}, T. Dado^{146a}, T. Dai⁹², O. Dale¹⁵, F. Dallaire⁹⁷, C. Dallapiccola⁸⁹, M. Dam³⁹, J.R. Dandoy¹²⁴, M.F. Daneri²⁹, N.P. Dang¹⁷⁶, A.C. Daniells¹⁹, N.S. Dann⁸⁷, M. Danninger¹⁷¹, M. Dano Hoffmann¹³⁸, V. Dao¹⁵⁰, G. Darbo^{53a}, S. Darmora⁸, J. Dassoulas³, A. Dattagupta¹¹⁸, T. Daubney⁴⁵, W. Davey²³, C. David⁴⁵, T. Davidek¹³¹, D.R. Davis⁴⁸, P. Davison⁸¹, E. Dawe⁹¹, I. Dawson¹⁴¹, K. De⁸, R. de Asmundis^{106a}, A. De Benedetti¹¹⁵, S. De Castro^{22a,22b}, S. De Cecco⁸³, N. De Groot¹⁰⁸, P. de Jong¹⁰⁹, H. De la Torre⁹³, F. De Lorenzi⁶⁷, A. De Maria⁵⁷, D. De Pedis^{134a}, A. De Salvo^{134a}, U. De Sanctis^{135a,135b}, A. De Santo¹⁵¹, K. De Vasconcelos Corga⁸⁸, J.B. De Vivie De Regie¹¹⁹, R. Debbe²⁷, C. Debenedetti¹³⁹, D.V. Dedovich⁶⁸, N. Dehghanian³, I. Deigaard¹⁰⁹, M. Del Gaudio^{40a,40b}, J. Del Peso⁸⁵, D. Delgove¹¹⁹, F. Deliot¹³⁸, C.M. Delitzsch⁷, A. Dell'Acqua³², L. Dell'Asta²⁴, M. Dell'Orso^{126a,126b}, M. Della Pietra^{106a,106b}, D. della Volpe⁵², M. Delmastro⁵, C. Delporte¹¹⁹, P.A. Delsart⁵⁸, D.A. DeMarco¹⁶¹, S. Demers¹⁷⁹, M. Demichev⁶⁸, A. Demilly⁸³, S.P. Denisov¹³², D. Denysiuk¹³⁸, D. Derendarz⁴², J.E. Derkaoui^{137d}, F. Derue⁸³, P. Dervan⁷⁷, K. Desch²³, C. Deterre⁴⁵, K. Dette¹⁶¹, M.R. Devesa²⁹, P.O. Deviveiros³², A. Dewhurst¹³³, S. Dhaliwal²⁵, F.A. Di Bello⁵², A. Di Ciaccio^{135a,135b}, L. Di Ciaccio⁵, W.K. Di Clemente¹²⁴, C. Di Donato^{106a,106b}, A. Di Girolamo³², B. Di Girolamo³², B. Di Micco^{136a,136b}, R. Di Nardo³², K.F. Di Petrillo⁵⁹, A. Di Simone⁵¹, R. Di Sipio¹⁶¹, D. Di Valentino³¹, C. Diaconu⁸⁸, M. Diamond¹⁶¹, F.A. Dias³⁹, M.A. Diaz^{34a}, J. Dickinson¹⁶, E.B. Diehl⁹², J. Dietrich¹⁷, S. Díez Cornell⁴⁵, A. Dimitrievska¹⁴, J. Dingfelder²³, P. Dita^{28b}, S. Dita^{28b}, F. Dittus³², F. Djama⁸⁸, T. Djobava^{54b}, J.I. Djuvsland^{60a}, M.A.B. do Vale^{26c}, M. Dobre^{28b}, D. Dodsworth²⁵, C. Doglioni⁸⁴, J. Dolejsi¹³¹, Z. Dolezal¹³¹, M. Donadelli^{26d}, S. Donati^{126a,126b}, J. Donini³⁷, J. Dopke¹³³, A. Doria^{106a}, M.T. Dova⁷⁴, A.T. Doyle⁵⁶, E. Drechsler⁵⁷, M. Dris¹⁰, Y. Du^{36b}, J. Duarte-Campderros¹⁵⁵, F. Dubinin⁹⁸, A. Dubreuil⁵², E. Duchovni¹⁷⁵, G. Duckeck¹⁰², A. Ducourthial⁸³, O.A. Ducu^{97,p}, D. Duda¹⁰⁹, A. Dudarev³², A.Ch. Dudder⁸⁶, E.M. Duffield¹⁶, L. Duflot¹¹⁹, M. Dührssen³², C. Dulsen¹⁷⁸, M. Dumancic¹⁷⁵, A.E. Dumitriu^{28b}, A.K. Duncan⁵⁶, M. Dunford^{60a}, A. Duperrin⁸⁸, H. Duran Yildiz^{4a}, M. Düren⁵⁵, A. Durglishvili^{54b}, D. Duschinger⁴⁷, B. Dutta⁴⁵, D. Duvnjak¹, M. Dyndal⁴⁵, B.S. Dziedzic⁴², C. Eckardt⁴⁵, K.M. Ecker¹⁰³, R.C. Edgar⁹², T. Eifert³², G. Eigen¹⁵, K. Einsweiler¹⁶, T. Ekelof¹⁶⁸, M. El Kacimi^{137c}, R. El Kosseifi⁸⁸, V. Ellajosyula⁸⁸, M. Ellert¹⁶⁸, S. Elles⁵, F. Ellinghaus¹⁷⁸, A.A. Elliot¹⁷², N. Ellis³², J. Elmsheuser²⁷, M. Elsing³², D. Emelianov¹³³, Y. Enari¹⁵⁷, J.S. Ennis¹⁷³, M.B. Epland⁴⁸, J. Erdmann⁴⁶, A. Ereditato¹⁸, M. Ernst²⁷, S. Errede¹⁶⁹, M. Escalier¹¹⁹, C. Escobar¹⁷⁰, B. Esposito⁵⁰, O. Estrada Pastor¹⁷⁰, A.I. Etienne¹³⁸, E. Etzion¹⁵⁵, H. Evans⁶⁴, A. Ezhilov¹²⁵, M. Ezzi^{137e}, F. Fabbri^{22a,22b}, L. Fabbri^{22a,22b}, V. Fabiani¹⁰⁸, G. Facini⁸¹, R.M. Fakhruudinov¹³², S. Falciano^{134a}, R.J. Falla⁸¹, J. Faltova³², Y. Fang^{35a}, M. Fanti^{94a,94b}, A. Farbin⁸, A. Farilla^{136a}, E.M. Farina^{123a,123b}, T. Farooque⁹³, S. Farrell¹⁶, S.M. Farrington¹⁷³, P. Farthouat³², F. Fassi^{137e}, P. Fassnacht³², D. Fassouliotis⁹, M. Faucci Giannelli⁴⁹, A. Favareto^{53a,53b}, W.J. Fawcett¹²², L. Fayard¹¹⁹, O.L. Fedin^{125,q}, W. Fedorko¹⁷¹, S. Feigl¹²¹, L. Feligioni⁸⁸, C. Feng^{36b}, E.J. Feng³², M. Feng⁴⁸, M.J. Fenton⁵⁶, A.B. Fenyuk¹³², L. Feremenga⁸, P. Fernandez Martinez¹⁷⁰, J. Ferrando⁴⁵, A. Ferrari¹⁶⁸, P. Ferrari¹⁰⁹, R. Ferrari^{123a}, D.E. Ferreira de Lima^{60b}, A. Ferrer¹⁷⁰, D. Ferrere⁵², C. Ferretti⁹², F. Fiedler⁸⁶, A. Filipčič⁷⁸, M. Filipuzzi⁴⁵, F. Filthaut¹⁰⁸, M. Fincke-Keeler¹⁷², K.D. Finelli²⁴, M.C.N. Fiolhais^{128a,128c,r}, L. Fiorini¹⁷⁰, A. Fischer², C. Fischer¹³, J. Fischer¹⁷⁸, W.C. Fisher⁹³, N. Flaschel⁴⁵, I. Fleck¹⁴³, P. Fleischmann⁹², R.R.M. Fletcher¹²⁴, T. Flick¹⁷⁸, B.M. Flierl¹⁰², L.R. Flores Castillo^{62a}, M.J. Flowerdew¹⁰³, G.T. Forcolin⁸⁷, A. Formica¹³⁸, F.A. Förster¹³, A. Forti⁸⁷, A.G. Foster¹⁹, D. Fournier¹¹⁹, H. Fox⁷⁵, S. Fracchia¹⁴¹, P. Francavilla^{126a,126b}, M. Franchini^{22a,22b}, S. Franchino^{60a}, D. Francis³², L. Franconi¹²¹, M. Franklin⁵⁹, M. Frate¹⁶⁶, M. Fraternali^{123a,123b}, D. Freeborn⁸¹, S.M. Fressard-Batraneanu³², B. Freund⁹⁷, W.S. Freund^{26a}, D. Froidevaux³², J.A. Frost¹²², C. Fukunaga¹⁵⁸, T. Fusayasu¹⁰⁴, J. Fuster¹⁷⁰, O. Gabizon¹⁵⁴, A. Gabrielli^{22a,22b}, A. Gabrielli¹⁶, G.P. Gach^{41a}, S. Gadatsch³², S. Gadomski⁸⁰, G. Gagliardi^{53a,53b},

L.G. Gagnon⁹⁷, C. Galea¹⁰⁸, B. Galhardo^{128a,128c}, E.J. Gallas¹²², B.J. Gallop¹³³, P. Gallus¹³⁰, G. Galster³⁹, K.K. Gan¹¹³, S. Ganguly³⁷, Y. Gao⁷⁷, Y.S. Gao^{145,g}, F.M. Garay Walls^{34a}, C. García¹⁷⁰, J.E. García Navarro¹⁷⁰, J.A. García Pascual^{35a}, M. Garcia-Sciveres¹⁶, R.W. Gardner³³, N. Garelli¹⁴⁵, V. Garonne¹²¹, A. Gascon Bravo⁴⁵, K. Gasnikova⁴⁵, C. Gatti⁵⁰, A. Gaudiello^{53a,53b}, G. Gaudio^{123a}, I.L. Gavrilenko⁹⁸, C. Gay¹⁷¹, G. Gaycken²³, E.N. Gazis¹⁰, C.N.P. Gee¹³³, J. Geisen⁵⁷, M. Geisen⁸⁶, M.P. Geisler^{60a}, K. Gellerstedt^{148a,148b}, C. Gemme^{53a}, M.H. Genest⁵⁸, C. Geng⁹², S. Gentile^{134a,134b}, C. Gentsos¹⁵⁶, S. George⁸⁰, D. Gerbaudo¹³, G. Geßner⁴⁶, S. Ghasemi¹⁴³, M. Ghneimat²³, B. Giacobbe^{22a}, S. Giagu^{134a,134b}, N. Giangiacomi^{22a,22b}, P. Giannetti^{126a}, S.M. Gibson⁸⁰, M. Gignac¹⁷¹, M. Gilchriese¹⁶, D. Gillberg³¹, G. Gilles¹⁷⁸, D.M. Gingrich^{3,d}, M.P. Giordani^{167a,167c}, F.M. Giorgi^{22a}, P.F. Giraud¹³⁸, P. Giromini⁵⁹, G. Giugliarelli^{167a,167c}, D. Giugni^{94a}, F. Giuli¹²², C. Giuliani¹⁰³, M. Giulini^{60b}, B.K. Gjølsten¹²¹, S. Gkaitatzis¹⁵⁶, I. Gkialas^{9,s}, E.L. Gkougkousis¹³, P. Gkoutoumis¹⁰, L.K. Gladilin¹⁰¹, C. Glasman⁸⁵, J. Glatzer¹³, P.C.F. Glaysheer⁴⁵, A. Glazov⁴⁵, M. Goblirsch-Kolb²⁵, J. Godlewski⁴², S. Goldfarb⁹¹, T. Golling⁵², D. Golubkov¹³², A. Gomes^{128a,128b,128d}, R. Gonçalves^{128a}, R. Goncalves Gama^{26a}, J. Goncalves Pinto Firmino Da Costa¹³⁸, G. Gonella⁵¹, L. Gonella¹⁹, A. Gongadze⁶⁸, F. Gonnella¹⁹, J.L. Gonski⁵⁹, S. González de la Hoz¹⁷⁰, S. Gonzalez-Sevilla⁵², L. Goossens³², P.A. Gorbounov⁹⁹, H.A. Gordon²⁷, B. Gorini³², E. Gorini^{76a,76b}, A. Gorišek⁷⁸, A.T. Goshaw⁴⁸, C. Gössling⁴⁶, M.I. Gostkin⁶⁸, C.A. Gottardo²³, C.R. Goudet¹¹⁹, D. Goudami^{137c}, A.G. Goussiou¹⁴⁰, N. Govender^{147b,t}, C. Goy⁵, E. Gozani¹⁵⁴, I. Grabowska-Bold^{41a}, P.O.J. Gradin¹⁶⁸, E.C. Graham⁷⁷, J. Gramling¹⁶⁶, E. Gramstad¹²¹, S. Grancagnolo¹⁷, V. Gratchev¹²⁵, P.M. Gravila^{28f}, C. Gray⁵⁶, H.M. Gray¹⁶, Z.D. Greenwood^{82,u}, C. Grefe²³, K. Gregersen⁸¹, I.M. Gregor⁴⁵, P. Grenier¹⁴⁵, K. Grevtsov⁵, J. Griffiths⁸, A.A. Grillo¹³⁹, K. Grimm⁷⁵, S. Grinstein^{13,v}, Ph. Gris³⁷, J.-F. Grivaz¹¹⁹, S. Groh⁸⁶, E. Gross¹⁷⁵, J. Grosse-Knetter⁵⁷, G.C. Grossi⁸², Z.J. Grout⁸¹, A. Grummer¹⁰⁷, L. Guan⁹², W. Guan¹⁷⁶, J. Guenther³², F. Guescini^{163a}, D. Guest¹⁶⁶, O. Gueta¹⁵⁵, B. Gui¹¹³, E. Guido^{53a,53b}, T. Guillemin⁵, S. Guindon³², U. Gul⁵⁶, C. Gumpert³², J. Guo^{36c}, W. Guo⁹², Y. Guo^{36a,w}, R. Gupta⁴³, S. Gurbuz^{20a}, G. Gustavino¹¹⁵, B.J. Gutelman¹⁵⁴, P. Gutierrez¹¹⁵, N.G. Gutierrez Ortiz⁸¹, C. Gutschow⁸¹, C. Guyot¹³⁸, M.P. Guzik^{41a}, C. Gwenlan¹²², C.B. Gwilliam⁷⁷, A. Haas¹¹², C. Haber¹⁶, H.K. Hadavand⁸, N. Haddad^{137e}, A. Hadeef⁸⁸, S. Hageböck²³, M. Hagihara¹⁶⁴, H. Hakobyan^{180,*}, M. Haleem⁴⁵, J. Haley¹¹⁶, G. Halladjian⁹³, G.D. Hallewell⁸⁸, K. Hamacher¹⁷⁸, P. Hamal¹¹⁷, K. Hamano¹⁷², A. Hamilton^{147a}, G.N. Hamity¹⁴¹, P.G. Hamnett⁴⁵, K. Han^{36a,x}, L. Han^{36a}, S. Han^{35a,35d}, K. Hanagaki^{69,y}, K. Hanawa¹⁵⁷, M. Hance¹³⁹, D.M. Handl¹⁰², B. Haney¹²⁴, P. Hanke^{60a}, J.B. Hansen³⁹, J.D. Hansen³⁹, M.C. Hansen²³, P.H. Hansen³⁹, K. Hara¹⁶⁴, A.S. Hard¹⁷⁶, T. Harenberg¹⁷⁸, F. Hariri¹¹⁹, S. Harkusha⁹⁵, P.F. Harrison¹⁷³, N.M. Hartmann¹⁰², Y. Hasegawa¹⁴², A. Hasib⁴⁹, S. Hassani¹³⁸, S. Haug¹⁸, R. Hauser⁹³, L. Hauswald⁴⁷, L.B. Havener³⁸, M. Havranek¹³⁰, C.M. Hawkes¹⁹, R.J. Hawkings³², D. Hayden⁹³, C.P. Hays¹²², J.M. Hays⁷⁹, H.S. Hayward⁷⁷, S.J. Haywood¹³³, T. Heck⁸⁶, V. Hedberg⁸⁴, L. Heelan⁸, S. Heer²³, K.K. Heidegger⁵¹, S. Heim⁴⁵, T. Heim¹⁶, B. Heinemann^{45,z}, J.J. Heinrich¹⁰², L. Heinrich¹¹², C. Heinz⁵⁵, J. Hejbal¹²⁹, L. Helary³², A. Held¹⁷¹, S. Hellman^{148a,148b}, C. Helsens³², R.C.W. Henderson⁷⁵, Y. Heng¹⁷⁶, S. Henkelmann¹⁷¹, A.M. Henriques Correia³², S. Henrot-Versille¹¹⁹, G.H. Herbert¹⁷, H. Herde²⁵, V. Herget¹⁷⁷, Y. Hernández Jiménez^{147c}, H. Heri⁸⁶, G. Herten⁵¹, R. Hertenberger¹⁰², L. Hervas³², T.C. Herwig¹²⁴, G.G. Hesketh⁸¹, N.P. Hessey^{163a}, J.W. Hetherly⁴³, S. Higashino⁶⁹, E. Higón-Rodríguez¹⁷⁰, K. Hildebrand³³, E. Hill¹⁷², J.C. Hill³⁰, K.H. Hiller⁴⁵, S.J. Hillier¹⁹, M. Hils⁴⁷, I. Hinchliffe¹⁶, M. Hirose⁵¹, D. Hirschbuehl¹⁷⁸, B. Hiti⁷⁸, O. Hladik¹²⁹, D.R. Hlaluku^{147c}, X. Hoad⁴⁹, J. Hobbs¹⁵⁰, N. Hod^{163a}, M.C. Hodgkinson¹⁴¹, P. Hodgson¹⁴¹, A. Hoecker³², M.R. Hoferkamp¹⁰⁷, F. Hoenig¹⁰², D. Hohn²³, T.R. Holmes³³, M. Holzbock¹⁰², M. Homann⁴⁶, S. Honda¹⁶⁴, T. Honda⁶⁹, T.M. Hong¹²⁷, B.H. Hooberman¹⁶⁹, W.H. Hopkins¹¹⁸, Y. Horii¹⁰⁵, A.J. Horton¹⁴⁴, J.-Y. Hostachy⁵⁸, A. Hostiuc¹⁴⁰, S. Hou¹⁵³, A. Hoummada^{137a}, J. Howarth⁸⁷, J. Hoya⁷⁴, M. Hrabovsky¹¹⁷, J. Hrdinka³², I. Hristova¹⁷, J. Hrivnac¹¹⁹, T. Hryn'ova⁵, A. Hrynevich⁹⁶, P.J. Hsu⁶³, S.-C. Hsu¹⁴⁰, Q. Hu²⁷, S. Hu^{36c}, Y. Huang^{35a}, Z. Hubacek¹³⁰, F. Hubaut⁸⁸, F. Huegging²³, T.B. Huffman¹²², E.W. Hughes³⁸,

M. Huhtinen³², R.F.H. Hunter³¹, P. Huo¹⁵⁰, N. Huseynov^{68,b}, J. Huston⁹³, J. Huth⁵⁹, R. Hyneman⁹², G. Iacobucci⁵², G. Iakovidis²⁷, I. Ibragimov¹⁴³, L. Iconomidou-Fayard¹¹⁹, Z. Idrissi^{137e}, P. Iengo³², O. Igonkina^{109,aa}, T. Iizawa¹⁷⁴, Y. Ikegami⁶⁹, M. Ikeno⁶⁹, Y. Ilchenko^{11,ab}, D. Iliadis¹⁵⁶, N. Ilic¹⁴⁵, F. Iltzsche⁴⁷, G. Introzzi^{123a,123b}, P. Ioannou^{9,*}, M. Iodice^{136a}, K. Iordanidou³⁸, V. Ippolito⁵⁹, M.F. Isacson¹⁶⁸, N. Ishijima¹²⁰, M. Ishino¹⁵⁷, M. Ishitsuka¹⁵⁹, C. Issever¹²², S. Istin^{20a}, F. Ito¹⁶⁴, J.M. Iturbe Ponce^{62a}, R. Iuppa^{162a,162b}, H. Iwasaki⁶⁹, J.M. Izen⁴⁴, V. Izzo^{106a}, S. Jabbar³, P. Jackson¹, R.M. Jacobs²³, V. Jain², K.B. Jakobi⁸⁶, K. Jakobs⁵¹, S. Jakobsen⁶⁵, T. Jakoubek¹²⁹, D.O. Jamin¹¹⁶, D.K. Jana⁸², R. Jansky⁵², J. Janssen²³, M. Janus⁵⁷, P.A. Janus^{41a}, G. Jarlskog⁸⁴, N. Javadov^{68,b}, T. Javůrek⁵¹, M. Javurkova⁵¹, F. Jeanneau¹³⁸, L. Jeanty¹⁶, J. Jejelava^{54a,ac}, A. Jelinskas¹⁷³, P. Jenni^{51,ad}, C. Jeske¹⁷³, S. Jézéquel⁵, H. Ji¹⁷⁶, J. Jia¹⁵⁰, H. Jiang⁶⁷, Y. Jiang^{36a}, Z. Jiang¹⁴⁵, S. Jiggins⁸¹, J. Jimenez Pena¹⁷⁰, S. Jin^{35b}, A. Jinaru^{28b}, O. Jinnouchi¹⁵⁹, H. Jivan^{147c}, P. Johansson¹⁴¹, K.A. Johns⁷, C.A. Johnson⁶⁴, W.J. Johnson¹⁴⁰, K. Jon-And^{148a,148b}, R.W.L. Jones⁷⁵, S.D. Jones¹⁵¹, S. Jones⁷, T.J. Jones⁷⁷, J. Jongmanns^{60a}, P.M. Jorge^{128a,128b}, J. Jovicevic^{163a}, X. Ju¹⁷⁶, A. Juste Rozas^{13,v}, M.K. Köhler¹⁷⁵, A. Kaczmarek⁴², M. Kado¹¹⁹, H. Kagan¹¹³, M. Kagan¹⁴⁵, S.J. Kahn⁸⁸, T. Kaji¹⁷⁴, E. Kajomovitz¹⁵⁴, C.W. Kalderon⁸⁴, A. Kaluza⁸⁶, S. Kama⁴³, A. Kamenshchikov¹³², N. Kanaya¹⁵⁷, L. Kanjir⁷⁸, V.A. Kantserov¹⁰⁰, J. Kanzaki⁶⁹, B. Kaplan¹¹², L.S. Kaplan¹⁷⁶, D. Kar^{147c}, K. Karakostas¹⁰, N. Karastathis¹⁰, M.J. Kareem^{163b}, E. Karentzos¹⁰, S.N. Karpov⁶⁸, Z.M. Karpova⁶⁸, V. Kartvelishvili⁷⁵, A.N. Karyukhin¹³², K. Kasahara¹⁶⁴, L. Kashif¹⁷⁶, R.D. Kass¹¹³, A. Kastanas¹⁴⁹, Y. Kataoka¹⁵⁷, C. Kato¹⁵⁷, A. Katre⁵², J. Katzy⁴⁵, K. Kawade⁷⁰, K. Kawagoe⁷³, T. Kawamoto¹⁵⁷, G. Kawamura⁵⁷, E.F. Kay⁷⁷, V.F. Kazanin^{111,c}, R. Keeler¹⁷², R. Kehoe⁴³, J.S. Keller³¹, E. Kellermann⁸⁴, J.J. Kempster⁸⁰, J. Kendrick¹⁹, H. Keoshkerian¹⁶¹, O. Kepka¹²⁹, B.P. Kerševan⁷⁸, S. Kersten¹⁷⁸, R.A. Keyes⁹⁰, M. Khader¹⁶⁹, F. Khalil-zada¹², A. Khanov¹¹⁶, A.G. Kharlamov^{111,c}, T. Kharlamova^{111,c}, A. Khodinov¹⁶⁰, T.J. Khoo⁵², V. Khovanskij^{99,*}, E. Khramov⁶⁸, J. Khubua^{54b,ae}, S. Kido⁷⁰, C.R. Kilby⁸⁰, H.Y. Kim⁸, S.H. Kim¹⁶⁴, Y.K. Kim³³, N. Kimura^{167a,167c}, O.M. Kind¹⁷, B.T. King⁷⁷, D. Kirchmeier⁴⁷, J. Kirk¹³³, A.E. Kiryunin¹⁰³, T. Kishimoto¹⁵⁷, D. Kisielewska^{41a}, V. Kitali⁴⁵, O. Kivernyk⁵, E. Kladiva^{146b}, T. Klapdor-Kleingrothaus⁵¹, M.H. Klein⁹², M. Klein⁷⁷, U. Klein⁷⁷, K. Kleinknecht⁸⁶, P. Klimek¹¹⁰, A. Klimentov²⁷, R. Klingenberg^{46,*}, T. Klingl²³, T. Klioutchnikova³², F.F. Klitzner¹⁰², E.-E. Kluge^{60a}, P. Kluit¹⁰⁹, S. Kluth¹⁰³, E. Kneringer⁶⁵, E.B.F.G. Knoops⁸⁸, A. Knue¹⁰³, A. Kobayashi¹⁵⁷, D. Kobayashi⁷³, T. Kobayashi¹⁵⁷, M. Kobel⁴⁷, M. Kocian¹⁴⁵, P. Kodys¹³¹, T. Koffas³¹, E. Koffeman¹⁰⁹, N.M. Köhler¹⁰³, T. Koi¹⁴⁵, M. Kolb^{60b}, I. Koletsou⁵, T. Kondo⁶⁹, N. Kondrashova^{36c}, K. Köneke⁵¹, A.C. König¹⁰⁸, T. Kono^{69,af}, R. Konoplich^{112,ag}, N. Konstantinidis⁸¹, B. Konya⁸⁴, R. Kopeliansky⁶⁴, S. Koperny^{41a}, K. Korcyl⁴², K. Kordas¹⁵⁶, A. Korn⁸¹, I. Korolkov¹³, E.V. Korolkova¹⁴¹, O. Kortner¹⁰³, S. Kortner¹⁰³, T. Kosek¹³¹, V.V. Kostyukhin²³, A. Kotwal⁴⁸, A. Koulouris¹⁰, A. Kourkoumeli-Charalampidi^{123a,123b}, C. Kourkoumelis⁹, E. Kourlitis¹⁴¹, V. Kouskoura²⁷, A.B. Kowalewska⁴², R. Kowalewski¹⁷², T.Z. Kowalski^{41a}, C. Kozakai¹⁵⁷, W. Kozanecki¹³⁸, A.S. Kozhin¹³², V.A. Kramarenko¹⁰¹, G. Kramberger⁷⁸, D. Krasnopevtsev¹⁰⁰, M.W. Krasny⁸³, A. Krasznahorkay³², D. Krauss¹⁰³, J.A. Kremer^{41a}, J. Kretzschmar⁷⁷, K. Kreutzfeldt⁵⁵, P. Krieger¹⁶¹, K. Krizka¹⁶, K. Kroeninger⁴⁶, H. Kroha¹⁰³, J. Kroll¹²⁹, J. Kroll¹²⁴, J. Kroseberg²³, J. Krstic¹⁴, U. Kruchonak⁶⁸, H. Krüger²³, N. Krumnack⁶⁷, M.C. Kruse⁴⁸, T. Kubota⁹¹, H. Kucuk⁸¹, S. Kудay^{4b}, J.T. Kuechler¹⁷⁸, S. Kuehn³², A. Kugel^{60a}, F. Kuger¹⁷⁷, T. Kuhl⁴⁵, V. Kukhtin⁶⁸, R. Kukla⁸⁸, Y. Kulchitsky⁹⁵, S. Kuleshov^{34b}, Y.P. Kulinich¹⁶⁹, M. Kuna¹¹, T. Kunigo⁷¹, A. Kupco¹²⁹, T. Kupfer⁴⁶, O. Kuprash¹⁵⁵, H. Kurashige⁷⁰, L.L. Kurchaninov^{163a}, Y.A. Kurochkin⁹⁵, M.G. Kurth^{35a,35d}, E.S. Kuwertz¹⁷², M. Kuze¹⁵⁹, J. Kvita¹¹⁷, T. Kwan¹⁷², D. Kyriazopoulos¹⁴¹, A. La Rosa¹⁰³, J.L. La Rosa Navarro^{26d}, L. La Rotonda^{40a,40b}, F. La Ruffa^{40a,40b}, C. Lacasta¹⁷⁰, F. Lacava^{134a,134b}, J. Lacey⁴⁵, D.P.J. Lack⁸⁷, H. Lacker¹⁷, D. Lacour⁸³, E. Ladygin⁶⁸, R. Lafaye⁵, B. Laforge⁸³, S. Lai⁵⁷, S. Lammers⁶⁴, W. Lampl⁷, E. Lançon²⁷, U. Landgraf⁵¹, M.P.J. Landon⁷⁹, M.C. Lanfermann⁵², V.S. Lang⁴⁵, J.C. Lange¹³, R.J. Langenberg³², A.J. Lankford¹⁶⁶, F. Lanni²⁷, K. Lantzsche²³, A. Lanza^{123a},

A. Lapertosa^{53a,53b}, S. Laplace⁸³, J.F. Laporte¹³⁸, T. Lari^{94a}, F. Lasagni Manghi^{22a,22b}, M. Lassnig³², T.S. Lau^{62a}, P. Laurelli⁵⁰, W. Lavrijsen¹⁶, A.T. Law¹³⁹, P. Laycock⁷⁷, T. Lazovich⁵⁹, M. Lazzaroni^{94a,94b}, B. Le⁹¹, O. Le Dortz⁸³, E. Le Guirriec⁸⁸, E.P. Le Quilleuc¹³⁸, M. LeBlanc⁷, T. LeCompte⁶, F. Ledroit-Guillon⁵⁸, C.A. Lee²⁷, G.R. Lee^{34a}, S.C. Lee¹⁵³, L. Lee⁵⁹, B. Lefebvre⁹⁰, G. Lefebvre⁸³, M. Lefebvre¹⁷², F. Legger¹⁰², C. Leggett¹⁶, G. Lehmann Miotto³², X. Lei⁷, W.A. Leight⁴⁵, M.A.L. Leite^{26d}, R. Leitner¹³¹, D. Lellouch¹⁷⁵, B. Lemmer⁵⁷, K.J.C. Leney⁸¹, T. Lenz²³, B. Lenzi³², R. Leone⁷, S. Leone^{126a}, C. Leonidopoulos⁴⁹, G. Lerner¹⁵¹, C. Leroy⁹⁷, R. Les¹⁶¹, A.A.J. Lesage¹³⁸, C.G. Lester³⁰, M. Levchenko¹²⁵, J. Levêque⁵, D. Levin⁹², L.J. Levinson¹⁷⁵, M. Levy¹⁹, D. Lewis⁷⁹, B. Li^{36a,w}, H. Li¹⁵⁰, L. Li^{36c}, Q. Li^{35a,35d}, Q. Li^{36a}, S. Li⁴⁸, X. Li^{36c}, Y. Li¹⁴³, Z. Li^{36b}, Z. Liang^{35a}, B. Liberti^{135a}, A. Liblong¹⁶¹, K. Lie^{62c}, W. Liebig¹⁵, A. Limosani¹⁵², C.Y. Lin³⁰, K. Lin⁹³, S.C. Lin¹⁸², T.H. Lin⁸⁶, R.A. Linck⁶⁴, B.E. Lindquist¹⁵⁰, A.E. Lioni⁵², E. Lipeles¹²⁴, A. Lipniacka¹⁵, M. Lisovyi^{60b}, T.M. Liss^{169,ah}, A. Lister¹⁷¹, A.M. Litke¹³⁹, B. Liu⁶⁷, H. Liu⁹², H. Liu²⁷, J.K.K. Liu¹²², J. Liu^{36b}, J.B. Liu^{36a}, K. Liu⁸⁸, L. Liu¹⁶⁹, M. Liu^{36a}, Y.L. Liu^{36a}, Y. Liu^{36a}, M. Livan^{123a,123b}, A. Lleres⁵⁸, J. Llorente Merino^{35a}, S.L. Lloyd⁷⁹, C.Y. Lo^{62b}, F. Lo Sterzo⁴³, E.M. Lobodzinska⁴⁵, P. Loch⁷, F.K. Loebinger⁸⁷, A. Loesle⁵¹, K.M. Loew²⁵, T. Lohse¹⁷, K. Lohwasser¹⁴¹, M. Lokajicek¹²⁹, B.A. Long²⁴, J.D. Long¹⁶⁹, R.E. Long⁷⁵, L. Longo^{76a,76b}, K.A. Looper¹¹³, J.A. Lopez^{34b}, I. Lopez Paz¹³, A. Lopez Solis⁸³, J. Lorenz¹⁰², N. Lorenzo Martinez⁵, M. Losada²¹, P.J. Lösel¹⁰², X. Lou^{35a}, A. Lounis¹¹⁹, J. Love⁶, P.A. Love⁷⁵, H. Lu^{62a}, N. Lu⁹², Y.J. Lu⁶³, H.J. Lubatti¹⁴⁰, C. Luci^{134a,134b}, A. Lucotte⁵⁸, C. Luedtke⁵¹, F. Luehring⁶⁴, W. Lukas⁶⁵, L. Luminari^{134a}, B. Lund-Jensen¹⁴⁹, M.S. Lutz⁸⁹, P.M. Luzzi⁸³, D. Lynn²⁷, R. Lysak¹²⁹, E. Lytken⁸⁴, F. Lyu^{35a}, V. Lyubushkin⁶⁸, H. Ma²⁷, L.L. Ma^{36b}, Y. Ma^{36b}, G. Maccarrone⁵⁰, A. Macchiolo¹⁰³, C.M. Macdonald¹⁴¹, B. Maček⁷⁸, J. Machado Miguens^{124,128b}, D. Madaffari¹⁷⁰, R. Madar³⁷, W.F. Mader⁴⁷, A. Madsen⁴⁵, N. Madysa⁴⁷, J. Maeda⁷⁰, S. Maeland¹⁵, T. Maeno²⁷, A.S. Maevskiy¹⁰¹, V. Magerl⁵¹, C. Maiani¹¹⁹, C. Maidantchik^{26a}, T. Maier¹⁰², A. Maio^{128a,128b,128d}, O. Majersky^{146a}, S. Majewski¹¹⁸, Y. Makida⁶⁹, N. Makovec¹¹⁹, B. Malaescu⁸³, Pa. Malecki⁴², V.P. Maleev¹²⁵, F. Malek⁵⁸, U. Mallik⁶⁶, D. Malon⁶, C. Malone³⁰, S. Maltezos¹⁰, S. Malyukov³², J. Mamuzic¹⁷⁰, G. Mancini⁵⁰, I. Mandić⁷⁸, J. Maneira^{128a,128b}, L. Manhaes de Andrade Filho^{26b}, J. Manjarres Ramos⁴⁷, K.H. Mankinen⁸⁴, A. Mann¹⁰², A. Manousos³², B. Mansoulie¹³⁸, J.D. Mansour^{35a}, R. Mantifel⁹⁰, M. Mantoani⁵⁷, S. Manzoni^{94a,94b}, L. Mapelli³², G. Marceca²⁹, L. March⁵², L. Marchese¹²², G. Marchiori⁸³, M. Marcisovsky¹²⁹, C.A. Marin Tobon³², M. Marjanovic³⁷, D.E. Marley⁹², F. Marroquim^{26a}, S.P. Marsden⁸⁷, Z. Marshall¹⁶, M.U.F. Martensson¹⁶⁸, S. Marti-Garcia¹⁷⁰, C.B. Martin¹¹³, T.A. Martin¹⁷³, V.J. Martin⁴⁹, B. Martin dit Latour¹⁵, M. Martinez^{13,v}, V.I. Martinez Outschoorn¹⁶⁹, S. Martin-Haugh¹³³, V.S. Martoiu^{28b}, A.C. Martyniuk⁸¹, A. Marzin³², L. Masetti⁸⁶, T. Mashimo¹⁵⁷, R. Mashinistov⁹⁸, J. Masik⁸⁷, A.L. Maslennikov^{111,c}, L.H. Mason⁹¹, L. Massa^{135a,135b}, P. Mastrandrea⁵, A. Mastroberardino^{40a,40b}, T. Masubuchi¹⁵⁷, P. Mättig¹⁷⁸, J. Maurer^{28b}, S.J. Maxfield⁷⁷, D.A. Maximov^{111,c}, R. Mazini¹⁵³, I. Maznas¹⁵⁶, S.M. Mazza^{94a,94b}, N.C. Mc Fadden¹⁰⁷, G. Mc Goldrick¹⁶¹, S.P. Mc Kee⁹², A. McCarn⁹², R.L. McCarthy¹⁵⁰, T.G. McCarthy¹⁰³, L.I. McClymont⁸¹, E.F. McDonald⁹¹, J.A. Mcfayden³², G. Mchedlidze⁵⁷, S.J. McMahon¹³³, P.C. McNamara⁹¹, C.J. McNicol¹⁷³, R.A. McPherson^{172,o}, S. Meehan¹⁴⁰, T.J. Megy⁵¹, S. Mehlhase¹⁰², A. Mehta⁷⁷, T. Meideck⁵⁸, K. Meier^{60a}, B. Meirose⁴⁴, D. Melini^{170,ai}, B.R. Mellado Garcia^{147c}, J.D. Mellenthin⁵⁷, M. Melo^{146a}, F. Meloni¹⁸, A. Melzer²³, S.B. Menary⁸⁷, L. Meng⁷⁷, X.T. Meng⁹², A. Mengarelli^{22a,22b}, S. Menke¹⁰³, E. Meoni^{40a,40b}, S. Mergelmeyer¹⁷, C. Merlassino¹⁸, P. Mermod⁵², L. Merola^{106a,106b}, C. Meroni^{94a}, F.S. Merritt³³, A. Messina^{134a,134b}, J. Metcalfe⁶, A.S. Mete¹⁶⁶, C. Meyer¹²⁴, J-P. Meyer¹³⁸, J. Meyer¹⁰⁹, H. Meyer Zu Theenhausen^{60a}, F. Miano¹⁵¹, R.P. Middleton¹³³, S. Miglioranza^{53a,53b}, L. Mijović⁴⁹, G. Mikenberg¹⁷⁵, M. Mikesikova¹²⁹, M. Mikuž⁷⁸, M. Milesi⁹¹, A. Milic¹⁶¹, D.A. Millar⁷⁹, D.W. Miller³³, C. Mills⁴⁹, A. Milov¹⁷⁵, D.A. Milstead^{148a,148b}, A.A. Minaenko¹³², Y. Minami¹⁵⁷, I.A. Minashvili^{54b}, A.I. Mincer¹¹², B. Mindur^{41a}, M. Mineev⁶⁸,

Y. Minegishi¹⁵⁷, Y. Ming¹⁷⁶, L.M. Mir¹³, A. Mirto^{76a,76b}, K.P. Mistry¹²⁴, T. Mitani¹⁷⁴, J. Mitrevski¹⁰², V.A. Mitsou¹⁷⁰, A. Miucci¹⁸, P.S. Miyagawa¹⁴¹, A. Mizukami⁶⁹, J.U. Mjörnmark⁸⁴, T. Mkrtchyan¹⁸⁰, M. Mlynarikova¹³¹, T. Moa^{148a,148b}, K. Mochizuki⁹⁷, P. Mogg⁵¹, S. Mohapatra³⁸, S. Molander^{148a,148b}, R. Moles-Valls²³, M.C. Mondragon⁹³, K. Mönig⁴⁵, J. Monk³⁹, E. Monnier⁸⁸, A. Montalbano¹⁵⁰, J. Montejó Berlingen³², F. Monticelli⁷⁴, S. Monzani^{94a}, R.W. Moore³, N. Morange¹¹⁹, D. Moreno²¹, M. Moreno Llácer³², P. Moretini^{53a}, M. Morgenstern¹⁰⁹, S. Morgenstern³², D. Mori¹⁴⁴, T. Mori¹⁵⁷, M. Morii⁵⁹, M. Morinaga¹⁷⁴, V. Morisbak¹²¹, A.K. Morley³², G. Mornacchi³², J.D. Morris⁷⁹, L. Morvaj¹⁵⁰, P. Moschovakos¹⁰, M. Mosidze^{54b}, H.J. Moss¹⁴¹, J. Moss^{145,aj}, K. Motohashi¹⁵⁹, R. Mount¹⁴⁵, E. Mountricha²⁷, E.J.W. Moyse⁸⁹, S. Muanza⁸⁸, F. Mueller¹⁰³, J. Mueller¹²⁷, R.S.P. Mueller¹⁰², D. Muenstermann⁷⁵, P. Mullen⁵⁶, G.A. Mullier¹⁸, F.J. Munoz Sanchez⁸⁷, W.J. Murray^{173,133}, H. Musheghyan³², M. Muškinja⁷⁸, C. Mwewa^{147a}, A.G. Myagkov^{132,ak}, M. Myska¹³⁰, B.P. Nachman¹⁶, O. Nackenhorst⁵², K. Nagai¹²², R. Nagai^{69,af}, K. Nagano⁶⁹, Y. Nagasaka⁶¹, K. Nagata¹⁶⁴, M. Nagel⁵¹, E. Nagy⁸⁸, A.M. Nairz³², Y. Nakahama¹⁰⁵, K. Nakamura⁶⁹, T. Nakamura¹⁵⁷, I. Nakano¹¹⁴, R.F. Naranjo Garcia⁴⁵, R. Narayan¹¹, D.I. Narrias Villar^{60a}, I. Naryshkin¹²⁵, T. Naumann⁴⁵, G. Navarro²¹, R. Nayyar⁷, H.A. Neal⁹², P.Yu. Nechaeva⁹⁸, T.J. Neep¹³⁸, A. Negri^{123a,123b}, M. Negrini^{22a}, S. Nektarijevic¹⁰⁸, C. Nellist⁵⁷, A. Nelson¹⁶⁶, M.E. Nelson¹²², S. Nemecek¹²⁹, P. Nemethy¹¹², M. Nessi^{32,al}, M.S. Neubauer¹⁶⁹, M. Neumann¹⁷⁸, P.R. Newman¹⁹, T.Y. Ng^{62c}, Y.S. Ng¹⁷, T. Nguyen Manh⁹⁷, R.B. Nickerson¹²², R. Nicolaidou¹³⁸, J. Nielsen¹³⁹, N. Nikiforou¹¹, V. Nikolaenko^{132,ak}, I. Nikolic-Audit⁸³, K. Nikolopoulos¹⁹, P. Nilsson²⁷, Y. Ninomiya⁶⁹, A. Nisati^{134a}, N. Nishu^{36c}, R. Nisius¹⁰³, I. Nitsche⁴⁶, T. Nitta¹⁷⁴, T. Nobe¹⁵⁷, Y. Noguchi⁷¹, M. Nomachi¹²⁰, I. Nomidis³¹, M.A. Nomura²⁷, T. Nooney⁷⁹, M. Nordberg³², N. Norjoharuddeen¹²², O. Novgorodova⁴⁷, M. Nozaki⁶⁹, L. Nozka¹¹⁷, K. Ntekas¹⁶⁶, E. Nurse⁸¹, F. Nuti⁹¹, K. O'connor²⁵, D.C. O'Neil¹⁴⁴, A.A. O'Rourke⁴⁵, V. O'Shea⁵⁶, F.G. Oakham^{31,d}, H. Oberlack¹⁰³, T. Obermann²³, J. Ocariz⁸³, A. Ochi⁷⁰, I. Ochoa³⁸, J.P. Ochoa-Ricoux^{34a}, S. Oda⁷³, S. Odaka⁶⁹, A. Oh⁸⁷, S.H. Oh⁴⁸, C.C. Ohm¹⁴⁹, H. Ohman¹⁶⁸, H. Oide^{53a,53b}, H. Okawa¹⁶⁴, Y. Okumura¹⁵⁷, T. Okuyama⁶⁹, A. Olariu^{28b}, L.F. Oleiro Seabra^{128a}, S.A. Olivares Pino^{34a}, D. Oliveira Damazio²⁷, M.J.R. Olsson³³, A. Olszewski⁴², J. Olszowska⁴², A. Onofre^{128a,128e}, K. Onogi¹⁰⁵, P.U.E. Onyisi^{11,ab}, H. Oppen¹²¹, M.J. Oreglia³³, Y. Oren¹⁵⁵, D. Orestano^{136a,136b}, N. Orlando^{62b}, R.S. Orr¹⁶¹, B. Osculati^{53a,53b,*}, R. Ospanov^{36a}, G. Otero y Garzon²⁹, H. Otono⁷³, M. Ouchrif^{137d}, F. Ould-Saada¹²¹, A. Ouraou¹³⁸, K.P. Oussoren¹⁰⁹, Q. Ouyang^{35a}, M. Owen⁵⁶, R.E. Owen¹⁹, V.E. Ozcan^{20a}, N. Ozturk⁸, K. Pachal¹⁴⁴, A. Pacheco Pages¹³, L. Pacheco Rodriguez¹³⁸, C. Padilla Aranda¹³, S. Pagan Griso¹⁶, M. Paganini¹⁷⁹, F. Paige²⁷, G. Palacino⁶⁴, S. Palazzo^{40a,40b}, S. Palestini³², M. Palka^{41b}, D. Pallin³⁷, E.St. Panagiotopoulou¹⁰, I. Panagoulas¹⁰, C.E. Pandini⁵², J.G. Panduro Vazquez⁸⁰, P. Pani³², S. Panitkin²⁷, D. Pantea^{28b}, L. Paolozzi⁵², Th.D. Papadopoulou¹⁰, K. Papageorgiou^{9,s}, A. Paramonov⁶, D. Paredes Hernandez¹⁷⁹, A.J. Parker⁷⁵, M.A. Parker³⁰, K.A. Parker⁴⁵, F. Parodi^{53a,53b}, J.A. Parsons³⁸, U. Parzefall⁵¹, V.R. Pascuzzi¹⁶¹, J.M. Pasner¹³⁹, E. Pasqualucci^{134a}, S. Passaggio^{53a}, Fr. Pastore⁸⁰, S. Patariaia⁸⁶, J.R. Pater⁸⁷, T. Pauly³², B. Pearson¹⁰³, S. Pedraza Lopez¹⁷⁰, R. Pedro^{128a,128b}, S.V. Peleganchuk^{111,c}, O. Penc¹²⁹, C. Peng^{35a,35d}, H. Peng^{36a}, J. Penwell⁶⁴, B.S. Peralva^{26b}, M.M. Perego¹³⁸, D.V. Perepelitsa²⁷, F. Peri¹⁷, L. Perini^{94a,94b}, H. Pernegger³², S. Perrella^{106a,106b}, R. Peschke⁴⁵, V.D. Peshekhonov^{68,*}, K. Peters⁴⁵, R.F.Y. Peters⁸⁷, B.A. Petersen³², T.C. Petersen³⁹, E. Petit⁵⁸, A. Petridis¹, C. Petridou¹⁵⁶, P. Petroff¹¹⁹, E. Petrolo^{134a}, M. Petrov¹²², F. Petrucci^{136a,136b}, N.E. Pettersson⁸⁹, A. Peyaud¹³⁸, R. Pezoa^{34b}, F.H. Phillips⁹³, P.W. Phillips¹³³, G. Piacquadio¹⁵⁰, E. Pianori¹⁷³, A. Picazio⁸⁹, M.A. Pickering¹²², R. Piegaia²⁹, J.E. Pilcher³³, A.D. Pilkington⁸⁷, M. Pinamonti^{135a,135b}, J.L. Pinfold³, H. Pirumov⁴⁵, M. Pitt¹⁷⁵, L. Plazak^{146a}, M.-A. Pleier²⁷, V. Pleskot⁸⁶, E. Plotnikova⁶⁸, D. Pluth⁶⁷, P. Podberezko¹¹¹, R. Poettgen⁸⁴, R. Poggi^{123a,123b}, L. Poggioli¹¹⁹, I. Pogrebnyak⁹³, D. Pohl²³, I. Pokharel⁵⁷, G. Polesello^{123a}, A. Poley⁴⁵, A. Policicchio^{40a,40b}, R. Polifka³², A. Polini^{22a}, C.S. Pollard⁵⁶, V. Polychronakos²⁷, K. Pommès³², D. Ponomarenko¹⁰⁰, L. Pontecorvo^{134a},

G.A. Popeneciu^{28d}, D.M. Portillo Quintero⁸³, S. Pospisil¹³⁰, K. Potamianos⁴⁵, I.N. Potrap⁶⁸, C.J. Potter³⁰, H. Potti¹¹, T. Poulsen⁸⁴, J. Poveda³², M.E. Pozo Astigarraga³², P. Pralavorio⁸⁸, A. Pranko¹⁶, S. Prell⁶⁷, D. Price⁸⁷, M. Primavera^{76a}, S. Prince⁹⁰, N. Proklova¹⁰⁰, K. Prokofiev^{62c}, F. Prokoshin^{34b}, S. Protopopescu²⁷, J. Proudfoot⁶, M. Przybycien^{41a}, A. Puri¹⁶⁹, P. Puzo¹¹⁹, J. Qian⁹², Y. Qin⁸⁷, A. Quadt⁵⁷, M. Queitsch-Maitland⁴⁵, D. Quilty⁵⁶, S. Raddum¹²¹, V. Radeka²⁷, V. Radescu¹²², S.K. Radhakrishnan¹⁵⁰, P. Radloff¹¹⁸, P. Rados⁹¹, F. Ragusa^{94a,94b}, G. Rahal¹⁸¹, J.A. Raine⁸⁷, S. Rajagopalan²⁷, C. Rangel-Smith¹⁶⁸, T. Rashid¹¹⁹, S. Raspopov⁵, M.G. Ratti^{94a,94b}, D.M. Rauch⁴⁵, F. Rauscher¹⁰², S. Rave⁸⁶, I. Ravinovich¹⁷⁵, J.H. Rawling⁸⁷, M. Raymond³², A.L. Read¹²¹, N.P. Readioff⁵⁸, M. Reale^{76a,76b}, D.M. Rebuzzi^{123a,123b}, A. Redelbach¹⁷⁷, G. Redlinger²⁷, R. Reece¹³⁹, R.G. Reed^{147c}, K. Reeves⁴⁴, L. Rehnisch¹⁷, J. Reichert¹²⁴, A. Reiss⁸⁶, C. Rembser³², H. Ren^{35a,35d}, M. Rescigno^{134a}, S. Resconi^{94a}, E.D. Resseguie¹²⁴, S. Rettie¹⁷¹, E. Reynolds¹⁹, O.L. Rezanova^{111,c}, P. Reznicek¹³¹, R. Rezvani⁹⁷, R. Richter¹⁰³, S. Richter⁸¹, E. Richter-Was^{41b}, O. Ricken²³, M. Ridel⁸³, P. Rieck¹⁰³, C.J. Riegel¹⁷⁸, J. Rieger⁵⁷, O. Rifki¹¹⁵, M. Rijssenbeek¹⁵⁰, A. Rimoldi^{123a,123b}, M. Rimoldi¹⁸, L. Rinaldi^{22a}, G. Ripellino¹⁴⁹, B. Ristic³², E. Ritsch³², I. Riu¹³, F. Rizatdinova¹¹⁶, E. Rizvi⁷⁹, C. Rizzi¹³, R.T. Roberts⁸⁷, S.H. Robertson^{90,o}, A. Robichaud-Veronneau⁹⁰, D. Robinson³⁰, J.E.M. Robinson⁴⁵, A. Robson⁵⁶, E. Rocco⁸⁶, C. Roda^{126a,126b}, Y. Rodina^{88,am}, S. Rodriguez Bosca¹⁷⁰, A. Rodriguez Perez¹³, D. Rodriguez Rodriguez¹⁷⁰, S. Roe³², C.S. Rogan⁵⁹, O. Röhne¹²¹, J. Roloff⁵⁹, A. Romanouk¹⁰⁰, M. Romano^{22a,22b}, S.M. Romano Saez³⁷, E. Romero Adam¹⁷⁰, N. Rompotis⁷⁷, M. Ronzani⁵¹, L. Roos⁸³, S. Rosati^{134a}, K. Rosbach⁵¹, P. Rose¹³⁹, N.-A. Rosien⁵⁷, E. Rossi^{106a,106b}, L.P. Rossi^{53a}, J.H.N. Rosten³⁰, R. Rosten¹⁴⁰, M. Rotaru^{28b}, J. Rothberg¹⁴⁰, D. Rousseau¹¹⁹, D. Roy^{147c}, A. Rozanov⁸⁸, Y. Rozen¹⁵⁴, X. Ruan^{147c}, F. Rubbo¹⁴⁵, F. Rühr⁵¹, A. Ruiz-Martinez³¹, Z. Rurikova⁵¹, N.A. Rusakovich⁶⁸, H.L. Russell⁹⁰, J.P. Rutherford⁷, N. Ruthmann³², E.M. Rüttinger⁴⁵, Y.F. Ryabov¹²⁵, M. Rybar¹⁶⁹, G. Rybkin¹¹⁹, S. Ryu⁶, A. Ryzhov¹³², G.F. Rzehorz⁵⁷, A.F. Saavedra¹⁵², G. Sabato¹⁰⁹, S. Sacerdoti²⁹, H.F.W. Sadrozinski¹³⁹, R. Sadykov⁶⁸, F. Safai Tehrani^{134a}, P. Saha¹¹⁰, M. Sahinsoy^{60a}, M. Saimpert⁴⁵, M. Saito¹⁵⁷, T. Saito¹⁵⁷, H. Sakamoto¹⁵⁷, Y. Sakurai¹⁷⁴, G. Salamanna^{136a,136b}, J.E. Salazar Loyola^{34b}, D. Salek¹⁰⁹, P.H. Sales De Bruin¹⁶⁸, D. Salihagic¹⁰³, A. Salnikov¹⁴⁵, J. Salt¹⁷⁰, D. Salvatore^{40a,40b}, F. Salvatore¹⁵¹, A. Salvucci^{62a,62b,62c}, A. Salzburger³², D. Sammel⁵¹, D. Sampsonidis¹⁵⁶, D. Sampsonidou¹⁵⁶, J. Sánchez¹⁷⁰, A. Sanchez Pineda^{167a,167c}, H. Sandaker¹²¹, R.L. Sandbach⁷⁹, C.O. Sander⁴⁵, M. Sandhoff¹⁷⁸, C. Sandoval²¹, D.P.C. Sankey¹³³, M. Sannino^{53a,53b}, Y. Sano¹⁰⁵, A. Sansoni⁵⁰, C. Santoni³⁷, H. Santos^{128a}, I. Santoyo Castillo¹⁵¹, A. Saponov⁶⁸, J.G. Saraiva^{128a,128d}, O. Sasaki⁶⁹, K. Sato¹⁶⁴, E. Sauvan⁵, G. Savage⁸⁰, P. Savard^{161,d}, N. Savic¹⁰³, C. Sawyer¹³³, L. Sawyer^{82,u}, C. Sbarra^{22a}, A. Sbrizzi^{22a,22b}, T. Scanlon⁸¹, D.A. Scannicchio¹⁶⁶, J. Schaarschmidt¹⁴⁰, P. Schacht¹⁰³, B.M. Schachtner¹⁰², D. Schaefer³³, L. Schaefer¹²⁴, J. Schaeffer⁸⁶, S. Schaepe³², S. Schaezel^{60b}, U. Schäfer⁸⁶, A.C. Schaffer¹¹⁹, D. Schaile¹⁰², R.D. Schamberger¹⁵⁰, V.A. Schegelsky¹²⁵, D. Scheirich¹³¹, F. Schenck¹⁷, M. Schernau¹⁶⁶, C. Schiavi^{53a,53b}, S. Schier¹³⁹, L.K. Schildgen²³, C. Schillo⁵¹, M. Schioppa^{40a,40b}, S. Schlenker³², K.R. Schmidt-Sommerfeld¹⁰³, K. Schmieden³², C. Schmitt⁸⁶, S. Schmitt⁴⁵, S. Schmitz⁸⁶, U. Schnoor⁵¹, L. Schoeffel¹³⁸, A. Schoening^{60b}, B.D. Schoenrock⁹³, E. Schopf²³, M. Schott⁸⁶, J.F.P. Schouwenberg¹⁰⁸, J. Schovancova³², S. Schramm⁵², N. Schuh⁸⁶, A. Schulte⁸⁶, M.J. Schultens²³, H.-C. Schultz-Coulon^{60a}, M. Schumacher⁵¹, B.A. Schumm¹³⁹, Ph. Schune¹³⁸, A. Schwartzman¹⁴⁵, T.A. Schwarz⁹², H. Schweiger⁸⁷, Ph. Schwemling¹³⁸, R. Schwienhorst⁹³, J. Schwindling¹³⁸, A. Sciandra²³, G. Sciolla²⁵, M. Scornajenghi^{40a,40b}, F. Scuri^{126a}, F. Scutti⁹¹, J. Searcy⁹², P. Seema²³, S.C. Seidel¹⁰⁷, A. Seiden¹³⁹, J.M. Seixas^{26a}, G. Sekhniaidze^{106a}, K. Sekhon⁹², S.J. Sekula⁴³, N. Semprini-Cesari^{22a,22b}, S. Senkin³⁷, C. Serfon¹²¹, L. Serin¹¹⁹, L. Serkin^{167a,167b}, M. Sessa^{136a,136b}, R. Seuster¹⁷², H. Severini¹¹⁵, T. Šfiligoj⁷⁸, F. Sforza¹⁶⁵, A. Sfyrila⁵², E. Shabalina⁵⁷, N.W. Shaikh^{148a,148b}, L.Y. Shan^{35a}, R. Shang¹⁶⁹, J.T. Shank²⁴, M. Shapiro¹⁶, P.B. Shatalov⁹⁹, K. Shaw^{167a,167b}, S.M. Shaw⁸⁷, A. Shcherbakova^{148a,148b}, C.Y. Shehu¹⁵¹, Y. Shen¹¹⁵, N. Sherafati³¹, A.D. Sherman²⁴, P. Sherwood⁸¹, L. Shi^{153,an}, S. Shimizu⁷⁰,

C.O. Shimmin¹⁷⁹, M. Shimojima¹⁰⁴, I.P.J. Shipsey¹²², S. Shirabe⁷³, M. Shiyakova^{68,ao}, J. Shlomi¹⁷⁵, A. Shmeleva⁹⁸, D. Shoaleh Saadi⁹⁷, M.J. Shochet³³, S. Shojaii^{94a,94b}, D.R. Shope¹¹⁵, S. Shrestha¹¹³, E. Shulga¹⁰⁰, M.A. Shupe⁷, P. Sicho¹²⁹, A.M. Sickles¹⁶⁹, P.E. Sidebo¹⁴⁹, E. Sideras Haddad^{147c}, O. Sidiropoulou¹⁷⁷, A. Sidoti^{22a,22b}, F. Siegert⁴⁷, Dj. Sijacki¹⁴, J. Silva^{128a,128d}, M. Silva Jr.¹⁷⁶, S.B. Silverstein^{148a}, V. Simak¹³⁰, L. Simic⁶⁸, S. Simion¹¹⁹, E. Simioni⁸⁶, B. Simmons⁸¹, M. Simon⁸⁶, P. Sinervo¹⁶¹, N.B. Sinev¹¹⁸, M. Sioli^{22a,22b}, G. Siragusa¹⁷⁷, I. Siral⁹², S.Yu. Sivoklokov¹⁰¹, J. Sjölin^{148a,148b}, M.B. Skinner⁷⁵, P. Skubic¹¹⁵, M. Slater¹⁹, T. Slavicek¹³⁰, M. Slawinska⁴², K. Sliwa¹⁶⁵, R. Slovak¹³¹, V. Smakhtin¹⁷⁵, B.H. Smart⁵, J. Smiesko^{146a}, N. Smirnov¹⁰⁰, S.Yu. Smirnov¹⁰⁰, Y. Smirnov¹⁰⁰, L.N. Smirnova^{101,ap}, O. Smirnova⁸⁴, J.W. Smith⁵⁷, M.N.K. Smith³⁸, R.W. Smith³⁸, M. Smizanska⁷⁵, K. Smolek¹³⁰, A.A. Snesarev⁹⁸, I.M. Snyder¹¹⁸, S. Snyder²⁷, R. Sobie^{172,o}, F. Socher⁴⁷, A. Soffer¹⁵⁵, A. Søggaard⁴⁹, D.A. Soh¹⁵³, G. Sokhrannyi⁷⁸, C.A. Solans Sanchez³², M. Solar¹³⁰, E.Yu. Soldatov¹⁰⁰, U. Soldevila¹⁷⁰, A.A. Solodkov¹³², A. Soloshenko⁶⁸, O.V. Solovyanov¹³², V. Solov'yev¹²⁵, P. Sommer¹⁴¹, H. Son¹⁶⁵, W. Song¹³³, A. Sopczak¹³⁰, D. Sosa^{60b}, C.L. Sotiropoulou^{126a,126b}, S. Sottocornola^{123a,123b}, R. Soualah^{167a,167c}, A.M. Soukharev^{111,c}, D. South⁴⁵, B.C. Sowden⁸⁰, S. Spagnolo^{76a,76b}, M. Spalla^{126a,126b}, M. Spangenberg¹⁷³, F. Spanò⁸⁰, D. Sperlich¹⁷, F. Spettel¹⁰³, T.M. Spieker^{60a}, R. Spighi^{22a}, G. Spigo³², L.A. Spiller⁹¹, M. Spousta¹³¹, R.D. St. Denis^{56,*}, A. Stabile^{94a,94b}, R. Stamen^{60a}, S. Stamm¹⁷, E. Stanecka⁴², R.W. Stanek⁶, C. Stanescu^{136a}, M.M. Stanitzki⁴⁵, B.S. Stapf¹⁰⁹, S. Stapnes¹²¹, E.A. Starchenko¹³², G.H. Stark³³, J. Stark⁵⁸, S.H. Stark³⁹, P. Staroba¹²⁹, P. Starovoitov^{60a}, S. Stärz³², R. Staszewski⁴², M. Stegler⁴⁵, P. Steinberg²⁷, B. Stelzer¹⁴⁴, H.J. Stelzer³², O. Stelzer-Chilton^{163a}, H. Stenzel⁵⁵, T.J. Stevenson⁷⁹, G.A. Stewart⁵⁶, M.C. Stockton¹¹⁸, M. Stoebe⁹⁰, G. Stoicea^{28b}, P. Stolte⁵⁷, S. Stonjek¹⁰³, A.R. Stradling⁸, A. Straessner⁴⁷, M.E. Stramaglia¹⁸, J. Strandberg¹⁴⁹, S. Strandberg^{148a,148b}, M. Strauss¹¹⁵, P. Strizenec^{146b}, R. Ströhmer¹⁷⁷, D.M. Strom¹¹⁸, R. Stroynowski⁴³, A. Strubig⁴⁹, S.A. Stucci²⁷, B. Stugu¹⁵, N.A. Styles⁴⁵, D. Su¹⁴⁵, J. Su¹²⁷, S. Suchek^{60a}, Y. Sugaya¹²⁰, M. Suk¹³⁰, V.V. Sulin⁹⁸, DMS Sultan^{162a,162b}, S. Sultansoy^{4c}, T. Sumida⁷¹, S. Sun⁵⁹, X. Sun³, K. Suruliz¹⁵¹, C.J.E. Suster¹⁵², M.R. Sutton¹⁵¹, S. Suzuki⁶⁹, M. Svatos¹²⁹, M. Swiatlowski³³, S.P. Swift², I. Sykora^{146a}, T. Sykora¹³¹, D. Ta⁵¹, K. Tackmann⁴⁵, J. Taenzer¹⁵⁵, A. Taffard¹⁶⁶, R. Tafirot^{163a}, E. Tahirovic⁷⁹, N. Taiblum¹⁵⁵, H. Takai²⁷, R. Takashima⁷², E.H. Takasugi¹⁰³, K. Takeda⁷⁰, T. Takeshita¹⁴², Y. Takubo⁶⁹, M. Talby⁸⁸, A.A. Talyshev^{111,c}, J. Tanaka¹⁵⁷, M. Tanaka¹⁵⁹, R. Tanaka¹¹⁹, R. Tanioka⁷⁰, B.B. Tannenwald¹¹³, S. Tapia Araya^{34b}, S. Tapprogge⁸⁶, S. Tarem¹⁵⁴, G.F. Tartarelli^{94a}, P. Tas¹³¹, M. Tasevsky¹²⁹, T. Tashiro⁷¹, E. Tassi^{40a,40b}, A. Tavares Delgado^{128a,128b}, Y. Tayalati^{137e}, A.C. Taylor¹⁰⁷, A.J. Taylor⁴⁹, G.N. Taylor⁹¹, P.T.E. Taylor⁹¹, W. Taylor^{163b}, P. Teixeira-Dias⁸⁰, D. Temple¹⁴⁴, H. Ten Kate³², P.K. Teng¹⁵³, J.J. Teoh¹²⁰, F. Tepel¹⁷⁸, S. Terada⁶⁹, K. Terashi¹⁵⁷, J. Terron⁸⁵, S. Terzo¹³, M. Testa⁵⁰, R.J. Teuscher^{161,o}, S.J. Thais¹⁷⁹, T. Thevenaux-Pelzer⁸⁸, F. Thiele³⁹, J.P. Thomas¹⁹, J. Thomas-Wilsker⁸⁰, P.D. Thompson¹⁹, A.S. Thompson⁵⁶, L.A. Thomsen¹⁷⁹, E. Thomson¹²⁴, Y. Tian³⁸, M.J. Tibbets¹⁶, R.E. Ticse Torres⁵⁷, V.O. Tikhomirov^{98,aq}, Yu.A. Tikhonov^{111,c}, S. Timoshenko¹⁰⁰, P. Tipton¹⁷⁹, S. Tisserant⁸⁸, K. Todome¹⁵⁹, S. Todorova-Nova⁵, S. Todt⁴⁷, J. Tojo⁷³, S. Tokár^{146a}, K. Tokushuku⁶⁹, E. Tolley¹¹³, L. Tomlinson⁸⁷, M. Tomoto¹⁰⁵, L. Tompkins^{145,ar}, K. Toms¹⁰⁷, B. Tong⁵⁹, P. Tornambe⁵¹, E. Torrence¹¹⁸, H. Torres⁴⁷, E. Torró Pastor¹⁴⁰, J. Toth^{88,as}, F. Touchard⁸⁸, D.R. Tovey¹⁴¹, C.J. Treado¹¹², T. Trefzger¹⁷⁷, F. Tresoldi¹⁵¹, A. Tricoli²⁷, I.M. Trigger^{163a}, S. Trincas-Duvold⁸³, M.F. Tripijana¹³, W. Trischuk¹⁶¹, B. Trocme⁵⁸, A. Trofymov⁴⁵, C. Troncon^{94a}, M. Trovatelli¹⁷², L. Truong^{147b}, M. Trzebinski⁴², A. Trzupek⁴², K.W. Tsang^{62a}, J.C.-L. Tseng¹²², P.V. Tsiarshka⁹⁵, N. Tsirintanis⁹, S. Tsiskaridze¹³, V. Tsiskaridze⁵¹, E.G. Tskhadadze^{54a}, I.I. Tsukerman⁹⁹, V. Tsulaia¹⁶, S. Tsuno⁶⁹, D. Tsybychev¹⁵⁰, Y. Tu^{62b}, A. Tudorache^{28b}, V. Tudorache^{28b}, T.T. Tulbure^{28a}, A.N. Tuna⁵⁹, S. Turchikhin⁶⁸, D. Turgeman¹⁷⁵, I. Turk Cakir^{4b,at}, R. Turra^{94a}, P.M. Tuts³⁸, G. Ucchielli^{22a,22b}, I. Ueda⁶⁹, M. Ughetto^{148a,148b}, F. Ukegawa¹⁶⁴, G. Unal³², A. Undrus²⁷, G. Unel¹⁶⁶, F.C. Ungaro⁹¹, Y. Unno⁶⁹, K. Uno¹⁵⁷, J. Urban^{146b}, P. Urquijo⁹¹, P. Urrejola⁸⁶, G. Usai⁸,

J. Usui⁶⁹, L. Vacavant⁸⁸, V. Vacek¹³⁰, B. Vachon⁹⁰, K.O.H. Vadla¹²¹, A. Vaidya⁸¹, C. Valderanis¹⁰², E. Valdes Santurio^{148a,148b}, M. Valente⁵², S. Valentinetti^{22a,22b}, A. Valero¹⁷⁰, L. Valéry¹³, A. Vallier⁵, J.A. Valls Ferrer¹⁷⁰, W. Van Den Wollenberg¹⁰⁹, H. van der Graaf¹⁰⁹, P. van Gemmeren⁶, J. Van Nieuwkoop¹⁴⁴, I. van Vulpen¹⁰⁹, M.C. van Woerden¹⁰⁹, M. Vanadia^{135a,135b}, W. Vandelli³², A. Vaniachine¹⁶⁰, P. Vankov¹⁰⁹, G. Vardanyan¹⁸⁰, R. Vari^{134a}, E.W. Varnes⁷, C. Varni^{53a,53b}, T. Varol⁴³, D. Varouchas¹¹⁹, A. Vartapetian⁸, K.E. Varvell¹⁵², J.G. Vasquez¹⁷⁹, G.A. Vasquez^{34b}, F. Vazeille³⁷, D. Vazquez Furelos¹³, T. Vazquez Schroeder⁹⁰, J. Veatch⁵⁷, V. Veeraraghavan⁷, L.M. Veloce¹⁶¹, F. Veloso^{128a,128c}, S. Veneziano^{134a}, A. Ventura^{76a,76b}, M. Venturi¹⁷², N. Venturi³², V. Vercesi^{123a}, M. Verducci^{136a,136b}, W. Verkerke¹⁰⁹, A.T. Vermeulen¹⁰⁹, J.C. Vermeulen¹⁰⁹, M.C. Vetterli^{144,d}, N. Viaux Maira^{34b}, O. Viazlo⁸⁴, I. Vichou^{169,*}, T. Vickey¹⁴¹, O.E. Vickey Boeriu¹⁴¹, G.H.A. Viehhauser¹²², S. Viel¹⁶, L. Vigani¹²², M. Villa^{22a,22b}, M. Villaplana Perez^{94a,94b}, E. Vilucchi⁵⁰, M.G. Vinciter³¹, V.B. Vinogradov⁶⁸, A. Vishwakarma⁴⁵, C. Vittori^{22a,22b}, I. Vivarelli¹⁵¹, S. Vlachos¹⁰, M. Vogel¹⁷⁸, P. Vokac¹³⁰, G. Volpi¹³, H. von der Schmitt¹⁰³, E. von Toerne²³, V. Vorobel¹³¹, K. Vorobev¹⁰⁰, M. Vos¹⁷⁰, R. Voss³², J.H. Vosseveld⁷⁷, N. Vranjes¹⁴, M. Vranjes Milosavljevic¹⁴, V. Vrba¹³⁰, M. Vreeswijk¹⁰⁹, R. Vuillermet³², I. Vukotic³³, P. Wagner²³, W. Wagner¹⁷⁸, J. Wagner-Kuhr¹⁰², H. Wahlberg⁷⁴, S. Währmund⁴⁷, K. Wakamiya⁷⁰, J. Walder⁷⁵, R. Walker¹⁰², W. Walkowiak¹⁴³, V. Wallangen^{148a,148b}, C. Wang^{35b}, C. Wang^{36b,au}, F. Wang¹⁷⁶, H. Wang¹⁶, H. Wang³, J. Wang⁴⁵, J. Wang¹⁵², Q. Wang¹¹⁵, R.-J. Wang⁸³, R. Wang⁶, S.M. Wang¹⁵³, T. Wang³⁸, W. Wang^{153,av}, W. Wang^{36a,aw}, Z. Wang^{36c}, C. Wanotayaroj⁴⁵, A. Warburton⁹⁰, C.P. Ward³⁰, D.R. Wardrope⁸¹, A. Washbrook⁴⁹, P.M. Watkins¹⁹, A.T. Watson¹⁹, M.F. Watson¹⁹, G. Watts¹⁴⁰, S. Watts⁸⁷, B.M. Waugh⁸¹, A.F. Webb¹¹, S. Webb⁸⁶, M.S. Weber¹⁸, S.M. Weber^{60a}, S.W. Weber¹⁷⁷, S.A. Weber³¹, J.S. Webster⁶, A.R. Weidberg¹²², B. Weinert⁶⁴, J. Weingarten⁵⁷, M. Weirich⁸⁶, C. Weiser⁵¹, P.S. Wells³², T. Wenaus²⁷, T. Wengler³², S. Wenig³², N. Wormes²³, M.D. Werner⁶⁷, P. Werner³², M. Wessels^{60a}, T.D. Weston¹⁸, K. Whalen¹¹⁸, N.L. Whallon¹⁴⁰, A.M. Wharton⁷⁵, A.S. White⁹², A. White⁸, M.J. White¹, R. White^{34b}, D. Whiteson¹⁶⁶, B.W. Whitmore⁷⁵, F.J. Wickens¹³³, W. Wiedenmann¹⁷⁶, M. WIELERS¹³³, C. Wigglesworth³⁹, L.A.M. Wiik-Fuchs⁵¹, A. Wildauer¹⁰³, F. Wilk⁸⁷, H.G. Wilkens³², H.H. Williams¹²⁴, S. Williams¹⁰⁹, C. Willis⁹³, S. Willocq⁸⁹, J.A. Wilson¹⁹, I. Wingerter-Seez⁵, E. Winkels¹⁵¹, F. Winklmeier¹¹⁸, O.J. Winston¹⁵¹, B.T. Winter²³, M. Wittgen¹⁴⁵, M. Wobisch^{82,u}, A. Wolf⁸⁶, T.M.H. Wolf¹⁰⁹, R. Wolff⁸⁸, M.W. Wolter⁴², H. Wolters^{128a,128c}, V.W.S. Wong¹⁷¹, N.L. Woods¹³⁹, S.D. Worm¹⁹, B.K. Wosiek⁴², J. Wotschack³², K.W. Wozniak⁴², M. Wu³³, S.L. Wu¹⁷⁶, X. Wu⁵², Y. Wu⁹², T.R. Wyatt⁸⁷, B.M. Wynne⁴⁹, S. Xella³⁹, Z. Xi⁹², L. Xia^{35c}, D. Xu^{35a}, L. Xu²⁷, T. Xu¹³⁸, W. Xu⁹², B. Yabsley¹⁵², S. Yacoob^{147a}, D. Yamaguchi¹⁵⁹, Y. Yamaguchi¹⁵⁹, A. Yamamoto⁶⁹, S. Yamamoto¹⁵⁷, T. Yamanaka¹⁵⁷, F. Yamane⁷⁰, M. Yamatani¹⁵⁷, T. Yamazaki¹⁵⁷, Y. Yamazaki⁷⁰, Z. Yan²⁴, H. Yang^{36c}, H. Yang¹⁶, Y. Yang¹⁵³, Z. Yang¹⁵, W.-M. Yao¹⁶, Y.C. Yap⁴⁵, Y. Yasu⁶⁹, E. Yatsenko⁵, K.H. Yau Wong²³, J. Ye⁴³, S. Ye²⁷, I. Yeletsikh⁶⁸, E. Yigitbasi²⁴, E. Yildirim⁸⁶, K. Yorita¹⁷⁴, K. Yoshihara¹²⁴, C. Young¹⁴⁵, C.J.S. Young³², J. Yu⁸, J. Yu⁶⁷, S.P.Y. Yuen²³, I. Yusufović^{30,ax}, B. Zabinski⁴², G. Zacharis¹⁰, R. Zaidan¹³, A.M. Zaitsev^{132,ak}, N. Zakharchuk⁴⁵, J. Zalieckas¹⁵, A. Zaman¹⁵⁰, S. Zambito⁵⁹, D. Zanzi⁹¹, C. Zeitnitz¹⁷⁸, G. Zemaityte¹²², A. Zemla^{41a}, J.C. Zeng¹⁶⁹, Q. Zeng¹⁴⁵, O. Zenin¹³², T. Ženiš^{146a}, D. Zerwas¹¹⁹, D. Zhang^{36b}, D. Zhang⁹², F. Zhang¹⁷⁶, G. Zhang^{36a,aw}, H. Zhang¹¹⁹, J. Zhang⁶, L. Zhang⁵¹, L. Zhang^{36a}, M. Zhang¹⁶⁹, P. Zhang^{35b}, R. Zhang²³, R. Zhang^{36a,au}, X. Zhang^{36b}, Y. Zhang^{35a,35d}, Z. Zhang¹¹⁹, X. Zhao⁴³, Y. Zhao^{36b,x}, Z. Zhao^{36a}, A. Zhemchugov⁶⁸, B. Zhou⁹², C. Zhou¹⁷⁶, L. Zhou⁴³, M. Zhou^{35a,35d}, M. Zhou¹⁵⁰, N. Zhou^{36c}, Y. Zhou⁷, C.G. Zhu^{36b}, H. Zhu^{35a}, J. Zhu⁹², Y. Zhu^{36a}, X. Zhuang^{35a}, K. Zhukov⁹⁸, A. Zibell¹⁷⁷, D. Zieminska⁶⁴, N.I. Zimine⁶⁸, C. Zimmermann⁸⁶, S. Zimmermann⁵¹, Z. Zinonos¹⁰³, M. Zinser⁸⁶, M. Ziolkowski¹⁴³, L. Živković¹⁴, G. Zobernig¹⁷⁶, A. Zoccoli^{22a,22b}, R. Zou³³, M. zur Nedden¹⁷, L. Zwalinski³².

¹ Department of Physics, University of Adelaide, Adelaide, Australia

- ² Physics Department, SUNY Albany, Albany NY, United States of America
- ³ Department of Physics, University of Alberta, Edmonton AB, Canada
- ⁴ ^(a) Department of Physics, Ankara University, Ankara; ^(b) Istanbul Aydin University, Istanbul; ^(c) Division of Physics, TOBB University of Economics and Technology, Ankara, Turkey
- ⁵ LAPP, CNRS/IN2P3 and Université Savoie Mont Blanc, Annecy-le-Vieux, France
- ⁶ High Energy Physics Division, Argonne National Laboratory, Argonne IL, United States of America
- ⁷ Department of Physics, University of Arizona, Tucson AZ, United States of America
- ⁸ Department of Physics, The University of Texas at Arlington, Arlington TX, United States of America
- ⁹ Physics Department, National and Kapodistrian University of Athens, Athens, Greece
- ¹⁰ Physics Department, National Technical University of Athens, Zografou, Greece
- ¹¹ Department of Physics, The University of Texas at Austin, Austin TX, United States of America
- ¹² Institute of Physics, Azerbaijan Academy of Sciences, Baku, Azerbaijan
- ¹³ Institut de Física d'Altes Energies (IFAE), The Barcelona Institute of Science and Technology, Barcelona, Spain
- ¹⁴ Institute of Physics, University of Belgrade, Belgrade, Serbia
- ¹⁵ Department for Physics and Technology, University of Bergen, Bergen, Norway
- ¹⁶ Physics Division, Lawrence Berkeley National Laboratory and University of California, Berkeley CA, United States of America
- ¹⁷ Department of Physics, Humboldt University, Berlin, Germany
- ¹⁸ Albert Einstein Center for Fundamental Physics and Laboratory for High Energy Physics, University of Bern, Bern, Switzerland
- ¹⁹ School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom
- ²⁰ ^(a) Department of Physics, Bogazici University, Istanbul; ^(b) Department of Physics Engineering, Gaziantep University, Gaziantep; ^(d) Istanbul Bilgi University, Faculty of Engineering and Natural Sciences, Istanbul; ^(e) Bahcesehir University, Faculty of Engineering and Natural Sciences, Istanbul, Turkey
- ²¹ Centro de Investigaciones, Universidad Antonio Narino, Bogota, Colombia
- ²² ^(a) INFN Sezione di Bologna; ^(b) Dipartimento di Fisica e Astronomia, Università di Bologna, Bologna, Italy
- ²³ Physikalisches Institut, University of Bonn, Bonn, Germany
- ²⁴ Department of Physics, Boston University, Boston MA, United States of America
- ²⁵ Department of Physics, Brandeis University, Waltham MA, United States of America
- ²⁶ ^(a) Universidade Federal do Rio De Janeiro COPPE/EE/IF, Rio de Janeiro; ^(b) Electrical Circuits Department, Federal University of Juiz de Fora (UFJF), Juiz de Fora; ^(c) Federal University of Sao Joao del Rei (UFSJ), Sao Joao del Rei; ^(d) Instituto de Fisica, Universidade de Sao Paulo, Sao Paulo, Brazil
- ²⁷ Physics Department, Brookhaven National Laboratory, Upton NY, United States of America
- ²⁸ ^(a) Transilvania University of Brasov, Brasov; ^(b) Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest; ^(c) Department of Physics, Alexandru Ioan Cuza University of Iasi, Iasi; ^(d) National Institute for Research and Development of Isotopic and Molecular Technologies, Physics Department, Cluj Napoca; ^(e) University Politehnica Bucharest, Bucharest; ^(f) West University in Timisoara, Timisoara, Romania
- ²⁹ Departamento de Física, Universidad de Buenos Aires, Buenos Aires, Argentina
- ³⁰ Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom
- ³¹ Department of Physics, Carleton University, Ottawa ON, Canada
- ³² CERN, Geneva, Switzerland
- ³³ Enrico Fermi Institute, University of Chicago, Chicago IL, United States of America
- ³⁴ ^(a) Departamento de Física, Pontificia Universidad Católica de Chile, Santiago; ^(b) Departamento de

Física, Universidad Técnica Federico Santa María, Valparaíso, Chile

³⁵ ^(a) Institute of High Energy Physics, Chinese Academy of Sciences, Beijing; ^(b) Department of Physics, Nanjing University, Jiangsu; ^(c) Physics Department, Tsinghua University, Beijing 100084; ^(d) University of Chinese Academy of Science (UCAS), Beijing, China

³⁶ ^(a) Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Anhui; ^(b) School of Physics, Shandong University, Shandong; ^(c) School of Physics and Astronomy, Key Laboratory for Particle Physics, Astrophysics and Cosmology, Ministry of Education; Shanghai Key Laboratory for Particle Physics and Cosmology, Tsung-Dao Lee Institute, Shanghai Jiao Tong University, China

³⁷ Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France

³⁸ Nevis Laboratory, Columbia University, Irvington NY, United States of America

³⁹ Niels Bohr Institute, University of Copenhagen, Kobenhavn, Denmark

⁴⁰ ^(a) INFN Gruppo Collegato di Cosenza, Laboratori Nazionali di Frascati; ^(b) Dipartimento di Fisica, Università della Calabria, Rende, Italy

⁴¹ ^(a) AGH University of Science and Technology, Faculty of Physics and Applied Computer Science, Krakow; ^(b) Marian Smoluchowski Institute of Physics, Jagiellonian University, Krakow, Poland

⁴² Institute of Nuclear Physics Polish Academy of Sciences, Krakow, Poland

⁴³ Physics Department, Southern Methodist University, Dallas TX, United States of America

⁴⁴ Physics Department, University of Texas at Dallas, Richardson TX, United States of America

⁴⁵ DESY, Hamburg and Zeuthen, Germany

⁴⁶ Lehrstuhl für Experimentelle Physik IV, Technische Universität Dortmund, Dortmund, Germany

⁴⁷ Institut für Kern- und Teilchenphysik, Technische Universität Dresden, Dresden, Germany

⁴⁸ Department of Physics, Duke University, Durham NC, United States of America

⁴⁹ SUPA - School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom

⁵⁰ INFN e Laboratori Nazionali di Frascati, Frascati, Italy

⁵¹ Fakultät für Mathematik und Physik, Albert-Ludwigs-Universität, Freiburg, Germany

⁵² Departement de Physique Nucleaire et Corpusculaire, Université de Genève, Geneva, Switzerland

⁵³ ^(a) INFN Sezione di Genova; ^(b) Dipartimento di Fisica, Università di Genova, Genova, Italy

⁵⁴ ^(a) E. Andronikashvili Institute of Physics, Iv. Javakhishvili Tbilisi State University, Tbilisi; ^(b) High Energy Physics Institute, Tbilisi State University, Tbilisi, Georgia

⁵⁵ II Physikalisches Institut, Justus-Liebig-Universität Giessen, Giessen, Germany

⁵⁶ SUPA - School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom

⁵⁷ II Physikalisches Institut, Georg-August-Universität, Göttingen, Germany

⁵⁸ Laboratoire de Physique Subatomique et de Cosmologie, Université Grenoble-Alpes, CNRS/IN2P3, Grenoble, France

⁵⁹ Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge MA, United States of America

⁶⁰ ^(a) Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg; ^(b) Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

⁶¹ Faculty of Applied Information Science, Hiroshima Institute of Technology, Hiroshima, Japan

⁶² ^(a) Department of Physics, The Chinese University of Hong Kong, Shatin, N.T., Hong Kong; ^(b) Department of Physics, The University of Hong Kong, Hong Kong; ^(c) Department of Physics and Institute for Advanced Study, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong, China

⁶³ Department of Physics, National Tsing Hua University, Taiwan, Taiwan

⁶⁴ Department of Physics, Indiana University, Bloomington IN, United States of America

⁶⁵ Institut für Astro- und Teilchenphysik, Leopold-Franzens-Universität, Innsbruck, Austria

- 66 University of Iowa, Iowa City IA, United States of America
- 67 Department of Physics and Astronomy, Iowa State University, Ames IA, United States of America
- 68 Joint Institute for Nuclear Research, JINR Dubna, Dubna, Russia
- 69 KEK, High Energy Accelerator Research Organization, Tsukuba, Japan
- 70 Graduate School of Science, Kobe University, Kobe, Japan
- 71 Faculty of Science, Kyoto University, Kyoto, Japan
- 72 Kyoto University of Education, Kyoto, Japan
- 73 Research Center for Advanced Particle Physics and Department of Physics, Kyushu University, Fukuoka, Japan
- 74 Instituto de Física La Plata, Universidad Nacional de La Plata and CONICET, La Plata, Argentina
- 75 Physics Department, Lancaster University, Lancaster, United Kingdom
- 76 ^(a) INFN Sezione di Lecce; ^(b) Dipartimento di Matematica e Fisica, Università del Salento, Lecce, Italy
- 77 Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom
- 78 Department of Experimental Particle Physics, Jožef Stefan Institute and Department of Physics, University of Ljubljana, Ljubljana, Slovenia
- 79 School of Physics and Astronomy, Queen Mary University of London, London, United Kingdom
- 80 Department of Physics, Royal Holloway University of London, Surrey, United Kingdom
- 81 Department of Physics and Astronomy, University College London, London, United Kingdom
- 82 Louisiana Tech University, Ruston LA, United States of America
- 83 Laboratoire de Physique Nucléaire et de Hautes Energies, UPMC and Université Paris-Diderot and CNRS/IN2P3, Paris, France
- 84 Fysiska institutionen, Lunds universitet, Lund, Sweden
- 85 Departamento de Física Teórica C-15, Universidad Autónoma de Madrid, Madrid, Spain
- 86 Institut für Physik, Universität Mainz, Mainz, Germany
- 87 School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom
- 88 CPPM, Aix-Marseille Université and CNRS/IN2P3, Marseille, France
- 89 Department of Physics, University of Massachusetts, Amherst MA, United States of America
- 90 Department of Physics, McGill University, Montreal QC, Canada
- 91 School of Physics, University of Melbourne, Victoria, Australia
- 92 Department of Physics, The University of Michigan, Ann Arbor MI, United States of America
- 93 Department of Physics and Astronomy, Michigan State University, East Lansing MI, United States of America
- 94 ^(a) INFN Sezione di Milano; ^(b) Dipartimento di Fisica, Università di Milano, Milano, Italy
- 95 B.I. Stepanov Institute of Physics, National Academy of Sciences of Belarus, Minsk, Republic of Belarus
- 96 Research Institute for Nuclear Problems of Byelorussian State University, Minsk, Republic of Belarus
- 97 Group of Particle Physics, University of Montreal, Montreal QC, Canada
- 98 P.N. Lebedev Physical Institute of the Russian Academy of Sciences, Moscow, Russia
- 99 Institute for Theoretical and Experimental Physics (ITEP), Moscow, Russia
- 100 National Research Nuclear University MEPhI, Moscow, Russia
- 101 D.V. Skobeltsyn Institute of Nuclear Physics, M.V. Lomonosov Moscow State University, Moscow, Russia
- 102 Fakultät für Physik, Ludwig-Maximilians-Universität München, München, Germany
- 103 Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), München, Germany
- 104 Nagasaki Institute of Applied Science, Nagasaki, Japan
- 105 Graduate School of Science and Kobayashi-Maskawa Institute, Nagoya University, Nagoya, Japan

- ¹⁰⁶ (a) INFN Sezione di Napoli; (b) Dipartimento di Fisica, Università di Napoli, Napoli, Italy
- ¹⁰⁷ Department of Physics and Astronomy, University of New Mexico, Albuquerque NM, United States of America
- ¹⁰⁸ Institute for Mathematics, Astrophysics and Particle Physics, Radboud University Nijmegen/Nikhef, Nijmegen, Netherlands
- ¹⁰⁹ Nikhef National Institute for Subatomic Physics and University of Amsterdam, Amsterdam, Netherlands
- ¹¹⁰ Department of Physics, Northern Illinois University, DeKalb IL, United States of America
- ¹¹¹ Budker Institute of Nuclear Physics, SB RAS, Novosibirsk, Russia
- ¹¹² Department of Physics, New York University, New York NY, United States of America
- ¹¹³ Ohio State University, Columbus OH, United States of America
- ¹¹⁴ Faculty of Science, Okayama University, Okayama, Japan
- ¹¹⁵ Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman OK, United States of America
- ¹¹⁶ Department of Physics, Oklahoma State University, Stillwater OK, United States of America
- ¹¹⁷ Palacký University, RCPTM, Olomouc, Czech Republic
- ¹¹⁸ Center for High Energy Physics, University of Oregon, Eugene OR, United States of America
- ¹¹⁹ LAL, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, Orsay, France
- ¹²⁰ Graduate School of Science, Osaka University, Osaka, Japan
- ¹²¹ Department of Physics, University of Oslo, Oslo, Norway
- ¹²² Department of Physics, Oxford University, Oxford, United Kingdom
- ¹²³ (a) INFN Sezione di Pavia; (b) Dipartimento di Fisica, Università di Pavia, Pavia, Italy
- ¹²⁴ Department of Physics, University of Pennsylvania, Philadelphia PA, United States of America
- ¹²⁵ National Research Centre "Kurchatov Institute" B.P.Konstantinov Petersburg Nuclear Physics Institute, St. Petersburg, Russia
- ¹²⁶ (a) INFN Sezione di Pisa; (b) Dipartimento di Fisica E. Fermi, Università di Pisa, Pisa, Italy
- ¹²⁷ Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh PA, United States of America
- ¹²⁸ (a) Laboratório de Instrumentação e Física Experimental de Partículas - LIP, Lisboa; (b) Faculdade de Ciências, Universidade de Lisboa, Lisboa; (c) Department of Physics, University of Coimbra, Coimbra; (d) Centro de Física Nuclear da Universidade de Lisboa, Lisboa; (e) Departamento de Física, Universidade do Minho, Braga; (f) Departamento de Física Teórica y del Cosmos, Universidad de Granada, Granada; (g) Dep Física and CEFITEC of Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Caparica, Portugal
- ¹²⁹ Institute of Physics, Academy of Sciences of the Czech Republic, Praha, Czech Republic
- ¹³⁰ Czech Technical University in Prague, Praha, Czech Republic
- ¹³¹ Charles University, Faculty of Mathematics and Physics, Prague, Czech Republic
- ¹³² State Research Center Institute for High Energy Physics (Protvino), NRC KI, Russia
- ¹³³ Particle Physics Department, Rutherford Appleton Laboratory, Didcot, United Kingdom
- ¹³⁴ (a) INFN Sezione di Roma; (b) Dipartimento di Fisica, Sapienza Università di Roma, Roma, Italy
- ¹³⁵ (a) INFN Sezione di Roma Tor Vergata; (b) Dipartimento di Fisica, Università di Roma Tor Vergata, Roma, Italy
- ¹³⁶ (a) INFN Sezione di Roma Tre; (b) Dipartimento di Matematica e Fisica, Università Roma Tre, Roma, Italy
- ¹³⁷ (a) Faculté des Sciences Ain Chock, Réseau Universitaire de Physique des Hautes Energies - Université Hassan II, Casablanca; (b) Centre National de l'Energie des Sciences Techniques Nucleaires, Rabat; (c) Faculté des Sciences Semlalia, Université Cadi Ayyad, LPHEA-Marrakech; (d) Faculté des

Sciences, Université Mohamed Premier and LPTPM, Oujda; ^(e) Faculté des sciences, Université Mohammed V, Rabat, Morocco

¹³⁸ DSM/IRFU (Institut de Recherches sur les Lois Fondamentales de l'Univers), CEA Saclay (Commissariat à l'Energie Atomique et aux Energies Alternatives), Gif-sur-Yvette, France

¹³⁹ Santa Cruz Institute for Particle Physics, University of California Santa Cruz, Santa Cruz CA, United States of America

¹⁴⁰ Department of Physics, University of Washington, Seattle WA, United States of America

¹⁴¹ Department of Physics and Astronomy, University of Sheffield, Sheffield, United Kingdom

¹⁴² Department of Physics, Shinshu University, Nagano, Japan

¹⁴³ Department Physik, Universität Siegen, Siegen, Germany

¹⁴⁴ Department of Physics, Simon Fraser University, Burnaby BC, Canada

¹⁴⁵ SLAC National Accelerator Laboratory, Stanford CA, United States of America

¹⁴⁶ ^(a) Faculty of Mathematics, Physics & Informatics, Comenius University, Bratislava; ^(b) Department of Subnuclear Physics, Institute of Experimental Physics of the Slovak Academy of Sciences, Kosice, Slovak Republic

¹⁴⁷ ^(a) Department of Physics, University of Cape Town, Cape Town; ^(b) Department of Physics, University of Johannesburg, Johannesburg; ^(c) School of Physics, University of the Witwatersrand, Johannesburg, South Africa

¹⁴⁸ ^(a) Department of Physics, Stockholm University; ^(b) The Oskar Klein Centre, Stockholm, Sweden

¹⁴⁹ Physics Department, Royal Institute of Technology, Stockholm, Sweden

¹⁵⁰ Departments of Physics & Astronomy and Chemistry, Stony Brook University, Stony Brook NY, United States of America

¹⁵¹ Department of Physics and Astronomy, University of Sussex, Brighton, United Kingdom

¹⁵² School of Physics, University of Sydney, Sydney, Australia

¹⁵³ Institute of Physics, Academia Sinica, Taipei, Taiwan

¹⁵⁴ Department of Physics, Technion: Israel Institute of Technology, Haifa, Israel

¹⁵⁵ Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv, Israel

¹⁵⁶ Department of Physics, Aristotle University of Thessaloniki, Thessaloniki, Greece

¹⁵⁷ International Center for Elementary Particle Physics and Department of Physics, The University of Tokyo, Tokyo, Japan

¹⁵⁸ Graduate School of Science and Technology, Tokyo Metropolitan University, Tokyo, Japan

¹⁵⁹ Department of Physics, Tokyo Institute of Technology, Tokyo, Japan

¹⁶⁰ Tomsk State University, Tomsk, Russia

¹⁶¹ Department of Physics, University of Toronto, Toronto ON, Canada

¹⁶² ^(a) INFN-TIFPA; ^(b) University of Trento, Trento, Italy

¹⁶³ ^(a) TRIUMF, Vancouver BC; ^(b) Department of Physics and Astronomy, York University, Toronto ON, Canada

¹⁶⁴ Faculty of Pure and Applied Sciences, and Center for Integrated Research in Fundamental Science and Engineering, University of Tsukuba, Tsukuba, Japan

¹⁶⁵ Department of Physics and Astronomy, Tufts University, Medford MA, United States of America

¹⁶⁶ Department of Physics and Astronomy, University of California Irvine, Irvine CA, United States of America

¹⁶⁷ ^(a) INFN Gruppo Collegato di Udine, Sezione di Trieste, Udine; ^(b) ICTP, Trieste; ^(c) Dipartimento di Chimica, Fisica e Ambiente, Università di Udine, Udine, Italy

¹⁶⁸ Department of Physics and Astronomy, University of Uppsala, Uppsala, Sweden

¹⁶⁹ Department of Physics, University of Illinois, Urbana IL, United States of America

¹⁷⁰ Instituto de Física Corpuscular (IFIC), Centro Mixto Universidad de Valencia - CSIC, Spain

- ¹⁷¹ Department of Physics, University of British Columbia, Vancouver BC, Canada
- ¹⁷² Department of Physics and Astronomy, University of Victoria, Victoria BC, Canada
- ¹⁷³ Department of Physics, University of Warwick, Coventry, United Kingdom
- ¹⁷⁴ Waseda University, Tokyo, Japan
- ¹⁷⁵ Department of Particle Physics, The Weizmann Institute of Science, Rehovot, Israel
- ¹⁷⁶ Department of Physics, University of Wisconsin, Madison WI, United States of America
- ¹⁷⁷ Fakultät für Physik und Astronomie, Julius-Maximilians-Universität, Würzburg, Germany
- ¹⁷⁸ Fakultät für Mathematik und Naturwissenschaften, Fachgruppe Physik, Bergische Universität Wuppertal, Wuppertal, Germany
- ¹⁷⁹ Department of Physics, Yale University, New Haven CT, United States of America
- ¹⁸⁰ Yerevan Physics Institute, Yerevan, Armenia
- ¹⁸¹ Centre de Calcul de l'Institut National de Physique Nucléaire et de Physique des Particules (IN2P3), Villeurbanne, France
- ¹⁸² Academia Sinica Grid Computing, Institute of Physics, Academia Sinica, Taipei, Taiwan
- ^a Also at Department of Physics, King's College London, London, United Kingdom
- ^b Also at Institute of Physics, Azerbaijan Academy of Sciences, Baku, Azerbaijan
- ^c Also at Novosibirsk State University, Novosibirsk, Russia
- ^d Also at TRIUMF, Vancouver BC, Canada
- ^e Also at Department of Physics & Astronomy, University of Louisville, Louisville, KY, United States of America
- ^f Also at Physics Department, An-Najah National University, Nablus, Palestine
- ^g Also at Department of Physics, California State University, Fresno CA, United States of America
- ^h Also at Department of Physics, University of Fribourg, Fribourg, Switzerland
- ⁱ Also at II Physikalisches Institut, Georg-August-Universität, Göttingen, Germany
- ^j Also at Departament de Física de la Universitat Autònoma de Barcelona, Barcelona, Spain
- ^k Also at Departamento de Física e Astronomia, Faculdade de Ciências, Universidade do Porto, Portugal
- ^l Also at Tomsk State University, Tomsk, and Moscow Institute of Physics and Technology State University, Dolgoprudny, Russia
- ^m Also at The Collaborative Innovation Center of Quantum Matter (CICQM), Beijing, China
- ⁿ Also at Università di Napoli Parthenope, Napoli, Italy
- ^o Also at Institute of Particle Physics (IPP), Canada
- ^p Also at Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest, Romania
- ^q Also at Department of Physics, St. Petersburg State Polytechnical University, St. Petersburg, Russia
- ^r Also at Borough of Manhattan Community College, City University of New York, New York City, United States of America
- ^s Also at Department of Financial and Management Engineering, University of the Aegean, Chios, Greece
- ^t Also at Centre for High Performance Computing, CSIR Campus, Rosebank, Cape Town, South Africa
- ^u Also at Louisiana Tech University, Ruston LA, United States of America
- ^v Also at Institutio Catalana de Recerca i Estudis Avancats, ICREA, Barcelona, Spain
- ^w Also at Department of Physics, The University of Michigan, Ann Arbor MI, United States of America
- ^x Also at LAL, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, Orsay, France
- ^y Also at Graduate School of Science, Osaka University, Osaka, Japan
- ^z Also at Fakultät für Mathematik und Physik, Albert-Ludwigs-Universität, Freiburg, Germany
- ^{aa} Also at Institute for Mathematics, Astrophysics and Particle Physics, Radboud University Nijmegen/Nikhef, Nijmegen, Netherlands
- ^{ab} Also at Department of Physics, The University of Texas at Austin, Austin TX, United States of

America

ac Also at Institute of Theoretical Physics, Ilia State University, Tbilisi, Georgia

ad Also at CERN, Geneva, Switzerland

ae Also at Georgian Technical University (GTU), Tbilisi, Georgia

af Also at Ochadai Academic Production, Ochanomizu University, Tokyo, Japan

ag Also at Manhattan College, New York NY, United States of America

ah Also at The City College of New York, New York NY, United States of America

ai Also at Departamento de Fisica Teorica y del Cosmos, Universidad de Granada, Granada, Portugal

aj Also at Department of Physics, California State University, Sacramento CA, United States of America

ak Also at Moscow Institute of Physics and Technology State University, Dolgoprudny, Russia

al Also at Departement de Physique Nucleaire et Corpusculaire, Université de Genève, Geneva, Switzerland

am Also at Institut de Física d'Altes Energies (IFAE), The Barcelona Institute of Science and Technology, Barcelona, Spain

an Also at School of Physics, Sun Yat-sen University, Guangzhou, China

ao Also at Institute for Nuclear Research and Nuclear Energy (INRNE) of the Bulgarian Academy of Sciences, Sofia, Bulgaria

ap Also at Faculty of Physics, M.V.Lomonosov Moscow State University, Moscow, Russia

aq Also at National Research Nuclear University MEPhI, Moscow, Russia

ar Also at Department of Physics, Stanford University, Stanford CA, United States of America

as Also at Institute for Particle and Nuclear Physics, Wigner Research Centre for Physics, Budapest, Hungary

at Also at Giresun University, Faculty of Engineering, Turkey

au Also at CPPM, Aix-Marseille Université and CNRS/IN2P3, Marseille, France

av Also at Department of Physics, Nanjing University, Jiangsu, China

aw Also at Institute of Physics, Academia Sinica, Taipei, Taiwan

ax Also at University of Malaya, Department of Physics, Kuala Lumpur, Malaysia

* Deceased