

# Photorespiration and nitrate assimilation: a major intersection between plant carbon and nitrogen

Arnold J. Bloom

Received: 19 August 2014 / Accepted: 27 October 2014 / Published online: 4 November 2014  
© Springer Science+Business Media Dordrecht 2014

**Abstract**  $C_3$  carbon fixation has a bad reputation, primarily because it is associated with photorespiration, a biochemical pathway thought to waste a substantial amount of the carbohydrate produced in a plant. This review presents evidence collected over nearly a century that (1) Rubisco when associated with  $Mn^{2+}$  generates additional reductant during photorespiration, (2) this reductant participates in the assimilation of nitrate into protein, and (3) this nitrate assimilation facilitates the use of a nitrogen source that other organisms tend to avoid. This phenomenon explains the continued dominance of  $C_3$  plants during the past 23 million years of low  $CO_2$  atmospheres as well as the decline in plant protein concentrations as atmospheric  $CO_2$  rises.

**Keywords** Photorespiration ·  $C_3$  carbon fixation · Nitrate assimilation · Photosynthesis · Plant evolution · Nitrogen sources

## Premise

Plants, by most accounts, convert less than 6 % of the incoming solar energy into useable chemical energy (Hall et al. 1999; Zhu et al. 2008). Efforts to improve this conversion rate have focused on the light-independent reactions of photosynthesis (e.g., Parry et al. 2013; Studer et al. 2014; Whitney et al. 2011; Zhu et al. 2010). “The light reactions are highly efficient, converting as much as 40–50 % of the captured solar energy into energy carriers. The dark reactions

are not developed for energy efficiency and it is here the energy is...lost” (Swedish Energy Agency 2003). In particular, Rubisco (ribulose-1,5-bisphosphate carboxylase/oxygenase; EC 4.1.1.39), the enzyme which catalyzes the first reaction of the  $C_3$  pathway and constitutes about half of the protein in leaves (Parry et al. 2003), has been identified as a target of opportunity.

## Competing reactions

Rubisco exhibits opposing tendencies in that it catalyzes two different chemical reactions: one reaction combines a five-carbon sugar RuBP (ribulose-1,5-bisphosphate) with  $CO_2$  (carboxylation), and the other reaction combines this same sugar with  $O_2$  (oxygenation).

- The carboxylation reaction of RuBP produces a six-carbon compound that quickly divides into two molecules of a three-carbon compound, PGA (3-phosphoglycerate), hence the name  $C_3$  carbon fixation. Six of these PGA molecules pass through an elaborate pathway that expends the energy of 18 ATP and 12 NADPH molecules, forms one molecule of fructose-6-phosphate, a six-carbon sugar, and regenerates six molecules of RuBP.
- The oxygenation reaction splits the RuBP into one molecule of a three-carbon PGA and one molecule of a two-carbon PG (2-phosphoglycolate), hence the name  $C_2$  pathway or, more commonly, photorespiration (Foyer et al. 2009). In total, photorespiration consumes 3.5 ATP and 2 NADPH per RuBP oxygenated and regenerated, but does not result in any net production of sugar (Bauwe et al. 2010; Tolbert 1994). Thus photorespiration seems to be largely a superfluous process, one thought to dissipate  $76.3 \text{ kcal mol}^{-1}$  as waste heat (Frank et al. 2000).

A. J. Bloom (✉)  
Department of Plant Sciences, University of California at Davis,  
Davis, USA  
e-mail: ajbloom@ucdavis.edu


The balance between  $C_3$  carbon fixation and photorespiration depends on the relative amounts of  $CO_2$  and  $O_2$  entering the active site of Rubisco and the specificity of the enzyme for each gas. Atmospheric concentrations of  $CO_2$  and  $O_2$  are currently 0.04 and 20.94 %, respectively, yielding a  $CO_2:O_2$  ratio of 0.0019. Gaseous  $CO_2$ , however, is much more soluble in water than  $O_2$ , and so the  $CO_2:O_2$  ratio near the chloroplast, the part of a cell where these reactions occur, is about 0.026 at 25 °C. Rubisco has about a 50-fold (cyanobacteria) to 100-fold (higher plants) greater specificity for  $CO_2$  than  $O_2$  (Galmes et al. 2005). Together, because of the relative concentrations of and specificity for  $CO_2$  over  $O_2$ , Rubisco catalyzes about two to three cycles of  $C_3$  carbon fixation for every cycle of photorespiration under current atmospheres (Sharkey 1988). Conditions that inhibit photorespiration—namely, high  $CO_2$ , or low  $O_2$  atmospheric concentrations—stimulate carbon fixation in the short term by about 35 %.

Temperature influences the balance between  $C_3$  carbon fixation and photorespiration in two ways. First, as temperature rises, the solubility of  $CO_2$  in water decreases more than the solubility of  $O_2$ , resulting in a lower  $CO_2:O_2$  ratio. Second, the enzymatic properties of Rubisco shift with increasing temperature, stimulating the reaction with  $O_2$  to a greater degree than the one with  $CO_2$ . Warmer temperatures, therefore, favor photorespiration over  $C_3$  carbon fixation, and photosynthetic conversion of absorbed light into sugars becomes less efficient (Ehleringer et al. 1997). Based on the temperature response of Rubisco carboxylation and oxygenation,  $C_4$  plants should be more competitive in regions where the mean monthly air temperature exceeds 22 °C (Collatz et al. 1998).

Overall, Rubisco seems a vestige of the high  $CO_2$  and low  $O_2$  atmospheres under which plants first evolved (Wingler et al. 2000). To compensate for the shortcomings of Rubisco, some plants employ  $CO_2$  pumping mechanisms such as  $C_4$  carbon fixation that elevate  $CO_2$  concentrations at the active site of the enzyme. The  $C_4$  pathway is one of the most convergent evolutionary adaptations in life with at least 66 independent origins (Sage et al. 2012). Extensive efforts are underway to emulate Mother Nature and transfer the  $C_4$  pathway into rice and other  $C_3$  crops (von Caemmerer et al. 2012).

Several observations, however, are inconsistent with the presumption that Rubisco is poorly suited to modern times.


- Earth's atmosphere has contained relatively low  $CO_2$  concentrations (lower than 0.04 %) for the past 23 million years (Fig. 1). During this period, the plant kingdom experienced major changes including the diversification of modern graminoids, especially grasses and sedges, and the appearance of many new  $C_4$  species, especially



**Fig. 1** A reconstruction of atmospheric  $CO_2$  concentrations based on boron isotope ratios of ancient planktonic foraminifer shells. (Data from Pearson and Palmer 2000)

when  $CO_2$  concentrations fell below 0.02 % (Sage et al. 2012). In a relatively short period of time (6 or 7 million years) (Osborne and Beerling 2006), the kinetics of Rubisco diverged between  $C_3$  and  $C_4$  plants (Studer et al. 2014). Rubisco in  $C_4$  plants operates under elevated  $CO_2$  conditions, and so the  $C_4$  enzyme has traded a lower specificity for  $CO_2$  relative to  $O_2$  ( $S_{c/o}$ ) for a higher catalytic efficiency ( $k_{cat}^c$ ) (Galmes et al. 2005; Sage 2002). Surprisingly, the kinetic properties of Rubisco do not differ greatly among higher  $C_3$  plants (Kane et al. 1994; Tcherkez et al. 2006). Thus, the kinetic properties of Rubisco were able to change when a species adopted the  $C_4$  pathway, but such changes were not warranted in  $C_3$  plants because Rubisco may already be “nearly perfectly optimized” for  $C_3$  carbon fixation (Tcherkez et al. 2006).

- Despite 23 million years of low atmospheric  $CO_2$  concentrations, 96 % of plant species depend solely on the  $C_3$  carbon fixation pathway (Sage et al. 1999).  $C_3$  species account for over 94 % of the Earth's biomass (Still et al. 2003). Species using other carbon fixation pathways are dominant only in hot and dry environments.
- The response of  $C_3$  species to elevated  $CO_2$  atmospheres is highly variable and often depends on plant N status (Cavagnaro et al. 2011; Duval et al. 2012; Finzi et al. 2007; Norby et al. 2010; Reich et al. 2006). Initially, elevated  $CO_2$  stimulates biomass accumulation by about 35 % (Fig. 2). This stimulation, however, tends to abate upon longer exposures in conjunction with a decline in plant protein concentrations (Cotrufo et al. 1998; Long et al. 2004).



**Fig. 2** Differences in biomass between elevated ( $\approx 567$  ppm) and ambient ( $\approx 365$  ppm) atmospheric  $\text{CO}_2$  after years of treatment. Shown are the data from seven different studies using the designated types of plants. (Data from Dukes et al. 2005; Kimball et al. 2007; Korner 2006; Norby et al. 2010; Rasse et al. 2005; Talhelm et al. 2014)


Explanations for the decline in plant protein concentrations at elevated  $\text{CO}_2$  include: (a) plants under elevated  $\text{CO}_2$  grow larger, diluting the protein within their tissues (Ellsworth et al. 2004; Taub and Wang 2008); (b) carbohydrates accumulate within leaves, down-regulating the amount of the most prevalent protein Rubisco (Long et al. 2004); (c) carbon enrichment of the rhizosphere leads to progressively greater limitations in the soil N available to plants (Reich et al. 2006); and (d) elevated  $\text{CO}_2$  directly inhibits plant N metabolism, especially the assimilation of  $\text{NO}_3^-$  into proteins in shoots of  $\text{C}_3$  plants (Bloom et al. 2012b). Recently, several independent meta-analyses conclude that this last explanation is the one most consistent with observations from hundreds of studies (Cheng et al. 2012; Myers et al. 2014; Pleijel and Uddling 2012).

### $\text{CO}_2$ inhibits $\text{NO}_3^-$ assimilation

Many independent methods for estimating  $\text{NO}_3^-$  assimilation confirm that elevated  $\text{CO}_2$  inhibits shoot  $\text{NO}_3^-$  assimilation in  $\text{C}_3$  plants. These methods include:


- 1  $^{15}\text{N}$ -labeling. Plants grown on  $\text{NO}_3^-$  containing N isotopes at natural abundance levels ( $\approx 0.366\%$   $^{15}\text{N}$ ) were exposed to a pulse of  $\text{NO}_3^-$  that was heavily enriched in  $^{15}\text{N}$ . The difference between the  $^{15}\text{N}$

- enrichment of total N and that of free  $\text{NO}_3^-$  provided an estimate of  $^{15}\text{N}$ - $\text{NO}_3^-$  assimilation, which decreased under  $\text{CO}_2$  enrichment (Bloom et al. 2010).
- 2  $^{14}\text{N}$ -labeling. Plants grown on 99.9 % enriched  $^{15}\text{N}$ - $\text{NO}_3^-$  were exposed to a pulse of  $\text{NO}_3^-$  containing N isotopes at natural abundance levels ( $\approx 0.366\%$   $^{15}\text{N}$ ); the difference between the  $^{14}\text{N}$  enrichment of total N and that of free  $\text{NO}_3^-$  provided an estimate of  $^{14}\text{N}$ - $\text{NO}_3^-$  assimilation, which decreased under  $\text{CO}_2$  enrichment (Bloom et al. 2010).
- 3 *Organic N accumulation*. Accumulation of organic N was followed in plants receiving  $\text{NO}_3^-$  as a sole N source, and this accumulation decreased under  $\text{CO}_2$  enrichment (Aranjuelo et al. 2013; Bloom et al. 2010; Lekshmy et al. 2013; Pleijel and Uddling 2012; Rachmilevitch et al. 2004).
- 4  *$\text{NO}_3^-$  depletion from a medium*. The decline of  $\text{NO}_3^-$  concentrations in a nutrient solution was monitored to calculate net plant  $\text{NO}_3^-$  absorption. The difference between this  $\text{NO}_3^-$  absorption and the accumulation of free  $\text{NO}_3^-$  within plant tissues estimated plant  $\text{NO}_3^-$  assimilation, which decreased under  $\text{CO}_2$  enrichment (Bloom et al. 2010; Rachmilevitch et al. 2004).
- 5 *Plant growth*.  $\text{C}_3$  species received either  $\text{NO}_3^-$  or  $\text{NH}_4^+$  as their sole N source.  $\text{CO}_2$  enrichment decreased growth of plants receiving  $\text{NO}_3^-$  (Fig. 3) but increased growth of those receiving  $\text{NH}_4^+$  (Bloom et al. 2012b, 2002; Carlisle et al. 2012).
- 6 *Isotopic discrimination by  $\text{NO}_3^-$  reductase*. Plants were grown under  $\text{NO}_3^-$  containing N isotopes at natural abundance levels ( $\approx 0.366\%$   $^{15}\text{N}$ ). Under  $\text{CO}_2$  enrichment, plant tissues became less enriched in  $^{15}\text{N}$ -organic N compounds presumably because (a)  $\text{CO}_2$  inhibited shoot  $\text{NO}_3^-$  assimilation, (b)  $\text{NO}_3^-$  availability became less limiting to assimilation, (c)  $\text{NO}_3^-$  reductase discriminated more against  $^{15}\text{N}$ - $\text{NO}_3^-$ , and (d) shoots assimilated relatively less  $^{15}\text{N}$ - $\text{NO}_3^-$  (Bloom et al. 2010, 2014).
- 7  $\Delta A_Q$ . Assimilatory quotient ( $A_Q$ ), the ratio of net  $\text{CO}_2$  consumption to net  $\text{O}_2$  evolution from shoots was measured in a plant receiving  $\text{NH}_4^+$  or  $\text{NO}_3^-$  as its sole N source (Fig. 4);  $A_Q$  decreased as  $\text{NO}_3^-$  assimilation increased because additional electrons generated from the light-dependent reactions of photosynthesis were transferred first to  $\text{NO}_3^-$  and then to  $\text{NO}_2^-$ . This stimulated net  $\text{O}_2$  evolution, but had little effect on  $\text{CO}_2$  consumption; therefore, the change in  $A_Q$  when a plant received  $\text{NH}_4^+$  instead of  $\text{NO}_3^-$  ( $\Delta A_Q$ ) provided an estimate of shoot  $\text{NO}_3^-$  assimilation (Bloom et al. 1989, 2002; Cen et al. 2001; Cramer and Myers 1948; Rachmilevitch et al. 2004; Van Niel et al. 1953; Warburg and Negelein




**Fig. 3** Relative growth rate in  $\text{g g}^{-1} \text{d}^{-1}$  of (A) loblolly pine *Pinus taeda* and (B) sweet gum *Liquidambar styraciflua* receiving  $\text{NO}_3^-$  nutrition in controlled environment chambers at subambient  $\text{CO}_2$  ( $310 \mu\text{mol mol}^{-1}$ , the level of about 50 years ago), ambient  $\text{CO}_2$  ( $400 \mu\text{mol mol}^{-1}$ , current level), or elevated  $\text{CO}_2$  ( $720 \mu\text{mol mol}^{-1}$ , the level anticipated in about 50 years).  $\text{CO}_2$  concentration had no

significant effect on the growth of plants receiving  $\text{NH}_4^+$  nutrition (data not shown). Time is in days after transplanting to a hydroponic solution. Shown are the predicted values and standard errors from mixed linear models with repeated measures on 6–10 individual plants. (Bloom et al. 2012b)



**Fig. 4** Shoot AQ (net  $\text{CO}_2$  consumed/net  $\text{O}_2$  evolved) as a function of internal  $\text{CO}_2$  concentrations ( $C_i$ ) for the 9  $C_3$  species in Fig. 5 when they received  $\text{NH}_4^+$  or  $\text{NO}_3^-$  as a sole N source (mean  $\pm$  SE; solid  $\pm$  shaded area). (Bloom, unpublished data)




**Fig. 5** Shoot  $\text{NO}_3^-$  assimilation as a function of shoot internal  $\text{CO}_2$  concentration ( $C_i$ ) for 9  $C_3$  species. Shoot  $\text{NO}_3^-$  assimilation is assessed by  $\Delta\text{AQ}$  (change in the ratio of shoot  $\text{CO}_2$  consumption to  $\text{O}_2$  evolution with a shift from  $\text{NO}_3^-$  to  $\text{NH}_4^+$  nutrition). (Bloom et al. 2012b; Searles and Bloom 2003)

1920). In nine taxonomically diverse  $C_3$  species,  $\Delta\text{AQ}$  decreased as shoot internal  $\text{CO}_2$  increased (Fig. 5).


- 8 *O<sub>2</sub> consumption.* Shoot  $\text{O}_2$  consumption in the light was estimated from the difference between gross  $\text{O}_2$  evolution via chlorophyll fluorescence and net  $\text{O}_2$  evolution via an  $\text{O}_2$  analyzer (Fig. 6). At ambient  $\text{CO}_2$ ,  $\text{O}_2$  consumption was lower when wheat plants received  $\text{NO}_3^-$  rather than  $\text{NH}_4^+$  because  $\text{NO}_3^-$  and

$\text{NO}_2^-$  were serving as electron acceptors. At elevated  $\text{CO}_2$ ,  $\text{O}_2$  consumption was not significantly different under the two N sources presumably because  $\text{NO}_3^-$  assimilation was negligible.

- 9 *Altered  $\text{NO}_3^-$  reductase capacity.* Shoot  $\text{CO}_2$  and  $\text{O}_2$  fluxes at ambient and elevated  $\text{CO}_2$  were contrasted between stages of plant development or genotypes that have greatly different  $\text{NO}_3^-$  reductase activities



**Fig. 6** Shoot  $O_2$  consumption in the light (gross  $O_2$ –net  $O_2$ ) as a function of  $C_i$  for wheat receiving  $NH_4^+$  or  $NO_3^-$  as a sole N source. Shown are the mean  $\pm$  SE for 5–7 replicates per treatment. (Cousins and Bloom 2004)



**Fig. 7**  $NO_3^-$  reductase activity ( $\mu\text{mol}$  of  $NO_2^-$  generated per g fresh mass per min) as a function of plant age (d) in leaves of a wild-type *A. thaliana* cv. Columbia (WT), a transgenic line harboring the chimeric gene *Lhch1\*3::Nia1\*2* (OE), and a genotype (*nia1 nia2*) with mutations in both structural genes for  $NO_3^-$  reductase (Mut). Because  $NO_3^-$  reductase is regulated through phosphorylation, leaf tissue was assayed under conditions that either dephosphorylated the enzyme (fully activated) or did not change its phosphorylation (in vivo). Shown are the mean  $\pm$  SE ( $n = 5$ –8 plants). (Rachmilevitch et al. 2004)

in situ. In particular, we contrasted 36- versus 48-d-old wild-type Arabidopsis, Arabidopsis  $NO_3^-$  reductase knockout mutants versus transgenic Arabidopsis overexpressing  $NO_3^-$  reductase (Fig. 7), and  $NO_3^-$

reductase-deficient barley mutants versus wild-type barley.  $\Delta AQ$  (change in the ratio of net  $CO_2$  consumption to net  $O_2$  evolution when a plant received  $NH_4^+$  instead of  $NO_3^-$ ) differed between these stages of development and genotypes under ambient  $CO_2$ , but not under elevated  $CO_2$  (Fig. 8). This indicates that none of the stages of development or genotypes were assimilating  $NO_3^-$  under elevated  $CO_2$  (Bloom et al. 1989; Rachmilevitch et al. 2004).


- 10  $NO_3^-$  reductase activity. Maximum in vitro  $NO_3^-$  reductase activity generally declined under  $CO_2$  enrichment (Lekshmy et al. 2013; Matt et al. 2001). Presumably, this reflected slower  $NO_3^-$  assimilation under  $CO_2$  enrichment.

### Physiological mechanisms

Three physiological mechanisms may be responsible for  $CO_2$  inhibition of shoot  $NO_3^-$  assimilation (Bloom et al. 2010).


- One mechanism is that elevated  $CO_2$  inhibits nitrite ( $NO_2^-$ ) transport into chloroplasts (Fig. 9). A chloroplast  $NO_2^-$  transporter from higher plants has only recently been identified (Maeda et al. 2014), and so the nature of this inhibition has yet to be determined. Nevertheless, this mechanism can be independent of photosynthesis and, thus, is probably responsible for  $CO_2$  inhibition of shoot  $NO_3^-$  assimilation in Arabidopsis and wheat during the nighttime (Rubio-Asensio, Rachmilevitch, and Bloom, unpublished data).
- Another mechanism is that processes in the chloroplast stroma compete for reduced ferredoxin ( $Fd_r$ ). FNR (ferredoxin-NADP reductase) has a higher affinity for  $Fd_r$  than NiR (nitrite reductase) (Knaff 1996), and so  $NO_3^-$  assimilation proceeds only if the availability of  $Fd_r$  exceeds that needed for NADPH formation (Backhausen et al. 2000; Robinson 1987). For most plants, this occurs when  $CO_2$  availability limits  $C_3$  carbon fixation (Bloom et al. 2010).
- A third mechanism involves photorespiration. Multiple lines of evidence link photorespiration with shoot  $NO_3^-$  assimilation in  $C_3$  plants. (a) Photorespiration stimulates the export of malate from chloroplasts (Backhausen et al. 1998; Taniguchi and Miyake 2012; Voss et al. 2013); this malate in the cytoplasm generates NADH (Igamberdiev et al. 2001; Taniguchi and Miyake 2012) that powers the first step of  $NO_3^-$  assimilation, the reduction of  $NO_3^-$  to  $NO_2^-$  (Quesada et al. 2000; Rathnam 1978; Robinson 1987). (b) Conditions that decrease photorespiration—namely, elevated  $CO_2$  and low  $O_2$ —decrease shoot  $NO_3^-$  reduction (Bloom et al. 2010; Rachmilevitch et al.





**Fig. 8** Changes in assimilatory quotient with the shift from  $\text{NO}_3^-$  to  $\text{NH}_4^+$  ( $\Delta\text{AQ}$ ) as a function of photosynthetic  $PFD$  (photon flux density) from shoots of *A. thaliana* cv. Columbia. (A) 36-day-old wild-type plants, (B) 48-d-old wild-type plants, (C) genotype with null mutations, and (D) overexpressing line. The plants were grown

under ambient  $\text{CO}_2$  (360  $\mu\text{mol mol}^{-1}$ ) and measured under ambient  $\text{CO}_2$  and  $\text{O}_2$  (360  $\mu\text{mol mol}^{-1}$   $\text{CO}_2$  and 21%  $\text{O}_2$ ; circles), elevated  $\text{CO}_2$  (720  $\mu\text{mol mol}^{-1}$   $\text{CO}_2$  and 21%  $\text{O}_2$ ; triangles), or low  $\text{O}_2$  (360  $\mu\text{mol mol}^{-1}$   $\text{CO}_2$  and 2%  $\text{O}_2$ ; squares). Shown are the mean  $\pm$  SE,  $n = 5\text{--}8$  plants. (Rachmilevitch et al. 2004)



**Fig. 9** Net  $\text{NO}_2^-$  uptake ( $\mu\text{mol mg}^{-1}$  chlorophyll  $\text{min}^{-1}$ ) by isolated chloroplasts as a function of  $\text{NO}_2^-$  concentration when the medium contained 0 (light symbols) or 300 (dark symbols)  $\mu\text{M}$   $\text{HCO}_3^-$ . Shown are the mean  $\pm$  SE ( $n = 3$ ) for wheat (circles) and pea (inverted triangles). (Bloom et al. 2002)


2004). (c) Mutants that alter malate transport or metabolism also alter both photorespiration and  $\text{NO}_3^-$  assimilation (Dutilleul et al. 2005; Schneiderei et al. 2006).

The first carboxylation reaction in the  $\text{C}_4$  carbon fixation pathway, by contrast, generates ample amounts of malate and NADH in the cytoplasm of mesophyll cells. This explains the  $\text{CO}_2$  independence of shoot  $\text{NO}_3^-$  assimilation in  $\text{C}_4$  plants (Bloom et al. 2010, 2012b).

### The Rubisco complex

Information about the biochemistry of RuBP oxygenation is limited. The stroma of the chloroplast contains similar amounts of  $\text{Mg}^{2+}$  (2 mM, Ishijima et al. 2003) and  $\text{Mn}^{2+}$  (2 mM, Burnell 1988; Robinson and Gibbs 1982). Rubisco may form a complex with either  $\text{Mg}^{2+}$  or  $\text{Mn}^{2+}$  (Pierce and Reddy 1986), but the affinity of Rubisco for  $\text{Mn}^{2+}$  is five times greater than that for  $\text{Mg}^{2+}$  (Christeller 1981). The stoichiometry of  $\text{CO}_2$  trapping (Miziorko and Sealy 1980) and  $^{31}\text{P}$  and  $^{13}\text{C}$  NMR measurements (Pierce and Reddy 1986) indicate that  $\text{Mn}^{2+}$  and  $\text{Mg}^{2+}$  share a common binding site in the large subunit of Rubisco. Nearly all of the biochemistry of Rubisco has been conducted in the presence of  $\text{Mg}^{2+}$  and in the absence of  $\text{Mn}^{2+}$  because Rubisco when associated with  $\text{Mn}^{2+}$  strongly favors RuBP oxygenation, whereas Rubisco when associated with  $\text{Mg}^{2+}$  favors RuBP carboxylation (Chen and Spreitzer 1992; Christeller and Laing 1979; Houtz et al. 1988; Jordan and Ogren 1981; Raghavendra et al. 1981; Wildner and Henkel 1979).

$\text{Mg}^{2+}$  has a pair of electrons in its outer shell, whereas  $\text{Mn}^{2+}$  has up to five unpaired electrons and thus participates more readily in redox reactions. In specific,  $\text{Mn}^{2+}$  participates in the catalytic process of RuBP oxygenation (Miziorko and Sealy 1984) during which it becomes excited and transfers an electron with every turnover (Lilley et al. 2003). One possibility is that  $\text{Mn}^{2+}$  transfers electrons to  $\text{NADP}^+$  (Fig. 11). The resultant NADPH activates Rubisco (Laing and Christeller 1976) and then converts OAA to malate for export to the cytoplasm. This malate in the cytoplasm generates NADH to convert  $\text{NO}_3^-$  to  $\text{NO}_2^-$ .



**Fig. 10** One possible scenario for the intermediates formed during RuBP oxygenation (Chen and Spreitzer 1992; Cleland et al. 1998; Lilley et al. 2003; Oliva et al. 2001; Tapia and Andrés 1992; Tcherkez et al. 2006)

Several additional observations are consistent with this hypothesis. RuBP oxygenation releases  $76.3 \text{ kcal mol}^{-1}$  (Frank et al. 2000), substantially more than the  $52 \text{ kcal mol}^{-1}$  required to reduce  $\text{NADP}^+$  to NADPH (Taiz and Zeiger 2010). NADPH complexes strongly with Rubisco and activates the enzyme, but only when  $\text{CO}_2$  and  $\text{Mg}^{2+}$  are present in suboptimal concentrations (Chollet and Anderson 1976; Chu and Bassham 1974; Matsumura et al. 2012; McCurry et al. 1981). NADPH binds to the catalytic site of Rubisco through metal-coordinated water molecules (Matsumura et al. 2012).

If Rubisco generates NADPH during RuBP oxygenation,  $\text{C}_3$  carbon fixation is more efficient than previously thought, and both  $\text{C}_3$  and  $\text{C}_4$  carbon fixation at moderate temperatures will expend the equivalent of about 11 ATPs per  $\text{CO}_2$  fixed. Indeed, the quantum yield of photosynthesis in an ambient  $\text{CO}_2$  and  $\text{O}_2$  atmosphere does not differ significantly between  $\text{C}_3$  and  $\text{C}_4$  species at temperatures between  $25^\circ$  and  $30^\circ \text{C}$  (Skillman 2008). Only under hotter and drier conditions does  $\text{C}_4$  carbon fixation become more efficient than  $\text{C}_3$  fixation. Therefore,  $\text{C}_3$  species continue to dominate in most locations.

### Why is photorespiration still prevalent?

Several phenomena are responsible for the persistence of photorespiration through 23 million years of low atmospheric  $\text{CO}_2$  concentrations.

- Rubisco oxygenation is inseparable from Rubisco carboxylation (Moroney et al. 2013; Tcherkez et al. 2006). Rubisco catalyzes the carboxylation reaction through


stabilizing the formation of the enediol conformation of RuBP (Fig. 10). This conformation, however, can react with either  $\text{CO}_2$  or  $\text{O}_2$ . The specificity of Rubisco for  $\text{CO}_2$  over  $\text{O}_2$  derives from stabilizing the six-carbon intermediate before it is cleaved to form two molecules of PGA. Consequently, any mutation that increases the specificity of Rubisco for  $\text{CO}_2$  over  $\text{O}_2$  slows the carboxylation reaction.

- Photorespiration maintains redox homeostasis within plant cells (Scheibe and Dietz 2012). Photosynthesis generates highly reactive compounds as it captures solar energy and converts it into energy-rich, but stable compounds such as carbohydrates. Metabolic pathways, especially under stressful conditions, may become unbalanced, and dangerous compounds such as reactive oxygen species (ROS) may accumulate (Voss et al. 2013). Photorespiration can dissipate many of these potentially dangerous compounds.
- Photorespiration produces  $\text{H}_2\text{O}_2$  in the peroxisome and thus serves as a mechanism for rapidly transferring a signal of photosynthesis to the entire plant cell (Foyer et al. 2009). This signal is involved in photoperiod detection and pathogen defense as well as responses to abiotic stress.
- Photorespiration serves as a mechanism for plants to use  $\text{NO}_3^-$  as a nitrogen source without diverting energy from  $\text{CO}_2$  fixation. The following provides details about this phenomenon.

### Nitrate as a nitrogen source

The element nitrogen is a constituent of many organic compounds including all amino acids and nucleic acids. As

**Fig. 11** Response of net CO<sub>2</sub> consumption (*left panels*) and net O<sub>2</sub> evolution (*right panels*) to photosynthetic photon flux density (PPFD) in wheat (*upper panels*) and tomato (*lower panels*) leaves when the plants received NH<sub>4</sub><sup>+</sup> (blue) or NO<sub>3</sub><sup>-</sup> (red) nutrition and were exposed to an atmosphere containing 720 (dark colors) or 360 (light colors) μmol mol<sup>-1</sup> CO<sub>2</sub>. Shown are the mean ± SE for six wheat plants and 6–9 tomato plants per treatment. Notice that in both species, CO<sub>2</sub> fluxes do not differ with N source, and that O<sub>2</sub> fluxes are faster under NO<sub>3</sub><sup>-</sup> nutrition than NH<sub>4</sub><sup>+</sup> nutrition, but only at higher light levels and 360 μmol mol<sup>-1</sup> CO<sub>2</sub>. (Cousins and Bloom 2004; Searles and Bloom 2003)



such, plants require a greater amount of nitrogen than any other mineral element, and its availability generally limits the productivity of natural and agricultural ecosystems (Epstein and Bloom 2005). Conversions among various nitrogen compounds are among the most energy-intensive reactions in life. Consider that plants are generally between 1 and 2 % organic nitrogen on a percentage dry weight basis, but that the conversion of NO<sub>3</sub><sup>-</sup> into organic nitrogen expends about 25 % of the total energy in shoots (Bloom et al. 1989) and roots (Bloom et al. 1992). These processes expend the energy equivalent of 12 ATP per NO<sub>3</sub><sup>-</sup> assimilated, whereas most biochemical reactions expend the energy equivalent of one or perhaps two ATP.


Most organisms prefer higher energy forms of nitrogen such as NH<sub>4</sub><sup>+</sup> or amino acids. Phytoplankton (Dortch 1990), fungi (Hodge et al. 2010), cyanobacteria (Ohashi et al. 2011), and bacteria (Luque-Almagro et al. 2011) absorb and assimilate NO<sub>3</sub><sup>-</sup> only in the absence of NH<sub>4</sub><sup>+</sup>. In many soils, microorganisms quickly absorb NH<sub>4</sub><sup>+</sup> and either assimilate it into amino acids or nitrify it to NO<sub>3</sub><sup>-</sup>. NH<sub>4</sub><sup>+</sup> also becomes adsorbed on the soil cation exchange matrix. Because soil microorganisms often ignore NO<sub>3</sub><sup>-</sup> and because NO<sub>3</sub><sup>-</sup> as an anion moves relatively freely

through the soil, NO<sub>3</sub><sup>-</sup> is often the predominant form of nitrogen available to plants (Epstein and Bloom 2005).

Nitrogen nutrition, NH<sub>4</sub><sup>+</sup> versus NO<sub>3</sub><sup>-</sup>, neither influences net CO<sub>2</sub> consumption (Fig. 11) nor cyclic electron flow around photosystem I at low light levels (Walker et al. 2014). This is consistent with the lack of competition for reductant between CO<sub>2</sub> fixation and NO<sub>3</sub><sup>-</sup> assimilation (Robinson 1988) because, as discussed previously, FNR has a higher affinity for Fd<sub>r</sub> than NiR. At high light levels and ambient CO<sub>2</sub> and O<sub>2</sub> concentrations, net O<sub>2</sub> evolution is faster (Figs. 11 and 12) and cyclic electron flow around photosystem I is higher (Walker et al. 2014) when plants receive NO<sub>3</sub><sup>-</sup> rather than NH<sub>4</sub><sup>+</sup> as a nitrogen source. Presumably, plants use reductant generated from the light-dependent reactions rather than mitochondrial respiration to assimilate NO<sub>3</sub><sup>-</sup> when CO<sub>2</sub> concentration limits CO<sub>2</sub> fixation.

When factors other than CO<sub>2</sub> limit CO<sub>2</sub> fixation, plants may delay assimilating the NO<sub>3</sub><sup>-</sup> that they have absorbed. Free NO<sub>3</sub><sup>-</sup> may comprise as much as 60 % of the total nitrogen in a plant (Maynard et al. 1976). This NO<sub>3</sub><sup>-</sup> serves as a metabolically benign osmoticant that balances other ions such as potassium in plant tissues and helps to





**Fig. 12** Responses of wheat shoots (mean  $\pm$  SE,  $n = 6$ ) to photosynthetic photon flux density (PPFD). **(A)** Changes in assimilatory quotient ( $AQ = \text{net CO}_2 \text{ consumed}/\text{net O}_2 \text{ evolved}$ ) with the shift from  $\text{NO}_3^-$  to  $\text{NH}_4^+$  as a N source. **(B)** Changes in the gross  $\text{O}_2$  consumed (gross  $\text{O}_2$  evolved minus net  $\text{O}_2$  evolved) with the shift from  $\text{NO}_3^-$  to  $\text{NH}_4^+$  as a N source. As light levels increased and  $360 \mu\text{mol mol}^{-1}$   $\text{CO}_2$  limited carbon fixation, exposure to  $\text{NO}_3^-$  stimulated the light-dependent reactions of photosynthesis to split water, evolve oxygen, and transfer electrons to  $\text{NO}_3^-$  and  $\text{NO}_2^-$  rather than to  $\text{CO}_2$ , and decreased gross  $\text{O}_2$  consumption (Cousins and Bloom 2004)

maintain a favorable cellular water status (Bloom et al. 2012a; Burns et al. 2010; Hanson and Hitz 1983; McIntyre 1997; Veen and Kleinendorst 1986).

In summary, the linkage between photorespiration and  $\text{NO}_3^-$  assimilation provides higher plants with a relatively abundant nitrogen source that other organisms cannot afford to use, but that  $\text{C}_3$  plants can use with little additional cost. Yes, photorespiration may sacrifice 20–35 % of  $\text{CO}_2$  fixation, but plants that are dependent on  $\text{NO}_3^-$  as a nitrogen source are spared the expense of either devoting 25 % of their photosynthate to  $\text{NO}_3^-$  assimilation or suffering protein deprivation. Apparently, over the last 23 million years, 96 % of higher plant species have adapted to this tradeoff.

**Acknowledgments** This work was funded in part by NSF IOS-13-58675. We thank J. Clark Lagarias and George H. Lorimer for their insights about redox reactions and the reviewers for their suggestions.

**Conflicts of interest** The author has no conflicts of interest with regards to this research.

## References

- Aranjuelo I, Cabrerizo PM, Arrese-Igor C, Aparicio-Tejo PM (2013) Pea plant responsiveness under elevated  $[\text{CO}_2]$  is conditioned by the N source ( $\text{N}_2$  fixation versus  $\text{NO}_3^-$  fertilization). *Environ Exp Bot*. doi:10.1016/j.envexpbot.2013.06.002
- Backhausen JE et al (1998) Transgenic potato plants with altered expression levels of chloroplast NADP-malate dehydrogenase: interactions between photosynthetic electron transport and malate metabolism in leaves and in isolated intact chloroplasts. *Planta* 207:105–114
- Backhausen JE, Kitzmann C, Horton P, Scheibe R (2000) Electron acceptors in isolated intact spinach chloroplasts act hierarchically to prevent over-reduction and competition for electrons. *Photosynth Res* 64:1–13
- Bauwe H, Hagemann M, Fernie AR (2010) Photorespiration: players, partners and origin. *Trends Plant Sci* 15:330–336. doi:10.1016/j.tplants.2010.03.006
- Bloom AJ, Caldwell RM, Finazzo J, Warner RL, Weissbart J (1989) Oxygen and carbon dioxide fluxes from barley shoots depend on nitrate assimilation. *Plant Physiol* 91:352–356
- Bloom AJ, Sukrapanna SS, Warner RL (1992) Root respiration associated with ammonium and nitrate absorption and assimilation by barley. *Plant Physiol* 99:1294–1301
- Bloom AJ, Smart DR, Nguyen DT, Searles PS (2002) Nitrogen assimilation and growth of wheat under elevated carbon dioxide. *Proc Natl Acad Sci USA* 99:1730–1735
- Bloom AJ, Burger M, Asensio JSR, Cousins AB (2010) Carbon dioxide enrichment inhibits nitrate assimilation in wheat and *Arabidopsis*. *Science* 328:899–903. doi:10.1126/science.1186440
- Bloom AJ, Randall L, Taylor AR, Silk WK (2012a) Deposition of ammonium and nitrate in the roots of maize seedlings supplied with different nitrogen salts. *J Exp Bot* 63:1997–2006. doi:10.1093/jxb/err410
- Bloom AJ, Rubio-Asensio JS, Randall L, Rachmilevitch S, Cousins AB, Carlisle EA (2012b)  $\text{CO}_2$  enrichment inhibits shoot nitrate assimilation in  $\text{C}_3$  but not  $\text{C}_4$  plants and slows growth under nitrate in  $\text{C}_3$  plants. *Ecology* 93:355–367
- Bloom AJ, Burger M, Kimball BA, Pinter PJ (2014) Nitrate assimilation is inhibited by elevated  $\text{CO}_2$  in field-grown wheat. *Nat Clim Change* 4:477–480. doi:10.1038/nclimate2183
- Burnell JN (1988) The biochemistry of manganese in plants. In: Graham RD, Hannam RJ, Uren NC (eds) Manganese in soils and plants. Kluwer Academic, Dordrecht, pp 125–137
- Burns IG, Zhang KF, Turner MK, Edmondson R (2010) Iso-osmotic regulation of nitrate accumulation in lettuce. *J Plant Nutr* 34:283–313. doi:10.1080/01904167.2011.533328
- Carlisle E, Myers SS, Raboy V, Bloom AJ (2012) The effects of inorganic nitrogen form and  $\text{CO}_2$  concentration on wheat yield and nutrient accumulation and distribution. *Front Plant Sci* 3:195. doi:10.3389/fpls.2012.00195
- Cavagnaro TR, Gleadow RM, Miller RE (2011) Plant nutrient acquisition and utilisation in a high carbon dioxide world. *Funct Plant Biol* 38:87–96. doi:10.1071/fp10124
- Cen Y-P, Turpin DH, Layzell DB (2001) Whole-plant gas exchange and reductive biosynthesis in white lupin. *Plant Physiol* 126:1555–1565
- Chen Z, Spreitzer RJ (1992) How various factors influence the  $\text{CO}_2/\text{O}_2$  specificity of ribulose-1,5-bisphosphate carboxylase/oxygenase. *Photosynth Res* 31:157–164

- Cheng L et al (2012) Arbuscular mycorrhizal fungi increase organic carbon decomposition under elevated CO<sub>2</sub>. *Science* 337:1084–1087. doi:10.1126/science.1224304
- Chollet R, Anderson LL (1976) Regulation of ribulose 1,5-bisphosphate carboxylase-oxygenase activities by temperature pretreatment and chloroplast metabolites. *Arch Biochem Biophys* 176:344–351. doi:10.1016/0003-9861(76)90173-9
- Christeller JT (1981) The effects of bivalent cations on ribulose bisphosphate carboxylase/oxygenase. *Biochem J* 193:839–844
- Christeller JT, Laing WA (1979) Effects of manganese ions and magnesium ions on the activity of soya-bean ribulose bisphosphate carboxylase/oxygenase. *Biochem J* 183:747–750
- Chu DK, Bassham JA (1974) Activation of ribulose 1,5-diphosphate carboxylase by nicotinamide adenine-dinucleotide phosphate and other chloroplast metabolites. *Plant Physiol* 54:556–559. doi:10.1104/pp.54.4.556
- Cleland WW, Andrews TJ, Gutteridge S, Hartman FC, Lorimer GH (1998) Mechanism of Rubisco: the carbamate as general base. *Chem Rev* 98:549–562. doi:10.1021/cr970010r
- Collatz GJ, Berry JA, Clark JS (1998) Effects of climate and atmospheric CO<sub>2</sub> partial pressure on the global distribution of C<sub>4</sub> grasses: present, past, and future. *Oecologia* 114:441–454. doi:10.1007/s004420050468
- Cotrufo MF, Ineson P, Scott A (1998) Elevated CO<sub>2</sub> reduces the nitrogen concentration of plant tissues. *Glob Change Biol* 4:43–54
- Cousins AB, Bloom AJ (2004) Oxygen consumption during leaf nitrate assimilation in a C<sub>3</sub> and C<sub>4</sub> plant: the role of mitochondrial respiration. *Plant, Cell Environ* 27:1537–1545
- Cramer M, Myers J (1948) Nitrate reduction and assimilation in *Chlorella*. *J Gen Physiol* 32:93–102
- Dortch Q (1990) The interaction between ammonium and nitrate uptake in phytoplankton. *Mar Ecol Prog* 61:183–201
- Dukes JS et al (2005) Responses of grassland production to single and multiple global environmental changes. *PLoS Biol* 3:1829–1837
- Dutilleul C, Lelarge C, Prioul J-L, De Paepe R, Foyer CH, Noctor G (2005) Mitochondria-driven changes in leaf NAD status exert a crucial influence on the control of nitrate assimilation and the integration of carbon and nitrogen metabolism. *Plant Physiol* 139:64–78
- Duval BD, Blankinship JC, Dijkstra P, Hungate BA (2012) CO<sub>2</sub> effects on plant nutrient concentration depend on plant functional group and available nitrogen: a meta-analysis. *Plant Ecol* 213:505–521. doi:10.1007/s11258-011-9998-8
- Ehleringer JR, Cerling TE, Helliker BR (1997) C<sub>4</sub> photosynthesis, atmospheric CO<sub>2</sub>, and climate. *Oecologia* 112:285–299
- Ellsworth DS, Reich PB, Naumburg ES, Koch GW, Kubiske ME, Smith SD (2004) Photosynthesis, carboxylation and leaf nitrogen responses of 16 species to elevated pCO<sub>2</sub> across four free-air CO<sub>2</sub> enrichment experiments in forest, grassland and desert. *Glob Change Biol* 10:2121–2138
- Epstein E, Bloom AJ (2005) Mineral nutrition of plants: principles and perspectives, 2nd edn. Sinauer Associates, Sunderland
- Finzi AC et al (2007) Increases in nitrogen uptake rather than nitrogen-use efficiency support higher rates of temperate forest productivity under elevated CO<sub>2</sub>. *Proc Natl Acad Sci USA* 104:14014–14019
- Foyer CH, Bloom AJ, Queval G, Noctor G (2009) Photorespiratory metabolism: genes, mutants, energetics, and redox signaling. *Annu Rev Plant Biol* 60:455–484
- Frank J, Kositzka MJ, Vater J, Holzwarth JF (2000) Microcalorimetric determination of the reaction enthalpy changes associated with the carboxylase and oxygenase reactions catalysed by ribulose 1,5-bisphosphate carboxylase/oxygenase (RUBISCO). *Phys Chem Chem Phys* 2:1301–1304
- Galmes J et al (2005) Rubisco specificity factor tends to be larger in plant species from drier habitats and in species with persistent leaves. *Plant, Cell Environ* 28:571–579
- Hall DO, Rao KK, Institute of Biology (1999) Photosynthesis. Studies in biology, 6th edn. Cambridge University Press, Cambridge
- Hanson AD, Hitz WD (1983) Water deficits and the nitrogen economy. In: Taylor HM, Jordan WR, Sinclair TR (eds) Limitations to efficient water use in crop production. ASA, Madison, pp 331–343
- Hodge A, Helgason T, Fitter AH (2010) Nutritional ecology of arbuscular mycorrhizal fungi. *Fungal Ecol* 3:267–273. doi:10.1016/j.funeco.2010.02.002
- Houtz RL, Nable RO, Cheniae GM (1988) Evidence for effects on the in vivo activity of ribulose-bisphosphate carboxylase/oxygenase during development of Mn toxicity in tobacco. *Plant Physiol* 86:1143–1149. doi:10.1104/pp.86.4.1143
- Igamberdiev AU, Bykova NV, Lea PJ, Gardstrom P (2001) The role of photorespiration in redox and energy balance of photosynthetic plant cells: a study with a barley mutant deficient in glycine decarboxylase. *Physiol Plant* 111:427–438
- Ishijima S, Uchibori A, Takagi H, Maki R, Ohnishi M (2003) Light-induced increase in free Mg<sup>2+</sup> concentration in spinach chloroplasts: measurement of free Mg<sup>2+</sup> by using a fluorescent probe and necessity of stromal alkalization. *Arch Biochem Biophys* 412:126–132. doi:10.1016/s0003-9861(03)00038-9
- Jordan DB, Ogren WL (1981) A sensitive assay procedure for simultaneous determination of ribulose-1,5-bisphosphate carboxylase and oxygenase activities. *Plant Physiol* 67:237–245
- Kane HJ, Viil J, Entsch B, Paul K, Morell MK, Andrews TJ (1994) An improved method for measuring the CO<sub>2</sub>/O<sub>2</sub> specificity of ribulosebisphosphate carboxylase-oxygenase. *Funct Plant Biol* 21:449–461
- Kimball BA, Idso SB, Johnson S, Rillig MC (2007) Seventeen years of carbon dioxide enrichment of sour orange trees: final results. *Glob Change Biol* 13:2171–2183
- Knaff DB (1996) Ferredoxin and ferredoxin-dependent enzymes. In: Ort DR, Yocum CF (eds) Oxygenic photosynthesis: the light reactions, vol 4., Advances in Photosynthesis. Kluwer Academic, Dordrecht, pp 333–361
- Korner C (2006) Plant CO<sub>2</sub> responses: an issue of definition, time and resource supply. *New Phytol* 172:393–411
- Laing WA, Christeller JT (1976) A model for the kinetics of activation and catalysis of ribulose 1,5-bisphosphate carboxylase. *Biochem J* 159:563–570
- Lekshmy S, Jain V, Khetarpal S, Pandey R (2013) Inhibition of nitrate uptake and assimilation in wheat seedlings grown under elevated CO<sub>2</sub>. *Indian J Plant Physiol* 18:23–29
- Lilley RMC, Wang XQ, Krausz E, Andrews TJ (2003) Complete spectra of the far-red chemiluminescence of the oxygenase reaction of Mn<sup>2+</sup>-activated ribulose-bisphosphate carboxylase/oxygenase establish excited Mn<sup>2+</sup> as the source. *J Biol Chem* 278:16488–16493
- Long SP, Ainsworth EA, Rogers A, Ort DR (2004) Rising atmospheric carbon dioxide: plants face the future. *Annu Rev Plant Biol* 55:591–628
- Luque-Almagro VM, Gates AJ, Moreno-Vivián C, Ferguson SJ, Richardson DJ, Roldán M (2011) Bacterial nitrate assimilation: gene distribution and regulation. *Biochem Soc Trans* 39:1838–1843
- Matsumura H et al (2012) Crystal structure of rice Rubisco and implications for activation induced by positive effectors NADPH and 6-phosphogluconate. *J Mol Biol* 422:75–86
- Matt P, Geiger M, Walch-Liu P, Engels C, Krapp A, Stitt M (2001) Elevated carbon dioxide increases nitrate uptake and nitrate reductase activity when tobacco is growing on nitrate, but increases ammonium uptake and inhibits nitrate reductase

- activity when tobacco is growing on ammonium nitrate. *Plant, Cell Environ* 24:1119–1137
- Maynard PN, Barker AV, Minotti PL, Peck NH (1976) Nitrate accumulation in vegetables. *Adv Agron* 28:71–119
- McCurry SD, Pierce J, Tolbert NE, Orme-Johnson WH (1981) On the mechanism of effector-mediated activation of ribulose biphosphate carboxylase/oxygenase. *J Biol Chem* 256:6623–6628
- McIntyre GI (1997) The role of nitrate in the osmotic and nutritional control of plant development. *Aust J Plant Physiol* 24:103–118
- Miziorko HM, Sealy RC (1980) Characterization of the ribulosebiphosphate carboxylase-carbon dioxide-divalent cation-carboxypentitol biphosphate complex. *Biochemistry* 19:1167–1171. doi:10.1021/bi00547a020
- Miziorko HM, Sealy RC (1984) Electron spin resonance studies of ribulose biphosphate carboxylase: identification of activator cation ligands. *Biochemistry* 23:479–485
- Moroney J, Jungnick N, DiMario R, Longstreth D (2013) Photorespiration and carbon concentrating mechanisms: two adaptations to high O<sub>2</sub>, low CO<sub>2</sub> conditions. *Photosynth Res* 117:121–131. doi:10.1007/s11220-013-9865-7
- Maeda S-i, Konishi M, Yanagisawa S, Omata T (2014) Nitrite transport activity of a novel HPP family protein conserved in cyanobacteria and chloroplasts. *Plant Cell Physiol* 55:1311–1324. doi:10.1093/pcp/pcu075
- Myers SS et al (2014) Increasing CO<sub>2</sub> threatens human nutrition. *Nature* 510:139–142. doi:10.1038/nature13179
- Norby RJ, Warren JM, Iversen CM, Medlyn BE, McMurtrie RE (2010) CO<sub>2</sub> enhancement of forest productivity constrained by limited nitrogen availability. *Proc Natl Acad Sci USA* 107:19368–19373. doi:10.1073/pnas.1006463107
- Ohashi Y et al (2011) Regulation of nitrate assimilation in cyanobacteria. *J Exp Bot* 62:1411–1424. doi:10.1093/jxb/erq427
- Oliva M, Safont VS, Andres J, Tapia O (2001) Transition structures for D-ribulose-1,5-bisphosphate carboxylase/oxygenase-catalyzed oxygenation chemistry: role of carbamylated lysine in a model magnesium coordination sphere. *J Phys Chem A* 105:4726–4736
- Osborne CP, Beerling DJ (2006) Nature's green revolution: the remarkable evolutionary rise of C<sub>4</sub> plants. *Philos Trans R Soc B Biol Sci* 361:173–194. doi:10.1098/rstb.2005.1737
- Parry MAJ, Andralojc PJ, Mitchell RAC, Madgwick PJ, Keys AJ (2003) Manipulation of Rubisco: the amount, activity, function and regulation. *J Exp Bot* 54:1321–1333
- Parry MAJ, Andralojc PJ, Scales JC, Salvucci ME, Carmo-Silva AE, Alonso H, Whitney SM (2013) Rubisco activity and regulation as targets for crop improvement. *J Exp Bot* 64:717–730. doi:10.1093/jxb/ers336
- Pearson PN, Palmer MR (2000) Atmospheric carbon dioxide concentrations over the past 60 million years. *Nature* 406:695–699
- Pierce J, Reddy GS (1986) The sites for catalysis and activation of ribulosebiphosphate carboxylase share a common domain. *Arch Biochem Biophys* 245:483–493
- Pleijel H, Uddling J (2012) Yield vs. quality trade-offs for wheat in response to carbon dioxide and ozone. *Glob Change Biol* 18:596–605. doi:10.1111/j.1365-2486.2011.2489.x
- Quesada A, Gomez-Garcia I, Fernandez E (2000) Involvement of chloroplast and mitochondria redox valves in nitrate assimilation. *Trends Plant Sci* 5:463–464
- Rachmilevitch S, Cousins AB, Bloom AJ (2004) Nitrate assimilation in plant shoots depends on photorespiration. *Proc Natl Acad Sci USA* 101:11506–11510
- Raghavendra AS, Carrillo NJ, Vallejos RH (1981) Differential modulation of carboxylase and oxygenase activities of ribulose 1, 5-bisphosphate carboxylase/oxygenase released from freshly ruptured spinach chloroplasts. *Plant Cell Physiol* 22:1113–1117
- Rasse DP, Peresta G, Drake BG (2005) Seventeen years of elevated CO<sub>2</sub> exposure in a Chesapeake Bay Wetland: sustained but contrasting responses of plant growth and CO<sub>2</sub> uptake. *Glob Change Biol* 11:369–377
- Rathnam CKM (1978) Malate and dihydroxyacetone phosphate-dependent nitrate reduction in spinach leaf protoplasts. *Plant Physiol* 62:220–223. doi:10.1104/pp.62.2.220
- Reich PB et al (2006) Nitrogen limitation constrains sustainability of ecosystem response to CO<sub>2</sub>. *Nature* 440:922–925
- Robinson JM (1987) Interactions of carbon and nitrogen metabolism in photosynthetic and non-photosynthetic tissues of higher plants: metabolic pathways and controls. In: Newman DW, Stuart KG (eds) *Models in Plant Physiology and Biochemistry*, vol 1. CRC Press, Boca Raton, pp 25–35
- Robinson JM (1988) Spinach leaf chloroplast CO<sub>2</sub> and NO<sub>2</sub><sup>-</sup> photoassimilations do not compete for photogenerated reductant manipulation of reductant levels by quantum flux density titrations. *Plant Physiol* 88:1373–1380
- Robinson JM, Gibbs M (1982) Hydrogen peroxide synthesis in isolated spinach chloroplast lamellae: an analysis of the Mehler reaction in the presence of NADP reduction and ATP formation. *Plant Physiol* 70:1249–1254
- Sage RF (2002) Variation in the *k*<sub>cat</sub> of Rubisco in C<sub>3</sub> and C<sub>4</sub> plants and some implications for photosynthetic performance at high and low temperature. *J Exp Bot* 53:609–620
- Sage RF, Li M, Monson RK (1999) The taxonomic distribution of C<sub>4</sub> photosynthesis. In: Sage RF, Monson RK (eds) *C<sub>4</sub> Plant Biology*. Academic Press, San Diego, pp 551–584
- Sage RF, Sage TL, Kocacinar F (2012) Photorespiration and the evolution of C<sub>4</sub> photosynthesis. *Annu Rev Plant Biol* 63:19–47
- Scheibe R, Dietz KJ (2012) Reduction-oxidation network for flexible adjustment of cellular metabolism in photoautotrophic cells. *Plant, Cell Environ* 35:202–216. doi:10.1111/j.1365-3040.2011.02319.x
- Schneiderei J, Hausler RE, Fiene G, Kaiser WM, Weber APM (2006) Antisense repression reveals a crucial role of the plastidic 2-oxoglutarate/malate translocator DiT1 at the interface between carbon and nitrogen metabolism. *Plant J* 45:206–224
- Searles PS, Bloom AJ (2003) Nitrate photoassimilation in tomato leaves under short-term exposure to elevated carbon dioxide and low oxygen. *Plant, Cell Environ* 26:1247–1255
- Sharkey TD (1988) Estimating the rate of photorespiration in leaves. *Physiol Plant* 73:147–152
- Skillman JB (2008) Quantum yield variation across the three pathways of photosynthesis: not yet out of the dark. *J Exp Bot* 59:1647–1661. doi:10.1093/jxb/Ern029
- Still CJ, Berry JA, Collatz GJ, DeFries RS (2003) Global distribution of C<sub>3</sub> and C<sub>4</sub> vegetation: carbon cycle implications. *Glob Biogeochem Cycles* 17:6-1–6-14
- Studer RA, Christin P-A, Williams MA, Orengo CA (2014) Stability-activity tradeoffs constrain the adaptive evolution of RubisCO. *Proc Natl Acad Sci* 111:2223–2228. doi:10.1073/pnas.1310811111
- Swedish Energy Agency (2003) *Artificial photosynthesis*. Eskilstuna, Sweden
- Taiz L, Zeiger E (2010) *Plant Physiology*, 5th edn. Sinauer Associates, Sunderland, MA
- Talhelm AF et al (2014) Elevated carbon dioxide and ozone alter productivity and ecosystem carbon content in northern temperate forests. *Glob Change Biol* 20:2492–2504. doi:10.1111/gcb.12564
- Taniguchi M, Miyake H (2012) Redox-shuttling between chloroplast and cytosol: integration of intra-chloroplast and extra-chloroplast metabolism. *Curr Opin Plant Biol* 15:252–260. doi:10.1016/j.pbi.2012.01.014

- Tapia O, Andrés J (1992) Towards an explanation of carboxylation/oxygenation bifunctionality in Rubisco. Transition structure for the carboxylation reaction of 2,3,4-pentanetriol. *Mol Eng* 2:37–41. doi:[10.1007/BF00999521](https://doi.org/10.1007/BF00999521)
- Taub DR, Wang XZ (2008) Why are nitrogen concentrations in plant tissues lower under elevated CO<sub>2</sub>? A critical examination of the hypotheses. *J Integr Plant Biol* 50:1365–1374. doi:[10.1111/j.1744-7909.2008.00754.x](https://doi.org/10.1111/j.1744-7909.2008.00754.x)
- Tcherkez GGB, Farquhar GD, Andrews TJ (2006) Despite slow catalysis and confused substrate specificity, all ribulose biphosphate carboxylases may be nearly perfectly optimized. *Proc Natl Acad Sci USA* 103:7246–7251
- Tolbert NE (1994) Role of photosynthesis and photorespiration in regulating CO<sub>2</sub> and O<sub>2</sub>. In: Tolbert NE, Preiss J (eds) *Regulation of CO<sub>2</sub> and O<sub>2</sub> by Photosynthetic Carbon Metabolism*. Oxford University Press, New York, pp 8–33
- Van Niel CB, Allen MB, Wright BE (1953) On the photochemical reduction of nitrate by algae. *Biochim Biophys Acta* 12:67–74. doi:[10.1016/0006-3002\(53\)90124-3](https://doi.org/10.1016/0006-3002(53)90124-3)
- Veen BW, Kleinendorst A (1986) The role of nitrate in osmoregulation of Italian ryegrass. *Plant Soil* 91:433–436. doi:[10.1007/bf02198139](https://doi.org/10.1007/bf02198139)
- von Caemmerer S, Quick WP, Furbank RT (2012) The development of C<sub>4</sub> rice: current progress and future challenges. *Science* 336:1671–1672
- Voss I, Sunil B, Scheibe R, Raghavendra AS (2013) Emerging concept for the role of photorespiration as an important part of abiotic stress response. *Plant Biol* 15:713–722. doi:[10.1111/j.1438-8677.2012.00710.x](https://doi.org/10.1111/j.1438-8677.2012.00710.x)
- Walker BJ, Strand DD, Kramer DM, Cousins AB (2014) The response of cyclic electron flow around photosystem I to changes in photorespiration and nitrate assimilation. *Plant Physiol* 165:453–462
- Warburg O, Negelein E (1920) Über die Reduktion der Salpetersäure in grünen Zellen. *Biochem Z* 110:66–115
- Whitney SM, Houtz RL, Alonso H (2011) Advancing our understanding and capacity to engineer nature's CO<sub>2</sub>-sequestering enzyme Rubisco. *Plant Physiol* 155:27–35. doi:[10.1104/pp.110.164814](https://doi.org/10.1104/pp.110.164814)
- Wildner GF, Henkel J (1979) The effect of divalent metal ions on the activity of Mg<sup>++</sup> depleted ribulose-1, 5-bisphosphate oxygenase. *Planta* 146:223–228
- Wingler A, Lea PJ, Quick WP, Leegood RC (2000) Photorespiration: metabolic pathways and their role in stress protection. *Philos Trans R Soc Lond Ser B Biol Sci* 355:1517–1529
- Zhu XG, Long SP, Ort DR (2008) What is the maximum efficiency with which photosynthesis can convert solar energy into biomass? *Curr Opin Biotechnol* 19:153–159
- Zhu XG, Long SP, Ort DR (2010) Improving photosynthetic efficiency for greater yield. *Annu Rev Plant Biol* 61:235–261. doi:[10.1146/annurev-arplant-042809-112206](https://doi.org/10.1146/annurev-arplant-042809-112206)