

RESEARCH ARTICLE


Expanding the genetic toolkit of *Tribolium castaneum*

Johnathan C. Rylee, Dylan J. Siniard[✉], Kaitlin Doucette, Gabriel E. Zentner*, Andrew C. Zelhof*

Department of Biology, Indiana University, Bloomington, IN, United States of America

✉ Current address: Division of Asthma Research, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, United States of America

* gzentner@indiana.edu (GEZ); azelhof@indiana.edu (ACZ)



OPEN ACCESS

Citation: Rylee JC, Siniard DJ, Doucette K, Zentner GE, Zelhof AC (2018) Expanding the genetic toolkit of *Tribolium castaneum*. PLoS ONE 13(4): e0195977. <https://doi.org/10.1371/journal.pone.0195977>

Editor: Subba Reddy Palli, University of Kentucky, UNITED STATES

Received: February 21, 2018

Accepted: April 3, 2018

Published: April 12, 2018

Copyright: © 2018 Rylee et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by Indiana University startup funds (to GEZ) and National Science Foundation grant IOS-1353267 (to ACZ). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Abstract

The red flour beetle, *Tribolium castaneum*, is an important model insect and agricultural pest. However, many standard genetic tools are lacking or underdeveloped in this system. Here, we present a set of new reagents to augment existing *Tribolium* genetic tools. We demonstrate a new GAL4 driver line that employs the promoter of a ribosomal protein gene to drive expression of a UAS responder in the fat body. We also present a novel dual fluorescent reporter that labels cell membranes and nuclei with different fluorophores for the analysis of cellular morphology. This approach also demonstrates the functionality of the viral T2A peptide for bicistronic gene expression in *Tribolium*. To facilitate classical genetic analysis, we created lines with visible genetic markers by CRISPR-mediated disruption of the *yellow* and *ebony* body color loci with a cassette carrying an attP site, enabling future φC31-mediated integration. Together, the reagents presented here will facilitate more robust genetic analysis in *Tribolium* and serve as a blueprint for the further development of this powerful model's genetic toolkit.

Introduction

The red flour beetle (*Tribolium castaneum*), a pest of stored agricultural products, has emerged as a promising system for biological research. It is a representative of the order Coleoptera, which comprises approximately 40% of known insect species and 25% of all known animals [1]. While *Drosophila melanogaster* is by far the most popular insect model system, many aspects of its development and physiology are not representative of insects in general, and so findings in *Tribolium* may be more broadly applicable to insects in many cases. Furthermore, Coleoptera includes significant agricultural pests such as the corn rootworm, Colorado potato beetle, and Asian longhorn beetle, and so using *Tribolium* as an insect model may lead to advances in pest control.

Advances in genetic tools have cemented the status of *Tribolium* as the second model insect of choice behind *Drosophila*. Transgenic *Tribolium* may be obtained using various transposons [2, 3] and more recently via CRISPR/Cas9-mediated genome editing [4]. Eye-specific fluorescent markers have also been developed to aid identification of transgenics [2]. Transposition

techniques have been used in for insertional mutagenesis, allowing the identification of essential genes as well as enhancer traps [5]. Another valuable tool for functional genomics, RNA interference (RNAi) via injection of eggs, larvae, or adults, has been implemented in *Tribolium*, both in targeted studies [6–8] and in a large-scale screen of the protein-coding genome [9]. Lastly, the GAL4/UAS system, a popular choice for spatiotemporally controlled expression of a gene of interest in *Drosophila*, has been demonstrated to function in *Tribolium* in the presence of a species-specific basal promoter [10].

Despite the proliferation of tools for genetic analysis and manipulation of *Tribolium*, notable gaps remain. In the case of the GAL4/UAS system, only two driver lines are available, one using a heat shock promoter [10] and the other making use of the odorant receptor co-receptor (*Orco*) regulatory regions [11]. Furthermore, the spectrum of *Tribolium* strains with visible phenotypic markers of known genetic location, which are staples of classical genetic analysis, is not as extensive as in other model organisms. Here, we present a set of reagents to address these issues and enhance the utility of *Tribolium* as a genetic model organism. We first present a GAL4 driver line that employs a ribosomal gene promoter to direct expression in the fat body and can serve as an second effective marker for transgenesis. In addition, we describe a GAL4-inducible cellular reporter in which the nucleus and endomembrane system are labeled with different fluorescent proteins, acting as a robust means by which to analyze cellular structure, particularly with respect to neurons. Furthermore, both the GAL4 and UAS cloning vectors are designed to accept any gene or genomic region of interest to generate new drivers and reporters. We also address the lack of visible phenotypic markers in *Tribolium* by using CRISPR to disrupt two genes involved in cuticle pigmentation via homologous recombination with cassettes containing an attP site to facilitate future genomic insertion of DNA of interest using the φC31 integrase. The tools presented here represent a valuable resource for the *Tribolium* research community and serve as a general template upon which further tools can be based.

Materials and methods

Tribolium husbandry and strains

All animals were raised at 28°C on a standard flour yeast mix. The following strains were utilized: *vermillion*^{white} (*v^W*), bearing a partial deletion of the *vermillion* gene with a concomitant white eyed phenotype [12], and *m26*, a *v^w* line with X-linked insertion of the piggyBac transposase marked with 3xP3-DsRed [3].

Vectors

All vectors will be made available through the *Drosophila* Genomics Resource Center at Indiana University (<https://dgrc.bio.indiana.edu/Home>).

P119der. 119der, a derivative of pSLfa[UAS-Tc'Hsp-p-tGFP-SV40] (kindly provided by Dr. Gregor Bucher) [10], was constructed by excision of tGFP with KpnI and NotI followed by replacement with the sequence ACTAGTGAATTCAAAGTACCACTCGAGAGCGGCCGCG. This replacement destroyed the KpnI site but preserved the unique NotI site and added a unique XhoI site. (DGRC # 1425)

P130der. p130der, a derivative of pSLfa[Hsp-p-Gal4Delta-SV40_attp] (kindly provided by Dr. Gregor Bucher) [10], was constructed by digestion with BamHI and addition of the sequence GGATCCAGGTACCAGCGGCCGAGGATCC, containing unique KpnI and NotI sites. (DGRC # 1426)

pGZ286 (pBac-3xP3-EGFP-pTC006550-GAL4Δ). The basal *hsp68* promoter of PCR-linearized p130der was replaced with the TC006550 promoter amplified from blaAmp-Tc6550Pro-GFPZeo-Luciferase-HSP-Orange-pIZT (kindly provided by Dr. Yoonseong Park)

[13] with NEBuilder HiFi assembly (NEB). The resulting pTC006550-GAL4Δ-SV40 polyA coding sequence was then amplified and inserted into PCR-linearized pBac[3xP3-EGFP] with NEBuilder HiFi assembly. (DGRC # 1427)

pTC241 (pBac-3XP3-EGFP-UAS-nls-EGFP-T2A-mCherry). The nls-EGFP-T2A-mCherryCAAX insert was amplified from pSYC-102 (a gift from Seok-Yong Choi, Addgene plasmid #74790) [14] as two fragments, and assembled into the NotI site of p119der using NEBuilder HiFi assembly (DGRC #1428). UAS-nls-EGFP-T2A-mCherryCAAX was then excised using flanking AscI sites and ligated into pBac[3xP3-GFP].

Immunofluorescence and imaging

Tribolium larvae were dissected in PBS and tissue was fixed in 4% paraformaldehyde/PBS. The tissue was washed 2X with PBT (PBS and 0.8% Triton-X 100) and incubated with DAPI (final concentration of 0.1 µg/ml) for 10 minutes and then washed with PBS before mounting for imaging (modified from [15]). Confocal images were captured on a Leica SP8 confocal utilizing a Leica 63X oil immersion objective with a numerical aperture of 1.40. Light and fluorescent whole animal images were collected on a LeicaMZ10F dissecting microscope. All pictures were processed in Adobe Photoshop.

CRISPR

gRNAs were designed using CRISPRdirect [16] using the Tcas3 genome assembly for the specificity check. Only high-quality gRNAs were selected. 20-mer protospacer sequences were cloned into BsaI-digested pU6b-BsaI-gRNA [4] by NEBuilder HiFi-mediated ssDNA oligo bridging as described (<https://www.neb.com/-/media/nebus/files/application-notes/construction-of-an-sgRNA-cas9-expression-vector-via-single-stranded-dna-oligo-bridging-of-double-stranded-dna-fragments.pdf>) using an ssDNA oligo consisting of the protospacer flanked by 25 bp regions of vector homology. Protospacer sequences used were CCGAAAATAATCTCCAGT (yellow, TC000802) (DGRC #1429) and TTTCGTAAAAGTTGAATCG (ebony, TC0011976) (DGRC #1430). Homology donors consisting of an attP site and 3xP3-DsRed-SV40 polyA flanked by loxP sites in the same orientation between 800 bp (yellow; left arm: ChLG2:7,663,877–7,664,676, right arm: ChLG2:7,664,677–7,665,476) or 726/739 bp (ebony; left arm: ChLG9:13,340,543–13,341,268, right arm: ChLG9:13,341,269–13,342,007) homology arms were synthesized by IDT and delivered in pUCIDT-amp. Mixtures consisting of 400–500 ng/mL each of Cas9 plasmid [4], sgRNA plasmid, and donor plasmid were injected into *v^w* embryos.

Results

Construction and implementation of a *ribo-GAL4* driver

Inducible expression of a transgene of interest is a key capability in any genetic model organism. In *Drosophila*, the bipartite *Saccharomyces cerevisiae* GAL4/UAS (upstream activating sequence) system, in which the GAL4 transcription factor binds to a 17-bp motif within the UAS to drive transcriptional activation, is frequently used for spatiotemporally controlled expression of a gene of interest [17]. In *Tribolium*, the GAL4-UAS system has been demonstrated to be functional when the UAS is coupled with the *hsp68* basal promoter and driven by heat shock-inducible GAL4 [10]. However, no ubiquitous GAL4 drivers have been reported for *Tribolium*. Such drivers are useful when screening for an organismal or developmental phenotype of overexpression or knockdown.

In an attempt to generate a ubiquitous and constitutive GAL4 driver line, we considered a previous study in which the promoter of a ribosomal protein gene (TC006550) was used to

drive high-level expression in the *Tribolium* TcA cell line [13]. We thus replaced the heat shock promoter of our p130der vector (see [Materials and Methods](#)), bearing the basal *hsp68* promoter and GAL4 Δ , with the *TC006550* promoter. GAL4 Δ is a variant of GAL4 in which the N- and C-termini, containing the DNA-binding and transcriptional activation domains, are directly fused [18]. This variant of GAL4 has been shown to increase transactivation by ~2-fold in *Drosophila* [19]. We then transferred the *TC006550* promoter and GAL4 Δ coding region to a piggyBac vector containing 3xP3-EGFP, enabling selection of transgenics by fluorescence in photoreceptors. This vector was injected into the m26 *Tribolium* line, which lacks eye pigmentation due to the *v^w* mutation and has 3xP3-DsRed-marked piggyBac transposase integrated into the X chromosome [3]. Resulting adults were outcrossed to *v^w* and progeny were assessed for GFP expression in the retina. *TC006550-GAL4 Δ* transformants (hereafter referred to as *ribo-GAL4*) were identified and the DsRed-marked transposase was removed through subsequent crosses and two independent insertions were generated. We note that while single-copy insertion of the *ribo-GAL4* construct can be detected via the associated EGFP fluorescence in the retina, this does not constitute or substitute for a genetically defined dominant marker.

To assay the functionality of the *ribo-GAL4* driver line, we crossed it to a previously described UAS-GFP responder line [10]. Larvae displayed strong whole-body fluorescence ([Fig 1A–1C](#)) and fluorescence is maintained throughout pupal development and into adulthood as expected for a ubiquitous expression. However, further examination revealed fluorescence was only detected in the putative fat body ([Fig 1D–1G](#)) and absent from other tissues (e.g. the gut, muscle, and CNS). We speculate that the lack of ubiquitous GAL4 expression in the *ribo-GAL4* line may reflect tissue-specific differences in ribosomal protein gene expression [20] and that, given the apparent *in vivo* expression profile of *TC006550*, the TcA cell line may be derived from fat cells.

Construction and implementation of a reporter for cell structure

There are numerous reporters available for highlighting cell structure and function. Our goal was to test whether these reporters could be simply swapped into a universal UAS cloning vector for *Tribolium* with zero or minimal changes to the already existing sequence. We chose nls-EGFP-T2A-mCherryCAAX [14] to test the utility of bicistronic fluorescent reporter expression for studying cell morphology as well as the use of the viral T2A peptide in *Tribolium*. When combined with our *ribo-GAL4* line, mCherry expression could easily be detected in whole larvae, mimicking the spatial and temporal pattern obtained with cytoplasmic GFP ([Fig 1A–1C](#)). To confirm the expression and localization of both the nuclear GFP and the endomembrane linked mCherry we examined the subcellular localization of each in fat cells. Colocalization with DAPI confirmed the subcellular localization of GFP in the nucleus with mCherryCAAX bound to membranes ([Fig 1D–1G](#)). These results indicate that existing fluorescent reporters can be easily implemented in *Tribolium* using our UAS vector and that the T2A peptide can be used for multicistronic gene expression in *Tribolium*.

CRISPR-based generation of lines with visible phenotypic markers

Defined phenotypic visible markers are essential for facilitating even basic tasks such as establishment of stocks of genetically modified organisms or tracking of specific chromosomes through multiple generations. While the scope of genetic reagents available in *Tribolium* has markedly increased over the past few decades, there are notable limitations and gaps in existing tools. For instance, the use of visible phenotypic markers for genetic mapping is quite limited, especially compared to *Drosophila*, where variations in wings, bristles, eye and body




Fig 1. Characterization of GAL4 and UAS vectors. (A-C) Tribolium larvae expressing either GFP (left) or nls-EGFP-T2A-mCherryCAAX (right) driven by *rbo-GAL4*. (A) White light illumination; (B) GFP illumination. The nls-EGFP signal is not detectable as compared to cytoplasmic GFP. (C) mCherry illumination. Scale bar = 1 mm. The mCherry illumination mimics the cytoplasmic GFP expression. (D-G) Images of fat cells expressing nls-EGFP (D) and mCherryCAAX (E), counterstained with DAPI (F), and the merge of the three labels (G). Each represents a single confocal section. Scale bar = 10 μ m.

<https://doi.org/10.1371/journal.pone.0195977.g001>

pigmentation, and body shape are available. Additionally, while dominant and recessive mutations giving rise to visible phenotypes in *Tribolium* have been documented [21–26], the majority have not been mapped to a tractable genetic interval and so cannot be used in such analyses. Notable exceptions to this include both dominant and recessive loci within the LG2 chromosome, particularly *Reindeer* (*Rd*), and several mutations in the homeotic complex (HOM-C) [26].

In order to create a general strategy for the expansion of a pool of visible phenotypic markers for *Tribolium*, we employed CRISPR based homologous recombination. We designed gRNAs against the coding regions of the *yellow* (TC000802) and *ebony* (TC011976) genes, as well as a disruption cassette with useful genetic features. In many insects, the disruption of




Fig 2. Construction and validation of the CRISPR gene disruption strategy. (A) Schematic representation of the gene disruption cassette used in this study. The positions and directionalities of primers used for genomic PCR validation are included in the schematic. (B) Positions of gRNAs used to disrupt the *yellow* and *ebony* loci. (C) Genomic PCR demonstrating the presence of each gene disruption cassette in the genome of transgenic but not parental beetles.

<https://doi.org/10.1371/journal.pone.0195977.g002>

either *ebony* and *yellow* leads to viable and fertile animals that are easily identifiable from their wild-type counterparts. With respect to our CRISPR procedure, we chose a homologous recombination strategy that would permit detection of a disruption in either locus regardless of whether a change in pigmentation resulted. For each targeting construct, we utilized 700–800 bp homology arms, and between them we enclosed an attP site to facilitate future φC31 integrase-mediated insertion of DNA of interest at a defined location [27]. More importantly, we included the DsRed Express fluorescent protein under the control of the eye-specific 3xP3 regulatory element to facilitate screening of CRISPR based recombinants; and two loxP sites in the same orientation, enabling future Cre-mediated excision of 3xP3-DsRed from the genome (Fig 2A). The positioning of the loxP sites is such that they flank only the fluorescent marker and so excision would leave the attP site intact, effectively recycling the DsRed marker for further use.

v^w embryos were injected with a mixture of Cas9 plasmids, gRNA expression plasmid, and repair template plasmids. Surviving adults were crossed to *v^w* and resulting progeny were screened upon eclosion for DsRed expression in the retina. For *yellow*, 308/356 injected larvae survived to adulthood and of these, two tested positive for germline transmission of the disrupted gene (0.6%). For *ebony*, 184/217 injected larvae survived to adulthood and of these, one tested positive for germline transmission of the disrupted gene (0.54%).

Characterization of *yellow*-edited adults revealed that cuticles of newly eclosed beetles were noticeably lighter than those of their *v^w* counterparts, but their color darkened over time until




Fig 3. Phenotypic characterization of yellow CRISPR beetles. Dorsal and ventral views of parental v^w and transgenic yellow CRISPR beetles at 2 and 4 days post-eclosion. Scale bar = 1 mm.

<https://doi.org/10.1371/journal.pone.0195977.g003>

they were difficult to distinguish from the parental line (Fig 3). In contrast, adults with homozygous disruption of *ebony* displayed substantially darker cuticular pigmentation than parental v^w individuals (Fig 4A and 4B) and DsRed fluorescence in their eyes (Fig 4C). While *ebony* CRISPR individuals were clearly phenotypically distinct from the parental line, their cuticle was not as dark as individuals injected with *ebony* dsRNA (<http://ibeetle-base.uni-goettingen.de/details/TC011976>). We speculate that this difference is due to the location of the gRNA target site. It lies within the seventh exon of *ebony*, which falls after the sequences encoding all but one predicted functional domain of the protein (Fig 5). However, this gRNA falls at the start of the last domain of the protein, and disruption of this domain likely explains the hypomorphic phenotype observed.

Conclusions

Here, we present a series of reagents aimed at increasing the genetic tractability of *Tribolium*. These are (1) a GAL4 driver plasmid and *Tribolium* line expressing GAL4 in fat cells; (2) a UAS plasmid and UAS-inducible dual-nuclear/endomembrane fluorescence reporter for analyzing cell structure; (3) a template gene disruption cassette for the generation of mutants and insertion of attP sites; and (4) *Tribolium* lines carrying disruptions of two body color loci, *yellow* and *ebony*.




Fig 4. Phenotypic characterization of ebony CRISPR beetles. (A) Dorsal and (B) ventral views of parental v^w and transgenic ebony CRISPR beetles. Scale bar = 1 mm. (C) DsRed fluorescence microscopic image of the eyes of v^w and transgenic ebony CRISPR beetles. Scale bar = 0.5 mm.

<https://doi.org/10.1371/journal.pone.0195977.g004>

We surmise that the failure of the *ribo*-GAL4 line to drive ubiquitous reporter expression is attributable to its late pupal origin [28], reflecting cell type variability in TC006550 expression. Indeed, heterogeneity in ribosomal protein expression across cell types has been widely reported [20, 29–31]. Other potential candidates for the establishment of a ubiquitous GAL4 line include the α -Tubulin1 promoter, which has been shown to drive ubiquitous GFP expression throughout the *Tribolium* life cycle [32], and the *Polyubiquitin* promoter [33]. The p130der plasmid permits efficient insertion of any potential genomic sequence for designing future GAL4 lines. Furthermore, our data demonstrates that any established reporter can be cloned into our modified version of pSLfa[UAS-Tc'Hsp-p-tGFP-SV40] [10], p119der, for direct expression in *Tribolium*. Future variations/deviations of both p130der and p119der will include an attB site for direct insertion into known genomic positions as well as a fluorescent marker to enable rapid screening of transgenics.

While we successfully disrupted the *yellow* and *ebony* loci via CRISPR, the visible phenotypes associated with these editing events were unexpected. In particular, disruption of *yellow* resulted in a slow-tanning phenotype, with young adults displaying a visibly lighter cuticle than the parental strain that then darkened until it was indistinguishable from that of non-edited beetles. In the case of *ebony*, we were able to achieve a marked darkening of the cuticle using our disruption strategy, but our mutation was potentially hypomorphic when compared to *ebony* RNAi, which yielded a darker black-body phenotype. In order to maximise the visible phenotypes obtainable with CRISPR disruption of an eye or body color locus, we therefore




Fig 5. NCBI conserved domain search results for the ebony protein. The location of the gRNA relative to the mature protein sequence is indicated.

<https://doi.org/10.1371/journal.pone.0195977.g005>

recommend pre-screening of candidate visible marker genes with RNAi prior to initiating genome editing. Several candidates for body eye and body color genes have been assessed by RNAi in the literature [34–37] and may serve as suitable targets for our CRISPR disruption approach. Moreover, our data demonstrate that the inclusion of an independent marker for CRISPR-based modifications is invaluable in recovering transformants and thus can mitigate the uncertainty associated with the targeting of other potential candidate loci for visible markers.

Supporting information

S1 Table. DNA oligonucleotides used in this work.
(DOCX)

Acknowledgments

We thank Drs. G. Bucher, S. Brown, and M. Lorenzen for *Tribolium* reagents. We thank Dr. J. Powers and the Indiana University Light Microscopy Imaging Center for assistance with image generation. We thank Dr. M. J. Wade for helpful discussions throughout the course of this work.

Author Contributions

Conceptualization: Gabriel E. Zentner, Andrew C. Zelhof.

Data curation: Johnathan C. Rylee, Dylan J. Siniard, Kaitlin Doucette, Gabriel E. Zentner, Andrew C. Zelhof.

Formal analysis: Gabriel E. Zentner, Andrew C. Zelhof.

Funding acquisition: Gabriel E. Zentner, Andrew C. Zelhof.

Investigation: Johnathan C. Rylee, Dylan J. Siniard, Kaitlin Doucette, Gabriel E. Zentner, Andrew C. Zelhof.

Methodology: Andrew C. Zelhof.

Project administration: Andrew C. Zelhof.

Resources: Dylan J. Siniard, Gabriel E. Zentner, Andrew C. Zelhof.

Supervision: Gabriel E. Zentner, Andrew C. Zelhof.

Visualization: Andrew C. Zelhof.

Writing – original draft: Gabriel E. Zentner, Andrew C. Zelhof.

Writing – review & editing: Johnathan C. Rylee, Gabriel E. Zentner, Andrew C. Zelhof.

References

1. Hunt T, Bergsten J, Levkanicova Z, Papadopoulou A, John OS, Wild R, et al. A Comprehensive Phylogeny of Beetles Reveals the Evolutionary Origins of a Superradiation. *Science*. 2007; 318(5858):1913–6. <https://doi.org/10.1126/science.1146954> PMID: 18096805
2. Berghammer AJ, Klingler M, Wimmer EA. A universal marker for transgenic insects. *Nature*. 1999; 402:370. <https://doi.org/10.1038/46463> PMID: 10586872
3. Lorenzen MD, Kimzey T, Shippy TD, Brown SJ, Denell RE, Beeman RW. *piggyBac*-based insertional mutagenesis in *Tribolium castaneum* using donor/helper hybrids. *Insect Mol Biol*. 2007; 16(3):265–75. <https://doi.org/10.1111/j.1365-2583.2007.00727.x> PMID: 17316329

4. Gilles AF, Schinko JB, Averof M. Efficient CRISPR-mediated gene targeting and transgene replacement in the beetle *Tribolium castaneum*. *Development*. 2015; 142(16):2832–9. <https://doi.org/10.1242/dev.125054> PMID: 26160901
5. Trauner J, Schinko J, Lorenzen MD, Shippy TD, Wimmer EA, Beeman RW, et al. Large-scale insertional mutagenesis of a coleopteran stored grain pest, the red flour beetle *Tribolium castaneum*, identifies embryonic lethal mutations and enhancer traps. *BMC Biol*. 2009; 7(1):73. <https://doi.org/10.1186/1741-7007-7-73> PMID: 19891766
6. Brown SJ, Mahaffey JP, Lorenzen MD, Denell RE, Mahaffey JW. Using RNAi to investigate orthologous homeotic gene function during development of distantly related insects. *Evolution & Development*. 1999; 1(1):11–5. <https://doi.org/10.1046/j.1525-142x.1999.99013.x>
7. Bucher G, Scholten J, Klingler M. Parental RNAi in *Tribolium* (Coleoptera). *Curr Biol*. 2009; 19(12):R85–R6. [https://doi.org/10.1016/S0960-9822\(09\)00666-8](https://doi.org/10.1016/S0960-9822(09)00666-8) PMID: 11839285
8. Tomoyasu Y, Denell RE. Larval RNAi in *Tribolium* (Coleoptera) for analyzing adult development. *Dev Genes Evol*. 2004; 214(11):575–8. <https://doi.org/10.1007/s00427-004-0434-0> PMID: 15365833
9. Schmitt-Engel C, Schultheis D, Schwirz J, Ströhlein N, Troelenberg N, Majumdar U, et al. The iBeetle large-scale RNAi screen reveals gene functions for insect development and physiology. *Nature Communications*. 2015; 6:7822. <https://doi.org/10.1038/ncomms8822> <https://www.nature.com/articles/ncomms8822-supplementary-information>. PMID: 26215380
10. Schinko JB, Weber M, Viktorinova I, Kiupakis A, Averof M, Klingler M, et al. Functionality of the GAL4/UAS system in *Tribolium* requires the use of endogenous core promoters. *BMC Dev Biol*. 2010; 10(1):53. <https://doi.org/10.1186/1471-213x-10-53> PMID: 20482875
11. Dippel S, Kollmann M, Oberhofer G, Montino A, Knoll C, Krala M, et al. Morphological and Transcriptomic Analysis of a Beetle Chemosensory System Reveals a Gnathal Olfactory Center. *BMC Biol*. 2016; 14(1):90. <https://doi.org/10.1186/s12915-016-0304-z> PMID: 27751175
12. Lorenzen MD, Brown SJ, Denell RE, Beeman RW. Cloning and Characterization of the *Tribolium castaneum* Eye-Color Genes Encoding Tryptophan Oxygenase and Kynurenine 3-Monoxygenase. *Genetics*. 2002; 160(1):225–34. PMID: 11805058
13. Silver K, Jiang H, Fu J, Phillips TW, Beeman RW, Park Y. The *Tribolium castaneum* cell line TcA: a new tool kit for cell biology. *Sci Rep*. 2014; 4:6840. <https://doi.org/10.1038/srep06840> <https://www.nature.com/articles/srep06840-supplementary-information>. PMID: 25354547
14. Kim JH, Lee S-R, Li L-H, Park H-J, Park J-H, Lee KY, et al. High Cleavage Efficiency of a 2A Peptide Derived from Porcine Teschovirus-1 in Human Cell Lines, Zebrafish and Mice. *PLOS ONE*. 2011; 6(4):e18556. <https://doi.org/10.1371/journal.pone.0018556> PMID: 21602908
15. Nie J, Mahato S, Zelhof AC. Imaging the *Drosophila* retina: zwitterionic buffers PIPES and HEPES induce morphological artifacts in tissue fixation. *BMC Dev Biol*. 2015; 15(1):10. <https://doi.org/10.1186/s12861-015-0056-y> PMID: 25645690
16. Naito Y, Hino K, Bono H, Ui-Tei K. CRISPRdirect: software for designing CRISPR/Cas guide RNA with reduced off-target sites. *Bioinformatics*. 2015; 31(7):1120–3. <https://doi.org/10.1093/bioinformatics/btu743> PMID: 25414360
17. Duffy JB. GAL4 system in drosophila: A fly geneticist's swiss army knife. *Genesis*. 2002; 34(1–2):1–15. <https://doi.org/10.1002/gen.10150> PMID: 12324939
18. Ma J, Ptashne M. Deletion analysis of GAL4 defines two transcriptional activating segments. *Cell*. 1987; 48(5):847–53. [https://doi.org/10.1016/0092-8674\(87\)90081-X](https://doi.org/10.1016/0092-8674(87)90081-X). PMID: 3028647
19. Viktorinová I, Wimmer EA. Comparative analysis of binary expression systems for directed gene expression in transgenic insects. *Insect Biochem Mol Biol*. 2007; 37(3):246–54. <https://doi.org/10.1016/j.ibmb.2006.11.010>. PMID: 17296499
20. Guimaraes JC, Zavolan M. Patterns of ribosomal protein expression specify normal and malignant human cells. *Genome Biology*. 2016; 17(1):236. <https://doi.org/10.1186/s13059-016-1104-z> PMID: 27884178
21. Brown E, Sokoloff A. LINKAGE STUDIES IN TRIBOLIUM CASTANEUM. XI. THE MAP POSITION OF CHARCOAL, A PSEUDOALLELE OF BLACK. *Canadian Journal of Genetics and Cytology*. 1978; 20(1):139–45. <https://doi.org/10.1139/g78-014>
22. Dawson PS. The "reindeer" mutation and a revision of linkage groups V and X in the flour beetle, *Tribolium castaneum*. *Canadian Journal of Genetics and Cytology*. 1984; 26(6):762–4. <https://doi.org/10.1139/g84-120>
23. Sokoloff A, Faustini D. Dachs, a mutant in *Tribolium* with effects analogous to engrailed in *Drosophila*. *Journal of Heredity*. 1987; 78(1):2–7. <https://doi.org/10.1093/oxfordjournals.jhered.a110299>
24. Beeman RW, Stuart JJ, Haas MS, Denell RE. Genetic analysis of the homeotic gene complex (HOM-C) in the beetle *Tribolium castaneum*. *Dev Biol*. 1989; 133(1):196–209. [https://doi.org/10.1016/0012-1606\(89\)90311-4](https://doi.org/10.1016/0012-1606(89)90311-4). PMID: 2565268

25. Mocelin G, Stuart JJ. Crossover Suppressors in *Tribolium castaneum*. *Journal of Heredity*. 1996; 87:27–34.
26. Beeman RW, Stuart JJ, Haas MS, Friesen KS. Chromosome Extraction and Revision of Linkage Group 2 in *Tribolium castaneum*. *Journal of Heredity*. 1996; 87:224–32. PMID: [8683098](https://doi.org/10.1007/s11626-012-9524-x)
27. Groth AC, Fish M, Nusse R, Calos MP. Construction of Transgenic Drosophila by Using the Site-Specific Integrase From Phage φC31. *Genetics*. 2004; 166(4):1775–82. <https://doi.org/10.1534/genetics.166.4.1775> PMID: [15126397](https://doi.org/10.1534/genetics.166.4.1775)
28. Goodman CL, Stanley D, Ringbauer JA, Beeman RW, Silver K, Park Y. A cell line derived from the red flour beetle *Tribolium castaneum* (Coleoptera: Tenebrionidae). *In Vitro Cell Dev Biol Anim*. 2012; 48(7):426–33. <https://doi.org/10.1007/s11626-012-9524-x> PMID: [22752637](https://doi.org/10.1007/s11626-012-9524-x)
29. Bortoluzzi S, d'Alessi F, Romualdi C, Danieli GA. Differential expression of genes coding for ribosomal proteins in different human tissues. *Bioinformatics*. 2001; 17(12):1152–7. <https://doi.org/10.1093/bioinformatics/17.12.1152> PMID: [11751223](https://doi.org/10.1093/bioinformatics/17.12.1152)
30. Thorrez L, Van Deun K, Tranchevent L-C, Van Lommel L, Engelen K, Marchal K, et al. Using Ribosomal Protein Genes as Reference: A Tale of Caution. *PLOS ONE*. 2008; 3(3):e1854. <https://doi.org/10.1371/journal.pone.0001854> PMID: [18365009](https://doi.org/10.1371/journal.pone.0001854)
31. Sauert M, Temmel H, Moll I. Heterogeneity of the translational machinery: Variations on a common theme. *Biochimie*. 2015; 114:39–47. <https://doi.org/10.1016/j.biochi.2014.12.011> PMID: [25542647](https://doi.org/10.1016/j.biochi.2014.12.011)
32. Siebert KS, Lorenzen MD, Brown SJ, Park Y, Beeman RW. Tubulin superfamily genes in *Tribolium castaneum* and the use of a Tubulin promoter to drive transgene expression. *Insect Biochem Mol Biol*. 2008; 38(8):749–55. <https://doi.org/10.1016/j.ibmb.2008.04.007> PMID: [18625397](https://doi.org/10.1016/j.ibmb.2008.04.007)
33. Lorenzen MD, Brown SJ, Denell RE, Beeman RW. Transgene expression from the *Tribolium castaneum* Polyubiquitin promoter. *Insect Mol Biol*. 2002; 11(5):399–407. <https://doi.org/10.1046/j.1365-2583.2002.00349.x> PMID: [12230539](https://doi.org/10.1046/j.1365-2583.2002.00349.x)
34. Osanai-Futahashi M, Tatematsu K-i, Yamamoto K, Narukawa J, Uchino K, Kayukawa T, et al. Identification of the Bombyx Red Egg Gene Reveals Involvement of a Novel Transporter Family Gene in Late Steps of the Insect Ommochrome Biosynthesis Pathway. *J Biol Chem*. 2012; 287(21):17706–14. <https://doi.org/10.1074/jbc.M111.321331> PMID: [22474291](https://doi.org/10.1074/jbc.M111.321331)
35. Broehan G, Kroeger T, Lorenzen M, Merzendorfer H. Functional analysis of the ATP-binding cassette (ABC) transporter gene family of *Tribolium castaneum*. *BMC Genomics*. 2013; 14(1):6. <https://doi.org/10.1186/1471-2164-14-6> PMID: [23324493](https://doi.org/10.1186/1471-2164-14-6)
36. Grubbs N, Haas S, Beeman RW, Lorenzen MD. The ABCs of Eye Color in *Tribolium castaneum*: Orthologs of the *Drosophila* white, scarlet, and brown Genes. *Genetics*. 2015; 199(3):749–59. <https://doi.org/10.1534/genetics.114.173971> PMC4349069. PMID: [25555987](https://doi.org/10.1534/genetics.114.173971)
37. Arakane Y, Lomakin J, Beeman RW, Muthukrishnan S, Gehrke SH, Kanost MR, et al. Molecular and Functional Analyses of Amino Acid Decarboxylases Involved in Cuticle Tanning in *Tribolium castaneum*. *J Biol Chem*. 2009; 284(24):16584–94. <https://doi.org/10.1074/jbc.M901629200> PMID: [19366687](https://doi.org/10.1074/jbc.M901629200)