

IMMersed CONCORDANCES OF LINKS AND HEEGAARD FLOER HOMOLOGY

MACIEJ BORODZIK AND EUGENE GORSKY

ABSTRACT. An immersed concordance between two links is a concordance with possible self-intersections. Given an immersed concordance we construct a smooth four-dimensional cobordism between surgeries on links. By applying d -invariant inequalities for this cobordism we obtain inequalities between the H -functions of links, which can be extracted from the link Floer homology package. As an application we show a Heegaard Floer theoretical criterion for bounding the splitting number of links. The criterion is especially effective for L-space links, and we present an infinite family of L-space links with vanishing linking numbers and arbitrary large splitting numbers. We also show a semicontinuity of the H -function under δ -constant deformations of singularities with many branches.

1. INTRODUCTION

1.1. Overview. An immersed cobordism between two links \mathcal{L}_1 and \mathcal{L}_2 in S^3 is a smoothly immersed surface in $S^3 \times [1, 2]$, whose boundary is $\mathcal{L}_1 \subset S^3 \times \{1\}$ and $\mathcal{L}_2 \subset S^3 \times \{2\}$. An immersed concordance is an immersed cobordism, whose all the components are immersed annuli. The notion of an immersed cobordism gives a unified approach for finding (usually lower) bounds on link invariants such as the smooth four genus, the clasp number, the splitting number and the unlinking number. Recently many papers using this technique appeared [3, 5, 19, 25, 27]. Generalizing the construction of [5] we can use an immersed concordance as a starting point in constructing a four-dimensional cobordism between large surgeries on \mathcal{L}_1 and \mathcal{L}_2 with precisely described surgery coefficients. Under some extra assumptions we can guarantee that the four-dimensional cobordism is negative definite. We apply the d -invariant inequality of Ozsváth and Szabó, see (4.3), to relate the d -invariants of the corresponding surgeries on \mathcal{L}_1 and \mathcal{L}_2 . These inequalities are best expressed in terms of the H -functions.

The H -function is a function that is used to calculate the d -invariants of large surgeries on links (see Theorem 4.10, which can be thought of as an informal definition of H). For knots it was first defined by Rasmussen in his thesis [31] (as an analogue of the Frøyshov invariant in Seiberg-Witten theory), who used it to obtain nontrivial bounds for the slice genus of knots. For L-space knots, the H -function can be easily reconstructed from the Alexander polynomial. For L-space links with several components (see Section 2.2), the H -function was introduced by the second author and Némethi [14] (denoted by small h there), who showed that for algebraic links it coincides with the Hilbert function defined by the valuations on the local ring of the corresponding singularity.

Unfortunately, apart from different notations of H in the literature, there are at least three different “natural” conventions on the definition of H , all differing by some shift of

2010 *Mathematics Subject Classification.* primary: 57M25, secondary: 14B07, 14H20.

Key words and phrases. L-space link, splitting number, immersed concordance, d -invariant, surgery, semigroup of a singular point, Hilbert function.

The second author was partially supported by the grants DMS-1559338 and RFBR-13-01-00755.

the argument. This can be seen in [4], where three different functions I , J and R denote very similar objects. In the link case the situation will be similar. The function called H will take as an argument the levels of the Alexander filtration in the chain complex CFL^- , that is, its arguments will be from some lattice. Shifting the argument of H by half the linking numbers will yield a function J from \mathbb{Z}^n to \mathbb{Z} . The normalization of the J -function makes it very suitable for studying link concordances. Finally, we will have a function R , defined for algebraic singularities, which most resembles the semigroup counting function from [4] and agrees with the Hilbert function from [14].

We define the H -function for general links and find inequalities between the H -functions of two links related by an immersed concordance (under some assumptions on the concordance). The following theorem is one of the main results of the paper. The statement is simpler in terms of the J -function than in terms of the H -function.

Theorem (Theorem 6.20). *Let \mathcal{L}_1 and \mathcal{L}_2 be two n -component links differing by a single positive crossing change, that is, \mathcal{L}_2 arises by changing a negative crossing of \mathcal{L}_1 into a positive one. Let J_1 and J_2 be the corresponding J -functions and let $\mathbf{m} \in \mathbb{Z}^n$, $\mathbf{m} = (m_1, \dots, m_n)$.*

(a) *If the crossing change is between two strands of the same component L_{1i} of \mathcal{L}_1 , then*

$$J_2(m_1, m_2, \dots, m_i + 1, \dots, m_n) \leq J_1(m_1, \dots, m_n) \leq J_2(m_1, \dots, m_i, \dots, m_n).$$

(b) *If the crossing change is between the i -th and j -th component of \mathcal{L}_1 , then*

$$J_2(m_1, m_2, \dots, m_n) \leq J_1(m_1, \dots, m_n) \leq J_2(m_1, \dots, m_i - 1, \dots, m_n)$$

and

$$J_2(m_1, m_2, \dots, m_n) \leq J_1(m_1, \dots, m_n) \leq J_2(m_1, \dots, m_j - 1, \dots, m_n)$$

As an application we provide new criteria for splitting numbers of links.

Theorem (Theorem 7.7).

(a) *Suppose that a two component link $\mathcal{L} = L_1 \cup L_2$ can be unlinked using b_+ positive and b_- negative crossing changes. Let g_i denote the slice genus of L_i . Define vectors*

$$\mathbf{g} := (g_1, g_2), \quad \tilde{\mathbf{g}} := \left(g_1 + \frac{1}{2} \text{lk}(L_1, L_2), g_2 + \frac{1}{2} \text{lk}(L_1, L_2) \right).$$

Define the region $R(b_+)$ by inequalities:

$$R(b_+) := \{(m_1, m_2) : m_1 + m_2 \geq b_+, m_1 \geq 0, m_2 \geq 0\}.$$

Then $J(\mathbf{m}) = \tilde{J}(\mathbf{m}) = 0$ for $\mathbf{m} \in R(b_+) + \mathbf{g}$.

(b) *If, in addition, \mathcal{L} is an L-space link, then*

$$\text{HFL}^-(\mathcal{L}, \mathbf{v}) = 0 \text{ for } \mathbf{v} \in R(b_+) + \tilde{\mathbf{g}} + (1, 1).$$

In particular, all coefficients of the Alexander polynomial vanish in $R(b_+) + \tilde{\mathbf{g}} + (\frac{1}{2}, \frac{1}{2})$.

In the examples we focus on a family of two-bridge links which were shown in [23] to be L-space links. Recall that the *splitting number* $sp(\mathcal{L})$ is the minimal number of crossings between *different* components of \mathcal{L} that should be changed to turn \mathcal{L} into a split link. Clearly, a linking number yields a lower bound for the splitting number, see Section 7 for the detailed discussion of splitting numbers and references.

Theorem (Theorem 7.12). *The splitting number of the two-component two-bridge link*

$$\mathcal{L}_n = b(4n^2 + 4n, -2n - 1)$$

equals $2n$, although the linking number between the components of \mathcal{L}_n vanishes.

We compare this theorem with the work of Batson and Seed [1] which provides a different bound for the splitting number in terms of Khovanov homology. It turns out that their lower bound is quite weak in this case and is at most three for all \mathcal{L}_n .

Remark 1.1. Almost at the same time as the first version of this paper was posted on the arXiv, a similar result was proven by Cimasoni, Conway and Zacharova [10] by completely different methods. In Section 7.6 we compute splitting numbers for a family of links where the methods of [10] cannot be applied directly.

Another application is a topological proof of semicontinuity of the Hilbert function of singularities under δ -constant deformations. The result was proved in [5] for unibranched singular points (there is also an algebraic proof of a more general version in [14] for one component links). Our result is for multibranched singularities under the assumption that the number of branches does not change.

1.2. Structure of the paper. The paper uses a lot of background facts about Heegaard Floer homology and L-space links, most of them were discussed in [14, 23] using slightly different sets of notations. For the reader's convenience, we repeat these facts and introduce the functions H and J in full generality in Sections 2 and 3. In Section 4, we relate the Ozsváth-Szabó d -invariants of large surgeries on a link to the H -function. Section 5 is the technical core of the paper: for an immersed cobordism between two links $\mathcal{L}_1, \mathcal{L}_2$ we construct a cobordism between the surgeries $S_{\mathbf{q}_1}^3(\mathcal{L}_1), S_{\mathbf{q}_2}^3(\mathcal{L}_2)$ of the 3-sphere on these links, and prove that under certain assumptions it is negative definite. In the negative definite case, we apply the classical inequality for d -invariants of $S_{\mathbf{q}_1}^3(\mathcal{L}_1), S_{\mathbf{q}_2}^3(\mathcal{L}_2)$, and obtain in Section 6 an inequality for H and J -functions for the links $\mathcal{L}_1, \mathcal{L}_2$ stated in Theorem 6.1. We use this result to prove Theorem 6.20.

In Section 7 we apply these results to obtain more concrete inequalities for two-component links, and prove Theorems 7.7 and 7.12. Finally, in the last section we apply the inequalities to algebraic links and compare them with the algebro-geometric results on deformations of singularities.

1.3. Notations and conventions. All links are assumed to be oriented. For a link \mathcal{L} , we denote by L_1, \dots , its components. This allows us to make a distinction between $\mathcal{L}_1, \mathcal{L}_2$ and L_1, L_2 . The former denotes two distinct links, the latter stands for two components of the same link \mathcal{L} .

We will mark vectors in n -dimensional lattices in bold, in particular, we will write $\mathbf{0} = (0, \dots, 0)$. Given $\mathbf{u}, \mathbf{v} \in \mathbb{Z}^n$, we write $\mathbf{u} \leq \mathbf{v}$ if $u_i \leq v_i$ for all i , and $\mathbf{u} < \mathbf{v}$ if $\mathbf{u} \leq \mathbf{v}$ and $\mathbf{u} \neq \mathbf{v}$. We write $\mathbf{u} \ll \mathbf{v}$ if $u_i \ll v_i$ for all i . We will write $\mathbf{w} = \max(\mathbf{u}, \mathbf{v})$ (resp. $\mathbf{w} = \min(\mathbf{u}, \mathbf{v})$) if $w_i = \max(u_i, v_i)$ (resp. $w_i = \min(u_i, v_i)$) for all i . We denote the i -th coordinate vector by \mathbf{e}_i .

For a subset $I = \{i_1, \dots, i_r\} \subset \{1, \dots, n\}$ and $\mathbf{u} \in \mathbb{Z}^n$, we denote by \mathbf{u}_I the vector $(u_{i_1}, \dots, u_{i_r})$. For a link $\mathcal{L} = \bigcup_{i=1}^n L_i$ we denote by $\mathcal{L}_I = L_{i_1} \cup \dots \cup L_{i_r}$ the corresponding sublink.

We will always work with $\mathbb{F} = \mathbb{Z}/2\mathbb{Z}$ coefficients.

Acknowledgements. The authors would like to thank to David Cimasoni, Anthony Conway, Stefan Friedl, Jennifer Hom, Yajing Liu, Charles Livingston, Wojciech Politarczyk and

Mark Powell for fruitful discussions. The project was started during a singularity theory conference in Edinburgh in July 2015. The authors would like to thank the ICMS for hospitality.

2. LINKS AND L-SPACES

2.1. Links and their Alexander polynomials. Let $\mathcal{L} \subset S^3$ be a link. Denote by L_1, \dots, L_n its components. Throughout the paper, the multivariable Alexander polynomial (see [17] for definition) will be symmetric up to sign:

$$\Delta(t_1^{-1}, \dots, t_n^{-1}) = \pm \Delta(t_1, \dots, t_n).$$

The sign of a multivariable Alexander polynomial can be fixed using the interpretation of the Alexander polynomial via the sign refined Reidemeister torsion; see [17, Section 4.9] for discussion and [36] for an introduction to Reidemeister torsion.

Example 2.1. The Alexander polynomial for the Whitehead link equals

$$\Delta_{Wh}(t_1, t_2) = -(t_1^{1/2} - t_1^{-1/2})(t_2^{1/2} - t_2^{-1/2}).$$

For the Borromean link the Alexander polynomial equals

$$\Delta_{Bor}(t_1, t_2, t_3) = (t_1^{1/2} - t_1^{-1/2})(t_2^{1/2} - t_2^{-1/2})(t_3^{1/2} - t_3^{-1/2}).$$

In some examples we will consider *algebraic links*, defined as intersections of complex plane curve singularities with a small 3-sphere. The Alexander polynomials of algebraic links were computed by Eisenbud and Neumann [12]. In Section 8 below we also discuss more recent results of Campillo, Delgado and Gusein-Zade [6], relating the Alexander polynomial to the algebraic invariants of a singularity, such as the multi-dimensional semigroup.

Example 2.2. The link of the singularity $x^2 = y^{2n}$ consists of 2 unknots with linking number n . The corresponding Alexander polynomial equals

$$\Delta_{2,2n}(t_1, t_2) = \frac{t_1^{n/2}t_2^{n/2} - t_1^{-n/2}t_2^{-n/2}}{t_1^{1/2}t_2^{1/2} - t_1^{-1/2}t_2^{-1/2}}.$$

For future reference we recall the Torres formula, proved first in [34]. It relates the Alexander polynomial of a link \mathcal{L} with the Alexander polynomial of its sublink.

Theorem 2.3 (Torres Formula). *Let $\mathcal{L} = L_1 \cup \dots \cup L_n$ be an n component link and let $\mathcal{L}' = L_1 \cup \dots \cup L_{n-1}$. The Alexander polynomials of \mathcal{L}' and of \mathcal{L} are related by the following formula.*

$$\Delta_{\mathcal{L}}(t_1, \dots, t_{n-1}, 1) = \begin{cases} \left(\prod_{i=1}^{n-1} t_i^{\frac{1}{2} \text{lk}(L_i, L_n)} - \prod_{i=1}^{n-1} t_i^{-\frac{1}{2} \text{lk}(L_i, L_n)} \right) \Delta_{\mathcal{L}'}(t_1, \dots, t_{n-1}) & \text{if } n > 2, \\ \frac{\left(t_1^{\frac{1}{2} \text{lk}(L_1, L_2)} - t_1^{-\frac{1}{2} \text{lk}(L_1, L_2)} \right)}{\left(t_1^{\frac{1}{2}} - t_1^{-\frac{1}{2}} \right)} \Delta_{\mathcal{L}'}(t_1) & \text{if } n = 2, \end{cases}$$

where $\text{lk}(L_i, L_n)$ is the linking number between L_i and L_n .

Theorem 2.3 shows that if $\text{lk}(L_i, L_j)$ are all nonzero then the Alexander polynomials of the sublinks are determined by the Alexander polynomial of the link. If some of the linking numbers vanish (as in Example 2.1) then this is no longer true.

2.2. L-spaces and L-space links. We will use the minus version of the Heegaard Floer link homology, defined in [30]. To fix the conventions, we assume that $\text{HF}^-(S^3) = \mathbb{F}[U]$ is supported in degrees $0, -2, -4, \dots$. To every 3-manifold M this theory associates a chain complex $CF^-(M)$ which naturally splits as a direct sum over Spin^c structures on M : $CF^-(M) = \bigoplus_s CF^-(M, s)$. The homology $\text{HF}^-(M) = \bigoplus_s \text{HF}^-(M, s)$, as a graded $\mathbb{F}[U]$ -module, is a topological invariant of M .

Definition 2.4.

- (a) A 3-manifold M is called an L-space if $b_1(M) = 0$ and its Heegaard Floer homology has minimal possible rank: $\text{HF}^-(M, s) \simeq \mathbb{F}[U]$ for all s .
- (b) A link \mathcal{L} is called an L-space link if $S_{\mathbf{q}}^3(\mathcal{L})$, the integral surgery of S^3 on the components of \mathcal{L} with coefficients $\mathbf{q} = (q_1, \dots, q_n)$, is an L-space for $\mathbf{q} \gg \mathbf{0}$.

For a link $\mathcal{L} = L_1 \cup \dots \cup L_n$ and a vector $\mathbf{m} \in \mathbb{Z}^n$ we define the framing matrix $\Lambda(\mathbf{m}) = (\Lambda_{ij}(\mathbf{m}))$:

$$(2.5) \quad \Lambda_{ij}(\mathbf{m}) = \begin{cases} \text{lk}(L_i, L_j) & \text{if } i \neq j, \\ m_i & \text{if } i = j. \end{cases}$$

It is well known that if $\det \Lambda \neq 0$ then $|H_1(S_{\mathbf{m}}^3(\mathcal{L}))| = |\det \Lambda(\mathbf{m})|$. We recall the following result of Liu.

Theorem 2.6 (see [23, Lemma 2.5]). *Suppose $\mathcal{L} = L_1 \cup \dots \cup L_n$ is a link. Let $\mathbf{m} = (m_1, \dots, m_n)$ be a framing such that*

- (a) *The framing matrix $\Lambda(\mathbf{m})$ is positive definite.*
- (b) *For every $I \subset \{1, \dots, n\}$ the \mathbf{m}_I -surgery on \mathcal{L}_I is an L-space.*

Then for any integer vector $\mathbf{m}' \geq \mathbf{m}$ the \mathbf{m}' -surgery on \mathcal{L} is an L-space. In particular, \mathcal{L} is an L-space link.

We will generalize this result for rational surgeries.

Proposition 2.7. *Suppose \mathcal{L} and \mathbf{m} are as in the statement of Theorem 2.6. Then for any rational framing vector $\mathbf{q} \geq \mathbf{m}$, the \mathbf{q} -surgery on \mathcal{L} is an L-space.*

Proof. For a surgery vector \mathbf{q} denote by $t(\mathbf{q})$ the number of non-integer entries in the vector \mathbf{q} .

Let us make the following statement.

$(I_{k,l})$ *For any $I \subset \{1, \dots, n\}$ with $|I| \leq l$, if $t(\mathbf{q}_I) \leq k$, then $S_{\mathbf{q}_I}^3(\mathcal{L}_I)$ is an L-space.*

The statement $(I_{0,l})$ is covered for all $l \geq 1$ by Theorem 2.6. Our aim is to show that $(I_{k,l})$ implies $(I_{k+1,l})$.

Choose $I \subset \{1, \dots, n\}$ with $|I| = l$. Take $\mathbf{q} \geq \mathbf{m}$ with $t(\mathbf{q}) = k + 1$. Suppose $j \in I$ is such that $q_j \notin \mathbb{Z}$ and let $I' = I \setminus \{j\}$. Let $Y = S_{\mathbf{q}_{I'}}^3(\mathcal{L}_{I'})$. As $t(\mathbf{q}_{I'}) = k$, the assumption $(I_{k,l-1})$ (which is contained in $(I_{k,l})$) implies that Y is an L-space. The component L_j can be regarded as a knot in Y . Let $\mathcal{A} \subset \mathbb{Q} \cup \{\infty\}$ be the set of surgery coefficients such that $a \in \mathcal{A}$ if and only if $Y_a(L_j)$ is an L-space. By the inductive assumption all integers $l \geq m_j$ belong to \mathcal{A} , indeed $Y_l(L_1)$ is the surgery on \mathcal{L} with coefficient \mathbf{q}'_I , where \mathbf{q}'_I is the vector \mathbf{q}_I with l at the j -th position. Furthermore $\infty \in \mathcal{A}$ as well, because Y itself is an L-space.

In [32] possible shapes of \mathcal{A} were classified. In short, \mathcal{A} is either empty, or it consists of one point or it is an interval in $\mathbb{Q} \cup \{\infty\} \setminus I$, where I is the set of surgery coefficients l such that $Y_l(L_1)$ is not a rational homology sphere. In the present situation, we have

shown that $m_1, m_1 + 1, \infty$ belong to \mathcal{A} . This implies all rational numbers greater than or equal to m_1 are in \mathcal{A} . This shows $(I_{k+1,l})$. \square

As an application of Proposition 2.7 we will prove the following result, which generalizes [16, Theorem 1.10].

Proposition 2.8. *Suppose $\mathcal{L} = L_1 \cup \dots \cup L_n$ is an L-space link. Let p, q be coprime positive integers and let $\mathcal{L}_{p,q}$ be the link $L_{cab} \cup L_2 \cup \dots \cup L_n$, where L_{cab} is the (p, q) cable on L_1 . If q/p is sufficiently large, then $\mathcal{L}_{p,q}$ is also an L-space link. More precisely, if \mathbf{m} is an integer vector satisfying the conditions of Theorem 2.6 then $\mathcal{L}_{p,q}$ is an L-space link if $q/p > m_1$.*

Proof. The proof is a direct generalization of [16, Proof of Theorem 1.10]. Choose p and q coprime and suppose that \mathbf{m} satisfies the conditions of Theorem 2.6 and $q/p > m_1$. First we will show that the \mathbf{m}' -surgery on $\mathcal{L}_{p,q}$ is an L-space, where $\mathbf{m}' = (pq, m_2, \dots, m_n)$. By [16, Section 2.4] we know that $S_{\mathbf{m}'}^3(\mathcal{L}_{p,q}) \simeq \text{Lens}(p, q) \# S_{\mathbf{m}''}^3(\mathcal{L})$, where we set $\mathbf{m}'' = (q/p, m_2, \dots, m_n)$ and $\text{Lens}(p, q)$ is the lens space. As $\text{Lens}(p, q)$ is an L-space and since a connected sum of L-spaces is an L-space, it is enough to show that $S_{\mathbf{m}''}^3(\mathcal{L})$ is an L-space. But $\mathbf{m}'' \geq \mathbf{m}$, so by Proposition 2.7 we conclude that $S_{\mathbf{m}''}^3(\mathcal{L})$ is an L-space. Hence the \mathbf{m}' -surgery on $\mathcal{L}_{p,q}$ is an L-space. The same proof applies to any sublink of $\mathcal{L}_{p,q}$ which contains L_{cab} , and for a sublink \mathcal{L}_I not containing L_{cab} the \mathbf{m}'_I -surgery is an L-space by assumption.

Let Λ_{pq} be the framing matrix for $\mathcal{L}_{p,q}$ with framing \mathbf{m}' , let Λ be the framing matrix for \mathcal{L} with framing \mathbf{m} . By assumption, Λ is positive definite. The matrix Λ_{pq} differs from Λ only at the first column and at the first row. As $\text{lk}(L_{cab}, L_j) = p \text{lk}(L_1, L_j)$ for $j = 2, \dots, n$, we conclude that Λ_{pq} can be obtained from Λ by multiplying the first row and the first column by p (the element in the top-left corner is multiplied by p^2) and then adding $qp - p^2 m_1$ to the element in the top-left corner. The first operation is a matrix congruence so it preserves positive definiteness of the matrix. Adding an element can be regarded as taking a sum with a matrix with all entries zero but $qp - p^2 m_1$ in the top-left corner. This matrix is positive semi-definite, because we assumed that $q/p > m_1$. Now a sum of a positive definite matrix and a positive semi-definite one is a positive definite matrix. Therefore Λ_{pq} is positive definite.

By Theorem 2.6 applied to $\mathcal{L}_{p,q}$ with framing \mathbf{m}' we conclude that $\mathcal{L}_{p,q}$ is an L-space link. \square

To make Proposition 2.8 more concrete, we have to present an explicit vector \mathbf{m} satisfying the conditions of Theorem 2.6. This is done in the following theorem.

Theorem 2.9. *Let D_i denote the maximal degree of t_i in the multivariable Alexander polynomial of an L-space link \mathcal{L} , $m_i \geq 2D_i + 2$. Assume that the number of components of \mathcal{L} equals $n > 1$ and $\text{lk}(L_i, L_j) \neq 0$ for all $i \neq j$. Then $\mathbf{m} = (m_1, \dots, m_n)$ satisfies the conditions of Theorem 2.6.*

Proof. By Theorem 2.6 it is sufficient to prove that $S_{\mathbf{m}}^3(\mathcal{L})$ is an L-space and the framing matrix $\Lambda(\mathbf{m})$ is positive definite. The former is proved below as Lemma 3.21. To prove the latter, remark that by Theorem 2.3 one has:

$$(2.10) \quad \frac{\Delta(L_1)}{t^{1/2} - t^{-1/2}} = \frac{\Delta(t, 1, \dots, 1)}{\prod_{j \neq 1} \left(t^{\frac{1}{2} \text{lk}(L_1, L_j)} - t^{-\frac{1}{2} \text{lk}(L_1, L_j)} \right)},$$

so

$$2D_i - \sum_{j \neq i} \text{lk}(L_i, L_j) + 1 \geq 2 \deg \Delta(L_i) \geq 0.$$

Now $\Lambda(\mathbf{m})$ is a sum of $\binom{n}{2}$ nonnegative definite matrices

$$\begin{pmatrix} \text{lk}(L_i, L_j) & \text{lk}(L_i, L_j) \\ \text{lk}(L_i, L_j) & \text{lk}(L_i, L_j) \end{pmatrix}$$

with the only nonzero block at i -th and j -th rows and columns, and a diagonal positive definite matrix with entries

$$m_i - \sum_{j \neq i} \text{lk}(L_i, L_j) \geq 2D_i + 2 - \sum_{j \neq i} \text{lk}(L_i, L_j) > 0,$$

so it is positive definite. \square

Remark 2.11. This bound is far from being optimal for links with many components. For example, it is proved in [13] that the point $(pq+1, \dots, pq+1)$ satisfies the conditions of Theorem 2.6 for the (pn, qn) torus link, while in the above bound one has $D_i = (npq - p - q)/2$ for $n > 2$. On the other hand, for the $(2, 2q)$ torus link we get $D_1 = D_2 = (q-1)/2$, so Theorem 2.9 gives $m_i \geq q+1$, and the two bounds agree.

3. HEEGAARD FLOER LINK HOMOLOGY AND THE H -FUNCTION FOR LINKS

In this section we define the H -function for links and collect some useful facts about it.

3.1. Alexander filtration. A knot K in a 3-manifold M induces a filtration on the Heegaard Floer complex $CF^-(M)$. Similarly, a link $\mathcal{L} = L_1 \cup \dots \cup L_n$ with n components in M induces n different filtrations on $CF^-(M)$, which can be interpreted as a filtration indexed by an n -dimensional lattice. For a link in S^3 , it is natural to make this lattice different from \mathbb{Z}^n .

Definition 3.1. Given an n -component oriented link $\mathcal{L} \subset S^3$, we define an affine lattice over \mathbb{Z}^n :

$$\mathbb{H}(\mathcal{L}) = \bigoplus_{i=1}^n \left[\mathbb{Z} + \frac{1}{2} \text{lk}(L_i, \mathcal{L} \setminus L_i) \right].$$

We also define the *linking vector*:

$$\boldsymbol{\ell} = \boldsymbol{\ell}(\mathcal{L}) = \frac{1}{2}(\text{lk}(L_1, \mathcal{L} \setminus L_1), \text{lk}(L_2, \mathcal{L} \setminus L_2), \dots, \text{lk}(L_n, \mathcal{L} \setminus L_n)).$$

We have $\mathbb{H}(\mathcal{L}) = \mathbb{Z}^n + \boldsymbol{\ell}$.

For $\mathbf{v} \in \mathbb{H}(\mathcal{L})$ define a subcomplex $A^-(\mathcal{L}, \mathbf{v}) \subset CF^-(S^3)$ corresponding to the filtration level \mathbf{v} . The filtration is ascending, so $A^-(\mathcal{L}, \mathbf{u}) \subset A^-(\mathcal{L}, \mathbf{v})$ for $\mathbf{u} \leq \mathbf{v}$. The Heegaard Floer link homology $HFL^-(\mathcal{L}, \mathbf{v})$ can be defined as the homology of the associated graded complex:

$$(3.2) \quad HFL^-(\mathcal{L}, \mathbf{v}) = H_* \left(A^-(\mathcal{L}; \mathbf{v}) / \sum_{\mathbf{u} < \mathbf{v}} A^-(\mathcal{L}; \mathbf{u}) \right).$$

The Euler characteristic of this homology was computed in [30]:

$$(3.3) \quad \tilde{\Delta}(t_1, \dots, t_n) := \sum_{\mathbf{v} \in \mathbb{H}(\mathcal{L})} \chi(\text{HFL}^-(\mathcal{L}, \mathbf{v})) t_1^{v_1} \cdots t_n^{v_n} =$$

$$= \begin{cases} (t_1 \cdots t_n)^{1/2} \Delta(t_1, \dots, t_n) & \text{if } n > 1, \\ \Delta(t)/(1 - t^{-1}) & \text{if } n = 1, \end{cases}$$

where, as above, $\Delta(t_1, \dots, t_n)$ denotes the symmetrized Alexander polynomial of \mathcal{L} .

One can forget a component L_n in \mathcal{L} and consider the $(n-1)$ -component link $\mathcal{L} \setminus L_n$. There is a natural forgetful map $\pi_n : \mathbb{H}(\mathcal{L}) \rightarrow \mathbb{H}(\mathcal{L} \setminus L_n)$ defined by the equation:

$$(3.4) \quad \pi_n(v_1, \dots, v_n) = \left(v_1 - \frac{1}{2} \text{lk}(L_1, L_n), \dots, v_{n-1} - \frac{1}{2} \text{lk}(L_{n-1}, L_n) \right).$$

In general, one defines a map $\pi_{\mathcal{L}'} : \mathbb{H}(\mathcal{L}) \rightarrow \mathbb{H}(\mathcal{L}')$ for every sublink $\mathcal{L}' \subset \mathcal{L}$:

$$(3.5) \quad [\pi_{\mathcal{L}'}(v)]_j = (v_j - \ell(\mathcal{L})_j + \ell(\mathcal{L}')_j) \quad \text{for } L_j \subset \mathcal{L}'.$$

For $v_n \gg 0$ the subcomplexes $A^-(\mathcal{L}; \mathbf{v})$ stabilize, and by [30, Proposition 7.1] one has a natural homotopy equivalence $A^-(\mathcal{L}; \mathbf{v}) \sim A^-(\mathcal{L} \setminus L_n; \pi_n(\mathbf{v}))$. More generally, for a sublink $\mathcal{L}' = L_{i_1} \cup \dots \cup L_{i_r}$, one gets:

$$(3.6) \quad A^-(\mathcal{L}'; \pi_{\mathcal{L}'}(\mathbf{v})) \sim A^-(\mathcal{L}; \mathbf{v}), \text{ if } v_i \gg 0 \text{ for all } i \notin \{i_1, \dots, i_r\}.$$

There is an action of commuting operators U_1, \dots, U_n on the complex $A^-(\mathcal{L})$. The action of U_i drops the homological grading by 2 and drops the i -th filtration level by 1. In particular, $U_i A^-(\mathcal{L}, \mathbf{v}) \subset A^-(\mathcal{L}, \mathbf{v} - \mathbf{e}_i)$. This action makes the complexes $A^-(\mathcal{L}, \mathbf{v})$ modules over the polynomial ring $\mathbb{F}[U_1, \dots, U_n]$. It is known [30] that $A^-(\mathcal{L}, \mathbf{v})$ is a finitely generated module over $\mathbb{F}[U_1, \dots, U_n]$, and the action of all the U_i on $A^-(\mathcal{L}, \mathbf{v})$ is homotopic to each other. In particular, all the U_i act in the same way in the homology $H_*(A^-(\mathcal{L}, \mathbf{v}))$, which can therefore be naturally considered as $\mathbb{F}[U]$ -module, where a single variable U acts as U_1 .

3.2. The H -function. It is known (see [24], this is also a consequence of the Large Surgery Theorem 4.7 below) that the homology of $A^-(\mathcal{L}, \mathbf{v})$ is isomorphic as an $\mathbb{F}[U]$ -module to the Heegaard Floer homology of a large surgery on \mathcal{L} equipped with a certain Spin^c structure. Therefore it always splits as a direct sum of a single copy of $\mathbb{F}[U]$ and some U -torsion. We begin with the following fact.

Lemma 3.7. *For $\mathbf{u} \leq \mathbf{v}$ the natural inclusion*

$$\iota_{\mathbf{u}, \mathbf{v}} : A^-(\mathcal{L}, \mathbf{u}) \hookrightarrow A^-(\mathcal{L}, \mathbf{v})$$

is injective on the free parts of the homology, hence it is a multiplication by a nonnegative power of U .

Proof. It is sufficient to prove that

$$\iota_{\mathbf{v} - \mathbf{e}_i, \mathbf{v}}^* : H_*(A^-(\mathcal{L}, \mathbf{v} - \mathbf{e}_i)) \hookrightarrow H_*(A^-(\mathcal{L}, \mathbf{v}))$$

is injective on the free parts. The latter holds because $A^-(\mathcal{L}, \mathbf{v} - \mathbf{e}_i)$ contains the image of $U_i \sim U$ acting on $A^-(\mathcal{L}, \mathbf{v})$. Indeed, if $H_*(A^-(\mathcal{L}, \mathbf{v})) \simeq \mathbb{F}[U] + T(\mathbf{v})$, where $T(\mathbf{v})$ is U -torsion, then $U\mathbb{F}[U] \subset H_*(U_i A^-(\mathcal{L}, \mathbf{v}))$. Consider the inclusions $U_i A^-(\mathcal{L}, \mathbf{v}) \subset A^-(\mathcal{L}, \mathbf{v} - \mathbf{e}_i) \subset A^-(\mathcal{L}, \mathbf{v})$. Since the composite inclusion of $U_i A^-(\mathcal{L}, \mathbf{v})$ into $A^-(\mathcal{L}, \mathbf{v})$ is injective on free parts, we conclude that $\iota_{\mathbf{v} - \mathbf{e}_i, \mathbf{v}}^*$ is injective and

$$(3.8) \quad U\mathbb{F}[U] \subset \iota_{\mathbf{v} - \mathbf{e}_i, \mathbf{v}}^* \mathbb{F}[U] \subset \mathbb{F}[U]$$

□

Definition 3.9. We define a function $H(\mathbf{v}) = H_{\mathcal{L}}(\mathbf{v})$ by saying that $-2H(\mathbf{v})$ is the maximal homological degree of the free part of $H_*(A^-(\mathcal{L}, \mathbf{v}))$.

We will gather now some important properties of the H -function.

Proposition 3.10. *The function $H(\mathbf{v})$ has nonnegative integer values. Furthermore, for all $\mathbf{v} \in \mathbb{H}(\mathcal{L})$ one has $H(\mathbf{v} - \mathbf{e}_i) = H(\mathbf{v})$ or $H(\mathbf{v} - \mathbf{e}_i) = H(\mathbf{v}) + 1$.*

Proof. By Lemma 3.7 the inclusion of $A^-(\mathcal{L}, \mathbf{v})$ in $CF^-(S^3)$ induces an injective map on the free parts of the homology, so it sends a generator of the free part to U^k times a generator of the free part for some $k \geq 0$. Since the inclusion preserves the homological grading (and the generator of $HF^-(S^3)$ has grading 0), the generator of the free part of $H_*(A^-(\mathcal{L}, \mathbf{v}))$ has grading $-2k$, and $k = H(\mathbf{v})$. The last statement immediately follows from (3.8). □

Proposition 3.11. *If \mathcal{L} is a split link then $H(\mathbf{v}) = \sum_{i=1}^n H_i(v_i)$, where H_i is the H -function for the i -th component of the link.*

Proof. For a split link by [30, Section 11] one has

$$A^-(\mathcal{L}, \mathbf{v}) \simeq A^-(L_1, v_1) \otimes_{\mathbb{F}[U]} \cdots \otimes_{\mathbb{F}[U]} A^-(L_n, v_n),$$

and the isomorphism preserves the homological gradings. Note that all the linking numbers of a split link vanish, so $\mathbb{H}(\mathcal{L}) = \mathbb{Z}^n$, and the projections to sublattices do not require any shifts as in (3.6). □

Proposition 3.12. *For a sublink $\mathcal{L}' = L_{i_1} \cup \dots \cup L_{i_r}$, one has*

$$(3.13) \quad H_{\mathcal{L}}(\mathbf{v}) = H_{\mathcal{L}'}(\pi_{\mathcal{L}'}(\mathbf{v})) \text{ if } v_i \gg 0 \text{ for } i \notin \{i_1, \dots, i_r\}.$$

Proof. Follows from (3.6). □

3.3. The H -function for L-space links. By Theorem 4.7 (see also [23]), a link is an L-space link if and only if $H_*(A^-(\mathbf{v})) \simeq \mathbb{F}[U]$ for all $\mathbf{v} \in \mathbb{H}(\mathcal{L})$. It turns out that for L-space links the H -function is determined by the Alexander polynomial.

Throughout Section 3.3 we will assume that \mathcal{L} is an L-space link. Since $H_*(A^-(\mathbf{v})) \simeq \mathbb{F}[U]$ for all $\mathbf{v} \in \mathbb{H}(\mathcal{L})$, by (3.2) and by the inclusion-exclusion formula one can write:

$$(3.14) \quad \chi(\text{HFL}^-(\mathcal{L}, \mathbf{v})) = \sum_{B \subset \{1, \dots, n\}} (-1)^{|B|-1} H_{\mathcal{L}}(\mathbf{v} - \mathbf{e}_B),$$

where \mathbf{e}_B denotes the characteristic vector of the subset $B \subset \{1, \dots, n\}$; see [13, formula (3.3)]. For $n = 1$ equation (3.14) has the form $\chi_{\mathcal{L}, \mathbf{v}} = H(\mathbf{v} - 1) - H(\mathbf{v})$, so $H(\mathbf{v})$ can be easily reconstructed from the Alexander polynomial: $H_{\mathcal{L}}(\mathbf{v}) = \sum_{\mathbf{u} \geq \mathbf{v}+1} \chi_{\mathcal{L}, \mathbf{u}}$. For $n > 1$, one can also show that equation (3.14) together with the boundary conditions (3.13) has a unique solution, which is given by the following theorem:

Theorem 3.15 ([14]). *The H -function of an L-space link is determined by the Alexander polynomials of its sublinks as following:*

$$(3.16) \quad H_{\mathcal{L}}(v_1, \dots, v_n) = \sum_{\mathcal{L}' \subseteq \mathcal{L}} (-1)^{\#\mathcal{L}'-1} \sum_{\substack{\mathbf{u}' \in \mathbb{H}(\mathcal{L}') \\ \mathbf{u}' \geq \pi_{\mathcal{L}'}(\mathbf{v}+1)}} \chi(\text{HFL}^-(\mathcal{L}', \mathbf{u}')),$$

where $\mathbf{1} = (1, \dots, 1)$.

There is a formula for the H -function in terms of the multivariable Alexander polynomial. Consider the generating function:

$$(3.17) \quad \mathbf{H}(t_1, \dots, t_n) := \sum_{v_1, \dots, v_n \in \mathbb{Z}^n} t_1^{v_1} \dots t_n^{v_n} H(v_1, \dots, v_n).$$

Note that \mathbf{H} is a Laurent series in $t_i^{-1/2}$: by (3.13) $H(\mathbf{v})$ vanishes if $v_i \gg 0$ for some i , but it does not vanish for $\mathbf{v} \ll \mathbf{0}$. Then [14, Theorem 3.4.3] implies

$$(3.18) \quad \mathbf{H}(t_1, \dots, t_n) = \prod_{i=1}^n \frac{1}{1 - t_i^{-1}} \sum_{\mathcal{L}' \subset \mathcal{L}} (-1)^{\#\mathcal{L}'-1} \tilde{\Delta}_{\mathcal{L}'}(t_{j_1}, \dots, t_{j_{\#\mathcal{L}'}}) \prod_{j: L_j \subset \mathcal{L}'} t_j^{\ell(\mathcal{L})_j - \ell(\mathcal{L}')_j - 1},$$

where $\tilde{\Delta}$ is defined by (3.3).

As above, let D_i denote the maximal t_i -degree of the Alexander polynomial of \mathcal{L} , $\mathbf{D} = (D_1, \dots, D_n)$.

Lemma 3.19. *Assume that $\text{lk}(L_i, L_j) \neq 0$, then $H(\mathbf{v}) = H(\min(\mathbf{v}, \mathbf{D}))$.*

Proof. By Theorem 2.3 the degree of the Alexander polynomial of a sublink $\mathcal{L}' = \mathcal{L}_I$ in variable t_i is less than or equal to $D_i - \frac{1}{2} \sum_{j \notin I} \text{lk}(L_i, L_j) = \pi_{\mathcal{L}'}(\mathbf{D})_i$. Therefore if $v_i > D_i$ then

$$\pi_{\mathcal{L}'}(\mathbf{v})_i > \pi_{\mathcal{L}'}(\mathbf{D})_i \geq \deg_{t_i} \Delta(\mathcal{L}'),$$

and if $\mathbf{u} \geq \pi_{\mathcal{L}'}(\mathbf{v}) + \mathbf{1}$ then $\chi(\text{HFL}^-(\mathcal{L}', \mathbf{u}')) = 0$. Therefore the summands contributing to (3.16) nontrivially correspond to subsets I such that $v_i \leq D_i$ for $i \in I$. Applying (3.16) to $\min(\mathbf{v}, \mathbf{D})$, one gets exactly the same summands. \square

Corollary 3.20. *For $\mathbf{v} \geq \mathbf{D}$ one has $H(\mathbf{v}) = 0$.*

Lemma 3.21. *Let \mathbf{D} be as above, then for $\mathbf{m} \geq 2\mathbf{D} + 2$ the surgery $S_{\mathbf{m}}^3(\mathcal{L})$ yields an L -space.*

Proof. Consider the parallelepiped P in \mathbb{Z}^n with opposite corners at \mathbf{D} and $-\mathbf{D}$. To compute the Heegaard Floer homology of $S_{\mathbf{m}}^3(\mathcal{L})$, we use the surgery complex of Manolescu-Ozsváth [24]. Every Spin^c structure on $S_{\mathbf{m}}^3(\mathcal{L})$ corresponds to an equivalence class of \mathbb{Z}^n modulo the lattice generated by the columns of $\Lambda(\mathbf{m})$. For $\mathbf{m} \geq 2\mathbf{D}$ this equivalence class has at most one point in P , and the whole surgery complex can be contracted to a single copy of $\mathbb{F}[U]$ supported at that point. For the precise description of the “truncation” procedure, we refer to [24, Section 8.3, Case I], where the constant b in [24, Lemma 8.8] can be chosen equal to \mathbf{D} by Lemma 3.19. \square

The following symmetry property of H , which generalizes the symmetry in the case of knots [26, 18], is proved in [23, Lemma 5.5].

Proposition 3.22. *For an L -space link one has*

$$(3.23) \quad H(-\mathbf{v}) = H(\mathbf{v}) + |\mathbf{v}|$$

The symmetry (3.23) and the projection formula (3.13) imply a useful “dual projection formula”.

Corollary 3.24. *Let \mathcal{L} be an L -space link, consider a set $I = \{i_1, \dots, i_r\} \subset \{1, \dots, n\}$ and the sublink \mathcal{L}_I . Then, as long as $v_j \ll 0$ for all $j \notin I$, the following holds:*

$$(3.25) \quad H_{\mathcal{L}}(\mathbf{v}) = H_{\mathcal{L}_I}(\pi_{\mathcal{L}_I}(\mathbf{v}) + 2\ell(\mathcal{L})_I - 2\ell(\mathcal{L}_I)) - \sum_{j \notin I} v_j + |\ell(\mathcal{L})_I - \ell(\mathcal{L}_I)|.$$

Proof. For $\mathbf{v} \in \mathbb{H}(\mathcal{L})$ set $\mathbf{v}_I = (v_{i_1}, \dots, v_{i_r})$. By (3.23), $H_{\mathcal{L}}(\mathbf{v}) = H_{\mathcal{L}}(-\mathbf{v}) - |\mathbf{v}|$. Since $-v_j \gg 0$ for $j \notin I$, the projection formula implies

$$H_{\mathcal{L}}(-\mathbf{v}) = H_{\mathcal{L}_I}(\pi_{\mathcal{L}_I}(-\mathbf{v})) = H_{\mathcal{L}_I}(-\mathbf{v}_I - \boldsymbol{\ell}(\mathcal{L})_I + \boldsymbol{\ell}(\mathcal{L}_I)),$$

and (3.23) for \mathcal{L}_I implies

$$\begin{aligned} H_{\mathcal{L}_I}(-\mathbf{v}_I - \boldsymbol{\ell}(\mathcal{L})_I + \boldsymbol{\ell}(\mathcal{L}_I)) &= H_{\mathcal{L}_I}(\mathbf{v}_I + \boldsymbol{\ell}(\mathcal{L})_I - \boldsymbol{\ell}(\mathcal{L}_I)) + |\mathbf{v}_I + \boldsymbol{\ell}(\mathcal{L})_I - \boldsymbol{\ell}(\mathcal{L}_I)| = \\ &= H_{\mathcal{L}_I}(\pi_{\mathcal{L}_I}(\mathbf{v}) + 2\boldsymbol{\ell}(\mathcal{L})_I - 2\boldsymbol{\ell}(\mathcal{L}_I)) + |\mathbf{v}_I| + |\boldsymbol{\ell}(\mathcal{L})_I - \boldsymbol{\ell}(\mathcal{L}_I)|. \end{aligned}$$

□

3.4. The J -function. The J -function of a link \mathcal{L} is essentially the same object as the H -function, only it differs from H by a shift in variables. This shift makes J a function on \mathbb{Z}^n instead of $\mathbb{H}(\mathcal{L})$. It is therefore more convenient to study changes of the J -function under some changes (like crossing changes) of the link \mathcal{L} : these changes might affect the lattice $\mathbb{H}(\mathcal{L})$. Yet another variant is the \tilde{J} -function, which turns out to be useful for bounding the splitting number of L-space links; see Section 7 for details.

Definition 3.26. The J -function of a link \mathcal{L} with n components is a function $J: \mathbb{Z}^n \rightarrow \mathbb{Z}$ given by

$$J(\mathbf{m}) = H(\mathbf{m} + \boldsymbol{\ell}), \quad \mathbf{m} \in \mathbb{Z}^n.$$

With this definition the projection formula (3.13) takes a particularly simple form.

Lemma 3.27. Let $\mathbf{m} \in \mathbb{Z}^n$ and $I \subset \{1, \dots, n\}$. Consider a sublink \mathcal{L}_I of \mathcal{L} and suppose that $m_i \gg 0$ for $i \notin I$. Then we have

$$J_{\mathcal{L}}(\mathbf{m}) = J_{\mathcal{L}_I}(\mathbf{m}_I).$$

Proof. Indeed, by (3.5) and (3.13):

$$J_{\mathcal{L}}(\mathbf{m}) = H_{\mathcal{L}}(\mathbf{m} + \boldsymbol{\ell}) = H_{\mathcal{L}_I}(\mathbf{m}_I + \boldsymbol{\ell}(\mathcal{L})_I - \boldsymbol{\ell}(\mathcal{L})_I + \boldsymbol{\ell}(\mathcal{L}_I)) = J_{\mathcal{L}_I}(\mathbf{m}_I).$$

□

In particular, the J -function of a component L_i can be reconstructed from the values of the J -function for \mathcal{L} evaluated on vectors whose all components but the i -th one are sufficiently large.

Definition 3.28. For $\mathbf{m} = (m_1, \dots, m_n) \in \mathbb{Z}^n$ define

$$\tilde{J}(\mathbf{m}) = J(\mathbf{m}) - \sum_{i=1}^n J_{L_i}(m_i).$$

The main feature of the \tilde{J} -function is the following corollary to Proposition 3.11.

Corollary 3.29. If \mathcal{L} is a split link, then $\tilde{J} = 0$.

For general L-space links the \tilde{J} -function can be calculated from the Alexander polynomial. We have the following result.

Proposition 3.30. Define the generating function

$$\tilde{\mathbf{J}}(t_1, \dots, t_n) = \sum_{m_1, \dots, m_n} t_1^{m_1} \dots t_n^{m_n} \tilde{J}(\mathbf{m}).$$

Then $\tilde{\mathbf{J}}$ is a Laurent series in $t_i^{-1/2}$ and the following equation holds:

$$(3.31) \quad \tilde{\mathbf{J}}(t_1, \dots, t_n) = \prod_{i=1}^n \frac{1}{1 - t_i^{-1}} \sum_{\substack{\mathcal{L}' \subset \mathcal{L} \\ \#\mathcal{L}' > 1}} (-1)^{\#\mathcal{L}' - 1} \Delta_{\mathcal{L}'}(t_{j_1}, \dots, t_{j_{\#\mathcal{L}'}}) \prod_{j: L_j \subset \mathcal{L}'} t_j^{-\ell(\mathcal{L}')_j - \frac{1}{2}}.$$

Proof. This is a consequence of previous definitions. The formula for the generating function for $J(\mathbf{m})$ immediately follows from (3.18). To get a generating function for $\tilde{\mathbf{J}}$ we need to subtract the sum of J -functions for components L_1, \dots, L_n of \mathcal{L} . We apply (3.18) again to calculate this contribution, and remark that for $r > 1$ one has $\tilde{\Delta}_{\mathcal{L}'} = \prod t_j^{1/2} \cdot \Delta'_{\mathcal{L}'}$. \square

Equation (3.31) takes a particularly simple form for a two-component link.

Corollary 3.32. *For a link with two components*

$$\tilde{\mathbf{J}}(t_1, t_2) = -\frac{(t_1 t_2)^{-\text{lk}(L_1, L_2)/2} \Delta(t_1, t_2)}{(t_1^{1/2} - t_1^{-1/2})(t_2^{1/2} - t_2^{-1/2})}.$$

3.5. Examples.

Example 3.33. Consider the Whitehead link. By [23, Example 3.1] it is an L-space link. The linking number vanishes. The symmetrized Alexander polynomial equals $\Delta(t_1, t_2) = -(t_1^{1/2} - t_1^{-1/2})(t_2^{1/2} - t_2^{-1/2})$ (see Example 2.1), so the nontrivial values of $\chi(\text{HFL}^-(\mathbf{v}))$ are

$$\chi(\text{HFL}^-(0, 0)) = -1, \quad \chi(\text{HFL}^-(1, 0)) = \chi(\text{HFL}^-(0, 1)) = 1, \quad \chi(\text{HFL}^-(1, 1)) = -1.$$

Furthermore, both components are unknots, so $\chi(\text{HFL}^-(\mathbf{v})) = 1$ for $\mathbf{v} = (v, 0)$ or $\mathbf{v} = (0, v)$ with $v \leq 0$, and $\chi(\text{HFL}^-(\mathbf{v})) = 0$ for $\mathbf{v} = (v, 0)$ or $\mathbf{v} = (0, v)$ for $v > 0$. The H -function of the components equals

$$H_k(v_k) = \sum_{j \geq v_k + 1} \chi(\text{HFL}^-(j)) = \max(-v_k, 0)$$

for $k = 1, 2$. By (3.16) we get

$$\begin{aligned} H(v_1, v_2) &= H_1(v_1) + H_2(v_2) - \sum_{\mathbf{u} \geq \mathbf{v} + \mathbf{1}} \chi(\text{HFL}^-(\mathbf{u})) = \\ &= \begin{cases} H_1(v_1) + H_2(v_2) + 1 = 1, & \text{if } \mathbf{v} = (0, 0) \\ H_1(v_1) + H_2(v_2), & \text{otherwise.} \end{cases} \end{aligned}$$

Example 3.34. Consider the Borromean link. By [23, Example 3.1] it is an L-space link. The linking number between components vanishes, so do all bivariate Alexander polynomials of sublinks. The trivariate symmetrized Alexander polynomial equals

$$\Delta(t_1, t_2, t_3) = (t_1^{1/2} - t_1^{-1/2})(t_2^{1/2} - t_2^{-1/2})(t_3^{1/2} - t_3^{-1/2}),$$

so

$$\chi(\text{HFL}^-(\mathcal{L}, \mathbf{v})) = \begin{cases} 1 & \text{if } \mathbf{v} = (1, 1, 1), (1, 0, 0), (0, 0, 1), (0, 1, 0) \\ -1 & \text{if } \mathbf{v} = (0, 1, 1), (1, 1, 0), (1, 0, 1), (0, 0, 0) \\ 0 & \text{otherwise} \end{cases}$$

By (3.16), we get

$$H(v_1, v_2, v_3) = \sum_{i=1}^3 H_i(v_i) + \sum_{\mathbf{u} \geq \mathbf{v}+1} \chi(\text{HFL}^-(\mathbf{u})) = \begin{cases} \sum_{i=1}^3 H_i(v_i) + 1 = 1 & \text{if } \mathbf{v} = (0, 0, 0) \\ \sum_{i=1}^3 H_i(v_i) & \text{otherwise.} \end{cases}$$

4. THE H -FUNCTION AND d -INVARIANTS

4.1. Ozsváth–Szabó d -invariants. Let Y be a rational homology three-sphere equipped with a Spin^c structure \mathfrak{s} . The d -invariant of (Y, \mathfrak{s}) is the maximal grading of an element $x \in HF^-(Y, \mathfrak{s})$, which maps non-trivially into $HF^\infty(Y, \mathfrak{s})$. It is a rational number. The usefulness of the d -invariant comes from two facts: firstly it behaves well under a negative definite Spin^c cobordism, secondly it can be calculated from the knot (or link) Floer chain complex; we will describe this in detail in Section 4.2 below. As for the behavior under cobordism, suppose that (Y_1, \mathfrak{s}_1) and (Y_2, \mathfrak{s}_2) are rational homology 3-spheres and W is a smooth 4-manifold with boundary $Y_2 \sqcup -Y_1$ endowed with a Spin^c structure \mathfrak{t} which restricts to \mathfrak{s}_2 on Y_2 and to \mathfrak{s}_1 on Y_1 , put differently, (W, \mathfrak{t}) is a Spin^c cobordism between (Y_1, \mathfrak{s}_1) and (Y_2, \mathfrak{s}_2) . The following result is proved in [28].

Theorem 4.1. *There exists a chain map $F_{(W, \mathfrak{t})}: CF^\infty(Y_1, \mathfrak{s}_1) \rightarrow CF^\infty(Y_2, \mathfrak{s}_2)$, commuting with multiplication by U and shifting the absolute grading by*

$$(4.2) \quad \deg F_{(W, \mathfrak{t})} = \frac{1}{4} (c_1^2(\mathfrak{t}) - 3\sigma(W) - 2\chi(W)),$$

The chain map descends to maps $HF^\infty(Y_1, \mathfrak{s}_1) \rightarrow HF^\infty(Y_2, \mathfrak{s}_2)$, $HF^-(Y_1, \mathfrak{s}_1) \rightarrow HF^-(Y_2, \mathfrak{s}_2)$ and $HF^+(Y_1, \mathfrak{s}_1) \rightarrow HF^+(Y_2, \mathfrak{s}_2)$ (which will still be denoted by $F_{(W, \mathfrak{t})}$). If additionally W is negative definite, that is, the intersection form on $H_2(W; \mathbb{Q})$ is negative definite, then $F_{(W, \mathfrak{t})}$ is an isomorphism on HF^∞ and

$$(4.3) \quad d(Y_2, \mathfrak{s}_2) \geq \deg F_{(W, \mathfrak{t})} + d(Y_1, \mathfrak{s}_1).$$

A standard way of interpreting $c_1^2(\mathfrak{t})$, provided $H_1(\partial W; \mathbb{Q}) = 0$, is to use an isomorphism $H^2(W; \mathbb{Q}) \cong H_2(W, \partial W; \mathbb{Q}) \cong H_2(W; \mathbb{Q})$ (the first part is Poincaré duality, the second is the long exact sequence of pair) to identify $c_1(\mathfrak{t})$ with an element in $H_2(W; \mathbb{Q})$. Then $c_1^2(\mathfrak{t})$ is a self-intersection of an element in $H_2(W; \mathbb{Q})$, so it is a rational number.

Remark 4.4. It follows from the definition that the degree is additive under the composition of Spin^c cobordisms.

The degree formula (4.2) will play an important role in this article. We will need the following fact, which is well known to the experts.

Proposition 4.5. *The degree $\deg F_{(W, \mathfrak{t})}$ is preserved under negative blow-ups. Namely, suppose $\pi: W' \rightarrow W$ is a blow-down map and E is the exceptional divisor. Let \mathfrak{t}' be the Spin^c structure of W' whose first Chern class is $\pi^*c_1(\mathfrak{t}) \pm PD[E]$, where PD stays for Poincaré dual. Then*

$$\deg F_{(W', \mathfrak{t}')} = \deg F_{(W, \mathfrak{t})}.$$

On the other hand, if $\pi: W' \rightarrow W$ is a blow-down of a $(+1)$ -sphere, then

$$\deg F_{(W', \mathfrak{t}')} = \deg F_{(W, \mathfrak{t})} - 1.$$

Proof. Let $s = [E]^2$, so $s = +1$ for the positive blow-up and $s = -1$ for the negative one. We have that $\chi(W') = \chi(W) + 1$, $c_1^2(\mathfrak{t}') = c_1^2(\mathfrak{t}) + s$ and $\sigma(W') = \sigma(W) + s$. The change of the degree is $\frac{1}{4}(s - 3s - 2) = -\frac{1}{2}(s + 1)$. \square

4.2. Large surgery theorem. The subcomplexes $A^-(\mathcal{L}, \mathbf{v})$ are naturally related to the surgeries of the 3-sphere on \mathcal{L} . Choose a framing vector $\mathbf{q} = (q_1, \dots, q_n)$ such that q_1, \dots, q_n are sufficiently large. Let Λ be the linking matrix of \mathcal{L} , that is $\Lambda_{ij} = \text{lk}(L_i, L_j)$ if $i \neq j$ and $\Lambda_{ii} = q_i$.

Form a four-manifold $X_{\mathbf{q}}$ by adding n two-handles to a ball B^4 : a handle with framing q_i is attached along the component L_i . The boundary $\partial X_{\mathbf{q}}$, denoted $Y_{\mathbf{q}}$, is the surgery on \mathcal{L} with framing \mathbf{q} . Let F_i be the surface obtained by gluing a core of the i -th handle to a Seifert surface for L_i . By construction the classes $[F_1], \dots, [F_n]$ freely generate $H_2(X_{\mathbf{q}})$. With this choice of generators, we identify $H_2(X_{\mathbf{q}})$ with \mathbb{Z}^n .

Suppose $\det \Lambda \neq 0$. In this case $Y_{\mathbf{q}}$ is a rational homology sphere. In [24, Section 8.5] there is given an enumeration of Spin^c structures on $Y_{\mathbf{q}}$, which we are now going to recall.

Fix $\zeta = (\zeta_1, \dots, \zeta_n)$, a sufficiently small real vector whose entries are linearly independent over \mathbb{Q} . Then let $P(\Lambda)$ be the hyper-parallelepiped with vertices

$$\zeta + \frac{1}{2}(\pm \Lambda_1 \pm \Lambda_2 \pm, \dots, \pm \Lambda_n),$$

where all combinations of the signs are used and $\Lambda_1, \dots, \Lambda_n$ are column vectors of the matrix Λ . Then $P(\Lambda)$ is the fundamental domain for $\mathbb{R}^n/H(\Lambda)$, where $H(\Lambda)$ is the lattice generated by the vectors $\Lambda_1, \dots, \Lambda_n$. Denote

$$P_{\mathbb{H}}(\Lambda) = P(\Lambda) \cap \mathbb{H}(\mathcal{L}),$$

where $\mathbb{H}(\mathcal{L})$ is the lattice for \mathcal{L} as described in Definition 3.1 above. The set $P_{\mathbb{H}}(\Lambda)$ might depend on the choice of ζ , but its cardinality is independent of ζ ; see [24, Section 8.5] for more details.

Proposition 4.6 (see [24, Equation (125)]). *For any $\mathbf{v} \in P_{\mathbb{H}}(\Lambda)$ there exists a unique Spin^c structure $\mathfrak{s}_{\mathbf{v}}$ on $Y_{\mathbf{q}}$ which extends to a Spin^c structure $\mathfrak{t}_{\mathbf{v}}$ on $X_{\mathbf{q}}$ with $c_1(\mathfrak{t}_{\mathbf{v}}) = 2\mathbf{v} - (\Lambda_1 + \dots + \Lambda_m)$.*

Theorem 4.7 (Large Surgery Theorem, see [24, Section 10.1]). *Assume that $\mathbf{q} \gg \mathbf{0}$. For any $\mathbf{v} \in P_{\mathbb{H}}(\Lambda)$, the homology of $A^-(\mathcal{L}; \mathbf{v})$ is isomorphic to the Heegaard Floer homology of $S^3_{\mathbf{q}}(\mathcal{L})$ with Spin^c structure $\mathfrak{s}_{\mathbf{v}}$. More precisely, we have an isomorphism over $\mathbb{F}[U]$:*

$$(4.8) \quad A^-(\mathcal{L}, \mathbf{v}) \simeq CF^-(S^3_{\mathbf{q}}(\mathcal{L}), \mathfrak{s}_{\mathbf{v}}),$$

where U acts as U_1 on the left hand side. In particular, the actions of all the U_i on $A^-(\mathcal{L}, \mathbf{v})$ are homotopic to each other and to the action of U .

It is important to note that the isomorphism (4.8) shifts the grading. The grading shift can be calculated explicitly from the linking matrix Λ and the vector \mathbf{v} . We present a more geometric way, which will suit best our applications.

Remove a small ball from $X_{\mathbf{q}}$ and call the resulting manifold $U_{\mathbf{q}}$. This is a cobordism between S^3 and $Y_{\mathbf{q}}$. Let $U'_{\mathbf{q}} = -U_{\mathbf{q}}$. The Spin^c structure $\mathfrak{t}_{\mathbf{v}}$ on $U'_{\mathbf{q}}$ gives a Spin^c cobordism between $(Y_{\mathbf{q}}, \mathfrak{s}_{\mathbf{v}})$ and S^3 (equipped with the unique Spin^c structure), so it induces a map $F_{(U'_{\mathbf{q}}, \mathfrak{t}_{\mathbf{v}})}$ between Heegaard Floer homologies of $(Y_{\mathbf{q}}, \mathfrak{s}_{\mathbf{v}})$ and S^3 .

Proposition 4.9 (see [24, Section 10]). *The isomorphism (4.8) shifts the degree by $-\deg F_{(U'_{\mathbf{q}}, \mathfrak{t}_{\mathbf{v}})}$.*

As a corollary we can give a formula for d -invariants of large surgery on links.

Theorem 4.10. *For $\mathbf{v} \in P_{\mathbb{H}}(\Lambda)$, the d -invariant of a surgery on \mathcal{L} is given by*

$$d(S^3_{\mathbf{q}}(\mathcal{L}), \mathfrak{s}_{\mathbf{v}}) = -\deg F_{(U'_{\mathbf{q}}, \mathfrak{t}_{\mathbf{v}})} - 2H(\mathbf{v}).$$

5. PSICs AND FOUR-MANIFOLDS

A positively self-intersecting cobordism (later shortened to: a PSIC) is a generalization of the notion of a positively self-intersecting annulus used in [5] as a way to translate the questions about the unknotting number of knots into questions about cobordisms of three-manifolds, where d -invariants can be used. The notion of a PSIC will play the same role for links.

5.1. PSICs in various guises. Let \mathcal{L}_1 and \mathcal{L}_2 be two links. Denote by L_{11}, \dots, L_{1n_1} and L_{21}, \dots, L_{2n_2} the components of \mathcal{L}_1 and \mathcal{L}_2 , respectively. A *positively self-intersecting cobordism* (in short: PSIC) from \mathcal{L}_1 to \mathcal{L}_2 is a surface A , smoothly immersed into $S^3 \times [1, 2]$ such that $\partial A = \mathcal{L}_2 \sqcup -\mathcal{L}_1$ and A is topologically a union of immersed punctured disks. We require that A has ordinary positive double points as singularities. Here and afterwards, whenever we write \mathcal{L}_1 or \mathcal{L}_2 it should be understood that $\mathcal{L}_1 \subset S^3 \times \{1\}$ and $\mathcal{L}_2 \subset S^3 \times \{2\}$. The same applies for the components of \mathcal{L}_1 and \mathcal{L}_2 .

If A is a PSIC, we denote by A_1, \dots, A_n its components. Each of the A_i is an immersed surface and $A = A_1 \cup \dots \cup A_n$. We define $\eta_{ij} = \#(A_i \cap A_j)$ for $i, j = 1, \dots, n$ and $i \neq j$; for $i = j$ we set η_{ii} to be the number of double points of A_i . The total number of double points of A is

$$p = \sum_{i \leq j} \eta_{ij}.$$

Furthermore, set

$$(5.1) \quad \mathbf{a} = (a_1, \dots, a_n), \text{ where } a_i = \sum_{j \neq i} \eta_{ij}.$$

The following specifications of the definition of a PSIC will be used in the present article:

- an *annular* PSIC, shortly APSIC, is a PSIC such that each of the A_i is an annulus such that $\partial A_i = L_{2i} \sqcup -L_{1i}$. For an APSIC $n = n_2 = n_1$. An exemplary APSIC is presented in Figure 1. An APIC is a concordance (in the standard sense) with positive self-intersections allowed. The notion of APSIC will be most important in the applications.
- a *sprouting* PSIC, shortly SPSIC, is a PSIC such that for every $i = 1, \dots, n$, the intersection $A_i \cap S^3 \times \{1\} = L_{1i}$. For a SPSIC we have $n_2 \geq n_1 = n$. Furthermore, for any $i = 1, \dots, n$ we define the subset

$$(5.2) \quad \Theta_i = \{j = 1, \dots, n : L_{2j} \subset \partial A_i\}.$$

- an *elementary sprouting* PSIC, shortly ESPSIC, is a SPSIC such that A is smooth and there exists $k \in \{1, \dots, n\}$ such that for $i \neq k$ we have $\Theta_i = \{i\}$ and $\Theta_k = \{k, n_2\}$. For an EPSIC we have $n_2 - 1 = n_1 = n$.

Let us introduce some useful terminology.

Definition 5.3. A double point z of A is called *multicolored* respectively *monochromatic*, if z belongs to two components of A (respectively, to one component of A).

For future reference we will need two simple facts. For simplicity, suppose A is an APSIC (analogous statement can be proved for general SPSIC, but we do not need it). Define a shorthand

$$l_{ij}^k = \text{lk}(L_{ki}, L_{kj}) \text{ for } i \neq j, k = 1, 2.$$




FIGURE 1. An APSIC between two two-component links. Two crossings are marked: one is a double point on an annulus, another one is a crossing between two annuli.




FIGURE 2. Step 2 of the construction in Section 5.2. The boundary of Z is the disjoint union of S^3 on the left and $S^3_{q_2}(\mathcal{L}_2)$ on the right.

As all the self-intersections of A are positive and a positive self-intersection between different link components increases the linking number by 1, we have that for $i \neq j$:

$$(5.4) \quad l_{ij}^2 = l_{ij}^1 + \eta_{ij}.$$

Equation (5.4) summed up over $j \neq i$ yields the following result.

Lemma 5.5. *Suppose that A is an APSIC and let ℓ_1 and ℓ_2 be the linking vectors of \mathcal{L}_1 and \mathcal{L}_2 respectively. Then*

$$\ell_2 - \ell_1 = \frac{1}{2}\mathbf{a}.$$

5.2. Topological constructions involving a PSIC. In the following we generalize the construction of [5] that based on a version of a PSIC for knots and as an output produced a cobordism between surgeries of the two knots involved. We begin with a rather general construction, later on we will specify its three variants. The construction is done in four steps.

Step 1. Begin with $A \subset S^3 \times [1, 2]$ and blow up all the double points of A (we do not specify yet, whether we perform positive or negative blow-ups). The exceptional divisors are denoted by E_1, \dots, E_p . For a component A_j of A , let \tilde{A}_j be its strict transform, that is, the closure in the blow-up of the inverse image of $A_j \setminus \text{Sing } A$.




FIGURE 3. The final step of constructing the cobordisms W_{01} , W_{12} and W_{02} . The shaded part is W_{01} . The unfilled part is W_{12} .

Set

$$(5.6) \quad d_{ij} := A_i \cdot E_j \text{ and } \tilde{d}_{ij} := \frac{A_i \cdot E_j}{E_j \cdot E_j}.$$

Step 2. Fix a framing vector $\mathbf{q}_2 = (q_{21}, \dots, q_{2n_2}) \in \mathbb{Z}^{n_2}$, where n_2 is the number of components of the link \mathcal{L}_2 . This turns \mathcal{L}_2 into a framed link and let Λ_2 be its framing matrix. Attach to the p -fold blow-up of $S^3 \times [1, 2]$ constructed in Step 1 n_2 two-handles along \mathcal{L}_2 with framings given by \mathbf{q}_2 . The resulting manifold is called Z . Its boundary is $Y_2 \sqcup -S^3 \times \{1\}$, where $Y_2 = S^3_{\mathbf{q}_2}(\mathcal{L}_2)$ is a surgery on \mathcal{L}_2 ; see Figure 2. Let C_1, \dots, C_{n_2} be the cores of the handles attached. For each component L_{2i} of \mathcal{L}_2 choose a Seifert surface $\Sigma_{2i} \subset S^3 \times \{2\}$. Let $F_{2i} = \Sigma_{2i} \cup C_i$. Then F_{21}, \dots, F_{2n_2} are closed connected surfaces.

Step 3. Form the union $D = C_1 \cup \dots \cup C_{n_2} \cup \tilde{A}_1 \cup \dots \cup \tilde{A}_n$. Denote by D_1, \dots, D_n the connected components of D as in Figure 2. Let N be a tubular neighbourhood of D in Z ; see Figure 3. Set $W_{12} = Z \setminus N$ and let $Y_1 = \partial W \setminus Y_2$. Then W_{12} is a cobordism between Y_1 and Y_2 .

Step 4. Take again Z and glue to it a four-ball B along $S^3 \times \{1\} \subset \partial Z$. Let $X_{\mathbf{q}_2}$ be the resulting manifold. Finally pick a small ball $B_0 \subset B$ and drill it out from $X_{\mathbf{q}_2}$. Set $W_{02} = X_{\mathbf{q}_2} \setminus B_0$ and $W_{01} = (B \setminus B_0) \cup N$, so that $W_{01} \cup W_{12} = W_{02}$. Let $Y_0 = S^3$. Then W_{01} is a cobordism between Y_0 and Y_1 and W_{02} is a cobordism between Y_0 and Y_2 . See Figure 3.

We have the following immediate observation.

Lemma 5.7.

- (a) *The cobordism W_{02} is a p -fold blow-up of the cobordism $U_{\mathbf{q}_2}(\mathcal{L}_2)$ defined before Proposition 4.9.*
- (b) *Suppose A is a SPSIC. Then D_1, \dots, D_n are disks and $n = n_1$ is the number of components of \mathcal{L}_1 . Furthermore Y_1 is a surgery on \mathcal{L}_1 with a framing vector $\mathbf{q}_1 = (q_{11}, \dots, q_{1n_1})$ depending on \mathbf{q}_2 and on the signs of blow-ups (we give a precise formula for \mathbf{q}_1 below). The cobordism W_{01} is identified with $U_{\mathbf{q}_1}(\mathcal{L}_1)$.*

From now on we will assume that A is a SPSIC.

Choose Seifert surfaces for L_{11}, \dots, L_{1n_1} and call them $\Sigma_{11}, \dots, \Sigma_{1n_1}$. Let F_{11}, \dots, F_{1n_1} be closed surfaces obtained by capping the disks D_1, \dots, D_{n_1} with $\Sigma_{11}, \dots, \Sigma_{1n_1}$. The

classes $[F_{11}], \dots, [F_{1n_1}]$ generate $H_2(W_{01}; \mathbb{Z})$. The map $W_{01} \hookrightarrow W_{02}$ induces a monomorphism on second homologies. We will not distinguish between the class $[F_{1i}] \in H_2(W_{01}; \mathbb{Z})$ and its image in $H_2(W_{02}; \mathbb{Z})$.

Lemma 5.8. *In $H_2(W_{02}, \mathbb{Z})$ we have the following relation.*

$$[F_{1i}] = \sum_{j \in \Theta_i} [F_{2j}] + \sum_{k=1}^p \tilde{d}_{ik} [E_k].$$

Proof. The class of $[F_{1i}]$ is the sum of a class $\sum_{j \in \Theta_i} [F_{2j}]$ and the class of a closed immersed surface

$$S_i = \Sigma_{1i} \cup \tilde{A}_i \cup \sum_{j \in \Theta_i} \Sigma_{2j}.$$

The surfaces S_i will usually be only immersed, because the Seifert surfaces $\Sigma_{21}, \dots, \Sigma_{2n_2}$ can intersect (we may assume that their intersection is transverse, but this is irrelevant for the present proof).

Notice that S_i can be regarded as a strict transform of a closed surface in $S^3 \times [0, 1]$, formed by capping the component A_i with the Seifert surfaces of corresponding links. This surface in $S^3 \times [0, 1]$ is homologically trivial, as $H_2(S^3 \times [0, 1]; \mathbb{Z}) = 0$. Therefore, the class of S_i in $H_2(W_{02}; \mathbb{Z})$ is a linear combination of classes generated by the exceptional divisors. The coefficients in this linear combination can be calculated by intersecting S_i with divisors E_1, \dots, E_p . More concretely $[S_i] = \sum \frac{S_i \cdot E_k}{E_k \cdot E_k} [E_k]$. But geometrically $S_i \cdot E_k = A_i \cdot E_k = d_{ik}$. The lemma follows. \square

Lemma 5.9. *If A is a SPSIC and all the blow-ups are of fixed sign (either all positive or all negative), then \mathbf{q}_1 and \mathbf{q}_2 are related by the following formula.*

If all the blow-ups are negative

$$(5.10) \quad q_{1i} = \sum_{j \in \Theta_i} q_{2j} - 4\eta_{ii} - a_i.$$

If all the blow-ups are positive

$$(5.11) \quad q_{1i} = \sum_{j \in \Theta_i} q_{2j} + a_i.$$

Proof. The coefficients q_{11}, \dots, q_{1n_1} are self-intersections of disks D_1, \dots, D_{n_1} . Here, by the word 'self-intersection' we mean the following: push slightly D_i to obtain another disk, called D'_i , intersecting D_i transversally and such that $\partial D'_i \subset S^3 \times \{1\}$ is disjoint from ∂D_i and the linking number $\text{lk}(\partial D_i, \partial D'_i)$ calculated on $S^3 \times \{1\}$ is equal to zero. Then the self-intersection of D_i is defined as the number of intersection points (counted with signs) of D_i and D'_i . In other words, the self-intersection of D_i is equal to the self-intersection of $[F_{1i}]$. On the other hand, the framing q_{2j} is interpreted in the same way as the self-intersection of $[F_{2j}]$.

As the classes $[F_{2j}]$ and $[E_k]$ are orthogonal, by Lemma 5.8 the difference of self-intersections

$$\left(\sum_{j \in \Theta_i} [F_{2j}] \right)^2 - [F_{1i}]^2$$

is equal to

$$\left(\sum_{k=1}^p \tilde{d}_{ik}[E_k] \right)^2 = \sum_{k=1}^p \tilde{d}_{ik}^2 [E_k]^2.$$

Now we have two cases. First suppose that all the blow-ups are negative. Then $[E_k]^2 = -1$ for all k . Moreover, $\tilde{d}_{ik}^2 = 0, 1$ or 4 is the square of the multiplicity of the double point z_k if $z_k \in A_i$ and $\tilde{d}_{ik} = 0$ if $z_k \notin A_i$. The sum $\sum \tilde{d}_{ik}^2$ can be calculated using the fact that a_i is equal to the number of multicolored double points on A_i and η_{ii} is the number of monochromatic double points on A_i . This proves (5.10).

The situation with positive blow-ups is analogous. There is one difference, though. If an exceptional divisor E_k is a blow-up of a monochromatic point on A_i , then $d_{ik} = 0$ (and not ± 2). This corresponds to the fact that in the blow-up the annulus \widetilde{A}_i will intersect the exceptional divisor E_k in two points with opposite orientations. \square

5.3. Homological properties of W_{12} . We will study some homological properties of W_{02} and W_{12} . They are synthesized in the following lemma, which is a direct generalization of results in [5, Section 2.2].

Lemma 5.12. *Suppose A is sprouting and \mathbf{q}_2 has all coordinates sufficiently large.*

- *The homology of $H_2(W_{12}; \mathbb{Z})$ is the coimage of the map $H_2(Z; \mathbb{Z}) \rightarrow \mathbb{Z}^{n_1}$ which takes x to the vector $(x \cdot D_1, \dots, x \cdot D_{n_1})$.*
- *If all the blow-ups are positive, the manifold W_{12} has positive definite intersection form on H_2 . If all the blow-ups are negative definite and A is an APSIC, then W_{12} has negative definite intersection form.*

Proof. The manifold W_{12} is built from Z by removing tubular neighborhoods of disks. As Z arises by gluing n_2 two-handles to the p -fold blow-up of $S^3 \times [0, 1]$ and the framing matrix is nondegenerate, we have $H_1(Z; \mathbb{Q}) = 0$, $H_2(Z; \mathbb{Q}) \cong \mathbb{Q}^{p+n_2}$ and $H_3(Z; \mathbb{Q}) = 0$, furthermore $H_3(Z; \mathbb{Z}) = 0$.

Consider the long exact sequence of homology (with \mathbb{Z} coefficients) of the pair (Z, W_{12}) . By excision we have $H_*(Z, W_{12}) \cong H_*(N, \partial_+ N)$. Here N is viewed as a D^2 bundle over D and N_+ is the associated S^1 bundle. Using Thom isomorphism we obtain that $H_3(N, \partial_+ N) = 0$ and that $H_2(N, \partial_+ N) \cong \mathbb{Z}^{n_1}$ is generated by classes $\alpha_j := [(t_j \times D^2, t_j \times S^1)] \in H_2(N, \partial_+ N; \mathbb{Z})$, $j = 1, \dots, n_1$, where t_1, \dots, t_{n_1} are some points in D_1, \dots, D_{n_1} respectively.

The latter implies that $H_2(W_{12}; \mathbb{Z})$ injects into $H_2(Z; \mathbb{Z})$. The map

$$\kappa: H_2(Z; \mathbb{Z}) \rightarrow H_2(Z, W_{12}; \mathbb{Z})$$

can be explicitly described. Namely, for $x \in H_2(Z; \mathbb{Z})$ we choose its representative as a union of cycles each intersecting D transversally. Then

$$\kappa(x) = (x \cdot D_1)\alpha_1 + (x \cdot D_2)\alpha_2 + \dots + (x \cdot D_{n_1})\alpha_{n_1}.$$

This proves the first part of the lemma.

The first part can also be rephrased in another way. As each class $x \in H_2(Z; \mathbb{Z})$ can be represented by a surface disjoint from $S^3 \times \{1\}$, the geometric intersection number $(x \cdot D_j)$ is equal to $x \cdot [F_{1j}]$. With this description it follows that $H_2(W_{12}; \mathbb{Z})$ is an orthogonal (with respect to the intersection form) complement to a submodule of $H_2(Z; \mathbb{Z})$ generated by $[F_{11}], \dots, [F_{1n_1}]$. The same applies for homologies with \mathbb{Q} coefficients. Therefore, the signature of the intersection form on W_{12} can be calculated as the difference of the

signature of the intersection form on Z and the signature of the intersection form on $[F_{11}], \dots, [F_{1n_1}]$. The proof of the second part follows now by a case by case analysis.

If all the blow-ups are positive, then Z has a positive definite intersection form, hence it restricts to a positive definite intersection form on W_{12} . If A is an APSIC and all the blow-ups are negative, then one readily computes that $b_2^+(Z) = n$ and $b_2^-(Z) = p$. Moreover, the intersection form on an n -dimensional subspace spanned by $[F_{11}], \dots, [F_{1n}]$ is positive definite. So its orthogonal complement is negative definite. \square

6. INEQUALITIES FOR THE H -FUNCTION UNDER THE CROSSING CHANGE

We will now assume that links \mathcal{L}_1 and \mathcal{L}_2 are connected by a PSIC. The inequality for d -invariants (4.3) will translate into the inequality between H -functions, or, equivalently, J -functions.

We are going to prove the following two results (we use the notation from Section 5.1).

Theorem 6.1. *Let A be an APSIC from \mathcal{L}_1 to \mathcal{L}_2 . Let J_1 and J_2 be the J -functions as in Definition 3.26. Set $\mathbf{r} = (\eta_{11}, \dots, \eta_{nn})$. Choose a presentation of η_{il} for $i < l$ as the sum*

$$\eta_{il} = \eta_{il}^1 + \eta_{il}^2,$$

where η_{il}^1, η_{il}^2 are non-negative integers. Set

$$k_i = \sum_{j < i} \eta_{ji}^2 + \sum_{j > i} \eta_{ij}^1.$$

Let $\mathbf{k} = (k_1, \dots, k_n)$. Then for any $\mathbf{m} \in \mathbb{Z}^n$ we have

$$(6.2) \quad J_1(\mathbf{m} + \mathbf{k}) \leq J_2(\mathbf{m}) \leq J_1(\mathbf{m} - \mathbf{r})$$

A counterpart of this result for an EPSIC is the following.

Theorem 6.3. *Suppose A is an EPSIC. Choose $\mathbf{m}_2 \in \mathbb{Z}^{n_2}$ and let $\mathbf{m}_1 \in \mathbb{Z}^{n_1}$ be given by $\mathbf{m}_{1i} = \mathbf{m}_{2i}$ if $i \neq k$ and $\mathbf{m}_{1k} = \mathbf{m}_{2k} + \mathbf{m}_{2,n_2}$. Then*

$$(6.4) \quad J_1(\mathbf{m}_1) \leq J_2(\mathbf{m}_2).$$

Theorem 6.1 is proved in Sections 6.1 and 6.2. Theorem 6.3 is proved in Section 6.3. In Section 6.4 we prove Theorem 6.20, which is a straightforward, but important, corollary of Theorem 6.1.

6.1. Proof of Theorem 6.1. Part 1. In this section we prove the part $J_2(\mathbf{m}) \leq J_1(\mathbf{m} - \mathbf{r})$.

Construct W_{02} by making negative blow-ups of the APSIC; see Section 5.2. Choose $\mathbf{m} \in \mathbb{Z}^n$. Pick \mathbf{q}_2 sufficiently large (we specify below the precise meaning of sufficiently large), but now we point out that \mathbf{q}_2 is chosen *after* \mathbf{m} . According to Lemma 5.9 define $\mathbf{q}_1 = \mathbf{q}_2 - 4\mathbf{r} - \mathbf{a}$.

Set $\mathbf{v}_2 = \mathbf{m} - \boldsymbol{\ell}_2$, where $\boldsymbol{\ell}_2$ is the linking vector for \mathcal{L}_2 . Let $\mathfrak{s}_{\mathbf{v}_2}$ be the Spin^c structure on Y_2 . It extends to a Spin^c structure $\mathbf{t}_{\mathbf{v}_2}$ on $U_{\mathbf{q}_2}(\mathcal{L}_2)$ (see Section 4.2 for definition of $U_{\mathbf{q}}(\mathcal{L})$). Recall that W_{02} is identified with a p -fold blow-up of $U_{\mathbf{q}_2}(\mathcal{L}_2)$. Let $\pi: W_{02} \rightarrow U_{\mathbf{q}_2}(\mathcal{L}_2)$ be the blow-down map. Define the Spin^c structure $\mathbf{t}'_{\mathbf{v}_2}$ on W_{02} as the Spin^c structure, whose first Chern class is equal to

$$\pi^* c_1(\mathbf{t}_{\mathbf{v}_2}) + PD[E_1] + \dots + PD[E_n].$$

Lemma 6.5. *The $Spin^c$ structure $\mathbf{t}'_{\mathbf{v}_2}$ on W_{02} restricts to the $Spin^c$ structure $\mathbf{s}_{\mathbf{v}_1}$ on Y_1 , where*

$$(6.6) \quad \mathbf{v}_1 = \mathbf{v}_2 - \mathbf{r} - \frac{1}{2}\mathbf{a}.$$

Proof of Lemma 6.5. By construction of W_{01} and by Lemma 5.9 we have $W_{01} = U_{\mathbf{q}_1}(\mathcal{L}_1)$. The $Spin^c$ structure $\mathbf{s}_{\mathbf{v}_1}$ on Y_1 extends to the $Spin^c$ structure $\mathbf{t}_{\mathbf{v}_1}$ on W_{01} . Our aim is to show that with the choice of \mathbf{v}_1 as in the statement of the lemma, $c_1(\mathbf{t}_{\mathbf{v}_1})$ and $c_1(\mathbf{t}'_{\mathbf{v}_2})$ evaluate in the same way on the classes $[F_{11}], \dots, [F_{1n}]$.

By definition of $\mathbf{t}'_{\mathbf{v}_2}$ we have

$$(6.7) \quad \begin{aligned} \langle c_1(\mathbf{t}'_{\mathbf{v}_2}), [F_{2i}] \rangle &= 2v_{2,i} - (\Lambda_{21} + \dots + \Lambda_{2n_2})_i \\ \langle c_1(\mathbf{t}'_{\mathbf{v}_2}), [E_j] \rangle &= E_j \cdot E_j = -1, \end{aligned}$$

where $\Lambda_{21}, \dots, \Lambda_{2n_2}$ are column vectors of the framing matrix Λ_2 for \mathcal{L}_2 . The subscript i in the first formula means that we take the i -th coordinate of the vector in the parentheses.

Combining (6.7) with Lemma 5.8 we obtain

$$(6.8) \quad \langle c_1(\mathbf{t}'_{\mathbf{v}_2}), [F_{1i}] \rangle = 2v_{2,i} - (\Lambda_{21} + \dots + \Lambda_{2n})_i + \sum_{j=1}^p d_{ij}.$$

The framing matrices Λ_2 and Λ_1 can be compared using (5.4) and (5.10).

$$(\Lambda_{21} + \dots + \Lambda_{2n})_i - (\Lambda_{11} + \dots + \Lambda_{1n})_i \stackrel{(5.4)}{=} \sum_{j \neq i} \eta_{ij} + q_{2i} - q_{1i} \stackrel{(5.10)}{=} 4\eta_{ii} + 2a_i.$$

Notice that $d_{ij} = 1$ for all multicolored double points that lie on A_i and $d_{ij} = 2$ for all monochromatic double points on A_i . Therefore (6.8) implies that

$$\langle c_1(\mathbf{t}'_{\mathbf{v}_2}), [F_{1i}] \rangle = 2v_{2,i} - (\Lambda_{21} + \dots + \Lambda_{2n})_i + a_i + 2\eta_{ii}.$$

The two above equations yield

$$\langle c_1(\mathbf{t}'_{\mathbf{v}_2}), [F_{1i}] \rangle = 2v_{2,i} - (\Lambda_{11} + \dots + \Lambda_{1n})_i - a_i - 2\eta_{ii}.$$

On the other hand, by Proposition 4.6

$$(6.9) \quad \langle c_1(\mathbf{t}_{\mathbf{v}_1}), [F_{1i}] \rangle = 2v_{1,i} - (\Lambda_{11} + \dots + \Lambda_{1n})_i.$$

Combining the two above formulae we conclude that $c_1(\mathbf{t}_{\mathbf{v}_1})$ and $c_1(\mathbf{t}'_{\mathbf{v}_2})$ evaluate to the same number on each of the $[F_{1i}]$. It follows that $c_1(\mathbf{t}_{\mathbf{v}_1}) = c_1(\mathbf{t}'_{\mathbf{v}_2})$ in $H^2(W_{01}; \mathbb{Z})$. By Proposition 4.6 (see also [24, Section 10.1]) this implies that $\mathbf{t}_{\mathbf{v}_1} = \mathbf{t}'_{\mathbf{v}_2}$. \square

We resume the proof of the inequality $J_2(\mathbf{m}) \leq J_1(\mathbf{m} - \mathbf{r})$. If \mathbf{q}_2 is large, then the statement of Theorem 4.10 holds for \mathbf{q}_2 -surgery on \mathcal{L}_2 and for \mathbf{q}_1 -surgery on \mathcal{L}_1 . Furthermore, we require that \mathbf{q}_2 and \mathbf{q}_1 are large enough so that $\mathbf{v}_2 \in P_{\mathbb{H}}(\Lambda_2)$ and $\mathbf{v}_1 \in P_{\mathbb{H}}(\Lambda_1)$.

Denote for simplicity $\mathbf{t} = \mathbf{t}'_{\mathbf{v}_2}$. By Lemma 6.5, (W_{12}, \mathbf{t}) is a $Spin^c$ cobordism between $(Y_1, \mathbf{s}_{\mathbf{v}_1})$ and $(Y_2, \mathbf{s}_{\mathbf{v}_2})$.

By Lemma 5.12 W_{12} is negative definite. By (4.2) we have:

$$(6.10) \quad d(Y_2, \mathbf{s}_{\mathbf{v}_2}) \geq \deg F_{(W_{12}, \mathbf{t})} + d(Y_1, \mathbf{s}_{\mathbf{v}_1}).$$

By Proposition 4.5 combined with Theorem 4.10:

$$(6.11) \quad \begin{aligned} d(Y_2, \mathbf{s}_{\mathbf{v}_2}) &= -\deg F(-W_{02}, \mathbf{t}) - 2J_2(\mathbf{v}_2 - \ell_2) - p \\ d(Y_1, \mathbf{s}_{\mathbf{v}_1}) &= -\deg F(-W_{01}, \mathbf{t}) - 2J_1(\mathbf{v}_1 - \ell_1). \end{aligned}$$

Notice that the first equation contains the term $-p$. This follows from the fact that W_{02} is not $U_{\mathbf{q}_2}(\mathcal{L}_2)$, but it is a negative blow-up of $U_{\mathbf{q}_2}(\mathcal{L}_2)$ with p blow-ups. If we reverse the orientation, the negative blow-up becomes a positive blow-up, so $\deg F_{(-W_{02}, t)} = \deg F_{(U_{\mathbf{q}_2}(\mathcal{L}_2), t)} - p$ by Proposition 4.5.

Substituting (6.11) into (6.10) we obtain:

$$(6.12) \quad -\deg F_{(-W_{02}, t)} + \deg F_{(-W_{01}, t)} - \deg F_{(W_{12}, t)} - p + J_1(\mathbf{v}_1 - \boldsymbol{\ell}_1) \geq J_2(\mathbf{v}_2 - \boldsymbol{\ell}_2).$$

Let us look at the expression

$$\Delta := \deg F_{(-W_{02}, t)} + \deg F_{(W_{12}, t)} - \deg F_{(-W_{01}, t)}.$$

Denote by c_{02}, c_{12} and c_{01} the evaluations of $c_1^2(t)$ on W_{02} , W_{12} and W_{01} respectively. Likewise denote by σ_{02}, σ_{12} and σ_{01} the corresponding signatures and $\chi_{02}, \chi_{12}, \chi_{01}$ the Euler characteristic. We have by (4.2):

$$\begin{aligned} 4\deg F_{(-W_{02}, t)} &= -c_{02} + 3\sigma_{02} - 2\chi_{02} \\ 4\deg F_{(W_{12}, t)} &= c_{12} - 3\sigma_{12} - 2\chi_{12} \\ -4\deg F_{(-W_{01}, t)} &= c_{01} - 3\sigma_{01} + 2\chi_{01}. \end{aligned}$$

Notice that in the above expression we switched signs of σ and c according to the orientation. Notice also that $\sigma_{02} = \sigma_{01} + \sigma_{12}$ and $\chi_{02} = \chi_{01} + \chi_{12}$ (additivity of the signature and of the Euler characteristic) and $c_{02} = c_{01} + c_{12}$ (functoriality of the Chern class). Summing up the three equations we obtain

$$4\Delta = 4\chi_{12} = -4\chi(W_{12}).$$

The Euler characteristic of W_{12} can be quickly calculated. Recall that in Section 5.2 the manifold W_{12} was constructed by taking $S^3 \times [0, 1]$, blowing up p times, gluing n two-handles and drilling out n disks. The original $S^3 \times [0, 1]$ has Euler characteristic 0. Each blow-up increases it by 1. A two-handle attachment increases it by 1 and drilling out a disk decreases it by 1. Finally $\chi(W_{12}) = p$ so $\Delta = -p$. Plugging the value of Δ into (6.12) we obtain.

$$J_1(\mathbf{v}_1 - \boldsymbol{\ell}_1) \geq J_2(\mathbf{v}_2 - \boldsymbol{\ell}_2).$$

By definition, $\mathbf{v}_2 - \boldsymbol{\ell}_2 = \mathbf{m}$. The last step is to calculate $\mathbf{v}_1 - \boldsymbol{\ell}_1$. We use Lemma 6.5. By Lemma 5.5, equation (6.6) can be rewritten as

$$\mathbf{v}_1 = \mathbf{v}_2 - \mathbf{k} - (\boldsymbol{\ell}_2 - \boldsymbol{\ell}_1).$$

This amounts to saying that $\mathbf{v}_1 - \boldsymbol{\ell}_1 = \mathbf{m} - \mathbf{r}$, so $J_1(\mathbf{m} - \mathbf{r}) \geq J_2(\mathbf{m})$.

6.2. Proof of Theorem 6.1. Part 2.

We are going to prove the part $J_1(\mathbf{m} + \mathbf{k}) \leq J_2(\mathbf{m})$.

The proof is analogous to the proof of $J_2(\mathbf{m}) \leq J_1(\mathbf{m} - \mathbf{r})$, although there are some differences. We construct W_{02} by making all blow-ups positive. Choose $\mathbf{m} \in \mathbb{Z}^n$ and let \mathbf{q}_2 be sufficiently large.

We begin with some combinatorics. Recall that the exceptional divisors of the blow-up are denoted E_1, \dots, E_p . We choose orientation of the divisors by requiring that if E_j is the exceptional divisor of the blow-up of the point of intersection $A_i \cap A_{i'}$ with $i < i'$, then $E_j \cap A_i = 1$ and $E_j \cap A_{i'} = -1$. The orientation of the exceptional divisors of blow-ups of monochromatic double points is irrelevant.

Choose now $\delta_1, \dots, \delta_p \in \{-1, +1\}$ in the following way. If E_j is the exceptional divisor of the blow-up of a monochromatic double point, then $\delta_j = 1$. Let now i and i' be the indices such that $i < i'$. Let $I_{ii'}$ be the set of indices $\{1, \dots, p\}$ such that $j \in I_{ii'}$ if and only if E_j is the exceptional divisor of the blow-up of a point in $A_i \cap A_{i'}$. We know that

$\#I_{ii'} = \eta_{ii'}$. Partition the set $I_{ii'}$ into two subsets $I_{ii'}^1$ and $I_{ii'}^2$ of cardinality $m_{ii'}^1$ and $m_{ii'}^2$ respectively. Set $\delta_j = -1$ for $j \in I_{ii'}^1$ and $\delta_j = 1$ for $j \in I_{ii'}^2$. Finally denote

$$\boldsymbol{\theta} = (\theta_1, \dots, \theta_n) = \left(\sum_{l=1}^p \delta_l d_{1l}, \dots, \sum_{l=1}^p \delta_l d_{nl} \right).$$

We have the following result

Lemma 6.13. *With the choice of $\delta_1, \dots, \delta_p$ as above and with \mathbf{k} as in the statement of Theorem 6.1 we have*

$$\boldsymbol{\ell}_2 - \boldsymbol{\ell}_1 - \mathbf{k} = \frac{1}{2} \boldsymbol{\theta}.$$

Proof. In view of Lemma 5.5 we need to prove that

$$\boldsymbol{\theta} = 2\mathbf{k} - \mathbf{a}.$$

By definition, $k_i = \sum_{j < i} \eta_{ji}^2 + \sum_{j > i} \eta_{ji}^1$. Using the definition of a_i in (5.1) and the fact that for $i < j$ $\eta_{ji} = \eta_{ij} = \eta_{ij}^1 + \eta_{ij}^2$ we transform the above equation into the following set of equations for $i = 1, \dots, n$:

$$(6.14) \quad \sum_{l=1}^p \delta_l d_{il} = \sum_{j < i} (\eta_{ji}^2 - \eta_{ji}^1) + \sum_{i' > i} (\eta_{ji}^1 - \eta_{ji}^2).$$

The way the exceptional divisors are oriented implies that $d_{il} = 1$ if $l \in I_{ii'}$ for some $i' > i$, $d_{il} = -1$ if $l \in I_{i'i}$ for some $i' < i$, and otherwise $d_{il} = 0$. The left hand side of (6.14) can be expressed as

$$\sum_{l=1}^p \delta_l d_{il} = \sum_{i' < i} \sum_{l \in I_{i'i}} \delta_l - \sum_{i' > i} \sum_{l \in I_{ii'}} \delta_l.$$

But $\sum_{l \in I_{ii'} \cup I_{i'i}} \delta_l = \eta_{i'i}^2 - \eta_{i'i}^1$ by definition, so

$$\sum_{l=1}^p \delta_l d_{il} = \sum_{j < i} (\eta_{ji}^2 - \eta_{ji}^1) + \sum_{i' > i} (\eta_{ji}^1 - \eta_{ji}^2).$$

This proves (6.14) and concludes the proof of the lemma. \square

We resume the proof of Theorem 6.1. The manifold W_{02} is a p -fold positive blow-up of $U_{\mathbf{q}_2}(\mathcal{L}_2)$, and let again π be the blow-down map. Choose $\mathbf{v}_2 = \mathbf{m} + \boldsymbol{\ell}_2$ and the Spin^c structure $\mathbf{t}'_{\mathbf{v}_2}$ on W_{02} given by

$$c_1(\mathbf{t}'_{\mathbf{v}_2}) = \pi^* c_1(\mathbf{t}_{\mathbf{v}_2}) + \delta_1 PD[E_1] + \dots + \delta_n PD[E_n].$$

We have the following result, which is a counterpart of Lemma 6.5.

Lemma 6.15. *The Spin^c structure $\mathbf{t}'_{\mathbf{v}_2}$ restricts to the Spin^c structure \mathbf{v}_1 on Y_1 , where*

$$\mathbf{v}_1 = \mathbf{v}_2 - \frac{1}{2} \boldsymbol{\theta}.$$

Proof of Lemma 6.15. The proof of Lemma 6.15 resembles the proof of Lemma 6.5. We will show that first Chern classes of the two Spin^c structures, $\mathbf{t}'_{\mathbf{v}_2}$ and $\mathbf{t}_{\mathbf{v}_1}$ evaluate in the same way on the generators of $H_2(W_{01}; \mathbb{Z})$ and conclude by Proposition 4.6.

By definition of $\mathbf{t}'_{\mathbf{v}_2}$ we obtain

$$(6.16) \quad \begin{aligned} \langle c_1(\mathbf{t}'_{\mathbf{v}_2}), [F_{2i}] \rangle &= 2v_{2,i} - (\Lambda_{21} + \dots + \Lambda_{2n_2})_i \\ \langle c_1(\mathbf{t}'_{\mathbf{v}_2}), [E_j] \rangle &= \delta_j E_j \cdot E_j = \delta_j. \end{aligned}$$

Combining (6.16) with Lemma 5.8 we obtain

$$(6.17) \quad \langle c_1(\mathbf{t}'_{\mathbf{v}_2}), [F_{1i}] \rangle = 2v_{2,i} - (\Lambda_{21} + \dots + \Lambda_{2n})_i + \sum_{j=1}^p \delta_j d_{ij}.$$

Notice that by Lemma 5.9 Y_1 is a \mathbf{q}_1 -surgery on \mathcal{L}_1 , where $\mathbf{q}_1 = \mathbf{q}_2 + \mathbf{a}$. Therefore a quick calculation using (5.4) yields

$$\begin{aligned} (\Lambda_{21} + \dots + \Lambda_{2n})_i - (\Lambda_{11} + \dots + \Lambda_{1n})_i &= \\ &(\Lambda_{21} + \dots + \Lambda_{2,i-1} + \Lambda_{2,i+1} + \dots + \Lambda_{2n})_i \\ &- (\Lambda_{21} + \dots + \Lambda_{2,i-1} + \Lambda_{2,i+1} + \dots + \Lambda_{2n})_i \\ &+ q_{2,i} - q_{1,i} = a_i - a_i = 0. \end{aligned}$$

Substituting this into (6.17) we obtain.

$$\langle c_1(\mathbf{t}'_{\mathbf{v}_2}), [F_{1i}] \rangle = 2v_{2,i} - (\Lambda_{11} + \dots + \Lambda_{1n})_i + \sum_{j=1}^p \delta_j d_{ij}.$$

The evaluation of $c_1(\mathbf{t}_{\mathbf{v}_1})$ on $[F_{1i}]$ is given by (6.9). We obtain that

$$\langle c_1(\mathbf{t}'_{\mathbf{v}_2}), [F_{1i}] \rangle = \langle c_1(\mathbf{t}_{\mathbf{v}_1}), [F_{1i}] \rangle, \text{ if } 2\mathbf{v}_2 - \boldsymbol{\theta} = 2\mathbf{v}_1.$$

□

We continue the proof of Theorem 6.1. With the choice of \mathbf{v}_1 as in Lemma 6.15, the manifold $(W_{12}, \mathbf{t}'_{\mathbf{v}_2})$ is a Spin^c cobordism between $(Y_1, \mathbf{s}_{\mathbf{v}_1})$ and $(Y_2, \mathbf{s}_{\mathbf{v}_2})$. By Lemma 5.12 W_{12} is positive definite. Then $-W_{12}$ is negative definite and (4.3) gives

$$d(Y_1, \mathbf{s}_{\mathbf{v}_1}) \geq \deg F_{(-W_{12}, \mathbf{t})} + d(Y_2, \mathbf{s}_{\mathbf{v}_2}).$$

Using again the formula for d -invariants of large surgeries we obtain

$$(6.18) \quad -\deg F_{(-W_{01}, \mathbf{t})} + \deg F_{(-W_{02}, \mathbf{t})} - \deg F_{(-W_{12}, \mathbf{t})} + J_2(\mathbf{v}_2 - \boldsymbol{\ell}_2) \geq J_1(\mathbf{v}_1 - \boldsymbol{\ell}_1).$$

Now the expression $\deg F_{(-W_{01}, \mathbf{t})} + \deg F_{(-W_{12}, \mathbf{t})} - \deg F_{(-W_{02}, \mathbf{t})}$ is much easier to handle than an analogous expression in Section 6.1 because $-W_{02} = -W_{01} \cup -W_{12}$. Therefore the map $F_{(-W_{02}, \mathbf{t})}$ is the composition of $F_{(-W_{01}, \mathbf{t})}$ and $F_{(-W_{12}, \mathbf{t})}$ so its degree is the sum of the degrees of the summands. The three degrees in (6.18) cancel out and we are left with

$$(6.19) \quad J_2(\mathbf{v}_2 - \boldsymbol{\ell}_2) \geq J_1(\mathbf{v}_1 - \boldsymbol{\ell}_1).$$

By definition $\mathbf{v}_2 - \boldsymbol{\ell}_2 = \mathbf{m}$. On the other hand, by Lemma 6.13 combined with Lemma 6.15:

$$\boldsymbol{\ell}_2 - \boldsymbol{\ell}_1 - \mathbf{k} = \frac{1}{2}\boldsymbol{\theta} = \mathbf{v}_2 - \mathbf{v}_1.$$

Plugging this into (6.19) yields $J_2(\mathbf{m}) \geq J_1(\mathbf{m} + \mathbf{k})$. This accomplishes the proof of Theorem 6.1.

6.3. Proof of Theorem 6.3. The construction is similar as in Section 6.2. Take $\mathbf{m}_2 \in \mathbb{Z}^{n_2}$ and let \mathbf{q}_2 be sufficiently large. The construction of W_{02} is as in the proof of Theorem 6.1, but there are no blow-ups, hence $W_{02} = U_{\mathbf{q}_2}(\mathcal{L}_2)$. We know that $W_{01} = U_{\mathbf{q}_1}(\mathcal{L}_1)$, where by Lemma 5.9 $q_{1i} = q_{2i}$ if $i \neq k$ and $q_{1k} = q_{2k} + q_{2n_2}$. Set $\mathbf{v}_2 = \mathbf{m}_2 - \boldsymbol{\ell}_2$ and let $\mathbf{t}_{\mathbf{v}_2}$ be the Spin^c structure on W_{02} extending the Spin^c structure $\mathbf{s}_{\mathbf{v}_2}$ on Y_2 . Evaluating $c_1(\mathbf{t}_{\mathbf{v}_2})$ on classes $[F_{11}], \dots, [F_{1n_1}]$ we show that $\mathbf{t}_{\mathbf{v}_2}$ restricts to $\mathbf{s}_{\mathbf{v}_1}$ on Y_1 , where $v_{1i} = v_{2i}$ if $i \neq k$ and $v_{1k} = v_{2k} + v_{2n_2}$. By Lemma 5.12, W_{12} is positive definite. Therefore $(-W_{12}, \mathbf{t}_{\mathbf{v}_2})$ is a negative definite Spin^c cobordism between $(Y_2, \mathbf{s}_{\mathbf{v}_2})$ and $(Y_1, \mathbf{s}_{\mathbf{v}_1})$. Acting exactly in the same way as in Section 6.2 we arrive at the inequality $J_2(\mathbf{v}_2 - \boldsymbol{\ell}_2) \geq J_1(\mathbf{v}_1 - \boldsymbol{\ell}_1)$. We have

$\mathbf{v}_2 - \boldsymbol{\ell}_2 = \mathbf{m}_2$. Moreover it is easy to see that with the definition of \mathbf{v}_1 and \mathbf{m}_1 , we have $\mathbf{v}_1 - \boldsymbol{\ell}_1 = \mathbf{m}_1$. This concludes the proof.

6.4. A variant of Theorem 6.1 for a single crossing change.

Theorem 6.20. *Let \mathcal{L}_1 and \mathcal{L}_2 be two n -component links differing by a single positive crossing change, that is, \mathcal{L}_2 arises by changing a negative crossing of \mathcal{L}_1 into a positive one. Let J_1 and J_2 be the corresponding J -functions and let $\mathbf{m} \in \mathbb{Z}^n$, $\mathbf{m} = (m_1, \dots, m_n)$.*

(a) *If the crossing change is between two strands of the same component L_{1i} of \mathcal{L}_1 , then*

$$J_2(m_1, m_2, \dots, m_i + 1, \dots, m_n) \leq J_1(m_1, \dots, m_n) \leq J_2(m_1, \dots, m_i, \dots, m_n).$$

(b) *If the crossing change is between the i -th and j -th component of \mathcal{L}_1 , then*

$$J_2(m_1, m_2, \dots, m_n) \leq J_1(m_1, \dots, m_n) \leq J_2(m_1, \dots, m_i - 1, \dots, m_n)$$

and

$$J_2(m_1, m_2, \dots, m_n) \leq J_1(m_1, \dots, m_n) \leq J_2(m_1, \dots, m_j - 1, \dots, m_n)$$

Proof. We begin with part (a). If \mathcal{L}_1 and \mathcal{L}_2 differ by a single positive crossing change involving the component L_{1i} , then there is an APSIC from \mathcal{L}_1 to \mathcal{L}_2 . The construction is a generalization of [5, Example 2.2]. We take a product cobordism between components L_{1j} and L_{2j} for $j \neq i$ and an annulus with a single positive double point connecting L_{1i} to L_{2i} . The cobordism has $\eta_{kl} = 0$ unless $k = l = i$ and $\eta_{ii} = 1$. In the notation of Theorem 6.1 we have $\mathbf{r} = \mathbf{e}_i$ and $\mathbf{k} = (0, \dots, 0)$. Part (a) of Theorem 6.20 follows immediately.

Part (b) is analogous. We construct an APSIC with $\eta_{kl} = 0$ with the exception that $\eta_{ij} = \eta_{ji} = 1$. We have $\mathbf{r} = (0, \dots, 0)$ and the splitting $1 = \eta_{ij} = \eta_{ij}^1 + \eta_{ij}^2$ can be done in two ways: $(\eta_{ij}^1, \eta_{ij}^2) = (0, 1)$ or $(1, 0)$. This gives two possibilities for choosing \mathbf{k} , namely $\mathbf{k} = \mathbf{e}_i$ or $\mathbf{k} = \mathbf{e}_j$. Applying Theorem 6.1 concludes the proof. \square

7. SPLITTING NUMBERS OF LINKS

Let us recall the following definition.

Definition 7.1. Let \mathcal{L} be a link with n components. The *splitting number* $sp(\mathcal{L})$ is the minimal number of *multicolored* crossing changes (that is, between different components) needed to turn \mathcal{L} into a split link.

Example 7.2. The splitting number of the Whitehead link is 2, even though the unlinking number is 1.

We will use the following terminology:

Definition 7.3. A *positive crossing change* is a change of a negative crossing of a link into a positive crossing. Likewise, a *negative crossing change* is a change of a positive crossing into a negative crossing.

7.1. Splitting number bound from the \tilde{J} -function. In Definition 3.28 we defined the \tilde{J} -function of a link. The following result gives a ready-to-use bound for the splitting number.

Theorem 7.4. *Suppose that \mathcal{L} can be turned into an unlink using t_+ positive and t_- negative multicolored crossing changes. Then $-t_- \leq \tilde{J}(\mathbf{m}) \leq t_+$ for all $\mathbf{m} \in \mathbb{Z}^n$.*

Proof. Use Theorem 6.20 together with Proposition 3.10 (the latter holds for the J -function as well, because J differs from H by an overall argument shift). We obtain that if two links \mathcal{L}_1 and \mathcal{L}_2 differ by a single positive multicolored crossing change, then for all $\mathbf{m} \in \mathbb{Z}^n$

$$J_2(\mathbf{m}) \leq J_1(\mathbf{m}) \leq J_2(\mathbf{m}) + 1.$$

Notice that a multicolored crossing change of a link \mathcal{L} does not affect the isotopy type of its components, in particular the J -functions of its components remain the same. This shows that if \mathcal{L}_1 and \mathcal{L}_2 differ by a single positive multicolored crossing change, then for all $\mathbf{m} \in \mathbb{Z}^n$:

$$\tilde{J}_2(\mathbf{m}) \leq \tilde{J}_1(\mathbf{m}) \leq \tilde{J}_2(\mathbf{m}) + 1.$$

Using this result repeatedly we show that if \mathcal{L}_1 and \mathcal{L}_2 differ by t_+ positive multicolored crossing changes and t_- negative multicolored crossing changes, then

$$\tilde{J}_2(\mathbf{m}) - t_- \leq \tilde{J}_1(\mathbf{m}) \leq \tilde{J}_2(\mathbf{m}) + t_+.$$

Suppose now that \tilde{J}_2 is a split link. Then by Corollary 3.29 we know that $\tilde{J}_2 = 0$. In particular

$$-t_- \leq \tilde{J}_1(\mathbf{m}) \leq t_+.$$

□

Theorem 7.4 is very useful for quick estimates of the splitting number of L-space links with two components, because then the \tilde{J} -function can be quickly determined from the Alexander polynomial.

Example 7.5. We continue the example of Whitehead link, see Example 3.33. As the Alexander polynomial is $\Delta = -(t_1 - 1)(t_2 - 1)t_1^{-1/2}t_2^{-1/2}$, by Corollary 3.32 we have $\tilde{J}(1, 1) = 1$. This shows that the splitting number of the Whitehead link is at least 1.

7.2. Two-component links. Theorem 6.20 can be used directly to obtain some bounds for splitting numbers for two-component links.

Theorem 7.6. *Let \mathcal{L} be an arbitrary link with two components, and a link \mathcal{L}' can be obtained from \mathcal{L} by changing a negative multicolored crossings to positive, and b positive multicolored crossings to negative. Write $a = a_1 + a_2$ and $b = b_1 + b_2$ for arbitrary nonnegative a_i, b_i , then the following inequalities hold for all m_1, m_2 :*

$$J'(m_1 + b_1, m_2 + b_2) \leq J(m_1, m_2) \leq J'(m_1 - a_1, m_2 - a_2).$$

Proof. It is sufficient to consider a single crossing change. If $a_1 = 1, a_2 = b_1 = b_2 = 0$, then by Theorem 6.20 one has

$$J'(m_1, m_2) \leq J(m_1, m_2) \leq J'(m_1 - 1, m_2).$$

If $b_1 = 1, a_1 = a_2 = b_2 = 0$, then by Theorem 6.20 one has

$$J(m_1, m_2) \leq J'(m_1, m_2) \leq J(m_1 - 1, m_2),$$


so

$$J'(m_1 + 1, m_2) \leq J(m_1, m_2) \leq J'(m_1, m_2).$$

□

The following corollary will be useful below:

Theorem 7.7.

FIGURE 4. Region $R(b_+) + \mathbf{g}$ where J and \tilde{J} vanish.

(a) Suppose that a two component link $\mathcal{L} = L_1 \cup L_2$ can be unlinked using b_+ positive and b_- negative crossing changes. Let g_i denote the slice genus of L_i . Define vectors

$$\mathbf{g} := (g_1, g_2), \quad \tilde{\mathbf{g}} := \left(g_1 + \frac{1}{2} \text{lk}(L_1, L_2), g_2 + \frac{1}{2} \text{lk}(L_1, L_2) \right).$$

Define the region $R(b_+)$ by inequalities:

$$R(b_+) := \{(m_1, m_2) : m_1 + m_2 \geq a, m_1 \geq 0, m_2 \geq 0\};$$

see Figure 4. Then $J(\mathbf{m}) = \tilde{J}(\mathbf{m}) = 0$ for $\mathbf{m} \in R(b_+) + \mathbf{g}$.

(b) If, in addition, \mathcal{L} is an L-space link, then

$$\text{HFL}^-(\mathcal{L}, \mathbf{v}) = 0 \text{ for } \mathbf{v} \in R(b_+) + \tilde{\mathbf{g}} + (1, 1).$$

In particular, all coefficients of the Alexander polynomial vanish in $R(a) + \tilde{\mathbf{g}} + (\frac{1}{2}, \frac{1}{2})$.

Proof. As above, let J_i denote the J -functions for the components of \mathcal{L} . For a split link $\mathcal{L}' = L_1 \sqcup L_2$ we get $J'(v_1, v_2) = J_1(v_1) + J_2(v_2)$. Furthermore, by [31, Corollary 7.4] we get $J_i(v_i) = 0$ for $v_i \geq g_i$.

Assume that $\mathbf{m} = (m_1, m_2)$ belongs to $R(a) + \mathbf{g}$. By definition, $J(m_1, m_2) \geq 0$. On the other hand, let us choose $b_1 = \min(m_1 - g_1, b_+)$ and $b_2 = b_+ - b_1$, then $m_1 - b_1 \geq g_1$ and

$$m_2 - a_2 = m_2 - a + \min(m_1 - g_1, b_+) \geq m_2 - b_+ + m_1 - g_1 \geq g_2.$$

Therefore by Theorem 7.6:

$$J(m_1, m_2) \leq J'(m_1 - b_1, m_2 - b_2) = J_1(m_1 - b_1) + J_2(m_2 - b_2) = 0.$$

Since $J_1(m_1) = J_2(m_2) = 0$, we get also get $\tilde{J}(\mathbf{m}) = 0$.

Suppose now that \mathcal{L} is an L-space link. By the above, H -vanishes in $\mathbf{v} \in R(b_+) + \tilde{\mathbf{g}}$. Corollary 3.32 implies the vanishing of the coefficients of the Alexander polynomial in $R(b_+) + \tilde{\mathbf{g}} + (\frac{1}{2}, \frac{1}{2})$. To show that $\text{HFL}^-(\mathbf{v}) = 0$ for $\mathbf{v} \in R(b_+) + \tilde{\mathbf{g}} + (1, 1)$, note that for such \mathbf{v} one has $H(\mathbf{v} - \mathbf{e}_i) = H(\mathbf{v}) = 0$, so the natural inclusions $A^-(\mathbf{v} - \mathbf{e}_i) \hookrightarrow A^-(\mathbf{v})$ induce isomorphisms on homology. By (3.2), $\text{HFL}^-(\mathbf{v}) = 0$. \square

Remark 7.8. Part (b) of Theorem 7.7 does not hold for non L-space links. For example, the link $L9a31$ in [9] has two components, one being an unknot and one being a trefoil. The linking number of the components is 1, so $\tilde{\mathbf{g}} = (\frac{1}{2}, \frac{3}{2})$. Now the Alexander polynomial is

$$t_1^{-1}t_2^{-2} - t_2^{-2} - 2t_1^{-1}t_2^{-1} + 4t_2^{-1} - t_1t_2^{-1} + 2t_1^{-1} - 5 + 2t_1 - t_1^{-1}t_2 + 4t_2 - 2t_1t_2 - t_2^2 + t_1t_2^2.$$




FIGURE 5. The link $b(24, -5)$. Its two components are unknots.

The term $t_1 t_2^2$ has exponents $(1, 2)$ which belong to $R(0) + \tilde{\mathbf{g}} + (\frac{1}{2}, \frac{1}{2})$. Therefore, Theorem 7.7 would imply that we need at least one positive crossing change to split $L9a31$. Such a crossing change will make the two components have linking number 2, so we will need at least two more crossing changes to make the linking number equal to 0. Altogether, we would need at least three crossing changes to split $L9a31$. However, we can split $L9a31$ in a single move.

7.3. Example: the two-bridge link $b(24, -5)$. We will discuss an example of the two-bridge link $b(24, -5)$ which was shown by Liu [23, Example 3.8] to be an L-space link. It is presented in Figure 5. The orientation of $b(24, -5)$ is as in [23]. The two components have linking number 0. In the notation of LinkInfo [9] it is the link $L9a40$. It was shown in [8, Section 7.1] that the splitting number of this link is 4. The tool was studying the smooth four genus of the link obtained by taking a double branch cover of one of the components of $b(24, -5)$. The splitting number of $b(24, -5)$ can be also detected by the signatures as in [10]. We will show that $sp(b(24, -5)) = 4$ using the J -function.

The Alexander polynomial of $b(24, -5)$ can be found on the LinkInfo web page [9] or calculated using the SnapPy package [11]. We have

$$\Delta(t_1, t_2) = -(t_1 + t_2 + 1 + t_1^{-1} + t_2^{-1})(t_1^{1/2} - t_1^{-1/2})(t_2^{1/2} - t_2^{-1/2}).$$

By Corollary 3.32 the generating function for the \tilde{J} -function equals

$$(7.9) \quad \tilde{\mathbf{J}}(t_1, t_2) = t_1 + t_2 + 1 + t_1^{-1} + t_2^{-1}.$$

Theorem 7.4 implies that we need to make at least one positive crossing change to unlink $b(24, -5)$. As the original linking number is zero and a positive crossing change increases the linking number, we have to compensate the positive crossing change with a negative crossing change, so the splitting number is at least 2. That is all we can deduce from Theorem 7.4.

On the other hand, $J(1, 0) = 1$, so by Theorem 7.7 one needs at least two positive crossing changes to split $b(24, -5)$. As each such crossing change increases the linking number between the two components of $b(24, -5)$, we also need two negative crossing changes. Therefore we have proved the following result.

Proposition 7.10. *The splitting number of $b(24, -5)$ is at least 4.*

It is quite easy to split the $b(24, -5)$ in four moves.




FIGURE 6. The general $b(rq - 1, -q)$ two-bridge link, where $r = 2n + 1$, $q = 2k + 1$. The linking number is $n - k$.




FIGURE 7. The E_1 page of the Batson-Seed spectral sequence for the link $b(48, -7)$ with splitting number 6. It can be shown that $E_2 = E_\infty$ is the Khovanov homology of the unknot. The calculations were made using the KnotKit program [33].

Remark 7.11. SnapPy and the LinkInfo webpage [9] give the Alexander polynomial of $b(24, -5)$ with opposite sign. To choose the sign we notice that the other choice of sign of the Alexander polynomial yields $\tilde{\mathbf{J}}$ with negative coefficients only, hence, for example $J(0, 0) = -1$. This contradicts the property of non-negativity of the J -function. Liu's algorithm in [22, Section 3.3] gives the proper sign of the Alexander polynomial.

7.4. More general two-bridge links. The arguments used in Section 7.3 can be easily generalized for the case of two-bridge links $\mathcal{L}_n = b(4n^2 + 4n, -2n - 1)$. The components of \mathcal{L}_n are unknots with linking number 0. For example, \mathcal{L}_1 is the Whitehead link. It is

proved in [23, Section 3] that all the \mathcal{L}_n are L-space links, and their Alexander polynomials were computed in [23, Section 6], see also [22, Section 3.3]:

$$\Delta_{\mathcal{L}_n}(t_1, t_2) = (-1)^n \sum_{|i+1/2|+|j+1/2|\leq n} (-1)^{i+j} t_1^{i+1/2} t_2^{j+1/2}.$$

Clearly,

$$\Delta(t_1, t_2) = -t_1^{n-1/2} \left(t_2^{1/2} - t_2^{-1/2} \right) + \text{terms of lower degree in } t_1,$$

Hence by Corollary 3.32:

$$\tilde{\mathbf{J}}_{\mathcal{L}_n}(t_1, t_2) = -\frac{\Delta_{\mathcal{L}_n}(t_1, t_2)}{\left(t_1^{1/2} - t_1^{-1/2} \right) \left(t_2^{1/2} - t_2^{-1/2} \right)} = t_1^{n-1} + \text{terms of lower degree in } t_1,$$

and $J(n-1, 0) = \tilde{J}(n-1, 0) = 1$. By Theorem 7.7 one needs at least n positive crossing changes to split \mathcal{L}_n , and the linking number argument from the previous section implies that one needs same number of negative crossing changes. We obtained the following result.

Theorem 7.12. *The splitting number of \mathcal{L}_n is at least $2n$.*

It is quite easy to split the \mathcal{L}_n in $2n$ moves using Figure 6 (where $k = n$).

7.5. Comparison with the Batson–Seed criterion. In [1] Batson and Seed proved an obstruction for splitting number. For the sake of simplicity we formulate the result for a link with two components and for homologies over \mathbb{Z}_2 .

Theorem 7.13. [1] *Suppose $\mathcal{L} = L_1 \cup L_2$ is a two component link and let \mathcal{L}' be a split link with components L_1, L_2 . Then there exists a spectral sequence, whose E_1 page is the Khovanov homology $Kh(L)$ and E_∞ page is the Khovanov homology $Kh(\mathcal{L}')$. If the splitting number of \mathcal{L} is k , then the E_k page is equal to the E_∞ page of this sequence.*

In Figure 7 there is shown the E_1 page of the Batson–Seed spectral sequence for $b(48, -7)$, whose splitting number was shown to be 6. The arrows in the figure correspond to the differential. We have $E_2 = E_\infty$, so Theorem 7.13 implies that $sp(b(48, -7)) \geq 2$. This means that the Batson–Seed criterion does not detect the splitting number of $b(48, -7)$.

For general two-bridge links $b(4n^2 + 4n, -2n - 1)$ we have the following observation, which limits the usage of the Batson–Seed criterion. It is well known to the experts.

Proposition 7.14. *Suppose \mathcal{L} is an alternating non-split link. Then the Batson–Seed spectral sequence collapses at most at the E_3 page.*

Proof. By [21] \mathcal{L} is Khovanov thin, that is, the Khovanov homology is supported on two diagonals. More precisely, if x is a non-trivial element of $Kh(\mathcal{L})$, then $q(x) = 2h(x) - \sigma(\mathcal{L}) \pm 1$, where $q(x)$ is the q -grading, $h(x)$ is the homological grading and $\sigma(\mathcal{L})$ is the signature of \mathcal{L} .

The differential in the Batson–Seed spectral sequence is $d = d_0 + d_1$, where d_0 is the standard differential in the Khovanov complex and d_1 decreases the homological grading by 1 and drops the q -grading by 2. A higher differential d_k changes the (h, q) bigrading by $(1 - 2k, -2k)$, and hence changes the difference $q - 2h$ by $2k - 2$. As \mathcal{L} is thin, the only potentially non-trivial differentials are d_0, d_1 and d_2 . \square

7.6. Cables on the Whitehead link. As a more complicated example, we calculate the splitting number of cables on the Whitehead link. Let $Wh_{p,q}$ denote the link consisting of the (p, q) cable on one of the component of the Whitehead link and the unchanged second component of it. It is clear that the linking number of $Wh_{p,q}$ vanishes. A $(1, 1)$ -surgery on the Whitehead link is an L-space [23, Example 3.1], hence by Proposition 2.8 $Wh_{p,q}$ is an L-space link as long as $1 < p < q$ and p, q are coprime.

The Whitehead link has Alexander polynomial $\Delta = -(t_1^{1/2} - t_1^{-1/2})(t_2^{1/2} - t_2^{-1/2})$. The Alexander polynomial of a cable link was calculated by Turaev in [35, Theorem 1.3.1], which we now state.

Theorem 7.15. *Let $\mathcal{L} = L_1 \cup \dots \cup L_n$ be a link and $\Delta_{\mathcal{L}}(t_1, \dots, t_n)$ its multivariable Alexander polynomial. Let $\mathcal{L}_{p,q}$ be as in the statement of Proposition 2.8 above. Set $T = t_1^q t_2^l \dots t_n^{l_n}$, where $l_j = \text{lk}(L_1, L_j)$. Then*

$$\Delta_{\mathcal{L}_{p,q}}(t_1, \dots, t_n) = \Delta_{\mathcal{L}}(t_1^p, t_2, t_3, \dots, t_n) \frac{T^{p/2} - T^{-p/2}}{T^{1/2} - T^{-1/2}}.$$

It follows from the theorem that the Alexander polynomial of $Wh_{p,q}$ is equal to

$$\Delta_{Wh_{p,q}} = - \left(t_1^{p/2} - t_1^{-p/2} \right) \left(t_2^{1/2} - t_2^{-1/2} \right) \frac{t_1^{pq/2} - t_1^{-pq/2}}{t_1^{q/2} - t_1^{-q/2}}.$$

From this we obtain by Corollary 3.32.

$$\tilde{\mathbf{J}}(t_1, t_2) = \frac{\left(t_1^{p/2} - t_1^{-p/2} \right) \left(t_1^{pq/2} - t_1^{-pq/2} \right)}{\left(t_1^{1/2} - t_1^{-1/2} \right) \left(t_1^{q/2} - t_1^{-q/2} \right)} = t_1^{\delta+(p-1)} + \text{terms of lower degree in } t_1,$$

where $\delta = \frac{1}{2}(p-1)(q-1)$. In particular, $\tilde{\mathbf{J}}(\delta + (p-1), 0) = 1$.

Now the genera of the components of $Wh_{p,q}$ are $g_1 = \delta, g_2 = 0$. By Theorem 7.7 we infer that we need to perform at least p positive multicolored crossing changes to transform $Wh_{p,q}$ into the disjoint sum of $T(p, q)$ and the unknot. The linking number argument shows that we also need p negative crossing changes, hence we obtain the following result.

Proposition 7.16. *The splitting number of the (p, q) -cable on the Whitehead link is at least $2p$.*

It is not hard to find a splitting sequence of length $2p$.

8. ALGEBRAIC LINKS

8.1. The H -function for links of singularities. Let C be a germ of a complex plane curve singularity with branches C_1, \dots, C_n . Its intersection with a small sphere is called an algebraic link. It is shown in [15] that all algebraic links are L-space links. For algebraic links the H -function admits the following description. Let $\gamma_i : (\mathbb{C}, 0) \rightarrow (C_i, 0)$ denote the uniformization of C_i . Define the set

$$\mathcal{J}(\mathbf{v}) := \{f \in \mathbb{C}[[x, y]] : \text{Ord}_0 f(\gamma_i(t)) \geq v_i\}$$

It is clear that $\mathcal{J}(\mathbf{v})$ is in fact a vector subspace of $\mathbb{C}[[x, y]]$. Define the *Hilbert function* of C as

$$(8.1) \quad R(\mathbf{v}) = \text{codim } \mathcal{J}(\mathbf{v}) = \dim_{\mathbb{C}} \mathbb{C}[x, y]/\mathcal{J}(\mathbf{v}).$$

Moreover, set

$$R_i(v_i) = R(0, \dots, 0, v_i, 0, \dots, 0).$$

Notice that for a singularity with one branch, $R(\mathbf{v}) = R_1(v_1)$ is the number of the elements of the semigroup of the singular point in the interval $[0, v_1]$, so (8.1) can be regarded as a generalization the definition of R -function in [4].

We can relate R to the H -function discussed above. Define

$$(8.2) \quad \mathbf{g} = (g_1, \dots, g_n); \quad \tilde{\mathbf{g}} = (\tilde{g}_1, \dots, \tilde{g}_n), \quad \tilde{g}_i = g_i + \frac{1}{2} \sum_{j \neq i} \text{lk}(L_i, L_j),$$

where g_i is the Seifert genus of L_i (or, equivalently, the delta-invariant of the singularity C_i). It is known that for algebraic links $2\tilde{\mathbf{g}}$ is the conductor of the multi-dimensional semigroup of C ; see for instance [20, Chapter 17]. Campillo, Delgado and Gusein-Zade related [6] the Alexander polynomial of an algebraic link to the semigroup of the corresponding curve. Based on their result and (3.16), the following formula for the Hilbert function was obtained in [14]:

Theorem 8.3 (see [14]). *For an algebraic link, one has*

$$(8.4) \quad H(\mathbf{v}) = R(\tilde{\mathbf{g}} - \mathbf{v}), \quad J(\mathbf{v}) = R(\mathbf{g} - \mathbf{v}).$$

Remark 8.5. It was proven in [7] that for all plane curve singularities the Hilbert function satisfies the following symmetry property:

$$(8.6) \quad R(2\tilde{\mathbf{g}} - \mathbf{v}) = R(\mathbf{v}) + |\tilde{\mathbf{g}}| - |\mathbf{v}|.$$

Indeed, this agrees with the symmetry property (3.23) of H .

Theorem 8.7. *We have the following inequality for the Hilbert function of a plane curve singularity.*

$$0 \geq R(\mathbf{v}) - \sum_{i=1}^n R_i(v_i) \geq - \sum_{i < j} \text{lk}(L_i, L_j).$$

Both inequalities are sharp.

Corollary 8.8. *For an algebraic link, for all \mathbf{v} :*

$$0 \geq \tilde{J}(\mathbf{v}) \geq - \sum_{i < j} \text{lk}(L_i, L_j)$$

Proof. By (8.4), $J(\mathbf{v}) = R(\mathbf{g} - \mathbf{v})$. Similarly, $J_i(\mathbf{v}_i) = R_i(g_i - v_i)$, so it remains to apply the theorem to the vector $\mathbf{g} - \mathbf{v}$. \square

Remark 8.9. Corollary 8.8 can be compared with Theorem 7.4. Indeed, all crossings in an algebraic link are positive, so $t_+ = 0$, and by the above corollary we get $t_- \geq \sum_{i < j} \text{lk}(L_i, L_j)$. In other words, to split an algebraic link one needs to change exactly $\sum_{i < j} \text{lk}(L_i, L_j)$ crossings from positive to negative. It is well known that the splitting number of an algebraic link is equal to $\sum_{i < j} \text{lk}(L_i, L_j)$.

The following two lemmas will be used in the proof of Theorem 8.7.

Lemma 8.10. *For $\mathbf{u}, \mathbf{v} \in \mathbb{Z}^n$, one has*

$$R(\mathbf{u}) + R(\mathbf{v}) \geq R(\min(\mathbf{u}, \mathbf{v})) + R(\max(\mathbf{u}, \mathbf{v})).$$

Proof. Indeed, $\mathcal{J}(\mathbf{u}), \mathcal{J}(\mathbf{v}) \subset \mathcal{J}(\min(\mathbf{u}, \mathbf{v}))$ and $\mathcal{J}(\mathbf{u}) \cap \mathcal{J}(\mathbf{v}) = \mathcal{J}(\max(\mathbf{u}, \mathbf{v}))$. One has $\dim \mathcal{J}(\min(\mathbf{u}, \mathbf{v}))/\mathcal{J}(\mathbf{u}) + \dim \mathcal{J}(\min(\mathbf{u}, \mathbf{v}))/\mathcal{J}(\mathbf{v}) \geq \dim \mathcal{J}(\min(\mathbf{u}, \mathbf{v}))/\mathcal{J}(\max(\mathbf{u}, \mathbf{v}))$, therefore

$$-R(\min(\mathbf{u}, \mathbf{v})) + R(\mathbf{u}) - R(\min(\mathbf{u}, \mathbf{v})) + R(\mathbf{v}) \geq -R(\min(\mathbf{u}, \mathbf{v})) + R(\max(\mathbf{u}, \mathbf{v})).$$

□

Lemma 8.11. *Suppose that $\mathbf{u}, \mathbf{v} \in \mathbb{Z}^n, 0 \leq \mathbf{u} \leq \mathbf{v}$. Then*

$$R(\mathbf{v}) - R(\mathbf{u}) \leq \sum_{i=1}^n (R_i(v_i) - R_i(u_i)).$$

Proof. Consider a sequence of lattice points $\mathbf{v}^{(i)} = (u_1, \dots, u_i, v_{i+1}, \dots, v_n)$, so that $\mathbf{u} = \mathbf{v}^{(n)}$ and $\mathbf{v} = \mathbf{v}^{(0)}$. Let \mathbf{e}_i denote the i -th coordinate vector. Then

$$\max(\mathbf{v}^{(i)}, v_i \mathbf{e}_i) = \mathbf{v}^{(i-1)}, \quad \min(\mathbf{v}^{(i)}, v_i \mathbf{e}_i) = u_i \mathbf{e}_i,$$

hence by Lemma 8.10:

$$R_i(v_i) + R(\mathbf{v}^{(i)}) \geq R(\mathbf{v}^{(i-1)}) + R_i(u_i),$$

so

$$R(\mathbf{v}^{(i-1)}) - R(\mathbf{v}^{(i)}) \leq R_i(v_i) - R_i(u_i),$$

and

$$R(\mathbf{v}) - R(\mathbf{u}) = R(\mathbf{v}^{(0)}) - R(\mathbf{v}^{(n)}) = \sum_{i=1}^n (R(\mathbf{v}^{(i-1)}) - R(\mathbf{v}^{(i)})) \leq \sum_{i=1}^n (R_i(v_i) - R_i(u_i)).$$

□

Proof of Theorem 8.7. By Lemma 8.11, one has

$$R(\mathbf{v}) = R(\mathbf{v}) - R(0) \leq \sum_{i=1}^n (R_i(v_i) - R_i(0)) = \sum_{i=1}^n R_i(v_i).$$

Furthermore, if $\mathbf{u} \gg \mathbf{0}$, then by (8.6) $R(\mathbf{u}) = |\mathbf{u}| - |\tilde{\mathbf{g}}|$. By Lemma 8.11, we get

$$\begin{aligned} R(\mathbf{u}) - R(\mathbf{v}) &\leq \sum_{i=1}^n (R_i(u_i) - R_i(v_i)), \text{ so} \\ |\mathbf{u}| - |\tilde{\mathbf{g}}| - R(\mathbf{v}) &\leq \sum_{i=1}^n (u_i - g_i - R_i(v_i)), \text{ that is} \\ R(\mathbf{v}) - \sum_{i=1}^n R_i(v_i) &\geq -|\tilde{\mathbf{g}}| + \sum_{i=1}^n g_i = -\sum_{i < j} l_{ij}. \end{aligned}$$

□

8.2. Semicontinuity of the Hilbert function. We can use Theorem 6.1 to give a topological proof of semigroup semicontinuity property under δ -constant deformation, generalizing the result of [5] for many components. We refer the reader to [2, 14] for other approaches to semicontinuity property of semigroups.

Suppose $F_t: (\mathbb{C}^2, 0) \rightarrow (\mathbb{C}, 0)$ is a family of polynomials depending on a parameter $t \in (-\varepsilon, \varepsilon) \subset \mathbb{R}$. We assume that for every t the curve $F_t^{-1}(0)$ has an isolated singularity with n branches.

Theorem 8.12. *Assume that the deformation is δ -constant. Then for any $\mathbf{m} \in \mathbb{Z}^n$ and $t \neq 0$ sufficiently close to 0 we have*

$$R_t(\mathbf{m}) \geq R_0(\mathbf{m}).$$

Proof. We follow the proof of [4, Theorem 2.15]. Take a ball $B \subset \mathbb{C}^2$ with center at 0 such that $F_0^{-1}(0) \cap \partial B$ is the link of the singularity of $F_0^{-1}(0)$ at 0. Denote this link by \mathcal{L}_2 .

Choose t sufficiently small. Then $F_t^{-1}(0) \cap \partial B$ is still isotopic to \mathcal{L}_2 . Choose a smaller ball B' with center at 0 such that $F_t^{-1}(0) \cap \partial B'$ is the link of the singularity of $F_t^{-1}(0)$ at 0. Denote this link by \mathcal{L}_1 . For $i = 1, 2$, let L_{i1}, \dots, L_{in} be the components of the link \mathcal{L}_i . Denote by g_{i1}, \dots, g_{in} the Seifert genus of the corresponding component. Let \mathbf{g} be as in (8.2).

Similarly to [5, Lemma 2.3], we construct an APSIC from \mathcal{L}_1 to \mathcal{L}_2 in the following way. As the deformation is δ -constant, we can find a complex parametrization ψ of $F_t^{-1}(0) \cap B''$ (where B'' is a ball slightly larger than B), whose domain is a disjoint union of n disks D_1, \dots, D_n . Set $D = D_1 \sqcup \dots \sqcup D_n$. Perturb ψ to a complex analytic map $\tilde{\psi}$ such that $\tilde{\psi}$ has only generic singularities, that is, positive double points. For small perturbation the links $\mathcal{L}'_2 = \tilde{\psi}(D) \cap \partial B$ and $\mathcal{L}'_1 = \tilde{\psi}(D) \cap \partial B'$ are isotopic to \mathcal{L}_2 and \mathcal{L}_1 respectively. The APSIC is the intersection $\tilde{\psi}(D) \cap (B \setminus B')$. The number of double monochromatic double points of the i -th component (denoted by η_{ii} in Section 5) is calculated as in [5, Lemma 2.3] and is equal to $g_{i2} - g_{i1}$.

We use Theorem 6.1 to obtain

$$(8.13) \quad J_2(\mathbf{m}') \leq J_1(\mathbf{m}' - \mathbf{k}),$$

where $\mathbf{k} = (\eta_{11}, \dots, \eta_{nn}) = \mathbf{g}_2 - \mathbf{g}_1$ and $\mathbf{m}' \in \mathbb{Z}^n$ is arbitrary. Substituting (8.4) into (8.13) we obtain

$$R_2(\mathbf{m}) = J_2(\mathbf{g}_2 - \mathbf{m}) \leq J_1(\mathbf{g}_2 - \mathbf{m} - \mathbf{k}) = J_1(\mathbf{g}_1 - \mathbf{m}) = R_1(\mathbf{m}).$$

□

REFERENCES

- [1] J. Batson, C. Seed, *A Link Splitting Spectral Sequence in Khovanov Homology*, Duke Math. J. **164**, no. 5 (2015), 801–841.
- [2] J. Bodnár, D. Celoria, M. Golla, *A note on cobordisms of algebraic knots*, preprint 2015, arxiv:1509.08821.
- [3] M. Borodzik, S. Friedl, M. Powell, *Blanchfield forms and Gordian distance*, J. Math. Soc. Japan **68** (2016), no. 3, 1047–1080.
- [4] M. Borodzik, C. Livingston, *Heegaard Floer homologies and rational cuspidal curves*, Forum of Math. Sigma, **2** (2014), e28, 23 pages.
- [5] M. Borodzik, C. Livingston, *Semigroups, d-invariants and deformations of cuspidal singular points of plane curves*, J. Lond. Math. Soc. **93** (2016), no. 2, 439–463.
- [6] A. Campillo, F. Delgado, S.M. Gusein-Zade, *The Alexander polynomial of a plane curve singularity via the ring of functions on it*, Duke Math. J. **117** (2003), no. 1, 125–156.
- [7] A. Campillo, F. Delgado, K. Kiyek, *Gorenstein property and symmetry for one-dimensional local Cohen-Macaulay rings*, Manuscripta Math. 83 (1994), no. 3-4, 405–423.
- [8] J. Cha, S. Friedl, M. Powell, *Splitting numbers of links*, to appear in Proc. Edin. Math. Soc., arxiv:1308.5638
- [9] J. Cha, C. Livingston, *LinkInfo: Table of Knot Invariants*, <http://www.indiana.edu/~linkinfo>, January 9, 2016.
- [10] D. Cimasoni, A. Conway, and L. Zacharova, *Splitting numbers and signatures*, Proc. Amer. Math. Soc. **144** (2016), no. 12, 5443–5455.
- [11] M. Culler, N. Dunfield, and J. Weeks, *SnapPy, a computer program for studying the geometry and topology of 3-manifolds*, <http://snappy.computop.org>.
- [12] D. Eisenbud, W. Neumann, *Three-dimensional link theory and invariants of plane curve singularities*, Annals Math. Studies **110**, Princeton University Press, Princeton, 1985.

- [13] E. Gorsky, J. Hom, *Cable links and L-space surgeries*, to appear in Quant. Topol., arxiv: 1502.05425.
- [14] E. Gorsky, A. Némethi. *Lattice and Heegaard Floer homologies of algebraic links*, Int. Math. Res. Not. 2015, no. 23, 12737–12780.
- [15] E. Gorsky, A. Némethi. *Links of plane curve singularities are L-space links*, Algebr. Geom. Topol. **16** (2016), no. 4, 1905–1912.
- [16] M. Hedden, *On knot Floer homology and cabling*, Algebr. Geom. Topol. **5** (2005), 1197–1222.
- [17] J. Hillman, *Algebraic invariants of links*, second edition, Series on Knots and Everything, 52. World Scientific Publishing Co. Inc, River Edge, NJ, (2012).
- [18] J. Hom, T. Lidman, N. Zufelt, *Reducible surgeries and Heegaard Floer homology*, Math. Res. Lett. **22** (2015), no. 3, 763–788.
- [19] A. Kawauchi, *The Alexander polynomials of immersed concordant links*, Bol. Soc. Mat. Mex. **20** (2014), no. 2, 559–578.
- [20] E. Kunz, *Introduction to plane algebraic curves*, Birkhäuser Boston Inc., Boston 2005.
- [21] E. Lee, *An endomorphism of the Khovanov invariant*, Adv. Math. **197** (2005), no. 2, 554–586.
- [22] Y. Liu, *Heegaard Floer homology and surgeries on two-bridge links*, preprint, arXiv:1402.5727.
- [23] Y. Liu, *L-space surgeries on links*, to appear in Quant. Topol., arXiv:1409.0075.
- [24] C. Manolescu, P. Ozsváth, *Heegaard Floer homology and integer surgeries on links*, preprint 2010, arXiv:1011.1317.
- [25] M. Nagel, B. Owens, *Unlinking information from 4-manifolds*, Bull. Lond. Math. Soc. **47** (2015), no. 6, 964–979.
- [26] Y. Ni, Z. Wu, *Heegaard Floer correction terms and rational genus bounds*, Adv. Math. **267** (2014), 360–380.
- [27] B. Owens, S. Strle, *Immersed disks, slicing numbers and concordance unknotting numbers*, preprint 2013, arxiv:1311.6702, to appear in Comm. Anal. Geom.
- [28] P. Ozsváth, Z. Szabó, *Absolutely graded Floer homologies and intersection forms for four-manifolds with boundary*, Adv. Math. **173** (2003), 179–261.
- [29] P. Ozsváth, Z. Szabó, *On knot Floer homology and lens space surgeries*, Topology, **44** (2005), 1281–1300.
- [30] P. Ozsváth, Z. Szabó. *Holomorphic discs, link invariants and the multi-variable Alexander polynomial*. Algebr. Geom. Topol. **8** (2008), no. 2, 615–692.
- [31] J. Rasmussen, *Floer homology and knot complements*, Ph.D. thesis, Harvard University, 2003.
- [32] J. Rasmussen, S. Rasmussen, *Floer Simple Manifolds and L-Space Intervals*, preprint 2015, arxiv: 1508.05900.
- [33] C. Seed, *Knotkit, a computer program to calculate Khovanov homology*, <https://github.com/cseed/knotkit/>.
- [34] G. Torres, *On the Alexander polynomial*. Ann. of Math. **57**(1953), 57–89.
- [35] V. Turaev, *Reidemeister torsion in knot theory*, Russian Math. Surveys 41 (1986), No. 1, 119–182.
- [36] V. Turaev, *An introduction to combinatorial torsion*, Notes taken by Felix Schlenk, Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, 2001.

INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCE, UL. ŚNIADECKICH 8, WARSAW,
POLAND

INSTITUTE OF MATHEMATICS, UNIVERSITY OF WARSAW, UL. BANACHA 2, 02-097 WARSAW,
POLAND

E-mail address: mchoro@mimuw.edu.pl

DEPARTMENT OF MATHEMATICS, UC DAVIS, ONE SHIELDS AVENUE DAVIS CA 95616 USA

NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS, VAVILOVA 7, MOSCOW, RUSSIA

E-mail address: egorskiy@math.ucdavis.edu