

LETTER

Magnetization pumping and dynamics in a Dzyaloshinskii-Moriya magnet

To cite this article: Alexey A. Kovalev and Utkan Güngörđü 2015 *EPL* **109** 67008

View the [article online](#) for updates and enhancements.

Related content

- [Thermomagnonic spin transfer and Peltier effects in insulating magnets](#)

Alexey A. Kovalev and Yaroslav Tserkovnyak

- [Pure spin currents in magnetically ordered insulator/normal metal heterostructures](#)

Matthias Althammer

- [Landau-Lifshitz theory of the magnon-drag thermopower](#)

B. Flebus, R. A. Duine and Y. Tserkovnyak

Recent citations

- [Pumping of magnons in a Dzyaloshinskii-Moriya ferromagnet](#)

Alexey A. Kovalev *et al*

- [Spin torque and Nernst effects in Dzyaloshinskii-Moriya ferromagnets](#)

Alexey A. Kovalev and Vladimir Zyuzin

- [Stability of skyrmion lattices and symmetries of quasi-two-dimensional chiral magnets](#)

Utkan Güngörđü *et al*

Magnetization pumping and dynamics in a Dzyaloshinskii-Moriya magnet

ALEXEY A. KOVALEV and UTKAN GÜNGÖRDÜ

Department of Physics and Astronomy and Nebraska Center for Materials and Nanoscience, University of Nebraska Lincoln, NE 68588, USA

received 22 January 2015; accepted in final form 22 March 2015
published online 10 April 2015

PACS 75.78.-n – Magnetization dynamics
PACS 75.60.Jk – Magnetization reversal mechanisms
PACS 85.80.Lp – Magnetothermal devices

Abstract – We formulate a phenomenological description of thin ferromagnetic layers with inversion asymmetry where the single-domain magnetic dynamics experiences magnon current-induced torques and leads to magnon-motive forces. We first construct a phenomenological theory based on irreversible thermodynamics, taking into account the symmetries of the system. Furthermore, we confirm that these effects originate from Dzyaloshinskii-Moriya interactions from the analysis based on the stochastic Landau-Lifshitz-Gilbert equation. Our phenomenological results generalize to a general form of Dzyaloshinskii-Moriya interactions and to other systems, such as pyrochlore crystals and chiral magnets. Possible applications include spin current generation, magnetization reversal and magnonic cooling.

Copyright © EPLA, 2015

Introduction. – Spincaloritronics studies various thermal effects relying on the spin degree of freedom [1]. The most prominent examples are the spin Seebeck effect [2], the spin Peltier effect [3,4], and thermally induced motion of domain walls [5,6]. Apart from fascinating physics, in some instances related to the appearance of spin-motive force [7], these studies might offer new ways for energy harvesting, cooling, and magnetization control [8,9]. Spincaloritronics might also help with the development of electronics relying on *pure* spin currents [3,10] which contrasts conventional electronics. The known ways to create pure spin currents include non-local spin injection in spin valves, optical injection by circularly polarized light, spin pumping, and spin Hall effect [11].

Pure spin currents in a form of magnon flow have attracted considerable attention recently as they can transfer signals [12] and even realize magnonic logic circuits with low dissipation and without generation of Oersted fields [13]. On the other hand, magnons can exhibit similar phenomena to electrons, *e.g.*, spin-transfer torque on magnetic textures such as domain walls [5] and skyrmions [14,15], Hall effect, and topologically protected edge states [16]. Such magnonic spin currents can be driven by radio-frequency fields or temperature gradients. In the ferrimagnetic insulator yttrium iron garnet (YIG)

magnons can travel over large distances without interruption due to remarkably low Gilbert damping [17].

Relativistic effects result in interesting physics in the context of spin currents. Discovery of spin-orbit torques [18] allows for magnetization control by charge currents in bilayers consisting of a layer with strong spin-orbit interactions, *e.g.*, metal or topological insulator, and a ferromagnet [19]. Spin-orbit torques are often interpreted in terms of a Rashba contribution [19] and a spin Hall contribution [20] while in a general scenario it is more useful to separate reactive and dissipative contributions [21]. Magnons, on the other hand, can be influenced by Dzyaloshinskii-Moriya interactions (DMI) similar to how electrons are influenced by spin-orbit interactions [22–24]. In particular, a spin-orbit-like torque generated by spin waves has been suggested [24]. DMI can result from spin-orbit interactions in systems with broken inversion symmetry [25] or from structural asymmetry in ultrathin magnetic bilayers [26]. Thus, one can naturally expect magnon analogs of spin-orbit torques and charge pumping observed in metal/ferromagnet bilayers (see fig. 1).

In this paper, we develop a general phenomenological description of the interplay between magnetization dynamics, magnon currents, and temperature gradients in




Fig. 1: (Color online) Single-domain magnetization dynamics induced by microwave field pumps magnon j and spin $j_s = -\hbar j$ currents by virtue of Dzyaloshinskii-Moriya interactions. This can develop a temperature gradient along the sample. Alternatively, a temperature gradient can result in magnon current and torque on uniform magnetization according to the Onsager reciprocity principle.

single-domain ferromagnetic layers lacking inversion symmetry. We accompany our analysis by a model based on the stochastic Landau-Lifshitz-Gilbert (LLG) equation with DMI characteristic to ultrathin magnetic layers with structural asymmetry. We obtain reactive and dissipative torques on uniform magnetization and discuss the possibility to reverse magnetization by magnon currents and temperature gradients. We discuss pumping of magnonic spin currents by precessing single-domain magnetization and analyze the feasibility of magnonic cooling.

Phenomenology of thermal magnons with DMI.

– In this section, we employ general principles of non-equilibrium thermodynamics [27] in order to formulate the phenomenology of thermal magnons applicable to a system with interfacial inversion asymmetry as in fig. 1. We begin by constructing a general phenomenological description of magnonic and thermal currents in a single-domain ferromagnet in which thermodynamic variables represent the direction of the reduced (averaged over the magnonic excitations) spin density \mathbf{m}_s (for convenience the index s is dropped in this section), density of magnons ρ and density of energy ρ_u . We assume that the single-domain ferromagnet is taken out of equilibrium by applying temperature and chemical potential gradients (non-zero chemical potential for magnons can be created by, *e.g.*, microwave pumping [28]). An appropriate equation of motion then determines how the ferromagnet evolves back towards equilibrium. We now write the rate of the entropy production [29]:

$$\dot{\mathbb{S}} = - \int d^3r \frac{\partial \cdot \mathbf{j}_u + \mu \dot{\rho} + \mathbf{H}_{\text{eff}} \cdot \dot{\mathbf{m}}}{T}, \quad (1)$$

where we introduced the magnon (\mathbf{j}) and energy (\mathbf{j}_u) currents, and the conjugate/force corresponding to the spin density direction defined as $-\delta_{\mathbf{m}} \dot{\mathbb{S}}|_{\mathbf{j}_u(\mathbf{j})=0} = \mathbf{H}_{\text{eff}}/T$. It is convenient to introduce the modified energy current $\mathbf{j}_q = \mathbf{j}_u - \mu \mathbf{j}$ in order to arrive at the more familiar equation for the rate of the entropy production [29]:

$$\dot{\mathbb{S}} = \int d^3r \left(-\frac{\partial T}{T^2} \cdot \mathbf{j}_q - \frac{\partial \mu}{T} \cdot \mathbf{j} - \frac{\mathbf{H}_{\text{eff}}}{T} \cdot \dot{\mathbf{m}} \right). \quad (2)$$

Here we integrated the term involving \mathbf{j}_q by parts, used the local conservation laws of energy and number of magnons, $\dot{\rho} = -\partial \cdot \mathbf{j} - \rho/\tau$ and $\dot{\rho}_u = -\partial \cdot \mathbf{j}_u$, and disregarded the term $\mu \rho/\tau$, where τ corresponds to the lifetime of magnons. This is possible when the number of magnons is approximately conserved. One can also consider the opposite limit in which magnons quickly relax to the local equilibrium without the build-up of large μ ($\mu \approx 0$) as, in this case, the term $\mu \rho/\tau$ can be also disregarded. The remaining conjugates/forces can be immediately identified as $-\delta_{\mathbf{j}_q} \dot{\mathbb{S}}|_{\mathbf{m}, \mathbf{j}_q=0} = -\partial(1/T)$ and $-\delta_{\mathbf{j}} \dot{\mathbb{S}}|_{\mathbf{m}, \mathbf{j}_q=0} = \partial \mu/T$.

We now relate the currents \mathbf{j} and \mathbf{j}_q and the time derivative of the spin density direction $\dot{\mathbf{m}}$ to the thermodynamic conjugates via kinetic coefficients. By accounting for the structural asymmetries defined by the \mathbf{n} -axis in the figure, we obtain the magnon/energy current expansion in terms of the chemical potential and temperature gradients as well as the magnetization dynamics responsible for fictitious fields on the magnons:

$$\begin{aligned} -\partial_\alpha \mu &= \hat{\varrho} \mathbf{j} + \hat{\Pi} \partial T/T - (\eta \mathbf{m} \times \mathcal{D}_\alpha \mathbf{m} + \vartheta \mathcal{D}_\alpha \mathbf{m}) \cdot \dot{\mathbf{m}}, \\ (j_q)_\alpha &= \hat{\Pi}^T \mathbf{j} - \hat{\kappa} \partial T - (\eta_1 \mathbf{m} \times \mathcal{D}_\alpha \mathbf{m} + \vartheta_1 \mathcal{D}_\alpha \mathbf{m}) \cdot \dot{\mathbf{m}}, \end{aligned} \quad (3)$$

where $\mathcal{D}_\alpha = (D/A)(\mathbf{n} \times \mathbf{e}_\alpha) \times$ is a part of the chiral derivative accounting for DMI, \mathbf{e}_α are basis vectors [21,30], η and ϑ are the so-called reactive (also referred to as spin-motive force) and dissipative coefficients [21] (generally $\mathbf{m} \cdot \mathbf{n}$ dependent), $\hat{\varrho}$, $\hat{\Pi}$ and $\hat{\kappa}$ are the resistivity, Peltier and thermal conductivity tensors, respectively, which are in general temperature dependent. As will become clear from the following discussion, it is convenient to invert the equation for the magnon current, as has been done in eq. (3). In the remaining part of the paper we will not consider corrections corresponding to η_1 and ϑ_1 as these contributions do not appear in our microscopic treatment. The axial symmetry around the \mathbf{n} -axis leads to the separation of the conductivity tensor $\hat{g} \equiv \hat{\varrho}^{-1}$ and the thermal conductivity tensor $\hat{\kappa}$ into the longitudinal g and κ , and the Hall g_H and κ_H contributions where $\hat{g} = g + g_H \mathbf{n} \times$ and $\hat{\kappa} = \kappa + \kappa_H \mathbf{n} \times$. The LLG equation becomes

$$\begin{aligned} \mathfrak{s}(1 + \alpha \mathbf{m} \times) \dot{\mathbf{m}} + \mathbf{m} \times \mathbf{H}_{\text{eff}} &= (\eta + \vartheta \mathbf{m} \times) j_\alpha \mathcal{D}_\alpha \mathbf{m} \\ &+ (\eta_1 + \vartheta_1 \mathbf{m} \times) (\partial_\alpha T/T) \mathcal{D}_\alpha \mathbf{m}, \end{aligned} \quad (4)$$

where \mathfrak{s} is the reduced spin density and the form of the torques on the right-hand side is dictated by the Onsager reciprocity principle. Note that in the simplest approximation \mathbf{H}_{eff} can be calculated from an appropriate functional expressed in terms of the direction of the spin density $\mathbf{m}(\mathbf{r}, t)$. In a more general setting, one may need to expand \mathbf{H}_{eff} in terms of small $\partial_i T$ and $\partial_i \mu$.

Torques by magnons from the stochastic LLG equation. – The phenomenological equation (4) can be derived from the LLG equation with DMI. To this end, we consider a ferromagnet with homogeneous magnetization well below the Curie temperature. We employ the

stochastic LLG equation:

$$s(1 + \alpha \mathbf{m} \times) \dot{\mathbf{m}} + \mathbf{m} \times (\mathbf{H}_{\text{eff}} + \mathbf{h}) = 0. \quad (5)$$

Here by s we denote the saturation spin density and by $\mathbf{m}(\mathbf{r}, t)$ a unit vector in the direction of the spin density, \mathbf{h} is the random Langevin field. According to the LLG phenomenology, the effective magnetic field can be found from the free energy, *i.e.* $\mathbf{H}_{\text{eff}} = -\delta_{\mathbf{m}} F$. In the discussion of thermal magnons, we disregard magnetostatic and magnetocrystalline anisotropies assuming sufficiently high temperatures, and the large k -limit. At very low temperatures, one needs to account for anisotropies as they lead to mixing of circular components of spin waves, and renormalization of phenomenological parameters in our theory. We consider the free-energy density $F = (A/2)(\partial_{\alpha} \mathbf{m})^2 + D \mathbf{m} \cdot ([\mathbf{n} \times \partial] \times \mathbf{m}) - \mathbf{m} \cdot \mathbf{H}$, where $A_{\text{ex}} = A/M_s$ is the exchange stiffness, D_{dm} describes DMI with $D \equiv D_{\text{dm}} M_s$, M_s is the saturation magnetization, and \mathbf{H}_e is the external magnetic field with $\mathbf{H} \equiv \mathbf{H}_e M_s$. This form of DMI can be derived for systems with the axial symmetry around the \mathbf{n} -axis and an interfacial inversion asymmetry along the \mathbf{n} -axis [26].

For simplicity, we assume that the slow-dynamics magnetization is static as we can account for the time-dependent effects by involving the Onsager reciprocity principle. Without loss of generality, we also assume a uniform temperature gradient along the x -axis. The vectors for the fast $\mathbf{m}_f(\mathbf{r}, t)$ and slow $\mathbf{m}_s(\mathbf{r}, t)$ magnetization dynamics are related by $\mathbf{m} = (1 - \mathbf{m}_f^2)^{1/2} \mathbf{m}_s + \mathbf{m}_f$, where $\mathbf{m}_s \cdot \mathbf{m}_f = 0$. The magnons are considered in a coordinate system in which the z' -axis points along the spin density of the slow dynamics (see fig. 1). In this coordinate system, small excitations will only have m'_x and m'_y components. We obtain the equation describing the fast magnetization dynamics by linearizing the LLG equation:

$$is\partial_t(1 - i\alpha)m_+ = [A(i\partial)^2 - 2iD\partial \cdot (\mathbf{n} \times \mathbf{m}_s) + H] m_+. \quad (6)$$

In the absence of anisotropy terms we can disregard the coupling between the circular components $m_{\pm} = m'_x(\mathbf{r}, t) \pm i m'_y(\mathbf{r}, t)$, where $m'_{x(y)}(\mathbf{r}, t)$ are the transverse components of the spin wave [31]. As was suggested in previous studies [23], the presence of DMI in eq. (6) leads to thermal magnons with shifted spectrum $\omega_k = [H + A(\mathbf{k} + \mathbf{k}_0)^2 - Ak_0^2]/s$, where $\mathbf{k}_0 = (D/A)(\mathbf{n} \times \mathbf{m}_s)$ describes the shift in the magnon momentum induced by DMI.

In eq. (5), we introduced the random Langevin field \mathbf{h} corresponding to thermal fluctuations at temperature T . According to the fluctuation dissipation theorem the random fields are described by the correlator [32]

$$\langle h_i(\mathbf{r}, t)h_j(\mathbf{r}', t') \rangle = 2\alpha k_B T(\mathbf{r})\delta_{ij}\delta(\mathbf{r} - \mathbf{r}')\delta(t - t'). \quad (7)$$

We treat the fast magnetization dynamics as a linear response to the fluctuating field. In addition, we consider the slow, single-domain magnetic dynamics with a long characteristic time-scale, *e.g.*, corresponding to a ferromagnetic resonance which is typically in GHz range.

We calculate the force that fast oscillations exert on the slow magnetization dynamics \mathbf{m}_s by employing the method developed in ref. [15]. The force due to rapid oscillations can only come from the second-order terms in $\mathbf{m}_f(\mathbf{r}, t)$. However, direct application of the expressions from ref. [15] relying on the exchange contributions results in vanishing force. In the effective field $\mathbf{H}_{\text{eff}} = \mathbf{H} + A\partial^2 \mathbf{m} - 2D[\mathbf{n} \times \partial] \times \mathbf{m}$ only the higher-order terms corresponding to DMI lead to a torque in the absence of magnetic textures:

$$\mathcal{T} = -\langle \mathbf{m}_f \times \mathbf{H}_{\text{eff}} \rangle = 2D\langle \mathbf{m}_f \times [(\mathbf{n} \times \partial) \times \mathbf{m}_f] \rangle, \quad (8)$$

where $\langle \dots \rangle$ stands for averaging over the fast oscillations induced by the random Langevin field. By analogy with the transverse spin accumulation in the context of spin-orbit torques [19], we introduce an auxiliary quantity with the meaning of the transverse spin accumulation, $\mathcal{S} = (1/A)\mathcal{T} \times \mathbf{m}_s$. By coarse-graining various contributions, we obtain the following expression for the transverse accumulation of magnon spins originating from DMI (*i.e.*, the exchange term leads to a vanishing contribution):

$$\mathcal{S} = \frac{2D}{A} \langle \mathbf{m}_f [\mathbf{n} \cdot (\mathbf{m}_s \cdot \partial) \mathbf{m}_f] \rangle. \quad (9)$$

We recall that we consider the reference frame with $\mathbf{z}' = \mathbf{m}_s$, in which vectors \mathcal{S} , \mathbf{m}_f , and $\partial_{\alpha} \mathbf{m}_f$ are in the x' - y' plane. We can simplify expressions by switching to complex notations $a \equiv a'_{x'} + ia'_{y'}$, where \mathbf{a} is an arbitrary vector in the x' - y' plane. In the simplified notations, we obtain the following expression for the spin accumulation $S = (D/A)\langle m_+(\mathbf{r}, t)[\mathbf{v} \cdot \partial]m_-(\mathbf{r}, t) \rangle$, where $\mathbf{v} \equiv (n_{z'}, in_{z'}, -n_{x'} - in_{y'})$ and $\partial = (\partial_{x'}, \partial_{y'}, \partial_{z'})$.

Since we are interested in the steady-state solution, we Fourier-transform $m_{\pm}(\mathbf{r}, t)$ with respect to time and transverse coordinate:

$$m_{\mp}(\mathbf{q}, \omega, x) = \int \frac{d^{d-1} \rho d\omega}{(2\pi)^d} e^{\pm i(\omega t - \mathbf{q} \cdot \rho)} m_{\mp}(\mathbf{r}, t), \quad (10)$$

which leads to the expression for the spin accumulation:

$$\mathcal{S} = \frac{D}{A} \int \frac{d^l \mathbf{q} d^l \mathbf{q}' d\omega' d\omega}{(2\pi)^{2d}} \langle m_+(\mathbf{q}, \omega, x) \mathbf{v} \cdot \partial m_-(\mathbf{q}', \omega', x) \rangle, \quad (11)$$

where $l = d - 1$, $d = 2$ or 3 depending on the dimensionality of the magnet and $\mathcal{S} = \mathcal{S}_{x'} + i\mathcal{S}_{y'}$ describes two components of the spin accumulation leading to the reactive and dissipative torques. As we are interested in the linear response to the random Langevin field, the LLG equation (5) takes the form of the following equation:

$$A[\partial_x^2 + \kappa^2] m_-(x, \mathbf{q}, \omega) = h(x, \mathbf{q}, \omega), \quad (12)$$

where the left-hand side of this equation coincides with eq. (6) and $\kappa^2 = [(1 + i\alpha)s\omega - H]/A - k_0^2 - q^2$. The momentum shift by \mathbf{k}_0 discussed after eq. (6) can be removed by a gauge transformation, thus the effect of DMI

on magnons can be accounted for by renormalizing κ^2 . We solve eq. (12) by employing the Green's function for the Helmholtz equation, $G(x - x_0) = ie^{ik|x-x_0|}/(2k)$. We substitute this solution in eq. (11) and carry through integrations over variables x and x_1 by employing the Fourier-transformed correlator for the stochastic fields [33]:

$$\frac{\langle h(x, \mathbf{q}, \omega)^* h(x_1, \mathbf{q}', \omega') \rangle}{4(2\pi)^d \alpha s k_B} = T(x) \delta(x - x_1) \delta(\mathbf{q} - \mathbf{q}') \delta(\omega - \omega'). \quad (13)$$

We arrive at the expression for the magnon spin torque:

$$\mathcal{T} = -i \frac{s\alpha D}{4A^2} \int \frac{d^{d-1}\mathbf{q}}{(2\pi)^d} \int_{\omega_0}^{\infty} d\omega \frac{[\mathbf{v} \cdot \partial](\hbar\omega \coth \frac{\hbar\omega}{2k_B T(\mathbf{r})})}{\kappa^* [\text{Im}(\kappa)]^2}, \quad (14)$$

where we had to limit the frequency integration by $\omega_0 = (H + Aq^2 + Ak_0^2)/s$ as within this description we are only interested in magnonic excitations with energies above the magnonic gap. In addition, we replaced $2k_B T \rightarrow \hbar\omega \coth(\hbar\omega/2k_B T)$ by employing the quantum fluctuation dissipation theorem in order to introduce a high-frequency cutoff at $\hbar\omega \gg k_B T$. Note that eq. (14) contains terms that can be identified as magnon currents obtained from the Boltzmann approach.

We can simplify eq. (14) after replacing integration over ω with integration over k and keeping only the first two orders in α , arriving at the following expression:

$$\mathcal{T} = -\frac{\hbar D}{A} (1 + i\beta) (\mathbf{v} \cdot \mathbf{j}). \quad (15)$$

Here $\mathbf{j} = \mathcal{J}_1 \partial T/T$ coincides with the expression for the magnon current calculated within the relaxation time approximation where $\mathcal{J}_n = - \int d^d\mathbf{k} / (2\pi)^d \tau(\varepsilon) \varepsilon^n v_x^2 \partial f_0 / \partial \varepsilon$ and $\tau(\varepsilon) = (2\alpha\omega)^{-1}$ account for the non-equilibrium distribution correction $\delta f = \tau\varepsilon(\partial f / \partial \varepsilon) v_\alpha (\partial_\alpha T/T)$ arising in the Boltzmann equation [34]. Here we also use the spectrum of magnons $\varepsilon(\mathbf{k}) = \hbar(Ak^2 + Ak_0^2 + H)/s$, the velocity $v_x = \partial\omega_k / \partial k_x$, and the Bose-Einstein equilibrium distribution $f_0 = \{\exp[\varepsilon/k_B T] - 1\}^{-1}$. Two terms in eq. (15) result in two torque components that are perpendicular to each other. Thus, the first term corresponds to the reactive torque and the second term corresponds to the dissipative torque, with the ratio between them given by the parameter β . As these corrections arise due to magnon spin dephasing, we expect that the parameter β should coincide with the one obtained in the context of magnonic torques in textured ferromagnets [15]. We can also confirm this by inspecting eq. (14) which results in the expression $\beta/\alpha = (d/2)B(x)/F_1(x) \sim d/2$, where $F_n(x) = \int_0^\infty d\epsilon \epsilon^{d/2} (\epsilon + x)^{n-1} e^{\epsilon+x} / (e^{\epsilon+x} - 1)^2$ and $B(x) = \int_0^\infty d\epsilon (\epsilon + x) \epsilon^{d/2-1} e^{\epsilon+x} / (e^{\epsilon+x} - 1)^2$ evaluated at the magnon gap $x = \hbar\omega_0/k_B T$, where $d = 2$ or 3 . We also recall that the magnon current density is given by [6]

$$j_\alpha = k_B \partial_\alpha T F_1 / (6\pi^2 \lambda \hbar \alpha), \quad (16)$$

where $d = 3$ and $\lambda = \sqrt{\hbar A/(sk_B T)}$ is the thermal magnon wavelength. In the case of $d = 2$ we obtain $j_\alpha = k_B \partial_\alpha T F_1 / (4\pi \hbar \alpha)$.

We express the result in eq. (15) in the form of the LLG equation which constitutes the main result of this section:

$$\mathfrak{s}(1 + \alpha^s \mathbf{m}_s \times) \dot{\mathbf{m}}_s + \mathbf{m}_s \times \mathbf{H}_{\text{eff}}^s = (\eta + \vartheta \mathbf{m}_s \times) j_\alpha \mathcal{D}_\alpha \mathbf{m}_s, \quad (17)$$

where $\eta = \hbar$, $\vartheta = \eta\beta$ [35], $\mathfrak{s} = |\langle \mathbf{m} \rangle| s$ is the renormalized spin density, $\mathbf{H}_{\text{eff}}^s = -(\mathfrak{s}/s) \langle \delta_{\mathbf{m}} F \rangle$ is the effective field, $\alpha^s = (\mathfrak{s}/s)\alpha$ is the renormalized Gilbert damping (in other sections we always skip the index s).

Critical current instability and switching. – The LLG equation (17) describes the magnon current-induced magnetic instabilities and magnetization switching. We can estimate the corresponding critical current and required temperature gradient by employing a simple stability analysis of the linearized LLG equation after transforming it to the Landau-Lifshitz form, *i.e.*, by multiplying eq. (17) by $(1 - \alpha \mathbf{m} \times)$. We consider the case in fig. 1 where a time-independent magnon current is $\mathbf{j} = j \mathbf{x}$, and the effective field is given as $\mathbf{H}_{\text{eff}}^s = H \mathbf{y} + K m_z \mathbf{z}$, where H is the strength of the external magnetic field and K is the easy-plane magnetic anisotropy, *e.g.*, corresponding to the shape anisotropy. When the temperature is uniform ($j = 0$) at equilibrium, the fixed point solution is $\mathbf{m}_s = -\mathbf{y}$. This solution becomes unstable when j reaches

$$j_c = \frac{A}{D} \frac{H + K/2}{|\vartheta/\alpha - \eta|}. \quad (18)$$

We assume that $H^2 + HK \geq 4K^2\alpha^2$ and $\vartheta > \eta\alpha$ (for $\vartheta < \eta\alpha$ eq. (18) gives only the upper bound for the critical current). For $\vartheta \sim \eta\beta$ one can see that our estimate of the critical current becomes large. When $\vartheta \sim \eta$ and $\alpha \ll 1$, we obtain a more favorable estimate $j_c \approx (H + K/2) \alpha A / \hbar D$ which will be used for our numerical estimates. The reason is that for a general form of DMI corresponding to pyrochlore crystals [16] or chiral magnets [36] we expect other mechanisms, not necessarily relying on the dissipative β -type correction [15], to contribute to the dissipative torque [35] by analogy with the current-induced torques in bilayers [21]. For estimates, we consider Cu_2OSeO_3 thin insulating layer in ferromagnetic phase. By taking the lattice spacing $a = 0.5 \text{ nm}$, $T = 50 \text{ K}$, $\alpha = 0.01$, $\mathfrak{s} = 0.5\hbar/a^3$, $A/(a^2 k_B) = 50 \text{ K}$, $K = 4\pi M_s$ and $D/(ak_B) = 3 \text{ K}$ we obtain $A/D \approx 80 \text{ nm}$ [36]. A temperature gradient comparable to experimentally accessible $\partial_\alpha T = 1 \text{ K}/\mu\text{m}$ [37] should be sufficient for the instability according to eq. (16).

Magnon pumping by magnetization dynamics. – The dissipative terms in eq. (3) lead to pumping of magnons by magnetization dynamics. This effect is analogous to charge pumping by a combination of the spin pumping and the inverse spin Hall effect in bilayers [38] where the magnetization dynamics is induced by external microwave fields. Under a simple circular precession of the magnetization the time-averaged magnon current is given by $j_{\text{pump}} = \sin^2 \theta \omega \vartheta / \varrho$ where θ is the cone angle of the magnetization dynamics. We can also define the corresponding spin current as $j_s = \hbar j_{\text{pump}}$. For

numerical estimates we use parameters characteristic to Pt/Co/AlO_x or Pt/CoFe/MgO thin films which we treat as two-dimensional magnets [39]. We use $A = 1.6 \times 10^{-11}$ J/m, $M_s = 8.3 \times 10^5$ A/m, $t = 0.6 \times 10^{-9}$ nm and $D = 4 \times 10^{-3}$ J/m² arriving at $j_s = \sin^2 \theta \hbar \omega D \vartheta / (4\pi A t \eta \alpha)$ which is reminiscent of the expression for spin pumping $j_s = \sin^2 \theta \hbar \omega g_{\uparrow\downarrow} / (4\pi)$ with $g_{\uparrow\downarrow} = D \vartheta / (A t \eta \alpha)$. Compared to spin pumping we recover an order of magnitude smaller spin current under equal pumping conditions.

We suppose that the power dissipated by magnetization dynamics, $P = \sin^2 \theta \omega^2 \alpha \mathfrak{s} V$, is equally divided between the cooled and heated reservoirs [9]. From eq. (3) the maximum cooling then corresponds to the regime in which the heat current carried by magnons from the cooled reservoir is exactly compensated by dissipation:

$$\frac{\Delta T}{TL} = \frac{\omega \Pi \vartheta / \varrho - \omega^2 \alpha \mathfrak{s} L / 2}{\Pi^2 / \varrho + \kappa T} \sin^2 \theta, \quad (19)$$

where κ could also include the thermal conductivity of phonons, L is the length of the magnet, and we assume that heat flows couple to magnetization only via magnon currents. The maximum is reached for $\omega = \Pi \vartheta / (\varrho \alpha \mathfrak{s} L)$. By analogy with the thermoelectric figure of merit we can define the figure of merit for magnonic cooling as $ZT = 2\Delta T_{max}/T$ which leads to the following expression:

$$\frac{ZT}{\sin^2 \theta} = \frac{(\Pi \vartheta / \varrho)^2 / (\alpha \mathfrak{s})}{\Pi^2 / \varrho + \kappa T} = \frac{D^2 (\vartheta / \eta)^2 \hbar}{6\pi^2 \mathfrak{s} A^2 \lambda \alpha^2} \frac{F_1^2}{F_2}, \quad (20)$$

where we assume sufficiently low temperature so that the effect of phonons can be disregarded, *e.g.*, at 3 K the thermal conductivity of magnons can become comparable to the thermal conductivity of phonons [17], $\Pi / \varrho = -\mathcal{J}_1$ and $\Pi^2 / \varrho + \kappa T = \mathcal{J}_2$ within the relaxation time approximation applied to thermal magnons, and $F_1^2 / F_2 \sim 1$ (for $d = 2$ we obtain $Z = D^2 (\vartheta / \eta)^2 \hbar F_1^2 / (4\pi \mathfrak{s} A^2 \alpha^2 F_2)$, where \mathfrak{s} is the spin surface density). For Pt/Co/AlO_x or Pt/CoFe/MgO thin films we obtain $ZT \sim 0.001$. The absolute cooling of the cold reservoir is relatively weak which is a consequence of the large dissipated power P . However, the relative temperature difference between reservoirs found without P can be large at typical ferromagnetic resonance frequencies, *i.e.*, $\Delta T / T \sim 0.05$, and it should be measurable.

Generalizations to arbitrary DMI. – The magnon spin torque in eq. (17) can be obtained from results in ref. [15] (see eq. (13)) by replacing the texture derivative ∂_i with a chiral derivative $\mathcal{D}_\alpha = \partial_\alpha + (D/A)(\mathbf{n} \times \mathbf{e}_\alpha) \times$ [21,30]. In general, such procedure does not guarantee the correct values for ϑ and η . Based on the phenomenological symmetry-based argument, we can describe the magnon spin torque and magnon pumping in eqs. (3) and (4) for the most general form of DMI by substituting the chiral derivative, $\mathcal{D}_\alpha = \partial_\alpha + (\mathbf{D}_\alpha / A) \times$, *i.e.*, for the reactive and dissipative torques we obtain

$$\mathcal{T} = (1/A) (\eta + \vartheta \mathbf{m} \times) j_\alpha \mathbf{D}_\alpha \times \mathbf{m}. \quad (21)$$

Here we sum over repeated indices, and the tensor $D_{\alpha\beta} = \mathbf{D}_\alpha \cdot \mathbf{e}_\beta$ describes the most general form of DMI, $F_D = D_{\alpha\beta} \varepsilon_{\beta\delta\gamma} m_\delta \partial_\alpha m_\gamma$, leading to the exchange contribution in the free energy, $F_{ex} = (A/2)(\mathcal{D}_\alpha \mathbf{m})^2$. By separating $D_{\alpha\beta}$ into symmetric and antisymmetric parts, $D_{\alpha\beta} = D_{\alpha\beta}^{sym} + \varepsilon_{\alpha\beta\gamma} D_\gamma^{ant}$, we identify $\mathbf{D}^{ant} = D \mathbf{n}$ for DMI due to structural asymmetry in the figure. The contribution $D_{\alpha\beta}^{sym} = D \delta_{\alpha\beta}$ arises in non-centrosymmetric crystals, *e.g.*, in Cu₂OSeO₃, resulting in $\mathcal{T} = (D/A) (\eta + \vartheta \mathbf{m} \times) \mathbf{j} \times \mathbf{m}$.

Conclusions. – We developed a phenomenological description for the interplay between magnetization dynamics and magnon currents in ferromagnets with DMI. Our theory describes: i) magnon current-induced instability and switching; ii) pure spin current pumping; and iii) cooling effects in single-domain magnets. The strength of all mentioned effects is related to the magnitude of the dissipative torque which is weakened by the Gilbert damping factor compared to the reactive torque for the simple form of DMI considered here. The general form of DMI, $\mathbf{D}_{ij} \cdot (\mathbf{S}_i \times \mathbf{S}_j)$, should in principle result in a situation where the dissipative and reactive torques are comparable [35] in analogy to spin-orbit torques in metal/ferromagnet bilayers. We thus expect magnonic dissipative torques and spin pumping in such systems as pyrochlore crystals (*e.g.*, Lu₂V₂O₇) and chiral magnets (*e.g.*, MnSi or Cu₂OSeO₃). The proposed pumping mechanism could potentially be useful for electronics relying on pure spin currents. Our results agree with ref. [24] where only the reactive thermal torque has been discussed in detail.

We are grateful to K. BELASHCHENKO and C. BINEK for discussions. This work was supported in part by the NSF under Grants Nos. Phy-1415600, No. DMR-1420645, and NSF-EPSCoR 1004094, and performed in part at the Central Facilities of the Nebraska Center for Materials and Nanoscience, supported by the Nebraska Research Initiative.

REFERENCES

- [1] GOENNENWEIN S. T. B. and BAUER G. E. W., *Nat. Nanotechnol.*, **7** (2012) 145; BAUER G. E. W., SAITO E. and VAN WEES B. J., *Nat. Mater.*, **11** (2012) 391.
- [2] UCHIDA K., TAKAHASHI S., HARII K., IEDA J., KOSHIBAE W., ANDO K., MAEKAWA S. and SAITO E., *Nature*, **455** (2008) 778; UCHIDA K., XIAO J., ADACHI H., OHE J., TAKAHASHI S., IEDA J., OTA T., KAJIWARA Y., UMEZAWA H., KAWAI H., BAUER G. E. W., MAEKAWA S. and SAITO E., *Nat. Mater.*, **9** (2010) 894; JAWORSKI C. M., YANG J., MACK S., AWSCHALOM D. D., HEREMANS J. P. and MYERS R. C., *Nat. Mater.*, **9** (2010) 898; XIAO J., BAUER G. E. W., UCHIDA K.-C., SAITO E. and MAEKAWA S., *Phys. Rev. B*, **81** (2010) 214418; ADACHI H., UCHIDA K.-I., SAITO E. and MAEKAWA S., *Rep. Prog. Phys.*, **76** (2013) 036501.

[3] SLACHTER A., BAKKER F. L., ADAM J.-P. and VAN WEES B. J., *Nat. Phys.*, **6** (2010) 879.

[4] DUBI Y. and DI VENTRA M., *Phys. Rev. B*, **79** (2009) 081302; ŚWIRKOWICZ R., WIERZBICKI M. and BARNAS J., *Phys. Rev. B*, **80** (2009) 195409; FLIPSE J., BAKKER F. L., SLACHTER A., DEJENE F. K. and VAN WEES B. J., *Nat. Nanotechnol.*, **7** (2012) 166.

[5] HINZKE D. and NOWAK U., *Phys. Rev. Lett.*, **107** (2011) 027205; YAN P., WANG X. S. and WANG X. R., *Phys. Rev. Lett.*, **107** (2011) 177207; TORREJON J., MALINOWSKI G., PELLOUX M., WEIL R., THIAVILLE A., CURIALE J., LACOUR D., MONTAIGNE F. and HEHN M., *Phys. Rev. Lett.*, **109** (2012) 106601; JIANG W., UPADHYAYA P., FAN Y., ZHAO J., WANG M., CHANG L.-T., LANG M., WONG K. L., LEWIS M., LIN Y.-T., TANG J., CHEREPOV S., ZHOU X., TSERKOVNYAK Y., SCHWARTZ R. N. and WANG K. L., *Phys. Rev. Lett.*, **110** (2013) 177202.

[6] KOVALEV A. A. and TSERKOVNYAK Y., *EPL*, **97** (2012) 67002.

[7] VOLOVIK G. E., *J. Phys. C: Solid State Phys.*, **20** (1987) L83; BARNES S. E. and MAEKAWA S., *Phys. Rev. Lett.*, **98** (2007) 246601; KIM K.-W., MOON J.-H., LEE K.-J. and LEE H.-W., *Phys. Rev. Lett.*, **108** (2012) 217202.

[8] HATAMI M., BAUER G. E. W., ZHANG Q. and KELLY P. J., *Phys. Rev. Lett.*, **99** (2007) 066603; BAUER G. E. W., BRETZEL S., BRATAAS A. and TSERKOVNYAK Y., *Phys. Rev. B*, **81** (2010) 024427; CAHAYA A. B., TRETIAKOV O. A. and BAUER G. E. W., *Appl. Phys. Lett.*, **104** (2014) 042402.

[9] KOVALEV A. A. and TSERKOVNYAK Y., *Solid State Commun.*, **150** (2010) 500.

[10] KOVALEV A. A. and TSERKOVNYAK Y., *Phys. Rev. B*, **80** (2009) 100408; BAUER G. E. W., BRETZEL S., BRATAAS A. and TSERKOVNYAK Y., *Phys. Rev. B*, **81** (2010) 024427.

[11] JOHNSON M. and SILSBEY R. H., *Phys. Rev. Lett.*, **55** (1985) 1790; JEDEMA F. J., FILIP A. T. and VAN WEES B. J., *Nature*, **410** (2001) 345; ŽUTIĆ I., FABIAN J. and DAS SARMA S., *Rev. Mod. Phys.*, **76** (2004) 323; TSERKOVNYAK Y., BRATAAS A. and BAUER G. E., *Phys. Rev. Lett.*, **88** (2002) 117601; DYAKONOV M. I. and PEREL V. I., *Phys. Lett. A*, **35** (1971) 459.

[12] KAJIWARA Y., HARII K., TAKAHASHI S., OHE J., UCHIDA K., MIZUGUCHI M., UMEZAWA H., KAWAI H., ANDO K., TAKANASHI K., MAEKAWA S. and SAITO E., *Nature*, **464** (2010) 262.

[13] KHITUN A. and WANG K. L., *J. Appl. Phys.*, **110** (2011) 034306.

[14] KONG L. and ZANG J., *Phys. Rev. Lett.*, **111** (2013) 067203; LIN S.-Z., BATISTA C. D., REICHARDT C. and SAXENA A., *Phys. Rev. Lett.*, **112** (2014) 187203.

[15] KOVALEV A. A., *Phys. Rev. B*, **89** (2014) 241101.

[16] ONOSE Y., IDEUE T., KATSURA H., SHIOMI Y., NAGAOSA N. and TOKURA Y., *Science*, **329** (2010) 297; MATSUMOTO R. and MURAKAMI S., *Phys. Rev. Lett.*, **106** (2011) 197202; MOOK A., HENK J. and MERTIG I., *Phys. Rev. B*, **90** (2014) 024412.

[17] DOUGLASS R. L., *Phys. Rev.*, **129** (1963) 1132; DEMOKRITOV S. O., DEMIDOV V. E., DZYAPKO O., MELKOV G. A., SERGA A. A., HILLEBRANDS B. and SLAVIN A. N., *Nature*, **443** (2006) 430; UCHIDA K.-I., ADACHI H., OTA T., NAKAYAMA H., MAEKAWA S. and SAITO E., *Appl. Phys. Lett.*, **97** (2010) 172505.

[18] FANG D., KUREBAYASHI H., WUNDERLICH J., VÝBORNÝ K., ZÁRBO L. P., CAMPION R. P., CASIRAGHI A., GALLAGHER B. L., JUNGWIRTH T. and FERGUSON A. J., *Nat. Nanotechnol.*, **6** (2011) 413.

[19] MIHAI MIRON I., GAUDIN G., AUFFRET S., RODMACQ B., SCHUHL A., PIZZINI S., VOGEL J. and GAMBARDELLA P., *Nat. Mater.*, **9** (2010) 230; LIU L., PAI C.-F., LI Y., TSENG H. W., RALPH D. C. and BUHRMAN R. A., *Science*, **336** (2012) 555; CHERNYSHOV A., OVERBY M., LIU X., FURDYNA J. K., LYANDA-GELLER Y. and ROKHINSON L. P., *Nat. Phys.*, **5** (2009) 656.

[20] LIU L., MORIYAMA T., RALPH D. and BUHRMAN R., *Phys. Rev. Lett.*, **106** (2011) 036601.

[21] TSERKOVNYAK Y. and BENDER S. A., *Phys. Rev. B*, **90** (2014) 014428.

[22] COSTA A. T., MUNIZ R. B., LOUNIS S., KLAUTAU A. B. and MILLS D. L., *Phys. Rev. B*, **82** (2010) 014428.

[23] MOON J.-H., SEO S.-M., LEE K.-J., KIM K.-W., RYU J., LEE H.-W., MCMICHAEL R. D. and STILES M. D., *Phys. Rev. B*, **88** (2013) 184404.

[24] MANCHON A., NDIAYE P. B., MOON J.-H., LEE H.-W. and LEE K.-J., *Phys. Rev. B*, **90** (2014) 224403.

[25] DZYALOSHINSKY I., *J. Phys. Chem. Solids*, **4** (1958) 241; MORIYA T., *Phys. Rev.*, **120** (1960) 91.

[26] CRÉPIEUX A. and LACROIX C., *J. Magn. & Magn. Mater.*, **182** (1998) 341; THIAVILLE A., ROHART S., JUÉ É., CROS V. and FERT A., *EPL*, **100** (2012) 57002.

[27] LANDAU L. and LIFSHITZ E., *Statistical Physics*, Vol. **5** (Pergamon, Oxford) 1980.

[28] DEMOKRITOV S. O., DEMIDOV V. E., DZYAPKO O., MELKOV G. A., SERGA A. A., HILLEBRANDS B. and SLAVIN A. N., *Nature*, **443** (2006) 430.

[29] LANDAU L. and LIFSHITZ E., *Electrodynamics of Continuous Media*, Vol. **8**, 2nd edition (Pergamon, Oxford) 1984.

[30] KIM K.-W., LEE H.-W., LEE K.-J. and STILES M. D., *Phys. Rev. Lett.*, **111** (2013) 216601.

[31] DUGAEV V. K., BRUNO P., CANALS B. and LACROIX C., *Phys. Rev. B*, **72** (2005) 024456.

[32] BROWN W. F., *Phys. Rev.*, **130** (1963) 1677.

[33] HOFFMAN S., SATO K. and TSERKOVNYAK Y., *Phys. Rev. B*, **88** (2013) 064408.

[34] ASHCROFT N. W. and MERMIN N. D., *Solid State Physics* (Cengage Learning) 1976.

[35] In our discussion the strength of the dissipative torque is proportional to the dephasing parameter β while it does not have to be so for a general magnonic band with non-trivial Berry curvature.

[36] SEKI S., YU X. Z., ISHIWATA S. and TOKURA Y., *Science*, **336** (2012) 198.

[37] BRANDL F. and GRUNDLER D., *Appl. Phys. Lett.*, **104** (2014) 172401.

[38] MOSENDZ O., VLAMINCK V., PEARSON J. E., FRADIN F. Y., BAUER G. E. W., BADER S. D. and HOFFMANN A., *Phys. Rev. B*, **82** (2010) 214403.

[39] MIROU M., MOORE T., SZAMBOLICS H., BUDA-PREJBEANU L. D., AUFFRET S., RODMACQ B., PIZZINI S., VOGEL J., BONFIM M., SCHUHL A. and GAUDIN G., *Nat. Mater.*, **10** (2011) 419.