

Lump solutions to the Kadomtsev–Petviashvili I equation with a self-consistent source

Xuelin Yong^{a,b,*}, Wen-Xiu Ma^{b,c,d,e}, Yehui Huang^a, Yong Liu^a

^a School of Mathematical Sciences and Physics, North China Electric Power University, Beijing 102206, China

^b Department of Mathematics and Statistics, University of South Florida, Tampa, FL 33620, USA

^c Department of Mathematics, Zhejiang Normal University, Jinhua 321004, Zhejiang, China

^d College of Mathematics and Systems Science, Shandong University of Science and Technology, Qingdao 266590, Shandong, China

^e Department of Mathematical Sciences, North-West University, Mafikeng Campus, Private Bag X 2046, Mmabatho 2735, South Africa

ARTICLE INFO

Article history:

Received 19 October 2017

Received in revised form 19 January 2018

Accepted 4 February 2018

Available online 2 March 2018

Keywords:

KPI equation with a self-consistent source

Hirota bilinear method

Lump solution

ABSTRACT

Based on symbolic computations, lump solutions to the Kadomtsev–Petviashvili I (KPI) equation with a self-consistent source (KPIESCS) are constructed by using the Hirota bilinear method and an ansatz technique. In contrast with lower-order lump solutions of the Kadomtsev–Petviashvili (KP) equation, the presented lump solutions to the KPIESCS exhibit more diverse nonlinear phenomena. The method used here is more natural and simpler.

© 2018 Elsevier Ltd. All rights reserved.

1. Introduction

Soliton equations with self-consistent sources are role models in many fields of physics, and important developments have been made in exploring their soliton phenomena [1–20]. The Kadomtsev–Petviashvili (KP) equation with a self-consistent source arose in describing the interaction of long and short waves and its soliton solutions were first found by Mel'nikov [21,22]. Later, N -soliton solutions of the KP equation with self-consistent sources were obtained through the Hirota method [23] and the generalized binary Darboux transformation [24]. Furthermore, the general high-order rogue waves which are given in terms of determinants whose matrix elements have plain algebraic expressions of the KP equation with self-consistent sources were derived via the Hirota method [25].

In contrast to soliton solutions, lump solutions are another kind of important exact solutions, which are rational analytical and localized in all directions in the space. General rational function solutions were presented for the Korteweg–de Vries (KdV) equation, the Boussinesq equation, the nonlinear Schrödinger (NLS) equation and the Toda lattice equation systematically through the Wronskian and Casoratian determinant techniques for integrable equations [26–30]. Special examples of lump solutions are also found for many integrable equations such as the Kadomtsev–Petviashvili I (KPI) equation [31,32], the three-dimensional three-wave resonant interaction [33], the B-Kadomtsev–Petviashvili (BKP) equation [34], the Davey–Stewartson-II equation [35,36] and the Ishimori-I equation [37].

In mathematical physics, the KP equation is usually written as

$$(u_t + 6uu_x + u_{xxx})_x - \sigma u_{yy} = 0, \sigma = \pm 1, \quad (1)$$

which is classified as the KPI equation when $\sigma = 1$ and the KP-II equation when $\sigma = -1$. Recently, one of the authors (Ma) has proposed a symbolic computation method to search for positive quadratic function solutions, particularly for the

* Corresponding author at: School of Mathematical Sciences and Physics, North China Electric Power University, Beijing 102206, China.

E-mail address: yongxuelin@ncepu.edu.cn (X. Yong).

(2 + 1)-dimensional bilinear KPI equation [38]. The obtained quadratic function solutions in the KPI case [38] contain a set of six free parameters, and taking special choices of the involved parameters covers a particular class of lump solutions generated from computing long wave limits of soliton solutions. The same idea was also adopted to derive lumps and interaction solutions between lumps and kinks for several equations such as KPI equation [39,40], BKP equation [41,42], (2 + 1)-dimensional Ito equation and (2 + 1)-dimensional Caudrey–Dodd–Gibbon–Sawada–Kotera (CDGKS) equation [43], (2 + 1)-dimensional Sawada–Kotera equation [44], dimensionally reduced gKP and gBKP equations [45], (3 + 1)-dimensional Jimbo–Miwa equation [46,47]. Especially, in the case of the (2 + 1)-dimensional Ito equation, an interesting characteristic that an arbitrary function is involved in the resulting interaction solutions was explored [48]. It should be noticed that the Hirota bilinear method plays an important role in these works and this technique has the advantage of being applicable directly upon the equations [49–51].

In this paper, we would like to discuss the KPI equation with a self-consistent source (KPIESCS)

$$\begin{cases} (u_t + 6uu_x + u_{xxx} + 8|\Phi|_x^2)_x - 3u_{yy} = 0, \\ i\Phi_y = \Phi_{xx} + u\Phi, \end{cases} \quad (2)$$

and want to determine its lump solutions and their dynamics. We begin with the Hirota bilinear form of the KPIESCS and make some ansatz by combination functions of quadratic functions to solve the corresponding bilinear counterpart equation. The resulting solutions exhibit more diverse nonlinear phenomena than lower-order lumps solutions to the KPI equation and provide supplements to the existing solutions in the literature. A few concluding remarks will be given in the last section.

2. Lump solutions to KPIESCS

Through the dependent variable transformation $u = 2(\ln F)_{xx}$ and $\Phi = G/F$, the KPIESCS is rewritten as

$$\begin{cases} \left[\frac{2(F_{xt}F - F_xF_t + F_{xxxx}F - 4F_{xxx}F_x + 3F_{xx}^2 - 3F_{yy}F + 3F_y^2) + 8GG^*}{F^2} \right]_{xx} = 0, \\ i(G_yF - GF_y) - G_{xx}F + 2G_xF_x - GF_{xx} = 0, \end{cases} \quad (3)$$

from which we obtain its bilinear equation

$$\begin{cases} 2(F_{xt}F - F_xF_t + F_{xxxx}F - 4F_{xxx}F_x + 3F_{xx}^2 - 3F_{yy}F + 3F_y^2) + 8GG^* = cF^2, \\ i(G_yF - GF_y) - G_{xx}F + 2G_xF_x - GF_{xx} = 0, \end{cases} \quad (4)$$

where c is a constant of integration. Here asterisk G^* means the complex conjugation and D is the famous Hirota bilinear operator [26]. Eq. (4), with $c = 8$, may also be written concisely in terms of the D -operators as

$$\begin{cases} (D_xD_t + D_x^4 - 3D_y^2)F \cdot F + 8(GG^* - F^2) = 0, \\ (iD_y - D_x^2)G \cdot F = 0. \end{cases} \quad (5)$$

To search for lump solutions to the above bilinear equation, putting $G = G_R + iG_I$ and supposing

$$\begin{cases} F = 1 + \xi_1^2 + \xi_2^2, \\ G_R = b_0 + b_1\xi_1 + b_2\xi_2 + b_3\xi_1^2 + b_4\xi_2^2, \\ G_I = c_0 + c_1\xi_1 + c_2\xi_2 + c_3\xi_1^2 + c_4\xi_2^2, \end{cases} \quad (6)$$

with

$$\begin{cases} \xi_1 = a_1x + a_2y + a_3t + a_4, \\ \xi_2 = a_5x + a_6y + a_7t + a_8, \end{cases} \quad (7)$$

where the parameters $a_i (1 \leq i \leq 8)$, b_j , $c_j (0 \leq j \leq 4)$ are all real constants to be determined. It is easy to see that the assumption for F here guarantees analyticity and rational localization of solutions. Substituting Eqs. (6) and (7) into Eq. (5) and equating all the coefficients of different polynomials of x, y, t to zero, we obtain a set of algebraic equations on the undetermined parameters. After careful discussions, we can generate the following set of constraining equations for the parameters

$$\begin{aligned} \{a_3 = \frac{(a_2^2 - a_1^4)[3(a_1^4 + a_2^2)^2 + 16a_1^4]}{a_1(a_1^4 + a_2^2)^2}, a_5 = 0, a_6 = a_1^2, \\ a_7 = \frac{2a_1a_2[3(a_1^4 + a_2^2)^2 - 16a_1^4]}{(a_1^4 + a_2^2)^2}, b_0 = \frac{b_3(a_2^2 - 3a_1^4)}{a_1^4 + a_2^2}, \\ b_1 = kc_1, b_2 = kc_2, b_4 = b_3, c_0 = -kb_0, c_1 = -\frac{4a_1^2a_2b_3}{a_1^4 + a_2^2}, \end{aligned} \quad (8)$$




Fig. 1. (Color online) Plots of lump solution for u with $a_1 = 1, a_2 = -2, a_4 = -1, a_8 = 0, b_3 = \frac{3}{5}, k = -\frac{4}{3}$ at $t = 0$.

$$c_2 = \frac{4a_1^4 b_3}{a_1^4 + a_2^2}, c_3 = -kb_3, c_4 = c_3\},$$

where k is an auxiliary arbitrary real constant which needs to satisfy the following condition

$$b_3^2(1 + k^2) = 1, \quad (9)$$

and the other parameters not expressed in the set are arbitrary, provided that all corresponding formulas are well defined. Therefore, the non-zero determinant condition discussed before in [38] now becomes $a_1 a_6 - a_2 a_5 = a_1^3 \neq 0$, which guarantees the localization of the associated solutions.

Actually, it is readily observed that at any given time t , the above lump solution $u \rightarrow 0$ when $x^2 + y^2 \rightarrow \infty$. We can also get all critical points of the function F at a fixed time t by solving the first-order derivative equations $(F_x, F_y) = 0$ to obtain

$$\begin{cases} x(t) = \frac{3(a_1^4 + a_2^2)^3 + 16(a_1^8 - 3a_1^4 a_2^2)}{a_1^2(a_1^4 + a_2^2)^2} t - \frac{a_1^2 a_4 - a_2 a_8}{a_1^3}, \\ y(t) = \frac{2a_2[16a_1^4 - 3(a_1^4 + a_2^2)^2]}{a_1(a_1^4 + a_2^2)^2} t - \frac{a_8}{a_1^2}, \end{cases} \quad (10)$$

and they are also critical points in the (x, y) -plane for the function $u = 2(\ln F)_{xx}$. Moreover, based on the second partial derivative test, we can know that the lump solution u has a peak at the critical point $(x(t), y(t))$ since

$$u_{xx} = -24a_1^4 < 0, u_{xx}u_{yy} - u_{xy}^2 = 192a_1^{10} > 0. \quad (11)$$

However, with respect to the consistent-source term $|\Phi|^2 = |G|^2/F^2 = (G_R^2 + G_I^2)/F^2$, it decays into $b_3^2 + c_3^2 = 1$ for $|x|, |y| \rightarrow \infty$. Its critical points do not always coincide with those functions F and G . For illustration, two special pairs of the lump solutions with specific values of the parameters are given as follows.

Firstly, if we take a special choice for the parameters

$$a_1 = 1, a_2 = -2, a_4 = -1, a_8 = 0, b_3 = \frac{3}{5}, k = -\frac{4}{3}, \quad (12)$$

we get the lump solution which was exactly presented in Ref. [24] as $\mu = 1/2, \nu = 1$. The corresponding three dimensional plots and density plots when $t = 0$ are shown in Figs. 1 and 2, respectively. The lump solutions are all centered at $(1, 0)$, but the source term $|\Phi|^2$ owns two maximal points and one minimal point as shown in the figures. Meanwhile, we can see a twist in Fig. 2(b) for the source term.

Secondly, the plots for another selection of the parameters

$$a_1 = 1, a_2 = -1, a_4 = -1, a_8 = 0, b_3 = 1, k = 0, \quad (13)$$

when $t = 1$ are depicted in Figs. 3 and 4, respectively. It can be seen that the potential u is similar, but the source term $|\Phi|^2$ exhibits different and diverse dynamics from the first case. It owns two peak points and two minimal points compared with Fig. 2.




Fig. 2. (Color online) Plots of lump solution for $|\Phi|^2$ with $a_1 = 1$, $a_2 = -2$, $a_4 = -1$, $a_8 = 0$, $b_3 = \frac{3}{5}$, $k = -\frac{4}{3}$ at $t = 0$.




Fig. 3. (Color online) Plots of lump solution for u with $a_1 = 1$, $a_2 = -1$, $a_4 = -1$, $a_8 = 0$, $b_3 = 1$, $k = 0$ at $t = 1$.

3. Conclusion

In this paper, the lump solutions to the KPI equation with a self-consistent source (KPIESCS) were presented by using the direct ansatz and the Hirota bilinear method. It is shown that the direct ansatz is a powerful means for seeking lump solutions to nonlinear equations, when it is combined with the Hirota bilinear method. We can see that the solutions obtained are consistent with those presented before, and the approach used here is more natural and simpler. It is hoped that the presented results would be helpful in understanding the propagation processes of nonlinear waves, particularly in fluid mechanics. Recently, interaction solutions between lumps and solitons have attracted a lot of attention and various kinds of interaction solutions have been obtained [52]. It will be interesting to discuss the existence conditions and solution methods for getting new interaction solutions to nonlinear wave equations. Those problems are left for future research publications.

Acknowledgments

This work is supported in part by the 13th Five Year National Key Research and Development Program of China with Grant No. 2016YFC0401406, the NSF of China with Grant Nos. 11301179, 11371326, 11271008, the Fundamental Research Funds of




Fig. 4. (Color online) Plots of lump solution for $|\Phi|^2$ with $a_1 = 1$, $a_2 = -1$, $a_4 = -1$, $a_8 = 0$, $b_3 = 1$, $k = 0$ at $t = 1$.

the Central Universities with Grant Nos. 2015MS56, 2016MS63, the State Scholarship Fund of China, the 111 Project of China (B16002), the NSF under the grant DMS-1664561, Natural Science Fund for Colleges and Universities of Jiangsu Province under the grant 17KJB110020, and the Distinguished Professorships at Shanghai University of Electric Power and Shanghai Second Polytechnic University. The authors would like to thank the reviewers for their comments to improve the paper.

References

- [1] V.K. Mel'nikov, Integration method of the Korteweg–de Vries equation with a self-consistent source, *Phys. Lett. A* 133 (1988) 493–496.
- [2] V.K. Mel'nikov, Capture and confinement of solitons in nonlinear integrable systems, *Comm. Math. Phys.* 120 (1989) 451–468.
- [3] V.K. Mel'nikov, New method for deriving nonlinear integrable systems, *J. Math. Phys.* 31 (1990) 1106–1113.
- [4] J. Leon, Nonlinear evolutions with singular dispersion laws and forced systems, *Phys. Lett. A* 144 (1990) 444–452.
- [5] V.K. Mel'nikov, Integration of the nonlinear Schrödinger equation with a self-consistent source, *Comm. Math. Phys.* 137 (1991) 359–381.
- [6] Y.B. Zeng, W.X. Ma, R.L. Lin, Integration of the soliton hierarchy with self-consistent sources, *J. Math. Phys.* 41 (2001) 5453–5489.
- [7] R.L. Lin, Y.B. Zeng, W.X. Ma, Solving the KdV hierarchy with self-consistent sources by inverse scattering method, *Physica A* 291 (2001) 287–298.
- [8] Y.B. Zeng, W.X. Ma, Y.J. Shao, Two binary Darboux transformations for the KdV hierarchy with self-consistent sources, *J. Math. Phys.* 42 (2001) 2113–2128.
- [9] Y.B. Zeng, Y.J. Shao, W.X. Ma, Integral-type Darboux transformations for the mKdV hierarchy with self-consistent sources, *Commun. Theor. Phys.* 38 (2002) 641–648.
- [10] D.J. Zhang, The N-soliton solutions for the modified KdV equation with self-consistent sources, *J. Phys. Soc. Japan* 71 (2002) 2649–2656.
- [11] W.X. Ma, Soliton, positon and negaton solutions to a Schrödinger self-consistent source equation, *J. Phys. Soc. Japan* 72 (2003) 3017–3019.
- [12] W.X. Ma, Complexiton solutions of the Korteweg–de Vries equation with self-consistent sources, *Chaos Solitons Fractals* 26 (2005) 1453–1458.
- [13] X.B. Hu, H.Y. Wang, Construction of dKP and BKP equations with self-consistent sources, *Inverse Problems* 22 (2006) 1903.
- [14] X.B. Hu, H.Y. Wang, New type of Kadomtsev–Petviashvili equation with self-consistent sources and its bilinear Bäcklund transformation, *Inverse Problems* 23 (2007) 1433.
- [15] Y.Q. Yao, Y.B. Zeng, Integrable Rosochatius deformations of higher-order constrained flows and the soliton hierarchy with self-consistent sources, *J. Phys. A: Math. Theor.* 41 (2008) 295205.
- [16] Y.Q. Yao, X.J. Liu, Y.B. Zeng, A new extended discrete KP hierarchy and a generalized dressing method, *J. Phys. A: Math. Theor.* 42 (2009) 454026.
- [17] W.X. Ma, An extended Harry Dym hierarchy, *J. Phys. A: Math. Theor.* 43 (2010) 165202.
- [18] Y.Q. Yao, Y.H. Huang, G.X. Dong, Y.B. Zeng, The new integrable deformations of a short pulse equation and sine–Gordon equation, and their solutions, *J. Phys. A: Math. Theor.* 44 (2011) 065201.
- [19] J.Y. Zhu, X.G. Geng, A hierarchy of coupled evolution equations with self-consistent sources and the dressing method, *J. Phys. A: Math. Theor.* 46 (2012) 035204.
- [20] H. Wang, T.C. Xia, Conservation laws and self-consistent sources for a super KN hierarchy, *Appl. Math. Comput.* 219 (2013) 5458–5464.
- [21] V.K. Mel'nikov, A direct method for deriving a multi-soliton solution for the problem of interaction of waves on the x, y plane, *Comm. Math. Phys.* 112 (1987) 639–652.
- [22] V.K. Mel'nikov, Interaction of solitary waves in the system described by the Kadomtsev–Petviashvili equation with a self-consistent source, *Comm. Math. Phys.* 126 (1989) 201–215.
- [23] S.F. Deng, D.Y. Chen, D.J. Zhang, The multisoliton solutions of the KP equation with self-consistent sources, *J. Phys. Soc. Japan* 72 (2003) 2184–2192.
- [24] T. Xiao, Y.B. Zeng, Generalized Darboux transformations for the KP equation with self-consistent sources, *J. Phys. A: Math. Gen.* 37 (2004) 7143.
- [25] Y. Zhang, Y.B. Sun, W. Xiang, The rogue waves of the KP equation with self-consistent sources, *Appl. Math. Comput.* 263 (2015) 204–213.
- [26] R. Hirota, *The Direct Method in Soliton Theory*, Cambridge University Press, New York, 2004.

- [27] N.C. Freeman, J.J.C. Nimmo, Soliton solutions of the Korteweg–de Vries and Kadomtsev–Petviashvili equations: the Wronskian technique, *Phys. Lett. A* 95 (1983) 1–3.
- [28] Y. Ohta, J.K. Yang, General high-order rogue waves and their dynamics in the nonlinear Schrödinger equation, *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* 468 (2011) 1716–1740.
- [29] B.L. Guo, L.M. Lin, Q.P. Liu, Nonlinear Schrödinger equation: generalized Darboux transformation and rogue wave solutions, *Phys. Rev. E* 85 (2012) 026607.
- [30] W.X. Ma, Y. You, Solving the Korteweg–de Vries equation by its bilinear form: Wronskian solutions, *Trans. Amer. Math. Soc.* 357 (2005) 1753–1778.
- [31] S.V. Manakov, V.E. Zakharov, L.A. Bordag, A.R. Its, V.B. Matveev, Two-dimensional solitons of the Kadomtsev–Petviashvili equation and their interaction, *Phys. Lett. A* 63 (1977) 205–206.
- [32] Z. Lu, E.M. Tian, R. Grimshaw, Interaction of two lump solitons described by the Kadomtsev–Petviashvili I equation, *Wave Motion* 40 (2004) 123–135.
- [33] D.J. Kaup, The lump solutions and the Bäcklund transformation for the three-dimensional three-wave resonant interaction, *J. Math. Phys.* 22 (1981) 1176–1181.
- [34] C.R. Gilson, J.J.C. Nimmo, Lump solutions of the BKP equation, *Phys. Lett. A* 147 (1990) 472–476.
- [35] J. Satsuma, M.J. Ablowitz, Two-dimensional lumps in nonlinear dispersive systems, *J. Math. Phys.* 20 (1979) 1496–1503.
- [36] A.S. Fokas, M.J. Ablowitz, On the inverse scattering transform of multidimensional nonlinear equations related to first-order systems in the plane, *J. Math. Phys.* 25 (1984) 2494–2505.
- [37] K. Imai, Dromion and lump solutions of the Ishimori-I equation, *Progr. Theoret. Phys.* 98 (1997) 1013–1023.
- [38] W.X. Ma, Lump solutions to the Kadomtsev–Petviashvili equation, *Phys. Lett. A* 379 (2015) 1975–1978.
- [39] J.Y. Yang, W.X. Ma, Abundant interaction solutions to the KP equation, *Nonlinear Dynam.* 89 (2017) 1539–1544.
- [40] H.Q. Zhao, W.X. Ma, Mixed lump-kink solutions to the KP equation, *Comput. Math. Appl.* 74 (2017) 1399–1405.
- [41] J.Y. Yang, W.X. Ma, Lump solutions of the BKP equation by symbolic computation, *Internat. J. Modern Phys. B* 30 (2016) 1640028.
- [42] J.B. Zhang, W.X. Ma, Mixed lump-kink solutions to the BKP equation, *Comput. Math. Appl.* 74 (2017) 591–596.
- [43] Y.N. Tang, S.Q. Tao, G. Qing, Lump solitons and the interaction phenomena of them for two classes of nonlinear evolution equations, *Comput. Math. Appl.* 72 (2016) 2334–2342.
- [44] H.Q. Zhang, W.X. Ma, Lump solutions to the (2+1)-dimensional Sawada-Kotera equation, *Nonlinear Dynam.* 87 (2017) 2305–2310.
- [45] W.X. Ma, Z.Y. Qin, X. Lv, Lump solutions to dimensionally reduced p-gKP and p-gBKP equations, *Nonlinear Dynam.* 84 (2016) 923–931.
- [46] J.Y. Yang, W.X. Ma, Abundant lump-type solutions of the Jimbo–Miwa equation in (3+1)-dimensions, *Comput. Math. Appl.* 73 (2017) 220–225.
- [47] H.Q. Sun, A.H. Chen, Lump and lump-kink solutions of the (3+1)-dimensional Jimbo–Miwa and two extended Jimbo–Miwa equations, *Appl. Math. Lett.* 68 (2017) 55–61.
- [48] W.X. Ma, X.L. Yong, H.Q. Zhang, Diversity of interaction solutions to the (2+1)-dimensional Ito equation, *Comput. Math. Appl.* (2017). <http://dx.doi.org/10.1016/j.camwa.2017.09.013>.
- [49] B.Q. Li, Y.L. Ma, L.P. Mo, Y.Y. Fu, The N-loop soliton solutions for (2+1)-dimensional Vakhnenko equation, *Comput. Math. Appl.* 74 (2017) 504–512.
- [50] B.Q. Li, Y.L. Ma, T.M. Yang, The oscillating collisions between the three solitons for a dual-mode fiber coupler system, *Superlattices Microstruct.* 110 (2017) 126–132.
- [51] B.Q. Li, Y.L. Ma, T.M. Yang, Stable optical soliton in the ring-cavity fiber system with carbon nanotube as saturable absorber, *Superlattices Microstruct.* (2017). <http://dx.doi.org/10.1016/j.spmi.2017.11.016>.
- [52] S.L. Jia, Y.T. Gao, L. Hu, Q.M. Huang, W.Q. Hu, Soliton-like, periodic wave and rational solutions for a (3+1)-dimensional Boiti–Leon–Manna–Pempinelli equation in the incompressible fluid, *Superlattices Microstruct.* 102 (2017) 273–283.