



Snowmelt Timing Regulates Community Composition, Phenology, and Physiological Performance of Alpine Plants

Daniel E. Winkler^{1,2*}, Ramona J. Butz^{3,4}, Matthew J. Germino⁵, Keith Reinhardt⁶ and Lara M. Kueppers⁷

¹ School of Natural Sciences, University of California, Merced, Merced, CA, United States, ² United States Geological Survey, Southwest Biological Science Center, Moab, UT, United States, ³ Department of Forestry & Wildland Resources, Humboldt State University, Arcata, CA, United States, ⁴ Pacific Southwest Region, United States Department of Agriculture Forest Service, Eureka, CA, United States, ⁵ United States Geological Survey, Forest and Rangeland Ecosystem Science Center, Boise, ID, United States, ⁶ Department of Biological Sciences, Idaho State University, Pocatello, ID, United States, ⁷ Energy & Resource Group, University of California, Berkeley, Berkeley, CA, United States

OPEN ACCESS

Edited by:

Hans J. De Boeck,
University of Antwerp, Belgium

Reviewed by:

Thomas Abeli,
University of Pavia, Italy
Roman Zweifel,
Swiss Federal Institute for Forest,
Snow and Landscape Research,
Switzerland

*Correspondence:

Daniel E. Winkler
winklerde@gmail.com

Specialty section:

This article was submitted to
Functional Plant Ecology,
a section of the journal
Frontiers in Plant Science

Received: 15 May 2018

Accepted: 13 July 2018

Published: 31 July 2018

Citation:

Winkler DE, Butz RJ, Germino MJ,
Reinhardt K and Kueppers LM (2018)
Snowmelt Timing Regulates
Community Composition, Phenology,
and Physiological Performance
of Alpine Plants.
Front. Plant Sci. 9:1140.
doi: 10.3389/fpls.2018.01140

The spatial patterning of alpine plant communities is strongly influenced by the variation in physical factors such as temperature and moisture, which are strongly affected by snow depth and snowmelt patterns. Earlier snowmelt timing and greater soil-moisture limitations may favor wide-ranging species adapted to a broader set of ecohydrological conditions than alpine-restricted species. We asked how plant community composition, phenology, plant water relations, and photosynthetic gas exchange of alpine-restricted and wide-ranging species differ in their responses to a ca. 40-day snowmelt gradient in the Colorado Rocky Mountains (*Lewisia pygmaea*, *Sibbaldia procumbens*, and *Hymenoxys grandiflora* were alpine-restricted and *Artemisia scopulorum*, *Carex rupestris*, and *Geum rossii* were wide-ranging species). As hypothesized, species richness and foliar cover increased with earlier snowmelt, due to a greater abundance of wide-ranging species present in earlier melting plots. Flowering initiation occurred earlier with earlier snowmelt for 12 out of 19 species analyzed, while flowering duration was shortened with later snowmelt for six species (all but one were wide-ranging species). We observed >50% declines in net photosynthesis from July to September as soil moisture and plant water potentials declined. Early-season stomatal conductance was higher in wide-ranging species, indicating a more competitive strategy for water acquisition when soil moisture is high. Even so, there were no associated differences in photosynthesis or transpiration, suggesting no strong differences between these groups in physiology. Our findings reveal that plant species with different ranges (alpine-restricted vs. wide-ranging) could have differential phenological and physiological responses to snowmelt timing and associated soil moisture dry-down, and that alpine-restricted species' performance is more sensitive to snowmelt. As a result, alpine-restricted species may serve as better indicator species than their wide-ranging heterospecifics. Overall, alpine community composition and peak % cover are

strongly structured by spatio-temporal patterns in snowmelt timing. Thus, near-term, community-wide changes (or variation) in phenology and physiology in response to shifts in snowmelt timing or rates of soil dry down are likely to be contingent on the legacy of past climate on community structure.

Keywords: distribution, elevation, flowering, Niwot Ridge, photosynthesis, snowmelt gradient, spatio-temporal dynamics, water relations

INTRODUCTION

Recent climate warming coupled with regional declines in winter precipitation have led to an advance in the timing of snowmelt, one of the fastest changing environmental factors in alpine systems worldwide (Dyer and Mote, 2006; Barnett et al., 2008; Clow, 2010; Stocker et al., 2013; but see Mote et al., 2018). This has potentially serious implications for alpine plant communities (Wipf et al., 2006; Wipf and Rixen, 2010) and the ecosystem services they provide (e.g., carbon sequestration, mountain resilience; Bowman and Fisk, 2001; Barni et al., 2007). Spatial variation in the timing of snowmelt generates hydroclimate gradients over very short distances, structuring alpine plant communities (e.g., moist meadows, dry meadows, snowbeds; Walker et al., 1993; Taylor and Seastedt, 1994; Choler et al., 2001; Bruun et al., 2006; Jonas et al., 2008; Litaor et al., 2008). These gradients within the alpine zone include differences in timing and duration of soil water- and nutrient-availability, pH levels, and soil organic matter content (Walker et al., 1993; Stanton et al., 1994). Hydroclimate gradients represent natural experiments that can be used to better understand ecological processes in alpine systems (e.g., Cabrera et al., 1998; Michalet et al., 2014) and can also potentially act as space-for-time substitutions (Dunne et al., 2004).

Combined with relatively short growing season lengths (<3 months), these hydroclimate and associated environmental factors result in a landscape mosaic of vegetation communities, which vary in species composition, productivity, and physiological performance (Billings and Bliss, 1959; Stanton et al., 1994; Galen and Stanton, 1995; Germino and Smith, 2001; Winkler et al., 2016a). For example, sites with earlier snowmelt are typically more productive (Kudo et al., 1999) and can have greater species richness corresponding with better soil fertility (Stanton et al., 1994). This higher richness also reflects the larger number of wide-ranging species in relatively lower elevation and/or early melting sites (Kammer and Möhl, 2002; Bruun et al., 2006; Lenoir et al., 2008; Erschbamer et al., 2009). Typically, later melting sites are where alpine-restricted or specialist species occur (Odland and Munkejord, 2008; Pickering et al., 2014). While as many as 25% of species in alpine zones can also be found below treeline (Rundel, 2011), this leaves 75% of alpine species susceptible to competition with wide-ranging species as conditions change (Bruun et al., 2006; Steinbauer et al., 2018). However, to date, no studies have compared the potentially distinct phenological and physiological sensitivities of alpine-restricted and wide-ranging species.

Changes in growing season length may negatively influence plant production and sexual reproduction indirectly via shifts

in phenological cues including snowmelt timing (Kudo et al., 1999; Hülber et al., 2006; Venn and Morgan, 2007; Baptist and Choler, 2008). Individual alpine species vary in their ability to initiate growth immediately following snow retreat as a result of variation in traits like bud preformation, tolerance to chilling and photoinhibition, ability to develop under or around snow, photoperiodism, seed dormancy, and speed of development (Amen, 1965; Hamerlynck and Smith, 1994; Meloche and Diggle, 2001; Germino and Smith, 2000; Keller and Körner, 2003). Thus, snowmelt timing is expected to have varied effects on alpine plant phenology, including flowering duration; yet, we can expect that the strongest effects can be observed at early melting sites where diversity is typically higher and composed of a relatively larger number of wide-ranging species (Holway and Ward, 1963; Kudo, 1991, 1992; Kudo and Hirao, 2006; Sherwood et al., 2017). For example, Venn and Morgan (2007) found that individual species varied in their response to snowmelt timing, with some species synchronously flowering regardless of snowmelt timing due to bud preformation and others producing flowers shortly after snowmelt, thus closely tracking snowmelt. However, it remains unknown how species respond to differences in snowmelt timing as a function of their distributions.

While plant physiological performance also likely responds to snowmelt timing, this relationship has rarely been explicitly quantified as it has in phenological research (but see Oberbauer and Billings, 1981; Germino and Smith, 2001). In some temperate alpine systems, where soils rapidly dry down following snowmelt, and as environmental conditions change throughout the growing season, plants are exposed to intense radiative forcing and drier atmospheric conditions that can lead to desiccation as the season progresses (Geller and Smith, 1982; Smith and Johnson, 2009). Species-specific ecophysiological responses to snowmelt timing and soil dry-down rates may further explain patterns of alpine community composition and productivity. Adaptation to water stress and photosynthetic capacity of individual species likely contribute to individual presence in a community, with species exhibiting the highest water potentials in the wettest sites with earliest snowmelt (Oberbauer and Billings, 1981), and could be linked to whether they are alpine-restricted or wide-ranging species. Species that are wide-ranging across elevation gradients (i.e., broad environmental tolerance) may also be relatively well suited to adjust performance across small-scale gradients generated by snowmelt timing. For example, the wide-ranging *Bistorta vivipara* successfully adjusted (i.e., exhibited increased photosynthetic rates) when water was experimentally added at the end of the growing season when soils were driest (Enquist and Ebersole, 1994) but was unresponsive to watering earlier in the season when soils were likely still saturated with snowmelt

runoff from higher slopes or when summer precipitation events were frequent. This may explain a similar lack of photosynthetic response observed during early- and mid-season sampling events in a similar experiment that added water above ambient precipitation levels (Bowman et al., 1995). We might further expect that alpine-restricted species are at risk of local extinction if they are not able to physiologically adjust in response to earlier snowmelt and associated exposure outside of their typical optima (Lenoir et al., 2008). Yet, whether these potential plant relationships with snowmelt gradients can be attributed to plant-soil water relationships remains an unanswered question in alpine ecology.

In this study, we utilized a ca. 40-day snowmelt gradient in the Colorado Rocky Mountains to assess the associations between snowmelt timing and alpine community composition and function. We asked how community diversity and richness varied due to snowmelt timing. We predicted, as others have shown at similar sites (Litaor et al., 2008), that community composition at this site would be tightly correlated with snowmelt patterns. We also asked how flowering phenology and plant peak % cover varied due to snowmelt date. Given the short growing season length in this and similar systems, we expected that species would initiate growth shortly after snowmelt, and develop earlier with earlier snowmelt. We expected earlier melting plots, therefore, to have longer growing seasons resulting in higher species richness and peak % cover. Finally, we asked if alpine-restricted species (*Lewisia pygmaea*, *Sibbaldia procumbens*, *Hymenoxys grandiflora*) and wide-ranging species (*Artemisia scopulorum*, *Carex rupestris*, *Geum rossii*) differed in the sensitivity of plant water potential and leaf-level gas exchange to snowmelt timing. We expected that plant water potential and gas exchange would remain high across the gradient in wide-ranging species due to these species' presumed broad environmental tolerances. Conversely, we expected that alpine-restricted species would experience lowered water potentials and gas exchange by the end of the growing season in early melting plots reflecting greater moisture limitation compared to wide-ranging species.

MATERIALS AND METHODS

Study Site

Our alpine research site is located at Niwot Ridge, at 3545 m in elevation in the Front Range of the Colorado Rocky Mountains (40° 03' 14.84'' N, 105° 35' 37.71'' W). The site is on a 15° south-southeast facing slope, 400 m above local treeline. This site is characterized by a short growing season that typically lasts from

June through September (Greenland, 1989), high inter-annual variability in monthly mean temperatures and precipitation, and low growing season precipitation (Walker et al., 1995). Climate data recorded at the nearby Niwot Ridge LTER Saddle weather station show mean annual air temperature was -2.15°C and mean annual precipitation was 966 mm from 1981 to 2009. Mean microclimate data were calculated across the summer growing season (1 June–30 September; Table 1). Approximately 80% of the precipitation falls as snow at Niwot Ridge (Blanken et al., 2009). Snow depth at the site is spatially variable and controlled by topography and westerly wind (Walker et al., 1995; Litaor et al., 2008). The soils, developed on glaciofluvial deposits or residuum derived from igneous and metamorphic rock, are mapped as Moran family and classified as lithic Cryorthents. Community composition at the site is similar to moist and dry meadow community types (May and Webber, 1982; see Winkler et al., 2016a for a detailed description of the community).

We established 20 3 m diameter plots stratified by local elevation and aspect, as well as total plant cover. We divided plots into 4 1 m² quadrants to account for any fine-scale microtopography within plots (Supplementary Figure S1). The average slope of individual plots was 16% with a range of 8.5–21.5% across plots. The first plot was snow free on May 29 and the last plot on June 22. Variation in snow depth and snowmelt timing across plots results from prevailing winds from the west, the south-easterly aspect of the site, as well as microtopography within the site.

Field Measurements

All data were recorded during the 2009 growing season, before the plots became part of the Alpine Treeline Warming Experiment. We defined the growing season as the time from the date of snowmelt until all plants had senesced in late September. Date of snowmelt was determined by manual snow surveys carried out three times per week starting at the onset of spring snowmelt. Digital SLR planimetric photographs were taken 1.5 m above each quadrant to determine the date of snowmelt, defined as the first snow free day when all subsequent snow cover events lasted less than four continuous days.

We conducted vegetation surveys at peak community productivity (determined by weekly inspection of plant flowering phenology in each plot; Negi et al., 1992). Surveys began in the middle of July and concluded by early August following the ca. 40-day snowmelt gradient from the lowest to highest elevation plots at the site. We visually estimated peak % cover as an indirect estimate of productivity (Winkler et al., 2016a), first for the entire community and then for individual species using a 1 m² survey

TABLE 1 | Summary of microclimate data for summer 2009 (Jun 1 – Sep 30).

Snowmelt dates	R_{solar} (W m ⁻²)	T_{air} (°C)	T_{min} (°C)	T_{max} (°C)	T_{soil} (°C)	# $T_{\text{night}} < 0^{\circ}\text{C}$	VPD (kPa)
149–173	441 (8)	7.9 (0.3)	3.5 (0.3)	13.4 (0.4)	6.7 (0.2)	16	0.43 (0.01)

Microclimate data except for snowmelt dates are from an LTER climate station located nearby (<500 m away and approximately at the same elevation). Snowmelt dates indicate the plot-level range in snowmelt in Julian days in the study site. Solar radiation (R_{solar} ; 300–3000 m), air (T_{air}), minimum air (T_{min}), maximum air (T_{max}), and soil (T_{soil}) temperatures, and air vapor pressure deficit (VPD) are summer daily means. # $T_{\text{night}} < 0^{\circ}\text{C}$ is the number of nights during the summer when air temperatures dropped below 0°C . Numbers in parentheses are SE.

grid divided into 10 cm² units. We also estimated % cover of all other surface types (i.e., solid rock, lichen cover, bryophyte cover, bare ground, fine litter, and woody debris). We measured 45 species across our site in 2009 (Supplementary Table S1). We conducted flowering phenology surveys for all species in each quadrant weekly beginning at snowmelt and continuing until flowering ceased toward the end of the growing season. These measures were used to determine the date of first flower and flowering duration for each species in each quadrant.

We selected six species for physiological measurements during the growing season, including three alpine-restricted species that only occur in the alpine zone (*L. pygmaea*, *S. procumbens*, and *H. grandiflora*) and three wide-ranging species that occur in the alpine zone but also occur at or below treeline (*A. scopulorum*,

C. rupestris, and *G. rossii*). We measured gas exchange in the field on intact leaves during three sampling events each lasting 2 days (early July, *n* = 19; early August, *n* = 63; and early September, *n* = 55). We randomly shuffled species and plots sampled to avoid effects of sampling time (none were detected). All measurements were conducted during the hours of peak sunlight (>90% of maximum light) using an infrared gas analyzer photosynthesis system (LI-6400XT; Li-Cor Biosciences, Lincoln, NE, United States) equipped with a 2 cm × 3 cm leaf chamber with an internal LED light source. During all measurements, chamber conditions were set to a photon flux density of 1500 μmol m⁻² s⁻¹ and CO₂ concentration of 400 μl mol⁻¹. Vapor pressure was kept at ambient levels during all measurements and temperature inside the chamber was set to match ambient air temperature using the flow of air inside the chamber. Gas exchange was calculated on a projected leaf area basis, with leaf area in the chamber determined using digital photographs of the portion of leaf area inside chamber gaskets and using image processing software (ImageJ; Scion Co., Frederick, MD, United States). During the July and September measurement events, conditions were partially cloudy; during the August measurements, conditions were sunny.

Net CO₂ assimilation (*A*_{net}), stomatal conductance (*g*), internal concentration of CO₂ (*C*_i), and transpiration (*E*) were calculated according to Ball (1987) and Farquhar and von Caemmerer (1982), and water-use efficiency (WUE) was

TABLE 2 | Species diversity of the entire alpine plant community at our site.

Diversity measure	Value
Alpha	17
Beta	2.65
Gamma	45
Shannon's <i>H</i>	2.10
Pielou's <i>J</i>	0.75

Diversity indices are defined in the "Materials and Methods" section of the text.




FIGURE 1 | % slope, snowmelt timing (Julian day), richness, and % cover gradients fit as splines across our site. Points indicate locations of plots at our site and black contours indicate elevation. Splines were generated using generalized cross-validation scores to portray relationships between elevation and each variable along contours.

calculated as A_{net}/E . Following an initial measurement of A_{net} under saturating light, the chamber was darkened (light intensity of $0 \mu\text{mol m}^{-2} \text{s}^{-1}$), and dark respiration (R_d) was recorded after CO_2 ceased to increase in the chamber and used as an indication of growth and maintenance respiration (Atkin et al., 2000). R_d under full sun conditions can be less than in what would be measured at night (Krömer, 1995), and consequently our calculated values may be overestimated. During the third sampling event (early September), we also increased CO_2 to $800 \mu\text{mol mol}^{-1}$ ($n = 55$) to generate information on stomatal limitation to photosynthesis (greater increases in photosynthesis with increased CO_2 at $800 \mu\text{mol mol}^{-1}$ indicate greater stomatal limitation to photosynthesis at $400 \mu\text{mol mol}^{-1}$; Sage, 1994). One individual per species was sampled for gas exchange measurements in each plot when possible (not all species occurred in all plots). Last, we measured plant water potential on detached leaves at pre-dawn (0400–0600 h local time) using a Scholander type pressure chamber (PMS-1000; PMS Instruments, Corvallis, OR, United States). Due to the long-term nature of our research plots, pre-dawn water potential measurements were made outside of the plots on 2–3 individuals of each species at each sampling date near early-, mid- and late-melting plots.

Statistical Analyses

We calculated alpha (average species richness per plot), beta (a measure of species turnover among plots calculated as the ratio of gamma over alpha), and gamma (total species richness across all plots) diversity, and Shannon's H and Pielou's J (Pielou, 1969) as measures of evenness. Diversity metrics are reported in Table 2. All analyses were carried out in R 3.3.2. (R Core Team, 2014). We displayed the relationships between %

slope of plots, snowmelt timing, richness, and % cover across the elevation gradient of our site using the Tps command from the fields package (Nychka et al., 2015). This command uses generalized cross-validation scores to portray relationships between elevation and each variable along contours. We also used non-metric multidimensional scaling (NMS) ordination to examine associations among snowmelt date and cover variables across plots. We did the same for presence/absence of individual species, log-transforming data to account for zero values and high skew (McCune and Grace, 2002). NMS ordinations were conducted using PC-ORD (MjM Software Design; Gleneden Beach, OR, United States).

Species that occurred in ten or more plots were selected for phenological analyses ($n = 19$ spp.). We used a linear mixed effects model to determine whether snowmelt timing explains date of first flower across all 19 species, with snowmelt timing as a fixed effect and plot, quadrant, and species as nested random effects to account for pseudoreplication across quadrants and species. We compared this model to a null (intercept-only model) and used the change in Akaike Information Criterion corrected for small sample sizes (ΔAICc ; Johnson and Omland, 2004; Aho et al., 2014). We subsequently used linear regression to test for correlations between date of first flower and snowmelt timing, and flowering duration and snowmelt timing for individual species. We used an alpha value with Bonferroni correction ($\alpha = 0.025$) to account for non-independence, but note marginally significant values ($\alpha = 0.05$) when appropriate to address our increased concern over Type II error.

We used linear mixed effects models to test for differences in A_{net} with snowmelt timing (i.e., early, mid, and late) and group (i.e., alpine-restricted vs. wide-ranging species), their interaction, and sampling event (i.e., July, August, and September) as fixed




FIGURE 2 | Non-metric multidimensional scaling (NMS) ordination of % cover measurements (i.e., vascular cover, bare ground, litter, gravel, and snowmelt) across the field site illustrating within community gradients. Numbers correspond to plot identifiers and colored triangles indicate snowmelt timing of each plot (warmer colors indicate later snowmelt).

effects. Similar to our phenology models, plot, quadrant, and species were included as nested random effects to account for pseudoreplication across quadrants and species. We compared the full model to simpler versions and used the same ΔAICc approach described above to determine the best-fit model. We carried out the same analysis for R_d , g , C_i , E , and WUE. When full models indicated a fixed effect was important for predicting performance, we compared least-squares means of groupings adjusted for repeated measures using the Tukey method (e.g., July vs. August sampling events, alpine-restricted vs. wide-ranging species, etc.). Last, we used the same mixed model strategy for our pre-dawn water potential measurements. Models were evaluated using the nlme and lsmeans packages (Lenth, 2016; Pinheiro et al., 2018).

RESULTS

Community Responses

Species richness and total % cover by vascular plants differed in their sensitivity to elevation and snowmelt timing (Figure 1), with richness more closely tracking elevation and % cover more closely tracking snowmelt timing. NMS ordination of plot level % cover achieved the greatest reduction in stress with just two axes (Figure 2). The proportion of variance explained by the first two axes was 0.812 and 0.093 (total 0.906 explained). Vascular plant cover, litter, snowmelt timing, and gravel were significantly associated with the ordination axes. Plots with earlier snowmelt exhibited a higher percentage of vascular cover, and lower percentages of litter and gravel cover. Species compositions of plots were determined by snowmelt timing, with ca. 20% of the 45 species occurring only in the earliest or latest melting plots (Figure 3). A subset of ca. 25 species were found across the entire snowmelt gradient and are clustered at the center of the NMS ordination (Figure 3).


Phenological Responses

Linear mixed models predicting date of first flower with snowmelt timing ($\text{AICc} = 2164.77$) outperformed a null model ($\text{AICc} = 2196.75$, $\Delta\text{AICc} = 31.99$), suggesting that snowmelt timing has predictive power in explaining the date of first flower across our community. The same was true for the duration of flowering across our site (full model $\text{AICc} = 2201.78$, null model $\text{AICc} = 2226.49$, $\Delta\text{AICc} = 24.71$). Of the nineteen species analyzed, twelve showed a significant positive correlation between date of first flower and snowmelt timing ($P \leq 0.025$; with *C. rupestris* and *Trifolium parryi* $P \leq 0.05$; Table 3), while only six species showed a significant negative correlation between flowering duration and snowmelt ($P \leq 0.025$). Of the twelve that showed significant correlations between date of first flower and snowmelt timing, a majority of these species are wide-ranging species (*A. scopulorum*, *B. bistortoides*, *C. rupestris*, *Castilleja occidentalis*, *Erigeron simplex*, *G. rossii*, *Lloydia serotina*, and *Potentilla diversifolia*, *T. parryi*; Table 3) whereas only three are alpine-restricted species (*Chionophila jamesii*, *L. pygmaea*, and *Silene acaulis*). Of the six species that showed a significant correlation between flowering duration and snowmelt, five are

wide-ranging species (*A. scopulorum*, *B. bistortoides*, *Castilleja occidentalis*, *E. simplex*, and *Lloydia serotina*) and only one is alpine-restricted (*L. pygmaea*; Table 3).

Physiological Responses

Plant gas exchange varied substantially through time but was invariant across species and the spatial snowmelt gradient



(Figure 4). The interaction of group and snowmelt timing with sampling event as an additional covariate best predicted photosynthesis ($w_i = 0.99$; Supplementary Table S2). However, photosynthesis did not differ between alpine-restricted and wide-ranging species and photosynthesis declined in all species > 50% throughout the growing season (Figure 4A). Compared to in July, A_{net} was significantly lower in August ($t = 5.41$, $P < 0.001$) and September ($t = 5.20$, $P < 0.001$). Photosynthesis increased when measured at $800 \mu\text{mol mol}^{-1} \text{CO}_2$ compared to $400 \mu\text{mol mol}^{-1} \text{CO}_2$ in September (Figure 5), especially in wide-ranging species (though not statistically different). Increases in photosynthesis were greater in plots that had melted out late in the summer (1.05-fold increase at 800 compared to $400 \mu\text{mol mol}^{-1}$) compared to plots that had melted mid-summer (0.89-fold increase) or in early summer (0.65 increase; Figure 5), though these differences were not statistically significant.

Dark respiration was also best predicted by the full model with the interaction of group and snowmelt timing, with sampling event as an additional covariate ($w_i = 0.70$; Supplementary Table S3). Although respiration decreased by ~50% in all species throughout the growing season, these decreases were not significantly explained by either species groupings or snowmelt timing (Figures 4B,H). Internal-concentration of leaf CO_2 was also predicted best by the full model ($w_i = 0.99$; Supplementary Table S4), but values for the two species groups tracked each other, slightly declining throughout the season. Transpiration was also best predicted by the full model ($w_i = 0.90$;

Supplementary Table S5). Transpiration similarly showed no difference between groups but significantly declined throughout the season; July and August values were 30–50% higher than September values (July: $t = 4.269$, $P < 0.01$; August: $t = 6.70$, $P < 0.001$; Figures 4E,K).

Stomatal conductance and WUE were best explained by group and sampling event (g: $w_i = 0.98$; WUE $w_i = 0.40$; Supplementary Tables S6, S7). Stomatal conductance was higher for wide-ranging species than alpine-restricted species, mostly due to differences early in the summer ($t = -2.05$, $P = 0.04$; Figures 4C,I), though all species exhibited declines in stomatal conductance from early to late in the growing season (all *post hoc* pairwise tests had $P < 0.001$). WUE models with only snowmelt timing and sampling event performed just as strongly ($w_i = 0.36$, $\Delta\text{AICc} = 0.21$; Supplementary Table S7), suggesting that sampling event is largely driving the model's predictive power. WUE values significantly declined from July to August ($t = -5.04$, $P < 0.001$; Figure 4F) but there were no differences between groups, nor were there differences associated with snowmelt timing (Figure 4L). Pre-dawn water potential was best explained by snowmelt timing and sampling event ($w_i = 0.60$; Supplementary Table S8). Last, pre-dawn leaf water potential for all species increased (became less negative) by ca. 50% from July to August ($t = -4.68$, $P < 0.001$; Figure 6) and then significantly decreased by ca. 150% from August to September ($t = 9.00$, $P < 0.001$). However, there were no statistically significant differences due to snowmelt date (early vs. late melt: $t = -2.69$,

TABLE 3 | Linear regressions of date of first flower (Julian day) and flowering duration (number of days) as functions of date of snowmelt (Julian day).

Species	<i>n</i>	First flower					Flowering duration			
		$\bar{x} \pm \text{SEM}$	r^2	Slope	<i>P</i>		$\bar{x} \pm \text{SEM}$	r^2	slope	<i>P</i>
<i>Arenaria fendleri</i>	W	20	198 ± 3	0.129	0.136	0.121	41 ± 5	0.099	0.201	0.176
<i>Artemisia scopulorum</i>	W	20	193 ± 8	0.510	0.657	<0.001***	24 ± 8	0.235	−0.449	0.030*
<i>Bistorta bistortoides</i>	W	19	194 ± 9	0.478	0.782	0.001***	35 ± 9	0.296	−0.579	0.016*
<i>Carex rupestris</i>	W	16	194 ± 12	0.315	0.760	0.037*	28 ± 17	0.100	−0.597	0.271
<i>Castilleja occidentalis</i>	W	14	183 ± 6	0.539	0.537	0.001***	37 ± 15	0.439	−1.144	0.005**
<i>Chionophila jamesii</i>	A	20	197 ± 8	0.326	0.574	0.009**	20 ± 9	0.183	−0.454	0.060
<i>Erigeron simplex</i>	W	20	197 ± 6	0.766	0.670	<0.001***	11 ± 6	0.394	−0.478	0.003**
<i>Festuca brachyphylla</i>	W	15	219 ± 4	0.169	0.169	0.128	12 ± 3	0.172	−0.147	0.124
<i>Geum rossii</i>	W	20	186 ± 7	0.838	0.781	<0.001***	39 ± 6	0.032	−0.142	0.449
<i>Lewisia pygmaea</i>	A	20	184 ± 7	0.604	0.678	<0.001***	21 ± 8	0.269	−0.524	0.019*
<i>Lloydia serotina</i>	W	16	188 ± 8	0.724	0.833	<0.001***	15 ± 8	0.549	−0.668	0.001***
<i>Luzula spicata</i>	W	18	217 ± 1	0.022	0.023	0.554	7 ± 3	0.001	0.015	0.890
<i>Minuartia obtusiloba</i>	A	20	189 ± 3	0.010	0.039	0.680	47 ± 8	0.072	0.264	0.252
<i>Potentilla diversifolia</i>	W	16	194 ± 7	0.411	0.473	0.007**	26 ± 14	0.114	−0.533	0.200
<i>Sedum lanceolatum</i>	W	14	221 ± 8	0.023	0.133	0.604	14 ± 5	0.235	−0.284	0.079
<i>Silene acaulis</i>	A	10	199 ± 5	0.544	0.484	0.015*	10 ± 6	0.003	0.044	0.879
<i>Sibbaldia procumbens</i>	A	12	187 ± 5	0.010	0.216	0.758	26 ± 11	0.124	1.567	0.261
<i>Trifolium parryi</i>	W	10	194 ± 7	0.414	0.539	0.045*	11 ± 7	0.241	−0.404	0.150
<i>Trisetum spicatum</i>	W	20	202 ± 7	0.089	0.269	0.201	32 ± 7	0.005	−0.055	0.773

Mean Julian day of first flower and standard error ($\bar{x} \pm \text{SEM}$) are calculated for each species across all plots it occurred in (*n*). (For $\bar{x}$, day 183 = 2 July; day 221 = 9 August). First flower: linear regression of date of first flower of each species in each plot as a function of date of snowmelt of each plot. Mean flowering duration (mean number of Julian days: $\bar{x}$) and standard deviations ($\pm s$) are calculated for each species across all plots it occurred in. Flowering duration: linear regression of flowering duration of each species in each plot as a function of date of snowmelt of each plot. *** ≤ 0.001 , ** ≤ 0.01 , * ≤ 0.05 . W near a species name indicates that it is a wide-ranging species. A near a species name indicates that it is an alpine-restricted species.




FIGURE 4 | Physiological measurements separated by alpine-restricted (closed circles and solid lines) and wide-ranging species (open circles and dotted lines) by sampling month (A–F) and snowmelt timing (G–L). Means and standard errors are presented for photosynthetic rates (A_{net} ; $\mu\text{mol m}^{-2} \text{s}^{-1}$), dark respiration (R_d ; $\mu\text{mol CO}_2 \text{m}^{-2} \text{s}^{-1}$), stomatal conductance (g ; $\text{mol m}^{-2} \text{s}^{-1}$), CO₂ assimilation (C_i ; PPM), transpiration (E ; $\text{mmol H}_2\text{O m}^{-2} \text{s}^{-1}$), and water-use efficiency (WUE; $\mu\text{mol mmol}^{-1}$).

$P = 0.14$; early vs. mid melt: $t = -2.38$, $P = 0.19$; late vs. mid melt: $t = 0.20$, $P = 0.98$; **Figure 6**).


DISCUSSION

Our findings reveal that spatial variation in the composition and peak % cover of this moist alpine meadow community is driven by snowmelt timing. Flowering phenology was also strongly related to snowmelt timing for a majority of species. Leaf physiology tracked seasonal soil dry-down similarly among species, regardless of whether they were alpine-restricted or wide-ranging species. While early season stomatal conductance was higher in wide-ranging species, there were no associated differences in photosynthesis or transpiration, suggesting no strong differences between these groups in physiology.

Previous research in the Mosquito Range in Colorado shows that species richness and total community cover were

significantly greater in early melting microsites (Stanton et al., 1994). Our results corroborate these findings but also suggest that richness can more closely follow elevation gradients than snowmelt. Species richness and peak % cover increased with earlier snowmelt, due to a larger number of wide-ranging species present in earlier melting plots. Interestingly, peak % cover typically declined in areas where topographic depressions led to later snowmelt.


Future climate scenarios project a decrease in the snow-to-rain ratio during winter as a result of warmer temperatures (Stocker et al., 2013). Decreases in snowpack combined with warmer temperatures will ultimately lead to an advance of snowmelt and this has already been observed in montane systems (Barnett et al., 2008; Clow, 2010). Changes in snowmelt timing across a gradient or within a patch, and the associated changes in soil dry-down rates have implications for community structure and function (Pickering et al., 2014). Not only can we expect shifts in productivity, perhaps leading to a more productive alpine



(Kullman, 2010; but see Baptist et al., 2010), but an overall compositional shift that will result in novel alpine communities is also possible (Alexander et al., 2015). Productivity in this system has previously been shown to be sensitive to changes in soil moisture (Winkler et al., 2016a). This was shown using a warming experiment at our site that advanced snowmelt and subsequently led to more rapid soil dry-down rates, thus influencing productivity. Our current study illustrates a similar

effect of snowmelt timing but also reveals spatial structuring of the community across the gradient that contributes to productivity differences. If species are able to locally shift upward in elevation (Lenoir et al., 2008; Kiełtyk, 2017) or seek refuge in nearby microclimates (e.g., local depressions) where snowmelt changes are less pronounced (Opedal et al., 2015), productivity changes will reflect both the immediate effects of earlier snowmelt and the longer term effects of community change.

The flowering phenology of many alpine species worldwide closely tracks environmental cues including snowmelt timing, presumably as a result of pressure from relatively short growing seasons (Kudo, 1991; Walker et al., 1995; Studer et al., 2005; Kudo and Hirao, 2006; Björk and Molau, 2007). Our study reveals that flowering phenology can vary depending on whether a species' range is restricted to the alpine or extends below treeline. We show that wide-ranging species have more flexible flowering phenologies and, in most cases, will flower longer with earlier snowmelt (nine out of twelve species with significant correlations between snowmelt timing and timing of flowering initiation were wide-ranging species) whereas alpine-restricted species appear more conservative with initiation (only 3 out of 12 species showed significant correlations with snowmelt). The same was true for duration of flowering with only one alpine-restricted species and five wide-ranging species showing a significant correlation with snowmelt. Although our results are within one community type, other studies have shown that entire communities can exhibit this relationship where more diverse communities flower earlier and more consistently (i.e., those similar to microsites where snowmelt occurs earlier in our community), while less diverse communities composed primarily of specialists (i.e., those similar to the microsites with



late snowmelt in our community), are largely at the whim of snowmelt timing (Kudo and Hirao, 2006; Venn and Morgan, 2007; Baptist et al., 2010).

Many of the trends we observed in flowering phenology across the snowmelt gradient were species-level responses to snowmelt timing and related abiotic drivers. That said, these plant-environment relationships can sometimes be carried up to functional groups (Iversen et al., 2009; but see Henry and Molau, 1997). For example, nearly all of the graminoid species in our site flowered irrespective of snowmelt timing, flowering later in the season than other species on average and highlighting their late-season, drought-avoidance strategies (Rosbakh et al., 2017). However, these strategies might not always be dependent on environmental conditions experienced during the current growing season. It is estimated that ca. 50% or more of alpine plant species pre-form leaf and flower buds before the growing season begins and oftentimes 1+ years prior (Theodose et al., 1996; Körner, 2003). Since much is theorized and little is known about the prevalence of bud preformation and whether or not it occurs in many of our study species, it is difficult to say to what extent previous-years climate plays a role in the initiation of growth and flowering. However, in order to take advantage of early snowmelt, early flowering species need to maintain a high metabolic readiness while under the snow, which is dependent upon bud preformation (Körner, 1999). *G. rossii* date of first flower had the strongest correlation with date of snowmelt. *G. rossii* is not only a dominant species in our plots, but also has a long period of growth for each leaf and inflorescence (3 years) from initiation through senescence as a result of bud preformation (Meloche and Diggle, 2001). Along with temperature, the role of photoperiodism is certainly a main driver in the initiation of seasonal growth in many alpine plants (Keller and Körner, 2003). However, our data suggest that wide-ranging species may be taking advantage of growth initiation immediately after snowmelt in order to establish themselves as dominants in the community (both *G. rossii* and *C. rupestris* alternate dominance or co-dominate in our plots in terms of cover). Further studies are needed to elucidate the role of bud preformation in species at Niwot Ridge in order to concretely say what the extent of the impact of earlier snowmelt will be on these species, as well as those species whose phenologies are not correlated with snowmelt date.

Advances in snowmelt timing may provide a longer period for growth, but may also expose plants to an increased number of spring frost events that hinder growth and performance (Wipf et al., 2009; Sierra-Almeida and Cavieres, 2010). Few studies have examined alpine plant physiological connections to snowmelt timing but several studies have looked at related environmental variables that track snowmelt gradients like the one in our study (e.g., early season temperatures, species temperature optima, soil moisture status; Germino and Smith, 2000, 2001; Shen et al., 2009; Shi et al., 2010). However, our results suggest that snowmelt timing does not have a large impact on plant performance and, instead, physiological adjustment occurs in both alpine-restricted and wide-ranging species as the season progresses and environmental conditions change.

It is likely that the species in our community are displaying some level of physiological compensation by altering assimilation rates and water use along the snowmelt gradient (Ren et al., 2010). In addition to influencing composition and phenology, microclimate can also regulate plant physiological performance (Germino and Smith, 2001). We did detect higher stomatal conductance rates in wide-ranging species compared to alpine-restricted species early in the season. Even so, species overwhelmingly adjusted photosynthetic, conductance, transpiration rates, and leaf water potentials as water availability declined during the season. Water-use efficiency similarly declined from early to mid-season but increased late in the growing season when soils were driest and plants switched to investing in reproductive structures. Together, our physiological data confirm that alpine plant species are highly adapted to large amounts of environmental variation (Körner, 1999) and that responses to short-term change may not be as apparent if physiological thresholds are not crossed (Walther et al., 2002).

In this sense, the potential for species replacement or invasion increases as temperatures warm or other driving factors (e.g., soil moisture status) change with shifts in snowmelt timing. For example, snowmelt changes and subsequent increases in soil dry-down rates have already led to the encroachment of an invasive subalpine dwarf bamboo in Hokkaido, Japan (Winkler et al., 2016b). Changes in microclimates in alpine settings have the potential to negatively impact current community members by creating unfavorable conditions and creating stepping stones for non-native species to invade (Lembrechts et al., 2017). Similarly, warmer temperatures may lead to shifts in the position of local treeline (Kueppers et al., 2017), though these shifts may largely depend on available soil moisture (Moyes et al., 2013, 2015). Nonetheless, our results suggest that alpine community composition and peak % cover are strongly structured by spatio-temporal patterns in snowmelt timing, and may not change as rapidly as climate due to long generation times.

AUTHOR CONTRIBUTIONS

LK, MG, KR, and RB conceived and designed the experiments. RB collected the data. DW and RB analyzed the data. DW, LK, and RB wrote the manuscript. MG and KR provided editorial advice.

FUNDING

This research was supported by the Office of Science, (BER), United States Department of Energy. Additional funding was provided by the Environmental Systems Graduate Group and Graduate Division at the University of California, Merced.

ACKNOWLEDGMENTS

Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the United States Government. We thank the Mountain Research

Station and Niwot Ridge LTER at the University of Colorado, Boulder for logistical support. Additionally, we thank all who assisted in data collection and maintenance of the experiment, especially M. Barlerin, S. Barlerin, S. Ferrenberg, A. Dixon, A. Faist, and A. Farnham.

REFERENCES

- Aho, K., Derryberry, D., and Peterson, T. (2014). Model selection for ecologists: the worldviews of AIC and BIC. *Ecology* 95, 631–636. doi: 10.1890/13-1452.1
- Alexander, J. M., Diez, J. M., and Levine, J. M. (2015). Novel competitors shape species' responses to climate change. *Nature* 525, 515–518. doi: 10.1038/nature14952
- Amen, R. D. (1965). Seed dormancy in the alpine rush, *Luzula Spicata* L. *Ecology* 46, 361–364. doi: 10.2307/1936346
- Atkin, O. K., Millar, A. H., Gardestrom, P., and Day, D. A. (2000). "Photosynthesis, carbohydrate metabolism and respiration in leaves of higher plants," in *Photosynthesis: Physiology and Metabolism*, eds R. C. Leegood, E. Thomas, and S. Von Caemmerer (Dordrecht: Kluwer), 153–175. doi: 10.1007/0-306-48137-5_7
- Ball, J. T. (1987). "Calculations related to gas exchange," in *Stomatal Function*, ed. I. R. Cowan (Palo Alto, CA: Stanford University Press), 445–476.
- Baptist, F., and Choler, P. (2008). A simulation of the importance of length of growing season and canopy functional properties on the seasonal gross primary production of temperate alpine meadows. *Ann. Bot.* 101, 549–559. doi: 10.1093/aob/mcm318
- Baptist, F., Flahaut, C., Streb, P., and Choler, P. (2010). No increase in alpine snowbed productivity in response to experimental lengthening of the growing season. *Plant Biol.* 12, 755–764. doi: 10.1111/j.1438-8677.2009.00286.x
- Barnett, T. P., Pierce, D. W., Hidalgo, H. G., Bonfils, C., Santer, B. D., Das, T., et al. (2008). Human-induced changes in the hydrology of the Western United States. *Science* 319, 1080–1083. doi: 10.1126/science.1152538
- Barni, E., Freppaz, M., and Siniscalco, C. (2007). Interactions between vegetation, roots, and soil stability in restored high-altitude ski runs in the Alps. *Arct. Antarct. Alp. Res.* 39, 25–33. doi: 10.1657/1523-0430(2007)39[25:IBVRAS]2.0.CO;2
- Billings, W. D., and Bliss, L. C. (1959). An alpine snowbank environment and its effects on vegetation, plant development, and productivity. *Ecology* 40, 388–397. doi: 10.2307/1929755
- Björk, R. G., and Molau, U. (2007). Ecology of alpine snowbeds and the impact of global change. *Arct. Antarct. Alp. Res.* 39, 34–43. doi: 10.1657/1523-0430(2007)39[34:EOASAT]2.0.CO;2
- Blanken, P. D., Williams, M. W., Burns, S. P., Monson, R. K., Knowles, J., Chowanski, K., et al. (2009). A comparison of water and carbon dioxide exchange at a windy alpine tundra and subalpine forest site near niwot ridge, Colorado. *Biogeochemistry* 95, 61–76. doi: 10.1007/s10533-009-9325-9
- Bowman, W. D., and Fisk, M. C. (2001). "Primary production," in *Structure and Function of an Alpine Ecosystem: Niwot Ridge, Colorado*, eds W. D. Bowman and T. R. Seastedt (Oxford: Oxford University Press), 177–197.
- Bowman, W. D., Theodose, T. A., and Fisk, M. C. (1995). Physiological and production responses of plant growth forms to increases in limiting resources in alpine tundra: implications for differential community response to environmental change. *Oecologia* 101, 217–227. doi: 10.1007/BF00317287
- Bruun, H. H., Moen, J., Virtanen, R., Grytnes, J. A., Oksanen, L., and Angerbjörn, A. (2006). Effects of altitude and topography on species richness of vascular plants, bryophytes and lichens in alpine communities. *J. Veg. Sci.* 17, 37–46. doi: 10.1111/j.1654-1103.2006.tb02421.x
- Cabrera, H. M., Rada, F., and Cavieles, L. (1998). Effects of temperature on photosynthesis of two morphologically contrasting plant species along an altitudinal gradient in the tropical high Andes. *Oecologia* 114, 145–152. doi: 10.1007/s004420050430
- Choler, P., Michalet, R., and Callaway, R. M. (2001). Facilitation and competition on gradients in alpine plant communities. *Ecology* 82, 3295–3308. doi: 10.1890/0012-9658(2001)082[3295:FACOGI]2.0.CO;2
- Clow, D. W. (2010). Changes in the timing of snowmelt and streamflow in Colorado: a response to recent warming. *J. Clim.* 23, 2293–2306. doi: 10.1175/2009JCLI2951.1
- Dunne, J. A., Saleska, S. R., Fischer, M. L., and Harte, J. (2004). Integrating experimental and gradient methods in ecological climate change research. *Ecology* 85, 904–916. doi: 10.1890/03-8003
- Dyer, J. L., and Mote, T. L. (2006). Spatial variability and trends in observed snow depth over North America. *Geophys. Res. Lett.* 33:L16503. doi: 10.1029/2006GL027258
- Enquist, B. J., and Ebersole, J. J. (1994). Effects of added water on photosynthesis of *Bistorta vivipara*: the importance of water relations and leaf nitrogen in 2 alpine communities, Pikes Peak, Colorado, USA. *Arct. Alp. Res.* 26, 29–34. doi: 10.2307/1551873
- Erschbamer, B., Kiebachner, T., Mallaun, M., and Unterluggauer, P. (2009). Short-term signals of climate change along an altitudinal gradient in the South Alps. *Plant Ecol.* 202, 79–89. doi: 10.1007/s11258-008-9556-1
- Farquhar, G. D., and von Caemmerer, S. (1982). "Modeling of photosynthetic response to environmental conditions," in *Encyclopedia of Plant Physiology*, eds O. L. Lange, P. S. Noble, C. B. Osmond, and H. Ziegler (Berlin: Springer-Verlag), 549–558.
- Galen, C., and Stanton, M. (1995). Responses of snowbed plant species to changes in growing-season length. *Ecology* 76, 1546–1557. doi: 10.2307/1938156
- Geller, G. N., and Smith, W. K. (1982). Influence of leaf size, orientation, and arrangement on temperature and transpiration in three high-elevation, large-leaved herbs. *Oecologia* 53, 227–234. doi: 10.1007/BF00545668
- Germiño, M. J., and Smith, W. K. (2000). Differences in microsite, plant form, and low-temperature photoinhibition in alpine plants. *Arct. Antarct. Alp. Res.* 32, 388–396. doi: 10.1080/15230430.2000.12003382
- Germiño, M. J., and Smith, W. K. (2001). Relative importance of microhabitat, plant form and photosynthetic physiology to carbon gain in two alpine herbs. *Funct. Ecol.* 15, 243–251. doi: 10.1046/j.1365-2435.2001.00506.x
- Greenland, D. (1989). The climate of niwot ridge, front range, Colorado. *Arct. Alp. Res.* 21, 380–391. doi: 10.2307/1551647
- Hamerlynck, E. P., and Smith, W. K. (1994). Subnivean and emergent microclimate, photosynthesis, and growth in *Erythronium grandiflorum* pursh, a snowbank geophyte. *Arct. Alp. Res.* 26, 21–28. doi: 10.2307/1551872
- Henry, G. H. R., and Molau, U. (1997). Tundra plants and climate change: the International Tundra Experiment (ITEX). *Glob. Change Biol.* 3, 1–9. doi: 10.1098/rstb.2012.0481
- Holway, J. G., and Ward, R. T. (1963). Snow and meltwater effects in an area of Colorado alpine. *Am. Midl. Nat.* 69, 189–197. doi: 10.2307/2422853
- Hülber, K., Gottfried, M., Pauli, H., Winkler, M., and Grabherr, G. (2006). Phenological responses of snowbed species to snow removal dates in the Central Alps: implications for climate warming. *Arct. Antarct. Alp. Res.* 38, 99–103. doi: 10.1657/1523-0430(2006)038[0099:PROSST]2.0.CO;2
- Iversen, M., Bräthen, K. A., Yoccoz, N. G., and Ims, R. A. (2009). Predictors of plant phenology in a diverse high-latitude alpine landscape: growth forms and topography. *J. Veg. Sci.* 20, 903–915. doi: 10.1111/j.1654-1103.2009.01088.x
- Johnson, J. B., and Omland, K. S. (2004). Model selection in ecology and evolution. *Trends Ecol. Evol.* 19, 101–108. doi: 10.1016/j.tree.2003.10.013
- Jonas, T., Rixen, C., Sturm, M., and Stoeckli, V. (2008). How alpine plant growth is linked to snow cover and climate variability. *J. Geophys. Res.* 113:G03013. doi: 10.1029/2007JG000680
- Kammer, P. M., and Möhl, A. (2002). Factors controlling species richness in alpine plant communities: an assessment of the importance of stress and disturbance. *Arct. Antarct. Alp. Res.* 34, 398–407. doi: 10.1080/15230430.2002.12003510
- Keller, F., and Körner, C. (2003). The role of photoperiodism in alpine plant development. *Arct. Antarct. Alp. Res.* 35, 361–368. doi: 10.1657/1523-0430(2003)035[0361:TROPIA]2.0.CO;2

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2018.01140/full#supplementary-material>

- Kieltyk, P. (2017). Variation of vegetative and floral traits in the alpine plant *solidago minuta*: evidence for local optimum along an elevational gradient. *Alp. Bot.* 128, 47–57. doi: 10.1007/s00035-017-0197-7
- Körner, C. (1999). “Alpine plants: stressed or adapted?,” in *Physiological Plant Ecology: 39th Symposium of the British Ecological Society*, eds M. C. Press, J. D. Scholes, and M. G. Barker (Cambridge: Cambridge University Press), 297–311.
- Körner, C. (2003). *Alpine Plant Life: Functional Plant Ecology of High Mountain Ecosystems*. Berlin: Springer Science & Business Media. doi: 10.1007/978-3-642-18970-8
- Krömer, S. (1995). Respiration during photosynthesis. *Annu. Rev. Plant Biol.* 46, 45–70. doi: 10.1146/annurev.pp.46.060195.000401
- Kudo, G. (1991). Effects of snow-free period on the phenology of alpine plants inhabiting snow patches. *Arct. Alp. Res.* 23, 436–443. doi: 10.2307/1551685
- Kudo, G. (1992). Performance and phenology of alpine herbs along a snow-melting gradient. *Ecol. Res.* 7, 297–304. doi: 10.1007/BF02347098
- Kudo, G., and Hirao, A. S. (2006). Habitat-specific responses in the flowering phenology and seed set of alpine plants to climate variation: implications for global-change impacts. *Popul. Ecol.* 48, 49–58. doi: 10.1007/s10144-005-0242-z
- Kudo, G., Nordenhäll, U., and Molau, U. (1999). Effects of snowmelt timing on leaf traits, leaf production, and shoot growth of alpine plants: comparisons along a snowmelt gradient in northern Sweden. *Ecoscience* 6, 439–450. doi: 10.1080/11956860.1999.11682543
- Kueppers, L. M., Conlisk, E., Castanha, C., Moyes, A. B., Germino, M. J., Valpine, P., et al. (2017). Warming and provenance limit tree recruitment across and beyond the elevation range of subalpine forest. *Glob. Change Biol.* 23, 2383–2395. doi: 10.1111/gcb.13561
- Kullman, L. (2010). A richer, greener and smaller alpine world: review and projection of warming-induced plant cover change in the Swedish scandes. *Ambio* 39, 159–169. doi: 10.1007/s13280-010-0021-8
- Lembrechts, J. J., Lenoir, J., Nuñez, M. A., Pauchard, A., Geron, C., Bussé, G., et al. (2017). Microclimate variability in alpine ecosystems as stepping stones for non-native plant establishment above their current elevational limit. *Ecography* 41, 900–909. doi: 10.1111/ecog.03263
- Lenoir, J., Gégout, J. C., Marquet, P. A., De Ruffray, P., and Brisse, H. (2008). A significant upward shift in plant species optimum elevation during the 20th century. *Science* 320, 1768–1771. doi: 10.1126/science.1156831
- Lenth, R. V. (2016). Least-squares means: the R package lsmeans. *J. Stat. Softw.* 69, 1–33. doi: 10.18637/jss.v069.i01
- Litaor, M. I., Williams, M., and Seastedt, T. R. (2008). Topographic controls on snow distribution, soil moisture, and species diversity of herbaceous alpine vegetation, niwot ridge, Colorado. *J. Geophys. Res. Biogeosci.* 113:G02008. doi: 10.1029/2007JG000419
- May, D. E., and Webber, P. J. (1982). “Spatial and temporal variation of vegetation and its productivity on niwot ridge, Colorado,” in *Ecological Studies in the Colorado Alpine, a Festschrift for John W. Marr*, ed. J. Halfpenny (Denver, CO: University of Colorado), 35–62.
- McCune, B., and Grace, J. B. (2002). *Analysis of Ecological Communities*. Gleneden Beach, OR: MjM Software Design.
- Meloche, C. G., and Diggle, P. K. (2001). Preformation, architectural complexity, and developmental flexibility in *Acomastylis rossii* (Rosaceae). *Am. J. Bot.* 88, 980–991. doi: 10.2307/2657079
- Michalet, R., Schöb, C., Lortie, C. J., Brooker, R. W., and Callaway, R. M. (2014). Partitioning net interactions among plants along altitudinal gradients to study community responses to climate change. *Funct. Ecol.* 28, 75–86. doi: 10.1111/1365-2435.12136
- Mote, P. W., Li, S., Lettenmaier, D. P., Xiao, M., and Engel, R. (2018). Dramatic declines in snowpack in the western US. *Clim. Atmos. Sci.* 1:2. doi: 10.1038/s41612-018-0012-1
- Moyes, A. B., Castanha, C., Germino, M. J., and Kueppers, L. M. (2013). Warming and the dependence of limber pine (*Pinus flexilis*) establishment on summer soil moisture within and above its current elevation range. *Oecologia* 171, 271–282. doi: 10.1007/s00442-012-2410-0
- Moyes, A. B., Germino, M. J., and Kueppers, L. M. (2015). Moisture rivals temperature in limiting photosynthesis by trees establishing beyond their cold-edge range limit under ambient and warmed conditions. *New Phytol.* 207, 1005–1014. doi: 10.1111/nph.13422
- Negi, G. C. S., Rikhari, H. C., and Singh, S. P. (1992). Phenological features in relation to growth forms and biomass accumulation in an alpine meadow of the Central Himalaya. *Vegetatio* 101, 161–170.
- Nychka, D., Furrer, R., and Sain, S. (2015). *Package ‘Fields’*. Available at: <http://cran.r-project.org/web/packages/fields/fields.pdf>
- Oberbauer, S. F., and Billings, W. D. (1981). Drought tolerance and water use by plants along an alpine topographic gradient. *Oecologia* 50, 325–331. doi: 10.1007/BF00344971
- Odland, A., and Munkejord, H. K. (2008). Plants as indicators of snow layer duration in southern Norwegian mountains. *Ecol. Indic.* 8, 57–68. doi: 10.1016/j.ecolind.2006.12.005
- Opedal, ØH., Armbruster, W. S., and Graae, B. J. (2015). Linking small-scale topography with microclimate, plant species diversity and intra-specific trait variation in an alpine landscape. *Plant Ecol. Divers.* 8, 305–315. doi: 10.1080/17550874.2014.987330
- Pickering, C., Green, K., Barros, A. A., and Venn, S. (2014). A resurvey of late-lying snowpatches reveals changes in both species and functional composition across snowmelt zones. *Alp. Bot.* 124, 93–103. doi: 10.1007/s00035-014-0140-0
- Pielou, E. C. (1969). *An Introduction to Mathematical Ecology*. New York, NY: John Wiley & Sons.
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., and R Core Team. (2018). *nlme: Linear and Nonlinear Mixed Effects Models. R Package Version 3, 1–137*. Available at: <https://CRAN.R-project.org/package=nlme>
- R Core Team (2014). *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- Ren, F., Zhou, H. K., Zhao, X. Q., Han, F., Shi, L. N., Duan, J. C., et al. (2010). Influence of simulated warming using OTC on physiological-biochemical characteristics of *Elymus nutans* in alpine meadow on Qinghai-Tibetan plateau. *Acta Ecol. Sin.* 30, 166–171. doi: 10.1016/j.chnaes.2010.04.007
- Rosbakh, S., Leingärtner, A., Hoiss, B., Krauss, J., Steffan-Dewenter, I., and Poschlod, P. (2017). Contrasting effects of extreme drought and snowmelt patterns on mountain plants along an elevation gradient. *Front. plant sci.* 8:1478. doi: 10.3389/fpls.2017.01478
- Rundel, P. W. (2011). The diversity and biogeography of the alpine flora of the Sierra Nevada, California. *Madroño* 58, 153–184. doi: 10.3120/0024-9637-58.3.153
- Sage, R. F. (1994). Acclimation of photosynthesis to increasing atmospheric CO₂: the gas exchange perspective. *Photosynth. Res.* 39, 351–368. doi: 10.1007/BF00014591
- Shen, H., Klein, J. A., Zhao, X., and Tang, Y. (2009). Leaf photosynthesis and simulated carbon budget of *Gentiana straminea* from a decade-long warming experiment. *J. Plant Ecol.* 2, 207–216. doi: 10.1093/jpe/rtp025
- Sherwood, J. A., Debinski, D. M., Caragea, P. C., and Germino, M. J. (2017). Effects of experimentally reduced snowpack and passive warming on montane meadow plant phenology and floral resources. *Ecosphere* 8:e01745. doi: 10.1002/ecs2.1745
- Shi, F. S., Wu, Y., Wu, N., and Luo, P. (2010). Different growth and physiological responses to experimental warming of two dominant plant species *Elymus nutans* and *Potentilla anserina* in an alpine meadow of the eastern Tibetan Plateau. *Photosynthetica* 48, 437–445. doi: 10.1007/s11099-010-0058-8
- Sierra-Almeida, A., and Cavieres, L. A. (2010). Summer freezing resistance decreased in high-elevation plants exposed to experimental warming in the central Chilean andes. *Oecologia* 163, 267–276. doi: 10.1007/s00442-010-1592-6
- Smith, W. K., and Johnson, D. M. (2009). “Biophysical effects of altitude on plant gas exchange,” in *Perspectives in Biophysical Plant Ecophysiology. A Tribute to Park S. Nobel*, ed. W. K. Smith (Mexico: Universidad Nacional Autonoma de Mexico), 257–280.
- Stanton, M. L., Rejmanek, M., and Galen, C. (1994). Changes in vegetation and soil fertility along a predictable snowmelt gradient in the Mosquito Range Colorado, USA. *Arct. Alp. Res.* 26, 364–374. doi: 10.2307/1551798
- Steinbauer, M. J., Grytnes, J. A., Jurasinski, G., Kulonen, A., and Lenoir, J. (2018). Accelerated increase in plant species richness on mountain summits is linked to warming. *Nature* 556, 231–234. doi: 10.1038/s41586-018-0005-6
- Stocker, T. F., Dahe, Q., and Plattner, G. K. (2013). *Climate Change 2013: The Physical Science Basis. In: Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press.

- Studer, S., Appenzeller, C., and Defila, C. (2005). Inter-annual variability and decadal trends in alpine spring phenology: a multivariate analysis approach. *Clim. Change* 73, 395–414. doi: 10.1007/s10584-005-6886-z
- Taylor, R. V., and Seastedt, T. R. (1994). Short- and long-term patterns of soil moisture in Alpine tundra. *Arct. Alp. Res.* 26, 14–20. doi: 10.2307/1551871
- Theodose, T. A., Jaeger, C. H. I. I. I., Bowman, W. D., and Schardt, J. C. (1996). Uptake and allocation of ^{15}N in alpine plants: implications for the importance of competitive ability in predicting community structure in a stressful environment. *Oikos* 75, 59–66. doi: 10.2307/3546321
- Venn, S. E., and Morgan, J. W. (2007). Phytomass and phenology of three alpine snowpatch species across a natural snowmelt gradient. *Aust. J. Bot.* 55, 450–456. doi: 10.1071/BT06003
- Walker, D. A., Halfpenny, J. C., Walker, M. D., and Wessman, C. A. (1993). Long-term studies of snow-vegetation interactions. *Bioscience* 43, 287–301. doi: 10.2307/1312061
- Walker, M. D., Ingersoll, R. C., and Webber, P. J. (1995). Effects of internannual climate variation on phenology and growth of two alpine forbs. *Ecology* 76, 1067–1083. doi: 10.2307/1940916
- Walther, G. R., Post, E., Convey, P., Menzel, A., and Parmesan, C. (2002). Ecological responses to recent climate change. *Nature* 416, 389–395. doi: 10.1038/416389a
- Winkler, D. E., Chapin, K. J., and Kueppers, L. M. (2016a). Soil moisture mediates alpine life form and community productivity responses to warming. *Ecology* 97, 1553–1563. doi: 10.1890/15-1197.1
- Winkler, D. E., Amagai, Y., Huxman, T. E., Kaneko, M., and Kudo, G. (2016b). Seasonal dry-down rates and high stress tolerance promote bamboo invasion above and below treeline. *Plant Ecol.* 217, 1219–1234. doi: 10.1007/s11258-016-0649-y
- Wipf, S., and Rixen, C. (2010). A review of snow manipulation experiments in Arctic and alpine tundra ecosystems. *Polar Res.* 29, 95–109. doi: 10.1111/j.1751-8369.2010.00153.x
- Wipf, S., Rixen, C., and Mulder, C. P. H. (2006). Advanced snowmelt causes shift towards positive neighbour interactions in a subarctic tundra community. *Glob. Change Biol.* 12, 1496–1506. doi: 10.1111/j.1365-2486.2006.01185.x
- Wipf, S., Stoeckli, V., and Bebi, P. (2009). Winter climate change in alpine tundra: plant responses to changes in snow depth and snowmelt timing. *Clim. Change* 94, 105–121. doi: 10.1007/s10584-009-9546-x

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Winkler, Butz, Germino, Reinhardt and Kueppers. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.