

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/320676952>

Elucidating the regional spatiotemporal characteristics of Laramide deformation through low-temperature thermochronology

Conference Paper · October 2017

DOI: 10.1130/abs/2017AM-299560

CITATIONS

0

READS

55

6 authors, including:



Jacob Oliver Thacker

University of New Mexico

15 PUBLICATIONS 2 CITATIONS

[SEE PROFILE](#)



Karl Karlstrom

University of New Mexico

380 PUBLICATIONS 7,264 CITATIONS

[SEE PROFILE](#)



Jerome j Kendall

University of New Mexico

10 PUBLICATIONS 12 CITATIONS

[SEE PROFILE](#)



Shari Kelley

New Mexico Institute of Mining and Technology

12 PUBLICATIONS 53 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Elasticity Olivine-Wadsleyite [View project](#)



fore land deformation controls [View project](#)



22-25 October
Seattle, Washington, USA

THE GEOLOGICAL SOCIETY
 OF AMERICA®

27-3: ELUCIDATING THE REGIONAL SPATIOTEMPORAL CHARACTERISTICS OF LARAMIDE DEFORMATION THROUGH LOW-TEMPERATURE THERMOCHRONOLOGY



Sunday, 22 October 2017

08:35 AM - 08:50 AM

📍 *The Conference Center - Tahoma 4*

Several tectonic models involving different geodynamic mechanisms have been proposed to explain the tectonic evolution of the western United States and the formation of Laramide basement-involved arches ~90 to 45 Ma. Flat slab eastward subduction of the Farallon plate remains the most viable model to account for this enigmatic deformation up to 1000 km from the plate margin, though alternative models do persist in the literature. These include west-dipping subduction, lithospheric buckling, and terrane accretion. These diverse models can be tested through timing of deformation, spatial distribution and timing of magmatism, kinematic analysis of faults, and paleostress estimates. We focus here on timing of deformation, as each model should produce markedly different kinematic signatures (e.g. sweeping vs. dynamic deformation). In order to elucidate the spatiotemporal characteristics of Laramide deformation, we utilize apatite fission track (AFT) and apatite (U-Th)/He (AHe) thermochronology for their ability to determine rock cooling histories from ~120 °C (AFT) to ~40 °C (AHe). These methods are ideal for interpreting the timing of uplift and exhumation of Laramide arches, and many Laramide thermochronology studies have been completed. These data are now compiled, and identified gaps have been sampled in order to supplement the regional dataset at key locations. We model these data from many of the Laramide ranges with the program HeFTy to produce continuous time-temperature path models. Models are constrained and evaluated through integration of various geologic data, and are used to help interpret the onset, duration, and cooling rates of deformation for individual Laramide arches. An automated script is used to calculate these characteristics, allowing for quick and statistically meaningful results. Our goal is to then analyze this information geospatially to produce regional cross sections and maps at various time slices to show the spatiotemporal evolution of Laramide structures across the foreland. Through this regional analysis, the plausibility of proposed Laramide geodynamic mechanisms may then be kinematically assessed, and will provide key details for moving forward with understanding the orogenic evolution of western North America and the formation of Laramide arches.

Authors

Jacob O. Thacker

University of New Mexico

Karl E. Karlstrom

University of New Mexico

Jerry Kendall

University of New Mexico

Shari A. Kelley

New Mexico Tech

Ryan S. Crow

U.S. Geological Survey

Brandon Schmandt

University of New Mexico

Final Paper Number 27-3

View Related Events

Day: Sunday, 22 October 2017