

# Metal Reactivity in Laboratory Burned Wood from a Watershed Affected by Wildfires

Asifur Rahman,<sup>†</sup> Eliane El Hayek,<sup>‡</sup> Johanna M. Blake,<sup>§</sup> Rebecca J. Bixby,<sup>||</sup> Abdul-Mehdi Ali,<sup>⊥</sup> Michael Spilde,<sup>⊥</sup> Amanda A. Otieno,<sup>#</sup> Keely Miltenberger,<sup>⊥</sup> Cyrena Ridgeway,<sup>¶</sup> Kateryna Artyushkova,<sup>▽</sup> Viorel Atudorei,<sup>⊥</sup> and José M. Cerrato<sup>\*,†,||</sup>

<sup>†</sup>Department of Civil Engineering, MSC01 1070, University of New Mexico, Albuquerque, New Mexico 87131, United States

<sup>‡</sup>Department of Chemistry, MSC03 2060, University of New Mexico, Albuquerque, New Mexico 87131, United States

<sup>§</sup>U.S. Geological Survey, 6700 Edith Blvd. NE, Albuquerque, New Mexico 87113, United States

<sup>||</sup>Department of Biology and Museum of Southwestern Biology, MSC03 2020, University of New Mexico, Albuquerque, New Mexico 87131, United States

<sup>⊥</sup>Department of Earth and Planetary Sciences, MSC03 2040, University of New Mexico, Albuquerque, New Mexico 87131, United States


<sup>#</sup>Water Resources Program, MSC05 3110, University of New Mexico, Albuquerque, New Mexico 87131, United States

<sup>¶</sup>Department of Civil Engineering, New Mexico State University, Las Cruces, New Mexico 88001, United States

<sup>▽</sup>Department of Chemical and Biological Engineering, MSC01 1120, University of New Mexico, Albuquerque, New Mexico 87131, United States

## Supporting Information

**ABSTRACT:** We investigated interfacial processes affecting metal mobility by wood ash under laboratory-controlled conditions using aqueous chemistry, microscopy, and spectroscopy. The Valles Caldera National Preserve in New Mexico experiences catastrophic wildfires of devastating effects. Wood samples of Ponderosa Pine, Colorado Blue Spruce, and Quaking Aspen collected from this site were exposed to temperatures of 60, 350, and 550 °C. The 350 °C Pine ash had the highest content of Cu ( $4997 \pm 262 \text{ mg kg}^{-1}$ ), Cr ( $543 \pm 124 \text{ mg kg}^{-1}$ ), and labile dissolved organic carbon (DOC,  $11.3 \pm 0.28 \text{ mg L}^{-1}$ ). Sorption experiments were conducted by reacting 350 °C Pine, Spruce, and Aspen ashes separately with  $10 \mu\text{M}$  Cu(II) and Cr(VI) solutions. Up to a 94% decrease in Cu(II) concentration was observed in solution while Cr(VI) concentration showed a limited decrease (up to 13%) after 180 min of reaction. X-ray photoelectron spectroscopy (XPS) analyses detected increased association of Cu(II) on the near surface region of the reacted 350 °C Pine ash from the sorption experiments compared to the unreacted ash. The results suggest that dissolution and sorption processes should be considered to better understand the potential effects of metals transported by wood ash on water quality that have important implications for postfire recovery and response strategies.



## INTRODUCTION

Postfire storm events in watersheds can cause the transport of wood ash into nearby streams which has detrimental impacts on water quality. In the United States, the forests in the southwestern regions have seen increased occurrences of large intensity wildfires because of worldwide effects of climate change.<sup>1–3</sup> The Valles Caldera National Preserve (VALL) in north central New Mexico is an example of a site with frequent wildfire activity in recent years.<sup>4</sup> For example, the following two major wildfires have affected the VALL watershed since 2011: (1) the Thompson Ridge wildfire burned 23 965 acres in VALL in 2013,<sup>5</sup> and (2) the catastrophic Las Conchas, one of the largest in New Mexico history, burned over 156 000 acres

in the Jemez Mountains in 2011.<sup>6</sup> Postfire runoff of debris and ash in the Rio Grande river, following thunderstorms over the burned area, caused turbidity peaks of over 1000 NTU, sags in dissolved oxygen (DO), and fluctuations in pH (7.5 to 9).<sup>7</sup> Total concentrations of Al and Cu in the Rio Grande following the Las Conchas fire were above the aquatic life criterion for both metals.<sup>8</sup> Additionally, postfire runoff caused transport of organic matter and nutrients (6× background levels for NO<sub>3</sub>–

Received: January 28, 2018

Revised: May 6, 2018

Accepted: July 5, 2018

Published: July 18, 2018

N and 100× background levels for  $\text{PO}_4$ ) with debris and ash in the VALL, which substantially affected the quality of water supplied to nearby communities.<sup>7,9</sup> Information on the composition and reactivity of metals associated with wood ash is important to better understand the potential impacts on water quality.

While previous studies have characterized metals and organic matter in ash and soil,<sup>10–13</sup> the specific mechanisms controlling postfire metal mobilization remain poorly understood. Elevated concentrations of metals have been observed in sediments and surface water in fire-affected watersheds, for several months after the fire events.<sup>1,14–17</sup> Recent studies have linked water extractable organic matter (WEOM) from burned soil<sup>10,18</sup> and ash<sup>19</sup> to the formation of disinfection byproducts (DBPs).<sup>20,21</sup> Temperature is a particularly important factor that affects the elemental composition and mineralogy of wood ash.<sup>22–24</sup> For example, a previous study from our group identified the presence of Ca, Mg, Al, Fe, and Mn as metal bearing carbonate and oxide phases in wood ash burned at 550 °C.<sup>12</sup> Results from laboratory batch experiments suggested that these metal-bearing phases are readily water-soluble, but the readsorption of these metals to ash can occur in later times of the experiments.<sup>12</sup> Although this study provides valuable insights into the presence of metal bearing phases in ash and their potential effects on metal readsorption, the knowledge on the specific processes affecting metal dissolution and sorption in wood ash is still limited.

More mechanistic investigations have been reported in the literature related to reactivity of biochar, a material similar to wood ash in composition.<sup>25,26</sup> Biochar is a natural sorbent and is increasingly applied in environmental remediation of organic and inorganic contaminants.<sup>27–29</sup> The sorption of Cu(II) to organic functional groups of biochar in water can occur at pH below 7.<sup>30</sup> However, Cu-associated phases like azurite ( $\text{Cu}_3(\text{CO}_3)_2(\text{OH})_2$ ) and tenorite ( $\text{CuO}$ ) precipitate within the biochar surface at pH higher than 7.<sup>30</sup> Sorption between positively charged ions and negatively charged biochar is an effective mechanism for immobilizing metals in soil.<sup>29,31,32</sup> The immobilization of Cr(VI) through biochar sorption is significantly reduced at pH 5 and above.<sup>33–35</sup> Enhanced mobility of oxyanions like As<sup>36,37</sup> and Sb<sup>38</sup> has been observed in biochar-treated soil. Similar mechanistic studies are necessary to better understand sorption, precipitation, and dissolution reactions facilitated by wood ash that can affect postfire metal mobility.

The main objective of this study is to investigate the interfacial processes affecting wood ash reactions with water by integrating laboratory experiments, spectroscopy, microscopy, and aqueous chemistry methods. Soil and surface water chemistry from burned areas of VALL provide environmental context for the study. The release of metals and dissolved organic carbon (DOC) was assessed in batch experiments reacting laboratory burned wood ash with water. Additional experiments were conducted to investigate sorption processes that affect ash–metal interactions. The focus of this study is to identify interfacial physical–chemical processes that have not been extensively studied in the existing wildfire literature. The results from this investigation have relevant implications for the improvement of postfire response in affected watersheds.

## MATERIALS AND METHODS

**Field Sampling and Ash Preparation.** The East Fork Jemez River and the adjacent area were affected by the Las

Conchas and Thompson Ridge wildfires in the past. There is a lack of information about metals in soils, water, and wood from the VALL. Water and soil sampling were done to assess the current availability of metals in this fire affected watershed. Wood was collected from an unburned area to experiment with laboratory burn temperatures. Additional details about sampling methods are provided in the [Supporting Information](#). The description and coordinates of the sampling sites in Valles Caldera for wood, water, and soil samples are shown in [Figure S1](#). The vegetation in the VALL is dominated by different species of Pine, Spruce, Aspen, and Oak.<sup>39</sup> Wood samples of Ponderosa Pine, Colorado Blue Spruce, and Quaking Aspen were collected from higher elevation mixed coniferous forest areas which have a fire disturbance history from Las Conchas and other fires.<sup>12,40</sup> This area in VALL is also densely forested which undergoes prescribed burns periodically and account for approximately 25% of the precipitation volume in the Caldera.<sup>41</sup> From this point on, we will call these tree species simply Pine, Spruce, and Aspen. The collected wood samples were crushed using a wood chipper (Sun Joe CJ601E). The crushed samples were ground to fine powders using a Powdertec 3090 sample mill and then oven-dried at 60 °C for 48 h before burning. The ground and dried samples for each tree were mixed in a container to homogenize the samples before burning. The homogenized samples were burned in a laboratory muffle furnace at 350 °C (moderate burn) and at 550 °C (high burn) for 4 h to prepare ash. [Table S1](#) contains data on the mass of samples before and after burning.

### Acid Digestion and Solution Chemistry Analyses.

Wood samples were acid digested in triplicates ( $n = 3$ ) at 95 °C for 4 h using Aqua Regia [2 mL of  $\text{HNO}_3$  (67–70%) + 6 mL of HCl (34–37%), trace metal grade]. Following heating, acid extracts were diluted with 18 MΩ water to 50 mL. Processing of all aqueous samples (water, soil, and wood) for this study was done by filtering through a 0.45 μm filter, acidifying with 2%  $\text{HNO}_3$ , and refrigerating at 4 °C until further solution chemistry analyses. Inductively coupled plasma-optical emission spectrometry (ICP-OES, PerkinElmer Optima 5300DV) was used for detection of concentrations of major elements (Ca, Mg, and K). Minor or trace elements were analyzed using inductively coupled plasma-mass spectrometry (ICP-MS, PerkinElmer NexION 300D). Both the ICP-OES and ICP-MS analyzed an internal indium standard and were calibrated using a 5-point calibration curve. The quality of the results was ensured with proper quality control and quality assurance standards. The detection limits for the ICP-OES and ICP-MS for specific elements are shown in [Table S2](#). The DOC in these samples was measured using a Tekmar-Dohrmann Phoenix 8000 TOC Analyzer.

### Batch Experiments for DOC Concentration and Metal

**Dissolution.** For DOC dissolution experiments, batch reactors were operated in triplicates by reacting a 0.1 g sample of 60, 350, and 550 °C Pine, Spruce, and Aspen with 30 mL of 18 MΩ deionized water. Replicates ( $n = 3$ ) were sampled at 0, 4, 24, and 72 h and were analyzed using a Tekmar-Dohrmann Phoenix 8000 TOC Analyzer, following the 5310-C persulfate-ultraviolet (UV) method.<sup>42</sup> For metal dissolution experiments, 0.1 g samples of 350 and 550 °C Pine ash were reacted with 30 mL of 18 mΩ water. Samples were collected at 5 min, 15 min, 30 min, 1 h, 2 h, and 3 h, centrifuged at 3000 rpm (1660g) for 15 min, and processed for further ICP analyses.

**Batch Sorption Experiments of Cu(II) and Cr(VI) onto 350 °C Ash.** Batch sorption experiments in triplicates were

**Table 1. Elemental Content of Water (Sites 1 to 5) and Soil (Sites V1 to V3) Samples from the East Fork Jemez River within the Valle Grande Area in VALL<sup>a</sup>**

site	pH	alkalinity (mg L <sup>-1</sup> )	TOC (mg C L <sup>-1</sup> )	water elemental content (μg L <sup>-1</sup> )				
				Cu total	Cr total	Fe total	Zn total	Mn total
Site 1	7.17–8.44	30.5–42.0	0.80–2.00	2.81–16.7	BDL <sup>b</sup> –105	289–932	52.2–103	6.90–57.2
Site 2	7.11–8.06	31.0–38.3	1.60–10.8	5.44–37.4	BDL–8.20	465–2650	41.4–335	8.83–146
Site 3	6.55–7.66	24.1–43.8	4.00–10.3	7.93–23.4	BDL–48.4	38.2–677	57.4–352	1.58–39.2
Site 4	6.48–8.01	38.7–43.7	4.60–11.3	5.31–21.7	BDL–16.5	358–757	60.7–241	6.56–22.4
Site 5	6.48–7.09	33.8–40.9	3.20–11.0	BDL–25.2	BDL–11.5	331–948	72.8–135	BDL–44.4
soil sampling sites				soil elemental content (mg kg <sup>-1</sup> )				
				Cu total	Cr total	Fe total	Zn total	Mn total
Site V1				59.4–86.9	10.9–35.3	7800–9620	13.0–38.5	219–319
Site V2				71.3–89.9	9.25–20.4	4980–9850	10.2–31.4	50.8–194
Site V3				44.2–261	11.0–17.2	5050–7940	14.9–37.9	92.7–139
Site V4 (high burn)				7.16–17.5	5.81–13.3	109–221	19.5–35.2	260–347
Site V5 (low burn)				5.97–18.6	4.96–15.2	86.4–190	22.2–54.2	179–344
Site V6 (unburned)				7.97–13.9	5.31–11.3	102–184	30.8–39.4	217–454

<sup>a</sup>Aqueous and soil elemental content was measured with ICP-OES and ICP-MS. Sites V4 to V6 contain soil samples from the hill slope of the Sierra de Los Valles dome located near the headwaters of the river. Ranges of concentrations (minimum to maximum) for each site are shown.

<sup>b</sup>BDL = below detection limit.

conducted to investigate the effect of 350 °C Pine, Spruce, and Aspen ashes on mobilization of Cu(II) and Cr(VI) in water. We selected Cu(II) and Cr(VI) as examples of a cation that acts as a Lewis Acid (e.g., Cu) and an oxyanion (e.g., CrO<sub>4</sub><sup>2-</sup>) that could negatively impact surface waters. Additionally, elevated concentrations of these metals were found in surface water following wildfire events near VALL.<sup>8,43</sup> Before the sorption experiments, ash samples were characterized by measuring Brunauer–Emmett–Teller (BET) specific surface area, zeta potential, and C, H, N, and O elemental contents (wt %). Detailed descriptions of these methods are in the [Supporting Information](#). For the sorption experiments, stock solutions (1000 ppm) of Cr(IV) and Cu(II) were prepared by dissolving analytical grade (>99% purity) K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> and CuCl<sub>2</sub>·2H<sub>2</sub>O in 18 MΩ water. For the experiment, a 0.1 g sample of 350 °C Pine, Spruce, and Aspen ashes was reacted separately with 10 μM of Cu(II) and Cr(VI) stock solutions mixed in 50 mL of 18 MΩ water. The concentrations of Cu(II) and Cr(VI) were chosen to reflect maximum levels measured in water samples collected during storm events after the Cerro Grande fire.<sup>43</sup> The pH was adjusted to 7.0 ± 0.2 using 2% HNO<sub>3</sub>. Control experiments were done at pH 7.0 ± 0.2 for Cu(II) and Cr(VI). Samples were collected at 0, 4, 24, and 72 h, centrifuged at 6000 rpm (6640g) for 3 min, and processed for further solution chemistry analyses using ICP-MS.

**Solid Phase Analyses (SEM/EDX, EPMA, XRD, XPS).** Solid phase analyses were performed on the unreacted and reacted 350 °C Pine ash from the batch sorption experiments by applying X-ray photoelectron spectroscopy (XPS), scanning electron microscopy coupled to energy dispersive X-ray spectroscopy (SEM/EDX), electron probe microanalysis (EPMA), and X-ray diffraction (XRD). Additional descriptions of these methods are in the [Supporting Information](#).


**Statistical Analysis.** Univariate data analyses were performed using the statistical software R.<sup>44</sup> The statistical package in OriginPro<sup>45</sup> was used for principal component analysis (PCA). Due to the non-normality of the data, nonparametric tests for differences in acid extractable metal concentrations (log<sub>10</sub> transformed to reduce skewness in distribution) were performed to differentiate among 3 tree

species (Pine, Spruce, and Aspen) and among 3 temperatures (60, 350, and 550 °C). The Kruskal–Wallis test was performed to determine if the tree species and the temperatures differ significantly (defined as *p* < 0.05) with respect to metal concentrations (log<sub>10</sub> transformed). Wilcoxon rank sum test was used to do pairwise comparisons of all the samples to test for significant differences between tree species and temperatures (defined as *p* < 0.05). PCA was performed to better understand the correlations among ash samples in triplicates with respect to acid extractable metal (Al, Ca, Cr, Cu, Fe, K, Mg, Mn, Ni, Si, Sr, Zn) concentrations for all tree species.


## RESULTS AND DISCUSSION

**Water Quality and Sediment Data from Valles Caldera.** The presence of Cu, Cr, Fe, Zn, and Mn was observed in water from the wildfire affected East Fork Jemez River and in soils exposed to varying burn severities upslope from the East Fork Jemez River headwaters ([Table 1](#)). Maximum total concentrations of Cu (37.4 μg L<sup>-1</sup>) and Fe (2650 μg L<sup>-1</sup>) in the water samples collected from site 2 and Zn (352 μg L<sup>-1</sup>) from site 3 were above the USEPA standards for acute or chronic exposure values for aquatic life in freshwater ([Tables 1](#) and [S3](#)). Total Cr (105 μg L<sup>-1</sup>) in the water samples collected from site 1 exceeded the USEPA standards for acute or chronic exposure for both Cr(III) (74 μg L<sup>-1</sup>) and Cr(VI) (11 μg L<sup>-1</sup>). In a 2001 study by the New Mexico Environment Department, Al, pH, DO, and turbidity in the East Fork Jemez river were listed as exceeding the Total Maximum Daily Load, while metals, such as Cu, Cr, and Zn, were found to be below the detection limit,<sup>46,47</sup> lower than found in this study. Average concentrations of major and trace elements (mg kg<sup>-1</sup>) in nonanthropogenically affected soils in the US estimated by Burt et al.<sup>48</sup> ([Table S3](#)) were used to evaluate the metal concentrations in the soil samples collected along the river reaches. Iron (Fe) was the most abundant metal with concentrations ranging from 4980 to 9850 mg kg<sup>-1</sup>. Copper (Cu) concentration in the collected soil samples from sites V1, V2, and V3 ranges from 44.2 to 261 mg kg<sup>-1</sup>, higher than the average Cu concentration (24.7 mg kg<sup>-1</sup>) of nonanthropogenic affected soils in the US ([Table S3](#)).





**Figure 1.** Acid extractable concentrations (mean  $\pm$  SD) of 9 metals varied across different tree species of (a) Pine, (b) Spruce, and (c) Aspen. The major elements (Ca, Mg, and K) were predominant in all tree species at 60, 350, and 550 °C (Table S2). Ash produced at both 350 °C (moderate burn) and 550 °C (high burn) contained higher metal concentrations than in samples dried at 60 °C (unburned) for all species.




**Figure 2.** Concentrations of (a) Fe, Cu, and Zn and (b) Cr and Ni at 0, 4, 24, and 72 h in reaction of 550 °C Pine ash with 18 MΩ water. In (c) and (d), metal concentrations are shown for reaction of 350 °C Pine ash with 18 MΩ water. DOC concentration (mg carbon L<sup>-1</sup>) at 0, 4, 24, and 72 h ( $n = 3$ ) is shown in (e) for 350 and 550 °C ash samples and in (f) for 60 °C crushed wood samples.

Concentrations of Cu, Cr, and Fe were found lower in soils collected from high, low, and unburned sites of V4 to V6 (Table 1) compared to soil in the channel and banks of the East Fork Jemez River. The higher concentrations of metals in water and soil in a river located near a burned watershed, such as the East Fork Jemez, can increase from the addition of ash and debris by stormwater runoff. For example, after the Cerro Grande fire in 2000, elevated concentrations of total Cu (610  $\mu\text{g L}^{-1}$ ) and total Cr (510  $\mu\text{g L}^{-1}$ ) were measured in postfire runoff water samples collected from the burned watersheds near Guaje Canyon which is close to the Los Alamos National Laboratory.<sup>43</sup> Additional laboratory experiments were done to determine the concentrations of metals in oven-dried wood and ash and assess the reactivity of wood ash upon reaction with 18 MΩ water.

**Acid Extractable Metal Content in Wood Exposed to 60, 350, and 550 °C.** We compared the acid extractable metal contents in 9 samples (Pine, Spruce, and Aspen at 60,

350, and 550 °C, Figure 1, Table S4). Median metal concentrations at 350 °C (moderate burn) and 550 °C (high burn) for all tree species (e.g., Pine, Spruce, and Aspen) were significantly higher ( $p < 0.05$ , Tables S5 and S6) compared to oven-dried wood at 60 °C (unburned). The acid extractable metal concentrations for all samples at 60, 350, and 550 °C are shown in Table S4. Pairwise comparisons suggest that metal contents in oven-dried wood at 60 °C for Pine, Spruce, and Aspen were not significantly different from each other ( $p > 0.05$ , Tables S5 and S6). Among the ash samples at 350 and 550 °C, Pine and Aspen were significantly different ( $p < 0.05$ , Table S6) from each other with respect to acid extractable metal concentrations. Principal component analysis (PCA, Figure S2) suggests that 350 and 550 °C Pine ash samples were associated with high concentrations of most of the metals (Cu, Cr, Si, Ni, Fe, K, and Mg). High concentrations of Ca, Sr, and Zn were associated with 350 and 550 °C Aspen ash samples, while 350 and 550 °C Spruce



**Figure 3.** Results from the metal sorption experiments ( $n = 3$ , sampling interval = 5 min, 15 min, 30 min, 1 h, 2 h, and 3 h) conducted by reacting 10  $\mu\text{M}$  of (a) Cu(II) and (b) Cr(VI) separately in a solution containing 0.1 g of 350  $^{\circ}\text{C}$  Pine ash with 50 mL of 18 M $\Omega$  water. Note that the control experiments were conducted by reacting 10  $\mu\text{M}$  of Cu(II) and Cr(VI) in 18 M $\Omega$  water without ash.

ash samples were associated with high concentrations of Mn, Al, and Fe.


Major elements (Ca, Mg, K) were found to be predominant in oven-dried wood (60  $^{\circ}\text{C}$ ) and in ash samples (350 and 550  $^{\circ}\text{C}$ ) for all tree species (Table S4), consistent with findings for wood<sup>49</sup> and wood ash<sup>12,50</sup> from previous studies. Pine ash showed higher concentrations with increasing temperatures for major elements (Ca, Mg, and K) and for heavy metals such as Al, Fe, Mn, and Ni (Table S4), consistent with findings from a previous study conducted on Lodgepole Pine.<sup>51</sup> The concentration of Cu in Pine ash ( $4997 \pm 262 \text{ mg kg}^{-1}$  at 350  $^{\circ}\text{C}$  and  $2765 \pm 302 \text{ mg kg}^{-1}$  at 550  $^{\circ}\text{C}$ ) was higher than previously reported values for Pine ash.<sup>12,51</sup> Due to the dominance of Pine tree species in the forests of the western United States, much of the existing literature has focused on the metal and DOC composition of ash produced from different species of Pine (e.g., Ponderosa and Lodgepole).<sup>12,18,52–54</sup> We conducted additional experiments with 350  $^{\circ}\text{C}$  Pine, Spruce, and Aspen ashes to assess the release of dissolved organic carbon and other metals over time.

**Metal and DOC Leachates from Ash (350 and 550  $^{\circ}\text{C}$ ) Reacted with Water.** Metal leaching experiments were conducted to observe the dissolution of selected metals (Cr, Ni, Fe, Cu, and Zn) in reaction with 350 and 550  $^{\circ}\text{C}$  Pine ash in 18 M $\Omega$  water (Figure 2a–d). Dissolution of metal bearing phases caused a rapid increase in the pH of water (Figure S3), measured at  $10.0 \pm 0.5$  for the duration of the experiment. Less than 3% by mass [determined in acid extractable analysis (Table S4)] of Cr, Ni, Fe, Cu, and Zn was released in solution after 72 h of reaction. Metals like Cu, Fe, Zn, and Ni showed an initial release followed by a decrease in metal concentration over time for both the 350 and 550  $^{\circ}\text{C}$  ash samples (Figure 2a–d). Limited fluctuations in the concentration of Cr released in solution were observed over time (Figure 2b,d). The high pH and alkalinity in these experiments (Figure S3) are likely due to the dissolution of metal bearing carbonate and oxide phases, such as calcite, quartz, and whewellite [ $\text{Ca}(\text{C}_2\text{O}_4) \cdot \text{H}_2\text{O}$ ], which were identified by XRD analysis in the unreacted 350  $^{\circ}\text{C}$  Pine ash (Figure S4). A previous study from our group also identified the presence of calcite and other metal bearing phases in ash burned at 550  $^{\circ}\text{C}$ .<sup>12</sup>

The DOC concentration decreased with increasing temperature in reaction of 60, 350, and 550  $^{\circ}\text{C}$  samples with 18 M $\Omega$

water. The 60  $^{\circ}\text{C}$  (unburned) samples released higher DOC concentrations (110.7 to 338 mg carbon  $\text{L}^{-1}$ ) compared to 350  $^{\circ}\text{C}$  (4.25 to 11.3 mg carbon  $\text{L}^{-1}$ ) and 550  $^{\circ}\text{C}$  ash samples (1.27 to 2.77 mg carbon  $\text{L}^{-1}$ ) (Figure 2e,f). In this study, different sections (i.e., leaves, twigs, needles) of the collected tree species were crushed and ground to powders and homogenized before conducting the experiments. This process could have contributed to the high DOC release from the unburned samples. Additionally, a variable release of DOC could result from contributions from different tree sections, obtaining DOC concentrations that are higher than the ones reported in this study.<sup>55,56</sup> The decrease in DOC concentration from 350 to 550  $^{\circ}\text{C}$  ash suggests that a greater loss of organic matter occurs at a higher burning temperature, consistent with findings from previous studies.<sup>10,12</sup> The range of DOC concentrations (1.27 to 11.3 mg carbon  $\text{L}^{-1}$ ) measured for 350 and 550  $^{\circ}\text{C}$  ash samples in this study are consistent with those reported in previous studies from field<sup>15,18</sup> and laboratory studies.<sup>10</sup> Previously, Wang et al.<sup>54</sup> observed decreased reactivity of the ash in forming DBPs such as trihalomethane (THM) and chloral hydrate (CHD) with increasing temperature from 50 to 400  $^{\circ}\text{C}$ . The DOC loss at higher temperatures for ash in this study is comparable to that observed in laboratory heated soil by Cawley et al.<sup>10</sup> Thus, the temperature dependent variability for both ash and soil can have implications in terms of variable loading of DOM and DBP precursors from different burn conditions associated with wildfires and prescribed fires, as suggested by others.<sup>10,54</sup>

**Sorption to 350  $^{\circ}\text{C}$  Ash.** We further explored the sorption of Cu(II) (a cationic metal) and Cr(VI) (an oxyanion) to 350  $^{\circ}\text{C}$  ash at pH  $7.0 \pm 0.2$  in batch sorption experiments. Zeta potential measurements for the 350  $^{\circ}\text{C}$  Pine ash showed increasingly negative surface charge with increasing solution pH (Figure S5), similar to another carbon-based material like biochar.<sup>57,58</sup> The surface area of Pine ash increased from 36.9  $\text{m}^2/\text{g}$  at 350  $^{\circ}\text{C}$  to 294.4  $\text{m}^2/\text{g}$  at 550  $^{\circ}\text{C}$  (Table S7). Previously, de Mendonça et al.<sup>57</sup> reported an increase in surface area due to the creation of micropores for biochar burned at 400 and 600  $^{\circ}\text{C}$ . A higher loss of C, H, and O occurred in ash with increasing temperature (Table S8). Lower H/C ratios in 350 and 550  $^{\circ}\text{C}$  can be a measure of a higher degree of aromaticity in the ash samples, as suggested by previous studies on biochar.<sup>58,59</sup> Higher O/C and (O + N)/C



**Figure 4.** Cu 2p high resolution spectra for the (a) reacted 350 °C Pine ash and (b) unreacted 350 °C Pine ash. (c) Percentages of different oxidation states in the Cu 2p spectra for the reacted ash determined by using reference Cu 2p spectra for CuO, CuCO<sub>3</sub>, Cu<sub>2</sub>O, and Cu metal.

ratios, for the 350 and 550 °C ash samples, are indicators of increased hydrophilicity and polarity.<sup>58–60</sup> We observed more than an 80% decrease initially in Cu(II) concentration in the solution reacting with 350 °C Pine ash (Figure 3a), as indicated by measured Cu(II) concentration ( $103.6 \pm 3.1 \mu\text{g L}^{-1}$ ) after 5 min of reaction and up to a 92% decrease after 180 min of reaction. Cu(II) concentration in the control decreased only 5%, to  $601.2 \pm 53.8 \mu\text{g L}^{-1}$  after 180 min of reaction from the initially added concentration of  $635 \mu\text{g L}^{-1}$ . In experiments with Cr(VI), a low decrease in Cr(VI) concentration was observed, as the measured concentration after 180 min was  $451.7 \pm 7.8 \mu\text{g L}^{-1}$ , representing only a 13% decrease from the initially added Cr(VI) of  $520 \mu\text{g L}^{-1}$  (Figure 3b). Cr(VI) showed a negligible decrease in concentration in the Cr(VI)–control experiment (Figure 3b). Similar results were observed for both Cu(II) and Cr(VI) in the sorption experiments with 350 °C Spruce and Aspen ashes (Figure S6). These results suggest that burning caused similar effects on metal reactivity of all three ashes (Pine, Spruce, and Aspen).

The different response of Cu(II) and Cr(VI) concentration during the sorption experiments suggests that high Cu(II) association to ash occurred due to possible electrostatic attraction of the positively charged Cu(II) to the negatively charged ash surface. The effect of sorption capacity of the carbonate phases in wood ash in reacting with cations such as Ca<sup>2+</sup>, Mg<sup>2+</sup>, Al<sup>3+</sup>, Mn<sup>2+</sup>, Fe<sup>2+</sup>, Pb<sup>2+</sup>, Cu<sup>2+</sup>, Zn<sup>2+</sup>, and Cd<sup>2+</sup> has been discussed in other studies.<sup>12,13</sup> This is a relevant property of wood ash that should be considered when investigating the persistence of metals such as Cu, Pb, Ni, Fe, and Zn associated with ash and debris in wildfire affected watersheds, as reported in several postfire investigations.<sup>61–63</sup> The low sorption of Cr(VI) observed in this study is consistent with other studies reporting low sorption rates for As(V), Cr(VI), and Se(VI) to carbonaceous materials at pH 5.0 and above.<sup>64–66</sup> For example,

a recent study by Alam et al. observed over 90% removal of Cd(II) and below 20% removal for Se(VI) at pH 6.0 and above, using biochar as an adsorbent.<sup>64</sup> At the experimental pH of  $7.0 \pm 0.2$  used in this study, Cr(VI) is expected to exist in the solution as stable oxyanion forms (e.g., CrO<sub>4</sub><sup>2−</sup>, HCrO<sub>4</sub><sup>−</sup>).<sup>67</sup> Therefore, the electrostatic repulsion between the negatively charged ash surface and Cr(VI) oxyanions can account for the low decrease in Cr(VI) concentration in solution. The association of Cu(II) in the unreacted and reacted 350 °C Pine ash solids was further analyzed using microscopy and spectroscopy.

**Solid Phase Analyses of Unreacted and Reacted 350 °C Pine Ash.** SEM analysis detected the presence of Cu on 350 °C Pine ash before and after exposure to batch experiments (Figure S7). For example, EDS spectra of a Cu grain showed 69.51 wt % of Cu for the unreacted ash and 63.55 wt % of Cu for the reacted ash. SEM/EDS results confirmed the high Cu concentration ( $4997 \pm 262 \text{ mg kg}^{-1}$ ) measured from the acid extraction analyses. Electron microprobe mapping detected a low level of Cu (0.012%) associated with the Ca<sup>2+</sup> minerals in the reacted ash (Figure S8), while it was below the detection limit for the unreacted ash. The predominant form of the Ca<sup>2+</sup> mineral is most likely calcite (CaCO<sub>3</sub>), given the presence of 76–78 wt % of calcite in the reacted sample (Figure S4). Given that Ca is a macronutrient in plants, Ca<sup>2+</sup> minerals in the form of CaO and CaCO<sub>3</sub> are abundant across a variety of plant cells.<sup>68,69</sup> XRD analyses on reacted and unreacted samples indicated presence of quartz and calcite (CaCO<sub>3</sub>) as predominant mineral phases (Figure S4). While microscopy analyses identified the presence of Cu, it was challenging to obtain specific information about the association of Cu on the reacted ash surface from these analyses. Thus, additional analyses using XPS were done to measure the signal of Cu 2p on the “near-surface” region to

identify the possible association of Cu to ash after reaction in batch sorption experiments.

Results from the XPS survey scan revealed that 0.11% Cu 2p was present in the reacted ash, suggesting that Cu is associated at the top 5–10 nm of the ash “near surface” region (Table S9). However, the Cu 2p % for the unreacted ash was below the detection limit (Table S9). Thus, the high resolution Cu 2p peak obtained for the unreacted ash was noisy and could not be used for curve fitting analyses (Figure 4). Curve fitting of high resolution XPS Cu 2p spectra obtained for the reacted sample was conducted using reference spectra for CuO, CuCO<sub>3</sub>, Cu<sub>2</sub>O, and Cu metal as indicated in the Materials and Methods section. Curve fitting analysis suggests that the main species of Cu present on the reacted ash are Cu(II) in the form of CuO (64.2%) and CuCO<sub>3</sub> (18.6%) and Cu(I) in the form of Cu<sub>2</sub>O (17.3%) (Figure 4). The presence of Cu(I) on the reacted ash surface suggests a possible reduction of some of the Cu(II) to Cu(I), as suggested by a previous study by Bogusz et al.<sup>60</sup>

The association of Cu on the reacted 350 °C Pine ash near surface region suggests a likely surface controlled process involved in the removal of Cu(II) in the sorption experiments. Curve fitting analysis of C 1s high resolution spectra showed an increase in the percentages of C\*–CO<sub>x</sub>, C=O, and C–OH bonds in the reacted ash (Figure S9). The presence of surface functional groups (–C=O, –COOH) in ash can act as negatively charged binding sites for positively charged cations. The increased polarity (higher O/C and (O + N)/C ratios) in the ash samples measured by the C, H, N, and O analyses (Table S8) is consistent with the increase in functional groups, identified by XPS. The removal of cationic metals such as Cu<sup>2+</sup> and Cd<sup>2+</sup> through associations with surface functional groups of biochar has been discussed in the literature.<sup>30,60,70</sup> These properties are also relevant to better understand the effect of burned soil and ash on postfire mobilization of heavy metals.

**Environmental Implications.** The results from this investigation indicate how metal and DOC content associated with ash burned at different temperatures (350 and 550 °C) can differ for Pine, Spruce, and Aspen. This outcome may have relevant implications when considering the wide variation in vegetation across large watersheds when assessing response to wildfire events. Ash burned at 350 °C had higher DOC concentration in water compared to ash burned at 550 °C. This observed increase may have important implications in terms of increased DOC fluxes in postfire watersheds from moderately burned ash and soil reported in previous studies.<sup>15,71</sup> The batch experiments conducted in 18 MΩ water indicate that metals such as Cr, Ni, Fe, Cu, and Zn were dissolved in the initial stages of the experiment, followed by a decrease in concentration over the duration of the experiment. This observation is consistent with a previous study suggesting that metal (Ca, Mg, Al, Fe, and Mn) dissolution occurred in initial stages of the batch experiments conducted with ash from Pine, Aspen, and Spruce trees from the Caldera, followed by reassociation of these metals to ash over time.<sup>12</sup> However, a new finding was obtained from the sorption experiments conducted in this study which indicate that up to 94% of Cu is removed from solution after 180 min of reaction due to association of this metal in the 350 °C Pine ash surface. A similar behavior is expected for other positive cations such as Ca<sup>2+</sup>, Mg<sup>2+</sup>, Zn<sup>2+</sup>, Al<sup>3+</sup>, Fe<sup>2+</sup>, Cd<sup>2+</sup>, and Pb<sup>2+</sup> among others, as suggested in other studies related to wood ash<sup>12,13</sup> and biochar<sup>29,72</sup> reactivity. The integration of results from the

metal dissolution and sorption experiments in this study provide novel insights about postfire mobilization of cationic metals in burned watersheds. However, oxyanions such as Cr(VI) are expected to have limited association to ash in natural pH conditions and are likely to have higher mobility in watersheds affected by wildfires. Future experiments are necessary to study metal reactivity in wood ash in dynamic flow conditions which allow interactions between ash and sediments in water. This study provides relevant insights on water quality that could be considered for postfire response and recovery strategies by local authorities.

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.est.8b00530.

Additional materials and methods; mass of wood samples; detection limits for analyses; maximum contaminant level and different standards for exposure limits; acid extractable elemental content for wood samples; Kruskal Wallis test; Wilcoxon rank sum test; specific surface areas and elemental composition and ratios of samples; atomic content of Pine ash; map of sampling locations; principal component analysis; pH and alkalinity; XRD patterns; zeta potential measurements; results from the metal and batch sorption experiments; microprobe mapping; XPS high resolution C 1s spectra and percent compositions; XPS high-resolution Cu 2p spectra and binding energies (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Author

\*E-mail: [jcerrato@unm.edu](mailto:jcerrato@unm.edu); telephone: (001) (505) 277-0870; fax: (001) (505) 277-1988.

### ORCID

Kateryna Artyushkova: 0000-0002-2611-0422

José M. Cerrato: 0000-0002-2473-6376

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

Thanks to Dr. Robert Parmenter for scientific advice and support for this study. Funding for this research was provided by the Oak Ridge Associated Universities (ORAU) Program Ralph E. Powe Jr. Junior Faculty Enhancement Award, the New Mexico Water Resources Research Institute, the National Science Foundation under New Mexico EPSCoR (Grant Number #IIA-1301346), and CREST (Grant Number 1345169). Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the views of the National Science Foundation. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

## ■ REFERENCES

- (1) Bladon, K. D.; Emelko, M. B.; Silins, U.; Stone, M. Wildfire and the Future of Water Supply. *Environ. Sci. Technol.* **2014**, *48* (16), 8936–8943.
- (2) Spracklen, D. V.; Mickley, L. J.; Logan, J. A.; Hudman, R. C.; Yevich, R.; Flannigan, M. D.; Westerling, A. L. Impacts of climate change from 2000 to 2050 on wildfire activity and carbonaceous



aerosol concentrations in the western United States. *J. Geophys. Res.* **2009**, *114*, 2156–2202.

(3) Westerling, A. L.; Hidalgo, H. G.; Cayan, D. R.; Swetnam, T. W. Warming and Earlier Spring Increase Western U.S. Forest Wildfire Activity. *Science* **2006**, *313* (5789), 940–943.

(4) *Fire Regimes of Montane Grasslands of the Valles Caldera National Preserve, New Mexico*; Highlights of JFSP: 06-3-1-27; Joint Fire Science Program (JFSP): Boise, ID, 2011; <http://digitalcommons.unl.edu/cgi/viewcontent.cgi?article=1082&context=jfस्पresearch>.

(5) *Distribution and Transport of Pyrogenic Black Carbon in Soils Affected by Wildfires, Valles Caldera, New Mexico, with Implications for Contaminant Transport*; Report Number: 365; New Mexico Water Resources Research Institute (NMWRI): Las Cruces, NM, 2015; <https://nmwri.nmsu.edu/wp-content/uploads/2015/technical-reports/tr365.pdf>.

(6) Parmenter, R. R.; Oertel, R. W.; Compton, T. S.; Kindschuh, S.; Peyton, M.; Meyer, W.; Caldwell, C.; Jacobi, G. Z.; Myers, O.; Zeigler, M. Fire and floods in the Valles Caldera National Preserve, New Mexico: The 2011 Las Conchas Fire impacts on montane species diversity and food webs. In *97th ESA Annual Convention*, Portland, Oregon, USA, 2012.

(7) Dahm, C. N.; Candelaria-Ley, R. I.; Reale, C. S.; Reale, J. K.; Van Horn, D. J. Extreme water quality degradation following a catastrophic forest fire. *Freshwater Biol.* **2015**, *60* (12), 2584–2599.

(8) *2011 Las Conchas Fire Impacts to Water Quality in the Rio Grande*; Surface Water Quality Bureau, New Mexico Environment Department: Santa Fe, NM, 2011; <https://www.env.nm.gov/swqb/Wildfire/6.LasConchas-ImpactsToRioGrande2011.pdf>.

(9) Sherson, L. R.; Van Horn, D. J.; Gomez-Velez, J. D.; Crossey, L. J.; Dahm, C. N. Nutrient dynamics in an alpine headwater stream: use of continuous water quality sensors to examine responses to wildfire and precipitation events. *Hydrol. Process.* **2015**, *29* (14), 3193–3207.

(10) Cawley, K. M.; Hohner, A. K.; Podgorski, D. C.; Cooper, W. T.; Korak, J. A.; Rosario-Ortiz, F. L. Molecular and spectroscopic characterization of water extractable organic matter from thermally altered soils reveal insight into disinfection byproduct precursors. *Environ. Sci. Technol.* **2017**, *51* (2), 771–779.

(11) Steenari, B.-M.; Karlsson, L.-G.; Lindqvist, O. Evaluation of the leaching characteristics of wood ash and the influence of ash agglomeration. *Biomass Bioenergy* **1999**, *16* (2), 119–136.

(12) Cerrato, J. M.; Blake, J. M.; Hirani, C.; Clark, A. L.; Ali, A.-M. S.; Artyushkova, K.; Peterson, E.; Bixby, R. J. Wildfires and water chemistry: effect of metals associated with wood ash. *Environ. Sci. Process. Impacts.* **2016**, *18* (8), 1078–1089.

(13) Chirenje, T.; Ma, L. Q.; Lu, L. Retention of Cd, Cu, Pb and Zn by wood ash, lime and fume dust. *Water, Air, Soil Pollut.* **2006**, *171* (1), 301–314.

(14) Costa, M. R.; Calvão, A. R.; Aranha, J. Linking wildfire effects on soil and water chemistry of the Marão River watershed, Portugal, and biomass changes detected from Landsat imagery. *Appl. Geochem.* **2014**, *44*, 93–102.

(15) Emelko, M. B.; Silins, U.; Bladon, K. D.; Stone, M. Implications of land disturbance on drinking water treatability in a changing climate: demonstrating the need for “source water supply and protection” strategies. *Water Res.* **2011**, *45* (2), 461–472.

(16) Smith, H. G.; Sheridan, G. J.; Lane, P. N.; Nyman, P.; Haydon, S. Wildfire effects on water quality in forest catchments: a review with implications for water supply. *J. Hydrol.* **2011**, *396* (1), 170–192.

(17) Biswas, A.; Blum, J. D.; Klaue, B.; Keeler, G. J. Release of mercury from Rocky Mountain forest fires. *Global Biogeochem. Cycles* **2007**, *21* (1), GB1002.

(18) Hohner, A. K.; Cawley, K.; Oropeza, J.; Summers, R. S.; Rosario-Ortiz, F. L. Drinking water treatment response following a Colorado wildfire. *Water Res.* **2016**, *105*, 187–198.

(19) Wang, J.-J.; Dahlgren, R. A.; Erşan, M. S.; Karanfil, T.; Chow, A. T. Wildfire altering terrestrial precursors of disinfection byproducts in forest detritus. *Environ. Sci. Technol.* **2015**, *49* (10), 5921–5929.

(20) Bond, T.; Huang, J.; Templeton, M. R.; Graham, N. Occurrence and control of nitrogenous disinfection by-products in drinking water—a review. *Water Res.* **2011**, *45* (15), 4341–4354.

(21) Bagthoth, S.; Sharma, S.; Amy, G. Tracking natural organic matter (NOM) in a drinking water treatment plant using fluorescence excitation–emission matrices and PARAFAC. *Water Res.* **2011**, *45* (2), 797–809.

(22) Bodí, M. B.; Martín, D. A.; Balfour, V. N.; Santín, C.; Doerr, S. H.; Pereira, P.; Cerdà, A.; Mataix-Solera, J. Wildland fire ash: production, composition and eco-hydro-geomorphic effects. *Earth-Sci. Rev.* **2014**, *130*, 103–127.

(23) Ulery, A.; Graham, R.; Amrhein, C. Wood-ash composition and soil pH following intense burning. *Soil Sci.* **1993**, *156* (5), 358–364.

(24) Misra, M. K.; Ragland, K. W.; Baker, A. J. Wood ash composition as a function of furnace temperature. *Biomass Bioenergy* **1993**, *4* (2), 103–116.

(25) Jones, D. L.; Quilliam, R. Metal contaminated biochar and wood ash negatively affect plant growth and soil quality after land application. *J. Hazard. Mater.* **2014**, *276*, 362–370.

(26) Lucchini, P.; Quilliam, R.; DeLuca, T. H.; Vamerli, T.; Jones, D. L. Increased bioavailability of metals in two contrasting agricultural soils treated with waste wood-derived biochar and ash. *Environ. Sci. Pollut. Res.* **2014**, *21* (5), 3230–3240.

(27) Saeed, A.; Akhter, M. W.; Iqbal, M. Removal and recovery of heavy metals from aqueous solution using papaya wood as a new biosorbent. *Sep. Purif. Technol.* **2005**, *45* (1), 25–31.

(28) Yang, X.-B.; Ying, G.-G.; Peng, P.-A.; Wang, L.; Zhao, J.-L.; Zhang, L.-J.; Yuan, P.; He, H.-P. Influence of biochars on plant uptake and dissipation of two pesticides in an agricultural soil. *J. Agric. Food Chem.* **2010**, *58* (13), 7915–7921.

(29) Ahmad, M.; Rajapaksha, A. U.; Lim, J. E.; Zhang, M.; Bolan, N.; Mohan, D.; Vithanage, M.; Lee, S. S.; Ok, Y. S. Biochar as a sorbent for contaminant management in soil and water: a review. *Chemosphere* **2014**, *99*, 19–33.

(30) Ippolito, J.; Strawn, D.; Scheckel, K.; Novak, J.; Ahmedna, M.; Niandou, M. Macroscopic and molecular investigations of copper sorption by a steam-activated biochar. *J. Environ. Qual.* **2012**, *41* (4), 1150–1156.

(31) Uchimiya, M.; Wartelle, L. H.; Klasson, K. T.; Fortier, C. A.; Lima, I. M. Influence of pyrolysis temperature on biochar property and function as a heavy metal sorbent in soil. *J. Agric. Food Chem.* **2011**, *59* (6), 2501–2510.

(32) Beesley, L.; Marmiroli, M. The immobilisation and retention of soluble arsenic, cadmium and zinc by biochar. *Environ. Pollut.* **2011**, *159* (2), 474–480.

(33) Hsu, N.-H.; Wang, S.-L.; Lin, Y.-C.; Sheng, G. D.; Lee, J.-F. Reduction of Cr (VI) by crop-residue-derived black carbon. *Environ. Sci. Technol.* **2009**, *43* (23), 8801–8806.

(34) Hsu, N.-H.; Wang, S.-L.; Liao, Y.-H.; Huang, S.-T.; Tzou, Y.-M.; Huang, Y.-M. Removal of hexavalent chromium from acidic aqueous solutions using rice straw-derived carbon. *J. Hazard. Mater.* **2009**, *171* (1), 1066–1070.

(35) Dong, X.; Ma, L. Q.; Li, Y. Characteristics and mechanisms of hexavalent chromium removal by biochar from sugar beet tailing. *J. Hazard. Mater.* **2011**, *190* (1), 909–915.

(36) Park, J. H.; Lamb, D.; Paneerselvam, P.; Choppala, G.; Bolan, N.; Chung, J.-W. Role of organic amendments on enhanced bioremediation of heavy metal (loid) contaminated soils. *J. Hazard. Mater.* **2011**, *185* (2), 549–574.

(37) Zhang, X.; Wang, H.; He, L.; Lu, K.; Sarmah, A.; Li, J.; Bolan, N. S.; Pei, J.; Huang, H. Using biochar for remediation of soils contaminated with heavy metals and organic pollutants. *Environ. Sci. Pollut. Res.* **2013**, *20* (12), 8472–8483.

(38) Uchimiya, M.; Bannon, D. I.; Wartelle, L. H.; Lima, I. M.; Klasson, K. T. Lead retention by broiler litter biochars in small arms range soil: impact of pyrolysis temperature. *J. Agric. Food Chem.* **2012**, *60* (20), 5035–5044.

(39) *A Vegetation Map of Valles Caldera National Preserve, New Mexico*; Highlights of 06-GTR-302; Natural Heritage: Albuquerque,



NM, 2006; <https://nhnm.unm.edu/sites/default/files/nonsensitive/publications/nhnm/U06MUL01NMUS.pdf>.

(40) A vegetation survey and preliminary ecological assessment of Valles Caldera National Preserve; Highlights of 03-GTR-272; Natural Heritage: Albuquerque, NM, 2003; [https://www.fs.usda.gov/Internet/FSE\\_DOCUMENTS/stelprd5384299.pdf](https://www.fs.usda.gov/Internet/FSE_DOCUMENTS/stelprd5384299.pdf).

(41) An Investigation into the Potential Impacts of Watershed Restoration and Wildfire on Water Yields and Water Supply Resilience in the Rio Grande Water Fund Project Area; Middle Rio Grande Conservancy District: Albuquerque, 2017; [http://riograndewaterfund.org/wp-content/uploads/2017/01/rgwf\\_stone\\_etal\\_2017.pdf](http://riograndewaterfund.org/wp-content/uploads/2017/01/rgwf_stone_etal_2017.pdf).

(42) Water Environment Federation; American Public Health Association. *Standard methods for the examination of water and wastewater*; American Public Health Association (APHA): Washington, DC, USA, 2005.

(43) Quality of storm water runoff at Los Alamos National Laboratory in 2000 with emphasis on the impact of the Cerro Grande Fire; Highlights of LA-13926; Los Alamos National Laboratory: Los Alamos, NM; <http://permalink.lanl.gov/object/tr?what=info:lanl-repo/lareport/LA-14177>.

(44) R Core Team R: A language and environment for statistical computing; R foundation for statistical computing: Vienna, Austria, 2015.

(45) Seifert, E. OriginPro 9.1: Scientific Data Analysis and Graphing Software-Software Review. *J. Chem. Inf. Model.* **2014**, *54* (5), 1552–1552.

(46) Water Quality Survey Summary for the Valles Caldera National Preserve Watershed; Surface Water Quality Bureau, New Mexico Environment Department: Santa Fe, NM, 2006; <https://www.env.nm.gov/swqb/Surveys/VallesCalderaWQS2001.pdf>.

(47) USEPA STORET Database; <https://www.waterqualitydata.us/portal/31EFkjem020.7> (accessed on Dec 22, 2017).

(48) Burt, R.; Wilson, M.; Mays, M.; Lee, C. Major and trace elements of selected pedons in the USA. *J. Environ. Qual.* **2003**, *32* (6), 2109–2121.

(49) Plant tissue analysis and interpretation for vegetable crops in Florida; Highlights of HS964; IFAS Extension, University of Florida: Gainesville, FL, 2003; <http://edis.ifas.ufl.edu/pdffiles/EP/EP08100.pdf>.

(50) Reimann, C.; Ottesen, R. T.; Andersson, M.; Arnoldussen, A.; Koller, F.; Englmaier, P. Element levels in birch and spruce wood ashes—green energy? *Sci. Total Environ.* **2008**, *393* (2), 191–197.

(51) Etiegni, L.; Campbell, A. Physical and chemical characteristics of wood ash. *Bioresour. Technol.* **1991**, *37* (2), 173–178.

(52) Tsai, K.-P.; Uzun, H.; Karanfil, T.; Chow, A. T. Dynamic Changes of Disinfection Byproduct Precursors following Exposures of *Microcystis aeruginosa* to Wildfire Ash Solutions. *Environ. Sci. Technol.* **2017**, *51* (15), 8272–8282.

(53) Gabet, E. J.; Bookter, A. Physical, chemical and hydrological properties of Ponderosa pine ash. *Int. J. Wildland Fire* **2011**, *20* (3), 443–452.

(54) Wang, J.-J.; Dahlgren, R. A.; Chow, A. T. Controlled burning of forest detritus altering spectroscopic characteristics and chlorine reactivity of dissolved organic matter: effects of temperature and oxygen availability. *Environ. Sci. Technol.* **2015**, *49* (24), 14019–14027.

(55) O'connell, M.; Baldwin, D. S.; Robertson, A.; Rees, G. Release and bioavailability of dissolved organic matter from floodplain litter: influence of origin and oxygen levels. *Freshwater Biol.* **2000**, *45* (3), 333–342.

(56) Moore, T.; Dalva, M. Some controls on the release of dissolved organic carbon by plant tissues and soils. *Soil Sci.* **2001**, *166* (1), 38–47.

(57) de Mendonça, F. G.; da Cunha, I. T.; Soares, R. R.; Tristão, J. C.; Lago, R. M. Tuning the surface properties of biochar by thermal treatment. *Bioresour. Technol.* **2017**, *246* (246), 28–33.

(58) Fang, Q.; Chen, B.; Lin, Y.; Guan, Y. Aromatic and hydrophobic surfaces of wood-derived biochar enhance perchlorate

adsorption via hydrogen bonding to oxygen-containing organic groups. *Environ. Sci. Technol.* **2014**, *48* (1), 279–288.

(59) von Gunten, K.; Alam, M. S.; Hubmann, M.; Ok, Y. S.; Konhauser, K. O.; Alessi, D. S. Modified sequential extraction for biochar and petroleum coke: Metal release potential and its environmental implications. *Bioresour. Technol.* **2017**, *236*, 106–110.

(60) Bogusz, A.; Oleszczuk, P.; Dobrowolski, R. Application of laboratory prepared and commercially available biochars to adsorption of cadmium, copper and zinc ions from water. *Bioresour. Technol.* **2015**, *196*, 540–549.

(61) Burton, C. A.; Hoefen, T. M.; Plumlee, G. S.; Baumberger, K. L.; Backlin, A. R.; Gallegos, E.; Fisher, R. N. Trace Elements in Stormflow, Ash, and Burned Soil following the 2009 Station Fire in Southern California. *PLoS One* **2016**, *11* (5), e0153372.

(62) Burke, M.; Hogue, T.; Kinoshita, A.; Barco, J.; Wessel, C.; Stein, E. Pre-and post-fire pollutant loads in an urban fringe watershed in Southern California. *Environ. Monit. Assess.* **2013**, *185* (12), 10131–10145.

(63) Stein, E. D.; Brown, J. S.; Hogue, T. S.; Burke, M. P.; Kinoshita, A. Stormwater contaminant loading following southern California wildfires. *Environ. Toxicol. Chem.* **2012**, *31* (11), 2625–2638.

(64) Alam, M. S.; Swaren, L.; von Gunten, K.; Cossio, M.; Bishop, B.; Robbins, L. J.; Hou, D.; Flynn, S. L.; Ok, Y. S.; Konhauser, K. O.; et al. Application of surface complexation modeling to trace metals uptake by biochar-amended agricultural soils. *Appl. Geochem.* **2018**, *88*, 103–112.

(65) Mohan, D.; Pittman, C. U. Arsenic removal from water/wastewater using adsorbents—a critical review. *J. Hazard. Mater.* **2007**, *142* (1), 1–53.

(66) Bhattacharya, A.; Naiya, T.; Mandal, S.; Das, S. Adsorption, kinetics and equilibrium studies on removal of Cr (VI) from aqueous solutions using different low-cost adsorbents. *Chem. Eng. J.* **2008**, *137* (3), 529–541.

(67) Kotaš, J.; Stasicka, Z. Chromium occurrence in the environment and methods of its speciation. *Environ. Pollut.* **2000**, *107* (3), 263–283.

(68) Bauer, P.; Elbaum, R.; Weiss, I. M. Calcium and silicon mineralization in land plants: transport, structure and function. *Plant Sci.* **2011**, *180* (6), 746–756.

(69) Franceschi, V. R.; Nakata, P. A. Calcium oxalate in plants: formation and function. *Annu. Rev. Plant Biol.* **2005**, *56*, 41–71.

(70) Harvey, O. R.; Herbert, B. E.; Rhue, R. D.; Kuo, L.-J. Metal interactions at the biochar-water interface: energetics and structure-sorption relationships elucidated by flow adsorption microcalorimetry. *Environ. Sci. Technol.* **2011**, *45* (13), 5550–5556.

(71) Myers-Pigg, A. N.; Louchouart, P.; Amon, R. M.; Prokushkin, A.; Pierce, K.; Rubtsov, A. Labile pyrogenic dissolved organic carbon in major Siberian Arctic rivers: Implications for wildfire-stream metabolic linkages. *Geophys. Res. Lett.* **2015**, *42* (2), 377–385.

(72) Lima, I. M.; Boateng, A. A.; Klasson, K. T. Physicochemical and adsorptive properties of fast-pyrolysis bio-chars and their steam activated counterparts. *J. Chem. Technol. Biotechnol.* **2010**, *85* (11), 1515–1521.