

Tuning the Formal Potential of Ferrocyanide over a 2.1 V Range

Brendon J. McNicholas,^a Robert H. Grubbs,^a Jay R. Winkler,^a Harry B. Gray,*^a and Emmanuelle Despagnet-Ayoub*^{a,b}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

We report the synthesis and characterization of homoleptic borane adducts of hexacyanoferrate(II). Borane coordination blueshifts d-d transitions and CN IR and Raman frequencies. Control over redox properties is established with respect to borane Lewis acidity, reflected in peak anodic potential shifts per borane of +250 mV for BPh₃ and +350 mV for B(C₆F₅)₃. Electron transfer from [Fe(CN-B(C₆F₅)₃)₆]⁴⁻ to photogenerated [Ru(2,2'-bipyridine)₃]³⁺ is very rapid, consistent with voltammetry data. Coordination by Lewis acids provides an avenue for selective modification of the electronic structures and electrochemical properties of cyanometalates, opening the way for the assembly of a library of new reversible redox couples for use in non-aqueous redox flow batteries.

Introduction

Duward Shriver pioneered the study of borane adducts of cyanometalates.^{1a-d} Although he established from analysis of vibrational spectroscopic data that borane coordination (CN-BR₃) enhanced M(d⁶)-CN π backbonding, surprisingly, to the best of our knowledge, no structures of d⁶ low-spin borane adducts have been hitherto reported.² Indeed, only three homoleptic borane adducts have been crystallographically characterized, (TEA)₃[Cr(NC-BPh₃)₆] (TEA = tetraethylammonium, BPh₃ = triphenylborane), (CPh₃)₂[Ni(CN-B(C₆F₅)₃)₄] and (CPh₃)₂[Pd(CN-B(C₆F₅)₃)₄] (CPh₃⁺ = trityl cation, B(C₆F₅)₃ = tris(pentafluorophenyl)borane).^{3,4} Also surprising is that very little is known about the electrochemistry of coordinatively-saturated borane adducts of hexacyanometalates, with the majority of previous work focused on the solvent dependence of cyanometalate redox potentials.^{1d,5-8} We show herein that the formal potentials of these boronated adducts span an unusually wide range (over 2.1 V), opening the way for use as oxidants and reductants in energy storage devices.

Results and Discussion

Tetrabutylammonium (TBA) and TEA hexacyanoferrate(II) compounds (**1a**, **1b**) were prepared using a modified literature procedure.⁹ Neutralization of H₄[Fe(CN)₆] with TBAOH (or TEAOH) in water generated the alkylammonium salt in quantitative yield. Bis(triphenylphosphine)iminium (PPN) hexacyanoferrate(II) (**1c**) was prepared by combining four equivalents of PPNCl with one equivalent of K₄[Fe(CN)₆] in water. ¹H NMR, ¹³C{¹H} NMR, UV-vis, solid-state IR, and voltammetry confirmed the purity of **1**, with a single, reversible redox couple with a formal potential of -1.25 V vs. Fc^{+/-} in MeCN.

The borane adducts of **1** were synthesized by combining six equivalents of borane with one equivalent of **1** dissolved in dichloromethane (DCM) in a nitrogen-filled glove box. Coordination of borane generated (TEA)₄[Fe(CN-BPh₃)₆] (**2**) and (PPN)₄[Fe(CN-B(C₆F₅)₃)₆] (**3**). Each complex was purified and subsequently analyzed by ¹¹B and ¹³C NMR spectroscopy, X-ray crystallography, UV-vis spectroscopy, elemental analysis, and voltammetry.

Full borane coordination for all species was confirmed by analysis of X-ray crystallographic data (see SI), with representative structures depicted in Figure 1. The average M-CN (1.91 Å) and C-N (1.17 Å) bond lengths for **2** compared to **1**, 1.93 Å and 1.17 Å (Figure S11), respectively, are consistent with competing σ and π interactions, where σ donation from nitrogen to boron weakens the M-CN bond, while π backbonding strengthens it. Thus, the contraction of the M-CN bond in **2** is negligibly small. The average M-C-N (174.6°) and C-N-B (172.3°) bond angles for complex **2** are not perfectly linear, likely due to the effects of steric clash among the aryl groups. The average M-C-N (176.9°) and C-N-B (173.9°) bond angles for complex **3** are similarly perturbed. There are slight contractions in average M-CN (1.90 Å) and C-N (1.15 Å) bond lengths for **3** compared to those in complex **2**.

In the ¹³C{¹H} NMR spectra, the cyanide carbon exhibits only one peak, indicating that boranes are bound to all six cyanides (δ_{CN} : 159 ppm for **2** and 162 ppm for **3**). The carbon chemical shift follows the expected downfield trend for a Lewis acid inductively withdrawing electron density through the terminal nitrogen. The change in shift between the BPh₃ and B(C₆F₅)₃ adducts is small due to cooperative σ donation from the nitrogen and π backdonation from the metal center. It is well understood that isocyanoborate complexes experience decreased σ -bonding relative to cyano parents due to reduced electron density on carbon. Conversely, isocyanoborate complexes experience greater π -backbonding relative to cyanide complexes due to lower $\pi^*(CN)$ energies. The ¹¹B NMR spectra of **2** and **3** are in line with increased electron-withdrawing by B(C₆F₅)₃ relative to BPh₃, with the ¹¹B signal for B(C₆F₅)₃ more upfield versus BPh₃ (δ = -5.3 ppm for **2** and -14.4 ppm for **3**).

^a Beckman Institute, and Division of Chemistry and Chemical Engineering, California Institute of Technology, 1200 East California Boulevard, Mail Code 139-74, Pasadena, California 91125, United States

^b Occidental College, Norris Hall of Chemistry, 1600 Campus Rd, Los Angeles, CA 90041, United States

Electronic Supplementary Information (ESI) available: [Materials and methods, additional data from NMR, X-ray crystallographic, laser-quench, and electrochemical experiments]. See DOI: 10.1039/x0xx00000x




Figure 1. **A.** Molecular structure of $(\text{TEA})_4[\text{Fe}(\text{CN-BPh}_3)_6]$ (**2**). Thermal ellipsoids set at 50% probability. Cations and protons omitted for clarity. Selected bond distances and angles: Fe1-C1: 1.908(7) Å, Fe1-C3: 1.913(8) Å, Fe1-C4: 1.908(8) Å, C1-N1: 1.170(9) Å, C3-N3: 1.164(10) Å, C4-N4: 1.166(10) Å, N1-B1: 1.559(9) Å, N3-B3: 1.573(10) Å, N4-B4: 1.557(10) Å, Fe1-C1-N1: 174.0(7)°, Fe1-C3-N3: 174.0(7)°, Fe4-C4-N4: 179.6(8)°, C1-N1-B1: 172.0(8)°, C3-N3-B3: 173.5(8)°, C4-N4-B4: 176.3(7)°, **B.** Molecular structure of $(\text{PPN})_4[\text{Fe}(\text{CN-B(C}_6\text{F}_5)_3)_6]$ (**3**). Thermal ellipsoids set at 50% probability. Cations omitted for clarity. Selected bond distances and angles (labels as in **2**): Fe1-C1: 1.899(4) Å, Fe1-C3: 1.904(3) Å, Fe1-C4: 1.897(3) Å, C1-N1: 1.146(4) Å, C3-N3: 1.152(4) Å, C4-N4: 1.160(4) Å, N1-B1: 1.545(5) Å, N3-B3: 1.550(5) Å, N4-B4: 1.551(4) Å, Fe1-C1-N1: 176.9(3)°, Fe1-C3-N3: 177.9(3)°, Fe1-C4-N4: 175.9(3)°, C1-N1-B1: 173.2(3)°, C3-N3-B3: 175.1(3)°, C4-N4-B4: 173.3(4)°.




Figure 2. **A.** Infrared spectra, **B.** Raman spectra.

A

The solid-state IR and Raman spectra of **1** and its various adducts are shown in **Figure 2**. As expected, the CN stretching frequency increases as the Lewis acidity of the borane or the oxidation state of the metal increases. Increased Lewis acidity causes increased stretching frequency, a result of lowering the absolute energies of the cyanide-based π -bonding orbitals.⁹

B

The lowest energy absorption band in the UV-vis spectra (**Figure 3**) of the adducts is attributable to a spin-allowed d-d transition ($^1\text{A}_{1g} \rightarrow ^1\text{T}_{1g}$).¹⁰ This band, 323 nm in water, redshifts to 357 nm in MeCN. Upon coordination of BPh₃ and B(C₆F₅)₃, **2** and **3** exhibit blueshifted absorbance maxima, indicating that increased backbonding outweighs decreased σ bonding in the octahedral ligand field.¹⁰ For **2** and **3**, bands below 270 nm are attributable to borane π to π^* transitions (**Figures S16-S18**).

Cyclic voltammograms of pure **1b** in DCM solution and one with two equivalents of BPh₃ added are shown in **Figure 4A**. Both CVs were corrected for the non-Faradaic charging current and integrated to ensure that the anodic charge passed was equal for both, which suggests that the only redox reaction taking place is the one of interest (Fe^{III/II}). By adding two equivalents of BPh₃, seven anodic peaks are visible, corresponding to a distribution of all possible numbers of boranes coordinated to hexacyanoferrate(II). Addition of a greater number of equivalents of borane increases the peak current for the coordinatively-saturated species (**Figure S7**). The seven anodic peaks corresponding to different coordination numbers are clearly seen in differential pulse voltammetry (**Figure 4B**). The differential current responses for the two- and three-coordinate species are broader, likely due to the existence of isomers being oxidized at slightly different potentials, which decreases the intensity and broadens the observed differential wave for these adducts.¹¹ As expected, the addition of borane results in a more positive peak anodic potential, likely due to lowering the absolute energies of the metal-based orbitals. The trend in peak anodic potential shift (from cyclic voltammetry) per borane added to hexacyanoferrate(II) is linear (**Figure 4C**), suggesting an electron withdrawing effect that is solely dependent on the Lewis acidity of the borane, with little to no attenuating effects as more boranes are added to the secondary coordination sphere.¹² Assuming the peak anodic potential correlates with the formal potential for each borane species, the linear behavior is consistent with,

$$E_n^o - E_{n-1}^o = \frac{RT}{F} \ln\left(\frac{K_n^{II}}{K_{n-1}^{III}}\right) \quad (1)$$

where the change in formal potential upon coordination of an additional borane, $E_n^o - E_{n-1}^o$, is proportional to the ratio of binding constants of BPh₃ to the Fe(II) and Fe(III) states of the complex, K_n^{II}/K_{n-1}^{III} .¹² The approximate $E_{1/2}$ value for **2** is 0.32 V vs. Fc^{+/-}.

Although full coordination of borane in complex **2** was confirmed by both X-ray crystallography and ¹³C NMR, the voltammetry of **2** in both DCM and MeCN is not electrochemically reversible (**Figure 4D**).




Figure 3. UV-vis absorption spectra of **1** and its borane adducts. Absorbance maxima (nm) and extinction coefficients ($M^{-1} \text{ cm}^{-1}$): **1b**: 357 (583), 255 (7,204), 227 (10,130); **2**: 324 (347); **3**: 317 (333), 265 (9,540), 230 (35,650).




Figure 4. **A.** Cyclic voltammograms of 5 mM **1b** (blue) and 5 mM **1b** with two equivalents of BPh_3 (black) at 100 mV s^{-1} in DCM with 0.1 M TBAPF_6 . **B.** Differential pulse voltammetry of **1b** with two equivalents of BPh_3 . **C.** Linear regression of peak anodic potential from cyclic voltammetry versus number of BPh_3 molecules coordinated to ferrocyanide. **D.** Normalized scan-rate dependence of **2** in MeCN with 0.2 M TBAPF_6 .

Cyclic voltammetry suggests that the complex undergoes electron transfer followed by a borane dissociation, with the metalloproduct undergoing reduction and subsequent re-oxidation (EC mechanism).¹³ The proposed mechanism is supported by a peak current ratio that is less than one for the six-coordinate species and by the presence of cathodic waves that correspond to reduction of the four-, five-, and six-coordinate species (Figure 4A/4D).¹³ We observe a 28 mV shift in peak anodic potential with a ten-fold increase in scan rate, consistent with the theoretical value of 30 mV for a purely kinetic EC mechanism.¹³ This mechanism also is supported by the appearance of anodic waves for the four- and five-coordinate species as a result of sweeping through multiple cycles at fast scan rates and by the decrease in anodic current for the six-coordinate species after the first scan (Figure S8). The peak cathodic current does not increase with increasing scan rate, consistent with

very rapid BPh_3 dissociation.¹³ Additionally, **2** was oxidized with dibenzo-1,4-dioxin radical cation in MeCN, and ^1H NMR in CD_3CN confirmed the presence of $\text{Ph}_3\text{B-NCMe}$ (Figure S9).

In contrast to **2**, complex **3** displays a single, electrochemically reversible redox event with a formal potential of 0.85 V vs. $\text{Fc}^{+}/0$ (Figure 5A), corresponding to a 2.1 V anodic shift in the $\text{Fe}^{\text{III}}/\text{II}$ couple, which is a 350 mV anodic shift per $\text{B}(\text{C}_6\text{F}_5)_3$ added to hexacyanoferrate(II). Similar to **2**, a linear potential shift per borane added to hexacyanoferrate(II) was observed for **3** (Figure S10). Borane adducts of $\text{Fe}(\text{phen})_2(\text{CN})_2$ in dichloromethane showed an approximately 300 mV increase in peak anodic potential per borane, with BBr_3 producing the largest shift.^{1d} Cyanorhenate(I) complexes, $\text{Re}(\text{R}_2\text{phen})(\text{CO})_3[\text{CN-B}(\text{C}_6\text{F}_5)_3]$, where $\text{R} = \text{H, Me}$, displayed a \sim 320 mV shift in peak anodic potential ($E_{\text{p,a}}$) in acetonitrile compared to the parent.¹⁴ Cyanoosmate(II) complexes, $[\text{Os}(4,4'\text{-R}_2\text{bpy})_2(\text{CNBL}_3)_2]$, where $\text{R} = \text{H, Me}$, showed \sim 420 and \sim 290 mV anodic shifts per borane for $\text{L} = (\text{C}_6\text{F}_5)_3\text{B}$ and BPh_3 in MeCN, respectively.¹³ As the $\text{Fe}^{\text{III}}/\text{II}$ peak anodic potential shifts are very near those observed for $\text{Re}(\text{I})$ and $\text{Os}(\text{II})$ complexes, there is minimal dependence on metal identity or oxidation state.

The peak current, i_p , of an electrochemically reversible, diffusion-controlled voltammogram is defined by,¹⁶

$$i_p = 0.4463n^{3/2}F^{3/2}AC_0^*\sqrt{\frac{D_0v}{RT}} \quad (2)$$

where n is the number of electrons, F is Faraday's constant, D_0 is the diffusion coefficient ($\text{cm}^2 \text{ s}^{-1}$), v is the scan rate (V s^{-1}), A is the surface area of the electrode ($A = 0.0707 \text{ cm}^2$) and C_0^* is the bulk concentration of the redox-active species (mol cm^{-3}). The diffusion coefficient for **3** in MeCN, obtained from a Randles-Sevcik plot of peak current vs. $v^{1/2}$, was found to be $4.9 \times 10^{-6} \text{ cm}^2 \text{ s}^{-1}$, which is similar to values for potassium ferrocyanide in aqueous electrolyte.^{17,18}




Figure 5. **A.** Cyclic voltammetry (scan rates from 25 mV s^{-1} to 1000 mV s^{-1}) of 3.6 mM **3** in acetonitrile with 0.2 M TBAPF_6 . Potentials relative to $\text{Fc}^{+}/0$. **B.** Randles-Sevcik plot of the CV data from **5A**.

Oxidation of **3** by flash-quench-generated $[\text{Ru}(2,2'\text{-bipyridine})_3]^{3+}$ was very rapid. From a linear fit of electron transfer rate versus concentration (Figure S15), the apparent second order rate constant, k_{ex} , was found to be $8.4 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$. Using an ion-pair preequilibrium model, K_0 , the ion-pair association constant, was found to be 0.19 M^{-1} , with k_{et} estimated to be $4.5 \times 10^9 \text{ s}^{-1}$.¹⁹ We conclude that electron tunneling from Fe^{II} to Ru^{III} in the ion-paired precursor complex is not inhibited by the wall of 90 fluorine atoms in the boronated cyanide complex.

ARTICLE

Journal Name

Conclusions

We have demonstrated that boranes can tune the formal potentials of reversible redox couples. This finding means that researchers can selectively alter the formal potentials of cyanide-based inorganic complexes, providing opportunities for creating new quenchers, oxidants, and single-electron transfer reagents. Importantly, the fluorinated “cage” surrounding the iron center of **3** does not hinder outer-sphere electron transfer, indicating that boronated cyanide complexes will likely be useful as electrolytes in non-aqueous redox flow batteries.²⁰

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors thank Mike Gray and Kimberly See for help in collecting solid-state Raman data. Mike Takase and Larry Henling provided invaluable assistance in collecting and refining X-ray crystallographic data. The authors also thank Brian Sanders and Wesley Kramer for helpful discussions. Supported by NSF (CCI CHE-1305124) and (CHE-1763429). Additional support was provided by the Arnold and Mabel Beckman Foundation.

Notes and references

- (a) D. F. Shriver, *J. Am. Chem. Soc.*, 1962, **84**, 4610-4611. (b) D. F. Shriver *J. Am. Chem. Soc.*, 1963, **85**, 1405-1408. (c) D. F. Shriver, *The Ambident Nature of Cyanide*. In *Structure and Bonding*. Springer: Berlin, Heidelberg, 1966; Vol. 1, p 32-58.
- (d) C. F. Woodcock and D. F. Shriver, *Inorg. Chem.*, 1986, **25**, 2137-2142.
- 2 J. S. Kristoff and D. F. Shriver, *Inorg. Chem.*, 1973, **12**, 1788-1793.
- 3 E. J. Schelter, M. Shatruk, R. A. Heintz, J. R. Galán-Mascarós, and K. R. Dunbar, *Chem. Commun.*, 2005, **11**, 1417-1419.
- 4 J. Zhou, S. J. Lancaster, D. A. Walker, S. Beck, M. Thornton-Pett, and M., *J. Am. Chem. Soc.*, 2001, **123**, 223-237.
- 5 V. Gutmann, G. Gritzner, and K. Danksagmuller, *Inorganica Chim. Acta*, 1976, **17**, 81-86.
- 6 P. K. Mascharak, *Inorg. Chem.*, 1986, **25**, 245-247.
- 7 H. E. Toma and M. S. Takasugi, *J. Solution Chem.*, 1989, **18**, 575-583.
- 8 J. Jiang, A. Acunzo, and S. A. Koch, *J. Am. Chem. Soc.*, 2001, **123**, 12109-12110.
- 9 D. F. Shriver, S. A. Shriver, and S. E. Anderson, *Inorg. Chem.*, 1965, **4**, 725-730.
- 10 J. J. Alexander and H. B. Gray, *J. Am. Chem. Soc.*, 1968, **90**, 4260-4271.
- 11 D. E. Richardson and H. Taube, *Inorg. Chem.*, 1981, **20**, 1278-1285.
- 12 FD. Lexa, P. Rentien, J. M. Savéant and F. Xu, *J. Electroanal. Chem.*, 1985, **191**, 253-279.
- 13 R. S. Nicholson and I. Shain, *Anal. Chem.*, 1964, **36**, 706.
- 14 W. -K. Chu, C. -C. Ko, K. -C. Chan, S. -M. Yiu, F. -L. Wong, C. -S. Ling, and V. A. L. Roy, *Chem. Mater.*, 2014, **26**, 2544-2550.
- 15 W. -K. Chu, S. -M. Yiu, and C. -C. Ko, *Organometallics*, 2014, **33**, 6771-6777.
- 16 A. J. Bard and L. R. Faulkner, *Electrochemical Methods*, Wiley, 2000.
- 17 S. J. Konopka and B. McDuffie, *Anal. Chem.*, 1970, **42**, 1741-1746.
- 18 G. Gerhardt and R. N. Adams, *Anal. Chem.*, 1982, **54**, 2618-2620.
- 19 G. M. Brown and N. Sutin, *J. Am. Chem. Soc.*, 1979, **101**, 883-892.
- 20 J. M. Stauber, S. Zhang, N. Gvozdik, Y. Jiang, L. Avena, K. J. Stevenson, and C. C. Cummins, *J. Am. Chem. Soc.*, 2018, **140**, 538-541.