



SYMPOSIUM

IUCN Conservation Status Does Not Predict Glucocorticoid Concentrations in Reptiles and Birds

Lynn B. Martin,^{1,*} Maren Vitousek,^{†,‡} Jeremy W. Donald,[§] Travis Flock,[¶] Matthew J. Fuxjager,^{||} Wolfgang Goymann,[#] Michaela Hau,^{*,**} Jerry Husak,^{††} Michele A. Johnson,^{‡‡} Bonnie Kircher,^{§§} Rosemary Knapp,^{¶¶} Eliot T. Miller,[†] Laura A. Schoenle,^{*,|||} Tony Williams^{##} and Clinton D. Francis^{***}

^{*}Department of Global Health, University of South Florida, Tampa, FL 33620, USA; [†]Cornell Lab of Ornithology, Ithaca, NY 14850, USA; [‡]Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, NY 14853, USA; [§]Coates Library, Trinity University, San Antonio, TX 78212, USA; [¶]Department of Integrative Biology, University of South Florida, Tampa, FL 33620, USA; ^{||}Department of Biology, Wake Forest University, Winston-Salem, NC 27109, USA; [#]Max Planck Institute for Ornithology, 82319 Starnberg, Germany; ^{**}Department of Biology, University of Konstanz, 78464 Konstanz, Germany; ^{††}Department of Biology, University of St. Thomas, St. Paul, MN 55105, USA; ^{‡‡}Department of Biology, Trinity University, San Antonio, TX 78212, USA; ^{§§}Department of Biology, University of Florida, Gainesville, FL 32608, USA; ^{¶¶}Department of Biology, University of Oklahoma, Norman, OK 73019, USA; ^{|||}Department of Biology, Hamilton College, Clinton, NY 13323, USA; ^{##}Department of Biological Sciences, Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6; ^{***}Biological Sciences Department, California Polytechnic State University, San Luis Obispo, CA 93407, USA

From the symposium “Understanding the Evolution of Endocrine System Variation through Large-scale Comparative Analyses” presented at the annual meeting of the Society for Integrative and Comparative Biology, January 3–7, 2018 at San Francisco, California.

¹E-mail: lbmartin@usf.edu

Synopsis Circulating glucocorticoids (GCs) are the most commonly used biomarkers of stress in wildlife. However, their utility as a tool for identifying and/or managing at-risk species has varied. Here, we took a very broad approach to conservation physiology, asking whether International Union for the Conservation of Nature (IUCN) listing status (concern versus no obvious concern) and/or location within a geographic range (edge versus non-edge) predicted baseline and post-restraint concentrations of corticosterone (CORT) among many species of birds and reptiles. Even though such an approach can be viewed as coarse, we asked in this analysis whether CORT concentrations might be useful to implicate species at risk. Indeed, our effort, relying on HormoneBase, a repository of data on wildlife steroids, complements several other large-scale efforts in this issue to describe and understand GC variation. Using a phylogenetically informed Bayesian approach, we found little evidence that either IUCN status or edge/non-edge location in a geographic distribution were related to GC levels. However, we did confirm patterns described in previous studies, namely that breeding condition and evolutionary relatedness among species predicted some GC variation. Given the broad scope of our work, we are reluctant to conclude that IUCN status and location within a range are unrelated to GC regulation. We encourage future more targeted efforts on GCs in at-risk populations to reveal how factors leading to IUCN listing or the environmental conditions at range edges impact individual performance and fitness, particularly in the mammals, amphibians, and fish species we could not study here because data are currently unavailable.

Introduction

The first efforts in ecology were largely physiological (Cooke et al. 2013). The father of animal ecology, Victor Shelford (Shelford 1911), comprehensively

identified factors affecting plant and animal distributions, predominantly focusing on energy balance, thermal relationships, and other physiological processes to explain why some species, and not others,

thrived in some areas. In modern times, efforts to use physiology to understand geographic distributions have morphed into at least two subdisciplines. One is the description of basic physiological patterns at large spatiotemporal scales, a discipline termed macrophysiology. Macrophysiology has been very insightful for some plants, invertebrates and ectothermic vertebrates (Chown et al. 2004). For instance, by comparing rare plant species to common ones within the same genus, researchers revealed traits that make rare species distinctive as conservation priorities (Dunbar-Co et al. 2009). This approach for vertebrates, especially in the context of conservation, has been rare (Jessop et al. 2013b). Explicit focus on conservation, however, is the basis of a second subfield: conservation physiology (Wikelski and Cooke 2006). Conservation physiologists seek to use physiological data to predict how natural and anthropogenic stressors pose a challenge to population viability and animal welfare. To date, almost all research in the discipline has involved vertebrates (Cooke et al. 2013), and the majority of studies have relied on a particular group of steroid hormones, the glucocorticoids (GCs).

GCs have been favored in conservation physiology because they often relate to individual health and even fitness (Dantzer et al. 2014), making them potential predictors of population viability. These hormones are also quite simple to measure, being easily detectable in the blood or even feathers, fur, urine, blow (in the case of cetaceans), and feces. GCs also have many functions relevant to conservation interests (Angelier and Wingfield 2013; Sorenson et al. 2017). GCs are integral to water balance (McCormick and Romero 2017) and coordinating key life history stage transitions (Crespi et al. 2013), however, they have mostly been studied because of their impacts on energy metabolism (i.e., baseline GCs). In particular, GCs are critical for how individuals respond to unfamiliar, unpredictable, and often adverse conditions (i.e., responses to acute stressors such as severe weather or short-term food unavailability [Romero and Wingfield 2015]). In conservation physiology, GCs are typically portrayed as biomarkers of stress. The implicit assumption has been that GC dysregulation implicates organisms in need of aid, although description of genuine dysregulation is quite difficult (Tarlow and Blumstein 2007; Lindenmayer et al. 2013; Dantzer et al. 2014; Kilvitis et al. 2017). A prime recent example involves Cape mountain zebras (Lea et al. 2018). There, zebra fecal GCs were higher where grass forage was low and inversely related to female fecundity and subsequent population growth in the

same sites. Overall, conservation physiology aspires to use GCs to indicate cause–effect relationship between stressors and population dynamics (Wikelski and Cooke 2006; Madliger et al. 2015), which could lead to suitable mitigation or remediation options (Cooke et al. 2013).

In many ways, a zeal for GCs as biomarkers of wildlife stress is justified (Martin et al. 2016a). GC measurements, in an appropriate context, have long been recognized as useful to determine exposure to various pesticides, herbicides, and other anthropogenic toxins (Martin et al. 2010; Rohr et al. 2013). Likewise, multiple large-scale analyses have revealed that anthropogenic disturbances are associated with altered GCs (Dickens and Romero 2013; Dantzer et al. 2014; Kleist et al. 2018). For instance, GC regulation in Magellanic penguins (*Spheniscus magellanicus*) (Walker et al. 2005), marine iguanas (*Amblyrhynchus cristatus* [Berger et al. 2007]), and hoatzins (*Opisthocomus hoazin* [Mullner et al. 2004]) is altered by tourism. Translocation into captive breeding programs or to more favorable parts of the range also affects GCs (Dickens et al. 2009), as do changes in conspecific density, with high densities and frequent territorial incursions increasing GCs in all vertebrate classes (Creel et al. 2013). The trouble with using GCs as biomarkers of stress for conservation aims is that relationships between GC concentrations and individual fecundity, survival, and recruitment are often complex (Breuner et al. 2008; Bonier et al. 2009; Sorenson et al. 2017). The implicit assumption of much conservation endocrinology, that “high” GCs indicate at-risk organisms (Dantzer et al. 2014; McCormick and Romero 2017), is probably untenable. Indeed, baseline GCs can increase, decrease, or remain unchanged in response to (chronic) stressors (Dickens and Romero 2013).

On the other hand, large-scale comparisons among species have revealed that simple descriptions of GC regulation can be insightful about challenges faced by populations. Hau et al. (2010) found that baseline GCs (i.e., those measured from blood taken quickly upon organism capture) were higher in species with shorter breeding seasons and smaller body mass. By contrast, GCs measured after a short physical restraint (i.e., stress responses) varied inversely with body mass and positively with annual adult survival rates. In another study, avian species that maintained high residual reproductive value after their first reproductive events mounted weaker GC responses than species with faster-paced reproductive life histories (Bokony et al. 2009). Finally, and most relevant to the present study, GCs have been revealed to have some value for understanding how species

might respond to global change (Jessop et al. 2013a): among 22 reptile and 66 bird species, combinations of a few variables (e.g., body mass, net primary productivity, and latitude) explained appreciable (14% and 33%, respectively) variation in GC responses to restraint.

The above broad comparative studies raise the possibility that a few measures of GCs might implicate at-risk species, which could help direct resource manager efforts to taxa most needing attention. Even though such a study could not indicate the causes of the distress that species experience, the availability of HormoneBase and the insight offered by prior broad comparative studies support effort to query the utility of GCs as a coarse bellwether of conservation risk. Ideally, as in the zebra study discussed above (Lea et al. 2018), one would characterize how individual variation in physiology predicts fitness in the same animals in response to adversity. However, if species prone to conservation concern generally regulate GCs differently than those that cope well with adversity, one might be able to implicate vulnerable species before concerted, expensive, and labor-intensive management efforts ensue.

To test the utility of GCs as such a proxy for conservation risk at the species level, we used HormoneBase (Vitousek et al. 2018), a database containing >6580 entries on glucocorticoid and androgen hormone levels in 474 species. First, we asked whether International Union for the Conservation of Nature (IUCN) status predicted baseline and post-restraint GCs in birds and reptiles, as these two vertebrate groups were the only groups with sufficient data for this analysis. The IUCN is a global organization charged with assigning a “concern” status to many of the world’s species. It categorizes extinction risk, including “red list” assignment to some, depending on traits such as a geographic distribution, demographic structure, and population size and trend (Baillie et al. 2004). According to the IUCN, 86–88% of all birds, mammals, and amphibians assessed in 2010 were somehow threatened, usually by habitat alteration and/or loss (IUCN 2012). We asked whether GCs differ systematically between species of least concern (i.e., the IUCN category for species with no known threats) versus species in all “concern” categories. There were insufficient species in various categories of concern to make comparisons at a higher resolution (see below). Also, we felt that multiple forms of selection and/or phenotypic plasticity (Patterson et al. 2011; Wada 2014) made directional predictions between GCs and risk status difficult at the level of our analysis. The most straightforward expectation would be high GCs in

species of concern. However, the reverse pattern was also plausible because (1) chronic stressors could lead to habituation of the hypothalamic–pituitary–adrenal axis (HPA), (2) exposure to adversity in early-life or selection could favor damped GC responses, or (3) adrenal insufficiency could lead to low GCs in at-risk populations. A third possibility was simply that phylogeny was the strongest driver of variation, an observation consistent with previous work (Jessop et al. 2013b). Our interest was solely to determine whether general patterns between IUCN status and GCs were detectable given the great enthusiasm for GCs in conservation physiology. Follow-up work involving data not available in HormoneBase would be necessary to discern why particular directional patterns occurred, which we describe in the “Discussion” section.

Our second interest was to reveal whether populations from the edges of geographic distributions regulate GCs differently than populations from range cores. This comparison was partly motivated by conservation interests, as at-risk populations commonly dwell at the margins of species distributions where habitat suitability is probably poor compared with the core. However, this part of our analysis was also motivated by work from some authors of the present paper (Martin et al. 2005; Liebl and Martin 2012, 2013; Martin and Liebl 2014; Martin et al. 2015, 2016a) and others (Atwell et al. 2012; Brown et al. 2015). Recurrently, GC responses to restraint have been found stronger at range edges than range cores. Although direct evidence does not yet exist regarding the eco-evolutionary reasons for these patterns, some evidence indicates that strong stress responses at range edges might facilitate particular behaviors conducive to success in novel and/or challenging areas (Martin and Fitzgerald 2005; Liebl and Martin 2012, 2014). In the present study, we expected post-restraint GCs to be higher if samples were collected from the margins versus the core of a species’ range. We did not make directional predictions for baseline GCs, but we conducted the exploratory analysis to search for patterns that might be of future value to conservation biologists.

We, as others in conservation endocrinology, chose to focus on GCs because of their salience to homeostasis (Sapolsky et al. 2000). When GCs are dysregulated, gluconeogenesis, water and electrolyte balance, and a host of other processes compromise nerve signaling, muscle contraction, immune defenses, and eventually reduce physical performance, often leading to death (McCormick and Romero 2017). We chose to study the two forms of GC data available in HormoneBase (baseline and

post-restraint), as GC effects vary depending on which receptors they engage. Baseline GC levels, the effects of which are predominantly mediated by mineralocorticoid receptors, are usually interpreted to reflect integrated energetic expenditures over long time periods (Crespi et al. 2013). That is, baseline GCs often increase moderately with energetic demands, including increases in physical workload, thermoregulatory, reproductive, or defensive (e.g., immunity) needs. Both high and low baseline GCs can therefore have negative health consequences depending on how long and by how much concentrations diverge from typical circadian/circannual rhythms (McEwen and Wingfield 2003; Romero et al. 2009; Martin et al. 2016c). By contrast, post-restraint GCs are mostly regulated through lower-affinity glucocorticoid receptors (GRs). Transient elevations of GCs work through GRs to facilitate emergency life processes that help animals avoid, endure, recover, or cope with stressors (Wingfield et al. 1998). If stressors continue or intensify, and individuals are not able to escape or habituate, persistent GC elevations can lead to pathology, as systems remain engaged beyond the point that they are protective (Korte et al. 2005; Martin 2009). Below, we describe how we used HormoneBase to discern whether IUCN status and point of sample collection within a species' range predict GCs in several species of wild birds and reptiles.

Methods

HormoneBase

Over the course of several years, we searched the literature to find all published accounts of circulating GC concentrations in wild organisms. The details about this search and the resultant database are explained in a recent publication (Vitousek et al. 2018). There as here, we analyzed only data on circulating corticosterone (CORT) concentrations, as comparisons of GC metabolites and/or fecal GCs were unavailable and would be difficult to interpret in any case due to interspecific variation in digestive physiology (Goymann 2012). We excluded mammal (M), amphibian (A), and fish (F) samples from our analysis for one or more of the following reasons: (1) too few species have an IUCN status (A), (2) marine/aquatic species ranges were impossible to describe in the same way as terrestrial birds and reptiles (F), and/or (3) entries in HormoneBase were too few to make meaningful comparisons (M and A). We also had to exclude some species of marine birds because their ranges could not be

described in a manner comparable to non-marine species. We focused on CORT as it is the most widely-measured GC in birds and reptiles.

Statistical analysis

We used Bayesian-informed generalized linear mixed-effect models in the R package MCMCglmm (Hadfield 2010) to ascertain whether (1) IUCN status and (2) location within a species range where samples were collected predicted baseline and post-restraint concentrations of CORT. For IUCN status, we used the IUCN Red List website (<http://www.iucnredlist.org>) to assign all species for which we had data to a category. There are four categories of concern (near-threatened, vulnerable, endangered, and critically endangered) and one of "least concern" assigned by IUCN, but there were too few data for species in "concern" categories for us to analyze data based on original assignments. Subsequently, all species of any level of concern were collapsed into a single "concern" category and these species compared with the remaining "least concern" species. This approach resulted in 1153 measures of baseline glucocorticoids from 139 species assigned to "least concern," and 228 from 22 species assigned to "concern" (Supplementary Table S1). For post-restraint CORT we had 410 glucocorticoid measures from 79 species assigned to "least concern" and 78 measures from 13 species assigned to "concern" (Supplementary Table S2).

To categorize location of capture within each species' range, we used IUCN distribution maps in conjunction with Google Earth (<https://earth.google.com/web>) and geographic coordinates (from the original papers reported in HormoneBase) to assign each HormoneBase entry to one of three categories. For each species, we determined whether samples were collected within the inner 10% of the area of a species' range (i.e., core), the outer 10% of a species' range (i.e., edge), or the intervening 80% (i.e., intermediate). However, as our expectations were that range edges would be most different (and demanding) from other parts of the range (and to maximize statistical power), we collapsed the core and intermediate samples into a single category (i.e., non-edge), and used this binary predictor in all final analyses. We conducted separate analyses for IUCN status and site within the geographic distribution. Although many species of concern (in regards to IUCN status) have narrow distributions, our interests in the two factors were distinct. This approach resulted in 326 "edge" and 526 "non-edge" baseline CORT data points from 78 and 43 species,

respectively (Supplementary Table S3). For post-restraint CORT, we had 175 “edge” and 146 “non-edge” data points from 51 and 28 species, respectively (Supplementary Table S4).

As the broad comparative approach we took might obscure subtle relationships, we took several steps to disentangle focal predictors from other variables that might influence CORT. First, to account for variability in assay methodology (Fanson et al. 2017), we used laboratory identity as a random effect. Although this factor might not capture all potential sources of variation in CORT data, it should encompass substantial variation in many aspects of methodology and importantly it was the only relevant methodological variable we had available for the majority of data. Second, because our analyses often included multiple observations of the same species, we included species as a random effect. Third, we included several variables as covariates in models that are known or expected to affect GCs. We included altitude and latitude at the site of collection, mean body mass of individuals in the collected population (Haase et al. 2016), and breeding status (breeding versus non-breeding, with breeding being the reference state in models [Romero 2002]). HormoneBase was the source of all of these covariates for the sake of consistency in analytical efforts among projects (Johnson et al. 2018). We also attempted to use ambient temperature (mean and standard deviation) in the month in which samples were collected, but as that approach greatly diminished our sample sizes, we did not include climate variables in our analysis. We included all samples from HormoneBase in our analysis as long as authors identified them as baseline.

Finally, we used a tree developed specifically for analyses of HormoneBase data in this SICB symposium (Johnson et al. 2018) as our phylogenetic hypothesis to account for potential phylogenetic structure in the relationship between GCs and the predictor variables, which was converted into an inverted phylogenetic covariance matrix prior to analyses. We allowed the model to estimate Pagel's λ , and we report the posterior mode from the MCMC chain. Pagel's λ ranges from 0 to 1 and reflects no and high phylogenetic signal, respectively. We used uninformed priors in the models; default priors for all fixed effects and, reflecting an inverse-Gamma distribution, we used $V=1$ and $\mu=0.02$ for the variance components of each random effect. We scaled all continuous predictors (i.e., z -transformations) to facilitate direct comparisons of variable effects in models to each other. We conducted the IUCN and geographic distribution analyses

separately, as inclusion of both predictors in models would have greatly reduced our sample sizes (due to missing IUCN or distribution data for some HormoneBase entries).

We used an all-subsets approach in model selection. Specifically, from our global model we ranked models using the Deviance Information Criteria (DIC) using the dredge function in the MuMIn R package (Barton 2009) and considered all models $DIC \leq 2$ from the top ranked model as competitive. All model runs were based on 2000 samples drawn from 50,000 MCMC iterations, a burn-in of 10,000, and a thinning rate of 20. For each global model, plus all strongly ranked models (i.e., $DIC \leq 2$), we verified that MCMC chains were mixing by visually inspecting trace and density plots and ensuring that autocorrelation of sampled iterations was less than 0.1. Finally, using the R package coda (Plummer et al. 2008), we verified that four independent chains converge using the Gelman–Rubin statistic. We also calculated marginal (fixed effects only) and conditional (fixed and random effects) R^2 for all posterior models (Nakagawa and Schielzeth 2013). We used two pieces of evidence to determine whether a predictor had a strong effect: (1) whether predictors occurred in well-supported models (i.e., $DIC \leq 2$) and (2) whether the 95% credible intervals, calculated from the posterior means, excluded zero (bolded terms in relevant tables).

Results

IUCN status

The top models for baseline CORT always included breeding status and latitude, but none included IUCN status, and all marginal R^2 values for all models were very low (Table 1). In all models, breeding birds had higher baseline CORT than non-breeding birds and Pagel's λ values were high (Table 2). Latitude was negatively related to baseline CORT in only one model, and CIs overlapped zero for all other terms in all other models. Models for post-restraint CORT also always included breeding status, and two models included IUCN category (Table 3); again though, marginal R^2 was low for all models. Most models also included body mass. However, only breeding status CIs did not overlap zero in the top models (Table 4); again, breeding status was related to post-restraint CORT concentrations. Although IUCN was included in two of the five top models, its influence was very modest. Pagel's λ values were also high for all models.

Table 1 Top models predicting variation among avian and reptilian baseline corticosterone entries in HormoneBase (IUCN status)

Model	Terms in model	df	Loglikelihood	DIC	Delta	Weight	Marginal R^2	Conditional R^2
1	Breeding + latitude + body mass	7	−1223.29	2614	0	0.246	0.01	0.92
2	Breeding + latitude	6	−1223.67	2614.5	0.41	0.200	0	0.93
3	Breeding + latitude + altitude + body mass	8	−1223.63	2615.3	1.26	0.131	0.01	0.93
4	Breeding + latitude + altitude	7	−1224	2615.7	1.68	0.106	0.01	0.93

Table 2 Composition of top models for baseline corticosterone, IUCN analysis

Model	Predictor	Post.mean	Lower 95%	Upper 95%	Effective sample size
1	(Intercept)	1.363	−0.390	3.293	2000
(λ=0.93)	Latitude (scaled)	−0.133	−0.244	−0.029	2000
	Body mass (log, scaled)	0.098	−0.030	0.221	2000
	Breeding (±)	−0.134	−0.228	−0.041	1936
2	(Intercept)	1.464	−0.333	3.351	2000
(λ=0.92)	Latitude (scaled)	−0.119	−0.236	−0.016	2000
	Breeding (±)	−0.134	−0.230	−0.041	2000
3	(Intercept)	1.409	−0.416	3.221	2000
(λ=0.93)	Latitude (scaled)	−0.113	−0.225	−0.010	2000
	Body mass (log, scaled)	0.097	−0.013	0.228	2000
	Altitude (scaled)	0.044	−0.020	0.107	2000
	Breeding (±)	−0.135	−0.225	−0.036	2276
4	(Intercept)	1.478	−0.371	3.279	2000
(λ=0.91)	Latitude (scaled)	−0.097	−0.207	0.013	2182
	Altitude (scaled)	0.042	−0.022	0.109	2138
	Breeding (±)	−0.134	−0.225	−0.039	2136

Bolded text highlights terms with credible intervals that do not overlap zero.

Table 3 Top models predicting variation among avian and reptilian post-restraint corticosterone entries in HormoneBase (IUCN status)

Model	Terms in model	df	Loglikelihood	DIC	Delta	Weight	Marginal R^2	Conditional R^2
1	Breeding + body mass	6	−264.888	619	0	0.172	0.005	0.93
2	Breeding + IUCN + body mass	7	−264.681	619.3	0.32	0.146	0.006	0.93
3	Breeding + latitude + body mass	7	−264.687	620	1.03	0.103	0.005	0.93
4	Breeding + IUCN + latitude + body mass	8	−264.739	620.3	1.32	0.089	0.007	0.93
5	Breeding	5	−265.583	620.6	1.63	0.076	0.002	0.93

Geographic distribution

Location in the range was included in two of the top models for baseline CORT, but breeding status and latitude were included in all models, and body mass and altitude were included in many models (Table 5). All marginal R^2 values were low. Only breeding status and altitude affected baseline CORT (Table 6). Range location CIs always overlapped zero when it was included as a predictor; the same was true for all other variables in the best-fit models. Pagel's λ was also high in all four models. For post-restraint CORT, only two models were

supported, both of which included breeding status and body mass (Table 7), but marginal R^2 values were again very low. Breeding animals had higher post-restraint CORT in all models, but no other effects, including range location, were strong based on CIs overlapping zero (Table 8). Pagel's λ values were high for both models.

Discussion

We investigated whether IUCN status and location within the geographic range predicted GC concentrations in birds and reptiles because GCs enable

Table 4 Composition of top models for post-restraint corticosterone, IUCN analysis

Model	Predictor	Post.mean	Lower 95%	Upper 95%	Effective sample size
1	(Intercept)	2.906	1.674	4.217	2000
($\lambda=0.93$)	Breeding ($\pm$)	-0.178	-0.280	-0.068	2000
	Body mass (log, scaled)	0.085	-0.027	0.192	2000
2	(Intercept)	2.941	1.685	4.164	2000
($\lambda=0.93$)	Breeding ($\pm$)	-0.183	-0.299	-0.082	2000
	IUCN	-0.102	-0.440	0.208	2146
	Body mass (log, scaled)	0.091	-0.023	0.198	2000
3	(Intercept)	2.883	1.529	4.152	2131
($\lambda=0.93$)	Latitude (scaled)	-0.010	-0.104	0.078	2223
	Breeding ($\pm$)	-0.178	-0.277	-0.062	2000
	Body mass (log, scaled)	0.089	-0.026	0.194	2000
4	(Intercept)	2.896	1.639	4.262	2000
($\lambda=0.92$)	IUCN	-0.112	-0.411	0.229	2000
	Latitude (scaled)	-0.013	-0.097	0.088	2000
	Breeding ($\pm$)	-0.181	-0.287	-0.062	2000
	Body mass (log, scaled)	0.097	-0.019	0.214	2000
5	(Intercept)	3.011	1.770	4.280	2000
($\lambda=0.92$)	Breeding ($\pm$)	-0.174	-0.280	-0.066	2000

Bolded text highlights terms with credible intervals that do not overlap zero.

Table 5 Top models predicting variation among avian and reptilian baseline corticosterone entries in HormoneBase (geographic range)

Model	Terms in model	df	Loglikelihood	DIC	Delta	Weight	Marginal R^2	Conditional R^2
1	Breeding + range + latitude + altitude	8	-708.403	1534.6	0	0.239	0.01	0.94
2	Breeding + range + latitude + altitude + body mass	9	-707.89	1534.9	0.28	0.208	0.01	0.95
3	Breeding + latitude + altitude + body mass	8	-708.817	1535.5	0.83	0.157	0	0.95
4	Breeding + latitude + body mass	7	-709.065	1535.7	1.03	0.142	0	0.94

individuals to adjust their phenotypes to variation in environmental conditions (Hau et al. 2016). Unlike most other biomolecules (Martin et al. 2011), GCs can enduringly or reversibly modulate the properties of many tissues, giving them a molecular form of functional pleiotropy that orients organismal phenotypes to a shared life priority (McGlothlin and Ketterson 2008; Ketterson et al. 2009; Cohen et al. 2012; Martin and Cohen 2014). Such extensive effects of GCs make them both potential facilitators and impediments to conservation interests, depending on their collective effects across tissues and contexts. Overall, we found little evidence that IUCN status (some level of concern versus least concern) or location within the geographic range (edge versus non-edge) predicted baseline and post-restraint CORT concentrations of several bird and reptile species.

Null results are always difficult to discuss and interpret, but our rediscovery of previously described interspecific patterns (e.g., effects of breeding status

on CORT) gives us confidence that our analysis was appropriately executed. As with previous work (Casagrande et al. 2018), we found seasonal changes in the regulation of GCs breeding (Romero 2002) even though we used a very coarse surrogate for breeding status (yes/no). Likewise, we have strong evidence for effects of phylogeny (all $\lambda > 0.90$) on both forms of CORT (Jessop et al. 2013b), and we found effects of latitude on BL CORT consistent with other studies (Hau 2010, 4767). In light of these rediscoveries and the exceptionally broad scope of our work, we think it is reasonable to conclude that IUCN listing for species and edge/non-edge status of populations are unrelated to CORT in birds and lizards. For these reasons, we discourage simple future efforts to use CORT as a biomarker of stress at such very broad scales. Below though, we discuss a few potential reasons why other studies of GCs in the service of conservation are warranted, and we propose some potentially useful ways forward at multiple levels of analysis.

Table 6 Composition of top models for baseline corticosterone, range analysis

Model	Predictor	Post.mean	Lower 95%	Upper 95%	Effective sample size
1 ($\lambda=0.95$)	(Intercept)	2.204	0.189	4.160	2024
	Range	−0.083	−0.242	0.084	2000
	Latitude (scaled)	−0.053	−0.159	0.043	1859
	Altitude (scaled)	0.092	0.018	0.166	2000
	Breeding ($\pm$)	−0.167	−0.277	−0.054	2150
2 ($\lambda=0.95$)	(Intercept)	2.156	0.084	4.256	2000
	Latitude (scaled)	−0.061	−0.165	0.039	2000
	Range	−0.091	−0.259	0.071	2000
	Altitude (scaled)	0.091	0.017	0.165	2000
	Breeding ($\pm$)	−0.166	−0.275	−0.047	2000
3 ($\lambda=0.94$)	Body mass (log, scaled)	0.060	−0.069	0.209	2000
	(Intercept)	2.038	0.077	4.202	2023
	Latitude (scaled)	−0.060	−0.164	0.037	1847
	Altitude (scaled)	0.104	0.031	0.179	2000
	Breeding ($\pm$)	−0.165	−0.282	−0.052	2000
4 ($\lambda=0.95$)	Body mass (log, scaled)	0.057	−0.085	0.196	2000
	(Intercept)	2.125	−0.085	3.991	2000
	Latitude (scaled)	−0.054	−0.143	0.051	2000
	Altitude (scaled)	0.103	0.027	0.173	1875
	Breeding ($\pm$)	−0.163	−0.280	−0.041	2245

Bolded text highlights terms with credible intervals that do not overlap zero.

Table 7 Top models predicting variation among avian and reptilian post-restraint corticosterone entries in HormoneBase (geographic range)

Model	Terms in model	df	Loglikelihood	DIC	Delta	Weight	Marginal R^2	Conditional R^2
1	Breeding + body mass	6	−170.157	406.5	0	0.244	0.01	0.94
2	Breeding + latitude + body mass	7	−170.26	407.1	0.65	0.176	0	0.94

Table 8 Composition of top models for post-restraint corticosterone, range analysis

Model	Predictor	Post.mean	Lower 95%	Upper 95%	Effective sample size
1 ($\lambda=0.95$)	(Intercept)	3.747	2.037	5.595	2299
	Breeding ($\pm$)	−0.247	−0.395	−0.121	2000
	Body mass (log, scaled)	0.066	−0.040	0.165	2303
2 ($\lambda=0.95$)	(Intercept)	3.694	1.974	5.440	2000
	Breeding ($\pm$)	−0.259	−0.405	−0.137	2000
	Latitude (scaled)	−0.031	−0.125	0.061	1829
	Body mass (log, scaled)	0.072	−0.030	0.177	2000

Bolded text highlights terms with credible intervals that do not overlap zero.

IUCN status

IUCN status did not predict baseline or post-restraint CORT in birds and reptiles well in spite of quite large sample sizes and phylogenetic coverage. Our care to include several covariates in our analyses

also should have helped tease out any relationships between IUCN status and CORT variation if they exist. The most likely reasons we did not reveal appreciable effects of IUCN status on CORT is that (1) variation in CORT is strongly context-dependent (Busch and Hayward 2009) and (2) IUCN status is

too imprecise an indicator of stress experienced by wildlife. In regards to the former, context-dependence is the rule more than the exception for HPA function, the neuroendocrine axis regulating the release of CORT into the bloodstream. CORT and other HPA regulatory elements fluctuate over several timescales (Woods and Wilson 2014), and these fluctuations are integral to CORT achieving its physiological functions. Baseline CORT largely works in a permissive manner, enhancing the actions of catecholamines, preparing the immune system for insult, stimulating lipolysis and gluconeogenesis, and increasing food consumption and deposition of energy stores (Sapolsky et al. 2000). Conversely, post-restraint CORT mediates emergency life history responses, inducing rapid and transient changes that help an individual flee, endure, actively cope with, and recover from adversity (Wingfield et al. 1998). Perhaps had we included life history stage, health, sex, and variation, and other traits of individuals (Korte et al. 2005), we might have revealed effects of IUCN status on CORT. Repeatedly, breeding status was one of the best predictors of CORT variation. We were unable to include more precise variables because HormoneBase does not include data at the individual animal level and because of the exceptional inherent diversity in life history strategies among the species we considered.

Even had we individual-level data or the ability to describe species in a more specific way, we might not have detected effects of IUCN status on CORT. Others have used GCs for conservation pursuits because of their utility as proxies for individual health and fitness (Strasser and Heath 2013), and the most informative studies, in terms of linking physiology to conservation risk or mitigation opportunity, have been at the level of populations (Martínez-Mota et al. 2007; Homyack 2010). For instance, within species, individuals with lower body condition often have higher baseline GCs (Lindström et al. 2005; Angelier et al. 2009). However, relationships between CORT and aspects of fitness are complex (Schoenle et al. 2018), so even population-level comparative work risks imprecision. For example, baseline CORT in tree swallows (*Tachycineta bicolor*) was only elevated in some populations when food restriction was imposed on females during the period when offspring were being fed (Madliger and Love 2014); outside this period and/or in other populations, baseline CORT did not change with food restriction. In house sparrows (*Passer domesticus*), the direction of the relationship between baseline CORT and reproductive success flipped depending on the breeding stage; before egg-laying, the relationship was

positive but during offspring provisioning, it was negative (Ouyang et al. 2011). These studies and the absence of intelligible patterns in our study highlight that for conservation purposes, we probably need to characterize better how HPA responses to adversity mediate fitness among individuals within populations (see below).

A second likely reason for no influence of IUCN status on CORT is the breadth of factors that leads to IUCN listing in the first place. Listing represents the integration of several different forms of information about a species as well as trends of focal populations (IUCN 2012). This complex algorithm for listing means that organisms can occupy the most dire rankings for quite different reasons. Many of the most threatened species are listed because of very narrow geographic ranges (i.e., endemics), whereas others are listed because their populations are declining over large spatial scales (e.g., habitat destruction), and still others are listed because of particularly vulnerable life stages (e.g., marine turtles). Conservation threats also equate to sources of selection in an evolutionary sense, so when selection acts at different points of life, CORT variation can be driven in a particular direction, obscuring any interspecific patterns. In white-crowned sparrows (*Zonotrichia leucophrys*), for example, survival selection favored high post-restraint CORT, but fecundity selection favored weaker CORT responses to restraint (Patterson et al. 2011). We focused on adult animals here to try to moderate age-dependency in our comparisons, but ideally, one would analyze data from multiple age classes (and other sensitive categories mediating listing) to reveal what about IUCN listing instigates GC variation.

Our perspective is that IUCN status, as a species-level designation, is probably just too coarse a category to relate to GCs or most any other form of physiological variation in the service of conservation. Moreover, as many data in HormoneBase will have come from samples available from the least sensitive parts of a species range (because managers are probably often reluctant to permit capture and handling of critically endangered species), it might be even less likely that coarse analyses would reveal patterns. By contrast, some populations of IUCN-listed species might have been restricted to protected but otherwise suboptimal habitats. Without sampling from multiple points across the range of an IUCN-listed species, it might be hard to link IUCN status and GC variation. Some populations will simply experience less stress than others. Although it is as yet unclear which of the two above factors predominate in our study, it is sensible that future conservation endocrinology efforts take a more focused approach.

Geographic range

Although the “edge” term appeared in the two top models for baseline CORT, its explanatory power was weak compared with other factors. Much like IUCN status, capture location (i.e., edge/non-edge) is probably too imprecise a way to describe adversity of conditions at a site. Our main motivation for conducting this comparison came from previous work on range-expanding populations; the roles of GCs in individual fitness is probably quite distinct in range-edge organisms relative to those enduring conditions for generations at the core. In light of the present results, if we are to discern whether and how location within a range influences GCs, or vice versa, we will need to investigate directly the forces inducing variation in the first place. Although repeatability of both stress-induced GC and baseline concentrations are high in birds (Taff et al. 2018), both measures are also strongly influenced by environmental conditions. Data in HormoneBase represent a mix of within- and among-individual variation, yet without repeated GC measures from individuals, one will remain unable to separate these sources of variation (Baugh et al. 2014). Environmentally-induced variation is important functionally, but it does not capture as well how GCs mediate fitness and thus success or failure in marginal areas (Hau et al. 2016). Moreover, because relationships between fitness and GCs can be non-linear and vary among populations (Martin et al. 2005; Busch and Hayward 2009; Crespi et al. 2013), it will be imperative to study some populations intensively to reveal how GCs mitigates population dynamics at various sites (Zanette et al. 2011; Dantzer et al. 2013; Kleist et al. 2018). Perhaps if we could have compared relationships between GCs and fitness among edge and non-edge sites, we would have revealed interesting patterns. Such data are not available in HormoneBase, however, largely because those datasets remain relatively rare. In the future, efforts to link GCs and geographic distribution should focus on how GCs support population viability, but perhaps expand their perspectives on the functions of GCs. So far, conservation and comparative endocrinologists have focused mostly on the role of CORT in energy balance (Vera et al. 2017). However, GCs have manifold effects (Dallman et al. 2007) including those that mitigate responses to infections (Gervasi et al. 2016; Martin et al. 2016b; Gervasi et al. 2017) and even water balance (Vera et al. 2017). These well-known effects of GCs have been little considered for affecting the viability or distribution of populations, even species in marginal habitats where desertification (Hofmeister and Rubenstein 2016), climate

change, or other challenges to water balance are becoming more common.

Suggestions for future conservation physiology involving GCs

Our results suggest that future work attend to the particular pathways by which GCs affect individual fitness instead of taking coarse comparative approaches, even with large databases. We do not mean to disparage all future large-scale comparative conservation endocrinology efforts, as they could be insightful. For instance, because GCs are deposited into fur, feather, feces, and other tissues at slower rates than they are released and metabolized in the blood, it might be informative to compare GCs from other tissues as proxies of IUCN listing or other forms of conservation risk at broad (i.e., species) levels. Typically though, nuanced approaches to GC conservation physiology will probably be more insightful (Chown and Gaston 2016). The work of Valladares et al. (2014) is a great example; they studied how relationships between phenotypic plasticity and the thermal niche could be used to improve forecasts of species responses to environmental change. We encourage that researchers interested in GCs as a biomarker of wildlife stress also consider the physiological functions of GCs and the context in which it is measured.

As an example of a future promising approach, consider how habitat degradation and destruction could work through GCs to affect mammalian extinction risk. Habitat degradation and destruction are the main threats to extinction for most vertebrate species (Drake and Griffen 2010), but risk changes depending on variation in environmental factors, body size, and intrinsic traits (e.g., life history schedules) among and within species/populations. Large mammals, particularly those from the tropics (Fritz et al. 2009), tend to have higher extinction risks because of their low reproductive rates. However, species above a 3-kg body size threshold are disproportionately (negatively) impacted by environmental (and intrinsic) factors including human population density (Cardillo et al. 2004). To discern whether GCs can serve as biomarkers with genuine conservation value, one could track serum (or possibly fecal) GCs in at-risk populations, expecting GCs to become increasingly dysregulated as populations experience critical slowing down, points on population growth trajectories at which rates of recovery from small perturbations decrease (Drake and Griffen 2010). These quite involved efforts would be prohibitive in many systems, but their execution in a

few species could provide valuable guidance for particularly threatened species.

Note too that for GCs to be supported as genuinely useful biomarkers, one will need to link concentrations in single measurements to the regulatory plasticity of the HPA that mediates GC effects on performance and fitness (Guindre-Parker 2018). Although aspects of GC regulation are sometimes inter-related (Liebl et al. 2013), one should not assume that single measures are proxies for the flexible changes in hormone concentrations that mediate function and fitness. GCs regulatory flexibility probably represents an important form of phenotypic flexibility (Martin and Cohen 2014; Martin and Liebl 2014; Martin et al. 2015; Hau et al. 2016; Taff and Vitousek 2016; Kilvitis et al. 2017), which is beneficial when it allows an individual to alter its phenotype to match a changing environment (DeWitt et al. 1998). In fluctuating environments flexible individuals may be more competitive and have higher reproductive success and survival than inflexible (canalized) individuals, although outcomes will be contingent on the time scale over which environments fluctuate and whether regulation can keep pace. Conversely, although more canalized individuals may have a lower potential to adapt to fluctuating environments, they may be more successful under stable conditions, especially if plasticity is costly (DeWitt et al. 1998). It would be insightful to determine whether HPA regulatory flexibility predicts stable and/or unstable population dynamics, ideally using state-based population models (Crespi et al. 2013).

A final lucrative consideration for conservation endocrinology is to account for the form of chronic stress that is leading populations or species to be of concern in the first place (Dickens and Romero 2013). In terms of their endocrinological effects, chronic stressors tend to take two forms (Martí and Armario 1998). In chronic continuous stress (e.g., some adverse social situations), stressor exposure is omnipresent. In these scenarios, HPA habituation often occurs and GC elevations subside or even decrease relative to levels measured prior to stressor exposure. The other form, chronic intermittent stress, involves exposure to a series of stressors on a consistent or rotating basis. Animals in these conditions do not habituate and maintain high levels of GCs for long periods. We already know from populations of no obvious concern that GCs can be low at certain times of year in challenging environments (e.g., high latitudes), presumably to allow individuals to continue to breed in spite of suboptimal conditions (Wingfield and Sapolsky 2003).

Growing evidence also suggests that selection often favors dampened GC stress responses in human-disturbed areas too (Partecke and Gwinner 2007; Atwell et al. 2012). We recognize that it will be very difficult to conduct involved endocrinological studies on many threatened and endangered wildlife, but more physiologically-nuanced approaches are critical if GCs are to serve as a broadly-useful conservation tool. The distinct research methods and lexicons of physiology and conservation biology alone make collaborations challenging (Lennox and Cooke 2014), but the payoff is mutual awareness and, ideally, realized conservation aspirations (Redpath et al. 2013).

Acknowledgments

The authors thank Samantha Murphy, Steffanie Munguia, Shawn Zamani, Urszula Komendi, Lauren Kenney, and Samantha Oakey for help in compiling data for the analysis.

Funding

This work was supported by the National Science Foundation [grants 1257773 and 1656618 to L.B.M. and grant 1457251 to M.V.], and the HormoneBase Consortium acknowledges the Company of Biologists as well as the DAB, DCE, DCPB, and DEE Divisions of SICB for support for the symposium.

Supplementary data

Supplementary data available at ICB online.

References

- Angelier F, Moe B, Blanc S, Chastel O. 2009. What factors drive prolactin and corticosterone responses to stress in a long-lived bird species (snow petrel *Pagodroma nivea*)? *Physiol Biochem Zool* 82:590–602.
- Angelier F, Wingfield JC. 2013. Importance of the glucocorticoid stress response in a changing world: theory, hypotheses and perspectives. *Gen Comp Endocrinol* 190:118–28.
- Atwell JW, Cardoso GC, Whittaker DJ, Campbell-Nelson S, Robertson KW, Ketterson ED. 2012. Boldness behavior and stress physiology in a novel urban environment suggest rapid correlated evolutionary adaptation. *Behav Ecol* 23:960–9.
- Baillie J, Hilton-Taylor C, Stuart SN. 2004. 2004 IUCN red list of threatened species: a global species assessment. IUCN.
- Barton K. 2009. MuMIn: multi-model inference. R package version 1.0.0 (<http://r-forge.r-project.org/projects/mumin/>).
- Baugh AT, van Oers K, Dingemanse NJ, Hau M. 2014. Baseline and stress-induced glucocorticoid concentrations

- are not repeatable but covary within individual great tits (*Parus major*). *Gen Comp Endocrinol* 208:154–63.
- Berger S, Wikelski M, Romero LM, Kalko EKV, Rödl T. 2007. Behavioral and physiological adjustments to new predators in an endemic island species, the Galapagos marine iguana. *Horm Behav* 52:653–63.
- Bokony V, Lendvai AZ, Liker A, Angelier F, Wingfield JC, Chastel O. 2009. Stress response and the value of reproduction: are birds prudent parents? *Am Nat* 173:589–98.
- Bonier F, Martin PR, Moore IT, Wingfield JC. 2009. Do baseline glucocorticoids predict fitness? *Trends Ecol Evol* 24:634–42.
- Breuner CW, Patterson SH, Hahn TP. 2008. In search of relationships between the acute adrenocortical response and fitness. *Gen Comp Endocrinol* 157:288–95.
- Brown GP, Kelehear C, Shilton CM, Phillips BL, Shine R. 2015. Stress and immunity at the invasion front: a comparison across cane toad (*Rhinella marina*) populations. *Biol J Linn Soc* 116:748–60.
- Busch DS, Hayward LS. 2009. Stress in a conservation context: a discussion of glucocorticoid actions and how levels change with conservation-relevant variables. *Biol Conserv* 142:2844–53.
- Cardillo M, Purvis A, Sechrest W, Gittleman JL, Bielby J, Mace GM. 2004. Human population density and extinction risk in the world's carnivores. *PLoS Biol* 2:e197.
- Casagrande S, Zsolt Garamszegi L, Goymann W, Donald J, Francis CD, Fuxjager MJ, Husak JF, Johnson MA, Kircher B, Knapp R, et al. 2018. Do seasonal glucocorticoid changes depend on reproductive investment? A comparative approach in birds. *Integr Comp Biol* published online (doi: 10.1093/icb/icy022).
- Chown S, Gaston K, Robinson D. 2004. Macrophysiology: large-scale patterns in physiological traits and their ecological implications. *Funct Ecol* 18:159–67.
- Chown SL, Gaston KJ. 2016. Macrophysiology—progress and prospects. *Funct Ecol* 30:330–44.
- Cohen AA, Martin LB, Wingfield JC, McWilliams SR, Dunne JA. 2012. Physiological regulatory networks: ecological roles and evolutionary constraints. *Trends Ecol Evol* 27:428–35.
- Cooke SJ, Sack L, Franklin CE, Farrell AP, Beardall J, Wikelski M, Chown SL. 2013. What is conservation physiology? Perspectives on an increasingly integrated and essential science. *Conserv Physiol* 1:cot001.
- Creel S, Dantzer B, Goymann W, Rubenstein DR. 2013. The ecology of stress: effects of the social environment. *Funct Ecol* 27:66–80.
- Crespi EJ, Williams TD, Jessop TS, Delehanty B. 2013. Life history and the ecology of stress: how do glucocorticoid hormones influence life-history variation in animals? *Funct Ecol* 27:93–106.
- Dallman MF, Warne JP, Foster MT, Pecoraro NC. 2007. Glucocorticoids and insulin both modulate caloric intake through actions on the brain. *J Physiol* 583:431–6.
- Dantzer B, Fletcher QE, Boonstra R, Sheriff MJ. 2014. Measures of physiological stress: a transparent or opaque window into the status, management and conservation of species? *Conserv Physiol* 2:cou023.
- Dantzer B, Newman AE, Boonstra R, Palme R, Boutin S, Humphries MM, McAdam AG. 2013. Density triggers maternal hormones that increase adaptive offspring growth in a wild mammal. *Science* 340:1215–7.
- DeWitt TJ, Sih A, Wilson DS. 1998. Costs and limits of phenotypic plasticity. *Trends Ecol Evol* 13:77–81.
- Dickens MJ, Delehanty DJ, Romero LM. 2009. Stress and translocation: alterations in the stress physiology of translocated birds. *Proc Biol Sci* 276:2051–6.
- Dickens MJ, Romero LM. 2013. A consensus endocrine profile for chronically stressed wild animals does not exist. *Gen Comp Endocrinol* 191:177–89.
- Drake JM, Griffen BD. 2010. Early warning signals of extinction in deteriorating environments. *Nature* 467:456.
- Dunbar-Co S, Sporck MJ, Sack L. 2009. Leaf trait diversification and design in seven rare taxa of the Hawaiian *Plantago* radiation. *Int J Plant Sci* 170:61–75.
- Fanson KV, Németh Z, Ramenofsky M, Wingfield JC, Buchanan KL. 2017. Inter-laboratory variation in corticosterone measurement: implications for comparative ecological and evolutionary studies. *Methods Ecol Evol* 8:1745–54.
- Fritz SA, Bininda-Emonds OR, Purvis A. 2009. Geographical variation in predictors of mammalian extinction risk: big is bad, but only in the tropics. *Ecol Lett* 12:538–49.
- Gervasi SS, Burgan SC, Hofmeister E, Unnasch TR, Martin LB. 2017. Stress hormones predict a host superspreader phenotype in the West Nile virus system. *Proc Biol Sci* 284:pii: 20171090.
- Gervasi SS, Burkett-Cadena N, Burgan SC, Schrey AW, Hassan HK, Unnasch TR, Martin LB. 2016. Host stress hormones alter vector feeding preferences, success, and productivity. *Proc Biol Sci* 283:pii: 20161278.
- Goymann W. 2012. On the use of non-invasive hormone research in uncontrolled, natural environments: the problem with sex, diet, metabolic rate and the individual. *Methods Ecol Evol* 3:757–65.
- Guindre-Parker S. 2018. The evolutionary endocrinology of circulating glucocorticoids in free-living vertebrates: recent advances and future directions across scales of study. *Integr Comp Biol* published online (doi: 10.1093/icb/icy048).
- Haase CG, Long AK, Gillooly JF. 2016. Energetics of stress: linking plasma cortisol levels to metabolic rate in mammals. *Biol Lett* 12:20150867.
- Hadfield JD. 2010. MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. *J Stat Softw* 33:1–22.
- Hau M, Casagrande S, Ouyang JQ, Baugh AT. 2016. Glucocorticoid-mediated phenotypes in vertebrates: multi-level variation and evolution. *Adv Study Behav* 48:41–115.
- Hau M, Ricklefs RE, Wikelski M, Lee KA, Brawn JD. 2010. Corticosterone, testosterone and life-history strategies of birds. *Proc Biol Sci* 277:3203–12.
- Hofmeister NR, Rubenstein DR. 2016. Environmental variability and the evolution of the glucocorticoid receptor (Nr3c1) in African starlings. *Ecol Lett* 19:1219–27.
- Homyack JA. 2010. Evaluating habitat quality of vertebrates using conservation physiology tools. *Wildlife Res* 37:332–42.
- IUCN. 2012. IUCN Red List Categories and Criteria: Version 3.1. Cambridge (UK): IUCN. p. 32.
- Jessop TS, Letnic M, Webb JK, Dempster T. 2013. Adrenocortical stress responses influence an invasive

- vertebrate's fitness in an extreme environment. *Proc Biol Sci* 280:20131444.
- Jessop TS, Woodford R, Symonds MR. 2013. Macrostress: do large-scale ecological patterns exist in the glucocorticoid stress response of vertebrates?. *Funct Ecol* 27:120–30.
- Johnson MA, Francis CD, Miller ET, Downs CJ, Vitousek MN. 2018. Detecting bias in large-scale comparative analyses: methods for expanding the scope of hypothesis-testing with HormoneBase. *Integr Comp Biol* published online (doi: 10.1093/icb/icy045).
- Ketterson ED, Atwell JW, McGlothlin JW. 2009. Phenotypic integration and independence: hormones, performance, and response to environmental change. *Integr Comp Biol* 49:365–79.
- Kilvitis HJ, Hanson H, Schrey AW, Martin LB. 2017. Epigenetic potential as a mechanism of phenotypic plasticity in vertebrate range expansions. *Integr Comp Biol* 57:385–95.
- Kleist NJ, Guralnick RP, Cruz A, Lowry CA, Francis CD. 2018. Chronic anthropogenic noise disrupts glucocorticoid signaling and has multiple effects on fitness in an avian community. *Proc Natl Acad Sci U S A* 115:E648–57.
- Korte SM, Koolhaas JM, Wingfield JC, McEwen BS. 2005. The Darwinian concept of stress: benefits of allostasis and costs of allostatic load and the trade-offs in health and disease. *Neurosci Biobehav Rev* 29:3–38.
- Lea J, Walker SL, Kerley GI, Jackson J, Matevich SC, Shultz S. 2018. Non-invasive physiological markers demonstrate link between habitat quality, adult sex ratio and poor population growth rate in a vulnerable species, the Cape mountain zebra. *Funct Ecol* 32:300–12.
- Lennox R, Cooke SJ. 2014. State of the interface between conservation and physiology: a bibliometric analysis. *Conserv Physiol* 2:cou003.
- Liebl AL, Martin LB. 2012. Exploratory behaviour and stressor hyper-responsiveness facilitate range expansion of an introduced songbird. *Proc Biol Sci* 279:4375–81.
- Liebl AL, Martin LB. 2013. Stress hormone receptors change as range expansion progresses in house sparrows. *Biol Lett* 9:20130181.
- Liebl AL, Martin LB. 2014. Living on the edge: range edge birds consume novel foods sooner than established ones. *Behav Ecol* 25:1089–96.
- Liebl AL, Shimizu T, Martin LB. 2013. Covariation among glucocorticoid regulatory elements varies seasonally in house sparrows. *Gen Comp Endocrinol* 183:32–7.
- Lindenmayer DB, Piggott MP, Wintle BA. 2013. Counting the books while the library burns: why conservation monitoring programs need a plan for action. *Front Ecol Environ* 11:549–55.
- Lindström KM, Hawley DM, Davis AK, Wikelski M. 2005. Stress responses and disease in three wintering house finch (*Carpodacus mexicanus*) populations along a latitudinal gradient. *Gen Comp Endocrinol* 143:231–9.
- Madliger CL, Love OP. 2014. The need for a predictive, context-dependent approach to the application of stress hormones in conservation. *Conserv Biol* 28:283–7.
- Madliger CL, Semeniuk CA, Harris CM, Love OP. 2015. Assessing baseline stress physiology as an integrator of environmental quality in a wild avian population: implications for use as a conservation biomarker. *Biol Conserv* 192:409–17.
- Martí O, Armario A. 1998. Anterior pituitary response to stress: time-related changes and adaptation. *Int J Dev Neurosci* 16:241–60.
- Martin LB. 2009. Stress and immunity in wild animals: timing is everything. *Gen Comp Endocrinol* 163:70–6.
- Martin LB, Brace AJ, Kilvitis HJ, Gervasi SS. 2016a. Invader endocrinology: the regulation of behaviour in pesky phenotypes. In: Weis JS, Sol D, editors. *Biological invasions and animal behaviour*. Cambridge, UK: Cambridge University Press. p. 26–46.
- Martin LB, Burgan S, Adelman JS, Gervasi SS. 2016b. Host competence: an organismal trait to integrate immunology and epidemiology. *Integr Comp Biol* 56:1225–37.
- Martin LB, Burgan SC, Adelman JS, Gervasi SS. 2016c. Host competence: an organismal trait to integrate immunology and epidemiology. *Integr Comp Biol* 56:1225–37.
- Martin LB, Cohen A. 2014. Physiological regulatory networks: the orchestra of life? In: Martin LB, Ghalambor CK, Woods HA, editors. *Integrative organismal biology*. Hoboken (NJ): John Wiley & Sons, Inc. p. 137–52.
- Martin LB, Fitzgerald L. 2005. A taste for novelty in invading house sparrows, *Passer domesticus*. *Behav Ecol* 16:702–7.
- Martin LB, Gilliam J, Han P, Lee K, Wikelski M. 2005. Corticosterone suppresses cutaneous immune function in temperate but not tropical house sparrows, *Passer domesticus*. *Gen Comp Endocrinol* 140:126–35.
- Martin LB, Hopkins WA, Mydlarz L, Rohr JR. 2010. The effects of anthropogenic global changes on immune functions and disease resistance. *Ann N Y Acad Sci* 1195:129–48.
- Martin LB, Liebl AL. 2014. Physiological flexibility in an avian range expansion. *Gen Comp Endocrinol* 206:227–34.
- Martin LB, Liebl AL, Kilvitis HJ. 2015. Covariation in stress and immune gene expression in a range expanding bird. *Gen Comp Endocrinol* 211:14–9.
- Martin LB, Liebl AL, Trotter JH, Richards CL, McCoy K, McCoy MW. 2011. Integrator networks: illuminating the black box linking genotype and phenotype. *Integr Comp Biol* 51:514–27.
- Martínez-Mota R, Valdespino C, Sánchez-Ramos M, Serio-Silva J. 2007. Effects of forest fragmentation on the physiological stress response of black howler monkeys. *Anim Conserv* 10:374–9.
- McCormick SD, Romero LM. 2017. Conservation endocrinology. *Bioscience* 67:429–42.
- McEwen BS, Wingfield JC. 2003. The concept of allostasis in biology and biomedicine. *Horm Behav* 43:2–15.
- McGlothlin JW, Ketterson ED. 2008. Hormone-mediated suites as adaptations and evolutionary constraints. *Philos Trans R Soc Lond B Biol Sci* 363:1611–20.
- Mullner A, Linsenmair KE, Wikelski M. 2004. Exposure to ecotourism reduces survival and affects stress response in hoatzin chicks (*Opisthocomus hoazin*). *Biol Conserv* 118:549–58.
- Nakagawa S, Schielzeth H. 2013. A general and simple method for obtaining R^2 from generalized linear mixed-effects models. *Methods Ecol Evol* 4:133–42.
- Ouyang JQ, Sharp PJ, Dawson A, Quetting M, Hau M. 2011. Hormone levels predict individual differences in reproductive success in a passerine bird. *Proc Roy Soc Lond B Biol Sci* 278:2537–45.

- Partecke J, Gwinner E. 2007. Increased sedentariness in European Blackbirds following urbanization: a consequence of local adaptation? *Ecology* 88:882–90.
- Patterson SH, Winkler DW, Breuner CW. 2011. Glucocorticoids, individual quality and reproductive investment in a passerine bird. *Anim Behav* 81:1239–47.
- Plummer M, Best N, Cowles K, Vines K. 2008. coda: Output analysis and diagnostics for MCMC. R package version 0.13-3 (<http://CRAN.R-project.org/package=coda>).
- Redpath SM, Young J, Evely A, Adams WM, Sutherland WJ, Whitehouse A, Amar A, Lambert RA, Linnell JDC, Watt A, et al. 2013. Understanding and managing conservation conflicts. *Trends Ecol Evol* 28:100–9.
- Rohr JR, Raffel TR, Halstead NT, McMahon TA, Johnson SA, Boughton RK, Martin LB. 2013. Early-life exposure to a herbicide has enduring effects on pathogen-induced mortality. *Proc Biol Sci* 280:20131502.
- Romero LM. 2002. Seasonal changes in plasma glucocorticoid concentrations in free-living vertebrates. *Gen Comp Endocrinol* 128:1–24.
- Romero LM, Dickens MJ, Cyr NE. 2009. The reactive scope model—a new model integrating homeostasis, allostasis, and stress. *Horm Behav* 55:375–89.
- Romero LM, Wingfield JC. 2015. Tempests, poxes, predators, and people: stress in wild animals and how they cope. Oxford University Press.
- Sapolsky RM, Romero LM, Munck AU. 2000. How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocr Rev* 21:55–89.
- Schoenle LA, Zimmer C, Vitousek MN. 2018. Understanding context dependence in glucocorticoid–fitness relationships: the role of the nature of the challenge, the intensity and frequency of stressors, and life history. *Integr Comp Biol* published online (doi: 10.1093/icb/icy046).
- Shelford VE. 1911. Physiological animal geography. *J Morphol* 22:551–618.
- Sorenson GH, Dey CJ, Madliger CL, Love OP. 2017. Effectiveness of baseline corticosterone as a monitoring tool for fitness: a meta-analysis in seabirds. *Oecologia* 183:353–65.
- Strasser EH, Heath JA. 2013. Reproductive failure of a human-tolerant species, the American kestrel, is associated with stress and human disturbance. *J Appl Ecol* 50:912–9.
- Taff CC, Schoenle LA, Vitousek MN. 2018. The repeatability of glucocorticoids: a review and meta-analysis. *Gen Comp Endocrinol* 260:136–45.
- Taff CC, Vitousek MN. 2016. Endocrine flexibility: optimizing phenotypes in a dynamic world? *Trends Ecol Evol* 31:476–88.
- Tarlow EM, Blumstein DT. 2007. Evaluating methods to quantify anthropogenic stressors on wild animals. *Appl Animal Behav Sci* 102:429–51.
- Valladares F, Matesanz S, Guilhaumon F, Araújo MB, Balaguer L, Benito-Garzon M, Cornwell W, Gianoli E, van Kleunen M, Naya DE, et al. 2014. The effects of phenotypic plasticity and local adaptation on forecasts of species range shifts under climate change. *Ecol Lett* 17:1351–64.
- Vera F, Zenuto R, Antenucci CD. 2017. Expanding the actions of cortisol and corticosterone in wild vertebrates: a necessary step to overcome the emerging challenges. *Gen Comp Endocrinol* 246:337–53.
- Vitousek MN, Johnson MA, Donald JW, Francis CD, Fuxjager MJ, Goymann W, Hau M, Husak JF, Kircher BK, Knapp R, et al. 2018. HormoneBase, a population-level database of steroid hormone levels across vertebrates. *Sci Data* 5:180097.
- Wada H. 2014. Developmental plasticity of individual variation in stress responses. In: Martin LB, Ghalambor CK, Woods HA, editors. *Integrative organismal biology*. Hoboken (NJ): John Wiley & Sons, Inc. p. 187–205.
- Walker BG, Boersma PD, Wingfield JC. 2005. Physiological and behavioral differences in Magellanic Penguin chicks in undisturbed and tourist-visited locations of a colony. *Conserv Biol* 19:1571–7.
- Wikelski M, Cooke SJ. 2006. Conservation physiology. *Trends Ecol Evol* 21:38–46.
- Wingfield JC, Maney DL, Breuner CW, Jacobs JD, Lynn S, Ramenofsky M, Richardson RD. 1998. Ecological bases of hormone–behavior interactions: the “emergency life history stage”. *Am Zool* 38:191–206.
- Wingfield JC, Sapolsky RM. 2003. Reproduction and resistance to stress: when and how. *J Neuroendocrinol* 15:711–24.
- Woods HA, Wilson JK. 2014. An elephant in the fog: unifying concepts of physiological stasis and change. In: Martin LB, Ghalambor CK, Woods HA, editors. *Integrative organismal biology*. Hoboken (NJ): John Wiley & Sons, Inc. p. 119–35.
- Zanette LY, White AF, Allen MC, Clinchy M. 2011. Perceived predation risk reduces the number of offspring songbirds produce per year. *Science* 334:1398–401.