

Anomalous elastic behavior of phase egg, $\text{AlSiO}_3(\text{OH})$, at high pressures

MAINAK MOOKHERJEE^{1,*}, WENDY R. PANERO², BERND WUNDER³, AND SANDRO JAHN⁴

¹Earth, Ocean and Atmospheric Sciences, Florida State University, Tallahassee, Florida 32306, U.S.A.

²School of Earth Sciences, Ohio State University, Columbus, Ohio 43210, U.S.A.

³Deutsches GeoForschungsZentrum GFZ, Telegrafenberg, 14473 Potsdam, Germany

⁴Institut für Geologie und Mineralogie, Zùlpicher Str. 49b, Universität zu Köln, 50674 Cologne, Germany

ABSTRACT

Phase egg, $[\text{AlSiO}_3(\text{OH})]$, is an aluminosilicate hydrous mineral that is thermodynamically stable in lithological compositions represented by $\text{Al}_2\text{O}_3\text{--SiO}_2\text{--H}_2\text{O}$ (ASH) ternary, i.e., a simplified ternary for the mineralogy of subducted sediments and continental crustal rocks. High-pressure and high-temperature experiments on lithological compositions resembling hydrated sedimentary layers in subducting slabs show that phase egg is stable up to pressures of 20–30 GPa, which translates to the transition zone to lower mantle depths. Thus, phase egg is a potential candidate for transporting water into the Earth's mantle transition zone. In this study, we use *first-principles* simulations based on density functional theory to explore the pressure dependence of crystal structure and how it influences energetics and elasticity. Our results indicate that phase egg exhibits anomalous behavior of the pressure dependence of the elasticity at mantle transition zone depths (~15 GPa). Such anomalous behavior in the elasticity is related to changes in the hydrogen bonding $\text{O-H}\cdots\text{O}$ configurations, which we delineate as a transition from a low-pressure to a high-pressure structure of phase egg. Full elastic constant tensors indicate that phase egg is very anisotropic resulting in a maximum anisotropy of compressional wave velocity, $\Delta v_p \approx 30\%$ and of shear wave velocity, $\Delta v_s \approx 17\%$ at zero pressures. Our results also indicate that the phase egg has one of the fastest bulk sound velocities (v_p and v_s) compared to other hydrous aluminous phases in the ASH ternary, which include topaz-OH, phase Pi, and $\delta\text{-AlOOH}$. However, the bulk sound velocity of phase egg is slower than that of stishovite. At depths corresponding to the base of mantle transition zone, phase egg decomposes to a mixture of $\delta\text{-AlOOH}$ and stishovite. The changes in compressional Δv_p and shear Δv_s velocity associated with the decomposition is ~0.42% and -1.23%, respectively. Although phase egg may be limited to subducted sediments, it could hold several weight percentages of water along a normal mantle geotherm.

Keywords: Phase egg, equation of state, elasticity, anisotropy, symmetric hydrogen bonding, subduction zone

INTRODUCTION

Based on estimates of planetary accretion, geochemistry, and the influx and outgassing of water, it is evident that our understanding of the size of the deep mantle reservoir of water is far from complete. Water is essential for the sustenance of planetary activities including melting (Hirschmann 2006) that eventually leads to large-scale geochemical differentiation. Water also helps in sustaining planetary geodynamics. In particular, water affects transport properties including the rheology (Mei and Kohlstedt 2000), viscosity (e.g., Ichikawa et al. 2015), and electrical conductivity (e.g., Wang et al. 2006a). Without water, the mantle convection processes for a dry silicate rock will be extremely sluggish. Therefore, it is important to estimate the size of such a water reservoir in the deep Earth. Knowing the extent of mantle hydration is also likely to enhance understanding of the role of solid Earth in influencing the sea level over long timescales (~10⁹ years). However, constraining the water content of the deep Earth is not straightforward. It is important to estimate how water is

transported into the deep Earth and the water-storage potential and stability of minerals and rocks at deep Earth conditions, which is an upper bound of possible water content. Estimation of actual water storage requires geophysical observations of elasticity and electrical conductivity of the deep Earth and the effect of water on those physical properties.

Most of the mantle consists of nominally anhydrous minerals (NAMs) with trace quantities of water, which are thermodynamically stable along a normal mantle geotherm, i.e., do not readily decompose at these temperatures. Hydrous mineral phases are distinct from the NAM phases and contain structurally bound hydroxyl groups, i.e., OH^- groups occur in well-defined crystallographic sites (Smyth 2006). Although, water is transported into the deep Earth via subduction of hydrated lithosphere containing hydrous mineral phases (Kawamoto 2006), a distinct limitation for hydrous phases as a potential reservoir for water over geological timescale is that they are not thermodynamically stable in a normal mantle geotherm. Hence, water is likely to be partitioned to NAMs whose water retention capacity is significantly lower. For example, while hydrous phases can host several weight percentages of water (Kawamoto 2006; Mookherjee et al. 2015),

* E-mail: mmookherjee@fsu.edu

at depths greater than 200 km, NAMs can host 70–700 ppm of H₂O as hydroxyl defects (Bolfan-Casanova 2005). However, based on high-pressure and high-temperature experiments it is known that aluminous hydrous phases have significantly greater thermal stability than the magnesio-silicate hydrous phases and are likely to be stable in normal mantle geotherms. The aluminous hydrous phases can be represented by the Al₂O₃-SiO₂-H₂O (ASH) ternary, i.e., a simplified ternary for the mineralogy of a subducted sedimentary layer (Peacock 1990; Ono 1999; Schmidt et al. 1998; Schreyer 1995; Wunder et al. 1993a, 1993b; Pamato et al. 2015). Several hydrous mineral phases are stable in the ASH ternary system. These include gibbsite [Al(OH)₃], diasporite (AlOOH), kaolinite [Al₂Si₂O₅(OH)₄], topaz-OH [Al₂SiO₄(OH)₂], phase Pi [Al₃Si₂O₇(OH)₃], phase egg [AlSiO₃(OH)], and dense high-pressure phases such as δ-AlOOH. In addition, recent experimental studies suggest that the dense hydrous magnesium silicate minerals phase-D [MgSi₂O₄(OH)₂] could have an aluminum-rich end-member with stoichiometry of Al₂SiO₄(OH)₂ (Pamato et al. 2015). It is estimated that the net water contribution from oceanic crust is likely to be greater than from the sediments (Peacock 1990). However, owing to the greater thermal stability of the minerals stabilized in the sedimentary layer, they may be effective in transporting water into the deep Earth in warmer subduction zones (Ono 1998). Among these hydrous phases, phase egg has the potential to carry water to mantle transition zone depths and, as a result, it has been the subject of numerous investigations that elucidate its high-pressure and temperature stability and compressibility. Phase egg was first synthesized above 10 GPa (Eggleton et al. 1978), and since then the thermodynamic stability have been investigated numerous times with the high-pressure limits ranging from 20–30 GPa where phase egg decomposes to δ-AlOOH and stishovite (Schmidt et al. 1998; Ono 1999; Sano et al. 2004; Fukuyama et al. 2017; Abe et al. 2018). Phase egg has also been found as inclusions in natural diamonds from Junia, Brazil, indicating that crustal materials are indeed deeply subducted to the mantle transition zone and upper part of lower mantle (Wirth et al. 2007). Although phase egg could potentially play an important role in transporting water into the deep Earth, its elasticity remains unknown. The latter is a crucial physical property that will help us to understand the mantle hydration caused by subduction of hydrated sediments or alumina rich crustal rocks.

In this study, we use *first-principles* simulations to explore how pressure affects the crystal structure, behavior of protons, equation of state, full elastic constant tensor, and elastic anisotropy of phase egg at high pressures.

METHODS

Phase egg, AlSiO₃(OH), is stable at pressures greater than 11 GPa and it has a monoclinic symmetry with space group *P*2₁/*n* (Schmidt et al. 1998). Based on the powder X-ray diffraction refinement of the crystal structure of phase egg, all Al and Si are located in the octahedral coordination with oxygen with one unique Si and Al site each in the unit cell (Schmidt et al. 1998). There are four crystallographically distinct positions for oxygen atoms: O(1), O(2), O(3), and O(4). The O(1) and O(2) oxygen atoms are bonded to two silicon (Si) and one aluminum (Al) atoms. The O(4) atom is bonded to the proton (H), two aluminum (Al) atoms, and a silicon (Si) atom. The structure consists of short chain segments connecting four edge-sharing octahedral units in the *a*-axis direction with Al-Si-Si-Al ordering in the *a*-*c* plane. These 4-member edge-sharing chains are linked in the *a*-axis direction by edge link between the AlO₆ and the neighboring SiO₆ octahedral unit (Fig. 1).

These two-dimensional sheets are then linked in the (011) direction through Al-Si corner linked octahedral unit. Hydrogen bonding bridges corners of AlO₆ units, also linking the 2D structures of the edge-sharing octahedral units. Based on ¹H-²⁹Si NMR spectra of phase egg, it has been suggested that there could be some disorder of Si and Al between the octahedral sites (Xue et al. 2006).

We performed first-principles quantum mechanical calculations based on the density functional theory (DFT) (Hohenberg and Kohn 1964; Kohn and Sham 1965). For the investigation of the phase egg, we employed a widely used approximation to the exchange-correlation functional: the generalized gradient approximation (GGA) and the highly accurate projector augmented wave method (PAW) as implemented in the Vienna ab initio simulation package (VASP) (Kresse and Hafner 1993; Kresse and Furthmüller 1996a, 1996b; Perdew et al. 1996; Kresse and Joubert 1999). First, we performed a series of convergence tests by varying the energy cutoff and *k*-points. We found that an energy cut-off $E_{\text{cut}} = 800$ eV and a *k*-point mesh of $6 \times 9 \times 6$ Monkhorst-Pack grid (Monkhorst and Pack 1976) with 100 irreducible *k*-points is sufficient for describing the energetics of phase-egg (Supplemental¹ Fig. S1). We also used the van der Waals corrections as implemented in VASP (Grimme 2006). All computations were performed on the primitive unit cell of phase egg with the starting guess from X-ray diffraction crystal structure (Schmidt et al. 1998). To test the assertion that the Al- and Si-octahedral sites may be disordered (Xue et al. 2006), we have also explored the effect of octahedral cation disorder in which we exchanged 25 and 50% of the Al and Si octahedral sites.

The elastic tensor was derived using the finite strain approach. Starting from the optimized cell at a given pressure, we have strained the lattice parameters by 1% and then relaxed the atomic positions and computed the corresponding stress tensor. Using Hooke's law, the elastic constants were derived from the stress-strain relations, as outlined in previous studies (Chheda et al. 2014). Finally, we computed the single-crystal azimuthal anisotropy for compressional velocity (A_{vp}) i.e., $[(v_p^{\text{max}} - v_p^{\text{min}}) \times 200 / (v_p^{\text{max}} + v_p^{\text{min}})]$ and shear velocity (A_{vs}) i.e., $[(v_s^{\text{max}} - v_s^{\text{min}}) \times 200 / (v_s^{\text{max}} + v_s^{\text{min}})]$ of phase-egg using the formulation for maximum polarization anisotropy (Mainprice 1990).

RESULTS

Crystal structure

The converged structure has a monoclinic symmetry and space group (no. 14) *P*2₁/*c*. The crystallographic space group *P*2₁/*n* is an alternate setting for *P*2₁/*c*. In space group *P*2₁/*n*, the translation is along a diagonal on the plane perpendicular to 2₁-screw axis, whereas the translation is along the *c*-axis on the plane perpendicular to the 2₁-screw axis. We note that the disordered phase has higher enthalpy than the ordered state by 0.5 eV, or a Gibbs free energy difference between the ordered and the disordered states at 1800 K of ~0.2 eV/molecular unit (19 kJ/mol). The enthalpy and Gibbs free energy difference remain largely insensitive to pressures (Supplemental¹ Fig. S2). Since the energetic cost of disordering is slightly greater than




FIGURE 1. Crystal structure of phase egg: (a) projection down the *b*-axis, and (b) projection down the *a*-axis. The octahedral units, i.e., AlO₆ and SiO₆, are represented by gray and light blue spheres, respectively. The hydrogen atoms and the oxygen atoms are shown as silver and red spheres, respectively. (Color online.)

the ordered state, we have used the ordered crystal structure for the determination of the full elastic constant tensor.

At ambient conditions, the crystal structure of phase egg has a single proton (H) with fractional coordinates of $x_H = 0.796$, $y_H = 0.553$, $z_H = 0.422$. The proton forms a hydroxyl O(4)-H bond with the O(4) atom with fractional coordinate of $x_{O(4)} = 0.760$, $y_{O(4)} = 0.215$, $z_{O(4)} = 0.129$. The proton also forms a hydrogen bond H...O with the O(3) atom with fractional coordinate of $x_{O(3)} = 0.880$, $y_{O(3)} = 0.201$, $z_{O(3)} = 0.512$ (Schmidt et al. 1998). Our results indicate that the O(4)-H...O(3) hydrogen bond is strengthened at high pressures, the O(4)-H bond is enhanced, and H...O(3) length decreases. At around ~15 GPa, the proton flips its position and is transferred to the O(3) atom forming an O(3)-H group and an O(4)...H bond (Fig. 2). This is slightly different from more conventional symmetrization of hydrogen bonds at high pressures where the proton is at the center of the O...O forming a O-H-O unit. At pressures lower than the proton

transfer ~15 GPa, we define the crystal structure as the phase egg (LP), and at higher pressure, we define the crystal structure as phase egg (HP), where “LP” and “HP” refers to low- and high-pressures, respectively. The energetics and properties of the phase egg (HP) extrapolated to pressures lower than 15 GPa differs significantly from that of phase egg (LP). At pressures >15 GPa the structures are identical (Fig. 2). Our pressure volume results and the evolution of lattice parameter upon compression are in good agreement with previous experiments (Vanpeteghem et al. 2003). High-pressure symmetrization of hydrogen bond has been reported in many hydrous minerals including ice phases (Goncharov et al. 1996); δ -AlOOH (Tsuchiya et al. 2002; Panero and Stixrude 2004; Tsuchiya and Tsuchiya 2009; Sano-Furukawa et al. 2009; Mashino et al. 2016); δ -CrOOH analog to the δ -AlOOH phase (Jahn et al. 2012); ϵ -FeOOH (Otte et al. 2009); phase D (Tsuchiya et al. 2005; Hushur et al. 2011); phase H (Tsuchiya and Mookherjee 2015; Panero and Caracas 2017) and phase Pi (Peng et al. 2017). For symmetric hydrogen bonds, often the O-H...O angle is ~180° or close to a linear arrangement. However, for phase egg we notice that the O(4)-H...O(3) angle to be around 160° at pressures where proton transfer occurs (~15 GPa), leading to a new configuration of O(4)...H-O(3) (Fig. 2). In a recently examined hydrous aluminosilicate phase Pi, a similar observation was made, few of the O-H...O configurations remained nonlinear even after compressions, however, the O-H...O configuration that exhibited symmetric hydrogen bonding at high pressure became linear (Peng et al. 2017). In addition, a new monoclinic phase of AlOOH has been predicted which has nonlinear and bent O-H-O linkages instead of symmetric hydrogen bonding at high pressures (Zhong et al. 2016).

Equation of state

The volume dependence of the total energy obtained using first-principles simulation are adequately described by a third-order Birch-Murnaghan equation of state (Birch 1978), where total energy, E is expressed as

$$E = E_0 + \frac{9}{2} K_0 V_0 \left[f_v^2 + (K'_0 - 4) f_v^3 \right] \quad (1)$$

where, E_0 , K_0 , V_0 , and K'_0 represents the ground state energy, bulk modulus at zero-pressure, unit-cell volume at zero pressure, and pressure derivative of zero-pressure bulk modulus, respectively. The Eulerian finite strain (f_v) and is defined as

$$f_v = \frac{1}{2} \left[\left(\frac{V_0}{V} \right)^{\frac{2}{3}} - 1 \right] \quad (2)$$

We find that the phase egg (LP) has lower energy than that of the phase egg (HP). Upon compression, the energies of the two crystal structures approach each other gradually and become indistinguishable at a volume ~196 Å³ corresponding to a pressure ~15 GPa. The energy vs. volume relationship of phase egg (LP) and phase egg (HP) suggest that at lower pressures the asymmetric arrangement of hydrogen, i.e., O(4)-H...O(3) is favored, at pressures greater than 15 GPa, the proton flips its position to O(4)...H-O(3). The energy vs. volume relationship suggests a second-order phase transition (Fig. 3). Similar second-order transitions characterized by proton order-disorder




FIGURE 2. (a) Plot of hydroxyl bond length: $d_{O(4)-H}$, hydrogen bond length: $d_{O(4)-H...O(3)}$, and separation of the oxygen atoms, i.e., O(4)...O(3), vs. unit-cell volume and unit-cell pressure. (b) Hydrogen-hydrogen repulsion denoted by H...H separation distances. (c) The hydrogen bond angle, O(4)-H...O(3). The blue and red symbols represent the low-pressure (LP) and high-pressure (HP) phase egg. (Color online.)

behavior and/or hydrogen bond symmetrization have been documented for condensed phases including several H₂O-ice phases (Schweizer and Stilling 1980; Lee et al. 1993), dense hydrous magnesium silicate minerals phase-D (Tsuchiya et al. 2005), phase-H (Tsuchiya and Mookherjee 2015), and δ -AlOOH aluminum oxy-hydroxide (Tsuchiya et al. 2002; Panero and Stixrude 2004; Tsuchiya and Tsuchiya 2009). Such second-order transitions are often associated with discontinuity in second-order derivatives of energy, such as bulk modulus (Tsuchiya et al. 2002; Sano-Furukawa et al. 2009; Hushur et al. 2011). In contrast, the pressure-volume relations for phases exhibiting second-order transitions related to symmetrization of hydrogen bonds are often continuous with slight or no difference in the zero-pressure volume between the low- and high-pressure phases (e.g., phase-D: Hushur et al. 2011).

The bulk modulus at zero pressure (K_0^{LP}), unit-cell volume at zero pressure (V_0^{LP}), and the pressure derivative of zero-pressure bulk modulus ($K_0'^{LP}$) for phase egg (LP), i.e., at pressures below the transfer of a proton, are 164.4 (± 1.8) GPa, 210.21 (± 0.14) Å³, and 7.14 (± 0.24), respectively. The bulk modulus at zero-pressure (K_0^{HP}) unit-cell volume at zero pressure (V_0^{HP}) and the pressure derivative of zero-pressure bulk modulus ($K_0'^{HP}$) for

phase egg (HP) at pressures above the transfer of proton are, 222.8 (± 0.2) GPa, 207.74 (± 0.01) Å³, and 4.44 (± 0.02), respectively. The zero-pressure volume of the LP phase egg is $\sim 1.2\%$ larger than the HP phase egg. In contrast, the zero pressure bulk modulus of LP phase egg is $\sim 35.5\%$ softer than the HP phase egg. This behavior is similar to what has been observed in the equation of state studies in phase-D where the zero pressure volume of the hydrogen off-centered (HOC) phase is $\sim 2.1\%$ larger than the hydrogen centered (HC) phase (Hushur et al. 2011). In contrast, the zero pressure bulk modulus of the HC phase is $\sim 18\%$ stiffer than the HOC phase (Hushur et al. 2011).

The experimentally determined equation of state parameters for phase egg, i.e., (K_0^{exp}), (V_0^{exp}), and ($K_0'^{exp}$) are 157.0 (± 4.0) GPa, 212.99 (± 0.01) Å³, and 6.5 (± 0.4), respectively (Vanpeteghem et al. 2003). The unit-cell volume at zero pressure for phase egg (LP) determined by first-principles simulation is 1.3% smaller than the experimental results. Whereas, the bulk modulus at zero-pressure determined by first-principles simulation is 4.7% greater than the experimental results. This is expected because the experiments are at 300 K and the first-principles simulations are at static conditions (0 K). The variation of the lattice parameters, i.e., a , b , c , β with respect to the unit-cell volume for the low-pressure and the high-pressure phase egg determined using first-principles simulations agree with the experimental results (Fig. 3, Supplemental¹ Fig. S3). The linear moduli at zero pressure for the lattice parameter: a , b , and $c^* = c \times \sin\beta$, i.e., K_a^{LP} , K_b^{LP} , and $K_{c^*}^{LP}$ for the phase egg (LP) are 975.6 (± 1.2), 344.6 (± 0.6), and 531.1 (± 0.8) GPa, respectively. The linear moduli at zero-pressure for the lattice parameter— a , b , and c^* , i.e., K_a^{HP} , K_b^{HP} , and $K_{c^*}^{HP}$ for the phase egg (HP) are 815.6 (± 2.8), 507.9 (± 0.7), and 789.6 (± 1.1) GPa (Supplemental¹ Fig. S4). The proton transfer stiffens with $K_b^{HP} > K_b^{LP}$ and $K_{c^*}^{HP} > K_{c^*}^{LP}$ by $\sim 32\%$. This stiffening is likely related to the proton transfer and the significant changes of compressibility along the b - and c^* -lattice directions are related to the orientation of the bent and nonlinear O-H $\cdots$ O bonds. Upon compression, of phase egg exhibit anomalous elastic behavior at ~ 15 GPa, most likely related to the observed proton transfer. Our results on equation of state and evolution of lattice parameters with pressure are in good agreement with previous experimental results (Vanpeteghem et al. 2003). The experimental results showed a strong anisotropic behavior with larger compressibility along the b -axis compared to the a - and c -axes directions. The measured change in the slope of compressibility in b -direction with a significant increase in stiffness above ~ 19 GPa, was identified as a phase transition due to a hydrogen bond symmetrization (Sikka 2007).




FIGURE 3. (a) Plot of total energy, (b) pressure, (c) lattice parameters a , (d) lattice parameter b , (e) lattice parameter c , and (f) lattice parameter β , for low-pressure (LP) (blue symbols) and high-pressure (HP) (red symbols) phase egg vs. unit-cell volume. The gray filled symbols are from X-ray diffraction results (Vanpeteghem et al. 2003). Unit-cell volume of (LP) (blue symbols) and high-pressure (HP) (red symbols) phase egg is also plotted as a function of pressure and compared with experimental data (gray filled symbols) (Supplemental¹ Fig. S3). The volume/pressure dependence of the lattice parameters is characteristics of a second-order transition and has been documented in several hydrous phases (Tsuchiya et al. 2002, 2005; Tsuchiya and Mookherjee 2015). (Color online.)

Elasticity

Monoclinic phase egg ($P2_1/c$) has 13 non-zero independent elastic constants (Nye 1985). Upon compression almost all the individual components of the full elastic constant tensor become stiffer except the off-diagonal components C_{25} , C_{35} , and C_{46} (Fig. 4). The pressure dependence of the elasticity data can be described with the finite strain formulation (Karki et al. 2001) (Supplemental¹ Table S1)

$$C_{ijkl} = (1 + 2f_v)^2 \left[C_{ijkl0} + b_1 f_v + 2b_2 f_v^2 \right] - P \Delta_{ijkl} \quad (3)$$

where f_e is the finite Eulerian strain as defined in Equation 2

$$b_1 = 3K_0(C'_{ijkl0} + \Delta_{ijkl}) - 7C_{ijkl0} \quad (4)$$

$$b_2 = 9K_0^2 C''_{ijkl0} + 3K_0'(b_1 + 7C_{ijkl0}) - 16b_1 - 49C_{ijkl0} \quad (5)$$

and

$$\Delta_{ijkl} = -\delta_{ij}\delta_{kl} - \delta_{ik}\delta_{jl} - \delta_{il}\delta_{jk} \quad (6)$$

where C'_{ijkl0} and C''_{ijkl0} are the first and second derivatives of C_{ijkl0} , with respect to pressure at ambient conditions. Δ_{ijkl} is equal to -3 for the principal constants (C_{iiii} in full tensor and C_{ii} in Voigt notation, with $i = 1, 2, 3$), -1 for the off-diagonal elastic constants (C_{ijij} in full tensor and C_{ij} in Voigt notation, with $i = 1, 2, 3$; $i \neq j$), -1 for the shear constants (C_{ijij} in full tensor notation with $i = 1, 2, 3$; $i \neq j$ and C_{ij} in Voigt notation with $i = 4, 5, 6$; $i = j$), and 0 otherwise. δ_{ij} is the Kronecker δ ($\delta_{ij} = 1$ for $i = j$, and $\delta_{ij} = 0$, for $i \neq j$).

At around ~ 15 GPa, i.e., the pressure corresponding to mantle transition zone, many components of the full elastic constant tensor exhibit anomalous behavior. At pressure greater than 15 GPa, all components of the elastic constants become stiffer. This anomalous behavior in elasticity is very likely related to the changes in the hydrogen bonding, in particular, the proton attached to O(4) atom flips to the O(3) atom. We analyzed the elasticity phase-egg (LP) for pressures below

the proton transfer (~ 15 GPa). And we also determined the elasticity of phase egg (HP) for pressure > 15 GPa. We note that at low pressures, there is a significant difference between several components of the elastic constant tensor between the phase-egg (LP) and the metastable extension of elasticity of phase egg (HP) to pressures below 15 GPa. At pressures greater than 15 GPa, elasticity of phase egg (LP) and phase egg (HP) are identical (Fig. 4). At zero-pressure conditions the principal elastic constants exhibit the relationship: $C_{11}^{\text{LP}} > C_{33}^{\text{LP}} > C_{22}^{\text{LP}}$. The same relationship holds for the zero-pressure metastable extension of high-pressure phase egg (HP), i.e., $C_{11}^{\text{HP}} > C_{33}^{\text{HP}} > C_{22}^{\text{HP}}$. We also note that the $C_{11}^{\text{HP}} \sim C_{11}^{\text{LP}}$, $C_{33}^{\text{HP}} > C_{33}^{\text{LP}}$ by 14% and $C_{22}^{\text{HP}} > C_{22}^{\text{LP}}$ by 30%. This is consistent with the fact that the orientation of the hydrogen bond is mostly aligned along the b -axis but is tilted such that it has a component along the c -axis of the crystal structure. The crystallographic a -axis direction remains largely unaffected by the proton transfer. We also note that at pressures lower than the proton transfer pressures (~ 15 GPa), the pressure derivatives for the principal elastic constants $C_{ii}^{\text{LP}} > C_{ii}^{\text{HP}}$ by 19, 29, and 58% for $i = 1, 2$, and 3, respectively.

For the shear elastic components i.e., $i = 4, 5, 6$, $i = j$, at zero pressure conditions, the following relations holds $C_{55}^{\text{LP}} > C_{66}^{\text{LP}} > C_{44}^{\text{LP}}$. Similar relationship holds for the zero-pressure metastable extension of phase egg (HP): $C_{55}^{\text{HP}} > C_{66}^{\text{HP}} > C_{44}^{\text{HP}}$. However, unlike the principal components, the shear components have different relationships between the low-pressure phase egg (LP) and the metastable extension of high-pressure phase egg (HP), $C_{44}^{\text{HP}} \sim$




FIGURE 4. Full elastic constant tensor (C_{ij}) components, for phase egg as a function of pressure (Supplemental¹ Table S1). There are 12 subplots and an inset, representing the 13 independent elastic constants for the monoclinic symmetry. The low-pressure (LP) and high-pressure (HP) phase egg is represented by blue and red symbols, respectively. (Color online.)

C_{44}^{LP} , $C_{66}^{HP} < C_{66}^{LP}$ by 1.8% and $C_{55}^{HP} < C_{55}^{LP}$ by 2.4%. We also note that at $P < 15$ GPa, the pressure derivatives for the shear elastic constants $C_{ii}^{LP} > C_{ii}^{HP}$ by 33, 28, and 28% for $i = 4, 5$, and 6, respectively.

Our results show that the phase egg exhibits significant elastic anisotropy (Fig. 5). The P-wave azimuthal and S-wave anisotropy for phase egg for the phase egg (LP) are $Av_p \sim 30\%$ and $Av_s \sim 17\%$ at 0 GPa. The anisotropy reduces at high pressure and then increases closer to the proton transition pressures ~ 15 GPa. At pressures greater than 15 GPa, $Av_p \sim 10\%$ and $Av_s \sim 10\%$ and remains unchanged at higher pressures (Fig. 5).

The isotropic bulk (K) and shear (G) moduli are determined using the relations

$$K_V = \left(\frac{1}{9} \right) [C_{11} + C_{22} + C_{33} + 2(C_{12} + C_{13} + C_{23})] \quad (7)$$

$$K_R = [s_{11} + s_{22} + s_{33} + 2(s_{12} + s_{13} + s_{23})]^{-1} \quad (8)$$

$$G_V = \left(\frac{1}{15} \right) [C_{11} + C_{22} + C_{33} - (C_{12} + C_{13} + C_{23}) + 3(C_{44} + C_{55} + C_{66})] \quad (9)$$

$$G_R = 15 [4(s_{11} + s_{22} + s_{33}) - (s_{12} + s_{13} + s_{23}) + 3(s_{44} + s_{55} + s_{66})]^{-1} \quad (10)$$

where s_{ijkl} represents the elastic compliance tensor and is related C_{ijkl} by the relation $s_{ijkl}C_{ijkl} = \delta_{im}\delta_{jn}$, subscript V , and R represents Voigt and Reuss bounds.

The Voigt limit of the bulk (K_V^{HP}) and shear (G_V^{HP}) moduli at zero pressure for phase egg (HP) are 229 and 159 GPa, respectively. In contrast, the Voigt limit of the bulk (K_V^{LP}) and shear (G_V^{LP}) moduli at zero-pressure for low-pressure phase egg (LP) are 201 and 151 GPa, respectively. Thus, the Voigt limit for the phase egg (HP) is greater than phase egg (LP) by 14 and 6%, respectively. And the Reuss limit of the bulk (K_R^{HP}) and shear (G_R^{HP}) moduli for the phase egg (HP) are 223 and 157 GPa, respectively. In contrast, the Reuss limit of the bulk (K_R^{LP}) and shear (G_R^{LP}) moduli for the phase egg (LP) are 165 and 142 GPa, respectively. Thus, the Reuss limit for the phase egg (HP) is greater than phase egg (LP) by $\sim 35\%$ and 10%, respectively.

DISCUSSION

The mineral phases in the Al_2O_3 - SiO_2 - H_2O (ASH) ternary show correlation between velocity and density. Stishovite and corundum are the densest phase and have the fastest bulk sound velocity. Among the hydrous phases, phase egg (~ 7.5 wt% H_2O) and δ - $AlOOH$ (~ 15 wt% H_2O) are the densest and also have fast sound wave velocity (Table 1; Fig. 6). Along the SiO_2 - $AlOOH$ join, we note that the zero-pressure density follows the trend: $\rho_0^0 < \rho_0^{top-o} < \rho_0^{\delta-LP} < \rho_0^{egg-LP} < \rho_0^{st}$. Also, along the SiO_2 - $AlOOH$ join, the compressional velocity (v_p) extrapolated to zero pressure, the following trend is observed $v_p^{st} < v_p^{top-o} < v_p^{\delta-LP} < v_p^{egg-LP} < v_p^{st}$. Similar trends for shear (v_s) velocity extrapolated to zero pressure i.e., $v_s^{st} < v_s^{top-o} < v_s^{\delta-LP} < v_s^{egg-LP} < v_s^{st}$ is also observed (Table 1; Fig. 6). The observed trend in the density and shear wave velocity could be explained in terms of the crystal structures. For instance, the crystal structure of phase-Pi that is stable at relatively low pressures between 2 and 7 GPa (Wunder et al. 1993a, 1993b) consists of layers of distorted eight-membered rings formed by AlO_6 units alternating with layers consisting of SiO_4 tetrahedral units (Peng et al. 2017). Crystal structure of topaz-OH that is stable up to 12 GPa (Ono 1998) also consists of edge-sharing AlO_6 octahedral units forming crankshaft chains and SiO_4 tetrahedral units sharing corners. In contrast, phase egg is formed at higher pressures and has both aluminum and silicon in octahedral coordination. The zero-pressure density of phase egg (LP) is slightly greater than the expected density based on an ideal solid solution of SiO_2 and δ - $AlOOH$ (LP). The δ - $AlOOH$ (LP) refers to the hydrogen in off-center position (HOC), i.e., asymmetric hydrogen bond at low-pressure condition (Tsuchiya and Tsuchiya 2009). The compressional velocity (v_p) and shear velocity (v_s) extrapolated to zero pressure for phase egg (LP) is lower than the expected v_p and v_s based on an ideal solid solution of SiO_2 and δ - $AlOOH$ (LP) by 6.3 and 6.5%, respectively (Fig. 6).

IMPLICATIONS

In a simplified ASH system, at the base of the mantle transition zone and along a warm subduction zone geotherm, phase-egg




FIGURE 5. (a) Plot of elastic anisotropy Av_p and Av_s for phase egg as a function of pressure. The low-pressure (LP) and high-pressure (HP) phase egg is represented by blue and red symbols, respectively. The stereographic projection down the X_3 axes for Av_p , Av_s , and v_s polarization planes at (b) ~ 0 GPa and (c) ~ 23 GPa. (Color online.)

TABLE 1. Density, compressional (v_p), and shear (v_s) wave velocity of mineral phases in Al_2O_3 - SiO_2 - H_2O ternary

Mineral	Abbreviation	Stoichiometry	ρ (g/cm ³)	v_p (km/s)	v_s (km/s)	H_2O (wt%)	Reference
Corundum	cor	Al_2O_3	3.95	10.94	6.41	0.00	Ohno et al. (1986)
Quartz	qz	SiO_2	2.64	6.09	4.12	0.00	Ohno et al. (2006)
Coesite	cs	SiO_2	2.93	8.17	4.58	0.00	Weidner and Carleton (1977)
Stishovite	st	SiO_2	4.30	11.91	7.28	0.00	Jiang et al. (2009)
Ice	ice-iii	H_2O	1.16	3.66	2.01	100.00	Tulk et al. (1994)
Andalusite	and	Al_2SiO_5	3.15	9.76	5.65	0.00	Vaughan and Weidner (1978)
Sillimanite	sil	Al_2SiO_5	3.24	9.65	5.42	0.00	Vaughan and Weidner (1978)
Kyanite	ky	Al_2SiO_5	3.76	9.68	5.87	0.00	Winkler et al. (2001)
Diaspore	dia	AlOOH	3.38	9.42	5.83	15.00	Jiang et al. (2008)
δ -AlOOH	δ -loP	AlOOH	3.39	9.82	6.19	15.00	Tsuchiya and Tsuchiya (2009)
δ -AlOOH	δ -hiP	AlOOH	3.47	10.87	6.60	15.00	Tsuchiya and Tsuchiya (2009)
Kaolinite	kl	$\text{Al}_2\text{Si}_2\text{O}_5(\text{OH})_4$	2.52	6.23	3.55	18.19	Katahara (1996)
Topaz-OH	top-m	$\text{Al}_2\text{SiO}_4(\text{OH})_2$	3.43	9.75	5.81	10.00	Mookherjee et al. (2016)
Topaz-OH	top-o	$\text{Al}_2\text{SiO}_4(\text{OH})_2$	3.39	9.51	5.51	10.00	Mookherjee et al. (2016)
Phase-pi	pi	$\text{Al}_3\text{Si}_2\text{O}_7(\text{OH})_3$	3.21	8.86	5.28	9.00	Peng et al. (2017)
Phase-egg	egg-LP	$\text{AlSiO}_3(\text{OH})$	3.79	9.99	6.22	7.50	This study
Phase-egg	egg-HP	$\text{AlSiO}_3(\text{OH})$	3.84	10.68	6.42	7.50	This study





FIGURE 6. (a) A ternary plot with Al_2O_3 - SiO_2 - H_2O as the end-member components, relevant for the mineral phases stable in subducted sediments. Also shown is the blue shaded region along the line joining the SiO_2 and AlOOH with a series of hydrous phases, δ - AlOOH , topaz-OH, phase egg, phase-pi, and stishovite. Plot of (b) v_p and (c) v_s as a function of density for the mineral phases in the ASH ternary (Table 1). The vertical gray-brown shaded line demonstrates that the density of low-pressure (LP) and high-pressure (HP) phase egg are very similar, but the velocities are distinct. This is related to the fact that second-order transition associated with the proton transfer affects the second derivative of energy, i.e., elastic constants but the volume and density remains mostly unaffected by the proton transfer. The blue and red dashed line represents linear regression fits for the compressional wave, v_p is given by $[2.77(\pm 0.49) \times \rho + 0.07(\pm 1.64)]$ and for the shear wave, v_s is given by $[1.74(\pm 0.27) \times \rho - 0.23(\pm 0.91)]$. The light blue and red lines represent 95% confidence level for the primary and shear velocity and density regressions. Plot of zero pressure (d) density (e) v_p and v_s for hydrous mineral phase along the SiO_2 and AlOOH join. (Color online.)

decomposes to a mixture of aluminum oxyhydroxide (δ - AlOOH) and stishovite (SiO_2) via the reaction-



This is likely to be associated with Δv_p and Δv_s of 0.42 and -1.23% , respectively. Among all the hydrous phases in the ASH ternary system, Phase egg has one of the fastest compressional

(v_p) and shear (v_s) velocity (Fig. 7). The velocity of phase egg is slower than stishovite but remains faster than δ - AlOOH phase at all the pressures within phase egg's thermodynamic stability field. The δ - AlOOH phase with symmetric hydrogen bonding has sound wave velocities faster than phase egg, however, that is likely to occur at pressures corresponding to the lower mantle. In our analysis, we have used v_p and v_s of δ - AlOOH (Tsuchiya and Tsuchiya 2009) and stishovite (Karki et al. 1997). The static sound wave velocity results from the first-principles studies are in very good agreement with experimental studies including static




FIGURE 7. (a) Plot of K_H and G_H , i.e., Hill averaged (average of Voigt and Reuss limits) for low-pressure (LP) and high-pressure (HP) phase egg represented by blue and red symbols, respectively. Also shown is an experimentally determined bulk modulus in gray symbol (Vanepetghem et al. 2003). (b) Plot of pressure dependence of v_p and v_s for low-pressure (LP) and high-pressure (HP) phase egg and other mineral phases including, stishovite (Karki et al. 1997) topaz-OH (Mookherjee et al. 2016), phase Pi (Peng et al. 2017), and δ -AlOOH (Tsuchiya and Tsuchiya 2009) that are likely stable in subducted sediments. (c) A pressure-temperature-depth phase diagram illustrating the stability of phase egg. The figure is modified from a previous study (Wirth et al. 2007). Also shown is the green-shaded area representing the mantle transition zone. The blue shaded band at ~15 GPa indicates the predicted pressure for the proton transfer, i.e., the boundary between phase egg (LP) and phase egg (LP) associated with the elastic anomaly as observed in our study. The dashed circle at the intersection of warm subduction geotherm with the high-pressure stability of phase egg (HP), where phase egg will decompose to δ -AlOOH and stishovite is likely to be associated with Δv_p and Δv_s of 0.42 and -1.23%, respectively. (Color online.)

Brillouin scattering experiments on stishovite (Jiang et al. 2009) and δ -AlOOH (Mashino et al. 2016).

The temperature of decomposition of phase egg at the base of the transition zone along a warmer subduction zone geotherm is ~1000 °C at ~22 GPa (Wirth et al. 2007). It is likely that the temperature will affect the bulk sound velocity of all the hydrous phases including phase egg. The effect of temperature on the elasticity and seismic velocities of the hydrous phases in ASH ternary remains mostly unknown, except for stishovite (Yang and Wu 2014). The effect of temperatures on v_p and v_s for natural topaz containing both hydrogen and fluorine have been recently constrained and are of the order of $dv_p/dT \sim -3.10 \times 10^{-4}$ and $dv_s/dT \sim -2.30 \times 10^{-4}$ (Tennakoon et al. 2018). The effect of temperature on the elasticity of phase egg (LP) and its high-pressure polymorph, phase egg (HP) across the proton transfer needs to be constrained in future work. Presence of minor amounts of fluorine (F⁻) has been speculated in mantle minerals in the transition zone (Grützner et al. 2018). Substitution of OH⁻ by fluorine (F⁻) occurs in natural minerals including topaz and recent study shows that such substitution does indeed influence the bulk sound velocity (Tennakoon et al. 2018).

The sound wave velocity of phase egg is significantly faster than the major mantle phases such as wadsleyite (Zha et al. 1997), ringwoodite (Li 2003), pyrope (Sinogeikin and Bass 2000), and majoritic garnet (Irifune et al. 2008) that are stable

in the transition zone. Few weight percentages of water tend to reduce the seismic velocity of wadsleyite (Mao et al. 2008) and ringwoodite (Panero 2010; Wang et al. 2006b). Phase egg could host almost 7.5 wt% of water and yet still have significantly faster velocity compared to hydrous wadsleyite (Mao et al. 2008) and ringwoodite (Wang et al. 2006b). Although the presence of phase egg is likely to be limited mostly to subducted sediments, it could store several weight percentages of water and be stable in a normal mantle geotherm and may not be readily detectable. Certainly, more research is warranted to have better constraints on the effect of temperature, anisotropy, petrographic fabric, or lattice preferred orientations on the seismic velocity of phase egg and other mineral phases in subducted sediments.

FUNDING

M.M. acknowledges the funding support from U.S. National Science Foundation grant (EAR-1634422; EAR-1753125), Ralph E. Powe Junior Faculty Enhancement Award, Oak Ridge Associated University (ORAU), and computing resources from the Extreme Science and Engineering Discovery Environment (XSEDE), which is supported by National Science Foundation grant number OCI-1053575. W.R.P. acknowledges the funding support from the National Science Foundation CAREER grant (EAR-60023026).

ACKNOWLEDGMENTS

W.R.P. acknowledges computational time from the Ohio Supercomputer Center (PAS-0238-1). Authors acknowledge constructive reviews/comments from Zhicheng Jing, Zhongqing Wu, and an anonymous reviewer.

REFERENCES CITED

- Abe, R., Shibazaki, Y., Ozawa, S., Ohira, I., Tobe, H., and Suzuki, A. (2018) In-situ X-ray diffraction studies of hydrous aluminosilicate at high pressure and temperature. *Journal of Mineralogical and Petrological Sciences*, 113, 106–111.
- Birch, F. (1978) Finite strain isotherm and velocities for single crystal and polycrystalline NaCl at high-pressures and 300 K. *Journal of Geophysical Research*, 83, 1257–1268.
- Bolfan-Casanova, N. (2005) Water in the Earth's mantle. *Mineralogical Magazine*, 69, 229–257.
- Chheda, T., Mookherjee, M., Mainprice, D., dos Santos, A.M., Molaison, J.J., Chantel, J., Manthilake, G., and Bassett, W.A. (2014) Structure and elasticity of phlogopite under compression: Geophysical implications. *Physics of the Earth and Planetary Interior*, 233, 1–12.
- Eggelton, R.A., Boland, J.N., and Ringwood, A.E. (1978) High pressure synthesis of a new aluminosilicate: $\text{Al}_2\text{Si}_2\text{O}_7(\text{OH})$. *Geochemical Journal*, 12, 191–194.
- Fukuyama, K., Ohtani, E., Shibazaki, Y., Kagi, H., and Suzuki, A. (2017) Stability field of phase egg, AlSiO_3OH at high pressure and high temperature: possible water reservoir in mantle transition zone. *Journal of Mineralogical and Petrological Sciences*, 112, 31–35.
- Goncharov, A.F., Struzhkin, V.V., Somayazulu, M.S., Hemley, R.J., and Mao, H.K. (1996) Compression of ice to 210 gigapascals: Infrared evidence for a symmetric hydrogen-bonded phase. *Science*, 273, 218–220.
- Grimme, S. (2006) Semiempirical GGA-type density functional constructed with a long-range dispersion correction. *Journal of Computational Chemistry*, 27, 1787.
- Grützner, T., Klemme, Rohrbach, A., Grevasoni, F., and Berndt, J. (2018) The effect of fluorine on the stability of wadsleyite: Implication for the nature and depths of the transition zone in the Earth's mantle. *Earth and Planetary Sciences Letters*, 482, 236–244.
- Hirschmann, M. (2006) Water, melting, and the deep Earth H_2O cycle. *Annual Reviews of Earth and Planetary Sciences*, 34, 629–653.
- Hohenberg, P., and Kohn, W. (1964) Inhomogeneous electron gas. *Physical Review B*, 136, B864–871.
- Hushur, A., Manghani, M.H., Smyth, J.R., Williams, Q., Hellebrand, E., Lonappan, D., Ye, Y., Dera, P., and Frost, D.J. (2011) Hydrogen bond symmetrization and equation of state of phase D. *Journal of Geophysical Research*, 116, B06203.
- Ichikawa, H., Kawai, K., Yamamoto, S., and Kameyama, M. (2015) Effect of water on subduction of continental materials to the deep Earth. In A. Khan and F. Deschamps, Eds., *The Earth's Heterogeneous Mantle*, p. 275–299. Springer.
- Irfune, T., Higo, Y., Inoue, T., Kono, Y., Ohfuji, H., and Funakoshi, K. (2008) Sound velocities of majorite garnet and composition of the mantle transition region. *Nature*, 451, 814–817.
- Jahn, S., Wunder, B., Koch-Müller, M., Tarrieu, L., Pöhle, M., Watenphul, A., and Taran, M.N. (2012) Pressure-induced hydrogen bond symmetrisation in guyanite, $\beta\text{-CrOOH}$: evidence from spectroscopy and ab initio simulations. *European Journal of Mineralogy*, 24, 839–850.
- Jiang, F., Majzlan, J., Speziale, S., He, D., and Duffy, T.S. (2008) Single-crystal elasticity of diaspore, AlOOH , to 12 GPa by Brillouin scattering. *Physics of the Earth and Planetary Interior*, 170, 221–228.
- Jiang, F., Gwanmesia, G.D., Dyuzheva, T.I., and Duffy, T.S. (2009) Elasticity of stishovite and acoustic mode softening under high pressure by Brillouin scattering. *Physics of the Earth and Planetary Interior*, 172, 235–240.
- Karki, B.B., Stixrude, L., and Crain, J. (1997) Ab initio elasticity of three high-pressure polymorphs of silica. *Geophysical Research Letters*, 24, 3269–3272.
- Karki, B.B., Stixrude, L., and Wentzcovitch, R.M. (2001) High-pressure elastic properties of major materials of Earth's mantle from first principles. *Reviews of Geophysics*, 39, 507–534.
- Katahara, K.W. (1996) Clay mineral elastic properties. SEG Annual Meeting Expanded abstracts, 1691–1694. DOI: <http://dx.doi.org/10.1190/1.1826454>.
- Kawamoto, T. (2006) Hydrous phases and water transport in the subducting slab. *Reviews of Mineralogy and Geochemistry*, 62, 85–116.
- Kohn, W., and Sham, L.J. (1965) Self-consistent equations including exchange and correlation effects. *Physical Review*, 140, A1133–A1138.
- Kresse, G., and Furthmüller, J. (1996a) Efficiency of ab-initio total energy calculations for metals and semiconductors. *Computational Material Sciences*, 6, 15–50.
- (1996b) Efficient iterative schemes for ab initio total-energy calculations using plane-wave basis set. *Physical Review B*, 54, 11169–11186.
- Kresse, G., and Hafner, J. (1993) Ab initio Molecular Dynamics for liquid-metals. *Physical Review B*, 47, 558–561.
- Kresse, G., and Joubert, D. (1999) From ultrasoft pseudopotentials to the projector augmented-wave method. *Physical Review B*, 59, 1758–1775.
- Lee, C., Vanderbilt, D., Laasonen, K., Car, R., and Parrinello, M. (1993) Ab initio studies on the structural and dynamical properties of ice. *Physical Review B*, 47, 4863–4872.
- Li, B. (2003) Compressional and shear wave velocities of ringwoodite $\gamma\text{-Mg}_2\text{SiO}_4$ to 12 GPa. *American Mineralogist*, 88, 1312–1317.
- Mainprice, D. (1990) An efficient FORTRAN program to calculate seismic anisotropy from the lattice preferred orientation of minerals. *Computational Geoscience*, 16, 385–393.
- Mao, Z., Jacobsen, S.D., Jiang, F., Smyth, J.R., Holl, C.M., and Duffy, T.S. (2008) Elasticity of hydrous wadsleyite to 12 GPa: Implications for Earth's transition zone. *Geophysical Research Letters*, 35, L21305. doi:10.1029/2008GL035618.
- Mashino, I., Murakami, M., and Ohtani, E. (2016) Sound velocities of $\delta\text{-AlOOH}$ up to core-mantle boundary pressures with implications for the seismic anomalies in the deep mantle. *Journal of Geophysical Research: Solid Earth*, 121, 595–609. doi:10.1002/2015JB012477.
- Mei, S., and Kohlstedt, D.L. (2000) Influence of water on plastic deformation of olivine aggregates 1. Diffusion creep regime. *Journal of Geophysical Research*, 105, 21,457–21,469.
- Monkhorst, H.J., and Pack, J.D. (1976) Special points for Brillouin-zone integrations. *Physical Review B*, 13, 5188–5192.
- Mookherjee, M., Speziale, S., Marquardt, H., Jahn, S., Wunder, B., Koch-Müller, M., and Liermann, H.-P. (2015) Equation of state and elasticity of the 3.65 Å phase: implications for the X-discontinuity. *American Mineralogist*, 100, 2199–2208.
- Mookherjee, M., Tsuchiya, J., and Hariharan, A. (2016) Crystal structure, equation of state, and elasticity of hydrous aluminosilicate phase, topaz-OH ($\text{Al}_2\text{SiO}_4(\text{OH})_2$) at high pressures. *Physics of the Earth and Planetary Interior*, 251, 24–35.
- Nye, J.F. (1985) *Physical Properties of Crystals*. Oxford University Press, Clarendon.
- Ohno, I., Yamamoto, S., Anderson, O.L., and Noda, J. (1986) Determination of elastic constants of trigonal crystals by the rectangular parallelepiped resonance method. *Journal of Physics and Chemistry of Solids*, 47, 1103–1108.
- Ohno, I., Harada, K., and Yoshitomi, C. (2006) Temperature variation of elastic constants of quartz across α - β transition. *Physics and Chemistry of Minerals*, 33, 1–9.
- Ono, S. (1998) Stability limits of hydrous minerals in sediment and mid-ocean ridge basalt compositions: Implications for water transport in subduction zones. *Journal of Geophysical Research*, 103, 18,253–18,267.
- (1999) High temperature stability of phase egg, $\text{AlSiO}_3(\text{OH})$. *Contributions to Mineralogy and Petrology*, 137, 83–89.
- Otte, K., Pentcheva, R., Schmal, W.W., and Rustad, J.R. (2009) Pressure-induced structural and electronic transitions in FeOOH from first principles. *Physical Review B*, 80, 205116.
- Pamato, M.G., Myhill, R., Boffa Ballaran, T., Frost, D.J., Heidelbach, F., and Miyajima, N. (2015) Lower-mantle water reservoir implied by the extreme stability of hydrous aluminosilicate. *Nature Geosciences*, 8, 75–79.
- Panero, W.R. (2010) First principles determination of the structure and elasticity of hydrous ringwoodite. *Journal of Geophysical Research*, 115, B03203. doi:10.1029/2008JB006282.
- Panero, W.R., and Caracas, R. (2017) Stability of phase H in the $\text{MgSiO}_3\text{H}_2\text{O}$ – AlOOH – SiO_2 system. *Earth and Planetary Science Letters*, 463, 171–177.
- Panero, W.R., and Stixrude, L. (2004) Hydrogen incorporation in stishovite at high pressure and symmetric hydrogen bonding in $\delta\text{-AlOOH}$. *Earth and Planetary Science Letters*, 221, 421–431.
- Peacock, S.M. (1990) Fluid processes in subduction zones. *Science*, 248, 329–337.
- Peng, Y., Mookherjee, M., Hermann, A., Bajgain, S., Liu, S., and Wunder, B. (2017) Elasticity of phase-Pi ($\text{Al}_2\text{Si}_2\text{O}_7(\text{OH})_2$)—A hydrous aluminosilicate phase. *Physics of the Earth and Planetary Interior*, 269, 91–97. DOI: <https://doi.org/10.1016/j.pepi.2017.05.016>.
- Perdew, J.P., Burke, K., and Ernzerhof, M. (1996) Generalized gradient approximation made simple. *Physical Review Letters*, 77, 3865–3868.
- Sano, A., Ohtani, E., Kubo, T., and Funakoshi, K.-i. (2004) In situ X-ray observation of decomposition of hydrous aluminum silicate AlSiO_3OH and aluminum oxide hydroxide $\delta\text{-AlOOH}$ at high pressure and temperature. *Journal of Physics and Chemistry of Solids*, 65, 1547–1554.
- Sano-Furukawa, A., Kagi, H., Nagai, T., Nakano, S., Ushijima, D., Iizuka, R., Ohtani, E., and Yagi, T. (2009) Change in compressibility of $\delta\text{-AlOOH}$ and $\delta\text{-AlOOD}$ at high pressure: A study of isotope effect and hydrogen-bond symmetrization. *American Mineralogist*, 94, 1255–1261.
- Schmidt, M.W., Finger, L.W., Angel, R.J., and Robert, E. (1998) Synthesis, crystal structure and phase relations of AlSi_2OH , a high-pressure hydrous phase. *American Mineralogist*, 83, 881–888.
- Schreyer, W. (1995) Ultradeep metamorphic rocks: The retrospective viewpoint. *Journal of Geophysical Research*, 100, 8353–8366.
- Schweizer, K.S., and Stillinger, F.H. (1980) High pressure phase transitions and hydrogen-bond symmetry in ice polymorphs. *Journal of Chemical Physics*, 80, 1230–1240.
- Sikka, S.K. (2007) On some bond correlations at high pressures. *High Pressure Research*, 27, 313–319.
- Sinogeikin, S.V., and Bass, J.D. (2000) Single-crystal elasticity of pyrope and MgO to 20 GPa by Brillouin scattering in the diamond cell. *Physics of the Earth and Planetary Interior*, 120, 43–62.
- Smyth, J. (2006) Hydrogen in high-pressure silicate and oxide mineral structures. *Reviews of Mineralogy and Geochemistry*, 62, 85–116.
- Tennakoon, S., Peng, Y., Mookherjee, M., Speziale, S., Manthilake, G., Besara, T., Andreu, L., and Rivera, F. (2018) Single crystal elasticity of natural topaz at high-temperatures. *Scientific Reports*, 8, 1372. doi:10.1038/s41598-017-

- 17856-3.
- Tsuchiya, J., and Mookherjee, M. (2015) Crystal structure, equation of state, and elasticity of phase H (MgSiO_3H_2) at Earth's lower mantle pressures. *Scientific Reports*, 5, 15534. DOI: <https://doi.org/10.1038/srep15534>.
- Tsuchiya, J., and Tsuchiya, T. (2009) Elastic properties of $\delta\text{-AlOOH}$ under pressure: First principles investigation. *Physics of the Earth and Planetary Interior*, 174, 122–127.
- Tsuchiya, J., Tsuchiya, T., Tsuneyuki, S., and Yamanaka, T. (2002) First principles calculation of high-pressure hydrous phase, $\delta\text{-AlOOH}$. *Geophysical Research Letters*, 29.
- Tsuchiya, J., Tsuchiya, T., and Tsuneyuki, S. (2005) First-principles study of hydrogen bond symmetrization of phase D under high pressure. *American Mineralogist*, 90, 44–49.
- Tulk, C.A., Gagnon, R.E., Kieft, H., and Clouter, M.J. (1994) Elastic constants of ice III by Brillouin spectroscopy. *Journal of Chemical Physics*, 101, 2350.
- Vanpeteghem, C.B., Ohtani, E., Kondo, T., Takemura, K., and Kikegawa, T. (2003) Compressibility of phase egg $\text{AlSiO}_3(\text{OH})$: Equation of state and role of water at high pressure. *American Mineralogist*, 88, 1408–1411.
- Vaughan, M.T., and Weidner, D.J. (1978) The relationship of elasticity and crystal structure in andalusite and sillimanite. *Physics and Chemistry of Minerals*, 3, 133–144.
- Wang, D.J., Mookherjee, M., Xu, Y., and Karato, S. (2006a) The effect of water on the electrical conductivity of olivine. *Nature*, 443, 977–980.
- Wang, J., Sinogeikin, S.V., Inoue, T., and Bass, J.D. (2006b) Elastic properties of hydrous ringwoodite at high-pressure conditions. *Geophysical Research Letters*, 33, L14308, doi:10.1029/2006GL026441.
- Weidner, D.J., and Carleton, H.R. (1977) Elasticity of coesite. *Journal of Geophysical Research*, 82, 1334–1346.
- Winkler, B., Hytha, M., Warren, M.C., Milman, V., Gale, J.D., and Schreyer, J. (2001) Calculation of the elastic constants of the Al_2SiO_5 polymorphs andalusite, sillimanite and kyanite. *Zeitschrift für Kristallographie*, 216, 67–70.
- Wirth, R., Vollmer, C., Brenker, F., Matsyuk, S., and Kaminsky, F. (2007) Inclusion of nanocrystalline hydrous aluminum silicate “Phase egg” in superdeep diamonds from Junia (Mato Grosso State, Brazil). *Earth and Planetary Science Letters*, 259, 384–399.
- Wunder, B., Medenbach, O., Krause, W., and Schreyer, W. (1993a) Synthesis, properties and stability of $\text{Al}_2\text{Si}_2\text{O}_7(\text{OH})_3$ (phase Pt), a hydrous high-pressure phase in the system $\text{Al}_2\text{O}_3\text{-SiO}_2\text{-H}_2\text{O}$ (ASH). *European Journal of Mineralogy*, 5, 637–649.
- Wunder, B., Rubie, D.C., Ross, C.R. II, Medenbach, O., Seifert, F., and Schreyer, W. (1993b) Synthesis, stability, and properties of $\text{Al}_2\text{SiO}_4(\text{OH})_2$: A fully hydrated analogue of topaz. *American Mineralogist*, 78, 285–297.
- Xue, X., Kanzaki, M., Fukui, H., Ito, E., and Hashimoto, T. (2006) Cation order and hydrogen bonding of high-pressure phases in the $\text{Al}_2\text{O}_3\text{-SiO}_2\text{-H}_2\text{O}$ system: An NMR and Raman study. *American Mineralogist*, 91, 850–861.
- Yang, R., and Wu, Z. (2014) Elastic properties of stishovite and the CaCl_2 -type silica at the mantle temperature and pressure: An ab initio investigation. *Earth and Planetary Science Letters*, 404, 14–21.
- Zhong, X., Hermann, A., Wang, Y., and Ma, Y. (2016) Monoclinic high-pressure polymorph of AlOOH predicted from first principles. *Physical Review B*, 94, 224110.

MANUSCRIPT RECEIVED JUNE 19, 2018

MANUSCRIPT ACCEPTED SEPTEMBER 17, 2018

MANUSCRIPT HANDLED BY ZHICHENG JING

Endnote:

¹Deposit item AM-19-16694, Supplemental Figures and Material. Deposit items are free to all readers and found on the MSA website, via the specific issue's Table of Contents (go to http://www.minsocam.org/MSA/AmMin/TOC/2019/Jan2019_data/Jan2019_data.html).