

Allyl-Nickel Catalysis Enables Carbonyl Dehydrogenation and Oxidative Cycloalkenylation of Ketones

David Huang, Suzanne M. Szewczyk, Pengpeng Zhang, and Timothy R. Newhouse*

Department of Chemistry, Yale University, 225 Prospect Street, New Haven, Connecticut 06520-8107, United States

S Supporting Information

ABSTRACT: We herein disclose the first report of a first-row transition metal-catalyzed α,β -dehydrogenation of carbonyl compounds using allyl-nickel catalysis. This development overcomes several limitations of previously reported allyl-palladium-catalyzed oxidation, and is further leveraged for the development of an oxidative cycloalkenylation reaction that provides access to bicycloalkenones with fused, bridged, and spirocyclic ring systems using unactivated ketone and alkene precursors.


Methodological advancements in the area of transition metal catalysis have enabled challenging C–C bond formations using unstabilized enolates. Among these advancements are α -alkenylation reactions¹ that employ a cross-coupling strategy (Figure 1A) to unite a vinyl halide and a metal enolate to provide synthetically versatile β,γ -enone products. As an alternative to cross coupling, seminal work by Saegusa² and Kende³ demonstrated that unactivated alkenes can be employed for the cycloalkenylation⁴ of ketones using palladium, but required the preactivation of the carbonyls as enoxysilanes (Figure 1B). Additionally, stabilized enolates⁵ such as 1,3-diketones can also be employed in cycloalkenylation reactions, including using palladium catalysis as reported by Widenhoefer.⁶

As existing strategies for α -alkenylation require either activated ketone or alkene partners,⁵ a direct alkenylation between unactivated ketones and alkenes by an oxidative approach would advantageously allow for the employment of more abundantly available precursors, and contribute to increased step-efficiency for cycloalkenylation reactions, which are broadly used in natural product synthesis for the construction of carbocycles.^{4b}


Given our previous success using allyl oxidants to promote β -hydride elimination in the context of palladium-catalyzed carbonyl dehydrogenation (10),^{7,8} we hypothesized that allyl-transition metal catalysis could also facilitate our desired dehydrogenative cycloalkenylation. In particular, we were interested in probing the possibility of using allyl-nickel catalysis to facilitate the desired cycloalkenylation reaction due to nickel's increased propensity to undergo migratory insertion.⁹

The envisioned oxidative cycloalkenylation reaction involves generation of allyl-nickel enolate 4 by deprotonation of 1 followed by transmetalation with allyl-nickel species 3. A subsequent migratory insertion event with a pendant alkene would generate alkyl-nickel species 5 that then undergoes β -


A. α -Alkenylation of Ketones via Cross-Coupling



B. α -Alkenylation via Enoxysilanes



C. Oxidative Cycloalkenylation Using Allyl-Nickel Catalysis



Mechanistic Strategy:




Figure 1. Strategies for α -alkenylation of ketones.

hydride elimination to give the desired cycloalkenylation product (2) and allyl-nickel-hydride (6). Catalyst turnover by reductive elimination of 6 gives propene as the stoichiometric byproduct and Ni(0), which completes the catalytic cycle via oxidative addition to allyl electrophile 9. Additional mechanistic support comes from Tsuji's observation that allyl-palladium enolates generated by the decarboxylation of allyl β -ketoesters could undergo migratory insertion and β -hydride

Received: March 7, 2019

Published: March 12, 2019

elimination with electron-deficient alkenes to give cycloalkenylation products.¹⁰

Despite the often more challenging β -hydride elimination of first-row transition metals,⁹ successful development of this methodology would provide a more sustainable and cost-effective alternative for allyl-metal-catalyzed β -hydride elimination in both cycloalkenylation and carbonyl α,β -dehydrogenation contexts. We also aimed to leverage the inherent differences of first- and second-row transition metals to address some of the current limitations of allyl-palladium-catalyzed dehydrogenation chemistry. In particular, for reluctant substrates that give predominantly Tsuji–Trost allylation byproducts, we expected the use of a less electrophilic allyl-nickel species to be more successful.

Unactivated carbonyl compounds can undergo α,β -dehydrogenation by several methods including using metal catalysis.¹¹ Stahl has reported an exciting process for ketone and aldehyde dehydrogenation using palladium catalysis that is beginning to see widespread adoption.¹² Dong has also reported a palladium-catalyzed approach for lactams,¹³ and more recently a method employing platinum to dehydrogenate several substrate classes.¹⁴ Methods utilizing first-row metals for dehydrogenation have either focused on activated systems that lead to products with extended conjugation, or these intermediates were not isolated but rather reacted further to form more stable products.¹⁵ This report describes the first practical first-row transition metal-catalyzed α,β -dehydrogenation of carbonyl compounds using nickel-catalysis. The discovery of nickel-catalyzed dehydrogenation was a starting point for the development of an oxidative cycloalkenylation of ketone enolates for the construction of diverse bicyclic scaffolds.

Ketone **11a** was used as our model substrate (Table 1) as we previously identified that acyclic ketones were poor substrates

Table 1. Optimization of α,β -Dehydrogenation


Entry	Base	Catalyst	Yield (%) ^a
1	Zn(TMP) ₂	[Pd(allyl)Cl] ₂	8
2	Zn(TMP) ₂	MnBr ₂	0
3	Zn(TMP) ₂	FeBr ₂	0
4	Zn(TMP) ₂	CoBr ₂	0
5	Zn(TMP) ₂	CuBr ₂	0
6	Zn(TMP) ₂	NiBr ₂ (dme)	7
7	Zn(TMP) ₂ , ZnBr ₂	NiBr ₂ (dme)	12
8	LiTMP, ZnBr ₂	NiBr ₂ (dme)	61
9	LiCyan, ZnBr ₂	NiBr ₂ (dme)	69

^a¹H-NMR yield was determined using 1,3,5-trimethoxybenzene as an internal standard.

for our reported allyl-palladium-catalyzed ketone dehydrogenation protocol (entry 1).⁸ Several first-row metal catalysts were investigated (entries 2–6), and encouragingly we found that a similar, albeit low yield of enone **10a** could be detected using NiBr₂(dme) (entry 6), demonstrating the feasibility of a nickel-mediated reaction. Investigation of the base (entries 7–9) revealed that our previously disclosed hindered lithium anilide base, LiCyan, proved to be most effective.^{7b}

We investigated the scope of the nickel-catalyzed dehydrogenation with particular interest in other substrates that were previously unsuccessful with the palladium-catalyzed system (Table 2). A variety of acyclic ketones, including aliphatic

Table 2. Scope of Carbonyl and Nitrile α,β -Dehydrogenation



^a1.2 equiv Zn(TMP)₂ and no ZnBr₂ was used. ^b1,4-Dioxane was used as solvent. ^cReaction conducted at 23 °C. ^d1.0 equiv diethyl allyl phosphate was used.

(**10a–c**) and electron-rich aryl ketones (**10d–e**) were efficiently dehydrogenated under the optimized conditions, and tolerated acid sensitive acetal and cyclopropane functionality. Primary nitriles (**10f**) and more hindered secondary nitriles (**10g–j**) were also excellent substrates for this nickel-catalyzed dehydrogenation. Notably, α -aryl nitriles (**10i–j**), which predominantly gave allylation byproducts when palladium was used, were efficiently dehydrogenated using the less electrophilic allyl-nickel system. Similarly, α -aryl ester

10k was also accessed in good yield. Unsaturated ester **10l** containing *N*-Piv functionality was also obtained in excellent yield. Amides (**10m–o**) were a viable substrate class for this nickel-catalyzed dehydrogenation, including sterically hindered β,β -disubstituted amide **10o**.

When we tried our optimized conditions on cyclic ketones, we saw significantly depreciated yields. Instead we found that commercially available $\text{Zn}(\text{TMP})_2$ was most effective for cyclic ketones, as previously demonstrated.⁸ β -ketoester **10p** could be dehydrogenated in good yield, despite the possibility of metal chelation by the 1,3-dicarbonyl suppressing reactivity. Ketone **10q** containing a pendant alkene and mutilin derivative **10r** could also be accessed. Interestingly, we found that $\text{Zn}(\text{TMP})_2$ could also be used to efficiently dehydrogenate 5-, 6-, and 7-membered lactones (**10s–u**) and lactams (**10v–x**) despite their decreased acidity. Notably, unsaturated lactam **10v** is a key intermediate needed to access an important class of IRAK4 inhibitors.¹⁶ While LiCyan is still an effective base for lactones and lactams, it was generally observed that $\text{Zn}(\text{TMP})_2$ outperformed LiCyan for cyclic carbonyl compounds (see [Supporting Information](#)). The use of $\text{Zn}(\text{TMP})_2$ also provides the advantage of a more operationally simple protocol. Under our reaction conditions, structural features of the molecule (acyclic or cyclic) seem to dictate the optimal base necessary for efficient dehydrogenation.


Furthermore, the development of a nickel-catalyzed system for carbonyl dehydrogenation overcomes previous limitations in scope, such as acyclic ketones (**10a–e**) and allylation-prone α -aryl substrates (**10i–k**) and provides an efficient and cost-effective alternative for carbonyl dehydrogenation, which has only been demonstrated previously with precious metals.

Using our newly established allyl-nickel-based system, we then explored the possibility of an oxidative cycloalkenylation of ketones. It was found that with only a minor modification to the dehydrogenation conditions using $\text{Zn}(\text{TMP})_2$ as base, the desired cycloalkenylation could be achieved. The addition of ZnBr_2 as an additive was crucial for attaining an efficient reaction and a broad substrate scope (see [Supporting Information](#) for a broad investigation of additives), and highlights the remarkable role of salt additives in transformations involving organozinc species.¹⁷

The substrate scope was explored with particular interest in the various carbocyclic architectures that could be accessed ([Table 3](#)). Acyclic ketones could efficiently cyclize with a pendant homoallyl group, providing β -methyl cyclohexenone products after alkene isomerization to the more thermodynamically stable α,β -enone products (**2a–d**). Importantly, ethyl ketones could also be used despite the possibility of dehydrogenation, providing access to tetrasubstituted enone **2b**. Enone product **2c** highlights that this protocol is selective for cyclization of 6-membered rings rather than 5-membered rings, which is in contrast to previous reports of palladium-mediated cycloalkenylations.⁴ Gratifyingly, sensitive furan functionality was also well tolerated (**2d**). Geminally substituted cyclic structures could be used to access spirocyclic motifs in good yields (**2e–h**) and tolerated the acid sensitive *N*-Boc (**2f**) and acetal functionality (**2g–h**).

Bicyclo[4.3.0]nonanes and bicyclo[4.4.0]decanes could also be accessed using this methodology as demonstrated by products **2i–q**. Access to piperitone derivative **2i** highlights that 5-membered ring formation is also possible, and **2i** was prepared on 1 g scale without any depreciation in yield. β,γ -enone **2j** can be used to access a variety of eudesmane-type

Table 3. Scope of Oxidative Cycloalkenylation



^a2.0 equiv $\text{Zn}(\text{TMP})_2$ was used. ^b20 mol % $\text{NiBr}_2(\text{dme})$ and no ZnBr_2 was used. ^c40 °C. ^d1.5 equiv $\text{Zn}(\text{TMP})_2$ was used.

sesquiterpene natural products, but was previously reported to be a challenging scaffold to access by existing cycloalkenylation approaches.¹⁸ Gratifyingly, pendant alkenes bearing electron withdrawing groups were also tolerated in the substrate scope and could provide α,β -unsaturated ester product **2k** as a 2:1 mixture of separable *E*- and *Z*-isomers. Cycloalkenylation with olefins bearing electron-donating substituents such as alkyl or ether groups were not tolerated. Other bicyclo[4.3.0]nonanes (**2l–n**), such as verbenone derivative **2l** bearing an acid-sensitive cyclobutane and Boc-protected indole **2n** were also excellent substrates under the reaction conditions. Cycloalkenylation initiated by cyclopentanones could also be achieved (**2o–q**) and demonstrated the tolerance of

benzonitrile (**2p**) and TBS-protected alcohol functionality (**2q**). Indanone **2o** was accessed using modified reaction conditions that employed 20 mol % catalyst loading and no ZnBr_2 additive, which was found to be an excellent alternative for reluctant substrates. Notably, while 6-membered ring formation under previously reported cycloalkenylation methods with palladium is challenging,^{2,4} this allyl-nickel system provides an efficient disconnection for forming cyclohexanes with unactivated alkenes.

Although cycloalkenylation to form fused ring systems was not possible when β -hydride elimination was a competing pathway, bridged bicyclic ring systems such as bicyclo[3.3.1]nonanes (**2r–t**) and bicyclo[3.2.1]octanes (**2u**), which are important core structures in a variety of natural products,¹⁹ could be accessed despite the possibility of β -hydride elimination. Access to bicyclo[3.3.1]nonane **2r** demonstrates the possibility of bifurcating reactivity using reagent control (Table 2, **10q**), although this was not found to be a general phenomenon. Wieland–Miescher ketone derivative **2t** containing two ketone functional groups selectively underwent cycloalkenylation via the thermodynamic enolate, and **2v** demonstrates selective cyclization with the axially disposed *O*-tethered alkene to give a bicyclo[3.3.1]nonane rather than with the all-carbon tethered alkene to give a bicyclo[3.2.1]octane.

Under the reported reaction conditions, all substrates were completely selective for the *exo*-mode of cyclization and gave *cis*-fused products. No alkene isomerization was observed for substrates that gave fused or bridged bicyclic products, which is an advantage over previously reported palladium-mediated cycloalkenylations of enoxysilanes, which gave isomeric mixtures of alkenes.⁴

We wanted to probe the identity of the metal responsible for C–C bond formation between the ketone enolate and the pendant olefin (Figure 2). Two mechanistic possibilities could be envisioned: one involves the carbometalation of the

pendant alkene²⁰ by a zinc enolate and the other involves the proposed migratory insertion of an alkene to an allyl-nickel enolate. Under the standard conditions, **1i** underwent efficient alkenylation (Figure 2a). When substrate **1i** was heated with $\text{Zn}(\text{TMP})_2$ and ZnBr_2 , we found that the zinc-mediated cyclization was indeed possible, albeit in low conversion. Subsequent quenching with D_2O afforded the fully saturated, deuterated species **12** in 32% yield with 68% deuterium incorporation (Figure 2b) and suggests the intermediacy of an alkylzinc intermediate. Related cyclizations of activated zinc enolates with unactivated olefins have been reported.²¹

Despite the possibility of a zinc-mediated cyclization, the subsequent addition of nickel and allyl oxidant to effect transmetalation and β -hydride elimination of the alkylzinc species did not provide a significant amount of alkenylation product **2i** (Figure 2c). Similarly, treatment of an alkylzinc intermediate prepared from the corresponding alkyl bromide to identical allyl-nickel conditions also provided low yields of alkenylation product (see Supporting Information). These results suggest that while a zinc-mediated anionic cyclization followed by an allyl-nickel-mediated β -hydride elimination is one possible mechanism for cycloalkenylation, the migratory insertion of a nickel enolate cannot be ruled out given additional experiments conducted in the absence of zinc (Figure 2d).

The nickel-mediated cyclization was demonstrated to be plausible by treatment of α -bromo ketone **13** with a stoichiometric quantity of $\text{Ni}(\text{cod})_2$. The low yield for this transformation may reflect the importance of the specific ligand sphere accessed through our catalytic manifold and emphasizes the synthetic utility of our methodology. Taken together, these mechanistic investigations highlight the possibility of both zinc and nickel to mediate C–C bond formation between unstabilized enolates and unactivated alkenes. Additional mechanistic investigations of the roles of zinc and nickel are ongoing and will be reported in due course.

We have effectively expanded allyl-metal catalysis for carbonyl α,β -dehydrogenation from palladium to nickel, thereby allowing for the use of a more sustainable and inexpensive first-row transition metal catalyst. Furthermore, key limitations to the substrate specificity and scope of the allyl-palladium-catalyzed α,β -dehydrogenation conditions were overcome using the more general nickel-catalyzed reaction conditions for the dehydrogenation of ketones, nitriles, esters, and amides. The use of allyl-nickel catalysis also enabled an oxidative cycloalkenylation reaction between unstabilized ketone enolates and unactivated alkenes, providing access to diverse bicyclic architectures and a new retrosynthetic C–C bond disconnection for the synthesis of complex polycyclic molecules.

■ ASSOCIATED CONTENT

● Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.9b02552.

Experimental procedures and spectroscopic data for all new compounds including ^1H - and ^{13}C NMR spectra (PDF)

X-ray crystallography of compound **SI-13** (CIF)




Figure 2. Mechanistic investigations.

AUTHOR INFORMATION

Corresponding Author

*timothy.newhouse@yale.edu

ORCID

Timothy R. Newhouse: 0000-0001-8741-7236

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We are grateful for financial support from Yale University, the Sloan Foundation, Nalas Engineering, Amgen, the NSF (CAREER, 1653793), and the NIH-funded Chemistry/Biology Interface Training Program (D.H., T32GM067543). Dr. Brandon Mercado is gratefully acknowledged for X-ray crystallography of compound SI-13.

REFERENCES

- (1) (a) For a seminal report of nickel-catalyzed alkenylation of enolates, see: Millard, A. A.; Rathke, M. W. A Nickel Catalyst for the Arylation and Vinylation of Lithium Ester Enolates. *J. Am. Chem. Soc.* **1977**, *99*, 4833–4835. (b) For a review on palladium- and nickel-catalyzed α -alkenylation via cross coupling, see: Ankner, T.; Cosner, C. C.; Helquist, P. Palladium- and Nickel-Catalyzed Alkenylation of Enolates. *Chem. - Eur. J.* **2013**, *19*, 1858–1871. (c) For a recent report of copper-catalyzed alkenylation of zinc enolates employing iodonium salts, see: Liu, C.; Wang, Q. Alkenylation of C(sp³)-H Bonds by Zincation/Copper-Catalyzed Cross Coupling with Iodonium Salts. *Angew. Chem., Int. Ed.* **2018**, *57*, 4727–4731.
- (2) Ito, Y.; Aoyama, H.; Hirao, T.; Mochizuki, A.; Saegusa, T. Cyclization Reactions via Oxo- π -allylpalladium(II) Intermediates. *J. Am. Chem. Soc.* **1979**, *101*, 494–496.
- (3) (a) Kende, A. S.; Roth, B.; Sanfilippo, S. J. Facile, Palladium(II)-Mediated Synthesis of Bridged and Spirocyclic Bicycloalkenones. *J. Am. Chem. Soc.* **1982**, *104*, 1784–1785. (b) Kende, A. S.; Roth, B.; Sanfilippo, P. J.; Blacklock, T. J. Mechanism and Regioisomeric Control in Palladium(II)-Mediated Cycloalkenylations. A Novel Total Synthesis of ($\pm$)-Quadron. *J. Am. Chem. Soc.* **1982**, *104*, 5808–5810.
- (4) (a) For a review on palladium-mediated cycloalkenylations, see: LeBras, J.; Muzart, J. Base-free palladium-mediated cycloalkenylations of olefinic enolic systems. *Tetrahedron* **2015**, *71*, 9035–9059. (b) Toyota, M.; Ihara, M. Development of Palladium-Catalyzed Cycloalkenylation and its Application to Natural Product Synthesis. *Synlett* **2002**, 1211–1222.
- (5) Dénès, F.; Pérez-Luna, A.; Chemla, F. Addition of Metal Enolate Derivatives to Unactivated Carbon-Carbon Multiple Bonds. *Chem. Rev.* **2010**, *110*, 2366–2447.
- (6) Pei, T.; Wang, X.; Widenhoefer, R. A. Palladium-Catalyzed Intramolecular Oxidative Alkylation of Unactivated Olefins. *J. Am. Chem. Soc.* **2003**, *125*, 648–649.
- (7) (a) Chen, Y.; Romaine, J. P.; Newhouse, T. R. Palladium-Catalyzed α,β -Dehydrogenation of Esters and Nitriles. *J. Am. Chem. Soc.* **2015**, *137*, 5875–5878. (b) Chen, Y.; Turlik, A.; Newhouse, T. R. Amide α,β -Dehydrogenation Using Allyl-Palladium Catalysis and a Hindered Monodentate Anilide. *J. Am. Chem. Soc.* **2016**, *138*, 1166–1169. (c) Zhao, Y.; Chen, Y.; Newhouse, T. R. Allyl-Palladium Catalyzed α,β -Dehydrogenation of Carboxylic Acids via Enediolates. *Angew. Chem., Int. Ed.* **2017**, *56*, 13122–13125. (d) Schuppe, A. W.; Huang, D.; Chen, Y.; Newhouse, T. R. Total Synthesis of (–)-Xylogranatopyridine B via a Palladium-Catalyzed Oxidative Stannylation of Enones. *J. Am. Chem. Soc.* **2018**, *140*, 2062–2066. (e) Szewczyk, S. M.; Zhao, Y.; Sakai, H. A.; Dube, P.; Newhouse, T. R. α,β -Dehydrogenation of esters with free O-H and N-H functionalities with allyl-palladium catalysis. *Tetrahedron* **2018**, *74*, 3293–3300.
- (8) For allyl-palladium-catalyzed dehydrogenation of ketones, see: (a) Chen, Y.; Huang, D.; Zhao, Y.; Newhouse, T. R. Allyl-Palladium-Catalyzed Ketone Dehydrogenation Enables Telescoping with Enone α,β -Vicinal Difunctionalization. *Angew. Chem., Int. Ed.* **2017**, *56*, 8258–8262. (b) Huang, D.; Zhao, Y.; Newhouse, T. R. Synthesis of Cyclic Enones by Allyl-Palladium-Catalyzed α,β -Dehydrogenation. *Org. Lett.* **2018**, *20*, 684–687.
- (9) (a) Tasker, S. Z.; Standley, E. A.; Jamison, T. F. Recent advances in homogeneous nickel catalysis. *Nature* **2014**, *509*, 299–309. (b) Zweig, J. E.; Kim, D. E.; Newhouse, T. R. Methods Utilizing First-Row Transition Metals in Natural Product Total Synthesis. *Chem. Rev.* **2017**, *117*, 11680–11752.
- (10) Nokami, J.; Watanabe, H.; Mandai, T.; Kawada, M.; Tsuji, J. The palladium-catalyzed Michael addition reaction of the ketone enolates generated by the decarboxylation of allyl- β -keto carboxylates under neutral conditions. *Tetrahedron Lett.* **1989**, *30*, 4829–4832.
- (11) For reviews on carbonyl dehydrogenation, see: (a) Muzart, J. One-Pot Syntheses of α,β -Unsaturated Carbonyl Compounds through Palladium-Mediated Dehydrogenation of Ketones, Aldehydes, Esters, Lactones, and Amides. *Eur. J. Org. Chem.* **2010**, *2010*, 3779–3790. (b) Stahl, S. S.; Diao, T. Oxidation Adjacent to C = X bonds by Dehydrogenation. *Comp. Org. Synth.* **2014**, *7*, 178–212. (c) Turlik, A.; Chen, Y.; Newhouse, T. R. Dehydrogenation Adjacent to Carbonyls Using Palladium-Allyl Intermediates. *Synlett* **2016**, *27*, 331–336. (d) Iosub, A. V.; Stahl, S. S. Palladium-Catalyzed Aerobic Dehydrogenation of Cyclic Hydrocarbons for the Synthesis of Substituted Aromatics and Other Unsaturated Products. *ACS Catal.* **2016**, *6*, 8201–8213. (e) Hirao, T. Synthetic Strategy: Palladium-Catalyzed Dehydrogenation of Carbonyl Compounds. *J. Org. Chem.* **2019**, *84*, 1687–1692.
- (12) (a) Diao, T.; Stahl, S. S. Synthesis of Cyclic Enones via Direct Palladium-Catalyzed Aerobic Dehydrogenation of Ketones. *J. Am. Chem. Soc.* **2011**, *133*, 14566–14569. (b) Diao, T.; Wadzinski, T. J.; Stahl, S. S. Direct aerobic α,β -dehydrogenation of aldehydes and ketones with a Pd(TFA)₂/4,5-diazafluorenone catalyst. *Chem. Sci.* **2012**, *3*, 887–891. (c) Diao, T.; Pun, D.; Stahl, S. S. Aerobic Dehydrogenation of Cyclohexanone to Cyclohexenone Catalyzed by Pd(DMSO)₂(TFA)₂: Evidence for Ligand-Controlled Chemoselectivity. *J. Am. Chem. Soc.* **2013**, *135*, 8205–8212.
- (13) Chen, M.; Dong, G. Direct Catalytic Desaturation of Lactams Enabled by Soft Enolization. *J. Am. Chem. Soc.* **2017**, *139*, 7757–7760.
- (14) Chen, M.; Rago, A.; Dong, G. Platinum-catalyzed Desaturation of Lactams, Ketones and Lactones. *Angew. Chem., Int. Ed.* **2018**, *57*, 16205–16209.
- (15) (a) Ueno, S.; Shimizu, R.; Kuwano, R. Nickel-Catalyzed Formation of a Carbon-Nitrogen Bond at the β -Position of Saturated Ketones. *Angew. Chem., Int. Ed.* **2009**, *48*, 4543–4545. (b) Tutkowski, B. M.; Grigalunas, M.; Wiest, O.; Helquist, P. A nickel-catalyzed α,β -coupling of saturated ketones. *Tetrahedron Lett.* **2015**, *56*, 3468–3472. (c) Jie, X.; Shang, Y.; Zhang, X.; Su, W. Cu-Catalyzed Sequential Dehydrogenation–Conjugate Addition for β -Functionalization of Saturated Ketones: Scope and Mechanism. *J. Am. Chem. Soc.* **2016**, *138*, 5623–5633. (d) Shang, Y.; Jie, X.; Jonnada, K.; Zafar, S. N.; Su, W. Dehydrogenative desaturation-relay via formation of multicenter-stabilized radical intermediates. *Nat. Commun.* **2017**, *8*, 2273. (e) Li, H.; Jiang, Q.; Jie, X.; Shang, Y.; Zhang, Y.; Goossen, L. J.; Su, W. Rh/Cu-Catalyzed Ketone β -Functionalization by Merging Ketone Dehydrogenation and Carboxyl-Directed C-H Alkylation. *ACS Catal.* **2018**, *8*, 4777–4782.
- (16) Lee, K. L.; Ambler, C. M.; Anderson, D. R.; Boscoe, B. P.; Bree, A. G.; Brodfuehrer, J. L.; Chang, J. S.; Choi, C.; Chung, S.; Curran, K. J.; Day, J. E.; Dehnhardt, C. M.; Dower, K.; Drozd, S. E.; Frisbie, R. K.; Gavrin, L. K.; Goldberg, J. A.; Han, S.; Hegen, M.; Hepworth, D.; Hope, H. R.; Kamtekar, S.; Kilty, I. C.; Lee, A.; Lin, L.-L.; Lovering, F. E.; Lowe, M. D.; Mathias, J. P.; Morgan, H. M.; Murphy, E. A.; Papaioannou, N.; Patny, J. P.; Pierce, B. S.; Rao, V. R.; Saiah, E.; Samardjiev, I. J.; Samas, B. M.; Shen, M. W. H.; Shin, J. H.; Soutter, H. H.; Strohsbach, J. W.; Symanowicz, P. T.; Thomason, J. R.; Trzupek, J. D.; Vargas, R.; Vincent, F.; Yan, J.; Zapf, C. W.; Wright, S. W. Discovery of Clinical Candidate 1-([2S,3S,4S])-3-Ethyl-4-fluoro-5-

oxopyrrolidin-2-yl]methoxy}-7-methoxyisoquinoline-6-carboxamide (PF-06650833), a Potent, Selective Inhibitor of Interleukin-1 Receptor Associated Kinase 4 (IRAK4), by Fragment-Based Drug Design. *J. Med. Chem.* **2017**, *60*, 5521–5542.

(17) (a) Achonduh, G. T.; Hadei, N.; Valente, C.; Avola, S.; O'Brien, C. J.; Organ, M. G. On the role of additives in alkyl–alkyl Negishi cross-couplings. *Chem. Commun.* **2010**, *46*, 4109–4111.

(b) McCann, L. M.; Organ, M. C. On the Remarkably Different Role of Salt in the Cross-Coupling of Arylzincs From That Seen with Alkylzincs. *Angew. Chem., Int. Ed.* **2014**, *53*, 4386–4389.

(18) (a) Chen, K.; Ishihara, Y.; Galán, M. M.; Baran, P. S. Total synthesis of eudesmane terpenes: cyclase phase. *Tetrahedron* **2010**, *66*, 4738–4744. (b) Thomas, A. F.; Ozainne, M.; Decorzant, R.; Naf, F.; Lukacs, G. 10-Epijuneol, a new *cis*-eudesmane sesquiterpenoid. *Tetrahedron* **1976**, *32*, 2261–2264. (c) Toyota, M.; Asakawa, Y. An eudesmane-type sesquiterpene alcohol from the liverwort *Frullania tamarisci*. *Phytochemistry* **1990**, *29*, 3664–3665.

(19) (a) Presset, M.; Coquerel, Y.; Rodriguez, J. Syntheses and Applications of Functionalized Bicyclo[3.2.1]octanes: Thirteen Years of Progress. *Chem. Rev.* **2013**, *113*, 525–595. (b) Ruiz, M.; López-Alvarado, P.; Giorgi, G.; Menéndez, J. C. Domino reactions for the synthesis of bridged bicyclic frameworks: fast access to bicyclo-[*n*.3.1]alkanes. *Chem. Soc. Rev.* **2011**, *40*, 3445–3454.

(20) (a) Stadtmüller, H.; Lentz, R.; Tucker, C. E.; Stüdemann, T.; Dörner, W.; Knochel, P. Palladium-Catalyzed Iodine-Zinc Exchange Reactions. A New Palladium-Mediated Intramolecular Carbozincation of Alkenes. *J. Am. Chem. Soc.* **1993**, *115*, 7027–7028. (b) Meyer, C.; Marek, I.; Courtemanche, G.; Normant, J.-F. Carbocyclization of Functionalized Zinc Organometallics. *Synlett* **1993**, *1993*, 266–268.

(21) Pérez-Luna, A.; Botuha, C.; Ferreira, F.; Chemla, F. Carbometalation of unactivated alkenes by zinc enolate derivatives. *New J. Chem.* **2008**, *32*, 594–606.