

SYMMETRIC AND ALTERNATING POWERS OF WEIL REPRESENTATIONS OF FINITE SYMPLECTIC GROUPS

ROBERT M. GURALNICK, KAY MAGAARD, AND PHAM HUU TIEP

ABSTRACT. Motivated by an earlier result of N. Katz, we establish all possible equalities between symmetric squares, alternating squares, and tensor products of complex irreducible Weil characters of finite symplectic groups in odd characteristic. We also construct an infinite series of examples of irreducible symmetric cubes and alternating cubes of complex representations of finite groups.

1. INTRODUCTION

The so-called *Weil representations* were introduced by A. Weil [W] for classical groups over local fields. Weil mentioned that the finite field case may be considered analogously. This was developed in detail by R. E. Howe [Hw] and P. Gérardin [Ger], for characteristic zero representations. The same complex representations were introduced independently by I. M. Isaacs [Is] and H. N. Ward [Wa] for finite symplectic groups $\mathrm{Sp}_{2n}(q)$ with q odd, and by G. M. Seitz [Se] for finite unitary groups. (These representations for $\mathrm{Sp}_{2n}(p)$ were also constructed in [BRW].) Weil representations attract much attention because of their many interesting features, cf. for instance [D], [DT], [Gow], [Gr], [ST1], [ST2], [T1], [T2], [Z].

The construction of Weil representations may be found in [Hw], [Ger], [Se], etc. Let us recall this construction in the case of $G = \mathrm{Sp}_{2n}(q)$, where q is a power of an odd prime $p > 2$ and $n \geq 1$. Let $V = \mathbb{F}_q^{2n}$ be endowed with a non-degenerate symplectic form $(\cdot, \cdot) : V \times V \rightarrow \mathbb{F}_q$. Then one checks that $Q := \{(\alpha, v) \mid \alpha \in \mathbb{F}_q, v \in V\}$, considered with the operation

$$(\alpha, v) \cdot (\beta, w) = (\alpha + \beta + (v, w), v + w)$$

is a p -group (of special type), with

$$\mathbf{Z}(Q) = [Q, Q] = \Phi(Q) = \{(\alpha, 0) \mid \alpha \in \mathbb{F}_q\}$$

2010 *Mathematics Subject Classification.* 20C33, 20C20.

Key words and phrases. Finite symplectic groups, Weil representations.

The authors are grateful to Nick Katz for bringing the problem to their attention. They are also grateful to the referee for careful reading and helpful comments on the paper.

The first and the third authors gratefully acknowledge the support of the NSF (grants DMS-1600056, DMS-1201374, and DMS-1665014).

being elementary abelian of order q . Then $G = \mathrm{Sp}(V)$ acts on Q as outer automorphisms that act trivially on $\mathbf{Z}(Q)$.

Fix a nontrivial linear character ψ of the additive group $(\mathbb{F}_q, +)$. Then ψ can be naturally viewed as a character of $\mathbf{Z}(Q)$. One can show, cf. [Gr, §13], that there is a unique (up to equivalence) complex irreducible representation Φ of Q of degree q^n that affords the character $q^n\psi$ of $\mathbf{Z}(Q)$. For any $g \in G$, the action of g on Q fixes ψ and so sends Φ to an equivalent representation $\Lambda(g)\Phi\Lambda(g)^{-1}$ for some $\Lambda(g) \in \mathrm{GL}_{q^n}(\mathbb{C})$. The map $g \mapsto \Lambda(g)$ is in fact a projective representation of G and can be lifted to a linear representation Ω , see [Gr, §13], which is called a *reducible Weil representation* of G , with character $\omega = \omega_n$. The latter character can be decomposed into the sum $\eta + \xi$ of two (irreducible) *Weil characters*, which correspond to the two eigenspaces of $\Omega(\mathbf{j})$, where $\mathbf{Z}(G) = \langle \mathbf{j} \rangle \cong C_2$. Here, η has degree $(q^n - 1)/2$ and ξ has degree $(q^n + 1)/2$.

The character ω remains unchanged in the above construction if one replaces ψ by another character $\psi_a : x \mapsto \psi(ax)$ as long as $a \in \mathbb{F}_q^{\times 2}$. If, however, we replace ψ by $\psi^* : x \mapsto \psi(ax)$ for some $a \in \mathbb{F}_q^{\times} \setminus \mathbb{F}_q^{\times 2}$, then we get a new character $\omega^* = \eta^* + \xi^*$, which is the sum of another pair of Weil characters, η^* of degree $(q^n - 1)/2$ and ξ^* of degree $(q^n + 1)/2$. It is also convenient to consider $H = \mathrm{CSp}(V)$, the group of conformal transformations of the symplectic space V :

$$H = \{f \in \mathrm{GL}(V) \mid \exists \kappa(f) \in \mathbb{F}_q^{\times}, (f(u), f(v)) = \kappa(f)(u, v), \forall u, v \in V\}$$

Then a conjugation by $f \in H$ with $\kappa(f) \in \mathbb{F}_q^{\times} \setminus \mathbb{F}_q^{\times 2}$ fuses ω with ω^* , η with η^* , and ξ with ξ^* , see [Gr, Lemma 13.4].

In the course of his study of finite monodromy groups on \mathbb{A}^1 , Nick Katz [K] noted that, when $n = 1$ the alternating square $\wedge^2(\xi)$ coincides with either $\mathrm{Sym}^2(\eta)$ or $\mathrm{Sym}^2(\eta^*)$. The goal of this paper is to show that in fact this phenomenon, as well as several other interesting equalities between symmetric square, alternating squares, and tensor products, of Weil representations indeed occur for all n .

Theorem 1.1. *Let q be an odd prime power, n any positive integer, and let $\{\eta, \xi\}$, $\{\eta^*, \xi^*\}$ be the two pairs of complex Weil characters of $\mathrm{Sp}_{2n}(q)$ defined above. Then the following statements hold.*

- (i) $\xi\eta = \xi^*\eta^*$. However, $\xi\eta^* \neq \xi^*\eta$, $\xi^2 \neq (\xi^*)^2$, and $\eta^2 \neq (\eta^*)^2$.
- (ii) Suppose $q \equiv \pm 1 \pmod{8}$. Then

$$\mathrm{Sym}^2(\eta) = \mathrm{Sym}^2(\eta^*) = \wedge^2(\xi) = \wedge^2(\xi^*).$$

However, $\mathrm{Sym}^2(\xi) \neq \mathrm{Sym}^2(\xi^*)$, $\wedge^2(\eta) \neq \wedge^2(\eta^*)$.

- (iii) Suppose $q \equiv \pm 3 \pmod{8}$. Then

$$\mathrm{Sym}^2(\eta) = \wedge^2(\xi^*), \mathrm{Sym}^2(\eta^*) = \wedge^2(\xi), \mathrm{Sym}^2(\xi) = \mathrm{Sym}^2(\xi^*), \wedge^2(\eta) = \wedge^2(\eta^*).$$

However, $\mathrm{Sym}^2(\eta) \neq \mathrm{Sym}^2(\eta^*)$ and $\wedge^2(\xi) \neq \wedge^2(\xi^*)$.

Let $\epsilon := (-1)^{(q-1)/2}$. Our proof of Theorem 1.1 relies on the following result which is of independent interest:

Theorem 1.2. *Let \mathbf{j} denote the central involution of $\mathrm{Sp}_{2n}(q)$. Then for any element $g \in \mathrm{Sp}_{2n}(q)$,*

$$\omega(g)\omega(\mathbf{j}g) = \epsilon^n\omega(g^2), \quad \omega^*(g)\omega^*(\mathbf{j}g) = \epsilon^n\omega^*(g^2)$$

if $q \equiv \pm 1 \pmod{8}$, and

$$\omega(g)\omega(\mathbf{j}g) = \epsilon^n\omega^*(g^2), \quad \omega^*(g)\omega^*(\mathbf{j}g) = \epsilon^n\omega(g^2)$$

if $q \equiv \pm 3 \pmod{8}$.

In the next theorem, we construct, for the first time, infinite series of examples of irreducible symmetric cubes and alternating cubes of complex representations (of finite groups of Lie type of degree > 1), as well as irreducible tensor products of complex representations, with one factor being non-Weil. Note that previous examples of the latter kind [MT] all involve only Weil representations. Furthermore, it was shown in [GT] that, if $k \geq 4$ and Φ is a complex irreducible representation of a finite group G of degree ≥ 5 then $\mathrm{Sym}^k(\Phi)$ cannot be irreducible, unless G is one of a few (explicitly known) groups of small order. In fact, we also exhibit examples of complex representations Φ of degree > 1 with $\mathsf{S}_{2,1}(\Phi)$ being irreducible, where $\mathsf{S}_{2,1}(\cdot)$ is the Schur functor labeled by the partition $(2, 1)$, cf. [FH, §6.1]. Questions about irreducibility of tensor products and symmetric/alternating powers (or more generally, of $\mathsf{S}_\lambda(\Phi)$ for Schur functors S_λ labeled by partitions λ of a small positive integer) of representations Φ of finite groups are of interest to the Aschbacher-Scott program [A], [AS] of classifying maximal subgroups of finite classical groups, cf. [MM], [MMT], [MRT], [Ma].

Theorem 1.3. *Suppose that $n \geq 3$. Then the following characters*

$$\begin{aligned} \wedge^3(\xi), \quad \mathrm{Sym}^3(\eta), \quad \mathrm{Sym}^3(\xi) - \bar{\xi}, \quad \wedge^3(\eta) - \bar{\eta}, \quad \mathrm{Sym}^2(\xi) \otimes \eta, \\ \wedge^2(\eta) \otimes \xi, \quad \wedge^2(\xi) \otimes \eta - \bar{\eta}, \quad \mathrm{Sym}^2(\eta) \otimes \xi - \bar{\xi}, \quad \mathsf{S}_{2,1}(\xi), \quad \mathsf{S}_{2,1}(\eta) \end{aligned}$$

of $\mathrm{Sp}_{2n}(3)$ are all irreducible and pairwise distinct.

Some (but not all) of our results can also be extended to the modular case. We restrict ourselves to formulate the result in some interesting cases. Let $G = \mathrm{Sp}_{2n}(q)$ with q odd as above, and let \mathbb{F} be an algebraically closed field of characteristic $\ell \nmid 2q$. It is well known, cf. [Gr, Proposition 13.6], that there are irreducible $\mathbb{F}G$ -modules U , respectively, \tilde{U} , W , and \tilde{W} , whose Brauer characters are the restriction of η , η^* , ξ , and ξ^* , respectively, to the ℓ' -elements of G . Recall [Zs] that if $k \geq 3$ then $q^k - 1$ admits a *primitive prime divisor* $\mathrm{ppd}(q, k)$, that is, a prime divisor of $q^k - 1$ that does not divide $\prod_{j=1}^{k-1}(q^j - 1)$. Note that $\mathrm{ppd}(q, 2n)$ divides $q^n + 1$.

Theorem 1.4. *Suppose that either $\ell \nmid |G|$, or $\ell = \text{ppd}(q, 2n)$ and $n \geq 2$, or $\ell = \text{ppd}(q, n)$ and $n \geq 3$ is odd, or $n = 1$ and $q \equiv 1 \pmod{4}$ and $\ell|(q+1)/2$, or $n = 1$ and $q \equiv 3 \pmod{4}$ and $\ell|(q-1)/2$. Then the following statements hold.*

(i) *The modules $U \otimes W$, $\tilde{U} \otimes \tilde{W}$, $\text{Sym}^2(U)$, $\text{Sym}^2(\tilde{U})$, $\wedge^2(W)$, $\wedge^2(\tilde{W})$ are all semisimple. Furthermore, $U \otimes W \cong \tilde{U} \otimes \tilde{W}$. If $q \equiv \pm 1 \pmod{8}$ then*

$$\text{Sym}^2(U) \cong \text{Sym}^2(\tilde{U}) \cong \wedge^2(W) \cong \wedge^2(\tilde{W}).$$

If $q \equiv \pm 3 \pmod{8}$, then

$$\text{Sym}^2(U) \cong \wedge^2(\tilde{W}), \quad \text{Sym}^2(\tilde{U}) \cong \wedge^2(W).$$

(ii) *Suppose that $q = 3$ and $n \geq 3$. Then the modules*

$$\wedge^3(W), \quad \text{Sym}^3(U), \quad \text{Sym}^2(W) \otimes U, \quad \wedge^2(U) \otimes W, \quad \mathbf{S}_{2,1}(U), \quad \mathbf{S}_{2,1}(W)$$

of $\text{Sp}_{2n}(3)$ are all irreducible.

2. WEIL CHARACTER VALUES

Recall that we have fixed a nontrivial linear character ψ of the additive group $(\mathbb{F}_q, +)$, and let $\psi^*(x) = \psi(ax)$ for some non-square element $a \in \mathbb{F}_q^\times$. Let the functions $\gamma, \gamma^* : \mathbb{F}_q \rightarrow \mathbb{C}$ be defined via

$$\gamma(b) = q^{-1/2} \sum_{x \in \mathbb{F}_q} \psi(bx^2), \quad \gamma^*(b) = q^{-1/2} \sum_{x \in \mathbb{F}_q} \psi^*(bx^2), \quad \forall b \in \mathbb{F}_q.$$

Also, let χ denote the unique linear character of order 2 of the multiplicative group \mathbb{F}_q^\times , and let

$$d(x) := \dim_{\mathbb{F}_q} \text{Ker}(x)$$

for any $x \in \text{End}(V)$.

A key role in our arguments is played by the following result [Th, Theorem 1A, Remark 1.3]:

Theorem 2.1. *There is a function $\sigma : \text{Sp}_{2n}(q) \rightarrow \mathbb{F}_q^\times$ such that the following statements hold.*

(i) *If $g \in \text{Sp}_{2n}(q)$ then*

$$\omega(g) = q^{(1/2)d(g-1_V)} \gamma(1)^{2n-d(g-1_V)} \chi(\sigma(g)).$$

(ii) *If $g \in \text{Sp}_{2n}(q)$ and $\det(g - 1_V) \neq 0$, then $\sigma(g) = \det(g - 1_V)$.*

First we record the values of $\gamma(1)$ and $\gamma^*(1)$:

Lemma 2.2. $\gamma^*(1) = -\gamma(1)$ and $\{\gamma(1), \gamma^*(1)\} = \{\epsilon^{1/2}, -\epsilon^{1/2}\}$.

Proof. This follows, for instance, from formula (ii) in [Th, §2] and the well-known character table of $\text{Sp}_2(q) \cong \text{SL}_2(q)$, which shows that $\{\omega(t), \omega^*(t)\} = \{(\epsilon q)^{1/2}, -(\epsilon q)^{1/2}\}$ for a transvection $t \in \text{Sp}_2(q)$ when $n = 1$, see e.g. [TZ1, Lemma 2.6(iii)]. \square

Fix a decomposition $V = A \oplus A^*$ into a direct sum of two maximal totally isotropic subspaces. Then the stabilizer of this decomposition is isomorphic to $\mathrm{GL}(A) \cong \mathrm{GL}_n(q)$, and the restriction of $\omega = \omega_n$ to $\mathrm{GL}_n(q)$ gives rise to the *reducible Weil character*

$$(2.1) \quad \tau_n : g \mapsto q^{\dim_{\mathbb{F}_q} \mathrm{Ker}(g-1_A)}$$

of $\mathrm{GL}(A)$ (which is just the permutation character of the action of $\mathrm{GL}(A)$ on the point set of $A \cong \mathbb{F}_q^n$). Similarly, one can embed $\mathrm{GU}(B) \cong \mathrm{GU}_n(q)$ in $\mathrm{Sp}_{2n}(q)$ (where $B = \mathbb{F}_{q^2}^n$ is endowed with a non-degenerate Hermitian form), and then the restriction of ω_n to $\mathrm{GU}_n(q)$ gives rise to the *reducible Weil character*

$$(2.2) \quad \zeta_n : g \mapsto (-1)^n (-q)^{\dim_{\mathbb{F}_{q^2}} \mathrm{Ker}(g-1_B)}$$

of $\mathrm{GU}(B)$, cf. [Ger].

On the other hand, $G = \mathrm{Sp}_{2n}(q)$ can also be naturally embedded in $\mathrm{GL}_{2n}(q)$ and $\mathrm{GU}_{2n}(q)$, and one has the following formulae for the restriction of τ_{2n} and ζ_{2n} to G :

Lemma 2.3. *The following statements hold.*

- (i) *For any $g \in \mathrm{Sp}_{2n}(q)$, $\omega^*(g) = (-1)^{d(g-1_V)} \omega(g)$.*
- (ii) *If $q \equiv 1 \pmod{4}$, then $(\omega_n)^2 = (\omega_n^*)^2 = (\tau_{2n})|_G$ and $\omega_n \omega_n^* = (\zeta_{2n})|_G$.*
- (iii) *If $q \equiv 3 \pmod{4}$, then $(\omega_n)^2 = (\omega_n^*)^2 = (\zeta_{2n})|_G$ and $\omega_n \omega_n^* = (\tau_{2n})|_G$.*

Proof. (i) follows from Theorem 2.1(i) and Lemma 2.2. By the same statements, we also have

$$\omega(g)^2 = q^{d(g-1_V)} \epsilon^{2n-d(g-1_V)} = (\epsilon q)^{d(g-1_V)},$$

and

$$\omega(g) \omega^*(g) = q^{d(g-1_V)} (-\epsilon)^{2n-d(g-1_V)} = (-\epsilon q)^{d(g-1_V)},$$

whence (ii) and (iii) follow. \square

Note that the function σ is multiplicative in the sense that if $V = A \oplus B$ is an orthogonal sum of two non-degenerate subspaces and $g = xy$ with $x \in \mathrm{Sp}(A)$ and $y \in \mathrm{Sp}(B)$, then $\sigma(g) = \sigma(x)\sigma(y)$, cf. [Th, §4.2.1]. It follows from Theorem 2.1(i) that

$$(2.3) \quad \omega_n(g) = \omega_k(x) \omega_{n-k}(y),$$

if $\dim A = 2k$ and $\dim B = 2n - 2k$.

Proposition 2.4. *For any $g \in \mathrm{Sp}_{2n}(q)$, the following statements hold.*

- (i) $d(g^2 - 1_V) = d(g - 1_V) + d(\mathbf{j}g - 1_V)$.
- (ii) $d(\mathbf{j}g^2 - 1_V)$ is even. In particular, $\omega(\mathbf{j}g^2) = \omega^*(\mathbf{j}g^2)$.

Proof. Without loss we may extend the symplectic form (\cdot, \cdot) to $\tilde{V} := V \otimes_{\mathbb{F}_q} \overline{\mathbb{F}}_q$ and then replace V by \tilde{V} (which we then denote by V again), and $\mathrm{Sp}_{2n}(\mathbb{F}_q)$ by $\mathrm{Sp}_{2n}(\overline{\mathbb{F}}_q)$.

Let s and u denote the semisimple and unipotent part of g , and let $V_\lambda := \text{Ker}(s - \lambda \cdot 1_V)$ for any $\lambda \in \overline{\mathbb{F}}_q^\times$. Also, let i and $-i$ denote the two square roots of -1 in $\overline{\mathbb{F}}_q$. Then

$$(2.4) \quad V = \bigoplus_{\lambda \in \overline{\mathbb{F}}_q^\times} V_\lambda.$$

(i) It suffices to show that $\text{Ker}(g^2 - 1_V) = \text{Ker}(g - 1_V) \oplus \text{Ker}(g + 1_V)$. One inclusion is obvious. For the other inclusion, let $v \in \text{Ker}(g^2 - 1_V)$. Then $s^2(v) = v$, and the decomposition (2.4) implies that $v = v_+ + v_-$ with $v_+ \in V_1$ and $v_- \in V_{-1}$. As $g^2 = s^2u^2$ fixes v and stabilizes each of V_1 and V_{-1} , we must have that

$$(2.5) \quad u^2(v_+) = v_+, \quad u^2(v_-) = v_-.$$

Note that the order of the unipotent element u is odd, so (2.5) implies that u fixes each of v_+ and v_- . Thus $v_+ \in \text{Ker}(g - 1_V)$ and $v_- \in \text{Ker}(g + 1_V)$, and the statement follows.

(ii) Note that $V_\lambda \perp V_\mu$ whenever $\lambda\mu \neq 1$; in particular, $V_{\pm i} \perp V_\lambda$ whenever $\lambda \neq \pm i$. It follows that $W := V_i \oplus V_{-i}$ is a g -invariant non-degenerate subspace of V . Furthermore, the subspaces V_i and V_{-i} are both totally isotropic, and of the same dimension since s and s^{-1} are conjugate in $\text{Sp}(V)$, see [TZ3, Proposition 3.1]. Thus both $V_{\pm i}$ are maximal totally isotropic subspaces of W , and we can identify V_{-i} with the dual space V_i^* . As u stabilizes each of $V_{\pm i}$, it follows that $\text{Ker}((u|_{V_i})^2 - 1_{V_i})$ and $\text{Ker}((u|_{V_{-i}})^2 - 1_{V_{-i}})$ have the same dimension. Arguing as in (i), we can show that

$$\text{Ker}(\mathbf{j}g^2 - 1_V) = \text{Ker}((u|_{V_i})^2 - 1_{V_i}) \oplus \text{Ker}((u|_{V_{-i}})^2 - 1_{V_{-i}}).$$

Hence $\text{Ker}(\mathbf{j}g^2 - 1_V)$ has even dimension, yielding the first part of the statement. The second part now follows from Lemma 2.3(i). \square

It suffices to prove just one of the two identities in Theorem 1.2 (in each of the two cases for q); the other identity then follows using a conjugation by some $x \in \text{CSp}_{2n}(q)$.

Proposition 2.5. *Theorem 1.2 holds if $g \in \text{Sp}_{2n}(q)$ is unipotent.*

Proof. (i) Since both ω and ω^* are multiplicative, see (2.3), it suffices to prove the statement in the case g fixes no proper non-degenerate subspace of V . Since g is unipotent, it follows from the proof of [TZ2, Proposition 7.1] that the Jordan canonical form of g on V is either a single block J_{2k} of even size $2k$, or a direct sum tJ_k of $t \geq 1$ Jordan blocks of the same odd size k (in fact $t = 2$, but we do not need it).

Note that $\det(\mathbf{j}g - 1_V) = (-2)^{2n}$ is a square in \mathbb{F}_q^\times , so by Theorem 2.1 and Lemma 2.2 we have that $\chi(\sigma(\mathbf{j}g)) = 1$ and

$$(2.6) \quad \omega(\mathbf{j}g) = \epsilon^n.$$

Suppose in addition that $q \equiv \pm 1 \pmod{8}$. Then 2 is a square in \mathbb{F}_q^\times by quadratic reciprocity. By [TZ2, Proposition 7.1] and its proof, g and g^2 are conjugate in $G =$

$\mathrm{Sp}_{2n}(q)$. In particular, $\omega(g) = \omega(g^2)$, and so we are done by (2.6). So it remains to consider the case $q \equiv \pm 3 \pmod{8}$, equivalently, 2 is *not* a square in \mathbb{F}_q^\times .

(ii) Consider the case g acts on V as J_{2k} , i.e. $k = n$ and g is a regular unipotent element. By [TZ2, Lemma 7.2], there is some $x \in H \setminus \mathbf{Z}(H)G$ conjugating g to g^2 , where $H = \mathrm{CSp}_{2n}(q)$. As mentioned above, H fuses ω and ω^* , and certainly $\mathbf{Z}(H)G$ fixes each of ω and ω^* . Hence $\omega^* = \omega^x$, and so

$$\omega^*(g^2) = \omega^x(g^x) = \omega(g),$$

and we are again done by (2.6).

(iii) Now we may assume that the Jordan canonical form of g on V is tJ_k for some odd k . By part (c) of the proof of [TZ2, Proposition 7.1], g and g^2 are conjugate in G , and so $\omega(g) = \omega(g^2)$ and also $d(g - 1_V) = d(g^2 - 1_V)$. As $tk = 2n$, we have that $d(g - 1_V) = t$ is even, whence $d(g^2 - 1_V)$ is even. The latter implies by Lemma 2.3(i) that $\omega^*(g^2) = \omega(g^2)$. Thus $\omega^*(g^2) = \omega(g)$, and we are done by (2.6). \square

Proof of Theorem 1.2. Let s denote the semisimple part of g . Then we can decompose V as an orthogonal sum $V = V' \oplus V_1 \oplus V_{-1}$ of g -invariant non-degenerate subspaces, where s acts as 1_{V_1} on V_1 , $-1_{V_{-1}}$ on V_{-1} , and neither 1 nor -1 is an eigenvalue for $s|_{V'}$. Again using the multiplicativity of ω and ω^* in the sense of (2.3), it suffices to prove the statement in the case g fixes no proper non-degenerate subspace of V . The latter condition on g implies that exactly one of the three summands V' , V_1 , V_{-1} is nonzero (and then equal to V).

Suppose $V = V_1$. Then g is unipotent, and we are done by Proposition 2.5.

Next suppose that $V = V_{-1}$. Then $\mathbf{j}g$ is unipotent and $g^2 = (\mathbf{j}g)^2$, and so we are again done by applying Proposition 2.5 to $\mathbf{j}g$.

Thus we may assume now that $V = V'$, i.e. $\det(g^2 - 1_V) \neq 0$ and $d(g^2 - 1_V) = 0$. By Proposition 2.4(i),

$$d(g - 1_V) = d(\mathbf{j}g - 1_V) = 0, \quad \det(g - 1_V) \neq 0, \quad \det(\mathbf{j}g - 1_V) \neq 0.$$

It follows by Theorem 2.1 and Lemma 2.2 that

$$\omega(g) = \epsilon^n \chi(\det(g - 1_V)), \quad \omega(\mathbf{j}g) = \epsilon^n \chi(\det(\mathbf{j}g - 1_V)) = \epsilon^n \chi(\det(g + 1_V)).$$

Hence

$$\omega(g)\omega(\mathbf{j}g) = \chi(\det(g^2 - 1_V)).$$

On the other hand, Theorem 2.1 and Lemma 2.2 applied to g^2 yield

$$\omega^*(g^2) = \epsilon^n \chi(\det(g^2 - 1_V)) = \omega(g^2),$$

and the statement follows. \square

3. SYMMETRIC AND ALTERNATING SQUARES

In this section, we set $\delta := \epsilon^n = (-1)^{n(q-1)/2}$. First we give a formula for the irreducible Weil characters in terms of ω and ω^* :

Lemma 3.1. *The following statements hold for any $g \in G = \mathrm{Sp}_{2n}(q)$:*

$$\begin{aligned}\xi(g) &= \frac{\omega(g) + \delta\omega(\mathbf{j}g)}{2}, \quad \eta(g) = \frac{\omega(g) - \delta\omega(\mathbf{j}g)}{2}, \\ \xi^*(g) &= \frac{\omega^*(g) + \delta\omega^*(\mathbf{j}g)}{2}, \quad \eta^*(g) = \frac{\omega^*(g) - \delta\omega^*(\mathbf{j}g)}{2}.\end{aligned}$$

Proof. By [TZ1, Lemma 2.6(i)], \mathbf{j} acts on any G -module affording the character ξ or ξ^* as the multiplication by δ . Similarly, \mathbf{j} acts on any G -module affording the character η or η^* as the multiplication by $-\delta$. Hence

$$\omega(g) = \xi(g) + \eta(g), \quad \omega(\mathbf{j}g) = \xi(\mathbf{j}g) + \eta(\mathbf{j}g) = \delta\xi(g) - \delta\eta(g),$$

and the statements follow. \square

Corollary 3.2. $\xi\eta = \xi^*\eta^*$.

Proof. By Lemma 3.1, for any $g \in \mathrm{Sp}_{2n}(q)$ we have

$$\xi(g)\eta(g) = \frac{\omega(g)^2 - \omega(\mathbf{j}g)^2}{4}, \quad \xi^*(g)\eta^*(g) = \frac{\omega^*(g)^2 - \omega^*(\mathbf{j}g)^2}{4}.$$

Since $\omega(h)^2 = \omega^*(h)^2$ for all $h \in \mathrm{Sp}_{2n}(q)$ by Lemma 2.3, the statement follows. \square

Lemma 3.3. (i) *If $q \equiv \pm 1 \pmod{8}$ then $\mathrm{Sym}^2(\xi) \neq \mathrm{Sym}^2(\xi^*)$ and $\wedge^2(\eta) \neq \wedge^2(\eta^*)$.*
(ii) *If $q \equiv \pm 3 \pmod{8}$ then $\wedge^2(\xi) \neq \wedge^2(\xi^*)$ and $\mathrm{Sym}^2(\eta) \neq \mathrm{Sym}^2(\eta^*)$.*
(iii) *In general, $\xi\eta^* \neq \xi^*\eta$.*

Proof. (i) Consider a transvection $t \in G = \mathrm{Sp}_{2n}(q)$. As mentioned in part (i) of the proof of Proposition 2.5, $q \equiv \pm 1 \pmod{8}$ implies that t^2 is conjugate to t in G . Hence [TZ1, Lemma 2.6(iii)] implies that

$$\mathrm{Sym}^2(\xi)(t) = \frac{3 + \epsilon q^{2n-1} + 4q^{n-1}\sqrt{\epsilon q}}{8}, \quad \mathrm{Sym}^2(\xi^*)(t) = \frac{3 + \epsilon q^{2n-1} - 4q^{n-1}\sqrt{\epsilon q}}{8}.$$

In particular, $\mathrm{Sym}^2(\xi) \neq \mathrm{Sym}^2(\xi^*)$. Similarly, $\wedge^2(\eta) \neq \wedge^2(\eta^*)$.

(ii) We again evaluate relevant characters at a transvection $t \in G$. Since $q \equiv \pm 3 \pmod{8}$, 2 is not a square in \mathbb{F}_q^\times . Hence by the proof of [TZ2, Proposition 7.1], there is some $x \in H \setminus \mathbf{Z}(H)G$ that conjugates t to t^2 , where $H = \mathrm{CSp}_{2n}(q)$ as usual. As mentioned above, $\eta^x = \eta^*$, hence

$$\eta^*(t^2) = \eta^x(t^2) = \eta(t).$$

Now using [TZ1, Lemma 2.6(iii)] we can see that

$$\text{Sym}^2(\eta^*) = \frac{\eta^*(t)^2 + \eta(t)}{2} = \frac{-1 + \epsilon q^{2n-1} + 4q^{n-1}\sqrt{\epsilon q}}{8}.$$

Similarly, one computes that

$$\text{Sym}^2(\eta) = \frac{\eta(t)^2 + \eta^*(t)}{2} = \frac{-1 + \epsilon q^{2n-1} - 4q^{n-1}\sqrt{\epsilon q}}{8},$$

and so $\text{Sym}^2(\eta) \neq \text{Sym}^2(\eta^*)$. The inequality $\wedge^2(\xi) \neq \wedge^2(\xi^*)$ is proved in the same way.

(iii) Again we use [TZ1, Lemma 2.6(iii)] to see

$$\xi(t)\eta^*(t) = \frac{-(1 + q^{n-1}\sqrt{\epsilon q})^2}{4} \neq \frac{-(1 - q^{n-1}\sqrt{\epsilon q})^2}{4} = \xi^*(t)\eta(t).$$

□

Proof of Theorem 1.1. By Lemma 3.1 and Lemma 2.3(i) we have

$$(3.1) \quad \begin{aligned} \wedge^2(\xi)(g) &= \frac{\omega(g)^2 + \omega(\mathbf{j}g)^2 + 2\delta\omega(g)\omega(\mathbf{j}g)}{8} - \frac{\omega(g^2) + \delta\omega(\mathbf{j}g^2)}{4}, \\ \wedge^2(\xi^*)(g) &= \frac{\omega(g)^2 + \omega(\mathbf{j}g)^2 + 2\delta\omega^*(g)\omega^*(\mathbf{j}g)}{8} - \frac{\omega^*(g^2) + \delta\omega^*(\mathbf{j}g^2)}{4}. \end{aligned}$$

Similarly,

$$(3.2) \quad \begin{aligned} \text{Sym}^2(\eta)(g) &= \frac{\omega(g)^2 + \omega(\mathbf{j}g)^2 - 2\delta\omega(g)\omega(\mathbf{j}g)}{8} + \frac{\omega(g^2) - \delta\omega(\mathbf{j}g^2)}{4}, \\ \text{Sym}^2(\eta^*)(g) &= \frac{\omega(g)^2 + \omega(\mathbf{j}g)^2 - 2\delta\omega^*(g)\omega^*(\mathbf{j}g)}{8} + \frac{\omega^*(g^2) - \delta\omega^*(\mathbf{j}g^2)}{4}. \end{aligned}$$

Also, by Proposition 2.4(ii) we have

$$(3.3) \quad \omega(\mathbf{j}g^2) = \omega^*(\mathbf{j}g^2).$$

(i) First we consider the case $q \equiv \pm 1 \pmod{8}$. By Theorem 1.2,

$$\omega(g)\omega(\mathbf{j}g) = \delta\omega(g^2), \quad \omega^*(g)\omega^*(\mathbf{j}g) = \delta\omega^*(g^2).$$

Together with (3.1), (3.2), and (3.3), this implies that

$$\wedge^2(\xi) = \wedge^2(\xi^*) = \text{Sym}^2(\eta) = \text{Sym}^2(\eta^*).$$

(ii) Assume now that $q \equiv \pm 3 \pmod{8}$. By Theorem 1.2,

$$(3.4) \quad \omega(g)\omega(\mathbf{j}g) = \delta\omega^*(g^2), \quad \omega^*(g)\omega^*(\mathbf{j}g) = \delta\omega(g^2).$$

Together with (3.1), (3.2), and (3.4), this implies that

$$\wedge^2(\xi) = \text{Sym}^2(\eta^*), \quad \wedge^2(\xi^*) = \text{Sym}^2(\eta).$$

We also have by Lemma 3.1 and Lemma 2.3(i) that

$$\begin{aligned}\text{Sym}^2(\xi)(g) &= \frac{\omega(g)^2 + \omega(\mathbf{j}g)^2 + 2\delta\omega(g)\omega(\mathbf{j}g)}{8} + \frac{\omega(g^2) + \delta\omega(\mathbf{j}g^2)}{4}, \\ \text{Sym}^2(\xi^*)(g) &= \frac{\omega(g)^2 + \omega(\mathbf{j}g)^2 + 2\delta\omega^*(g)\omega^*(\mathbf{j}g)}{8} + \frac{\omega^*(g^2) + \delta\omega^*(\mathbf{j}g^2)}{4}, \\ \wedge^2(\eta)(g) &= \frac{\omega(g)^2 + \omega(\mathbf{j}g)^2 - 2\delta\omega(g)\omega(\mathbf{j}g)}{8} - \frac{\omega(g^2) - \delta\omega(\mathbf{j}g^2)}{4}, \\ \wedge^2(\eta^*)(g) &= \frac{\omega(g)^2 + \omega(\mathbf{j}g)^2 - 2\delta\omega^*(g)\omega^*(\mathbf{j}g)}{8} - \frac{\omega^*(g^2) - \delta\omega^*(\mathbf{j}g^2)}{4}.\end{aligned}$$

Using (3.3) and (3.4), we arrive at

$$(3.5) \quad \text{Sym}^2(\xi) = \text{Sym}^2(\xi^*), \quad \wedge^2(\eta) = \wedge^2(\eta^*).$$

(iii) The remaining statements in Theorem 1.1 now follow from Corollary 3.2 and Lemma 3.3. \square

Note that in the case $q = 3, 5$ formula (3.5) and Corollary 3.2 were established in [MT, Proposition 5.5], but by different means, along the lines of §4. We also note that, even though special unitary groups $\text{SU}_n(q)$ admit complex irreducible Weil characters of degree k and $k + 1$ for a suitable k , the symmetric square of the smaller-degree character does not coincide with the alternating square of another Weil character of the larger degree.

A further remark is that Theorem 1.1 establishes the existence of *some* isomorphisms between certain symmetric and alternating squares of complex Weil modules. It remains a question if one can construct *natural* isomorphisms between these pairs of complex modules in those cases.

4. SYMMETRIC AND ALTERNATING CUBES

Recall the Weil character τ_n of $\text{GL}_n(q)$ as defined in (2.1).

Lemma 4.1. *Fix a positive integer k and let*

$$s(k, n) := [(\tau_{2n})^k|_{\text{Sp}_{2n}(q)}, 1_{\text{Sp}_{2n}(q)}]_{\text{Sp}_{2n}(q)}.$$

Then $s(k, n) = s(k, k)$ for all $n \geq k$.

Proof. Note that $s(k, n)$ is just the number of G -orbits on the set V^k of ordered k -tuples (v_1, \dots, v_k) , with $v_i \in V = \mathbb{F}_q^{2n}$ and $G = \text{Sp}(V)$. Fix a symplectic basis $(e_1, \dots, e_n, f_1, \dots, f_n)$ of V , i.e. $(e_i, e_j) = 0 = (f_i, f_j)$ and $(e_i, f_j) = \delta_{i,j}$. Also, let

$$U_j := \langle e_1, \dots, e_j, f_1, \dots, f_j \rangle_{\mathbb{F}_q}$$

for $1 \leq j \leq k$.

(i) First we show that any G -orbit on V^k intersects $(U_k)^k$, which implies that $s(k, n) \leq s(k, k)$. In fact, we will prove by induction on $1 \leq j \leq k$ that the G -orbit of any $\alpha = (v_1, \dots, v_j) \in V^j$ intersects $(U_j)^j$.

For the induction base $j = 1$, we can certainly find $g \in G$ such that $g(v_1) = e_1 \in U_1$.

For the induction step $j \geq 2$, we may assume that there is some $g \in G$ such that $g(v_i) \in U_{j-1}$ for all $1 \leq i \leq j-1$. Now write $g(v_j) = u + w$ with

$$u \in U_{j-1}, w \in W := \langle e_j, \dots, e_n, f_j, \dots, f_n \rangle_{\mathbb{F}_q}.$$

Then we can find $h \in \mathrm{Sp}(W) \leq G$ such that h acts trivially on U_{j-1} and $h(w) = e_j$, and observe that

$$hg(\alpha) = (g(v_1), \dots, g(v_{j-1}), u + e_j) \in U_j^j.$$

(ii) Conversely, suppose that $(u_1, \dots, u_k), (v_1, \dots, v_k) \in U^k$ belong to the same G -orbit, i.e. there is some $g \in G$ such that $g(u_i) = v_i$ for all i . Then $g|_M$ is an isometry between the subspaces $M := \langle u_1, \dots, u_k \rangle_{\mathbb{F}_q}$ and $\langle v_1, \dots, v_k \rangle_{\mathbb{F}_q}$ of U_k . By Witt's lemma, $g|_M$ extends to some $h \in \mathrm{Sp}(U_k)$, whence (u_1, \dots, u_k) and (v_1, \dots, v_k) belong to the same $\mathrm{Sp}(U_k)$ -orbit. Thus $s(k, k) \leq s(k, n)$. \square

Corollary 4.2. *For $G = \mathrm{Sp}_{2n}(3)$ and $n \geq 3$,*

$$[\omega^3, \omega^3]_G = 80.$$

Proof. By Lemma 2.3(iii),

$$[\omega^3, \omega^3]_G = [(\omega\bar{\omega})^3, 1_G]_G = [(\tau_6)^3|_G, 1_G]_G = s(3, n).$$

Direct computation using [GAP] shows that $s(3, 3) = 80$, and so we are done by Lemma 4.1. \square

Proof of Theorem 1.3. From now on, let $G = \mathrm{Sp}_{2n}(3)$ with $n \geq 3$. By [Gr, Lemma 13.4],

$$\omega^* = \bar{\omega}, \xi^* = \bar{\xi}, \eta^* = \bar{\eta}.$$

(i) First we note by Theorem 1.1 and [MT, Proposition 5.4] that the characters

$$(4.1) \quad \begin{aligned} \mathrm{Sym}^2(\xi) &= \mathrm{Sym}^2(\bar{\xi}), \wedge^2(\eta) = \wedge^2(\bar{\eta}), \xi\eta = \bar{\xi}\bar{\eta} \\ &\mathrm{Sym}^2(\eta), \mathrm{Sym}^2(\bar{\eta}), \wedge^2(\xi), \wedge^2(\bar{\xi}), \xi\bar{\eta}, \bar{\xi}\eta \end{aligned}$$

are all irreducible, but $\wedge^2(\xi) \neq \wedge^2(\bar{\xi})$. It follows that

$$(4.2) \quad \begin{aligned} [\mathrm{Sym}^2(\xi) \otimes \xi, \bar{\xi}]_G &= [\mathrm{Sym}^2(\xi), \mathrm{Sym}^2(\bar{\xi}) + \wedge^2(\bar{\xi})]_G = 1, \\ [\wedge^2(\xi) \otimes \xi, \bar{\xi}]_G &= [\wedge^2(\xi), \mathrm{Sym}^2(\bar{\xi}) + \wedge^2(\bar{\xi})]_G = 0. \end{aligned}$$

On the other hand,

$$(4.3) \quad \mathrm{Sym}^2(\xi) \otimes \xi = \mathrm{Sym}^3(\xi) + \mathsf{S}_{2,1}(\xi), \wedge^2(\xi) \otimes \xi = \wedge^3(\xi) + \mathsf{S}_{2,1}(\xi),$$

where $S_{2,1}(\cdot)$ is the Schur functor labeled by the partition $(2, 1)$ of 3, see [FH, (6.8), (6.9)]. Hence, (4.2) implies that $[S_{2,1}(\xi), \bar{\xi}]_G = 0$ and

$$(4.4) \quad [Sym^3(\xi), \bar{\xi}]_G = 1;$$

in particular, $Sym^3(\xi)$ is *not* irreducible. Similarly,

$$\begin{aligned} [Sym^2(\eta) \otimes \eta, \bar{\eta}]_G &= [Sym^2(\eta), Sym^2(\bar{\eta}) + \wedge^2(\bar{\eta})]_G = 0, \\ [\wedge^2(\eta) \otimes \eta, \bar{\eta}]_G &= [\wedge^2(\eta), Sym^2(\bar{\eta}) + \wedge^2(\bar{\eta})]_G = 1, \end{aligned}$$

implying that

$$(4.5) \quad [\wedge^3(\eta), \bar{\eta}]_G = 1.$$

In particular, $\wedge^3(\eta)$ is *not* irreducible. Also, by Theorem 1.1(iii) we have

$$(4.6) \quad \begin{aligned} [\wedge^2(\xi) \otimes \eta, \bar{\eta}]_G &= [Sym^2(\bar{\eta}) \otimes \eta, \bar{\eta}]_G = [Sym^2(\bar{\eta}), Sym^2(\bar{\eta}) + \wedge^2(\bar{\eta})]_G = 1, \\ [Sym^2(\eta) \otimes \xi, \bar{\xi}]_G &= [\wedge^2(\bar{\xi}) \otimes \xi, \bar{\xi}]_G = [\wedge^2(\bar{\xi}), Sym^2(\bar{\xi}) + \wedge^2(\bar{\xi})]_G = 1. \end{aligned}$$

Now using (4.3), (4.4), (4.5), and (4.6), we can decompose

$$(4.7) \quad \begin{aligned} \omega^3 &= (\xi + \eta)^3 = \xi^3 + \eta^3 + 3\xi^2 \otimes \eta + 3\eta^2 \otimes \xi \\ &= Sym^3(\xi) + \wedge^3(\xi) + 2S_{2,1}(\xi) + Sym^3(\eta) + \wedge^3(\eta) + 2S_{2,1}(\eta) \\ &\quad + 3Sym^2(\xi) \otimes \eta + 3\wedge^2(\xi) \otimes \eta + 3Sym^2(\eta) \otimes \xi + 3\wedge^2(\eta) \otimes \xi \\ &= (Sym^3(\xi) - \bar{\xi}) + \wedge^3(\xi) + 2S_{2,1}(\xi) + Sym^3(\eta) + (\wedge^3(\eta) - \bar{\eta}) + 2S_{2,1}(\eta) \\ &\quad + 3Sym^2(\xi) \otimes \eta + 3(\wedge^2(\xi) \otimes \eta - \bar{\eta}) + 3(Sym^2(\eta) \otimes \xi - \bar{\xi}) + 3\wedge^2(\eta) \otimes \xi \\ &\quad + 4\bar{\xi} + 4\bar{\eta} \end{aligned}$$

into a sum of *true* characters of G . This decomposition implies in particular that $[\omega^3, \omega^3]_G \geq 80$. On the other hand, Corollary 4.2 states that $[\omega^3, \omega^3]_G = 80$. It follows that the summands in the last expression given in (4.7) for ω^3 must be all irreducible and pairwise distinct. \square

5. THE MODULAR CASE

In this section, we prove Theorem 1.4. The statements are obvious if $\ell \nmid |G|$, so we will assume that $\ell \mid |G|$. Now the assumption on ℓ implies that $\ell \mid (q^{2n} - 1)$, and that any irreducible character θ of G has ℓ -defect 0 if $\ell \mid (q^n - 1)$ but $(q^n - 1)/(q - 1) \mid \theta(1)$, or if $\ell \mid (q^n + 1)$ but $(q^n + 1)/2 \mid \theta(1)$.

(a) Now, in the case of Theorem 1.4(ii), note by Theorem 1.3 that the Brauer character φ of any module in the list is the reduction modulo ℓ of some irreducible character $\theta \in \text{Irr}(G)$ of degree divisible by $(q^{2n} - 1)/4$. By the aforementioned remark, θ has ℓ -defect 0, and so φ is irreducible as claimed.

(b) We will now work on the proof of Theorem 1.4(i). As mentioned in [MT, §5], the restriction of the character τ_{2n} in Lemma 2.3 to G is the sum of twice the

trivial character, an irreducible character α_n of degree $(q^n - 1)(q^n + q)/2(q - 1)$ (of multiplicity 1), an irreducible character β_n of degree $(q^n + 1)(q^n - q)/2(q - 1)$ (of multiplicity 1) if $n > 1$, two irreducible characters of degree $(q^{2n} - 1)/2(q - 1)$ (each of multiplicity 1), and $(q - 3)/2$ irreducible characters of degree $(q^{2n} - 1)/(q - 1)$ (each of multiplicity 2).

Similarly, the restriction of the character ζ_{2n} in Lemma 2.3 to G is the sum of an irreducible character γ_n of degree $(q^n + 1)(q^n + q)/2(q + 1)$ (of multiplicity 1), an irreducible character δ_n of degree $(q^n - 1)(q^n - q)/2(q + 1)$ (of multiplicity 1) if $n > 1$, two irreducible characters of degree $(q^{2n} - 1)/2(q + 1)$ (each of multiplicity 1), and $(q - 1)/2$ irreducible characters of degree $(q^{2n} - 1)/(q + 1)$ (each of multiplicity 2).

(c) Here we consider the case where $q \equiv 1 \pmod{4}$ and moreover $\ell|(q + 1)$ if $n = 1$. Set $e := (q^{2n} - 1)/2(q - 1)$. By Theorem 1.1(i) and Lemma 2.3(ii), $\xi\eta = \xi^*\eta^*$ and $(\tau_{2n})|_G - 2\xi\eta$ is a *true* character. In particular, every irreducible constituent of $\xi\eta$ must occur in $(\tau_{2n})|_G$ with multiplicity at least 2. Clearly, 1_G is not an irreducible constituent of $\xi\eta$. It follows from the remarks in (b) that every irreducible constituent in $\xi\eta$ has degree $2e$, whence it has ℓ -defect 0 (and occurs with multiplicity 1). Hence the module $U \otimes W$ is semisimple, and similarly for $\tilde{U} \otimes \tilde{W}$. As these two modules have the same Brauer character, they are isomorphic.

It is well known, cf. [Gr, Corollary 13.7], that ξ has Frobenius-Schur indicator +1 and η has Frobenius-Schur indicator -1. Thus 1_G is neither an irreducible constituent of $\wedge^2(\xi)$, nor of $\text{Sym}^2(\eta)$. Next, the common degree of $\wedge^2(\xi)$ and $\text{Sym}^2(\eta)$ is divisible by e . On the other hand, none of the integers $\alpha_n(1)$, $\beta_n(1)$, and $\alpha_n(1) + \beta_n(1)$ is divisible by e . It follows from the remarks in (b) that every irreducible constituent in $\wedge^2(\xi)$ and in $\text{Sym}^2(\eta)$ has degree divisible by e , whence they all have ℓ -defect 0. Hence the modules $\text{Sym}^2(U)$ and $\wedge^2(W)$ are semisimple, and similarly for $\text{Sym}^2(\tilde{U})$ and $\wedge^2(\tilde{W})$. In particular, if any two of these four modules have the same Brauer character, then they are also isomorphic. Hence we are done by Theorem 1.1.

(d) Here we consider the case where $q \equiv 3 \pmod{4}$ and moreover $\ell|(q - 1)$ if $n = 1$. Set $f := (q^{2n} - 1)/2(q + 1)$. By Theorem 1.1(i) and Lemma 2.3(iii), $\xi\eta = \xi^*\eta^*$ and $(\zeta_{2n})|_G - 2\xi\eta$ is a *true* character. In particular, every irreducible constituent of $\xi\eta$ must occur in $(\zeta_{2n})|_G$ with multiplicity at least 2. It follows from the remarks in (b) that every irreducible constituent in $\xi\eta$ has degree $2f$, whence it has ℓ -defect 0 (and occurs with multiplicity 1). Hence the module $U \otimes W$ is semisimple, and similarly for $\tilde{U} \otimes \tilde{W}$. As these two modules have the same Brauer character, they are isomorphic.

As above, the common degree of $\wedge^2(\xi)$ and $\text{Sym}^2(\eta)$ is divisible by f . On the other hand, none of the integers $\gamma_n(1)$, $\delta_n(1)$, and $\gamma_n(1) + \delta_n(1)$ is divisible by f . It follows from the remarks in (b) that every irreducible constituent in $\wedge^2(\xi)$ and in $\text{Sym}^2(\eta)$ has degree divisible by f , whence they all have ℓ -defect 0. Hence the modules $\text{Sym}^2(U)$ and $\wedge^2(W)$ are semisimple, and similarly for $\text{Sym}^2(\tilde{U})$ and $\wedge^2(\tilde{W})$. In particular, if any two of these four modules have the same Brauer character, then

they are also isomorphic. Hence we are again done by Theorem 1.1, and the proof of Theorem 1.4 is completed.

REFERENCES

- [A] M. Aschbacher, On the maximal subgroups of the finite classical groups, *Invent. Math.* **76** (1984), 469–514.
- [AS] M. Aschbacher and L. Scott, Maximal subgroups of finite groups, *J. Algebra* **92** (1985), 44–80.
- [BRW] B. Bolt, T. G. Room and G. E. Wall, On the Clifford collineation, transform and similarity groups. I, *J. Austral. Math. Soc.* **2** (1961–62), 60–79.
- [D] N. Dummigan, Symplectic group lattices as Mordell-Weil sublattices, *J. Number Theory* **61** (1996), 365–387.
- [DT] N. Dummigan and Pham Huu Tiep, Lower bounds for the minima of certain symplectic and unitary group lattices, *Amer. J. Math.* **121** (1999), 889–918.
- [FH] W. Fulton and J. Harris, ‘*Representation Theory*’, Springer-Verlag, New York, 1991.
- [GAP] The GAP group, ‘*GAP - groups, algorithms, and programming*’, Version 4.4, 2004, <http://www.gap-system.org>.
- [Ger] P. Gérardin, Weil representations associated to finite fields, *J. Algebra* **46** (1977), 54–101.
- [Gow] R. Gow, Even unimodular lattices associated with the Weil representation of finite symplectic group, *J. Algebra* **122** (1989), 510–519.
- [Gr] B. H. Gross, Group representations and lattices, *J. Amer. Math. Soc.* **3** (1990), 929–960.
- [GT] R. M. Guralnick and Pham Huu Tiep, Symmetric powers and a conjecture of Kollar and Larsen, *Invent. Math.* **174** (2008), 505–554.
- [Hw] R. E. Howe, On the characters of Weil’s representations, *Trans. Amer. Math. Soc.* **177** (1973), 287–298.
- [Is] I. M. Isaacs, Characters of solvable and symplectic groups, *Amer. J. Math.* **95** (1973), 594–635.
- [K] N. Katz, Rigid local systems on \mathbb{A}^1 with finite monodromy, (preprint).
- [MM] K. Magaard and G. Malle, Irreducibility of alternating and symmetric squares, *Manuscripta Math.* **95** (1998), 169–180.
- [MMT] K. Magaard, G. Malle, and Pham Huu Tiep, Irreducibility of tensor squares, symmetric squares and alternating squares, *Pacific J. Math.* **202** (2002), 379–427.
- [MRT] K. Magaard, G. Röhrle, and D. Testerman, On the irreducibility of symmetrizations of cross-characteristic representations of finite classical groups, *J. Pure Appl. Alg.* **217** (2013), 1427–1446.
- [MT] K. Magaard and Pham Huu Tiep, Irreducible tensor products of representations of quasi-simple finite groups of Lie type, in: ‘*Modular Representation Theory of Finite Groups*’, M. J. Collins, B. J. Parshall, L. L. Scott, eds., Walter de Gruyter, Berlin et al, 2001, pp. 239–262.
- [Ma] G. Malle, Almost irreducible tensor squares, *Comm. Algebra* **27** (1999), 1033–1051.
- [ST1] R. Scharlau and Pham Huu Tiep, Symplectic groups, symplectic spreads, codes, and unimodular lattices, *J. Algebra* **194** (1997), 113–156.
- [ST2] R. Scharlau and Pham Huu Tiep, Symplectic group lattices, *Trans. Amer. Math. Soc.* **351** (1999), 2101–2139.
- [Se] G. M. Seitz, Some representations of classical groups, *J. London Math. Soc.* **10** (1975), 115–120.

- [Th] T. Thomas, The character of the Weil representation, *J. Lond. Math. Soc.* **77** (2008), 221–239.
- [T1] Pham Huu Tiep, Globally irreducible representations of finite groups and integral lattices, *Geom. Dedicata* **64** (1997), 85–123.
- [T2] Pham Huu Tiep, Weil representations as globally irreducible representations, *Math. Nachr.* **184** (1997), 313–327.
- [TZ1] Pham Huu Tiep and A. E. Zalesskii, Some characterizations of the Weil representations of the symplectic and unitary groups, *J. Algebra* **192** (1997), 130–165.
- [TZ2] Pham Huu Tiep and A. E. Zalesskii, Unipotent elements of finite groups of Lie type and realization fields of their complex representations, *J. Algebra* **271** (2004), 327–390.
- [TZ3] Pham Huu Tiep and A. E. Zalesskii, Real conjugacy classes in algebraic groups and finite groups of Lie type, *J. Group Theory* **8** (2005), 291–315.
- [Wa] H. N. Ward, Representations of symplectic groups, *J. Algebra* **20** (1972), 182–195.
- [W] A. Weil, Sur certains groupes d’opérateurs unitaires, *Acta Math.* **111** (1964), 143–211.
- [Z] A. E. Zalesskii, Eigenvalues of matrices of complex representations of finite groups of Lie type, Lecture Note Math. no. **1352**, Springer, Berlin et al, 1988, pp. 206–218.
- [Zs] K. Zsigmondy, Zur Theorie der Potenzreste, *Monatsh. Math. Phys.* **3** (1892), 265–284.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA 90089-2532, USA

E-mail address: guralnic@usc.edu

SCHOOL OF MATHEMATICS, UNIVERSITY OF BIRMINGHAM, EDGBASTON, BIRMINGHAM, B15 2TT, UK

E-mail address: kymgrd@yahoo.com

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATAWAY, NJ 08854-8019, USA

E-mail address: tiep@math.rutgers.edu