

## Revision of *Batrachospermum* Section *Virescentia* (Batrachospermales, Rhodophyta) with the Establishment of the New Genus, *Virescentia* Stat. nov.

Author(s): Orlando Necchi Jr, Douglas C. Agostinho & Morgan L. Vis

Source: Cryptogamie, Algologie, 39(3):313-338.

Published By: Association des Amis des Cryptogames

https://doi.org/10.7872/crya/v39.iss3.2018.313

URL: http://www.bioone.org/doi/full/10.7872/crya/v39.iss3.2018.313

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms of use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

# Revision of *Batrachospermum* section *Virescentia* (Batrachospermales, Rhodophyta) with the establishment of the new genus, *Virescentia* stat. nov.

Orlando NECCHI Jra\*, Douglas C. AGOSTINHOa & Morgan L. VISb

<sup>o</sup>São Paulo State University, Zoology and Botany Department, Rua Cristóvão Colombo, 2265, S. José Rio Preto, SP 15054-000 Brazil

<sup>b</sup>Ohio University, Department of Environmental and Plant Biology, Porter Hall, Athens, OH 45701 USA

Abstract – In recent years, sections of the paraphyletic genus Batrachospermum have been investigated using DNA sequence data, as well as morphology and some have been raised to genera in order to resolve this paraphylly. The species of Batrachospermum section Virescentia form a well-supported clade and we propose the raising of this section to the genus Virescentia stat. nov. In addition, we re-evaluated the characters used to circumscribe species by reexamination of type specimens and new collections in historically important areas. Of the eleven previously accepted species, we recognize Dve species and a new species is proposed as follows: V. crispata, V. gulbenkiana (synomyms V. azeredoi and V. ferreri), V. helminthosa (type species), V. viride-americana sp. nov., V. viride-brasiliensis and V. vogesiaca. The species are circumscribed on the basis of morphological characteristics (shape of fascicles, occurrence of secondary fascicles, disposition of carpogonial branches and size of carpogonia), geographic distribution, and DNA sequence data (rbcL and COI-5P). Descriptions, identi cation key and photomicrographs are presented for each recognized species. Two potential species from Japan based on DNA sequence data remained undescribed due to lack of voucher specimens for morphological observation. Other species previously assigned to the section Virescentia are referred to other genera of Batrachospermales based on morphology (disposition of carpogonial branches, shape of trichogynes, types of gonimoblast \(\bar{\pi}\) aments and arrangement of carposporophytes).

COI-5P / Batrachospermales / biogeography / morphology / phylogeny / rbcL / revision / Rhodophyta / taxonomy

### INTRODUCTION

The genus *Batrachospermum* has been shown to be paraphyletic in previous phylogenetic studies of the freshwater red algal order Batrachospermales (Vis *et al.*, 1998; Entwisle *et al.*, 2009). The genus *Batrachospermum* included nine sections (*Acarposporophytum, Aristata, Batrachospermum, Gonimopropagulum, Helminthoidea, Macrospora, Setacea, Turfosa* and *Virescentia*) and was provisionally kept by Entwisle *et al.* (2009) until further detailed taxonomic research could be conducted; an informal group (named 'Australasica') was also recognized. The

<sup>\*</sup> Corresponding author: orlando@ibilce.unesp.br

genus *Kumanoa* was established for species formerly classi ☐ed in sections *Contorta* and *Hybrida* (Entwisle *et al.*, 2009). Subsequently, species of the monophyletic section *Helminthoidea* were investigated and the genus *Sheathia* was proposed with the transfer of previously described species and proposal of new species (Salomaki *et al.*, 2014). More recently, section *Setacea* was raised to the genus level as *Atrophycus* (Rossignolo & Necchi, 2016; Rossignolo *et al.*, 2016). However, Entwisle *et al.* (2016) adopted an alternative scheme combining section *Setacea*, all taxa of the 'Australasica' group and previously described species of *Nothocladus* into the genus *Nothocladus* "lato sensu" based on phylogenetic analyses of ☐ve genes.

Virescentia is one of the sections within the paraphyletic genus Batrachospermum that has yet to be revised. DNA sequence (plastid-encoded RuBisCO large-subunit gene, rbcL and nuclear small subunit ribosomal DNA, SSU) evidence has shown this section to be monophyletic (Vis et al., 1998, 2001; Hanyuda et al., 2004; Entwisle et al., 2009; Agostinho & Necchi, 2014). Section Virescentia was proposed by Sirodot (1873) to include species of the genus with greenish thalli and carpogonia having cylindrical and stalked trichogynes. Additional characters have been added to the section including well-developed whorls, carpogonial branches composed of short cells that are well-differentiated from the fascicle cells. carposporophytes inserted centrally in the whorl, densely arranged and large (Necchi, 1990; Sheath et al., 1994; Kumano, 2002; Agostinho & Necchi, 2014). The monograph of Batrachospermum by Sirodot (1884) represents the starting point for the studies on this section with seven new species described and illustrated. Subsequent taxonomic treatments for species of this section have been mostly in more general studies, in which species of the genus Batrachospermum for various regions of the world were described such as Israelson (1942) for Sweden, Mori (1975) for Japan, Necchi (1990) for Brazil and Eloranta et al. (2011) for Central Europe. Only two studies have focused exclusively on species of the section, Sheath et al. (1994) for North America and Agostinho & Necchi (2014) for Brazil.

Kumano (2002) recognized eleven species within the section: B. azeredoi Reis (1967), B. bakarense Kumano & Ratnas. (Ratnasabapathy & Kumano, 1984), B. crispatum Kumano & Ratnas. (Ratnasabapathy & Kumano, 1982a), B. desikacharvi Sankaran (1984), B. elegans Sirodot (1884), B. ferreri Reis (1967), B. gombakense Kumano & Ratnas. (Ratnasabapathy & Kumano, 1982b), B. gulbenkianum Reis (1965), B. helminthosum Bory (1808), B. transtaganum Reis (1970) and B. vogesiacum T.G. Schultz ex Skuja (1938). The following criteria have been applied to distinguish species within the section: sexuality (monoecious, dioecious or polyoecious), position and length of the carpogonial branch, size of carpogonia, number of cell storeys in primary fascicles, density of secondary fascicles and the presence of terminal hairs (Sirodot, 1884; Mori, 1975; Starmach, 1977; Necchi, 1990). The morphometric analyses by Sheath et al. (1994) of type specimens and North American populations resulted in the proposal of many synonymies and recognition of only two species, B. helminthosum Bory (synonyms B. bruziense Sirodot, B. graibussoniense Sirodot, B. sirodotii Reis – as B. virgatum (Kützing) Sirodot – B. testale Sirodot and B. vogesiacum Schultz – as B. □agelliforme Sirodot) and B. elegans Sirodot (synonym B. coerulescens Sirodot). Sheath et al. (1994) distinguished the two species based on the presence in B. elegans or absence in B. helminthosum of basal knobs or protuberances on the trichogynes. However, Vis et al. (2001) showed that most qualitative features are universally present in B. helminthosum and B. elegans and morphometric characteristics are highly variable, with signi cant overlap among specimens of the two species. Hanyuda et

al. (2004) also concluded that characters generally used to distinguish species (sexuality, origin of the carpogonium bearing branch and trichogynes with or without basal knobs or branching) are not  $\Box$ xed genetically, but are affected by environmental conditions (seasonally or environmentally induced changes in the gametophyte) and proposed that B. elegans should be placed in synonymy with B. helminthosum. Agostinho & Necchi (2014) noted considerable divergence in morphometric characters (size of carpogonia, carposporophytes and carposporangia) among samples from Brazil and North America and type specimens from Europe indicating the existence of at least three distinct species with non-overlapping geographic distributions: Brazil, USA and Europe. The recognition of these groups was supported by the sequence analyses of rbcL and barcode region of the mitochondrial encoded cytochrome c oxidase subunit 1, COI-5P (Hanyuda et al., 2004; Rueness, 2010; Agostinho & Necchi, 2014). Sequence data of the *rbc*L also indicated the occurrence of more than one species within the section *Virescentia* from Japan (Hanyuda et al., 2004). Chiasson et al. (2014) revisited historically important areas in France, where several species were originally collected by Bory (1808) and Sirodot (1884). They presented rbcL sequenced data clearly showing B. vogesiacum being a species distinct from B. helminthosum.

While the morphological study of Sheath *et al.* (1994) reduced the number of species in the section, DNA sequence data have suggested that there is considerable genetic variation among populations and this variation is partitioned geographically (Hanyuda *et al.*, 2004; Agostinho & Necchi, 2014). We re-analyzed the DNA sequence data to evaluate the taxonomic status of the section. In addition, we have examined the morphology of type specimens, new collections from historically important and specimens for which there is DNA sequence data to provide updated species information.

#### MATERIALS AND METHODS

Sixteen type specimens of species attributed to section *Virescentia* were examined (Table 1). Type specimens were borrowed from the following herbaria (acronyms according to Thiers *et al.*, 2018): COI, PC and TNS-AL. In addition, taxonomically relevant specimens from which sequence data were reported were examined as follows: two specimens from France (Chiasson *et al.*, 2014) and ten specimens from USA (Vis *et al.*, 2001; House *et al.*, 2008; Sherwood *et al.*, 2008) in BHO. The synonyms listed for each species are all designated in this study.

All morphological and morphometric characters referred to in relevant taxonomic literature of section *Virescentia* (Sirodot, 1884; Israelson, 1942; Mori, 1975; Necchi, 1990; Sheath *et al.*, 1994; Kumano, 2002; Agostinho & Necchi, 2014) were evaluated. For observations and measurements, we used an Olympus SZH10 stereoscope and an Olympus BH-2 light microscope (Olympus Corp., Tokyo, Japan). Photomicrographs were taken with a Leica DFC 320 digital camera using LAS capture and image analysis software coupled to a Leica DM5000 microscope (Leica Microsystems, Wetzlar, Germany).

Tree reconstruction analyses were conducted using DNA sequence data from previously published studies (Tables S1-S2) available in GenBank (Benson *et al.*, 2013). For COI-5P data, distance analysis using Neighbor Joining (NJ) and Kimura two-parameter distances was performed using a plugin of Geneious version

**Table 1.** Type and historically important specimens of *Batrachospermum* section *Virescentia* examined in this study.

Species	Type and herbarium number <sup>1</sup>	Locality, collector and date
B. azeredoi Reis	Holotype, COI	Portugal, Vila da Feira, River Caster, M.P. Reis & A. Santos, no. 482, 17.vi.1966
B. bakarense Kumano & Ratnas.	Isotype, TNS-AL 169085	Malaysia, Kelantan, Sungai Bakar, M. Ratnasabapathy no 3, 03.vi.1982
B. bruziense Sirodot	Lectotype, PC0452411	France, town of Bruz near Rennes, Fontaine de Chatillon; S. Sirodot, 27.iii.1882
B. coerulescens Sirodot	Not type, PC0452372	France, Le Meu at the top of Montfort near Tréguil; S. Sirodot, 29.vii.1874
B. crispatum Kumano & Ratnas.	Isotype, TNS-AL 169089	Malaysia, Pulau Tioman, Sungai Ayer Besar; M. Ratnasabapathy no. 21, 24.v.1974
B. elegans Sirodot	Lectotype, PC0452233	France, Morbihan near Compénéac, Fontaine de Pont- Garnier; S. Sirodot, 02.v.1869
B. ferreri Reis	Holotype, COI	Portugal, Vale de Cambra, Mina do Pintor, M.P. Reis & A. Santos, no. 485, 17.vi.1966
B. gibbosum Reis	Holotype, COI	Portugal, São Pedro do Sul, River Arões; M.P. Reis & A. Santos, no. 449, 07.vii.1965
B. gombakense Kumano & Ratnas.	Isotype, TNS-AL 169097	Malaysia, Selangor, Sungai Gombak; M. Ratnasabapathy no. 1221, 31.v.1976
B. gulbenkianum Reis	Holotype, COI	Portugal, Confulcos, River Águeda; M.P. Reis & A. Santos, no. 456, 07.vii.1965
B. helminthosum Bory	Lectotype, PC0591734	France, Fougéres; Bory de St. Vincent, no collection date
B. testale Sirodot	Lectotype, PC0451869	France, near Montfort, Fontaine de Gaillardon; S. Sirodot, 24.iv.1882
B. transtaganum Reis	Holotype, COI 542	Portugal, Odemira, River Torgal; M.P. Reis no. 542A, 19.iv.1968
B. viride Sirodot	Lectotype, PC0451877	France, near Rennes, Ruisseaux de Gallet; S. Sirodot, 02.v.1882
B. viride-brasiliense Necchi & D.C. Agostinho	Holotype, SJRP 29845	Brazil, São Paulo State: Campos do Jordão, 'Ducha da Prata', M.O. Paiano, 15.xii.2009
B. vogesiacum T. G. Schultz ex Skuja	Lectotype, PC0591499	France, Laugerie-Haute; S. Sirodot, 01.viii.1880

<sup>1.</sup> Herbarium acronyms according to Thiers (2018).

10 (http://www.geneious.com, Kearse et al., 2012). For phylogenetic analyses of the rbcL data, a GTR+I+G was determined to be the best-□t model of sequence evolution by the Akaike Information Criterion using jModelTest 2.1.4 (Darriba et al., 2012). Two separate analyses were conducted based on: 1) a general matrix with sequences of representative members of the Batrachospermales; 2) a matrix with all sequences available for members of section Virescentia only. Maximum-likelihood (ML) topologies and bootstrap values from 1,000 replicates were inferred using Randomized Accelerated Maximum Likelihood graphic user interface (RAxMLGUI version 1.2;

Silvestro & Michalak, 2011). Bayesian analysis (BA) was performed in MrBayes using Geneious plugin, with three runs of □ve chains of Metropolis coupled Markov Chain Monte Carlo for 10 x10<sup>6</sup> generations. 500,000 chains were removed as burnin prior to determining the posterior probabilities. The COI-5P and *rbc*L sequence data were not combined because there were few specimens in common between the two data sets of section *Virescentia* (Table S1), which would have considerably limited the number of specimens in the analysis.

#### RESULTS AND DISCUSSION

## Molecular analyses

BA and ML analyses of the rbcL sequences revealed strong support for the section Virescentia as a monophyletic group (Fig. S1, Supporting information), corroborating results of previous studies (Vis et al., 1998; Hanyuda et al., 2004; Entwisle et al., 2009; Agostinho & Necchi, 2014; Chiasson et al., 2014). Within the section, there were Due clades with high support, corresponding to four geographic regions (Fig. S2, Supporting information): Asia, Europe (two clades), North America (USA) and South America (Brazil). Each clade represented a species, as follows: Asia (Batrachospermum sp.), Europe (B. helminthosum and B. vogesiacum), North America (Batrachospermum sp.) and South America (B. viride-brasiliense). There was a remaining sequence from Asia (Japan) that did not form a close association with other specimens and may also represent a distinct species. This analysis corroborates previous studies with the recognition of two species from Asia (Japan), as found by Hanyuda et al. (2004), two species from Europe as reported by Chiasson et al. (2014), one species from North America (USA) and another one from South America (Brazil), as reported by Agostinho & Necchi (2014). The sequences from USA formed a clade with two groups showing sequence divergences within a group of 0-19 bp (0-1.5%) and between groups of 15-30 bp (1.2-2.4%).

The COI-5P NJ tree (Fig. S3, Supporting information) showed three well-supported groups, with clear geographical disjunction between clades of sequences from Europe (Norway), North America (USA) and South America (Brazil); no data for Asia were available. As shown by *rbcL* analysis, sequences of each clade represent a species, as follows: Europe (*B. helminthosum*), North America (*Batrachospermum* sp.) and South America (*B. viride-brasiliense*). These results match those reported by Agostinho & Necchi (2014). The sequences from USA formed a group with two minor groups, showing sequence divergences within each minor group of 0-28 bp (0-4.3%) and between groups of 36-46 bp (5.5-7.0%).

## Morphological analyses

On the basis of the high support of *Batrachospermum* section *Virescentia* in the phylogenetic analyses (Vis *et al.*, 1998; Hanyuda *et al.*, 2004; Vis *et al.*, 2005; Entwisle *et al.*, 2009; Agostinho & Necchi, 2014; Chiasson *et al.*, 2014; this study) and to continue to rectify the paraphyly of the genus *Batrachospermum* by raising sections that form distinct clades within the genus *Batrachospermum* to the generic level, we propose this section as a genus.

Virescentia (Sirodot) Necchi, D.C.Agostinho & M.L.Vis, gen. et stat. nov.

**Basionym:** section *Virescentia* Sirodot, *Comptes-rendus hebdomadaires des séances de l'académie des sciences* 76: 1219 (1873).

**Synonyms:** section *Viridia* De Toni, *Sylloge Floridearum*, 60 (1897); section *Verts* Sirodot, *Batrachospermes*, 269 (1884); section *Claviformia* Reis, *Boletim da Sociedade Broteriana* 46: 209 (1972).

Revised description: Plants monoecious, dioecious or polyoecious, greenish, bluegreenish or less often brownish; branching irregular or pseudo-dichotomous; whorls well-developed, contiguous or separated, obconic, pear-shaped, spherical or barrelshaped; cortical \(\sigma\) aments of the main axis well-developed, formed by one or two layers of cylindrical cells; main axis sometimes brownish colored; primary fascicles formed by cells variable in shape along the fascicle, ranging from cylindrical to elliptical, obovoid or pear-shaped or audouinelloid (same shape along the fascicle, cylindrical or barrel-shaped), with 6-19 cell storeys; secondary fascicles few and sparse to abundant and covering the entire internode; spermatangia spherical or obovoid, on primary or secondary fascicles, 4-9 µm wide; carpogonial branches well differentiated from the fascicles, straight, rarely curved, developing from the periaxial, proximal cells of primary fascicles, rarely from distal cells of primary fascicles, secondary fascicles or cortical \( \square\) aments, short or long, composed of 1-12(-20) disc- or barrel-shaped cells, 3.0-37.5(-68) μm long; involucral flaments, short, composed 1-4 cylindrical, elliptical or spherical cells; carpogonia 20-110 µm long, with stalked, rarely shortly stalked, cylindrical, sub-cylindrical, club-shaped or elliptical trichogynes; carposporophytes axial, hemispherical or spherical, inserted within the whorls, 1-2, rarely 3, per whorl, large, (55-)120-550 µm wide and (30-)100-420 µm in height; gonimoblast □aments densely arranged, composed of 2-7 cylindrical, elliptic or barrel-shaped cells; carposporangia obovoid, pear-shaped, club-shaped or elliptic, 10-35 µm long, 6.5-24 µm wide.

Type species: V. helminthosa (Bory) Necchi, D.C.Agostinho & M.L.Vis, comb. nov.

Remarks: members of the genus can be recognized by the combinations of characters: including presence of well-developed whorls; carpogonial branches well differentiated from the fascicles, straight, rarely curved; carpogonia with cylindrical, club-shaped or elliptical, stalked trichogynes; axial, large and dense carposporophytes. Members of *Virescentia* are comparable with those of section *Turfosa* in gross vegetative morphology and presence of large carposporophytes, but differ having only one type of gonimoblast \(\percap{\text{\text{daments}}}\), which grow radially from the fertilized carpogonia forming a hemispherical or spherical structure. Members of the genus *Kumanoa* also have the same type of carposporophytes, but they have strongly curved or twisted carpogonial branches.

## Key to the species of the genus Virescentia

3b.	Carpogonia long, (30-)40-110 µm in length, known distribution	
	in the Americas	
	4a. Secondary fascicles few and sparse	.V. gulbenkiana
	4b. Secondary fascicles abundant and covering ≥ half	
	of the internode length	V. vogesiaca
5a.	Known distribution in North America (USA)V. vi	ride-americana
	Known distribution in South America (Brazil) V. vir	

Virescentia crispata (Kumano & Ratnas.) Necchi, D.C.Agostinho & M.L.Vis, comb. nov. Figs 1-7

**Basionym:** Batrachospermum crispatum Kumano & Ratnas., Japanese journal of phycology 30: 18 (1982a).

Revised description: Plants dioecious or polyoecious; branching pseudo-dichotomous; whorls contiguous, obconic or pear-shaped, 150–350 μm wide; primary fascicles curved, unilaterally branched, with 6–13 cell storeys; secondary fascicles abundant and covering the entire internode; spermatangia spherical or obovoid, on primary or secondary fascicles, 6–8 μm wide; carpogonial branches slightly curved, composed of 3-4 disc- or barrel-shaped cells, developing from the periaxial or proximal cells of primary fascicles, short, 20-28 μm long; involucral □aments, short, composed of 2-4 cylindrical or elliptic cells; carpogonia 54-75 μm long, with stalked and cylindrical trichogynes; carposporophytes 1-2 per whorl, 140-300 μm wide and 100-250 μm high; gonimoblast μm long and 8-10 μm wide. Cells; carposporangia club-shaped or obovoid, 14-30 μm long and 8-10 μm wide.

**Distribution:** Asia (Malaysia), known from the type locality only (Ratnasabapathy & Kumano, 1982a).

**Remarks:** This is the most easily distinguished species of the genus due to the presence of audouinelloid and curved primary and secondary fascicles, a character lacking in the remaining species.


Virescentia gulbenkiana (Reis) Necchi, D.C.Agostinho & M.L.Vis, comb. nov. Figs 8-12

**Basionym:** Batrachospermum gulbenkianum Reis, Anuário da Sociedade Broteriana 31: 31, 1965).

**Synonyms:** Batrachospermum azeredoi Reis, Boletim da sociedade Broteriana 41: 167, (1967); B. ferreri Reis, Boletim da sociedade Broteriana 41: 174 (1967); B. gibbosum Reis, Boletim da sociedade Broteriana 43: XXX (1969).


Revised description: plants monoecious or polyoecious; branching irregular; whorls contiguous or separated, barrel-shaped or spherical, 430-810 μm wide; primary fascicles straight, with 8-13 cell storeys; secondary fascicles few and sparse; spermatangia spherical, on primary fascicles, 4-6 μm wide; carpogonial branches straight, composed of 3-12 disc-shaped cells, developing from the periaxial or proximal cells of primary fascicles, short or long, 9-35 μm long; involucral □laments, short, composed of 1-3 cylindrical or elliptic cells; carpogonia 20-39 μm long, with stalked, club-shaped or cylindrical trichogynes; carposporophytes 1-3 per whorl, 120-300 μm wide and 60-150 μm high; gonimoblast □laments with 2-5 cylindrical cells; carposporangia obovoid or pear-shaped, 10-18 μm long, 6.5-10 μm wide.

**Distribution:** Europe, several localities in Portugal (Reis 1965, 1967, 1972, 1974). **Remarks:** the distinction of *V. gulbenkiana* from *B. azeredoi*, *B. ferreri* and *B. gibbosum* was very unclear, broadly overlapping for most morphometric and morphological characters. Thus, we propose here the three later species to be synonyms of *V. gulbenkiana*. The proposal of section *Claviformia* to classify



Figs 1-7. Virescentia crispata (Kumano & Ratnas.) Necchi, D.C.Agostinho & M.L.Vis comb. nov. 1-2. Whorls with carposporophytes (arrows); 3. Detail of a primary fascicle showing audouinelloid cells and curved disposition; 4-6. Details of carpogonia with curved carpogonial branches; 4-5. Young carpogonia; 6. Mature (fertilized) carpogonium with attached spermatium (arrow); 7. Distal cells of gonimoblast □aments with carposporangia. Scale bars: 1-2 = 100 μm; 3-7 = 10 μm. Figs 2-7 redrawn from Ratnasabapathy & Kumano (1982a). Fig. 1 from the isotype (Table 1).


V. azeredoi and V. ferreri (Reis, 1972) has no basis, because the proposed characters (carpogonial branches short or long; carpogonia with club-shaped or cylindrical trichogynes; carposporophytes large, 1-2 per whorl) □ easily within the circumscription of Virescentia.



Figs 8-12. Virescentia gulbenkiana (Reis) Necchi, D.C.Agostinho & M.L.Vis comb. Nov. 8–9. Whorls with carposporophytes (arrows); 10-11. Details of carpogonia; 10. Young carpogonium with shortly stalked trichogyne (arrow); 11. Mature (fertilized) carpogonium with attached spermatium (arrow); 12. Gonimoblast □aments with carposporangia. Scale bars: 250 μm for □gure 8; 100 μm for □gure 9; 10 μm for □gures 10-12. Figure 10 modi□ed from Reis (1967). Figs 8, 12 from the holotype of V. ferreri; Figs 9, 11 from the holotype of V. gulbenkiana (Table 1).

Virescentia helminthosa (Bory) Necchi, D.C.Agostinho & M.L.Vis, comb. nov. Figs 13-17

**Basionym:** Batrachospermum helminthosum Bory, Annales du muséum d'histoire naturelle 12: 316 (1808).



Figs 13-17. Virescentia helminthosa (Bory) Necchi, D.C.Agostinho & M.L.Vis comb. nov. 13. Whorls with carposporophytes (arrows); 14. Detail of a curved young carpogonial branch; 15-16. Details of carpogonia; 15. Curved young carpogonial branch (arrowhead) and a mature carpogonium with stalked trichogyne (arrow); 16. Mature carpogonium with curved carpogonial branch; 17. Gonimoblast □aments with carposporangia. Scale bars: 13 = 250 μm; 14-17 = 10 μm. Fig. 13 from the lectotype of V. helminthosa (Table 1); □gs. 14-17 from BHO A-0936.

**Revised description:** Plants monoecious or polyoecious; branching irregular; whorls contiguous or separated, barrel-shaped, spherical or pear-shaped, 300-800  $\mu$ m wide; primary fascicles straight, with 8-15 cell storeys; secondary fascicles varying from

few and sparse to abundant and covering the entire internode; spermatangia spherical, on primary or secondary fascicles, 5-7 μm wide; carpogonial branches slightly curved, composed of 3-11 disc-shaped cells, developing from the periaxial, proximal or distal cells of primary fascicles or cortical □aments of the main axis, short or long, 12-35 μm long; involucral □aments, short, composed of 1-3 cylindrical, elliptical or spherical cells; carpogonia 40-79 μm long, with stalked, cylindrical trichogynes, sometimes bifurcated or with knobs; carposporophytes 1-2 per whorl, 150-420 μm wide, 100-320 μm high; gonimoblast □aments with 3-7 cylindrical cells; carposporangia obovoid or pear-shaped, 14-20.5 μm long, 8.5-12.5 μm wide. **Distribution:** Europe - Finland, France, Norway, Poland, Portugal, Spain, Sweden (Sirodot, 1884; Israelson, 1942; Reis, 1974; Eloranta *et al.* 2011; Chiasson *et al.*, 2014).

**Representative DNA sequences:** HQ412541, HQ412542 (COI-5P); KJ825955, KM593807, KM593841 (*rbc*L).

Additional specimens examined: France, Dirt road to Chappelle Saint Clair-de-Gout, off of D433, 44.237472, -0.058806, M. Vis, W. Chiasson & K. Chiasson 29.x.2011, BHO A-0936. Remarks: this is the most widely reported species in the genus; however, recent studies based on molecular data (Hanyuda *et al.*, 2004; Rueness, 2010; Agostinho & Necchi, 2014; Chiasson *et al.*, 2014) indicate that it is an European species and all records outside Europe most likely represent other species. All species described by Sirodot (1884) were based on criteria that are currently regarded as having no taxonomic importance (color, sexuality, arrangement of primary fascicles, presence of hairs on fascicles, characters of the asexual 'Chantransia' stage). Thus, those species were placed as synonyms of *V. helminthosa*, reinforcing the taxonomic interpretations of previous authors (Sheath *et al.*, 1994; Vis *et al.*, 2001; Hanyuda *et al.*, 2004).


Virescentia viride-americana Necchi, D.C.Agostinho & M.L.Vis, sp. nov.

Figs 18-22

**Description**: Plants monoecious or dioecious or polyoecious; branching irregular; whorls contiguous or separated, barrel-shaped, spherical or pear-shaped, 280-940 μm side; primary fascicles straight, with 8-19 cell storeys; secondary fascicles varying from few and sparse to abundant and covering the entire internode; spermatangia spherical, on primary or secondary fascicles, 4-8 μm wide; carpogonial straight, composed of 1-7 disc-shaped cells, developing from the periaxial or proximal cells of primary fascicles, short or long, 6-30 μm long; involucral □aments, short, composed of 1-3 cylindrical or elliptical cells; carpogonia (30-)40-77 μm long, with stalked, club-shaped or cylindrical trichogynes; carposporophytes 1-2 per whorl, (130-)140-390 μm wide, 70-200(-240) μm high; gonimoblast □aments with 3-5 cylindrical cells; carposporangia obovoid or pear-shaped, 15-24 μm long, 8-12 μm wide.

Additional specimens examined: USA: Indiana, Otter Creek, M.L. Vis & M.M. Hall, 29.v.1999., BHO A-1177; Massachusetts, Poquoy Brook, M.L. Vis & M.M. Hall, 16.iv.1999, BHO A-1175; Michigan, Flemming Creek, M.L. Vis & W.B. Chiasson, 30.v.2000, BHO A-1473; Ohio, Five Mile Creek, M.L. Vis & R.G. Verb, 11.v.1998, BHO A-1179; Ohio, Sun □sh Creek, M. Vis & D. House, 20.iv.2007, BHO A-0102; Rhode Island, Wood River tributary, M.L. Vis & M.M. Hall, 16.iv.1999, BHO A-1178; Rhode Island, Chipuxet River, no collector, no date, BHO A-1180; Tennessee, Scarboro Creek, M.L. Vis & M.M. Hall, 30.v.1999, BHO A-1176; Virginia, Carvin Creek, M. Vis & D. House, 22.iii.2007, BHO A-0128.

**Holotype:** USA, Big Run, Ohio. 4.2 miles east on Big Run Road (CR8S) from SR 329, 39.353813 N 81.842615 W, M.L. Vis & R.G. Verb, 29.v.1998, BHO A-1174.



Figs 18-22. Virescentia viride-americana Necchi, D.C.Agostinho & M.L.Vis sp. nov. 18. Whorls with carposporophyte (arrow); 19. Detail of an entire carposporophyte; 20-21. Details of carpogonia; 20. Young carpogonium with stalked trichogyne (arrow); 21. Mature carpogonium with stalked trichogyne (arrow); 22. Gonimoblast □aments with carposporangia. Scale bars: 18 = 100 μm; 19 = 25 μm; 20-22 = 10 μm. All □gures from the holotype (Table 1).

Etymology: the species epithet indicates that the alga is a member of the *Virescentia* (from Latin *viride*) from North America (from Latin *americana*).

**Distribution:** North America – eastern, southern and mid-western United States (Sheath *et al.*, 1994; Vis *et al.*, 2001).

**Representative DNA sequences:** EU636733, EU636736, EU073836, EU073839, EU073845, EU073846, EU073847 (COI-5P); AF244110, AF244111, AF244115, AF244117, AF244118 (*rbc*L).

**Remarks:** this species has been previously reported as V. helminthosa (Sheath et al., 1994, and references therein), which is now regarded an exclusively European species. Based on recent studies including molecular data (Hanyuda et al., 2004; Agostinho & Necchi, 2014; Chiasson et al., 2014), the sequences of North American specimens are genetically divergent from the those of Asia, Europe and South America, representing a distinct species, that is here described as new to science. The species is highly divergent from the South American V. viride-brasiliensis in sequence data of rbcL (36-57 bp, 2.9-4.5%) and COI-5P (57-69 bp, 8.6-10.4%). However, these two species are morphologically very similar with considerable overlap for all morphological characters. In addition to the unique rbcL and COI-5P sequences for these species, they have a disjunct geographic pattern that can be applied as an additional criterion. The North American specimens formed a clade with two subgroups in both the rbcL and COI-5P tree as noted above, but the sequence variation within and among these two groups overlapped. The morphology in all taxonomic characters among the populations was similar and there was no geographic disjunction between the two groups. Thus, we have chosen to describe a single species that is somewhat variable in DNA sequence.

Virescentia viride-brasiliensis (Necchi & Agostinho) Necchi, M.L.Vis & D.C.Agostinho, comb. nov. Figs 23-26

**Basionym:** Batrachospermum viride-brasiliense Necchi & Agostinho, Phycologia 53: 566 (2014).

**Distribution:** South America – southeastern and southern Brazil (Espírito Santo, Paraná, Rio Grande do Sul, São Paulo; Agostinho & Necchi, 2014).


**Representative DNA sequences:** KX452188 to KX452196 (COI-5P); KM078031 to KM078042 (*rbc*L).

Revised description: plants monoecious, dioecious or polyoecious; branching irregular; whorls contiguous or separated, barrel-shaped, spherical, pear-shaped or obconic, 280-1100 μm wide; primary fascicles straight, with 7-18 cell storeys; secondary fascicles varying from few and sparse to abundant and covering the entire internode; spermatangia spherical or obovoid, on primary or secondary fascicles, 4-9 μm wide; carpogonial straight, short, composed of 1-7 disc-shaped cells, developing from the periaxial or proximal cells of primary fascicles, short or long, 4-37.5 μm long; involucral □aments, short, composed of 1-3 cylindrical or spherical cells; carpogonia 40-110 μm long, with stalked, club-shaped or sub-cylindrical trichogynes; carposporophytes axial, inserted within the whorls, 1-2 per whorl, 200-550 μm wide and 100-300 μm in height; gonimoblast □aments with 3-6 cells cylindrical or barrel-shaped cells; carposporangia obovoid, pear-shaped or elliptic, 19-35 μm long, 10-24 μm wide.

**Remarks:** this species resembles *V. viride-americana* in all morphological features. However, based on the divergence in sequences of *rbc*L and COI-5P and the disjunct geographic distribution it was proposed a new species by Agostinho & Necchi (2014).


Virescentia vogesiaca (T.G. Schultz ex Skuja) Necchi, D.C.Agostinho & M.L.Vis, comb. nov. Figs 27-32

**Basionym:** Batrachospermum vogesiacum T. G. Schultz ex Skuja, Archiv für Hydrobiologie Suppl 15: 623 (1938).



Figs 23-26. Virescentia viride-brasiliensis (Necchi & D.C. Agostinho) Necchi, D.C.Agostinho & M.L.Vis comb. nov. 23. Whorls with carposporophyte (arrow); 24-25. Details of carpogonia showing long stalk trichogyne (arrow); 26. Gonimoblast □aments with carposporangia. Scale bars: 23 = 250 μm; 24-26 = 10 μm. All □gures from the holotype (Table 1).

**Description:** Plants monoecious; branching irregular; whorls contiguous or separated, barrel-shaped, obconic, pear-shaped or spherical, 340-700 μm wide; primary fascicles straight, with 7-14 cell storeys; secondary fascicles abundant, covering the entire internode; spermatangia spherical, on primary or secondary fascicles, 5-8 μm wide; carpogonial branches straight, composed of (4-)8-20 disc-shaped cells, developing from the periaxial or proximal cells of primary fascicles, long or short, (13-)25-68 μm long; involucral □aments, short, composed of 1-3 elliptical or spherical cells; carpogonia 20-45 μm long, with stalked or shortly stalked, club-shaped or elliptical



Figs 27-32. Virescentia vogesiaca (T.G. Schultz) Necchi, D.C.Agostinho & M.L.Vis comb. nov. 27-28. Whorls with carposporophytes (arrows); 29. Detail of a young carpogonial branch; 30-31. Details of mature carpogonia; 30. Carpogonium with short carpogonial branch and stalked trichogyne (arrow); 31. Carpogonium with long carpogonial branch and unstalked trichogyne; 32. Gonimoblast □aments with carposporangia. Scale bars: 27-28 = 250 μm; 29-32 = 10 μm. Figs. 29, 31 from the lectotype (Table 1); □gs 27-28, 30, 32 from BHO A-0926.

trichogynes; carposporophytes 1-2 per whorl, 140-330  $\mu m$  wide, 95-175  $\mu m$  high; gonimoblast  $\square$ aments with 4-7 cylindrical or elliptical cells; carposporangia obovoid, 13-19  $\mu m$  long, 8.5-13  $\mu m$  wide.

**Distribution:** Europe - Belgium, France, Spain, Sweden (Sirodot, 1884; Israelson, 1942; Eloranta *et al.*, 2011; Chiasson *et al.*, 2014).

**Representative DNA sequences:** KJ825954, KU754497 (*rbc*L).

Additional specimens examined: France, River at St Jacques, 43.929083, -0.731194, M. Vis, W. Chiasson & K. Chiasson 26.x.2011, BHO A-0926.

**Remarks:** we con ☐rmed the distinction of this species from *V. helminthosa* based on genetic divergences among sequences of *rbc*L (121-122 bp, 9.4-9.5%), that corroborates the previous ☐nding by Chiasson *et al.* (2014). In addition, these species can be differentiated by the disposition of the carpogonial branches: slightly curved in *V. helminthosa* and straight in *V. vogesiaca*. This morphological character is consistent with the original description of each species by Sirodot (1884).

#### Excluded taxa

Batrachospermum bakarense Kumano & Ratnas., Japanese journal of phycology 32: 119 (1984)

Based on the characters described in the protologue and observed in the type specimen, i.e. club-shaped and unstalked trichogynes, loosely arranged carposporophytes and two types of gonimoblast  $\square$ aments (prostrate and erect), we concluded that this species belongs to section *Turfosa*, which will be treated in a separate paper.

Batrachospermum desickacharyi Sankaran, Phykos 23: 169 (1984)

Based on the characters described in the protologue and by Kumano (2002), i.e. carpogonia short, carposporophytes stalked and small, we concluded that this species belongs to either *Batrachospermum* sensu stricto or *Sheathia*.

Batrachospermum gombakense Kumano & Ratnas., Japanese journal of phycology 30: 119 (1982b)

Based on the characters described in the protologue and observed in the type specimen, i.e. club-shaped and unstalked trichogynes, loosely arranged carposporophytes and two types of gonimoblast  $\square$ aments (prostrate and erect), we concluded that this species belongs to section *Turfosa*, which will be treated in a separate paper.

Batrachospermum transtaganum Reis, Memórias da sociedade Broteriana 21: 23 (1970)

Based on the characters described in the protologue and observed in the type specimen, we concluded that this species is a synonym of *Kumanoa virgato-decaisneana* as described in Necchi & Vis (2012). Thus, it is here formally transferred to the genus *Kumanoa* as a synonym of *K. virgato-decaisneana*: *Kumanoa transtagana* (Reis) Necchi, D.C. Agostinho & M.L.Vis. *Virescentia* spp.

These taxa were reported as *B. helminthosum* by Hanyuda *et al.* (2004) from Japan and based on genetic data (*rbc*L sequence analysis) and most likely represent two distinct, probably new, species. The authors described we distinct haplotypes, four belonging to a single species and a thone to a separate species. However, according to Hanyuda (pers. comm.) there are no voucher for those specimens required to describe morphological characters to propose new species. The status of these Japanese materials remains unresolved.

Acknowledgments. This study was supported by Foundation for Research Support of São Paulo State - FAPESP (Brazil) grants (2012/12016-6, 2016/07808-1) and National Science Foundation grants DEB 0235676, 0936855, and 1655230. Authors are grateful to the herbarium curators (COI, PC and TNS-AL) to allow access to the type specimens. Latin and nomenclatural advice by Kanchi Natarajan Gandhi, Harvard University, is greatly appreciated.

Some of the research was conducted while MLV was an invited Professor at PC and on a Fulbright Scholarship visiting ON. Help in  $\Box$ eldwork by Ciro Branco, Wayne Chiasson, Kyle Chiasson, Melissa Hall, Denise House, Rogério Krupek, Monica Paiano, Eric Salomaki, and Bob Verb is gratefully acknowledged.

#### REFERENCES

- AGOSTINHO, D.C. & NECCHI, O. Jr., 2014 Systematics of the section *Virescentia* of the genus *Batrachospermum* (Batrachospermales, Rhodophyta) in Brazil. *Phycologia* 53: 561-570.
- BENSON, D. A., CAVANAUGH, M., CLARK, K., OSTELL, J., WHEELER, D.L., 2013 GenBank. Nucleic acids research 41: 36-42.
- BORY DE SAINT-VINCENT J.B.G.M., 1808 Mémoire sur le genre Batrachosperma, de la familie des Conferves. *Annales du muséum d'histoire naturelle* 12: 310-332, pl. 29-31.
- DARRIBA, D., TABOADA, G.L., DOALLO, R. & POSADA, D, 2012 jModelTest 2: more models, new heuristics and parallel computing. *Nature methods* 9: 772.
- ELORANTA P., KWANDRÂNS J. & KUSEL-FETZMANN E., 2011 Rhodophyceae and Phaeophyceae; S Bwasser on Mitteleuropa Band 7. Heidelberg, Spectrum Akademischer Verlag, 155 p.
- ENTWISLE T.J, VIS M.L., CHIASSON W.B., NECCHI O. JR. & SHERWOOD A.R., 2009 Systematics of the Batrachospermales A Synthesis. *Journal of phycology* 45: 704-715.
- ENTWISLE, T.J., JOHNSTON, E.T., LAM, D.W., STEWART, S.A. & VIS, M.L., 2016 *Nocturama* gen. nov., *Nothocladus* s. lat. and other taxonomic novelties resulting from the further resolution of paraphyly in Australasian members of *Batrachospermum* (Batrachospermales, Rhodophyta). *Journal of phycology* 52: 384-396.
- HANYUDA, T., SUZAWA, Y., SUZAWA, T., ARAI, S., SATO, H., UEDA, K. & KUMANO, S., 2004
   Biogeography and taxonomy of Batrachospermum helminthosum (Batrachospermales, Rhodophyta) in Japan inferred from rbcL gene sequences. Journal of phycology 40: 581-588.
- HOUSE, D.L., SHERWOOD, AR. & VIS, M.L., 2008 Comparison of three organelle markers for phylogeographic inference in *Batrachospermum helminthosum* (Batrachospermales, Rhodophyta) from North America. *Phycological research* 56: 69-75.
- ISRAELSON, G., 1942 The freshwater Florideae of Sweden. Symbolae botanicae Upsalienses 6: 1-134.
- KEARSE, M., MOIR, R., WILSON, A., STONES-HAVAS, S., CHEUNG, M., STURROCK, S., BUXTON, S., COOPER, A., MARKOWITZ, S., DURAN, C., THIERER, T., ASHTON, B., MENTJIES, P., & DRUMMOND, A., 2012 Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28: 1647-1649.
- KUMANO S., 2002 Freshwater red algae of the world. Bristol, Biopress Ltd., 375 p.
- KUMANO, S. & RATNASABAPATHY, M., 1984 Studies on freshwater red algae of Malaysia IV. Batrachospermum bakarense, sp. nov. from Sungai Bakar, Kelantan, West Malaysia. Japanese journal of phycology 32: 19-23.
- MORI, M., 1975 Studies of the genus Batrachospermum in Japan. Japanese Journal of Botany 20: 461-484.
- NECCHI, O. Jr., 1989 Rhodophyta de água doce do estado de São Paulo: levantamento taxonômico. Boletim de botânica, Universidade de São Paulo 11: 11-69.
- NECCHI, O. Jr., 1990 Revision of the genus Batrachospermum Roth (Rhodophyta, Batrachospermales) in Brazil. Stuttgart, J. Cramer, [Bibliotheca Phycologica 84] 201 p.
- NECCHI, O. Jr. & VIS, M.L., 2012 Monograph of the genus Kumanoa (Rhodophyta, Batrachospermales. Stuttgart, J. Cramer, [Bibliotheca Phycologica 116] 79 p.
- RATNASABAPATHY, M. & KUMANO, S., 1982a Studies on freshwater red algae of Malaysia I. Some taxa of the genera *Batrachospermum*, *Ballia* and *Caloglossa* from Pulau Tioman, West Malaysia. *Japanese journal of phycology* 30: 15-22.
- RATNASABAPATHY, M. & KUMANO, S., 1982b Studies on freshwater red algae of Malaysia II.

  Three species of *Batrachospermum* from Sungai Gombak and Sungai Pusu, Selangor, West Malaysia. *Japanese journal of phycology* 30: 119-124.
- REIS, M. P., 1965 Batrachospermum gulbenkianum, sp. nov. Anuário da sociedade Broteriana 31: 31-45.

- REIS, M.P., 1967 Duas espécies novas de Batrachospermum Roth: B. azeredoi e B. ferreri. Boletim da sociedade Broteriana 41: 167-189.
- REIS, M.P., 1970 Rhodophyceae novae. Memórias da sociedade Broteriana 21: 23-26.
- REIS, M.P., 1972 Estudo comparativo de Batrachospermum helminthosum, Bory, B. coerulescens Sirod. e B. azeredoi P. Reis e descrição de uma variedade nova. Boletim da sociedade Broteriana 46: 191-217.
- REIS, M.P., 1974 Chaves para a identi □cação das espécies portuguesas de *Batrachospermum* Roth.

  \*\*Anuário da sociedade Broteriana 40: 37-129.
- ROSSIGNOLO, N.L. & NECCHI, O. Jr., 2016 Revision of section Setacea of the genus Batrachospermum (Batrachospermales, Rhodophyta) with emphasis on specimens from Brazil. Phycologia 55: 337-346.
- ROSSIGNOLO, N.L. & NECCHI, O. Jr. & GUIRY, M.D., 2016 Atrophycus, a new genus name for "Setacea (De Toni) Necchi & Rossignolo". Notulae algarum 26: 1-2.
- RUENESS, J., 2010 DNA barcoding of select freshwater and marine red algae (Rhodophyta). Cryptogamie, Algologie 31: 377-386.
- SALOMAKI, E.D., KWANDRANS, J., ELORANTA, P. & VIS, M.L., 2014 Molecular and morphological evidence for *Sheathia* gen. nov. (Batrachospermales, Rhodophyta) and three new species. *Journal of phycology* 50: 531-547.
- SANKARAN, V., 1984 Batrachospermum desikacharyi sp. nov. (Rhodophyta) from Valparai, Anamalais, Tamil Nadu. Phykos 23: 163-170.
- SHEATH, R.G., VIS, M.L. & COLE, K.M., 1994 Distribution and systematics of Batrachospermum (Batrachospermales, Rhodophyta) in North America. 4. Section Virescentia. Journal of phycology 30: 108-17.
- SHERWOOD, A.R., VIS, M.L., ENTWISLE, T.J., NECCHI, O. Jr. & PRESTING, G.G., 2008 Contrasting intra versus interspecies DNA sequence variation for representatives of the Batrachospermales (Rhodophyta): Insights from a DNA barcoding approach. *Phycological research* 56: 269-279.
- SILVESTRO, D. & MICHALAK, I, 2011 RAXML GUI: a graphical front-end for RAXML. Organism diversity and evolution 12: 335-337.
- SIRODOT S., 1873 Nouvelle classi □cation des algues d'eau douce du genre Batrachospermum: développement; générations alternates. Comptes-rendus hebdomadaires des séances de l'Académie des Sciences 76: 1216-1120.
- SIRODOT S., 1884 Les Batrachospermes: Organisation, Fonctions, Développement, Classi Cation. Paris, Librairie de l'Académie de Médecine, G. Masson, 299 p., 50 pl.
- SKUJA, H., 1938 Die Süsswasser rhodophyceen der Deutschen Limnologischen Sunda-Expedition. Archiv für Hydrobiologie 15: 603-637.
- STARMACH, K., 1977 Phaeophyta Brunatnice, Rhodophyta Krasnorosty. Flora słodkowodna Polski 14. Warszawa, Panstwowe Wydawnictwo Naukowe. 445 p.
- THIERS, B., 2018 (continuously updated) Index Herbariorum: a global directory of public herbaria and associated staff: New York Botanical Garden's Virtual Herbarium. http://sweetgum.nybg.org/ih/; searched on (February 14, 2018).
- VIS, M.L., SAUNDERS, G.W., SHEATH, R.G., DUNSE, K. & ENTWISLE, T.J., 1998 Phylogeny of the Batrachospermales (Rhodophyta) as inferred from rbcL and 18S ribosomal RNA gene DNA sequences. Journal of phycology 34: 341-350.
- VIS, M.L., MILLER, E.J. & HALL, M.M., 2001 Biogeographic analyses of Batrachospermum helminthosum (Batrachospermales, Rhodophyta) in North America using molecular and morphological data. Phycologia 40: 2-9.

## SUPPORTING INFORMATION

Table S1. rbcL sequence data from additional taxa of Batrachospermales used for phylogenetic analyses.

Taxon	GenBank Accession Number
Atrophycus ater	KT802842
Atrophycus puiggarianus	FJ386462
Balliopsis prieurii	AY960688
Batrachospermum brasiliense	FJ386458
Batrachospermum cayennense	KM055245
Batrachospermum cayennense	AY423393
Batrachospermum dapsile	KM593855
Batrachospermum gelatinosum	GU810833
Batrachospermum macrosporum	EU106060
Batrachospermum macrosporum	EU106056
Batrachospermum naiadis	KM593857
Batrachospermum pozoazulense	KM593863
Batrachospermum shanxiense	KM593851
Batrachospermum turfosum	KJ825960
Kumanoa ambigua	AY423390
Kumanoa americana	JN589995
Kumanoa australica	JN589998
Kumanoa gudjewga	JN590002
Kumanoa holtonii	JN590004
Kumanoa louisianae	JN590005
Kumanoa nodi⊡ora	AY423398
Китапоа ргосагра	FJ386464
Kumanoa vittata	JN590010
Lemanea 🗆 wiatilis	KM055243
Lemanea fucina	KJ825958
Lympha mucosa	KM593865
Nemalionopsis shawii	KM005141
Nocturama antipodites	FJ386456
Nothocladus campyloclonus	FJ386459
Nothocladus diatyches	KT802848
Nothocladus discors	KT802850
Nothocladus nodosus	AF029152
Nothocladus theaquus	KT802863
Nothocladus wattsii	KT802866
Paralemanea sp.	GQ285124
Petrohua bernabei	AY960690

**Table S1**. *rbc*L sequence data from additional taxa of Batrachospermales used for phylogenetic analyses *(continued)*.

Taxon	GenBank Accession Number
Psilosiphon scoparium	AF029155
Sheathia americana	JX669757
Sheathia confusa	DQ393133
Sheathia exigua	GU457345
Sheathia heterocortica	DQ393136
Sheathia involuta	KU672395
Sirodotia delicatula	DQ646476
Sirodotia huillensis	AF029157
Sirodotia suecica	JF344718
Thorea riekei	KM005140
Tuomeya americana	KM055244
Volatus carrionii	KM593864
Volatus personatus	KM593856
Volatus ulterior	KM593852

**Table S2.** Sequences of *rbc*L and COI-5P of *Batrachospermum* section *Virescentia* from GenBank used in the analyses.

Canaina	Location	GenBank Acession Number		D - f
Species		rbcL	COI-5P*	— Reference
Batrachospermum sp.	U.S.A., Michigan, Spring Brook	AF244109	EU636727 EU073847	Vis et al. (2001)
Batrachospermum sp.	U.S.A., Michigan, Flemming Creek	AF244110	EU636727 EU636736 EU073847	Vis et al. (2001)
Batrachospermum sp.	U.S.A., Ohio, Big Run Creek	AF244111	EU636733 EU073833 EU073838	Vis et al. (2001)
Batrachospermum sp.	U.S.A., Ohio, Browning Creek	AF244112	EU073834 EU073837 EU073848 EU636732 EU636734	Vis et al. (2001)
Batrachospermum sp.	U.S.A., Connecticut, Mill Brook	AF244113	EU073834 EU636732	Vis et al. (2001)
Batrachospermum sp.	U.S.A., Louisiana, unnamed stream	AF244114	EU636729 EU073845	Vis et al. (2001)

<sup>\*</sup> the COI-5P was sequenced from multiple specimens within a location such that there may be multiple sequences per location and the same COI-5P haplotype can occur in more than one location. These data are from either Sherwood *et al.* (2008) or House *et al.* (2008).

	Location	GenBank Acession Number		Deference.
Species		rbcL	COI-5P*	— Reference
Batrachospermum sp.	U.S.A., Massachusetts, Poquoy Brook	AF244115	EU073839 EU636735	Vis et al. (2001)
Batrachospermum sp.	U.S.A., North Carolina, Cedar Fork	AF244116	EU073834 EU073836 EU073838 EU073840	Vis et al. (2001)
Batrachospermum sp.	U.S.A., Tennessee, Scarboro Creek	AF244117	EU073842 EU073843 EU073844 EU073846 EU636728 EU636730 EU636731	Vis et al. (2001)
Batrachospermum sp.	U.S.A., Indiana, Otter Creek	AF244118	EU073845 EU636729	Vis et al. (2001)
Batrachospermum sp.	U.S.A., Rhode Island, Wood River tributary	AF244119	EU073833 EU073836	Vis et al. (2001)
Batrachospermum sp.	U.S.A., Ohio, Five Mile Creek	AF244120	EU073834 EU073838 EU636736	Vis et al. (2001)
Batrachospermum sp.	U.S.A., Rhode Island, Chipuxet River	AF029142		Vis et al. (1998)
Batrachospermum sp.	U.S.A., Ohio, Yellow Springs	AY198417	EU073847 EU073848 EU636727	Chiasson et al. (2003)
Batrachospermum sp.	U.S.A., Ohio, Big Run		EU073833	Vis et al. (2001)
Batrachospermum sp.	U.S.A., Virgina, Carvin Creek		EU073833 EU073835 EU073840 EU073843	
Batrachospermum sp.	U.S.A., Ohio, Sun⊑sh Creek		EU073841	
Batrachospermum sp.	Japan, Kagoshima, Oguchi	AB114642		Hanyuda et al. (2004)
Batrachospermum sp.	Japan, Kyoto, Tanba	AB114643		Hanyuda <i>et al</i> . (2004)
Batrachospermum sp.	Japan, Ehime, Komatsu	AB114644		Hanyuda <i>et al</i> . (2004)
Batrachospermum sp.	Japan, Iwate, Takizawa	AB114645		Hanyuda <i>et al</i> . (2004)

<sup>\*</sup> the COI-5P was sequenced from multiple specimens within a location such that there may be multiple sequences per location and the same COI-5P haplotype can occur in more than one location. These data are from either Sherwood et al. (2008) or House et al. (2008).

**Table S2.** Sequences of *rbc*L and COI-5P of *Batrachospermum* section *Virescentia* from GenBank used in the analyses *(continued)*.

Species	Location	GenBank Acession Number		Dafanan aa
		rbcL	COI-5P*	— Reference
Batrachospermum sp.	Japan, Okinawa, Nago	AB114646		Hanyuda <i>et al</i> . (2004)
B. helminthosum	Norway, Askerelva, county of Akershus		HQ412541	Rueness (2010)
B. helminthosum	Norway, Askerelva, county of Akershus		HQ412542	Rueness (2010)
B. helminthosum	France, Le Boulet Prioult, near Le Petit Boulet	KJ825955		Chiasson <i>et al</i> . (2014)
B. helminthosum	Portugal, Portalegere, Termas de Nisa	KM593807		Chapuis <i>et al</i> . unpublished
B. helminthosum	Spain, Galicia, A Coruña, Riachuelo das Brañas	KM593841		Chapuis <i>et al</i> . unpublished
B. viride-brasiliense	Brasil, SP, Maringá, Pavão River	FJ386460		Entwisle <i>et al</i> . (2009)
B. viride-brasiliense	SP: Campos do Jordão, Ducha da Prata	KM259993	KM078048	Agostinho & Necchi (2014)
B. viride-brasiliense	SP: Paraibuna, Pousada Alto da Serra, Rio Negro	KM259994	KM097031	Agostinho & Necchi (2014)
B. viride-brasiliense	PR: Guarapuava, Rio das Pedras	KM259996	KM097033	Agostinho & Necchi (2014)
B. viride-brasiliense	PR: Guarapuava, Rio Lageado, Bacia do Rio Marrecas	KM259995	KM097032	Agostinho & Necchi (2014)
B. viride-brasiliense	ES: Domingos Martins, Pedra Azul, Rio Tucú	KM259998	KM097035	Agostinho & Necchi (2014)
B. viride-brasiliense	PR: Teixeira Soares, Floresta Nacional de Irati	KM259999	KM097036	Agostinho & Necchi (2014)
B. viride-brasiliense	PR: Teixeira Soares, Floresta Nacional de Irati	KM260000	KM097037	Agostinho & Necchi (2014)
B. viride-brasiliense	SC: São Domingos, Parque Estadual das Araucárias, rio Jacutinga	KM260001	KM097038	Agostinho & Necchi (2014)

<sup>\*</sup> the COI-5P was sequenced from multiple specimens within a location such that there may be multiple sequences per location and the same COI-5P haplotype can occur in more than one location. These data are from either Sherwood *et al.* (2008) or House *et al.* (2008).

a ·	Location	GenBank Acession Number		D (
Species		rbcL	COI-5P*	— Reference
B. viride-brasiliense	SC: São Domingos, Parque Estadual das Araucárias	KM259997	KM097034	Agostinho & Necchi (2014)
B. viride-brasiliense	SC: São Domingos, Parque Estadual das Araucárias	KM260002	KM097039	Agostinho & Necchi (2014)
B. viride-brasiliense	SP: Brotas, Parque dos Saltos	KM260003	KM097040	Agostinho & Necchi (2014)
B. viride-brasiliense	RS: Cambará do Sul, Parque Nacional de Aparados da Serra, Arroio Camisas	KM260004	KM097041	Agostinho & Necchi (2014)
B. viride-brasiliense	RS: Cambará do Sul, Parque Aparados da Serra, Arroio Perdizes	KM260005	KM097042	Agostinho & Necchi (2014)
B. vogesiacum	France, river at St. Jacques	KJ825954		Chiasson <i>et al.</i> (2014)
B. vogesiacum	Spain, Zamora, Río Tera	KU754497		Chapuis <i>et al</i> . unpublished

<sup>\*</sup> the COI-5P was sequenced from multiple specimens within a location such that there may be multiple sequences per location and the same COI-5P haplotype can occur in more than one location. These data are from either Sherwood *et al.* (2008) or House *et al.* (2008).




Fig. S1. Phylogenetic tree (maximum likelihood) showing relationship of *Virescentia* stat. nov. with other genera and sections of Batrachospermales based on rbcL sequence data. \* = bootstrap support > 80 and posterior probability > 0.90; nodes without values indicate bootstrap values  $\leq$  80 and posterior probability  $\leq$  0.90. Information on the sequences are listed in Table S1 (Supporting Information). Scale represents substitutions per site.




Fig. S2. Phylogenetic tree (maximum likelihood) based on rbcL sequences. \* = bootstrap values > 90 and posterior probability > 0.90; nodes without values indicate bootstrap values 70 and posterior probability 0.70. Information on the sequences are listed in Table S2 (Supporting Information). Scale represents substitutions per site.




Fig. S3. Distance tree (neighbor-joining) based on COI-5P sequences. \* = bootstrap values > 95; nodes without values indicate bootstrap values  $\leq$  70. Information on the sequences are listed in Table S2 (Supporting Information). Scale represents substitutions per site.