

Asymptotically compatible discretization of multidimensional nonlocal diffusion models and approximation of nonlocal Green's functions

QIANG DU, YUNZHE TAO AND XIAOCHUAN TIAN

Department of Applied Physics and Applied Mathematics, Columbia University, New York, NY 10027

AND

JIANG YANG*

*Department Mathematics, Southern University of Science and Technology, Shenzhen, 518055, China
and Department of Applied Physics and Applied Mathematics, Columbia University,
New York, NY 10027*

*Corresponding author: yangj7@sustc.edu.cn

[Received on 2 September 2017; revised on 17 January 2018]

Nonlocal diffusion equations and their numerical approximations have attracted much attention in the literature as nonlocal modeling becomes popular in various applications. This paper continues the study of robust discretization schemes for the numerical solution of nonlocal models. In particular, we present quadrature-based finite difference approximations of some linear nonlocal diffusion equations in multidimensions. These approximations are able to preserve various nice properties of the nonlocal continuum models such as the maximum principle and they are shown to be asymptotically compatible in the sense that as the nonlocality vanishes, the numerical solutions can give consistent local limits. The approximation errors are proved to be of optimal order in both nonlocal and asymptotically local settings. The numerical schemes involve a unique design of quadrature weights that reflect the multidimensional nature and require technical estimates on nonconventional divided differences for their numerical analysis. We also study numerical approximations of nonlocal Green's functions associated with nonlocal models. Unlike their local counterparts, nonlocal Green's functions might become singular measures that are not well defined pointwise. We demonstrate how to combine a splitting technique with the asymptotically compatible schemes to provide effective numerical approximations of these singular measures.

Keywords: nonlocal models; nonlocal diffusion; peridynamics; nonlocal gradient; asymptotic compatibility; quadrature collocation approximations; nonlocal Green's function.

1. Introduction

In this paper, we study numerical approximations of the following multidimensional linear nonlocal diffusion equation:

$$-\mathcal{L}_\delta u^\delta(\mathbf{x}) = f(\mathbf{x}), \quad \mathbf{x} \in \Omega \subset \mathbb{R}^d, \quad (1.1)$$

with Ω being a given domain in $\mathbb{R}^d$ ($d \geq 2$), f a given right-hand side, u^δ the solution to be sought and the nonlocal operator $\mathcal{L}_\delta$ defined by

$$\mathcal{L}_\delta u(\mathbf{x}) = 2 \int_{B_\delta(\mathbf{0})} \rho_\delta(|\mathbf{z}|) (u(\mathbf{x} + \mathbf{z}) - u(\mathbf{x})) \, d\mathbf{z}. \quad (1.2)$$

The operator $\mathcal{L}_\delta$ is parametrized by a positive horizon parameter δ measuring the range of nonlocal interactions. The specific form of such nonlocal interactions is prescribed by a non-negative kernel function $\rho_\delta = \rho_\delta(|z|)$. The integral in (1.2) is interpreted in the principal value sense whenever needed (Mengesha & Du, 2014). More discussions on the nonlocal model are given in Section 2. A main task of this work is to develop quadrature-based finite difference approximation schemes and associated Green's functions for (1.1). The approximation schemes are able to preserve the maximum principle at the discrete level and they are shown to be asymptotically compatible in the sense that as the nonlocality vanishes, the numerical solutions can give consistent local limits. They are also shown to be of optimal order in both nonlocal and asymptotically local settings. Previous studies of schemes enjoying such optimal-order error estimates, discrete maximum principles and asymptotical compatibility have been mostly confined to the one-dimensional space (Tian & Du, 2013). While the algorithmic development is in a similar spirit to that for the one-dimensional case, the multidimensional extension made in this work involves both new design elements that reflect the multidimensional nature and more technical convergence analysis relying on new interpolation error estimates of nonconventionally defined weighted divided differences.

The motivation for our work is rooted in the need to develop robust approximations of nonlocal models that serve as alternatives to classical partial differential equations (PDEs). Recently, there have been many studies on the application of nonlocal modeling to problems in mechanics, physics and materials, biological and social sciences (Bates & Chmaj, 1999; Applebaum, 2004; Gilboa & Osher, 2008; Bobaru & Duangpanya, 2010; Buades *et al.*, 2010; Lou *et al.*, 2010; Tadmor & Tan, 2014). The integral formulations of spatial interactions in nonlocal models can account for nonlocal effects and allow more singular solutions. An example is the nonlocal peridynamics (PD) theory (Silling, 2000) and its applications in studying cracks and materials failure, as well as other mechanical properties and physical processes (Askari *et al.*, 2008; Silling & Lehoucq, 2010; Silling *et al.*, 2010a; Palatucci *et al.*, 2012). Rigorous mathematics of nonlocal models has proved to be necessary in order to gain fundamental insights and to guide the modeling and simulation efforts (Silling & Askari, 2005; Macek & Silling, 2007; Bobaru *et al.*, 2009; Andreu *et al.*, 2010; Kilic & Madenci, 2010; Zhou & Du, 2010; Chen & Gunzburger, 2011; Du & Zhou, 2011; Du *et al.*, 2012, 2013, 2017b).

For nonlocal models like (1.1), as $\delta \rightarrow 0$, the nonlocal interactions that define the model can become localized so that the zero-horizon (or local) limit of the nonlocal operator, when valid both physically and mathematically, can be represented by a local differential operator. Given such a scenario the corresponding nonlocal model naturally converges to a conventional differential equation model in the local limit; see for instance, the convergence analysis of linear state-based PD models to the classical Navier equation of linear elasticity (Mengesha & Du, 2014). There have been concerns about whether such consistency can be preserved at the discrete level when the nonlocal models are discretized numerically. Discrete schemes of nonlocal models that preserve the correct limiting behavior are called asymptotically compatible (AC) schemes, a notion developed first in the studies by Tian & Du (2013, 2014). In other words, the numerical approximation given by an AC scheme can reproduce the correct local limiting solution as the horizon δ and the mesh spacing (denoted by h) approach zero, if the convergence to such a limiting local solution is valid on the continuum level. While Tian & Du (2013) presented results concerning the AC property for a one-dimensional scalar model solved by a number of different discretization methods, Tian & Du (2014) managed to develop more general results that are applicable to linear systems in multidimensions based on Galerkin finite element approximations on unstructured meshes. A few subsequent studies have been carried out. For example, Tao *et al.* (2017) extended the study of AC schemes to nonlocal diffusion equations with Neumann-type volume constraints, Du & Yang (2016) discussed AC schemes based on Fourier spectral methods for problems

defined on periodic cells and [Du & Yang \(2017\)](#) proposed an efficient and accurate hybrid algorithm to implement the Fourier spectral methods, while [Du *et al.* \(2017b\)](#) developed discontinuous-Galerkin-based AC schemes. Moreover, AC schemes were proposed in the study by [Chen *et al.* \(2017\)](#) for nonlocal time–space models and in the study by [Du *et al.* \(2016\)](#) for the robust recovery of the nonlocal gradient of the solutions to nonlocal models. For technical reasons, much of the numerical analysis in these works, with the exception of the study by [Tian & Du \(2014\)](#) on conforming Galerkin finite element approximations, has focused on nonlocal operators defined for a one-dimensional spatial variable. In comparison with other AC schemes such as those based on the Galerkin finite element and spectral methods, quadrature-based collocation-type finite difference schemes not only offer optimal-order convergent approximations with simpler implementations but also preserve the discrete maximum principle and asymptotic compatibility; see details in Section 3. Furthermore, these quadrature-based schemes are closely related to other discretizations based on strong forms, such as the original mesh-free discretization developed in the study by [Silling \(2000\)](#).

As an application of the discretization scheme, we consider in Section 4 the numerical computation of nonlocal Green's functions $G_\delta = G_\delta(\mathbf{x}, \mathbf{y})$ for the nonlocal diffusion model (1.1). Nonlocal Green's functions are important tools for nonlocal continuum models ([Weckner *et al.*, 2009](#); [Silling *et al.*, 2010b](#); [Wang *et al.*, 2016, 2017](#)), just like their local versions, i.e., conventional Green's functions associated with PDEs. Since analytical expressions can be expected only in simple cases, the numerical study of nonlocal Green's functions has particular significance. We show here that while the local/conventional and nonlocal Green's functions enjoy many similarities and are intimately connected through the local limiting process, there are also some fundamental differences among them. In particular, for a large class of nonlocal interaction kernels the associated Green's functions $G_\delta = G_\delta(\mathbf{x}, \mathbf{y})$ on the continuum level for $\delta > 0$ are measure-valued distributions, which cannot be interpreted as conventional functions defined pointwise for $\mathbf{x}$ and $\mathbf{y}$. These differences are essential features that, if improperly handled, could present undesirable effects on their numerical approximations. This serves as another reminder of the subtlety involved in the nonlocal modeling. To better deal with these complications we discuss a splitting approach that utilizes the developed AC scheme to get effective and robust approximations of nonlocal Green's functions.

2. Multidimensional nonlocal diffusion model

For the sake of a clear illustration we let $u = u(\mathbf{x})$ denote a scalar deformation field and consider (1.1) subject to a nonlocal constraint of the Dirichlet type,

$$u = 0 \quad \text{on } \Omega_{\mathcal{I}}, \quad (2.1)$$

where $\Omega_{\mathcal{I}}$ is a boundary layer defined as

$$\Omega_{\mathcal{I}} = \{\mathbf{x} \notin \Omega \mid \text{dist}(\mathbf{x}, \Omega) \leq \delta\}.$$

For more discussions on nonlocal constraints defined on a domain with a nonzero volume we refer to [Du *et al.* \(2012, 2013\)](#).

Furthermore, let us consider suitably scaled kernels so that as δ goes to zero, the local limit of the nonlocal operator $\mathcal{L}_\delta$ is exactly the Laplace operator Δ (which is denoted by $\mathcal{L}_0$). To this end, it

requires that the kernel is a non-negative and nonincreasing function and has a finite second moment in $\mathbf{z}$ independent of the horizon parameter δ :

$$\int_{\mathcal{B}_\delta(\mathbf{0})} |\mathbf{z}|^2 \rho_\delta(|\mathbf{z}|) \, d\mathbf{z} = d.$$

The limiting local model of (1.1) is then given by

$$-\mathcal{L}_0 u^0 = f \quad \text{in } \Omega \quad \text{and} \quad u^0 = 0 \quad \text{on } \partial\Omega, \quad (2.2)$$

where, without loss of generality, $\mathcal{L}_0 = \Delta$ for the particular case studied here.

We note that a popular choice of ρ_δ is a rescaled kernel given by

$$\rho_\delta(|\mathbf{z}|) = \frac{1}{\delta^{2+d}} \rho\left(\frac{|\mathbf{z}|}{\delta}\right) \quad \forall \mathbf{z} \in \mathcal{B}_\delta(\mathbf{0}), \quad (2.3)$$

where $\rho = \rho(\xi)$ is a non-negative and nonincreasing function with compact support in $[0, 1]$ and a normalized moment

$$\int_0^1 \rho(\xi) \xi^{1+d} \, d\xi = c_d \quad (2.4)$$

for a given constant c_d , though our discussion here is not restricted to such a rescaled form.

3. Quadrature-based finite difference schemes and asymptotic compatibility

AC schemes provide robust numerical approximations of nonlocal models (Tian & Du, 2013, 2014) since the convergence of such schemes is insensitive to the choices of modeling and discretization parameters, in particular with respect to either a sufficiently small horizon δ or a sufficiently refined mesh spacing (denoted by h).

3.1 The discretization scheme

We now develop an AC quadrature-based finite difference discretization to the nonlocal diffusion equation (1.1) that extends a similar scheme presented first in the study by Tian & Du (2014) and also studied by Du & Tian (2014) for the one-dimensional case.

For simplicity, we let $\{\mathbf{x}_j \in \Omega \cup \Omega_T\}$ be the set of nodes (grid points) of a uniform Cartesian mesh $\mathcal{T}_h$ with mesh size h . Here, $\mathbf{j}$ denotes a multiindex corresponding to $\mathbf{x}_j = h\mathbf{j}$. We note that the particular ordering of the nodes affects the matrix structure corresponding to the discrete system but does not affect the numerical solution. First, at any node $\mathbf{x}_i \in \Omega$, the nonlocal operator can be rewritten as

$$\mathcal{L}_\delta u(\mathbf{x}_i) = 2 \int_{\mathcal{B}_\delta(\mathbf{0})} \frac{u(\mathbf{z} + \mathbf{x}_i) - u(\mathbf{x}_i)}{W(\mathbf{z})} W(\mathbf{z}) \rho_\delta(|\mathbf{z}|) \, d\mathbf{z}, \quad (3.1)$$

where $W(\mathbf{z})$ represents a weight function. Then a quadrature-based finite difference scheme of the nonlocal operator (1.2) can be given as

$$\mathcal{L}_{h,\delta} u(\mathbf{x}_i) = 2 \int_{\mathcal{B}_\delta(\mathbf{0})} \mathcal{I}_h \left(\frac{u(\mathbf{z} + \mathbf{x}_i) - u(\mathbf{x}_i)}{W(\mathbf{z})} \right) W(\mathbf{z}) \rho_\delta(|\mathbf{z}|) d\mathbf{z}, \quad (3.2)$$

where $\mathcal{I}_h(\cdot)$ represents the piecewise d -multilinear interpolation operator in $\mathbf{z}$ associated with the mesh $\mathcal{T}_h$, that is, $\mathcal{I}_h(u)$ is piecewise linear with respect to each component of the spatial variable. The nonlocal constraint with $u = 0$ is imposed at nodes in $\Omega_{\mathcal{T}}$. For the one-dimensional case, in the study by [Tian & Du \(2014\)](#) the function $W(z) = |z|$ was used for $z \in \mathbb{R}$. A key ingredient in its extension to the high-dimensional case is given by

$$W(\mathbf{z}) = \frac{|\mathbf{z}|^2}{|\mathbf{z}|_1}, \quad (3.3)$$

where the notation $|\cdot|_1$ stands for the ℓ_1 norm in the d -dimensional vector space while $|\cdot|$ denotes the standard Euclidean norm.

The auxiliary function $W = W(\mathbf{z})$ plays an important role in ensuring asymptotic compatibility for the multidimensional case. The resulting quadrature-based finite difference scheme for solving the nonlocal diffusion problem can be more conveniently written as the following: find $\{u_h^\delta(\mathbf{x}_i)\}$ such that $u_h^\delta(\mathbf{x}_i) = 0$ for $\mathbf{x}_i \in \Omega_{\mathcal{T}}$ and

$$-\mathcal{L}_{h,\delta} u_h^\delta(\mathbf{x}_i) = \sum_{\mathbf{x}_j \in \mathcal{B}_\delta(\mathbf{x}_i)} \left(u_h^\delta(\mathbf{x}_j) - u_h^\delta(\mathbf{x}_i) \right) p_{\mathbf{x}_i, \mathbf{x}_j}^\delta = f(\mathbf{x}_i) \quad \forall \mathbf{x}_i \in \Omega, \quad (3.4)$$

where the sum over $\mathbf{x}_j$ is for grid points in $\mathcal{B}_\delta(\mathbf{x}_i)$ but not including $\mathbf{x}_i$, $p_{\mathbf{x}_i, \mathbf{x}_j}^\delta = 2\beta_{\mathbf{j}\mathbf{i}}/W(\mathbf{x}_j - \mathbf{x}_i)$ with

$$\beta_{\mathbf{j}\mathbf{i}} = \int_{\mathcal{B}_\delta(\mathbf{0})} \phi_{\mathbf{j}}(\mathbf{z} + \mathbf{x}_i) W(\mathbf{z}) \rho_\delta(|\mathbf{z}|) d\mathbf{z}, \quad (3.5)$$

and $\phi_{\mathbf{j}}$ is the piecewise multilinear basis function satisfying $\phi_{\mathbf{j}}(\mathbf{x}_i) = 0$ when $\mathbf{i} \neq \mathbf{j}$ and $\phi_{\mathbf{j}}(\mathbf{x}_j) = 1$.

REMARK 3.1 The discrete scheme (3.4) is valid for general kernels $\rho_\delta = \rho_\delta(\mathbf{z})$ with bounded second moment in $\mathbf{z}$. This is another benefit of using the weight $W = W(\mathbf{z})$ in (3.3). Indeed, consider a kernel $\rho_\delta(|\mathbf{z}|) = c_{\delta,\alpha,d} |\mathbf{z}|^{-\alpha}$ with $\alpha \in [d, d+2)$ for $\mathbf{z} \in \mathcal{B}_\delta(\mathbf{0})$; then the integral in (3.5) may be unbounded for $|\mathbf{i} - \mathbf{j}| = 1$ if we do not use such a weight, or say, if $W(\mathbf{z}) = 1$ is taken instead.

REMARK 3.2 By the symmetry and translation invariance of the uniform mesh it can be observed that $\beta_{\mathbf{j}\mathbf{i}} = \beta_\delta(\mathbf{x}_i - \mathbf{x}_j)$ for some function β_δ . In fact, let ϕ_0 be the piecewise multilinear basis function located at the origin satisfying $\phi_0(\mathbf{x}_i) = 0$ with $\mathbf{x}_i \neq \mathbf{0}$ and $\phi_0(0) = 1$; then

$$\beta_{\mathbf{j}\mathbf{i}} = \beta_\delta(\mathbf{x}_i - \mathbf{x}_j), \quad \beta_\delta(\mathbf{x}) = \int_{\mathcal{B}_\delta(\mathbf{0})} \phi_0(\mathbf{z} + \mathbf{x}) W(\mathbf{z}) \rho_\delta(|\mathbf{z}|) d\mathbf{z}. \quad (3.6)$$

Moreover, it is easy to see that β_δ is symmetric with respect to any axis hyperplane. Also, $\beta_{\mathbf{j}\mathbf{i}} = 0$ if the support of $\phi_{\mathbf{j}}$ does not overlap with the ball $\mathcal{B}_\delta(\mathbf{x}_i)$. In comparison, we give the conventional central

finite difference scheme for the local diffusion equation (2.2):

$$-\mathcal{L}_{h,0}u_h^0(\mathbf{x}_i) = - \sum_{|\mathbf{x}_j - \mathbf{x}_i| = h} \left(u_h^0(\mathbf{x}_j) - u_h^0(\mathbf{x}_i) \right) \frac{1}{|\mathbf{x}_j - \mathbf{x}_i|^2} = f(\mathbf{x}_i) \quad \forall \mathbf{x}_i \in \Omega, \quad (3.7)$$

that is, it involves only the nearest neighbors along the axes.

3.2 Convergence analysis

Due to the symmetry of the kernel we may rewrite the nonlocal operator as

$$\mathcal{L}_\delta u(\mathbf{x}) = \int_{\mathcal{B}_\delta(\mathbf{0})} \rho_\delta(|\mathbf{z}|) (u(\mathbf{x} + \mathbf{z}) + u(\mathbf{x} - \mathbf{z}) - 2u(\mathbf{x})) \, d\mathbf{z}$$

and the discrete operator as

$$\mathcal{L}_{h,\delta} u(\mathbf{x}_i) = \sum_{\mathbf{j} \neq \mathbf{0}} \frac{u(\mathbf{x}_i + \mathbf{z}_j) + u(\mathbf{x}_i - \mathbf{z}_j) - 2u(\mathbf{x}_i)}{|\mathbf{z}_j|^2} |\mathbf{z}_j|_1 \beta_\delta(\mathbf{z}_j).$$

We first state a few technical lemmas concerning the nonlocal operators and the nonlocal discrete schemes that are nonlocal analogs of their well-known local versions.

First, by the fact that $W = W(\mathbf{z})$, the kernel ρ_δ , as well as the multilinear basis functions ϕ_j , is non-negative, we see that β_δ , $\beta_{j\mathbf{i}}$ and thus $p_{\mathbf{x}_i\mathbf{x}_i}^\delta$ are non-negative. This implies the M-matrix property of the coefficient matrix, just like the case corresponding to the central difference approximations of the diffusion operator.

LEMMA 3.3 (M-matrix). The coefficient matrix, also denoted by $\mathcal{L}_{h,\delta}$, corresponding to the linear system (3.4) is an M-matrix.

An important stability property can be derived via the discrete maximum principle that follows from the M-matrix property of the coefficient matrix. By constructing a suitable quadratic barrier function similar to the one-dimensional case by [Tao et al. \(2017\)](#), as well as the local counterpart, we can get the following lemma.

LEMMA 3.4 (Uniform stability). For any $\delta > 0$, $\mathcal{L}_{h,\delta}^{-1}$ is bounded. Furthermore, for any $0 < \delta < \delta_0$, we have

$$\left\| \mathcal{L}_{h,\delta}^{-1} \right\|_\infty \leq C(\delta_0) \quad (3.8)$$

for some constant $C(\delta_0)$ independent of h and δ as $h \rightarrow 0$.

Next we consider the uniform consistency derived via the truncation error calculation. Here the uniform consistency means that the truncation error is independent of δ for small δ . We first state the following useful and vital lemma, which is a nonlocal analog of the fact that the central difference approximation to the Laplacian, as given by (3.7), is exact for quadratic polynomials.

LEMMA 3.5 (Quadratic exactness). For any quadratic polynomial in $\mathbb{R}^d$ given as $u(\mathbf{x}) = \mathbf{x} \otimes \mathbf{x} : M$, where $M = (m_{kj})$ is a constant matrix, we have

$$\mathcal{L}_{h,\delta} u(\mathbf{x}_i) = \mathcal{L}_\delta u(\mathbf{x}_i) = \sum_k m_{kk} \quad \forall \mathbf{i}. \quad (3.9)$$

Proof. The fact that

$$\int_{B_\delta(\mathbf{0})} \mathbf{z} \otimes \mathbf{z} \rho_\delta(|\mathbf{z}|) d\mathbf{z} = I_d,$$

where I_d is the $d \times d$ identity matrix, yields

$$\mathcal{L}_\delta u(\mathbf{x}_i) \equiv \sum_k m_{kk}.$$

Meanwhile, when $u(\mathbf{x}) = \mathbf{x} \otimes \mathbf{x} : M$, it is obvious that

$$u(\mathbf{x}_i + \mathbf{z}_j) + u(\mathbf{x}_i - \mathbf{z}_j) - 2u(\mathbf{x}_i) = \mathbf{z}_j \otimes \mathbf{z}_j : M.$$

Thus,

$$\mathcal{L}_{h,\delta} u(\mathbf{x}_i) = \sum_{\mathbf{j} \neq \mathbf{0}} \frac{\mathbf{z}_j \otimes \mathbf{z}_j : M}{|\mathbf{z}_j|^2} |\mathbf{z}_j|_1 \beta_\delta(\mathbf{z}_j) = Q : M.$$

We now need another observation on the coefficient matrix, that is, $\beta_\delta(\mathbf{z}_j)$ is an even function; moreover, it depends only on the absolute values of the individual components of $\mathbf{z}_j$. Thus, when doing the summation over $\mathbf{j}$, it is easy to check that $Q_{kl} = 0$ for $k \neq l$. We can further check that Q_{kk} is independent of k , and

$$\begin{aligned} \sum_k Q_{kk} &= \int_{B_\delta(\mathbf{0})} \left(\sum_{\mathbf{j} \neq \mathbf{0}} \frac{|\mathbf{z}_j|^2}{|\mathbf{z}_j|^2} |\mathbf{z}_j|_1 \phi_j(\mathbf{z}) \right) \frac{|\mathbf{z}|^2}{|\mathbf{z}|_1} \rho_\delta(|\mathbf{z}|) d\mathbf{z} \\ &= \int_{B_\delta(\mathbf{0})} |\mathbf{z}|^2 \rho_\delta(|\mathbf{z}|) d\mathbf{z} = d, \end{aligned}$$

where in the last step we have used the property that $|\mathbf{z}|_1$ is piecewise linear, and thus

$$\sum_{\mathbf{j} \neq \mathbf{0}} |\mathbf{z}_j|_1 \phi_j(\mathbf{z}) = \sum_{\mathbf{j}} |\mathbf{z}_j|_1 \phi_j(\mathbf{z}) = |\mathbf{z}|_1.$$

Hence, we can derive that $Q = I_d$, which gives

$$\mathcal{L}_\delta u(\mathbf{x}_i) = \mathcal{L}_{h,\delta} u(\mathbf{x}_i) \equiv \sum_k m_{kk}.$$

This completes the proof. $\square$

Before proceeding to the main theorem of this work we introduce some notation for simplicity of presentation. For any function $u = u(\mathbf{x}) : \mathbb{R}^d \mapsto \mathbb{R}$ and $\alpha \in \mathbb{N}^d$ let

$$\mathbf{x}^\alpha = \prod_{i=1}^d x_i^{\alpha_i}, \quad \partial^\alpha u(\mathbf{x}) = \partial_{\alpha_1} \cdots \partial_{\alpha_d} u(\mathbf{x}),$$

and $D^k u(\mathbf{x})$ be the set containing all the k th-order derivatives in the form of $\partial^\alpha u(\mathbf{x})$ with $|\alpha|_1 = k$, and

$$|D^k u|_\infty = \max_{|\alpha|_1=k} \sup_{\mathbf{x}} \partial^\alpha u(\mathbf{x}).$$

We further use D_2 to denote the Hessian and define

$$H^{\mathbf{x}}(\mathbf{z}) = u(\mathbf{x} + \mathbf{z}) + u(\mathbf{x} - \mathbf{z}) - 2u(\mathbf{x}), \quad H_1^{\mathbf{x}}(\mathbf{z}) = \mathbf{z} \otimes \mathbf{z} : D_2 u(\mathbf{x})$$

and

$$J^{\mathbf{x}}(\mathbf{z}) = \frac{H^{\mathbf{x}}(\mathbf{z})}{|\mathbf{z}|^2} |\mathbf{z}|_1, \quad J_1^{\mathbf{x}}(\mathbf{z}) = \frac{H_1^{\mathbf{x}}(\mathbf{z})}{|\mathbf{z}|^2} |\mathbf{z}|_1,$$

and let $H_2^{\mathbf{x}}(\mathbf{z}) = H^{\mathbf{x}}(\mathbf{z}) - H_1^{\mathbf{x}}(\mathbf{z})$ and $J_2^{\mathbf{x}}(\mathbf{z}) = J^{\mathbf{x}}(\mathbf{z}) - J_1^{\mathbf{x}}(\mathbf{z})$. By Taylor's expansion and the mean value theorem we can show that

$$\left\{ \begin{array}{l} H_2^{\mathbf{x}}(\mathbf{z}) = \sum_{|\alpha|_1=4} C_\alpha^1 \mathbf{z}^\alpha \partial^\alpha u(\mathbf{x}_1(\mathbf{z})), \\ \frac{\partial}{\partial z_i} H_2^{\mathbf{x}}(\mathbf{z}) = \sum_{|\alpha|_1=3} C_\alpha^2 \mathbf{z}^\alpha \partial^{\alpha+\mathbf{e}_i} u(\mathbf{x}_2(\mathbf{z})), \\ \frac{\partial^2}{\partial z_i \partial z_j} H_2^{\mathbf{x}}(\mathbf{z}) = \sum_{|\alpha|_1=2} C_\alpha^3 \mathbf{z}^\alpha \partial^{\alpha+\mathbf{e}_i+\mathbf{e}_j} u(\mathbf{x}_3(\mathbf{z})), \end{array} \right. \quad (3.10)$$

where $\mathbf{e}_i$ and $\mathbf{e}_j$ are unit vectors with only one nonzero element at the i th and j th positions, respectively, and $\mathbf{x}_k(\mathbf{z})$ are some points dependent on both $\mathbf{x}$ and $\mathbf{z}$. The following is an elementary but technical result to be used later.

LEMMA 3.6 For $J_2^{\mathbf{x}}(\mathbf{z})$ defined above, if $u \in C^4(\overline{\Omega})$, for any $\mathbf{z}$ and $\mathbf{x}$, we have

$$|D^2 J_2^{\mathbf{x}}|_\infty \leq C |D^4 u|_\infty |\mathbf{z}|_1, \quad (3.11)$$

where C is generic constant.

Proof. To carry out the estimates we first note that

$$\frac{\partial |\mathbf{z}|^2}{\partial z_i} = 2z_i \quad \text{and} \quad \frac{\partial |\mathbf{z}|^2}{\partial z_i \partial z_j} = 2\delta_{ij}.$$

Now, by the definition of $J_2^{\mathbf{x}}(\mathbf{z})$, we have

$$\frac{\partial^2}{\partial z_i \partial z_j} J_2^{\mathbf{x}}(\mathbf{z}) = \text{sign}(z_i) \frac{\partial}{\partial z_j} \frac{H_2^{\mathbf{x}}(\mathbf{z})}{|\mathbf{z}|^2} + \text{sign}(z_j) \frac{\partial}{\partial z_i} \frac{H_2^{\mathbf{x}}(\mathbf{z})}{|\mathbf{z}|^2} + |\mathbf{z}|_1 \frac{\partial^2}{\partial z_i \partial z_j} \frac{H_2^{\mathbf{x}}(\mathbf{z})}{|\mathbf{z}|^2},$$

where $\text{sign}(\cdot)$ is the sign function. For the first term in the above, we have from (3.10) that

$$\left| \text{sign}(z_i) \frac{\partial}{\partial z_j} \frac{H_2^x(\mathbf{z})}{|\mathbf{z}|^2} \right| \leq C \left(\left| \frac{\mathbf{z}^{\beta_1}}{|\mathbf{z}|^2} \right| + \left| \frac{\mathbf{z}^{\beta_2}}{|\mathbf{z}|^4} \right| \right) |D^4 u|_\infty,$$

where $|\beta_1|_1 = 3$ and $|\beta_2|_1 = 5$. The second term satisfies a similar estimate. Moreover, by (3.10) again, we get

$$\left| \frac{\partial^2}{\partial z_i \partial z_j} \frac{H_2^x(\mathbf{z})}{|\mathbf{z}|^2} \right| \leq C \left(\left| \frac{\mathbf{z}^{\alpha_1}}{|\mathbf{z}|^4} \right| + \left| \frac{\mathbf{z}^{\alpha_2}}{|\mathbf{z}|^8} \right| \right) |D^4 u|_\infty,$$

where $|\alpha_1|_1 = 4$ and $|\alpha_2|_1 = 8$. For any $\alpha \in \mathbb{N}^d$ it is obvious that

$$\left| \frac{\mathbf{z}^\alpha}{|\mathbf{z}|^{|\alpha|_1}} \right| \leq 1, \quad \left| \frac{\mathbf{z}^\alpha}{|\mathbf{z}|^{|\alpha|_1-1}} \right| \leq |\mathbf{z}|_1.$$

Combining the above estimates we can get the desired result (3.11). $\square$

We now present the truncation error analysis for the quadrature-based difference approximation of the nonlocal model.

LEMMA 3.7 (Uniform nonlocal truncation error). Assume that $u \in C^4(\overline{\Omega} \cup \overline{\Omega_T})$. Then it holds that

$$\max_{1 \leq i \leq N_s} |\mathcal{L}_{h,\delta} u(\mathbf{x}_i) - \mathcal{L}_\delta u(\mathbf{x}_i)| \leq C |D^4 u|_\infty h^2, \quad (3.12)$$

where C is a constant independent of δ and h .

Proof. Without loss of generality we take $\mathbf{x}_i = 0$ to be the origin. Note that

$$\mathcal{L}_\delta u(\mathbf{0}) = \int_{\mathcal{B}_\delta(\mathbf{0})} J^0(\mathbf{z}) \frac{|\mathbf{z}|^2}{|\mathbf{z}|_1} \rho_\delta(|\mathbf{z}|) \, d\mathbf{z}$$

and

$$\mathcal{L}_{h,\delta} u(\mathbf{0}) = \int_{\mathcal{B}_\delta(\mathbf{0})} \mathcal{I}_h(J^0(\mathbf{z})) \frac{|\mathbf{z}|^2}{|\mathbf{z}|_1} \rho_\delta(|\mathbf{z}|) \, d\mathbf{z}.$$

Simple subtraction gives us

$$\begin{aligned} |\mathcal{L}_{h,\delta} u(\mathbf{0}) - \mathcal{L}_\delta u(\mathbf{0})| &= \left| \int_{\mathcal{B}_\delta(\mathbf{0})} (\mathcal{I}_h(J^0(\mathbf{z})) - J^0(\mathbf{z})) \frac{|\mathbf{z}|^2}{|\mathbf{z}|_1} \rho_\delta(|\mathbf{z}|) \, d\mathbf{z} \right|, \\ &\leq \left| \int_{\mathcal{B}_\delta(\mathbf{0})} (\mathcal{I}_h(J_1^0(\mathbf{z})) - J_1^0(\mathbf{z})) \frac{|\mathbf{z}|^2}{|\mathbf{z}|_1} \rho_\delta(|\mathbf{z}|) \, d\mathbf{z} \right| \\ &\quad + \left| \int_{\mathcal{B}_\delta(\mathbf{0})} (\mathcal{I}_h(J_2^0(\mathbf{z})) - J_2^0(\mathbf{z})) \frac{|\mathbf{z}|^2}{|\mathbf{z}|_1} \rho_\delta(|\mathbf{z}|) \, d\mathbf{z} \right| \\ &:= E_1 + E_2, \end{aligned}$$

where

$$J_1^0(\mathbf{z}) = \frac{\mathbf{z} \otimes \mathbf{z} : D_2 u(\mathbf{0})}{|\mathbf{z}|^2} |\mathbf{z}|_1 \quad \text{and} \quad J_2^0(\mathbf{z}) = J^0(\mathbf{z}) - J_1^0(\mathbf{z}).$$

From Lemma 3.5, we know that $E_1 = 0$. For E_2 , we have

$$E_2 \leq Ch^2 \int_{B_\delta(\mathbf{0})} \left| D^2 J_2^0 \right|_\infty \frac{|\mathbf{z}|^2}{|\mathbf{z}|_1} \rho_\delta(|\mathbf{z}|) d\mathbf{z}.$$

Thus, by using Lemma 3.6 and the moment condition on the kernel that

$$\int_{B_\delta(\mathbf{0})} |\mathbf{z}|^2 \rho_\delta(|\mathbf{z}|) d\mathbf{z} = d,$$

we complete the proof. $\square$

Combining the stability result in Lemma 3.4 and the consistency result in Lemma 3.7, we immediately get the convergence of the quadrature-based finite difference scheme.

THEOREM 3.8 (Convergence to nonlocal solution). Assume that the nonlocal exact solution u^δ of (1.1) is smooth enough, such as $u^\delta \in C^4(\overline{\Omega \cup \Omega_{\mathcal{I}}})$, and $u^{\delta,h}$ is the numerical solution obtained by the scheme (3.4). Then for any fixed δ there exists some constant C_δ independent of h as $h \rightarrow 0$ such that

$$\left\| u^{\delta,h} - I_h u^\delta \right\|_\infty \leq Ch^2, \quad (3.13)$$

where $I_h u^\delta$ is the interpolation of the exact solution on the grid points. For any $\delta < \delta_0$ the constant C_δ is uniformly bounded by some generic constant $C(\delta_0)$.

Next we show the asymptotic compatibility of the quadrature-based finite difference scheme to the local limiting model. We assume that the local equation (2.2) is discretized by the classical central finite difference scheme denoted by $\mathcal{L}_{h,0}$, whose truncation error is

$$\max_{1 \leq i \leq N_s} |\mathcal{L}_{h,0} u(\mathbf{x}_i) - \mathcal{L}_0 u(\mathbf{x}_i)| = \mathcal{O}(h^2), \quad (3.14)$$

by the standard numerical PDE analysis, if $u \in C^4(\overline{\Omega})$.

We now estimate the modeling error at the discrete level, that is, the error between the discrete nonlocal operator $\mathcal{L}_{h,\delta}$ and $\mathcal{L}_{h,0}$.

LEMMA 3.9 (Discrete model error). Assume that $u \in C^4(\overline{\Omega \cup \Omega_{\mathcal{I}}})$. Then it holds that

$$\max_{1 \leq i \leq N_s} |\mathcal{L}_{h,\delta} u(\mathbf{x}_i) - \mathcal{L}_{h,0} u(\mathbf{x}_i)| = \mathcal{O}(\delta^2) + \mathcal{O}(h^2) \quad (3.15)$$

when δ and h are sufficiently small.

Proof. The triangle inequality gives

$$\begin{aligned} |\mathcal{L}_{h,\delta}u(\mathbf{x}_i) - \mathcal{L}_{h,0}u(\mathbf{x}_i)| &\leq |\mathcal{L}_{h,\delta}u(\mathbf{x}_i) - \mathcal{L}_\delta u(\mathbf{x}_i)| + |\mathcal{L}_0 u(\mathbf{x}_i) - \mathcal{L}_\delta u(\mathbf{x}_i)| \\ &\quad + |\mathcal{L}_0 u(\mathbf{x}_i) - \mathcal{L}_{h,0}u(\mathbf{x}_i)| \\ &:= R_1 + R_2 + R_3. \end{aligned}$$

By Lemma 3.7 we have

$$R_1 = \mathcal{O}(h^2).$$

For $u \in C^4(\overline{\Omega})$ the continuum property of nonlocal operators gives

$$R_2 = \mathcal{O}(\delta^2).$$

It is well known that

$$R_3 = \mathcal{O}(h^2).$$

The three equalities above together complete the proof. $\square$

Combining the consistency in Lemma 3.9 with the stability Lemma 3.4 again, we get convergence (and asymptotic compatibility) to the local limit.

THEOREM 3.10 (Asymptotic compatibility). Assume that the local exact solution u^0 of (2.2) is smooth enough, such as $u^0 \in C^4(\overline{\Omega \cup \Omega_T})$. Then for any $\delta < \delta_0$ there is some constant C independent of δ and h as $h \rightarrow 0$ such that

$$\|u^{\delta,h} - I_h u^0\|_\infty \leq C(h^2 + \delta^2), \quad (3.16)$$

where $I_h u^0$ is the interpolation of the exact solution on the grid points.

REMARK 3.11 For problems on the same domain but with periodic boundary conditions (PBCs) all similar estimates also hold, which requires only that the exact solution of the local model is a $C_{\text{per}}^4(\overline{\Omega})$ function.

REMARK 3.12 We now give a more explicit form of the algebraic system in two dimensions. We let $u_{j,k}^{\delta,h}$ denote the nodal value of the nonlocal difference solution at the mesh point $(x_j, y_k) \in \Omega = (-\pi, \pi)^2 \subset \mathbb{R}^2$. Then let $r = \lceil \delta/h \rceil$ be the smallest integer larger than or equal to δ/h ; we have

$$\mathcal{L}_{h,\delta} u_{j,k}^{\delta,h} = \sum_{p=0}^r \sum_{q=0}^r c_{p,q} \left(u_{j+p,k+q}^{\delta,h} + u_{j-p,k+q}^{\delta,h} + u_{j+p,k-q}^{\delta,h} + u_{j-p,k-q}^{\delta,h} - 4u_{j,k}^{\delta,h} \right),$$

where $c_{0,0} = 0$ and

$$c_{p,q} = \frac{ph + qh}{p^2 h^2 + q^2 h^2} \iint_{B_+(0,\delta)} \phi_{p,q}(s,t) \rho_\delta \left(\sqrt{s^2 + t^2} \right) \frac{s^2 + t^2}{s+t} \, ds \, dt,$$

with $B_+(0, \delta)$ denoting the first quadrant of the disc at the origin with radius δ . One may also observe that $c_{p,q} = c_{q,p}$ for any p and q .

4. Green's functions of a nonlocal operator with integrable kernels

By definition, the Green's function of the nonlocal equation (corresponding to the particular nonlocal operator $\mathcal{L}_\delta$) solves the nonlocal equation with right-hand side given by the Dirac delta measure $\delta_{\mathbf{y}}(\mathbf{x}) = \delta(\mathbf{x} - \mathbf{y})$ of the variable $\mathbf{x}$ for a given $\mathbf{y}$. Discussions of Green's functions for nonlocal models such as PD have been presented in a number of earlier studies (Weckner *et al.*, 2009; Silling *et al.*, 2010b; Wang *et al.*, 2016, 2017). Here we pay particular attention to the case where the nonlocal interaction kernel is given to be an integrable one. A consequence is that, unlike Green's functions for local diffusion equations, the nonlocal analog is generically a measure function. Hence, we focus on effective constructions in this case.

4.1 Periodic case

For the case of PBCs with $\Omega = [-\pi, \pi]^d$, we may impose the zero mean compatibility condition for both the right-hand side and the solution over Ω . Thus, we formally have the following modified equation of (1.1):

$$-\mathcal{L}_\delta G_\delta^\delta(\mathbf{x}) = \delta_{\mathbf{y}} - \frac{1}{|\Omega|},$$

where $|\Omega|$ is the area of the domain. Taking the L^2 -inner product of the above equation with $u^\delta(\mathbf{x})$ over $\mathbf{x}$ yields

$$\left(-\mathcal{L}_\delta G_\delta^\delta, u^\delta\right) = u^\delta(\mathbf{y}) - \frac{1}{|\Omega|} \int_\Omega u^\delta \, d\mathbf{x} = u^\delta(\mathbf{y}).$$

By the nonlocal Green's identity (Du *et al.*, 2012, 2013), which remains valid for the periodic case, we have $\left(-\mathcal{L}_\delta G_\delta^\delta, u^\delta\right) = \left(G_\delta^\delta, -\mathcal{L}_\delta u^\delta\right)$. We then get

$$u(\mathbf{y}) = \left(f, G_\delta^\delta\right),$$

which demonstrates how Green's function G_δ^δ can be used to represent solutions to the nonlocal equation (1.1).

Now, for the case where the kernel is integrable we denote

$$C_\delta = \int_{\mathcal{B}_\delta(\mathbf{0})} \rho_\delta(|\mathbf{z}|) \, d\mathbf{z}.$$

Then the nonlocal operator $-\mathcal{L}_\delta$ can be written as

$$-\mathcal{L}_\delta u = -\rho_\delta * u + C_\delta u.$$

Let us perform the splitting $G_\delta^\delta = S_1 + g_0$ with S_1 and g_0 to be determined; we then arrive at

$$-\mathcal{L}_\delta S_1(\mathbf{x}) - \rho_\delta * g_0 + C_\delta g_0 = \delta_{\mathbf{y}} - \frac{1}{|\Omega|}.$$

Setting

$$g_0 = C_\delta^{-1} \left(\delta_{\mathbf{y}} - \frac{1}{|\Omega|} \right) \quad \text{and} \quad g_1 = C_\delta^{-1} \rho_\delta * g_0 \quad (4.1)$$

leads to

$$-\mathcal{L}_\delta S_1(\mathbf{x}) = \rho_\delta * g_0(\mathbf{x}) = C_\delta g_1. \quad (4.2)$$

REMARK 4.1 Note that while g_0 is a singular measure we have generically that $g_1 \in L^1$ for an integrable ρ_δ . If additional regularity of ρ_δ can be assumed then we also get g_1 of the same regularity class as ρ_δ . For example, if ρ_δ is of the class C^∞ (as a function in the whole space having compact support in the ball of radius δ) then we also have g_1 and S_1 in C^∞ . Another example, in the one-dimensional case, is that if ρ_δ is a bounded variation (BV) function then so are g_1 and S_1 .

In turn, if we perform a similar splitting once again as $S_1 = S_2 + g_1$, we have

$$-\mathcal{L}_\delta S_2(\mathbf{x}) = \rho_\delta * g_1.$$

Similarly, repeating this splitting technique n times we get recursively that

$$g_n = C_\delta^{-1} \rho_\delta * g_{n-1}, \quad -\mathcal{L}_\delta S_{n+1}(\mathbf{x}) = \rho_\delta * g_n,$$

and generically, the later terms obtained by the recursion are more regular than the preceding terms, similarly to the observation in Remark 4.1. In the end, we get

$$G_{\mathbf{y}}^\delta = \sum_{n=0}^{\infty} g_n$$

with the following recursive relation:

$$g_0 = C_\delta^{-1} \left(\delta_{\mathbf{y}} - \frac{1}{|\Omega|} \right), \quad g_n = C_\delta^{-1} \rho_\delta * g_{n-1}, \quad n = 1, 2, \dots$$

Thus, we obtain an explicit series form of $G_{\mathbf{y}}^\delta(\mathbf{x})$, which is given below.

THEOREM 4.2 For an integrable kernel ρ_δ the nonlocal Green's function $G_{\mathbf{y}}^\delta$ for equation (1.1) with PBCs can be represented by

$$G_{\mathbf{y}}^\delta = \sum_{n=0}^{\infty} C_\delta^{-(n+1)} (\rho_\delta *)^n * \left(\delta_{\mathbf{y}} - \frac{1}{|\Omega|} \right). \quad (4.4)$$

Moreover, the solution u^δ of the nonlocal diffusion equation (1.1) is given by

$$u^\delta(\mathbf{y}) = (f, G_{\mathbf{y}}^\delta).$$

REMARK 4.3 Due to the lack of elliptic smoothing of the nonlocal diffusion equation corresponding to the integrable kernels, the nonlocal Green's function remains a composition of a singular part in the form of a Dirac delta measure and an integrable part. In this case and unlike local Green's functions for elliptic PDEs, G_y^δ should not be viewed as a function defined almost everywhere. Moreover, the regularity of the solution of u^δ is in general the same as the right-hand side f , since the regularity (f, G_y^δ) is dominated by the first term (g_0, f) , which is proportional to f . One could also split the solution into two parts, one given by (g_0, f) while the other accounts for a more regular part.

REMARK 4.4 It is worth mentioning that the series expression for the Green's function is a Neumann series expansion for the equation $(I - A)G = f$ where $Au = C_\delta^{-1} \rho_\delta * u$ and $f = C_\delta^{-1} \left(\delta_y - \frac{1}{|\Omega|} \right)$.

4.2 Other cases of nonlocal constraints

The approach of constructing Green's function through a series expansion is also applicable to other nonlocal constraints. For instance, we consider the Dirichlet-type volume-constrained problems given by (1.1) and (2.1). For any $y \in \Omega$ let us define $g^0 = \delta(x - y)$, $x \in \Omega \cup \Omega_T$. It is easy to see that, with g^n recursively defined by

$$\begin{cases} g^n(x) = C_\delta^{-1} \rho_\delta * g^{n-1}(x), & x \in \Omega, \\ g^n(x) = 0, & x \in \Omega_T, \end{cases}$$

the nonlocal Green's function is given by

$$G_y^\delta(x) = \sum_{n=0}^{\infty} g^n(x).$$

As in the case with PBC, we can also use the above expansion to get a series expression for a general solution, though the regularity pickup in the later terms of the series gets more delicate due to the dependence on matching the boundary data with the right-hand side and remains an interesting topic to be further studied.

REMARK 4.5 Other types of series expansions, in particular Fourier series expansion, of the nonlocal Green's functions have been presented in the literature by Weckner *et al.* (2009), Silling *et al.* (2010b) and Wang *et al.* (2016, 2017). Such expansions are valid formally, but for integrable kernels the series do not converge pointwisely.

4.3 Discrete nonlocal Green's function

Following the discussion above on the continuum level we can present a discrete analog with the AC quadrature-based finite difference scheme.

Given the uniform mesh $\{x_j\}$ the discrete Green's function (matrix) $G_{x_i}^{h,\delta}(x_j)$ is given by

$$-\mathcal{L}_{h,\delta} G_{x_i}^{h,\delta}(x_j) = \frac{1}{h^d} \delta_{ij} - \frac{1}{|\Omega|},$$

where δ_{ij} is the Kronecker delta function, that is, $\delta_{ii} = 1$ and $\delta_{ij} = 0$ for $i \neq j$. It follows that the discrete solution to the nonlocal model with a discrete right-hand side $\{f_j\}$ is given by

$$u^{h,\delta}(x_i) = \sum_{j \neq i} f_j G_{x_i}^{h,\delta}(x_j) \quad \text{for} \quad \sum_j f_j = 0.$$




FIG. 1. The function $u_1^{h,\delta}$ of the regular part of the nonlocal Green's function.

The study of discrete nonlocal Green's functions can be seen as a nonlocal analog of similar works on discrete Green's functions for local PDEs (Chung & Yau, 2000). Although the nonlocal Green's function at the continuum level is measure valued for integrable interaction kernels the discrete nonlocal Green's function given above is well defined at all the grid points for any finite δ and h . Due to the lack of regularity, however, we do not expect strong (or pointwise) convergence of $G_y^{h,\delta}(\mathbf{x})$ to $G_y^\delta(\mathbf{x})$ as $h \rightarrow 0$ for a given δ . Instead, we could attempt to get better approximations by the splitting technique, that is, we numerically solve for the regular part while invoking the analytical solution of singular part.

Therefore, we obtain a hybrid approximation of $G_{\mathbf{x}_j}^\delta(\mathbf{x}_i)$ by

$$G_{\mathbf{x}_j}^{h,\delta}(\mathbf{x}_i) = C_\delta^{-1} \left(\delta_{\mathbf{x}_j}(\mathbf{x}_i) - \frac{1}{|\Omega|} \right) + u_1^{h,\delta},$$

where the approximation $u_1^{h,\delta}$ of the regular part of the nonlocal Green's function S_1 in (4.2) satisfies

$$-\mathcal{L}_{h,\delta} u_1^{h,\delta}(\mathbf{x}_i) = \rho_\delta * g_0(\mathbf{x}_i).$$

Given the higher regularity of S_1 we expect to have good convergence of $u_1^{h,\delta}$ to S_1 for any given δ , which is illustrated in Fig. 1 for a kernel that is a positive constant over the horizon and vanishes outside (thus piecewise continuous). Moreover, due to the AC property of the discrete scheme, we also have convergence of $G_{\mathbf{x}_j}^{h,\delta}(\mathbf{x}_i)$ to the local Green's function as both h and δ approach zero. Notice that both Theorems 3.8 and 3.10 require that the solutions to be approximated are $C^4(\overline{\Omega \cup \Omega_I})$. Thus, if we




FIG. 2. Numerical comparison. Left: numerical solutions; right: zoom in around π .

subtract enough terms of the Green's function series expansion from the right-hand side, in principle we should be left with a smooth enough right-hand side that the results of both theorems are applicable. More discussions on this splitting technique are to be presented later. We thus can expect a robust approximation of the local and nonlocal Green's functions.

4.4 More discussion on the splitting technique

We next discuss a potential application of the splitting technique presented above in the numerical solution of nonlocal equations. By avoiding using derivatives, nonlocal models allow more singular solutions, which is a distinct advantage in physical modeling such as the study of material cracks. As an illustration we present a typical example showing how the splitting technique improves the performance of Fourier spectral methods for numerically solving nonlocal equations with discontinuous solutions. For simplicity, we take a one-dimensional nonlocal diffusion model (1.1) over a periodic cell $\Omega = [0, 2\pi]$ with a constant kernel $\rho_\delta(s) = \frac{3}{\delta^3} \chi_{[-\delta, \delta]}$, which is of the rescaled form given in (2.3), and a discontinuous right-hand side

$$f(x) = \text{sign}(x - \pi), \quad x \in [0, 2\pi],$$

where $\text{sign}(\cdot)$ is the sign function. In this setting, the exact solution has a jump at $x = \pi$, e.g., like studied in Du & Yang (2016). One numerical solution is obtained by applying the Fourier spectral method directly. It is not surprising that the well-known Gibbs phenomena occur around $x = \pi$ when standard Fourier spectral methods are directly used to approximate discontinuous solutions. On the other hand, if we use the splitting technique described above once, by first constructing the discontinuous part of the solution analytically, and then solving a new nonlocal diffusion equation with a smoother right-hand side by Fourier spectral methods, we can eliminate the Gibbs phenomena, as demonstrated in Fig. 2.

5. Conclusion

In this work, a quadrature-based finite difference scheme for a nonlocal diffusion model in multidimensions is proposed. Our analysis is given for problems with Dirichlet volumetric constraints. The case of a Neumann-type nonlocal volumetric constraint may involve additional complications, due to the need for ghost points, which require further attention to ensure second-order accuracy. One may adapt the scheme developed here to more general models. In fact, an extension based on the idea presented in a

draft version of this work has been given for nonlocal convection diffusion equations in the study by Tian *et al.* (2018) (where only a partial analysis of the diffusion term has been documented and one should refer to the more complete analysis presented in the current work). In deriving the error analysis our attention is focused on the case where the underlying solutions are smooth. Since nonlocal models are effective in describing physical processes with solution singularities, extending the analysis to cases with minimal regularity will be of interest. One can of course also consider extensions to systems involving vector fields such as the bond-based and state-based PD models.

Our study here also illustrated an important difference between the nonlocal Green's function and their local analog as the former may take on a possibly singular measure form. Nonlocal Green's functions can be useful in various applications such as the mean exit time computation of jump processes and analysis of mechanical responses to point loads in nonlocal mechanics (Weckner *et al.*, 2009; Silling *et al.*, 2010b; Du *et al.*, 2012; Wang *et al.*, 2016, 2017). The splitting technique presented here offers an effective algorithm for computing singular Green's functions. Moreover, one may further explore the applications of such techniques to the computation of singular solutions to more general nonlocal models as alluded to in Section 4.4.

In comparison with the Galerkin finite element AC scheme developed in the study by Tian & Du (2014) that works for linear systems in multidimensions using unstructured meshes, we see that the quadrature-based finite difference AC schemes given here are developed only for uniform Cartesian meshes with the same mesh size in all directions. However, the schemes given here are simpler to implement as they avoid the evaluation of integrals in $\mathbb{R}^{2d}$. Furthermore, they lead to M-matrices and the discrete maximum principle that are in general not the case for the finite element discretization. Hence, an interesting future work is to develop hybrid schemes and particle discretization (Liu *et al.*, 1996; Silling, 2000; Askari *et al.*, 2008; Bessa *et al.*, 2014) that can preserve the nice properties of the quadrature-based finite difference AC schemes while applicable to more general meshes.

Funding

United States National Science Foundation (DMS-1719699), Air Force Office of Scientific Research MURI center and the Army Research Office MURI Grant (W911NF-15-1-0562).

REFERENCES

- ANDREU, F., MAZÓN, J. M., ROSSI, J. D. & TOLEDO, J. (2010) *Nonlocal Diffusion Problems*. Mathematical Surveys and Monographs, vol. 165. Providence, RI: AMS.
- APPLEBAUM, D. (2004) *Lévy Processes and Stochastic Calculus*. Cambridge Studies in Advanced Mathematics, vol. 93. Cambridge, UK: Cambridge University Press.
- ASKARI, E., BOBARU, F., LEHOUCQ, R. B., PARKS, M. L., SILLING, S. A. & WECKNER, O. (2008) Peridynamics for multiscale materials modeling. *J. Phys. Conf. Ser.*, **125**, 12–78.
- BATES, P. W. & CHMAJ, A. (1999) An integrodifferential model for phase transitions: stationary solutions in higher space dimensions. *J. Stat. Phys.*, **95**, 1119–1139.
- BESSA, M., FOSTER, J., BELYTSCHKO, T. & LIU, W. K. (2014) A meshfree unification: reproducing kernel peridynamics. *Comput. Mech.*, **53**, 1251–1264.
- BOBARU, F. & DUANGPANYA, M. (2010) The peridynamic formulation for transient heat conduction. *Int. J. Heat Mass Transf.*, **53**, 4047–4059.
- BOBARU, F., YANG, M., ALVES, L. F., SILLING, S. A., ASKARI, E. & XU, J. (2009) Convergence, adaptive refinement, and scaling in 1D peridynamics. *Int. J. Numer. Methods Eng.*, **77**, 852–877.

- BUADES, A., COLL, B. & MOREL, J. M. (2010) Image denoising methods. A new nonlocal principle. *SIAM Rev.*, **52**, 113–147.
- CHEN, A., DU, Q., LI, C. & ZHOU, Z. (2017) Asymptotically compatible schemes for space–time nonlocal diffusion equations. *Chaos Solitons Fractals* (in press).
- CHEN, X. & GUNZBURGER, M. (2011) Continuous and discontinuous finite element methods for a peridynamics model of mechanics. *Comput. Methods Appl. Mech. Eng.*, **200**, 1237–1250.
- CHUNG, F. & YAU, S-T. (2000) Discrete Green's functions. *J. Comb. Theory A*, **91**, 191–214.
- DU, Q., GUNZBURGER, M., LEHOUCQ, R. B. & ZHOU, K. (2012) Analysis and approximation of nonlocal diffusion problems with volume constraints. *SIAM Rev.*, **56**, 676–696.
- DU, Q., GUNZBURGER, M., LEHOUCQ, R. B. & ZHOU, K. (2013) A nonlocal vector calculus, nonlocal volume-constrained problems, and nonlocal balance laws. *Math. Models Methods Appl. Sci.*, **23**, 493–540.
- DU, Q., JU, L. & LU, J. (2017a) A discontinuous Galerkin method for one-dimensional time-dependent nonlocal diffusion problems, *Preprint*.
- DU, Q., TAO, Y., TIAN, X. & YANG, J. (2016) Robust a posteriori stress analysis for quadrature collocation approximations of nonlocal models via nonlocal gradients. *Comput. Methods Appl. Mech. Eng.*, **310**, 605–627.
- DU, Q. & TIAN, X. (2014) Asymptotically compatible schemes for peridynamics based on numerical quadratures. *ASME 2014 International Mechanical Engineering Congress and Exposition (IMECE2014)*, vol. 1, Paper No. IMECE2014–39620, pp. V001T01A058.
- DU, Q. & YANG, J. (2016) Asymptotic compatible Fourier spectral approximations of nonlocal Allen–Cahn equations. *SIAM J. Numer. Anal.*, **54**, 1899–1919.
- DU, Q. & YANG, J. (2017) Fast and accurate implementation of Fourier spectral approximations of nonlocal diffusion operators and its applications. *J. Comput. Phys.*, **332**, 118–134.
- DU, Q., YANG, J. & ZHOU, Z. (2017b) Analysis of a nonlocal-in-time parabolic equation. *Discrete Continuous Dyn. Syst. B.*, **22**, 339–368.
- DU, Q. & ZHOU, K. (2011) Mathematical analysis for the peridynamic nonlocal continuum theory. *Math. Model. Numer. Anal.*, **45**, 217–234.
- GILBOA, G. & OSHER, S. (2008) Nonlocal operators with applications to image processing. *Multiscale Model. Simul.*, **7**, 1005–1028.
- KILIC, B. & MADENCI, E. (2010) Coupling of peridynamic theory and the finite element method. *J. Mech. Mater. Struct.*, **5**, 707–733.
- LIU, W. K., CHEN, Y., JUN, S., CHEN, J., BELYTSCHKO, T., PAN, C., URAS, R. & CHANG, C. (1996) Overview and applications of the reproducing kernel particle methods. *Arch. Comput. Methods Eng.*, **3**, 3–80.
- LOU, Y., ZHANG, X., OSHER, S. & BERTOZZI, A. (2010) Image recovery via nonlocal operators. *J. Sci. Comput.*, **42**, 185–197.
- MACEK, R. & SILLING, S. A. (2007) Peridynamics via finite element analysis. *Finite Elem. Anal. Des.*, **43**, 1169–1178.
- MENGESHA, T. & DU, Q., (2014) Nonlocal constrained value problems for a linear peridynamic Navier equation. *J. Elast.*, **116**, 27–51.
- PALATUCCI, G., SAVIN, O. & VALDINOCI, E. (2012) Local and global minimizers for a variational energy involving a fractional norm. *Ann. Mat. Pura Appl.*, **192**, 673–718.
- SILLING, S. A. (2000) Reformulation of elasticity theory for discontinuities and long-range forces. *J. Mech. Phys. Solids*, **48**, 175–209.
- SILLING, S. A. & ASKARI, E. (2005) A meshfree method based on the peridynamic model of solid mechanics. *Comput. Struct.*, **83**, 1526–1535.
- SILLING, S. A. & LEHOUCQ, R. B. (2010) Peridynamic theory of solid mechanics. *Adv. Appl. Mech.*, **44**, 73–168.
- SILLING, S. A., WECKNER, O., ASKARI, E. & BOBARU, F. (2010a) Crack nucleation in a peridynamic solid. *Int. J. Fract.*, **162**, 219–227.
- SILLING, S. A., ZIMMERMANN, M. & ABEYARATNE, R. (2010b) Deformation of a peridynamic bar. *J. Elast.*, **73**, 173–190.

- TADMOR, E. & TAN, C. (2014) Critical thresholds in flocking hydrodynamics with non-local alignment. *Philos. Trans. R. Soc. Lond. A Math. Phys. Eng. Sci.*, **372**: 20130401.
- TAO, Y., TIAN, X. & DU, Q. (2017) Nonlocal diffusion and peridynamic models with Neumann type constraints and their numerical approximations. *Appl. Math. Comput.*, **305**, 282–298.
- TIAN, H., JU, L. & DU, Q. (2018) A conservative nonlocal convection-diffusion model and asymptotically compatible finite difference discretization. *Comput. Methods Appl. Mech. Eng.*, **320**, 46–67.
- TIAN, X. & DU, Q. (2013) Analysis and comparison of different approximations to nonlocal diffusion and linear peridynamic equations. *SIAM J. Numer. Anal.*, **51**, 3458–3482.
- TIAN, X. & DU, Q. (2014) Asymptotically compatible schemes and applications to robust discretization of nonlocal models. *SIAM J. Numer. Anal.*, **52**, 1641–1665.
- WANG, L., XU, J. & WANG, J. (2016) The Green's functions for peridynamic non-local diffusion. *Proc. R. Soc. A*, **472**, 20160185.
- WANG, L., XU, J. & WANG, J. (2017) Static and dynamic Green's functions in peridynamics. *J. Elast.*, **126**, 95–125.
- WECKNER, O., BRUNK, G., EPTON, M., SILLING, S. & ASKARI, E. (2009) Green's functions in non-local three-dimensional linear elasticity. *Proc. R. Soc. Lond. A Math. Phys. Eng. Sci.*, **53**, 705–728.
- ZHOU, K. & DU, Q. (2010) Mathematical and numerical analysis of linear peridynamic models with nonlocal boundary conditions. *SIAM J. Numer. Anal.*, **48**, 1759–1780.