

Very high resolution mapping of coral reef state using airborne bathymetric LiDAR surface-intensity and drone imagery

Antoine Collin, Camille Ramambason, Yves Pastol, Elisa Casella, Alessio Rovere, Lauric Thiault, Benoît Espiau, Gilles Siu, Franck Lerouvreur, Nao Nakamura, James L. Hench, Russell J. Schmitt, Sally J. Holbrook, Matthias Troyer & Neil Davies

To cite this article: Antoine Collin, Camille Ramambason, Yves Pastol, Elisa Casella, Alessio Rovere, Lauric Thiault, Benoît Espiau, Gilles Siu, Franck Lerouvreur, Nao Nakamura, James L. Hench, Russell J. Schmitt, Sally J. Holbrook, Matthias Troyer & Neil Davies (2018) Very high resolution mapping of coral reef state using airborne bathymetric LiDAR surface-intensity and drone imagery, International Journal of Remote Sensing, 39:17, 5676-5688, DOI: [10.1080/01431161.2018.1500072](https://doi.org/10.1080/01431161.2018.1500072)

To link to this article: <https://doi.org/10.1080/01431161.2018.1500072>



Published online: 24 Jul 2018.



Submit your article to this journal




Article views: 201



View Crossmark data



Citing articles: 2 [View citing articles](#)



Very high resolution mapping of coral reef state using airborne bathymetric LiDAR surface-intensity and drone imagery

Antoine Collin^{a,b}, Camille Ramambason^a, Yves Pastol^c, Elisa Casella^d, Alessio Rovere^{d,e}, Lauric Thiault^{b,f}, Benoît Espiau^f, Gilles Siu^f, Franck Lerouvreur^f, Nao Nakamura^{b,f}, James L. Hench^g, Russell J. Schmitt^h, Sally J. Holbrook^h, Matthias Troyer^{i,j} and Neil Davies^{k,l}

^aEcole Pratique des Hautes Etudes (EPHE), PSL Research University, Dinard, Brittany, France; ^bLABoratoire d'EXcellence CORAIL, Perpignan, France; ^cService Hydrographique et Océanographique de la Marine, Brest, Brittany, France; ^dZMT, Leibniz Centre for Tropical Marine Research, Bremen, Germany; ^eMarum, University of Bremen, Germany; ^fPSL Research University: EPHE-UPVD-CNRS, USR 3278 CRIOBE, Papetoai, Moorea, French Polynesia; ^gNicholas School of the Environment, Duke University, Beaufort, North Carolina, USA; ^hDepartment of Ecology, Evolution and Marine Biology, and Marine Science Institute, University of California Santa Barbara, Santa Barbara, California, USA; ⁱDepartment of Physics, Institute for Theoretical Physics, ETH Zurich, Zurich, Switzerland; ^jMicrosoft Quantum Research, Redmond, Washington, USA; ^kGump South Pacific Research Station, University of California, Moorea, French Polynesia; ^lBerkeley Institute for Data Science, University of California, Berkeley, California, USA

ABSTRACT

Very high resolution (VHR) airborne data enable detection and physical measurements of individual coral reef colonies. The bathymetric LiDAR system, as an active remote sensing technique, accurately computes the coral reef ecosystem's surface and reflectance using a single green wavelength at the decimetre scale over 1-to-100 km² areas. A passive multispectral camera mounted on an airborne drone can build a blue-green-red (BGR) orthorectified mosaic at the centimetre scale over 0.01-to-0.1 km² areas. A combination of these technologies is used for the first time here to map coral reef ecological state at the submeter scale. Airborne drone BGR values (0.03 m pixel size) serve to calibrate airborne bathymetric LiDAR surface and intensity data (0.5 m pixel size). A classification of five ecological states is then mapped through an artificial neural network (ANN). The classification was developed over a small area (0.01 km²) in the lagoon of Moorea Island (French Polynesia) at VHR (0.5 m pixel size) and then extended to the whole lagoon (46.83 km²). The ANN was first calibrated with 275 samples to determine the class of coral state through LiDAR-based predictors; then, the classification was validated through 135 samples, reaching a satisfactory performance (overall accuracy = 0.75).

ARTICLE HISTORY

Received 30 October 2017
Accepted 22 June 2018

1. Introduction

Coral reefs host 25% of the marine biodiversity but are increasingly subject to global ocean-climate changes and local anthropogenic activities. Fine-scale monitoring of coral reef ecosystems and associated ecosystem services are needed for their management and spatial planning. Coral reef mapping usually relies on remote sensing for cost-effectively identifying their structural complexity, benthic composition, and regime surrogates over large areas (Goodman, Samuel, and Stuart 2013; Hedley et al. 2016). Spaceborne multispectral imagery demonstrates great spatial potential to accurately map coral reef colonies (Collin, Hench, and Planes 2012), habitats (Collin et al. 2016), health (Collin and Planes 2012; Collin, Archambault, and Planes 2014) and resilience (Rowlands et al. 2012; Knudby et al. 2013; Collin, Nadaoka, and Bernardo 2015). Airborne passive hyperspectral imagery, provided with dozens of spectral bands, enables coral reef benthos, substrates, and bathymetry to be significantly improved (Leiper et al. 2014). Airborne (usually on manned aircraft) active light detection and ranging (LiDAR) is now the reference system for measuring bathymetry, outperforming waterborne sound detection and ranging (SoNAR) devices, which are strongly impeded by shallow features, specifically in the coastal realm where coral reefs thrive (Costa, Battista, and Pittman 2009). LiDAR-derived morphometry indices can reveal efficient proxies for ecosystem characteristics, for example, estimates of reef fish assemblages (Wedding et al. 2008). Yet despite the increase in discrimination power showed over benthic habitats bathed with turbid waters, LiDAR indices have not been used to date to exploit the spectral information associated with water-penetrating green LiDAR wavelength for coral reef monitoring (Collin, Archambault, and Long 2008, 2011; Collin, Long, and Archambault 2011).

Unmanned airborne vehicles (UAVs or simply 'drones') are becoming an integral component of the scientific toolbox for coral reef research and management. Equipped with blue-green-red (BGR) spectral cameras, drones are able to measure coral reef bathymetry and derived terrain roughness at very high resolution (VHR) using the photogrammetry approach (Leon et al. 2015; Casella et al. 2017). The 3D point cloud, permitting 2D orthorectified BGR mosaics and 2.5D digital surface models (DSM), results from the multi-angle information of a single scene made possible by the spatially-even acquisition of BGR imagery from a moving airborne drone flying at low altitude (from 30 to 150 m): so-called 'structure-from-motion'. The images and by-products yield spatial resolution at the centimetre scale (i.e. 0.03 m pixel size). Coral reef states can be significantly distinguished using the resulting 0.03 m BGR orthomosaic drone dataset, enabling classification of reef ecological states.

Here we describe a methodology for creating the first coral reef ecological state map at VHR based solely on regional airborne LiDAR 'predictors' trained with local 'response' imagery from the drone. The bathymetric LiDAR Riegl VQ-820-G, mounted on a small plane or helicopter, serves as the remotely sensed 1-to-100 km² predictors with four measurements of surface and intensity (green) per m². The BGR GoPro, mounted on a consumer-grade airborne drone (DJI phantom 2), is used as the remotely sensed 0.01-to-0.1 km² response. Spearheading machine learners in satellite-based coastal prediction (Collin, Etienne, and Feunteun 2017), an artificial neural network (ANN) classifier is developed to provide a robust, yet simple, algorithm linking the two datasets. Our study takes

place on one of the best-studied islands in the world (Cressey 2015): Moorea (French Polynesia, Figure 1), a volcanic island with fringing, barrier and outer coral reefs in the central South Pacific Ocean. It contributes to efforts to build a 4D model—an Island Digital Ecosystem Avatar (IDEA)—of Moorea and to simulate of future states of the social-ecological system in support of scenario-based planning (Davies et al. 2016). We follow a drone-based assessment of ecological state (coral reef state classification; Table 1) and combine it with LiDAR-based data to spatially classify the coral reef state at VHR over a small area and then extend this to the whole island. Findings are discussed with a view to how this approach could advance an automated workflow for coral reef mapping.

2. Materials and methods

2.1. Study site

The study site is located in the northern lagoon of Moorea Island ($17^{\circ}33'S$, $149^{\circ}50'W$) in the Society Archipelago (French Polynesia, Figure 1(a)). Moorea demonstrates a highly resilient coral reefs (Adjeroud et al. 2009), especially its outer slope, which following the extremely low coral cover (2%) due to 2007–2010 outbreak of corallivore crown-of-thorns sea star (*Acanthaster planci*) and 2010 Oli cyclone strike, is recovering to record rates close to 70% (Chancerelle, pers. comm.). Located inside the 46.83 km^2 Moorea lagoon, the study site covers $11\ 710\text{ m}^2$ with a maximum depth of 2 m. It is bathed in oligotrophic, thus clear, seawater including various taxa of reef-building corals (*Porites*, *Acropora*, *Pocillopora*, *Montipora*), red calcareous algae (*Lithothamnium*), fleshy algae (red, brown and green) and a diversity of geomorphic features (rubble, sand and pavement).

2.2. Drone visible response

A drone-based spectral survey (Figure 1(b)) was carried out on 17 August 2015 using a BGR camera (GoPro Hero 4) mounted on a consumer-grade drone (DJI Phantom 2). Calm sea and low sun elevation angle were optimal conditions for this survey. A series of 360 geolocated BGR photographs, acquired at 30 m altitude at nadir, were mosaicked then processed using the photogrammetry software Agisoft Photoscan (<http://www.agisoft.ru>). Constrained by nine ground control points and three scale bars, the resulting orthorectified mosaic (WGS 84 datum and UTM 6S projection) has 0.03 m resolution (see Casella et al. 2017 for further details) and was therefore deemed as precise enough to be used as air-truth (Figure 1(c), Collin, Lambert, and Etienne 2018). A total of 410 sampling points over the BGR orthomosaic, corresponding to as many LiDAR soundings, were visually interpreted by an expert and classified into five ecological states (Figure 2 (a) and Table 1), each one composed of 55 training and 27 validation sub-datasets.

2.3. Lidar surface and intensity predictors

The airborne LiDAR campaign was conducted from 10 to 26 June 2015 (one month before the drone flight) using a Riegl VQ-820-G hydrographic laser scanner mounted on a small plane. The sensor was operated at 251 kHz, providing a minimum sounding density of four points per m^2 (0.5 m) and vertical accuracy of 0.15 m, computed from




Figure 1. (a) Moorea Island (French Polynesia) was surveyed by bathymetric LiDAR at island scale (10–26 June 2015) and over a small study area by airborne drone (17 August 2015; red rectangle). (b) Natural-coloured (blue-green-red) drone survey provides spectral information at 0.03 m pixel size (2133×6095 pixels), enabling resolution of coral reef (c) assembled colonies, (d) single colonies on sand/pavement, or (e) anthropogenic features.

43,798 comparisons (Pastol, Chamberlain, and Sinclair 2016). This bathymetric LiDAR pulses an electromagnetic radiation (532 nm wavelength, namely green) from the aircraft and records its travel time in air and water by means of a waveform (Collin, Archambault, and Long 2008). LiDAR surface and intensity are computed on-the-fly for each sounding by converting the time between sea surface and bottom green echoes into distance (knowing the light speed into water), and by recording the peak of bottom green echo, respectively. Maximum depth ever recorded by bathymetric LiDAR reached 76.1 m in Moorea Island during the studied survey (Pastol, Chamberlain, and Sinclair 2016) given the water clarity due to oligotrophic waters. Since our Moorea study limits to the shallow waters (< 10 m depth), LiDAR intensity has been directly processed with no water correction. As each LiDAR surface and intensity sounding is duly located by the




Figure 2. Maps of the (a) natural-coloured imagery with 410 air-truth sampling sites (red transparent disks), (b) bathymetric LiDAR surface soundings, and (c) bathymetric LiDAR intensity (532 nm wavelength) soundings. (a) is at 0.03 m, whereas (b) and (c) are at 0.5 m pixel size.

combination of HR global navigation satellite system and inertial measurement unit, digital surface and intensity models (DSM and DIM, Figure 2(b, c)) can be calculated using ordinary kriging method applied to LiDAR sounding clouds. LiDAR points and rasters were geographically referenced to WGS84 UTM 6S and altimetrically zeroed as the mean sea level (SHOM 2016). Drone-derived imagery was registered with LiDAR data using a 1st degree polynomial function and resampled with cubic convolution.

2.4. Artificial neural network classification

Given their performance in a comparative analysis (Collin, Etienne, and Feunteun 2017), we use an ANN approach as a classification procedure binding the drone-based air-truth and LiDAR-based variables.

The ANN builds non-linear classifications by minimizing least squares using a multi-layer perceptron classifying ecological state response, $h(X)$ (Table 1) with the LiDAR surface and intensity predictors, X , through a constant, k , and intermediate weighted, w_i , functions called neurons, n_i (Heermann and Khazenie 1992):

$$h(X) = k \left(\sum_i w_i n_i(X) \right) \quad (1)$$

Table 1. Ecological description of the five coral reefscape states identified on airborne drone blue-green-red imagery (0.03 m spatial resolution) enabling a coral reef state classification to be created and colour-coded.

Drone-based state					
Ecological composition	<i>Acropora/ Pocillopora/ Montipora</i> stony corals	<i>Acropora/ Pocillopora/ Montipora</i> stony corals with red calcareous algae	<i>Porites</i> stony corals	Microalgae on rubble	Sand on pavement
Structural complexity	Very High roughness	High roughness	Medium roughness	Low roughness	Very low roughness
Coral reef state	1	2	3	4	5
Colour class					

Neurons n_i are hereinafter based on hyperbolic tangents. ANN constrained by a single hidden layer provided with two neurons so the number of neurons to be in synergy with the number of inputs (predictors, Figure 3). Trained by the 275 calibration samples, the ANN will be validated by the remaining 135 validation samples.




Figure 3. Conceptual flowchart explaining how the combination of LiDAR surface and intensity can predict the ecological state class through an intermediate hidden layer provided with two neurons.

2.5. Performance analysis

The agreement between validation and classified pixels in the five ecological states was quantified using the confusion matrix, from which overall, producer's and user's accuracies (OA, PA and UA, respectively) were computed (Congalton and Green 2009). PA and UA were calculated as the correctly classified pixels in each coral state divided by the number of calibration pixels of the corresponding state, and the total number of pixels that were classified in that state, respectively. OA was reckoned as the correctly classified pixels in all states divided by the total number of pixels.

3. Results

3.1. Local coral reef state at very high resolution

The OA of the ANN classification reached a satisfactory performance (OA = 0.75), showing that the dual combination of LiDAR surface and intensity variables had a robust explanatory power of the variability of coral reef states (Table 2). Contrary to coral reef states 1, 5 and 3 that were adequately assigned (UA = 0.84, 0.80 and 0.74, respectively), intermediate coral reef states 2 and 4 were moderately classified with UA of 0.68 and 0.68, respectively (Table 2). Contrary to UA statistics, PA measures were evenly correct (From 0.81 to 0.71, Table 2). The ANN classifier was applied to each pixel of LiDAR DSM and DIM (Figure 2(b, c), respectively) in order to continuously map ecological state (Figure 4(b)) provided with 0.5 m spatial resolution (142 × 422 pixels).

3.2. Moorea coral reef state at very high resolution

Insofar as the ANN prediction was adequate enough to be extended, the digital ecological classification was mapped at the island scale. Moorea LiDAR DSM and DIM were first rasterized at 0.5 m spatial resolution (Figure 5(a, b)) and then entered as inputs to the ANN classification, which produced a digital model of coral reef ecological state over the whole island (Figure 5(c), 40,364 × 34,588 pixels). Moorea classes are dominated by sand on pavement (56.8%), followed by *Porites* stony corals (14.1%) and Microalgae

Table 2. Confusion matrix synthesizing the quality of the artificial neural network classification applied to the independent 135 validation pixels (27 pixels per coral reef state).

STATE	Reference class					Total	UA
	1	2	3	4	5		
Classified class							
1	21	2	2	0	0	25	0.84
2	4	17	3	1	0	25	0.68
3	1	3	20	2	1	27	0.74
4	0	1	1	19	7	28	0.68
5	0	1	2	3	24	30	0.80
Total	26	24	28	25	32	135	
PA	0.80	0.71	0.71	0.76	0.75		




Figure 4. (a) Natural-coloured (blue-green-red) airborne drone orthomosaic (2133×6095 pixels, 0.03 m pixel size), along with (b) digital coral reef state classification based on the drone response, LiDAR surface and intensity predictors and two-neuroned artificial neural network classifier (142×422 pixels with 0.5 m pixel size).

on rubble (13.8%), then *Acropora/Pocillopora/Montipora* stony corals with red calcareous algae (10.9%), and finally *Acropora/Pocillopora/Montipora* stony corals (4.4%). Overall, the coverage of hard corals (from state 1 to 3) appears significantly greater in the leeward side than the windward side.

4. Discussion

4.1. Airborne drone as 'air-truth'

The five coral reef ecological states were based on VHR BGR orthorectified mosaic derived from a consumer-grade multispectral camera driven by an airborne drone. This innovative procedure is supported by our knowledge of in situ coral reef features that can be discriminated at the centimetre scale. Insofar as both ecological composition and structural complexity are easily deduced from the BGR dataset, relatively inexpensive drone deployment can be used to obtain air-truth data directly even in places with little technical capacity. The geolocated photographs can be remotely processed and analysed in the cloud, given a suitable internet connection. With an easy-to-implement flight planning mobile application, rapid surveys could be conducted at even very remote locations with little infrastructure/capacity after short-term events such as cyclone/storm and bleaching. The number of states could be increased by either flying




Figure 5. Digital (a) surface, (b) intensity (532 nm wavelength), and (c) coral reef state classification derived from bathymetric LiDAR soundings (40,364 × 34,588 pixels at 0.5 m pixel size).

at lower altitude (to gain in spatial resolution) or using drone-mounted LiDAR that can enhance the vertical accuracy, for example, to differentiate coral from macroalgae (Leiper et al. 2014).

The use of this air-truth, in the form of a cost-efficient UAV-borne BGR orthomosaic, has a strong potential to be applicable to other worldwide coral lagoons and even to a large panel of coastal and aquatic areas, provided with relatively clear waters. This air-

truth leverages a high ratio of covered space unit per time unit while collecting centimetre-scale data, considerably outperforming submerged acquisitions, hindered by the very high viscosity of water.

4.2. Airborne lidar surface and intensity

The gradient of ecological states (from 1, well-developed hard coral, to 5, sand) was positively correlated with both surface ($r = 0.93$) and intensity ($r = 0.93$), showing that coral coverage decreases with depth and LiDAR green reflectance. The coral shrinkage with depth can be explained by the coral growth and structural complexification towards the surface (as a photosynthetic symbiont), what corroborates results derived from a space-borne reef health proxy (Collin, Hench, and Planes 2012). The negative trend between coral state and green reflectance coincides with in situ spectral measurements, making explicit a greater reflectance of increasingly depigmented blue- and brown-mode coral reefs in the coral health chart (Leiper et al. 2009). This increase in green reflectance (decrease in green absorbance) is linked to the loss of peridinin pigments contained in symbiotic zooxanthellae living in coral tissues (Collin and Planes 2012). Even if most bathymetric LiDAR systems use the single green wavelength, this electromagnetic radiation is relevant to distinguish coral reef state as highlighted in the elaboration of both the green-purple and the 'red edge'-green normalized difference ratios (Collin, Hench, and Planes 2012; Collin, Archambault, and Planes 2014; respectively).

Inner classification results (UA) revealed that coral- and sand-dominant states (1, 3 and 5) were successfully recognized, contrary to both assemblages of corals and rubble colonized by calcareous and micro-algae (2 and 4), respectively. We could assume that the spectral mixing due to the presence of algae on relatively 'pure' states was not very effectively resolved by the ANN classifier built from only LiDAR surface and intensity. We advocate the experiment of a coral reef state classification using an innovative bathymetric LiDAR, augmented by an added spectral wavelength (i.e. 355 nm, as the third harmonic of the 1064-nm laser), likely to detect the coral fluorescence as well as intermediate states (Sasano et al. 2012).

4.3. Moorea coral reef states' spatial patterns

The coral reef state classification, spatially classified at VHR, is a strong asset to outline hotspots of healthy coral reefs, thus of associated biodiversity and ecosystem services. The centimetre and decimetre scales targeted in this study greatly enhance the spatial resolution of coral reefs' diagnoses and prognoses, surpassing other recent studies using object-based image analysis, which bottom at 2 m or 10 m (Phinn, Roelfsema, and Mumby 2012; Roelfsema et al. 2013). LiDAR-based spatially explicit classification, provided with decimetre sounding density over 100 km², offers an unpublished map of Moorea coral reefs' health. Two main spatial patterns emerged from the spatially explicit classification: westward polarization of healthy fringing reefs and northward polarization of healthy barrier reefs.

Wide healthy fringing reefs along west shorelines strongly contrast with thin ones along the eastern coast. This outstanding geographic difference is very susceptible to be the consequence of the dominant easterly winds (i.e. Southeast trade winds), which entail significantly

greater amounts of rain then carried sediment, which, in turn, deposit onto and stress coral colonies (Fabricius 2005), impeding the development of eastern fringing coral reefs.

More extended barrier reefs are obvious in the northern compared to the southern lagoon. This patterning might be explained by the two dominant swell systems originating from South: 40% SE and 25% SSW (Etienne 2012). Swell average height tends to be higher than 4 m during Austral winter, what creates, at the reef, significant wave height greater than 8 m (e.g. Teahupoo spot in Southern Tahiti Iti, Etienne 2012). The exposure to this high to very high energy flow hinders the efficient settlement of coral larvae and breaks the coral assemblage structure (Madin and Connolly 2006). This interpretation is corroborated by the third dominant swell system (22% NE, Etienne 2012), which constrains NE lagoon to exhibit slightly less extended barrier reefs compared to NW.

5. Conclusion

This original research has demonstrated that airborne bathymetric LiDAR data are able to reliably map five ecological states in coral reef systems at VHR over shallow, clear waters. Reef state information can be gleaned from an airborne drone equipped with a multispectral imaging sensor. Novel findings can be summarized as follows:

- (1) Coral reef state at the colony-scale (pixel size = 0.03 m) can be sourced from a BGR camera mounted on an airborne low-altitude drone;
- (2) LiDAR surface and intensity are powerful predictors of coral reef ecological state at the colony-scale (pixel size = 0.03 m);
- (3) ANN is an efficient classification approach to predict ecological state based only LiDAR surface and intensity (OA = 0.75);
- (4) LiDAR surface and intensity are powerful predictors of ecological state at the landscape scale (pixel size = 0.5 m);
- (5) Healthy fringing and barrier coral reefs in Moorea are located on the western and northern parts of the lagoon, respectively.

Acknowledgments

Authors gratefully thank *Service Hydrographique et Océanographique de la Marine* for the LiDAR acquisition control and the IDEA Consortium for sparking this collaborative research. This work was partly supported by French Polynesia government for LiDAR acquisition, the ETH Zurich for purchasing satellite imagery, and the National Science Foundation through the Moorea Coral Reef LTER (OCE-1236905 and 1637396) and Physical Oceanography (OCE-143133) programs. Two valuable referees and the editor are deeply acknowledged for the manuscript improvement.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This work was supported by the National Science Foundation [OCE-1236905,OCE-143133,OCE-1637396].

References

Adjeroud, M., F. Michonneau, P. J. Edmunds, Y. Chancerelle, T. L. De Loma, L. Penin, L. Thibaut, J. Vidal-Dupiol, B. Salvat, and R. Galzin. 2009. "Recurrent Disturbances, Recovery Trajectories, and Resilience of Coral Assemblages on a South Central Pacific Reef." *Coral Reefs* 28 (3): 775–780. doi:10.1007/s00338-009-0515-7.

Casella, E., A. Collin, D. Harris, S. Ferse, S. Bejarano, V. Parravicini, J. L. Hench, and A. Rovere. 2017. "Mapping Coral Reefs Using Consumer-Grade Drones and Structure from Motion Photogrammetry Techniques." *Coral Reefs* 36 (1): 269–275. doi:10.1007/s00338-016-1522-0.

Collin, A., B. Long, and P. Archambault. 2011. "Benthic Classifications Using Bathymetric LiDAR Waveforms and Integration of Local Spatial Statistics and Textural Features." *Journal of Coastal Research* 62: 86–98. doi:10.2112/SI_62_9.

Collin, A., J. Laporte, B. Koetz, F. R. Martin-Lauzer, and Y. L. Desnos. 2016. "Mapping Bathymetry, Habitat, and Potential Bleaching of Coral Reefs Using Sentinel-2." In *Proceedings of the 13th International Coral Reef Symposium*, Honolulu, 373–387.

Collin, A., J. L. Hench, and S. Planes. 2012. "A Novel Spaceborne Proxy for Mapping Coral Cover." In *Proceedings of the 12th International Coral Reef Symposium*, Cairns, 1–5.

Collin, A., K. Nadaoka, and L. Bernardo. 2015. "Mapping the Socio-Economic and Ecological Resilience of Japanese Coral Reefs across a Decade." *ISPRS International Journal of Geo-Information* 4 (2): 900–927. doi:10.3390/ijgi4020900.

Collin, A., N. Lambert, and S. Etienne. 2018. "Satellite-Based Salt Marsh Elevation, Vegetation Height, and Species Composition Mapping Using the Superspectral WorldView-3 Imagery." *International Journal of Remote Sensing* 1–19. doi:10.1080/01431161.2018.1466084.

Collin, A., P. Archambault, and B. Long. 2008. "Mapping the Shallow Water Seabed Habitat with the SHOALS." *IEEE Transactions on Geoscience and Remote Sensing* 46 (10): 2947–2955. doi:10.1109/TGRS.2008.920020.

Collin, A., P. Archambault, and B. Long. 2011. "Predicting Species Diversity of Benthic Communities within Turbid Nearshore Using Full-Waveform Bathymetric LiDAR and Machine Learners." *PLoS one* 6 (6): e21265. doi:10.1371/journal.pone.0021265.

Collin, A., P. Archambault, and S. Planes. 2014. "Revealing the Regime of Shallow Coral Reefs at Patch Scale by Continuous Spatial Modeling." *Frontiers in Marine Science* 1: 65. doi:10.3389/fmars.2014.00065.

Collin, A., S. Etienne, and E. Feunteun. 2017. "VHR Coastal Bathymetry Using WorldView-3: Colour versus Learner." *Remote Sensing Letters* 8 (11): 1072–1081. doi:10.1080/2150704X.2017.1354261.

Collin, A., and S. Planes. 2012. "Enhancing Coral Health Detection Using Spectral Diversity Indices from Worldview-2 Imagery and Machine Learners." *Remote Sensing* 4 (10): 3244–3264. doi:10.3390/rs4103244.

Congalton, R. G., and K. Green. 2009. *Assessing the Accuracy of Remotely Sensed Data: Principles and Practices*. Boca Raton, FL: CRC /Taylor Francis press.

Costa, B. M., T. A. Battista, and S. J. Pittman. 2009. "Comparative Evaluation of Airborne LiDAR and Ship-Based Multibeam SoNAR Bathymetry and Intensity for Mapping Coral Reef Ecosystems." *Remote Sensing of Environment* 113 (5): 1082–1100. doi:10.1016/j.rse.2009.01.015.

Cressey, D. 2015. "Tropical Paradise Inspires Virtual Ecology Lab." *Nature* 517: 255–256. doi:10.1038/517255a.

Davies, N., D. Field, D. Gavaghan, S. J. Holbrook, S. Planes, M. Troyer, M. Bonsall, et al. 2016. "Simulating Social-Ecological Systems: The Island Digital Ecosystem Avatars (IDEA) Consortium." *Gigascience* 5: 14. doi:10.1186/s13742-016-0118-5.

Etienne, S. 2012. "Marine Inundation Hazards in French Polynesia: Geomorphic Impacts of Tropical Cyclone Oli in February 2010." *Geological Society, London, Special Publications* 361 (1): 21–39. doi:10.1144/SP361.4.

Fabricius, K. E. 2005. "Effects of Terrestrial Runoff on the Ecology of Corals and Coral Reefs: Review and Synthesis." *Marine Pollution Bulletin* 50 (2): 125–146. doi:10.1016/j.marpolbul.2004.11.028.

Goodman, J. A., J. P. Samuel, and R. P. Stuart. 2013. *Coral Reef Remote Sensing. A Guide for Mapping, Monitoring and Management*. Netherlands: Springer.

Hedley, J. D., C. M. Roelfsema, I. Chollett, A. R. Harborne, S. F. Heron, S. Weeks, W. J. Skirving, et al. 2016. "Remote Sensing of Coral Reefs for Monitoring and Management: A Review." *Remote Sens* 8: 118. doi:10.3390/rs8020118.

Heermann, P. D., and N. Khazenie. 1992. "Classification of Multispectral Remote Sensing Data Using a Back-Propagation Neural Network." *IEEE Transactions on Geoscience and Remote Sensing* 30 (1): 81–88. doi:10.1109/36.124218.

Knudby, A., S. Jupiter, C. Roelfsema, M. Lyons, and S. Phinn. 2013. "Mapping Coral Reef Resilience Indicators Using Field and Remotely Sensed Data." *Remote Sensing* 5 (3): 1311–1334. doi:10.3390/rs5031311.

Leiper, I. A., S. R. Phinn, C. M. Roelfsema, K. E. Joyce, and A. G. Dekker. 2014. "Mapping Coral Reef Benthos, Substrates, and Bathymetry, Using Compact Airborne Spectrographic Imager (CASI) Data." *Remote Sensing* 6 (7): 6423–6445. doi:10.3390/rs6076423.

Leiper, I. A., U. E. Siebeck, N. J. Marshall, and S. R. Phinn. 2009. "Coral Health Monitoring: Linking Coral Colour and Remote Sensing Techniques." *Canadian Journal of Remote Sensing* 35 (3): 276–286. doi:10.5589/m09-016.

Leon, J. X., C. M. Roelfsema, M. I. Saunders, and S. R. Phinn. 2015. "Measuring Coral Reef Terrain Roughness Using 'Structure-From-Motion' Close-Range Photogrammetry." *Geomorphology* 242: 21–28. doi:10.1016/j.geomorph.2015.01.030.

Madin, J. S., and S. R. Connolly. 2006. "Ecological Consequences of Major Hydrodynamic Disturbances on Coral Reefs." *Nature* 444 (7118): 477–480. doi:10.1038/nature05328.

Pastor, Y., L. Chamberlain, and M. Sinclair. 2016. "Airborne Bathymetric LiDAR and Coastal Zone Management in French Polynesia." In *Proc. of International Federation of Surveyors (FIG) Working Week*, Christchurch, May 2–6.

Phinn, S. R., C. M. Roelfsema, and P. J. Mumby. 2012. "Multi-Scale, Object-Based Image Analysis for Mapping Geomorphic and Ecological Zones on Coral Reefs." *International Journal of Remote Sensing* 33 (12): 3768–3797. doi:10.1080/01431161.2011.633122.

Roelfsema, C., S. Phinn, S. Jupiter, J. Comley, and S. Albert. 2013. "Mapping Coral Reefs at Reef to Reef-System Scales, 10s–1000s Km², Using Object-Based Image Analysis." *International Journal of Remote Sensing* 34 (18): 6367–6388. doi:10.1080/01431161.2013.800660.

Rowlands, G., S. Purkis, B. Riegl, L. Metsamaa, A. Bruckner, and P. Renaud. 2012. "Satellite Imaging Coral Reef Resilience at Regional Scale. A Case-Study from Saudi Arabia." *Marine Pollution Bulletin* 64 (6): 1222–1237. doi:10.1016/j.marpolbul.2012.03.003.

Sasano, M., H. Yamanouchi, A. Matsumoto, N. Kiriya, K. Hitomi, and K. Tamura. 2012. "Development of Boat-Based Fluorescence Imaging Lidar for Coral Monitoring." In *Proc. of 12th Internaitonal Coral Reef Symposium*, Cairns, 5A–7.

Service Hydrographique et Océanographique de la Marine (SHOM). 2016. *Lidar Polynésie française 2015 Produit Moorea SAU V*. 20160630. Brest, France: SHOM press.

Wedding, L. M., A. M. Friedlander, M. McGranaghan, R. S. Yost, and M. E. Monaco. 2008. "Using Bathymetric Lidar to Define Nearshore Benthic Habitat Complexity: Implications for Management of Reef Fish Assemblages in Hawaii." *Remote Sensing of Environment* 112 (11): 4159–4165. doi:10.1016/j.rse.2008.01.025.