

On the Global Dynamics of an Electroencephalographic Mean Field Model of the Neocortex*

Farshad Shirani[†], Wassim M. Haddad[‡], and Rafael de la Llave[‡]

Abstract. This paper investigates the global dynamics of a mean field model of the electroencephalogram developed by Liley, Cadusch, and Dafilis [*Network*, 13 (2002), pp. 67–113]. The model is presented as a system of coupled ordinary and partial differential equations with periodic boundary conditions. Existence, uniqueness, and regularity of weak and strong solutions of the model are established in appropriate function spaces, and the associated initial-boundary value problems are proved to be well-posed. Sufficient conditions are developed for the phase spaces of the model to ensure nonnegativity of certain quantities in the model, as required by their biophysical interpretation. It is shown that the semigroups of weak and strong solution operators possess bounded absorbing sets for the entire range of biophysical values of the parameters of the model. Challenges involved in establishing a global attractor for the model are discussed and it is shown that there exist parameter values for which the constructed semidynamical systems do not possess a compact global attractor due to the lack of the compactness property. Finally, using the theoretical results of the paper, instructive insights are provided into the complexity of the behavior of the model and computational analysis of the model.

Key words. mathematical neuroscience, neocortical dynamics, mean field model, evolution equations, global attractors

AMS subject classifications. 35Q92, 37L30, 35B40, 92C20

DOI. 10.1137/16M1098577

1. Introduction. Inspired by the seminal work of Alan Hodgkin and Andrew Huxley on modeling the flow of ionic currents through the membrane of a giant nerve fiber, numerous biophysical and mathematical models have been developed towards understanding the neurophysiology of the central nervous system and the underlying mechanism of the various phenomena that emerge during its vital operation in the body, many of which still remain a mystery to researchers [17, 26, 42, 54]. In particular, in exploring the core component of the central nervous system—the brain—substantial effort has been devoted to developing models at different levels of scope, from the *molecular and intercellular* level, dealing with the transportation of ions and the enzymatic kinetics of neurotransmitter-receptor binding at ion channels, to the *single cell and intracellular* level, dealing with the creation and transmission of action potential, to the *population and neuronal network* level, dealing with the average

*Received by the editors October 12, 2016; accepted for publication (in revised form) by J. van den Berg May 15, 2017; published electronically October 24, 2017.

<http://www.siam.org/journals/siads/16-4/M109857.html>

Funding: The work of the third author was supported in part by NSF grant DMS 1500943.

[†]School of Aerospace Engineering, Georgia Institute of Technology, Atlanta, GA 30332 (f.shirani@gatech.edu, wm.haddad@aerospace.gatech.edu).

[‡]School of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332 (rafael.delallave@math.gatech.edu).

behavior and synchronized activity of neuronal ensembles, to the *system level*, dealing with the systematic operation and interaction between cortical and subcortical components of the brain, and finally to the *behavioral and cognitive* level, dealing with the integrated mental activity and the creation of mind [1, 15, 22, 29, 30, 46, 48, 55].

As an effective methodology for developing models at the population and network level, mean field theory has been employed to construct approximate models for interconnected populations of neurons by averaging the effect of all other neurons on a given individual neuron inside a population. The resulting *averaged neuron* can be used to analyze the overall temporal behavior of a single population of neurons—leading to a *neural mass* model—or can be considered as a locally averaged component of a continuum of neural populations—leading to a spatio-temporal *mean field* model. These models are particularly useful in analyzing the electrophysiological activity of neuronal ensembles using local field potentials and electroencephalograms (EEGs) [10, 40, 43, 45].

The evolution equations that describe a mean field model of neural activity in the cortex are in the form of a system of partial differential equations (PDEs), or a system of coupled ordinary and partial differential equations. The theory of infinite-dimensional dynamical systems is hence used to analyze the global dynamics and long-term behavior of these systems. The classical approach to this problem follows several steps. First, existence, uniqueness, and regularity of solutions are established for all positive time in appropriately chosen problem-dependent function spaces, and the well-posedness of the problem is confirmed. Second, a semidynamical framework is constructed over a positively invariant complete normed space—the phase space for the evolution of the solutions—and is shown to possess bounded absorbing sets. Asymptotic compactness of the semigroup of solution operators is then ensured to guarantee the existence of a global attractor, which is a compact strictly invariant attracting set and which contains all the information regarding the asymptotic behavior of the model. Third, the Hausdorff or fractal dimension of the global attractor is estimated to show that the attractor is finite-dimensional, so that the asymptotic dynamics of the system is determined by a finite number of degrees of freedom. Fourth, the existence of an inertial manifold is established, which is a smooth finite-dimensional invariant manifold containing the global attractor. Consequently, the dynamics on the attractor can be presented by a finite set of ordinary differential equations (ODEs) and further characterized to give the overall picture of the long-term behavior of the system [7, 24, 25, 44, 51].

In this paper, we investigate the mean field model proposed in [36] for an understanding of the electrical activity in the neocortex as observed in the EEG. This model, which is comprised of a system of coupled ordinary and partial differential equations in a two-dimensional space, has been widely used in the literature to study the alpha- and gamma-band rhythmic activity in the cortex [4, 5], phase transition and burst suppression in cortical neurons during general anesthesia [6, 37, 49], the effect of anesthetic drugs on the EEG [2, 19], and epileptic seizures [31, 32, 33, 35]. Open-source tools for numerical implementation of the model and computation of equilibria and time-periodic solutions are developed in [23]. Complexity of the dynamics of the model, including periodic and pseudoperiodic solutions, chaotic behavior, multistability, and bifurcation are studied in [11, 12, 13, 20, 21, 52, 53].

The above results, however, are mainly computational or use approximate versions of the model. A rigorous analysis of the dynamics of the model in an infinite-dimensional dynamical

system framework as outlined above is not available in the literature. In particular, the basic problems of well-posedness of the initial-boundary value problem associated with the model and regularity of the solutions remain uninvestigated. It is not known under what conditions, if any, the components of the solutions of the model that are associated with nonnegative biophysical quantities remain nonnegative for all time. The solutions that take negative values for such quantities—even for a small interval of time in the distant future—cannot represent a biophysically plausible dynamics of the electrical activity in the neocortex.

The aim of this paper is to study the global dynamics of the mean field model discussed above, to ensure its biophysical plausibility, and to provide the basic analytical results required for characterization of the long-term dynamics of the model. Specifically, we follow the first two steps of the classical analysis approach to investigate the problem of existence or nonexistence of a global attractor.

This paper is organized as follows. In section 2, we introduce notation and recall key definitions that are necessary for developing the results in this paper. In section 3, we give a description of the anatomical structure of the neocortex and the physiological interactions that underlie the construction of the model. Moreover, we present the mathematical structure of the model as a system of coupled ordinary and partial differential equations with initial values and periodic boundary conditions. In section 4, following the first step of the classical analysis approach, we prove the existence and uniqueness of weak and strong solutions for the proposed initial value problem and analyze the regularity of these solutions.

As in the second step of the classical analysis approach, in section 5 we define semigroups of weak and strong solution operators and show their continuity properties. Moreover, we establish sufficient conditions on the phase spaces as to ensure biophysical plausibility of the evolution of the solutions under the associated semidynamical systems. In section 6, we show that the semigroups of solution operators possess bounded absorbing sets for all possible values of the biophysical parameters of the model. In section 7, we discuss challenges involved in establishing a global attractor for the model, and in particular, we show that there exist sets of values for the biophysical parameters of the model such that the associated semigroups of solution operators do not possess a compact global attractor. We conclude the paper in section 8 with a discussion on the results developed in the paper and their application to computational analysis of the model.

2. Notation and preliminaries. The notation used in this paper is fairly standard. Specifically, $\mathbb{R}^n$ denotes the n -dimensional real Euclidean space and $\mathbb{R}^{m \times n}$ denotes the space of real $m \times n$ matrices. A point $x \in \mathbb{R}^n$ is presented by the n -tuple $x = (x_1, \dots, x_n)$ or, when it appears in matrix operations, by the column vector $x = \begin{bmatrix} x_1 & \cdots & x_n \end{bmatrix}^T$, where $(\cdot)^T$ denotes the transpose. The nonnegative cone $\{x \in \mathbb{R}^n : x_j \geq 0 \text{ for } j = 1, \dots, n\}$ is denoted by $\mathbb{R}_+^n$. A sequence of points in $\mathbb{R}^n$ is denoted by $\{x^{(l)}\}_{l=1}^\infty$, with the j th component of $x^{(l)}$ denoted by $x_j^{(l)}$. Moreover, the trace of a square matrix $A \in \mathbb{R}^{n \times n}$ is denoted by $\text{tr } A$, and a block-diagonal matrix D with k blocks $D_1, \dots, D_k$ is denoted by $\text{diag}(D_1, \dots, D_k)$. For $x, y \in \mathbb{R}^n$, we write $x \geq y$ to denote componentwise inequality, that is, $x_j \geq y_j$, $j = 1, \dots, n$. For $A, B \in \mathbb{R}^{n \times n}$, we write $A \geq B$ to denote that $A - B$ is positive semidefinite. Finally, we denote by $0_{n \times n}$ and $I_{n \times n}$ the zero and identity matrices in $\mathbb{R}^{n \times n}$, respectively. We write I for the identity operator in other vector spaces.

For an inner product space $\mathcal{U}$, we denote the associated inner product by $(\cdot, \cdot)_{\mathcal{U}}$ and the norm generated by the inner product by $\|\cdot\|_{\mathcal{U}}$. For a Hilbert space $\mathcal{U}$, we denote the pairing of $\mathcal{U}$ with its dual space $\mathcal{U}^*$ by $\langle \cdot, \cdot \rangle_{\mathcal{U}}$. In particular, for $\mathcal{U} = \mathbb{R}^n$, we write $(\cdot, \cdot)_{\mathbb{R}^n}$ and $\|\cdot\|_{\mathbb{R}^n}$ for the standard inner product and the Euclidean norm, respectively. Similarly, for $\mathcal{U} = \mathbb{R}^{m \times n}$, we write $(\cdot, \cdot)_{\mathbb{R}^{m \times n}}$ for the standard inner product and $\|\cdot\|_{\mathbb{R}^{m \times n}}$ for the associated inner product norm. Moreover, we denote the vector 1-, 2-, and ∞ -norms in $\mathbb{R}^n$ by $\|\cdot\|_1$, $\|\cdot\|_2 = \|\cdot\|_{\mathbb{R}^n}$, and $\|\cdot\|_{\infty}$, respectively. The matrix 1-, 2-, and ∞ -norms in $\mathbb{R}^{m \times n}$ induced, respectively, by the vector 1-, 2-, and ∞ -norms in $\mathbb{R}^n$ are denoted by $\|\cdot\|_1$, $\|\cdot\|_2$, and $\|\cdot\|_{\infty}$.

Let Ω be an open subset of $\mathbb{R}^n$ denoting the *space* domain of a given dynamical system, with $x \in \Omega$ denoting a *spatial* point in Ω . The *time* domain of the system is given by the closed interval $[0, T] \subset \mathbb{R}$, $T > 0$, with the *temporal* point t . For a function $u : [0, T] \rightarrow \mathbb{R}$, the k th-order total derivative with respect to t at t_0 is denoted by $d_t^k u(t_0)$. For $k = 1$, we write $d_t u(t_0)$. For a function $u(x, t) : \Omega \times [0, T] \rightarrow \mathbb{R}$, the k th-order partial derivative with respect to t at (x_0, t_0) is denoted by $\partial_t^k u(x_0, t_0)$ and the k th-order partial derivative with respect to x_j at (x_0, t_0) is denoted by $\partial_{x_j}^k u(x_0, t_0)$, $j = 1, \dots, n$. For $k = 1$, we write $\partial_t u(x_0, t_0)$ and $\partial_{x_j} u(x_0, t_0)$. The gradient of u in Ω is denoted by $\partial_x u$ and is given by $\partial_x u := (\partial_{x_1} u, \dots, \partial_{x_n} u)$. The Laplacian of u in Ω is denoted by Δu and is given by $\Delta u := (\partial_{x_1}^2 + \dots + \partial_{x_n}^2)$. For a vector-valued function $u(x, t) : \Omega \times [0, T] \rightarrow \mathbb{R}^m$, we interpret $u(x, t)$ as the m -tuple $u(x, t) = (u_1(x, t), \dots, u_m(x, t))$, where each component $u_j(x, t)$, $j = 1, \dots, m$, is a scalar-valued function on $\Omega \times [0, T]$. In this case, $\partial_x u(x, t) \in \mathbb{R}^{m \times n}$ is the gradient of u and the vector Laplacian Δu is given by $\Delta u(x, t) := (\Delta u_1(x, t), \dots, \Delta u_m(x, t)) \in \mathbb{R}^m$, assuming Cartesian coordinates.

For every integer $k \geq 0$, the space of k -times continuously differentiable real-valued functions on Ω is denoted by $C^k(\Omega)$. The space $C^k(\overline{\Omega})$ consists of all functions in $C^k(\Omega)$ that, together with all of their partial derivatives up to the order k , are uniformly continuous in bounded subsets of Ω . Moreover, for $0 < \lambda \leq 1$, the Hölder space $C^{k, \lambda}(\overline{\Omega})$ is a subspace of $C^k(\overline{\Omega})$ consisting of functions whose partial derivatives of order k are Hölder continuous with exponent λ ; see [9, sect. 1.18] for details. We use $C_c^\infty(\Omega)$ to denote the space of infinitely differentiable real-valued functions with compact support in Ω . Moreover, we denote by $L_{\text{loc}}^1(\Omega)$ the space of locally integrable real-valued functions on Ω . Then, for every function $u \in L_{\text{loc}}^1(\Omega)$ and any multi-index α with $|\alpha| \geq 1$, the *weak partial derivative* of u in $L_{\text{loc}}^1(\Omega)$, of order $|\alpha|$, is defined by the distribution u^α that satisfies

$$\int_{\Omega} u^\alpha \phi \, dx = (-1)^{|\alpha|} \int_{\Omega} u \partial^\alpha \phi \, dx \quad \text{for all } \phi \in C_c^\infty(\Omega),$$

where $dx = dx_1 \cdots dx_n$ is the Lebesgue measure on $\mathbb{R}^n$; see [9, sect. 6.3] for details. With a minor abuse of notation, we use ∂_t^k and ∂_x^k to denote the k th-order weak—as well as classical—partial derivatives with respect to t and x , respectively. The distinction will be clear from the context, or will otherwise be explicitly specified.

The Hilbert space of vector-valued Lebesgue measurable functions $u : \Omega \rightarrow \mathbb{R}^m$ with finite L^2 -norm is denoted by $L^2(\Omega; \mathbb{R}^m)$, with the associated inner product and norm given by

$$(u, v)_{L^2(\Omega; \mathbb{R}^m)} := \int_{\Omega} (u(x), v(x))_{\mathbb{R}^m} \, dx, \quad \|u\|_{L^2(\Omega; \mathbb{R}^m)} := \left[\int_{\Omega} \|u(x)\|_{\mathbb{R}^m}^2 \, dx \right]^{\frac{1}{2}}.$$

The Banach space of vector-valued Lebesgue measurable functions $u : \Omega \rightarrow \mathbb{R}^m$ with finite L^∞ -norm is denoted by $L^\infty(\Omega; \mathbb{R}^m)$, with the norm

$$\|u\|_{L^\infty(\Omega; \mathbb{R}^m)} := \operatorname{ess\,sup}_{x \in \Omega} \|u(x)\|_\infty.$$

The Sobolev space of vector-valued functions $u \in L^p(\Omega; \mathbb{R}^m)$, whose all l th-order weak derivatives $\partial_x^l u$, $l \leq k$, exist and belong to $L^p(\Omega; \mathbb{R}^{m \times n^l})$, is denoted by $W^{k,p}(\Omega; \mathbb{R}^m)$. When $p = 2$, the Sobolev spaces $W^{k,2}(\Omega; \mathbb{R}^m)$ are Hilbert spaces for all $k \in [0, \infty)$, and are denoted by $H^k(\Omega; \mathbb{R}^m) := W^{k,2}(\Omega; \mathbb{R}^m)$. Specifically, $H^0(\Omega; \mathbb{R}^m) = L^2(\Omega; \mathbb{R}^m)$, and $H^1(\Omega; \mathbb{R}^m)$ is a Hilbert space with the inner product

$$(u, v)_{H^1(\Omega; \mathbb{R}^m)} := (u, v)_{L^2(\Omega; \mathbb{R}^m)} + (\partial_x u, \partial_x v)_{L^2(\Omega; \mathbb{R}^{m \times n})}.$$

Moreover, $H^2(\Omega; \mathbb{R}^m)$ is a Hilbert space with the inner product

$$(u, v)_{H^2(\Omega; \mathbb{R}^m)} := (u, v)_{L^2(\Omega; \mathbb{R}^m)} + (\partial_x u, \partial_x v)_{L^2(\Omega; \mathbb{R}^{m \times n})} + (\partial_x^2 u, \partial_x^2 v)_{L^2(\Omega; \mathbb{R}^{m \times n^2})}.$$

Let $\Omega = (0, \omega_1) \times \cdots \times (0, \omega_n)$, $\omega_j > 0$, $j = 1, \dots, n$, be an open rectangle in $\mathbb{R}^n$. A function $u : \mathbb{R}^n \rightarrow \mathbb{R}$ is called Ω -periodic if it is periodic in each direction, that is,

$$u(x + \omega_j e_j) = u(x), \quad j = 1, \dots, n, \quad x \in \mathbb{R}^n,$$

where e_j is the unit vector in the j th direction. Define the space $C_{\text{per}}^\infty(\Omega)$ as the restriction to Ω of the space of infinitely differentiable Ω -periodic functions. Then, the Sobolev space $H_{\text{per}}^k(\Omega)$, $k \geq 0$, is defined by the completion of $C_{\text{per}}^\infty(\Omega)$ in $H^k(\Omega)$; see [44, Def. 5.37] or, for an equivalent definition, [51, p. 50]. A vector-valued function $u : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is Ω -periodic if each of its components $u_j : \mathbb{R}^n \rightarrow \mathbb{R}$, $j = 1, \dots, m$, is Ω -periodic. The spaces $C_{\text{per}}^\infty(\Omega; \mathbb{R}^m)$ and $H_{\text{per}}^k(\Omega; \mathbb{R}^m)$ are then defined accordingly. It follows from Green's formula that

$$\begin{aligned} (1) \quad & (-\Delta u, v)_{L_{\text{per}}^2(\Omega; \mathbb{R}^m)} = (\partial_x u, \partial_x v)_{L_{\text{per}}^2(\Omega; \mathbb{R}^{m \times n})}, \\ & ((-\Delta + I)u, v)_{L_{\text{per}}^2(\Omega; \mathbb{R}^m)} = (u, v)_{H_{\text{per}}^1(\Omega; \mathbb{R}^m)}, \\ & (-\Delta u, (-\Delta + I)u)_{L_{\text{per}}^2(\Omega; \mathbb{R}^m)} = \|u\|_{H_{\text{per}}^2(\Omega; \mathbb{R}^m)}^2 - \|u\|_{L_{\text{per}}^2(\Omega; \mathbb{R}^m)}^2, \\ & \|(-\Delta + I)u\|_{L_{\text{per}}^2(\Omega; \mathbb{R}^m)}^2 = \|u\|_{H_{\text{per}}^2(\Omega; \mathbb{R}^m)}^2 + \|\partial_x u\|_{L_{\text{per}}^2(\Omega; \mathbb{R}^{m \times n})}^2 \\ & \quad = \|u\|_{H_{\text{per}}^1(\Omega; \mathbb{R}^m)}^2 + \|\partial_x u\|_{H_{\text{per}}^1(\Omega; \mathbb{R}^{m \times n})}^2. \end{aligned}$$

In this paper, we interchangeably view the function $u(x, t)$, $x \in \Omega$, $t \in [0, T]$, as a composite function of x and t , as well as a mapping u of t to a function of x , that is,

$$[u(t)](x) := u(x, t), \quad x \in \Omega, \quad t \in [0, T].$$

With a minor abuse of notation, the same symbol is used to denote both the original form of the function and the mapping. The distinction becomes evident in the way we define the space of such mappings or, equivalently, Banach space-valued functions; see, for example,

[16, App. E.5]. For a Banach space $\mathcal{U}$, the space $L^2(0, T; \mathcal{U})$ is composed of all strongly measurable Banach space-valued functions $u : [0, T] \rightarrow \mathcal{U}$ with the finite L^2 -norm defined by

$$\|u\|_{L^2(0, T; \mathcal{U})} := \left[\int_0^T \|u(t)\|_{\mathcal{U}}^2 dt \right]^{\frac{1}{2}}.$$

The space $C^0([0, T]; \mathcal{U})$ is composed of all continuous Banach space-valued functions $u : [0, T] \rightarrow \mathcal{U}$ with the finite uniform norm defined by

$$\|u\|_{C^0([0, T]; \mathcal{U})} := \max_{t \in [0, T]} \|u(t)\|_{\mathcal{U}}.$$

Accordingly, the spaces $C^k([0, T]; \mathcal{U})$ and $C^{k, \lambda}([0, T]; \mathcal{U})$, $k \geq 0$, $0 < \lambda \leq 1$, are defined as the space of k -times continuously differentiable Banach space-valued functions and its Hölder-continuous subspace. The Sobolev spaces $H^k(0, T; \mathcal{U})$, $k \geq 0$, are composed of all functions $u \in L^2(0, T; \mathcal{U})$, whose l th-order weak derivatives $d_t^l u$ exist for $l \leq k$ and belong to $L^2(0, T; \mathcal{U})$. In particular, for $k = 1$, we have

$$\|u\|_{H^1(0, T; \mathcal{U})} := \left[\int_0^T (\|u(t)\|_{\mathcal{U}}^2 + \|d_t u(t)\|_{\mathcal{U}}^2) dt \right]^{\frac{1}{2}}.$$

For further details on these spaces, see [16, sect. 5.9.2] and [44, sect. 7.1].

When $P : \mathcal{U} \rightarrow \mathcal{Y}$ is a mapping between the Banach spaces $\mathcal{U}$ and $\mathcal{Y}$, we denote the k th-order Fréchet derivative of P at u_0 by $d_u P(u_0)$. The space $C^k(\mathcal{U}; \mathcal{Y})$ is then composed of all k -times continuously differentiable mappings from $\mathcal{U}$ into $\mathcal{Y}$. For a mapping $P : \mathcal{U}_1 \times \cdots \times \mathcal{U}_m \rightarrow \mathcal{Y}$, where $\mathcal{Y}$ and $\mathcal{U}_j$, $j = 1, \dots, m$, are Banach spaces, $\partial_{u_j} P(u_0)$ is the j th partial Fréchet derivative of P at $u_0 = (u_{01}, \dots, u_{0m})$. The gradient of P at u_0 is then written as $\partial_u P(u_0)$; see [9, sect. 7.1] for details.

Finally, we denote the symmetric difference of two sets $\mathcal{X}$ and $\mathcal{Y}$ by $\mathcal{X} \triangle \mathcal{Y}$. In a topological space $\mathcal{X}$, we denote the closure of a set $\mathcal{X} \subset \mathcal{X}$ by $\overline{\mathcal{X}}$, its interior by $\mathcal{X}^\circ$, and its boundary by $\partial \mathcal{X}$. The characteristic function of $\mathcal{X}$ is denoted by $\chi_{\mathcal{X}}$. When $\mathcal{X}$ is a measure space, $|\mathcal{X}|$ denotes the measure of the set $\mathcal{X} \subset \mathcal{X}$. For normed vector spaces $\mathcal{X}$ and $\mathcal{Y}$, we write $\mathcal{X} \hookrightarrow \mathcal{Y}$ for continuous embedding of $\mathcal{X}$ in $\mathcal{Y}$, and $\mathcal{X} \Subset \mathcal{Y}$ for compact embedding of $\mathcal{X}$ in $\mathcal{Y}$; see [9, sect. 6.6] for details. When $\mathcal{X}$ is a metric space and the topology on $\mathcal{X}$ is induced by the given metric, $B(x, R)$ denotes the open ball centered at $x \in \mathcal{X}$ with radius $R > 0$, which is a basis element for the topology. For every bounded measurable set in $\mathcal{X}$, and in particular for $B(x, R)$, we denote by $f_{B(x, R)}$ the averaging operator over $B(x, R)$, that is, $f_{B(x, R)} := \frac{1}{|B(x, R)|} \int_{B(x, R)}$.

3. Model description. The neocortex has a layered columnar structure consisting mostly of six distinct layers. Neurons in the neocortex are organized in vertical columns, usually referred to as *cortical columns* or *macrocolumns*, which are a fraction of a millimeter wide and traverse all the layers of the neocortex from the white matter to the pial surface [27, 28, 41]. Depending on their type of action, neurons are mainly classified as *excitatory* or *inhibitory*, wherein this distinction depends on whether they increase the firing rate in the destination neurons they are communicating with, or they suppress them. Inhibitory neurons




Figure 1. Schematic of the structure of the neocortex with intracortical and corticocortical connections.

are located in all layers and usually have axons that remain within the same area as their cell body resides, and hence they have a local range of action. Layers III, V, and VI contain pyramidal excitatory neurons whose axons can provide long-range communication (projection) throughout the neocortex. Layer IV contains primarily star-shaped excitatory interneurons that receive sensory inputs from the thalamus. Figure 1 shows a schematic of the structure of the neocortex, including the intracortical and corticocortical neuronal connections; see [28, Chap. 15] for further details.

On a local scale, within a cortical column, neurons are densely interconnected and involve all types of feedforward and feedback intracortical connections. Such a dense and relatively

homogeneous local structure of the neocortex suggests modeling a local population of functionally similar neurons by a single *space-averaged neuron*, which preserves enough physiological information to understand the temporal patterns observed in spatially smoothed (averaged) EEG signals without creating excessive theoretical complications in the mathematical analysis of the model. On a global scale, in the exclusively excitatory corticocortical communication throughout the neocortex, two major patterns of connectivity are observed, namely, a homogeneous, symmetrical, and translation-invariant pattern of connections versus a heterogeneous, patchy, and asymmetrical distribution of connections. For modeling simplicity and due to the unavailability of detailed anatomical data, in the model that we investigate in this paper the corticocortical connectivity is assumed to be isotropic, homogeneous, symmetric, and translation invariant [36].

To establish the mathematical framework of the model, let $\Omega = (0, \omega) \times (0, \omega)$, $\omega > 0$, be an open rectangle in $\mathbb{R}^2$ that defines the domain of the neocortex. Each point $x = (x_1, x_2) \in \Omega$ indicates the location of a local network—possibly representing a cortical column—modeled by a space-averaged excitatory neuron and a space-averaged inhibitory neuron. Let E denote a population of excitatory neurons and let I denote a population of inhibitory neurons. For $x \in \Omega$, $t \in [0, T]$, $T > 0$, and $x, y \in \{E, I\}$, we denote by $v_x(x, t)$, measured in mV, the spatially mean soma membrane potential of a population of type x centered at x . Moreover, we denote by $i_{xy}(x, t)$, measured in mV, the spatially mean postsynaptic activation of synapses of a population of type x centered at x onto a population of type y centered at the same point x . In addition, we denote by $w_{ex}(x, t)$, measured in s^{-1} , the mean rate of corticocortical excitatory input pulses from the entire domain of the neocortex to a population of type x centered at x . Finally, we denote by $g_{xy}(x, t)$, measured in s^{-1} , the mean rate of subcortical input pulses of type x to a population of type y centered at x . Note that, by definition, $i_{xy}(x, t)$, $w_{ex}(x, t)$, and $g_{xy}(x, t)$ are nonnegative quantities.

Then, as developed in [36], the system of coupled ordinary and partial differential equations

$$\begin{aligned}
 (2) \quad & (\tau_E \partial_t + 1)v_E(x, t) = \frac{V_{EE} - v_E(x, t)}{|V_{EE}|} i_{EE}(x, t) + \frac{V_{IE} - v_E(x, t)}{|V_{IE}|} i_{IE}(x, t), \\
 & (\tau_I \partial_t + 1)v_I(x, t) = \frac{V_{EI} - v_I(x, t)}{|V_{EI}|} i_{EI}(x, t) + \frac{V_{II} - v_I(x, t)}{|V_{II}|} i_{II}(x, t), \\
 & (\partial_t + \gamma_{EE})^2 i_{EE}(x, t) = e \Upsilon_{EE} \gamma_{EE} [N_{EE} f_E(v_E(x, t)) + w_{EE}(x, t) + g_{EE}(x, t)], \\
 & (\partial_t + \gamma_{EI})^2 i_{EI}(x, t) = e \Upsilon_{EI} \gamma_{EI} [N_{EI} f_E(v_E(x, t)) + w_{EI}(x, t) + g_{EI}(x, t)], \\
 & (\partial_t + \gamma_{IE})^2 i_{IE}(x, t) = e \Upsilon_{IE} \gamma_{IE} [N_{IE} f_I(v_I(x, t)) + g_{IE}(x, t)], \\
 & (\partial_t + \gamma_{II})^2 i_{II}(x, t) = e \Upsilon_{II} \gamma_{II} [N_{II} f_I(v_I(x, t)) + g_{II}(x, t)], \\
 & \left[(\partial_t + \nu \Lambda_{EE})^2 - \frac{3}{2} \nu^2 \Delta \right] w_{EE}(x, t) = \nu^2 \Lambda_{EE}^2 M_{EE} f_E(v_E(x, t)), \\
 & \left[(\partial_t + \nu \Lambda_{EI})^2 - \frac{3}{2} \nu^2 \Delta \right] w_{EI}(x, t) = \nu^2 \Lambda_{EI}^2 M_{EI} f_E(v_E(x, t)), \quad (x, t) \in \Omega \times (0, T],
 \end{aligned}$$

Table 1

Definition and range of values for the biophysical parameters of the mean field model (2). All electric potentials are given with respect to the mean resting soma membrane potential $v_{\text{rest}} = -70 \text{ mV}$ [3].

Parameter	Definition	Range	Unit
τ_E	Passive excitatory membrane decay time constant	[0.005, 0.15]	s
τ_I	Passive inhibitory membrane decay time constant	[0.005, 0.15]	s
V_{EE}, V_{EI}	Mean excitatory Nernst potentials	[50, 80]	mV
V_{IE}, V_{II}	Mean inhibitory Nernst potentials	[-20, -5]	mV
γ_{EE}, γ_{EI}	Excitatory postsynaptic potential rate constants	[100, 1000]	s^{-1}
γ_{IE}, γ_{II}	Inhibitory postsynaptic potential rate constants	[10, 500]	s^{-1}
$\Upsilon_{EE}, \Upsilon_{EI}$	Amplitude of excitatory postsynaptic potentials	[0.1, 2.0]	mV
$\Upsilon_{IE}, \Upsilon_{II}$	Amplitude of inhibitory postsynaptic potentials	[0.1, 2.0]	mV
N_{EE}, N_{EI}	Number of intracortical excitatory connections	[2000, 5000]	—
N_{IE}, N_{II}	Number of intracortical inhibitory connections	[100, 1000]	—
ν	Corticocortical conduction velocity	[100, 1000]	cm/s
$\Lambda_{EE}, \Lambda_{EI}$	Decay scale of corticocortical excitatory connectivities	[0.1, 1.0]	cm^{-1}
M_{EE}, M_{EI}	Number of corticocortical excitatory connections	[2000, 5000]	—
F_E	Maximum mean excitatory firing rate	[50, 500]	s^{-1}
F_I	Maximum mean inhibitory firing rate	[50, 500]	s^{-1}
μ_E	Excitatory firing threshold potential	[15, 30]	mV
μ_I	Inhibitory firing threshold potential	[15, 30]	mV
σ_E	Standard deviation of excitatory firing threshold potential	[2, 7]	mV
σ_I	Standard deviation of inhibitory firing threshold potential	[2, 7]	mV

with periodic boundary conditions provides a mean field model for the electrocortical activity in the neocortex. Here, e is the Napier constant and $f_x(\cdot)$ is the mean firing rate function of a population of type x and is given by

$$(3) \quad f_x(v_x(x, t)) := \frac{F_x}{1 + \exp\left(-\sqrt{2} \frac{v_x(x, t) - \mu_x}{\sigma_x}\right)}, \quad x \in \{E, I\}.$$

The definition of the biophysical parameters of the model and the ranges of the values they may take are given in Table 1. For the range of values given in Table 1, we have $|V_{EE}| = V_{EE}$, $|V_{EI}| = V_{EI}$, $|V_{IE}| = -V_{IE}$, and $|V_{II}| = -V_{II}$, which we use to simplify (2). Note that, in addition to the notational changes to the original equations given in [36], we have changed the reference of the electric potential to the *resting potential* to avoid the constant terms that would otherwise appear in (2). Figure 2 shows a schematic of intracortical, corticocortical, and subcortical inputs to two local networks located at points x and y , along with their contribution to the global corticocortical activation as modeled by (2). The specific coupling between the equations of the model is depicted by the block diagram shown in Figure 3.

The first two equations in (2), that is, the v -equations, model the dynamics of the resistive-capacitive membrane of the space-averaged neurons located at x . In the absence




Figure 2. Cortical inputs to two local networks located at points x and y as modeled by (2).




Figure 3. Block diagram of the mean field model (2). The operators $P_1, \dots, P_8$ represent the eight equations in (2), respectively. As in Figures 1 and 2, the blocks associated with excitatory populations are shown in red, and the blocks associated with inhibitory populations are shown in blue.

of postsynaptic i -inputs, the mean membrane potential decays exponentially to the resting potential. The fractions appearing in the equations weight the postsynaptic inputs to incorporate the effect of transmembrane diffusive ion flows into the model. Specifically, the depolarizing effect of excitatory inputs on the membrane is linearly decreased by the weights as the membrane potential rises to the Nernst (reversal) potential. When the membrane potential exceeds the Nernst potential, the effect is reversed and further excitation tends to hyperpolarize the membrane. The weights associated with the inhibitory postsynaptic inputs have opposite signs at the resting potential, and hence they have an opposite reversal effect.

The critically damped second-order dynamics of the four i -equations in (2) generates a synaptic α -function—as in the classical dendritic cable theory—in response to an impulse.

As shown in Figure 2, these second-order dynamical systems are driven by three different sources of presynaptic spikes, namely, the inputs $N_{XY}f_X(v_X)$ from local neuronal populations, the excitatory inputs w_{EX} from corticocortical fibers, and the inputs g_{XY} from subcortical regions. As a result, these four equations generate the postsynaptic responses that modulate the polarization of the cell membranes according to the v -equations discussed before.

Unlike the conduction through short-range intracortical fibers, the conduction through long-range corticocortical fibers cannot be assumed to be instantaneous. The w -equations in (2) form a system of telegraph equations that effectively models the propagation of the excitatory axonal pulses through corticocortical fibers. To derive these equations, it is assumed in [36] that the strength of corticocortical connections onto a local population decays exponentially with distance, with the characteristic scale Λ_{EX} . Moreover, it is assumed that the spatial distribution of connections is isotropic and homogeneous all over the neocortex.

In practical applications, the key variable in the model presented by (2) is the mean membrane potential of excitatory populations $v_E(x, t)$, which is presumed to be linearly proportional to EEG recordings from the scalp [36, 37]. For further details of the model see [36], or the introductory sections of [6, 21, 37].

Now, let

$$\begin{aligned} v(x, t) &:= (v_E(x, t), v_I(x, t)) \in \mathbb{R}^2, \\ i(x, t) &:= (i_{EE}(x, t), i_{EI}(x, t), i_{IE}(x, t), i_{II}(x, t)) \in \mathbb{R}^4, \\ w(x, t) &:= (w_{EE}(x, t), w_{EI}(x, t)) \in \mathbb{R}^2, \\ g(x, t) &:= (g_{EE}(x, t), g_{EI}(x, t), g_{IE}(x, t), g_{II}(x, t)) \in \mathbb{R}^4, \end{aligned}$$

and note that (2) can be represented in vector form in $\Omega \times (0, T]$ as

$$(4) \quad \Phi \partial_t v + v - J_1 i + J_2 v i^T \Psi J_4 + J_3 v i^T \Psi J_5 = 0,$$

$$(5) \quad \partial_t^2 i + 2\Gamma \partial_t i + \Gamma^2 i - e\Upsilon \Gamma J_6 w - e\Upsilon \Gamma N J_7 f(v) = e\Upsilon \Gamma g,$$

$$(6) \quad \partial_t^2 w + 2\nu \Lambda \partial_t w - \frac{3}{2} \nu^2 \Delta w + \nu^2 \Lambda^2 w - \nu^2 \Lambda^2 M J_8 f(v) = 0,$$

where v , i , and w are Ω -periodic vector-valued functions with the initial values

$$(7) \quad v|_{t=0} = v_0, \quad i|_{t=0} = i_0, \quad (\partial_t i)|_{t=0} = i'_0, \quad w|_{t=0} = w_0, \quad (\partial_t w)|_{t=0} = w'_0,$$

and

$$(8) \quad \begin{aligned} \Phi &= \text{diag}(\tau_E, \tau_I), & \Psi &= \text{diag}\left(\frac{1}{|V_{EE}|}, \frac{1}{|V_{EI}|}, \frac{1}{|V_{IE}|}, \frac{1}{|V_{II}|}\right), \\ \Gamma &= \text{diag}(\gamma_{EE}, \gamma_{EI}, \gamma_{IE}, \gamma_{II}), & \Upsilon &= \text{diag}(\Upsilon_{EE}, \Upsilon_{EI}, \Upsilon_{IE}, \Upsilon_{II}), \\ N &= \text{diag}(N_{EE}, N_{EI}, N_{IE}, N_{II}), & M &= \text{diag}(M_{EE}, M_{EI}), \\ \Lambda &= \text{diag}(\Lambda_{EE}, \Lambda_{EI}), & J_1 &= [I_{2 \times 2} \quad -I_{2 \times 2}], \\ J_2 &= \text{diag}(1, 0), & J_3 &= \text{diag}(0, 1), \\ J_4 &= [1 \quad 0 \quad 1 \quad 0]^T, & J_5 &= [0 \quad 1 \quad 0 \quad 1]^T, \end{aligned}$$

$$\begin{aligned}
J_6 &= \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}^T, & J_7 &= \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix}^T, \\
J_8 &= \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}, & f(v) &= \begin{bmatrix} f_E([1 \ 0] v) \\ f_I([0 \ 1] v) \end{bmatrix}.
\end{aligned}$$

For simplicity of exposition, the dependence of the functions v , i , w , and g on the arguments (x, t) is not explicitly shown in (4)–(6). Note that (4) and (5), which model the local dynamics of the neocortex, are essentially systems of ODEs. These equations do not possess any spatial smoothing components, and hence their solutions are expected to evolve in less regular function spaces [39, 47]. The system of PDEs (6) consists of two telegraph equations coupled indirectly through (4) and (5); see Figure 3.

Remark 3.1 (Variations in parameter values). In the analysis that follows in the rest of the paper, we assume that all the parameters of the model are constant. However, in practical applications, certain parameters may be considered to vary in time or space to model specific physiological situations in the brain. The variations can occur independently, or can be modeled using additional ODEs or PDEs coupled with the existing equations. We give all the details of the results—some of which may, however, be considered fairly standard—along with a careful inclusion of all parameters. Therefore, in applications it should be possible to easily observe where the parameters of interest appear in the analysis, and whether or not their particular variations can affect the validity of the results.

4. Existence and uniqueness of solutions. In this section, we investigate the problem of existence, uniqueness, and regularity of solutions for (4)–(6) with the initial values (7) and periodic boundary conditions. We set appropriate spaces of Ω -periodic functions as the functional framework of the problem by which we include the boundary conditions in the solution spaces. We view $v(x, t)$, $i(x, t)$, and $w(x, t)$ as Banach space-valued functions and follow the standard technique of Galerkin approximations [16, 44, 51] to construct weak and strong solutions in Theorems 4.5 and 4.7. The details of the proof of these results can be skipped if the reader is proficient in the analysis of the Galerkin method.

First, define the function spaces

$$\begin{aligned}
(9) \quad \mathcal{L}_v^2 &:= L_{\text{per}}^2(\Omega; \mathbb{R}^2), & \mathcal{L}_i^2 &:= L_{\text{per}}^2(\Omega; \mathbb{R}^4), & \mathcal{L}_w^2 &:= L_{\text{per}}^2(\Omega; \mathbb{R}^2), \\
\mathcal{L}_v^\infty &:= L_{\text{per}}^\infty(\Omega; \mathbb{R}^2), & \mathcal{L}_i^\infty &:= L_{\text{per}}^\infty(\Omega; \mathbb{R}^4), & \mathcal{L}_w^\infty &:= L_{\text{per}}^\infty(\Omega; \mathbb{R}^2), \\
\mathcal{H}_w^1 &:= H_{\text{per}}^1(\Omega; \mathbb{R}^2), & \mathcal{H}_w^2 &:= H_{\text{per}}^2(\Omega; \mathbb{R}^2), \\
\mathcal{L}_{\partial w}^2 &:= L_{\text{per}}^2(\Omega; \mathbb{R}^{2 \times 2}), & \mathcal{H}_{\partial w}^1 &:= H_{\text{per}}^1(\Omega; \mathbb{R}^{2 \times 2}), \\
\mathcal{W}_w^{1,\infty} &:= W_{\text{per}}^{1,\infty}(\Omega; \mathbb{R}^2),
\end{aligned}$$

and denote by $\mathcal{L}_v^{2*}$, $\mathcal{L}_i^{2*}$, and $\mathcal{H}_w^{1*}$ the dual spaces of $\mathcal{L}_v^2$, $\mathcal{L}_i^2$, and $\mathcal{H}_w^1$, respectively. Note that $\mathcal{L}_v^2$ and $\mathcal{L}_i^2$ are, respectively, isometrically isomorphic to $\mathcal{L}_v^{2*}$ and $\mathcal{L}_i^{2*}$ [18, Thm. 6.15], which we denote by $\mathcal{L}_v^{2*} = \mathcal{L}_v^2$ and $\mathcal{L}_i^{2*} = \mathcal{L}_i^2$. By the Rellich–Kondrachov compact embedding theorems we have $\mathcal{H}_w^{1*} \subseteq \mathcal{L}_w^2 \hookrightarrow \mathcal{H}_w^{1*}$; see [9, Thm. 6.6-3] and [44, Thm. A.4]. Moreover, there

exists a dual orthogonal basis of $\mathcal{H}_w^1$ and $\mathcal{L}_w^2$ given by the following lemma. The proof of this lemma is fairly standard and follows the general results given in [51, sect. II.2.1].

Lemma 4.1 (Dual orthogonal basis). *There exists an orthonormal basis of $\mathcal{L}_w^2$ that is also an orthogonal basis of $\mathcal{H}_w^1$, and can be constructed by the eigenfunctions of the linear operator $A := -\Delta + I$.*

Now, before proceeding to the main results of this section, we define the notions of *weak* and *strong* solutions of (4)–(7) as used in this paper.

Definition 4.2 (Weak solution). *A solution (v, i, w) is called an Ω -periodic weak solution of the initial value problem (4)–(7) if it solves the weak version of the problem wherein the equations are understood as equalities in the space of duals $L^2(0, T; \mathcal{L}_v^{2*} \times \mathcal{L}_i^{2*} \times \mathcal{H}_w^{1*})$. That is, the functions*

$$v \in L^2(0, T; \mathcal{L}_v^2), \quad i \in L^2(0, T; \mathcal{L}_i^2), \quad w \in L^2(0, T; \mathcal{H}_w^1)$$

with

$$\begin{aligned} d_t v &\in L^2(0, T; \mathcal{L}_v^{2*}), & d_t i &\in L^2(0, T; \mathcal{L}_i^2), & d_t^2 i &\in L^2(0, T; \mathcal{L}_i^{2*}), \\ d_t w &\in L^2(0, T; \mathcal{L}_w^2), & d_t^2 w &\in L^2(0, T; \mathcal{H}_w^{1*}) \end{aligned}$$

construct an Ω -periodic weak solution for (4)–(7) if, for every $\ell_v \in \mathcal{L}_v^2$, $\ell_i \in \mathcal{L}_i^2$, $h_w \in \mathcal{H}_w^1$, and almost every $t \in [0, T]$, $T > 0$,

$$(10) \quad \langle \Phi d_t v, \ell_v \rangle_{\mathcal{L}_v^2} + (v, \ell_v)_{\mathcal{L}_v^2} - (J_1 i, \ell_v)_{\mathcal{L}_v^2} + \left(J_2 v i^T \Psi J_4 + J_3 v i^T \Psi J_5, \ell_v \right)_{\mathcal{L}_v^2} = 0,$$

$$(11) \quad \langle d_t^2 i, \ell_i \rangle_{\mathcal{L}_i^2} + 2(\Gamma d_t i, \ell_i)_{\mathcal{L}_i^2} + (\Gamma^2 i, \ell_i)_{\mathcal{L}_i^2} - e(\Upsilon \Gamma J_6 w, \ell_i)_{\mathcal{L}_i^2} \\ - e(\Upsilon \Gamma N J_7 f(v), \ell_i)_{\mathcal{L}_i^2} = e(\Upsilon \Gamma g, \ell_i)_{\mathcal{L}_i^2},$$

$$(12) \quad \langle d_t^2 w, h_w \rangle_{\mathcal{H}_w^1} + 2\nu(\Lambda d_t w, h_w)_{\mathcal{L}_w^2} + \frac{3}{2}\nu^2(\partial_x w, \partial_x h_w)_{\mathcal{L}_{\partial w}^2} \\ + \nu^2(\Lambda^2 w, h_w)_{\mathcal{L}_w^2} - \nu^2(\Lambda^2 M J_8 f(v), h_w)_{\mathcal{L}_w^2} = 0$$

with the initial values

$$(13) \quad v(0) = v_0, \quad i(0) = i_0, \quad d_t i(0) = i'_0, \quad w(0) = w_0, \quad d_t w(0) = w'_0.$$

Definition 4.3 (Strong solution). *A solution (v, i, w) is called an Ω -periodic strong solution of the initial value problem (4)–(7) if it solves the strong version of the problem wherein the equations are understood as equalities in $L^2(0, T; \mathcal{L}_v^2 \times \mathcal{L}_i^2 \times \mathcal{L}_w^2)$. That is, the functions*

$$v \in H^1(0, T; \mathcal{L}_v^2), \quad i \in H^2(0, T; \mathcal{L}_i^2), \quad w \in L^2(0, T; \mathcal{H}_w^2)$$

with

$$\begin{aligned} d_t v &\in L^2(0, T; \mathcal{L}_v^2), & d_t i &\in H^1(0, T; \mathcal{L}_i^2), & d_t^2 i &\in L^2(0, T; \mathcal{L}_i^2), \\ d_t w &\in L^2(0, T; \mathcal{H}_w^1), & d_t^2 w &\in L^2(0, T; \mathcal{L}_w^2) \end{aligned}$$

construct an Ω -periodic strong solution for (4)–(7) wherein they solve the equations for almost every $x \in \Omega$ and almost every $t \in [0, T]$, $T > 0$.

Now, let $\mathcal{B}_v = \{\ell_v^{(l)}\}_{l=1}^\infty$ be a basis of $\mathcal{L}_v^2$ such that $\{\Phi^{\frac{1}{2}}\ell_v^{(l)}\}_{l=1}^\infty$ is orthonormal in $\mathcal{L}_v^2$. Note that (8), with the range of values given in Table 1, implies that Φ is a positive-definite diagonal matrix, and hence such a basis exists. Moreover, let $\mathcal{B}_i = \{\ell_i^{(l)}\}_{l=1}^\infty$ be an orthonormal basis of $\mathcal{L}_i^2$, and let $\mathcal{B}_w = \{h_w^{(l)}\}_{l=1}^\infty$ be an orthogonal basis of $\mathcal{H}_w^1$ that is orthonormal in $\mathcal{L}_w^2$; see Lemma 4.1 for the existence and structure of $\mathcal{B}_w$. Finally, construct the set $\mathcal{B} = \{b^{(k)}\}_{k=1}^\infty \subset \mathcal{L}_v^2 \times \mathcal{L}_i^2 \times \mathcal{H}_w^1$ as

$$(14) \quad \mathcal{B} := \mathcal{B}_v \times \mathcal{B}_i \times \mathcal{B}_w = \left\{ b^{(k)} = \left(\ell_v^{(k)}, \ell_i^{(k)}, h_w^{(k)} \right) : \ell_v^{(k)} \in \mathcal{B}_v, \ell_i^{(k)} \in \mathcal{B}_i, h_w^{(k)} \in \mathcal{B}_w \right\}_{k=1}^\infty.$$

For each positive integer m , we seek approximations $v^{(m)} : [0, T] \rightarrow \mathcal{L}_v^2$, $i^{(m)} : [0, T] \rightarrow \mathcal{L}_i^2$, and $w^{(m)} : [0, T] \rightarrow \mathcal{H}_w^1$ of the form

$$(15) \quad v^{(m)}(t) := \sum_{k=1}^m c_{v_k}^{(m)}(t) \ell_v^{(k)},$$

$$(16) \quad i^{(m)}(t) := \sum_{k=1}^m c_{i_k}^{(m)}(t) \ell_i^{(k)},$$

$$(17) \quad w^{(m)}(t) := \sum_{k=1}^m c_{w_k}^{(m)}(t) h_w^{(k)},$$

constructed by the first m components of $\mathcal{B}$ and sufficiently smooth scalar-valued functions $c_{v_k}^{(m)}$, $c_{i_k}^{(m)}$, and $c_{w_k}^{(m)}$ on $[0, T]$ such that these approximations satisfy

$$(18) \quad \left(\Phi d_t v^{(m)}, \ell_v^{(k)} \right)_{\mathcal{L}_v^2} + \left(v^{(m)}, \ell_v^{(k)} \right)_{\mathcal{L}_v^2} - \left(J_1 i^{(m)}, \ell_v^{(k)} \right)_{\mathcal{L}_v^2} \\ + \left(J_2 v^{(m)} i^{(m)T} \Psi J_4 + J_3 v^{(m)} i^{(m)T} \Psi J_5, \ell_v^{(k)} \right)_{\mathcal{L}_v^2} = 0,$$

$$(19) \quad \left(d_t^2 i^{(m)}, \ell_i^{(k)} \right)_{\mathcal{L}_i^2} + 2 \left(\Gamma d_t i^{(m)}, \ell_i^{(k)} \right)_{\mathcal{L}_i^2} + \left(\Gamma^2 i^{(m)}, \ell_i^{(k)} \right)_{\mathcal{L}_i^2} \\ - e \left(\Upsilon \Gamma J_6 w^{(m)}, \ell_i^{(k)} \right)_{\mathcal{L}_i^2} - e \left(\Upsilon \Gamma N J_7 f(v^{(m)}), \ell_i^{(k)} \right)_{\mathcal{L}_i^2} = e \left(\Upsilon \Gamma g, \ell_i^{(k)} \right)_{\mathcal{L}_i^2},$$

$$(20) \quad \left(d_t^2 w^{(m)}, h_w^{(k)} \right)_{\mathcal{L}_w^2} + 2\nu \left(\Lambda d_t w^{(m)}, h_w^{(k)} \right)_{\mathcal{L}_w^2} + \frac{3}{2} \nu^2 \left(\partial_x w^{(m)}, \partial_x h_w^{(k)} \right)_{\mathcal{L}_{\partial w}^2} \\ + \nu^2 \left(\Lambda^2 w^{(m)}, h_w^{(k)} \right)_{\mathcal{L}_w^2} - \nu^2 \left(\Lambda^2 M J_8 f(v^{(m)}), h_w^{(k)} \right)_{\mathcal{L}_w^2} = 0$$

for all $t \in [0, T]$ and $k = 1, \dots, m$, subject to the initial conditions

$$(21) \quad c_{v_k}^{(m)}(0) = \left(v_0, \ell_v^{(k)} \right)_{\mathcal{L}_v^2}, \quad c_{i_k}^{(m)}(0) = \left(i_0, \ell_i^{(k)} \right)_{\mathcal{L}_i^2}, \quad d_t c_{i_k}^{(m)}(0) = \left(i'_0, \ell_i^{(k)} \right)_{\mathcal{L}_i^2}, \\ c_{w_k}^{(m)}(0) = \left(w_0, h_w^{(k)} \right)_{\mathcal{L}_w^2}, \quad d_t c_{w_k}^{(m)}(0) = \left(w'_0, h_w^{(k)} \right)_{\mathcal{L}_w^2}$$

on the coefficients $c_k^{(m)}(t) := (c_{v_k}^{(m)}(t), c_{i_k}^{(m)}(t), c_{w_k}^{(m)}(t)) \in \mathbb{R}^3$.

Equations (18)–(21) are equivalent to a system of nonlinear $3m$ -dimensional ODEs on coefficients $c^{(m)}(t) = (c_1^{(m)}(t), \dots, c_m^{(m)}(t)) \in \mathbb{R}^{3m}$. Therefore, by the standard theory of ODEs [50, Thm. 2.1], there exists a unique function $c^{(m)}(t)$ that solves (18)–(21) for $t \in [0, T_m)$, $T_m > 0$, with the approximations (15)–(17). Moreover, $T_m = T$ for all positive integers m , which follows from Proposition 4.4.

The standard Galerkin approximation method involves providing energy estimates that are uniform in m for all the approximations $(v^{(m)}, i^{(m)}, w^{(m)})$. Such a priori energy estimates then allow construction of solutions by passing to the limits as $m \rightarrow \infty$. The following proposition gives the desired estimates for the approximations (15)–(17).

Proposition 4.4 (Energy estimates). *Suppose $g \in L^2(0, T; \mathcal{L}_i^2)$ and, for every positive integer m , let $v^{(m)}$, $i^{(m)}$, and $w^{(m)}$ be functions of the form (15)–(17), respectively, satisfying (18)–(20) with the initial conditions (21). Then there exist positive constants α_v , β_v , α_i , and α_w , dependent only on the parameters of the model, such that for every positive integer m ,*

$$(22) \quad \sup_{t \in [0, T]} \left(\|v^{(m)}(t)\|_{\mathcal{L}_v^2}^2 \right) + \|d_t v^{(m)}\|_{L^2(0, T; \mathcal{L}_v^{2*})}^2 \leq \kappa_v,$$

$$(23) \quad \sup_{t \in [0, T]} \left(\|d_t i^{(m)}(t)\|_{\mathcal{L}_i^2}^2 + \|i^{(m)}(t)\|_{\mathcal{L}_i^2}^2 \right) + \|d_t^2 i^{(m)}\|_{L^2(0, T; \mathcal{L}_i^{2*})}^2 \leq \kappa_i,$$

$$(24) \quad \sup_{t \in [0, T]} \left(\|d_t w^{(m)}(t)\|_{\mathcal{L}_w^2}^2 + \|w^{(m)}(t)\|_{\mathcal{H}_w^1}^2 \right) + \|d_t^2 w^{(m)}\|_{L^2(0, T; \mathcal{H}_w^{1*})}^2 \leq \kappa_w,$$

where κ_v , κ_i , and κ_w are positive constants given, independently of m , by

$$(25) \quad \kappa_v := \alpha_v \left((1 + (1 + \sqrt{\kappa_i})^2 T) \exp(\beta_v \sqrt{\kappa_i} T) \left[\|v_0\|_{\mathcal{L}_v^2}^2 + \sqrt{\kappa_i} \right] + \kappa_i T \right),$$

$$(26) \quad \kappa_i := \alpha_i \left((1 + T) \left[\|i'_0\|_{\mathcal{L}_i^2}^2 + \|i_0\|_{\mathcal{L}_i^2}^2 \right] + (2 + T) \left[T(\kappa_w + |\Omega|) + \|g\|_{L^2(0, T; \mathcal{L}_i^2)}^2 \right] \right),$$

$$(27) \quad \kappa_w := \alpha_w \left((1 + T) \left[\|w'_0\|_{\mathcal{L}_w^2}^2 + \|w_0\|_{\mathcal{H}_w^1}^2 \right] + (2 + T) T |\Omega| \right).$$

Proof. Multiplying (20) by $d_t c_{w_k}^{(m)}$ and summing over $k = 1, \dots, m$ yields

$$\begin{aligned} & \left(d_t^2 w^{(m)}, d_t w^{(m)} \right)_{\mathcal{L}_w^2} + 2\nu \left(\Lambda d_t w^{(m)}, d_t w^{(m)} \right)_{\mathcal{L}_w^2} + \frac{3}{2} \nu^2 \left(\partial_x w^{(m)}, d_t \partial_x w^{(m)} \right)_{\mathcal{L}_{\partial w}^2} \\ & + \nu^2 \left(\Lambda^2 w^{(m)}, d_t w^{(m)} \right)_{\mathcal{L}_w^2} - \nu^2 \left(\Lambda^2 M J_8 f(v^{(m)}), d_t w^{(m)} \right)_{\mathcal{L}_w^2} = 0, \end{aligned}$$

or, equivalently,

$$\begin{aligned} & \frac{1}{2} d_t \left[\|d_t w^{(m)}\|_{\mathcal{L}_w^2}^2 + \frac{3}{2} \nu^2 \|\partial_x w^{(m)}\|_{\mathcal{L}_{\partial w}^2}^2 + \nu^2 \|\Lambda w^{(m)}\|_{\mathcal{L}_w^2}^2 \right] \\ & + 2\nu \left\| \Lambda^{\frac{1}{2}} d_t w^{(m)} \right\|_{\mathcal{L}_w^2}^2 - \nu^2 \left(\Lambda^2 M J_8 f(v^{(m)}), d_t w^{(m)} \right)_{\mathcal{L}_w^2} = 0. \end{aligned}$$

Now, Young's inequality implies that, for every $\varepsilon_1 > 0$,

$$\begin{aligned} \nu^2 \left(\Lambda^2 M J_8 f(v^{(m)}), d_t w^{(m)} \right)_{\mathcal{L}_w^2} &\leq \varepsilon_1 \nu^2 \|d_t w^{(m)}\|_{\mathcal{L}_w^2}^2 + \frac{\nu^2}{4\varepsilon_1} \|\Lambda^2 M J_8 f(v^{(m)})\|_{\mathcal{L}_w^2}^2 \\ &= \varepsilon_1 \nu^2 \|d_t w^{(m)}\|_{\mathcal{L}_w^2}^2 + \frac{\nu^2}{4\varepsilon_1} \operatorname{tr}(\Lambda^4 M^2) \int_{\Omega} |f_E(v_E^{(m)})|^2 dx \\ &\leq \varepsilon_1 \nu^2 \|d_t w^{(m)}\|_{\mathcal{L}_w^2}^2 + \frac{\nu^2}{4\varepsilon_1} |\Omega| F_E^2 \operatorname{tr}(\Lambda^4 M^2). \end{aligned}$$

Therefore,

$$\begin{aligned} d_t \left[\|d_t w^{(m)}\|_{\mathcal{L}_w^2}^2 + \frac{3}{2} \nu^2 \|\partial_x w^{(m)}\|_{\mathcal{L}_{\partial w}^2}^2 + \nu^2 \|\Lambda w^{(m)}\|_{\mathcal{L}_w^2}^2 \right] &+ 2\nu(2\Lambda_{\min} - \varepsilon_1 \nu) \|d_t w^{(m)}\|_{\mathcal{L}_w^2}^2 \\ &\leq \frac{\nu^2}{2\varepsilon_1} |\Omega| F_E^2 \operatorname{tr}(\Lambda^4 M^2), \end{aligned}$$

where $\Lambda_{\min} := \min\{\Lambda_{EE}, \Lambda_{EI}\}$ is the smallest eigenvalue of Λ .

Next, setting $\varepsilon_1 = \frac{2}{\nu} \Lambda_{\min}$ and integrating with respect to time over $[0, t]$ yields

$$\begin{aligned} \|d_t w^{(m)}(t)\|_{\mathcal{L}_w^2}^2 + \frac{3}{2} \nu^2 \|\partial_x w^{(m)}(t)\|_{\mathcal{L}_{\partial w}^2}^2 + \nu^2 \|\Lambda w^{(m)}(t)\|_{\mathcal{L}_w^2}^2 \\ \leq \left(\|d_t w^{(m)}\|_{\mathcal{L}_w^2}^2 + \frac{3}{2} \nu^2 \|\partial_x w^{(m)}\|_{\mathcal{L}_{\partial w}^2}^2 + \nu^2 \|\Lambda w^{(m)}\|_{\mathcal{L}_w^2}^2 \right) \Big|_{t=0} + \frac{1}{4} \frac{\nu^3}{\Lambda_{\min}} |\Omega| F_E^2 \operatorname{tr}(\Lambda^4 M^2) t, \end{aligned}$$

which, using (21), implies

$$\|d_t w^{(m)}(t)\|_{\mathcal{L}_w^2}^2 + \|w^{(m)}(t)\|_{\mathcal{H}_w^1}^2 \leq \hat{\alpha}_w \left(\|w_0'\|_{\mathcal{L}_w^2}^2 + \|w_0\|_{\mathcal{H}_w^1}^2 + \frac{1}{4} \frac{\nu^3}{\Lambda_{\min}} |\Omega| F_E^2 \operatorname{tr}(\Lambda^4 M^2) t \right)$$

for all $t \in [0, T]$ and some $\hat{\alpha}_w > 0$. Since this inequality holds for all $t \in [0, T]$, it follows that

$$(28) \quad \sup_{t \in [0, T]} \left(\|d_t w^{(m)}(t)\|_{\mathcal{L}_w^2}^2 + \|w^{(m)}(t)\|_{\mathcal{H}_w^1}^2 \right) \leq \hat{\kappa}_w,$$

where

$$\hat{\kappa}_w := \hat{\alpha}_w \left(\|w_0'\|_{\mathcal{L}_w^2}^2 + \|w_0\|_{\mathcal{H}_w^1}^2 + \frac{1}{4} \frac{\nu^3}{\Lambda_{\min}} |\Omega| F_E^2 \operatorname{tr}(\Lambda^4 M^2) T \right).$$

Now, fix $\bar{h} \in \mathcal{H}_w^1$ such that $\|\bar{h}\|_{\mathcal{H}_w^1} \leq 1$ and decompose $\bar{h}$ as $\bar{h} = h + h^\perp$, where $h \in \operatorname{span}\{h_w^{(k)}\}_{k=1}^m$ and $(h_w^{(k)} h^\perp)_{\mathcal{L}_w^2} = 0$, $k = 1, \dots, m$. Since the basis $\mathcal{B}_w$ used to construct $\mathcal{B}$ in (14) is orthonormal in $\mathcal{L}_w^2$, it follows from (17) that

$$\langle d_t^2 w^{(m)}, \bar{h} \rangle_{\mathcal{H}_w^1} = (d_t^2 w^{(m)}, \bar{h})_{\mathcal{L}_w^2} = (d_t^2 w^{(m)}, h)_{\mathcal{L}_w^2},$$

where the first equality holds since $d_t^2 w^{(m)} \in \mathcal{H}_w^1$; see the proof of [16, Thm. 5.9-1]. Therefore, (20) gives

$$\begin{aligned} \langle d_t^2 w^{(m)}, \bar{h} \rangle_{\mathcal{H}_w^1} &= -2\nu (\Lambda d_t w^{(m)}, h)_{\mathcal{L}_w^2} - \frac{3}{2} \nu^2 (\partial_x w^{(m)}, \partial_x h)_{\mathcal{L}_{\partial w}^2} \\ &\quad - \nu^2 (\Lambda^2 w^{(m)}, h)_{\mathcal{L}_w^2} + \nu^2 (\Lambda^2 M J_8 f(v^{(m)}), h)_{\mathcal{L}_w^2}. \end{aligned}$$

Since $\mathcal{B}_w$ is orthogonal in $\mathcal{H}_w^1$, we have $\|h\|_{\mathcal{H}_w^1} \leq \|\bar{h}\|_{\mathcal{H}_w^1} \leq 1$, and hence the Cauchy–Schwarz inequality gives

$$\begin{aligned} \left| \langle d_t^2 w^{(m)}, \bar{h} \rangle_{\mathcal{H}_w^1} \right| &\leq 2\nu \|d_t w^{(m)}\|_{\mathcal{L}_w^2} + \frac{3}{2} \nu^2 \|\partial_x w^{(m)}\|_{\mathcal{L}_{\partial w}^2} + \nu^2 \|\Lambda^2 w^{(m)}\|_{\mathcal{L}_w^2} + \nu^2 \|\Lambda^2 M J_8 f(v^{(m)})\|_{\mathcal{L}_w^2} \\ &\leq \alpha_1 \left(\|d_t w^{(m)}\|_{\mathcal{L}_w^2} + \|w^{(m)}\|_{\mathcal{H}_w^1} + \nu^2 (|\Omega| F_E^2 \operatorname{tr}(\Lambda^4 M^2))^{\frac{1}{2}} \right) \end{aligned}$$

for some $\alpha_1 > 0$. Therefore, there exists $\alpha_2 > 0$ such that

$$\int_0^T \|d_t^2 w^{(m)}\|_{\mathcal{H}_w^{1*}}^2 dt \leq \alpha_2 \int_0^T \left(\|d_t w^{(m)}\|_{\mathcal{L}_w^2}^2 + \|w^{(m)}\|_{\mathcal{H}_w^1}^2 + \nu^4 |\Omega| F_E^2 \operatorname{tr}(\Lambda^4 M^2) \right) dt,$$

which, using (28), yields

$$\|d_t^2 w^{(m)}\|_{L^2(0,T;\mathcal{H}_w^{1*})}^2 \leq \alpha_2 (\hat{\kappa}_w + \nu^4 |\Omega| F_E^2 \operatorname{tr}(\Lambda^4 M^2)) T.$$

This inequality, together with (28), establishes the bound (24) with (27) for some $\alpha_w > 0$.

Next, multiplying (19) by $d_t c_{i_k}^{(m)}$ and summing over $k = 1, \dots, m$ yields

$$\begin{aligned} (29) \quad (d_t^2 i^{(m)}, d_t i^{(m)})_{\mathcal{L}_i^2} &+ 2(\Gamma d_t i^{(m)}, d_t i^{(m)})_{\mathcal{L}_i^2} + (\Gamma^2 i^{(m)}, d_t i^{(m)})_{\mathcal{L}_i^2} - e(\Upsilon \Gamma J_6 w^{(m)}, d_t i^{(m)})_{\mathcal{L}_i^2} \\ &- e(\Upsilon \Gamma N J_7 f(v^{(m)}), d_t i^{(m)})_{\mathcal{L}_i^2} = e(\Upsilon \Gamma g, d_t i^{(m)})_{\mathcal{L}_i^2}. \end{aligned}$$

For the second term, we have

$$(\Gamma d_t i^{(m)}, d_t i^{(m)})_{\mathcal{L}_i^2} \geq \gamma_{\min} \|d_t i^{(m)}\|_{\mathcal{L}_i^2}^2,$$

where $\gamma_{\min} := \min\{\gamma_{EE}, \gamma_{EI}, \gamma_{IE}, \gamma_{II}\}$ is the smallest eigenvalue of Γ . Now, using Young's inequality and recalling (24) we obtain, for every $\varepsilon_2, \dots, \varepsilon_4 > 0$,

$$\begin{aligned} e(\Upsilon \Gamma J_6 w^{(m)}, d_t i^{(m)})_{\mathcal{L}_i^2} &\leq \varepsilon_2 \|d_t i^{(m)}\|_{\mathcal{L}_i^2}^2 + \frac{e^2}{4\varepsilon_2} \|\Upsilon \Gamma J_6 w^{(m)}\|_{\mathcal{L}_i^2}^2 \\ &\leq \varepsilon_2 \|d_t i^{(m)}\|_{\mathcal{L}_i^2}^2 + \frac{e^2}{4\varepsilon_2} \|\Upsilon \Gamma J_6\|_2^2 \|w^{(m)}\|_{\mathcal{L}_w^2}^2 \\ &\leq \varepsilon_2 \|d_t i^{(m)}\|_{\mathcal{L}_i^2}^2 + \frac{e^2 \kappa_w}{4\varepsilon_2} \|\Upsilon \Gamma J_6\|_2^2, \end{aligned}$$

$$\begin{aligned}
e(\Upsilon \Gamma N J_7 f(v^{(m)}), d_t i^{(m)})_{\mathcal{L}_i^2} &\leq \varepsilon_3 \|d_t i^{(m)}\|_{\mathcal{L}_i^2}^2 + \frac{e^2}{4\varepsilon_3} \|\Upsilon \Gamma N J_7 f(v^{(m)})\|_{\mathcal{L}_i^2}^2 \\
&\leq \varepsilon_3 \|d_t i^{(m)}\|_{\mathcal{L}_i^2}^2 + \frac{e^2}{4\varepsilon_3} \|\Upsilon \Gamma N J_7\|_2^2 \|f(v^{(m)})\|_{\mathcal{L}_v^2}^2 \\
&\leq \varepsilon_3 \|d_t i^{(m)}\|_{\mathcal{L}_i^2}^2 + \frac{e^2 |\Omega|}{4\varepsilon_3} (F_E^2 + F_I^2) \|\Upsilon \Gamma N J_7\|_2^2, \\
e(\Upsilon \Gamma g, d_t i^{(m)})_{\mathcal{L}_i^2} &\leq \varepsilon_4 \|d_t i^{(m)}\|_{\mathcal{L}_i^2}^2 + \frac{e^2}{4\varepsilon_4} \|\Upsilon \Gamma g\|_{\mathcal{L}_i^2}^2 \\
&\leq \varepsilon_4 \|d_t i^{(m)}\|_{\mathcal{L}_i^2}^2 + \frac{e^2}{4\varepsilon_4} \|\Upsilon \Gamma\|_2^2 \|g\|_{\mathcal{L}_i^2}^2.
\end{aligned}$$

Hence, with the above inequalities, (29) implies

$$\begin{aligned}
&d_t \left[\|d_t i^{(m)}\|_{\mathcal{L}_i^2}^2 + \|\Gamma i^{(m)}\|_{\mathcal{L}_i^2}^2 \right] + 2(2\gamma_{\min} - \varepsilon_2 - \varepsilon_3 - \varepsilon_4) \|d_t i^{(m)}\|_{\mathcal{L}_i^2}^2 \\
&\leq \frac{e^2 \kappa_w}{2\varepsilon_2} \|\Upsilon \Gamma J_6\|_2^2 + \frac{e^2 |\Omega|}{2\varepsilon_3} (F_E^2 + F_I^2) \|\Upsilon \Gamma N J_7\|_2^2 + \frac{e^2}{2\varepsilon_4} \|\Upsilon \Gamma\|_2^2 \|g\|_{\mathcal{L}_i^2}^2.
\end{aligned}$$

Now, setting $\varepsilon_2 = \varepsilon_3 = \frac{1}{2}\gamma_{\min}$ and $\varepsilon_4 = \gamma_{\min}$, integrating with respect to time over $[0, t]$, and taking the supremum over $t \in [0, T]$, we have

$$(30) \quad \sup_{t \in [0, T]} \left(\|d_t i^{(m)}(t)\|_{\mathcal{L}_i^2}^2 + \|i^{(m)}(t)\|_{\mathcal{L}_i^2}^2 \right) \leq \hat{\kappa}_i,$$

where, for some $\hat{\alpha}_i > 0$,

$$\begin{aligned}
\hat{\kappa}_i &= \hat{\alpha}_i \left(\|i_0'\|_{\mathcal{L}_i^2}^2 + \|i_0\|_{\mathcal{L}_i^2}^2 + \left[\frac{e^2 \kappa_w}{\gamma_{\min}} \|\Upsilon \Gamma J_6\|_2^2 + \frac{e^2 |\Omega|}{\gamma_{\min}} (F_E^2 + F_I^2) \|\Upsilon \Gamma N J_7\|_2^2 \right] T \right. \\
&\quad \left. + \frac{e^2}{2\gamma_{\min}} \|\Upsilon \Gamma\|_2^2 \|g\|_{L^2(0, T; \mathcal{L}_i^2)}^2 \right).
\end{aligned}$$

Fix $\bar{\ell} \in \mathcal{L}_i^2$ such that $\|\bar{\ell}\|_{\mathcal{L}_i^2} \leq 1$ and decompose $\bar{\ell}$ as $\bar{\ell} = \ell + \ell^\perp$, where $\ell \in \text{span}\{\ell_i^{(k)}\}_{k=1}^m$ and $(\ell_i^{(k)}, \ell^\perp)_{\mathcal{L}_i^2} = 0$, $k = 1, \dots, m$. Using (16) and (19), we obtain

$$\begin{aligned}
\langle d_t^2 i^{(m)}, \bar{\ell} \rangle_{\mathcal{L}_i^2} &= (d_t^2 i^{(m)}, \bar{\ell})_{\mathcal{L}_i^2} = (d_t^2 i^{(m)}, \ell)_{\mathcal{L}_i^2} \\
&= -2(\Gamma d_t i^{(m)}, \ell)_{\mathcal{L}_i^2} - (\Gamma^2 i^{(m)}, \ell)_{\mathcal{L}_i^2} + e(\Upsilon \Gamma J_6 w^{(m)}, \ell)_{\mathcal{L}_i^2} \\
&\quad + e(\Upsilon \Gamma N J_7 f(v^{(m)}), \ell)_{\mathcal{L}_i^2} + e(\Upsilon \Gamma g, \ell)_{\mathcal{L}_i^2}.
\end{aligned}$$

The orthogonality of the basis $\mathcal{B}_i$ in (14) implies $\|\ell\|_{\mathcal{L}_i^2} \leq 1$, and hence

$$\begin{aligned}
\left| \langle d_t^2 i^{(m)}, \bar{\ell} \rangle_{\mathcal{L}_i^2} \right| &\leq 2\|\Gamma\|_2 \|d_t i^{(m)}\|_{\mathcal{L}_i^2} + \|\Gamma^2\|_2 \|i^{(m)}\|_{\mathcal{L}_i^2} \\
&\quad + e\|\Upsilon \Gamma J_6 w^{(m)}\|_{\mathcal{L}_i^2} + e\|\Upsilon \Gamma N J_7 f(v^{(m)})\|_{\mathcal{L}_i^2} + e\|\Upsilon \Gamma g\|_{\mathcal{L}_i^2}.
\end{aligned}$$

Therefore, it follows from the same inequalities used to derive (30) that, for some $\alpha_3 > 0$,

$$\begin{aligned} \|d_t^2 i^{(m)}\|_{L^2(0,T;\mathcal{L}_i^{2*})}^2 &\leq \alpha_3 \left([\hat{\kappa}_i + e^2 \kappa_w \|\Upsilon \Gamma J_6\|_2^2 + e^2 |\Omega| (F_E^2 + F_I^2) \|\Upsilon \Gamma N J_7\|_2^2] T \right. \\ &\quad \left. + e^2 \|\Upsilon \Gamma\|_2^2 \|g\|_{L^2(0,T;\mathcal{L}_i^2)}^2 \right). \end{aligned}$$

This, together with (30), establishes the bound (23) with (26) for some $\alpha_i > 0$.

Finally, multiplying (18) by $c_{v_k}^{(m)}$ and summing over $k = 1, \dots, m$ yields

$$(31) \quad \begin{aligned} &(\Phi d_t v^{(m)}, v^{(m)})_{\mathcal{L}_v^2} + (v^{(m)}, v^{(m)})_{\mathcal{L}_v^2} - (J_1 i^{(m)}, v^{(m)})_{\mathcal{L}_v^2} \\ &\quad + \left(J_2 v^{(m)} i^{(m)\top} \Psi J_4 + J_3 v^{(m)} i^{(m)\top} \Psi J_5, v^{(m)} \right)_{\mathcal{L}_v^2} = 0. \end{aligned}$$

Now, using Young's inequality and recalling (23), we obtain, for every $\varepsilon_5 > 0$,

$$\begin{aligned} (J_1 i^{(m)}, v^{(m)})_{\mathcal{L}_v^2} &\leq \varepsilon_5 \|v^{(m)}\|_{\mathcal{L}_v^2}^2 + \frac{1}{4\varepsilon_5} \|J_1 i^{(m)}\|_{\mathcal{L}_v^2}^2 \\ &\leq \varepsilon_5 \|v^{(m)}\|_{\mathcal{L}_v^2}^2 + \frac{1}{2\varepsilon_5} \|i^{(m)}\|_{\mathcal{L}_v^2}^2 \\ &\leq \varepsilon_5 \|v^{(m)}\|_{\mathcal{L}_v^2}^2 + \frac{\kappa_i}{2\varepsilon_5}. \end{aligned}$$

Moreover, using Hölder's inequality in $\mathbb{R}^2$ and the Cauchy–Schwarz inequality in $\mathbb{R}^4$, we obtain

$$\begin{aligned} &-\left(J_2 v^{(m)} i^{(m)\top} \Psi J_4 + J_3 v^{(m)} i^{(m)\top} \Psi J_5, v^{(m)} \right)_{\mathcal{L}_v^2} \\ &= - \int_{\Omega} \left[\left(v_1^{(m)} \right)^2 i^{(m)\top} \Psi J_4 + \left(v_2^{(m)} \right)^2 i^{(m)\top} \Psi J_5 \right] dx \\ &\leq \int_{\Omega} \|v^{(m)}\|_{\mathbb{R}^2}^2 \max \left\{ \left| i^{(m)\top} \Psi J_4 \right|, \left| i^{(m)\top} \Psi J_5 \right| \right\} dx \\ &\leq \int_{\Omega} \|v^{(m)}\|_{\mathbb{R}^2}^2 \|i^{(m)}\|_{\mathbb{R}^4} \max \{ \|\Psi J_4\|_{\mathbb{R}^4}, \|\Psi J_5\|_{\mathbb{R}^4} \} dx \\ &\leq \sqrt{2\kappa_i} \|\Psi\|_2 \|v^{(m)}\|_{\mathcal{L}_v^2}^2. \end{aligned}$$

Therefore, (31) implies

$$d_t \left\| \Phi^{\frac{1}{2}} v^{(m)} \right\|_{\mathcal{L}_v^2}^2 + 2 \left(1 - \varepsilon_5 - \sqrt{2\kappa_i} \|\Psi\|_2 \right) \|v^{(m)}\|_{\mathcal{L}_v^2}^2 \leq \frac{\kappa_i}{\varepsilon_5}.$$

Next, setting $\varepsilon_5 = 1$ and using Grönwall's inequality [51, sect. III.1.1.3.] yields

$$(32) \quad \sup_{t \in [0, T]} \left(\|v^{(m)}(t)\|_{\mathcal{L}_v^2}^2 \right) \leq \hat{\kappa}_v,$$

where, for some $\hat{\alpha}_v > 0$ and $\hat{\beta}_v > 0$,

$$\hat{\kappa}_v = \hat{\alpha}_v \exp \left(\hat{\beta}_v \sqrt{2\kappa_i} \|\Psi\|_2 T \right) \left(\|v_0\|_{\mathcal{L}_v^2}^2 + \frac{\kappa_i}{\sqrt{2\kappa_i} \|\Psi\|_2} \right).$$

Now, fix $\bar{\ell} \in \mathcal{L}_v^2$ such that $\|\bar{\ell}\|_{\mathcal{L}_v^2} \leq 1$ and decompose $\bar{\ell}$ as $\bar{\ell} = \ell + \ell^\perp$, where $\ell \in \text{span}\{\ell_v^{(k)}\}_{k=1}^m$ and $(\Phi \ell_v^{(k)}, \ell^\perp)_{\mathcal{L}_v^2} = 0$, $k = 1, \dots, m$. Note that this decomposition exists due to the way in which we construct the basis $\mathcal{B}_v$ in (14), wherein the elements, weighted by $\Phi^{\frac{1}{2}}$, are orthonormal in $\mathcal{L}_v^2$. Then, it follows from (15) and (18) that

$$\begin{aligned} \langle \Phi d_t v^{(m)}, \bar{\ell} \rangle_{\mathcal{L}_v^2} &= (\Phi d_t v^{(m)}, \bar{\ell})_{\mathcal{L}_v^2} = (\Phi d_t v^{(m)}, \ell)_{\mathcal{L}_v^2} \\ &= -(v^{(m)}, \ell)_{\mathcal{L}_v^2} + (J_1 i^{(m)}, \ell)_{\mathcal{L}_v^2} - \left(J_2 v^{(m)} i^{(m)\top} \Psi J_4 + J_3 v^{(m)} i^{(m)\top} \Psi J_5, \ell \right)_{\mathcal{L}_v^2}. \end{aligned}$$

Since $\mathcal{B}_v$ is a $\Phi^{\frac{1}{2}}$ -weighted orthonormal set in $\mathcal{L}_v^2$, it follows that

$$\|\ell\|_{\mathcal{L}_v^2} \leq \|\Phi^{-\frac{1}{2}}\|_2 \|\Phi^{\frac{1}{2}} \ell\|_{\mathcal{L}_v^2} \leq \|\Phi^{-\frac{1}{2}}\|_2 \|\Phi^{\frac{1}{2}} \bar{\ell}\|_{\mathcal{L}_v^2} \leq \|\Phi^{-\frac{1}{2}}\|_2 \|\Phi^{\frac{1}{2}}\|_2 \|\bar{\ell}\|_{\mathcal{L}_v^2} \leq \|\Phi^{-\frac{1}{2}}\|_2 \|\Phi^{\frac{1}{2}}\|_2$$

and hence, letting $\alpha_4 := \|\Phi^{-\frac{1}{2}}\|_2 \|\Phi^{\frac{1}{2}}\|_2$ and using the Cauchy–Schwarz inequality, we have

$$\begin{aligned} \left| \langle \Phi d_t v^{(m)}, \bar{\ell} \rangle_{\mathcal{L}_v^2} \right| &\leq \alpha_4 \left(\|v^{(m)}\|_{\mathcal{L}_v^2} + \|J_1 i^{(m)}\|_{\mathcal{L}_v^2} + \|J_2 v^{(m)} i^{(m)\top} \Psi J_4 + J_3 v^{(m)} i^{(m)\top} \Psi J_5\|_{\mathcal{L}_v^2} \right) \\ &\leq \alpha_4 \left(\|v^{(m)}\|_{\mathcal{L}_v^2} + \sqrt{2} \|i^{(m)}\|_{\mathcal{L}_i^2} + 2\sqrt{2} \|v^{(m)}\|_{\mathcal{L}_v^2} \|i^{(m)}\|_{\mathcal{L}_i^2} \|\Psi\|_2 \right) \\ &\leq \alpha_4 \left((1 + 2\sqrt{2\kappa_i} \|\Psi\|_2) \|v^{(m)}\|_{\mathcal{L}_v^2} + \sqrt{2\kappa_i} \right), \end{aligned}$$

which along with (32) implies that, for some $\alpha_5 > 0$,

$$\|d_t v^{(m)}\|_{L^2(0,T;\mathcal{L}_v^{2*})}^2 \leq \alpha_5 \left((1 + 2\sqrt{2\kappa_i} \|\Psi\|_2)^2 \hat{\kappa}_v + 2\kappa_i \right) T.$$

This, together with (32), establishes the bound (22) with (25) for some $\alpha_v > 0$. Note that constants $\alpha_1, \dots, \alpha_5, \hat{\alpha}_v, \hat{\beta}_v, \hat{\alpha}_i$, and $\hat{\alpha}_w$ depend only on the parameters of the model, which further implies that the constants $\alpha_v, \beta_v, \alpha_i$, and α_w also depend only on the parameters of the model and completes the proof. $\blacksquare$

Theorem 4.5 (Existence and uniqueness of weak solutions). *Suppose that $g \in L^2(0, T; \mathcal{L}_i^2)$, $v_0 \in \mathcal{L}_v^2$, $i_0 \in \mathcal{L}_i^2$, $i'_0 \in \mathcal{L}_i^2$, $w_0 \in \mathcal{H}_w^1$, and $w'_0 \in \mathcal{L}_w^2$. Then there exists a unique Ω -periodic weak solution (v, i, w) of the initial value problem (4)–(7).*

Proof. The energy estimate (22) implies that the sequence $\{v^{(m)}\}_{m=1}^\infty$ is bounded in $L^2(0, T; \mathcal{L}_v^2)$ and the sequence $\{d_t v^{(m)}\}_{m=1}^\infty$ is bounded in $L^2(0, T; \mathcal{L}_v^{2*})$. Since $\mathcal{L}_v^{2*} = \mathcal{L}_v^2$, it follows that $\{v^{(m)}\}_{m=1}^\infty$ is bounded in $H^1(0, T; \mathcal{L}_v^2)$ and $\{d_t v^{(m)}\}_{m=1}^\infty$ is bounded in $L^2(0, T; \mathcal{L}_v^2)$. Similarly, since $\mathcal{L}_i^{2*} = \mathcal{L}_i^2$, the energy estimate (23) implies that the sequence $\{i^{(m)}\}_{m=1}^\infty$ is bounded in $H^2(0, T; \mathcal{L}_i^2)$, the sequence $\{d_t i^{(m)}\}_{m=1}^\infty$ is bounded in $H^1(0, T; \mathcal{L}_i^2)$, and the sequence $\{d_t^2 i^{(m)}\}_{m=1}^\infty$ is bounded in $L^2(0, T; \mathcal{L}_i^2)$. Finally, the energy estimate (24) implies that the sequence $\{w^{(m)}\}_{m=1}^\infty$ is bounded in $L^2(0, T; \mathcal{H}_w^1)$, the sequence $\{d_t w^{(m)}\}_{m=1}^\infty$ is bounded in $L^2(0, T; \mathcal{L}_w^2)$, and the sequence $\{d_t^2 w^{(m)}\}_{m=1}^\infty$ is bounded in $L^2(0, T; \mathcal{H}_w^{1*})$. Now, it follows from the Rellich–Kondrachov compact embedding theorems [9, Thm. 6.6–3] that $H^1(0, T; \mathcal{L}_v^2) \Subset L^2(0, T; \mathcal{L}_v^2)$ and $H^1(0, T; \mathcal{L}_i^2) \Subset L^2(0, T; \mathcal{L}_i^2)$. Therefore, by [9, Thm. 2.10-1b], there exist subsequences $\{v^{(m_k)}\}_{k=1}^\infty$, $\{i^{(m_k)}\}_{k=1}^\infty$, and $\{d_t i^{(m_k)}\}_{k=1}^\infty$ such that

$$(33) \quad \begin{aligned} v^{(m_k)} &\rightarrow v \quad \text{strongly in } L^2(0, T; \mathcal{L}_v^2), \\ i^{(m_k)} &\rightarrow i \quad \text{strongly in } L^2(0, T; \mathcal{L}_i^2), \\ d_t i^{(m_k)} &\rightarrow i' \quad \text{strongly in } L^2(0, T; \mathcal{L}_i^2). \end{aligned}$$

Moreover, by the Banach–Eberlein–Šmulian theorem [9, Thm. 5.14-4], there exist subsequences $\{d_t v^{(m_k)}\}_{k=1}^\infty$, $d_t^2 \{i^{(m_k)}\}_{k=1}^\infty$, $\{w^{(m_k)}\}_{k=1}^\infty$, $\{d_t w^{(m_k)}\}_{k=1}^\infty$, and $\{d_t^2 w^{(m_k)}\}_{k=1}^\infty$ such that

$$(34) \quad \begin{aligned} d_t v^{(m_k)} &\rightharpoonup v' \quad \text{weakly in } L^2(0, T; \mathcal{L}_v^2), \\ d_t^2 i^{(m_k)} &\rightharpoonup i'' \quad \text{weakly in } L^2(0, T; \mathcal{L}_i^2), \\ w^{(m_k)} &\rightharpoonup w \quad \text{weakly in } L^2(0, T; \mathcal{H}_w^1), \\ d_t w^{(m_k)} &\rightharpoonup w' \quad \text{weakly in } L^2(0, T; \mathcal{L}_w^2), \\ d_t^2 w^{(m_k)} &\rightharpoonup w'' \quad \text{weakly in } L^2(0, T; \mathcal{H}_w^{1*}), \end{aligned}$$

where the time derivatives in the above analysis are derivatives in the weak sense.

Next, we show that

$$v' = d_t v, \quad i' = d_t i, \quad i'' = d_t^2 i, \quad w' = d_t w, \quad w'' = d_t^2 w.$$

Since $L^2(0, T; \mathcal{H}_w^1)$ is reflexive, the weak and weak* convergences coincide. Recalling the definitions of weak* convergence and weak derivatives, it follows that, for every $h \in \mathcal{H}_w^1$ and $\phi \in C_c^\infty([0, T])$,

$$\begin{aligned} \left\langle \int_0^T w'' \phi dt, h \right\rangle_{\mathcal{H}_w^1} &= \int_0^T \langle w'' \phi, h \rangle_{\mathcal{H}_w^1} dt = \lim_{k \rightarrow \infty} \int_0^T \langle d_t^2 w^{(m_k)} \phi, h \rangle_{\mathcal{H}_w^1} dt \\ &= \lim_{k \rightarrow \infty} \left\langle \int_0^T d_t^2 w^{(m_k)} \phi dt, h \right\rangle_{\mathcal{H}_w^1} = \lim_{k \rightarrow \infty} \left\langle (-1)^2 \int_0^T w^{(m_k)} d_t^2 \phi dt, h \right\rangle_{\mathcal{H}_w^1} \\ &= \lim_{k \rightarrow \infty} (-1)^2 \int_0^T \langle w^{(m_k)} d_t^2 \phi, h \rangle_{\mathcal{H}_w^1} dt = (-1)^2 \int_0^T \langle w d_t^2 \phi, h \rangle_{\mathcal{H}_w^1} dt \\ &= \left\langle (-1)^2 \int_0^T w d_t^2 \phi dt, h \right\rangle_{\mathcal{H}_w^1}, \end{aligned}$$

which implies $w'' = d_t^2 w$ in the weak sense. The other identities are proved similarly.

Now, recall (3) and (8) and note that the nonlinear map $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is bounded and smooth, and in particular is Lipschitz continuous. Therefore, it follows from the strong convergence of $\{v^{(m_k)}\}_{k=1}^\infty$ in (33) that

$$(35) \quad f(v^{(m_k)}) \rightarrow f(v) \quad \text{strongly in } L^2(0, T; \mathcal{L}_v^2).$$

For the bilinear term $J_2 v i^T \Psi J_4$, use (22) and (23) to write

$$\begin{aligned} &\left\| J_2 \left(v i^T - v^{(m_k)} i^{(m_k)T} \right) \Psi J_4 \right\|_{L^2(0, T; \mathcal{L}_v^2)} \\ &\leq \left\| J_2 (v - v^{(m_k)}) i^T \Psi J_4 \right\|_{L^2(0, T; \mathcal{L}_v^2)} + \left\| J_2 v^{(m_k)} (i - i^{(m_k)})^T \Psi J_4 \right\|_{L^2(0, T; \mathcal{L}_v^2)} \end{aligned}$$

$$\begin{aligned}
&\leq \sqrt{2}\|\Psi\|_2 \left[\|v - v^{(m_k)}\|_{L^2(0,T;\mathcal{L}_v^2)} \|i\|_{L^2(0,T;\mathcal{L}_i^2)} + \|v^{(m_k)}\|_{L^2(0,T;\mathcal{L}_v^2)} \|i - i^{(m_k)}\|_{L^2(0,T;\mathcal{L}_i^2)} \right] \\
&\leq \sqrt{2}\|\Psi\|_2 \left[\sqrt{\kappa_i} \|v - v^{(m_k)}\|_{L^2(0,T;\mathcal{L}_v^2)} + \sqrt{\kappa_v} \|i - i^{(m_k)}\|_{L^2(0,T;\mathcal{L}_i^2)} \right].
\end{aligned}$$

The same inequality holds for the bilinear term $J_3 v i^T \Psi J_5$ as well. Therefore, (33) gives

$$\begin{aligned}
(36) \quad &J_2 v^{(m_k)} i^{(m_k)T} \Psi J_4 \rightarrow J_2 v i^T \Psi J_4 \quad \text{strongly in } L^2(0, T; \mathcal{L}_v^2), \\
&J_3 v^{(m_k)} i^{(m_k)T} \Psi J_5 \rightarrow J_3 v i^T \Psi J_5 \quad \text{strongly in } L^2(0, T; \mathcal{L}_v^2).
\end{aligned}$$

Next, fix a positive integer K and choose the functions

$$\begin{aligned}
\hat{v} &= \sum_{k=1}^K c_{v_k}(t) \ell_v^{(k)} \in C^1([0, T]; \mathcal{L}_v^2), \\
\hat{i} &= \sum_{k=1}^K c_{i_k}(t) \ell_i^{(k)} \in C^1([0, T]; \mathcal{L}_i^2), \\
\hat{w} &= \sum_{k=1}^K c_{w_k}(t) h_w^{(k)} \in C^1([0, T]; \mathcal{H}_w^1),
\end{aligned}$$

where c_{v_k} , c_{i_k} , and c_{w_k} are sufficiently smooth functions on $[0, T]$, and $(\ell_v^{(k)}, \ell_i^{(k)}, h_w^{(k)})$, $k = 1, \dots, K$, are the first K components of $\mathcal{B}$ given by (14). Set $m = m_k$ in (18)–(20) and choose $m_k \geq K$. Then, multiplying (18)–(20) by c_{v_k} , c_{i_k} , and c_{w_k} , respectively, summing over $k = 1, \dots, K$, and integrating over $t \in [0, T]$ yields

$$\begin{aligned}
(37) \quad &\int_0^T \left[\langle \Phi d_t v^{(m_k)}, \hat{v} \rangle_{\mathcal{L}_v^2} + (v^{(m_k)}, \hat{v})_{\mathcal{L}_v^2} - (J_1 i^{(m_k)}, \hat{v})_{\mathcal{L}_v^2} \right. \\
&\quad \left. + \left(J_2 v^{(m_k)} i^{(m_k)T} \Psi J_4 + J_3 v^{(m_k)} i^{(m_k)T} \Psi J_5, \hat{v} \right)_{\mathcal{L}_v^2} \right] dt = 0, \\
&\int_0^T \left[\langle d_t^2 i^{(m_k)}, \hat{i} \rangle_{\mathcal{L}_i^2} + 2(\Gamma d_t i^{(m_k)}, \hat{i})_{\mathcal{L}_i^2} + (\Gamma^2 i^{(m_k)}, \hat{i})_{\mathcal{L}_i^2} \right. \\
&\quad \left. - e(\Upsilon \Gamma J_6 w^{(m_k)}, \hat{i})_{\mathcal{L}_i^2} - e(\Upsilon \Gamma N J_7 f(v^{(m_k)}), \hat{i})_{\mathcal{L}_i^2} - e(\Upsilon \Gamma g, \hat{i})_{\mathcal{L}_i^2} \right] dt = 0, \\
&\int_0^T \left[\langle d_t^2 w^{(m_k)}, \hat{w} \rangle_{\mathcal{L}_w^2} + 2\nu(\Lambda d_t w^{(m_k)}, \hat{w})_{\mathcal{L}_w^2} + \frac{3}{2}\nu^2(\partial_x w^{(m_k)}, \partial_x \hat{w})_{\mathcal{L}_{\partial w}^2} \right. \\
&\quad \left. + \nu^2(\Lambda^2 w^{(m_k)}, \hat{w})_{\mathcal{L}_w^2} - \nu^2(\Lambda^2 M J_8 f(v^{(m)}), \hat{w})_{\mathcal{L}_w^2} \right] dt = 0.
\end{aligned}$$

Note that the families of functions $\hat{v}$, $\hat{i}$, and $\hat{w}$ chosen above are dense in the spaces $L^2(0, T; \mathcal{L}_v^2)$, $L^2(0, T; \mathcal{L}_i^2)$, and $L^2(0, T; \mathcal{H}_w^1)$, respectively. Therefore, (37) holds for all functions $\hat{v} \in L^2(0, T; \mathcal{L}_v^2)$, $\hat{i} \in L^2(0, T; \mathcal{L}_i^2)$, and $\hat{w} \in L^2(0, T; \mathcal{H}_w^1)$. Now, use (33)–(36) to pass to the limits in (37), which implies that (10)–(12) hold for all $\ell_v \in \mathcal{L}_v^2$, $\ell_i \in \mathcal{L}_i^2$, $h_w \in \mathcal{H}_w^1$, and almost every $t \in [0, T]$.

It remains to verify the initial conditions (13). Choose the functions

$$\hat{v} \in C^1([0, T]; \mathcal{L}_v^2), \quad \hat{i} \in C^2([0, T]; \mathcal{L}_i^2), \quad \hat{w} \in C^2([0, T]; \mathcal{H}_w^1)$$

such that these functions vanish at the end point $t = T$. Integrating by parts in (37) yields

$$(38) \quad \begin{aligned} \int_0^T \left[-(\Phi v^{(m_k)}, d_t \hat{v})_{\mathcal{L}_v^2} + \dots \right] dt &= (\Phi v^{(m_k)}(0), \hat{v}(0))_{\mathcal{L}_v^2}, \\ \int_0^T \left[(i^{(m_k)}, d_t^2 \hat{i})_{\mathcal{L}_i^2} + \dots \right] dt &= \dots + (d_t i^{(m_k)}(0), \hat{i}(0))_{\mathcal{L}_i^2} - (i^{(m_k)}(0), d_t \hat{i}(0))_{\mathcal{L}_i^2}, \\ \int_0^T \left[(w^{(m_k)}, d_t^2 \hat{w})_{\mathcal{H}_w^1} + \dots \right] dt &= (d_t w^{(m_k)}(0), \hat{w}(0))_{\mathcal{L}_w^2} - (w^{(m_k)}(0), d_t \hat{w}(0))_{\mathcal{L}_w^2}, \end{aligned}$$

where “...” denotes terms that are not pertinent to the analysis. Similarly, integrating by parts in the limit of (37) yields

$$(39) \quad \begin{aligned} \int_0^T \left[-(\Phi v, d_t \hat{v})_{\mathcal{L}_v^2} + \dots \right] dt &= (\Phi v(0), \hat{v}(0))_{\mathcal{L}_v^2}, \\ \int_0^T \left[(i, d_t^2 \hat{i})_{\mathcal{L}_i^2} + \dots \right] dt &= \dots + (d_t i(0), \hat{i}(0))_{\mathcal{L}_i^2} - (i(0), d_t \hat{i}(0))_{\mathcal{L}_i^2}, \\ \int_0^T \left[(w, d_t^2 \hat{w})_{\mathcal{H}_w^1} + \dots \right] dt &= (d_t w(0), \hat{w}(0))_{\mathcal{L}_w^2} - (w(0), d_t \hat{w}(0))_{\mathcal{L}_w^2}. \end{aligned}$$

Now, consider the initial conditions (21), pass to the limits in (38) through (33)–(36), and compare the results with (39). Since $\hat{v}$, $\hat{i}$, and $\hat{w}$ are arbitrary, the initial condition (13) holds and this completes the proof of existence.

To prove uniqueness, assume by contradiction that there exist two weak solutions $(\tilde{v}, \tilde{i}, \tilde{w})$ and $(\hat{v}, \hat{i}, \hat{w})$ for (2), initiating from the same initial values, such that $(\tilde{v}, \tilde{i}, \tilde{w}) \neq (\hat{v}, \hat{i}, \hat{w})$. Then, $(v, i, w) := (\tilde{v}, \tilde{i}, \tilde{w}) - (\hat{v}, \hat{i}, \hat{w})$ is a weak solution initiating from the zero initial condition $(v_0, i_0, i'_0, w_0, w'_0) = 0$. Now, fix $s \in [0, T]$ and define, for $0 \leq t \leq T$, the functions

$$(40) \quad p(t) := \int_0^t w(r) dr, \quad q(t) := \begin{cases} \int_t^s w(r) dr & \text{if } 0 \leq t \leq s, \\ 0 & \text{if } s < t \leq T. \end{cases}$$

Note that $p(t) \in \mathcal{H}_w^1$ and $q(t) \in \mathcal{H}_w^1$ for all $t \in [0, T]$, and hence p and q are regular enough to be used as the test function h_w in (12). Moreover, $q(s) = 0$, $q(0) = p(s)$, and $p(0) = 0$. Let $\tilde{u}$ and $\hat{u}$ satisfy (10)–(12) with the same test functions $\ell_v = v(t)$, $\ell_i = d_t i(t)$, and $h_w = q(t)$. Subtracting the two sets of equations and integrating over $t \in [0, s]$ yields

$$(41) \quad \begin{aligned} \int_0^s \left[\langle \Phi d_t v, v \rangle_{\mathcal{L}_v^2} + (v, v)_{\mathcal{L}_v^2} - (J_1 i, v)_{\mathcal{L}_v^2} \right. \\ \left. + \left(J_2 (\tilde{v} \tilde{i}^T - \hat{v} \hat{i}^T) \Psi J_4 + J_3 (\tilde{v} \tilde{i}^T - \hat{v} \hat{i}^T) \Psi J_5, v \right)_{\mathcal{L}_v^2} \right] dt = 0, \end{aligned}$$

$$(42) \quad \begin{aligned} \int_0^s \left[\langle d_t^2 i, d_t i \rangle_{\mathcal{L}_i^2} + 2(\Gamma d_t i, d_t i)_{\mathcal{L}_i^2} + (\Gamma^2 i, d_t i)_{\mathcal{L}_i^2} - e(\Upsilon \Gamma J_6 w, d_t i)_{\mathcal{L}_i^2} \right. \\ \left. - e(\Upsilon \Gamma N J_7 (f(\tilde{v}) - f(\hat{v})), d_t i)_{\mathcal{L}_i^2} \right] dt = 0, \end{aligned}$$

$$(43) \quad \int_0^s \left[\langle d_t^2 w, q \rangle_{\mathcal{H}_w^1} + 2\nu(\Lambda d_t w, q)_{\mathcal{L}_w^2} + \frac{3}{2}\nu^2(\partial_x w, \partial_x q)_{\mathcal{L}_{\partial w}^2} + \nu^2(\Lambda^2 w, q)_{\mathcal{L}_w^2} \right. \\ \left. - \nu^2(\Lambda^2 M J_8(f(\tilde{v}) - f(\hat{v})), q)_{\mathcal{L}_w^2} \right] dt = 0.$$

Next, integrating by parts in the first and second terms in (43) yields

$$\int_0^s \left[- (d_t w, d_t q)_{\mathcal{L}_w^2} - 2\nu(\Lambda w, d_t q)_{\mathcal{L}_w^2} + \frac{3}{2}\nu^2(\partial_x w, \partial_x q)_{\mathcal{L}_{\partial w}^2} + \nu^2(\Lambda^2 w, q)_{\mathcal{L}_w^2} \right] dt \\ = \int_0^s \nu^2(\Lambda^2 M J_8(f(\tilde{v}) - f(\hat{v})), q)_{\mathcal{L}_w^2} dt.$$

Note that $\langle d_t w, d_t q \rangle_{\mathcal{H}_w^1} = (d_t w, d_t q)_{\mathcal{L}_w^2}$, since $d_t w \in \mathcal{L}_w^2$ for almost every $t \in [0, T]$; see the proof of [16, Thm. 5.9-1]. Now, it follows from the definition of $q(t)$ that $d_t q = -w$ for all $t \in [0, s]$. Therefore,

$$(44) \quad \int_0^s \left[\frac{1}{2} d_t \left(\|w\|_{\mathcal{L}_w^2}^2 - \frac{3}{2}\nu^2 \|\partial_x q\|_{\mathcal{L}_{\partial w}^2}^2 \right) + 2\nu \|\Lambda^{\frac{1}{2}} w\|_{\mathcal{L}_w^2}^2 + \nu^2(\Lambda^2 w, q)_{\mathcal{L}_w^2} \right] dt \\ = \int_0^s \nu^2(\Lambda^2 M J_8(f(\tilde{v}) - f(\hat{v})), q)_{\mathcal{L}_w^2} dt.$$

Using Young's inequality,

$$\nu^2(\Lambda^2 M J_8(f(\tilde{v}) - f(\hat{v})), q)_{\mathcal{L}_w^2} \leq \frac{1}{4}\nu^2 \|q\|_{\mathcal{L}_w^2}^2 + \nu^2 \operatorname{tr}(\Lambda^4 M^2) \left[\sup_{v_E(x,t) \in \mathbb{R}} |\partial_{v_E} f_E(v_E)| \right]^2 \|v\|_{\mathcal{L}_v^2}^2 \\ \leq \frac{1}{4}\nu^2 \|q\|_{\mathcal{L}_w^2}^2 + \frac{1}{8}\nu^2 \frac{F_E^2}{\sigma_E^2} \operatorname{tr}(\Lambda^4 M^2) \|v\|_{\mathcal{L}_v^2}^2, \\ -\nu^2(\Lambda^2 w, q)_{\mathcal{L}_w^2} \leq \frac{1}{4}\nu^2 \|q\|_{\mathcal{L}_w^2}^2 + \nu^2 \|\Lambda\|_2^4 \|w\|_{\mathcal{L}_w^2}^2,$$

where the second inequality follows, for $x = E$, from differentiating (3) as

$$(45) \quad \partial_{v_x} f_x(v_x) = \frac{\sqrt{2}}{\sigma_x} F_x \exp \left(-\sqrt{2} \frac{v_x - \mu_x}{\sigma_x} \right) \left[1 + \exp \left(-\sqrt{2} \frac{v_x - \mu_x}{\sigma_x} \right) \right]^{-2}, \quad x \in \{E, I\},$$

which implies $\sup_{v_x(x,t) \in \mathbb{R}} |\partial_{v_x} f_x(v_x)| \leq \frac{F_x}{2\sqrt{2}\sigma_x}$.

Now, (44) implies

$$\frac{1}{2} \|w(s)\|_{\mathcal{L}_w^2}^2 + \frac{3}{4}\nu^2 \|q(0)\|_{\mathcal{H}_w^1}^2 \leq \int_0^s \left[\left(-2\nu \Lambda_{\min} + \nu^2 \|\Lambda\|_2^4 \right) \|w\|_{\mathcal{L}_w^2}^2 + \frac{1}{2}\nu^2 \|q\|_{\mathcal{L}_w^2}^2 \right. \\ \left. + \frac{1}{8}\nu^2 \frac{F_E^2}{\sigma_E^2} \operatorname{tr}(\Lambda^4 M^2) \|v\|_{\mathcal{L}_v^2}^2 \right] dt + \frac{3}{4}\nu^2 \|q(0)\|_{\mathcal{L}_w^2}^2,$$

where $\Lambda_{\min} := \min\{\Lambda_{EE}, \Lambda_{EI}\}$ is the smallest eigenvalue of Λ . Noting from (40) that $q(t) = p(s) - p(t)$ for all $t \in [0, s]$, it follows that the above inequality can be written as

$$\begin{aligned} \frac{1}{2}\|w(s)\|_{\mathcal{L}_w^2}^2 + \frac{3}{4}\nu^2\|p(s)\|_{\mathcal{H}_w^1}^2 &\leq \int_0^s \left[\left(-2\nu\Lambda_{\min} + \nu^2\|\Lambda\|_2^4 \right) \|w(t)\|_{\mathcal{L}_w^2}^2 + \frac{1}{2}\nu^2\|p(s) - p(t)\|_{\mathcal{L}_w^2}^2 \right. \\ &\quad \left. + \frac{1}{8}\nu^2\frac{F_E^2}{\sigma_E^2} \operatorname{tr}(\Lambda^4 M^2) \|v(t)\|_{\mathcal{L}_v^2}^2 \right] dt + \frac{3}{4}\nu^2\|p(s)\|_{\mathcal{L}_w^2}^2. \end{aligned}$$

Using the Cauchy–Schwarz inequality, it follows from the definition of $p(t)$ given by (40) that $\|p(s)\|_{\mathcal{L}_w^2}^2 \leq s \int_0^s \|w(t)\|_{\mathcal{L}_w^2}^2 dt$. Moreover,

$$\|p(s) - p(t)\|_{\mathcal{L}_w^2}^2 \leq 2\|p(s)\|_{\mathcal{L}_w^2}^2 + 2\|p(t)\|_{\mathcal{L}_w^2}^2 \leq 2\|p(s)\|_{\mathcal{H}_w^1}^2 + 2\|p(t)\|_{\mathcal{H}_w^1}^2.$$

Therefore,

$$(46) \quad \frac{1}{2}\|w(s)\|_{\mathcal{L}_w^2}^2 + \nu^2 \left(\frac{3}{4} - s \right) \|p(s)\|_{\mathcal{H}_w^1}^2 \leq \int_0^s \left[\left(-2\nu\Lambda_{\min} + \nu^2\|\Lambda\|_2^4 + \frac{3}{4}\nu^2 s \right) \|w(t)\|_{\mathcal{L}_w^2}^2 \right. \\ \left. + \nu^2\|p(t)\|_{\mathcal{H}_w^1}^2 + \frac{1}{8}\nu^2\frac{F_E^2}{\sigma_E^2} \operatorname{tr}(\Lambda^4 M^2) \|v(t)\|_{\mathcal{L}_v^2}^2 \right] dt.$$

Next, recalling (22) and (23) and using the Cauchy–Schwarz and Young inequalities, it follows that the fourth term in (41) satisfies, for every $\varepsilon_1 > 0$,

$$\begin{aligned} \left(J_2(\tilde{v}i^T - \hat{v}i^T)\Psi J_4, v \right)_{\mathcal{L}_v^2} &= (J_2\tilde{v}i^T\Psi J_4, v)_{\mathcal{L}_v^2} + (J_2\hat{v}i^T\Psi J_4, v)_{\mathcal{L}_v^2} \\ &\geq -\sqrt{2\kappa_{\hat{v}}} \|\Psi\|_2 \|v\|_{\mathcal{L}_v^2}^2 - \varepsilon_1 \|v\|_{\mathcal{L}_v^2}^2 - \frac{2\kappa_{\hat{v}}}{4\varepsilon_1} \|\Psi\|_2^2 \|i\|_{\mathcal{L}_i^2}^2, \end{aligned}$$

where $\kappa_{\hat{v}}$ and $\kappa_{\hat{i}}$ are in the form of (25) and (26), respectively. The same inequality holds for $(J_3(\tilde{v}i^T - \hat{v}i^T)\Psi J_5, v)_{\mathcal{L}_v^2}$. Similarly, using Young's inequality and (45),

$$\begin{aligned} e(\Upsilon\Gamma N J_7(f(\tilde{v}) - f(\hat{v})), d_t i)_{\mathcal{L}_i^2} &\leq \varepsilon_2 \|d_t i\|_{\mathcal{L}_i^2}^2 + \frac{e^2}{4\varepsilon_2} \|\Upsilon\Gamma N J_7\|_2^2 \sup_{v(x,t) \in \mathbb{R}^2} \|\partial_v f(v)\|_2^2 \|v\|_{\mathcal{L}_v^2}^2 \\ &\leq \varepsilon_2 \|d_t i\|_{\mathcal{L}_i^2}^2 + \frac{e^2}{32\varepsilon_2} \|\Upsilon\Gamma N J_7\|_2^2 \max \left\{ \frac{F_E^2}{\sigma_E^2}, \frac{F_I^2}{\sigma_I^2} \right\} \|v\|_{\mathcal{L}_v^2}^2 \end{aligned}$$

for every $\varepsilon_2 > 0$. Moreover, for every $\varepsilon_3 > 0$ and $\varepsilon_4 > 0$,

$$\begin{aligned} (J_1 i, v)_{\mathcal{L}_v^2} &\leq \varepsilon_4 \|v\|_{\mathcal{L}_v^2}^2 + \frac{1}{2\varepsilon_4} \|i\|_{\mathcal{L}_i^2}^2, \\ e(\Upsilon\Gamma J_6 w, d_t i)_{\mathcal{L}_i^2} &\leq \varepsilon_4 \|d_t i\|_{\mathcal{L}_i^2}^2 + \frac{e^2}{4\varepsilon_4} \|\Upsilon\Gamma J_6\|_2^2 \|w\|_{\mathcal{L}_w^2}^2. \end{aligned}$$

Substituting the above inequalities into (41) and (42), and adding the resulting inequalities to (46) yields, for some $\alpha > 0$,

$$\begin{aligned} &\|\Phi^{\frac{1}{2}}v(s)\|_{\mathcal{L}_v^2}^2 + \|d_t i(s)\|_{\mathcal{L}_i^2}^2 + \|\Gamma i(s)\|_{\mathcal{L}_i^2}^2 + \|w(s)\|_{\mathcal{L}_w^2}^2 + \nu^2 \left(\frac{3}{2} - 2s \right) \|p(s)\|_{\mathcal{H}_w^1}^2 \\ &\leq \alpha \int_0^s \left[\|v(t)\|_{\mathcal{L}_v^2}^2 + \|d_t i(t)\|_{\mathcal{L}_i^2}^2 + \|i(t)\|_{\mathcal{L}_i^2}^2 + \|w(t)\|_{\mathcal{L}_w^2}^2 + \|p(t)\|_{\mathcal{H}_w^1}^2 \right] dt. \end{aligned}$$

Now, setting $T_1 = \frac{3}{4}$, it follows from the integral form of Grönwall's inequality [16, App. B.2] that $(v(s), i(s), w(s)) = 0$ for all $s \in [0, T_1]$. Repeating the same arguments for intervals $[T_1, 2T_1]$, $[2T_1, 3T_1]$, $\dots$, we deduce $(v(t), i(t), w(t)) = 0$ for all $t \in [0, T]$, and hence $(\tilde{v}, \tilde{i}, \tilde{w}) = (\hat{v}, \hat{i}, \hat{w})$ for all $t \in [0, T]$, which is a contradiction and completes the proof of uniqueness. ■

Proposition 4.6 (Regularity of weak solutions). *Suppose that the assumptions of Theorem 4.5 hold, namely, $g \in L^2(0, T; \mathcal{L}_i^2)$, $v_0 \in \mathcal{L}_v^2$, $i_0 \in \mathcal{L}_i^2$, $i'_0 \in \mathcal{L}_i^2$, $w_0 \in \mathcal{H}_w^1$, and $w'_0 \in \mathcal{L}_w^2$. Then the Ω -periodic weak solution (v, i, w) of the initial value problem (4)–(7) satisfies*

$$(47) \quad \begin{aligned} & \operatorname{ess\,sup}_{t \in [0, T]} \left(\|v(t)\|_{\mathcal{L}_v^2}^2 \right) + \|d_t v\|_{L^2(0, T; \mathcal{L}_v^2)}^2 \leq \kappa_v, \\ & \operatorname{ess\,sup}_{t \in [0, T]} \left(\|d_t i(t)\|_{\mathcal{L}_i^2}^2 + \|i(t)\|_{\mathcal{L}_i^2}^2 \right) + \|d_t^2 i\|_{L^2(0, T; \mathcal{L}_i^2)}^2 \leq \kappa_i, \\ & \operatorname{ess\,sup}_{t \in [0, T]} \left(\|d_t w(t)\|_{\mathcal{L}_w^2}^2 + \|w(t)\|_{\mathcal{H}_w^1}^2 \right) + \|d_t^2 w\|_{L^2(0, T; \mathcal{H}_w^{1*})}^2 \leq \kappa_w, \end{aligned}$$

$$(48) \quad \begin{aligned} v & \in H^3(0, T; \mathcal{L}_v^2) \cap C^{2, \frac{1}{2}}([0, T]; \mathcal{L}_v^2), \\ i & \in H^2(0, T; \mathcal{L}_i^2) \cap C^{1, \frac{1}{2}}([0, T]; \mathcal{L}_i^2), & d_t i & \in H^1(0, T; \mathcal{L}_i^2) \cap C^{0, \frac{1}{2}}([0, T]; \mathcal{L}_i^2), \\ w & \in H^1(0, T; \mathcal{L}_w^2) \cap C^0([0, T]; \mathcal{H}_w^1), & d_t w & \in C^0([0, T]; \mathcal{L}_w^2), \end{aligned}$$

where κ_v , κ_i , and κ_w are given by (25)–(27). Moreover, if $g \in C^0([0, T]; \mathcal{L}_i^2)$, then

$$(49) \quad v \in C^3([0, T]; \mathcal{L}_v^2), \quad i \in C^2([0, T]; \mathcal{L}_i^2), \quad d_t i \in C^1([0, T]; \mathcal{L}_i^2),$$

and if $g \in C^1([0, T]; \mathcal{L}_i^2)$, then

$$(50) \quad v \in C^4([0, T]; \mathcal{L}_v^2), \quad i \in C^3([0, T]; \mathcal{L}_i^2), \quad d_t i \in C^2([0, T]; \mathcal{L}_i^2).$$

Proof. First, recall that $\mathcal{L}_v^2 = \mathcal{L}_v^{2*}$ and $\mathcal{L}_i^2 = \mathcal{L}_i^{2*}$. Assertion (47) follows immediately from (22)–(24) by setting $m = m_k$ and passing to the limits through (33) and (34). The inclusions in H^1 , H^2 , and H^3 in assertion (48) are immediate from (47) and twice differentiation of (4). The Sobolev embedding theorems [9, Thm. 6.6-1] applied to Banach space-valued functions on $[0, T] \subset \mathbb{R}$ imply that $v \in C^{2, \frac{1}{2}}([0, T]; \mathcal{L}_v^2)$, $i \in C^{1, \frac{1}{2}}([0, T]; \mathcal{L}_i^2)$, and $d_t i \in C^{0, \frac{1}{2}}([0, T]; \mathcal{L}_i^2)$.

Consider the time-independent self-adjoint linear operator $A := (-\frac{3}{2}\nu^2\Delta + I) : \mathcal{H}_w^1 \rightarrow \mathcal{H}_w^{1*}$. Note that $f(v) \in C^{2, \frac{1}{2}}([0, T]; \mathcal{L}_v^\infty)$, since f is a bounded smooth function and $v \in C^{2, \frac{1}{2}}([0, T]; \mathcal{L}_v^2)$. Then, it follows from (6) and (47) that $d_t^2 w + Aw \in L^2(0, T; \mathcal{L}_w^2)$. Therefore, by [51, Lem. II.4.1], we have $w \in C^0([0, T]; \mathcal{H}_w^1)$ and $d_t w \in C^0([0, T]; \mathcal{L}_w^2)$, which completes the proof of (48). Assertions (49) and (50) are now immediate from (4), (5), and (48). ■

Theorem 4.7 (Existence and uniqueness of strong solutions). *Suppose that $g \in L^2(0, T; \mathcal{L}_i^2)$, $v_0 \in \mathcal{L}_v^2$, $i_0 \in \mathcal{L}_i^2$, $i'_0 \in \mathcal{L}_i^2$, $w_0 \in \mathcal{H}_w^2$, and $w'_0 \in \mathcal{H}_w^1$. Then there exists a unique Ω -periodic strong solution (v, i, w) of the initial value problem (4)–(7).*

Proof. Uniqueness follows immediately from Theorem 4.5, since every strong solution of (4)–(7) is also a weak solution. Moreover, Proposition 4.6 implies that the weak solutions

$v \in H^1(0, T; \mathcal{L}_w^2)$ and $i \in H^2(0, T; \mathcal{L}_i^2)$ are indeed strong solutions as given in Definition 4.3. It remains to prove the regularity required for w by Definition 4.3.

Consider (20) with the approximation (17), let $\mathcal{B}_w = \{h_w^{(k)}\}_{k=1}^\infty$ be the orthogonal basis of $\mathcal{H}_w^1$ consisting of the eigenfunctions of $A := -\Delta + I$ as given by Lemma 4.1, and let λ_k denote the eigenvalue corresponding to the eigenfunction $h_w^{(k)}$. Multiplying (20) by $\lambda_k c_{w_k}^{(m)}$ and summing over $k = 1, \dots, m$ yields

$$\begin{aligned} & (\mathrm{d}_t^2 w^{(m)}, Aw^{(m)})_{\mathcal{L}_w^2} + 2\nu(\Lambda \mathrm{d}_t w^{(m)}, Aw^{(m)})_{\mathcal{L}_w^2} + \frac{3}{2}\nu^2(\partial_x w^{(m)}, A\partial_x w^{(m)})_{\mathcal{L}_{\partial w}^2} \\ & + \nu^2(\Lambda^2 w^{(m)}, Aw^{(m)})_{\mathcal{L}_w^2} - \nu^2(\Lambda^2 \mathrm{MJ}_8 f(v^{(m)}), Aw^{(m)})_{\mathcal{L}_w^2} = 0. \end{aligned}$$

Now, Young's inequality implies that, for every $\varepsilon_1, \dots, \varepsilon_4 > 0$,

$$\begin{aligned} & -(\mathrm{d}_t^2 w^{(m)}, Aw^{(m)})_{\mathcal{L}_w^2} \leq \varepsilon_1 \|Aw^{(m)}\|_{\mathcal{L}_w^2}^2 + \frac{1}{4\varepsilon_1} \|\mathrm{d}_t^2 w^{(m)}\|_{\mathcal{L}_w^2}^2, \\ & -(\Lambda \mathrm{d}_t w^{(m)}, Aw^{(m)})_{\mathcal{L}_w^2} \leq \varepsilon_2 \|Aw^{(m)}\|_{\mathcal{L}_w^2}^2 + \frac{1}{4\varepsilon_2} \|\Lambda \mathrm{d}_t w^{(m)}\|_{\mathcal{L}_w^2}^2, \\ & -(\Lambda^2 w^{(m)}, Aw^{(m)})_{\mathcal{L}_w^2} \leq \varepsilon_3 \|Aw^{(m)}\|_{\mathcal{L}_w^2}^2 + \frac{1}{4\varepsilon_3} \|\Lambda^2 w^{(m)}\|_{\mathcal{L}_w^2}^2, \\ & (\Lambda^2 \mathrm{MJ}_8 f(v^{(m)}), Aw^{(m)})_{\mathcal{L}_w^2} \leq \varepsilon_4 \|Aw^{(m)}\|_{\mathcal{L}_w^2}^2 + \frac{1}{4\varepsilon_4} \|\Lambda^2 \mathrm{MJ}_8 f(v^{(m)})\|_{\mathcal{L}_w^2}^2 \\ & \leq \varepsilon_4 \|Aw^{(m)}\|_{\mathcal{L}_w^2}^2 + \frac{1}{4\varepsilon_4} |\Omega| F_E^2 \operatorname{tr}(\Lambda^4 M^2). \end{aligned}$$

Therefore, using (1),

$$\begin{aligned} \frac{3}{2}\nu^2 \|w^{(m)}\|_{\mathcal{H}_w^2}^2 & \leq (\varepsilon_1 + 2\nu\varepsilon_2 + \nu^2\varepsilon_3 + \nu^2\varepsilon_4) \left(\|w^{(m)}\|_{\mathcal{H}_w^2}^2 + \|\partial_x w^{(m)}\|_{\mathcal{L}_{\partial w}^2}^2 \right) \\ & + \frac{3}{2}\nu^2 \|w^{(m)}\|_{\mathcal{L}_w^2}^2 + \frac{1}{4\varepsilon_1} \|\mathrm{d}_t^2 w^{(m)}\|_{\mathcal{L}_w^2}^2 + \frac{\nu}{2\varepsilon_2} \|\Lambda \mathrm{d}_t w^{(m)}\|_{\mathcal{L}_w^2}^2 \\ & + \frac{\nu^2}{4\varepsilon_3} \|\Lambda^2 w^{(m)}\|_{\mathcal{L}_w^2}^2 + \frac{\nu^2}{4\varepsilon_4} |\Omega| F_E^2 \operatorname{tr}(\Lambda^4 M^2). \end{aligned}$$

Next, set $\varepsilon_1 = \frac{\nu^2}{8}$, $\varepsilon_2 = \frac{\nu}{16}$, $\varepsilon_3 = \frac{1}{8}$, and $\varepsilon_4 = \frac{1}{8}$, and note that, for some constant $\beta > 0$,

$$(51) \quad \|w^{(m)}\|_{\mathcal{H}_w^2}^2 \leq \beta \left(\|\mathrm{d}_t^2 w^{(m)}\|_{\mathcal{L}_w^2}^2 + \|\mathrm{d}_t w^{(m)}\|_{\mathcal{L}_w^2}^2 + \|w^{(m)}\|_{\mathcal{H}_w^1}^2 + |\Omega| F_E^2 \operatorname{tr}(\Lambda^4 M^2) \right).$$

Bounds on $\|\mathrm{d}_t w^{(m)}\|_{\mathcal{L}_w^2}$ and $\|w^{(m)}\|_{\mathcal{H}_w^1}$ are given by the energy estimate (24). To establish bounds on $\|\mathrm{d}_t^2 w^{(m)}\|_{\mathcal{L}_w^2}$ and $\|\mathrm{d}_t w^{(m)}\|_{\mathcal{H}_w^1}$, consider (20) with the initial values given in (21). Differentiating (20) with respect to t , multiplying the result by $\mathrm{d}_t^2 c_{w_k}^{(m)}$, and summing over $k = 1, \dots, m$ yields

$$\begin{aligned} & (\mathrm{d}_t^2 \dot{w}^{(m)}, \mathrm{d}_t \dot{w}^{(m)})_{\mathcal{L}_w^2} + 2\nu(\Lambda \mathrm{d}_t \dot{w}^{(m)}, \mathrm{d}_t \dot{w}^{(m)})_{\mathcal{L}_w^2} + \frac{3}{2}\nu^2(\partial_x \dot{w}^{(m)}, \mathrm{d}_t \partial_x \dot{w}^{(m)})_{\mathcal{L}_{\partial w}^2} \\ & + \nu^2(\Lambda^2 \dot{w}^{(m)}, \mathrm{d}_t \dot{w}^{(m)})_{\mathcal{L}_w^2} - \nu^2(\Lambda^2 \mathrm{MJ}_8 \mathrm{d}_t f(v^{(m)}), \mathrm{d}_t \dot{w}^{(m)})_{\mathcal{L}_w^2} = 0, \end{aligned}$$

where $\dot{w} := d_t w$ and $d_t f_E(v_E^{(m)}) = \partial_{v_E} f_E(v_E^{(m)}) d_t v_E^{(m)}$. Now, (45) with $x = E$ gives

$$(52) \quad \begin{aligned} \|\Lambda^2 M J_8 d_t f(v^{(m)})\|_{\mathcal{L}_w^2}^2 &= \text{tr}(\Lambda^4 M^2) \int_{\Omega} |d_t f_E(v_E^{(m)})|^2 dx \\ &\leq \text{tr}(\Lambda^4 M^2) \frac{F_E^2}{8\sigma_E^2} \int_{\Omega} |d_t v_E^{(m)}|^2 dx \leq \text{tr}(\Lambda^4 M^2) \frac{F_E^2}{8\sigma_E^2} \|d_t v^{(m)}\|_{\mathcal{L}_v^2}^2. \end{aligned}$$

Using similar arguments as in the proof of Proposition 4.4, it follows from the above inequality and Young's inequality that, for every $\varepsilon > 0$,

$$\begin{aligned} d_t \left[\|d_t \dot{w}^{(m)}\|_{\mathcal{L}_w^2}^2 + \frac{3}{2} \nu^2 \|\partial_x \dot{w}^{(m)}\|_{\mathcal{L}_{\partial w}^2}^2 + \nu^2 \|\Lambda \dot{w}^{(m)}\|_{\mathcal{L}_w^2}^2 \right] &+ 2\nu(2\Lambda_{\min} - \varepsilon \nu) \|d_t \dot{w}^{(m)}\|_{\mathcal{L}_w^2}^2 \\ &\leq \frac{\nu^2}{2\varepsilon} \frac{F_E^2}{8\sigma_E^2} \text{tr}(\Lambda^4 M^2) \|d_t v^{(m)}\|_{\mathcal{L}_v^2}^2, \end{aligned}$$

where $\Lambda_{\min} := \min\{\Lambda_{EE}, \Lambda_{EI}\}$ is the smallest eigenvalue of Λ . Next, setting $\varepsilon = \frac{2}{\nu} \Lambda_{\min}$, replacing $\dot{w} = d_t w$, and using Grönwall's inequality yields

$$(53) \quad \begin{aligned} \|d_t^2 w^{(m)}(t)\|_{\mathcal{L}_w^2}^2 + \frac{3}{2} \nu^2 \|d_t \partial_x w^{(m)}(t)\|_{\mathcal{L}_{\partial w}^2}^2 + \nu^2 \|\Lambda d_t w^{(m)}(t)\|_{\mathcal{L}_w^2}^2 \\ \leq \left(\|d_t^2 w^{(m)}\|_{\mathcal{L}_w^2}^2 + \frac{3}{2} \nu^2 \|d_t \partial_x w^{(m)}\|_{\mathcal{L}_{\partial w}^2}^2 + \nu^2 \|\Lambda d_t w^{(m)}\|_{\mathcal{L}_w^2}^2 \right) \Big|_{t=0} \\ + \frac{1}{32} \frac{\nu^3}{\Lambda_{\min} \sigma^2} F_E^2 \text{tr}(\Lambda^4 M^2) \|d_t v^{(m)}\|_{L^2(0,T;\mathcal{L}_v^2)}^2. \end{aligned}$$

Finally, it follows from (20) and (21) that, for some $\alpha_1 > 0$,

$$\|d_t^2 w^{(m)}\|_{\mathcal{L}_w^2}^2 \Big|_{t=0} \leq \alpha_1 \left(\|w'_0\|_{\mathcal{H}_w^1}^2 + \|w_0\|_{\mathcal{H}_w^2}^2 + \nu^2 |\Omega| F_E^2 \text{tr}(\Lambda^4 M^2) \right).$$

Now, using the energy estimate (22) and the above inequality in (53), it follows that

$$\|d_t^2 w^{(m)}(t)\|_{\mathcal{L}_w^2}^2 + \|d_t w^{(m)}(t)\|_{\mathcal{H}_w^1}^2 \leq \alpha_2 \left(\|w'_0\|_{\mathcal{H}_w^1}^2 + \|w_0\|_{\mathcal{H}_w^2}^2 + (|\Omega| + \kappa_v) F_E^2 \right)$$

for some $\alpha_2 > 0$ and all $t \in [0, T]$. Since this inequality and (51) hold for all $t \in [0, T]$, it follows that

$$(54) \quad \sup_{t \in [0, T]} \left(\|d_t^2 w^{(m)}(t)\|_{\mathcal{L}_w^2}^2 + \|d_t w^{(m)}(t)\|_{\mathcal{H}_w^1}^2 + \|w^{(m)}(t)\|_{\mathcal{H}_w^2}^2 \right) \leq \hat{\beta}_w,$$

where

$$\hat{\beta}_w := \alpha \left(\|w'_0\|_{\mathcal{H}_w^1}^2 + \|w_0\|_{\mathcal{H}_w^2}^2 + (|\Omega| + \kappa_v) F_E^2 \right)$$

for some $\alpha > 0$. Now, using the above estimate and passing to the limits, the result follows by similar arguments as in the proof of Theorem 4.5. ■

Proposition 4.8 (Regularity of strong solutions). *Suppose that the assumptions of Theorem 4.7 hold, namely, $g \in L^2(0, T; \mathcal{L}_i^2)$, $v_0 \in \mathcal{L}_v^2$, $i_0 \in \mathcal{L}_i^2$, $i'_0 \in \mathcal{L}_i^2$, $w_0 \in \mathcal{H}_w^2$, and $w'_0 \in \mathcal{H}_w^1$. Then, in addition to the properties of the weak solution given in Proposition 4.6, the Ω -periodic strong solution (v, i, w) of the initial value problem (4)–(7) satisfies*

$$(55) \quad \operatorname{ess\,sup}_{t \in [0, T]} \left(\|d_t^2 w(t)\|_{\mathcal{L}_w^2}^2 + \|d_t w(t)\|_{\mathcal{H}_w^1}^2 + \|w(t)\|_{\mathcal{H}_w^2}^2 \right) + \|d_t^3 w\|_{L^2(0, T; \mathcal{H}_w^{1*})}^2 \leq \beta_w,$$

$$(56) \quad w \in H^2(0, T; \mathcal{L}_w^2) \cap H^1(0, T; \mathcal{H}_w^1) \cap C^{1, \frac{1}{2}}([0, T]; \mathcal{L}_w^2) \cap C^{0, \frac{1}{2}}([0, T]; \mathcal{H}_w^1) \\ \cap C^0([0, T]; \mathcal{H}_w^2) \cap C^0([0, T]; C_{\text{per}}^{0, \lambda}(\overline{\Omega}, \mathbb{R}^2)),$$

$$d_t w \in H^1(0, T; \mathcal{L}_w^2) \cap C^{0, \frac{1}{2}}([0, T]; \mathcal{L}_w^2) \cap C^0([0, T]; \mathcal{H}_w^1),$$

$$d_t^2 w \in C^0([0, T]; \mathcal{L}_w^2)$$

for all $\lambda \in (0, 1)$ and some $\beta_w > 0$.

Proof. Differentiate (20) with respect to t and denote $\dot{w} := d_t w$. Use (52) and follow the same steps used to prove (24) in Proposition 4.4 to show $\|d_t^2 \dot{w}^{(m)}\|_{L^2(0, T; \mathcal{H}_w^{1*})}^2 \leq \tilde{\beta}_w$ for every positive integer m , all $t \in [0, T]$, and some $\tilde{\beta}_w > 0$ proportional to $\hat{\beta}_w$ in (54). Replacing $\dot{w} = d_t w$, adding the result to (54), and passing to the limits establishes (55) for some $\beta_w > 0$ proportional to $\hat{\beta}_w$.

The inclusions in H^1 and H^2 in assertion (56) follow immediately from (55). The inclusions in the Hölder spaces $C^{0, \frac{1}{2}}$ and $C^{1, \frac{1}{2}}$ are implied by the Sobolev embedding theorems [9, Thm. 6.6-1] applied to Banach space-valued functions on $[0, T] \subset \mathbb{R}$.

To show $d_t w \in C^0([0, T]; \mathcal{H}_w^1)$ and $d_t^2 w \in C^0([0, T]; \mathcal{L}_w^2)$, consider the time-independent self-adjoint linear operator $A := (-\frac{3}{2}\nu^2 \Delta + I) : \mathcal{H}_w^1 \rightarrow \mathcal{H}_w^{1*}$. Differentiate (6) with respect to t and denote $\dot{w} := d_t w$. Note that $d_t f(v) \in C^1([0, T]; \mathcal{L}_v^\infty)$, since $\partial_v f$ is a bounded smooth function and $d_t v \in C^1([0, T]; \mathcal{L}_v^2)$, given by Proposition 4.6. Then, it follows from (6) and (55) that $d_t^2 \dot{w} + A \dot{w} \in L^2(0, T; \mathcal{L}_w^2)$. Therefore, by [51, Lem. II.4.1], we have $\dot{w} \in C^0([0, T]; \mathcal{H}_w^1)$ and $d_t \dot{w} \in C^0([0, T]; \mathcal{L}_w^2)$.

Next, noting that $f(v) \in C^2([0, T]; \mathcal{L}_v^\infty)$, $w \in C^{1, \frac{1}{2}}([0, T]; \mathcal{L}_w^2)$, $d_t w \in C^{0, \frac{1}{2}}([0, T]; \mathcal{L}_w^2)$, and $d_t^2 w \in C^0([0, T]; \mathcal{L}_w^2)$, it follows from (6) that $(-\Delta + I)w \in C^0([0, T]; \mathcal{L}_w^2)$, and hence $w \in C^0([0, T]; \mathcal{H}_w^2)$. Moreover, using the Sobolev embedding theorems applied to Ω -periodic functions in $\mathbb{R}^2$, this further implies that $w \in C^0([0, T]; C_{\text{per}}^{0, \lambda}(\overline{\Omega}, \mathbb{R}^2))$. ■

Other than the regularity properties given in Propositions 4.6 and 4.8, boundedness of weak and strong solutions associated with bounded input functions g can also be established. We defer this result to section 5.1, as a corollary of Proposition 5.3.

In the remainder of the paper, as suggested in [44, sect. 11.1.2], we give formal arguments for some of the proofs, in the sense that we take the inner product of (6) with functions that belong to $\mathcal{L}_w^2$, instead of functions belonging to $\mathcal{H}_w^1$ as required for the test functions h_w in (12). However, the proofs can be made rigorous using the Galerkin approximation technique based on the dual orthogonal basis of $\mathcal{H}_w^1 \subseteq \mathcal{L}_w^2$ and then passing to the limits, as in the proofs of Theorems 4.5 and 4.7. See the discussion and results in [44, sect. 11.1.2] for further details.

5. Semidynamical systems and biophysical plausibility of the evolution. In this section, we establish a semidynamical system framework for the initial value problem presented in section 4. Assume $g \in L^2(0, \infty; \mathcal{L}_i^2)$ and let $u(t) := (v(t), i(t), d_t i(t), w(t), d_t w(t))$ denote a solution of (4)–(6) with the initial value $u_0 := u(0) = (v_0, i_0, i'_0, w_0, w'_0)$. Recall Definitions 4.2 and 4.3 and the results of Theorems 4.5 and 4.7 to note that the Hilbert spaces

$$(57) \quad \begin{aligned} \mathcal{U}_w &:= \mathcal{L}_v^2 \times \mathcal{L}_i^2 \times \mathcal{L}_i^2 \times \mathcal{H}_w^1 \times \mathcal{L}_w^2, \\ \mathcal{U}_s &:= \mathcal{L}_v^2 \times \mathcal{L}_i^2 \times \mathcal{L}_i^2 \times \mathcal{H}_w^2 \times \mathcal{H}_w^1 \end{aligned}$$

construct, respectively, the phase spaces associated with the weak and strong solutions. Now, for every $t \in [0, \infty)$, define the mappings

$$\begin{aligned} S_w(t) : \mathcal{U}_w &\rightarrow \mathcal{U}_w, & S_w(t)u_0 &:= u(t), \\ S_s(t) : \mathcal{U}_s &\rightarrow \mathcal{U}_s, & S_s(t)u_0 &:= u(t). \end{aligned}$$

The existence and uniqueness of the solutions given by Theorems 4.5 and 4.7 along with the time-continuity of the solutions given by Propositions 4.6 and 4.8 imply that the above mappings are well defined for all $t \in [0, \infty)$. Then, $\{S_w(t)\}_{t \in [0, \infty)}$ and $\{S_s(t)\}_{t \in [0, \infty)}$ form semigroups of operators which give the weak and strong solutions of (2), respectively. The following propositions show that these semigroups are continuous, which also ensures that the initial value problems of finding weak and strong solutions for (2) are well-posed.

Proposition 5.1 (Continuity of the semigroup $\{S_w\}$). *The semigroup $\{S_w(t)\}_{t \in [0, \infty)}$ of weak solution operators is continuous for all $g \in L^2(0, \infty; \mathcal{L}_i^2)$.*

Proof. Continuity of the semigroup with respect to t follows immediately from the continuity of the weak solutions given in Proposition 4.6. It remains to prove continuous dependence of the solution on the initial values. Let $\tilde{u}_0$ and $\hat{u}_0$ be any two initial values in $\mathcal{U}_w$ that give the solutions $\tilde{u}(t) = S_w(t)\tilde{u}_0$ and $\hat{u}(t) = S_w(t)\hat{u}_0$ for all $t \in [0, T]$, $T > 0$. Let $u(t) := \tilde{u}(t) - \hat{u}(t)$ be the weak solution with the initial value $u_0 := \tilde{u}_0 - \hat{u}_0$. Now, consider (4)–(6) satisfied by $\tilde{u}$ and $\hat{u}$, and take the inner product of (4)–(6) in each set with v , $d_t i$, and $d_t w$, respectively. Subtracting the resulting two sets of equations yields

$$(58) \quad (\Phi d_t v, v)_{\mathcal{L}_v^2} + (vv)_{\mathcal{L}_v^2} - (J_1 i, v)_{\mathcal{L}_v^2} + \left(J_2(\tilde{v} \tilde{i}^T - \hat{v} \hat{i}^T) \Psi J_4 + J_3(\tilde{v} \tilde{i}^T - \hat{v} \hat{i}^T) \Psi J_5, v \right)_{\mathcal{L}_v^2} = 0,$$

$$(59) \quad (d_t^2 i, d_t i)_{\mathcal{L}_i^2} + 2(\Gamma d_t i, d_t i)_{\mathcal{L}_i^2} + (\Gamma^2 i, d_t i)_{\mathcal{L}_i^2} - e(\Upsilon \Gamma J_6 w, d_t i)_{\mathcal{L}_i^2} - e(\Upsilon \Gamma N J_7(f(\tilde{v}) - f(\hat{v})), d_t i)_{\mathcal{L}_i^2} = 0,$$

$$(60) \quad (d_t^2 w, d_t w)_{\mathcal{L}_w^2} + 2\nu(\Lambda d_t w, d_t w)_{\mathcal{L}_w^2} + \frac{3}{2}\nu^2(\partial_x w, d_t \partial_x w)_{\mathcal{L}_{\partial w}^2} + \nu^2(\Lambda^2 w, d_t w)_{\mathcal{L}_w^2} - \nu^2(\Lambda^2 M J_8(f(\tilde{v}) - f(\hat{v})), d_t w)_{\mathcal{L}_w^2} = 0.$$

As in the proof of uniqueness given in Theorem 4.5,

$$(61) \quad \begin{aligned} -(J_2(\tilde{v} \tilde{i}^T - \hat{v} \hat{i}^T) \Psi J_4, v)_{\mathcal{L}_v^2} &\leq \sqrt{2\kappa_i} \|\Psi\|_2 \|v\|_{\mathcal{L}_v^2}^2 + \|v\|_{\mathcal{L}_v^2}^2 + \frac{1}{2}\kappa_{\tilde{v}} \|\Psi\|_2^2 \|\tilde{i}\|_{\mathcal{L}_i^2}^2, \\ -(J_3(\tilde{v} \tilde{i}^T - \hat{v} \hat{i}^T) \Psi J_5, v)_{\mathcal{L}_v^2} &\leq \sqrt{2\kappa_i} \|\Psi\|_2 \|v\|_{\mathcal{L}_v^2}^2 + \|v\|_{\mathcal{L}_v^2}^2 + \frac{1}{2}\kappa_{\tilde{v}} \|\Psi\|_2^2 \|\tilde{i}\|_{\mathcal{L}_i^2}^2, \end{aligned}$$

$$\begin{aligned}
e(\Upsilon \Gamma N J_7(f(\tilde{v}) - f(\hat{v})), d_t i)_{\mathcal{L}_i^2} &\leq \|d_t i\|_{\mathcal{L}_i^2}^2 + \frac{1}{32} e^2 \|\Upsilon \Gamma N J_7\|_2^2 \max \left\{ \frac{F_E^2}{\sigma_E^2}, \frac{F_I^2}{\sigma_I^2} \right\} \|v\|_{\mathcal{L}_v^2}^2, \\
\nu^2 (\Lambda^2 M J_8(f(\tilde{v}) - f(\hat{v})), d_t w)_{\mathcal{L}_w^2} &\leq \nu^2 \|d_t w\|_{\mathcal{L}_w^2}^2 + \frac{1}{32} \nu^2 \frac{F_E^2}{\sigma_E^2} \operatorname{tr}(\Lambda^4 M^2) \|v\|_{\mathcal{L}_v^2}^2, \\
(J_1 i, v)_{\mathcal{L}_v^2} &\leq \|v\|_{\mathcal{L}_v^2}^2 + \frac{1}{2} \|i\|_{\mathcal{L}_i^2}^2, \\
e(\Upsilon \Gamma J_6 w, d_t i)_{\mathcal{L}_i^2} &\leq \|d_t i\|_{\mathcal{L}_i^2}^2 + \frac{1}{4} e^2 \|\Upsilon \Gamma J_6\|_2^2 \|w\|_{\mathcal{L}_w^2}^2,
\end{aligned}$$

where $\kappa_{\hat{v}}$ and $\kappa_{\tilde{v}}$ are in the form of (25) and (26). Now, substituting the above inequalities into (58)–(60), adding the resulting inequalities together, and using Grönwall's inequality yield, for some $\alpha, \beta > 0$,

$$(62) \quad \|u(t)\|_{\mathcal{U}_w}^2 \leq \beta e^{\alpha T} \|u_0\|_{\mathcal{U}_w}^2 \quad \text{for all } t \in [0, T],$$

which completes the proof. ■

Proposition 5.2 (Continuity of the semigroup $\{S_s\}$). *The semigroup $\{S_s(t)\}_{t \in [0, \infty)}$ of strong solution operators is continuous for all $g \in L^2(0, \infty; \mathcal{L}_i^2)$.*

Proof. Continuity of the semigroup with respect to t follows immediately from the time continuity of the strong solutions given by Proposition 4.8. To prove continuous dependence on the initial values, consider any two initial values $\tilde{u}_0$ and $\hat{u}_0$ in $\mathcal{U}_s$ and construct the solutions $\tilde{u}(t) = S_s(t)\tilde{u}_0$ and $\hat{u}(t) = S_s(t)\hat{u}_0$, $t \in [0, T]$, $T > 0$, for (4)–(6). Let $u := \tilde{u} - \hat{u}$ and $A := -\Delta + I$, and take the inner product of (4)–(6) for each solution with v , $d_t i$, and $Ad_t w$, respectively. Subtracting the resulting two sets of equations gives (58), (59), and

$$\begin{aligned}
(63) \quad &\frac{1}{2} d_t \|d_t w\|_{\mathcal{H}_w^1}^2 + 2\nu \|\Lambda^{\frac{1}{2}} d_t w\|_{\mathcal{H}_w^1}^2 + \frac{3}{4} \nu^2 d_t \|\partial_x w\|_{\mathcal{H}_{\partial w}^1}^2 + \frac{1}{2} \nu^2 d_t \|\Lambda w\|_{\mathcal{H}_w^1}^2 \\
&= \nu^2 (\Lambda^2 M J_8(f(\tilde{v}) - f(\hat{v})), Ad_t w)_{\mathcal{L}_w^2}.
\end{aligned}$$

Note that (62) also holds since $\mathcal{U}_s \subset \mathcal{U}_w$, and since (58) and (59) remain unchanged, the continuity of v and i holds.

Now, it follows from (63) by integrating over $[0, t]$ that

$$\begin{aligned}
\|d_t w\|_{\mathcal{H}_w^1}^2 + \nu^2 \left[\frac{3}{2} \|\partial_x w\|_{\mathcal{H}_{\partial w}^1}^2 + \|\Lambda w\|_{\mathcal{H}_w^1}^2 \right] &\leq \left(\|d_t w\|_{\mathcal{H}_w^1}^2 + \nu^2 \left[\frac{3}{2} \|\partial_x w\|_{\mathcal{H}_{\partial w}^1}^2 + \|\Lambda w\|_{\mathcal{H}_w^1}^2 \right] \right) \Big|_{t=0} \\
&\quad + 2\nu^2 \int_0^t (\Lambda^2 M J_8(f(\tilde{v}) - f(\hat{v})), Ad_s w)_{\mathcal{L}_w^2} ds,
\end{aligned}$$

which, using (1), can be written equivalently for some $\alpha_1, \beta_1 > 0$ as

$$(64) \quad Q(w(t), d_t w(t)) \leq \alpha_1 Q(w(0), d_t w(0)) + \beta_1 \int_0^t (\Lambda^2 M J_8(f(\tilde{v}) - f(\hat{v})), Ad_s w)_{\mathcal{L}_w^2} ds,$$

where

$$(65) \quad Q(w(t), d_t w(t)) := \|d_t w(t)\|_{\mathcal{H}_w^1}^2 + \|Aw(t)\|_{\mathcal{L}_w^2}^2.$$

Integrating by parts in the second term of the right-hand side of (64) yields

$$(66) \quad \beta_1 \int_0^t (\Lambda^2 M J_8(f(\tilde{v}) - f(\hat{v})), Ad_s w)_{\mathcal{L}_w^2} ds = \beta_1 (\Lambda^2 M J_8(f(\tilde{v}) - f(\hat{v})), Aw)_{\mathcal{L}_w^2} \\ - \beta_1 (\Lambda^2 M J_8(f(\tilde{v}_0) - f(\hat{v}_0)), Aw_0)_{\mathcal{L}_w^2} \\ - \beta_1 \int_0^t (\Lambda^2 M J_8 d_s(f(\tilde{v}) - f(\hat{v})), Aw)_{\mathcal{L}_w^2} ds.$$

Next, recalling that $\sup_{v_X(x,t) \in \mathbb{R}} |\partial_{v_X} f_X(v_X)| \leq \frac{F_X}{2\sqrt{2}\sigma_X}$ by (45) and using Young's inequality, we obtain

$$(67) \quad \beta_1 (\Lambda^2 M J_8(f(\tilde{v}) - f(\hat{v})), Aw)_{\mathcal{L}_w^2} \leq \frac{1}{2} \|Aw\|_{\mathcal{L}_w^2}^2 + \frac{\beta_1^2 F_E^2}{16 \sigma_E^2} \text{tr}(\Lambda^4 M^2) \|v\|_{\mathcal{L}_v^2}^2, \\ -\beta_1 (\Lambda^2 M J_8(f(\tilde{v}_0) - f(\hat{v}_0)), Aw_0)_{\mathcal{L}_w^2} \leq \frac{1}{2} \|Aw_0\|_{\mathcal{L}_w^2}^2 + \frac{\beta_1^2 F_E^2}{16 \sigma_E^2} \text{tr}(\Lambda^4 M^2) \|v_0\|_{\mathcal{L}_v^2}^2.$$

Moreover,

$$-\beta_1 (\Lambda^2 M J_8 d_s(f(\tilde{v}) - f(\hat{v})), Aw)_{\mathcal{L}_w^2} \\ = -\beta_1 (\Lambda^2 M J_8(\partial_{\tilde{v}} f(\tilde{v}) d_s \tilde{v} - \partial_{\hat{v}} f(\hat{v}) d_s \hat{v}), Aw)_{\mathcal{L}_w^2} \\ \leq \frac{1}{2} \|Aw\|_{\mathcal{L}_w^2}^2 + \frac{1}{2} \beta_1^2 \|\Lambda^2 M J_8(\partial_{\tilde{v}} f(\tilde{v}) d_s \tilde{v} - \partial_{\hat{v}} f(\hat{v}) d_s \hat{v})\|_{\mathcal{L}_w^2}^2 \\ = \frac{1}{2} \|Aw\|_{\mathcal{L}_w^2}^2 + \frac{1}{2} \beta_1^2 \text{tr}(\Lambda^4 M^2) \int_{\Omega} |\partial_{\tilde{v}_E} f(\tilde{v}_E) d_s \tilde{v}_E - \partial_{\hat{v}_E} f(\hat{v}_E) d_s \hat{v}_E|^2 dx,$$

where, noting that $\sup_{v_E(x,t) \in \mathbb{R}} |\partial_{v_E}^2 f_E(v_E)| < \frac{1}{5} \frac{F_E}{\sigma_E^2}$ by direct computation of the derivative of (45), we can write

$$|\partial_{\tilde{v}_E} f(\tilde{v}_E) d_s \tilde{v}_E - \partial_{\hat{v}_E} f(\hat{v}_E) d_s \hat{v}_E|^2 dx = |\partial_{\tilde{v}_E} f(\tilde{v}_E) d_s v_E + (\partial_{\tilde{v}_E} f(\tilde{v}_E) - \partial_{\hat{v}_E} f(\hat{v}_E)) d_s \hat{v}_E|^2 \\ \leq 2 |\partial_{\tilde{v}_E} f(\tilde{v}_E)|^2 |d_s v_E|^2 + 2 |\partial_{\tilde{v}_E} f(\tilde{v}_E) - \partial_{\hat{v}_E} f(\hat{v}_E)|^2 |d_s \hat{v}_E|^2 \\ \leq \frac{1}{4} \frac{F_E^2}{\sigma_E^2} |d_s v_E|^2 + 2 \left[\sup_{v_E(x,t) \in \mathbb{R}} |\partial_{v_E}^2 f_E(v_E)| \right]^2 |v_E|^2 |d_s \hat{v}_E|^2 \\ \leq \frac{1}{4} \frac{F_E^2}{\sigma_E^2} |d_s v_E|^2 + \frac{2}{25} \frac{F_E^2}{\sigma_E^4} |v_E|^2 |d_s \hat{v}_E|^2.$$

Therefore, it follows that

$$(68) \quad -\beta_1 (\Lambda^2 M J_8 d_s(f(\tilde{v}) - f(\hat{v})), Aw)_{\mathcal{L}_w^2} \leq \frac{1}{2} \|Aw\|_{\mathcal{L}_w^2}^2 + \frac{\beta_1^2 F_E^2}{8 \sigma_E^2} \text{tr}(\Lambda^4 M^2) \|d_s v\|_{\mathcal{L}_v^2}^2 \\ + \frac{\beta_1^2 F_E^2}{25 \sigma_E^4} \text{tr}(\Lambda^4 M^2) \|d_s \hat{v}\|_{C^1([0,T]; \mathcal{L}_v^2)}^2 \|v\|_{\mathcal{L}_v^2}^2.$$

Moreover, (4) implies that, for some $\alpha_2 > 0$,

$$(69) \quad \|d_s v(s)\|_{\mathcal{L}_v^2}^2 \leq \alpha_2 \left(\|v(s)\|_{\mathcal{L}_v^2}^2 + \|i(s)\|_{\mathcal{L}_i^2}^2 + \|v(s)\|_{\mathcal{L}_v^2}^2 \|i(s)\|_{\mathcal{L}_i^2}^2 \right) \quad \text{for all } s \in [0, T].$$

Now, substituting (67), (68), and (69) into (66) and using (62), it follows that there exist some $\beta_2, \dots, \beta_6 > 0$ such that

$$\begin{aligned} & \beta_1 \int_0^t (\Lambda^2 M J_8(f(\tilde{v}) - f(\hat{v})), d_s A w)_{\mathcal{L}_w^2} ds \\ & \leq \frac{1}{2} \int_0^t \|A w\|_{\mathcal{L}_w^2}^2 ds + \beta_2 \int_0^t \left(\|v\|_{\mathcal{L}_v^2}^2 + \|i\|_{\mathcal{L}_i^2}^2 + \|v\|_{\mathcal{L}_v^2}^2 \|i\|_{\mathcal{L}_i^2}^2 \right) ds \\ & \quad + \frac{1}{2} \|A w\|_{\mathcal{L}_w^2}^2 + \beta_3 \|v\|_{\mathcal{L}_v^2}^2 + \frac{1}{2} \|A w_0\|_{\mathcal{L}_w^2}^2 + \beta_4 \|v_0\|_{\mathcal{L}_v^2}^2 \\ & \leq \frac{1}{2} \int_0^t \|A w\|_{\mathcal{L}_w^2}^2 ds + \beta_5 \|u_0\|_{\mathcal{U}_w}^2 (1 + \|u_0\|_{\mathcal{U}_w}^2) t + \frac{1}{2} \|A w\|_{\mathcal{L}_w^2}^2 \\ & \quad + \frac{1}{2} \|A w_0\|_{\mathcal{L}_w^2}^2 + \beta_6 \|u_0\|_{\mathcal{U}_w}^2. \end{aligned}$$

Substituting this inequality into (64) yields

$$(70) \quad \begin{aligned} \frac{1}{2} Q(w(t), d_t w(t)) & \leq \frac{1}{2} \int_0^t Q(w(s), d_s w(s)) ds + \beta_5 \|u_0\|_{\mathcal{U}_w}^2 (1 + \|u_0\|_{\mathcal{U}_w}^2) t \\ & \quad + \alpha_1 Q(w(0), d_t w(0)) + \frac{1}{2} \|A w_0\|_{\mathcal{L}_w^2}^2 + \beta_6 \|u_0\|_{\mathcal{U}_w}^2, \end{aligned}$$

where, using Grönwall's inequality for the function $\frac{1}{2} \int_0^t Q(w(s), d_s w(s)) ds$, we can write

$$\begin{aligned} \frac{1}{2} \int_0^t Q(w(s), d_s w(s)) ds & \leq \beta_5 \|u_0\|_{\mathcal{U}_w}^2 (1 + \|u_0\|_{\mathcal{U}_w}^2) (e^t - (t+1)) \\ & \quad + \left[\alpha_1 Q(w(0), d_t w(0)) + \frac{1}{2} \|A w_0\|_{\mathcal{L}_w^2}^2 + \beta_6 \|u_0\|_{\mathcal{U}_w}^2 \right] (e^t - 1). \end{aligned}$$

This inequality, along with (70) and the definition of Q given by (65), implies that, for some $\beta_7 > 0$,

$$Q(w(t), d_t w(t)) \leq \beta_7 e^T [Q(w(0), d_t w(0)) + \|u_0\|_{\mathcal{U}_w}^2 (1 + \|u_0\|_{\mathcal{U}_w}^2)] \quad \text{for all } t \in [0, T].$$

Now, noting that $Q(w(0), d_t w(0)) = \|w'_0\|_{\mathcal{H}_w^1}^2 + \|A w_0\|_{\mathcal{L}_w^2}^2$, it follows from the above inequality and (62) that, for some $\hat{\alpha}, \hat{\beta} > 0$,

$$\|u(t)\|_{\mathcal{U}_s}^2 \leq \hat{\beta} e^{\hat{\alpha} T} \|u_0\|_{\mathcal{U}_s}^2 (1 + \|u_0\|_{\mathcal{U}_w}^2) \quad \text{for all } t \in [0, T],$$

which completes the proof. ■

5.1. Biophysically plausible phase spaces. Although the spaces $\mathcal{U}_w$ and $\mathcal{U}_s$ constructed in (57) provide the theoretical phase spaces of the problem for the solutions constructed in section 4, the evolution of the dynamics of the model is not biophysically plausible on the entire spaces $\mathcal{U}_w$ and $\mathcal{U}_s$. As described in section 3, $i(x, t)$, $w(x, t)$, and $g(x, t)$ represent nonnegative biophysical quantities. In fact, initial functions $i'_0 \in \mathcal{L}_i^2$ and $w'_0 \in \mathcal{L}_w^2$ can be constructed such that the solutions $i(x, t)$ and $w(x, t)$, despite starting from nonnegative initial values $i_0 \in \mathcal{L}_i^2$ and $w_0 \in \mathcal{H}_w^1$, take negative values over a set $\mathcal{X} \subset \Omega$ of positive measure for a time interval of positive length. In the following propositions, we establish conditions under which the dynamics of the model is guaranteed to evolve in biophysically plausible subsets of $\mathcal{U}_w$ and $\mathcal{U}_s$.

Proposition 5.3 (Nonnegativity of the solution $w(x, t)$). *Suppose that $w \in L^2(0, T; \mathcal{H}_w^1)$ is the w -component of an Ω -periodic weak solution $u(t) = S_w(t)u_0$ of (4)–(7) and define the set $\mathcal{D}_w \subset \mathcal{H}_w^1 \times \mathcal{L}_w^2$ as*

$$(71) \quad \mathcal{D}_w := \{(w_0, w'_0) \in \mathcal{W}_w^{1,\infty} \times \mathcal{L}_w^\infty : w'_0 + \nu \Lambda w_0 \geq 0 \text{ a.e. in } \Omega, \\ \text{and } w_0(y) + \partial_y w_0(y)(y - x) \geq 0 \text{ for almost every } x \in \Omega, y \in B(x, t), t \in (0, T]\}.$$

Then, for every initial value $(w_0, w'_0) \in \mathcal{D}_w$, the solution $w(x, t)$ remains nonnegative almost everywhere in Ω for all $t \in (0, T]$.

Proof. First, note that the weak and strong solutions coincide for $v(t)$ and they satisfy (4) and (5) almost everywhere in Ω for all $t \in [0, T]$, $T > 0$; see the proof of Theorem 4.7. Substituting $v(t)$ into f , we can interpret $f(v)$ in (6) as a function $\hat{f}(x, t) := f(v(x, t))$ for almost every $x \in \Omega$ and all $t \in [0, T]$. Next, using (3), (8), and Proposition 4.6, it is implied that $\hat{f} \in L^\infty(0, T; \mathcal{L}_v^\infty)$ and $\hat{f} > 0$ in $\Omega \times [0, T]$. Now, replace $f(v)$ in (6) by $\hat{f}$ and scale x by the factor $\sqrt{\frac{3}{2}}\nu$ to obtain

$$\begin{aligned} \partial_t^2 \tilde{w} + 2\nu \Lambda \partial_t \tilde{w} - \Delta \tilde{w} + \nu^2 \Lambda^2 \tilde{w} - \tilde{f} &= 0 & \text{in } \tilde{\Omega} \times (0, T], \\ \tilde{w} = \tilde{w}_0, \quad \partial_t \tilde{w} = \tilde{w}'_0 & & \text{on } \tilde{\Omega} \times \{0\}, \end{aligned}$$

where $\tilde{\Omega} := \sqrt{\frac{3}{2}}\nu\Omega$, and $\tilde{w}$, $\tilde{w}_0$, $\tilde{w}'_0$, and $\tilde{f}$ denote w , w_0 , w'_0 , and $\nu^2 \Lambda^2 M J_8 \hat{f}$ in the scaled domain $\tilde{\Omega}$, respectively. Note that, with the new interpretation of f , the above equation is a system of two decoupled telegraph equations. Therefore, applying the same arguments to each of the two equations independently, in what follows we assume without loss of generality that the above equation is a scalar equation.

Using the change of variable $q := e^{\nu \Lambda t} \tilde{w}$, the problem can be transformed to the initial value problem of the standard wave equation given by

$$(72) \quad \begin{aligned} \partial_t^2 q - \Delta q &= e^{\nu \Lambda t} \tilde{f} & \text{in } \mathbb{R}^2 \times (0, T], \\ q = \tilde{w}_0, \quad \partial_t q &= \tilde{w}'_0 + \nu \Lambda \tilde{w}_0 & \text{on } \mathbb{R}^2 \times \{0\}. \end{aligned}$$

Here, the extension from $\tilde{\Omega}$ to $\mathbb{R}^2$ is done periodically due to the $\tilde{\Omega}$ -periodicity of the functions. Let $\tilde{w}_{0\varepsilon}$, $\tilde{w}'_{0\varepsilon}$, and $\tilde{f}_\varepsilon$ denote, respectively, $\tilde{w}_0$, $\tilde{w}'_0$, and $\tilde{f}$ after mollification by the standard positive mollifier $\phi_\varepsilon \in C_c^\infty(\mathbb{R}^2)$; see [9, sect. 2.6]. Using Poisson's formula for the homogeneous wave equation in $\mathbb{R}^2$, along with Duhamel's principle for the nonhomogeneous problem [16, sect. 2.4], it follows that the function

$$\begin{aligned}
(73) \quad q_\varepsilon(x, t) &:= \frac{1}{2} \int_{B(x, t)} \frac{t[\tilde{w}_{0\varepsilon}(y) + (\partial_y \tilde{w}_{0\varepsilon}(y), y - x)_{\mathbb{R}^2}] + t^2[\tilde{w}'_{0\varepsilon}(y) + \nu \Lambda \tilde{w}_{0\varepsilon}(y)]}{\left[t^2 - \|y - x\|_{\mathbb{R}^2}^2\right]^{\frac{1}{2}}} dy \\
&\quad + \frac{1}{2} \int_0^t (t-s)^2 e^{\nu \Lambda s} \int_{B(x, t-s)} \frac{\tilde{f}_\varepsilon(y, s)}{\left[(t-s)^2 - \|y - x\|_{\mathbb{R}^2}^2\right]^{\frac{1}{2}}} dy ds
\end{aligned}$$

solves (72) classically for the forcing term $e^{\nu \Lambda t} \tilde{f}_\varepsilon$ and initial values $\tilde{w}_{0\varepsilon}$ and $\tilde{w}'_{0\varepsilon}$.

The second term in this solution is nonnegative for all $t \in [0, T]$, since $\tilde{f}$ and consequently, $\tilde{f}_\varepsilon$ are nonnegative on $B(x, t)$ for all $x \in \Omega$ and all $t \in [0, T]$. Moreover, by [9, Thm. 2.6-1] and the definition of the weak derivative, we can write

$$\begin{aligned}
(\partial_y \tilde{w}_{0\varepsilon}(y), y - x)_{\mathbb{R}^2} &= \left(\int_{B(y, \varepsilon)} \partial_y \phi_\varepsilon(y - z) \tilde{w}_0(z) dz, y - x \right)_{\mathbb{R}^2} \\
&= \left(- \int_{B(y, \varepsilon)} \partial_z \phi_\varepsilon(y - z) \tilde{w}_0(z) dz, y - x \right)_{\mathbb{R}^2} \\
&= \left(\int_{B(y, \varepsilon)} \phi_\varepsilon(y - z) \partial_z \tilde{w}_0(z) dz, y - x \right)_{\mathbb{R}^2} \\
&= \int_{B(y, \varepsilon)} \phi_\varepsilon(y - z) (\partial_z \tilde{w}_0(z), z - x)_{\mathbb{R}^2} dz \\
&\quad + \int_{B(y, \varepsilon)} \phi_\varepsilon(y - z) (\partial_z \tilde{w}_0(z), y - z)_{\mathbb{R}^2} dz,
\end{aligned}$$

where, using Hölder's inequality and the property $\int_{B(0, \varepsilon)} \phi_\varepsilon(x) dx = 1$, we have

$$\begin{aligned}
\left| \int_{B(y, \varepsilon)} \phi_\varepsilon(y - z) (\partial_z \tilde{w}_0(z), y - z)_{\mathbb{R}^2} dz \right| &\leq \|\partial_x \tilde{w}_0\|_{\mathcal{L}_{\partial w}^\infty} \int_{B(y, \varepsilon)} \phi_\varepsilon(y - z) \|y - z\|_1 dz \\
&\leq \sqrt{2} \|\partial_x \tilde{w}_0\|_{\mathcal{L}_{\partial w}^\infty} \varepsilon.
\end{aligned}$$

Therefore, it follows that

$$\begin{aligned}
&\int_{B(x, t)} \frac{t[\tilde{w}_{0\varepsilon}(y) + (\partial_y \tilde{w}_{0\varepsilon}(y), y - x)_{\mathbb{R}^2}]}{\left[t^2 - \|y - x\|_{\mathbb{R}^2}^2\right]^{\frac{1}{2}}} dy \\
&\geq \int_{B(x, t)} t \left[\frac{\int_{B(y, \varepsilon)} \phi_\varepsilon(y - z) [\tilde{w}_0(z) + (\partial_z \tilde{w}_0(z), z - x)_{\mathbb{R}^2}] dz}{\left[t^2 - \|y - x\|_{\mathbb{R}^2}^2\right]^{\frac{1}{2}}} - \frac{\sqrt{2} \|\partial_x w_0\|_{\mathcal{L}_{\partial w}^\infty} \varepsilon}{\left[t^2 - \|y - x\|_{\mathbb{R}^2}^2\right]^{\frac{1}{2}}} \right] dy \\
&\geq -\sqrt{2} \|\partial_x \tilde{w}_0\|_{\mathcal{L}_{\partial w}^\infty} \varepsilon \quad \text{for all } (\tilde{w}_0, \tilde{w}'_0) \in \tilde{\mathcal{D}}_w,
\end{aligned}$$

where $\tilde{\mathcal{D}}_w$ denotes $\mathcal{D}_w$ in the scaled domain $\tilde{\Omega}$. Note that the last inequality holds since the first term in the integration on the right-hand side is nonnegative by (71), and $t[t^2 - \|y - x\|_{\mathbb{R}^2}^2]^{-\frac{1}{2}}$

takes the average value 1 over the ball $B(x, t)$. Finally, note that $\tilde{w}'_{0\varepsilon}(y) + \nu\Lambda\tilde{w}_{0\varepsilon}(y)$ in (73) is nonnegative on $B(x, t)$ when $(\tilde{w}_0, \tilde{w}'_0) \in \tilde{\mathcal{D}}_w$. Therefore, it follows that

$$(74) \quad q_\varepsilon(x, t) \geq -\sqrt{2}\|\partial_x \tilde{w}_0\|_{\mathcal{L}^\infty_{\partial w}} \varepsilon \quad \text{for all } (\tilde{w}_0, \tilde{w}'_0) \in \tilde{\mathcal{D}}_w.$$

Now, taking the limits as $\varepsilon \rightarrow 0$, it follows from [9, Thm. 2.6-3] that $\tilde{w}_{0\varepsilon} \rightarrow \tilde{w}_0$, $\tilde{w}'_{0\varepsilon} \rightarrow \tilde{w}'_0$, and $\tilde{f}_\varepsilon \rightarrow \tilde{f}$ in $L^2(\tilde{\Omega}_t)$, where $\tilde{\Omega}_t := \{y \in \mathbb{R}^2 : y \in B(x, t), x \in \Omega\}$. Therefore, there exists a subsequence $\{\varepsilon_n\}_{n=1}^\infty$, convergent to 0, such that $\tilde{w}_{0\varepsilon_n} \rightarrow \tilde{w}_0$, $\tilde{w}'_{0\varepsilon_n} \rightarrow \tilde{w}'_0$, and $\tilde{f}_{\varepsilon_n} \rightarrow \tilde{f}$ almost everywhere on Ω_t as $n \rightarrow \infty$ [18, Thm. 2.30]. Moreover, since $(\tilde{w}_0, \tilde{w}'_0) \in \mathcal{W}_w^{1,\infty} \times \mathcal{L}_w^\infty$ in $\tilde{\mathcal{D}}_w$, $\tilde{f} \in L^\infty(0, T; \mathcal{L}_v^\infty)$, and the function $[t^2 - \|y - x\|_{\mathbb{R}^2}^2]^{-\frac{1}{2}}$ is integrable over $B(x, t)$, it follows that the integrands in (73) are uniformly bounded with respect to ε by integrable functions over $B(x, t)$. Lebesgue's dominated convergence theorem then implies that $q(x, t) := \lim_{n \rightarrow \infty} q_{\varepsilon_n}(x, t)$ exists on $\tilde{\Omega}_t$ and, by uniqueness of the weak solution, is a weak solution of the wave equation (72). Now, letting $\varepsilon = \varepsilon_n \rightarrow 0$ in (74), it follows that if $(\tilde{w}_0, \tilde{w}'_0) \in \tilde{\mathcal{D}}_w$, then $q(x, t) \geq 0$ for almost every $x \in \tilde{\Omega}$ and all $t \in (0, T]$. This completes the proof, since the change of variable $\tilde{w} = e^{-\nu\Lambda t}q$ and space rescaling $\Omega = \sqrt{\frac{2}{3}}\nu^{-1}\tilde{\Omega}$ do not change the sign of solutions. $\blacksquare$

Corollary 5.4 (Boundedness of the weak solutions). *Suppose $g \in L^\infty(0, T; \mathcal{L}_i^\infty)$, $v_0 \in \mathcal{L}_v^\infty$, $i_0 \in \mathcal{L}_i^\infty$, $i'_0 \in \mathcal{L}_i^\infty$, $w_0 \in \mathcal{W}_w^{1,\infty}$, and $w'_0 \in \mathcal{L}_w^\infty$. Then, in addition to the regularities given by Proposition 4.6, the weak solution $(v(t), i(t), w(t))$ of (4)–(7) satisfies*

$$\begin{aligned} v &\in W^{2,\infty}(0, T; \mathcal{L}_v^\infty) \cap C^{1,1}([0, T]; \mathcal{L}_v^\infty), \\ i &\in W^{1,\infty}(0, T; \mathcal{L}_i^\infty) \cap C^{0,1}([0, T]; \mathcal{L}_i^\infty), \\ w &\in L^\infty(0, T; \mathcal{L}_w^\infty). \end{aligned}$$

Proof. The boundedness of w follows immediately from the proof of Proposition 5.3, since under the assumption $w_0 \in \mathcal{W}_w^{1,\infty}$ and $w'_0 \in \mathcal{L}_w^\infty$ the integrands in (73) are integrable and each component of the weak solution $w(t)$ is achieved almost everywhere in Ω as the limit of (73) when $\varepsilon \rightarrow 0$, followed by the space rescaling from $\tilde{\Omega}$ to Ω .

Now, to prove boundedness of v , i , and $d_t i$, let $x_0 \in \Omega$ be any Lebesgue point¹ of the initial functions v_0 , i_0 , i'_0 , w_0 , and $g(0)$. Take the $\mathbb{R}^4$ -inner product of (5) at x_0 with $d_t i(x_0, t)$ for every $t \in (0, T]$ to obtain

$$\begin{aligned} (d_t^2 i_{x_0}, d_t i_{x_0})_{\mathbb{R}^4} + 2(\Gamma d_t i_{x_0}, d_t i_{x_0})_{\mathbb{R}^4} + (\Gamma^2 i_{x_0}, d_t i_{x_0})_{\mathbb{R}^4} \\ - e(\Upsilon \Gamma J_6 w_{x_0}, d_t i_{x_0})_{\mathbb{R}^4} - e(\Upsilon \Gamma N J_7 f(v_{x_0}), d_t i_{x_0})_{\mathbb{R}^4} = e(\Upsilon \Gamma g_{x_0}, d_t i_{x_0})_{\mathbb{R}^4}, \end{aligned}$$

where $v_{x_0}(t) := v(x_0, t)$, $i_{x_0}(t) := i(x_0, t)$, $w_{x_0}(t) := w(x_0, t)$, and $g_{x_0}(t) := g(x_0, t)$. This equality is similar to (29) in the proof of Proposition 4.4, with the $\mathcal{L}_i^2$ -inner product being replaced by the $\mathbb{R}^4$ -inner product, and $v^{(m)}$, $i^{(m)}$, and $w^{(m)}$ being replaced by v_{x_0} , i_{x_0} , and w_{x_0} , respectively. Therefore, similar arguments as in the proof of Proposition 4.4 imply that

$$(75) \quad \sup_{t \in [0, T]} (\|d_t i_{x_0}(t)\|_{\mathbb{R}^4}^2 + \|i_{x_0}(t)\|_{\mathbb{R}^4}^2) \leq \kappa_i,$$

¹The choice of a Lebesgue point is for the sake of definiteness. Almost every point in Ω can be used as x_0 .

where, with $\kappa_w := \|w\|_{L^\infty(0,T;\mathcal{L}_w^\infty)}^2$ and for some $\alpha_1 > 0$ independent of x_0 ,

$$\kappa_i = \alpha_1 \left(\|i_0'\|_{\mathcal{L}_i^\infty}^2 + \|i_0\|_{\mathcal{L}_i^\infty}^2 + \left[\frac{e^2 \kappa_w}{\gamma_{\min}} \|\Upsilon \Gamma J_6\|_2^2 + \frac{e^2 |\Omega|}{\gamma_{\min}} (F_E^2 + F_I^2) \|\Upsilon \Gamma N J_7\|_2^2 \right] T + \frac{e^2}{2\gamma_{\min}} \|\Upsilon \Gamma\|_2^2 \|g\|_{L^\infty(0,T;\mathcal{L}_i^\infty)}^2 \right),$$

and $\gamma_{\min}$ is the smallest eigenvalue of Γ .

Similarly, taking the $\mathbb{R}^2$ -inner product of (4) at x_0 with $v_{x_0}(t)$ and using the arguments following (31) in the proof of Proposition 4.4 yields

$$(76) \quad \sup_{t \in [0,T]} \left(\|v_{x_0}(t)\|_{\mathcal{L}_v^2}^2 \right) \leq \kappa_v,$$

where, for some $\alpha_2, \beta > 0$ independent of x_0 ,

$$\kappa_v = \alpha_2 \exp(\beta \sqrt{2\kappa_i} \|\Psi\|_2 T) \left(\|v_0\|_{\mathcal{L}_v^\infty}^2 + \frac{\kappa_i}{\sqrt{2\kappa_i} \|\Psi\|_2} \right).$$

Now, note that almost every point $x_0 \in \Omega$ is a Lebesgue point for the locally integrable initial functions, and the estimates κ_v and κ_i are independent of x_0 . Therefore, taking the supremum over all Lebesgue points $x_0 \in \Omega$ in (75) and (76) implies $v \in L^\infty(0,T;\mathcal{L}_v^\infty)$ and $i \in W^{1,\infty}(0,T;\mathcal{L}_i^\infty)$, which, recalling (4), further imply $v \in W^{2,\infty}(0,T;\mathcal{L}_v^\infty)$. Finally, it follows by using Morrey's inequality [16, Thm. 5.6-4 and Thm. 5.6-5] that $v \in C^{1,1}([0,T];\mathcal{L}_v^\infty)$ and $i \in C^{0,1}([0,T];\mathcal{L}_i^\infty)$, which completes the proof. $\blacksquare$

Next, we recall and use the following standard result in the theory of ODEs to establish conditions that guarantee nonnegativity of $i(x,t)$ for all biophysically plausible values of the input g , that is, for all $g \in L^2(0,T;\mathcal{D}_g)$, where

$$(77) \quad \mathcal{D}_g := \{\ell \in \mathcal{L}_i^2 : \ell \geq 0 \text{ a.e. in } \Omega\}.$$

Proposition 5.5 (Invariance of the nonnegative cone [7, Prop. I.1.1]). *Let $\{S(t)\}_{t \in [0,\infty)}$ be the semigroup of solution operators associated with the ODE*

$$d_t q(t) = P(q(t)), \quad q(t) \in \mathbb{R}^n, \quad t \in [0,\infty),$$

where $P : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a continuous locally Lipschitz mapping. Then the nonnegative cone $\mathbb{R}_+^n$ is invariant for $\{S(t)\}_{t \in [0,\infty)}$ if and only if $P(q)$ is quasi-positive, that is, for every $j \in \{1, \dots, n\}$,

$$P_j(q_1, \dots, q_n) \geq 0 \text{ whenever } q_j = 0 \text{ and } q_k \geq 0 \text{ for all } k \neq j.$$

Proposition 5.6 (Positively invariant region for the solution $i(x,t)$). *Suppose $g \in L^2(0,T;\mathcal{D}_g)$ and let $u(t) = S_w(t)u_0$ be an Ω -periodic weak solution of (4)–(7). Suppose the w -component of the weak solution, $w(x,t)$, is nonnegative for almost every $x \in \Omega$ and all $t \in [0,T]$, $T > 0$, and define the set*

$$(78) \quad \mathcal{D}_i := \{(\ell, \ell') \in \mathcal{L}_i^2 \times \mathcal{L}_i^2 : \ell \geq 0 \text{ and } \ell' + \Gamma \ell \geq 0 \text{ a.e. in } \Omega\}.$$

Then, for every $(i_0, i'_0) \in \mathcal{D}_i$, we have $(i(t), d_t i(t)) \in \mathcal{D}_i$ almost everywhere in Ω for all $t \in [0, T]$. An identical result holds for strong solutions $u(t) = S_s(t)u_0$ of (4)–(7) with nonnegative w -component.

Proof. Let $b := d_t i + \Gamma i$ and rewrite (5) as the first-order system of equations

$$(79) \quad \begin{aligned} d_t i &= -\Gamma i + b, \\ d_t b &= -\Gamma b + e\Upsilon\Gamma J_6 w + e\Upsilon\Gamma N J_7 f(v) + e\Upsilon\Gamma g. \end{aligned}$$

Let $x_0 \in \Omega$ be a Lebesgue point of the initial functions v_0 , i_0 , i'_0 , w_0 , and $g(0)$, and define $v_{x_0}(t)$, $i_{x_0}(t)$, $w_{x_0}(t)$, and $g_{x_0}(t)$ as given in the proof of Corollary 5.4. Accordingly, let $b_{x_0}(t) := b(x_0, t) = d_t i_{x_0}(t) + \Gamma i_{x_0}(t)$.

Now, (79) implies that the function $q_{x_0} := (i_{x_0}, b_{x_0})$ satisfies the ODE $d_t q_{x_0}(t) = P(q_{x_0}(t))$, $t \in [0, T]$, where the mapping $P : \mathbb{R}^8 \rightarrow \mathbb{R}^8$ given by

$$P(q_{x_0}) = P(i_{x_0}, b_{x_0}) := (-\Gamma i_{x_0} + b_{x_0}, -\Gamma b_{x_0} + e\Upsilon\Gamma J_6 w_{x_0} + e\Upsilon\Gamma N J_7 f(v_{x_0}) + e\Upsilon\Gamma g_{x_0})$$

is Lipschitz continuous. Moreover, note that by assumption we have $w_{x_0} \geq 0$ and $g_{x_0} \geq 0$, which, along with the definitions of f , Υ , Γ , N , J_6 , and J_7 given by (3) and (8), implies $e\Upsilon\Gamma J_6 w_{x_0}(t) \geq 0$, $e\Upsilon\Gamma N J_7 f(v_{x_0}(t)) \geq 0$, and $e\Upsilon\Gamma g_{x_0}(t) \geq 0$ for all $t \in [0, T]$. Therefore, it follows that P is quasi-positive, and hence by Proposition 5.5 we have $q_{x_0}(t) \geq 0$ for all $t \in [0, T]$. This completes the proof, since x_0 is an arbitrary Lebesgue point of the initial functions and almost every point in Ω is a Lebesgue point for these functions.² ■

Remark 5.7 (Biophysically plausible set of initial values). Propositions 5.3 and 5.6 ensure that if $g \in L^2(0, \infty; \mathcal{D}_g)$, where $\mathcal{D}_g$ is given by (77), and the initial values lie in the set

$$(80) \quad \mathcal{D}_{\text{Bio}} := \mathcal{L}_v^2 \times \mathcal{D}_i \times \mathcal{D}_w,$$

where $\mathcal{D}_w$ and $\mathcal{D}_i$ are given by (71) and (78), respectively, then $i(x, t)$ and $w(x, t)$ always remain nonnegative at almost every point in Ω as they evolve in time. However, it should be noted that this does not imply that the set $\mathcal{D}_{\text{Bio}} \subset \mathcal{U}_w$ is positively invariant, since Proposition 5.3 does not imply positive invariance of the set $\mathcal{D}_w$. Therefore, $\mathcal{D}_{\text{Bio}}$ cannot serve as a phase space for the semidynamical system framework of the problem.

In the analysis of the following sections, nonnegativity of the solution $i(x, t)$ is essential. Moreover, it would be of no practical value to analyze the dynamics of the model out of the biophysical regions of the phase space. Therefore, we define

²Note that there are fairly standard results in the literature that ensure the positivity of a $C^1(\overline{\Omega} \times [0, T]; \mathbb{R}^m)$ function as it evolves in time, provided its time derivative satisfies certain conditions on the boundary of the positive cone; see, for example, [34, Lem. 6] and [8]. The proofs of these results are relatively geometrical and usually use continuity of the functions and the compactness of $\overline{\Omega}$. However, these proofs are by no means applicable to functions in $C^1([0, T]; L^2(\Omega; \mathbb{R}^m))$. In fact, functions in $C^1([0, T]; L^2(\Omega; \mathbb{R}^m))$ are allowed to *leak* through the boundary of the positive cone on sets of measure zero at every $t \in [0, T]$. Since any subinterval of $[0, T]$ is uncountable, it is not guaranteed that the uncountable union of such leakage sets will have measure zero over a subinterval. In the proof of Proposition 5.6, we use the additional property that the functions are governed by a system of ODEs. Therefore, for all $t \in (0, T]$, the Banach space-valued function $i(t)$ is defined at the same almost every points in Ω as it is initially defined at $t = 0$. In other words, the leakage set remains unchanged for all $t \in (0, T]$.

$$(81) \quad \begin{aligned} \mathcal{D}_w &:= \{u_0 \in \mathcal{U}_w : i(t) \geq 0, w(t) \geq 0 \text{ a.e. in } \Omega \text{ for all } t \in [0, \infty), u(t) = S_w(t)u_0\}, \\ \mathcal{D}_s &:= \{u_0 \in \mathcal{U}_s : i(t) \geq 0, w(t) \geq 0 \text{ a.e. in } \Omega \text{ for all } t \in [0, \infty), u(t) = S_s(t)u_0\} \end{aligned}$$

as the maximal closed subsets of $\mathcal{U}_w$ and $\mathcal{U}_s$ for the initial values of the weak and strong solutions, respectively, such that i and w initiating from the points in these sets evolve non-negatively over time. Note that $\mathcal{D}_w$ and $\mathcal{D}_s$ are nonempty since $\mathcal{D}_{\text{Bio}} \subset \mathcal{D}_w$ and $\mathcal{D}_{\text{Bio}} \cap \mathcal{U}_s \subset \mathcal{D}_s$ when $g \in L^2(0, \infty, \mathcal{D}_g)$. Moreover, $\mathcal{D}_w$ and $\mathcal{D}_s$ are closed sets since $\{S_w(t)\}_{t \in [0, \infty)}$ and $\{S_s(t)\}_{t \in [0, \infty)}$ are continuous semigroups, as given by Propositions 5.1 and 5.2. Moreover, it follows immediately from the definitions given by (81) that $\mathcal{D}_w$ and $\mathcal{D}_s$ are positively invariant sets. Therefore, endowed with the metric induced by the norm in $\mathcal{U}_w$ and $\mathcal{U}_s$, the sets $\mathcal{D}_w$ and $\mathcal{D}_s$ form positively invariant complete metric spaces and can be considered as biophysically plausible phase spaces of the model, based on which we construct the semidynamical systems

$$\left(\mathcal{D}_w, \{S_w(t)\}_{t \in [0, \infty)} \right), \quad \left(\mathcal{D}_s, \{S_s(t)\}_{t \in [0, \infty)} \right)$$

associated with the weak and strong solutions of (4)–(7), respectively, and investigate their global dynamics in the remainder of the paper.

6. Existence of absorbing sets. In this section, we prove the existence of bounded absorbing sets for the semigroups $\{S_w(t)\}_{t \in [0, \infty)}$ and $\{S_s(t)\}_{t \in [0, \infty)}$ acting on $\mathcal{D}_w$ and $\mathcal{D}_s$, respectively. First, we recall the following definition of an absorbing set for an operator semigroup.

Definition 6.1 (Absorbing set [7, Def. II.2.3]). A set $\mathcal{B}_0$ in a complete metric space $\mathcal{D}$ is called an absorbing set for the semigroup $\{S(t) : \mathcal{D} \rightarrow \mathcal{D}\}_{t \in [0, \infty)}$ if for every bounded set $\mathcal{B} \in \mathcal{D}$ there exists $t_0(\mathcal{B}) \in (0, \infty)$ such that $S(t)\mathcal{B} \subset \mathcal{B}_0$ for all $t \geq t_0(\mathcal{B})$.

Theorem 6.2 (Existence of absorbing sets in $\mathcal{D}_w$). Assume that $g \in L^\infty(0, \infty; \mathcal{D}_g)$ and that there exists $\theta > 2\gamma_{\min}^{-3}$ such that

- (i) $\frac{4}{3}\theta e^2 \Upsilon_{\text{EE}}^2 \gamma_{\max}(\nu \Lambda_{\text{EE}})^{-3} < 1$,
- (ii) $\frac{4}{3}\theta e^2 \Upsilon_{\text{EI}}^2 \gamma_{\max}(\nu \Lambda_{\text{EI}})^{-3} < 1$,

where $\gamma_{\min} := \min\{\gamma_{\text{EE}}, \gamma_{\text{EI}}, \gamma_{\text{IE}}, \gamma_{\text{II}}\}$ and $\gamma_{\max} := \max\{\gamma_{\text{EE}}, \gamma_{\text{EI}}, \gamma_{\text{IE}}, \gamma_{\text{II}}\}$ are the smallest and largest eigenvalues of Γ , respectively. Then the semigroup $\{S_w(t) : \mathcal{D}_w \rightarrow \mathcal{D}_w\}_{t \in [0, \infty)}$ associated with the weak solutions of (4)–(7) has a bounded absorbing set $\mathcal{B}_w$. Specifically, consider the functions $Q_w^- : \mathcal{D}_w \rightarrow [0, \infty)$ and $Q_w^+ : \mathcal{D}_w \rightarrow [0, \infty)$ defined by

$$(82) \quad \begin{aligned} Q_w^-(u) &:= \|\Phi^{\frac{1}{2}}v\|_{\mathcal{L}_v^2}^2 + \theta \left\| d_t i + \frac{3}{2}\Gamma i \right\|_{\mathcal{L}_i^2}^2 + \frac{1}{4}\theta \|\Gamma i\|_{\mathcal{L}_i^2}^2 + \left\| d_t w + \frac{3}{2}\nu \Lambda w \right\|_{\mathcal{L}_w^2}^2 \\ &\quad + \frac{1}{4}\nu^2 \min\{6, \Lambda_{\min}^2\} \|w\|_{\mathcal{H}_w^1}^2, \\ Q_w^+(u) &:= \|\Phi^{\frac{1}{2}}v\|_{\mathcal{L}_v^2}^2 + \theta \left\| d_t i + \frac{3}{2}\Gamma i \right\|_{\mathcal{L}_i^2}^2 + \frac{1}{4}\theta \|\Gamma i\|_{\mathcal{L}_i^2}^2 + \left\| d_t w + \frac{3}{2}\nu \Lambda w \right\|_{\mathcal{L}_w^2}^2 \\ &\quad + \frac{1}{4}\nu^2 \max\{6, \Lambda_{\max}^2\} \|w\|_{\mathcal{H}_w^1}^2, \end{aligned}$$

and a scalar ε such that

$$(83) \quad \max \left\{ \frac{4}{3} \theta e^2 \Upsilon_{\text{EE}}^2 \gamma_{\text{max}} (\nu \Lambda_{\text{EE}})^{-3}, \frac{4}{3} \theta e^2 \Upsilon_{\text{EI}}^2 \gamma_{\text{max}} (\nu \Lambda_{\text{EI}})^{-3} \right\} < 2\gamma_{\text{max}} \varepsilon < 1.$$

Let $\tau_{\text{max}} := \max\{\tau_{\text{E}}, \tau_{\text{I}}\}$ denote the largest eigenvalue of Φ , and let $\Lambda_{\text{min}} := \min\{\Lambda_{\text{EE}}, \Lambda_{\text{EI}}\}$ and $\Lambda_{\text{max}} := \max\{\Lambda_{\text{EE}}, \Lambda_{\text{EI}}\}$ denote the smallest and largest eigenvalues of Λ , respectively. Let $\rho_{\text{w}}^2 := \frac{\beta_{\text{w}}}{\alpha_{\text{w}}}$, where

$$(84) \quad \alpha_{\text{w}} := \min \left\{ \frac{2}{3} \tau_{\text{max}}^{-1}, \left(\frac{1}{2} \gamma_{\text{max}}^{-1} - \varepsilon \right) \gamma_{\text{min}}^2, 3\theta^{-1} (\theta \gamma_{\text{min}} - 2\gamma_{\text{min}}^{-2}), \frac{1}{2} \nu \Lambda_{\text{min}}, \right. \\ \left. 3\nu \Lambda_{\text{max}}^{-2} \min \left\{ \Lambda_{\text{EE}}^3 - \frac{2}{3} \frac{\theta e^2}{\nu^3 \varepsilon} \Upsilon_{\text{EE}}^2, \Lambda_{\text{EI}}^3 - \frac{2}{3} \frac{\theta e^2}{\nu^3 \varepsilon} \Upsilon_{\text{EI}}^2 \right\} \right\},$$

$$(85) \quad \beta_{\text{w}} := \frac{4\theta e^2}{\gamma_{\text{max}}^{-1} - 2\varepsilon} \left[|\Omega| (\text{F}_{\text{E}}^2 + \text{F}_{\text{I}}^2) \|\Upsilon \text{N} J_7\|_2^2 + \|\Upsilon\|_2^2 \|g\|_{L^\infty(0, \infty; \mathcal{L}_i^2)}^2 \right] + 2\nu^3 |\Omega| \text{F}_{\text{E}}^2 \text{tr}(\Lambda^3 \text{M}^2).$$

Then, for all $\rho > \rho_{\text{w}}$, the bounded sets $\mathcal{B}_{\text{w}} := \{u \in \mathcal{D}_{\text{w}} : Q_{\text{w}}^-(u) \leq \rho^2\}$ are absorbing in $\mathcal{D}_{\text{w}}$. Moreover, for every bounded set $\mathcal{B} \subset \mathcal{D}_{\text{w}}$, there exists $R > 0$ such that $Q_{\text{w}}^+(u_0) \leq R^2$ for all $u_0 \in \mathcal{B}$, and $S(t)\mathcal{B} \subset \mathcal{B}_{\text{w}}$ for all $t \geq t_{\text{w}}(\mathcal{B})$, where

$$(86) \quad t_{\text{w}}(\mathcal{B}) = t_{\text{w}}(R) := \max \left\{ 0, \frac{1}{\alpha_{\text{w}}} \log \frac{R^2}{\rho^2 - \rho_{\text{w}}^2} \right\}.$$

Proof. First, taking the inner product of (4) with v yields

$$\frac{1}{2} \text{d}_t \|\Phi^{\frac{1}{2}} v\|_{\mathcal{L}_v^2}^2 + \|v\|_{\mathcal{L}_v^2}^2 - (J_1 i, v)_{\mathcal{L}_v^2} + \int_{\Omega} (v_1^2 i^{\text{T}} \Psi J_4 + v_2^2 i^{\text{T}} \Psi J_5) \, \text{d}x = 0.$$

The integral term in this equation is nonnegative in $\mathcal{D}_{\text{w}}$ for all $t \in [0, \infty)$; see (8) and (81). Therefore, dropping the integral term and using Young's inequality yields, for every $\varepsilon_1 > 0$,

$$(87) \quad \text{d}_t \|\Phi^{\frac{1}{2}} v\|_{\mathcal{L}_v^2}^2 \leq -2(1 - \varepsilon_1) \|v\|_{\mathcal{L}_v^2}^2 + \frac{1}{\varepsilon_1} \|i\|_{\mathcal{L}_i^2}^2 \\ \leq -2(1 - \varepsilon_1) \tau_{\text{max}}^{-1} \|\Phi^{\frac{1}{2}} v\|_{\mathcal{L}_v^2}^2 + \frac{1}{\varepsilon_1 \gamma_{\text{min}}^2} \|\Gamma i\|_{\mathcal{L}_i^2}^2.$$

Next, let $b := \text{d}_t i + \frac{3}{2} \Gamma i$ and rewrite (5) as

$$\text{d}_t b + \frac{1}{2} \Gamma b + \frac{1}{4} \Gamma^2 i - e \Upsilon \Gamma J_6 w - e \Upsilon \Gamma \text{N} J_7 f(v) = e \Upsilon \Gamma g.$$

Taking the inner product of the above equality with b yields

$$\frac{1}{2} \text{d}_t \|b\|_{\mathcal{L}_i^2}^2 + \frac{1}{2} (\Gamma b, b)_{\mathcal{L}_i^2} + \frac{1}{8} \text{d}_t \|\Gamma i\|_{\mathcal{L}_i^2}^2 + \frac{3}{8} \|\Gamma^{\frac{3}{2}} i\|_{\mathcal{L}_i^2}^2 \\ - e (\Upsilon \Gamma J_6 w, b)_{\mathcal{L}_i^2} - e (\Upsilon \Gamma \text{N} J_7 f(v), b)_{\mathcal{L}_i^2} = e (\Upsilon \Gamma g, b)_{\mathcal{L}_i^2}.$$

Note that

$$\begin{aligned} (\Gamma b, b)_{\mathcal{L}_i^2} &\geq \gamma_{\max}^{-1} \|\Gamma b\|_{\mathcal{L}_i^2}^2, \\ \|\Gamma^{\frac{3}{2}} i\|_{\mathcal{L}_i^2}^2 &\geq \gamma_{\min} \|\Gamma i\|_{\mathcal{L}_i^2}^2, \end{aligned}$$

and, using similar arguments as in the proof of Proposition 4.4, it follows that for every $\varepsilon_2, \varepsilon_3, \varepsilon_4 > 0$,

$$\begin{aligned} e(\Upsilon \Gamma J_6 w, b)_{\mathcal{L}_i^2} &\leq \varepsilon_2 \|\Gamma b\|_{\mathcal{L}_i^2}^2 + \frac{e^2}{4\varepsilon_2} \|\Upsilon J_6 w\|_{\mathcal{L}_i^2}^2 \\ e(\Upsilon \Gamma N J_7 f(v), b)_{\mathcal{L}_i^2} &\leq \varepsilon_3 \|\Gamma b\|_{\mathcal{L}_i^2}^2 + \frac{e^2 |\Omega|}{4\varepsilon_3} (F_E^2 + F_I^2) \|\Upsilon N J_7\|_2^2, \\ e(\Upsilon \Gamma g, b)_{\mathcal{L}_i^2} &\leq \varepsilon_4 \|\Gamma b\|_{\mathcal{L}_i^2}^2 + \frac{e^2}{4\varepsilon_4} \|\Upsilon\|_2^2 \|g\|_{\mathcal{L}_i^2}^2. \end{aligned}$$

Therefore,

$$\begin{aligned} (88) \quad d_t \left[\|b\|_{\mathcal{L}_i^2}^2 + \frac{1}{4} \|\Gamma i\|_{\mathcal{L}_i^2}^2 \right] &\leq -(\gamma_{\max}^{-1} - 2(\varepsilon_2 + \varepsilon_3 + \varepsilon_4)) \|\Gamma b\|_{\mathcal{L}_i^2}^2 - \frac{3}{4} \gamma_{\min} \|\Gamma i\|_{\mathcal{L}_i^2}^2 \\ &\quad + \frac{e^2}{2\varepsilon_2} \|\Upsilon J_6 w\|_{\mathcal{L}_i^2}^2 + \frac{e^2}{2\varepsilon_3} |\Omega| (F_E^2 + F_I^2) \|\Upsilon N J_7\|_2^2 + \frac{e^2}{2\varepsilon_4} \|\Upsilon\|_2^2 \|g\|_{\mathcal{L}_i^2}^2. \end{aligned}$$

Next, let $q := d_t w + \frac{3}{2} \nu \Lambda w$ and rewrite (6) as

$$(89) \quad d_t q + \frac{1}{2} \nu \Lambda q - \frac{3}{2} \nu^2 \Delta w + \frac{1}{4} \nu^2 \Lambda^2 w - \nu^2 \Lambda^2 M J_8 f(v) = 0.$$

Taking the inner product of this equality with q yields

$$\begin{aligned} \frac{1}{2} d_t \|q\|_{\mathcal{L}_w^2}^2 + \frac{1}{2} \nu \|\Lambda^{\frac{1}{2}} q\|_{\mathcal{L}_w^2}^2 + \frac{3}{4} \nu^2 d_t \|\partial_x w\|_{\mathcal{L}_{\partial w}^2}^2 + \frac{9}{4} \nu^3 \|\Lambda^{\frac{1}{2}} \partial_x w\|_{\mathcal{L}_{\partial w}^2}^2 + \frac{1}{8} \nu^2 d_t \|\Lambda w\|_{\mathcal{L}_w^2}^2 \\ + \frac{3}{8} \nu^3 \|\Lambda^{\frac{3}{2}} w\|_{\mathcal{L}_w^2}^2 - \nu^2 (\Lambda^2 M J_8 f(v), q)_{\mathcal{L}_w^2} = 0. \end{aligned}$$

Using similar arguments as in the proof of Proposition 4.4, we can write, for every $\varepsilon_5 > 0$,

$$(\Lambda^2 M J_8 f(v^{(m)}), q)_{\mathcal{L}_w^2} \leq \varepsilon_5 \|\Lambda^{\frac{1}{2}} q\|_{\mathcal{L}_w^2}^2 + \frac{1}{4\varepsilon_5} |\Omega| F_E^2 \operatorname{tr}(\Lambda^3 M^2),$$

and hence it follows that

$$\begin{aligned} (90) \quad d_t \left[\|q\|_{\mathcal{L}_w^2}^2 + \frac{3}{2} \nu^2 \|\partial_x w\|_{\mathcal{L}_{\partial w}^2}^2 + \frac{1}{4} \nu^2 \|\Lambda w\|_{\mathcal{L}_w^2}^2 \right] &\leq -\nu(1 - 2\nu\varepsilon_5) \|\Lambda^{\frac{1}{2}} q\|_{\mathcal{L}_w^2}^2 \\ &\quad - 3\nu \left(\frac{3}{2} \nu^2 \|\Lambda^{\frac{1}{2}} \partial_x w\|_{\mathcal{L}_{\partial w}^2}^2 + \frac{1}{4} \nu^2 \|\Lambda^{\frac{3}{2}} w\|_{\mathcal{L}_w^2}^2 \right) \\ &\quad + \frac{\nu^2}{2\varepsilon_5} |\Omega| F_E^2 \operatorname{tr}(\Lambda^3 M^2). \end{aligned}$$

Now, set $\varepsilon_1 = \frac{2}{3}$ in (87), set $\varepsilon_3 = \varepsilon_4 = \frac{1}{8}(\gamma_{\max}^{-1} - 2\varepsilon)$ in (88) with $\varepsilon := \varepsilon_2$, and set $\varepsilon_5 = \frac{1}{4\nu}$ in (90). Then, multiplying (88) by $\theta > 0$ and adding the result to (87) and (90) yields

$$\begin{aligned} d_t Q_w &\leq -\frac{2}{3}\tau_{\max}^{-1}\|\Phi^{\frac{1}{2}}v\|_{\mathcal{L}_v^2}^2 - \theta\left(\frac{1}{2}\gamma_{\max}^{-1} - \varepsilon\right)\|\Gamma b\|_{\mathcal{L}_i^2}^2 \\ &\quad - \frac{3}{4}(\theta\gamma_{\min} - 2\gamma_{\min}^{-2})\|\Gamma i\|_{\mathcal{L}_i^2}^2 - \frac{1}{2}\nu\|\Lambda^{\frac{1}{2}}q\|_{\mathcal{L}_w^2}^2 \\ &\quad - 3\nu\left(\frac{3}{2}\nu^2\|\Lambda^{\frac{1}{2}}\partial_x w\|_{\mathcal{L}_{\partial w}^2}^2 + \frac{1}{4}\nu^2\left(\left[\Lambda^3 - \frac{2\theta e^2}{3\nu^3\varepsilon}J_6^T\Upsilon^2J_6\right]w, w\right)_{\mathcal{L}_w^2}\right) + \beta_w, \end{aligned}$$

where β_w is given by (85) and

$$(91) \quad Q_w(u) = \|\Phi^{\frac{1}{2}}v\|_{\mathcal{L}_v^2}^2 + \theta\|b\|_{\mathcal{L}_i^2}^2 + \frac{1}{4}\theta\|\Gamma i\|_{\mathcal{L}_i^2}^2 + \|q\|_{\mathcal{L}_w^2}^2 + \frac{3}{2}\nu^2\|\partial_x w\|_{\mathcal{L}_{\partial w}^2}^2 + \frac{1}{4}\nu^2\|\Lambda w\|_{\mathcal{L}_w^2}^2.$$

Note that for $\theta > 2\gamma_{\min}^{-3}$, we have $\theta\gamma_{\min} - 2\gamma_{\min}^{-2} > 0$, and for the range of values of ε given by (83), we have $\frac{1}{2}\gamma_{\max}^{-1} - \varepsilon > 0$. Moreover, assumptions (i) and (ii) along with (83) ensure that $\Lambda^3 - \frac{2\theta e^2}{3\nu^3\varepsilon}J_6^T\Upsilon^2J_6 > 0$. Therefore, with the decay rate α_w given by (84),

$$(92) \quad d_t Q_w(u) \leq -\alpha_w Q_w(u) + \beta_w,$$

and hence, using Grönwall's inequality [51, sect. III.1.1.3.],

$$(93) \quad Q_w^-(u(t)) \leq Q_w^+(u(0))e^{-\alpha_w t} + \rho_0^2(1 - e^{-\alpha_w t}),$$

where Q_w^- and Q_w^+ are given in (82) and $\limsup_{t \rightarrow \infty} Q_w^-(u(t)) \leq \rho_0^2 := \frac{\beta_w}{\alpha_w}$. Now, since the mapping

$$(94) \quad (v, i, i', w, w') \mapsto \left(\Phi^{\frac{1}{2}}v, \frac{1}{2}\theta^{\frac{1}{2}}\Gamma i, \theta^{\frac{1}{2}}\left[i' + \frac{3}{2}\Gamma i\right], \frac{1}{2}\nu[\max\{6, \Lambda_{\max}^2\}]^{\frac{1}{2}}w, w' + \frac{3}{2}\nu\Lambda w\right)$$

is a linear isomorphism over $\mathcal{U}_w$, for every bounded set $\mathcal{B} \subset \mathcal{D}_w$ there exists $R > 0$ such that $Q_w^+(u_0) \leq R^2$ for all $u_0 \in \mathcal{B}$. Hence, it is immediate from (93) that $S_w(t)\mathcal{B} \subset \mathcal{B}_w$ for all $t \geq t_w(\mathcal{B})$, where $t_w(\mathcal{B})$ is given by (86). ■

Theorem 6.3 (Existence of absorbing sets in $\mathcal{D}_s$). *Suppose the assumptions of Theorem 6.2 hold, namely, assume that $g \in L^\infty(0, \infty; \mathcal{D}_g)$ and that there exists $\theta > 2\gamma_{\min}^{-3}$ such that the biophysical parameters of the model satisfy*

- (i) $\frac{4}{3}\theta e^2 \Upsilon_{EE}^2 \gamma_{\max}(\nu\Lambda_{EE})^{-3} < 1$,
- (ii) $\frac{4}{3}\theta e^2 \Upsilon_{EI}^2 \gamma_{\max}(\nu\Lambda_{EI})^{-3} < 1$,

where $\gamma_{\min}$ and $\gamma_{\max}$ are the smallest and largest eigenvalues of Γ , respectively. Then the semigroup $\{S_s(t) : \mathcal{D}_s \rightarrow \mathcal{D}_s\}_{t \in [0, \infty)}$ associated with the strong solutions of (4)–(7) has a bounded absorbing set $\mathcal{B}_s$. Specifically, consider the function $Q_s^- : \mathcal{D}_s \rightarrow [0, \infty)$ defined by

$$\begin{aligned} (95) \quad Q_s^-(u) &:= \|\Phi^{\frac{1}{2}}v\|_{\mathcal{L}_v^2}^2 + \theta\left\|d_t i + \frac{3}{2}\Gamma i\right\|_{\mathcal{L}_i^2}^2 + \frac{1}{4}\theta\|\Gamma i\|_{\mathcal{L}_i^2}^2 + \left\|d_t w + \frac{3}{2}\nu\Lambda w\right\|_{\mathcal{H}_w^1}^2 \\ &\quad + \frac{1}{8}\nu^2 \min\{6, \Lambda_{\min}^2\}\|(-\Delta + I)w\|_{\mathcal{L}_w^2}^2, \end{aligned}$$

and denote by $\Lambda_{\min}$ and $\Lambda_{\max}$ the smallest and largest eigenvalues of Λ , respectively, and by $\tau_{\max}$ the largest eigenvalue of Φ . Let $\rho_s^2 := \frac{2\beta_s}{\alpha_s}$ with

$$(96) \quad \alpha_s := \min \left\{ \frac{2}{3} \tau_{\max}^{-1}, \left(\frac{1}{2} \gamma_{\max}^{-1} - \varepsilon \right) \gamma_{\min}^2, 3\theta^{-1} (\theta \gamma_{\min} - 2\gamma_{\min}^{-2}), \nu \Lambda_{\min}, \right. \\ \left. 3\nu \Lambda_{\max}^{-2} \min \left\{ \Lambda_{\text{EE}}^3 - \frac{2\theta e^2}{3\nu^3 \varepsilon} \Upsilon_{\text{EE}}^2, \Lambda_{\text{EI}}^3 - \frac{2\theta e^2}{3\nu^3 \varepsilon} \Upsilon_{\text{EI}}^2 \right\} \right\},$$

$$(97) \quad \beta_s := \frac{4\theta e^2}{\gamma_{\max}^{-1} - 2\varepsilon} \left[|\Omega| (F_{\text{E}}^2 + F_{\text{I}}^2) \|\Upsilon \text{N} J_7\|_2^2 + \|\Upsilon\|_2^2 \|g\|_{L^\infty(0, \infty; \mathcal{L}_i^2)}^2 \right] \\ + 2\nu^2 \left[\frac{1}{32\varepsilon_1} \frac{F_{\text{E}}^2}{\sigma_{\text{E}}^2} \text{tr}(\Lambda^4 \text{M}^2) \eta \rho_w^2 (1 + \rho_w^2) + \frac{1}{4} |\Omega| F_{\text{E}}^2 \text{tr}(\Lambda^4 \text{M}^2) \left(\frac{1}{\varepsilon_1} + \frac{\alpha_s}{\varepsilon_2} \right) \right],$$

where η is a positive constant, $\rho_w^2 := \frac{\beta_w}{\alpha_w}$ is the same constant as given in Theorem 6.2, the scalar ε takes values within the same range as given by (83), and

$$(98) \quad \varepsilon_1 := \frac{1}{32} \alpha_s \min \{6, \Lambda_{\min}^2\} \left(1 + \left\| \frac{3}{2} \nu \Lambda - \alpha I \right\|_2^2 \right)^{-1}, \quad \varepsilon_2 := \frac{1}{16} \min \{6, \Lambda_{\min}^2\}.$$

Then, for all $\rho > \rho_s$, the bounded sets $\mathcal{B}_s := \{u \in \mathcal{D}_s : Q_s^-(u) \leq \rho^2\}$ are absorbing in $\mathcal{D}_s$.

Proof. Let $A := -\Delta + I$ and take the inner product of (89) with Aq to obtain

$$\frac{1}{2} \text{d}_t \|q\|_{\mathcal{H}_w^1}^2 + \frac{1}{2} \nu \|\Lambda^{\frac{1}{2}} q\|_{\mathcal{H}_w^1}^2 + \frac{3}{4} \nu^2 \text{d}_t \|\partial_x w\|_{\mathcal{H}_{\partial w}^1}^2 + \frac{9}{4} \nu^3 \|\Lambda^{\frac{1}{2}} \partial_x w\|_{\mathcal{H}_{\partial w}^1}^2 + \frac{1}{8} \nu^2 \text{d}_t \|\Lambda w\|_{\mathcal{H}_w^1}^2 \\ + \frac{3}{8} \nu^3 \|\Lambda^{\frac{3}{2}} w\|_{\mathcal{H}_w^1}^2 - \nu^2 (\Lambda^2 \text{M} J_8 f(v), Aq)_{\mathcal{L}_w^2} = 0.$$

This equality, along with the inequalities (87) and (88) derived in the proof of Theorem 6.2 and the same values of $\varepsilon_1, \dots, \varepsilon_4$ therein, implies that

$$\text{d}_t Q_s \leq -\frac{2}{3} \tau_{\max} \|\Phi^{\frac{1}{2}} v\|_{\mathcal{L}_v^2}^2 - \theta \left(\frac{1}{2} \gamma_{\max}^{-1} - \varepsilon \right) \|\Gamma b\|_{\mathcal{L}_i^2}^2 - \frac{3}{4} (\theta \gamma_{\min} - 2\gamma_{\min}^{-2}) \|\Gamma i\|_{\mathcal{L}_i^2}^2 - \nu \|\Lambda^{\frac{1}{2}} q\|_{\mathcal{H}_w^1}^2 \\ - 3\nu \left(\frac{3}{2} \nu^2 \|\Lambda^{\frac{1}{2}} \partial_x w\|_{\mathcal{H}_{\partial w}^1}^2 + \frac{1}{4} \nu^2 \left(\left[\Lambda^3 - \frac{2\theta e^2}{3\nu^3 \varepsilon} J_6^T \Upsilon^2 J_6 \right] w, w \right)_{\mathcal{H}_w^1} \right) \\ + 2\nu^2 (\Lambda^2 \text{M} J_8 f(v), Aq)_{\mathcal{L}_w^2} + \beta,$$

where

$$Q_s(u) := \|\Phi^{\frac{1}{2}} v\|_{\mathcal{L}_v^2}^2 + \theta \|b\|_{\mathcal{L}_i^2}^2 + \frac{1}{4} \theta \|\Gamma i\|_{\mathcal{L}_i^2}^2 + \|q\|_{\mathcal{H}_w^1}^2 + \frac{3}{2} \nu^2 \|\partial_x w\|_{\mathcal{H}_{\partial w}^1}^2 + \frac{1}{4} \nu^2 \|\Lambda w\|_{\mathcal{H}_w^1}^2, \\ \beta := \frac{4\theta e^2}{\gamma_{\max}^{-1} - 2\varepsilon} \left[|\Omega| (F_{\text{E}}^2 + F_{\text{I}}^2) \|\Upsilon \text{N} J_7\|_2^2 + \|\Upsilon\|_2^2 \|g\|_{L^\infty(0, \infty; \mathcal{L}_i^2)}^2 \right],$$

and ε takes values within the range given by (83). Now, using similar arguments as in the proof of Theorem 6.2, it follows from assumptions (i) and (ii) with $\theta > 2\gamma_{\min}^{-3}$ that

$$(99) \quad d_t Q_s(u) \leq -\alpha_s Q_s(u) + 2\nu^2 (\Lambda^2 M J_8 f(v), Aq)_{\mathcal{L}_w^2} + \beta,$$

where the decay rate α_s is given by (96). Then, Grönwall's inequality [51, sect. III.1.1.3.] implies

$$(100) \quad Q_s(u(t)) \leq Q_s(u(0))e^{-\alpha_s t} + 2\nu^2 \int_0^t (\Lambda^2 M J_8 f(v), Aq)_{\mathcal{L}_w^2} e^{\alpha_s(s-t)} ds + \frac{\beta}{\alpha_s} (1 - e^{-\alpha_s t}).$$

Replacing $q := d_t w + \frac{3}{2}\nu\Lambda w$ in the integral term in the above inequality and integrating by parts yields

$$\begin{aligned} \int_0^t (\Lambda^2 M J_8 f(v), Aq)_{\mathcal{L}_w^2} e^{\alpha_s(s-t)} ds &= - \int_0^t (\Lambda^2 M J_8 d_s f(v), Aw)_{\mathcal{L}_w^2} e^{\alpha_s(s-t)} ds \\ &\quad + \int_0^t \left(\Lambda^2 M J_8 f(v), \left(\frac{3}{2}\nu\Lambda - \alpha_s I \right) Aw \right)_{\mathcal{L}_w^2} e^{\alpha_s(s-t)} ds \\ &\quad + (\Lambda^2 M J_8 f(v), Aw)_{\mathcal{L}_w^2} - (\Lambda^2 M J_8 f(v_0), Aw_0)_{\mathcal{L}_w^2} e^{-\alpha_s t}. \end{aligned}$$

Next, noting that $d_s f(v) = \partial_v f(v) d_s v$ and $\sup_{v_E(x,t) \in \mathbb{R}} |\partial_{v_E} f_E(v_E)| \leq \frac{F_E}{2\sqrt{2}\sigma_E}$ by (45), it follows that, for every $\varepsilon_1, \varepsilon_2 > 0$,

$$\begin{aligned} \int_0^t (\Lambda^2 M J_8 f(v), Aq)_{\mathcal{L}_w^2} e^{\alpha_s(s-t)} ds &\leq \varepsilon_1 \left(1 + \left\| \frac{3}{2}\nu\Lambda - \alpha_s I \right\|_2^2 \right) \int_0^t \|Aw\|_{\mathcal{L}_w^2}^2 e^{\alpha_s(s-t)} ds \\ &\quad + \frac{1}{32\varepsilon_1} \frac{F_E^2}{\sigma_E^2} \text{tr}(\Lambda^4 M^2) \int_0^t \|d_s v\|_{\mathcal{L}_v^2}^2 e^{\alpha_s(s-t)} ds \\ &\quad + \varepsilon_2 \|Aw\|_{\mathcal{L}_w^2}^2 + \frac{1}{4} |\Omega| F_E^2 \text{tr}(\Lambda^4 M^2) \left(\frac{1}{\alpha_s \varepsilon_1} + \frac{1}{\varepsilon_2} \right) \\ &\quad - (\Lambda^2 M J_8 f(v_0), Aw_0)_{\mathcal{L}_w^2} e^{-\alpha_s t}. \end{aligned}$$

Moreover, it follows from Theorem 6.2 that for every bounded set $\mathcal{B} \subset \mathcal{D}_s$ there exists a time $t_w(\mathcal{B})$, given by (86), and positive constants η_1 and η_2 such that $\|v(t)\|_{\mathcal{L}_v^2}^2 \leq \eta_1 \rho_w^2$ and $\|i(t)\|_{\mathcal{L}_i^2}^2 \leq \eta_2 \rho_w^2$ for all $t \geq t_w(\mathcal{B})$. Therefore, using the estimate (69), we can write

$$(101) \quad \begin{aligned} \int_0^t \|d_s v\|_{\mathcal{L}_v^2}^2 e^{\alpha_s(s-t)} ds &\leq \int_0^{t_w(\mathcal{B})} \|d_s v\|_{\mathcal{L}_v^2}^2 e^{\alpha_s(s-t)} ds + \frac{1}{\alpha_s} \eta \rho_w^2 (1 + \rho_w^2) \\ &\leq \kappa_0(\mathcal{B}) e^{-\alpha_s t} + \frac{1}{\alpha_s} \eta \rho_w^2 (1 + \rho_w^2), \end{aligned}$$

where η is a positive constant and, for some $\alpha > 0$,

$$\kappa_0(\mathcal{B}) := \alpha \int_0^{t_w(\mathcal{B})} \left(\|v(s)\|_{\mathcal{L}_v^2}^2 + \|i(s)\|_{\mathcal{L}_i^2}^2 + \|v(s)\|_{\mathcal{L}_v^2}^2 \|i(s)\|_{\mathcal{L}_i^2}^2 \right) e^{\alpha_s s} ds < \infty.$$

Now, using the above estimate for the integral term in (100) with ε_1 and ε_2 given by (98) yields

$$(102) \quad Q_s^-(u)e^{\alpha_s t} \leq \frac{1}{2}\alpha_s \int_0^t Q_s^-(u)e^{\alpha_s s} ds + \kappa(\mathcal{B}) + \frac{\beta_s}{\alpha_s} e^{\alpha_s t},$$

where $\beta_s := \beta + 2\nu^2 \left[\frac{1}{32\varepsilon_1} \frac{F_E^2}{\sigma_E^2} \text{tr}(\Lambda^4 M^2) \eta \rho_w^2 (1 + \rho_w^2) + \frac{1}{4} |\Omega| F_E^2 \text{tr}(\Lambda^4 M^2) \left(\frac{1}{\varepsilon_1} + \frac{\alpha_s}{\varepsilon_2} \right) \right]$ as given in (97), $Q_s^-(u)$ is given in (95), and

$$\begin{aligned} \kappa(\mathcal{B}) &:= Q_s^+(u(0)) + 2\nu^2 \left[\frac{1}{32\varepsilon_1} \frac{F_E^2}{\sigma_E^2} \text{tr}(\Lambda^4 M^2) \kappa_0(\mathcal{B}) - (\Lambda^2 M J_8 f(v_0), Aw_0)_{\mathcal{L}_w^2} \right] - \frac{\beta}{\alpha_s}, \\ Q_s^+(u) &:= \|\Phi^{\frac{1}{2}} v\|_{\mathcal{L}_v^2}^2 + \theta \|b\|_{\mathcal{L}_i^2}^2 + \frac{1}{4} \theta \|\Gamma i\|_{\mathcal{L}_i^2}^2 + \|q\|_{\mathcal{H}_w^1}^2 + \frac{1}{4} \nu^2 \max\{6, \Lambda_{\max}^2\} \|Aw\|_{\mathcal{L}_w^2}^2. \end{aligned}$$

Next, using Grönwall's inequality for the function $\int_0^t Q_s^-(u)e^{\alpha_s s} ds$ in (102) gives

$$\int_0^t Q_s^-(u)e^{\alpha_s s} ds \leq \frac{1}{\frac{1}{2}\alpha_s} \left[\kappa(\mathcal{B}) (e^{\frac{1}{2}\alpha_s t} - 1) + \frac{\beta_s}{\alpha_s} (e^{\alpha_s t} - e^{\frac{1}{2}\alpha_s t}) \right],$$

which along with (102) implies

$$(103) \quad Q_s^-(u) \leq \kappa(\mathcal{B}) e^{-\frac{1}{2}\alpha_s t} + \rho_s^2 \left(1 - \frac{1}{2} e^{-\frac{1}{2}\alpha_s t} \right),$$

where $\limsup_{t \rightarrow \infty} Q_s^-(u(t)) \leq \rho_s^2 := \frac{2\beta_s}{\alpha_s}$.

Finally, considering the linear isomorphism (94) over $\mathcal{U}_s$, it follows that for every bounded set $\mathcal{B} \subset \mathcal{D}_s$ there exists $R > 0$ such that $\kappa(\mathcal{B}) \leq R^2$ for all $u_0 \in \mathcal{B}$. Therefore, (103) implies that $S_s(t)\mathcal{B} \subset \mathcal{B}_s$ for all $t \geq t_s(\mathcal{B})$ and some $t_s(\mathcal{B}) > 0$, which completes the proof. $\blacksquare$

Note that an estimate similar to (86) given in Theorem 6.2 can also be obtained for $t_s(\mathcal{B})$ in the proof of Theorem 6.3. However, this would be of limited practical value since the bound (101) is very conservative for times $t \ll t_w(\mathcal{B})$.

Remark 6.4 (Conditions on parameter sets). For the range of values given in Table 1, the maximum value that the left-hand side of the inequalities in assumptions (i) and (ii) of Theorems 6.2 and 6.3 may take is 39.4083θ , which is achieved when $\Upsilon_{EE} = 2$, $\Upsilon_{EI} = 2$, $\Lambda_{EE} = 0.1$, $\Lambda_{EI} = 0.1$, $\nu = 100$, and $\gamma_{\max} = 1000$. Assumptions (i) and (ii) then require that $\theta < \frac{1}{39.4083} = 0.0254$. Moreover, Theorems 6.2 and 6.3 allow for $\theta > 2\gamma_{\min}^{-3} \geq 0.002$, in accordance with Table 1. This implies that—for the entire range of values that the biophysical parameters of the model may take—the conditions imposed by Theorems 6.2 and 6.3 are satisfied at least for any $0.002 < \theta < 0.0254$, and the model (2) possesses bounded absorbing sets as given by these theorems.

7. Existence and nonexistence of a global attractor. In this section, we investigate the problem of existence of a global attractor for the semigroups $\{S_w(t) : \mathcal{D}_w \rightarrow \mathcal{D}_w\}_{t \in [0, \infty)}$ and $\{S_s(t) : \mathcal{D}_s \rightarrow \mathcal{D}_s\}_{t \in [0, \infty)}$ of solution operators of (4)–(7). First, we recall the definition of a global attractor, and a widely used theorem for establishing the existence of a global attractor. See [25, Chap. 1] for the motivation behind this definition, and [25, Chap. 3] for further results.

Definition 7.1 (Attracting set [7, Def. II.2.4]). A set $\mathcal{P}$ in a complete metric space $\mathcal{D}$ is called an attracting set for a semigroup $\{S(t)\}_{t \in [0, \infty)}$ acting in $\mathcal{D}$ if, for every bounded set $\mathcal{B} \in \mathcal{D}$, $\text{dist}_{\mathcal{D}}(S(t)\mathcal{B}, \mathcal{P}) \rightarrow 0$ as $t \rightarrow \infty$. Here, $\text{dist}_{\mathcal{D}}(\mathcal{G}, \mathcal{H}) := \sup_{g \in \mathcal{G}} \inf_{h \in \mathcal{H}} \|g - h\|_{\mathcal{D}}$ is the Hausdorff distance between the two sets $\mathcal{G}, \mathcal{H} \subset \mathcal{D}$.

Definition 7.2 (Global attractor [7, Def. II.3.1]). A bounded set $\mathcal{A}$ in a complete metric space $\mathcal{D}$ is called a global attractor for a semigroup $\{S(t)\}_{t \in [0, \infty)}$ acting in $\mathcal{D}$ if it satisfies the following conditions:

- (i) $\mathcal{A}$ is compact in $\mathcal{D}$,
- (ii) $\mathcal{A}$ is an attracting set for $\{S(t)\}_{t \in [0, \infty)}$,
- (iii) $\mathcal{A}$ is strictly invariant with respect to $\{S(t)\}_{t \in [0, \infty)}$, that is, $S(t)\mathcal{A} = \mathcal{A}$ for all $t \in [0, \infty)$.

Definition 7.3 (Asymptotic compactness [7, Def. II.2.5]). The semigroup $\{S(t)\}_{t \in [0, \infty)}$ acting in a complete metric space $\mathcal{D}$ is called asymptotically compact if it possesses a compact attracting set $\mathcal{K} \in \mathcal{D}$.

Theorem 7.4 (Global attractor [7, Thm. II.3.1]). Let $\{S(t)\}_{t \in [0, \infty)}$ be an asymptotically compact continuous semigroup in a complete metric space $\mathcal{D}$, possessing a compact attracting set $\mathcal{K} \in \mathcal{D}$. Then $\{S(t)\}_{t \in [0, \infty)}$ has a global attractor $\mathcal{A} \subset \mathcal{K}$ given by $\mathcal{A} = \omega(\mathcal{K})$, where $\omega(\mathcal{K})$ is the ω -limit set of $\mathcal{K}$.

7.1. Challenges in establishing a global attractor. In this section, we discuss some of the standard approaches available in the literature for establishing a global attractor based on Theorem 7.4, and identify reasons that make these approaches rather unpromising for the model (4)–(6).

Continuity of $\{S_w(t)\}_{t \in [0, \infty)}$ and $\{S_s(t)\}_{t \in [0, \infty)}$, as required by Theorem 7.4, is established in Propositions 5.1 and 5.2, respectively. To prove asymptotic compactness of a semigroup $\{S(t)\}_{t \in [0, \infty)}$ acting in $\mathcal{D}$, a general approach is to first show that the semigroup possesses a bounded absorbing set and then to show that the semigroup is κ -contracting, meaning that $\lim_{t \rightarrow \infty} \kappa(S(t)\mathcal{B}) = 0$ for any bounded set $\mathcal{B} \in \mathcal{D}$, where κ denotes the Kuratowski measure of noncompactness; see [38, 56] and [25, Chap. 3]. An effective way to establish the latter property is through a decomposition $S(t) = S_1(t) + S_2(t)$ such that for every bounded set $\mathcal{B} \in \mathcal{D}$ the component $S_1(t)\mathcal{B}$ converges uniformly to 0 as $t \rightarrow \infty$, and the component $S_2(t)\mathcal{B}$ is κ -contractive or is precompact in $\mathcal{D}$ for large t [47, 51].

As the first step towards proving the asymptotic compactness property stated above, existence of bounded absorbing sets for $\{S_w(t)\}_{t \in [0, \infty)}$ and $\{S_s(t)\}_{t \in [0, \infty)}$ is established in Theorems 6.2 and 6.3, respectively. However, it turns out that the κ -contracting property is hard to achieve for the model (4)–(6) with parameter values in the range given in Table 1, due to the lack of space-dissipative terms in the ODEs (4) and (5), the nature of nonlinear couplings in (4) and (5), and the range of values of the biophysical parameters of the model.

The uniform compactness of the component $S_2(t)$ in the decomposition approach stated above is usually verified by establishing energy estimates in more regular function spaces and then deducing compactness from compact embedding theorems. This approach, although successfully used in [39] to prove existence of a global attractor for a coupled ODE-PDE reaction-diffusion system, is not very promising here. In [39], the ODE subsystem is linear

and the energy estimates in a higher regular space are achieved by taking space derivatives of the ODEs and constructing energy functionals for the resulting equations. As seen in the proof of Theorem 6.2, the nonnegativity of $i(x, t)$ is a key property that permits elimination of the sign-indefinite quadratic term in the energy equation of (4), which results in the energy variation inequality (87). This nonnegativity property, however, is not preserved in the derivative or any other variations of $i(x, t)$, leaving some sign-indefinite quadratic terms in the analysis. Moreover, it can be observed from the range of parameter values given in Table 1 that the sign-indefinite nonlinear terms that would appear in the energy equations of any variations of (4) and (5) have significantly larger coefficients compared with the sign-definite dissipative terms. This makes it challenging to balance the terms in the energy functional in order to absorb the nondissipative terms into dissipative ones. Finally, the nonlinear terms appearing in (4) and (5) do not satisfy the usual assumptions, as in, e.g., [14, sect. 11.1], that enable shaping the energy functional to eliminate the nondissipative terms that would otherwise appear in the equations.

Some other techniques are available in the literature to avoid energy estimations in higher regular spaces. In [38], for instance, the notion of ω -limit compactness is used to develop necessary and sufficient conditions for existence of a global attractor. This is accomplished by decomposing the phase space into two spaces, one of which is finite-dimensional, and then showing that for every bounded set $\mathcal{B} \subset \mathcal{D}$ the canonical projection of $S(t)\mathcal{B}$ onto the finite-dimensional space is bounded, and the canonical projection on the complement space remains arbitrarily small for sufficiently large $t \geq t_0$ for some $t_0 = t_0(\mathcal{B}) > 0$. These decomposition techniques, however, rely on the spectral decomposition of the space-acting operators to construct the desired phase space decomposition. Such operators do not exist in the ODE subsystems (4) and (5) in our problem.

7.2. Nonexistence of a global attractor. As discussed in section 7.1, establishing a global attractor for (4)–(6) is a challenging problem. In fact, in this section we show that there exist sets of parameter values, leading to physiologically reasonable behavior in the model, for which the semigroups $\{S_w(t)\}_{t \in [0, \infty)}$ and $\{S_s(t)\}_{t \in [0, \infty)}$ do not possess a global attractor.

We first use [14, Prop. 11.11] to prove Theorem 7.5 below, which gives sufficient conditions for noncompactness of the equilibrium sets of (4)–(6) in $\mathcal{U}_w$ and $\mathcal{U}_s$. However, before embarking on the technical details of this theorem, we delineate the main idea using the following intuitive discussion.

Assume that the ODE components (4) and (5) are decoupled from the PDE component (6) by freezing $w(x, t)$ in space and time in (5). In this case, (4) and (5) can be viewed *pointwise* as an uncountable set of dynamical systems governed by ODEs that are enumerated by points $x \in \Omega$. To distinguish this pointwise view, let $(v_x(t), i_x(t))$ denote the solution of the dynamical system located at $x \in \Omega$, in contrast with $(v(x, t), i(x, t))$ that denotes the solution of the decoupled ODEs (4) and (5) defined over Ω . Note that the pointwise-defined dynamical systems are fully decoupled from each other, which means the solutions $(v_x(t), i_x(t))$ and $(v_y(t), i_y(t))$ evolve totally independently in time for every $x \neq y \in \Omega$.

Now, assume further that the decoupled ODE system (4) and (5) possesses more than one equilibrium, two of which are denoted by (v_e, i_e) and (v_0, i_0) . Then, all pointwise defined dynamical systems correspondingly possess more than one equilibrium, in particular $(v_{x_e}, i_{x_e}) = (v_e(x), i_e(x))$ and $(v_{x_0}, i_{x_0}) = (v_0(x), i_0(x))$ for the system located at x . This implies that

the solutions $(v_x(t), i_x(t))$ can converge independently to different values at different points $x \in \Omega$. Therefore, when composed together, they form a solution $(v(x, t), i(x, t))$ for the decoupled ODE system (4) and (5), which can possibly develop drastic discontinuities over Ω as it evolves in time. Note that such discontinuities in the solutions can occur even though the initial values are smooth. Moreover, it follows in particular that the ODE system (4) and (5) possesses an uncountable discrete equilibrium set. In fact, any function composed arbitrarily of either values (v_{x_e}, i_{x_e}) and (v_{x_0}, i_{x_0}) at each point $x \in \Omega$ would be an equilibrium.

The idea of Theorem 7.5 is to prove that the space-smoothing effect of the coupling with the PDE component (6) is not sufficiently strong to rule out the discontinuities of the above nature in (v, i) and, in particular, having a noncompact equilibrium set. Define the mappings

$$(104) \quad \begin{aligned} P_v(v, i) &:= v - J_1 i + J_2 v^T \Psi J_4 + J_3 v^T \Psi J_5, \\ P_i(v, i) &:= (e\Upsilon)^{-1} \Gamma i - N J_7 f(v) - g, \end{aligned}$$

and let (v_e, i_e, w_e) be an equilibrium of (4)–(6), that is, $P_v(v_e, i_e) = 0$ and $P_i(v_e, i_e) = J_6 w_e$. Assume that there exists $(v_0, i_0) \neq (v_e, i_e)$ such that $P_v(v_0, i_0) = 0$ and $P_i(v_0, i_0) = P_i(v_e, i_e)$. In this case, (v_e, i_e) and (v_0, i_0) are both equilibria of the system (4) and (5) if we assume that it is decoupled from (6) by freezing w at $w = w_e$. Therefore, motivated by the discussion above, we can construct a new equilibrium $(\bar{v}, \bar{i})$ for this decoupled system by letting $(\bar{v}, \bar{i}) = (v_0, i_0)$ over an arbitrary set Ω_0 , and $(\bar{v}, \bar{i}) = (v_e, i_e)$ over the complement set Ω_e . This construction is illustrated in Figure 4.

Since w is not actually frozen at $w = w_e$, the function $(\bar{v}, \bar{i})$ is not necessarily a component of a new equilibrium of the coupled system (4)–(6). However, if it is certain that w remains close to w_e , then we can expect that there exists a new equilibrium (v^*, i^*, w^*) of (4)–(6) whose component (v^*, i^*) is close to $(\bar{v}, \bar{i})$. Since the w -component of an equilibrium of (4)–(6) is continuous over Ω , we may postulate that, provided the sets Ω_0 are sufficiently small, updating (v_e, i_e) by $(\bar{v}, \bar{i})$ in the equilibrium equations would not greatly deviate the w -component from




Figure 4. Illustrative construction of new equilibria as given by Theorem 7.5. To avoid unnecessary complexity in the graph, only one representative curve out of the six curves in the (v, i) components of the solutions is shown.

w_e and the above expectation is satisfied. This postulation is indeed true, and it is proved in Theorem 7.5 that under certain conditions a new equilibrium (v^*, i^*, w^*) exists such that (v^*, i^*) are arbitrarily close to $(\bar{v}, \bar{i})$ provided Ω_0 is sufficiently small. The proof is relatively involved and constitutes the core part of the proof of Theorem 7.5. It relies strongly on the $\mathcal{L}_w^\infty$ -boundedness of the space-acting operator A^{-1} that appears in the equilibrium equations, and on assumption (iv) of Theorem 7.5. Figure 4 gives an illustration of the component (v^*, i^*) lying uniformly closer than ε to $(\bar{v}, \bar{i})$.

Finally, the noncompactness of the equilibrium set of (4)–(6) follows if we show that the existence of equilibria (v^*, i^*, w^*) is uniform with respect to the shape of the sets Ω_0 , that is, as long as only the size of Ω_0 is smaller than a uniform bound. In this case, we take ε small enough such that the distance between $(v_e(x), i_e(x))$ and $(v_0(x), i_0(x))$ is larger than 3ε . Then, for any two sufficiently small sets $\tilde{\Omega}_0$ and $\hat{\Omega}_0$, we can construct new equilibria as discussed above, having components closer than ε to their associated estimates $(\bar{v}, \bar{i})$. It can be observed from Figure 4 that the associated components (v^*, i^*) of these two equilibria would certainly be at a distance larger than ε from each other, at least on the difference between the two sets $\tilde{\Omega}_0$ and $\hat{\Omega}_0$. Therefore, since this construction is independent of the shape of the sets $\tilde{\Omega}_0$ and $\hat{\Omega}_0$ and we have uncountably different choices for these sets, it follows that we can construct an uncountable set of disjoint equilibria. This implies noncompactness of the equilibrium set of (4)–(6). Theorem 7.5 below gives rigorous arguments for the above discussion.

Theorem 7.5 (Noncompactness of equilibrium sets). *Suppose g is bounded and constant in time, that is, $g(x, t) = g(x)$ for all $(x, t) \in \Omega \times [0, \infty)$ and $g \in \mathcal{L}_i^\infty$. Let $u_e := (v_e, i_e, 0, w_e, 0)$ be an equilibrium of (4)–(6) such that $v_e \in \mathcal{L}_v^\infty$, $i_e \in \mathcal{L}_i^\infty$, and $w_e \in \mathcal{H}_w^2$. Define the mapping $P = (P_v, P_i) : \mathcal{L}_v^\infty \times \mathcal{L}_i^\infty \rightarrow \mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$ as in (104) and let $A := -\frac{3}{2}\Delta + \Lambda^2 I$. Assume that the following conditions hold.*

- (i) Λ_{EE} and Λ_{EI} take the same values, that is, $\Lambda = \Lambda_{EE}I_{2 \times 2} = \Lambda_{EI}I_{2 \times 2}$.
- (ii) There exists $(v_0, i_0) \in \mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$ such that

$$\operatorname{ess\,inf}_{x \in \Omega} \|(v_e(x), i_e(x)) - (v_0(x), i_0(x))\|_\infty > 0$$

and

$$(105) \quad P_v(v_0, i_0) = 0, \quad P_i(v_0, i_0) = P_i(v_e, i_e).$$

- (iii) $\partial_{(v,i)} P(v_e, i_e)$ and $\partial_{(v,i)} P(v_0, i_0)$ are nonsingular almost everywhere in Ω .
- (iv) There exists $\alpha > 0$ such that, for every $b = (b_v, b_i) \in \mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$, the system of equations

$$(106) \quad \partial_{(v,i)} P_v(v_e, i_e) \phi = b_v,$$

$$\partial_{(v,i)} P_i(v_e, i_e) \phi - J_6 A^{-1} \Lambda^2 M J_8 \partial_v f(v_e) \phi_v = b_i$$

has a unique solution $\phi = (\phi_v, \phi_i) \in \mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$ that satisfies

$$(107) \quad \|\phi\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} \leq \alpha \|b\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty}.$$

Then, for a measurable partition $\Omega = \Omega_e \cup \Omega_0$ and

$$(108) \quad \bar{v} := v_e \chi_{\Omega_e} + v_0 \chi_{\Omega_0}, \quad \bar{i} := i_e \chi_{\Omega_e} + i_0 \chi_{\Omega_0},$$

the following assertions hold.

- (I) For every $\varepsilon > 0$ there exists $\delta > 0$ and an equilibrium $u^* := (v^*, i^*, 0, w^*, 0)$ of (4)–(6) such that

$$\|(v^*, i^*) - (\bar{v}, \bar{i})\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} \leq \varepsilon \quad \text{whenever } |\Omega_0| \leq \delta.$$

- (II) The equilibrium sets of (4)–(6) are noncompact in $\mathcal{U}_s$ and $\mathcal{U}_w$.

Proof. The proof is organized in three steps.

Step 1. We show that there exists $\bar{\alpha} > 0$ such that, for every $b = (b_v, b_i) \in \mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$, the system of equations

$$(109) \quad \begin{aligned} \partial_{(v,i)} P_v(\bar{v}, \bar{i}) \phi &= b_v, \\ \partial_{(v,i)} P_i(\bar{v}, \bar{i}) \phi - J_6 A^{-1} \Lambda^2 M J_8 \partial_v f(\bar{v}) \phi_v &= b_i \end{aligned}$$

has a unique solution $\phi \in \mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$ that satisfies $\|\phi\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} \leq \bar{\alpha} \|b\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty}$. This provides the required conditions of the implicit function theorem that is used in Step 2 to prove the existence of the equilibrium u^* . The proof proceeds by iteratively constructing a solution by starting from the solution of (106) and applying certain corrections at each iteration.

Let $\phi^{(0)} = (\phi_v^{(0)}, \phi_i^{(0)})$ be the solution of (106) for a given $b \in \mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$ and construct an approximate solution for (109) of the form $\phi^{(1)} := \phi^{(0)} + \phi_r^{(1)}$, where $\phi_r^{(1)} = (\phi_{r_v}^{(1)}, \phi_{r_i}^{(1)})$ is the unique solution of

$$(110) \quad \partial_{(v,i)} P(v_0, i_0) \phi_r^{(1)} = (\partial_{(v,i)} P(v_e, i_e) - \partial_{(v,i)} P(v_0, i_0)) \phi^{(0)} \chi_{\Omega_0}.$$

Note that by assumption (iii) the unique solution $\phi_r^{(1)}$ exists and belongs to $\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$. The approximate solution $\phi^{(1)}$ solves

$$\begin{aligned} \partial_{(v,i)} P_v(\bar{v}, \bar{i}) \phi^{(1)} &= b_v, \\ \partial_{(v,i)} P_i(\bar{v}, \bar{i}) \phi^{(1)} - J_6 A^{-1} \Lambda^2 M J_8 \partial_v f(\bar{v}) \phi_v^{(1)} &= b_i + b_{r_i}^{(1)}, \end{aligned}$$

where $b_r^{(1)} = (0, b_{r_i}^{(1)})$,

$$(111) \quad b_{r_i}^{(1)} := J_6 A^{-1} \Lambda^2 M J_8 \left[(\partial_v f(v_e) - \partial_v f(v_0)) \phi_v^{(0)} - \partial_v f(v_0) \phi_{r_v}^{(1)} \right] \chi_{\Omega_0}$$

is the remainder resulting from the approximation error in $\phi^{(1)}$.

Now, note that by assumption (iv) there exists $\alpha_0 := \alpha > 0$ such that

$$(112) \quad \|\phi^{(0)}\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} \leq \alpha_0 \|b\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty}.$$

Moreover, since by assumption (ii) we have $(v_0, i_0) \in \mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$, it is immediate from the definition of P_v and P_i , given by (104), that $\partial_{(v,i)} P(v_0, i_0)$ is bounded. This, along with assumption (iii) and (112), implies that the solution $\phi_r^{(1)}$ of (110) satisfies

$$(113) \quad \|\phi_r^{(1)}\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} \leq \zeta_1 \|\phi^{(0)}\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} \leq \alpha_1 \|b\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty}$$

for some $\zeta_1, \alpha_1 > 0$.

Next, note that since $A^{-1} : \mathcal{L}_w^2 \rightarrow \mathcal{H}_w^2$ is a bounded operator and f is smooth, the definition of $b_{r_i}^{(1)}$, given by (111), implies that $b_{r_i}^{(1)} \in H_{\text{per}}^2(\Omega; \mathbb{R}^4)$. Moreover, it further implies by the Sobolev embedding theorems [9, Thm. 6.6-1] that $b_{r_i}^{(1)} \in C_{\text{per}}^{0,\lambda}(\overline{\Omega}, \mathbb{R}^4)$ for all $\lambda \in (0, 1)$ and, in particular, $\|b_{r_i}^{(1)}\|_{\mathcal{L}_i^\infty} \leq \zeta_2 \|b_{r_i}^{(1)}\|_{H_{\text{per}}^2(\Omega; \mathbb{R}^4)}$ for some $\zeta_2 > 0$. Therefore, using (112) and (113), there exist $\zeta_3, \zeta_4, \zeta_5, \beta_1 > 0$ such that

$$(114) \quad \begin{aligned} \|b_r^{(1)}\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} &\leq \zeta_2 \|b_{r_i}^{(1)}\|_{H_{\text{per}}^2(\Omega; \mathbb{R}^4)} \leq \zeta_3 \left(\|\phi_v^{(0)}\|_{\mathcal{L}_v^2} + \|\phi_{r_v}^{(1)}\|_{\mathcal{L}_v^2} \right) \\ &\leq \zeta_4 \|\phi^{(0)}\|_{\mathcal{L}_v^2 \times \mathcal{L}_i^2} \leq \zeta_5 |\Omega_0|^{\frac{1}{2}} \|\phi^{(0)}\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} \\ &\leq \beta_1 |\Omega_0|^{\frac{1}{2}} \|b\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty}. \end{aligned}$$

Now, for $m = 2, 3, \dots$, let $\phi^{(m)} := \phi^{(m-1)} + \phi_r^{(m)}$, where $\phi_r^{(m)}$ is the unique solution of

$$\partial_{(v,i)} P(v_0, i_0) \phi_r^{(m)} = -b_r^{(m-1)} \chi_{\Omega_0}.$$

It follows immediately that, for some $\eta > 0$,

$$(115) \quad \|\phi_r^{(m)}\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} \leq \eta \|b_r^{(m-1)}\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty}, \quad m = 2, 3, \dots$$

Moreover, $\phi_r^{(m)}$ solves the system of equations

$$\begin{aligned} \partial_{(v,i)} P_v(\bar{v}, \bar{i}) \phi^{(m)} &= b_v, \\ \partial_{(v,i)} P_i(\bar{v}, \bar{i}) \phi^{(m)} - J_6 A^{-1} \Lambda^2 M J_8 \partial_v f(\bar{v}) \phi_v^{(m)} &= b_i + b_{r_i}^{(m)}, \end{aligned}$$

where

$$b_{r_i}^{(m)} := -J_6 A^{-1} \Lambda^2 M J_8 \partial_v f(v_0) \phi_{r_v}^{(m)} \chi_{\Omega_0}, \quad m = 2, 3, \dots$$

Using the Sobolev embedding theorems and (115), the remainder $b_r^{(m)} = (0, b_{r_i}^{(m)})$ satisfies, for some $\zeta_6, \zeta_7, \zeta_8, \beta > 0$,

$$\begin{aligned} \|b_r^{(m)}\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} &\leq \zeta_6 \|b_{r_i}^{(m)}\|_{H_{\text{per}}^2(\Omega; \mathbb{R}^4)} \leq \zeta_7 \|\phi_r^{(m)}\|_{\mathcal{L}_v^2 \times \mathcal{L}_i^2} \leq \zeta_8 |\Omega_0|^{\frac{1}{2}} \|\phi_r^{(m)}\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} \\ &\leq \beta |\Omega_0|^{\frac{1}{2}} \|b_r^{(m-1)}\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty}, \quad m = 2, 3, \dots, \end{aligned}$$

which, letting $\kappa := \beta |\Omega_0|^{\frac{1}{2}}$ and recalling (114), implies

$$(116) \quad \|b_r^{(m)}\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} \leq \beta_1 |\Omega_0|^{\frac{1}{2}} \kappa^{(m-1)} \|b\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty}, \quad m = 2, 3, \dots$$

Now, let $|\Omega_0| < \bar{\delta}$, $\bar{\delta} > 0$, and choose $\bar{\delta}$ such that $\kappa < 1$. Note that β , and consequently the choice of $\bar{\delta}$ and the value of κ , do not depend on b and the specific form of the partition $\Omega = \Omega_e \cup \Omega_0$. Therefore, it follows that $\|b_r^{(m)}\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} \rightarrow 0$ as $m \rightarrow \infty$, and hence $\phi^{(m)}$ converges to a solution ϕ for (109) when $|\Omega_0| < \bar{\delta}$. Moreover, (112)–(116) imply

$$\begin{aligned} \|\phi^{(m)}\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} &\leq \|\phi^{(0)}\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} + \|\phi_r^{(1)}\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} + \sum_{l=2}^m \|\phi_r^{(l)}\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} \\ &\leq \left[\alpha_0 + \alpha_1 + \eta\beta_1 |\Omega_0|^{\frac{1}{2}} \sum_{l=2}^m \kappa^{(l-2)} \right] \|b\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty}, \end{aligned}$$

and hence, taking the limit as $m \rightarrow \infty$, there exists $\bar{\alpha} > 0$, independent of the form of the partition, such that

$$(117) \quad \|\phi\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} \leq \bar{\alpha} \|b\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty}.$$

To prove that the solution constructed above for (109) is unique, first note that by assumption (i) the operator A becomes a scalar operator given by $A = (-\frac{3}{2}\Delta + \Lambda_{EE}^2 I)$. Then, considering the structure of the matrix parameters given by (8) and reinspect the expanded form (2), the system of equations (109) can be transformed to a system composed of five algebraic equations and one PDE by pre-multiplying the second equation in (109) by the elementary matrix

$$\left[\begin{array}{cc|c} 1 & 0 & \\ -\frac{M_{EI}}{M_{EE}} & 1 & 0_{2 \times 2} \\ \hline 0_{2 \times 2} & & I_{2 \times 2} \end{array} \right].$$

This follows from the fact that the scalar operator $(-\frac{3}{2}\Delta + \Lambda_{EE}^2 I)^{-1}$ acts only on one of the unknowns, namely, ϕ_{v_E} . Now, since $\partial_{(v,i)} P(\bar{v}, \bar{i})$ is nonsingular by assumption (iii), the five unknowns $\phi_i = (\phi_{i_{EE}}, \phi_{i_{EI}}, \phi_{i_{iE}}, \phi_{i_{iI}})$ and ϕ_{v_i} can be uniquely determined in terms of ϕ_{v_E} by elementary algebraic operations. Consequently, (109) is reduced to a scalar PDE of the form

$$p(\bar{v}, \bar{i}) \phi_{v_E} - \left(-\frac{3}{2}\Delta + \Lambda_{EE}^2 I \right)^{-1} \Lambda_{EE}^2 M_{EE} \partial_{v_E} f(\bar{v}_E) \phi_{v_E} = \hat{h},$$

where $\hat{h} \in L_{\text{per}}^\infty(\Omega, \mathbb{R})$ is given by the same elementary operations on b , and $p(\bar{v}, \bar{i})$ is nonzero almost everywhere in Ω , since elementary operations do not disrupt the nonsingularity of $\partial_{(v,i)} P(\bar{v}, \bar{i})$.

Next, dividing by $p(\bar{v}, \bar{i})$, the above equation can be written as

$$(118) \quad (I - K) \phi_{v_E} = h,$$

where $K := p(\bar{v}, \bar{i})^{-1} \Lambda_{EE}^2 M_{EE} \partial_{v_E} f(\bar{v}_E) (-\frac{3}{2}\Delta + \Lambda_{EE}^2 I)^{-1}$ and $h := p(\bar{v}, \bar{i})^{-1} \hat{h}$. The operator $K : L_{\text{per}}^2(\Omega, \mathbb{R}) \rightarrow L_{\text{per}}^2(\Omega, \mathbb{R})$ is linear, self-adjoint, and compact by the Rellich–Kondrachov compact embedding theorems [9, Thm. 6.6-3]. The existence of solutions of (109) proved above guarantees the existence of a solution $\phi_{v_E} \in L_{\text{per}}^\infty(\Omega, \mathbb{R})$ for every $h \in L_{\text{per}}^\infty(\Omega, \mathbb{R})$, which implies $L_{\text{per}}^\infty(\Omega, \mathbb{R}) \subset \text{Range}(I - K)$. However, $\text{Range}(I - K) = \text{Kernel}(I - K^*)^\perp = \text{Kernel}(I - K)^\perp$ by the Fredholm alternative [16, Thm. 5, App. D], and hence $L_{\text{per}}^\infty(\Omega, \mathbb{R}) \cap \text{Kernel}(I - K) = \{0\}$. This proves the uniqueness of bounded solutions of (118), and consequently the uniqueness of solutions of (109) for every $b = (b_v, b_i) \in \mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$.

Step 2. We prove assertion (I) using the implicit function theorem. Note that since $u_e := (v_e, i_e, 0, w_e, 0)$ is an equilibrium of (4)–(6), we have

$$(119) \quad P_v(v_e, i_e) = 0, \quad P_i(v_e, i_e) = J_6 w_e, \quad w_e = A^{-1} \Lambda^2 M J_8 f(v_e).$$

We seek an equilibrium $u^* := (v^*, i^*, 0, w^*, 0)$ such that

$$v^* = \bar{v} + \phi_v, \quad i^* = \bar{i} + \phi_i,$$

where $\phi := (\phi_v, \phi_i) \in \mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$ is a small corrector function that satisfies

$$(120) \quad P_v(v^*, i^*) = 0, \quad P_i(v^*, i^*) = J_6 w^*, \quad w^* = A^{-1} \Lambda^2 M J_8 f(v^*).$$

Note that (105), (108), and (119) imply

$$P_v(\bar{v}, \bar{i}) = 0, \quad P_i(\bar{v}, \bar{i}) = J_6 w_e, \quad v_e = \bar{v} - (v_0 - v_e) \chi_{\Omega_0}.$$

Therefore, the system of equations (120) is equivalent to

$$(121) \quad \begin{aligned} P_v(\bar{v} + \phi_v, \bar{i} + \phi_i) - P_v(\bar{v}, \bar{i}) &= 0, \\ P_i(\bar{v} + \phi_v, \bar{i} + \phi_i) - P_i(\bar{v}, \bar{i}) &= J_6 A^{-1} \Lambda^2 M J_8 (f(\bar{v} + \phi_v) - f(\bar{v} - (v_0 - v_e) \chi_{\Omega_0})), \end{aligned}$$

which, by the implicit function theorem [9, Thm. 7.13-1], has a unique solution $\phi \in \mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$, since (109) has a unique solution in $\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$ for every $b \in \mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$, as proved in Step 1. Moreover, it is immediate from the definition of the Fréchet derivative of the mappings P_i and P_v that the solution of (121) is arbitrarily close to the solution of (109) with

$$b := (0, J_6 A^{-1} \Lambda^2 M J_8 \partial_v f(\bar{v})(v_0 - v_e)) \chi_{\Omega_0},$$

provided these solutions are sufficiently small. This is ensured by (117) for small $|\Omega_0|$, since $\|b\|_{\mathcal{L}_v^\infty \times \mathcal{L}_i^\infty} \leq \zeta |\Omega_0|^{\frac{1}{2}}$ for some $\zeta > 0$. Therefore, it follows that assertion (I) holds for some $\delta = \delta(\varepsilon) \leq \bar{\delta}$.

Step 3. We prove assertion (II) using the fact that $\delta = \delta(\varepsilon) > 0$ in assertion (I) is independent of the specific form of the partition $\Omega = \Omega_e \cup \Omega_0$. Figure 4 can be used to visualize the arguments of the proof.

Let

$$(122) \quad \varepsilon := \frac{1}{3} \operatorname{ess\,inf}_{x \in \Omega} \|(v_e(x), i_e(x)) - (v_0(x), i_0(x))\|_\infty > 0$$

in assertion (I), and let $\delta = \delta(\varepsilon) > 0$ be the corresponding bound on the size of the partitions that satisfies the result of assertion (I). Note that $\varepsilon > 0$ by assumption (ii). Moreover, let $\mathcal{M}(\Omega)$ denote the set of all measurable subsets of Ω and define

$$\mathcal{P}_\delta(\Omega) := \{(\Omega_e, \Omega_0) \in \mathcal{M}(\Omega) \times \mathcal{M}(\Omega) : \Omega_e = \Omega \setminus \Omega_0, |\Omega_0| \leq \delta\}.$$

Let $\Theta_\delta(\Omega) \subset \mathcal{P}_\delta(\Omega)$ such that, for every $\tilde{\theta} = (\tilde{\Omega}_e, \tilde{\Omega}_0) \in \Theta_\delta(\Omega)$ and $\hat{\theta} = (\hat{\Omega}_e, \hat{\Omega}_0) \in \Theta_\delta(\Omega)$, we have $|\hat{\Omega}_0 \triangle \tilde{\Omega}_0| > \frac{1}{2}\delta$. Note that $\Theta_\delta(\Omega)$ is an uncountable set that can be viewed as an index

set enumerating all measurable partitions $\Omega = \Omega_e \cup \Omega_0$, $|\Omega_0| \leq \delta$, which are distinct in the sense of measure by a factor of at least $\frac{1}{2}\delta$.

Now, it follows from assertion (I) that, for every $\tilde{\theta} \neq \hat{\theta} \in \Theta_\delta(\Omega)$, there exist equilibria $u_{\tilde{\theta}} := (v_{\tilde{\theta}}, i_{\tilde{\theta}}, 0, w_{\tilde{\theta}}, 0)$ and $u_{\hat{\theta}} := (v_{\hat{\theta}}, i_{\hat{\theta}}, 0, w_{\hat{\theta}}, 0)$ such that

$$\begin{aligned} \operatorname{ess\,sup}_{x \in (\tilde{\Omega}_e \cap \hat{\Omega}_0)} \|(v_{\tilde{\theta}}(x), i_{\tilde{\theta}}(x)) - (v_0(x), i_0(x))\|_\infty &\leq \varepsilon, \\ \operatorname{ess\,sup}_{x \in (\tilde{\Omega}_0 \cap \hat{\Omega}_e)} \|(v_{\tilde{\theta}}(x), i_{\tilde{\theta}}(x)) - (v_e(x), i_e(x))\|_\infty &\leq \varepsilon, \\ \operatorname{ess\,sup}_{x \in (\tilde{\Omega}_e \cap \hat{\Omega}_0)} \|(v_{\hat{\theta}}(x), i_{\hat{\theta}}(x)) - (v_e(x), i_e(x))\|_\infty &\leq \varepsilon, \\ \operatorname{ess\,sup}_{x \in (\tilde{\Omega}_0 \cap \hat{\Omega}_e)} \|(v_{\hat{\theta}}(x), i_{\hat{\theta}}(x)) - (v_0(x), i_0(x))\|_\infty &\leq \varepsilon. \end{aligned}$$

Therefore, noting that $\tilde{\Omega}_0 \triangle \hat{\Omega}_0 = (\tilde{\Omega}_0 \cap \hat{\Omega}_e) \cup (\tilde{\Omega}_e \cap \hat{\Omega}_0)$ and recalling the definition of ε given by (122),

$$\operatorname{ess\,inf}_{x \in (\tilde{\Omega}_0 \triangle \hat{\Omega}_0)} \|(v_{\tilde{\theta}}(x), i_{\tilde{\theta}}(x)) - (v_{\hat{\theta}}(x), i_{\hat{\theta}}(x))\|_\infty \geq \varepsilon,$$

which further implies

$$\|(v_{\tilde{\theta}}, i_{\tilde{\theta}}) - (v_{\hat{\theta}}, i_{\hat{\theta}})\|_{\mathcal{L}_v^2 \times \mathcal{L}_i^2} \geq |\tilde{\Omega}_0 \triangle \hat{\Omega}_0|^{\frac{1}{2}} \operatorname{ess\,inf}_{x \in (\tilde{\Omega}_0 \triangle \hat{\Omega}_0)} \|(v_{\tilde{\theta}}(x), i_{\tilde{\theta}}(x)) - (v_{\hat{\theta}}(x), i_{\hat{\theta}}(x))\|_\infty > \left(\frac{1}{2}\delta\right)^{\frac{1}{2}} \varepsilon.$$

Since $\tilde{\theta}$ and $\hat{\theta}$ are arbitrary, it follows that the set $\mathcal{E} := \{u_\theta\}_{\theta \in \Theta_\delta(\Omega)}$ composed of the equilibria u_θ constructed as above is an uncountable discrete subset of the equilibrium sets of (4)–(6) in $\mathcal{U}_s$ and $\mathcal{U}_w$. This completes the proof. $\blacksquare$

Remark 7.6 (Alternative assumptions for Theorem 7.5). According to the proof of Theorem 7.5, some of the assumptions of this theorem can be relaxed or replaced by alternative assumptions as follows.

- Assumption (i) is used to prove the uniqueness of solutions of (109). Without this assumption, the operator A is not a scalar operator and (109) cannot be reduced to a scalar PDE using elementary algebraic operations. The operator K representing the system of PDEs in this case would not be self-adjoint, and hence application of the Fredholm alternative would not immediately imply uniqueness of the solutions. However, an alternative assumption to assumption (i) can be made on the adjoint of the operator K , so that the uniqueness of the solutions of (109) is still ensured using the Fredholm alternative. We avoid this complication, since the fiber decay scale constants Λ_{EE} and Λ_{EI} are always assumed to be equal in the practical applications of the model [3].
- In assumption (ii), it suffices to have $\operatorname{ess\,inf}_{x \in \mathcal{X}} \|(v_e(x), i_e(x)) - (v_0(x), i_0(x))\|_\infty > 0$, where $\mathcal{X}$ is any measurable subset of Ω with positive measure. Correspondingly, it suffices that the nonsingularity in assumption (iii) holds almost everywhere on an open

subset $\mathcal{Y} \supset \mathcal{X}$ of Ω . In this case, the proof is modified by restricting $\mathcal{P}_\delta(\Omega)$ to its subset consisting of partitions with $\Omega_0 \subset \mathcal{X}$. The index set $\Theta_\delta(\Omega)$ remains uncountable, and the noncompactness result of the theorem holds with no change.

Remark 7.7 (Nonexistence of a global attractor). Suppose that the assumptions of Theorem 7.5 hold for an input g and an equilibrium u_e that further satisfy $i_e, w_e > \varepsilon_1, \varepsilon_1 > 0$, almost everywhere in Ω and $g \in \mathcal{D}_g$, where $\mathcal{D}_g$ is given by (77). Note that u_e then belongs to $\mathcal{D}_s$. Then, the equation $P_i(v_e, i_e) = J_6 w_e$ in the equilibrium equations (119) implies that $P_i(v_e, i_e) \geq 0$, and hence $P_i(v_0, i_0) \geq 0$ in (105). Therefore, it follows from the definition of P_i given by (104) that every solution i_0 of (105) satisfies $i_0 > \varepsilon_2, \varepsilon_2 > 0$, almost everywhere in Ω . Then, by definition of $(\bar{v}, \bar{i})$, given by (108), all equilibria u^* constructed by assertion (I) of Theorem 7.5 satisfy $i^* > 0$ almost everywhere in Ω when δ is sufficiently small. Also, the equilibrium equations $w_e = A^{-1}\Lambda^2 M J_8 f(v_e)$ and $w^* = A^{-1}\Lambda^2 M J_8 f(v^*)$ imply that

$$\|w^* - w_e\|_{\mathcal{L}_w^\infty} \leq \beta_1 \|w^* - w_e\|_{\mathcal{H}_w^2} \leq \beta_2 \|v^* - v_e\|_{\mathcal{L}_v^2}$$

for some $\beta_1, \beta_2 > 0$, and hence $w^* > 0$ almost everywhere in Ω , when δ is sufficiently small. Therefore, assertion (II) of Theorem 7.5 ensures existence of a biophysically plausible noncompact set of equilibria $\mathcal{E} \subset \mathcal{D}_s \subset \mathcal{D}_w$. This, in particular, implies that in the case where the assumptions of Theorem 7.5 are satisfied for some u_e and g as given above, the semigroups $\{S_w(t) : \mathcal{D}_w \rightarrow \mathcal{D}_w\}_{t \in [0, \infty)}$ and $\{S_s(t) : \mathcal{D}_s \rightarrow \mathcal{D}_s\}_{t \in [0, \infty)}$ are not asymptotically compact, and hence they do not possess a global attractor.

The assumptions of Theorem 7.5 are relatively straightforward to check for the space-homogeneous equilibria of (4)–(6). Consider the set of values given in Table 2 for the parameters of the model, which are suggested in [3, Table VI, col. 2] as a set of parameter values leading to physiologically reasonable behavior of the model. The parameters $\bar{g}_{EE}$, $\bar{g}_{EI}$, $\bar{g}_{IE}$, and $\bar{g}_{II}$ are the mean values of the physiologically shaped random signals used in [3] as the subcortical inputs g_{EE} , g_{EI} , g_{IE} , and g_{II} , respectively. Here, we set $g(t, x) = (\bar{g}_{EE}, \bar{g}_{EI}, \bar{g}_{IE}, \bar{g}_{II})$ for all x and t , and check the assumptions of Theorem 7.5 for a space-homogeneous equilibrium of (4)–(6).

Table 2

A set of biophysically plausible parameter values for the model (2) for which Theorem 7.5 implies nonexistence of a global attractor [3, Table VI, col. 2]. The parameters $\bar{g}_{EE}$, $\bar{g}_{EI}$, $\bar{g}_{IE}$, and $\bar{g}_{II}$ are, respectively, the mean values of the physiologically shaped random inputs g_{EE} , g_{EI} , g_{IE} , and g_{II} used in [3].

Parameter	τ_E	τ_I	V_{EE}	V_{EI}	V_{IE}	V_{II}	γ_{EE}	γ_{EI}
Value	11.787×10^{-3}	138.25×10^{-3}	61.264	51.703	-7.127	-12.679	816.04	261.29
Parameter	γ_{IE}	γ_{II}	Υ_{EE}	Υ_{EI}	Υ_{IE}	Υ_{II}	N_{EE}	N_{EI}
Value	219.09	40.575	0.92695	1.3012	0.19053	0.94921	3893.0	3326.8
Parameter	N_{IE}	N_{II}	ν	$\Lambda_{EE}, \Lambda_{EI}$	M_{EE}	M_{EI}	F_E	F_I
Value	839.39	682.41	101.78	0.96545	4013.5	1544.3	266.44	300.65
Parameter	μ_E	μ_I	σ_E	σ_I	$\bar{g}_{EE}$	$\bar{g}_{EI}$	$\bar{g}_{IE}$	$\bar{g}_{II}$
Value	30.628	19.383	5.6536	3.3140	83.190	6407.5	0	0

Assumption (i) holds with $\Lambda_{\text{EE}} = \Lambda_{\text{EE}} = 0.96545$, as given in Table 2. Solving the equations $P_v(v_e, i_e) = 0$, $P_i(v_e, i_e) = J_6 w_e$, and $w_e = M J_8 f(v_e)$, a space-homogeneous equilibrium is calculated as

$$v_e = (1.9629, 6.5150), \quad i_e = (5.2552, 100.2372, 2.4493, 53.5665), \quad w_e = (821.7136, 316.1760).$$

Note that the numbers given here should actually be regarded as constant functions over Ω . Assumption (ii) then holds by finding a solution $(v_0, i_0) \neq (v_e, i_e)$ for (105) as

$$v_0 = (10.9417, 7.7148), \quad i_0 = (25.9005, 177.5837, 4.0757, 89.1352).$$

Assumption (iii) also holds with the following nonsingular matrix-valued functions:

$$\begin{aligned} \partial_{(v,i)} P(v_e, i_e) &= \begin{bmatrix} 1.4294 & 0 & -0.9680 & 0 & 1.2754 & 0 \\ 0 & 7.1635 & 0 & -0.8740 & 0 & 1.5138 \\ -199.2222 & 0 & 323.8625 & 0 & 0 & 0 \\ -170.2472 & 0 & 0 & 73.8727 & 0 & 0 \\ 0 & -440.3409 & 0 & 0 & 423.0237 & 0 \\ 0 & -357.9898 & 0 & 0 & 0 & 15.7254 \end{bmatrix}, \\ \partial_{(v,i)} P(v_0, i_0) &= \begin{bmatrix} 1.9946 & 0 & -0.8214 & 0 & 2.5352 & 0 \\ 0 & 11.4648 & 0 & -0.8508 & 0 & 1.6085 \\ -1858.395 & 0 & 323.8625 & 0 & 0 & 0 \\ -1588.109 & 0 & 0 & 73.8727 & 0 & 0 \\ 0 & -730.7260 & 0 & 0 & 423.0237 & 0 \\ 0 & -594.0680 & 0 & 0 & 0 & 15.7254 \end{bmatrix}. \end{aligned}$$

To check assumption (iv), note that, for every $b = (b_v, b_i) \in \mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$, elementary algebraic operations reduce (106) to

$$(123) \quad \begin{aligned} \phi_{v_e} &= 0.6287\phi_{i_{\text{EE}}} + h_{v_e}, & \phi_{v_i} &= 0.0521\phi_{i_{\text{EE}}} + h_{v_i}, \\ \phi_{i_{\text{EI}}} &= 2.4834\phi_{i_{\text{EE}}} + h_{i_{\text{EI}}}, & \phi_{i_{\text{IE}}} &= 0.0543\phi_{i_{\text{EE}}} + h_{i_{\text{IE}}}, & \phi_{i_{\text{II}}} &= 1.1870\phi_{i_{\text{EE}}} + h_{i_{\text{II}}} \end{aligned}$$

and the scalar PDE

$$(124) \quad (I - D)\phi_{i_{\text{EE}}} = h_{i_{\text{EE}}}, \quad D := 0.6060 \left(-\frac{3}{2}\Delta + 0.96545^2 I \right)^{-1},$$

where $h = (h_v, h_i) \in \mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$ is the result of the same algebraic operations on b . Now, note that since $-\Delta$ is a nonnegative operator in $H_{\text{per}}^2(\Omega; \mathbb{R})$, it follows from the spectral theory of bounded linear self-adjoint operators [16, App. D.6] that the spectrum of the operator $(I - D) : L_{\text{per}}^2(\Omega; \mathbb{R}) \rightarrow L_{\text{per}}^2(\Omega; \mathbb{R})$ lies entirely above $1 - 0.6060 \times 0.96545^{-2} = 0.3498 > 0$. Therefore, the PDE (124) has a unique solution $\phi_{i_{\text{EE}}} \in L_{\text{per}}^2(\Omega; \mathbb{R})$ for every $h_{i_{\text{EE}}} \in L_{\text{per}}^2(\Omega; \mathbb{R}) \supset L_{\text{per}}^\infty(\Omega; \mathbb{R})$, and hence it follows from (123) that (106) has a unique solution $\phi = (\phi_v, \phi_i) \in \mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$ for every $b \in \mathcal{L}_v^\infty \times \mathcal{L}_i^\infty$.

It remains to check (107). Using the spectral theory of bounded linear self-adjoint operators and the Cauchy–Schwarz inequality, we can write

$$\begin{aligned}\|\phi_{i_{EE}}\|_{L^2_{\text{per}}(\Omega;\mathbb{R})}^2 &\leq \frac{1}{0.3498}((I-D)\phi_{i_{EE}}, \phi_{i_{EE}})_{L^2_{\text{per}}(\Omega;\mathbb{R})} = \frac{1}{0.3498}(h_{i_{EE}}, \phi_{i_{EE}})_{L^2_{\text{per}}(\Omega;\mathbb{R})} \\ &\leq \frac{1}{0.3498}\|h_{i_{EE}}\|_{L^2_{\text{per}}(\Omega;\mathbb{R})}\|\phi_{i_{EE}}\|_{L^2_{\text{per}}(\Omega;\mathbb{R})}.\end{aligned}$$

Therefore, there exists $\alpha_1 = \frac{1}{0.3498} > 0$ such that

$$\|\phi_{i_{EE}}\|_{L^2_{\text{per}}(\Omega;\mathbb{R})} \leq \alpha_1 \|h_{i_{EE}}\|_{L^2_{\text{per}}(\Omega;\mathbb{R})}.$$

Now, using (124) and the Sobolev embedding theorems, we can write, for some $\alpha_2, \alpha_3 > 0$,

$$\begin{aligned}\|\phi_{i_{EE}}\|_{L^{\infty}_{\text{per}}(\Omega;\mathbb{R})} &\leq \|h_{i_{EE}}\|_{L^{\infty}_{\text{per}}(\Omega;\mathbb{R})} + \|D\phi_{i_{EE}}\|_{L^{\infty}_{\text{per}}(\Omega;\mathbb{R})} \leq \|h_{i_{EE}}\|_{L^{\infty}_{\text{per}}(\Omega;\mathbb{R})} + \alpha_2 \|D\phi_{i_{EE}}\|_{H^2_{\text{per}}(\Omega;\mathbb{R})} \\ &\leq \|h_{i_{EE}}\|_{L^{\infty}_{\text{per}}(\Omega;\mathbb{R})} + \alpha_3 \|\phi_{i_{EE}}\|_{L^2_{\text{per}}(\Omega;\mathbb{R})} \leq \|h_{i_{EE}}\|_{L^{\infty}_{\text{per}}(\Omega;\mathbb{R})} + \alpha_1 \alpha_3 \|h_{i_{EE}}\|_{L^2_{\text{per}}(\Omega;\mathbb{R})} \\ &\leq \left(1 + \alpha_1 \alpha_3 |\Omega|^{\frac{1}{2}}\right) \|h_{i_{EE}}\|_{L^{\infty}_{\text{per}}(\Omega;\mathbb{R})},\end{aligned}$$

which, along with the algebraic equalities (123), implies (107). Hence, assumption (iv) holds.

It is now implied by Theorem 7.5 that the equilibrium sets of (4)–(6) are noncompact in $\mathcal{U}_s$ and $\mathcal{U}_w$. Moreover, it follows immediately from the equilibrium equations (119) and the definition of P_i given by (104) that, in general, all space-homogeneous equilibria i_e and w_e are positive and, in particular, belong to $\mathcal{D}_{\text{Bio}} \cap \mathcal{D}_s$. Therefore, by Remark 7.7, the semigroups $\{S_w(t) : \mathcal{D}_w \rightarrow \mathcal{D}_w\}_{t \in [0, \infty)}$ and $\{S_s(t) : \mathcal{D}_s \rightarrow \mathcal{D}_s\}_{t \in [0, \infty)}$ associated with (4)–(6) with parameter values given by Table 2 do not possess a global attractor.

It can be shown by similar calculations as above that the assumptions of Theorem 7.5 are satisfied by space-homogeneous equilibria of the model for 3 other sets of parameter values out the 24 sets available in [3, Tables V and VI], namely, the sets given in [3, Table V, col. 2] and [3, Table VI, col. 10 and col. 12]. Moreover, it is likely that these assumptions or their possible alternatives suggested in Remark 7.6 would also hold for other sets of parameter values if we consider equilibria u_e and inputs g that are not homogeneous over Ω . Checking the assumptions of Theorem 7.5 in this case is, however, not straightforward.

8. Discussion and conclusion. In this paper, we developed basic analytical results to establish a global attractor theory for the mean field model of the electroencephalogram (EEG) proposed by Liley, Cadusch, and Dafilis [36]. We showed that the initial-boundary value problem associated with the model is well-posed in the weak and strong sense, and established sufficient conditions for the nonnegativity of the $i(x, t)$ and $w(x, t)$ components of the solution over the entire time horizon. Moreover, we proved existence of bounded absorbing sets for semigroups of weak and strong solutions, and discussed the challenges involved in proving the asymptotic compactness property for these semigroups. Finally, we showed that the equilibrium sets of the model are noncompact for some physiologically reasonable sets of parameter values which, in particular, implies nonexistence of a global attractor.

The conditions developed in this paper for ensuring nonnegativity of the solution components $i(x, t)$ and $w(x, t)$ over the entire infinite time horizon can be useful for computational analysis of the model. Without using such mathematical analysis, it is impossible to ensure that the solutions computed numerically over a finite time horizon are biophysically plausible since, evidently, negativity might occur for time intervals beyond the finite time horizon of numerical computations. This fact has been overlooked in most of the available computational

analysis of the model. However, in these computational studies, the initial values are usually set equal to a numerically computed space-homogeneous equilibrium of the model, or equal to zero when no equilibrium is found numerically. In both cases, the preset initial values satisfy the sufficient conditions developed in section 5.1 of this paper for biophysical plausibility of the solutions. It is perhaps an intractable problem to specify a set of biophysical initial values for a model of the EEG; however, analyzing a more diverse set of reasonable initial values satisfying the sufficient conditions developed in section 5.1 can be beneficial in observing different behaviors of the model.

Existence of bounded absorbing sets is a desirable global property for a model of electrical activity in the neocortex. As stated in Remark 6.4, the EEG model investigated in this paper possesses this global property for its entire range of parameter values given in Table 1. Moreover, this property holds independently of the parameters of the firing rate functions, the number of intracortical and corticocortical connections, the mean Nernst potentials, and the membrane time constants, as observed in assumptions (i) and (ii) of Theorems 6.2 and 6.3.

The lack of space-dissipative terms in the ODE components (4) and (5) of the model is one of the major sources of difficulty in establishing a global attractor. Indeed, as discussed in section 7.2, the $v(x, t)$ and $i(x, t)$ components of the solution can evolve discontinuously in space despite continuous evolution of the $w(x, t)$ component. Other than disrupting the asymptotic compactness property of the semigroups of solution operators, these space irregularities can predict sharp transitions in the $v(x, t)$ and $i(x, t)$ components of the solution, which can potentially be problematic in numerical computation of the solutions.

Slight modifications to the model that result in the presence of additional space-dissipative terms in the ODEs can improve the regularity of the solutions and can be of particular advantage in numerical computations. The fact that some of the equations of the model appear as ODEs is partially due to the simplifying assumption of *instantaneous* conduction through short-range fibers. Removing such simplifying assumptions, or considering a singularly perturbed version of (4) and (5) by artificially including additional diffusion terms $\varepsilon \Delta$, with sufficiently small ε , can be considered as potential modifications. Any such modifications should, however, maintain the neurophysiological plausibility of the model.

The regularization made by appropriate modifications to the model may result in the possibility of establishing the asymptotic compactness property. However, the analysis in section 7.2 suggests that the resulting compact attractor would be of very high dimension for some sets of parameter values. Based on this observation, we speculate that the noncompactness of the attracting sets shown in this paper can provide an explanation for the possibility of having a rich variety of behaviors for this model, some of which are already shown by computational analysis in the literature; see, for example, [4, 5, 11, 12, 13, 20, 21, 52, 53]. Such diversity of complicated behaviors is indeed what one would expect from a model of the neocortex, the part of the brain that is presumed to be responsible for the extremely complicated perceptual and cognitive functionality of the brain.

Acknowledgment. The authors would like to thank Professor Andrzej Świąch from the School of Mathematics at Georgia Institute of Technology for his helpful suggestions on some of the proofs appearing in this paper.

REFERENCES

- [1] M. F. BEAR, B. W. CONNORS, AND M. A. PARADISO, *Neuroscience: Exploring the Brain*, 4th ed., Wolters Kluwer, Philadelphia, PA, 2016.
- [2] I. BOJAK, H. C. DAY, AND D. T. J. LILEY, *Ketamine, propofol and the EEG: A neural field analysis of HCN1-mediated interactions*, *Front. Comput. Neurosci.*, 7 (2013), 22, doi:10.3389/fncom.2013.00022.
- [3] I. BOJAK AND D. T. J. LILEY, *Modeling the effects of anesthesia on the electroencephalogram*, *Phys. Rev. E*, 71 (2005), 041902, doi:10.1103/PhysRevE.71.041902.
- [4] I. BOJAK AND D. T. J. LILEY, *Self-organized 40 Hz synchronization in a physiological theory of EEG*, *Neurocomputing*, 70 (2007), pp. 2085–2090, doi:10.1016/j.neucom.2006.10.087.
- [5] I. BOJAK, D. T. J. LILEY, P. CADUSCH, AND K. CHENG, *Electrorhythmogenesis and anaesthesia in a physiological mean field theory*, *Neurocomputing*, 58–60 (2004), pp. 1197–1202, https://doi.org/10.1016/j.neucom.2004.01.185.
- [6] I. BOJAK, Z. V. STOYANOV, AND D. LILEY, *Emergence of spatially heterogeneous burst suppression in a neural field model of electrocortical activity*, *Front. Comput. Neurosci.*, 9 (2015), doi:10.3389/fncom.2015.00018.
- [7] V. V. CHEPYZHOV AND M. I. VISHIK, *Attractors for Equations of Mathematical Physics*, American Mathematical Society, Providence, RI, 2002.
- [8] K. CHUEH, C. CONLEY, AND J. SMOLLER, *Positively invariant regions for systems of nonlinear diffusion equations*, *Indiana Univ. Math. J.*, 26 (1977), pp. 373–392, doi:10.1512/iumj.1977.26.26029.
- [9] P. G. CIARLET, *Linear and Nonlinear Functional Analysis with Applications*, Society for Industrial and Applied Mathematics, Philadelphia, PA, 2013.
- [10] S. COOMBES, P. BEIM GRABEN, R. POTTHAST, AND J. WRIGHT, EDS., *Neural Fields: Theory and Applications*, Springer-Verlag, New York, 2014.
- [11] M. P. DAFILIS, F. FRASCOLI, P. J. CADUSCH, AND D. T. J. LILEY, *Four dimensional chaos and intermittency in a mesoscopic model of the electroencephalogram*, *Chaos*, 23 (2013), 023111, doi:10.1063/1.4804176.
- [12] M. P. DAFILIS, F. FRASCOLI, P. J. CADUSCH, AND D. T. J. LILEY, *Extensive four-dimensional chaos in a mesoscopic model of the electroencephalogram*, *J. Math. Neurosci.*, 5 (2015), doi:10.1186/s13408-015-0028-3.
- [13] M. P. DAFILIS, D. T. J. LILEY, AND P. J. CADUSCH, *Robust chaos in a model of the electroencephalogram: Implications for brain dynamics*, *Chaos*, 11 (2001), pp. 474–478, doi:10.1063/1.1394193.
- [14] M. EFENDIEV, *Attractors for Degenerate Parabolic Type Equations*, American Mathematical Society, Providence, RI, 2013.
- [15] G. B. ERMENTROUT AND D. H. TERMAN, *Mathematical Foundations of Neuroscience*, Springer-Verlag, New York, 2010.
- [16] L. C. EVANS, *Partial Differential Equations*, 2nd ed., American Mathematical Society, Providence, RI, 2010.
- [17] R. FITZHUGH, *Impulses and physiological states in theoretical models of nerve membrane*, *Biophys. J.*, 1 (1961), pp. 445–466, doi:10.1016/S0006-3495(61)86902-6.
- [18] G. B. FOLLAND, *Real Analysis: Modern Techniques and Their Applications*, 2nd ed., John Wiley & Sons, New York, 1999.
- [19] B. L. FOSTER, I. BOJAK, AND D. T. J. LILEY, *Population based models of cortical drug response: Insights from anaesthesia*, *Cogn. Neurodyn.*, 2 (2008), pp. 283–296, doi:10.1007/s11571-008-9063-z.
- [20] F. FRASCOLI, M. P. DAFILIS, L. VAN VEEN, I. BOJAK, AND D. T. J. LILEY, *Dynamical complexity in a mean-field model of human EEG*, *Proc. SPIE*, 7270 (2008), 72700V, doi:10.1117/12.813966.
- [21] F. FRASCOLI, L. VAN VEEN, I. BOJAK, AND D. T. J. LILEY, *Metabifurcation analysis of a mean field model of the cortex*, *Phys. D*, 240 (2011), pp. 949–962, doi:10.1016/j.physd.2011.02.002.
- [22] W. GERSTNER AND W. M. KISTLER, *Spiking Neuron Models: Single Neurons, Populations, Plasticity*, Cambridge University Press, New York, 2002.
- [23] K. R. GREEN AND L. VAN VEEN, *Open-source tools for dynamical analysis of Liley’s mean-field cortex model*, *J. Comput. Sci.*, 5 (2014), pp. 507–516, doi:10.1016/j.jocs.2013.06.001.
- [24] J. GUCKENHEIMER, *Review: Roger Temam, Infinite-dimensional dynamical systems in mechanics and physics*, *Bull. Amer. Math. Soc. (N.S.)*, 21 (1989), pp. 196–198, doi:10.1090/S0273-0979-1989-15819-9.

- [25] J. K. HALE, *Asymptotic Behavior of Dissipative Systems*, American Mathematical Society, Providence, RI, 1988.
- [26] A. L. HODGKIN AND A. F. HUXLEY, *A quantitative description of membrane current and its application to conduction and excitation in nerve*, J. Physiol., 117 (1952), pp. 500–544, doi:10.1113/jphysiol.1952.sp004764.
- [27] J. C. HORTON AND D. L. ADAMS, *The cortical column: A structure without a function*, Philos. Trans. R. Soc. Lond. Bio. Sci., 360 (2005), pp. 837–862, doi:10.1098/rstb.2005.1623.
- [28] E. R. KANDEL, J. H. SCHWARTZ, T. M. JESSELL, S. A. SIEGELBAUM, AND A. J. HUDSPETH, *Principles of Neural Science*, 5th ed., McGraw-Hill, New York, 2013.
- [29] J. KEENER AND J. SNEYD, *Mathematical Physiology*, 2nd ed., Springer-Verlag, New York, 2009.
- [30] C. KOCH, *Biophysics of Computation: Information Processing in Single Neurons*, Oxford University Press, New York, 2004.
- [31] M. A. KRAMER, H. E. KIRSCH, AND A. J. SZERI, *Pathological pattern formation and cortical propagation of epileptic seizures*, J. R. Soc. Interface, 2 (2005), pp. 113–127, doi:10.1098/rsif.2005.0028.
- [32] M. A. KRAMER, A. J. SZERI, J. W. SLEIGH, AND H. E. KIRSCH, *Mechanisms of seizure propagation in a cortical model*, J. Comput. Neurosci., 22 (2006), pp. 63–80, doi:10.1007/s10827-006-9508-5.
- [33] M. A. KRAMER, W. TRUCCOLO, U. T. EDEN, K. Q. LEPAGE, L. R. HOCHBERG, E. N. ESKANDAR, J. R. MADSEN, J. W. LEE, A. MAHESHWARI, E. HALGREN, C. J. CHU, AND S. S. CASH, *Human seizures self-terminate across spatial scales via a critical transition*, Proc. Natl. Acad. Sci. USA, 109 (2012), pp. 21116–21121, doi:10.1073/pnas.1210047110.
- [34] K. H. W. KÜFNER, *Invariant regions for quasilinear reaction-diffusion systems and applications to a two population model*, NoDEA Nonlinear Differential Equations and Appl., 3 (1996), pp. 421–444, doi:10.1007/BF01193829.
- [35] D. T. J. LILEY AND I. BOJAK, *Understanding the transition to seizure by modeling the epileptiform activity of general anesthetic agents*, J. Clin. Neurophysiol., 22 (2005), pp. 300–313.
- [36] D. T. J. LILEY, P. J. CADUSCH, AND M. P. DAFILIS, *A spatially continuous mean field theory of electrocortical dynamics*, Network, 13 (2002), pp. 67–113, doi:10.1080/net.13.1.67.113.
- [37] D. T. J. LILEY AND M. WALSH, *The mesoscopic modeling of burst suppression during anesthesia*, Front. Comput. Neurosci., 7 (2013), 46, doi:10.3389/fncom.2013.00046.
- [38] Q. MA, S. WANG, AND C. ZHONG, *Necessary and sufficient conditions for the existence of global attractors for semigroups and applications*, Indiana Univ. Math. J., 51 (2002), pp. 1541–1559, doi:10.1512/iumj.2002.51.2255.
- [39] M. MARION, *Finite-dimensional attractors associated with partly dissipative reaction-diffusion systems*, SIAM J. Math. Anal., 20 (1989), pp. 816–844, doi:10.1137/0520057.
- [40] R. J. MORAN, D. A. PINOTSIS, AND K. J. FRISTON, *Neural masses and fields in dynamic causal modeling*, Front. Comput. Neurosci., 7 (2013), 57, doi:10.3389/fncom.2013.00057.
- [41] V. B. MOUNTCASTLE, *The columnar organization of the neocortex*, Brain, 120 (1997), pp. 701–722, doi:10.1093/brain/120.4.701.
- [42] J. NAGUMO, S. ARIMOTO, AND S. YOSHIKAWA, *An active pulse transmission line simulating nerve axon*, Proc. IRE, 50 (1962), pp. 2061–2070, doi:10.1109/JRPROC.1962.288235.
- [43] D. A. PINOTSIS, P. ROBINSON, P. GRABEN, AND K. J. FRISTON, *Neural masses and fields: Modeling the dynamics of brain activity*, Front. Comput. Neurosci., 8 (2014), 149, doi:10.3389/fncom.2014.00149.
- [44] J. C. ROBINSON, *Infinite-Dimensional Dynamical Systems: An Introduction to Dissipative Parabolic PDEs and the Theory of Global Attractors*, Cambridge University Press, Cambridge, 2001.
- [45] P. A. ROBINSON, C. J. RENNIE, D. L. ROWE, S. C. O’CONNOR, J. J. WRIGHT, E. GORDON, AND R. W. WHITEHOUSE, *Neurophysical modeling of brain dynamics*, Neuropsychopharmacology, 28 (2003), pp. S74–S79, doi:10.1038/sj.npp.1300143.
- [46] P. SANZ-LEON, S. A. KNOCK, A. SPIEGLER, AND V. K. JIRSA, *Mathematical framework for large-scale brain network modeling in the virtual brain*, NeuroImage, 111 (2015), pp. 385–430, doi:10.1016/j.neuroimage.2015.01.002.
- [47] G. R. SELL AND Y. YOU, *Dynamics of Evolutionary Equations*, Springer-Verlag, New York, 2002.
- [48] O. SPORNS, *Contributions and challenges for network models in cognitive neuroscience*, Nat. Neurosci., 17 (2014), pp. 652–660, doi:10.1038/nn.3690.

- [49] M. L. STEYN-ROSS, D. A. STEYN-ROSS, AND J. W. SLEIGH, *Modelling general anaesthesia as a first-order phase transition in the cortex*, Prog. Biophys. Mol. Biol., 85 (2004), pp. 369–385, doi:10.1016/j.pbiomolbio.2004.02.001.
- [50] M. E. TAYLOR, *Partial Differential Equations I: Basic Theorey*, 2nd ed., Springer-Verlag, New York, 2011.
- [51] R. TEMAM, *Infinite-Dimensional Dynamical Systems in Mechanics and Physics*, Springer-Verlag, New York, 1997.
- [52] L. VAN VEEN AND K. R. GREEN, *Periodic solutions to a mean-field model for electrocortical activity*, Eur. Phys. J. Spec. Top., 223 (2014), pp. 2979–2988, doi:10.1140/epjst/e2014-02311-y.
- [53] L. VAN VEEN AND D. T. J. LILEY, *Chaos via Shilnikov’s saddle-node bifurcation in a theory of the electroencephalogram*, Phys. Rev. Lett., 97 (2006), 208101, doi:10.1103/PhysRevLett.97.208101.
- [54] H. R. WILSON AND J. D. COWAN, *Excitatory and inhibitory interactions in localized populations of model neurons*, Biophys. J., 12 (1972), pp. 1–24, doi:10.1016/S0006-3495(72)86068-5.
- [55] M. T. WILSON, P. A. ROBINSON, B. O’NEILL, AND D. A. STEYN-ROSS, *Complementarity of spike- and rate-based dynamics of neural systems*, PLoS Comput. Biol., 8 (2012), doi:10.1371/journal.pcbi.1002560.
- [56] Y. YOU, *Global dynamics of the Brusselator equations*, Dyn. Partial Differ. Equ., 4 (2007), pp. 167–196, doi:10.4310/DPDE.2007.v4.n2.a4.