


Using Structurally Well-Defined Norbornyl-Bridged Acene Dimers to Map a Mechanistic Landscape for Correlated Triplet Formation in Singlet Fission

Alexander T. Gilligan, Ethan G. Miller, Tarek Sammakia,^{ip} and Niels H. Damrauer*^{ip}

Department of Chemistry and Biochemistry, University of Colorado Boulder, Boulder, Colorado 80309, United States

Supporting Information

ABSTRACT: Structurally well-defined TIPS-acetylene substituted tetracene (TIPS-BT1') and pentacene (TIPS-BP1') dimers utilizing a [2.2.1] bicyclic norbornyl bridge have been studied—primarily using time-resolved spectroscopic methods—to uncover mechanistic details about primary steps in singlet fission leading to formation of the biexcitonic ¹TT state as well as decay pathways to the ground state. For TIPS-BP1' in room-temperature toluene, ¹TT formation is rapid and complete, occurring in 4.4 ps. Decay to the ground state in 100 ns is the primary loss pathway for ¹TT in this system. For TIPS-BT1', the ¹TT is also observed to form rapidly (with a time constant of 5 ps), but in this case it occurs in concert with establishment of an excited-state equilibrium ($K \sim 1$) with the singlet exciton state S_1 at an energy of 2.3 eV above the ground state. The equilibrated states survive for 36 ns and are lost to ground state through both radiative and nonradiative pathways via the S_1 and nonradiative pathways via the ¹TT. The rapidity of ¹TT formation in TIPS-BT1' is at first glance surprising. However, our analysis suggests that the few-parameter rate constant expression of Marcus theory explains both individual and comparative findings in the set of systems, thus establishing benchmarks for diabatic coupling and reorganization energy needed for efficient ¹TT formation. Finally, a comparison of TIPS-BT1' with previous results obtained for a close constitutional isomer (TIPS-BT1) differing in the placement of TIPS-acetylene side groups suggests that the magnitude of exchange interaction in the correlated triplet manifold plays a critical role dictating ¹TT yield in the tetracenic systems.



INTRODUCTION

Molecular dimers have emerged as key platforms for the mechanistic exploration of singlet fission (SF),^{1,2} and in particular initial photophysics wherein a photoinduced singlet exciton is transformed into a multiexciton state, which is characterized as a singlet-coupled pair of triplets (¹TT). Understanding how to control such dynamics is motivated by the premise that SF may serve as a means to down-convert higher energy solar photons into multiple electronic excitations rather than into a single excitation plus waste heat.³ Dimer and small oligomer systems using acenes^{4–20} and also diimides²¹ and isobenzofurans^{22,23} are enabling the interrogation of numerous fundamental issues affecting SF rates and yields, including reaction thermodynamics,^{6,16–18,24} state couplings,^{8,9,22,25,26} charge-transfer (CT) intermediates,^{10,19,21,22} the role of entropy,⁸ spin dynamics,^{15,27} and exciton binding.^{8,23,28}

Within the overall body of dimer work in the literature, a leading role has been played by pentacene-based systems^{5–7,10–12,14,15,17–19} where the $S_1 \rightarrow$ ¹TT reaction driving force is significant at -200 to -300 meV and where ¹TT yields are commonly high, even in the first systems reported.^{5–7} A variety of structural motifs have been explored which fall loosely into two groups. In one of these, dimer connectivity

occurs via the chromophore ends either using single bonds through the acene 2 position^{5,11} or using bicyclic moieties that connect simultaneously through the 2 and 3 positions.^{14,17,18} This latter group includes the [2.2.1]-bridge dimer TIPS-BP1' (see Figure 1) discussed herein whose synthesis and preliminary photophysics were recently reported by us.¹⁸ In the second group, connectivity occurs at the pentacene middle, through the 6 position directly^{6,10} or via acetylene substituents that then link to a common bridge.^{7,12,15,19} While the scope of systems is relatively large and growing, there is not yet consensus about factors controlling important mechanistic details, such as the rate constant for the $S_1 \rightarrow$ ¹TT forward process. For example, there remain questions about electronic coupling for the photoreaction and whether it is dominated by terms that (a) directly connect the single and double exciton states²⁹ or (b) demand participation by virtual CT states, as is the more common assumption, or (c) entirely system specific. We believe that structurally well-defined dimer systems, including our [2.2.1]-bridge approach and the [2.2.2] and spiropyran approaches of Campos and Sfeir,¹⁴ can play an important clarifying role in the field. By reducing conforma-

Received: January 24, 2019

Published: March 19, 2019




Figure 1. Norbornyl-bridged acene dimers discussed in text. BT1 is the conceptual parent.⁹ The photophysics of triisopropylsilyl (TIPS)-acetylene substituted bis-tetracene TIPS-BT1 has been explored extensively elsewhere.^{9,13} This current work focuses on the constitutional isomer TIPS-BT1' and the bis-pentacene dimer TIPS-BP1'.¹⁸

tional freedom, such systems limit configuration interaction with low-energy singlet excimer states.^{8,16} As well, they limit uncertainties about state coupling magnitudes and mechanisms that depend on relative chromophore orientation and orientation with respect to bridging moieties. From this vantage point, we would argue that structural definition in dimers provides an opportunity to connect with theory through powerful few-parameter rate expressions such as Marcus theory.^{30,31} If this is the case, and if computational tools can be employed to accurately predict physical quantities such as diabatic state couplings, then unifying design principles may have a better chance of emerging.

Although to a lesser extent than the pentacenic systems discussed above, tetracenic dimers have also been explored and contribute to an overall mechanistic understanding. Early work by Bardeen and co-workers considered phenylene-spaced tetracene dimers.⁴ They saw evidence in delayed fluorescence for involvement of the $S_1 \rightarrow {}^1\text{TT}$ photoreaction (and its reverse), although they concluded that the ${}^1\text{TT}$ yield was low, of order 3%; notably, that yield can be substantially increased in related systems by introduction of small oligomers such as trimers and tetramers.^{32,33} By contrast, Bradforth, Thompson, and co-workers studying highly through-space coupled tetracene dimer systems saw quantitative conversion of the singlet exciton to a new state that bears both excimer and multieexcitonic (${}^1\text{TT}$) character.⁸ In a related system modified to engage only through-bond coupling, they later report rapid formation of a triplet signature that lives for ~ 100 ps, consistent with ${}^1\text{TT}$ participation in the overall photoreaction.²⁰ In more rigid and weakly coupled dimers, we initially reported photoluminescence dynamics in room-temperature toluene for our [2.2.1]-bridge parent BT1⁹ (Figure 1) and like Bardeen and co-workers concluded that the ${}^1\text{TT}$ yield was low. Our subsequent photophysical studies

of a more soluble dimer TIPS-BT1 (Figure 1) in toluene showed single-exponential singlet-exciton loss concomitant with ground state recovery on the 24 ns time scale, and we concluded that the $S_1 \rightarrow {}^1\text{TT}$ photoreaction was not operable in that system.¹³ We understood this as being a manifestation of point group symmetry properties in the dimer, and specifically a plane of symmetry that passes through both chromophores of the dimer, that limits electronic coupling in the photoreaction (this symmetry plane can be understood easily in the diabatic state picture as obviating nonhorizontal electron-transfer couplings between virtual CT states and the ${}^1\text{TT}$).^{25,34,35} Interestingly, Saito and co-workers recently studied a bent cyclooctatetraene-bridged TIPS-tetracenic dimer with comparable symmetry called FLAP2, and while it has poor photostability compared to its anthracenic and pentacenic analogues, it offers compelling evidence for engaging the $S_1 \rightarrow {}^1\text{TT}$ photoreaction on a ps time scale.¹⁷ Those workers note that FLAP2 would have substantially more conformational flexibility about the bridge compared to TIPS-BT1 and suggest that this could lead to the stark dynamical differences between the two dimer systems.

In the work that follows, we explore excited-state dynamics for a constitutional isomer of TIPS-BT1 called TIPS-BT1' (Figure 1), where the acetylene substitution pattern on each chromophore is moved outward by a ring relative to the bridge, comparable to what is seen in FLAP2. Transient spectral data offer compelling evidence for the $S_1 \rightarrow {}^1\text{TT}$ photoreaction as part of a picosecond time scale equilibration between these states. These data then suggest that the photoreaction energetics are highly sensitive to subtle changes in substitution patterns, for example, between TIPS-BT1 and TIPS-BT1' and lead to marked changes in ${}^1\text{TT}$ yield. Overall, Marcus theory offers a unifying explanation of dynamics in the full set of substituted dimers—TIPS-BT1, TIPS-BT1', and the pentacenic TIPS-BP1'—with vibronic coupling derived from symmetry-breaking motions being sufficient to engender fast dynamics.

RESULTS AND DISCUSSION

Pentacene Dimer. It is useful to start by characterizing the photoinduced dynamics of TIPS-BP1' (Figure 1), a molecule whose reaction driving force is expected to facilitate rapid formation of ${}^1\text{TT}$ based on results from a growing number of pentacene-based systems in the literature.^{3-7,10-12,14,15,17-19} For example, in phenylene-bridged TIPS-pentacene dimers studied by Guldi, Tykwinski, and co-workers, phosphorescence measurements identified a T_1 energy of 0.8 eV.⁷ Given the S_1 energy of 1.9 eV (ours is measured at 1.93 eV, *vide infra*), their systems were thermodynamically competent for singlet fission with a driving force of -0.3 eV. It is noted that in the communication of our synthetic methodology, we showed preliminary spectral evidence for ${}^1\text{TT}$ in TIPS-BP1' at 10 ps following photoexcitation.¹⁸ However, that work did not establish time constants or yields. Beginning with ground-state absorption, Figure 2a shows a normalized spectrum collected for TIPS-BP1' in room-temperature toluene in a wavelength region that is coincident with our TA measurements described below. To the red is a vibronic progression characteristic of TIPS-Pentacene (TIPS-Pc) moieties, with the 0–0 band peaking at 638 nm. As we have previously described for related systems, the symmetry of this dimer and the fact that the $S_1 \leftarrow S_0$ is acene short-axis polarized, means that only the higher-energy excitonic transition in a Davydov-split pair is




Figure 2. (a) Steady-state electronic absorption spectrum of TIPS-BP1' in toluene at room temperature. (b) TA spectra of TIPS-BP1' in room-temperature toluene (normalized at $\Delta t = 530$ ps). The region surrounding the excitation wavelength of 588 nm is removed due to pump scatter. (c) Selected single wavelength kinetics traces (data points) for TIPS-BP1' with applied model fits (lines) retrieved from global analysis. (d) Kinetic model of decay pathways of TIPS-BP1' after initial excitation.

bright.^{9,13,18} In other words, this system exhibits H-type coupling with respect to the $S_1 \leftarrow S_0$ transition of each chromophore arm. To the blue and peaked at 444 nm is a second progression that is also observed in monomer models such as TIPS-Pc.¹⁴ Not observed in toluene due to its UV cutoff is the characteristic Davydov splitting associated with coupling the individual-chromophore long axis transitions. As we have shown elsewhere,¹⁸ this feature is seen for the molecule in chloroform with intense absorption bands at 308 and 333 nm indicating a peak splitting of 0.30 eV. The relative intensity of these two features is readily understood^{9,13,18} given the geometry of the dimer (and in particular the obtuse angle of 113° between chromophores) where the more intense lower energy transition at 333 nm arises due to the in-phase addition of the transition dipole moments (*x*-polarization; where dimer is oriented in the *xz* plane), whereas the less-intense higher energy transition at 308 nm is due to the subtraction of the transition dipole moments (*z*-polarization; where the *z*-axis coincides with the *C*₂ symmetry element.) A molar extinction spectrum collected in chloroform is shown in Figure S1. The dimer exhibits very weak photoluminescence (Figures S2 and S16) with an emissive quantum yield of ~0.5% (compared with 72% for TIPS-BT1 and TIPS-BT1'). Given rate constant and driving force arguments for ¹TT formation in TIPS-BP1' (vide infra), we suspect this emission arises from an impurity not detected in the ¹H NMR baseline¹⁸ where one of the two

arms has been oxidized, while the second arm remains acene-like and photoluminescent.

Transient absorption (TA) dynamics were collected for TIPS-BP1' in room-temperature toluene following photoexcitation with ~50 fs laser pulses at a center wavelength of 588 nm (Figure 2b). The early transient spectrum resembles the lowest energy singlet exciton in a monomer model TIPS-Pc³⁶ (see Figure S17) including the excited-state absorption (ESA) at ~440–470 nm. That spectrum rapidly gives way to a new one that is characterized by the strong ESA at 517 nm along with a vibronic shoulder at 480 nm. These features, which do not further evolve out to the ~1 ns limit of this experiment, herald a state with triplet electronic character as seen in a number of SF-active systems involving TIPS-acetylene substituted pentacene chromophores.^{5,7} In line with other studies, the speed of the reaction is highly suggestive that the product state is not T_1 produced through intersystem crossing, but rather the ¹TT produced with spin conserving internal conversion. The data over the time range from 0.5 to 1500 ps are readily fit with a global A → B model with a time constant of 4.4 ps (Figure S4). It is noted that the strong ESA feature shows a small ~1 nm blueshift over the course of its formation. Although not definitively assigned at this point, it is our expectation that the reactant singlet exciton (state A) is delocalized over both acene arms as was indicated in detailed time-resolved emission studies of TIPS-BT1.¹³ In order to estimate the yield of ¹TT (state B), a sensitization experiment was undertaken to determine the molar extinction of the triplet in TIPS-BP1', using photoexcited (360 nm) anthracene as a collisional triplet-triplet energy transfer partner (see details in the SI and Figure S11). Here, the assumption is made that the spectral character of T_1 (observed lifetime $\tau_{\text{obs}} = 55 \mu\text{s}$ in room-temperature toluene; see Figure S13) is a suitable surrogate for each of the two chromophores in the ¹TT of TIPS-BP1'. This situation is enabled by the structural rigidity of this dimer, which limits conformational relaxation that might permit significant admixture by other states in the singlet manifold such as excimers.^{8,10} With this analysis (see details in the SI) we find a yield of $97 \pm 11\%$ from the perspective of the ¹TT or $194 \pm 22\%$ from the perspective of triplet excitons (see SI for a description of how error was propagated). These values are in line with those seen in other pentacene dimer systems.^{5,10,12,14}

A longer time resolution TA spectroscopy was used to interrogate the fate of the transient described above that was produced in 4.4 ps. As shown in Figure S6, the large majority of the signal decays toward baseline with single exponential character and a lifetime of 102 ns. This shortened lifetime for a species that has triplet spectral character (vide supra) is further support for the assignment to ¹TT.^{5,6,15} It is noted that a minor 3.5% shelf is observed in the time window whose eventual decay to baseline requires 56 μs , thus suggesting assignment to T_1 . Power-dependent studies did not show a percentage change in the magnitude of the shelf thereby arguing against production of T_1 by collision between ¹TT and ground-state species (Figure S10). It is possible that the shelf manifests as the spin-entangled ¹TT mixes with the ³TT and eventually undergoes decoherence within the dimer into pairs of uncorrelated triplets.^{15,27} If this is the case, the shelf would correspond to a dissociated triplet yield of 7% of a possible 200%. A full assignment will require spin-sensitive measurements such as time-resolved EPR.^{15,27,37,38} Regarding the 102 ns lifetime tied to the ¹TT → GS decay, it is acknowledged

that this time scale is considerably lengthened compared to observations in initial highly coupled pentacene dimer systems (e.g., BP0 from Campos/Sfeir,⁵ DP-Mes, and DP-TIPS from Musser and co-workers,¹⁰ and the pheny-ethynyl-bridged systems of Guldi/Tykwinski⁷). However, a number of systems have now been reported with >100 ns TT lifetimes in a variety of solvents.^{5,14,15}

Discussion of a Common Model. As these TIPS-BP1' studies will help us to understand data in the full series of molecules (Figure 1), it is useful to present a common framework for analyzing kinetics at this point in the paper. Because of the structural definition of these types of dimers, along with the weakly polar solvent environment that precludes significant participation by CT states (vide infra), a relatively simple three-state model can be utilized (Figure 2d).^{8,9,21} This includes a singlet exciton state, the TT, and the ground state. The singlet exciton state is coupled directly to the ground state via both radiative and nonradiative pathways (k_r and k_{nr}), and it can also be lost due to formation of the TT via k_{fiss} or reformed via the fusion process encompassed in k_{fus} . The last rate constant component in this model is the loss pathway linking the TT directly to the ground state, which is referred to as k_{TT} . In our understanding of these systems at this time, we assume that TT is primarily the pure singlet ^1TT produced in the spin-allowed k_{fiss} process, but recognize that this is not an eigenstate of the system^{1,39} and that spin mixing with the ^5TT will begin to occur during the TT lifetime. In a related vein, the model ignores processes leading to the singlet fission product $T_1 + T_1$, which is presumed to occur in conjunction with spin mixing and decoherence, via the ^5TT . As a common model for each of the dimers explored, this is reasonable given that for TIPS-BP1' the long-time shelf corresponding to this product is relatively small (<3.5%) and for TIPS-BT1' it is nearly undetectable.

With this model we can now establish rate constants for the photophysical behavior in TIPS-BP1'. Recalling that the measured ^1TT yield determined using sensitization experiments is approximately quantitative, a large equilibrium constant $K = k_{fiss}/k_{fus}$ ($K \geq 100$) is expected such that the observed exponential decay of 4.4 ps reflects $1/k_{fiss}$ with little contamination (<1%) from k_{fus} . Note that $K = 100$ at room temperature for a system with a modest $S_1 \rightarrow ^1\text{TT}$ reaction driving force of -0.12 eV. If the driving force were -0.34 eV as estimated in a related system,⁷ the equilibrium constant K would be $>5 \times 10^5$. The large equilibrium constant K also means that the observed 102 ns lifetime of the TA signal has little contamination from k_r and k_{nr} and rather reflects, almost exclusively, $1/k_{TT}$. The values of k_{fiss} and k_{TT} obtained for TIPS-BP1' are listed in Table 1.

Tetracene Dimers. We next consider the photoinduced dynamics of TIPS-BT1' whose synthesis follows the same general approach used to prepare the larger acene dimer TIPS-BP1'.¹⁸ As described in the Introduction, we had previously concluded that the close tetracene dimer analog TIPS-BT1 is inactive toward ^1TT formation as studied in weakly polar toluene.¹³ As such, our assumption at the outset was that TIPS-BT1' would also be inactive toward these photophysics due to their structural similarity. This assumption is called into question below.

Steady-state absorption for TIPS-BT1' in room-temperature toluene is shown in Figure 3 in a spectral region highlighting properties of the lowest energy allowed vibronic transition. TIPS-BT1', like TIPS-BP1', exhibits H-type coupling with an

Table 1. Summary of Room Temperature Photophysical Properties for Dimer Species in Toluene

| | TIPS-BT1 ^a | TIPS-BT1' ^b | TIPS-BP1' ^b |
|--------------------------------------|-----------------------|--------------------------------|--------------------------------|
| $\Phi_{\text{em}}^{\text{c}}$ | 0.72 ± 0.09 | 0.72 ± 0.09 | <0.01 |
| $\tau_{\text{obs-fast}}$ (ps) | 0.85 | 2.5 ± 0.3 | 4.4 ± 0.2 |
| k_{fiss} (s ⁻¹) | 1.1×10^{11} | $(2.0 \pm 0.2) \times 10^{11}$ | $(2.3 \pm 0.1) \times 10^{11}$ |
| k_{fus} (s ⁻¹) | 1.1×10^{12} | $(2.0 \pm 0.2) \times 10^{11}$ | $<(2.2 \pm 0.1) \times 10^9$ |
| τ_{obs} (ns) | 24.3 | 36 ± 3 | 102 ± 3^d |
| k_{TT} (s ⁻¹) | — | — | $(9.8 \pm 0.3) \times 10^6$ |
| $\phi(^1\text{TT})$ | ≤ 0.1 | 0.50 ± 0.08 | $\geq 0.97 \pm 0.11$ |
| S_1 (eV) | 2.33 | 2.32 | 1.93 |
| $K = k_{\text{fiss}}/k_{\text{fus}}$ | 0.1 | 1.0 ± 0.1 | $10^2-10^5^e$ |

^aTIPS-BT1 taken from known value.¹³ ^bReported error is 2σ of three independent measurements. ^cTIPS-BT1' measured relative to coumarin 540A (coumarin 153) in methanol ($\Phi_{\text{em}} = 0.45$),⁴⁰ and TIPS-BP1' measured relative to oxazine 720 (oxazine 170) in methanol ($\Phi_{\text{em}} = 0.63$).⁴¹ ^dLifetime represents decay of 97% of initial signal. The remaining signal decays with a lifetime of 56 ± 10 μs . ^eSee text for discussion of this range of K .




Figure 3. Normalized steady-state electronic absorption (solid) and emission (dashed) spectra for TIPS-BT1 (red) and TIPS-BT1' (green) in room-temperature toluene.

optically allowed higher energy transition and a dark energetically lower but proximal transition. Also shown in Figure 3 is the emission spectrum collected for TIPS-BT1' in the same solvent. The spectrum mirrors the absorption and shows Stokes shifting of 8 nm. From the average of the 0–0 absorption and emission peaks, the value of the optically bright S_1 is determined to be 2.32 eV (see Table 1).

As was also the case for TIPS-BP1', the toluene solvent UV absorption cutoff precludes observation of Davydov coupling between chromophore long-axis transitions. An absorption spectrum collected for TIPS-BT1' in room-temperature chloroform that does show this splitting is presented elsewhere¹⁸ (the molar extinction spectrum is also presented in Figure S1). In those data, the splitting is 0.47 eV; that is, a value substantially larger than what is observed for TIPS-BP1' (0.30 eV, vide supra). It is understood that a significant fraction of the Davydov splitting occurs via Coulomb interaction between individual chromophore transition dipole moments^{34,42} and in the case of TIPS-BT1' those moments have a smaller separation than in TIPS-BP1'.

We next consider a comparison of steady-state photophysical data collected for TIPS-BT1' versus the constitutional isomer TIPS-BT1. Of note, there is very little wavelength shift between these two molecules. The 0–0 transition in TIPS-BT1' is red-shifted relative to TIPS-BT1 in both absorption and emission data with the bathochromic shift being small (3

and 4 nm, respectively). Averaging 0–0 absorption and emission peaks, the optically bright S_1 in TIPS-BT1 was determined to be 2.33 eV¹³ (see Table 1) or 10 meV higher than what is found in TIPS-BT1'. There are subtle spectral differences between these two molecules that are also worth noting. For TIPS-BT1', the ratio of 0–0 to 0–1 peak heights in *both* absorption and emission experiments is larger than what is found in TIPS-BT1. For the S_1 manifold, this is an indication that the two chromophores in TIPS-BT1' are more weakly interacting than what is seen in TIPS-BT1.⁴³ Stated a different way, it can be said that in TIPS-BT1', where the silyl-acetylene groups of the two chromophores are further separated from one-another, the absorptive and emissive transitions are more characteristic of monomer-like line shapes.

Time-correlated single photon measurements at 539 and 584 nm, the primary spectral features in Figure 3, were employed to determine photoluminescence lifetime properties for TIPS-BT1'. The data sets can be modeled using a single exponential decay function with time constant $\tau_{\text{obs}} = 36 \pm 3$ ns (see Figure S3). Notably this observed lifetime is larger than the value recorded for TIPS-BT1 ($\tau_{\text{obs}} = 24.3$ ns¹³) in the same solvent and temperature. Both values are larger than the lifetime collected for the monomer TIPS-Tc ($\tau_{\text{obs}} = 12.5$ ns¹³). We will come back to the lifetime difference between TIPS-BT1' and TIPS-BT1 as it relates to interpretation of an overall decay model for these types of systems.

In our previous communication of the synthetic approach to TIPS-BT1' and TIPS-BP1', we reported initial TA spectra collected for these dimers at $\Delta t = 1$ and 10 ps after photoexcitation over a probe spectral range of 450–650 nm chosen to interrogate the larger dimer TIPS-BP1'.¹⁸ In that probe range, no substantial changes were observed for TIPS-BT1', and this lead us to a preliminarily conclusion that SF dynamics are inactive, in line with our interpretation of photophysics for TIPS-BT1.¹³ However, that TA experiment has now been revisited with finer time resolution and using a bluer probe spectrum inspired by the band shape changes observed for TIPS-BP1' in Figure 2.

TA dynamics for TIPS-BT1' following ~50 fs pulse excitation at a center wavelength of 530 nm are shown in Figure 4b. Unlike previous measurements for TIPS-BT1 where spectral dynamics were not observed,¹³ these new data for TIPS-BT1' show striking evolution within the first ~15 ps in spectral regions blue of 450 nm. In particular, rapid loss of intensity is seen for a band in the vicinity of 425 nm, whose line shape is modified by ground-state bleach features (see comparison with Figure 4a), but otherwise heralds the S_1 . Dynamics are seen at other wavelengths as well including significant modification of the magnitude of stimulated emission monitored at ~584 nm. Single wavelength kinetic traces extracted from the full spectral data indicate changes in the first 15 ps followed by a lack of further evolution on the 100 ps time scale. The full data set for TIPS-BT1' inclusive of spectra from $\Delta t = 500$ fs to 1.5 ns can be modeled using two single exponentially decaying basis functions, one of which has a time constant of 2.5 ps, while the second is longer but poorly determined given the time limit of this TA experiment (see modeling discussion in SI and species associated spectra in Figure S5). We will return to the faster dynamics later and discuss the slower decay first.

To better resolve the slower dynamics, the second TA spectrometer with longer time resolution was again employed. Transient spectral features of TIPS-BT1' decay to <1% of




Figure 4. (a) Steady-state electronic absorption (solid) and emission (dashed) spectra of dimer TIPS-BT1' in toluene at room temperature. (b) TA spectra of TIPS-BT1' in toluene following ultrafast excitation at 530 nm. The spectral region around the excitation wavelength is removed due to pump scatter. (c) Selected single wavelength kinetics traces (data points) taken from the full-spectrum data with applied model fits (lines) retrieved from global analysis.

baseline and are globally modeled using a single exponential decay with a time constant of 35.8 ns (see Figure S8). The spectral profile is identical to the second retrieved global fit basis spectrum. This time constant matches the 36 ns lifetime determined from the time-correlated single photon counting studies well (vide supra) and represents ground-state recovery.

Returning to the faster 2.5 ps spectral dynamics in Figure 4, it is noted that the observed changes *cannot* be rationalized by invoking the participation of an intramolecular CT state formally reducing one chromophore arm of the dimer while oxidizing the other. Whereas population of such a state was previously observed in TIPS-BT1, that measurement required solvation in a polar benzonitrile medium, and the results highlighted that excited-state equilibrium is established between the CT and a dimer-delocalized singlet exciton at 2.29 eV above ground state.¹³ For the same molecule in less polar toluene, where the singlet exciton state is at a similar energy of 2.33 eV, no CT excited-state properties are observed.¹³ From the perspective of TA spectral changes, the observation of CT for TIPS-BT1 in benzonitrile was very clearly indicated by a transient increase in the magnitude of features tied to the ground-state bleach. This was particularly noticeable at probe wavelengths between ~460 and 525 nm

where singlet exciton ESA features overlap strongly with loss of $S_1 \leftarrow S_0$ absorption: As the singlet exciton ESA is lost in populating the CT, the bleach-related features grow in magnitude with large $-\Delta A$ variations. Such changes are absent in TIPS-BT1' in toluene (Figure 4b), and in fact at a wavelength of 515 nm, we observe a small positive change in ΔA as the dynamics unfold.

On the other hand, it is possible to rationalize the transient spectral changes observed for TIPS-BT1' in Figure 4b if the state being populated has triplet electronic character. Figure 5a




Figure 5. (a) Triplet $\Delta\epsilon$ spectrum for TIPS-BT1' from sensitization experiment in toluene (see SI for sensitization experiment details and Figure S13). (b) Selected spectral slices for TIPS-BT1' at 1 ps (blue) and 120 ps (red) along with a reconstructed TA spectrum (green) that is comprised of a superposition between the 1 ps TA spectrum and the sensitized triplet $\Delta\epsilon$ spectrum from (a).

presents the $\Delta\epsilon$ spectrum collected for TIPS-BT1' following triplet sensitization (see SI for experimental details and Figure S13) which shows two important qualities: first, weak ESA to the blue of 450 nm, and second, stronger ESA between 450 and 550 nm that is highly modulated with ground-state bleach features leading to the appearance of several positive and negative TA features. The importance of the former is tied to that fact that in TIPS-BT1' and in other acetylene-substituted tetracene dimers, the singlet exciton state produced by visible light absorption has a strong ESA in the 400–450 nm region. As time evolves and population leaves this state, a weak ESA in the product can accommodate observation of transient loss in ΔA , consistent with what is seen in the first 10 ps (Figure 4b). The importance of the latter ties to our ΔA observations between 460 and 550 nm, where changes during the dynamics are actually muted. In TIPS-BT1', both the nascent singlet exciton and the triplet observed in this region have a strong ESA that is highly modulated by negative peaks associated with

ground-state bleach (see $\Delta t = 1$ ps in Figure 4b). Thus, during interconversion from excited-state reactant to product, overall changes in ΔA in this spectral region may in principle be subtle.

A more quantitative analysis begins by treating later-time spectra—described by the second component retrieved from the global analysis—in terms of two basis functions. The first is a ΔA spectrum collected at early time ($\Delta t = 1$ ps) where the dominant contribution is from the singlet exciton whose excited-state concentration can be quantified by taking into account the laser power, spot size, and sample absorbance (see SI for details). The second is the triplet $\Delta\epsilon$ spectrum discussed above (Figure 5a). Using a superposition of these two basis functions (50% singlet exciton and 100% triplet), we are able to recreate the $\Delta t = 120$ ps spectrum with high fidelity as shown in Figure 5b. There are two clear implications. The first is that the early dynamics serve to establish an equilibrium between the singlet exciton and a state with triplet character. Given the time scale for the dynamics, that product state cannot be the T_1 and rather, is very likely, the ^1TT where the structural integrity of the bridging norbornyl group enables the two chromophores to essentially preserve their triplet electronic character. This behavior in a tetracene dimer is different than a case where face-to-face interchromophore contact is more intimate leading to significant electronic perturbations.⁸ On the other hand, it is similar to observations by Saito and co-workers where the chromophores are separated by a bridge derived from cyclooctatetraene.^{17,44} Note that in ref 44 we discuss our conclusion that these workers overestimate their triplet yield and that their results are more closely aligned with ours than has been reported.^{17,44} The second implication has to do with the basis function percentages needed to reproduce the later-time spectra. The 100% triplet yield needed should be thought of as a 50% TT, so the overall population remains conserved in the experiment (50% S_1 and 50% ^1TT). Thus, the equilibrium constant established with the 2.5 ps time scale is $K = 1$. In terms of energetics, this means that the ^1TT lies in close energetic proximity to 2.32 eV where we measure the optically bright S_1 (Table 1).

Disentangling Dynamics in TIPS-BT1'. Kinetic modeling using the framework presented in Figure 2d was undertaken for the TIPS-BT1' data. There are too few independent measurements to uniquely determine each of the rate constants, and we choose to draw from information obtained with the other dimers TIPS-BP1' and TIPS-BT1 in order to gain insight. A starting point is the final decay rate constant k_{TT} . In TIPS-BT1', ground-state recovery is strongly influenced by the three rate constants k_r , k_{nr} , and k_{TT} , such that lifetime measurement—even with inclusion of radiative quantum yield information—is insufficient for independent determination of k_{TT} . We thus rely on insight from the larger dimer TIPS-BP1' where k_{TT} was determined to be $1 \times 10^7 \text{ s}^{-1}$ (Table 1). While useful for modeling purposes, this value is likely an overestimate for TIPS-BT1'. First, $^1\text{TT} \rightarrow \text{GS}$ is highly exergonic for both dimers (>-1.5 eV) with values that are significantly larger than what would be expected for the reorganization energy of the electronic transformation in each corresponding system. Thus, $^1\text{TT} \rightarrow \text{GS}$ for either TIPS-BP1' or TIPS-BT1' is expected to take place in the Marcus inverted region where the reaction should slow as the driving force is increased from TIPS-BP1' to TIPS-BT1'. Indeed Sanders et al. have observed energy gap law behavior for this decay process

in a series of heterodimers.²⁴ Unfortunately, estimating the extent of the effect in TIPS-BT1' is further challenged because reorganization energy is also impacted as the acene size is changed. Notably, however, the conclusions reached below are relatively insensitive to the precise value of k_{TT} , and we are comfortable setting the value for TIPS-BT1' at the value measured for TIPS-BP1'.

The next consideration is k_r and k_{nr} . Here the dimer system TIPS-BT1 is useful as it is a close structural analog to TIPS-BT1', but one where ${}^1\text{TT}$ formation is minor (K is small) such that the previously reported values of k_r and k_{nr} ($k_r = 3.0 \times 10^7 \text{ s}^{-1}$ and $k_{\text{nr}} = 1.2 \times 10^7 \text{ s}^{-1}$) are the dominant decay paths. A minor point about these rate constants is discussed in ref 45. Using these k_r and k_{nr} values along with k_{TT} obtained from TIPS-BP1', the three-state model predicts an observed lifetime for TIPS-BT1'—that of the $S_1 \rightleftharpoons {}^1\text{TT}$ equilibrium—of 38 ns. This is, in our view, remarkably similar to the kinetic observation of 36 ns (Table 1), thus providing strong support that we understand this TIPS-BT1' system and that a simple three-state model is appropriate.

The final consideration is k_{fiss} and k_{fus} . The observed 2.5 ps dynamics in TIPS-BT1' represents establishment of the $S_1 \rightleftharpoons {}^1\text{TT}$ equilibrium, which then decays in 36 ns. Because of the large separation in these time scales, the rate constant for establishing the equilibrium is simply the sum of k_{fiss} and k_{fus} ($k_{\text{obs-fast}} = k_{\text{fiss}} + k_{\text{fus}} = 4.0 \times 10^{11} \text{ s}^{-1}$). Given $K = 1$, ${}^1\text{TT}$ is both formed and lost with a time constant of 5 ps ($k_{\text{fiss}} = k_{\text{fus}} = 2.0 \times 10^{11} \text{ s}^{-1}$). These rate constants were able to accurately reproduce the dynamics of the S_1 and ${}^1\text{TT}$ populations present in TIPS-BT1' (see example in Figure S14). The large k_{fiss} in TIPS-BT1' was initially surprising to us given aforementioned symmetry issues for this class of dimers.²⁵ However, diabatic coupling arguments can serve as a basis for understanding this rate constant magnitude. In theoretical explorations of vibronic coupling in BT1—explored because many vibrations break the aforementioned plane of symmetry—we predicted diabatic couplings (V_{eff}) between a singlet exciton state and the ${}^1\text{TT}$ of order 5.5 meV.²⁵ Such a quantity is not insignificant inasmuch as it approximately matches what is predicted²⁵ for tetracene dimer pairs (7.3 meV) germane to the crystal environment where singlet fission is known to take place on the picosecond time scale and be quantitative.³⁹ Although we have not calculated a comparable V_{eff} value for a TIPS-BT1' model, we apply the 5.5 meV from the structurally similar BT1 to make rate constant estimates with nonadiabatic Marcus theory (see SI for the equation and a schematic in Figure S15 showing the parameters). Using this 5.5 meV V_{eff} as the state coupling, along with a reaction driving force $\Delta G = 0$ meV that is appropriate for a system where $K = 1$, one matches the $k_{\text{fiss}} = 2 \times 10^{11} \text{ s}^{-1}$ of TIPS-BT1' when the reorganization energy of the reaction is small, but not unreasonable, at $\lambda = 0.18$ eV (note that $\lambda = 0.13$ eV has been used in the description of ${}^1\text{TT}$ formation in solid-state acene systems).³¹ These V_{eff} and λ parameter values justify the use of nonadiabatic Marcus theory. First the electronic coupling between reactant and product states (V_{eff}) is weak and less than kT . More importantly, its value is significantly smaller (by a factor of >30) than the nuclear reorganization in the reaction (λ), meaning that structural and solvent fluctuations are required to bring the singlet and ${}^1\text{TT}$ states into resonance for rare electron tunneling events transforming reactant to product states. In other words, electron/phonon coupling is large by comparison

to electron/electron couplings between states such that $S_1 \rightarrow {}^1\text{TT}$ is best considered as an incoherent hopping process.

There is an issue that should be discussed at this point for the sake of completeness. Namely, we have previously argued for BT1 that λ for the diabatic $S_0S_1 \rightarrow {}^1\text{TT}$ may be larger, of order 0.5 eV.⁹ The origin of this prediction is in calculations we made using structures from DFT and TD-DFT with gradients, that predicted a significant intramolecular (inner-sphere) reorganization energy $\lambda_i = 0.43$ eV ($S_1 \rightarrow Q$). Given the current results, this may be an overestimation. A potential origin of this overestimation can be understood in the following way. In our hands, TD-DFT as applied to BT1 and related systems, including use of a toluene solvent continuum model, finds an optimized singlet excited state that is arm-localized. This is true not only for BT1 but also when acetylene substituents are added in respective TIPS-BT1 and TIPS-BT1' models. However, arm localization contradicts spectroscopic findings for TIPS-BT1¹³ where it is apparent that the singlet exciton state for the molecule in toluene is dimer delocalized. We surmise that λ_i would be smaller for a dimer-delocalized exciton compared to the arm-localized state found using TD-DFT, and additional theory is needed to explore this point. If a lower value of λ is operative as is now expected, then we also need to rationalize biexponential photoluminescence behavior observed for BT1.⁹ One reasonable explanation, given the poor solubility of BT1 that precluded exploration with TA in the first place, is that aggregation effects contribute to multiexponential decay behavior.

Comparing TIPS-BT1' with TIPS-BP1'. We were initially rather surprised by the overall finding that k_{fiss} for TIPS-BT1' ($2.0 \times 10^{11} \text{ s}^{-1}$) is similar to that of the larger and more exoergic TIPS-BP1' ($2.3 \times 10^{11} \text{ s}^{-1}$; *vide supra*). As noted earlier for TIPS-BP1', the $S_1 \rightarrow {}^1\text{TT}$ reaction driving force is substantial and expected to be in the -0.2 to -0.35 eV range.⁷ However, at the same time the reaction reorganization energy is expected to be smaller in TIPS-BP1' than the $\lambda = 0.18$ eV suggested above for TIPS-BT1' given the larger and more delocalized chromophores of the pentacene dimer. Thus, for TIPS-BP1', $S_1 \rightarrow {}^1\text{TT}$ conversion is likely to take place in the Marcus inverted region in contrast to the analogous reaction for TIPS-BT1', and this should contribute to reaction slowing, contrary to our initial assumption. Additionally, whereas the vibronic coupling theory mentioned above predicted diabatic coupling values of order $V_{\text{eff}} = 5.5$ meV for BT1, there is reason to expect it would be smaller in pentacene-based systems where exciton location from the perspective of the individual chromophores of the dimer is moved further away from the bridge linking the two. Qualitatively in support of this, we note our previous observation (*vide supra*) that Davydov splitting manifest in the UV is smaller for TIPS-BP1' (0.30 eV) than it is for TIPS-BT1' (0.47 eV). Factoring each of these things for TIPS-BP1', inverted region reactivity and smaller V_{eff} compared to TIPS-BT1', it is straightforward to come up with reasonable conditions that give $k_{\text{fiss}} = 2.3 \times 10^{11} \text{ s}^{-1}$ (an example of possible values is given in ref 46). However, given that each Marcus theory parameter is expected to change on going from TIPS-BT1' to TIPS-BP1', it is difficult to make specific predictions without further constraints that may come from theory and experiment. Nonetheless, we can emphasize at this point that Marcus theory readily describes the set of behaviors seen in these types of dimer systems. It is also our hope that such a parametric rate theory can be useful

comparing the behavior of other known systems. To this end, consistent treatments of diabatic coupling, reorganization energy, and driving force are needed.

Revisiting TIPS-BT1. As discussed in the Introduction, our published interpretation of TIPS-BT1 photophysics in toluene was that it did not engage in ^1TT formation and only decayed to ground state via k_r and k_{nr} .¹³ This was based primarily the lack of spectral evolution in the TA region (~ 420 nm) where there is a strong ESA attributed to the singlet exciton. In that published work, however, we did note a subtle (<10%) exponential decay of the singlet exciton feature in single-wavelength data ($\lambda_{\text{probe}} = 429$ nm) that was fit with an 850 fs time constant. While the chance of ^1TT involvement was discussed, it was ultimately dismissed given the stark time scale difference to our BT1 data⁹ and because the absence of spectral evolution argued against it. However, based on the findings herein for TIPS-BT1', it seems prudent to revisit these conclusions for TIPS-BT1. With the findings for TIPS-BT1' as a quantitative guide (vide supra), the ~10% decay of the S_1 magnitude in TIPS-BT1 at $\lambda_{\text{probe}} = 429$ nm is consistent with establishment of a $S_1 \rightleftharpoons ^1\text{TT}$ equilibrium, but one where the equilibrium constant is small at $K \sim 0.1$. Using this value in the framework of the three-state kinetic model (Figure 2d), S_1 would decay in 850 fs (~10% of signal) as observed if $k_{fiss} = 1.07 \times 10^{11} \text{ s}^{-1}$ (9.3 ps). This corresponds to an expected slowing relative to TIPS-BT1' ($k_{fiss} = 2.0 \times 10^{11} \text{ s}^{-1}$; 5 ps), consistent with the more endergonic driving force of 59 meV (to accommodate $K = 0.1$). Again Marcus theory is adequate for understanding these results. For example, if λ and V_{eff} are, respectively, held fixed at the previously discussed values of 0.18 eV and 5.5 meV, the time scale for ^1TT formation in TIPS-BT1 is predicted to be 18 ps; that is, of the right order of magnitude compared with the 9.3 ps time constant discussed above. Full agreement is achieved if V_{eff} is increased to 7.7 meV. An increase in V_{eff} for TIPS-BT1 relative to TIPS-BT1' appears to us reasonable, given that the position of the TIPS-acetylene groups influences where the exciton resides, from the perspective of each chromophore relative to the bridge. Qualitative support for a coupling increase is the stronger excitonic interaction observed in TIPS-BT1 compared to TIPS-BT1' based on vibronic features in the $S_1 \leftarrow S_0$ manifold (vide supra; Figure 3). It is also worth noting the possibility that these subtle side-group perturbations impact electronic coupling for SF (increase it in TIPS-BT1 compared to TIPS-BT1') in ways similar to those observed by Lukman, Musser, and co-workers.¹⁰ In their pentacene dimer systems, swapping TIPS-acetylene for mesityl side groups serves to increase the CT-state character in the adiabatic reactant singlet state with profound impacts on dynamics and mechanism. In our systems, it is possible that TIPS-BT1 has slightly more CT character in the reactant adiabatic singlet compared to TIPS-BT1'. As noted earlier (Table 1), the S_1 in TIPS-BT1 is modestly higher in energy, and this could facilitate more mixing with an otherwise isoenergetic higher-lying CT state. As well, the CT state could be lower in TIPS-BT1 compared to TIPS-BT1' if, as expected, the cation and anion charge densities exploit the acetylene side groups. In TIPS-BT1, the side groups on the opposing chromophores are physically closer to each other, and this would increase the Coulombic attraction between that cation and anion, thus stabilizing the CT. These issues may be factored in dimer design, but care must be taken to control relative reactant versus product energetics, as discussed extensively in the next section.

TIPS-BT1 versus TIPS-BT1'. As a final point of discussion, we consider how the subtle structural side-group changes that have been implemented manifest in the equilibrium shift from TIPS-BT1 ($K = 0.1$) to TIPS-BT1' ($K = 1$), recalling that this corresponds to a 59 meV exoergic shift for the $S_1 \rightarrow ^1\text{TT}$ photoreaction between these two dimers. Some of this could come from state energetics based on observations already discussed. As shown in Figure 3 and Table 1, the S_1 in TIPS-BT1 is slightly higher in energy compared to TIPS-BT1', by 10 meV. One potential origin of this has to do with electronic perturbations to the acene chromophores that arise from linear attachment to the bicyclic alkyl bridge. In the consideration of monomer models, we have previously shown that the electron-rich bridge serves to modestly destabilize S_1 and T_1 states relative to pure tetracene, presumably due to electron-donating properties of the bridge and their preferential impact on the acene LUMO.³⁴ In the context of the current dimers, it is reasonable to expect that the position of the TIPS-acetylene substituents will impact the S_1 energy and that this state will be higher for TIPS-BT1 because the acetylene substituents, which participate in determining the average position of the exciton, are closer to the destabilizing bridge. At first glance, the higher S_1 might appear to suggest that K would be larger in TIPS-BT1. Importantly, however, the same argument applies to the T_1 states; that is, more destabilization in TIPS-BT1 compared to TIPS-BT1'. Assuming the energy perturbation in the triplet manifold is similar to that of the S_1 ,³⁴ the $S_1 \rightarrow ^1\text{TT}$ photoreaction is expected to be more uphill for TIPS-BT1 compared to TIPS-BT1', given that the energy of the ^1TT is approximately twice the energy of the T_1 . However, the extent should be small, of order 10 meV, and while it can contribute, it does not appear significant enough to explain the equilibrium shift observations in total. We have also briefly considered an explanation based on S_1 energies. In principle, observed differences in excitonic interactions for TIPS-BT1 versus TIPS-BT1' would manifest in larger energy splitting (ΔE_{S_1}) between the higher energy optically bright S_1 and the lower energy dark S_1 . Given that the photoreaction of interest will occur primarily from the lower energy dark S_1 , the more excitonically coupled TIPS-BT1 could be preferentially disadvantaged. However, in order for this effect to meaningfully lower the equilibrium constant of interest, the difference in ΔE_{S_1} for TIPS-BT1 versus TIPS-BT1' (i.e., $\Delta \Delta E_{S_1}$) needs to be a substantial percentage of 59 meV. We do not think this is the case for these dimers. When considering Davydov splitting in the $S_3 \leftarrow S_0$ region, TIPS-BT1 exhibits a larger value (0.499 eV¹³) compared to TIPS-BT1' (0.472 eV) (see SI), but this represents a 5% difference out of ~0.5 eV of splitting. If we apply this percentage difference to the much smaller Davydov splitting expected for the $S_1 \leftarrow S_0$ transition in the visible (of order 30 meV³⁴), we find only 1.5 meV to work with. This is not enough to substantively impact the $S_1 \rightleftharpoons ^1\text{TT}$ equilibrium.

A final source of energy perturbation that intrigues us has to do with the biexcitonic TT manifold. As discussed recently by Greenham, Behrends, and co-workers in their electron spin resonance studies of singlet fission in TIPS-tetracene films, triplet interactions in biexciton states are dominated, not by dipolar coupling, but by exchange interactions.³⁸ The perturbation to the energies of the different state multiplicities that emerge, including the ^1TT , ^3TT , and ^5TT , depends on the extent to which relevant orbitals in the individual chromophore triplets share common space. Unlike dipolar coupling, exchange interactions can account for significant amounts of

energy, of order eV, when the extent of common orbital space is extensive as it is in individual acenes; that is, the reason they are useful for SF problems. Thus, even if common orbital space is not large, as one might expect for two acene chromophores juxtaposed relative to one another across a bridge, we suspect that it is not unreasonable to obtain the 10s of meV contributions needed to shift the $S_1 \rightleftharpoons ^1\text{TT}$ equilibrium between the two dimers. This would occur by utilizing a combination of through-space interactions as well as through-bond pathways mediated by the norbornyl-bridge σ and σ^* system. Such pathways are known to be effective for coupling π -chromophore systems in both electron-and energy-transfer problems.^{42,47–49} In order for this exchange effect to contribute to the observations in our dimer systems, the sign of the TT exchange interaction needs to be controlled such that ^1TT is destabilized at the same time that ^5TT is stabilized. This is the same direction one would expect for Hund's rule. In TIPS-BT1' where the acetylene substituents draw the two triplet excitons further away from one another, exchange interactions would decrease, leading to smaller energy splitting between ^1TT and ^5TT and less energetic cost to populating the ^1TT from the S_1 as we have observed. On the other hand, in TIPS-BT1 where the position of the acetylene substituents favors stronger exchange interactions in the TT manifold, the ^1TT would be pushed to higher energy thus decreasing its relative population within $S_1 \rightleftharpoons ^1\text{TT}$ equilibrium, again consistent with our observations. High level electronic structure theory is now needed to confirm the sign of the exchange interaction and to determine the magnitude of the effect in these systems.

CONCLUSION

In these studies we have focused on two structurally well-defined acene dimers for exploration of excited-state dynamics tied to singlet fission. Our emphasis has been on understanding time scales for formation of the multiexcitonic ^1TT state as well as its loss to the ground state either directly or via pathways involving reformation and decay of the singlet exciton state. The first dimer system, TIPS-BP1', is pentacenic in nature such that ^1TT formation is exoergic and seen to be efficient with \sim unit quantum yield. The second of these systems, TIPS-BT1', is tetracenic and is a close constitutional isomer of a dimer recently studied by our group called TIPS-BT1. The two differ only in the placement of solubilizing TIPS-acetylene side groups. They are energetically quite similar, as borne out using static absorption and emission spectroscopies, and yet they exhibit markedly different evolution of TA features including strong evidence in TIPS-BT1' for the rapid emergence of significant ^1TT population.

There are several notable individual findings that are summarized below. However, we first emphasize the general conclusion that in this class of pentacenic and tetracenic dimer systems, where structural definition is by design, we have achieved a unifying understanding of dynamics in terms of the few-parameter rate constant expression of Marcus theory. This allows us to assess appropriate magnitudes for diabatic coupling, reorganization energy λ , and driving force that enables efficient ^1TT formation in these and related systems. The overall mechanistic understanding means that these systems can provide benchmarks upon which subsequent variations that alter structure, energetics, and symmetry can be judged.

The first notable specific finding concerns TIPS-BT1' where we observe rapid formation of the ^1TT ($\tau_{\text{fiss}} = 5$ ps) in concert with establishment of an excited-state equilibrium of equal proportions ($K \sim 1$) with the singlet exciton state S_1 that resides 2.3 eV above the ground state. The established equilibrium means that the ^1TT resides at a highly similar energy. This speed is initially surprising given the absence of reaction driving force and given the unfavorable structural symmetry in this dimer (a long-axis reflection plane) expected to limit diabatic coupling between reactant and product.^{25,34} However, we conclude that we have the framework to rationalize this time constant. Theory we previously applied to the parent norbornyl-bridged tetracene dimer BT1, that factors vibronic coupling through symmetry-breaking vibrational motions (normal modes within the A_2 and B_2 irreducible representations), predicts an effective diabatic coupling V_{eff} of order 5.5 meV.²⁵ Such an amount, while appearing to be small, can accommodate $\tau_{\text{fiss}} = 5$ ps without a driving force (appropriate because $K \sim 1$) when the reorganization energy is low, but entirely reasonable, at $\lambda = 0.18$ eV. Subsequent theory would be useful to refine these numbers, but it is becoming clear that only modest diabatic couplings are needed to enable efficient ^1TT formation in competition to other radiative and nonradiative decay pathways, in large part because of the small reorganization energies associated with highly delocalized acetylene-substituted acene chromophores engaging in SF. A final point is made about TIPS-BT1' in relation to the lifetime of the ^1TT that might be relied upon for subsequent generation of states like the ^5TT or separated triplets. In this tetracenic system, the ^1TT energy is poised to limit the nonradiative decay to ground state (encompassed in the rate constant k_{TT}) compared to pentacenic systems like TIPS-BP1' that exhibit ^1TT lifetimes of order 100 ns. Unfortunately, excited-state equilibrium with the singlet exciton state undermines this potential gain.

The second notable specific finding concerns the observation that ^1TT formation in the pentacenic TIPS-BP1' (4.4 ps) is not substantially faster than in TIPS-BT1' (5.0 ps) despite the significantly larger (exergonic) reaction driving force of 200–350 meV (giving the ^1TT an energy above the ground state of \sim 1.58–1.73 eV). This can be partially understood now in the context of Marcus theory where the reaction in TIPS-BP1' should be slowed by placement in the inverted region. However, other effects are also expected to be in play. Namely, we anticipate reductions in both λ and V_{eff} for the more π -delocalized and excitonically separated TIPS-BP1' relative to TIPS-BT1' to contribute to the observed similarity in ^1TT formation rate constants.

The final notable specific finding concerns the comparison between TIPS-BT1' and the close constitutional isomer TIPS-BT1 and the fact that despite nearly identical singlet exciton energies, these two molecules exhibit markedly different ^1TT yields. We are intrigued by the possibility that we are observing the effect of exchange interactions between triplets in the multiexcitonic TT manifold where subtle structural changes, that is, the placement of the TIPS-acetylene substituents in TIPS-BT1' versus TIPS-BT1, are controlling its magnitude and where the comparative observation is revealing its sign. The ^1TT yields in TIPS-BT1' versus TIPS-BT1 are consistent with a scenario where exchange interactions raise the energy of the ^1TT relative to higher multiplicities ^3TT and ^5TT . In TIPS-BT1', the relative placement of the acetylene side groups draws the triplet excitons further away from one another,

thereby lowering the overall energy of the ^1TT and enabling its substantial participation ($K \sim 1$) in equilibrium with the S_1 singlet exciton state. The mechanistic details revealed in these comparative studies can be used in the design and interpretation of new systems and architectures to exploit the ^1TT as a gateway to the ^5TT or separate triplets.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/jacs.9b00904](https://doi.org/10.1021/jacs.9b00904).

Experimental methods and analysis details along with additional data Figures (S1–S15) and Table S1 ([PDF](#))

AUTHOR INFORMATION

Corresponding Author

*niels.damrauer@colorado.edu

ORCID

Tarek Sammakia: [0000-0003-3703-5672](https://orcid.org/0000-0003-3703-5672)

Niels H. Damrauer: [0000-0001-8337-9375](https://orcid.org/0000-0001-8337-9375)

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We acknowledge support from the National Science Foundation through grant CHE-1665375. In certain cases A.T.G. utilized resources at the National Renewable Energy Laboratory, and we thank Dr. Justin Johnson and Dr. Melissa Gish for assistance. This manuscript is based upon work conducted while T.S. was supported by, and serving at, the National Science Foundation.

REFERENCES

- Smith, M. B.; Michl, J. Singlet Fission. *Chem. Rev.* **2010**, *110*, 6891–6936.
- Smith, M. B.; Michl, J. Recent Advances in Singlet Fission. *Annu. Rev. Phys. Chem.* **2013**, *64*, 361–386.
- Hanna, M. C.; Nozik, A. J. Solar Conversion Efficiency of Photovoltaic and Photoelectrolysis Cells with Carrier Multiplication Absorbers. *J. Appl. Phys.* **2006**, *100*, 074510.
- Müller, A. M.; Avlasevich, Y. S.; Schoeller, W. W.; Müllen, K.; Bardeen, C. J. Exciton Fission and Fusion in Bis(Tetracene) Molecules with Different Covalent Linker Structures. *J. Am. Chem. Soc.* **2007**, *129*, 14240–14250.
- Sanders, S. N.; Kumarasamy, E.; Pun, A. B.; Trinh, M. T.; Choi, B.; Xia, J. L.; Taffet, E. J.; Low, J. Z.; Miller, J. R.; Roy, X.; Zhu, X. Y.; Steigerwald, M. L.; Sfeir, M. Y.; Campos, L. M. Quantitative Intramolecular Singlet Fission in Bipentacenes. *J. Am. Chem. Soc.* **2015**, *137*, 8965–8972.
- Lukman, S.; Musser, A. J.; Chen, K.; Athanasopoulos, S.; Yong, C. K.; Zeng, Z. B.; Ye, Q.; Chi, C. Y.; Hodgkiss, J. M.; Wu, J. S.; Friend, R. H.; Greenham, N. C. Tuneable Singlet Exciton Fission and Triplet-Triplet Annihilation in an Orthogonal Pentacene Dimer. *Adv. Funct. Mater.* **2015**, *25*, 5452–5461.
- Zirzlmeier, J.; Lehnher, D.; Coto, P. B.; Chernick, E. T.; Casillas, R.; Basel, B. S.; Thoss, M.; Tykwienski, R. R.; Guldi, D. M. Singlet Fission in Pentacene Dimers. *Proc. Natl. Acad. Sci. U. S. A.* **2015**, *112*, 5325–5330.
- Korovina, N. V.; Das, S.; Nett, Z.; Feng, X.; Joy, J.; Haiges, R.; Krylov, A. I.; Bradforth, S. E.; Thompson, M. E. Singlet Fission in a Covalently Linked Cofacial Alkynyltetracene Dimer. *J. Am. Chem. Soc.* **2016**, *138*, 617–627.
- Cook, J.; Carey, T. J.; Damrauer, N. H. Solution-Phase Singlet Fission in a Structurally Well-Defined Norbornyl-Bridged Tetracene Dimer. *J. Phys. Chem. A* **2016**, *120*, 4473–4481.
- Lukman, S.; Chen, K.; Hodgkiss, J. M.; Turban, D. H. P.; Hine, N. D. M.; Dong, S. Q.; Wu, J. S.; Greenham, N. C.; Musser, A. J. Tuning the Role of Charge-Transfer States in Intramolecular Singlet Exciton Fission through Side-Group Engineering. *Nat. Commun.* **2016**, *7*, 13622.
- Sakuma, T.; Sakai, H.; Araki, Y.; Mori, T.; Wada, T.; Tkachenko, N. V.; Hasobe, T. Long-Lived Triplet Excited States of Bent-Shaped Pentacene Dimers by Intramolecular Singlet Fission. *J. Phys. Chem. A* **2016**, *120*, 1867–1875.
- Zirzlmeier, J.; Casillas, R.; Reddy, S. R.; Coto, P. B.; Lehnher, D.; Chernick, E. T.; Papadopoulos, I.; Thoss, M.; Tykwienski, R. R.; Guldi, D. M. Solution-Based Intramolecular Singlet Fission in Cross-Conjugated Pentacene Dimers. *Nanoscale* **2016**, *8*, 10113–10123.
- Cook, J. D.; Carey, T. J.; Arias, D. H.; Johnson, J. C.; Damrauer, N. H. Solvent-Controlled Branching of Localized Versus Delocalized Singlet Exciton States and Equilibration with Charge Transfer in a Structurally Well-Defined Tetracene Dimer. *J. Phys. Chem. A* **2017**, *121*, 9229–9242.
- Kumarasamy, E.; Sanders, S. N.; Tayebjee, M. J. Y.; Asadpoordarvish, A.; Hele, T. J. H.; Fuemmeler, E. G.; Pun, A. B.; Yablon, L. M.; Low, J. Z.; Paley, D. W.; Dean, J. C.; Choi, B.; Scholes, G. D.; Steigerwald, M.; Ananth, N.; McCamey, D. R.; Sfeir, M. Y.; Campos, L. M. Tuning Singlet Fission in Pi-Bridge-Pi Chromophores. *J. Am. Chem. Soc.* **2017**, *139*, 12488–12494.
- Basel, B. S.; Zirzlmeier, J.; Hetzer, C.; Phelan, B. T.; Krzyaniak, M. D.; Reddy, S. R.; Coto, P. B.; Horwitz, N. E.; Young, R. M.; White, F. J.; Hampel, F.; Clark, T.; Thoss, M.; Tykwienski, R. R.; Wasielewski, M. R.; Guldi, D. M. Unified Model for Singlet Fission within a Non-Conjugated Covalent Pentacene Dimer. *Nat. Commun.* **2017**, *8*, 15171.
- Dean, J. C.; Zhang, R.; Hallani, R. K.; Pensack, R. D.; Sanders, S. N.; Oblinsky, D. G.; Parkin, S. R.; Campos, L. M.; Anthony, J. E.; Scholes, G. D. Photophysical Characterization and Time-Resolved Spectroscopy of a Anthradithiophene Dimer: Exploring the Role of Conformation in Singlet Fission. *Phys. Chem. Chem. Phys.* **2017**, *19*, 23162–23175.
- Yamakado, T.; Takahashi, S.; Watanabe, K.; Matsumoto, Y.; Osuka, A.; Saito, S. Conformational Planarization versus Singlet Fission: Distinct Excited-State Dynamics of Cyclooctatetraene-Fused Acene Dimers. *Angew. Chem., Int. Ed.* **2018**, *57*, 5438–5443.
- Carey, T. J.; Miller, E. G.; Gilligan, A. T.; Sammakia, T.; Damrauer, N. H. Modular Synthesis of Rigid Polyacene Dimers for Singlet Fission. *Org. Lett.* **2018**, *20*, 457–460.
- Basel, B. S.; Zirzlmeier, J.; Hetzer, C.; Reddy, S. R.; Phelan, B. T.; Krzyaniak, M. D.; Volland, M. K.; Coto, P. B.; Young, R. M.; Clark, T.; Thoss, M.; Tykwienski, R. R.; Wasielewski, M. R.; Guldi, D. M. Evidence for Charge-Transfer Mediation in the Primary Events of Singlet Fission in a Weakly Coupled Pentacene Dimer. *Chem.* **2018**, *4*, 1092–1111.
- Korovina, N. V.; Joy, J.; Feng, X. T.; Feltenberger, C.; Krylov, A. I.; Bradforth, S. E.; Thompson, M. E. Linker-Dependent Singlet Fission in Tetracene Dimers. *J. Am. Chem. Soc.* **2018**, *140*, 10179–10190.
- Margulies, E. A.; Miller, C. E.; Wu, Y.; Ma, L.; Schatz, G. C.; Young, R. M.; Wasielewski, M. R. Enabling Singlet Fission by Controlling Intramolecular Charge Transfer in π -Stacked Covalent Terrylenediimide Dimers. *Nat. Chem.* **2016**, *8*, 1120–1125.
- Johnson, J. C.; Akdag, A.; Zamadar, M.; Chen, X.; Schwerin, A. F.; Paci, I.; Smith, M. B.; Havlas, Z.; Miller, J. R.; Ratner, M. A.; Nozik, A. J.; Michl, J. Toward Designed Singlet Fission: Solution Photophysics of Two Indirectly Coupled Covalent Dimers of 1,3-Diphenylisobenzofuran. *J. Phys. Chem. B* **2013**, *117*, 4680–4695.
- Schrauben, J. N.; Akdag, A.; Wen, J.; Havlas, Z.; Ryerson, J. L.; Smith, M. B.; Michl, J.; Johnson, J. C. Excitation Localization/ Delocalization Isomerism in a Strongly Coupled Covalent Dimer of 1,3-Diphenylisobenzofuran. *J. Phys. Chem. A* **2016**, *120*, 3473–3483.

(24) Sanders, S. N.; Kumarasamy, E.; Pun, A. B.; Steigerwald, M. L.; Sfeir, M. Y.; Campos, L. M. Intramolecular Singlet Fission in Oligoacene Heterodimers. *Angew. Chem., Int. Ed.* **2016**, *55*, 3373–3377.

(25) Alguire, E. C.; Subotnik, J. E.; Damrauer, N. H. Exploring Non-Condon Effects in a Covalent Tetracene Dimer: How Important Are Vibrations in Determining the Electronic Coupling for Singlet Fission? *J. Phys. Chem. A* **2015**, *119*, 299–311.

(26) Sanders, S. N.; Kumarasamy, E.; Pun, A. B.; Appavoo, K.; Steigerwald, M. L.; Campos, L. M.; Sfeir, M. Y. Exciton Correlations in Intramolecular Singlet Fission. *J. Am. Chem. Soc.* **2016**, *138*, 7289–7297.

(27) Tayebjee, M. J. Y.; Sanders, S. N.; Kumarasamy, E.; Campos, L. M.; Sfeir, M. Y.; McCamey, D. R. Quintet Multiexciton Dynamics in Singlet Fission. *Nat. Phys.* **2017**, *13*, 182–188.

(28) Trinh, M. T.; Pinkard, A.; Pun, A. B.; Sanders, S. N.; Kumarasamy, E.; Sfeir, M. Y.; Campos, L. M.; Roy, X.; Zhu, X. Y. Distinct properties of the triplet pair state from singlet fission. *Sci. Adv.* **2017**, *3*, e1700241.

(29) Fuemmeler, E. G.; Sanders, S. N.; Pun, A. B.; Kumarasamy, E.; Zeng, T.; Miyata, K.; Steigerwald, M. L.; Zhu, X. Y.; Sfeir, M. Y.; Campos, L. M.; Ananth, N. A Direct Mechanism of Ultrafast Intramolecular Singlet Fission in Pentacene Dimers. *ACS Cent. Sci.* **2016**, *2*, 316–324.

(30) Berkelbach, T. C.; Hybertsen, M. S.; Reichman, D. R. Microscopic Theory of Singlet Exciton Fission. II. Application to Pentacene Dimers and the Role of Superexchange. *J. Chem. Phys.* **2013**, *138*, 114103.

(31) Yost, S. R.; Lee, J.; Wilson, M. W. B.; Wu, T.; McMahon, D. P.; Parkhurst, R. R.; Thompson, N. J.; Congreve, D. N.; Rao, A.; Johnson, K.; Sfeir, M. Y.; Bawendi, M. G.; Swager, T. M.; Friend, R. H.; Baldo, M. A.; Van Voorhis, T. A Transferable Model for Singlet-Fission Kinetics. *Nat. Chem.* **2014**, *6*, 492–497.

(32) Liu, H. Y.; Wang, R.; Shen, L.; Xu, Y. Q.; Xiao, M.; Zhang, C. F.; Li, X. Y. A Covalently Linked Tetracene Trimer: Synthesis and Singlet Exciton Fission Property. *Org. Lett.* **2017**, *19*, 580–583.

(33) Liu, H. Y.; Wang, Z. W.; Wang, X. M.; Shen, L.; Zhang, C. F.; Xiao, M.; Li, X. Y. Singlet exciton fission in a linear tetracene tetramer. *J. Mater. Chem. C* **2018**, *6*, 3245–3253.

(34) Vallett, P. J.; Snyder, J. L.; Damrauer, N. H. Tunable Electronic Coupling and Driving Force in Structurally Well-Defined Tetracene Dimers for Molecular Singlet Fission: A Computational Exploration Using Density Functional Theory. *J. Phys. Chem. A* **2013**, *117*, 10824–10838.

(35) Damrauer, N. H.; Snyder, J. L. Symmetry-Directed Control of Electronic Coupling for Singlet Fission in Covalent Bis-Acene Dimers. *J. Phys. Chem. Lett.* **2015**, *6*, 4456–4462.

(36) Pace, N. A.; Arias, D. H.; Granger, D. B.; Christensen, S.; Anthony, J. E.; Johnson, J. C. Dynamics of Singlet Fission and Electron Injection in Self-Assembled Acene Monolayers on Titanium Dioxide. *Chem. Sci.* **2018**, *9*, 3004–3013.

(37) Bayliss, S. L.; Weiss, L. R.; Rao, A.; Friend, R. H.; Chepelianskii, A. D.; Greenham, N. C. Spin signatures of exchange-coupled triplet pairs formed by singlet fission. *Phys. Rev. B: Condens. Matter Mater. Phys.* **2016**, *94*, 045204.

(38) Weiss, L. R.; Bayliss, S. L.; Krafft, F.; Thorley, K. J.; Anthony, J. E.; Bittl, R.; Friend, R. H.; Rao, A.; Greenham, N. C.; Behrends, J. Strongly exchange-coupled triplet pairs in an organic semiconductor. *Nat. Phys.* **2017**, *13*, 176–181.

(39) Burdett, J. J.; Piland, G. B.; Bardeen, C. J. Magnetic field effects and the role of spin states in singlet fission. *Chem. Phys. Lett.* **2013**, *585*, 1–10.

(40) Lewis, J. E.; Maroncelli, M. On the (Uninteresting) Dependence of the Absorption and Emission Transition Moments of Coumarin 153 on Solvent. *Chem. Phys. Lett.* **1998**, *282*, 197.

(41) Sens, R.; Drexhage, K. H. Fluorescence Quantum Yield of Oxazine and Carbazine Laser-Dyes. *J. Lumin.* **1981**, *24–25*, 709–712.

(42) Scholes, G. D.; Ghiggino, K. P.; Oliver, A. M.; Paddon-Row, M. N. Through-Space and Through-Bond Effects on Exciton Interactions in Rigidly Linked Dinaphthyl Molecules. *J. Am. Chem. Soc.* **1993**, *115*, 4345–4349.

(43) Spano, F. C. The Spectral Signatures of Frenkel Polarons in H- and J-Aggregates. *Acc. Chem. Res.* **2010**, *43*, 429–439.

(44) It is noted that Saito and co-workers report a nearly quantitative triplet yield of 180%. In our view, this is difficult to reconcile with the reported spectra that show a significant ESA band at ~ 460 nm as well as significant stimulated emission at ~ 610 nm, thus signifying the presence of a large amount of excitonic singlet. These workers mention that to account for the dissociation of triplets into independent species from the ^1TT , the calculated excited-state concentration must be doubled. We believe that this step is unnecessary and that it leads to a triplet yield overestimation while significant amounts of singlets remain. We suspect that the triplet yield in Saito's system is likely closer to the 100% that we report here.

(45) As described later, we find $K \sim 0.1$ in TIPS-BT1, and this means that we are underestimating these k_r and k_{nr} values by a small amount in this molecule by assuming K is very small, as per our original interpretation (cf. 13). We recover the observed decay with the $K = 0.1$ in place by setting $k_{\text{tot}} = k_r + k_{nr} = 4.5 \times 10^7$ rather than 4.2×10^7 . But this translates to a modest difference in lifetime (22.2 ns versus 23.8 ns) and the overall conclusion holds.

(46) For example, setting $\Delta G = -0.34$ eV (cf. 7) while leaving $\lambda = 0.18$ eV, one obtains this value of k_{fiss} when V_{eff} is modestly decreased from 5.5 to 4.8 meV.

(47) Paddon-Row, M. N.; Shephard, M. J. Through-bond orbital coupling, the parity rule, and the design of "superbridges" which exhibit greatly enhanced electronic coupling: A natural bond orbital analysis. *J. Am. Chem. Soc.* **1997**, *119*, 5355–5365.

(48) Clayton, A. H. A.; Scholes, G. D.; Ghiggino, K. P.; Paddon-Row, M. N. Through-Bond and Through-Space Coupling in Photoinduced Electron and Energy Transfer: An ab Initio and Semiempirical Study. *J. Phys. Chem.* **1996**, *100*, 10912–10918.

(49) Scholes, G. D.; Ghiggino, K. P.; Oliver, A. M.; Paddon-Row, M. N. Intramolecular Electronic-Energy Transfer between Rigidly Linked Naphthalene and Anthracene Chromophores. *J. Phys. Chem.* **1993**, *97*, 11871–11876.