

KOSZUL PROPERTIES OF THE MOMENT MAP OF SOME CLASSICAL REPRESENTATIONS

ALDO CONCA, HANS-CHRISTIAN HERBIG, AND SRIKANTH B. IYENGAR

ABSTRACT. This work concerns the moment map μ associated with the standard representation of a classical Lie algebra. For applications to deformation quantization it is desirable that $S/(\mu)$, the coordinate algebra of the zero fibre of μ , be Koszul. The main result is that this algebra is not Koszul for the standard representation of \mathfrak{sl}_n , and of \mathfrak{sp}_n . This is deduced from a computation of the Betti numbers of $S/(\mu)$ as an S -module, which are of interest also from the point of view of commutative algebra.

1. INTRODUCTION

If one squeezes a nice and regular object into a mold that is too small, the object will crack and fold and exhibit irregularities. This is precisely what happens when one is representing a reductive complex Lie algebra \mathfrak{g} on a low dimensional complex vector space V . The irregularities become apparent when studying the *moment map* $\mu: V \times V^* \rightarrow \mathfrak{g}^*$ of the representation; see Section 2. When $\dim(V)$ is small the zero fibre $\mu^{-1}(0)$ cannot have codimension $\dim(\mathfrak{g})$, which translates to the statement that the moment map cannot be a complete intersection in $V \times V^*$.

When V is spacious enough to properly host \mathfrak{g} , the moment map has a lot of good features. Tools to measure the size of the representation have been developed by Schwarz [22, 21] in connection with the *modularity*, also known as modality [25]. Specifically, the moment map is a complete intersection if and only if the representation is *0-modular*. A stronger condition is *1-largeness*; then $\mu^{-1}(0)$ is irreducible and the ideal (μ) generated in $S := \mathbb{C}[V \times V^*]$ by the components of μ is radical. If the representation is even *2-large*, then $\mu^{-1}(0)$ is also a normal variety. When \mathfrak{g} is simple, all but finitely many representations are 2-large; for details, see [12].

This work reported in this paper concerns the Koszul property for the algebra $S/(\mu)$ and the following question:

For which representations $\mathfrak{g} : V$ is the algebra $S/(\mu)$ Koszul?

The motivation to study this question comes from the Batalin-Fradkin-Vilkovisky approach to symplectic reduction [11]. It has been observed by the second author that the BFV-quantization scheme of [5] can be generalized to the situation when $S/(\mu)$ is a Koszul algebra. The main reason for this is that in the Tate model [24] of $S/(\mu)$ over S , the internal degree is linked to homological degree; see [3]. In the non-Koszul case however it appears that the BFV-quantization is unfeasible as the degrees proliferate which makes it impossible to control anomalous terms.

Date: May 13, 2018.

2010 Mathematics Subject Classification. 13D02, 16S37, 53D20.

Key words and phrases. Betti number, classical Lie algebra, Koszul algebra, moment map, Poincaré series, standard representation.

When $\mathfrak{g} : V$ is 0-modular, the ideal (μ) is a complete intersection of quadrics and then it is well known that $S/(\mu)$ is a Koszul algebra; see, for example, [18]. So the question is moot only for representations that are not 0-modular. For any representation $\mathfrak{g} : V$, the ideal (μ) is homogeneous quadratic, which is a necessary for $S/(\mu)$ to be Koszul. However, this condition is far from sufficient, and detecting when an algebra is Koszul is a notoriously difficult problem.

In this paper we develop methods to test when the algebra $S/(\mu)$ is *not* Koszul. We focus on the smallest (that is to say, the most ‘squeezed’) nontrivial representations: the standard representations of the classical Lie algebras $\mathfrak{gl}_n, \mathfrak{sl}_n, \mathfrak{so}_n, \mathfrak{sp}_n$. Here is summary of our results.

Theorem 1.1. *Let $\mathfrak{g} : V$ denote the standard representation of the Lie algebras considered above, and let $R = S/(\mu)$ be the coordinate algebra of the zero fibre. The following statements hold for each $n \geq 1$.*

- (1) *For $\mathfrak{g} = \mathfrak{gl}_n$ the algebra R is Koszul, $\dim R = n$ and $\operatorname{depth} R = 1$.*
- (2) *For $\mathfrak{g} = \mathfrak{so}_n$ the algebra R is Koszul and $\dim R = n + 1 = \operatorname{depth} R$.*
- (3) *For $\mathfrak{g} = \mathfrak{sl}_n$ the algebra R is not Koszul, $\dim R = n$ and $\operatorname{depth} R = 0$.*
- (4) *For $\mathfrak{g} = \mathfrak{sp}_n$ the algebra R is not Koszul, $\dim R = 2n$ and $\operatorname{depth} R = 0$.*

For $\mathfrak{gl}_n : \mathbb{C}^n$ the ideal (μ) is generated by quadratic monomials, and this implies the Koszul property. For $\mathfrak{so}_n : \mathbb{C}^n$ for $n \geq 3$ the components of the moment map (also known as angular momentum) are the 2×2 minors of a generic $2 \times n$ matrix and it is well-known that the corresponding algebra $S/(\mu)$ is Koszul. These assertions, and the claims about the dimension and depth, are justified in Section 4.

The proofs of the assertion that $S/(\mu)$ is *not* Koszul for $\mathfrak{sl}_n : \mathbb{C}^n$ and $\mathfrak{sp}_n : \mathbb{C}^{2n}$ for each $n \geq 2$ take up the bulk of this paper and require computing the Betti numbers of the corresponding algebra $S/(\mu)$, viewed as a module over the polynomial ring S ; see Sections 5 and 6. Another crucial input in the proofs are certain obstructions to the Koszul property in terms of Betti tables, described in Section 3.

The Betti tables of these algebras are also of interest from a purely commutative algebra point of view, for they exhibit intriguing combinatorial patterns; this is discussed in Section B of the Appendix.

There is another family of small representations for which $S/(\mu)$ is known not to be Koszul: the adjoint representations $\mathfrak{gl}_n : \mathfrak{gl}_n$ for $n \geq 3$. Here the zero locus of the moment map is essentially the *commuting variety*, that is to say, the variety of pairs of commuting matrices in $\mathfrak{gl}_n \times \mathfrak{gl}_n$. The first few syzygies have been calculated by F. Hreinsdottir [13] and it follows from Remark 3.6 that the corresponding algebra is not Koszul. A similar consideration also rules out $\mathfrak{so}_4 : 2k^4$ and $\mathfrak{sl}_3 : 2k^3$, for their Betti tables can be computed using, for example, Macaulay 2.

2. MOMENT MAPS

Let \mathfrak{g} be a finite dimensional reductive Lie algebra over the field k of characteristic zero and $\mathfrak{g} \rightarrow \mathfrak{gl}(V), \xi \mapsto (v \mapsto \xi.v)$ a finite dimensional representation of \mathfrak{g} ; we write $\mathfrak{g} : V$ to indicate this situation. The *moment map* associated to this representation is the bilinear form

$$\begin{aligned} \mu : V \times V^* &\longrightarrow \mathfrak{g}^* && \text{given by} \\ \mu(v, \alpha)(\xi) &:= \langle \alpha, \xi.v \rangle && \text{for } v \in V, \alpha \in V^* \text{ and } \xi \in \mathfrak{g}. \end{aligned}$$

Here $\langle \cdot, \cdot \rangle$ denotes the dual pairing between V^* and V . Its dual is the map

$$\begin{aligned} \mathfrak{g} &\longrightarrow V \times V^* \quad \text{with } \xi \mapsto \mu_\xi, \text{ where} \\ \mu_\xi(v, \alpha) &:= \langle \alpha, \xi \cdot v \rangle. \end{aligned}$$

The map $\xi \mapsto \mu_\xi$ extends to a Lie algebra homomorphism $\mathfrak{g} \rightarrow S := k[V \times V^*]$ when S is equipped with the unique Poisson bracket $\{ \cdot, \cdot \}$ such that

$$\{v, \alpha\} := \langle \alpha, v \rangle \quad \text{and} \quad \{v, v'\} := 0 =: \{\alpha, \alpha'\}$$

for all $v, v' \in V$ and $\alpha, \alpha' \in V^*$. We are interested in the ideal of S generated by the components of the moment map:

$$I := (\mu) = \{\mu_\xi \mid \xi \in \mathfrak{g}\}.$$

This is a subalgebra with respect to the Poisson bracket $\{ \cdot, \cdot \}$, but typically not an ideal with respect to the Poisson bracket. Since the μ_ξ are quadratic forms, I is homogenous, when S is viewed as a graded ring with $S_1 = V \times V^*$; in fact, there is a natural \mathbb{N}^2 -grading on S , and I is homogenous also with respect to this finer grading; this will prove to be useful in what follows.

Let us explain the coordinates expressions used for $V^* \times V$, \mathfrak{g} , S and μ .

2.1. Standard representation of \mathfrak{gl}_n . The Lie algebra \mathfrak{gl}_n is the space of $n \times n$ matrices over k , with bracket $[A, B] = AB - BA$. Let $\mathfrak{gl}_n : k^n$ be the representation given by the obvious action of \mathfrak{gl}_n on k^n .

With respect to the canonical bases e_1, e_2, \dots, e_n for $V = k^n$ and its dual basis f_1, f_2, \dots, f_n a point in $V \times V^*$ is identified with its linear coordinates

$$(p_1, p_2, \dots, p_n, q_1, q_2, \dots, q_n) = \sum q_i e_i + p_i f_i.$$

Accordingly we identify $k[V \times V^*]$ with

$$S := k \begin{bmatrix} p_1 & \dots & p_n \\ q_1 & \dots & q_n \end{bmatrix}.$$

For $i, j \in \{1, 2, \dots, n\}$, let E_{ij} denote the $n \times n$ -matrix with 1 in the i th row and j th column, and zero elsewhere. These matrices form a basis for \mathfrak{gl}_n , and it is not hard to verify that the components of the moment map are $p_i q_j$ for $i, j = 1, 2, \dots, n$.

2.2. Standard representation of \mathfrak{sl}_n . The Lie algebra \mathfrak{sl}_n is the Lie subalgebra of \mathfrak{gl}_n consisting of matrices of trace zero. The representation $\mathfrak{gl}_n : k^n$ from 2.1 thus restricts to one of \mathfrak{sl}_n . A basis for \mathfrak{sl}_n is given by $E_{i,j}$ where $1 \leq i, j \leq n$ with $i \neq j$, and $E_{i,i} - E_{i+1,i+1}$ for $i = 1, 2, \dots, n-1$. Hence the components of the moment map are $p_i q_j$ for $(1 \leq i, j \leq n \text{ with } i \neq j)$ and $p_i q_i - p_{i+1} q_{i+1}$ for $(i = 1, 2, \dots, n-1)$.

2.3. Standard representation of \mathfrak{so}_n . Recall that \mathfrak{so}_n is the Lie subalgebra of \mathfrak{gl}_n consisting of anti-symmetric matrices. The matrices $E_{i,j} - E_{j,i}$ where $1 \leq i < j \leq n$ are a basis for \mathfrak{so}_n . Restricting the standard representation of \mathfrak{gl}_n to \mathfrak{so}_n yields a moment map with components $p_i q_j - p_j q_i$ ($1 \leq i < j \leq n$), that is to say, the 2×2 -minors of the generic $2 \times n$ -matrix.

2.4. Standard representation of \mathfrak{sp}_n . The Lie algebra \mathfrak{sp}_n consists of $2n \times 2n$ -matrices A such that

$$A^t S = S A \quad \text{where } S = \begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix}$$

Here I_n is the $n \times n$ identity matrix. A basis for \mathfrak{sp}_n is given by the $2n \times 2n$ matrices

$$\begin{aligned} E_{i,j} - E_{n+j,n+i} & \quad \text{for } i, j = 1, \dots, n \\ E_{i,n+i} \text{ and } E_{n+i,i} & \quad \text{for } i = 1, \dots, n \\ E_{i,n+j} + E_{j,n+i} \text{ and } E_{n+i,j} + E_{n+j,i} & \quad \text{for } 1 \leq i < j \leq n \end{aligned}$$

We use $(p_{11}, \dots, p_{1n}, p_{21}, \dots, p_{2n}, q_{11}, \dots, q_{1n}, q_{21}, \dots, q_{2n})$ for linear coordinates for $V^* \times V$ and identify $k[V \times V^*]$ with

$$S := k \begin{bmatrix} p_{11} & \dots & p_{1n} & p_{21} & \dots & p_{2n} \\ q_{11} & \dots & q_{1n} & q_{21} & \dots & q_{2n} \end{bmatrix}$$

The corresponding components of the moment map are

$$\begin{aligned} p_{1i}q_{1j} - p_{2j}q_{1i} & \quad \text{for } i, j = 1, \dots, n \\ p_{1i}q_{2i} \text{ and } p_{2i}q_{1i} & \quad \text{for } i = 1, \dots, n \\ p_{1i}q_{2j} + p_{1j}q_{2i} \text{ and } p_{2i}q_{1j} + p_{2j}q_{1i} & \quad \text{for } 1 \leq i < j \leq n \end{aligned}$$

3. KOSZUL ALGEBRAS

In this section we recall the definition of certain invariants of modules over graded rings. The focus will be on Koszul algebras and some obstructions to this property.

3.1. Fine grading. The algebras and modules that we study will be equipped with a \mathbb{N}^2 -grading and the arguments exploit this structure. We prepare for this by introducing notation concerning \mathbb{N}^c -graded objects, for a positive integer $c \geq 1$.

Let X be an \mathbb{N}^c -graded set. When we speak of an element x in X , it should be understood that x is homogeneous: $x \in X_v$ for some $v \in \mathbb{N}^c$. Then v is the *degree* of x , that we denote $\deg x$. For any v in \mathbb{N}^c set

$$\|v\| := \sum_{i=1}^c v_i \quad \text{where } v = (v_1, \dots, v_c).$$

This is the *total degree* of v . When k is a field X is a \mathbb{N}^c -graded k -vector space, set

$$\text{Hilb}_X(s_1, \dots, s_c) := \sum_{v \in \mathbb{N}^c} \text{rank}_k(X_v) s_1^{v_1} \cdots s_c^{v_c},$$

the *Hilbert series* of X , viewed as an formal power series over \mathbb{Z} in indeterminates s_1, \dots, s_c . We will also need the version based on total degrees

$$\text{Hilb}_X(s) := \sum_{j \in \mathbb{N}} \text{rank}_k(X_j) s^j = \text{Hilb}_X(s, \dots, s).$$

3.2. Graded rings. Let k be a field (of arbitrary characteristic) and R a standard \mathbb{N}^c -graded k -algebra, meaning:

- (1) $R^0 = k$;
- (2) R is an \mathbb{N}^c -graded commutative k -algebra;
- (3) $\text{rank}_k(\bigoplus_{\|v\|=1} R_v)$ is finite
- (4) R is generated by elements in R_v with $\|v\| = 1$.

These conditions imply R is noetherian and that $\text{rank}_k R_v$ is finite for each $v \in \mathbb{N}^c$. This ring is also local, in the graded sense: The set $\{r \in R \mid \|\deg(r)\| \geq 1\}$ is the unique homogeneous maximal ideal of R .

Any finitely generated \mathbb{N}^c -graded R -module M is noetherian, and hence $\text{rank}_k M_v$ is finite for each $v \in \mathbb{N}^c$. The *socle* of M is the k -vector subspace

$$\{m \in M \mid r \cdot m = 0 \text{ for all } r \in R \text{ with } \|r\| \geq 1\}.$$

Let M, N be \mathbb{N}^c -graded R -modules. Each $\text{Tor}_i^R(M, N)$ is endowed with a \mathbb{N}^c -grading compatible with the action of R . Thus $\text{Tor}^R(M, N)$ is $(\mathbb{N} \times \mathbb{N}^c)$ -graded. We write $\| \cdot \|$ for the homological grading. For example, for any a in $\text{Tor}^R(M, N)$

$$|a| = i \quad \text{and} \quad \deg(a) = v \text{ means } a \in \text{Tor}_i^R(M, N)_v$$

where $v \in \mathbb{N}^c$. Then $\|\deg(a)\|$ is the total (internal) degree of a .

3.3. Betti numbers. Let M be a finitely generated \mathbb{N}^c -graded R -module. The *graded Betti numbers* of M are the integers

$$\beta_{i,v}^R(M) := \text{rank}_k \text{Tor}_i^R(k, M)_v \quad \text{for } i \in \mathbb{N} \text{ and } v \in \mathbb{N}^c$$

and the associated generating function

$$P_M^R(s_1, \dots, s_c, u) := \sum_{i,v} \beta_{i,v}^R(M) s_1^{v_1} \cdots s_c^{v_c} u^i,$$

viewed as an element in $\mathbb{Z}[s_1, \dots, s_c][[u]]$, is the *graded Poincaré series* of M . Often, one is interested only in homological degrees and the associated Poincaré series:

$$\beta_i^R(M) := \sum_{v \in \mathbb{N}^c} \beta_{i,v}^R(M) \quad \text{and} \quad P_M^R(u) := \sum_{i,j} \beta_i^R(M) u^i$$

In what follows, we often compute the Poincaré series of M as the Hilbert series of $\text{Ext}_R(M, k)$. The natural grading on this k -vector space is cohomological and so, when dealing with them, we will tacitly switch to this one, by setting

$$\text{Ext}_R^i(M, k)^v := \text{Ext}_R^i(M, k)_{-v},$$

for any finitely generated \mathbb{N}^c -graded R -module M .

We will also need to consider the following integers:

$$t_i^R(M) := \sup\{\|v\| \in \mathbb{N} \mid \text{Tor}_i^R(k, M)_v \neq 0\} \quad \text{for each } i \in \mathbb{N}.$$

This is the highest total degree of a Betti number of M in homological degree i .

3.4. Koszul algebras. When R is regular (that is to say, isomorphic to the symmetric k -algebra on R_1) one has $t_i^R(k) = i$ for $0 \leq i \leq \dim R$. Even when R is singular (meaning, not regular) it is not hard to see that

$$t_i^R(k) \geq i \quad \text{for } i \geq 0.$$

When equality holds for each i one says that the k -algebra R is *Koszul*. In this case, it is not hard to verify that there is an equality of formal power series

$$P_k^R(-u) \text{Hilb}_R(u) = 1.$$

It is known that the converse also holds; see, for example, [8, Theorem 1].

When R is Koszul, the relations defining R must be quadratic; in other words, there is an isomorphism of k -algebra $R \cong S/I$, where S is a standard graded polynomial ring over k and I can be generated by quadratic forms. The converse holds if I is monomial—see, for example, [7, Theorem 15]—but not in general.

Remark 3.5. If for some integer $s \geq 0$ one has $t_i^S(R) \leq i + 1$ for each $i \leq s$, then

$$t_i^R(k) \leq i \quad \text{for } i \leq s + 1.$$

Moreover, if $s = \text{pd}_S R$, the projective dimension of R over S , then R is Koszul.

These results are immediate from the graded version of a coefficientwise inequality [2, Proposition 3.3.2] due to Serre, relating the Poincaré series of k over R , and of k and R over S .

The next result is sometimes helpful in detecting when a k -algebra is *not* Koszul.

Remark 3.6. If R is Koszul, then $\beta_1^S(R) = \beta_1^S(R)_2$ and

$$\beta_i^S(R)_{2i} \leq \binom{\beta_1^S(R)}{i} \quad \text{for each integer } i \geq 1.$$

This inequality is due to Avramov, Conca, and Iyengar [4, Theorem 5.1].

A different obstruction to the Koszul property involves Massey products on the Koszul homology of R .

3.7. Matric Massey products. Let R be a standard \mathbb{N}^c -graded k -algebra. Let K be the Koszul complex of R and set $H := H(K)$. The canonical DG (= Differential Graded) R -algebra structure on K is compatible with the \mathbb{N}^c -grading. More precisely, for elements a, b in K one has

$$\begin{aligned} |ab| &= |a| + |b| \quad \text{and} \quad \deg(ab) = \deg(a) + \deg(b) \\ |d(a)| &= |a| - 1 \quad \text{and} \quad \deg(d(a)) = \deg(a). \end{aligned}$$

We recall the notion of Massey products on H , for which Kraines [15] is a good reference, keeping track of the finer grading involved. To begin with, for any element a in K , one sets

$$\bar{a} := (-1)^{|a|+1}a.$$

Let w_1, \dots, w_s be elements in $H_{\geq 1}$; as always, these are assumed to be homogenous with respect to the $(\mathbb{N} \times \mathbb{N}^c)$ -grading on H . The Massey product $\langle w_1, \dots, w_s \rangle$ is said to be *defined* if there is a collection of elements a_{ij} in K , for $1 \leq i < j \leq s$

and $(i, j) \neq (1, s)$ such that, for each i the element a_{ii} is cycle representing w_i and for each pair (i, j) as above, one has

$$(3.7.1) \quad d(a_{ij}) = \sum_{r=i}^{j-1} \bar{a}_{ir} a_{r+1,j}.$$

Such a collection $\{a_{ij}\}$ of elements is a *defining system* for $\langle w_1, \dots, w_s \rangle$. Given a defining system, it is readily verified that the element

$$(3.7.2) \quad a_{1s} := \sum_{r=1}^{s-1} \bar{a}_{1r} a_{r+1,s}$$

is a cycle in K . The Massey product $\langle w_1, \dots, w_s \rangle$, when it is defined, is the collection of all elements in H that can be represented by cycles of the form a_{1s} for some defining system $\{a_{ij}\}$ as above. Observe that these lie in $H_{p,v}$ where

$$p = \sum_{r=1}^s |w_i| + (s-2) \quad \text{and} \quad v = \sum_{r=1}^s \deg(w_i).$$

We need an extension of all this to matrices with coefficients in H . These *matric Massey products* were introduced by May [16] in the context of general DG algebras and brought to bear to study commutative rings (specifically, Koszul homology) by Avramov [1]. In this context, one considers matrices W_1, \dots, W_s with entries in H with appropriate restrictions on their sizes and the degrees (both homological and internal) of the entries in them to ensure that the products involved in (3.7.1) and (3.7.2) are permissible; each a_{ij} is replaced by a matrix A_{ij} with entries in K .

Elements of the matric Massey product $\langle W_1, \dots, W_s \rangle$, when it is defined, are matrices with entries in $H_{\geq 1}$. For any matrix W in $\langle W_1, \dots, W_s \rangle$, the degrees of $W_{i,j}$, the entry in position (i, j) of W , are given by

$$\begin{aligned} |W_{ij}| &= \sum_{r=1}^s |(W_r)_{i_{r-1}i_r}| + (s-2) \\ \deg(W_{ij}) &= \sum_{r=1}^s \deg((W_r)_{i_{r-1}i_r}) \end{aligned}$$

where $i_0 = i$ and $i_r = s$, and i_1, \dots, i_{s-1} are any permissible choice of integers; for example, $1, \dots, 1$.

An element w in H is *decomposable* as a matric Massey product if it belongs to $\langle W_1, \dots, W_s \rangle$ for some system of matrices with entries in $H_{\geq 1}$; necessarily, W_1 is then a row matrix and W_s is a column matrix.

Lemma 3.8. *Let R be a standard \mathbb{N}^c -graded k -algebra and H its Koszul homology. If s is an integer such that $t_i^R(k) \leq i$ for each $i \leq s$, then the elements of $H_{i,v}$ are decomposable as matric Massey products when $2 \leq i \leq s-1$ and $\|\deg v\| \geq i+2$.*

Proof. In what follows, we set

$$T := \text{Tor}^R(k, k)/(\text{Tor}_1^R(k, k)).$$

This inherits from $\text{Tor}^R(k, k)$ a structure of a $(\mathbb{N} \times \mathbb{N}^c)$ -graded k -algebra, and our hypothesis translates to $T_{i,v} = 0$ when $i \leq s$ and $\|\deg v\| \geq i+1$. The crux of the argument is that there is a k -linear map $\sigma: H_{\geq 1} \rightarrow T$ of homological degree one,

called the suspension; see, for example, [10, Definition 3.7]. This map respects the internal gradings, that is to say, for each $i \geq 1$ and $v \in \mathbb{N}$ one has

$$\sigma_{i,v}: H_{i,v} \longrightarrow T_{i+1,v}.$$

Thus, the hypothesis on then $t_i^R(k)$ entails $T_{i+1,v} = 0$ whenever $1 \leq i \leq s-1$ and $\|\deg v\| \geq i+2$. Therefore for these i, v one has $H_{i,v} \subseteq \text{Ker}(\sigma)$. It remains to note that $\text{Ker}(\sigma)$ consists precisely of elements decomposable as matric Massey products in $H_{\geq 1}$; see [10, Corollary 5.13]. \square

4. THE GENERAL LINEAR AND THE SPECIAL ORTHOGONAL CASE

In this section we describe the Betti numbers of the moment maps associated to canonical representation of the general linear, and the special orthogonal, Lie algebra. The moment maps in question are described in 2.1 and 2.3. These results are special cases of general results in commutative algebra. The main reason for describing the computations carefully is that they get used in the later sections.

4.1. The general linear group. Let k be a field and S the polynomial ring over k in indeterminates $\underline{p} := p_1, \dots, p_n$ and $\underline{q} := q_1, \dots, q_n$. Thus

$$S := k \begin{bmatrix} p_1 & \dots & p_n \\ q_1 & \dots & q_n \end{bmatrix}$$

We view S as an \mathbb{N}^2 -graded k -algebra with each p_i of degree $[1, 0]$ and each q_i of degree $[0, 1]$. Set

$$I := (\underline{p}) \cap (\underline{q}) \quad \text{and} \quad A = S/I.$$

Since I is generated by quadratic monomials, the k -algebra A is Koszul; see Definition 3.4. Next we compute the Betti numbers of A as an S -module.

As I is homogenous, the k -algebra A inherits the \mathbb{N}^2 -grading of S . Noting that $(\underline{p}) + (\underline{q})$ is the maximal ideal of S , the standard Mayer-Vietoris sequence for the ideals (\underline{p}) and (\underline{q}) reads

$$(4.1.1) \quad 0 \longrightarrow A \xrightarrow{\iota} C \xrightarrow{\pi} k \longrightarrow 0, \quad \text{where } C = S/(\underline{p}) \bigoplus S/(\underline{q}).$$

From this it follows that the bigraded Hilbert series of A is

$$(4.1.2) \quad \text{Hilb}_A(s, t) = \frac{1}{(1-s)^n} + \frac{1}{(1-t)^n} - 1$$

Next we compute the Poincaré series of A , viewed as an S -module.

4.2. Structure of $\text{Ext}_S(A, k)$ as module over $\text{Ext}_S(k, k)$. Set $\mathfrak{m} = (\underline{p}, \underline{q})$; this is the homogenous maximal ideal of S . The residue classes modulo \mathfrak{m}^2 of the sequence $\underline{p}, \underline{q}$ is a basis for the k -vector space $\mathfrak{m}/\mathfrak{m}^2$. Let $e_1, \dots, e_n, f_1, \dots, f_n$ be the dual basis, of the k -vector space

$$V := \text{Hom}_k(\mathfrak{m}/\mathfrak{m}^2, k).$$

We view this as a \mathbb{N}^2 -graded k -vector space with (upper grading) $\deg(e_i) = [1, 0]$ and $\deg(f_i) = [0, 1]$, in cohomological degree one. Then one has

$$\Lambda := \text{Ext}_S(k, k) \cong \wedge V \quad \text{and} \quad \text{Ext}_S(C, k) \cong \Lambda/(\underline{f}) \oplus \Lambda/(\underline{e}),$$

and the map $\text{Ext}_S(\pi, k)$, with π as in (4.1.1), is the canonical map

$$\Lambda \longrightarrow \Lambda/(\underline{f}) \oplus \Lambda/(\underline{e}).$$

It is easy to check that this map is surjective in homological degrees ≥ 1 ; for example, in degree one the map is an isomorphism and, as Λ -modules, the source and target are generated by their components in homological degree zero. It follows that (4.1.1) induces an exact sequence of graded k -vector spaces

$$(4.2.1) \quad 0 \longrightarrow \mathrm{Ext}_S^{\geq 1}(A, k) \longrightarrow \mathrm{Ext}_S^{\geq 2}(k, k)[0, 0, 1] \longrightarrow \mathrm{Ext}_S^{\geq 2}(C, k)[0, 0, 1] \longrightarrow 0.$$

Noting that $\mathrm{Ext}_S(k, k)$ is an exterior algebra on $2n$ generators, half of which have (upper) degree $(1, 0, 1)$ and the rest have degree $(0, 1, 1)$. Thus one gets

$$(4.2.2) \quad P_k^S(s, t, u) := (1+su)^n(1+tu)^n \quad \text{and} \quad P_C^S(s, t, u) := (1+su)^n + (1+tu)^n$$

This discussion leads to the following result.

Theorem 4.3. *Let k be a field and let S and A be the k -algebras introduced in 4.1. The trigraded Poincaré series of A over S is*

$$P_A^S(s, t, u) := 1 + u^{-1}((1+su)^n - 1)((1+tu)^n - 1).$$

The nonzero graded Betti numbers of A over S are

$$\beta_{i,v}^S(A) = \begin{cases} 1 & \text{for } i = 0 \text{ and } v = [0, 0] \\ \binom{n}{v_1} \binom{n}{v_2} & \text{for } i \geq 1, v_i \geq 1, \text{ and } v_1 + v_2 = i + 1 \end{cases}$$

The Poincaré series is $1 + u^{-1}((1+u)^n - 1)^2$. The Betti numbers of A over S are

$$\beta_i^S(A) = \begin{cases} 1 & \text{for } i = 0 \\ \binom{2n}{i+1} - 2\binom{n}{i+1} & \text{for } i \geq 1. \end{cases}$$

Moreover, $\dim A = n$ and $\mathrm{depth} A = 1$.

Proof. The claims about the Poincaré series are immediate from (4.2.1) and (4.2.2). The ones about the dimension and the depth of A are from (4.1.1). \square

For later use, we record that following result that is immediate from (4.2.1).

Lemma 4.4. *As a Λ -module, $\mathrm{Ext}_S^{\geq 1}(A, k)[0, 0, -1]$ is isomorphic to the ideal*

$$(e_j \wedge f_j \mid 1 \leq i, j \leq n) \text{ of } \Lambda$$

\square

This completes our discussion on the general linear Lie algebra.

4.5. Special orthogonal Lie algebra. Let k be a field and S the polynomial ring over k in indeterminates $\underline{p} := p_1, \dots, p_n$ and $\underline{q} := q_1, \dots, q_n$. Thus

$$S := k \begin{bmatrix} p_1 & \dots & p_n \\ q_1 & \dots & q_n \end{bmatrix}$$

As before, S is to be viewed as an \mathbb{N}^2 -graded k -algebra with each p_i of degree $[1, 0]$ and each q_i of degree $[0, 1]$. Let I be the 2×2 minors of the matrix above and set $A = S/I$. The bigraded Hilbert series of A is

$$\frac{1 - st \sum_{i=0}^{n-2} (-1)^i \binom{n}{2+i} h_i(s, t)}{(1-s)^n(1-t)^n}$$

where $h_i(s, t)$ is the sum of all monomials of degree i in s and t . In particular

$$\mathrm{Hilb}_A(s) = \frac{1 + s(n-1)}{(1-s)^{n+1}}.$$

The minimal free resolution of A over S is given by the Eagon-Northcott complex; see, for example, [6, Section 2C]. It yields $t_i^S(A) \leq i + 1$ for each i , and hence A is Koszul; see Remark 3.5. From this one gets that the Poincaré series of k over A is

$$P_k^A(u) = \frac{(1+u)^{n+1}}{1-u(n-1)}$$

5. THE SPECIAL LINEAR CASE

In this section we study the moment map associated to the canonical representation of the special linear Lie algebra, \mathfrak{sl}_n ; see 2.2.

Let k be a field; for what follows, its characteristic has to be zero, or at least larger than $(n+1)/2$, for we need Lemma A.1. As in Section 4, let S be the polynomial ring over k in indeterminates $\underline{p} := p_1, \dots, p_n$ in degree $[1, 0]$, and $\underline{q} := q_1, \dots, q_n$, in degree $[0, 1]$. Set

$$B = S/I \quad \text{where } I = (\{p_i q_j, p_i q_i - p_j q_j \mid 1 \leq i, j \leq n \text{ with } i \neq j\}).$$

The ideal I is homogenous, with respect to the \mathbb{N}^2 grading on S , and hence the k -algebra B inherits this grading.

It is straightforward to check that $p_1 q_1$ is in the socle of B and that $B/(p_1 q_1) = A$, the k -algebra encountered in Section 4. There is thus an exact sequence of \mathbb{N}^2 -graded S -modules

$$(5.0.1) \quad 0 \longrightarrow k[-1, -1] \xrightarrow{1 \mapsto p_1 q_1} B \longrightarrow A \longrightarrow 0$$

and from this one and (4.1.2) one obtains that the bigraded Hilbert series of B :

$$\text{Hilb}_B(s, t) = \frac{1}{(1-s)^n} + \frac{1}{(1-t)^n} - 1 + st.$$

Next we determine the Poincaré series of B , as an S -module. To this end we compute $\text{Ext}_S(B, k)$, exploiting the fact that this is a \mathbb{N}^3 -graded module over the \mathbb{N}^3 -graded k -algebra

$$\Lambda = \text{Ext}_S(k, k).$$

By (4.2.2) one gets

$$\text{Hilb}_\Lambda(s, t, u) = (1+su)^n(1+tu)^n$$

where u is the cohomological degree. The exact sequence (5.0.1) induces an exact sequence of graded Λ -modules

$$\longrightarrow \Lambda[-1, -1, -1] \xrightarrow{\beta} \text{Ext}_S(A, k) \longrightarrow \text{Ext}_S(B, k) \longrightarrow \Lambda[-1, -1, 0] \xrightarrow{\beta[1]} \longrightarrow$$

This gives an equality of formal power series

$$P_B^S(s, t, u) = \text{Hilb}_{\text{Coker}(\beta)}(s, t, u) + u^{-1} \cdot \text{Hilb}_{\text{Ker}(\beta)}(s, t, u).$$

It thus remains to compute the Hilbert series of $\text{Ker}(\beta)$ and $\text{Coker}(\beta)$.

Let C be the S -module $S/(p) \oplus S/(q)$, encountered in Section 4; see (4.1.1). The exact sequence (4.2.1), reproduced in the top row of the diagram below, is one of graded Λ -modules:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Ext}_S^{\geq 1}(A, k) & \xrightarrow{\gamma} & \Lambda[0, 0, 1] & \longrightarrow & \text{Ext}_S(C, k)[0, 0, 1] \longrightarrow 0 \\ & & \beta \uparrow & & \gamma \beta \uparrow & & \\ & & \Lambda[-1, -1, -1] & = & \Lambda[-1, -1, -1] & & \end{array}$$

The Snake Lemma then gives $\text{Ker}(\beta) = \text{Ker}(\gamma\beta)$ and an exact sequence

$$0 \longrightarrow \text{Coker}(\beta) \longrightarrow \text{Coker}(\gamma\beta) \longrightarrow \text{Ext}_S(C, k)[0, 0, 1] \longrightarrow 0,$$

of graded Λ -modules. It thus remains to study the map

$$\Lambda[-1, -1, -1] \xrightarrow{\gamma\beta} \Lambda[0, 0, 1]$$

This map is Λ -linear, so it is determined by $\gamma\beta(1)$, where $1 \in \Lambda$. To identify this element, as in Section 4, let $e_1, \dots, e_n, f_1, \dots, f_n$ be the basis of

$$\Lambda^1 = \text{Hom}_k(\mathfrak{m}/\mathfrak{m}^2, k)$$

dual to that of the images of $\underline{p}, \underline{q}$ in $\mathfrak{m}/\mathfrak{m}^2$; here $\mathfrak{m} = (\underline{p}, \underline{q})$.

Claim. $\gamma\beta(1) = \sum_{1 \leq i \leq n} e_i f_i$ in Λ^2 .

Indeed, $\beta(1)$ is the element in $\text{Ext}_S^1(A, k)$ corresponding to the exact sequence (5.0.1), and γ is the map defined by concatenation with the sequence (4.1.1). It follows that $\gamma\beta(1)$ is the element of Λ^2 corresponding to the exact sequence

$$(5.0.2) \quad 0 \longrightarrow k[-1, -1] \xrightarrow{1 \mapsto p_1 q_1} B \xrightarrow{\iota} C \xrightarrow{\pi} k \longrightarrow 0$$

To identify this element in Λ , let E be the Koszul complex of $\underline{p}, \underline{q}$. We view it as the exterior algebra $\bigwedge \bigoplus_i (SP_i \oplus SQ_i)$ with $P_i \mapsto p_i$ and $Q_i \mapsto q_i$. The first steps of a lifting of the natural morphism $E \rightarrow k$ to the complex above is given below:

$$\begin{array}{ccccccc} 0 & \longrightarrow & k[-1, -1] & \xrightarrow{1 \mapsto p_1 q_1} & B & \xrightarrow{\iota} & C \xrightarrow{\pi} k \longrightarrow 0 \\ & & \kappa_2 \uparrow & & \kappa_1 \uparrow & & \kappa_0 \uparrow \quad \parallel \\ \cdots & \longrightarrow & \bigoplus_{i,j} SP_i Q_j & \longrightarrow & \bigoplus_i (SP_i \oplus SQ_i) & \longrightarrow & S \longrightarrow k \longrightarrow 0 \end{array}$$

where the maps κ_i are given by $\kappa_0(1) = (0, 1)$, and

$$\begin{aligned} \kappa_1(P_i) &= p_i \quad \text{and} \quad \kappa_1(Q_i) = 0 \quad \text{for each } i \\ \kappa_2(P_i P_j) &= 0, \quad \kappa_2(Q_i Q_j) = 0, \quad \text{and} \quad \kappa_2(P_i Q_j) = 1 \quad \text{for each } i, j. \end{aligned}$$

The claim follows as (5.0.2) corresponds to the class of κ_2 in $\Lambda^2 = H_{-2}(\text{Hom}_S(E, k))$.

From the preceding claim and Proposition A.2, it follows that the map $\gamma\beta$ has maximal rank, and hence that

$$\begin{aligned} \text{Hilb}_{\text{Ker } \beta}(s, t, u) &= u[(stu^2 - 1)(1 + su)^n(1 + tu)^n]_+ \\ \text{Hilb}_{\text{Coker } \beta}(s, t, u) &= u[(1 - stu^2)(1 + su)^n(1 + tu)^n]_+ - u((1 + su)^n + (1 + tu)^n) \end{aligned}$$

where $[h(s, t, u)]_+$ denotes positive part of $h(s, t, u)$; namely, those terms whose coefficients are negative integers are omitted.

Summing up one gets the result below.

Theorem 5.1. *Let k be a field, and let S and B the k -algebras introduced at the beginning of this section; in particular, the characteristic of k is zero, or greater than $(n + 1)/2$. The trigraded Poincaré series of B over S is*

$$\begin{aligned} 2 - (1 + su)^n - (1 + tu)^n + & [(1 - stu^2)(1 + su)^n(1 + tu)^n]_+ \\ & + u^{-2}[(stu^2 - 1)(1 + su)^n(1 + tu)^n]_+ \end{aligned}$$

The Poincaré series is thus

$$1 - 2[(1 + u)^n - 1] + u^{-1}[(1 - u^2)(1 + u)^{2n} - 1]_+ + u^{-2}[(u^2 - 1)(1 + u)^{2n}]_+$$

So the Betti numbers of B over S are

$$\beta_i^S(B) = \begin{cases} 1 & \text{for } i = 0 \\ \binom{2n}{i+1} - \binom{2n}{i-1} - 2\binom{n}{i+1} & \text{for } 1 \leq i \leq n-1 \\ \binom{2n}{i} - \binom{2n}{i+2} & \text{for } n \leq i \leq 2n \end{cases}$$

Moreover, $\dim B = n$ and $\operatorname{depth} B = 0$. □

Next we focus on the (lack of) the Koszul property of B .

Proposition 5.2. *One has $t_i^B(k) = i$ for $0 \leq i \leq n$ and $t_{n+1}^B(k) = n+2$; in particular, the k -algebra B is not Koszul.*

Proof. From Theorem 5.1 yields $t_i^S(B) = i+1$ for $i \leq n-1$, and hence $t_i^B(k) = i$ for $0 \leq i \leq n$, by Remark 3.5. It remains to verify the assertion about $t_{n+1}^B(k)$.

Let H denote the Koszul homology algebra of B . This is a $(\mathbb{N} \times \mathbb{N}^2)$ -graded algebra, with

$$H_{i,v} \cong \operatorname{Tor}_i^S(B, k)_v \quad \text{for } i \in \mathbb{N} \text{ and } v \in \mathbb{N}^2.$$

Thus $\operatorname{Hilb}_H(s, t, u)$ is the Poincaré series of B over S , computed in Theorem 5.1. It follows from that result if $v_1 = 0$ or $v_2 = 0$, then $H_{i,v} = 0$. Thus, for any element $h \in H_{i,v}$ that is decomposable as a matric Massey product, one must have $v_1 \geq 2$ and $v_2 \geq 2$. On the other hand, one has

$$H_{n+1,v} \cong k \quad \text{for } v = [n+1, 1] \text{ and } v = [1, n+1].$$

Thus, elements in these subspaces of H are indecomposable as matric Massey products. It now remains to apply Lemma 3.8 to deduce that $t_{n+1}^B(k) > n+1$. □

6. THE SYMPLECTIC CASE

In this section we study the moment map associated to the canonical representation of the symplectic Lie algebra, \mathfrak{sp}_n ; see 2.4. Let k be a field of characteristic not equal to two and S the polynomial ring in $4n$ indeterminates:

$$S := k \begin{bmatrix} p_{11} & \dots & p_{1n} & p_{21} & \dots & p_{2n} \\ q_{11} & \dots & q_{1n} & q_{21} & \dots & q_{2n} \end{bmatrix}$$

Set $\underline{p} = \{p_{ij}\}_{i,j}$ and $\underline{q} = \{q_{ij}\}_{i,j}$. As before, we view S as an \mathbb{N}^2 -graded k -algebra with the \underline{p} of degree $[1, 0]$ and the \underline{q} of degree $[0, 1]$. Let I be the ideal generated by the $2n^2 + n$ homogenous quadratic forms

$$(6.0.1) \quad p_{1i}q_{1j} - p_{2i}q_{2j} \quad 1 \leq i, j \leq n$$

$$(6.0.2) \quad p_{1i}q_{2i}, p_{2i}q_{1i} \quad 1 \leq i \leq n$$

$$(6.0.3) \quad p_{1i}q_{2j} + p_{1j}q_{2i} \quad 1 \leq i < j \leq n$$

$$(6.0.4) \quad p_{2i}q_{1j} + p_{2j}q_{1i} \quad 1 \leq i < j \leq n$$

The focus of this section is on the k -algebra $Z := S/I$, which is also \mathbb{N}^2 -graded.

Lemma 6.1. *The socle of Z is $J := (\underline{p})(\underline{q})$, viewed as an ideal in Z . It is a vector space of rank $2n^2 - n$, concentrated in degree $[1, 1]$.*

Proof. The main task is to verify that J is in the socle of Z ; then, since the ring Z/J is reduced (it is isomorphic to $S/(\underline{p})(\underline{q})$) it would follow that J is equal to the socle of Z . Moreover, then the assertion about the rank of J is clear.

For the proof, it is convenient to relabel the variables \underline{p} and \underline{q} as follows:

$$a_i := p_{1i}, \quad b_i := q_{1i}, \quad c_i := p_{2i}, \quad d_i := q_{2i} \quad \text{for each } i.$$

Since the characteristic of k is not two, it is easy to verify that the ideal I can be generated by the elements

- (1) $a_i b_j - c_i d_j$
- (2) $a_i d_j + d_i a_j$
- (3) $c_i b_j + b_i c_j$

where $1 \leq i, j \leq n$. Recall that Z is \mathbb{N}^2 -graded, with each a_i and c_i have degree $[1, 0]$ while each b_i and d_i has degree $[0, 1]$. The desired result is that the component $Z_{[1,1]}$ is the socle of Z . This is equivalent to the assertion that $Z_{[2,1]} = 0 = Z_{[1,2]}$. By symmetry it suffices to prove that $Z_{[2,1]} = 0$, that is to say, that the monomials of the type

$$aab, \quad aad, \quad acb, \quad acd, \quad ccb, \quad ccd$$

are all zero in Z . By virtue of equation (1) above, monomials of the type aab , respectively acb , are equal to those of the type acd , respectively ccd . It thus suffices to consider the following types of monomials: aad , acd , ccb , and ccd .

Types aad and ccb: Using equation (2), one gets that

$$a_i a_j d_k = -a_i d_j a_k = -a_k a_i d_j.$$

Note that the indices on the right can be obtained from the left by applying the three cycle (123) , and this involves a change of sign. So, applying the same rule three time one gets that

$$a_i a_j d_k = -a_i a_j d_k$$

and hence that $a_i a_j d_k = 0$; recall that the characteristic of k is not two.

A similar argument, now using (3), shows that $c_i c_j b_k = 0$ for all i, j, k .

Type acd: Now using equation (1) twice and then equation (2), one gets

$$a_i c_j d_k = a_i a_j b_k = c_i a_j d_k = -c_i d_j a_k = -a_k c_i d_j$$

Once again, the indices on the right are obtained from those on the left by applying a three cycle, along with a change of sign. So applying this thrice we deduce that $a_i c_j d_k = 0$, as desired.

Type ccd: Applying (1), then (3), and then again (1) yields

$$c_i c_j d_k = c_i a_j b_k = -b_i a_j c_k = -d_i c_j c_k = -c_i c_j d_k$$

so arguing as before one deduces that $c_i c_j d_k = 0$. □

Set $A = S/(p)(q)$. By Lemma 6.1 one has an exact sequence

$$(6.1.1) \quad 0 \longrightarrow J \longrightarrow Z \xrightarrow{\varepsilon} A \longrightarrow 0$$

of \mathbb{N}^2 -graded S -modules. Thus the bigraded Hilbert series of Z is

$$\text{Hilb}_Z(s, t) = \frac{1}{(1-s)^{2n}} + \frac{1}{(1-t)^{2n}} - 1 + (2n^2 - n)st.$$

We compute the Poincaré series of Z . The crucial computation is the following.

Lemma 6.2. *The exact sequence (6.1.1) induces the following isomorphisms and exact sequences of \mathbb{N}^2 -graded k -vector spaces:*

$$\begin{aligned} \mathrm{Ext}_S^0(A, k) &\xrightarrow{\cong} \mathrm{Ext}_S^0(Z, k) \\ 0 \longrightarrow \mathrm{Hom}_k(J, k) &\longrightarrow \mathrm{Ext}_S^1(A, k) \longrightarrow \mathrm{Ext}_S^1(Z, k) \longrightarrow 0 \\ \mathrm{Ext}_S^1(J, k) &\xrightarrow{\cong} \mathrm{Ext}_S^2(A, k) \\ 0 \longrightarrow \mathrm{Ext}_S^i(Z, k) &\longrightarrow \mathrm{Ext}_S^i(J, k) \longrightarrow \mathrm{Ext}_S^{i+1}(A, k) \longrightarrow 0 \end{aligned}$$

for each $i \geq 2$.

Proof. Indeed, the first isomorphism is clear. Given this, applying $\mathrm{Hom}_S(-, k)$ to the exact sequence (6.1.1) yields an exact sequence

$0 \longrightarrow \mathrm{Hom}_k(J, k) \longrightarrow \mathrm{Ext}_S^1(A, k) \longrightarrow \mathrm{Ext}_S^1(Z, k) \xrightarrow{\eta} \mathrm{Ext}_S^1(J, k) \xrightarrow{\partial} \mathrm{Ext}_S^2(A, k)$ of \mathbb{N}^2 -graded k -vector spaces. We claim $\eta = 0$ and ∂ is an isomorphism, which justifies all but the last assertion in the statement. To verify the claim, note that

$$\begin{aligned} \mathrm{rank}_k \eta &= \mathrm{rank}_k \mathrm{Ext}_S^1(Z, k) - \mathrm{rank}_k \mathrm{Ext}_S^1(A, k) + \mathrm{rank}_k \mathrm{Hom}_k(J, k) \\ &= (2n^2 + n) - 4n^2 + (2n^2 - n) \\ &= 0 \end{aligned}$$

where the second equality comes from Lemma 6.1. Thus $\eta = 0$ and hence the map ∂ is injective. As $\mathrm{Ext}_S^1(J, k)$ is isomorphic to $\mathrm{Ext}_S^1(k, k) \otimes_k \mathrm{Hom}_k(J, k)$, one gets the first equality below

$$\begin{aligned} \mathrm{rank}_k \mathrm{Ext}_S^1(J, k) &= (2n^2 - n) \mathrm{rank}_k \mathrm{Ext}_S^1(k, k) = (2n^2 - n)4n \\ \mathrm{rank}_k \mathrm{Ext}_S^2(A, k) &= \binom{4n}{3} - 2\binom{2n}{3} = (2n^2 - n)4n \end{aligned}$$

The second equality is from Theorem 4.3; keep in mind that the \underline{p} and \underline{q} have been doubled. Thus it follows that ∂ is an isomorphism, as desired.

To verify the exactness of the last family of sequences, it suffices to prove that the map

$$\mathrm{Ext}_S^{\geq 1}(J, k) \longrightarrow \mathrm{Ext}_S^{\geq 2}(A, k)$$

induced by (6.1.1) is surjective. This map is linear with respect to the action of $\Lambda := \mathrm{Ext}_S(k, k)$. Since $\mathrm{Ext}_S^{\geq 2}(A, k)$ is generated by $\mathrm{Ext}_S^2(A, k)$ as a Λ -module, by Lemma 4.4, it suffices to verify that the map of k -vector spaces

$$\mathrm{Ext}_S^1(J, k) \longrightarrow \mathrm{Ext}_S^2(A, k)$$

is surjective. This is the map ∂ that we already know is an isomorphism. \square

Lemma 6.2 implies that the minimal resolution of Z over S is pure, and in fact linear after the second syzygy. From the previous claim and Theorem 4.3 (for $2n$) one gets the following result.

Theorem 6.3. *Let k be a field, and let S and A the k -algebras introduced at the beginning of this section; in particular, the characteristic of k is zero, or odd. The nonzero graded Betti numbers of A over S are*

$$\beta_{i,v}^S(A) = \begin{cases} 1 & i = 0 \text{ and } v = [0, 0] \\ (2n^2 + n) & i = 1 \text{ and } v = [1, 1] \\ (2n^2 - n) \binom{2n}{v_1-1} \binom{2n}{v_2-1} - \binom{2n}{v_1} \binom{2n}{v_2} & i \geq 2, v_i \geq 1, \text{ and } v_1 + v_2 = i + 2 \end{cases}$$

The Betti numbers of Z over S are

$$\beta_i^S(A) = \begin{cases} 1 & \text{for } i = 0 \\ 2n^2 + n & \text{for } i = 1 \\ (2n^2 - n)\binom{4n}{i} - \binom{4n}{i+2} + 2\binom{2n}{i+2} & \text{for } i \geq 2 \end{cases}$$

Moreover, $\dim Z = 2n$ and $\operatorname{depth} Z = 0$.

Proof. The assertions about the (graded) Betti numbers have already been justified. The dimension and the depth of Z are from (6.1.1). \square

This has the following corollary.

Proposition 6.4. *The k -algebra Z is not Koszul.*

Proof. From Theorem 6.3 one gets the first equality below

$$\beta_2^S(Z)_4 = \beta_2^S(Z) = \frac{5}{3}n^2(4n^2 - 1) > \frac{1}{2}(2n^2 + n)(2n^2 + n - 1) = \binom{\beta_1^S(Z)}{2}$$

The inequality is readily verified. Now apply Remark 3.6. \square

APPENDIX A. QUADRATIC EXTERIOR FORMS AND MAXIMAL RANK

The main task of this section is to prove Proposition A.2, which was used in the proof of Theorem 5.1. Throughout k will be a field. For elements $\mathbf{x} = x_1, \dots, x_u$ in a commutative ring, we write $s_i(\mathbf{x})$ for the i -th elementary symmetric polynomials in \mathbf{x} with the convention that $s_0(\mathbf{x}) = 1$ and $s_i(\mathbf{x}) = 0$ for $i < 0$ or $i > u$.

Lemma A.1. *In $k[x_1, \dots, x_u, y_1, \dots, y_v]/J$, where $J = (y_1^2, \dots, y_v^2, x_1^2, \dots, x_u^2)$, for every $d \geq 0$ there is an equality*

$$(s_1(\mathbf{x}) + s_1(\mathbf{y})) \sum_{k=0}^d (-1)^k k! (d-k)! s_k(\mathbf{x}) s_{d-k}(\mathbf{y}) = (d+1)! (s_{d+1}(\mathbf{y}) + (-1)^d s_{d+1}(\mathbf{x})).$$

Proof. It is not hard to verify that there is an equality

$$s_1(\mathbf{x}) s_k(\mathbf{x}) = (k+1) s_{k+1}(\mathbf{x});$$

by symmetry, this implies also that $s_1(\mathbf{y}) s_{d-k}(\mathbf{y}) = (d-k+1) s_{d-k+1}(\mathbf{y})$. Using these one gets equalities

$$\begin{aligned} (s_1(\mathbf{x}) + s_1(\mathbf{y})) \sum_{k=0}^d (-1)^k k! (d-k)! s_k(\mathbf{x}) s_{d-k}(\mathbf{y}) \\ = \sum_{k=0}^d (-1)^k k! (d+1-k)! s_k(\mathbf{x}) s_{d-k+1}(\mathbf{y}) \\ + \sum_{k=0}^d (-1)^k (k+1)! (d-k)! s_{k+1}(\mathbf{x}) s_{d-k}(\mathbf{y}) \\ = (d+1)! (s_{d+1}(\mathbf{y}) + (-1)^d s_{d+1}(\mathbf{x})) \end{aligned}$$

\square

Let V be a k -vector space of dimension $2n$, and Λ the exterior algebra on V . Let $e_1, \dots, e_n, f_1, \dots, f_n$ be a basis of V and set

$$w := \sum_{i=1}^n e_i f_i;$$

this is an element in Λ_2 . The result below is well known when $\text{char } k = 0$ and can be deduced as a special case of the Hard Lefschetz Theorem; see [9, page 122]. However, we have been unable to find an argument that covers also the case of positive characteristics in the literature.

Proposition A.2. *If $\text{char } k = 0$ or $\text{char } k > (n+1)/2$, then the multiplication map $w: \Lambda_i \rightarrow \Lambda_{i+2}$ has maximal rank for every i ; in other words, $w: \Lambda_i \rightarrow \Lambda_{i+2}$ is injective for $i \leq n-1$ and surjective for $i \geq n-1$.*

Proof. To begin with, there is an isomorphism $\text{Hom}_k(\Lambda, k) \cong \Lambda(2n)$ of Λ -modules and from this it follows that it suffices to verify that multiplication by w is injective in degrees $\leq n-1$. Moreover, it suffices to verify injectivity for $i = n-1$.

Indeed, assume $w: \Lambda_i \rightarrow \Lambda_{i+2}$ is injective for some $i \leq n-1$. For any nonzero element λ of degree $i-1$, there exists an element ν in Λ_1 for which $\lambda\nu \neq 0$; this is because the socle of Λ is Λ_{2n} . Then $w\lambda\nu \neq 0$ implies $w\lambda \neq 0$. An iteration yields the desired result.

To reiterate: It suffices to verify that $w: \Lambda_{n-1} \rightarrow \Lambda_{n+1}$ is injective. Since the ranks of the source and target coincide, this is equivalent to verifying that the map is surjective. We prove this by an induction on n , the base case $n = 1$ being obvious.

Suppose $n \geq 2$. Then Λ_{n+1} has a k -basis consisting of monomials

$$\mu = e_{i_1} \dots e_{i_a} f_{j_1} \dots f_{j_b}$$

where $1 \leq a, b \leq n$ with $a+b = n+1$, and $i_1 < \dots < i_a$ and $j_1 < \dots < j_b$,

If for some h we have $i_h \notin \{j_1, \dots, j_b\}$, then $e_{i_h}w = e_{i_h}w_1$ with

$$w_1 = \sum_{v=1, v \neq i_h}^n e_v f_v$$

and, by induction on n , there exists $\lambda \in \Lambda_{n-2}(U)$ such that

$$\lambda w_1 = e_{i_1} \dots \hat{e}_{i_h} \dots e_{i_a} f_{j_1} \dots f_{j_b}.$$

Here U is the k -subspace of V generated by $\{e_1, \dots, e_n, f_1, \dots, f_n\} \setminus \{e_{i_h}, f_{i_h}\}$. Then

$$\nu e_{i_h}w = \nu e_{i_h}w_1 = \pm e_{i_h}\nu w_1 = \pm \mu.$$

A similar argument settles the case when there exists h with $j_h \notin \{i_1, \dots, i_a\}$.

It remains to consider the case $a = b$ and $\{i_1, \dots, i_a\} = \{j_1, \dots, j_b\}$. Set $m = a = b$ so that $n = 2m-1$. We can assume (renaming the indices) that $i_h = j_h = h$ for $h = 1, \dots, m$, that is to say, $\mu = \prod_{i=1}^m e_i f_i$.

We apply Lemma A.1 as follows: we set $v = m$, $u = m-1$ and $d = m-1$ and

$$\begin{aligned} x_i &\rightarrow e_{m+i} f_{m+i} & \text{for } i = 1, \dots, m-1 \\ y_i &\rightarrow e_i f_i & \text{for } i = 1, \dots, m. \end{aligned}$$

Such a specialization makes sense because the elements $x_i y_i$ have square zero and commute among themselves. Since $s_1(s') + s_1(t') = w$, $s_m(t') = \prod_{i=1}^m s_i f_i$ and $s_m(s') = 0$ we obtain:

$$w \sum_{k=0}^{m-1} (-1)^k k! (m-1-k)! s_k(s') s_{m-1-k}(t') = m! \prod_{i=1}^m s_i f_i.$$

Since we are assuming that $\text{char } k = 0$ or $\text{char } k > (n+1)/2$ we have that $m!$ is invertible and hence we may conclude that $w\nu = \prod_{i=1}^m s_i f_i$ with

$$\nu = \frac{1}{(m!)} \sum_{k=0}^{m-1} (-1)^k k! (m-1-k)! s_k(s') s_{m-1-k}(t').$$

This completes the proof. \square

APPENDIX B. BETTI TABLES AND NUMEROLOGY

In this section we collect some observations, and questions, concerning the algebras studied in this work. To begin with, consider the following Betti tables for the representations $\mathfrak{sl}_n : k^n$, computed in Theorem 5.1, for the first few values of n .

	0	1	2	3	4	5	6	7	8	9	10	11	12
$\mathfrak{sl}_2 : k^2$	0	1	—	—	—	—	—	—	—	—	—	—	—
	1	—	3	—	—	—	—	—	—	—	—	—	—
	2	—	—	5	4	1	—	—	—	—	—	—	—
$\mathfrak{sl}_3 : k^3$	0	1	—	—	—	—	—	—	—	—	—	—	—
	1	—	8	12	—	—	—	—	—	—	—	—	—
	2	—	—	—	14	14	6	1	—	—	—	—	—
$\mathfrak{sl}_4 : k^4$	0	1	—	—	—	—	—	—	—	—	—	—	—
	1	—	15	40	40	—	—	—	—	—	—	—	—
	2	—	—	—	—	42	48	27	8	1	—	—	—
$\mathfrak{sl}_5 : k^5$	0	1	—	—	—	—	—	—	—	—	—	—	—
	1	—	24	90	155	130	—	—	—	—	—	—	—
	2	—	—	—	—	—	132	165	110	44	10	1	—
$\mathfrak{sl}_6 : k^6$	0	1	—	—	—	—	—	—	—	—	—	—	—
	1	—	35	168	399	560	427	—	—	—	—	—	—
	2	—	—	—	—	—	—	429	572	429	208	65	12

The framed numbers in the table are the *Catalan numbers*; see [17, A000108]. This is true for each n , and what is more, the *Catalan triangle* introduced by Shapiro [20], appears as well.

Indeed, the sequence $(C_n)_{n \geq 0}$ of Catalan numbers, which has numerous combinatorial interpretations [23, 14], can be defined as follows:

$$C_n := \frac{1}{n+1} \binom{2n}{n} = \binom{2n}{n} - \binom{2n}{n+1}.$$

The *Segner's recursion formula* $C_{n+1} = \sum_{i,j: i+j=n} C_i C_j$ can be rewritten in terms of the generating function $C(x) = \sum_{n \geq 0} C_n x^n$ as follows: $x C^2(x) = C(x) - C_0$.

The entries $B(N, r)$ of the Catalan triangle, with $N, r \geq 1$, are defined by the higher moments $\gamma^r(x) =: \sum_{N \geq 1} B(N, r) x^N$ of the generating function $\gamma(x) = C(x) - C_0 = C(x) - 1$. In other words,

$$B(N, r) = \sum_{\substack{i_1, i_2, \dots, i_r \geq 1 \\ \sum_j i_j = N}} C_{i_1} C_{i_2} \cdots C_{i_r}.$$

It has been shown in [20] that $B(N, r) = \frac{r}{N} \binom{2N}{N-r}$, which also makes sense for $r < 0$.

The Catalan triangle can be recovered when taking differences of columns in the Pascal triangle:

$$\begin{aligned}
\binom{2n}{i} - \binom{2n}{i+2} &= \left(1 - \frac{(2n-i)(2n-i-1)}{(i+1)(i+2)}\right) \binom{2n}{i} \\
&= 2 \frac{(i+1-n)(2n+1)}{(i+1)(i+2)} \binom{2n}{i} \\
&= \frac{i+1-n}{n+1} \frac{(2n+1)(2n+2)}{(i+1)(i+2)} \binom{2n}{i} \\
&= \frac{i+1-n}{n+1} \binom{2n+2}{i+2} \\
&= B(n+1, i+1-n)
\end{aligned}$$

This proves the occurrence of the Catalan triangle in the Betti table of $\mathfrak{sl}_n : k^n$; compare Theorem 5.1. The many combinatorial interpretations of the Catalan triangle [14, 20] raise the question: *Is there a minimal free resolution of the moment map of $\mathfrak{sl}_n : k^n$ that underlies the occurrence of the Catalan triangles?*

The Betti tables of the moment map for the representations $\mathfrak{sp}_n : k^{2n}$, computed in Theorem 6.3, are also worth pointing to. For $n = 1$ the Betti table is the same as that of $\mathfrak{sl}_2 : k^2$. For $n = 2, 3$ one gets the following.

		0	1	2	3	4	5	6	7	8	9	10	11	12
$\mathfrak{sp}_2 : k^4$	0	1	—	—	—	—	—	—	—	—	—	—	—	—
	1	—	10	—	—	—	—	—	—	—	—	—	—	—
	2	—	—	100	280	392	328	167	48	6	—	—	—	—
$\mathfrak{sp}_3 : k^6$	0	1	—	—	—	—	—	—	—	—	—	—	—	—
	1	—	21	—	—	—	—	—	—	—	—	—	—	—
	2	—	—	525	2520	6503	11088	13365	11660	7359	3288	989	180	15

The noteworthy feature here is that the Betti table has only two strands, and the jump from one to the next occurs after the second step.

Poincaré series. Finally we discuss the Poincaré series of k over the coordinate algebra of the zero fibre. Let R be a standard graded algebra, as in Section 3. For simplicity, we focus on the \mathbb{N} -graded case. We are interested in the Betti numbers of k as an R -module. A basic question is when the corresponding Poincaré series $P_k^R(u)$ is rational; it is not so for a general graded ring R ; see [2, §4.3]. *What about for algebras of the form $S/(\mu)$, for a moment map μ ?*

The Poincaré series for a complete intersection is rational [2, Theorem 9.2.1], so this takes care of 0-modular representations. Thus, as for the Koszul property, the question is moot only for small representations. Rationality also holds when R is Koszul, for then there is an equality $P_k^R(s, u) = \text{Hilb}_R(-su)^{-1}$. This settles the case of the standard representations $\mathfrak{gl}_n : k^n$ and $\mathfrak{so}_n : k^n$.

We do not know the answer for $\mathfrak{sl}_n : k^n$ and $\mathfrak{sp}_n : k^n$. For $\mathfrak{sl}_2 : k^2$ Gröbner basis calculations (using Macaulay2) suggest that the Betti table of the minimal free resolution of k is upper triangular, with a new strand appearing after every 3 steps in the homological degree. The following formula for the Poincaré series best fits the available numerical data

$$P_k^{S/(\mu)}(s, u) = \frac{(1+us)^2}{(1-us)^3(1+us) - 2u^3s^4}.$$

For $\mathfrak{sl}_3 : k^3$ we also encounter a triangular shape but we have been unable to guess what the Poincaré series might be.

Following the arXiv posting of an earlier version of this manuscript, Jan-Erik Roos wrote to us (email, dated 17 June 2017) that he could answer some of the questions posed in the preceding paragraphs. Pointing to his paper [19], he writes “I have in principle solved “all” cases of the Homological Behaviour of families of quadratic forms in four variables...” Roos notes that for $\mathfrak{sl}_2 : k^2$ the ring $S/(\mu)$ is isomorphic to the one in [19, Case 7, pp. 427], and that the corresponding Poincaré series is precisely the one we proposed above. Furthermore, he proposes the following formula for the Poincaré series for the case of $\mathfrak{sl}_3 : k^3$

$$P_k^{S/(\mu)}(s, u) = \frac{(1+us)^3}{(1-us)(1-2us-4u^2s^2-2u^3s^3+u^4s^4)-2u^4s^5}.$$

and suggests a method to tackle $\mathfrak{sl}_n : k^n$, for arbitrary n ; we hope to develop these ideas in due course.

Acknowledgements. Our thanks to Lucho Avramov for helpful conversations regarding this work; in particular, for pointing out Lemma 3.8, and the work of Hreinsdóttir [13]. Part of this article is based on work supported by the National Science Foundation under Grant No. 0932078000, while AC and SBI were in residence at the Mathematical Sciences Research Institute in Berkeley, California, during the 2012–2013 Special Year in Commutative Algebra. AC was supported by INdAM-GNSAGA and PRIN “Geometry of Algebraic Varieties” 2015EYPTSB_008. SBI was partly supported by NSF grants DMS-1503044.

REFERENCES

- [1] L. L. Avramov, *The Hopf algebra of a local ring*, Izv. Akad. Nauk SSSR Ser. Mat., **38** (1974), 253–277.
- [2] L. L. Avramov, *Infinite free resolutions*, Six lectures on commutative algebra (Bellaterra, 1996), Progr. Math. **166**, Birkhäuser, Basel, 1998; 1–118.
- [3] L. L. Avramov, A. Conca and S. B. Iyengar, *Free resolutions over commutative Koszul algebras*, Math. Res. Lett. **17** (2010), 197–210.
- [4] L. L. Avramov, A. Conca and S. B. Iyengar, *Subadditivity of syzygies of Koszul algebras*, Math. Ann. **361** (2015), 511–534.
- [5] M. Bordemann, H.-C. Herbig and M. J. Pflaum, *A homological approach to singular reduction in deformation quantization*, in Ed: S. Sternberg, ‘Singularity theory’, World Sci. Publ., Hackensack, NJ, 2007, 443–461.
- [6] W. Bruns and U. Vetter, *Determinantal rings*, Lect. Notes Math. **1327**, Springer 1988.
- [7] A. Conca, E. De Negri and M. E. Rossi, *Koszul Algebras and regularity*, Commutative algebra, 285–315, Springer, New York, 2013.
- [8] R. Fröberg, *Koszul algebras*, Advances in Commutative Ring Theory, Proceedings of Fez Conference 1997. In: Lectures Notes in Pure and Applied Mathematics, vol. **205**, Marcel Dekker, New York (1999)
- [9] P. Griffiths and J. Harris, *Principles of algebraic geometry*, Reprint of the 1978 original. Wiley Classics Library. John Wiley & Sons, Inc., New York, 1994.
- [10] V. K. A. M. Gugenheim and J. P. May, *On the theory and applications of differential torsion products*, Memoirs of the American Mathematical Society, **142**, American Mathematical Society, Providence, R.I., 1974.
- [11] M. Henneaux and C. Teitelboim, *Quantization of gauge systems*, Princeton University Press, Princeton, NJ, 1992.
- [12] H.-C. Herbig and G. W. Schwarz, *The Koszul complex of a moment map*, J. Symplectic Geom., **11** (2013), 497–508.
- [13] F. Hreinsdóttir, *The Koszul dual of the ring of commuting matrices*, Communications in Algebra, **26** (2011), 3807–3819.

- [14] T. Koshy, *Catalan numbers with applications*, Oxford University Press. xiv, 422 p.
- [15] D. Kraines, *Massey higher products*, Trans. Amer. Math. Soc. **124** (1966), 431–449.
- [16] J. P. May, *Matrix Massey products*, J. Algebra **12** (1969), 533–568.
- [17] *The On-Line Encyclopedia of Integer Sequences*, published electronically at <http://oeis.org>, 2014.
- [18] A. Polischuk, L. Positselski *Quadratic Algebras*, AMS University Lecture Series, 2005.
- [19] J.-E. Roos, *Homological properties of the homology algebra of the Koszul complex of a local ring: examples and questions*, J. Algebra **465** (2016), 399–436.
- [20] L. W. Shapiro, *A Catalan triangle*, Discrete Math. **14** (1976), 83–90.
- [21] G. W. Schwarz, *Differential operators on quotients of simple groups*, J. Algebra **169** (1994), 248–273.
- [22] G. W. Schwarz, *Lifting differential operators from orbit spaces*, Ann. Sci. École Norm. Sup. (4), **28** (1995), 253–305.
- [23] R. P. Stanley, *Enumerative combinatorics. Volume 2.*, Cambridge Studies in Advanced Mathematics. **62**. Cambridge University Press. xii, 581 p.
- [24] J. Tate, *Homology of Noetherian rings and local rings*, Illinois J. Math. **1** (1957), 14–27.
- [25] È. B. Vinberg, *Complexity of actions of reductive groups*, Funktsional. Anal. i Prilozhen. **20** (1986), 1–13, 96.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÁ DI GENOVA, VIA DODECANESO 35, I-16146 GENOVA, ITALY

E-mail address: `conca@dima.unige.it`

DEPARTAMENTO DE MATEMÁTICA APLICADA, UNIVERSIDADE FEDERAL DO RIO DE JANEIRO (UFRJ), AV. ATHOS DA SILVEIRA RAMOS 149, CENTRO DE TECNOLOGIA - BLOCO C, CEP: 21941-909 - RIO DE JANEIRO, BRAZIL

E-mail address: `herbig@labma.ufrj.br`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY, UT 84112, U.S.A.

E-mail address: `iyengar@math.utah.edu`