



PAPER

General soliton solution to a nonlocal nonlinear Schrödinger equation with zero and nonzero boundary conditions

Recent citations

- [Integrable nonlocal asymptotic reductions of physically significant nonlinear equations](#)
Mark J Ablowitz and Ziad H Musslimani

To cite this article: Bao-Feng Feng *et al* 2018 *Nonlinearity* **31** 5385

View the [article online](#) for updates and enhancements.

General soliton solution to a nonlocal nonlinear Schrödinger equation with zero and nonzero boundary conditions

Bao-Feng Feng¹, Xu-Dan Luo², Mark J Ablowitz³
and Ziad H Musslimani⁴

¹ School of Mathematical and Statistical Sciences, The University of Texas Rio Grande Valley, Edinburg, TX 78539-2999, United States of America

² Department of Mathematics, State University of New York at Buffalo, Buffalo, NY 14260, United States of America

³ Department of Applied Mathematics, University of Colorado, Boulder, CO 80309-0526, United States of America

⁴ Department of Mathematics, Florida State University, Tallahassee, FL 32306-4510, United States of America

E-mail: baofeng.feng@utrgv.edu

Received 30 December 2017, revised 28 August 2018

Accepted for publication 11 September 2018

Published 30 October 2018



CrossMark

Recommended by Professor Tamara Grava

Abstract

General soliton solutions to a nonlocal nonlinear Schrödinger (NLS) equation with PT-symmetry for both zero and nonzero boundary conditions are considered via the combination of Hirota's bilinear method and the Kadomtsev–Petviashvili (KP) hierarchy reduction method. First, general N -soliton solutions with zero boundary conditions are constructed. Starting from the tau functions of the two-component KP hierarchy, it is shown that they can be expressed in terms of either Gramian or double Wronskian determinants. On the contrary, from the tau functions of single component KP hierarchy, general soliton solutions to the nonlocal NLS equation with nonzero boundary conditions are obtained. All possible soliton solutions to nonlocal NLS with Parity (PT)-symmetry for both zero and nonzero boundary conditions are found in the present paper.

Keywords: nonlocal nonlinear Schrödinger equation, Hirota's bilinear method, the KP hierarchy reduction method, soliton solutions with zero and nonzero boundary conditions

Mathematics Subject Classification numbers: 37K15, 37Q55, 35Q51

(Some figures may appear in colour only in the online journal)

1. Introduction

The study of the nonlinear Schrödinger (NLS) equation and its multi-component generalization lies at the forefront of research in applied mathematics and mathematical physics. This is the case since they have been recognized as generic models for describing the evolution of slowly varying wave packets in general nonlinear wave systems [1–3]. In 1972 Zakharov and Shabat found that the NLS equation possesses a Lax pair and can be solved/linearized by inverse scattering [4]. Motivated by these results in 1973 Ablowitz, Kaup, Newell and Segur [5] generalized the linear operators used by Zakharov and Shabat; they showed that the NLS, sine-Gordon [6], modified-KdV, KdV all could be solved/linearized by inverse scattering. Soon afterwards in 1974, AKNS [7] developed a general framework to find integrable systems solvable by what they termed the Inverse Scattering Transform (IST). The method was associated with classes of equations (later called recursion operators) and was applied to find solutions to the initial value problem with rapidly decaying data on the line.

The NLS and coupled NLS equations arise in a variety of physical contexts such as nonlinear optics [8–12], Bose–Einstein condensates [13], water waves [14, 15] and plasma physics [16], amongst many others.

The idea in [7] was to consider the scattering problem

$$v_x = Xv = \begin{pmatrix} -ik & q(x, t) \\ r(x, t) & ik \end{pmatrix} v, \quad (1)$$

$$v_t = Tv = \begin{pmatrix} 2ik^2 + iqr & -2kq - iq_x \\ -2kr + ir_x & -2ik^2 - iqr \end{pmatrix} v. \quad (2)$$

The compatibility condition of the linear spectral problem (1) and (2) leads to

$$\begin{cases} iq_t = q_{xx} - 2rq^2, \\ -ir_t = r_{xx} - 2qr^2. \end{cases} \quad (3)$$

Under the standard symmetry reduction

$$r(x, t) = \sigma q^*(x, t), \quad \sigma = \pm 1, \quad (4)$$

the system (3) gives the classical NLS equation

$$iq_t(x, t) = q_{xx}(x, t) - 2\sigma q^*(x, t) q^2(x, t). \quad (5)$$

Recently, an integrable nonlocal NLS equation

$$iq_t(x, t) = q_{xx}(x, t) - 2\sigma q^*(-x, t) q^2(x, t) \quad (6)$$

was found in [17] under a new symmetry reduction

$$r(x, t) = \sigma q^*(-x, t), \quad \sigma = \pm 1. \quad (7)$$

This nonlocal equation is shown to be related to an unconventional magnetic system [18]. It is parity-time (PT) symmetric, i.e. it is invariant under the joint transformations of $x \rightarrow -x$, $t \rightarrow t$ and complex conjugation, thus is related to a hot research area of contemporary physics [19, 20]. Due to this potential application, it has been shown that different symmetry reductions from the linear problem of general AKNS hierarchy and other integrable hierarchies can also lead to various types of new nonlocal equations, which have been extensively studied [21–57]. Typical examples are the reverse space-time nonlocal NLS equation and the reverse time nonlocal NLS equation, the complex/real space-time sine-Gordon equation [21–24], the

complex/real reverse space-time mKdV equation [21, 22], the nonlocal derivative NLS equation [34], and the multi-dimensional nonlocal Davey–Stewartson equation [22, 27]. On the other hand, corresponding to the classical semi-discrete NLS equation [58, 59], the nonlocal NLS equation also admits its semi-discrete version [26] and multi-component generalizations [22, 36].

It has been known that the inverse scattering transform (IST) has been successfully developed for the NLS, coupled NLS, as well as their semi-discrete analogues [60–70]. Moreover, the IST to solve the initial-value problem with vanishing boundary conditions for the nonlocal NLS equation has been developed in [17, 21, 22]. For pure soliton solutions corresponding to reflectionless potentials, one obtains one- and two-soliton solutions; these solutions can have singularities. In particular, the decaying one-soliton solution is found to be [17]

$$q = \frac{2(\eta + \bar{\eta})e^{-2\bar{\eta}x - 4i\bar{\eta}^2t + i\theta}}{1 - e^{-2(\eta + \bar{\eta})x + 4i(\eta^2 - \bar{\eta}^2)t + i(\theta + \bar{\theta})}}. \quad (8)$$

This solution is singular at $x = 0$ when $\eta \neq \bar{\eta}$ and $4(\eta^2 - \bar{\eta}^2)t + (\theta + \bar{\theta}) = 2n\pi, n \in \mathbb{Z}$. When $\eta = \bar{\eta}, \theta + \bar{\theta} \neq 2n\pi, n \in \mathbb{Z}$ the solution is not singular. It is also noted that the same solution has been found in [48, 52] via Darboux transformations and AKNS reductions, respectively.

The inverse scattering transform (IST) to solve the initial-value problem with nonzero boundary conditions has also been developed in [23]. To be specific, under the following boundary conditions

$$q(x, t) \rightarrow q_{\pm}(t) = q_0 e^{i(\alpha t + \theta_{\pm})}, \text{ as } x \rightarrow \pm\infty, \quad (9)$$

where $q_0 > 0, 0 \leq \theta_{\pm} < 2\pi$, the following cases are studied.

- $\sigma = -1, \Delta\theta = \theta_+ - \theta_- = \pi$: a one-soliton solution is found:

$$q = q_0 e^{i(2q_0^2 t + \theta_+ - \pi)} \tanh[q_0 x - i\theta_*] \quad (10)$$

where $\theta_* = \frac{1}{2}(\theta_+ + \theta_- + \pi)$, where θ_1 is related to the associated scattering data (normalization constant). This solution is stationary in space and oscillating in time.

- $\sigma = -1, \Delta\theta = 0$: in this case, there is no ‘proper exponentially decaying’ pure one soliton solution. The simplest decaying pure reflectionless potential generates a two-soliton solution.
- $\sigma = 1, \Delta\theta = 0$: in this case only an even number of soliton solutions arise; they are related to an even number ($2N$) of eigenvalues.
- $\sigma = 1, \Delta\theta = \pi$: in this case, there is no eigenvalues/solitons.

Regardless of the success of the IST for the nonlocal NLS equation, it is still not easy to come up with its general soliton solution in closed form by this method. On the other hand, Hirota’s bilinear method is a direct and powerful method in finding multi-soliton solutions to soliton equations [71]. In the early 1980s, suggested by Hirota’s direct method, Sato discovered that the soliton solutions to the Kadomtsev–Petviashvili (KP) equation and its hierarchy (named tau-functions afterwards) satisfy the so-called Plücker relations, and found that the totality of solutions to the KP hierarchy forms an infinite dimensional Grassmann manifold [72]. This is called Sato theory later on. Soon afterwards, Ueno and Takasaki extended Sato theory to include two-dimensional Toda lattice (2DTL) hierarchy [73]; Date, Jimbo and Miwa further

developed Sato's idea with transformation group theory, and presented Lie algebraic classification of soliton equations [74, 75, 76].

The KP-hierarchy reduction method was first developed by the Kyoto school [74], and later was used to obtain soliton solutions in the NLS equation, the modified KdV equation, the Davey–Stewartson equation and the coupled higher order NLS equations [77, 78]. Recently, this method has been applied to derive dark–dark soliton solution in two-coupled NLS equation of the mixed type [79]. Unlike the inverse scattering transform method [60], and Hirota's bilinear method [71], the KP-hierarchy reduction method starts with the general KP hierarchy including the two-dimensional Toda-hierarchy [80] and derives the general soliton solution in either determinant or pfaffian form reduced directly by the tau functions of the KP hierarchy. In spite of many successes of the KP-hierarchy reduction method, constructing general soliton solutions by this method to the nonlocal NLS equation remains an unsolved and challenging problem, especially for the nonzero boundary conditions. The difficulty lies in the simultaneous realization of both the nonlocal and complex conjugate reductions. Therefore, it motivates the present work, which intends to construct general soliton solutions in determinant forms to the nonlocal NLS equation with both the zero and nonzero boundary conditions.

In the present paper, we find solitons that are both decaying and have nonzero boundary values. For the decaying boundary condition, we construct the general N -soliton solutions expressed in both the Gram-type and the double Wronski-type determinants. The explicit one-soliton solution agrees with the solution found by the inverse scattering transform [17, 57], which is also listed in (8). For the nonzero boundary condition, the general soliton solutions are constructed for all cases except for the case: $\sigma = 1, \Delta = \pi$ where there is no soliton solution. To be specific, for the case of $\sigma = -1, \Delta = \pi$, the solution (97) is the same as above; for the case of $\sigma = 1, \Delta = 0$, it is true that only the determinant soliton solution $2N \times 2N$ can be constructed, which is consistent with the fact that only the even number of soliton solutions with an even number of eigenvalues exist. The simplest solution based on the tau-functions found here corresponds to (5.90) in [23] but with a simpler form. Finally, for the case of $\sigma = -1, \Delta = 0$, the soliton solution (108) found in this paper is the same as the one found in [23].

The outline of the present paper is organized as follows. In section 2, we construct the bright soliton solution to the nonlocal NLS equation with a zero boundary condition. Both the Gram-type and double Wronski-type solutions start from the tau functions in Gram-type or double Wronski-type of two-component KP hierarchy. In section 3, general multi-soliton solutions for the nonlocal NLS equation of $\sigma = 1$ and of $\sigma = -1$ are constructed, which recover all possible solutions found in [23]. The paper concludes with comments and remarks in section 4.

2. General soliton solution with zero boundary condition

In this section, we consider general soliton solutions to the nonlocal NLS equation (6) under the boundary condition $q(x, t) \rightarrow 0$ as $x \rightarrow \pm\infty$. This type of soliton solution is usually called bright soliton solution. We first give the bilinear forms for the nonlocal NLS equation (6). By introducing the dependent variable transformation

$$q(x, t) = \frac{g(x, t)}{f(x, t)}, \quad (11)$$

the nonlocal NLS equation (6) is converted into the following bilinear equations

$$\begin{cases} (iD_t - D_x^2)g(x, t) \cdot f(x, t) = 0, \\ D_x^2 f(x, t) \cdot f(x, t) + 2\sigma g(x, t)g^*(-x, t) = 0, \end{cases} \quad (12)$$

where Hirota's bilinear operator is defined as

$$D_x^n D_t^l f(x, t) \cdot g(x, t) = \left(\frac{\partial}{\partial x} - \frac{\partial}{\partial x'} \right)^n \left(\frac{\partial}{\partial t} - \frac{\partial}{\partial t'} \right)^l f(x, t)g(x', t') \Big|_{x=x', t=t'}.$$

Similar to the classical NLS equation, the bright soliton solutions to the nonlocal NLS can be obtained from the tau functions of two-component KP hierarchy. It is known that the tau functions have two different forms: one is the Gram-type, the other is the Wronski-type. They are basically equivalent, however, each form has advantage and disadvantage in view of different symmetries. In what follows, we will derive two types of soliton solutions via KP hierarchy reduction method.

2.1. General bright soliton solution expressed by Gramian determinant

In this subsection, we will construct soliton solution in terms of Gram-type determinant. To this end, let us start with a Gram-type determinant expression of the tau functions for two-component KP hierarchy,

$$\tau_0 = |A|, \quad (13)$$

$$\tau_1 = \begin{vmatrix} A & \Phi^T \\ -\bar{\Psi} & 0 \end{vmatrix}, \quad \tau_{-1} = \begin{vmatrix} A & \Psi^T \\ -\bar{\Phi} & 0 \end{vmatrix} \quad (14)$$

where the elements of matrix A are

$$a_{ij} = \frac{1}{p_i + \bar{p}_j} e^{\xi_i + \bar{\xi}_j} + \frac{1}{q_i + \bar{q}_j} e^{\eta_i + \bar{\eta}_j}$$

with

$$\xi_i = p_i x_1 + p_i^2 x_2 + \xi_{i0}, \quad \bar{\xi}_j = \bar{p}_j x_1 - \bar{p}_j^2 x_2 + \bar{\xi}_{j0},$$

$$\eta_i = q_i y_1 + \eta_{i0}, \quad \bar{\eta}_j = \bar{q}_j y_1 + \bar{\eta}_{j0},$$

and the superscript T represent the transpose, $\Phi, \bar{\Phi}, \Psi, \bar{\Psi}$ are row vectors defined by

$$\Phi = (e^{\xi_1}, \dots, e^{\xi_N}), \quad \Psi = (e^{\eta_1}, \dots, e^{\eta_N}),$$

$$\bar{\Phi} = (e^{\bar{\xi}_1}, \dots, e^{\bar{\xi}_N}), \quad \bar{\Psi} = (e^{\bar{\eta}_1}, \dots, e^{\bar{\eta}_N}).$$

It can be shown below that the tau functions given above satisfy the following two bilinear equations

$$\begin{cases} (D_{x_2} - D_{x_1}^2)\tau_1 \cdot \tau_0 = 0, \\ D_{x_1} D_{y_1} \tau_0 \cdot \tau_0 = -2\tau_1 \tau_{-1}. \end{cases} \quad (15)$$

The proof is given as follows. Since

$$\partial_{x_1} \tau_0 = \begin{vmatrix} A & \Phi^T \\ -\bar{\Phi} & 0 \end{vmatrix}, \quad \partial_{x_1} \partial_{y_1} \tau_0 = \begin{vmatrix} A & \Phi^T & \Psi^T \\ -\bar{\Phi} & 0 & 0 \\ -\bar{\Psi} & 0 & 0 \end{vmatrix},$$

then based on the Jacobi identity for the determinant, we have

$$(2\partial_{x_1} \partial_{y_1}) \tau_0 \times \tau_0 = 2(\partial_{x_1} \tau_0) \times (\partial_{x_1} \tau_y) - 2\tau_1 \times \tau_{-1},$$

which is exactly the second bilinear equation.

Moreover, we can easily verify the following relations

$$\begin{aligned} \partial_{x_2} \tau_0 &= \begin{vmatrix} A & \Phi_{x_1}^T \\ -\bar{\Phi} & 0 \end{vmatrix} - \begin{vmatrix} A & \Phi_{x_1}^T \\ -\bar{\Phi}_{x_1} & 0 \end{vmatrix}, \\ \partial_{x_1}^2 \tau_0 &= \begin{vmatrix} A & \Phi_{x_1}^T \\ -\bar{\Phi} & 0 \end{vmatrix} + \begin{vmatrix} A & \Phi_{x_1}^T \\ -\bar{\Phi}_{x_1} & 0 \end{vmatrix}, \quad \partial_{x_1} \tau_1 = \begin{vmatrix} A & \Phi_{x_1}^T \\ -\bar{\Psi} & 0 \end{vmatrix}, \end{aligned}$$

and

$$\begin{aligned} \partial_{x_1}^2 \tau_1 &= \begin{vmatrix} A & \Phi^T & \Phi_{x_1}^T \\ -\bar{\Phi} & 0 & 0 \\ -\bar{\Psi} & 0 & 0 \end{vmatrix} + \begin{vmatrix} A & \Phi_{x_1}^T \\ -\bar{\Psi} & 0 \end{vmatrix}, \\ \partial_{x_2} \tau_1 &= - \begin{vmatrix} A & \Phi^T & \Phi_{x_1}^T \\ -\bar{\Phi} & 0 & 0 \\ -\bar{\Psi} & 0 & 0 \end{vmatrix} + \begin{vmatrix} A & \Phi_{x_1}^T \\ -\bar{\Psi} & 0 \end{vmatrix}, \end{aligned}$$

where Φ_{x_1} and $\Phi_{x_1 x_1}$ represent the first and second derivatives of Φ with respect to x_1 , respectively. Therefore we have

$$\begin{aligned} (\partial_{x_2} - \partial_{x_1}^2) \tau_0 &= 2 \begin{vmatrix} A & \Phi_{x_1}^T \\ -\bar{\Psi} & 0 \end{vmatrix}, \\ (\partial_{x_2} - \partial_{x_1}^2) \tau_1 &= -2 \begin{vmatrix} A & \Phi^T & \Phi_{x_1}^T \\ -\bar{\Phi} & 0 & 0 \\ -\bar{\Psi} & 0 & 0 \end{vmatrix}. \end{aligned}$$

From the Jacobi identity for the determinant, we have

$$((\partial_{x_2} - \partial_{x_1}^2) \tau_1) \times \tau_0 = ((\partial_{x_2} - \partial_{x_1}^2) \tau_0) \times \tau_1 - 2(\partial_{x_1} \tau_1) \times (\partial_{x_1} \tau_{f0})$$

which gives the first bilinear equation. The proof is done.

Next, we show the reduction processes. By row operations τ_0 can be rewritten as

$$\tau_0 = \prod_{j=1}^N e^{\xi_j + \bar{\xi}_j} |a'_{ij}|_{1 \leq i, j \leq N} = \prod_{j=1}^N e^{\xi_j + \bar{\xi}_j} \left| \frac{1}{p_i + \bar{p}_j} + \frac{1}{q_i + \bar{q}_j} e^{(\eta_i - \xi_i) + (\bar{\eta}_j - \bar{\xi}_j)} \right|$$

where

$$\eta_i - \xi_i = q_i y_1 - p_i x_1 + \cdots, \quad \bar{\eta}_j - \bar{\xi}_j = \bar{q}_j y_1 - \bar{p}_j x_1 + \cdots.$$

Thus if we impose constraints on parameters

$$q_j = p_j, \quad \bar{q}_j = \bar{p}_j, \quad j = 1, \dots, N, \quad (16)$$

then the following relation holds

$$(\partial_{x_1} + \partial_{y_1})a'_{ij} = 0$$

which implies

$$(\partial_{x_1} + \partial_{y_1})\tau_0 = \sum_{j=1}^N (p_j + \bar{p}_j)\tau_0. \quad (17)$$

Consequently, the second bilinear equation in (3) becomes

$$D_{x_1}^2 \tau_0 \cdot \tau_0 = 2\tau_1 \tau_{-1}. \quad (18)$$

Due to the dimension reduction (17), y_1 becomes a dummy variable, which can be treated as zero. We then have the following Lemma.

Lemma 2.1. Assume $x_1 = x$ is real and $x_2 = -it$ is a purely imaginary variable. If p_j and $\bar{p}_j$ ($j = 1, \dots, N$) are real, η_{j0} , $\bar{\eta}_{j0}$ are purely imaginary, or the subset of (p_j, η_{j0}) or $(\bar{p}_j, \bar{\eta}_{j0})$ occurs in pairs such that $p_k = p_{k'}^*$, $\eta_{k0} = \eta_{k'0}^*$ or $\bar{p}_k = \bar{p}_{k'}^*$, $\bar{\eta}_{k0} = \bar{\eta}_{k'0}^*$, then

$$\tau_0(x, t) = C\tau_0^*(-x, t) \quad \tau_{-1}(x, t) = C\tau_1^*(-x, t), \quad (19)$$

where $C = C_1 C_2$, $C_1 = \prod_{j=1}^N e^{(\xi_j + \bar{\xi}_j)}$, $C_2 = \prod_{j=1}^N e^{(\eta_j + \bar{\eta}_j)}$.

Proof. First, we prove the case when p_i and $\bar{p}_i$ are all real, η_{i0} , $\bar{\eta}_{i0}$ are all purely imaginary. In this case,

$$\begin{aligned} \xi_i &= p_i x - i p_i^2 t, & \eta_i &= i \theta_i, \\ \bar{\xi}_j &= \bar{p}_j x + i \bar{p}_j^2 t, & \bar{\eta}_j &= i \bar{\theta}_j, \end{aligned}$$

it then follows

$$\xi_i^*(-x, t) = -\xi_i(x, t), \quad \bar{\xi}_i^*(-x, t) = -\bar{\xi}_i(x, t).$$

We note that $\tau_0(x, t)$ can be rewritten as

$$\begin{aligned} \tau_0(x, t) &= \prod_{j=1}^N e^{(\xi_j + \bar{\xi}_j)} \left| \frac{1}{p_i + \bar{p}_j} (1 + e^{-(p_i + \bar{p}_j)x + i(p_i^2 - \bar{p}_j^2)t + i(\theta_i + \bar{\theta}_j)}) \right|, \\ &= \prod_{j=1}^N e^{i(\theta_j + \bar{\theta}_j)} \left| \frac{1}{p_i + \bar{p}_j} (1 + e^{(p_i + \bar{p}_j)x - i(p_i^2 - \bar{p}_j^2)t - i(\theta_i + \bar{\theta}_j)}) \right|, \end{aligned} \quad (20)$$

which implies

$$\tau_0^*(-x, t) = \prod_{j=1}^N e^{-(\xi_j + \bar{\xi}_j)} \left| \frac{1}{p_i + \bar{p}_j} (1 + e^{(p_i + \bar{p}_j)x - i(p_i^2 - \bar{p}_j^2)t - i(\theta_i + \bar{\theta}_j)}) \right|. \quad (21)$$

Therefore $\tau_0(x, t) = C\tau_0^*(-x, t)$. On the other hand,

$$\begin{aligned}
\tau_1(x, t) &= \begin{vmatrix} \frac{1}{p_i + \bar{p}_j} (e^{(p_i + \bar{p}_j)x - i(p_i^2 - \bar{p}_j^2)t} + e^{i(\theta_i + \bar{\theta}_j)}) & e^{p_i x - i p_i^2 t} \\ -e^{i\bar{\theta}_j} & 0 \end{vmatrix}, \\
&= \prod_{j=1}^N e^{i(\theta_j + \bar{\theta}_j)} \begin{vmatrix} \frac{1}{p_i + \bar{p}_j} (1 + e^{\xi_i + \bar{\xi}_j - i(\theta_i + \bar{\theta}_j)}) & e^{\xi_i - i\theta_i} \\ -1 & 0 \end{vmatrix}, \\
&= \prod_{j=1}^N e^{(\xi_j + \bar{\xi}_j)} \begin{vmatrix} \frac{1}{p_i + \bar{p}_j} (1 + e^{-\xi_i - \bar{\xi}_j + i(\theta_i + \bar{\theta}_j)}) & 1 \\ -e^{-\xi_j + i\bar{\theta}_j} & 0 \end{vmatrix}, \tag{22}
\end{aligned}$$

$$\begin{aligned}
\tau_{-1}(x, t) &= \begin{vmatrix} \frac{1}{p_i + \bar{p}_j} (e^{(p_i + \bar{p}_j)x - i(p_i^2 - \bar{p}_j^2)t} + e^{i(\theta_i + \bar{\theta}_j)}) & e^{i\theta_i} \\ -e^{\bar{p}_j x + i\bar{p}_j^2 t} & 0 \end{vmatrix}, \\
&= \prod_{j=1}^N e^{(\xi_j + \bar{\xi}_j)} \begin{vmatrix} \frac{1}{p_i + \bar{p}_j} (1 + e^{-\xi_i - \bar{\xi}_j + i(\theta_i + \bar{\theta}_j)}) & e^{-\xi_i + i\theta_i} \\ -1 & 0 \end{vmatrix}. \tag{23}
\end{aligned}$$

Therefore

$$\begin{aligned}
\tau_1^*(-x, t) &= \prod_{j=1}^N e^{(\xi_j + \bar{\xi}_j)^*(-x, t)} \begin{vmatrix} \frac{1}{p_i + \bar{p}_j} (1 + e^{\xi_i + \bar{\xi}_j - i(\theta_i + \bar{\theta}_j)}) & 1 \\ -e^{\bar{\xi}_j - i\bar{\theta}_j} & 0 \end{vmatrix}, \\
&= \prod_{j=1}^N e^{-i(\theta_j + \bar{\theta}_j)} \begin{vmatrix} \frac{1}{p_i + \bar{p}_j} (1 + e^{-\xi_i - \bar{\xi}_j + i(\theta_i + \bar{\theta}_j)}) & e^{-\xi_i + i\theta_i} \\ -1 & 0 \end{vmatrix}. \tag{24}
\end{aligned}$$

Obviously $\tau_{-1}(x, t) = C\tau_1^*(-x, t)$. Next we prove the case where there are pairs of wave numbers $p_k, p_{k'}, \bar{p}_k, \bar{p}_{k'}$ such that $p_k = p_{k'}^*, \bar{p}_k = \bar{p}_{k'}^*$. Moreover, $\eta_{k'0} = -\eta_{k0}^*, \bar{\eta}_{k'0} = -\bar{\eta}_{k0}^*$. Note that $p_k, p_{k'}$ or $\bar{p}_k, \bar{p}_{k'}$ being real; $\eta_{k'0}, \eta_{k0}$ or $\bar{\eta}_{k'0}, \bar{\eta}_{k0}$ being purely imaginary is simply a special case. Under this case, we also have $\xi_k^*(-x, t) = -\xi_{k'}(x, t), \bar{\xi}_k^*(-x, t) = -\bar{\xi}_{k'}(x, t)$, thus

$$\begin{aligned}
\tau_0(x, t) &= \begin{vmatrix} \frac{1}{p_k + \bar{p}_{k'}} (e^{\xi_k + \bar{\xi}_{k'}} + e^{\eta_{k0} + \bar{\eta}_{k'0}}) & \dots \\ \dots & \frac{1}{p_{k'} + \bar{p}_k} (e^{\xi_{k'} + \bar{\xi}_k} + e^{\eta_{k'0} + \bar{\eta}_{k0}}) \end{vmatrix} \\
&= C_2 \begin{vmatrix} \frac{1}{p_k + \bar{p}_{k'}} (1 + e^{\xi_k + \bar{\xi}_{k'} - \eta_{k0} - \bar{\eta}_{k'0}}) & \dots \\ \dots & \frac{1}{p_{k'} + \bar{p}_k} (1 + e^{\xi_{k'} + \bar{\xi}_k - \eta_{k'0} - \bar{\eta}_{k0}}) \end{vmatrix} \\
&= C_1 \begin{vmatrix} \frac{1}{p_k + \bar{p}_{k'}} (1 + e^{-(\xi_k + \bar{\xi}_{k'} - \eta_{k0} - \bar{\eta}_{k'0})}) & \dots \\ \dots & \frac{1}{p_{k'} + \bar{p}_k} (1 + e^{-(\xi_{k'} + \bar{\xi}_k - \eta_{k'0} - \bar{\eta}_{k0})}) \end{vmatrix} \\
&= C_1 \begin{vmatrix} \frac{1}{p_{k'} + \bar{p}_k} (1 + e^{-(\xi_{k'} + \bar{\xi}_k - \eta_{k'0} - \bar{\eta}_{k0})}) & \dots \\ \dots & \frac{1}{p_k + \bar{p}_{k'}} (1 + e^{-(\xi_k + \bar{\xi}_{k'} - \eta_{k0} - \bar{\eta}_{k'0})}) \end{vmatrix}. \tag{25}
\end{aligned}$$

Therefore

$$\tau_0^*(-x, t) = C_1^{-1} \begin{vmatrix} \frac{1}{p_k + \bar{p}_{k'}} (1 + e^{(\xi_k + \bar{\xi}_{k'} - \eta_{k0} - \bar{\eta}_{k'0}))} & \dots & \dots \\ \dots & \dots & \dots \\ \dots & \dots & \frac{1}{p_{k'} + \bar{p}_k} (1 + e^{-(\xi_k + \bar{\xi}_{k'} - \eta_{k0} - \bar{\eta}_{k'0}))} \end{vmatrix}. \quad (26)$$

We again found $\tau_0(x, t) = C\tau_0^*(-x, t)$. On the other hand,

$$\begin{aligned} \tau_1(x, t) &= \begin{vmatrix} \frac{1}{p_k + \bar{p}_{k'}} (e^{\xi_k + \bar{\xi}_{k'}} + e^{\eta_{k0} + \bar{\eta}_{k'0}}) & \dots & e^{\xi_k} \\ \dots & \frac{1}{p_{k'} + \bar{p}_k} (e^{\xi_{k'} + \bar{\xi}_k} + e^{\eta_{k'0} + \bar{\eta}_{k0}}) & e^{\xi_{k'}} \\ -e^{\bar{\eta}_{k'0}} & -e^{\bar{\eta}_{k0}} & 0 \end{vmatrix} \\ &= C_2 \begin{vmatrix} \frac{1}{p_k + \bar{p}_{k'}} (1 + e^{\xi_k + \bar{\xi}_{k'} - \eta_{k0} - \bar{\eta}_{k'0}}) & \dots & e^{\xi_k - \eta_{k0}} \\ \dots & \frac{1}{p_{k'} + \bar{p}_k} (1 + e^{\xi_{k'} + \bar{\xi}_k - \eta_{k'0} - \bar{\eta}_{k0}}) & e^{\xi_{k'} - \eta_{k'0}} \\ -1 & -1 & 0 \end{vmatrix} \\ &= C_1 \begin{vmatrix} \frac{1}{p_k + \bar{p}_{k'}} (1 + e^{-(\xi_k + \bar{\xi}_{k'} - \eta_{k0} - \bar{\eta}_{k'0}))} & \dots & 1 \\ \dots & \frac{1}{p_{k'} + \bar{p}_k} (1 + e^{-(\xi_{k'} + \bar{\xi}_k - \eta_{k'0} - \bar{\eta}_{k0}))} & 1 \\ -e^{\bar{\eta}_{k'0}} & -e^{\bar{\eta}_{k0}} & 0 \end{vmatrix} \\ &= C_1 \begin{vmatrix} \frac{1}{p_{k'} + \bar{p}_k} (1 + e^{-(\xi_{k'} + \bar{\xi}_k - \eta_{k'0} - \bar{\eta}_{k0}))} & \dots & 1 \\ \dots & \frac{1}{p_k + \bar{p}_{k'}} (1 + e^{-(\xi_k + \bar{\xi}_{k'} - \eta_{k0} - \bar{\eta}_{k'0}))} & 1 \\ -e^{\bar{\eta}_{k0}} & -e^{\bar{\eta}_{k'0}} & 0 \end{vmatrix}, \end{aligned}$$

$$\begin{aligned} \tau_{-1}(x, t) &= \begin{vmatrix} \frac{1}{p_k + \bar{p}_{k'}} (e^{\xi_k + \bar{\xi}_{k'}} + e^{\eta_{k0} + \bar{\eta}_{k'0}}) & \dots & e^{\eta_{k0}} \\ \dots & \frac{1}{p_{k'} + \bar{p}_k} (e^{\xi_{k'} + \bar{\xi}_k} + e^{\eta_{k'0} + \bar{\eta}_{k0}}) & e^{\eta_{k'0}} \\ -e^{\bar{\xi}_{k'}} & -e^{\bar{\xi}_k} & 0 \end{vmatrix} \\ &= C_1 \begin{vmatrix} \frac{1}{p_{k'} + \bar{p}_k} (1 + e^{-(\xi_{k'} + \bar{\xi}_k - \eta_{k'0} - \bar{\eta}_{k0}))} & \dots & e^{-\xi_{k'} + \eta_{k'0}} \\ \dots & \frac{1}{p_k + \bar{p}_{k'}} (1 + e^{-(\xi_k + \bar{\xi}_{k'} - \eta_{k0} - \bar{\eta}_{k'0}))} & e^{-\xi_k + \eta_{k0}} \\ -1 & -1 & 0 \end{vmatrix}. \end{aligned}$$

Thus

$$\tau_{-1}^*(-x, t) = C_1^{-1} \begin{vmatrix} \frac{1}{p_k + \bar{p}_{k'}} (1 + e^{\xi_k + \bar{\xi}_{k'} - \eta_{k0} - \bar{\eta}_{k'0}}) & \dots & e^{\xi_k - \eta_{k0}} \\ \dots & \frac{1}{p_{k'} + \bar{p}_k} (1 + e^{\xi_{k'} + \bar{\xi}_k - \eta_{k'0} - \bar{\eta}_{k0}}) & e^{\xi_{k'} - \eta_{k'0}} \\ -1 & -1 & 0 \end{vmatrix}.$$

We again found $\tau_{-1}(-x, t) = C\tau_{-1}^*(x, t)$. This completes the proof. $\square$

Following above Lemma, if we define $\tau_0(x, t) = \sqrt{C}f(x, t)$, $\tau_1(x, t) = \sqrt{C}g(x, t)$, $\tau_{-1}(x, t) = \sqrt{C}\bar{g}(x, t)$, then we have $f(x, t) = f^*(-x, t)$, $\bar{g}(x, t) = g^*(-x, t)$, which lead to the following bilinear equations

$$\begin{cases} (iD_t - D_x^2)g(x, t) \cdot f(x, t) = 0, \\ D_x^2 f(x, t) \cdot f(x, t) - 2g(x, t)g^*(-x, t) = 0. \end{cases} \quad (27)$$

Moreover, if we define

$$q(x, t) = \frac{g(x, t)}{f(x, t)}, \quad r(x, t) = \frac{-\bar{g}(x, t)}{f(x, t)}, \quad (28)$$

we then have

$$r(x, t) = -q^*(-x, t). \quad (29)$$

In summary, we have the general N -bright soliton solution (28) to the nonlocal NLS equation (6) with $\sigma = -1$.

$$f(x, t) = \frac{1}{\sqrt{C}} \left| \frac{1}{p_i + \bar{p}_j} (e^{(p_i + \bar{p}_j)x - i(p_i^2 - \bar{p}_j^2)t} + e^{(\eta_i + \bar{\eta}_j)}) \right|_{N \times N}, \quad (30)$$

$$g(x, t) = \frac{1}{\sqrt{C}} \left| \begin{array}{cc} \frac{1}{p_i + \bar{p}_j} (e^{(p_i + \bar{p}_j)x - i(p_i^2 - \bar{p}_j^2)t} + e^{(\eta_i + \bar{\eta}_j)}) & e^{p_i x - i p_i^2 t} \\ -e^{\bar{\eta}_j} & 0 \end{array} \right|_{(N+1) \times (N+1)}. \quad (31)$$

In what follows, we list one- and two-soliton solutions:

One-soliton solution:

$$f(x, t) = \frac{1}{p_1 + \bar{p}_1} e^{-\frac{1}{2}(\xi_1 + \bar{\xi}_1 - i\theta_1 - i\bar{\theta}_1)} \left(1 + e^{(p_1 + \bar{p}_1)x - i(p_1^2 - \bar{p}_1^2)t - i(\theta_1 + \bar{\theta}_1)} \right) \quad (32)$$

$$g(x, t) = e^{-\frac{1}{2}(\xi_1 + \bar{\xi}_1 + i\theta_1 + i\bar{\theta}_1)} e^{p_1 x - i p_1^2 t + i\bar{\theta}_1}. \quad (33)$$

So

$$q(x, t) = \frac{(p_1 + \bar{p}_1) e^{p_1 x - i p_1^2 t - i\theta_1}}{1 + e^{(p_1 + \bar{p}_1)x - i(p_1^2 - \bar{p}_1^2)t - i(\theta_1 + \bar{\theta}_1)}} \quad (34)$$

if we let $p_1 = -2\bar{\eta}$, $\bar{p}_1 = -2\eta$, $\theta_1 = -\bar{\theta} + \pi$, $\bar{\theta}_1 = -\theta$, then above solution exactly recovers the one-soliton solution found in [21], also mentioned in the introduction.

Two-soliton solution:

$$f = \left| \begin{array}{cc} \frac{1}{p_1 + \bar{p}_1} (e^{\xi_1 + \bar{\xi}_1} + e^{\eta_{10} + \bar{\eta}_{10}}) & \frac{1}{p_1 + \bar{p}_2} (e^{\xi_1 + \bar{\xi}_2} + e^{\eta_{10} + \bar{\eta}_{20}}) \\ \frac{1}{p_2 + \bar{p}_1} (e^{\xi_2 + \bar{\xi}_1} + e^{\eta_{20} + \bar{\eta}_{10}}) & \frac{1}{p_2 + \bar{p}_2} (e^{\xi_2 + \bar{\xi}_2} + e^{\eta_{20} + \bar{\eta}_{20}}) \end{array} \right|$$

$$= D \left(1 + e^{\xi_1 + \bar{\xi}_1 + \xi_2 + \bar{\xi}_2 - \eta_{10} - \bar{\eta}_{10} - \eta_{20} - \bar{\eta}_{20}} + \frac{(p_1 + \bar{p}_2)(p_2 + \bar{p}_1)}{(p_1 - p_2)(\bar{p}_1 - \bar{p}_2)} (e^{\xi_1 + \bar{\xi}_1 - \eta_{10} - \bar{\eta}_{10}} + e^{\xi_2 + \bar{\xi}_2 - \eta_{20} - \bar{\eta}_{20}}) \right.$$

$$\left. + \frac{(p_1 + \bar{p}_1)(p_2 + \bar{p}_2)}{(p_1 - p_2)(\bar{p}_1 - \bar{p}_2)} (e^{\xi_1 + \bar{\xi}_2 - \eta_{10} - \bar{\eta}_{20}} + e^{\xi_2 + \bar{\xi}_1 - \eta_{20} - \bar{\eta}_{10}}) \right), \quad (35)$$

$$g = \left| \begin{array}{cc} \frac{1}{p_1 + \bar{p}_1} (e^{\xi_1 + \bar{\xi}_1} + e^{\eta_{10} + \bar{\eta}_{10}}) & \frac{1}{p_1 + \bar{p}_2} (e^{\xi_1 + \bar{\xi}_2} + e^{\eta_{10} + \bar{\eta}_{20}}) & e^{\xi_1} \\ \frac{1}{p_2 + \bar{p}_1} (e^{\xi_2 + \bar{\xi}_1} + e^{\eta_{20} + \bar{\eta}_{10}}) & \frac{1}{p_2 + \bar{p}_2} (e^{\xi_2 + \bar{\xi}_2} + e^{\eta_{20} + \bar{\eta}_{20}}) & e^{\xi_2} \\ -e^{\bar{\eta}_{10}} & -e^{\bar{\eta}_{20}} & 0 \end{array} \right|$$

$$= D \left(\frac{(p_1 + \bar{p}_1)(p_2 + \bar{p}_1)}{\bar{p}_1 - \bar{p}_2} e^{\xi_1 + \xi_2 + \bar{\xi}_2 - \eta_{10} - \eta_{20} - \bar{\eta}_{20}} + \frac{(p_1 + \bar{p}_1)(p_1 + \bar{p}_2)}{p_1 - p_2} e^{\xi_1 - \eta_{10}} \right.$$

$$\left. - \left(\frac{(p_1 + \bar{p}_2)(p_2 + \bar{p}_2)}{\bar{p}_1 - \bar{p}_2} e^{\xi_1 + \xi_2 + \bar{\xi}_2 - \eta_{10} - \eta_{20} - \bar{\eta}_{20}} + \frac{(p_2 + \bar{p}_1)(p_2 + \bar{p}_2)}{p_1 - p_2} e^{\xi_2 - \eta_{20}} \right) \right), \quad (36)$$

where

$$D = \sqrt{\frac{C_2}{C_1}} \frac{(p_1 - p_2)(\bar{p}_1 - \bar{p}_2)}{(p_1 + \bar{p}_1)(p_1 + \bar{p}_2)(p_2 + \bar{p}_1)(p_2 + \bar{p}_2)}.$$

Two soliton solution could have the following cases:

- (i) $p_1, p_2, \bar{p}_1, \bar{p}_2$ are all real and $\eta_{10}, \eta_{20}, \bar{\eta}_{10}, \bar{\eta}_{20}$ are all purely imaginary.
- (ii) $p_1, p_2, \bar{p}_1, \bar{p}_2, \eta_{10}, \eta_{20}, \bar{\eta}_{10}, \bar{\eta}_{20}$ are all complex, $p_1 = p_2^*, \bar{p}_1 = \bar{p}_2^*, \eta_{10} = -\eta_{20}^*, \bar{\eta}_{10} = \bar{\eta}_{20}^*$.
- (iii) $p_1, p_2, \eta_{10}, \eta_{20}$ are complex number, $p_1 = p_2^*, \eta_{10} = -\eta_{20}^*$; $\bar{p}_1, \bar{p}_2$ are real and $\bar{\eta}_{10}, \bar{\eta}_{20}$ are all purely imaginary.
- (iv) $\bar{p}_1, \bar{p}_2, \bar{\eta}_{10}, \bar{\eta}_{20}$ are complex number, $\bar{p}_1 = \bar{p}_2^*, \bar{\eta}_{10} = \bar{\eta}_{20}^*$; p_1, p_2 are real and η_{10}, η_{20} are all purely imaginary.

It is noted that all above four cases regarding the bright soliton solution to the nonlocal NLS equation have been recognized in [57], their dynamics and singularities have been discussed in details there.

2.2. General bright soliton solution expressed by double Wronskian determinant

Alternatively, we can also present the general bright soliton solution to the nonlocal NLS equation (6) in terms of the double Wronskian determinant. To this end, we start with the tau functions for two-component KP hierarchy expressed in double Wronskian determinant

$$\tau_{N,M} = \begin{vmatrix} \phi_1^{(0)} & \cdots & \phi_1^{(N-1)} & \psi_1^{(0)} & \cdots & \psi_1^{(M-1)} \\ \phi_2^{(0)} & \cdots & \phi_2^{(N-1)} & \psi_2^{(0)} & \cdots & \psi_2^{(M-1)} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \phi_{N+M}^{(0)} & \cdots & \phi_{N+M}^{(N-1)} & \psi_{N+M}^{(0)} & \cdots & \psi_{N+M}^{(M-1)} \end{vmatrix}_{(N+M) \times (N+M)}, \quad (37)$$

here $\phi_i^{(n)}$ and $\psi_i^{(n)}$ take the form

$$\phi_i^{(n)} = p_i^n e^{\xi_i}, \quad \psi_i^{(n)} = q_i^n e^{\eta_i},$$

with

$$\xi_i = p_i x_1 + p_i^2 x_2 + \xi_{0i} + \cdots, \quad \eta_i = q_i y_1 + \eta_{0i} + \cdots.$$

The above tau functions satisfy the following bilinear equations

$$\begin{cases} (D_{x_2} - D_{x_1}^2) \tau_{N+1,N-1} \cdot \tau_{N,N} = 0, \\ (D_{x_2} - D_{x_1}^2) \tau_{N,N} \cdot \tau_{N-1,N+1} = 0, \\ D_{x_1} D_{y_1} \tau_{N,N} \cdot \tau_{N,N} - 2\tau_{N+1,N-1} \tau_{N-1,N+1} = 0. \end{cases} \quad (38)$$

The proof of above equations can be done by using the determinant technique [71], which is omitted here. In what follows, we will perform reduction procedure from bilinear equation (38) to the bilinear equation (12). For the sake of convenience, we take the following abbreviation

$$\tau_{N,M} = |0, \dots, N-1, 0', \dots, (M-1)'|.$$

First, we consider the dimension reduction. Imposing the condition $N = M$ and $q_i = p_i$ ($i = 1, \dots, 2N$), we then have

$$(\partial_{x_1} + \partial_{y_1})\tau_{N,N} = \left(\sum_{j=1}^{2N} p_j \right) \tau_{N,N}. \quad (39)$$

Under this reduction, y_1 becomes a dummy variable, which can be taken as zero. Applying variable transformations

$$x_1 = x, \quad x_2 = -it, \quad (40)$$

which implies

$$\partial_{x_1} = \partial_x, \quad \partial_{x_2} = i\partial_t, \quad (41)$$

then the bilinear equation (38) can be rewritten as

$$\begin{cases} (iD_t - D_x^2)\tau_{N+1,N-1} \cdot \tau_{N,N} = 0, \\ (iD_t - D_x^2)\tau_{N,N} \cdot \tau_{N-1,N+1} = 0, \\ D_x^2\tau_{N,N} \cdot \tau_{N,N} + 2\tau_{N+1,N-1}\tau_{N-1,N+1} = 0. \end{cases} \quad (42)$$

Lemma 2.2. Assume $x_1 = x$ is real and $x_2 = -it$ is a purely imaginary variable. Suppose there are K pairs of wave numbers (p_k, η_{k0}) and $(\bar{p}_k, \bar{\eta}_{k0})$ for $k = 1, \dots, K$ such that $p_k = p_k^*$, $\eta_{k0} = \eta_{k'0}^*$ or $\bar{p}_k = \bar{p}_{k'}^*$, $\bar{\eta}_{k0} = \bar{\eta}_{k'0}^*$; and the rest of wave numbers p_j and $\bar{p}_j$ are real, $\eta_{j0}, \bar{\eta}_{j0}$ are purely imaginary. Then

$$\tau_{N,N}(x, t) = C'(-1)^{N+K}\tau_{N,N}^*(-x, t), \quad (43)$$

$$\tau_{N-1,N+1}(x, t) = C'(-1)^{N+K+1}\tau_{N+1,N-1}^*(-x, t), \quad (44)$$

where $C' = \prod_{i=1}^N e^{\xi_i(x,t) - \eta_i}$.

Proof. We only give the proof when p_j and $\bar{p}_j$ are real and $\eta_{j0}, \bar{\eta}_{j0}$ are purely imaginary. Under this case, $\xi_i(x, t) = p_i x - ip_i^2 t = -\xi_i^*(-x, t)$, then it can be shown

$$\begin{aligned} \tau_{N-1,N+1}(x, t) &= |0, \dots, N-2, 0', \dots, N'|, \\ &= \left| e^{\xi_i(x,t)}, \dots, p_i^{N-2} e^{\xi_i(x,t)}, e^{-i\theta_i}, \dots, p_i^N e^{-i\theta_i} \right|, \\ &= \prod_{i=1}^N e^{\xi_i(x,t) - i\theta_i} \left| e^{i\theta_i}, \dots, p_i^{N-2} e^{i\theta_i}, e^{-\xi_i(x,t)}, \dots, p_i^N e^{-\xi_i(x,t)} \right|, \\ &= (-1)^N \prod_{i=1}^N e^{\xi_i(x,t) - i\theta_i} \left| e^{-\xi_i(x,t)}, \dots, p_i^N e^{-\xi_i(x,t)}, e^{i\theta_i}, \dots, p_i^{N-2} e^{i\theta_i} \right|, \\ &= (-1)^N \prod_{i=1}^N e^{\xi_i(x,t) - i\theta_i} \left| e^{\xi_i^*(-x,t)}, \dots, p_i^N e^{\xi_i^*(-x,t)}, e^{i\theta_i}, \dots, p_i^{N-2} e^{i\theta_i} \right|, \\ &= (-1)^{N+1} C' \tau_{N+1,N-1}^*(-x, t). \end{aligned} \quad (45)$$

$$\begin{aligned}
\tau_{N,N}(x,t) &= |0, \dots, N-1, 0', \dots, (N-1)'|, \\
&= \left| e^{\xi_i(x,t)}, \dots, p_i^{N-1} e^{\xi_i(x,t)}, e^{-i\theta_i}, \dots, p_i^{N-1} e^{-i\theta_i} \right|, \\
&= \prod_{i=1}^N e^{\xi_i(x,t)-i\theta_i} \left| e^{i\theta_i}, \dots, p_i^{N-1} e^{i\theta_i}, e^{-\xi_i(x,t)}, \dots, p_i^{N-1} e^{-\xi_i(x,t)} \right|, \\
&= (-1)^{N+1} \prod_{i=1}^N e^{\xi_i(x,t)-i\theta_i} \left| e^{-\xi_i(x,t)}, \dots, p_i^{N-1} e^{-\xi_i(x,t)}, e^{i\theta_i}, \dots, p_i^{N-1} e^{i\theta_i} \right|, \\
&= (-1)^N \prod_{i=1}^N e^{\xi_i(x,t)-i\theta_i} \left| e^{\xi_i^*(-x,t)}, \dots, p_i^{N-1} e^{\xi_i^*(-x,t)}, e^{i\theta_i}, \dots, p_i^{N-1} e^{i\theta_i} \right|, \\
&= (-1)^{N+1} C' \tau_{N,N}^*(-x,t).
\end{aligned} \tag{46}$$

For the case of complex conjugate wavenumber pairs, it can be approved the same way as the Lemma in previous subsection. So we omit the proof here. $\square$

Therefore if we define $\tau_{N,N}(x,t) = \sqrt{C'}f(x,t)$, $\tau_{N+1,N-1}(x,t) = \sqrt{C'}g(x,t)$, $\tau_{N-1,N+1}(x,t) = -\sqrt{C'}\bar{g}(x,t)$, then the following bilinear equations follow

$$\begin{cases} (iD_t - D_x^2)g(x,t) \cdot f(x,t) = 0, \\ D_x^2 f(x,t) \cdot f(x,t) - 2g(x,t)g^*(-x,t) = 0. \end{cases} \tag{47}$$

Moreover, if we define

$$q(x,t) = \frac{g(x,t)}{f(x,t)}, \quad r(x,t) = \frac{\bar{g}(x,t)}{f(x,t)}, \tag{48}$$

we then have

$$r(x,t) = -q^*(-x,t). \tag{49}$$

To summarize, we have the general N -bright soliton solution expressed by

$$q(x,t) = \frac{g'(x,t)}{f'(x,t)}, \tag{50}$$

where

$$f'(x,t) = \begin{vmatrix} \tilde{\phi}_1^{(0)} & \dots & \tilde{\phi}_1^{(N-1)} & \tilde{\psi}_1^{(0)} & \dots & \tilde{\psi}_1^{(N-1)} \\ \tilde{\phi}_2^{(0)} & \dots & \tilde{\phi}_2^{(N-1)} & \tilde{\psi}_2^{(0)} & \dots & \tilde{\psi}_2^{(N-1)} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \tilde{\phi}_{2N}^{(0)} & \dots & \tilde{\phi}_{2N}^{(N-1)} & \tilde{\psi}_{2N}^{(0)} & \dots & \tilde{\psi}_{2N}^{(N-1)} \end{vmatrix}_{2N \times 2N}, \tag{51}$$

$$g'(x,t) = \begin{vmatrix} \tilde{\phi}_1^{(0)} & \dots & \tilde{\phi}_1^{(N)} & \tilde{\psi}_1^{(0)} & \dots & \tilde{\psi}_1^{(N-2)} \\ \tilde{\phi}_2^{(0)} & \dots & \tilde{\phi}_2^{(N)} & \tilde{\psi}_2^{(0)} & \dots & \tilde{\psi}_2^{(N-2)} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \tilde{\phi}_{2N}^{(0)} & \dots & \tilde{\phi}_{2N}^{(N)} & \tilde{\psi}_{2N}^{(0)} & \dots & \tilde{\psi}_{2N}^{(N-2)} \end{vmatrix}_{2N \times 2N}, \tag{52}$$

here $\tilde{\phi}_i^{(n)}$ and $\tilde{\psi}_i^{(n)}$ take the form

$$\tilde{\phi}_i^{(n)} = p_i^n e^{p_i x - i p_i^2 t}, \quad \tilde{\psi}_i^{(n)} = p_i^n e^{-i \theta_i}.$$

In what follows, we will only illustrate one-soliton solution and make a comment. By taking $N = 1$, we get the tau functions for one-soliton solution,

$$f'(x, t) = \begin{vmatrix} e^{\xi_1} & e^{-i\theta_1} \\ e^{\xi_2} & e^{-i\theta_2} \end{vmatrix} = e^{\xi_1 - i\theta_2} (1 - e^{i\theta_2 - i\theta_1} e^{\xi_2 - \xi_1}) \quad (53)$$

$$g'(x, t) = \begin{vmatrix} e^{\xi_1} & p_1 e^{\xi_1} \\ e^{\xi_2} & p_2 e^{\xi_2} \end{vmatrix} = (p_2 - p_1) e^{\xi_2 + \xi_1}. \quad (54)$$

The above tau functions lead to the one-soliton as follows

$$q(x, t) = \frac{(p_2 - p_1) e^{\xi_2 + \xi_1}}{e^{\xi_1 - i\theta_2} (1 - e^{i\theta_2 - i\theta_1} e^{\xi_2 - \xi_1})} = \frac{(p_1 - p_2) e^{\xi_1 + i\theta_1}}{(1 - e^{i\theta_1 - i\theta_2} e^{\xi_1 - \xi_2})} \quad (55)$$

By having $p_2 \rightarrow -\bar{p}_1$, $\theta_1 \rightarrow -\theta_1$ and $\theta_2 \rightarrow \pi + \theta_2$, above one soliton solution (55) coincides with the one-soliton solution (34) obtained in previous subsection, thus the same solution found in [21].

3. Soliton solution to the nonlocal NLS equation with nonzero boundary condition

In this section, we consider the general soliton solution to the nonlocal NLS equation (6) with the same nonzero boundary condition as considered in [23]

$$q(x, t) \rightarrow \rho e^{i\theta_{\pm}}, \quad (\rho > 0) \quad \text{as } x \rightarrow \pm\infty, \quad (56)$$

where $\Delta\theta = \theta_+ - \theta_-$ is either 0 or π . Similar to the classical NLS equation, to construct soliton solutions of the nonlocal NLS equation with NZBCs, we need to start with the tau functions for single-component KP hierarchy expressed in Gram-type determinants

$$\tau_k = |m_{ij}(k)|_{1 \leq i, j \leq M} \quad (57)$$

where

$$m_{ij}(k) = c_{ij} + \frac{1}{p_i + \bar{p}_j} \left(-\frac{p_i}{\bar{p}_j} \right)^k e^{\xi_i + \bar{\xi}_j} \quad (58)$$

$$\xi_i = p_i^{-1} x_{-1} + p_i x_1 + p_i^2 x_2 + \xi_{0i} + \cdots$$

$$\bar{\xi}_i = \bar{p}_i^{-1} x_{-1} + \bar{p}_i x_1 - \bar{p}_i^2 x_2 + \bar{\xi}_{0i} + \cdots$$

Based on the Sato theory [72, 74, 75] or the Jacobi identity of determinants, the above tau functions satisfies the following bilinear equations

$$\begin{cases} (D_{x_2} - D_{x_1}^2) \tau_{k+1} \cdot \tau_k = 0, \\ (\frac{1}{2} D_{x_1} D_{x_{-1}} - 1) \tau_k \cdot \tau_k = -\tau_{k+1} \tau_{k-1}. \end{cases} \quad (59)$$

As stated in the subsequent sections, above two set of bilinear equations will be the key in constructing soliton solutions to the nonlocal NLS equation. We also assume $c_{ij} = c_i \delta_{ij}$ hereafter.

3.1. General soliton solution to the nonlocal NLS equation of $\sigma = 1$, $\Delta\theta = 0$ with nonzero boundary condition

The nonlocal NLS equation (6) of $\sigma = 1$ is converted into a set of bilinear equations

$$\begin{cases} (iD_t - D_x^2)g \cdot f = 0, \\ (D_x^2 - 2\rho^2)f \cdot f = -2\rho^2 g(x, t)g^*(-x, t), \end{cases} \quad (60)$$

via a variable transformation

$$q(x, t) = \rho \frac{g(x, t)}{f(x, t)} e^{i2\rho^2 t} \quad (61)$$

under the condition $f(x, t) = f^*(-x, t)$. Here ρ is a positive constant representing the amplitude of the nonzero plane wave background as $x \rightarrow \infty$. In what follows, we will show how to reduce the bilinear equation (59) to the bilinear equation (60) by the KP hierarchy reduction method. We remark here that similar reductions starting from the (constraint) KP hierarchy have been done to find various soliton solutions to the NLS equation in the literature [81–84].

Firstly, we perform the dimension reduction. Note that, by row operations, τ_k can be rewritten as

$$\tau_k = \prod_{j=1}^N e^{\xi_j + \bar{\xi}_j} |m'_{ij}| = \prod_{j=1}^N e^{\xi_j + \bar{\xi}_j} \left| c_i \delta_{ij} e^{-(\xi_i + \bar{\xi}_i)} + \frac{1}{p_i + \bar{p}_j} \left(-\frac{p_i}{\bar{p}_j} \right)^k \right|. \quad (62)$$

Since

$$(\partial_{x_1} - \rho^2 \partial_{x_{-1}})m'_{ij} = (p_i + \bar{p}_j) \left(1 - \frac{\rho^2}{p_i \bar{p}_j} \right) c_i \delta_{ij} e^{-(\xi_i + \bar{\xi}_i)}, \quad (63)$$

thus, if we impose the constraints

$$p_j \bar{p}_j = \rho^2, \quad j = 1, \dots, N, \quad (64)$$

then

$$\partial_{x_1} m'_{ij} = \rho^2 \partial_{x_{-1}} m'_{ij}, \quad (65)$$

which leads to

$$\partial_{x_1} \tau_k = \rho^2 \partial_{x_{-1}} \tau_k. \quad (66)$$

Therefore, the bilinear equation (59) become

$$\begin{cases} (D_{x_2} - D_{x_1}^2)\tau_{k+1} \cdot \tau_k = 0, \\ (D_{x_1}^2 - 2\rho^2)\tau_k \cdot \tau_k = -2\rho^2 \tau_{k+1} \tau_{k-1}. \end{cases} \quad (67)$$

Furthermore, we assume $x_1 = x$, $x_2 = -it$ and define

$$\tau_0(x, t) = Cf(x, t), \quad \tau_1(x, t) = Cg(x, t), \quad \tau_{-1}(x, t) = C\bar{g}(x, t), \quad (68)$$

with $C = \prod_{j=1}^N e^{\xi_j + \bar{\xi}_j}$, then the bilinear equations become

$$\begin{cases} (iD_t - D_x^2)g(x, t) \cdot f(x, t) = 0, \\ (D_x^2 - 2\rho^2)f(x, t) \cdot f(x, t) = -2\rho^2 g(x, t)\bar{g}(x, t). \end{cases} \quad (69)$$

Lemma 3.3. Consider $2N \times 2N$ matrices for $f(x, t)$, $g(x, t)$ and $\bar{g}(x, t)$. If $\bar{p}_j = p_j^*$ to be the complex conjugate of p_j , let $c_{N+j} = -c_j^*$, $p_{N+j} = -p_j$ to be complex, for $j = 1, \dots, N$, and $\xi_{0,N+j} = \xi_{0j}$ to be real, then we have

$$f(x, t) = f^*(-x, t), \quad \bar{g}(x, t) = g^*(-x, t). \quad (70)$$

Proof. Since

$$(\xi_i + \bar{\xi}_i)(x, t) = (p_i + p_i^*)x - (p_i^2 - (p_i^*)^2)it + \xi_{0i},$$

and

$$\begin{aligned} (\xi_{N+i} + \bar{\xi}_{N+i})(x, t) &= (p_{N+i} + p_{N+i}^*)x - (p_{N+i}^2 - (p_{N+i}^*)^2)it + \xi_{0, N+i} \\ &= -(p_i + p_i^*)x - (p_i^2 - (p_i^*)^2)it + \xi_{0i}, \end{aligned}$$

so we have

$$(\xi_{N+i} + \bar{\xi}_{N+i})(x, t) = (\xi_i + \bar{\xi}_i)^*(-x, t), \quad (\xi_i + \bar{\xi}_i)(x, t) = (\xi_{N+i} + \bar{\xi}_{N+i})^*(-x, t).$$

Note that

$$\begin{aligned} f(x, t) &= \left| \begin{array}{cc} c_i \delta_{ij} e^{-(\xi_i + \bar{\xi}_j)} + \frac{1}{p_i + p_j^*} & \frac{1}{p_i + p_{N+j}^*} \\ \frac{1}{p_{N+i} + p_j^*} & c_{N+i} \delta_{N+i, N+j} e^{-(\xi_{N+i} + \bar{\xi}_{N+j})} + \frac{1}{p_{N+i} + p_{N+j}^*} \end{array} \right|_{1 \leq i, j \leq N} \\ &= \left| \begin{array}{cc} c_{N+i} \delta_{N+i, N+j} e^{-(\xi_{N+i} + \bar{\xi}_{N+j})} + \frac{1}{p_{N+i} + p_{N+j}^*} & \frac{1}{p_{N+i} + p_j^*} \\ \frac{1}{p_i + p_{N+j}^*} & c_i \delta_{ij} e^{-(\xi_i + \bar{\xi}_j)} + \frac{1}{p_i + p_j^*} \end{array} \right| \\ &= \left| \begin{array}{cc} -c_i^* \delta_{ij} e^{-(\xi_{N+i} + \bar{\xi}_{N+j})} - \frac{1}{p_i + p_j^*} & -\frac{1}{p_i + p_{N+j}^*} \\ -\frac{1}{p_{N+i} + p_j^*} & -c_{N+i}^* \delta_{i, j} e^{-(\xi_i + \bar{\xi}_j)} - \frac{1}{p_{N+i} + p_{N+j}^*} \end{array} \right| \\ &= \left| \begin{array}{cc} c_i^* \delta_{ij} e^{-(\xi_{N+i} + \bar{\xi}_{N+j})} + \frac{1}{p_i + p_j^*} & \frac{1}{p_i + p_{N+j}^*} \\ \frac{1}{p_{N+i} + p_j^*} & c_{N+i}^* \delta_{i, j} e^{-(\xi_i + \bar{\xi}_j)} + \frac{1}{p_{N+i} + p_{N+j}^*} \end{array} \right|. \end{aligned} \quad (71)$$

On the other hand,

$$\begin{aligned} f(x, t) &= \left| \begin{array}{cc} c_i \delta_{ij} e^{-(\xi_i + \bar{\xi}_j)} + \frac{1}{p_i + p_j^*} & \frac{1}{p_i + p_{N+j}^*} \\ \frac{1}{p_{N+i} + p_j^*} & c_{N+i} \delta_{N+i, N+j} e^{-(\xi_{N+i} + \bar{\xi}_{N+j})} + \frac{1}{p_{N+i} + p_{N+j}^*} \end{array} \right| \\ &= \left| \begin{array}{cc} c_j \delta_{ji} e^{-(\xi_j + \bar{\xi}_i)} + \frac{1}{p_j + p_i^*} & \frac{1}{p_i^* + p_{N+j}} \\ \frac{1}{p_{N+i}^* + p_j} & c_{N+j} \delta_{N+j, N+i} e^{-(\xi_{N+j} + \bar{\xi}_{N+i})} + \frac{1}{p_{N+i}^* + p_{N+j}} \end{array} \right|, \end{aligned} \quad (72)$$

thus

$$f^*(-x, t) = \left| \begin{array}{cc} c_j \delta_{ji} e^{-(\xi_j + \bar{\xi}_i)^*(-x, t)} + \frac{1}{p_j + p_i^*} & \frac{1}{p_i^* + p_{N+j}} \\ \frac{1}{p_{N+i}^* + p_j} & c_{N+j} \delta_{N+j, N+i} e^{-(\xi_{N+j} + \bar{\xi}_{N+i})^*(-x, t)} + \frac{1}{p_{N+i}^* + p_{N+j}} \end{array} \right|. \quad (73)$$

Obviously $f(x, t) = f^*(-x, t)$. Next, let us proceed to prove $g(x, t) = g^*(-x, t)$. Note $g(x, t)$ can be written as

$$\begin{aligned}
g(x, t) &= \left| c_i \delta_{ij} \left(-\frac{p_i}{p_j^*} \right) e^{-(\xi_i + \bar{\xi}_j)} + \frac{1}{p_i + p_j^*} \frac{1}{p_i + p_{N+j}^*}, \right. \\
&\quad \left. \frac{1}{p_{N+i} + p_j^*} c_{N+i} \delta_{N+i, N+j} \left(-\frac{p_{N+i}}{p_{N+j}^*} \right) e^{-(\xi_{N+i} + \bar{\xi}_{N+j})} + \frac{1}{p_{N+i} + p_{N+j}^*} \right|, \\
&= \left| c_{N+i} \delta_{N+i, N+j} \left(-\frac{p_{N+i}}{p_{N+j}^*} \right) e^{-(\xi_{N+i} + \bar{\xi}_{N+j})} + \frac{1}{p_{N+i} + p_{N+j}^*} \frac{1}{p_{N+i} + p_j^*}, \right. \\
&\quad \left. \frac{1}{p_i + p_{N+j}^*} c_i \delta_{ij} \left(-\frac{p_i}{p_j^*} \right) e^{-(\xi_i + \bar{\xi}_j)} + \frac{1}{p_i + p_j^*} \right|, \\
&= \left| -c_i^* \delta_{ij} \left(-\frac{p_i}{p_j^*} \right) e^{-(\xi_{N+i} + \bar{\xi}_{N+j})} - \frac{1}{p_i + p_j^*} \frac{1}{p_i + p_{N+j}^*}, \right. \\
&\quad \left. -\frac{1}{p_{N+i} + p_j^*} -c_{N+i}^* \left(-\frac{p_{N+i}}{p_{N+j}^*} \right) \delta_{N+i, N+j} e^{-(\xi_i + \bar{\xi}_j)} - \frac{1}{p_{N+i} + p_{N+j}^*} \right|, \\
&= \left| c_i^* \delta_{ij} \left(-\frac{p_i}{p_j^*} \right) e^{-(\xi_{N+i} + \bar{\xi}_{N+j})} + \frac{1}{p_i + p_j^*} \frac{1}{p_i + p_{N+j}^*}, \right. \\
&\quad \left. \frac{1}{p_{N+i} + p_j^*} c_{N+i}^* \left(-\frac{p_{N+i}}{p_{N+j}^*} \right) \delta_{N+i, N+j} e^{-(\xi_i + \bar{\xi}_j)} + \frac{1}{p_{N+i} + p_{N+j}^*} \right|, \quad (74)
\end{aligned}$$

thus

$$g^*(-x, t) = \left| c_i \delta_{ij} \left(-\frac{p_i^*}{p_j} \right) e^{-(\xi_{N+i} + \bar{\xi}_{N+j})^*(-x, t)} + \frac{1}{p_i^* + p_j} \frac{1}{p_i^* + p_{N+j}}, \right. \\
\left. \frac{1}{p_{N+i}^* + p_j} c_{N+i} \left(-\frac{p_{N+i}^*}{p_{N+j}} \right) \delta_{ij} e^{-(\xi_i + \bar{\xi}_j)^*(-x, t)} + \frac{1}{p_{N+i}^* + p_{N+j}} \right|. \quad (75)$$

On the other hand,

$$\begin{aligned}
\bar{g}(x, t) &= \left| c_i \delta_{ij} \left(-\frac{p_i}{p_j^*} \right)^{-1} e^{-(\xi_i + \bar{\xi}_j)} + \frac{1}{p_i + p_j^*} \frac{1}{p_i + p_{N+j}^*}, \right. \\
&\quad \left. \frac{1}{p_{N+i} + p_j^*} c_{N+i} \delta_{N+j, N+i} \left(-\frac{p_{N+i}}{p_{N+j}^*} \right)^{-1} e^{-(\xi_{N+i} + \bar{\xi}_{N+j})} + \frac{1}{p_{N+i} + p_{N+j}^*} \right|, \\
&= \left| c_j \delta_{ji} \left(-\frac{p_j}{p_i^*} \right)^{-1} e^{-(\xi_j + \bar{\xi}_i)} + \frac{1}{p_j + p_i^*} \frac{1}{p_j + p_{N+i}^*}, \right. \\
&\quad \left. \frac{1}{p_j + p_{N+i}^*} c_{N+j} \delta_{N+j, N+i} \left(-\frac{p_{N+j}}{p_{N+i}^*} \right)^{-1} e^{-(\xi_{N+j} + \bar{\xi}_{N+i})} + \frac{1}{p_{N+j} + p_{N+i}^*} \right|.
\end{aligned}$$

Obviously $\bar{g}(x, t) = g^*(-x, t)$. $\square$

Based on above Lemma, two sets of bilinear equations (60) and (69) coincide. In summary, the nonlocal NLS equation admits the following general soliton solution with nonzero boundary conditions.

$$q(x, t) = \rho \frac{g(x, t)}{f(x, t)} e^{i2\rho^2 t}, \quad (76)$$

where

$$f(x, t) = \left| c_i \delta_{ij} e^{-(\xi_i + \bar{\xi}_j)} + \frac{1}{p_i + p_j^*} \right|_{0 \leq i, j \leq 2N}, \quad (77)$$

$$g(x, t) = \left| c_i \delta_{ij} \left(-\frac{p_i}{p_j^*} \right) e^{-(\xi_i + \bar{\xi}_j)} + \frac{1}{p_i + p_j^*} \right|_{0 \leq i, j \leq 2N}, \quad (78)$$

with $c_{N+i} = -c_i^*$, $p_{N+i} = -p_i$, $\xi_{0, N+i} = \xi_{0i}$ for $i = 1, \dots, N$. In what follows, we give explicit form of the solution for $N = 1$ and detailed analysis. By taking $N = 1$ in (77) and (78) and $p_1 = \rho e^{-i\theta}$ we get the tau functions

$$\begin{aligned}
f &= \left| c_1 e^{-(\xi_1 + \bar{\xi}_1)} + \frac{1}{p_1 + p_1^*} \frac{1}{\frac{1}{p_2 + p_2^*}} - c_1^* e^{-(\xi_2 + \bar{\xi}_2)} + \frac{1}{p_2 + p_2^*} \right| \\
&= -\rho^{-2} (|c_1|^2 \rho^2 e^{4\rho^2 t \sin 2\theta} + \frac{1}{2} \rho c_1 \sec \theta e^{-2\rho x \cos \theta + 2\rho^2 t \sin 2\theta} \\
&\quad + \frac{1}{2} \rho c_1^* \sec \theta e^{2\rho x \cos \theta + 2\rho^2 t \sin 2\theta} + \csc^2 2\theta), \tag{79}
\end{aligned}$$

$$\begin{aligned}
g &= \left| c_1 e^{-(\xi_1 + \bar{\xi}_1)} + \frac{1}{p_1 + p_1^*} \left(-\frac{p_1}{p_1^*} \right) \frac{1}{\frac{1}{p_2 + p_2^*}} \left(-\frac{p_1}{p_2^*} \right) \right. \\
&\quad \left. - c_1^* e^{-(\xi_2 + \bar{\xi}_2)} + \frac{1}{p_2 + p_2^*} \left(-\frac{p_2}{p_2^*} \right) \right| \\
&= -e^{2i\theta} \rho^{-2} (|c_1|^2 \rho^2 e^{4\rho^2 t \sin 2\theta - 2i\theta} - \frac{1}{2} \rho c_1 \sec \theta e^{-2\rho x \cos \theta + 2\rho^2 t \sin 2\theta} \\
&\quad - \frac{1}{2} \rho c_1^* \sec \theta e^{2\rho x \cos \theta + 2\rho^2 t \sin 2\theta} + \csc^2 2\theta e^{2i\theta}). \tag{80}
\end{aligned}$$

It can be easily shown that the resulting solution $q(x, t) = \rho \frac{g(x, t)}{f(x, t)} e^{i2\rho^2 t}$ corresponds to the solution (5.90) in [23] with $\sigma = 1$ and $\Delta\theta = 0$, and also the solution (12) in [31]. As analyzed in [31], there are three different cases which include dark–dark, dark–antidark (the amplitude of soliton is greater than the amplitude of the plane wave background) and antidark–antidark soliton interactions. This is the new phenomena in compared with the classical NLS equation. It is also noted that the nonsingular condition for the solution was also given by (13) in [31]. These three cases are illustrated in figures 1–3 below. Next, we investigate the asymptotic behavior of above two-soliton solution. To this end, we assume $0 < \theta < \pi/2$ without loss of generality, and $\kappa = 2\rho \cos \theta > 0$. We define the right-moving soliton along the line $\eta_1 = x - 2\rho t \sin \theta$ as soliton 1, and the left-moving soliton along the line $\eta_2 = x + 2\rho t \sin \theta$ as soliton 2. For the above choice of parameters, we have (i) $\eta_1 \approx 0$, $\eta_2 \rightarrow \mp\infty$ as $t \rightarrow \mp\infty$ for soliton 1 and (ii) $\eta_2 \approx 0$, $\eta_1 \rightarrow \pm\infty$ as $t \rightarrow \mp\infty$ for soliton 2. This leads to the following asymptotic forms for the two-soliton solution.

(i) Before collision ($t \rightarrow -\infty$)

Soliton 1 ($\eta_1 \approx 0$, $\eta_2 \rightarrow -\infty$):

$$\begin{aligned}
q &\rightarrow \rho \frac{\csc^2 \theta - \frac{1}{2} \rho c_1 \sec \theta e^{-\kappa \eta_1 - 2i\theta}}{\csc^2 \theta + \frac{1}{2} \rho c_1 \sec \theta e^{-\kappa \eta_1}} e^{i2\rho^2 t + 4i\theta}, \\
&= \rho \frac{1 - 2\rho |c_1| \sin^2 \theta \cos \theta e^{-\kappa \eta_1 - i(2\theta - \phi)}}{1 + 2\rho |c_1| \sin^2 \theta \cos \theta e^{-\kappa \eta_1 + i\phi}} e^{i2\rho^2 t + 4i\theta}, \tag{81}
\end{aligned}$$

$$|q|^2 \rightarrow \rho^2 \frac{\cosh(\eta_1 - \eta_{10}^-) - \cos(2\theta - \phi)}{\cosh(\eta_1 - \eta_{10}^-) + \cos \phi}. \tag{82}$$

Soliton 2 ($\eta_2 \approx 0$, $\eta_1 \rightarrow \infty$):

$$\begin{aligned}
q &\rightarrow \rho \frac{\csc^2 2\theta - \frac{1}{2} \rho c_1^* \sec \theta e^{-\kappa \eta_2 - 2i\theta}}{\csc^2 2\theta + \frac{1}{2} \rho c_1^* \sec \theta e^{-\kappa \eta_2}} e^{i2\rho^2 t + 4i\theta}, \\
&= \rho \frac{1 - 2\rho |c_1| \sin^2 \theta \cos \theta e^{-\kappa \eta_1 - i(2\theta + \phi)}}{1 + 2\rho |c_1| \sin^2 \theta \cos \theta e^{-\kappa \eta_1 - i\phi}} e^{i2\rho^2 t + 4i\theta}, \tag{83}
\end{aligned}$$




Figure 1. Dark–dark soliton with $\rho = 2.0$, $c_1 = 1.0 + 0.4i$ and $\theta = \pi/6$.




Figure 2. Dark–antidark soliton with $\rho = 2.0$, $c_1 = 0.4 + i$ and $\theta = \pi/3$.




Figure 3. Antidark–antidark soliton with $\rho = 2.0$, $c_1 = -1.0 + 0.3i$ and $\theta = \pi/3$.

$$|q|^2 \rightarrow \rho^2 \frac{\cosh(\eta_1 - \eta_{10}^-) - \cos(2\theta + \phi)}{\cosh(\eta_1 - \eta_{10}^-) + \cos \phi}, \quad (84)$$

where $e^{\eta_{10}^-} = 2\rho|c_1|\sin^2 \theta \cos \theta$.

(ii) After collision ($t \rightarrow \infty$)

Soliton 1 ($\eta_1 \approx 0, \eta_2 \rightarrow \infty$):

$$\begin{aligned} q &\rightarrow \rho \frac{|c_1|^2 \rho^2 - \frac{1}{2} \rho c_1^* \sec \theta e^{\kappa \eta_1 + 2i\theta}}{|c_1|^2 \rho^2 + \frac{1}{2} \rho c_1^* \sec \theta e^{\kappa \eta_1}} e^{i2\rho^2 t}, \\ &= \rho \frac{2\rho|c_1| \cos \theta e^{-\kappa \eta_1 - i(2\theta + \phi)} - 1}{2\rho|c_1| \cos \theta e^{-\kappa \eta_1 + i\phi} + 1} e^{i2\rho^2 t + 4i\theta} \end{aligned} \quad (85)$$

$$|q|^2 \rightarrow \rho^2 \frac{\cosh(\eta_1 - \eta_{10}^+) - \cos(2\theta - \phi)}{\cosh(\eta_1 - \eta_{10}^+) + \cos \phi}. \quad (86)$$

Soliton 2 ($\eta_2 \approx 0, \eta_1 \rightarrow -\infty$):

$$\begin{aligned} q &\rightarrow \rho \frac{|c_1|^2 \rho^2 - \frac{1}{2} \rho c_1 \sec \theta e^{\kappa \eta_2 + 2i\theta}}{|c_1|^2 \rho^2 + \frac{1}{2} \rho c_1 \sec \theta e^{\kappa \eta_2}} e^{i2\rho^2 t}, \\ &= \rho \frac{2\rho|c_1| \cos \theta e^{-\kappa \eta_1 - i(2\theta + \phi)} - 1}{2\rho|c_1| \cos \theta e^{-\kappa \eta_1 - i\phi} + 1} e^{i2\rho^2 t + 4i\theta} \end{aligned} \quad (87)$$

$$|q|^2 \rightarrow \rho^2 \frac{\cosh(\eta_1 - \eta_{10}^+) - \cos(2\theta + \phi)}{\cosh(\eta_1 - \eta_{10}^+) + \cos \phi}, \quad (88)$$

where $e^{\eta_{10}^+} = 2\rho|c_1| \cos \theta$.

3.2. Soliton solution with nonzero boundary condition to the nonlocal NLS equation with $\sigma = -1, \Delta\theta = \pi$

In this subsection, we consider the soliton solutions to the nonlocal NLS equation with $\sigma = -1$

$$iq_t(x, t) = q_{xx}(x, t) + 2q^*(-x, t) q^2(x, t) \quad (89)$$

under boundary condition $q(x, t) \rightarrow \rho$ as $x \rightarrow \pm\infty$. To this end, we introduce a variable transformation

$$q(x, t) = \rho \frac{g(x, t)}{f(x, t)} e^{2i\epsilon\rho^2 t},$$

with the condition $f(x, t) = \pm f^*(-x, t)$, then the nonlocal NLS equation is converted into the following bilinear equations

$$\begin{cases} (iD_t - D_x^2)g(x, t) \cdot f(x, t) = 0, \\ (D_x^2 - 2\epsilon\rho^2)f(x, t) \cdot f(x, t) = \pm 2\rho^2 g(x, t)g^*(-x, t), \end{cases} \quad (90)$$

We recall the bilinear equation (59) satisfied by the tau functions of single component KP hierarchy τ_k in previous section. We take the simplest case $p_1 = \bar{p}_1 = \rho$, then the second

bilinear equation in (59) reduces to the second equation in (67). Furthermore, if we let $c_1 = \frac{1}{2\rho}$, it then follows

$$\tau_0 = \frac{1}{2\rho} \left(1 + e^{\xi_1 + \bar{\xi}_1} \right), \quad (91)$$

$$\tau_1 = \frac{1}{2\rho} \left(1 - e^{\xi_1 + \bar{\xi}_1} \right), \quad (92)$$

$$\tau_{-1} = \frac{1}{2\rho} \left(1 - e^{\xi_1 + \bar{\xi}_1} \right), \quad (93)$$

where $e^{\xi_1 + \bar{\xi}_1} = e^{2\rho x}$ then if we define $f(x, t) = e^{\rho x} \tau_0$, $g(x, t) = e^{\rho x} \tau_1$, $\bar{g}(x, t) = e^{\rho x} \tau_{-1}$, we have

$$f(x, t) = f^*(-x, t), \quad \bar{g}(x, t) = -g^*(-x, t). \quad (94)$$

Furthermore, if we let $x_1 = x$, $x_2 = -it$ and $\epsilon = 1$, the bilinear equation (67) reduces to the bilinear equation (90). In this case, if we define

$$q(x, t) = \frac{g(x, t)}{f(x, t)} e^{i2\rho^2 t}, \quad r(x, t) = \frac{\bar{g}(x, t)}{f(x, t)} e^{-i2\rho^2 t}, \quad (95)$$

it is obvious that

$$r(x, t) = -q^*(-x, t), \quad (96)$$

which leads to the 1-soliton solution

$$q(x, t) = \rho \frac{1 - e^{\xi_1 + \bar{\xi}_1}}{1 + e^{\xi_1 + \bar{\xi}_1}} e^{2i\rho^2 t} = \rho e^{2i\rho^2 t} \tanh(\rho x). \quad (97)$$

This solution coincides with the solution (3.200) and (3.201) found in [23] for the case: $\sigma = -1$, $\Delta\theta = \pi$.

3.3. Soliton solution with nonzero boundary condition to the nonlocal NLS equation with $\sigma = -1$, $\Delta\theta = 0$

Again for the tau functions of single component KP hierarchy τ_k and the bilinear equations the tau functions satisfy in previous section. If we impose the condition $p_j \bar{p}_j = -\rho^2$, $j = 1, \dots, N$, then similarly, we have

$$\partial_{x_1} \tau_k = -\rho^2 \partial_{x_{-1}} \tau_k. \quad (98)$$

Furthermore, if we assume $x_1 = x$, $x_2 = -it$, then the bilinear equation (59) reduces to

$$\begin{cases} (iD_t - D_x^2) \tau_1(x, t) \cdot \tau_0(x, t) = 0, \\ (D_x^2 + 2\rho^2) \tau_0(x, t) \cdot \tau_0(x, t) = 2\rho^2 \tau_1(x, t) \tau_{-1}(x, t). \end{cases} \quad (99)$$

Consider a 2×2 matrix for $\tau_0(x, t)$, $\tau_1(x, t)$ and $\tau_{-1}(x, t)$, let $p_1 = v_1$, $\bar{p}_1 = -\rho^2/v_1$ where $v_1 > \rho$ is real, $p_2 = \bar{p}_1$, $\bar{p}_2 = p_1$, $c_2 = -c_1$ and

$$c_1^2 = \frac{1}{(p_1 + \bar{p}_1)^2} - \frac{1}{4p_1 \bar{p}_1}.$$

Since

$$\begin{aligned}
(\xi_1 + \bar{\xi}_1)(x, t) &= (v_1 - \rho^2/v_1)x + (v_1^2 - \rho^4/v_1^2)it, \\
(\xi_2 + \bar{\xi}_2)(x, t) &= (v_1 - \rho^2/v_1)x - (v_1^2 - \rho^4/v_1^2)it, \\
\tau_0(x, t) &= \begin{vmatrix} c_1 + \frac{1}{p_1 + \bar{p}_1} e^{\xi_1 + \bar{\xi}_1} & \frac{1}{p_1 + \bar{p}_2} e^{\xi_1 + \bar{\xi}_2} \\ \frac{1}{p_2 + \bar{p}_1} e^{\xi_2 + \bar{\xi}_1} & c_2 + \frac{1}{p_2 + \bar{p}_2} e^{\xi_2 + \bar{\xi}_2} \end{vmatrix} \\
&= -c_1^2 + \frac{c_1}{p_2 + \bar{p}_2} e^{\xi_2 + \bar{\xi}_2} - \frac{c_1}{p_1 + \bar{p}_1} e^{\xi_1 + \bar{\xi}_1} + c_1^2 e^{\xi_1 + \bar{\xi}_1 + \xi_2 + \bar{\xi}_2} \\
&= 2c_1 e^{(v_1 - \rho^2/v_1)x} \left\{ c_1 \sinh \left[(v_1 - \rho^2/v_1)x \right] - \frac{i}{p_1 + \bar{p}_1} \sin \left[(v_1^2 - \rho^4/v_1^2)t \right] \right\}. \quad (100)
\end{aligned}$$

Therefore

$$\tau_0^*(-x, t) = -e^{-2(v_1 - \rho^2/v_1)x} \tau_0(x, t). \quad (101)$$

Moreover

$$\begin{aligned}
\tau_1(x, t) &= \begin{vmatrix} c_1 + \frac{1}{p_1 + \bar{p}_1} \left(-\frac{p_1}{\bar{p}_1} \right) e^{\xi_1 + \bar{\xi}_1} & \frac{1}{p_1 + \bar{p}_2} \left(-\frac{p_1}{\bar{p}_2} \right) e^{\xi_1 + \bar{\xi}_2} \\ \frac{1}{p_2 + \bar{p}_1} \left(-\frac{p_2}{\bar{p}_1} \right) e^{\xi_2 + \bar{\xi}_1} & c_2 + \frac{1}{p_2 + \bar{p}_2} \left(-\frac{p_2}{\bar{p}_2} \right) e^{\xi_2 + \bar{\xi}_2} \end{vmatrix} \\
&= -c_1^2 + \frac{c_1}{p_2 + \bar{p}_2} \left(-\frac{p_2}{\bar{p}_2} \right) e^{\xi_2 + \bar{\xi}_2} - \frac{c_1}{p_1 + \bar{p}_1} \left(-\frac{p_1}{\bar{p}_1} \right) e^{\xi_1 + \bar{\xi}_1} + c_1^2 e^{\xi_1 + \bar{\xi}_1 + \xi_2 + \bar{\xi}_2} \\
&= c_1 e^{(v_1 - \rho^2/v_1)x} \left\{ 2c_1 \sinh \left[\left(v_1 - \frac{\rho^2}{v_1} \right) x \right] + \frac{1}{p_1 + \bar{p}_1} \left[\frac{p_1}{\bar{p}_1} e^{(v_1^2 - \frac{\rho^4}{v_1^2})it} - \frac{\bar{p}_1}{p_1} e^{-(v_1^2 - \frac{\rho^4}{v_1^2})it} \right] \right\}.
\end{aligned}$$

$$\begin{aligned}
\tau_{-1}(x, t) &= \begin{vmatrix} c_1 + \frac{1}{p_1 + \bar{p}_1} \left(-\frac{\bar{p}_1}{p_1} \right) e^{\xi_1 + \bar{\xi}_1} & \frac{1}{p_1 + \bar{p}_2} \left(-\frac{\bar{p}_2}{p_1} \right) e^{\xi_1 + \bar{\xi}_2} \\ \frac{1}{p_2 + \bar{p}_1} \left(-\frac{\bar{p}_1}{p_2} \right) e^{\xi_2 + \bar{\xi}_1} & c_2 + \frac{1}{p_2 + \bar{p}_2} \left(-\frac{\bar{p}_2}{p_2} \right) e^{\xi_2 + \bar{\xi}_2} \end{vmatrix} \\
&= -c_1^2 + \frac{c_1}{p_2 + \bar{p}_2} \left(-\frac{\bar{p}_2}{p_2} \right) e^{\xi_2 + \bar{\xi}_2} - \frac{c_1}{p_1 + \bar{p}_1} \left(-\frac{\bar{p}_1}{p_1} \right) e^{\xi_1 + \bar{\xi}_1} + c_1^2 e^{\xi_1 + \bar{\xi}_1 + \xi_2 + \bar{\xi}_2} \\
&= c_1 e^{(v_1 - \rho^2/v_1)x} \left\{ 2c_1 \sinh \left[\left(v_1 - \frac{\rho^2}{v_1} \right) x \right] + \frac{1}{p_1 + \bar{p}_1} \left[\frac{\bar{p}_1}{p_1} e^{(v_1^2 - \frac{\rho^4}{v_1^2})it} - \frac{p_1}{\bar{p}_1} e^{-(v_1^2 - \frac{\rho^4}{v_1^2})it} \right] \right\}.
\end{aligned}$$

Therefore

$$\tau_1^*(-x, t) = -e^{-2(v_1 - \rho^2/v_1)x} \tau_{-1}(x, t). \quad (102)$$

Consequently, if we define

$$\tau_0(x, t) = Cf(x, t), \quad \tau_1(x, t) = Cg(x, t), \quad \tau_{-1}(x, t) = C\bar{g}(x, t), \quad (103)$$

with $C = e^{(v_1 - \rho^2/v_1)x}$, then we can easily show that

$$f(x, t) = -f^*(-x, t), \quad \bar{g}(x, t) = -g^*(-x, t), \quad (104)$$

which reduces the bilinear equation (99) to

$$\begin{cases} (iD_t - D_x^2)g(x, t) \cdot f(x, t) = 0, \\ (D_x^2 + 2\rho^2)f(x, t) \cdot f(x, t) = 2\rho^2 g(x, t)g^*(-x, t). \end{cases} \quad (105)$$

Above bilinear equation coincides with the bilinear equation (90) with $\epsilon = -1$ in the case $f(x, t) = -f^*(-x, t)$. Moreover, for

$$q(x, t) = \frac{g(x, t)}{f(x, t)} e^{-i2\rho^2 t}, \quad r(x, t) = -\frac{\bar{g}(x, t)}{f(x, t)} e^{-i2\rho^2 t}, \quad (106)$$

it is obvious that

$$r(x, t) = -q^*(-x, t), \quad (107)$$

which leads to the 1-soliton solution

$$q(x, t) = \rho \frac{(v_1^2 + \rho^2) \sinh \left[(v_1 - \frac{\rho^2}{v_1})x \right] + \frac{2\rho^3}{v_1} e^{-(v_1^2 - \frac{\rho^4}{v_1^2})it} - \frac{2v_1^3}{\rho} e^{(v_1^2 - \frac{\rho^4}{v_1^2})it}}{(v_1^2 + \rho^2) \sinh [(v_1 - \rho^2/v_1)x] - 4i\rho v_1 \sin [(v_1^2 - \rho^4/v_1^2)t]} e^{-2i\rho^2 t}. \quad (108)$$

This solution coincides with the solution (4.130) in [23] for the case: $\sigma = -1$, $\Delta\theta = 0$, which is actually a stationary breather solution.

4. Summary and concluding remarks

In this paper, via the combination of Hirota's bilinear method and the KP hierarchy reduction method, we have constructed various general soliton solutions in the form of determinants for the nonlocal NLS equation with both zero and nonzero boundary conditions. The solutions obtained here coincide with the solutions found in the nonlocal NLS equation so far and reformulate them in compact form by using tau functions of the KP hierarchy.

A natural extension of the present work will be the construction of the general soliton solutions to the coupled nonlocal NLS equation

$$\{iq_{k,t}(x, t) = q_{k,xx}(x, t) - 2 \left(\sum_{j=1}^2 \sigma_j q_j(x, t) q_j^*(-x, t) \right) q_k(x, t), \quad k = 1, 2. \quad (109)$$

It is expected that the soliton solution to the above coupled nonlocal NLS equation is even richer and more complicated than the single component nonlocal NLS equation. We will explore this interesting topic and report the results elsewhere in the near future.

Acknowledgments

MJA was partially supported by NSF under Grant No. DMS-1712793. BF was partially supported by NSF Grant under Grant No. (DMS-1715991) and NSF of China for Overseas Scholar Collaboration Research under Grant No. 11728103.

ORCID iDs

Bao-Feng Feng  <https://orcid.org/0000-0002-2529-897X>

References

- [1] Benney D J and Newell A C 1967 *Stud. Appl. Math.* **46** 52–63

- [2] Ablowitz M J and Clarkson P A 1991 *Solitons, Nonlinear Evolution Equations and Inverse Scattering* (London Mathematical Society Lecture Notes Series vol 149) (Cambridge: Cambridge University Press)
- [3] Ablowitz M J, Prinari B and Trubatch A D 2004 *Discrete and Continuous Nonlinear Schrödinger Systems* (Cambridge: Cambridge University Press)
- [4] Zakharov V E and Shabat A B 1972 *J. Exp. Theor. Phys.* **34** 62–9
- [5] Ablowitz M J, Kaup D J, Newell A C and Segur H 1973 *Phys. Rev. Lett.* **31** 125–7
- [6] Ablowitz M J, Kaup D J, Newell A C and Segur H 1972 *Phys. Rev. Lett.* **30** 1262–4
- [7] Ablowitz M J, Kaup D J, Newell A C and Segur H 1974 *Stud. Appl. Math.* **53** 249–315
- [8] Hasegawa A and Tappert F 1973 *Appl. Phys. Lett.* **23** 142
- [9] Hasegawa A and Tappert F 1973 *Appl. Phys. Lett.* **23** 171
- [10] Agrawal G P 1995 *Nonlinear Fiber Optics* (New York: Academic)
- [11] Hasegawa A and Kodama Y 1995 *Solitons in Optical Communications* (Oxford: Clarendon)
- [12] Kivshar Y S and Agrawal G P 2003 *optical Solitons: from Fibers to Photonic Crystals* (New York: Academic)
- [13] Dalfó F, Giorgini S and Stringari L P 1999 *Rev. Mod. Phys.* **71** 463
- [14] Zakharov V E 1968 *J. Appl. Mech. Tech. Phys.* **4** 190
- [15] Benney D J and Roskes G K 1969 *Stud. Appl. Math.* **48** 377
- [16] Zakharov V E 1972 *Sov. Phys.—JETP* **35** 908
- [17] Ablowitz M J and Musslimani Z H 2013 *Phys. Rev. Lett.* **110** 064105
- [18] Gadzhimuradov T A and Agalarov A M 2016 *Phys. Rev. A* **93** 062124
- [19] Bender C M and Boettcher S 1998 *Phys. Rev. Lett.* **80** 5243
- [20] Konotop V V, Yang J and Zezyulin D A 2016 *Rev. Mod. Phys.* **88** 035002
- [21] Ablowitz M J and Musslimani Z H 2016 *Nonlinearity* **29** 915
- [22] Ablowitz M J and Musslimani Z H 2017 *Stud. Appl. Math.* **139** 7
- [23] Ablowitz M J, Luo X D and Musslimani Z H 2018 *J. Math. Phys.* **59** 011501
- [24] Ablowitz M J, Feng B F, Luo X D and Musslimani Z H 2018 *Stud. Appl. Math.* **141** 267–307
- [25] Ablowitz M J, Feng B F, Luo X D and Musslimani Z H 2018 *Theor. Math. Phys.* **196** 1241
- [26] Ablowitz M J and Musslimani Z H 2014 *Phys. Rev. E* **90** 032912
- [27] Fokas A S 2016 *Nonlinearity* **29** 319
- [28] Musslimani Z H, Makris K G, El Ganainy R and Christodoulides D N 2008 *Phys. Rev. Lett.* **100** 030402
- [29] Ma L Y and Zhu Z N 2016 *Appl. Math. Lett.* **59** 115
- [30] Ma L Y and Zhu Z N 2016 *J. Math. Phys.* **57** 083507
- [31] Li M and Xu T 2015 *Phys. Rev. E* **91** 033202
- [32] Huang X and Ling L 2016 *Euro. Phys. J. Plus* **131** 16148
- [33] Wen X Y, Yan Z and Yang Y 2016 *Chaos* **26** 063123
- [34] Zhou Z X 2018 *Nonlinear Sci. Numer. Simul.* **62** 480
- [35] Yan Z, Wen Z and Hang C 2015 *Phys. Rev. E* **92** 022913
- [36] Yan Z 2015 *Appl. Math. Lett.* **47** 61
- [37] Gerdjikov V S, Grahovski G G and Ivanov R I 2016 *Theor. Math. Phys.* **188** 1305
- [38] Xu Z X and Chow K W 2016 *Appl. Math. Lett.* **56** 72
- [39] Rao J G, Cheng Y and He J S 2017 *Stud. Appl. Math.* **139** 568
- [40] Yang B and Yang J 2018 *Stud. Appl. Math.* **140** 178
- [41] Gürses M 2017 *Phys. Lett. A* **381** 1791
- [42] Yang B and Chen Y 2017 (arXiv:1710.07061 [math-ph])
- [43] Chen K, Deng X, Lou S Y and Zhang D J 2018 *Stud. Appl. Math.* **141** 113
- [44] Khare A and Saxena A 2015 *J. Math. Phys.* **56** 032104
- [45] Sinha D and Ghosh P K 2015 *Phys. Rev. E* **91** 042908
- [46] Stalina S, Senthilvelana M and Lakshmanan M 2017 *Phys. Lett. A* **381** 2380
- [47] Gerdjikov V S and Saxena A 2017 *J. Math. Phys.* **58** 013502
- [48] Chen K and Zhang D J 2018 *Appl. Math. Lett.* **75** 82
- [49] Lou S Y and Huang F 2017 *Sci. Rep.* **7** 869
- [50] Ji J L and Zhu Z N 2017 *Commun. Nonlinear Sci. Numer. Simul.* **42** 699
- [51] Ji J L and Zhu Z N 2017 *J. Math. Anal. Appl.* **453** 973
- [52] Ma L Y, Shen S F and Zhu Z N 2017 *J. Math. Phys.* **58** 103501
- [53] Song C, Xiao D and Zhu Z N 2017 *J. Phys. Soc. Japan* **86** 054001

- [54] Song C, Xiao D and Zhu Z N 2017 *Commun. Nonlinear Sci. Numer. Simul.* **45** 13
- [55] Gürses M and Pekcan A 2018 *J. Math. Phys.* **59** 051501
- [56] Yang B and Yang J 2017 General rogue waves in the nonlocal PT-symmetric nonlinear Schrödinger equation and (arXiv:[1711.05930](#) [nlin.SI])
- [57] Yang J 2017 General N-solitons and their dynamics in several nonlocal nonlinear Schrödinger equations (arXiv:[1712.01181](#) [nlin.SI])
- [58] Ablowitz M J and Ladik J F 1975 *J. Math. Phys.* **16** 598
- [59] Ablowitz M J and Ladik J F 1976 *J. Math. Phys.* **17** 1011
- [60] Prinari B, Ablowitz M J and Biondini G 2006 *J. Math. Phys.* **47** 063508
- [61] Demontis F, Prinari B, van der Mee C and Vitale F 2012 *Stud. Appl. Math.* **131** 1
- [62] Demontis F, Prinari B, van der Mee C and Vitale F 2014 *J. Math. Phys.* **55** 101505
- [63] Biondini G and Kovacic G 2014 *J. Math. Phys.* **55** 031506
- [64] Biondini G and Kraus D 2015 *SIAM J. Math. Anal.* **47** 706
- [65] Biondini G, Kovacic G and Kraus D K 2015 *Nonlinearity* **28** 3101
- [66] Pichler M and Biondini G 2017 *IMA J. Appl. Math.* **82** 131
- [67] Vekslerchik V E and Konotop V V 1992 *Inverse Problems* **8** 889
- [68] Ablowitz M J, Biondini G and Prinari B 2007 *Inverse Problems* **23** 1711
- [69] Van Der Mee C 2015 *J. Nonlinear Math. Phys.* **22** 233
- [70] Prinari B and Vitale F 2015 *Stud. Appl. Math.* **137** 28
- [71] Hirota R 2004 *The Direct Method in Soliton Theory* (Cambridge: Cambridge University Press)
- [72] Sato M 1981 *RIMS Kyoto Univ. Kokyuroku* **439** 30
- [73] Ueno K and Takasaki K 1984 *Adv. Study Pure Math.* **4** 1
- [74] Jimbo M and Miwa T 1983 *Publ. RIMS Kyoto Univ.* **19** 943
- [75] Date E, Kashiwara M, Jimbo M and Miwa T 1983 Transformation groups for soliton equations *Nonlinear Integrable Systems Classical Theory and Quantum Theory* vol 39 (Singapore: World Scientific)
- [76] Miwa T, Jimbo M and Date E 2000 *Solitons, Differential Equations, Symmetries and Infinite Dimensional Algebras* (Cambridge: Cambridge University Press)
- [77] Ohta Y 1989 *RIMS Kokyuroku* **684** 1
- [78] Gilson C, Hietarinta J, Nimmo J and Ohta Y 2003 *Phys. Rev. E* **68** 016614
- [79] Ohta Y, Wang D S and Yang J 2011 *Stud. Appl. Math.* **127** 345
- [80] Miyake S, Ohta Y and Satsuma J 1990 *J. Phys. Soc. Japan* **59** 48
- [81] Pelinovsky D 1995 *Physica D* **87** 115
- [82] Loris I and Willox R 1997 *J. Math. Phys.* **38** 283–91
- [83] Loris I and Willox R 1997 *J. Phys. A: Math. Gen.* **30** 6925–38
- [84] Pelinovsky D 1998 *J. Math. Phys.* **39** 5377–95