
Contents lists available at ScienceDirect

Chemical Geology

journal homepage: www.elsevier.com/locate/chemgeo



ConwayGEOTRACES

The GEOTRACES Intermediate Data Product 2017[★]

Reiner Schlitzer^{a,*}, Robert F. Anderson^b, Elena Masferrer Dodas^c, Maeve Lohan^{aw}, Walter Geibert^a, Alessandro Tagliabue^e, Andrew Bowie^f, Catherine Jeandel^c, Maria T. Maldonado^h, William M. Landing^{bj}, Donna Cockwell^{ak}, Cyril Abadie^c, Wafa Abouchami^{ck}, Eric P. Achterberg^k, Alison Agather^{de}, Ana Aguliar-Islas^{cd}, Hendrik M. van Aken^g, Morten Andersen^{ds}, Corey Archer^{bp}, Maureen Auro^p, Hein J. de Baar^g, Oliver Baars^{k,ar}, Alex R. Baker^{cr}, Karel Bakker^g, Chandranath Basak^{bh}, Mark Baskaranⁱ, Nicholas R. Bates^j, Dorothea Bauch^k, Pieter van Beek^c, Melanie K. Behrens^{bh}, Erin Black^p, Katrin Bluhm^{cp}, Laurent Bopp^{br}, Heather Bouman^{ab}, Katlin Bowmanⁿ, Johann Bown^{l,bz} Philip Boyd^f, Marie Boye^{bc,l}, Edward A. Boyle^m, Pierre Branellec^{df}, Luke Bridgestock^{ab,bf}, Guillaume Brissebrat^{dd}, Thomas Browning^{ab,k}, Kenneth W. Bruland^{n,cb}, Hans-Jürgen Brumsack^{ap}, Mark Brzezinski^o, Clifton S. Buck^{ce}, Kristen N. Buck^{bo,j}, Ken Buesseler^p, Abby Bull^{ak}, Edward Butler^{q,be}, Pinghe Cai^r, Patricia Cámara Mor^{au}, Damien Cardinal^{bc}, Craig Carlson^o, Gonzalo Carrasco^{m,cz}, Núria Casacuberta^{bk}, Karen L. Casciotti^{cy}, Maxi Castrillejo^{au,bk,cl} Elena Chamizo^{dl}, Rosie Chance^{cr}, Matthew A. Charette^p, Joaquin E. Chaves^s, Hai Cheng^{t,ad}, Fanny Chever^l, Marcus Christl^{bk}, Thomas M. Church^v, Ivia Closset^{bc,o}, Albert Colman^w, Tim M. Conway^{cj}, Daniel Cossa^{bl}, Peter Croot^y, Jay T. Cullen^{bs}, Gregory A. Cutter^{du}, Chris Daniels^{ak}, Frank Dehairs^{aa}, Feifei Deng^{ab}, Huong Thi Dieu^{ac}, Brian Duggan^x, Gabriel Dulaquais¹, Cynthia Dumousseaud^{aw}, Yolanda Echegoyen-Sanz^m, R. Lawrence Edwards^{ad}, Michael Ellwood^{dy}, Eberhard Fahrbach^a, Jessica N. Fitzsimmons^{bw,bx}, A. Russell Flegal^{cb}, Martin Q. Fleisher^b, Tina van de Flierdt^{bf}, Martin Frank^k, Jana Friedrich^{a,ae}, Francois Fripiat^{aa}, Henning Fröllje^{bh}, Stephen J.G. Galer^{cj}, Toshitaka Gamo^{af}, Raja S. Ganeshram^{ax}, Jordi Garcia-Orellana^{au,cl}, Ester Garcia-Solsona^{cu}, Melanie Gault-Ringold^{al,f}, Ejin George^{al}, Loes J.A. Gerringa^g, Melissa Gilbert^{at}, Jose M. Godoy^{bd}, Steven L. Goldstein^b, Santiago R. Gonzalez^g, Karen Grissom^{at}, Chad Hammerschmidt^{de}, Alison Hartman^b, Christel S. Hassler^{db}, Ed C. Hathorne^k, Mariko Hatta^{ag}, Nicholas Hawco^p, Christopher T. Hayes^{at}, Lars-Eric Heimbürger^{cf}, Josh Helgoe^x, Maija Hellerⁿ, Gideon M. Henderson^{ab}, Paul B. Henderson^p, Steven van Heuven^{g,ah}, Peng Ho^{at}, Tristan J. Horner^p, Yu-Te Hsieh^{ab}, Kuo-Fang Huang^{ai,cq}, Matthew P. Humphreys^{aw,cr}, Kenji Isshiki^{dr}, Jeremy E. Jacquot^{bz}, David J. Janssen^{bs}, William J. Jenkins^p, Seth John^{bv}, Elizabeth M. Jones^{g,ah,dx}, Janice L. Jones^o, David C. Kadko^{bn}, Rick Kayser^m, Timothy C. Kenna^b, Roulin Khondoker^{bf}, Taejin Kim^{af,bq}, Lauren Kipp^p, Jessica K. Klar^{aw,c}, Maarten Klunder^g, Sven Kretschmer^a, Yuichiro Kumamoto^{aj}, Patrick Laan^{bz}, Marie Labatut^c, Francois Lacan^c, Phoebe J. Lamⁿ, Myriam Lambelet^{bf}, Carl H. Lamborgⁿ, Frédéric A.C. Le Moigne^k, Emilie Le Roy^c, Oliver J. Lechtenfeld^{cv}, Jong-Mi Leeⁿ, Pascale Lherminier^{df}, Susan Little^{bf}, Mercedes López-Lora^{dl}, Yanbin Lu^{ad}, Pere Masque^{au,ca,cl}, Edward Mawji^{ak,dc}, Charles R. Mcclain^s, Christopher Measures^{ag},

https://doi.org/10.1016/j.chemgeo.2018.05.040

Available online 01 June 2018 0009-2541/ © 2018 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (http://creativecommons.org/licenses/BY/4.0/).



^{*} This article is part of a special issue entitled: "Cycles of trace elements and isotopes in the ocean – GEOTRACES and beyond" - edited by Tim M. Conway, Tristan Horner, Yves Plancherel, and Aridane G. González.

^{*} Corresponding author at: Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, Am Handelshafen 12, Bremerhaven 27570, Germany. *E-mail address*: Reiner.Schlitzer@awi.de (R. Schlitzer).

Sanjin Mehicⁿ, Jan-Lukas Menzel Barraqueta^k, Pier van der Merwe^f, Rob Middag^g, Sebastian Mieruch^a, Angela Milne^d, Tomoharu Minami^{ba}, James W. Moffett^{bv}, Gwenaelle Moncoiffe^{bi}, Willard S. Moore^x, Paul J. Morris^p, Peter L. Morton^{c3} Yuzuru Nakaguchi^{di}, Noriko Nakayama^{af}, John Niedermillerⁱ, Jun Nishioka^{am}, Akira Nishiuchi^{di}, Abigail Noble^{an}, Hajime Obata^{af}, Sven Ober^g, Daniel C. Ohnemus^{ay}, Jan van Ooijen^g, Jeanette O'Sullivan^{be}, Stephanie Owens^p, Katharina Pahnke^{bh}, Maxence Paul^{bf}, Frank Pavia^b, Leopoldo D. Pena^{cu,b}, Brian Peters^{cy}, Frederic Planchon¹, Helene Planquette¹, Catherine Pradoux^c, Viena Puigcorbé^{ca}, Paul Quay^{ao}, Fabien Queroue¹, Amandine Radic^c, S. Rauschenberg^{ay}, Mark Rehkämper^{bf}, Robert Rember^{dh}, Tomas Remenyi^f, Joseph A. Resing^{co}, Joerg Rickli^{bp}, Sylvain Rigaud^{v,cm}, Micha J.A. Rijkenberg^g, Stephen Rintoul^{f,dp,dq}, Laura F. Robinson^{p,aq}, Montserrat Roca-Martí^{au}, Valenti Rodellas^{bt}, Tobias Roeske^a, John M. Rolison^{al}, Mark Rosenberg^f, Saeed Roshan^{as,cc}, Michiel M. Rutgers van der Loeff^a, Evgenia Ryabenko^k, Mak A. Saito^p, Lesley A. Salt^g, Virginie Sanial^p, Geraldine Sarthou¹, Christina Schallenberg^f, Ursula Schauer^a, Howie Scher^x, Christian Schlosser^{aw,k}, Bernhard Schnetger^{ap}, Peter Scott^{ab,cw}, Peter N. Sedwick^z, Igor Semiletov^{cg,ch}, Rachel Shelley^{l,bj}, Robert M. Sherrell^{bx,ct}, Alan M. Shiller^{at}, Daniel M. Sigman^u, Sunil Kumar Singh^{dm,dn}, Hans A. Slagter^g, Emma Slater^{bi}, William M. Smethie^b, Helen Snaith^{ak}, Yoshiki Sohrin^{ba}, Bettina Sohst^z, Jeroen E. Sonke^{dg}, Sabrina Speich^{av,br}, Reiner Steinfeldt^{bm}, Gillian Stewart^{dt}, Torben Stichel^{aw}, Claudine H. Stirling^{al}, Johnny Stutsman^{ao}, Gretchen J. Swarr^p, James H. Swift^{by}, Alexander Thomas^{ax}, Kay Thorne^{bi}, Claire P. Till^{do,n}, Ralph Till^{cb}, Ashley T. Townsend^{da}, Emily Townsend^x, Robyn Tuerena^{ax}, Benjamin S. Twining^{ay}, Derek Vance^{bp}, Sue Velazquez^{bs}, Celia Venchiarutti^a, Maria Villa-Alfageme^{dk}, Sebastian M. Vivancos^b, Antje H.L. Voelker^{az}, Bronwyn Wake^l, Mark J. Warner^{ao}, Ros Watson^{be}, Evaline van Weerlee^g, M. Alexandra Weigand^u, Yishai Weinstein^{dv}, Dominik Weiss^{bf}, Andreas Wisotzki^a, E. Malcolm S. Woodward^{bg}, Jingfeng Wu^{as,bb}, Yingzhe Wu^b, Kathrin Wuttig^f, Neil Wyatt^{aw}, Yang Xiangⁿ, Ruifang C. Xie^{k,cj}, Zichen Xue^{bf}, Hisayuki Yoshikawa^{ci,ch}, Jing Zhang^{cs,cr}, Pu Zhang^{ad}, Ye Zhao^{dw}, Linjie Zheng^{ba}, Xin-Yuan Zheng^{ab,bu}, Moritz Zieringer^k, Louise A. Zimmer^{cn}, Patrizia Ziveri^{au,dj}, Patricia Zunino^{df}, Chervl Zurbrick^m

^b Lamont-Doherty Earth Observatory of Columbia University, PO Box 1000, 61 Route 9W, Palisades 10964-1000, USA

^c LEGOS, University of Toulouse, CNRS, IRD, CNES, UPS, Toulouse, France

- ^d School of Geography, Earth and Environmental Sciences, University of Plymouth, Drake Circus, Plymouth PL4 8AA, United Kingdom
- ^e Dept. of Earth, Ocean and Ecological Sciences, School of Environmental Sciences, University of Liverpool, Liverpool, United Kingdom
- ^f Antarctic Climate and Ecosystems CRC and Institute for Marine and Antarctic Studies, University of Tasmania, Private Bag 80, Hobart 7001, Australia
- ⁸ NIOZ Royal Netherlands Institute For Sea Research and Utrecht University, PO Box 59, Den Burg 1790 AB, the Netherlands
- h University of British Columbia, Department of Earth, Ocean and Atmospheric Sciences, Earth Science Bldg., 2207 Main Mall, Vancouver, BC V6T 124, Canada
- ⁱ Department of Geology, Wayne State University, 0224 Old Main, 4841 Cass Avenue, Detroit 48202, USA
- ^j Bermuda Institute of Ocean Sciences, 17 Biological Lane, Ferry Reach, St. Georges, GE01, Bermuda
- ^k GEOMAR, Helmholtz Centre for Ocean Research Kiel, Wischhofstraße 1-3, Kiel 24148, Germany
- ¹ Laboratory of Marine Environmental Science (LEMAR, UMR CNRS UBO IRD Ifremer 6539), Institut Universitaire Européen de la Mer (IUEM), Place Nicolas Copernic, Technopôle Brest Iroise, Plouzane 29280, France
- ^m Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology (MIT), Building E25-619, 77 Massachusetts Avenue, Cambridge 02139, USA
- ⁿ University of California, Santa Cruz, Department of Ocean Sciences, 1156 High St, Santa Cruz, CA 95064, USA
- ^o Marine Science Institute, University of California, Santa Barbara, UC Santa Barbara, Santa Barbara 93106-9620, USA
- ^p Woods Hole Oceanographic Institution, Department of Marine Chemistry and Geochemistry, 266 Woods Hole Road, Woods Hole 02543, USA
- ⁹ Australian Institute of Marine Science, Darwin, PO Box 41775, Casuarina, NT 0811, Australia
- r State Key Laboratory of Marine Environmental Science, Xiamen University, 422 Siming South Road, Xiamen 361005, China
- ^s NASA Goddard Space Flight Center, Ocean Ecology Laboratory, Code 616, Greenbelt 20771, USA
- ^t Institute of Global Environmental Change, Xi'an Jiao Tong University, 99 Yanxiang Road, Western No. 1 Building, Xi'an 710049, China
- ^u Department of Geosciences, Princeton University, Princeton, NJ 08544, USA
- v College of Earth, Ocean, and Environment, University of Delaware, 111 Robinson Hall, Newark 19716-3501, USA
- ^w Department of the Geophysical Sciences, University of Chicago, 5734 S. Ellis Avenue, Chicago 60637, USA
- * Department of Earth and Ocean Sciences, University of South Carolina, 701 Sumter Street, EWS 617, Columbia 29208, USA
- ^y Department of Earth and Ocean Sciences, National University of Ireland Galway, University Road, Galway, Ireland
- ² Department of Ocean, Earth and Atmospheric Sciences, Old Dominion University, 4600 Elkhorn Avenue, Norfolk 23529, USA
- ^{aa} Analytical, Environmental and Geo-Chemistry Department, Vrije Universiteit Brussel, Pleinlaan 2, B-1050 Brussels, Belgium
- ab Department of Earth Sciences, University of Oxford, South Parks Road, Oxford OX1 3AN, United Kingdom
- ac Kyoto University, Institute for Chemical Research, Gokasho, Uji 611-0011, Japan

- ae Helmholtz Zentrum Geesthacht Center for Materials and Coastal Research, Max-Planck Str. 1, 21502 Geesthacht, Germany
- ^{af} Atmosphere and Ocean Research Institute, The University of Tokyo, 5-1-5, Kashiwanoha, Kashiwa 277-8564, Japan
- ^{ag} Department of Oceanography, University of Hawai'i at Manoa, 1000 Pope Road, Honolulu 96822-3324, USA
- ah Energy and Sustainability Research Institute Groningen, University of Groningen, Nijenborgh 4, 9747, AG, Groningen, the Netherlands
- ^{ai} Institute of Earth Sciences, Academia Sinica, 128, Sec. 2, Academia Road, Nangang, Taipei 11529, Taiwan

^a Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, Am Handelshafen 12, Bremerhaven 27570, Germany

^{ad} Department of Earth Sciences, University of Minnesota, 116 Church St. SE, Minneapolis 55455-0231, USA

^{aj} Research and Development Center for Global Change, Japan Agency for Marine-Earth Science and Technology, 2-15 Natsushima-Cho, Yokosuka 237-0061, Japan

- ak British Oceanographic Data Centre, National Oceanography Centre, Southampton, European Way, Southampton SO14 3ZH, United Kingdom
- ^{a1} Department of Chemistry, NIWA/University of Otago Research Centre for Oceanography, PO BOX 56, Dunedin 9054, New Zealand

^{am} Institute of Low Temperature Sciences, Hokkaido University, Kita-19, Nishi-8, Kita-ku, Sapporo 060-0819, Japan

- ^{an} Environmental Chemistry Group, Gradient, 20 University Road, Cambridge, MA 02138, USA
- ^{ao} School of Oceanography. University of Washington, PO Box 357940, Seattle 98195-7940, USA
- ^{ap} Institut für Chemie und Biologie des Meeres (ICBM), Universität Oldenburg, Postfach 2503, D-26111 Oldenburg, Germany
- aq Department of Earth Sciences, University of Bristol, Wills Memorial Building, Queen's Road, Bristol BS8 1RJ, United Kingdom
- ar NC State University, Department of Entomology & Plant Pathology, Raleigh, NC 27601, USA

as University of Miami, Rosenstiel School of Marine and Atmospheric Science (RSMAS) Marine and Atmospheric Chemistry (MAC), 4600 Rickenbacker Causeway, Miami 33149-1098 JISA

- ^{at} Division of Marine Science, University of Southern Mississippi, 1020 Balch Boulevard, Stennis Space Center, MS 39529, USA
- ^{au} Institute of Environmental Science and Technology, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain
- av Ocean Physics Laboratory, University of Western Brittany, 6 avenue Victor-Le-Gorgeu, BP 809, Brest 29285, France
- aw Ocean and Earth Science, National Oceanography Centre Southampton, University of Southampton, Southampton SO14 3ZH, United Kingdom
- ax University of Edinburgh, School of GeoSciences, Grant Institute, James Hutton Road, Edinburgh EH9 3FE, United Kingdom
- ^{ay} Bigelow Laboratory for Ocean Sciences, 60 Bigelow Drive, P.O. Box 380, East Boothbay, ME 04544, USA
- az Portuguese Institute of the Sea and the Atmosphere, Rua Alfredo Magalhães Ramalho 6, Lisbon 1495-006, Portugal
- ba Institute for Chemical Research, Kyoto University, Gokasho, Uji, Kyoto 611-0011, Japan
- ^{bb} School of Biology and Marine Sciences, Shenzhen University, Shenzhen, China
- ^{bc}LOCEAN, Sorbonne Université, 4 Place Jussieu, 75252 Paris, France
- ^{bd} Department of Chemistry, Pontifical Catholic University of Rio de Janeiro, Rua Marqués de Sao Vicente, 225, Sala 772-A Prédio Cardial Leme, Bloco Leopoldo Hainberger SJ, Gávea, Rio de Janeiro 22453-900, Brazil

be CSIRO Marine and Atmospheric Research, Hobart, Castray Esplanade, Hobart 7000, Australia

^{bf} Department of Earth Science and Engineering, Imperial College London, London SW7 2AZ, United Kingdom

^{bg} Plymouth Marine Laboratory, Prospect Place, The Hoe, Plymouth PL1 3DH, United Kingdom

^{bh} Max Planck Research Group for Marine Isotope Geochemistry, Institute for Chemistry and Biology of the Marine Environment (ICBM), University of Oldenburg, Carlvon-Ossietzky-Str. 9-11, 26129 Oldenburg, Germany

bi British Oceanographic Data Centre, National Oceanography Centre, Joseph Proudman Building, 6 Brownlow Street, Liverpool L3 5DA, United Kingdom

^{bj} Earth, Ocean and Atmospheric Science, Florida State University, Tallahassee, FL 32306, USA

^{bk} Laboratory of Ion Beam Physics, ETH Zurich, Otto-Stern-Weg 5, 8093 Zurich, Switzerland

^{b1} ISTerre, Université Grenoble Alpes, CS 40700, 38058 Grenoble Cedex 9, France

^{bm} Institute for Environmental Physics, University of Bremen, Otto-Hahn-Allee, Bremen 28359, Germany

^{bn} Applied Research Center, Florida International University, Miami, FL 33174, USA

^{bo} College of Marine Science, University of South Florida, St Petersburg, FL 33701, USA

^{bp} Institute for Geochemistry and Petrology, Department of Earth Sciences, ETH Zürich, Clausiusstrasse 25, 8092 Zürich, Switzerland

^{bq} School of Earth and Environmental Sciences, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Republic of Korea

^{br} Department of Geosciences, LMD-IPSL, Ecole normale supérieure & Paris Sciences Lettres, Paris, France

^{bs} School of Earth and Ocean Sciences, University of Victoria, Victoria, British Columbia, Canada

^{bt} Aix-Marseille Université, CNRS, IRD, INRA, Coll France, CEREGE, 13545 Aix-en-Provence, France

^{bu} Department of Geoscience, University of Wisconsin-Madison, WI 53706, USA

^{bv} Department of Earth Sciences, University of Southern California, 3651 Trousdale Parkway, Los Angeles, CA 90089, USA

bw Department of Oceanography, Texas A&M University, TX 77843, USA

bx Department of Marine and Coastal Sciences, Rutgers University, New Brunswick, NJ 08901, USA

by Scripps Institution of Oceanography, University of California, San Diego, 9500 Gilman Dr., MC-0236, La Jolla, CA 92093-0236, USA

bz Department of Biological Sciences, University of Southern California, 3616 Trousdale Parkway, Los Angeles, CA 90089, USA

- ca School of Science, Centre for Marine Ecosystems Research, Edith Cowan University, 270 Joondalup Drive, Joondalup, WA 6025, Australia
- cb Institute of Marine Sciences, University of California, Santa Cruz, 1156 High St., Santa Cruz, CA 95064, USA

^{cc} Department of Geography, University of California, Santa Barbara, CA 93106, USA

^{cd} College of Fisheries and Ocean Sciences, University of Alaska Fairbanks, Fairbanks, AK 99775, USA

- ce Skidaway Institute of Oceanography, University of Georgia, Savannah, GA 31411, USA
- cf Aix Marseille Université, CNRS/INSU, Université de Toulon, IRD, Mediterranean Institute of Oceanography (MIO) UM 110, 13288 Marseille, France
- ^{cg} Pacific Oceanological Institute, Far Eastern Branch of the Russian Academy of Sciences, 43 Baltic street, Vladivostok 690041, Russia

^{ch} National Tomsk Polytechnic University, 30 Prospect Lenina, Tomsk, Russia

- ci Faculty of Environmental Earth Science, Hokkaido University, Kita-10, Nishi-5, Kita-ku, Sapporo 060-0810, Japan
- cj College of Marine Science & School of Geosciences, University of South Florida, USA
- ck Max Planck Institute for Chemistry, Climate Geochemistry Department, Hahn-Meitner-Weg 1, 55128 Mainz, Germany

^{c1} Department of Physics, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain

^{cm} Univ. Nîmes, EA 7352 CHROME, rue du Dr Georges Salan, 30021 Nimes, France

^{cn} Danish Technological Institute, Kongsvang Alle 29, 8000 Aarhus C, Denmark

^{co} Joint Institute for the Study of the Atmosphere and the Ocean, University of Washington and NOAA Pacific Marine Environmental Laboratory, 7600 Sand Point Way NE, Seattle, WA 98115, USA

- ^{cp} Akvaplan-niva AS, Framsenteret, Postboks 6606, 9296 Tromsø, Norway
- ^{cq} Department of Geology and Geophysics, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA
- cr Centre for Ocean and Atmospheric Sciences, School of Environmental Sciences, University of East Anglia, Norwich NR4 7TJ, United Kingdom
- ^{cs} Graduate School of Science and Engineering, University of Toyama, 3190 Gofuku, Toyama 9308555, Japan
- ^{ct} Department of Earth and Planetary Sciences, Rutgers University, Piscataway, NJ 08854, USA
- ^{cu} Department of Earth and Ocean Dynamics, Universitat de Barcelona, 08028 Barcelona, Spain
- ev Department of Analytical Chemistry, Helmholtz-Centre for Environmental Research UFZ, Permoserstr. 15, Leipzig 04318, Germany
- ^{cw} CEOAS, Oregon State University, Corvallis, OR 97331-5503, USA
- $^{
 m cx}$ Geochemistry, National High Magnetic Field Laboratory, Tallahassee, FL 32310, USA
- ^{cy} Department of Earth System Science, Stanford University, Stanford, CA 94305, USA
- ^{cz} Singapore-MIT Alliance for Research and Technology, Singapore 138602, Singapore
- da Central Science Laboratory, University of Tasmania, Hobart, Tasmania, Australia
- db Department F.-A. Forel for Environmental and Aquatic Sciences, University of Geneva, 66 Bvd Carl-Vogt, 1211 Geneva 4, Switzerland
- d^c Ocean Biogeochemistry and Ecosystems, National Oceanography Centre, Southampton, European Way, Southampton SO14 3ZH, United Kingdom
- dd Observatoire Midi-Pyrenées, Université de Toulouse, CNRS, CNES, IRD, Météo France, UPS, France
- de Department of Earth & Environmental Sciences, Wright State University, Dayton, OH 45435, USA
- df Ifremer, Univ. Brest, CNRS, IRD, Laboratoire d'Océanographie Physique et Spatiale (LOPS), IUEM, F-29280 Plouzané, France
- ^{dg} Geosciences Environnement Toulouse, CNRS/IRD/Universite de Toulouse 3, France

^{dh} International Arctic Research Center, University of Alaska Fairbanks, USA

di School of Science and Engineering, Kindai University, 3-4-1 Kowakae, Higashiosaka, Osaka 5778502, Japan

dj ICREA, Pg. Lluís Companys 23, 08010 Barcelona, Spain

dk Universidad de Sevilla, Department of Applied Physics, Av. Reina Mercedes 4A, 41004 Sevilla, Spain

^{d1} Universidad de Sevilla-CSIC-JA, Centro Nacional de Aceleradores, 41092 Sevilla, Spain

^{dm} Physical Research Laboratory, Navrangpura, Ahmedabad 380001, India

^{dn} CSIR-National Institute of Oceanography, Dona Paula, Goa, India

do Chemistry Department, Humboldt State University, Arcata, CA 95521, USA

^{dp} CSIRO Oceans & Atmosphere, Hobart, Tasmania 7000, Australia

dq Centre for Southern Hemisphere Ocean Research, Hobart, Tasmania 7000, Australia

dr The Community Center for the Advancement of Education and Research, University of Kochi, 2-22, Eikokuji-cho, Kochi 780-8515, Japan

ds Cardiff University, School of Earth & Ocean Sciences, Cardiff CF10 3AT, United Kingdom

dt School of Earth and Environmental Sciences, Queens College, CUNY, Flushing, New York 11217, USA

du Department of Ocean, Earth, and Atmospheric Sciences, Old Dominion University, Norfolk, VA 23529, USA

^{dv} Bar-Ilan University, Ramat-Gan 5290002, Israel

dw Nu Instruments Ltd, Unit 74, Clywedog Road South, Wrexham Industrial Estate, LL13 9XS, United Kingdom

dx Institute of Marine Research, Sykehusveien 23, 9019 Tromsø, Norway

dy Research School of Earth Sciences, Australian National University, Canberra, ACT 2601, Australia

ARTICLE INFO

Keywords: GEOTRACES

Isotopes

IDP2017

Trace elements

Electronic atlas

ABSTRACT

The GEOTRACES Intermediate Data Product 2017 (IDP2017) is the second publicly available data product of the international GEOTRACES programme, and contains data measured and quality controlled before the end of 2016. The IDP2017 includes data from the Atlantic, Pacific, Arctic, Southern and Indian oceans, with about twice the data volume of the previous IDP2014. For the first time, the IDP2017 contains data for a large suite of biogeochemical parameters as well as aerosol and rain data characterising atmospheric trace element and isotope (TEI) sources. The TEI data in the IDP2017 are quality controlled by careful assessment of intercalibration results and multi-laboratory data comparisons at crossover stations. The IDP2017 consists of two parts: (1) a compilation of digital data for more than 450 TEIs as well as standard hydrographic parameters, and (2) the eGEOTRACES Electronic Atlas providing an on-line atlas that includes more than 590 section plots and 130 animated 3D scenes. The digital data are provided in several formats, including ASCII, Excel spreadsheet, netCDF, and Ocean Data View collection. Users can download the full data packages or make their own custom selections with a new on-line data extraction service. In addition to the actual data values, the IDP2017 also contains data quality flags and 1-o data error values where available. Quality flags and error values are useful for data filtering and for statistical analysis. Metadata about data originators, analytical methods and original publications related to the data are linked in an easily accessible way. The eGEOTRACES Electronic Atlas is the visual representation of the IDP2017 as section plots and rotating 3D scenes. The basin-wide 3D scenes combine data from many cruises and provide quick overviews of large-scale tracer distributions. These 3D scenes provide geographical and bathymetric context that is crucial for the interpretation and assessment of tracer plumes near ocean margins or along ridges. The IDP2017 is the result of a truly international effort involving 326 researchers from 25 countries. This publication provides the critical reference for unpublished data, as well as for studies that make use of a large cross-section of data from the IDP2017.

This article is part of a special issue entitled: "Cycles of trace elements and isotopes in the ocean – GEOTR-ACES and beyond" - edited by Tim M. Conway, Tristan Horner, Yves Plancherel, and Aridane G. González.

1. Introduction

In 2014, the international GEOTRACES programme (Anderson et al., 2014a, 2014b; SCOR Working Group, 2007; GEOTRACES, 2006; Anderson and Henderson, 2005; Frank et al., 2003; http://www.geotraces.org/) released its first *Intermediate Data Product 2014* (IDP2014, Mawji et al., 2015). The main motivation was to not wait until the end of the programme to issue a final data product. Instead, GEOT-RACES wants to create and release a series of intermediate data products at times when the programme is still very active and expanding, both in terms of observational activities as well as the scientific analysis and synthesis of the data produced so far. By releasing and sharing data at early stages, GEOTRACES intends to strengthen and intensify collaboration within the geochemical community itself, but also to attract and invite colleagues from other communities, such as physical, biological and paleo-oceanography, as well as modelling, to apply their unique knowledge and skills to marine biogeochemical research questions.

The release of the IDP2014 was a big success and was widely covered by international news media as well as a broad range of scientific journals (e.g., Morrison, 2014). The data product resulted from a significant effort to combine data from 15 cruises conducted by seven countries. The IDP2014 data cover the Atlantic, Arctic, Southern and Indian oceans and span the 2007 to 2012 period. There are data for 237 hydrographic parameters as well as trace elements and isotopes (TEIs) contributed by 133 scientists from 16 countries. Having such a large group of researchers collaborate on the project and submit high-quality data, sometimes unpublished, was a remarkable achievement.

The IDP2014 is being used widely and has stimulated collaborative research that would not have been possible without such a large, aggregated dataset. Since its release, users worldwide have downloaded the IDP2014 dataset 1410 times. Users of the data product are encouraged to cite the original papers written by the data originators, but the IDP contains significant unpublished data. The publications describing the IDPs thus provide the critical reference for unpublished data, as well as for studies that make use of a large cross-section of data from the IDP. The publication describing the IDP2014 (Mawji et al., 2015) has been cited 43 times, indicating that there is a significant number of scientific studies, such as large-scale modelling and basin-scale to global TEI evaluations, that make use of large portions of the IDP2014 data and could not have been performed otherwise (e.g., Abadie et al., 2017; Chien et al., 2016; Frants et al., 2016; Lerner et al., 2016; Schlitzer, 2016). In particular, the aggregated dissolved iron datasets from IDP2014 facilitated the first rigorous intercomparison of dissolved iron cycling from 13 global ocean models (Tagliabue et al., 2016).

Building on the success of the IDP2014 and following the long-term data product release plan, GEOTRACES released its second intermediate data product (IDP2017) at the Goldschmidt Conference 2017 in Paris. As with the previous product, IDP2017 consists of two parts: (1) the digital data compilation of TEIs as well as standard hydrographic parameters; and (2) the *eGEOTRACES Electronic Atlas* providing section plots and animated 3D scenes of the data. As described in detail below, the IDP2017 contains twice as much data compared to the previous IDP2014. For the first time, the IDP2017 contains significant amounts of biogeochemistry data as well as data for aerosols and rain. All data in the IDP2017 have passed the GEOTRACES standardisation and intercalibration protocols.

2. Intercalibration of data for IDP2017

The direct comparability of GEOTRACES TEI data from any cruise is a prerequisite for assessing global-scale distributions of TEIs, for identifying and quantifying sources and sinks as well as rates of internal cycling, and for providing a baseline against which future changes can be measured. This is also essential for our ability to model natural processes affected by TEIs in the ocean. Therefore, the standardisation and quality control of data sets has always been a cornerstone of the GEOTRACES programme. The importance of intercalibration was illustrated by the U.S. National Science Foundation (NSF)-funded 2003 SAFe iron intercomparison cruise (Johnson et al., 2007), which resulted in widely used consensus material for dissolved trace metals and rare earth elements. Through the GEOTRACES programme, two additional intercalibration cruises were conducted for all the main TEIs and documented in a special issue of Limnology and Oceanography Methods in 2012 (Vol. 10 issue 6). Moreover, a cookbook detailing recommended sample collection methods was produced to support intercalibration (http://www.geotraces.org/images/Cookbook.pdf). This document was updated prior to IDP2017 with new intercalibration procedures for TEIs not included in the IDP2014.

While the IDP2014 contained some data that were not quality controlled (identified as tier 2 data), IDP2017 is the first GEOTRACES intermediate data product in which all TEI data have passed the intercalibration procedures and been approved by the Standards and Intercalibration Committee (S&I Committee). This committee is currently a group of eight members approved by the GEOTRACES Scientific Steering Committee. Its members cover a broad range of analytical expertise for the TEIs in IDP2017. In addition, there are element co-ordinators for each group of TEIs who can guide new investigators in developing sample collection and analytical methods (http://www.geotraces.org/sic/s-i-committee/elemental-coordinators).

The intercalibration assessment of the TEI parameters for IDP2017 differed depending on several criteria. For example, the committee had to consider the maturity of the available analytical techniques for a given TEI, the type of TEI in GEOTRACES, the possibly transient nature of the signal, the nature of the data acquisition (e.g., sensor vs. bottle), and the participation in other programmes (such as CLIVAR) that have their own intercalibration procedures.


Irrespective of the quality criteria for individual TEIs, all data were expected to follow certain minimum standards, as shown in Fig. 1. First, written documentation of sampling, measurement and intercalibration procedures was required, provided directly to the S&I Committee as an intercalibration report. This report included details on how samples were collected, how they were processed on board, and how they were stored prior to analyses. This assessment must be carried out for each individual cruise leg, not just for a given laboratory, since the sampling equipment, analytical techniques and analysts may change between cruise legs. The actual assessment was based on the information in these reports and took place during meetings of the S&I Committee.

Second, the methods were assessed for suitability, which included (for example) a check if the procedures were following the cookbook or equivalent, if there were sufficient blank assessments, if detection limits were adequate for the target, and if the laboratory had systems for checking the internal consistency of data, for example replicate analyses, analyses of certified reference materials, or analyses of consensus materials produced from GEOTRACES intercalibration cruises.

Third, the external comparability of the data was assessed. This crucial step comprised an assessment of the crossover stations for key TEIs, that is, those TEIs considered to be of such widespread interest that they should be measured on every GEOTRACES section, and for other TEIs whenever possible. If no crossover stations were possible (e.g., only one cruise had taken place in this region), external comparability had to be demonstrated by participation in a laboratory intercalibration exercise (if such an exercise was available), by the analysis of replicate samples (e.g., where samples were exchanged with another laboratory), and by analyses of certified reference materials or consensus materials. For some TEIs it could also include a comparison to other data in the region of interest. External validation for certain parameters with a core user group outside the GEOTRACES community (e.g., DIC & Alkalinity data, CFCs, sensor data) could also be demonstrated via some other programme (e.g., GO-SHIP, CLIVAR). The assessment of external comparability had to consider the state of the art for any given TEI, with the recognition that the state of the art is changing rapidly, in large part due to GEOTRACES intercalibration activities.

Finally, the S&I Committee assessed jointly if the information provided had demonstrated that the analytical methods reflected the state of the art, and if the data provided had satisfied the quality requirements. If information was missing, the committee contacted the analysts to see if additional information could be provided that would satisfy the need for documentation and quality assurance.

Several parameters have been intercalibrated through new intercalibration exercises (e.g., Si isotopes: Grasse et al., 2017, REE: Behrens et al., 2016; ⁷Be, particulate TEIs and leachable particulate trace metals; Hg speciation), and new consensus materials have become available for the use of the GEOTRACES community (e.g., Arizona Test Dust for aerosols; Morton et al., 2013). More recently, a sea-ice intercalibration has begun. Results from these on-going intercalibration exercises will be publicised by GEOTRACES as they become available.



Dialogue between S&I committee and analyst to obtain adequate quality control for IDP2017

Fig. 1. Flow chart of data assessment for IDP2017.

3. IDP2017 digital data

Creation of the IDP2017 was coordinated and overseen by the GEOTRACES Data Management Committee (DMC). Collation of the cruise data and linkage with extensive metadata was carried out at the GEOTRACES Data Assembly Centre (GDAC) located at the British Oceanographic Data Centre. GDAC received data submissions from four national data centres (Biological & Chemical Oceanography Data Management Office (BCO-DMO; https://www.bco-dmo.org/), Japan Oceanographic Data Centre (JODC; http://www.jodc.go.jp/jodcweb/), LEFE CYBER France (http://www.obs-vlfr.fr/proof/index2.php), NIOZ - Netherlands Data Centre (https://www.nioz.nl/en/research/research-data)) or from GEOTRACES data originators directly. The lead author of this publication carried out the integration of the cruise data into global datasets.

The IDP2017 digital data package consists of three datasets: (1) discrete water sample data; (2) CTD sensor data; and, as a new dataset, (3) aerosol and rain data. The discrete sample and aerosol/rain datasets contain the GEOTRACES TEI data as well as data for a large suite of standard hydrographic data (discrete sample dataset only). The CTD sensor dataset contains high-resolution data from a variety of electronic sensors that are useful for TEI data interpretation and evaluation.

The discrete sample datasets include data from 39 cruises conducted by 11 countries during the 7-year period from 2007 to 2014 (Table 1). Twenty-four of the 39 cruises are new in the IDP2017. The dataset covers the Arctic, Atlantic, Southern, Indian oceans and, the Pacific

Ocean (Fig. 2). The best coverage and highest station density is found in the Atlantic, but the new data from the Pacific have already allowed accurate mapping of TEI distributions in parts of the South and North Pacific. In addition to twelve GEOTRACES sections (GA01, GA02, GA03, GA04, GA06, GA10, GA11, GI04, GP02, GP13, GP16, and GP18), which eventually will produce measurements of the large set of GEO-TRACES key TEIs (Table 2 in GEOTRACES, 2006), the IDP2017 also includes data from six cruises conducted as part of the International Polar Year (GIPY2, GIPY4, GIPY5, GIPY6, GIPY11, and GIPY13; for an overview of IPY activities see: https://www.icsu.org/publications/ understanding-earths-polar-challenges-international-polar-year-2007-2008). For the first time, the IDP2017 also includes GEOTRACES Compliant Data from four cruises (GAc02, GPc01, GPc02, and GPc03) and six GEOTRACES Process Studies (GPpr01, GPpr02, GPpr04, GPpr05, GPpr07, and GPpr10). Typically, these activities produce smaller sets of TEI measurements and sometimes have limited geographical coverage. Nevertheless, compliant data and process studies fill gaps in the overall sampling scheme and provide invaluable data for the quantification of TEI sources and sinks as well as the study of the internal cycling of TEIs. Links to the cruise reports of all cruises in the IDP2017 are provided in Table 2.

In total, the IDP2017 discrete sample dataset contains data for 1810 stations. Of these stations, 817 provide full-depth coverage of the water column. There are data for a total of 458 parameters, including (1) classical hydrographic parameters and tracers such as temperature, salinity, oxygen, nutrients, CFCs, SF₆, Tritium, and He-3, (2) dissolved

Table 1

List of cruises included in the GEOTRACES Intermediate Data Product 2017. Section suffixes denote individual parts of a section. A lower case "c" in the section name (as in GAc01) indicates compliant data while a lower case "pr" (as in GPpr01) indicates a process study. A y in the New column indicates new sections in the IDP2017. Many of the already existing sections had new data added since IDP2014. Cruise locations are illustrated in Fig. 2.

| Section | Cruise | Chief scientist | Country | Start date | End date | New |
|-----------|---------------|------------------------|-------------|-------------|-------------|-----|
| GA01 | GEOVIDE | Sarthou, Geraldine | France | 15-May-2014 | 30-Jun-2014 | У |
| GA02 (n) | PE319 | Gerringa, Loes | Netherlands | 28-Apr-2010 | 26-May-2010 | |
| GA02 (c) | PE321 | Rijkenberg, Micha | Netherlands | 11-Jun-2010 | 08-Jul-2010 | |
| GA02 (s) | JC057 | Rijkenberg, Micha | Netherlands | 01-Mar-2011 | 07-Apr-2011 | |
| GA03 (e) | KN199-4 | Jenkins, William | USA | 15-Oct-2010 | 04-Nov-2010 | |
| GA03 (w) | KN204-1 | Boyle, Edward | USA | 06-Nov-2011 | 11-Dec-2011 | |
| GA04 (n1) | PE370 | Rijkenberg, Micha | Netherlands | 14-May-2013 | 05-Jun-2013 | у |
| GA04 (bs) | PE373 | Rijkenberg, Micha | Netherlands | 13-Jul-2013 | 25-Jul-2013 | y |
| GA04 (n2) | PE374 | Rijkenberg, Micha | Netherlands | 25-Jul-2013 | 11-Aug-2013 | y |
| GA04 (s) | MedSeA | Garcia Orellana, Jordi | Spain | 05-May-2013 | 01-Jun-2013 | y |
| GA06 | D361 | Achterberg, Eric | ŮK | 07-Feb-2011 | 19-Mar-2011 | y |
| GA10 (e) | D357 | Henderson, Gideon | UK | 18-Oct-2010 | 22-Nov-2010 | 5 |
| GA10 (w) | JC068 | Henderson, Gideon | UK | 24-Dec-2011 | 27-Jan-2012 | |
| GA11 | M81 1 | Frank, Martin | Germany | 04-Feb-2010 | 08-Mar-2010 | |
| GAc01 | KN192-5 | Saito, Mak | USA | 16-Nov-2007 | 13-Dec-2007 | |
| GAc02 | AE1410 | Conte, Maureen | USA | 31-May-2014 | 08-Jun-2014 | у |
| GI04 | KH09-05 | Gamo, Toshitaka | Japan | 06-Nov-2009 | 10-Jan-2010 | 5 |
| GIPY02 | AU0703 | Griffiths, Brian | Australia | 21-Jan-2007 | 19-Feb-2007 | |
| GIPY04 | MD166 | Speich, Sabrina | France | 08-Feb-2008 | 24-Mar-2008 | |
| GIPY05 | ANT_XXIV_3 | Fahrbach, Eberhard | Germany | 06-Feb-2008 | 16-Apr-2008 | |
| GIPY06 | AU0806 | Rintoul, Steve | Australia | 22-Mar-2008 | 17-Apr-2008 | |
| GIPY11 | ARK_XXII_2 | Schauer, Ursula | Germany | 29-Jul-2007 | 07-Oct-2007 | |
| GIPY13 | ISSS-08 | Semiletov, Igor | Sweden | 18-Aug-2008 | 18-Sep-2008 | у |
| GP02 | KH12-4 | Gamo, Toshitaka | Japan | 23-Aug-2012 | 03-Oct-2012 | y |
| GP13 | SS2011-1 | Bowie, Andrew | Australia | 13-May-2011 | 05-Jun-2011 | y |
| GP13 | TAN1109-2 | Boyd, Philip | New Zealand | 06-Jun-2011 | 30-Jun-2011 | y |
| GP16 | TN303-EPZT | Moffett, James | USA | 25-Oct-2013 | 20-Dec-2013 | y |
| GP18 | KH11-7 | Zhang, Jing | Japan | 16-Jul-2011 | 04-Aug-2011 | y |
| GPc01 | SO202 | Gersonde, Rainer | Germany | 07-Jul-2009 | 29-Aug-2009 | y |
| GPc02 | ANT_XXVI_2 | Gersonde, Rainer | Germany | 27-Nov-2009 | 27-Jan-2010 | y |
| GPc03 | KM1128 | Lamborg, Carl | USA | 03-Oct-2011 | 24-Oct-2011 | y |
| GPpr01 | TAN0811 | Boyd, Philip | New Zealand | 15-Sep-2008 | 04-Oct-2008 | y |
| GPpr02 | SS01/10 | Hassler, Christel | Australia | 23-Jan-2010 | 15-Feb-2010 | у |
| GPpr04 | SO223T | Mohtadi, Mahyar | Germany | 09-Sep-2012 | 08-Oct-2012 | у |
| GPpr05 | KM1107 | Taylor, Brian | USA | 23-Feb-2011 | 25-Feb-2011 | y |
| GPpr07 | LineP_2012-13 | Robert, Marie | Canada | 14-Aug-2012 | 30-Aug-2012 | y |
| GPpr07 | LineP_2013-18 | Robert, Marie | Canada | 20-Aug-2013 | 05-Sep-2013 | y |
| GPpr07 | LineP_2014-19 | Robert, Marie | Canada | 19-Aug-2014 | 04-Sep-2014 | у |
| GPpr10 | TAN1212 | Boyd, Philip | New Zealand | 23-Sep-2012 | 23-Sep-2012 | y |

R. Schlitzer et al.




Fig. 2. Map of discrete sample stations included in the GEOTRACES Intermediate Data Product 2017. A lower case "c" in the section name (as in GAc01) indicates compliant data while a lower case "pr" (as in GPpr01) indicates a process study. Different colours and symbols are used to help distinguish between close-by sections. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

| Table 2 |
|--|
| Links to cruise reports of cruises included in the GEOTRACES Intermediate Data Product 2017. |

| Cruise | Cruise report |
|------------|---|
| AE1410 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/atlanticexplorer_ae1410.pdf |
| ANT_XXIV_3 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/polarstern_antxxiv3.pdf |
| ANT_XXVI_2 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/polarstern_ps75.pdf |
| ARK_XXII_2 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/polarstern_arkxxii2_07.pdf |
| AU0703 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/auroraaustralis0703.pdf |
| AU0806 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/auroraaustralis0806.pdf |
| D357 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/d357.pdf |
| D361 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/d361.pdf |
| TN303-EPZT | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/thomasgthompson_tn303.pdf |
| GEOVIDE | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/pourquoipas_geovide.pdf |
| ISSS-08 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/isss08.pdf |
| JC057 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/jc057.pdf |
| JC068 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/jc068.pdf |
| KH09-05 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/hakuhomaru_kh-09-5.pdf |
| KH12-4 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/hakuhomaru_kh12.pdf |
| KN199-4 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/kn199-4.pdf |
| KN204-1 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/knorr_kn204_1.pdf |
| KN192-5 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/kn192-5.pdf |
| MD166 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/mariondufresne166.pdf |
| MedSeA | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/angelesalvarino_medsea.pdf |
| M81_1 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/meteor81_1.pdf |
| PE319 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/pe319.pdf |
| PE321 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/pe321.pdf |
| PE370 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/pelagia_pe370.pdf |
| PE373 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/pe373.pdf |
| PE374 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/pe374.pdf |
| SO202 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/sonne_so202.pdf |
| SO223T | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/sonne_so223t.pdf |
| SS01/10 | https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/southernsurveyor01_2010.pdf |
| SS2011-1 | https://www.bodc.ac.uk/resources/inventories/cruise inventory/reports/ss2011.pdf |
| TAN1109-2 | https://www.bodc.ac.uk/resources/inventories/cruise inventory/reports/tangaroa1109.pdf |

and particulate trace elements such as Al, Ba, Cd, Cu, Fe, Mn, Mo, Ni, Pb, Zn and Rare Earth Elements (REEs), (3) stable isotopes such as H-2, C-13, N-15, O-18, Si-30, Fe-56, Cd-114, and Nd-143 as well as (4)

radioactive isotopes such as Pb-210, Po-210, Th-230, Pa-231, and Th-234. The IDP2017 discrete sample dataset also contains data for a wide range of biogeochemistry parameters, such as HPLC pigments,

Table 3

Number of measurements of selected GEOTRACES parameters in the discrete sample dataset of the IDP2017. Numbers in parentheses indicate the percentage of discrete samples that contain data for that parameter. The "All forms" values include dissolved as well as particulate measurements. For Fe this also includes data for Fe_II and soluble Fe.

| Parameter | Number of observations |
|-----------------|---|
| Trace elements | |
| Fe | All forms: 12,050 (25.8%); dissolved: 7690 (16.4%) |
| Mn | All forms: 10,375 (22.2%); dissolved: 6984 (14.9%) |
| Al | All forms: 10,656 (22.8%); dissolved: 7262 (15.5%) |
| Zn | All forms: 8787 (18.8%); dissolved: 6932 (14.8%) |
| Cd | All forms: 10,564 (22.6%); dissolved: 7197 (15.4%) |
| Pb | All forms: 9181 (19.6%); dissolved: 6157 (13.2%) |
| Cu | All forms: 7081 (15.1%); dissolved: 3996 (8.5%) |
| Stable isotopes | |
| Si-30 | All forms: 246 (0.5%); silicate: 246 (0.5%) |
| 0-18 | All forms: 1926 (4.1%); water: 1926 (4.1%) |
| N-15 | All forms: 1972 (4.2%); nitrate: 1972 (4.2%) |
| C-13 | All forms: 1113 (2.4%); DIC: 1113 (2.4%) |
| Radioactive iso | topes |
| Th-234 | All forms: 3768 (8.1%); dissolved plus total particulate: 2520 (5.4%) |
| Th-230 | All forms: 1805 (3.9%); dissolved: 1389 (3.0%) |
| Pa-231 | All forms: 1684 (3.6%); dissolved: 1292 (2.8%) |
| Pb-210 | All forms: 684 (1.5%); dissolved: 493 (0.9%) |
| Radiogenic isot | topes |
| Nd-143 | All forms: 696 (1.5%); dissolved: 684 (1.5%) |

metalloproteomics on filtered particles and metal content of single cells.

A total of 46,794 discrete samples were analysed from the 1810 stations. The average number of depths sampled at each station was 33 but reached up to 182 depths at heavily sampled "super" stations. Table 3 summarises the number of observations for selected parameters, including micronutrients essential to life in the ocean (e.g., Fe, Zn, Cd, Cu), tracers of modern processes in the ocean (e.g., Al, Mn, N-

15), tracers significantly perturbed by human activities (e.g., Hg, Pb), and tracers used as proxies to reconstruct the past (e.g., Th-230, Pa-231, Nd isotopes). Data for the micronutrients are most abundant, with the total number of Fe measurements totalling 12,050; of these, 7690 are for dissolved Fe alone. There are 3768 data values for the radioactive isotope Th-234 and around 1800 values for Th-230 and Pa-231.

The CTD sensor dataset contains temperature, salinity, oxygen, fluorescence, transmissometer, turbidity, and photosynthetically active radiation (PAR) data at 1827 stations at 1 m vertical resolution. The fluorescence and transmissometer data provide information on phytoplankton abundance and suspended particle concentrations and are thus important for the interpretation of TEI data. Where calibrated data were not available, raw values are provided. These uncalibrated data are still useful as they reveal the horizontal and vertical extent of phytoplankton patches and suspended particle layers.

For the first time, the IDP2017 contains TEI aerosol and rain data sampled from GEOTRACES cruises. Such data were collected at 243 locations in the Atlantic, Pacific, Mediterranean, and Black Sea (Fig. 3). Data are provided for 99 aerosol parameters, including total TEI concentrations as well as soluble TEI after strong or mild leaching. Also included are size-fractionated TEI concentrations on fine and coarse aerosols. The rain data consist of 68 parameters, including dissolved and total dissolvable TEI concentrations.

In addition to the actual data values, the IDP2017 also contains data quality flags and 1- σ data error values where available. Quality flags and error values are useful for data filtering and statistical data analysis. Quality flags are single character codes reflecting the quality of the respective data value. The IDP2017 uses the IODE quality flag set that is a standard flagging scheme for the exchange of oceanographic and marine meteorological data (www.iode.org/mg54_3). The IODE flagging scheme is generic and simple, only containing the five flags listed in Table 4.

The IDP2017 is an "intermediate" product, and there is clearly a significant amount of further data to come from GEOTRACES cruises,




Fig. 3. Map of aerosol and rain stations included in the GEOTRACES Intermediate Data Product 2017.

Table 4

The IODE quality flagging scheme used for the IDP2017.

| Value | Flag short name | Definition |
|-------|--|--|
| 1 | Good | Passed required QC tests |
| 2 | Not evaluated, not available or unknown | Used for data when no QC test performed or the information on quality is not available |
| 3 | Questionable/suspect | Failed non-critical metric or subjective test (s) |
| 4 | Bad | Failed critical QC test(s) or as assigned by the data provider |
| 9 | Missing data | Used as place holder when data are missing |

both those represented in the IDP2017, and those sections more recently completed or planned. The IDP2017 contains only those data that were completed and submitted before a cut-off date of December 2016. Further data will be included in subsequent intermediate products (as detailed below) and will significantly augment the data coverage represented in IDP2017.

4. Parameter naming conventions

The overall structure of the databases combined in IDP2017,

Table 5

including a single convention for naming all parameters (variables), was organised by a six-person Parameter Naming Committee (PNC) whose members interacted regularly with the Data Management Committee and with the Standards and Intercalibration Committee. With well over 400 parameters in IDP2017, and with the expectation that the number of parameters could eventually exceed 1000, a structure was sought that would allow users to search intuitively for data, using either tools incorporated into IDP2017 (see "Obtaining IDP2017 Data") or other search engines, utilising a common set of keywords or commands. The structure was designed to accommodate hydrographic and biogeochemical variables as well as TEIs, and to span a range of sampling environments, including seawater, aerosols and rain, while also anticipating the future addition of data from sea ice and sediments. With this in mind, the PNC devised a six-token parameter naming scheme, described in the next two paragraphs, that would encompass all of these characteristics as well as information about operationally defined chemical speciation and physical form of the substance of interest. It is hoped that incorporating all of this information into each parameter name will facilitate searches for highly specific types of data.

The IDP2017 employs the following parameter naming scheme. Standard hydrographic parameters, such as temperature, salinity and oxygen use names as defined in the WOCE/CLIVAR naming convention (CTDTMP, CTDSAL and CTDOXY for temperature, salinity and oxygen

| | 2 | 3 | 4 | 5 | 6 |
|---------------------|--|-------------------|---------------|--|------------------------|
| lement/ compound | [_Oxidation State] | [_Atomic Mass] | _Phase | _DataType | _Sampling System |
| Explanation | | | | Example | |
| Element or | compound (mandatory) | | | Fe, Th, DIC, NO3, L1Fe | |
| Oxidation s | ate as roman number (optional) | | | _II, _IV, _III_V_ where III and V ar | e combined |
| Atomic mas | s (optional); two entries for isotope ratios | | | _228, _208_204 | |
| | nich element or compound was measured | | | _A (aerosol) | |
| | Total Dissolvable concentration of a cons | | the dissolved | _C (colloidal) | |
| concentratio | on of the monomethyl form of a constituen | it) | | _D (dissolved) | |
| | | | | _DL (dissolved labile) | |
| | | | | _F (free (un-complexed)) | |
| | | | | _LPT (large particulate, total (unl | leached)) |
| | | | | _R (rain) | |
| | | | | _S (soluble) | |
| | | | | _SML (soluble mild leach) | |
| | | | | _SSL (soluble strong leach) | |
| | | | | _SP (small particulate) | |
| | | | | _SPL (small particulate, labile fra | |
| | | | | _SPR (small particulate, refractor | • |
| | | | | _SPT (small particulate, total (un | leached)) |
| | | | | _T (total) | |
| | | | | _TD (total dissolvable) | |
| | | | | _TP (total particulate) | ation) |
| | | | | _TPL (total particulate, labile frac | |
| DeteTrate (| n an datama) | | | _TPR (total particulate, refractory | (fraction) |
| DataType (1 | nandatory) | | | _CONC (concentration) | station) |
| | | | | _DELTA (isotope ratio in delta no _EPSILON (isotope ratio in epsilo | |
| | | | | _LogK (log of binding constant of | |
| | | | | _RATIO (atomic abundance ratio | |
| Sampling es | stem (mandatory) | | | _BOTTLE (Niskin or similar water | · · |
| compring sy | com (mandatory) | | | _FISH (trace-metal clean towed s | |
| | | | | _PUMP (either in-situ pump or or | |
| | | | | _UWAY (ship's underway surface | |
| | | | | _HIVOL (high-volume aerosol san | |
| | | | | _LOWVOL (low-volume aerosol s | - |
| | | | | FINE_IMPACTOR (size-fractionat | |
| | | | | fraction) | -, |
| | | | | _COARSE_IMPACTOR (size-fraction | onated aerosols, large |
| | | | | fraction) | , . 0 |
| | | | | _AUTO (automated aerosol samp | ler) |
| | | | | _MAN (aerosol sampler with man | |

Chemical Geology 493 (2018) 210-223

from CTD sensors; https://exchange-format.readthedocs.io/en/latest/ parameters.html). Other hydrographic parameters use names defined intuitively. Examples are PRESSURE for the CTD pressure at the bottle sample depth, SALINITY, PHOSPHATE, NITRATE, and SILICATE for salinity, phosphate, nitrate and silicate measured on bottle samples. Biogeochemistry parameters in the IDP2017 use names defined by SCOR naming conventions (e.g., HPLC pigments; Roy et al., 2011) or names that intuitively define the parameters (e.g., nifH_UCYN-

click on the info symbol () is required to open the respective info file in the web browser and obtain detailed information about the data originator and the analytical methods for the clicked parameter and cruise. One more mouse click shows the references of the original publications associated with the given parameter and cruise. Fig. 4 shows an ex-

| <u> </u> | I I I I I I I I I I I I I I I I I I I | , | | | |
|------------------|---------------------------------------|----------------|--------|-----------|------------------|
| 1 | 2 | 3 | 4 | 5 | 6 |
| Element/compound | [_Oxidation State] | [_Atomic Mass] | _Phase | _DataType | _Sampling System |
| | | | | | |

A_DNA_P_CONC_BOTTLE; concentration of nifH genes from uncultured unicellular cyanobacteria (UCYN-A) particles (P) in a bottle sample).

All other trace elements and isotope names are composed of up to six separate tokens, as follows:

Tokens 2 and 3 are optional, while all other tokens are mandatory. The meaning and possible values for all the six tokens are described in Table 5. Example parameter names can be found in Table 6.

The PNC sought to verify that parameter names supplied by contributing investigators complied with the convention described above. In cases where reported data did not comply with a master list of parameters, the PNC would examine the metadata accompanying the original data submission and rename the parameter if appropriate. If there were any question about the correct parameter name, then the PNC would contact the data originator to verify that the parameter had been renamed correctly.

5. Metadata and publication references

The IDP2017 digital datasets include the cruise reports of all the cruises (Table 2). These cruise reports provide detailed documentation of the ship operations, including descriptions of sampling procedures and gear as well as information on the laboratories and principal investigators involved. Access to the cruise reports is very easy. When using the ODV collection version of the IDP2017, a simple mouse click on the Cruise Report meta-variable opens the given cruise report and allows viewing in the web browser.

In addition, the IDP2017 also contains, for every parameter and every cruise, a data info file containing information about data originators, sample preparation and analytical methods as well as links to original publications related to the data. These info files are delivered with all IDP2017 output formats and can be viewed easily in the web browser. Access is particularly easy in ODV, where only one mouse

Table 6

| Example IDP2017 | parameter names. |
|-----------------|------------------|
|-----------------|------------------|

ample publication list for parameter Fe_D_CONC_BOTTLE along GP16. Proper linkage of the originator and publication information with the actual data is an important feature of the IDP2017 that makes it easy for users to identify, contact, and acknowledge originators.

The publication links in the IDP2017 info files refer to the reference database of original publications maintained at the GEOTRACES International Programme Office (IPO). This reference database is dynamic and updated whenever new papers are published. Clicking on a reference link in the IDP2017 will always show the up-to-date publication list at the time of the click. Future requests of the publication list related to, for instance, Fe D CONC BOTTLE along GP16 will, in addition to what is shown in Fig. 4, also include new papers published since the Fig. 4 creation date of December 2017. This dynamic inclusion of papers published after the release of the data product was a required feature for the IDP2017, because many datasets were unpublished at the time of data submission.

As a novelty for the IDP2017, the GEOTRACES IPO has made the publication database into a searchable on-line database available on the following GEOTRACES web page: http://www.geotraces.org/library-88/scientific-publications/peer-reviewed-papers. This database is not limited to the IDP2017 as it also includes other publications that are relevant for GEOTRACES research along with Master and PhD dissertations. Three types of search functionalities are available:

- (1) Simple search: users can search publications by "author", "title" or "journal" entering the desired term into a search box,
- (2) Advanced search: by means of dropdown menus, users can select publications by "author", "title", "GEOTRACES cruise", "year" or "type of document", and.

| Parameter name | Parameter description |
|-------------------------------|---|
| Fe_D_CONC_BOTTLE | Concentration of dissolved Fe |
| Fe_II_D_CONC_BOTTLE | Concentration of dissolved Fe(II) |
| Fe_II_TP_CONC_BOTTLE | Concentration of total particulate Fe(II) determined by filtration from a water sampling bottle |
| Fe_TPL_CONC_BOTTLE | Concentration of labile particulate iron determined by filtration from a water sampling bottle |
| Nd_143_144_D_RATIO_BOTTLE | Atom ratio of given isotopes for dissolved Nd |
| Nd_143_144_D_EPSILON_BOTTLE | Atom ratio of dissolved Nd isotopes expressed in conventional EPSILON notation |
| Cd_114_110_D_DELTA_BOTTLE | Atom ratio of dissolved Cd isotopes expressed in conventional DELTA notation |
| Cu_Cu'_D_CONC_BOTTLE | Concentration of dissolved inorganic Cu |
| Pb_206_204_D_RATIO_BOTTLE | Atom ratio of given isotopes for dissolved Pb |
| DIC_13_12_D_DELTA_BOTTLE | Atom ratio of given isotopes for dissolved C as DIC in delta notation |
| DIC_14_12_D_DELTA_BOTTLE | Atom ratio of radiocarbon as dissolved C in DIC in DELTA notation |
| NITRATE_15_14_D_DELTA_BOTTLE | Atom ratio of given isotopes for dissolved N as nitrate in delta notation |
| L1_Fe_D_CONC_BOTTLE | Concentration of dissolved L1 Fe-binding ligand |
| L1_Fe_D_LogK_BOTTLE | Log of the stability constant of L1 Fe |
| HOMOCYS_D_CONC_BOTTLE | Concentration of dissolved homocysteine |
| Chl a_HPLC_P_CONC_BOTTLE | Concentration of particulate Chlorophyll a measured using HPLC method |
| nifH_UCYN-A_DNA_P_CONC_BOTTLE | Abundance nifH Uncultured unicellular cyanobacteria (UCYN-A) |

GEOTRACES References

Cruise: GP16 Parameter: Fe_D_CONC_BOTTLE

6 publications found. Export -

2017

Fitzsimmons, J. N., John, S. G., Marsay, C. M., Hoffman, C. L., Nicholas, S. L., Toner, B. M., German, C. R., & Sherrell, R. M. (2017). Iron persistence in a distal hydrothermal plume supported by dissolved-particulate exchange. *Nature Geoscience*, 10(3), 195–201. doi:10.1038/ngeo2900

Heller, M. I., Lam, P. J., Moffett, J. W., Till, C. P., Lee, J.-M., Toner, B. M., & Marcus, M. A. (2017). Accumulation of Fe oxyhydroxides in the Peruvian oxygen deficient zone implies non-oxygen dependent Fe oxidation. *Geochimica et Cosmochimica Acta*, 211, 174–193. doi:10.1016/j.gca.2017.05.019

John, S. G., Helgoe, J., Townsend, E., Weber, T., DeVries, T., Tagliabue, A., Moore, K., Lam, P., Marsay, C. M., & Till, C. (2017). Biogeochemical cycling of Fe and Fe stable isotopes in the Eastern Tropical South Pacific. *Marine Chemistry*. doi:10.1016/j.marchem.2017.06.003

Sanial, V., Kipp, L. E., Henderson, P. B., van Beek, P., Reyss, J.-L., Hammond, D. E., Hawco, N. J., Saito, M. A., Resing, J. A., Sedwick, P., Moore, W. S., & Charette, M. A. (2017). Radium-228 as a tracer of dissolved trace element inputs from the Peruvian continental margin. *Marine Chemistry*. doi:10.1016/j.marchem.2017.05.008

2016

Boiteau, R. M., Mende, D. R., Hawco, N. J., McIlvin, M. R., Fitzsimmons, J. N., Saito, M. A., Sedwick, P. N., DeLong, E. F., & Repeta, D. J. (2016). Siderophorebased microbial adaptations to iron scarcity across the eastern Pacific Ocean. *Proceedings of the National Academy of Sciences of the United States of America*, 113(50), 14237–14242. doi:10.1073/pnas.1608594113

2015

Resing, J. A., Sedwick, P. N., German, C. R., Jenkins, W. J., Moffett, J. W., Sohst, B. M., & Tagliabue, A. (2015). Basin-scale transport of hydrothermal dissolved metals across the South Pacific Ocean. *Nature*, 523(7559), 200–203. doi:10.1038/nature14577

Fig. 4. Example list of publications for parameter Fe_D_CONC_BOTTLE along GP16 as of December 2017.

(3) Parameter search: allows users to access a list of publications by specific TEI. In addition users can also retrieve publications by group of parameters (e.g., Aerosols, Dissolved TEIs, etc.) or by predefined subgroups (e.g., dissolved trace elements, etc.).

In each case, search queries for "parameter" or "GEOTRACES cruise" will only list those publications linked to data included in the IDP2017.

6. Obtaining IDP2017 data

The IDP2017 digital data are available in two forms: (1) as full package downloads, or, (2) as customised data subsets using a new online data extraction service. Both methods require users to register (or login if already registered) and agree to IDP2017 usage rules before being able to access and download IDP2017 digital data. The usage rules ask for proper citation of the relevant original papers associated with the particular data used, as well as citation of the IDP2017 data product itself (this paper). Users are also asked to describe the purpose of the IDP2017 data download.

Full packages of the three IDP2017 datasets are available for download at https://www.bodc.ac.uk/geotraces/data/idp2017/. The data are provided in four formats: (1) ASCII text files suitable for usage in standard software, (2) Excel spreadsheet files for Microsoft Excel or similar software, (3) netCDF files suitable for access by models and netCDF readers, and (4) as ODV collections for use with the popular *Ocean Data View* software (https://odv.awi.de).

Users who only need data for a smaller subset of parameters and/or smaller geographical domain can use the new data subsetting and extraction service provided at https://webodv.awi.de/geotraces. After registration and login the user is guided through a three-step procedure. Step 1 allows for subsetting the set of stations to be downloaded by selecting one or more entries from the cruise list, zooming into a specific map domain and/or specifying one or more required parameters (variables). Only stations containing data for all the selected required parameters are included in the output dataset. Step 2 lets users customise the set of parameters (variables) to be included in the download file. This is done using a hierarchical tree of parameter groups and individual parameters. Users open/close parameter groups by clicking the +/- symbols. All parameters of a given group are selected/unselected by clicking the specific group box; individual parameters are selected/unselected by clicking the box of the individual parameter. A *Selection status* box always shows the currently selected numbers of stations and parameters (variables) to be included in the download file. Step 3 lets users choose among four data output formats (ASCII, ODV collection, netCDF, or WOCE WHP exchange) and initiate the actual data download. Selection settings are remembered when a user exits the session and are restored when logging in again later.

7. eGEOTRACES electronic atlas

The *eGEOTRACES Electronic Atlas* is the visual component of the IDP2017 and provides 593 section plots (Fig. 5) and 132 animated 3D scenes (Fig. 6) for many (but not all) of the parameters in the IDP2017. All plots are based on the digital data in the IDP2017, but data values flagged as *Questionable/suspect* or *Bad* (see Table 4) were filtered out and not used for the plots. The *eGEOTRACES* website http://egeotraces.org/ provides a dynamic map, where users start by selecting a data group and a tracer of interest. Sections containing a plot for the selected tracer are highlighted in red in the map, and basins containing a 3D animation for the selected tracer are highlighted in blue. Clicking on a red section label or a blue basin label will show the respective section plot or play the respective 3D scene. All section plots and 3D scenes show the names of scientists who produced or are responsible for the data. This makes it easy for users to identify and acknowledge data producers.




Fig. 5. Example eGEOTRACES section page.

Further clicking on a section plot loads a high-resolution version of the image, which can be saved for use in publications and presentations. The browser's *Back* button is used to return to the original section page. When viewing a rotating 3D scene clicking the *Larger-size Video* link produces a blown-up version of the animation. Clicking the *Normalsize Video* link at the bottom of the blown-up animation returns to the original size. An options bar appears when the mouse is over the 3D animation. Elements on the options bar can be used to stop the animation at arbitrary angles and quickly choose other viewing angles. Some browsers also allow download of the 3D movie file.

All section and 3D animation pages contain groups of links near the bottom of the page. These include (a) links to other tracers along this section or in this scene, (b) other 3D scenes with this tracer, and (c) other sections with this tracer. These links greatly facilitate switching between and comparing of different tracers, sections, and 3D scenes. All section plots use the same window layout, and the different section plots perfectly match when switching between tracers. The links under category (c) allow easy transitions between section plots and 3D animations.

Section and 3D scene pages also contain links to the original publications associated with the given tracer and section. Clicking on these links shows the current list of publications from the dynamically updated reference database maintained at the GEOTRACES IPO (see above).

eGEOTRACES provides quick overviews of the distributions of many geochemically relevant tracers. The 3D scenes provide geographical

and bathymetric context crucial for correctly assessing the extent and origin of tracer plumes as well as for inferring processes acting on the tracers and shaping their distribution. The numerous links to other tracers, sections, and basins found on section plots and 3D animations allow quick switching between tracers and domains, and facilitate comparisons between tracers. In addition to the benefit for scientific research, *eGEOTRACES* and its visual material can also help in teaching and outreach activities. The *eGEOTRACES* visuals can also help convey societally relevant scientific results to interested non-scientists and policy makers.

Images or 3D movies from the *eGEOTRACES* Atlas can be used free of charge for non-commercial purposes, such as in scientific publications, posters, presentations and teaching activities, as long as the source is cited as follows: *Schlitzer*, *R.*, *eGEOTRACES* - *Electronic Atlas of GEOTRACES* Sections and Animated 3D Scenes, http://egeotraces.org, 2017. Users must not remove the names of data producers and graphics creator. High-resolution images of the 3D scenes are available on request.

8. Summary

The new IDP2017 is a significant improvement over the earlier IDP2014 and roughly doubles the number of included cruises, stations, samples and parameters. The IDP2017 is a truly international product containing data from 326 researchers from 25 countries. The IDP2017




Fig. 6. Example eGEOTRACES 3D scene.

provides data for the Pacific Ocean, and the Mediterranean and Black seas, in addition to Atlantic, Arctic and Indian Oceans that were already represented in the previous data product. For the first time, the IDP2017 contains significant amounts of biogeochemistry data as well as TEI data for aerosols and rain. As before, users can obtain complete IDP2017 data sets as bulk downloads. Alternatively, there is now a customisable online data extraction service that allows data selections by domain, GEOTRACES sections, as well as parameters of interest. The extractor delivers smaller, more manageable data packages.

GEOTRACES invites and promotes use of the IDP2017 in the widest possible sense and envisages intensified collaboration within the marine geochemical community and beyond. Availability of large integrated and quality-controlled datasets, such as the IDP2017, enables a much wider range of studies than would be possible with individual single-cruise data alone.

The new, updated *eGEOTRACES* electronic atlas now contains more than 590 section plots (compared to 330 in IDP2014) and more than 130 animated 3D scenes (95 in IDP2014). Section and 3D scene pages are interlinked, and switching between different GEOTRACES sections, ocean basins and parameters is achieved with simple mouse clicks. *eGEOTRACES* section and 3D scene pages are now connected to the

GEOTRACES publication database, easily providing with a simple mouse click up-to-date reference lists to the original publications related to the displayed data. This feature makes identification of data originators easy and encourages proper citation or initiation of collaborative research.

The animated 3D scenes in the *eGEOTRACES Atlas* show large amounts of data in an intuitive way and with geographic and bathymetric context, thereby providing quick large-scale overviews of TEI distributions and helping the scientific interpretation of TEI data. In addition, these animations are also appealing to a wider target community, including scientists from other disciplines or policy makers, as well as interested members of the general public. GEOTRACES encourages wide usage of *eGEOTRACES* visuals for all purposes, including teaching and outreach.

The IDP2017 is the second in a series of planned intermediate data products, with the next scheduled for release in 2021. Future data products will extend the geographical coverage by including data from new GEOTRACES cruises, as well as providing additional data from existing cruises for parameters that take longer to measure and complete. GEOTRACES invites user feedback (ipo@geotraces.org) on the IDP2017 to help make the next IDP an even more useful product.

Acknowledgements

We gratefully acknowledge financial support by the Scientific Committee on Oceanic Research (SCOR) through grants from the U.S. National Science Foundation, including grants OCE-0608600, OCE-0938349, OCE-1243377, and OCE-1546580. Financial support was also provided by the UK Natural Environment Research Council (NERC), the Ministry of Earth Science of India, the Centre National de Recherche Scientifique, l'Université Paul Sabatier de Toulouse, the Observatoire Midi-Pyrénées Toulouse, the Universitat Autònoma de Barcelona, the Kiel Excellence Cluster The Future Ocean, the Swedish Museum of Natural History, The University of Tokyo, The University of British Columbia. The Royal Netherlands Institute for Sea Research, the GEOMAR-Helmholtz Centre for Ocean Research Kiel, and the Alfred Wegener Institute. We thank Edward Urban for his sustained strong support of the GEOTRACES programme and for all his assistance during the creation of the intermediate data products and the preparation of this publication.

References

- Abadie, C., Lacan, F., Radic, A., Pradoux, C., Poitrasson, F., 2017. Iron isotopes reveal distinct dissolved iron sources and pathways in the intermediate versus deep Southern Ocean. Proc. Natl. Acad. Sci. 114 (5), 858–863. http://dx.doi.org/10.1073/ pnas.1603107114.
- Anderson, R.F., Henderson, G., 2005. GEOTRACES. A global study of the marine biogeochemical cycles of trace elements and their isotopes. Oceanography 18 (3), 76–79. Anderson, R.F., Jeandel, C., Schlitzer, R., 2014a. GEOTRACES - marine biogeochemical
- cycles of trace elements and their isotopes. GSSA Geo. Bull. 57 (4), 37-40. Anderson, R.F., Mawji, E., Cutter, G.A., Measures, C.I., Jeandel, C., 2014b. GEOTRACES:
- changing the way we explore ocean chemistry. Oceanography 27 (1), 50–61. http:// dx.doi.org/10.5670/oceanog.2014.07.
- Behrens, M.K., Muratli, J., Pradoux, C., Wu, Y., Böning, P., Brumsack, H.-J., Goldstein, S.L., Haley, B., Jeandel, C., Paffrath, R., Pena, L.D., Schnetger, B., Pahnke, K., 2016. Rapid and precise analysis of rare earth elements in small volumes of seawater method and intercomparison. Mar. Chem. 186, 110–120.
- Chien, C.-T., Mackey, K.R.M., Dutkiewicz, S., Mahowald, N.M., Prospero, J.M., Paytan, A., 2016. Effects of African dust deposition on phytoplankton in the western tropical Atlantic Ocean off Barbados. Glob. Biogeochem. Cycles 30 (5), 716–734. http://dx. doi.org/10.1002/2015GB005334.
- Frank, M., Jeandel, C., Anderson, R.F., Henderson, G., Francois, R., Sharma, M., 2003. GEOTRACES: studying the global marine biogeochemistry of trace elements and

isotopes. EOS Trans. AGU 84 (34).

- Frants, M., Holzer, M., DeVries, T., Matear, R., 2016. Constraints on the global marine iron cycle from a simple inverse model. J. Geophys. Res. Biogeosci. 121 (1), 28–51. http://dx.doi.org/10.1002/2015JG003111.
- GEOTRACES, 2006. GEOTRACES (An International Study of the Marine Biogeochemical Cycles of Trace Elements and Their Isotopes): Science Plan. ISBN 1932-794. http:// www.geotraces.org/science/science-plan.
- Grasse, P., Brzezinski, M.A., Cardinal, D., de Souza, G.F., Andersson, P., Closset, I., Cao, Z., Dai, M., Ehlert, C., Estrade, N., 2017. GEOTRACES inter-calibration of the stable silicon isotope composition of dissolved silicic acid in seawater. J. Anal. At. Spectrom. 32, 562–578.
- Johnson, K.S., Elrod, V., Fitzwater, S., Plant, J., Boyle, E., Bergquist, B., Bruland, K., Aguilar-Islas, A., Buck, K., Lohan, M., Smith, G.J., Sohst, B., Coale, K., Gordon, M., Tanner, S., Measures, C., Moffett, J., Barbeau, K., King, A., Bowie, A., Chase, Z., Cullen, J., Laan, P., Landing, W., Mendez, J., Milne, A., Obata, H., Doi, T., Ossiander, L., Sarthou, G., Sedwick, P., Van den Berg, S., Laglera-Baquer, L., Wu, J.-f., Cai, Y., 2007. Developing standards for dissolved iron in seawater. EOS Trans. Am. Geophys. Union 88, 131–132.
- Lerner, P., Marchal, O., Lam, P.J., Anderson, R.F., Buesseler, K., Charette, M.A., Edwards, R.L., Hayes, C.T., Huang, K.-F., Lu, Y., Robinson, L.F., Solow, A., 2016. Testing models of thorium and particle cycling in the ocean using data from station GT11-22 of the U.S. GEOTRACES North Atlantic section. Deep-Sea Res. I Oceanogr. Res. Pap. 113 (Supplement C), 57–79. http://dx.doi.org/10.1016/j.dsr.2016.03.008.
- Mawji, E., Schlitzer, R., Dodas, E.M., et al., 2015. The GEOTRACES intermediate data product 2014. Mar. Chem. 177 (Part 1), 1–8. http://dx.doi.org/10.1016/j.marchem. 2015.04.005.
- Morrison, J., 2014. Digital atlas shows ocean' iron levels. Nature. http://dx.doi.org/10. 1038/nature.2014.14774.
- Morton, P.L., Landing, W.M., Hsu, S.-C., Milne, A., Aguilar-Islas, A.M., Baker, A.R., Bowie, A.R., Buck, C.S., Gao, Y., Gichuki, S., Hastings, M.G., Hatta, M., Johansen, A.M., Losno, R., Mead, C., Patey, M.D., Swarr, G., Vandermark, A., Zamora, L.M., 2013. Methods for the sampling and analysis of marine aerosols: results from the 2008 GEOTRACES aerosol intercalibration experiment. Limnol. Oceanogr. Methods 11, 62–78.
- Roy, S., Llewellyn, C.A., Egeland, E.S., Johnsen, G., 2011. Phytoplankton Pigments. Characterization, Chemotaxonomy and Applications in Oceanography. Cambridge University Press (ISBN: 9781107000667).
- Schlitzer, R., 2016. Quantifying He fluxes from the mantle using multi-tracer data assimilation. Philos. Trans. R. Soc. A Math. Phys. Eng. Sci. 374 (2081). http://dx.doi. org/10.1098/rsta.2015.0288.
- SCOR Working Group, 2007. GEOTRACES an international study of the global marine biogeochemical cycles of trace elements and their isotopes. Chemie Erde-Geochemistry. http://dx.doi.org/10.1016/j.chemer.2007.02.001.
- Tagliabue, A., Aumont, O., DeAth, R., Dunne, J.P., Dutkiewicz, S., Galbraith, E., Misumi, K., Moore, J.K., Ridgwell, A., Sherman, E., Stock, C., Vichi, M., Völker, C., Yool, A., 2016. How well do global ocean biogeochemistry models simulate dissolved iron distributions? Glob. Biogeochem. Cycles 30 (2), 149–174. http://dx.doi.org/10. 1002/2015GB005289.