

1 Extreme oxygen isotope zoning in garnet and zircon from a
2 metachert block in mélangé reveals metasomatism at the peak of
3 subduction metamorphism

4 **F. Zeb Page¹, Emilia M. Cameron^{1,2}, Clara Margaret Flood¹, Jeffrey W. Dobbins¹, Michael**
5 **J. Spicuzza², Kouki Kitajima², Ariel Strickland², Takayuki Ushikubo^{2***}, Christopher G.**
6 **Mattinson³, and John W. Valley²**

7 *¹ Geology Department, Oberlin College, Oberlin, Ohio 44074, USA*

8 *² WiscSIMS, Department of Geoscience, University of Wisconsin -Madison, Wisconsin 53706,*
9 *USA*

10 *³Department of Geological Sciences, Central Washington University, Ellensburg, Washington*
11 *98926*

12 **** Present Address: Kochi Institute for Core Sample Research, JAMSTEC 200 Monobe-otsu,*
13 *Nankoku, Kochi 783-8502 Japan*

14
15
16 Post-Review revision

17 April 17, 2019

ABSTRACT

A tectonic block of garnet quartzite in the amphibolite-facies mélange of the Catalina Schist (Santa Catalina Island, California, USA) records the metasomatic pre-treatment of high- $\delta^{18}\text{O}$ sediments as they enter the subduction zone. The block is primarily quartz, but contains two generations of garnet that record extreme oxygen isotope disequilibrium and inverse fractionations between garnet cores and matrix quartz. Rare mm-scale garnet crystals record prograde cation zoning patterns, whereas more abundant $\sim 200\text{-}\mu\text{m}$ diameter crystals have the same composition as rims on the larger garnets. Garnets of both generations have high $\delta^{18}\text{O}$ cores ($20.8\text{--}26.3\text{‰}$, VSMOW) that require an unusually high- $\delta^{18}\text{O}$ protolith, and lower- $\delta^{18}\text{O}$, less variable rims ($10.0\text{--}11.2\text{‰}$). Matrix quartz values are homogeneous (13.6‰). Zircon crystals contain detrital cores ($\delta^{18}\text{O} = 4.7\text{--}8.5\text{‰}$, $124.6 \pm 1.4\text{--}2.9$ Ma), with characteristic igneous trace-element composition likely sourced from arc-volcanics, surrounded by zircon with metamorphic age (115.1 ± 2.5 Ma) and trace-element compositions that suggest growth in the presence of garnet. Metamorphic zircon decreases in $\delta^{18}\text{O}$ from near-core (24.1‰) to rim (12.4‰), in equilibrium with zoned garnets.

Collectively, the data document the subduction of a mixed high- $\delta^{18}\text{O}$ siliceous ooze/volcanic ash protolith reaching temperatures of $550\text{--}625^\circ\text{C}$ prior to the nucleation of small garnets without influence from external fluids. Metasomatism is recorded in rims of both garnet and zircon populations as large volumes of broadly homogeneous subduction fluids stripped matrix quartz of its extremely high oxygen isotope signature. Zoned garnet and zircon in high- $\delta^{18}\text{O}$ subducted sediments offer a detailed window into subduction fluids.

INTRODUCTION

The nature and timing of mass transfer between the subducting plate and the sub-arc mantle is critical to our understanding of crustal formation at convergent margins and its geochemical signatures. Chemical and mechanical hybridization within subduction mélange plays an important role in these processes (e.g., Bebout and Penniston-Dorland, 2016), giving rise to models suggesting that partial melting of diapirs of hybridized mélange rocks are responsible for the classic trace element signature of arc rocks (Marschall and Schumacher, 2012) and the diversity of magma series found at convergent margins (Cruz-Uribe et al., 2018). Adding to these complications is the recent discovery that some sediments have entered the mantle and melted without mixing or hybridization, preserving extreme oxygen isotope signatures of surface weathering in their neoformed igneous zircon (Spencer et al., 2017). If subducted sediment can regularly carry its characteristically enriched oxygen isotope signature ($\delta^{18}\text{O} = \sim 7 - 42\text{‰}$, VSMOW [Vienna standard mean ocean water]; Kolodny and Epstein, 1976; Eiler et al., 2001; Payne et al., 2015) into the mantle ($\delta^{18}\text{O}_{\text{OI}} = 5.1\text{‰}$, Eiler et al., 2000), it is surprising that oxygen isotope variability within the subarc mantle is so subtle and challenging to measure (Eiler et al., 1998). A solution to this discrepancy may be found in the fluid metasomatism of subducted sediments.

The first and perhaps most dramatic illustrations of a high degree of fluid flow within subduction mélange were studies of the oxygen isotope ratios of quartz and carbonate in veins within the Catalina Schist subduction complex (California, USA) suggesting km-scale oxygen isotope homogenization driven by large fluid fluxes (Bebout and Barton, 1989; Bebout, 1991). Over the last quarter century, the Catalina Schist has served as a laboratory for the study of subduction mélange, with numerous studies detailing fluid metasomatism and mechanical

65 mixing processes in the subduction channel by means of stable isotopes (e.g., Bebout, 1991;
66 Penniston-Dorland et al., 2012), major and trace elements (e.g., Sorenson and Barton, 1987;
67 Hickmott et al., 1992; Penniston-Dorland et al., 2014) and radiogenic isotopes (King et al.,
68 2006).

69 The in-situ analysis of oxygen isotopes in garnet is a powerful tool with which to
70 decipher complex or extremely subtle fluid histories and tie them to the metamorphic record. In
71 rocks that have experienced significant metasomatism, the extremely slow intragranular
72 diffusion of oxygen in garnet allows it to preserve a robust geochemical record through all but
73 the hottest and longest of metamorphic events (Vielzeuf et al., 2005). Oxygen isotope variability
74 in garnets from eclogite has illustrated signals of infiltration by mantle (Russell et al., 2013) and
75 supracrustal (e.g., Page et al., 2014; Martin et al., 2014; Rubatto and Angiboust, 2015) fluids that
76 were previously undetectable using bulk methods.

77 Chert and siliceous schist are high- $\delta^{18}\text{O}$ lithologies (Eiler et al., 2001) that are found
78 within the amphibolite-facies Catalina Schist mélange (Platt, 1975). In this contribution, we
79 explore the metasomatism of a high- $\delta^{18}\text{O}$ garnet- and zircon- bearing metachert from a classic
80 subduction mélange, in order to better understand the timing and metamorphic conditions of
81 subduction fluid metasomatism, and to gain a more complete picture of how fluids mitigate the
82 influence of high- $\delta^{18}\text{O}$ subduction inputs.

84 **CATALINA GARNET QUARTZITE**

85 Although much less numerous than the better-studied garnet-hornblende lithology, tectonic
86 blocks of garnet quartzite are also found within the amphibolite-facies metasedimentary mélange
87 of the Catalina Schist (Santa Catalina Island, California, USA), as well as in more coherent, fault-

bounded sheets (Platt, 1975; Bebout, 1991). In this study, we report on one exceptional sample of garnet quartzite collected from a meter-scale tectonic block hosted in a shale-matrix *mélange* from Upper Cottonwood Canyon (33°23'46.20"N, 118°24'52.80"W, Fig. 1A). The quartzite is composed primarily of quartz (93%), garnet (6%), and chlorite (<0.5%), with trace rutile, apatite, amphibole, and zircon (Fig. 1B). Garnet is present in two populations: copious fine-grained (<200 μ m-diameter) crystals dispersed throughout the sample and a smaller number of larger garnets (1-3mm-diameter, Fig. 1B). The larger crystals have abundant inclusions, which are primarily quartz and apatite. X-ray mapping and major element traverses show that the larger garnets display classic prograde cation-zoning profiles with decreasing Mn and increasing Mg# from core to rim, and with rim compositions similar to smaller, more homogenous (in cations) garnets in the same rock (Figs. 2A, B).

Oxygen Isotopes of Quartz and Garnet

Ion microprobe analysis of garnets (Page et al., 2010, see GSA Data Repository for full data, Tables DR-1, DR-2, and methods) shows extreme oxygen isotope zoning; values of $\delta^{18}\text{O}$ are 20.8-26.3‰ in garnet cores and 10.0-11.2‰ in garnet rims (Fig. 2A). Both large and small garnets in this sample show a similar range in $\delta^{18}\text{O}$, despite the difference in cation zoning and crystal size. Zoning in oxygen isotopes is sharp, with up to a 7‰ drop in $\delta^{18}\text{O}$ over a few micrometers, whereas cation zonation is much more gradual with slightly increased Ca and Mg in the rims of larger garnets (Fig. 2). Smaller garnets are nearly homogeneous with a slight increase in Mg# from core to rim. Matrix quartz has no systematic zoning in cathodoluminescence imaging (CL) and is homogeneous in $\delta^{18}\text{O}$ with ion microprobe analyses (13.5‰) identical (within uncertainty) to bulk (~2mg) analysis by laser fluorination (13.6‰).

Garnet-core and quartz pairs yield reversed fractionations ($\delta^{18}\text{O}_{\text{grt}} > \delta^{18}\text{O}_{\text{qz}}$), indicating profound disequilibrium. Eleven analyses of quartz inclusions in large garnet cores yield $\delta^{18}\text{O} = 13.8\text{--}16.2\text{‰}$, higher than matrix quartz, but not in equilibrium with host garnet. Inclusions were generally $>50\mu\text{m}$, and commonly along cracks and so are unlikely to preserve their original values.

Oxygen Isotopes in Zircon

Zircons were separated from the sample and mounted in epoxy for analysis (see GSA Data Repository). CL imaging (Fig. 3A) reveal oscillatory-zoned cores, often as fragments of crystals, containing inclusions of quartz, K-feldspar, and biotite. These detrital cores are surrounded by annuli of variable intensity, somewhat mottled zircon, containing inclusions of quartz, biotite, sphene, and rutile. Outside of this mottled zone, zircons typically have darker-intensity-CL oscillatory-zoned rims, with rare crystals containing a brighter outer rim with faint oscillatory zoning.

Zircons were analyzed for their oxygen isotope ratios by ion microprobe using both a $\sim 15\text{-}\mu\text{m}$ and a sub- $1\text{-}\mu\text{m}$ diameter beam (Tables DR-4, -5). Highly precise and accurate oxygen isotope ratios from the larger analysis pits are correlated with CL zonation and inclusion population. Zircon cores ($n=7$) have $\delta^{18}\text{O}$ from 4.7 to 8.4‰ (Figs. 3A, 3B). Zircon with mottled CL immediately outside of detrital cores ($n=17$) has an extremely high $\delta^{18}\text{O}$ (Fig. 3A, 3B) of $22.6 \pm 3.3\text{‰}$ (2 SD, sample) if one anomalously low analysis is discounted. Intermediate-intensity oscillatory-zoned rims ($n=20$) have lower $\delta^{18}\text{O}$ values ($17.3 \pm 3.9\text{‰}$), and rare bright outer rims have lower-still $\delta^{18}\text{O}$ values ($12.9 \pm 3.3\text{‰}$). To further determine if there is a systematic zoning pattern in zircon like that found in garnet, 29 sub- $1\text{-}\mu\text{m}$ analyses (following

the method of Page et al., 2007) were made in traverses across a single zircon (Fig. 3A). These high-spatial resolution (but less precise, ± 0.9 - 1.7% , 2S.D.) analyses confirm the presence of a low $\delta^{18}\text{O}$ core ($6.3 \pm 1.1\%$, 2S.D., $n=6$), surrounded by an extremely high- $\delta^{18}\text{O}$ mottled CL region ($22.6 \pm 2.4\%$, $n=15$), indistinguishable within the uncertainty of the sub- $1\text{-}\mu\text{m}$ data from the $15\text{-}\mu\text{m}$ -diameter analyses of the same zones. An outer, darker oscillatory-zoned rim has $\delta^{18}\text{O}$ of $17.0 \pm 2.5\%$, $n=8$). The zircon chosen for this analysis does not have an outermost, lighter rim.

PRESSURE, TEMPERATURE, AND TIME HISTORY

The limited mineralogy of this sample coupled with its metasomatic history and zoned minerals, makes thermobarometry challenging. However, an equilibrium assemblage diagram calculated using an estimate of the bulk composition and the computer package Perple_X (Connolly, 2009; Fig. DR-1) yields reasonable results. The observed assemblage (qz+grt+ru $\pm$ chl) is predicted to form at pressures greater than 8kbar and temperatures greater than 550°C . The core to rim increase of Mg# observed in the large garnets is consistent with growth during increasing temperatures in the presence of chlorite, and is predicted by the model to have taken place at ~ 550 - 650°C , at pressures of greater than 11kbar, consistent with existing pressure and temperature estimates of amphibolite blocks in the same *mélange* and [Zr]-in-rutile thermometry from this same sample (Sorenson and Barton, 1987; Hartley et al., 2016; Penniston-Dorland et al., 2018). The closeness between the conditions predicted by the model and existing thermobarometry from Catalina suggests that the metasomatism of this block did not involve substantial change in cation composition. Regardless of the precise conditions of metamorphism, the concomitant decrease in $\delta^{18}\text{O}$ with increasing Mg# in garnet requires metasomatism as the sample increased in temperature within the subduction environment.

Zircons were additionally analyzed by SHRIMP-RG for U-Pb isotopes and select trace elements (GSA Data Repository). Detrital zircon cores have more elevated Th/U ratios (0.36-0.89), and are older than rims; 8 of 9 analyses yield a coherent ^{204}Pb -corrected $^{206}\text{Pb}^*/^{238}\text{U}$ age of $124.6 \pm 1.4/-2.9$ Ma (Fig. 3C). Th/U ratios of rims are lower (0.02-0.13) and yield an age of 115.1 ± 2.4 Ma consistent with an igneous origin for zircon cores, and a metamorphic one for the rims (Fig. 3C). Zircon rims also have smaller Eu-anomalies (Eu/Eu* close to 1) and flatter HREE patterns, consistent with a metamorphic origin in a garnet-present, plagioclase-absent high-pressure environment (Fig. 3D).

DISCUSSION

Taken together, the P-T-t-fluid data preserved in garnet and zircon from this sample provide a detailed record of metasomatic events within the subduction channel. A mixed-lithology protolith containing both extremely high- $\delta^{18}\text{O}$ siliceous material intermixed with intermediate/mafic igneous material including detrital igneous zircon grains was subducted in the Catalina trench. The most plausible interpretation is that the protolith was a mixture of chert or siliceous ooze mixed with ~124 Ma arc volcanoclastic material. The relative purity of the quartzite and the narrow range of zircon core ages seems to preclude weathering of plutonic source material as an origin for the inherited cores. This mixed sediment was subducted and metamorphosed initially as a closed system, with larger, prograde garnet cores having high and unchanging $\delta^{18}\text{O}$ values. The extreme oxygen isotope ratio of this sample (quartz in equilibrium with garnet cores at 550°C would have been greater than 30‰, Valley et al., 2003) makes it highly sensitive to infiltration from external fluids with lower $\delta^{18}\text{O}$. A second generation of garnets nucleated near the peak of metamorphism, but their growth was not initiated by an

external fluid, as core $\delta^{18}\text{O}$ compositions are identical to larger garnets. As metamorphic temperatures reached their peak, an external fluid permeated the sample, perhaps due to introduction of the block into the subduction mélange, shifting matrix quartz $\delta^{18}\text{O}$ from $\sim 30\text{‰}$ to 13.6‰ . Slow rates of intragranular diffusion preserve a record of the original high- $\delta^{18}\text{O}$ composition of garnet and zircon, and their continued growth documents decreasing $\delta^{18}\text{O}$ from $\sim 24\text{‰}$ to $\sim 17\text{‰}$ to $\sim 11\text{‰}$ VSMOW, possibly in two discrete pulses. Fractionation between matrix quartz and garnet rim compositions yield temperatures of $\sim 600\text{--}750^\circ\text{C}$ (Valley et al., 2003), consistent with estimates of peak metamorphic temperatures for the block and the region. Likewise, garnet cation composition records increasing temperature (pressure is not well constrained) during metasomatism. Perhaps upwelling within the subduction channel stopped quartz recrystallization and garnet growth simultaneously, effectively ending the record preserved in this sample.

The limited range of $\delta^{18}\text{O}$ in quartz and calcite veins within the Catalina Schist first reported by Bebout and Barton (1989) suggests that the entire package of subduction rocks on Catalina Island interacted with a remarkably homogeneous supracrustal fluid reservoir derived from metamorphic dehydration of minerals deeper along the subducting slab with an oxygen isotope composition of $13 \pm 1.0\text{‰}$ VSMOW. The quartz $\delta^{18}\text{O}$ value for the block in mélange in this study (13.6‰ VSMOW) yields a calculated water $\delta^{18}\text{O}$ value of 12.3‰ VSMOW (650°C , Friedman and O'Neil, 1977) in close agreement with the range reported by Bebout and Barton.

Although high- $\delta^{18}\text{O}$ sediments make up a volumetrically small portion of subducted material, the extreme contrast between their isotope ratios and those of the mantle make them likely candidates for introducing fine-scale isotope anomalies in the sub-arc mantle. Indeed, the recent discovery of high- $\delta^{18}\text{O}$ S-type granite within supra-subduction-zone mantle as well as

this contribution show that this can happen (Spencer et al., 2017). The sample documented in this study is an example of the most extreme contrast in $\delta^{18}\text{O}$ that one might expect to be subducted, with an estimated protolith $\delta^{18}\text{O}$ of 30‰. However, the metasomatic processes documented by garnet and zircon zonation in this metachert from Catalina show that subduction fluids can all but wipe out extremely high- $\delta^{18}\text{O}$ inputs to subduction zones. Given the modest modal proportion of garnet (7%) with respect to quartz (93%) in this sample, and assuming $\delta^{18}\text{O}$ values of 24‰ for garnet and 14‰ for quartz, the whole rock $\delta^{18}\text{O}$ of this rock must be less than 15‰, a value that can also be found in the much more abundant subducted metabasalts with protoliths enriched in ^{18}O by low-temperature interaction with sea water (Eiler, 2001). Subduction fluids play a vital role in the generation of arc magmatism and continental growth, but it also seems that they play an important role in buffering the $\delta^{18}\text{O}$ of rocks that are recycled into the mantle by subduction, with only strongly refractory (and volumetrically minor) phases such as zircon and garnet able to carry extreme oxygen isotope ratios into the mantle.

ACKNOWLEDGMENTS

We thank Eric Essene, who provided mentorship, assistance in the field, and partial funding for SIMS analysis, S. Penniston-Dorland, E. Walsh and the 2012 Keck Catalina Project (NSF-REU-1062720) students for assistance with fieldwork on Catalina Island, and the Catalina Island Conservancy for access and field support. Comments from Christopher Spencer, two anonymous reviewers and editor Chris Clarke improved this manuscript and are gratefully acknowledged. Assistance with EPMA and laser fluorination analyses were provided by G. Moore (University of Michigan) and Mike Spicuzza (University of Wisconsin). The WiscSIMS ion microprobe laboratory is supported by National Science Foundation (EAR-1355590,

226 1658823) and the University of Wisconsin–Madison. FZP, EMC, CMF and JWD gratefully
227 acknowledge financial support from Oberlin College and NSF (EAR-1249778). JWV is
228 supported by NSF (EAR-1524336).

229 REFERENCES CITED

- 230 Bebout, G. E., 1991, Field-based evidence for devolatilization in subduction zones: implications
231 for arc magmatism: *Science*, v. 251, p. 413-416.
- 232 Bebout, G. E., and Barton, M. D., 1989, Fluid flow and metasomatism in a subduction zone
233 hydrothermal system: Catalina Schist terrane, California: *Geology*, v. 17, p. 976-980.
- 234 Bebout, G. E., and Penniston-Dorland, S. C., 2016, Fluid and mass transfer at subduction
235 interfaces—The field metamorphic record: *Lithos*, v. 240-243, p. 228–258, doi:
236 10.1016/j.lithos.2015.10.007.
- 237 Connolly, J. A. D. (2009) The geodynamic equation of state: what and how: *Geochemistry,*
238 *Geophysics, Geosystems*, v. 10:Q10014, doi:10.1029/2009GC002540.
- 239 Cruz-Uribe, A.M., Marschall, H.R., Gaetani, G.A., and Le Roux, V., 2018, Generation of
240 alkaline magmas in subduction zones by partial melting of mélange diapirs—An
241 experimental study: *Geology*, v. 46, p. 343–346, doi: 10.1130/G39956.1.
- 242 Eiler, J.M., 2001, Oxygen isotope variations of basaltic lavas and upper mantle rocks, *in* Valley,
243 J.W. and Cole, D.R. eds., *Reviews in Mineralogy & Geochemistry Vol. 43: Stable Isotope*
244 *Geochemistry: Mineralogical Society of America*, p. 319–364,
245 <https://doi.org/10.2138/gsrmg.43.1.319>.
- 246 Eiler, J.M., McInnes, B., Valley, J.W., Graham, C.M., and Stolper, E.M., 1998, Oxygen isotope
247 evidence for slab-derived fluids in the sub-arc mantle: *Nature*, v. 393, p. 777–781,
248 <https://doi.org/10.1038/31679>.
- 249 Eiler, J.M., Schiano, P., Kitchen, N., and Stolper, E.M., 2000, Oxygen-isotope evidence for
250 recycled crust in the sources of mid-ocean-ridge basalts: *Nature*, v. 403, p. 530–534,
251 <https://doi.org/10.1038/35000553>.
- 252 Friedman, I. and O’Neil, J. R., 1977, Compilation of stable isotope fractionation factors of
253 geochemical interest: U.S. Geological Survey Professional Paper, v. 440-KK.
- 254 Hartley, E.S., Pereira, I., Moreira, H., Page, F.Z., and Storey, C.D., 2016, Petrology and trace
255 element thermometry of garnet-quartzite from the Catalina Schist, *Geological Society of*
256 *America Abstracts with Programs*, v.48, n.7. Abst. 258-3.
- 257 Hickmott, D., Sorensen, S. S., and Rogers, P., 1992, Metasomatism in a subduction complex;
258 constraints from microanalysis of trace elements in minerals from garnet amphibolite from
259 the Catalina Schist: *Geology*, v. 20, p. 347–350.
- 260 King, R. L., Bebout, G. E., Moriguti, T., and Nakamura, E., 2006, Elemental mixing systematics
261 and Sr-Nd isotope geochemistry of melange formation: Obstacles to identification of fluid
262 sources to arc volcanics: *Earth and Planetary Science Letters*, v. 246, p. 288–304.
- 263 Kolodny, Y., and Epstein, S., 1976, Stable isotope geochemistry of deep sea cherts: *Geochimica*
264 *et Cosmochimica Acta*, v. 40, p. 1195-1209.
- 265 Marschall, H.R., and Schumacher, J.C., 2012, Arc magmas sourced from mélange diapirs in
266 subduction zones: *Nature Geoscience*, v. 5, p. 862–867, doi: 10.1038/ngeo1634.
- 267 Martin, L.A., Rubatto, D., Crépeyron, C., Hermann, J., Putlitz, B., and Vitale Brovarone, A.,
268 2014, Garnet oxygen analysis by SHRIMP-SI: Matrix corrections and application to high-
269 pressure metasomatic rocks from Alpine Corsica: *Chemical Geology*, v. 374-375, p. 25–36,
270 doi: 10.1016/j.chemgeo.2014.02.010.
- 271 Page, F. Z., Ushikubo, T., Kita, N. T., Riciputi, L. R., and Valley, J. W., 2007, High-precision
272 oxygen isotope analysis of picogram samples reveals 2 μm gradients and slow diffusion in
273 zircon: *American Mineralogist*, v. 92, p. 1772-1775.



- Page, F. Z., Kita, N. T., and Valley, J. W., 2010, Ion microprobe analysis of oxygen isotopes in garnets of complex chemistry: *Chemical Geology*, v. 270, p. 9-19.
- Page, F.Z., Essene, E.J., Mukasa, S.B., and Valley, J.W., 2014, A garnet-zircon oxygen isotope record of subduction and exhumation fluids from the Franciscan Complex, California: *Journal of Petrology*, v. 55, p. 103–131.
- Payne, J.L., Hand, M., Pearson, N.J., Barovich, K.M., and McInerney, D.J., 2015, Crustal thickening and clay: Controls on O isotope variation in global magmatism and siliciclastic sedimentary rocks: *Earth and Planetary Science Letters*, v. 412, p. 70–76, doi: 10.1016/j.epsl.2014.12.037.
- Penniston-Dorland, S. C., Bebout, G. E., Pogge von Strandmann, P. A. E., Elliott, T., and Sorensen, S. S., 2012, Lithium and its isotopes as tracers of subduction fluids and metasomatic processes: Evidence from the Catalina Schist, California, USA: *Geochimica et Cosmochimica Acta*, v. 77, p. 530-545.
- Penniston-Dorland, S. C., Gorman, J. K., Bebout, G. E., Piccoli, P. M., and Walker, R. J., 2014, Reaction rind formation in the Catalina Schist: Deciphering a history of mechanical mixing and metasomatic alteration: *Chemical Geology*, v. 384, p. 47–61.
- Penniston-Dorland, S.C., Kohn, M.J., and Piccoli, P.M., 2018, A mélange of subduction temperatures: Evidence from Zr-in-rutile thermometry for strengthening of the subduction interface: *Earth and Planetary Science Letters*, v. 482, p. 525–535, doi: 10.1016/j.epsl.2017.11.005.
- Platt, J., 1975, Metamorphic and deformational processes in the Franciscan Complex, California: some insights from the Catalina Schist terrane: *Bulletin of the Geological Society of America*, v. 86, p. 1337–1347.
- Rubatto, D., and Angiboust, S., 2015, Oxygen isotope record of oceanic and high-pressure metasomatism: a P–T–time–fluid path for the Monviso eclogites (Italy): *Contributions to Mineralogy and Petrology*, v.170:44, p. 1–16, doi: 10.1007/s00410-015-1198-4.
- Russell, A. K., Kitajima, K., Strickland, A., Medaris, Jr, L. G., Schulze, D. J., and Valley, J.W., 2013, Eclogite-facies fluid infiltration: constraints from $\delta^{18}\text{O}$ zoning in garnet: *Contributions to Mineralogy and Petrology*, v. 165, p. 103-116.
- Spencer, C.J., Cavosie, A.J., Raub, T.D., Rollinson, H., Jeon, H., Searle, M.P., Miller, J., McDonald, B.J., Evans, N.J., and EIMF, 2017, Evidence for melting mud in Earth's mantle from extreme oxygen isotope signatures in zircon: *Geology*, v. 45, p. 975-978, doi: 10.1130/G39402.1.
- Valley, J. W., 2003, Oxygen isotopes in zircon: *Reviews in Mineralogy and Geochemistry*, v. 53, p. 343-385.
- Vielzeuf, D., Veschambre, M., and Brunet, F., 2005, Oxygen isotope heterogeneities and diffusion profile in composite metamorphic-magmatic garnets from the Pyrenees: *American Mineralogist*, v. 90, p. 463–472.



FIGURE CAPTIONS

Figure 1. A) Geologic sketch map of Santa Catalina Island, California (after Platt, 1975) showing sample locations. B) Polished thick section of a garnet quartzite showing two garnet sizes. (grt1 – larger, cation-zoned garnet, grt2 – smaller, garnet crystals, homogeneous in cations, qz-quartz, ru – rutile, ap – apatite, chl – chlorite)


Figure 2. A) $\delta^{18}\text{O}$ and cation traverse, rim to rim, of a single $\sim 2.5\text{mm}$ dia. garnet. The core region is generally homogeneous at $\sim 25\text{‰}$ (aquamarine points), transitions to intermediate values (orange) and low, $\sim 10\text{‰}$ rims (purple) over short intervals, although zoning is asymmetric. $\text{Mg}/(\text{Mg}+\text{Fe})$ ($\text{Mg\#}$, solid line) increases continuously core to rim. B) Ternary diagram of garnet cation compositions, mm-scale garnets are shown as solid circles, $\sim 100\mu\text{m}$ -scale garnets shown as open circles. Analysis location (core, intermediate, rim) is also correlated with $\delta^{18}\text{O}$, and indicated by color, as in A. Larger garnets have greater cation zoning than smaller garnets (dashed arrow), and all oxygen isotope zonation takes place at the most pyrope-rich compositions for both sizes (alm - almandine, pyp – pyrope, grs -- grossular, sps – spessartine).

Figure 3. Catalina quartzite zircon chemistry and age A) CL images ($25\mu\text{m}$ scale bars) of three zircons showing different CL domains (see text for details). Analysis of $\delta^{18}\text{O}$ are shown with two spot sizes and labeled with values in VSMOW ($\sim 15\mu\text{m}$, $\pm 0.2\text{--}0.4\text{‰}$ 2S.D., Table DR-5; $< 1\mu\text{m}$, $\pm 2.5\text{‰}$ 2S.D., Table DR-6). B) Histogram of analyses of zircon and garnet with a $15\mu\text{m}$ diameter spot from sample 05C-09 grouped by CL domain and/or location across a traverse, colors as in Figure 2.





A



B

