

Regioselective Metal-free Aza-Heck Reactions of Terminal Alkenes Catalyzed by Phosphine Selenides

Tianyi Zheng, John R. Tabor, Zackary L. Stein, and Forrest E. Michael*


Department of Chemistry, University of Washington, Box 351700, Seattle, Washington 98195-1700

ABSTRACT: Phosphine selenides are introduced as an alternate class of selenium-based catalysts for the aza-Heck reaction of alkenes. Using these catalysts, a range of terminal alkenes react with NFBS to give oxidative amination products. Judicious choice of phosphine ligands gives greater regio- and stereoselectivity than with diphenyl diselenide, enabling the selective formation of *E* terminal enimides in high yields. Isotope labeling experiments and measurements of kinetic isotope effects reveal that the reaction occurs stereospecifically via irreversible anti addition, followed by rate-determining syn elimination.

The aza-Heck¹ and aza-Wacker² reactions are powerful methods for the formation of new C-N bonds from alkenes. The vast majority of these reactions are catalyzed by palladium, proceeding via aminopalladation of the alkene, followed by beta-hydride elimination. Though many intramolecular aza-Wacker and aza-Heck reactions have been reported, only a few intermolecular versions are known.³⁻⁸ One of the major challenges in the development of an intermolecular aza-Wacker or aza-Heck reaction is the need to control the regioselectivity of both the aminopalladation and the beta-hydride elimination steps. The combination of these steps can give up to 4 different isomeric products. This regioselectivity issue is further compounded by the propensity of palladium complexes to promote alkene isomerization reactions, resulting in even more isomeric products.⁹ In addition to reducing the yield of the desired product, these isomers are often difficult to separate from it. Of the four potential products arising from unactivated terminal alkenes, only two have been reported to be accessed in reasonable selectivity. Stahl^{5,6} has reported the selective formation of the internal enamide, and Liu⁷ and White¹⁰ have reported the selective formation of terminal allylamine (Scheme 1, eqs 1 and 2).


Recently, Breder¹¹ and Zhao¹² have shown that diphenyl diselenide can function as a promising alternative catalyst for the aza-Heck reaction of alkenes, presumably via a mechanism conceptually similar to palladium (eqs 3 and 4). The regioselectivity is high for styrenes and allylic alcohols and a few cyclic alkenes. However, control of regioselectivity was again a problem for simple terminal alkenes; complex mixtures of four regio- and stereoisomers were formed under both sets of conditions (see below). To date, understanding of the mechanistic details that control this selectivity is rather limited.

Scheme 1. Intermolecular Aza-Wacker/Aza-Heck Reactions.




Though promising, further development of catalysts based on diphenyl diselenide is hindered by the difficulty of rapidly synthesizing an electronically and sterically diverse library of diselenides. Though Breder proposes that the Se-Se bond is crucial for the catalytic activity, we wondered whether other monoselenides might also be effective catalysts. Along these lines, we hypothesized that the selectivity of this and other reactions catalyzed by selenium compounds could be improved by conceiving of the phenyl group as a ligand for selenium rather than a substituent (Scheme 2). Replacement of this group by common ligands such as phosphines and *N*-heterocyclic carbenes (NHC) would suggest that phosphine selenides and selenoureas might also function as catalysts for these transformations.¹³ An advantage of this strategy is that phosphine selenides and selenoureas are easily prepared in a single step from the corresponding phosphine¹⁴ or NHC,¹⁵ providing convenient access to a large library of sterically and electronically diverse catalysts that could enable more control over reactivity, regioselectivity, and stereoselectivity. Herein, we report that phosphine selenides indeed serve as active catalysts for the aza-Heck reaction of terminal alkenes, and provide higher levels of regio- and stereoselectivity than diphenyl diselenide.

Scheme 2. Phosphine Selenides as Catalysts via Analogy to Arylselenides.



Treatment of terminal alkene **1a** under the conditions previously reported by Breder (Table 1, entry 1) and Zhao (entry 2) gave similar results to those reported by Zhao for 1-hexene, i.e., a mixture of four regio- and stereoisomers was formed in roughly equal amounts. When Ph₃PSe was used in place of diphenyl diselenide, the *E/Z* ratio improved slightly, but notably, the formation of internal regioisomer **4a** was suppressed (entry 3). A screen of other phosphine selenides revealed that a sterically hindered trialkylphosphine slightly improved the isomer ratios (entry 5). Ultimately, tri(*o*-tolyl)phosphine selenide was found to give the highest yield and selectivity for (*E*)-**2a** (entry 6). Unfortunately, the reactivity of the phosphine selenide catalysts under these conditions was substantially lower, giving only ~50% conversion after 65 h. A brief screen of reaction conditions revealed that simply adding 1 equivalent of a benzene-sulfonamide salt restored the reactivity, giving full conversion in 24 h (entry 7). As a final improvement to the procedure, we found that extending the reaction time for an additional 24 h further aminated all of the allyl sulfonamide isomer **3a** to bis(sulfonamide) **5a**¹⁶ (entry 8), without affecting the yield of **2a**. This allowed isolation of the desired terminal vinylamine product **2a** in 74% yield with only minor isomeric products. We also found that the phosphine selenide catalyst could be generated *in situ* by stirring Se powder with the phosphine for 3 hours prior to addition of the remaining reagents (entry 9). Selenium powder in the absence of phosphine did not catalyze the reaction, emphasizing the need for a stabilizing ligand (entry 10).

Table 1. Effects of Selenium Catalyst and Additives




entry	conditions	2a / ^a (<i>E/Z</i>)	3a / ^a	4a / ^a
1	5% PhSeSePh, THF (Breder, ref 11)	47 (1.8:1)	17	13
2	5% PhSeSePh, NaF, pyr, THF (Zhao, ref 12)	41 (1.9:1)	14	13
3	10% Ph ₃ PSe	18 (2.3:1)	7	tr
4	10% IPrSe	tr	tr	tr
5	10% Cy ₃ PSe	16 (2.6:1)	6	tr
6	10% (<i>o</i> -Tol) ₃ PSe	31 (4.1:1)	9	tr
7	10% (<i>o</i> -Tol) ₃ PSe, 1.0 equiv Et ₃ NHN(SO ₂ Ph) ₂ , 48 h	74 (5.7:1)	12	4
8	10% (<i>o</i> -Tol) ₃ PSe, 1.0 equiv Et ₃ NHN(SO ₂ Ph) ₂ , 48 h	74 (5.7:1)	tr	4

9	15% Se, 10% (<i>o</i> -Tol) ₃ P, 1.0 equiv, Et ₃ NHN(SO ₂ Ph) ₂ , 48 h	70 (4.8:1)	14	5
10	10% Se, 1.0 equiv Et ₃ NHN(SO ₂ Ph) ₂ , 48 h	0	0	0

^aNMR yields using 1,3-dinitrobenzene as internal standard. tr = trace

These optimized conditions were applied to the amination of a variety of terminal alkenes (Scheme 3). Yields and *E/Z* selectivities were generally high. The reaction is compatible with esters, ethers, electron-rich aromatics, silyl ethers, sulfonamides, and nitriles. Notably, aliphatic and aromatic halides could also be successfully used as substrates (**2c**, **2d**, **2j**). The reaction of **1a** was performed under an ambient atmosphere on 4 mmol scale and gave **2a** with no significant loss of yield or selectivity.

Scheme 3. Alkene Scope^a




^aIsolated yields.

To investigate the mechanism of this transformation further, the stereoisomeric deuterium labeled substrates (*E*)-**1t-d** and (*Z*)-**1t-d** were subjected to both our reaction conditions and the diphenyl diselenide-catalyzed conditions reported by Breder (Scheme 4). In all cases, we found that the hydrogen at the terminal carbon atom was substituted stereospecifically, which is consistent with an anti addition, followed by a syn elimination (see Scheme 5). Furthermore, we found that the *E/Z* ratio of the


products was strongly influenced by the geometry of the starting alkene, with the *Z* deuterated alkene giving substantially higher *E/Z* product ratios and the *E* deuterated alkene giving substantially lower *E/Z* ratios. By comparing these ratios with that of the closely related undeuterated substrate **1q** (7.6:1), it is apparent that the *E/Z* ratios increase by about a factor of 3 to 4 when using (*Z*)-**1t-d**, while they decrease by a roughly equivalent amount when using (*E*)-**1t-d**.

Scheme 4. Deuterium Labeling Experiment.



We propose the following mechanism based on these results (Scheme 5). The mechanistic experiments in Scheme 4 reveal that the basic mechanistic outline is the same for both diphenyl diselenide and phosphine selenide catalysts. Oxidative addition of NFBS to the phosphine selenide pre-catalyst gives selenium reagent **A**. Similar structures have been isolated and characterized by X-ray crystallography¹⁷ and may be thought of as phosphine complexes of Se(II). Dissociation of benzenesulfonimide and addition to the alkene generates seleniranium ion **B**. Attack of benzenesulfonimide on this species gives the alkylselenium fluoride **C**, which can eliminate in a process that is isoelectronic to the well-known selenoxide elimination to give the product **2**. This anti addition/syn elimination would result in the deuterium labeling patterns observed in the experiments in Scheme 4. Furthermore, the changes in *E/Z* ratios can be rationalized by a primary kinetic isotope effect in the elimination step ($k_H/k_D \sim 3-4$), which slows the formation of (*Z*)-**2** from the *Z* deuterated alkene (see structure **C'** in Scheme 5), and equivalently slows the formation of (*E*)-**2** from the *E* deuterated alkene. This isotope effect is comparable to that measured for the selenoxide elimination ($k_H/k_D = 5.2$).¹⁸


Scheme 5. Proposed Catalytic Cycle.



To learn more about the individual steps of the catalytic cycle, we measured the kinetic isotope effect (KIE) of deuterium substitution at the terminal position in two different ways (Scheme 6). First, we performed a competition experiment between unlabeled compound **1t** and dideuterated **1t-d₂**. Under

these conditions, we found a small inverse KIE ($k_H/k_D \sim 0.95$, see Supporting Information for details). We also monitored the consumption of starting alkene in separate reactions of compound **1t** and compound **1t-d₂**. Under these conditions, the deuterated compound was consumed significantly more slowly, indicating a primary KIE ($k_H/k_D \sim 2$).¹⁹ The inverse isotope effect in the competition experiment is typical for alkene addition reactions,²⁰ and indicates that the product-determining step is the addition to the alkene, i.e., the initial addition of selenium to the alkene is irreversible. The primary isotope effect observed on the overall rate, however, indicates that the elimination of fluoroselenide **C** is the overall rate-determining step.

Scheme 6. Kinetic Isotope Effects.



In summary, we have demonstrated that phosphine selenides may be used as a new class of catalysts in place of diaryldiselenide catalysts for the functionalization of alkenes. Proper choice of the phosphine allows enhanced regio- and stereoselectivity in the aza-Heck reaction of terminal unactivated alkenes with *N*-fluorobenzenesulfonimide. Mechanistic studies have revealed that the mechanism proceeds by an irreversible stereospecific anti addition, followed by a rate-determining syn elimination and that the syn elimination occurs with substantial C-H bond-breaking in the transition state.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Experimental details, characterization data, and copies of ¹H and ¹³C NMR spectra (PDF).

AUTHOR INFORMATION

Corresponding Author

* michael@chem.washington.edu

Notes

The authors declare no competing financial interests.

ACKNOWLEDGMENT

We thank the University of Washington and the National Science Foundation (CHE-1764450) for funding.

REFERENCES

- (1) Aza-Heck review: Race, N. J.; Hazelden, I. R.; Faulkner, A.; Bower, J. F. Recent Developments in the Use of Aza-Heck Cyclizations for the Synthesis of Chiral N-Heterocycles *Chem. Sci.* **2017**, 8, 5248-5260.

(2) Aza-Wacker review: McDonald, R. I.; Liu, G.; Stahl, S. S. Palladium(II)-Catalyzed Alkene Functionalization via Nucleopalladation: Stereochemical Pathways and Enantioselective Catalytic Applications. *Chem. Rev.* **2011**, *111*, 2981-3019.

(3) Hosokawa, T.; Takano, M.; Kuroki, Y.; Murahashi, S.-I. Palladium(II)-Catalyzed Amidation of Alkenes. *Tetrahedron Lett.* **1992**, *33*, 6643-6646.

(4) Timokhin, V. I.; Anastasi, N. R.; Stahl, S. S. Dioxygen-Coupled Oxidative Amination of Styrene. *J. Am. Chem. Soc.* **2003**, *125*, 12996-1997.

(5) Brice, J. L.; Harang, J. E.; Timokhin, V. I.; Anastasi, N. R.; Stahl, S. S. Aerobic Oxidative Amination of Unactivated Alkenes Catalyzed by Palladium. *J. Am. Chem. Soc.* **2005**, *127*, 2868-2869.

(6) Rogers, M. M.; Kotov, V.; Chatwichien, J.; Stahl, S. S. Palladium-Catalyzed Oxidative Amination of Alkenes: Improved Catalyst Reoxidation Enables the Use of Alkene as the Limiting Reagent. *Org. Lett.* **2007**, *9*, 4331-4334.

(7) Liu, G.; Yin, G.; Wu, L. Palladium-Catalyzed Intermolecular Aerobic Oxidative Amination of Terminal Alkenes: Efficient Synthesis of Linear Allylamine Derivatives. *Angew. Chem. Int. Ed.* **2008**, *47*, 4733-4736.

(8) Copper can promote a similar reaction for a limited selection of styrenes: Lisosz, T. W.; Chemler, S. R. Copper-Catalyzed Oxidative Amination and Allylic Amination of Alkenes. *Chem. Eur. J.* **2013**, *19*, 12771-12777.

(9) The regioselectivity of the elimination step can be controlled by choosing substrates that will isomerize to a single thermodynamically preferred product, see: Kohler, D. G.; Gockel, S. N.; Kennemur, J. L.; Waller, P. J.; Hull, K. L. Palladium-catalysed Anti-Markovnikov Selective Oxidative Amination. *Nature Chemistry* **2018**, *10*, 333-340.

(10) Though it does not proceed via an aza-Wacker mechanism, Pd-catalyzed allylic amination via a π -allyl intermediate gives the same product, see: Pattillo, C. C.; Strambeanu, I. I.; Calleja, P.; Vermeulen, N. A.; Mizuno, T.; White, M. C. Aerobic Linear Allylic C-H Amination: Overcoming Benzoquinone Inhibition. *J. Am. Chem. Soc.* **2016**, *138*, 1265-1272 and references therein.

(11) Trenner, J.; Depken, C.; Weber, T.; Breder, A. Direct Oxidative Allylic and Vinylic Amination of Alkenes through Selenium Catalysis. *Angew. Chem. Int. Ed.* **2013**, *52*, 8952-8956. Orgies, S.; Breder, A. Selenium-Catalyzed Oxidative C(sp²)-H Amination of Alkenes Exemplified in the Expedient Synthesis of (Aza-)Indoles. *Org. Lett.* **2015**, *17*, 2748-2751.


(12) Deng, Z.; Wei, J.; Liao, L.; Huang, H.; Zhao, X. Organoselenium-Catalyzed, Hydroxy-Controlled Regio- and Stereoselective Amination of Terminal Alkenes: Efficient Synthesis of 3-Amino Allylic Alcohols. *Org. Lett.* **2015**, *17*, 1834-1837. Zhang, X.; Guo, R.; Zhao, X. Organoselenium-catalyzed Synthesis of Indoles through Intramolecular C-H Amination. *Org. Chem. Front.* **2015**, *2*, 1334-1337.

(13) Phosphine selenides have previously been employed as catalysts for the enantioselective transfer of sulfonyl and selenenyl groups to alkenes: (a) Denmark, S. E.; Hartmann, E.; Kornfilt, D. J. P.; Wang, H. Mechanistic, Crystallographic, and Computational Studies on the

Catalytic, Enantioselective Sulfonylfunctionalization of Alkenes. *Nature Chemistry* **2014**, *6*, 1056-1064. (b) Denmark, S. E.; Chi, H. M. Lewis Base Catalyzed, Enantioselective, Intramolecular Sulfonylaminations of Olefins. *J. Am. Chem. Soc.* **2014**, *136*, 8915-8918. (c) Denmark, S. E.; Jaunet, A. Catalytic, Enantioselective, Intramolecular Carbosulfonylation of Olefins. *J. Am. Chem. Soc.* **2013**, *135*, 6419-6422. (d) Denmark, S. E.; Kalyani, D.; Collins, W. R. Preparative and Mechanistic Studies toward the Rational Development of Catalytic, Enantioselective Selenoetherification Reactions. *J. Am. Chem. Soc.* **2010**, *132*, 15752-15765. (e) Denmark, S. E.; Kornfilt, D. J. P.; Vogler, T. Catalytic Asymmetric Thiofunctionalization of Unactivated Alkenes. *J. Am. Chem. Soc.* **2011**, *133*, 15308-15311. (f) Denmark, S. E.; Collins, W. R. Lewis Base Activation of Lewis Acids: Development of a Lewis Base Catalyzed Selenolactonization. *Org. Lett.* **2007**, *9*, 3801-3804.

(14) Schmidpeter, A.; Brecht, H. Verschiebung der 31P-Resonanz von Phosphinchalkogoniden in Substitutionsreihen und bei Chalkogenmethylierung. *Z. Naturforschung*, **1969**, *24b*, 179-192.

(15) Vummaleti, S. C. C.; Nelson, D. J.; Poater, A.; Gomez-Suarez, A.; Cordes, D. B.; Slawin, A. M. Z.; Nolan, S. P.; Cavallo, L. What Can NMR Spectroscopy of Selenoureas and Phosphinidenes Teach Us about the π -Accepting Abilities of N-Heterocyclic Carbenes? *Chem. Sci.* **2015**, *6*, 1895-1904.



(17) (a) Aragoni, M. C.; Arca, M.; Demartin, F.; Devillanova, F. A.; Garau, A.; Grimaldi, P.; Isaia, F.; Lelj, F.; Lippolis, V.; Verani, G. First ICN Adduct with a Selenium Donor (R = Se): Is It an Ionic [RSeCN]⁺T⁻ or a "T-Shaped" R(I)SeCN Hypervalent Compound? *Eur. J. Inorg. Chem.* **2004**, 2363-2368. (b) Godfrey, S. M.; Jackson, S. L.; McAuliffe, C. A.; Pritchard, R. G. Reaction of Tertiary Phosphine Selenides, R₃PSe (R = Me₂N, Et₂N or C₆H₁₁), with Dibromine. The First Reported Examples of 1 : 1 Addition. *J. Chem. Soc., Dalton Trans.* **1998**, 4201-4204.

(18) Kwart, L. D.; Horgan, A. G.; Kwart, H. Structure of the Reaction Barrier in the Selenoxide-Mediated Formation of Olefins. *J. Am. Chem. Soc.* **1981**, *103*, 1232-1234.

(19) The isotope effect measured by the competition experiment in Scheme 5 is not exactly the same as that measured by labeling in Scheme 3 because neither experiment measures a pure single-step isotope effect for the elimination step. The labeling experiment incorporates a secondary isotope effect in the elimination of a proton, while the competition experiment may incorporate some of the secondary deuterium isotope effect from the addition steps.

(20) (a) Denney, D. B.; Tunkel, N. Deuterium Isotope Effects During Additions to Deuterated and Nondeuterated Olefins. *Chem. Ind.* **1959**, 1383. (b) Merrigan, S. R.; Singleton, D. A. ¹³C and ²H Kinetic Isotope Effects and the Mechanism of Bromination of 1-Pentene under Synthetic Conditions. *Org. Lett.* **1999**, *1*, 327-329.

Insert Table of Contents artwork here

